

Improved representation of soil moisture processes through incorporation of cosmic-ray neutron count measurements in a large-scale hydrologic model

Eshrat Fatima^{1,2}, Rohini Kumar¹, Sabine Attinger^{1,2}, Maren Kaluza¹, Oldrich Rakovec^{1,3}, Corinna Rebmann¹, Rafael Rosolem⁴, Sascha Oswald², Luis Samaniego^{1,2}, Steffen Zacharias⁵, and Martin Schrön⁵

¹Dep. Computational Hydrosystems, UFZ - Helmholtz Centre for Environmental Research GmbH, Leipzig, Germany

²Institute of Environmental Science and Geography, University of Potsdam, Potsdam, Germany

³Faculty of Environmental Sciences, Czech University of Life Sciences Prague, Praha-Suchdol 16500, Czech Republic

⁴Department of Civil Engineering, University of Bristol, Bristol

⁵Dep. Monitoring and Exploration Technologies, UFZ - Helmholtz Centre for Environmental Research GmbH, Leipzig, Germany

Correspondence: Eshrat Fatima (eshrat.fatima@uni-potsdam.de), Rohini Kumar (rohini.kumar@ufz.de), Martin Schrön (martin.schroen@ufz.de)

Abstract. Profound knowledge of soil moisture and its variability plays a crucial role in hydrological modeling to support agricultural management, flood and drought monitoring and forecasting, and groundwater recharge estimation. Cosmic-ray neutron sensing (CRNS) has been recognized as a promising tool for soil moisture monitoring due to its hectare-scale footprint and decimeter-scale measurement depth. Different approaches exist that could be the basis for incorporating CRNS data into distributed hydrologic models, but largely still need to be implemented, thoroughly compared, and tested across different soil and vegetation types. This study establishes a framework to accommodate neutron count measurements and assess the accuracy of soil water content simulated by the mesoscale Hydrological Model (mHM) for the first time. It covers CRNS observations across different vegetation types in Germany ranging from agricultural areas to forest. We include two different approaches to estimate CRNS neutron counts in mHM based on the simulated soil moisture: a method based on the Desilets equation and another one based on the Cosmic-ray Soil Moisture Interaction Code (COSMIC). Within the Desilets approach, we further test two different averaging methods for the vertically layered soil moisture, namely uniform vs. non-uniform weighting scheme depending on the CRNS penetrating depth. We use a Monte Carlo simulation method, specifically the Latin hypercube sampling approach with a large sample size ($S = 100\,000$) to explore and constrain the (behavioral) mHM parameterizations against observed CRNS neutron counts. Overall, the three methods perform well with Kling-Gupta efficiency > 0.8 and percent bias $< 1\%$ across the majority of investigated sites. We find that the non-uniform weighting scheme in the Desilets method and COSMIC method provides the most reliable performance, whereas the more commonly used approach with uniformly weighted average soil moisture overestimates the observed CRNS neutron counts. We then also demonstrate the usefulness of incorporating CRNS measurements into mHM for the simulations of both soil moisture and evapotranspiration and add a broader discussion on the potential and guidelines of incorporating CRNS measurements in large-scale hydrological and land surface models.

1 Introduction

Soil moisture is a key terrestrial climate variable because it controls the mass and energy exchange between the Earth's surface, the groundwater, the vegetation, and the atmosphere. Understanding soil moisture levels with changes in temperature is crucial for enhancing the predictability of climate patterns on inter-seasonal and annual time scales, as highlighted in previous studies (Santanello Jr et al., 2011; Seneviratne et al., 2006). Moreover, soil moisture variability also plays a significant role in a wide range of applications, including flood forecasting, weather forecasting, climate modeling, agricultural management, and groundwater recharge (Van Steenberg and Willems, 2013; Albergel et al., 2010; Jablonowski, 2004; Wahbi et al., 2018; Samaniego et al., 2019; Barbosa et al., 2021). In hydrological modeling, soil moisture is a key variable controlling the partitioning of precipitation into evapotranspiration, infiltration, and runoff (Fuamba et al., 2019; Zhuo et al., 2020). Proper initialization and modeling of soil moisture are crucial for predicting other hydrologic processes (e.g., runoff, evapotranspiration, etc). Nevertheless, uncertainties in input data and model parameters, along with limitations in the representation of subsurface processes, can impede the reliability of soil moisture estimation (Chen et al., 2011). Obtaining accurate soil moisture measurements at a field scale is challenging due to current measurement limitations and subsurface complexity (Dong and Ochsner, 2018). Estimating average soil moisture at a mesoscale ($\approx 1-100$ km) is particularly difficult due to measurement technique limitations in terms of their "footprint" and measurement methods to bridge the scale gap between point-scale and areal average measurements for hydrologic modeling (Chan et al., 2018). One promising approach to infer soil moisture at a field scale is the cosmic-ray neutron sensing (CRNS) technique (Zreda et al., 2008; Desilets et al., 2010; Köhli et al., 2015; Schrön et al., 2017).

Previous studies, such as Barbosa et al. (2021) and Brunetti et al. (2019), have recognized the importance of CRNS over traditional invasive point-scale techniques and have utilized the HYDRUS-1D model to simulate soil moisture at the field scale. HYDRUS-1D offers a valuable framework for modeling soil moisture dynamics and has been particularly addressing the subsurface processes. The studies incorporated a COSMIC operator to simulate neutron count rates of a CRNS measurement (Shuttleworth et al., 2013). They inversely calibrated soil hydraulic parameters by comparing observed and simulated neutron count rates, whereas beforehand this was limited to be done via comparison of depth-averaged soil moisture values (Rivera Villarreyes et al., 2014). The potential utility of using CRNS data to calculate volumetric soil water content (SWC) and improve soil hydraulic parameters within land surface models has also been observed earlier, as highlighted by Rosolem et al. (2014). Furthermore, depth-weighting schemes and hydrogen pools' effects on measurement depth revealed valuable insights. Shallow wetting fronts in sandy soils significantly impact measurement depth Franz et al. (2012). Baroni and Oswald (2015) assessed three weighting techniques, resulting in depths varying from 23 to 28 cm, optimal estimates were achieved using vertically varying weights and considering additional hydrogen pools. In Iwema et al. (2017), a Land Surface Model investigated the impact of reducing scale mismatch between energy flux and soil moisture observations using CRNS data. Patil et al. (2021) employed a distributed Land Surface Model, Data Assimilation Research Testbed (DART) with CRNS time series, and Ensemble Adjustment Kalman Filter to simulate water and energy balance. Both studies focused on analyzing land surface water and energy balance, exploring data assimilation and calibration techniques.

55 The Hydrologiska Bryans Vattenbalansavdelning (HBV) model, as studied by Dimitrova-Petrova et al. (2020), employed CRNS data in a mixed-agricultural landscape to explore water balance on the land surface. While, Beck et al. (2021) used remote sensing products and groundwater level measurements to temporally calibrate the HBV model, emphasizing the challenge of comparing satellite-derived soil moisture with point-scale in-situ measurements. Additionally, Baatz et al. (2017) was the first study that utilized spatially distributed hydrological modeling, integrating CRNS data, FAO and BK50 soil maps, and other
60 soil data in the Community Land Model (CLM). They demonstrated that assimilating CRNS data improved catchment-scale soil water content characterization by updating spatially distributed soil hydraulic parameters. Furthermore, Zhao et al. (2021) assessed the significance of CRNS data in CLM version 3.5, conducting simulations based on 13 CRNS stations over 2017-2018. Despite employing a simplified Richards equation, limitations included the absence of lateral flows and groundwater representation.

65 The mesoscale Hydrological Model (Samaniego et al., 2010b; Kumar et al., 2013b, mHM;) is known for its spatially distributed hydrologic predictions at a large scale incorporating scale-aware regionalized parameterization technique. Therefore, by including a CRNS neutron count framework, the mHM model becomes a useful tool for improving simulated soil water content and furthering our understanding of the water cycle. This is made possible by the availability of observed CRNS data, which opens up new opportunities for research into novel hypotheses, improving model performance, and developing hydro-
70 logical modeling methods. All the mentioned studies either compared the simulated and observed soil moisture products or incorporated the first COSMIC version to compare neutron counts directly. As argued by Shuttleworth et al. (2013), the usage of neutron counts is the favorable way to compare simulations with data, since the CRNS sensor intrinsically averages over soil moisture layers while the measurement depth varies with soil moisture and consequently over time. Hence, the direct usage of CRNS neutron counts avoids the question of which modeled SWC layer the observations should be compared to and at
75 what time scales. The COSMIC method enables a comprehensive representation of the neutron generation process, which is computationally more demanding than using an analytical formulation (e.g., Desilets et al., 2010; Köhli et al., 2021).

In this study, we established a framework to incorporate CRNS data into the mesoscale Hydrological Model (mHM) to compare empirical and physics-based approaches for neutron count estimation to improve soil water content parameters in mHM across different vegetation types in Germany. To do this, we compared modelled with measured neutron counts to
80 infer soil hydraulic parameters. Here, we test three approaches, (i) the direct calculation of neutrons from the equal-averaged SWC profiles based on Desilets et al. (2010), (ii) the same with a weighted-average profile SWC based on Schrön et al. (2017), and (iii) the physics-based model COSMIC by Shuttleworth et al. (2013). We evaluate the simulation of neutron counts at scales of $1.2 \text{ km} \times 1.2 \text{ km}$, comparing the results to observed neutron counts from three different sites including agriculture, deciduous forests, and grasslands. The goal of this study is to investigate the potential of using CRNS probes
85 and measured neutron counts to improve soil moisture predictions through simulations in mHM across different land covers and soil properties and to evaluate the feasibility of incorporating neutron count measurements into the modeling scheme. We employ a (calibration) framework by applying a Monte Carlo experiment to account for parameter uncertainties. We further cross-evaluate our simulations and test the reliability of the CRNS incorporated soil-moisture scheme in mHM for simulating other variables by utilising time series of observed evapotranspiration from an eddy covariance station available. Finally, we

90 discuss and provide guidelines (challenges and limitations) for incorporating CRNS measurements in a large-scale hydrologic model. In summary, the present paper aims to answer the following research questions:

- What is the best approach to simulate CRNS neutron counts in a hydrological model considering the heterogeneity of vertical soil moisture profiles?
- What is the impact of model calibration with CRNS observations on simulated evapotranspiration at Hohes Holz?
- 95 – Is the mHM at approx. 1 km resolution capable of capturing the dynamics of hectare-scale CRNS measurements at different landcover sites in a grid including 2 agriculture sites, 1 forest site, and 1 meadow site?

2 Materials and Methods

2.1 Experimental Site Description

For this study, we select four sites with CRNS sensors, namely *Grosses Bruch*, *Hohes Holz*, *Hordorf*, and *Cunnersdorf* in
100 Northern Germany, as provided already within COSMOS EU (Bogena et al., 2022) with particularly long time series and with different land cover, i.e., agriculture, forest, and meadow (see Tab. 1). The first three sites belong to the TERENO observatory “Harz/Central Germany lowland” (Zacharias et al., 2011) while the fourth site is part of an agricultural research farm operated by the German Weather Service (DWD). The *Grosses Bruch* site is a meadow/grassland that is usually flooded naturally once or twice a year. The meadows have sandy loam fluvisol-gleysol soil, which is 1.5 meters deep and partially covered with a
105 layer of peat (Wollschläger et al., 2017). Meteorological conditions like soil moisture and temperature at various depths are continuously monitored by a wireless soil moisture monitoring network (Schrön, 2017). *Hohes Holz* is a deciduous forest site and the performance of the CRNS sensor there is highly dependent on dynamic effects such as tree canopy water or seasonal fluctuations in wet biomass. Water trapped in leaves and litter can present a particular challenge for CRNS measurements, especially at forest stations (Bogena et al., 2013). Also, Bogena et al. (2022) indicated that the influence of seasonal changes
110 of biomass on the CRNS signal is much less important than the influence of changing soil moisture, even in Hohes Holz, as changes in soil moisture are the much larger source of variation represented by the CRNS measurements. The mean annual air temperature for each site ranges from 10.0 to 10.9 °C and the average yearly precipitation ranges from 458 to 535 mm.

Table 1. Geographical characteristics of study sites: Site Names, Geographic Coordinates, Climatic Data (Annual Precipitation in mm/year, Annual Mean Temperature in °C), and the Periods Covered in Observed and Simulated Datasets.

Site	Latitude [°N]	Longitude [°E]	Altitude [m]	Land Cover	Precipitation [mm/year]	Temperature [°C]	Period
<i>Grosses Bruch</i>	52.02	11.10	80	Pasture, grassland	458	10.1	24/06/2014–31/01/2021
<i>Hohes Holz</i>	52.09	11.22	217	Forest, hilltop	469	10.3	27/08/2014–31/01/2021
<i>Hordorf</i>	51.99	11.17	82	Cropland	463	10.3	29/09/2016–31/01/2021
<i>Cunnersdorf</i>	51.36	12.55	140	Cropland	535	10.9	23/06/2016–31/01/2021




Figure 1. Study area map of Germany, highlighting the four test sites where observed neutron count rates from CRNS are utilized to evaluate the performance of mHM. The figure utilizes OSM basemap layers from (© OpenStreetMap contributors 2021; distributed under the Open Data Commons Open Database License (ODbL) v1.0) OpenStreetMap contributors (2020).

2.2 The mesoscale Hydrological Model (mHM)

mHM is a spatially distributed process-based hydrologic model (www.ufz.de/mhm) representing processes such as canopy
 115 interception, snow accumulation and melting, soil moisture dynamics, infiltration and surface runoff, evaporation, underground storage, and runoff generation, deep infiltration and baseflow, as well as runoff attenuation and flood routing (Samaniego et al., 2010a; Kumar et al., 2013a). The mHM model is flexible for hydrological simulations at different spatial scales due to its novel Multi-scale Parameter Regionalization approach (MPR; Samaniego et al., 2010b); and has demonstrated applicability in

diverse settings (Samaniego et al., 2010a; Kumar et al., 2013a; Rakovec et al., 2016; Samaniego et al., 2017). The MPR's basic
120 concept is to estimate parameters (e.g., porosity) based on soil properties (e.g., sand and clay content) using transfer functions at
a fine spatial resolution (e.g. 100 m) and upscaling them to modelling resolutions (e.g., 1 km). In MPR, transfer functions (e.g.,
pedo-transfer functions to estimate soil parameters) are combined with morphological inputs (e.g., soil texture properties) and
thus lead to model hydrologic parameters (e.g., porosity or hydraulic conductivity of the soil) (Livneh et al., 2015; Zacharias
and Wessolek, 2007). In mHM, the soil moisture horizons/profile can be divided into several horizons, all of which are sensitive
125 to root water uptake and evapotranspiration processes. mHM simulates the daily dynamics of soil moisture at different depths
considering the incoming water (e.g., rainfall plus snow melt for the topmost layer and infiltration from above layers for
other layers) and outgoing ET and ex-filtration fluxes. Further details on mHM code can be found at <https://mhm-ufz.org> and
underlying modelling concepts at Samaniego et al. (2010a); Kumar et al. (2013a).

2.3 Model Set-up

130 The latest version 5.12 of mHM is used in this study (see Samaniego et al., 2023, and <https://github.com/mhm-ufz>). The model
is executed over six years (2014–2020) with a daily time step, and the spatial resolution of the mHM grid cells is fixed at
L1 and L2: $0.01562^\circ \times 0.01562^\circ$ is eq. $\sim 1.2 \text{ km} \times 1.2 \text{ km}$ using the WGS84 Coordinate Systems. Level 1 (L1) describes
the spatial resolution, as which dominant hydrological processes are modelled and Level 2 (L2) describes the resolution of
the meteorological forcing data. L0: $0.001953125^\circ \times 0.001953125^\circ$. Level 0 (L0) describes the subgrid variability of relevant
135 basin characteristics, which includes information on the soil as well as land use, topography, and geology.

Figure 2 shows the flow diagram depicting the basic methodology of our study, which includes the calculation of CRNS
neutron count rates based on daily soil moisture values simulated with mHM. The model boundary conditions such as pre-
cipitation and temperature for the mHM model are acquired from the German Weather Service (DWD) station closest to the
test site. The potential evapotranspiration required by mHM is estimated using the Hargreaves-Samani method (Hargreaves
and Samani, 1985). The model setup and parameterization for the soil moisture module use the scheme optimized by Boeing
et al. (2022). A raster dataset describing the distribution of the soils in the model area and a corresponding lookup table with
the attributes depth, soil texture (sand and clay fraction), and bulk density are required as soil input data and are derived from
national digital soil maps provided by the Federal Institute for Geosciences and Natural Resources (BGR, 2020). The data
set contains physical and chemical properties for soil at different layers and the available at a resolution of 1:250,000 (BUEK
145 200; BGR, 2020). mHM uses three dominant land cover classes (forest, permeable, and impervious) that were retrieved by a
GLOBCOVER database ESA (2009). Furthermore, vegetation characteristics like Leaf Area Index (LAI) and fraction of roots
for different vegetation types are prescribed in the model. The mHM soil domain is divided into three horizons with depths of
0–5 cm, 5–25 cm, and 25–60 cm. The upper two model layers are parameterized using the topsoil layer properties while for
the lower model layer, the subsoil properties are used. More details on the underlying input data for mHM can be obtained
150 from Boeing et al. (2022).

In our study, we utilized three distinct modules of parameters: Snow, Soil Moisture, and Neutrons, with a total of 29 param-
eters employed for the Desilets method and 31 parameters for the COSMIC method. The simulation of soil water content is

processed through these three modules to estimate neutron counts. To comprehensively cover the parameter set ranges, we employed 100 000 iterations. Finally, we selected the top 10 optimized parameter sets based on the objective function, $KGE_{\alpha,\beta}$, for further analysis and evaluation.




Figure 2. Flowchart depicting the methodology employed for calculating CRNS neutron counts through the utilization of the LHS technique for parameterization in mHM. The computation of CRNS neutron count is carried out through three distinct approaches: $N_{Des,U}$, $N_{Des,W}$, and N_{COSMIC} .

2.4 Conversion of soil moisture to neutron count rate

In this study, we compare observed neutron counts from CRNS data with simulated neutron counts estimated from modeled soil moisture with the goal of optimizing the parameterization of soil water content from mHM shown in Fig. 3. By coupling the approaches from Desilets et al. (2010) and Shuttleworth et al. (2013) each directly with the mHM model, we are able to account also for the uncertainty in the model predictions and test their feasibility across four distinct sites. We analyzed the soil water content data at different soil layers (0–5 cm, 5–25 cm, and 25–60 cm) in mHM, as utilized in the study by Boeing et al. (2022). The accuracy of numerical calculations (such as Shuttleworth et al. (2013) set up) would benefit from higher resolved soil profiles, however, our experiments demonstrated that varying soil depths from 3 to 6 layers did not have a substantial impact on the simulated neutron count results in mHM. Our main objective is to optimize the parameterization of soil hydraulic properties in mHM based on the comparison between measurement and modelled neutron counts.




Figure 3. Daily time series of soil water content ($\text{cm}^3\text{cm}^{-3}$) at the *Cunnersdorf* site. The graph shows a comparison between the measured SWC from CRNS data representing an integral over the first decimeters and the simulated data derived from the mHM for three distinct soil depths, at 0–5 cm (green), 5–25 cm (purple), and 25–60 cm (brown).

2.4.1 Desilets based method

In the present study, we utilize the soil moisture information from the mHM model to convert it into neutron counts using the Desilets et al. (2010) empirical-based approach by calculating neutron counts from soil moisture, three constant parameters, which is further improved by adding lattice water and bulk density following the approaches by Dong et al. (2014) and Hawdon et al. (2014), respectively. Theoretically, the N_0 parameter, which represents the neutron count rate level of the particular CRNS probe used for rather dry soil at the local conditions, should be site-specific but does not change over time, as noted

by Franz et al. (2013) and Hawdon et al. (2014). In order to obtain accurate measurements of soil moisture using CRNS data in the mHM model, N_0 has to be estimated through calibration and is crucial as it directly affects the accuracy of the mHM neutron counts results. This calibration parameter is specific to each site environment and reference condition. This parameter primarily depends on site-specific environmental factors and reference conditions. This coefficient is specific to the particular CRNS detector and may be impacted by factors such as terrain (topography) but also local soil, vegetation characteristics, and additional hydrogen pools (e.g., from organic material) at each observation site Schrön et al. (2021). Therefore, calibration of N_0 is necessary for each CRNS data set at a site to ensure realistic model output. Neutrons are sensitive to all kinds of hydrogen in the footprint, hence the variable θ denotes not only soil moisture, θ_{sm} , but is rather assumed to also include lattice water, θ_{lw} , as well as water equivalent from soil organic carbon, θ_{org} , and vegetation biomass, θ_{bio} .

Soil moisture for three vertical mHM soil layers is used to drive both the Desilets method and the COSMIC operator. To improve comparability between measurements and modeling techniques, Schrön et al. (2017) propose the depth-weighted approach. This approach incorporates the contributions of different soil layers by calculating depth-weighted average SWC, θ_{avg} , resulting in a more comprehensive representation of soil moisture dynamics. This gives CRNS neutron count rates, after being corrected for incoming neutron flux, pressure, and air humidity variations, to be

$$N_{Des} = N_{0,Des} \left(\frac{a_0}{(\theta_{avg} + \theta_{lw}) / (\rho_b / \rho_w) + a_2} + a_1 \right) \quad (1)$$

Among the four parameters, three of which are coefficients parameters a_i were determined empirically by (Desilets et al., 2010) who derived $a_0 = 0.0808$, $a_1 = 0.372$, and $a_2 = 0.115$, and are considered as constants for values of $\theta > 0.02 \text{ gg}^{-1}$. The fourth parameter, $N_{0,Des}$ is N_0 when using the Desilet's equation, and here is a free calibration parameter. Whereas the parameter θ_{lw} is the grid average volumetric water content of the equivalent lattice water content of the CRNS area ($\text{cm}^3 \text{ cm}^{-3}$), ρ_b (g cm^{-3}) is the bulk density of the dry soil, usually determined from soil samples, and $\rho_w = 1 \text{ g cm}^{-3}$ is the density of water. Regarding the variables of Soil Organic Carbon (SOC) and biomass, it's important to note that these variables are often not readily available, especially when it comes to biomass data. To address this, the free parameter N_0 is utilized to account for these unknowns. For lattice water, we assume a linear relationship to clay content (Avery et al., 2016):

$$\theta_{lw} = \theta_{lw0} \cdot C + \theta_{lw1} , \quad (2)$$

C denotes the clay fraction in % (Greacen, 1981). The derived quantity lattice water, θ_{lw} , is regionalized based on C and varies between 0.0 and $0.1 \text{ m}^3/\text{m}^3$. In order to obtain the average soil moisture for a layered soil moisture profile within mHM, the following averaging equation is employed:

$$\theta_{avg}(w, \theta) = \frac{\sum_{i=1}^n w_i \theta_i}{\sum_{i=1}^n w_i} \quad (3)$$

where the volumetric soil water content at a specific layer of mHM in a given profile is denoted by θ_i ($\text{m}^3 \text{ m}^{-3}$). The total number of layers in all soil sampling profiles is represented by the variable n , and the weight assigned to layer i is denoted by w_i . In the uniformly weighted approach, all weights equal one:

$$N_{Des,U} = N_{Des}(w_i = 1) \quad \forall i. \quad (4)$$

In the weighted-averaging approach, the weights are determined based on Schrön et al. (2017):

$$205 \quad N_{\text{Des},W} = N_{\text{Des}}(\theta_{\text{avg}}(w, \theta)), \quad (5)$$

$$\text{where } w_i = \int_{z_{i,\min}}^{z_{i,\max}} e^{-2z/D} dz \propto e^{-2z_{i,\min}/D} - e^{-2z_{i,\max}/D}$$

$$\text{and } D = \varrho_b^{-1} \left(p_0 + p_1 (p_2 + e^{-p_3 r}) \frac{p_4 + \theta}{p_5 + \theta} \right). \quad (6)$$

Here, the integral goes through each horizon from $z_{i,\min}$ to $z_{i,\max}$ in 1 mm steps and sums up the weight over the whole layer. z_i is the depth of the given soil moisture layer i , D is the average vertical footprint depth of the neutrons, p_i are numerical
 210 parameters presented in Schrön et al. (2017), and r (m) represents the distance from the sensor. It should be noted that the equation for D is valid for $\varrho_b > 1.0 \text{ g cm}^{-3}$ and soil moisture contents above $\theta > 2 \%$ Kasner et al. (2022). In our model, we set $r = 1$ m which is sufficient to represent the average depth across the footprint radius within the model grid. The soil moisture profile is converted to a single average neutron count per grid cell using Eqs. 1–5.

2.4.2 Cosmic Ray Soil Moisture Interaction Code (COSMIC)

215 The Cosmic Ray Soil Moisture Interaction Code (COSMIC) is an analytical, physics-based model that is well-suited for data assimilation applications. It includes descriptions of the degradation of incoming high-energy neutron flux with soil depth, the production of fast neutrons at each soil depth, and the scattering of resulting fast flux neutrons before reaching the soil surface. These processes have a parametric dependence on soil chemistry and moisture content. The COSMIC method solves this inverse problem by calculating neutron count rate based on soil water profiles, which could then be compared with observed
 220 neutrons without the need to deal with dynamic sensing depths or weightings.

$$N_{\text{COSMIC}} = N_{0,\text{COSMIC}} \sum A_{\text{high}}(z) X_{\text{eff}}(z) A_{\text{fast}}(z), \quad (7)$$

$$\text{where } A_{\text{high}}(z) = e^{-\Lambda_{\text{high}}(z)},$$

$$A_{\text{fast}}(z) = \frac{2}{\pi} \int_0^{\pi/2} e^{-\Lambda_{\text{fast}}(z)} (\cos \varphi)^{-1} d\varphi,$$

$$X_{\text{eff}}(z) = \alpha_{\text{COSMIC}} X_{\text{soil}} + X_{\text{water}}.$$

225 In this model, A_{high} represents the high-energy neutron attenuation, A_{fast} represents the fast neutron attenuation, and X_{eff} represents the production of fast neutrons from high-energy neutrons in the soil-water composite. It takes into account the different mechanisms in both, water and soil, where the soil is typically less effective in producing fast neutrons by a factor of $\alpha_{\text{COSMIC}} \approx 0.24 \text{ (g cm}^3 \text{ g}^{-1})$, depending on bulk density.

$$X_{\text{soil}}(z) = \Delta z \varrho_b, \quad (8)$$

$$230 \quad X_{\text{water}}(z) = \Delta z \varrho_{\text{water}} (\theta_z + \theta_{1w}), \quad (9)$$

The total attenuation lengths of high and fast neutrons in the soil water composite are described using physically motivated length scales L_i .

$$\Lambda_{\text{high}}(z) = \frac{X_{\text{soil}}(z)}{L_1} + \frac{X_{\text{water}}(z)}{L_2}, \quad (10)$$

$$\Lambda_{\text{fast}}(z) = \frac{X_{\text{soil}}(z)}{L_3} + \frac{X_{\text{water}}(z)}{L_4}. \quad (11)$$

235 The COSMIC function considers the attenuation of incoming high-energy neutrons (A_{high}) and their interaction with soil to produce effective neutrons (X_{eff}) in addition to the attenuation of isotropically propagating fast neutrons. The parameter α_{COSMIC} represents the soil's relative efficiency of forming fast neutrons, and length constants L_1 , L_2 , L_3 , and L_4 (in g cm^{-2}) are related to local soil properties. COSMIC uses several time-invariant, site-independent, and site-specific parameters, including $L_1 = 162.0$ (g cm^{-2}), $L_2 = 129.1$ (g cm^{-2}), and $L_4 = 3.16$ (g cm^{-2}), as reported by Shuttleworth et al. (2013), regardless
 240 of location. However, the L_3 (g cm^{-2}) parameter vary with soil bulk density ρ_b which change with depth. The parameter L_3 is correlated with the soil bulk density and according to the model code, L_{30} and L_{31} nomenclature are given as per the model code in mHM (<https://github.com/mhm-ufz>).

$$L_3 = L_{30}\rho_b - L_{31}. \quad (12)$$

The regional formulation of the COSMIC method has been revised to include the θ_w lattice water content as well.

245 Besides the addition of lattice water to the code, the original version of COSMIC has also been numerically optimized to substantially increase the computational performance. This includes the calculation of the geometric integral (Eq. 7) based on lookup tables.

2.5 Constraining of model parameterization

In this study, we employ a model calibration technique to identify the most suitable parameter values for the mHM model.
 250 Specifically, we utilize a total of 29 parameters for the Desilets based method and 31 parameters for the COSMIC method which includes hydrologic processes related to: snow, soil moisture, and neutron counts dynamics. The process of model calibration involves modifying the parameter values of the model to achieve a satisfactory standard for an objective function by comparing the predicted output with the observed data (James, 1982). We use the general concept of the KGE as a weighted combination of the three components (bias, variability, and correlation terms) to evaluate our simulation (Gupta et al., 2009).
 255 We excluded the correlation component from (Eq. 13) as our simulation already exhibited satisfactory correlation due to strong seasonality, we opted not to consider it in our assessment (objective function), as it accounted for 33% of the total weighting in the overall KGE score. Seasonality is an inherent characteristic in the northern hemisphere where precipitation minus evaporation is mostly driven by evapotranspiration. Even if a random parameter is selected correlation will always be higher because the meteorological forcing is the precipitation - evaporation is seasonal. This modified $\text{KGE}_{\alpha\beta}$ (Eq. 14)
 260 only depends on variability (α) and bias (β) and variants of it have been used also in other studies (see, e.g., Martinez and Gupta, 2010; Mai, 2023). We utilize observed neutron count data from CRNS and estimated neutron count data from the mHM

model to calculate various metrics such as the Kling-Gupta efficiency coefficient ($KGE_{\alpha\beta}$), Nash-Sutcliffe efficiency (NSE) by Nash and Sutcliffe (1970), coefficient of determination (R^2) by Kvålseth (1985), and percentage bias (PBIAS) by Gupta et al. (1999). The optimal PBIAS value is 0, with lower values indicating more accurate model simulations. Positive values indicate underestimation by the model, while negative values indicate overestimation. This approach allows us to minimize uncertainty in the simulated neutron count data by comparing it to observed data and determining the optimal parameter values for the mHM model. A summary of the individual parameters and their ranges can be found in Supplementary Table S1 and model performance measures are shown in Table 2.

$$KGE = 1 - \sqrt{(\alpha - 1)^2 + (\beta - 1)^2 + (\rho - 1)^2}, \quad (13)$$

$$KGE_{\alpha\beta} = 1 - \sqrt{(\alpha - 1)^2 + (\beta - 1)^2}, \quad (14)$$

$$\text{with Variability } \alpha = \sigma_{\text{sim}} / \sigma_{\text{obs}}, \quad (15)$$

$$\text{Bias } \beta = \mu_{\text{sim}} / \mu_{\text{obs}}, \quad (16)$$

$$\text{Correlation } \rho = \rho(\text{sim}, \text{obs}),$$

$$NSE = 1 - \frac{\sum_{i=1}^n (y_{\text{sim},i} - y_{\text{obs},i})^2}{\sum_{i=1}^n ((y_{\text{obs},i} - \bar{y}_{\text{obs},i})^2)}, \quad (17)$$

$$R^2 = \left(\frac{\sum_{i=1}^n (y_{\text{obs},i} - \bar{y}_{\text{obs},i})(y_{\text{sim},i} - \bar{y}_{\text{sim},i})}{\sum_{i=1}^n \sqrt{(y_{\text{obs},i} - \bar{y}_{\text{obs},i})^2} \sqrt{(y_{\text{sim},i} - \bar{y}_{\text{sim},i})^2}} \right)^2, \quad (18)$$

$$PBIAS = 100[\%](1 - \beta). \quad (19)$$

Table 2. Performance evaluations for the daily neutron counts simulation with observed CRNS dataset.

Indices	$KGE_{\alpha\beta}$	KGE	NSE	R^2	PBIAS
Range	$-\infty$ to 1	$-\infty$ to 1	$-\infty$ to 1	0 to 1	$-\infty$ to ∞
Optimal Value	1	1	1	1	0
Satisfactory Value	> 0.70	> 0.80	> 0.50	> 0.65	$< \pm 5$

3 Results

3.1 Constraining of the parameter distribution N_0 / Sensitivity Analysis

The sensitivity and uncertainty analysis performed in this study use a Latin Hypercube Sampling (LHS) approach, resulting in parameter distributions that large sample size was chosen to comprehensively explore the parameter sets and capture a wide range of possible parameter combinations in the prior range. The LHS approach creates a random value between the min and max values of the parameter set. Initial parameter ranges and exploratory model runs are set based on literature values (Boeing et al., 2022; Kumar et al., 2013b). Supplementary Table S1 shows the values for the parameters in all 100 000 simulations and the selected 29 parameters for the Desilets method and 31 parameters for the COSMIC method, which include snow, soil

285 moisture, and neutrons modules as behavioral simulations, with the posterior mean of the top 10 best parameters set. For further information and additional details about the calibrated parameters for each site, refer to Supplementary Table S2. Among the calibrated parameters, the N_0 parameters are different in each method since this parameter does not exactly have the same physical meaning in the Desilets and the COSMIC methods.

Fig. 4 displays the x-axis in gray, representing the original parameter range (600–1500) prior distribution for the Desilets method and (100–400) for COSMIC method. Meanwhile, the colored sections in brown, green, and purple indicate the parameter values of the calibration from the posterior distribution taken from the top-performing parameter sets for each study site. The most sensitive parameters during the calibration period are $N_{0,Des}$, $N_{0,COSMIC}$, `rootFractionCoefficient_pervious`, and `rootFractionCoefficient_forest` of land cover classes are employed: class 1 = forest which consisted of permeable areas covered by coniferous, deciduous, and mixed forests; class 2 = impervious cover with land uses like settlements, industrial parks, roads, airport runways, and railway tracks; and class 3 = permeable cover covered by fallow lands, or those surfaces covered by crops, grass, and orchards. The calibration process notably sharpens the Probability Density Function (PDF) of these significant parameters by eliminating some of the uncertainty linked to the variance in the prior probability distributions.

For agricultural sites such as *Cunnersdorf* and *Hordorf*, the $N_{0,Des}$ best estimate parameter results lie between 1000 and 1400 cph. Meanwhile, for *Hohes Holz*, the $N_{0,Des}$ parameter range lies between 800 and 1000 cph, the lowest value for the calibration parameter $N_{0,Des}$ is found between 800 and 900 cph due to the highest wet above biomass in the forest area. Similarly, for *Grosses Bruch* sites, the lowest value for the calibration parameter $N_{0,Des}$ is found between 800 and 900. The prior parameter distribution $N_{0,Des}$ 600 and 1500 cph is the same for all four experiments.

The estimated values of $N_{0,Des}$ and $N_{0,COSMIC}$ obtained in our study are close to the optimal values, indicating that the model has the potential to generate accurate cosmic-ray soil moisture estimates even under dry conditions. In contrast, some hydrological models, such as HBV and PREVAH (PREecipitation Runof EVApotranspiration Hydrological response unit model; Viviroli et al. 2009), have demonstrated weaker performance in simulating soil moisture, particularly during dry conditions (Orth et al., 2015), with slightly better agreement with observations observed during wet conditions. Table 3 provides detailed information on the mean and 95% confidence interval (CI) of the parameter values of the prior and posterior simulated results of N_0 . One of the important addition of this work is incorporating the lattice water account and we used the regionalization equation to calculate the lattice water which depends on the clay content with free parameters in Eqs. 2. The optimized parameter of the mHM shows the variation of the θ_{lw} ranges between (0.02–0.04) $\text{cm}^3\text{cm}^{-3}$ for different sites (see Supporting Information Table S3). This behaviour and the way it was defined in Eq. 1 indicate that θ_{lw} likely represents not only soil lattice water itself, but rather the total offset of all hydrogen pools in the vicinity of the sensor (see e.g., Schrön et al., 2017; Iwema et al., 2021).




Figure 4. Probability Density Function (PDF) of the mHM parameter N_0 [cph] for two different approaches: (a) the Desilets method, and (b) the COSMIC method. The prior PDF of the original sample, consisting of 100 000 data points, is represented by the grey color. The behavioral PDF, obtained after applying the objective function, is shown for weighted (brown), uniform (green), and COSMIC (purple). The black dashed line represents the one N_0 [cph] value that best fits the data.

Table 3. The four most right columns are the posterior ones of size 100 000 and are the same for all sites. The posterior distributions correspond to the distributions in the behavioral sample obtained after the application of the objective function. The values correspond to the median (Q_{50}) and the lower and upper bound of the 95% confidence interval ($Q_{2.5}$ and $Q_{97.5}$, respectively).

Parameter	CI	Prior	<i>Grosses Bruch</i>	<i>Hohes Holz</i>	<i>Hordorf</i>	<i>Cunnersdorf</i>
$N_{0\text{(Des,U)}}$	Q_{50}	1050	889	882	1092	1255
	$Q_{2.5}$	623	866	855	1069	1224
	$Q_{97.5}$	1477	945	908	1126	1291
	mean	1050	899	883	1096	1257
	sd	260	27	21	21	25
$N_{0\text{(Des,W)}}$	Q_{50}	1050	942	906	1158	1209
	$Q_{2.5}$	623	876	871	1116	1168
	$Q_{97.5}$	1477	954	915	1176	1262
	mean	1050	925	899	1152	1216
	sd	260	34	16	21	34
$N_{0\text{(COSMIC)}}$	Q_{50}	250	225	234	287	316
	$Q_{2.5}$	108	202	216	265	281
	$Q_{97.5}$	392	249	241	302	339
	mean	250	226	232	285	312
	sd	87	18	9	12	19

315 3.2 Time series analysis of simulated neutron counts

The study conducts simulations of neutron counts in mHM using soil moisture parameterizations, with results presented in Figs. 5–6 across different sites. The simulated neutron counts were based on the simulated soil moisture content at the modeled soil horizons i.e., 0–5 cm, 5–25 cm, and 25–60 cm. The results of the ensemble runs show that the precision is higher for the behavioral simulation ensembles 0.1 % (represented by dark gray shaded areas) than in the unconstrained simulated data 1 % (represented by light gray shaded areas). We select the best 0.1 % with the highest KGE from 100 000 model runs, and the results are presented in Tab. 4.

The N_{COSMIC} method performs best at the forest site (*Hohes Holz*), whereas at the agricultural sites (*Hordorf* and *Cunnersdorf*), the $N_{\text{Des,W}}$ method performs slightly better. Only for the grassland site (*Grosses Bruch*), the uniform method $N_{\text{Des,U}}$ slightly outperforms the other two methods i.e., $N_{\text{Des,W}}$ and N_{COSMIC} , while overestimating the observations at all the other sites. In general, we observe good model performance for all methods indicated by a correlation coefficient greater than 0.80 and a percent bias (PBIAS) below 2 % across the majority of investigated sites and methods. These results suggest that the neutron-forward models match the observed neutron counts well. However, the mean ensemble had difficulties reproducing the neutron counts for the *Grosses Bruch* site in all three methods. The incorporation of dynamic vegetation in models is important

as it can impact the model parameter LAI, which in turn can affect root water uptake and soil water content. Currently, these
330 factors are not considered in the models, leading to a permanent and systematic shift in these variables each year (Zink et al.,
2017; Massoud et al., 2019).

The results also highlight the uncertainties associated with model simulations and the sensitivity of the objective function.
We find that three soil moisture-related parameters, namely N_0 , *rotfrcoeffpre*, and *rotfrcoeffforest*, have the most significant impact
on the objective function $KGE_{\alpha\beta}$, compared to the other parameters of mHM. The parameter N_0 directly affects the neutron
335 count simulations, while the parameters *rotfrcoeffpre* and *rotfrcoeffforest* correspond to the fractions of vegetation roots in different
soil layers that directly affect the water availability related stress for the estimation of actual evapotranspiration, and thereby
the soil-water dynamics (Samaniego et al., 2010b; Kumar et al., 2013b). The best parameter set values in mHM across all sites
and methods are given in (Supporting Information Table S2).




Figure 5. Simulated daily time series of black for $N_{Des,W}$, red for $N_{Des,U}$ for the four sites. The black lines represent the median of the behavioural simulation ensembles that satisfy the objective function which is LHS10 ensemble members. The light grey shaded areas represent the 95% CI of the simulation ensembles corresponding to different levels of constraining which is LHS1000 ensemble members, and the observation is shown in grey points. Precipitation is shown in blue color on the top.




Figure 6. Simulated daily time series of N_{COSMIC} for the four sites. The black lines represent the median of the behavioural simulation ensembles that satisfy the objective function which is LHS10 ensemble members. The light grey shaded areas represent the 95% CI of the simulation ensembles corresponding to different levels of constraining which is LHS1000 ensemble members, and the observation is shown in grey points.

3.3 Model calibration statistics and evaluation

340 In addition to $\text{KGE}_{\alpha\beta}$, the four metrics KGE, NSE, R, and PBIAS are used to evaluate further the mHM neutron counts simulated with observed CRNS data. We employ LHS to generate a parameter sample of 100 000 for the three methods, namely $N_{\text{Des,U}}$, $N_{\text{Des,W}}$ and N_{COSMIC} , by uniformly distributing the ranges provided in the (supplementary Table S1). The top 10 parameter sets are found to perform satisfactorily with a KGE range of 0.80 to 0.93, as demonstrated in Table 4. The

calibrated parameter sets obtained from different objective functions are also evaluated and compared using various statistical indices, as shown in Figure 7, with most objective functions performing better than satisfactory based on the criteria in Table 2. The results for the COSMIC method indicate that the main contribution to poorer results during the evaluation period was due to the variability term (α). The boxplot displayed in Figure 7 illustrates the threshold achieved by the top 1000, 100, and 10 LHS members, along with the corresponding percentage of the best 10 LHS parameter sets that meet the threshold, as specified in (see Tab. 2). Among the 31 parameters selected to simulate neutron counts, this plot provides an overview of the distribution of results and their variability with respect to the threshold criteria.

Table 4. Performance metrics for model calibration (2014-2021) using various methods: Percent bias (PBIAS), coefficient of determination (R^2), Nash-Sutcliffe efficiency (NSE), and Kling-Gupta Efficiency (KGE) across different sites. Bold values indicate the best performance measures.

Sites	<i>Grosses Bruch</i>			<i>Hohes Holz</i>			<i>Hordorf</i>			<i>Cunnersdorf</i>		
Methods:	$N_{Des,U}$	$N_{Des,W}$	N_{COSMIC}	$N_{Des,U}$	$N_{Des,W}$	N_{COSMIC}	$N_{Des,U}$	$N_{Des,W}$	N_{COSMIC}	$N_{Des,U}$	$N_{Des,W}$	N_{COSMIC}
KGE $_{\alpha\beta}$	0.97	0.96	0.93	0.96	0.99	0.99	0.99	0.99	0.98	0.93	0.99	0.99
KGE	0.85	0.83	0.81	0.81	0.80	0.85	0.87	0.87	0.86	0.88	0.93	0.90
NSE	0.69	0.60	0.29	0.54	0.60	0.70	0.75	0.75	0.73	0.57	0.85	0.79
R^2	0.73	0.69	0.66	0.66	0.65	0.73	0.76	0.76	0.75	0.86	0.86	0.81
R	0.85	0.83	0.81	0.81	0.80	0.85	0.87	0.87	0.86	0.93	0.93	0.90
PBIAS	0.7%	-1.3%	-3.3%	2.0%	0.3%	0.4%	0.1%	0.1%	0.3%	6.0%	0.7%	0.5%




Figure 7. Evaluation of model performance using boxplots constraining of 1000 to the best 10 parameters set at four different sites, using three different methods, namely $N_{Des,W}$ in brown, $N_{Des,U}$ in green, and N_{COSMIC} in purple. The figure presents four subplots, where (a) represents Alpha, (b) Beta, (c) $KGE_{\alpha\beta}$, and (d) Kling-Gupta efficiency (KGE) and its components, i.e., the variability term (perfect value: 1), and bias term (perfect value: 1), respectively.

3.4 Comparing evapotranspiration at Hohes-Holz: eddy covariance observed data vs mHM simulation

The ensemble model of (10 members) simulations, is further examined with the evapotranspiration (ETa) to cross-evaluate and assess the model's ability to represent other fluxes and states next to neutron counts by using the ETa observational data from eddy covariance measurements Integrated Carbon Observation System (ICOS) at *Hohes Holz* Warm Winter (2022). In terms

355 of temporal dynamics, the model is able to capture the observed ETa quite well at the study site, as shown in Figure 8. Panel
(c) displays the scatter plot incorporating linear regression models to quantify the relationships between observed and mHM-
simulated ETa during both the growing and non-growing seasons. This plot provides insights into the seasonal variations in the
relationship between observed and simulated ET. It suggests that the model performs best during winter, while its performance
during summer is comparatively weaker. The correlation coefficients (r values) for each season are as follows: autumn [SON]
360 ($r = 0.79$), spring [MAM] ($r = 0.77$), summer [JJA] ($r = 0.42$), and winter [DJF] ($r = 0.87$). It is worth noting that winter shows
the highest correlation between observed and simulated ET, while summer exhibits the lowest correlation. The most significant
deviation in terms of RMSE is evident during the summer, when evapotranspiration is highest, while the smallest difference is
in winter when evapotranspiration has less impact. The model slightly overestimates evapotranspiration in summer and spring,
possibly because of the absence of a dynamic vegetation growth module in the mHM, also discussed for evapotranspiration in
365 Zink et al. (2017). The temporal dynamics of the model-simulated evapotranspiration are in good agreement with the observed
data from the *Hohes Holz* forest eddy covariance site, taken from Warm Winter (2022), as illustrated in Figure 8a. The daily
correlation between observed and simulated evapotranspiration is observed high in the growing season at $r = 0.84$, whereas
the lowest correlation is found in the non-growing season at $r = 0.65$ in Figure 8c. The highest deviation in terms of RMSE
is observed during summer when the highest fluxes occur, and the lowest during winter, in which the contribution of ETa is
370 lowest.

In Figure 8b, the prior and posterior parameter distributions of evapotranspiration for *Hohes Holz* are displayed. The prior
distribution represents the 100 000 parameters set utilized for the neutron counts simulation under Latin Hypercube Sampling
(LHS). The results demonstrate that the ensemble model of 10-member simulations (posterior) for neutron counts can also
effectively capture evapotranspiration, exhibiting a root mean square error (RMSE) of 0.76 mmd^{-1} of the growing season and
375 0.25 mmd^{-1} for non-growing when compared to observed ICOS data and simulated mHM. When compared to the model
simulations with prior parameter sets, we notice a substantial improvement in ET simulations (mean RMSE of 0.86 mmd^{-1} to
 0.76 mmd^{-1}). Furthermore, the RMSE range is also narrower for the posterior simulations compared to the prior ones which
further demonstrated the additional value of incorporating CRNS measurements in improving the consistency of both modeled
soil moisture and evapotranspiration estimates. Nevertheless, the overall agreement between modeled and observed ETa is
380 reasonably good; and the analysis reveals further improvement of model performance in the growing season.




Figure 8. (a) Time series of actual evapotranspiration (ET_a) from mHM (black), ICOS measurement (red), and the prior range of 100 000 realizations in (orange) color. (b) Boxplot of daily actual evapotranspiration (ET_a) differences between the growing and non-growing seasons, comparing two selected prior with 100 000 simulations, the values represent the mean of the statistical metrics and posterior with 10 ensemble member distributions using the root mean square error (RMSE) as the evaluation metric ($\mu\text{g m}^3$). (c) scatterplots of modeled vs. observed ET on a daily basis from ICOS during the growing season from March to August (green) and non-growing season from September to February (brown) at *Hohes Holz* eddy covariance station in a forest.

4 Discussion

This study assessed the suitability of CRNS observations at four sites to enhance soil moisture representation in mHM. The theoretical measurement depth for the cosmic-ray probe varies, ranging from ~ 12 cm in wet soils to ~ 76 cm in dry soils (Zreda et al. (2008, 2012a); Rosolem et al. (2014)).

385 To improve the soil moisture profile representation within mHM it is a major challenge to use a single vertically integrated
CRNS measurement. In order to have a fair comparison between the model and observed CRNS data, two conceptually dif-
ferent approaches were integrated into mHM to calculate neutron counts from different SWC horizon depths i.e., an empirical
method based on Desilets et al. (2010), and a physics-based method (COSMIC) based on Shuttleworth et al. (2013). Since
the empirical method is described by an analytical expression, taking into account the uniform average of the soil moisture
390 layers, it is straightforward to implement and therefore most commonly used (Zreda et al., 2012b; Rivera Villarreyes et al.,
2011; Andreasen et al., 2017; Bogena et al., 2022). However, the method comes with the risk of missing a representation of
the vertical profile of soil properties and water content. Therefore, we extended this uniform-averaging scheme with a vertical
weighting scheme to mimic the sensitivity of the neutrons to the upper layers both weighted and non-weighted soil moisture
approaches in the context of CRNS have been discussed (Rivera Villarreyes et al., 2014; Baroni and Oswald, 2015; Schreiner-
395 McGraw et al., 2016; Zreda, 2016; Schrön et al., 2017; Vather et al., 2019; Barbosa et al., 2021). The COSMIC operator also
accounts for the full soil moisture profile, but in a more physically behaved manner, following the track and attenuation of the
neutrons in and out of the soil column. The mHM model is now able to simulate neutrons directly with all three approaches.
The presented results confirmed general consistency with CRNS observations at four sites in Germany (Figs. 5 and 6).

Agricultural land presents a valuable opportunity to examine the interaction between soil moisture dynamics, crop growth,
400 irrigation methods, and vegetation dynamics. *Hordorf* and *Cunnerdorf* are specific agricultural sites where seasonal changes
in aboveground biomass are expected to be larger due to crop growth and harvest compared to grassland and forest sites. The
study by Schrön et al. (2017) found that the revised weighting strategy for CRNS data improved the accuracy of soil moisture
predictions at agriculture sites, but there is still room for improvement in capturing local dynamics through revised parameters
in the CRNS model. Our results showed that at the agriculture site, the $N_{Des,W}$ methods in mHM slightly out-performed the
405 other methods.

We also investigated *Hohes Holz*, a forest site, and observed an early simulation of approximately 28 days in the simulation
of neutron counts compared to the observations. The early simulation phase could be attributed to the limitation of mHM in
simulating the dynamics of detailed vegetation mechanisms Zink et al. (2017). One specific limitation is that the model does
not fully account for the fact that trees at the site have access to deeper water sources, which can result in water stress being
410 experienced at later times. Still, we get very good results in terms of KGE, for instance, indicating that these issues are of
minor importance and that all three methods in mHM representation of the forest are already performing quite well. While
CRNS and TDR generally agree at this site, the discrepancy shown in our results could be attributed to issues related to process
representation in mHM Boeing et al. (2022). Simulation of neutron data within the mHM model and subsequently comparing
it with observed counts can enhance the accuracy and precision of soil moisture measurements. Future research can focus on
415 exploring the potential relationships between CRNS data and soil moisture anomalies, thus furthering our understanding of the
dynamics of drought and assisting in the development of efficient drought monitoring and mitigation strategies.

To cross-evaluate our results, we generated and filtered the 100 000 regionalized parameter sets based on observed neutron
counts for behavioral solutions. After selecting the most effective solutions, we conduct cross-validation by comparing the
mHM simulations of evapotranspiration against observational data from eddy covariance measurements ICOS (Warm Winter,

420 2022; Pohl et al., 2023) at the *Hohes Holz*. Figure 8 shows the scatter including the seasonal correlation coefficient at the forest site. The results indicate low correlations in summer, likely due to mHM’s limitations in capturing evapotranspiration values with mHM’s static vegetation module. However, the model performs well in winter, with a high correlation between observed and simulated values of evapotranspiration, the results confirm the findings from Zink et al. (2017), who used mHM to estimate evapotranspiration, groundwater recharge, soil moisture, and runoff with 4 km spatial and daily temporal resolutions (1951–
425 2010). They utilized soil moisture observations from eddy covariance stations employing Time-Domain Reflectometer (TDR) or Frequency-Domain Reflectometer (FDR) sensors. Due to disparities in spatial representativeness and sampling depth, a direct comparison between observed and simulated soil moisture was not feasible, their findings revealed deviations in evapotranspiration during spring and in cropland areas, while soil moisture estimations exhibited good agreement with observed dynamics. The study highlights the importance of considering seasonal variations when analyzing the results. Discrepancies,
430 such as low correlations in summer, indicate the need for improvements in capturing evapotranspiration dynamics under varying environmental conditions. Refining vegetation dynamics representation could enhance the simulation of evapotranspiration processes. Additionally, the agreement between mHM and observed soil moisture dynamics suggests variable model performance for different hydrological variables, emphasizing the need for a comprehensive assessment of its capabilities across various environmental conditions and spatiotemporal scales. The accuracy of modeled evapotranspiration is linked to soil pa-
435 rameterization because soil water is the main source of evaporative water. During the growing season (summer), the model exhibited the largest variability in modeled ETa (Figure 8c). This can be associated among other things with a lack of a dynamic vegetation growth module in mHM, which may not capture the onset of the vegetation period adequately. This variability could also be attributed to seasonal changes in vapor pressure difference (VPD) or more localized processes occurring at the forest site (e.g., under-story vegetation), which are currently not considered in the model. The comparison of observed neutron
440 counts with simulated counts from mHM improved not only soil moisture estimation but also evapotranspiration estimation in the model. This provides evidence that CRNS data has the potential to improve hydrological process understanding as a whole. The *Grosses Bruch* site stands out as a mesophilic grassland site with a nearby water channel, shallow ground water, regular cattle grazing, and seasonal flooding (Hermanns et al., 2021). We find a large ensemble-related uncertainty at this site for all three methods, while the uniformly weighted approach $N_{Des,U}$ shows a slightly better performance than the other two methods
445 $N_{Des,W}$ and N_{COSMIC} (see Table. 4). The behaviour may result in a missing representation of locally significant hydrological components, such as dynamic biomass, snow, shallow ground water, or nearby surface ponding (Schrön et al., 2017). Moreover, in the middle of September, many cows had been present at this site, which could have led to a non-negligible variation of the neutron signal and thus to a non-meaningful expression of correlation-related measures Schrön et al. (2017). Döpfer et al. (2022) mentioned thigh impact of grazing on the plant traits and soil properties at this site. Additionally, the use of one
450 grid cell measurements by mHM in our study may have limited the accuracy of our results, as the depth of measurement may not be representative of the entire soil profile. Notably, neutron counts were found to provide a more accurate representation of soil water content during June, July, August, and September, when levels tend to be lower. Further exploration of neutron counts may yield additional improvements to model performance.

Overall, the three methods ($N_{\text{Des,U}}$, $N_{\text{Des,W}}$, and N_{COSMIC}) in mHM were able to consistently simulate the neutron count
455 variability throughout the available data period. However, a broader confidence interval is observed, indicating a greater range
of variations, which implies a higher degree of uncertainty in the N_{COSMIC} . The COSMIC approach explicitly accounts for water
content snow, vegetation interception, and root-zone soil processes that may likely lead to a better representation of observed
neutron count variation compared to Desilets that empirically represent such processes. The simulated time series tended to
slightly underestimate the CRNS neutron count rate, particularly during the dry season. This effect could be explained by the
460 known limitations of the equations under very dry conditions, while recent approaches exist (Köhli et al., 2021) that could lead
to further improvement in future studies. Nevertheless, the results generally confirmed the better performance of the $N_{\text{Des,W}}$
than $N_{\text{Des,U}}$, because of its more realistic representation of neutron propagation with depth. After optimizing the soil hydraulic
properties based on CRNS data, the integrated signal was reproduced very well (Fig. 5). The better performance of N_{COSMIC}
and $N_{\text{Des,W}}$ over $N_{\text{Des,U}}$ demonstrates the benefits of explicitly resolving individual soil moisture profiles, bulk densities, and
465 lattice water, as opposed to a uniform average across the layers. This perception, however, might depend on site-specific soil
profile characteristics and be less prominent if profiles are largely uniform or incorrectly resembled by the model structure.
We also included offset hydrogen pools in the form of lattice water to the N_0 calibration function, which was important for
more accurate soil moisture estimates, confirming initial suggestions by Bogena et al. (2013). Moreover, a strong correlation
between biomass and the N_0 parameter was reported in several studies (Franz et al., 2013; Hawdon et al., 2014; Baatz et al.,
470 2014, 2015). In our study, we pass the N_0 parameter as a calibration parameter set in mHM. In using the CRNS soil moisture
measurement the drier locations show larger deviations than the wetter locations (Iwema et al., 2015). The possibility of using
simulated high-resolution soil moisture profiles instead of a few measurements at different soil depths could further increase the
accuracy of the model predictions (Brunetti et al., 2019). One of the primary sources of uncertainty at the *Grosses Bruch* site is
surface ponding and shallow groundwater, as well as the loamy texture of the soil. Those factors contribute to the formation of
475 permanent water ponds in the area and may introduce uniform or even inverse soil moisture profiles which directly influence
the neutron emissions, but cannot be captured by the mHM model. Another factor is the time-variable effect of crowding
cows near the station, which may influence the CRNS signal, but is challenging to correct in the CRNS measurement (Schrön
et al., 2017). We incorporate the CRNS parameter set in mHM, and some parameters related to soil moisture and neutron
counts are effectively constrained based on the objective function using $\text{KGE}_{\alpha\beta}$. However, there is still room for improvement,
480 particularly with regard to the coefficient in root fractions distributed across soil layers.

According to Beck et al. (2021), model calibration provides more overall benefits than data assimilation. Furthermore, model
calibration can be advantageous for regions with both sparse and dense rain gauge networks, whereas data assimilation is more
beneficial for regions with sparse rain gauge networks. In this paper, the Latin Hypercube Sampling (LHS) method McKay
and Conover (1979) is adopted to generate input variable samples, which is a stratified sampling method that reduces the
485 number of simulations required compared to the conventional Monte Carlo method Iman and Helton (1988). LHS divides the
range of each input into N intervals and selects one representative value from each interval to ensure full coverage of the input
variables range and representation of all possible values in the simulation. Previous studies by Smith et al. (2019) and Liu et al.
(2022) address the challenges of using the original KGE in Markov chain Monte Carlo (MCMC) methods, offering insights for

accurate parameter estimation and posterior distribution exploration. To address this issue, it is recommended to use adaptations
490 to the LHS method instead of directly using the original KGE to improve the exploration of the posterior distributions. Our
approach can estimate the posterior distributions of model parameters based on the objective function $KGE_{\alpha\beta}$ by taking the
variance and bias.

This paper provides a framework to incorporate CRNS data into the mHM to assess the accuracy of soil water content on dif-
ferent land cover types, including agricultural land, deciduous forest, and grassland. The integration of methods from Desilets
495 et al. (2010) and Shuttleworth et al. (2013) in mHM, using climatic data and soil physical parameters, the parameterization of
evapotranspiration is effectively improved (see Fig. 9). Additionally, Supplementary Fig. S5 illustrates the improved represen-
tation of soil moisture for Hohes Holz. This framework lies in its ability to utilize observed neutron counts data, allowing for a
comprehensive assessment of the model's performance and enhancing its reliability in hydrological modeling.




Figure 9. Comparison of weekly observed actual evapotranspiration (grey dots) and simulated actual evapotranspiration using the default mHM parameters by Boeing et al. (2022) (red line) and the calibrated simulation (blue dots) over the *Hohes Holz* site.

5 Conclusion and future outlook

500 This study evaluates the potential of the mHM a large-scale hydrological model for simulating neutron counts at the 1.2
km \times 1.2 km scale across different land cover sites for the period 2014–2021. Two empirical and one physical model approaches
are evaluated for deriving neutrons from the soil moisture profile. Neutron measurement data from four sites in Germany are
integrated, and the influence on hydrological model parameters, as well as simulated soil moisture and evapotranspiration are
analyzed. The parameter sample of 100 000 realizations for neutron counts was taken, which are analyzed regarding their
505 uncertainty caused by the parameter estimation. The parameter sets are filtered based on the KGE of observed vs simulated
neutron counts. The best 1% member ensemble simulations are evaluated with neutron counts, evapotranspiration, and soil
moisture observations.

The evaluation of neutron counts at four different sites yields a KGE value of > 0.8 , indicating a satisfactory representation
of the neutron observations. The 1% ensemble parameter set is found more representative of 100 000 realizations, suggesting

510 a reliable model performance. The performance of the neutron counting methods varies across different land cover types. The $N_{Des,W}$ method generally demonstrates good performance, particularly at the agricultural sites. While the N_{COSMIC} method performs slightly better at forest site and the $N_{Des,U}$ method shows slightly better results at the grassland site.

However, there is still room for improvement in some areas. Specifically, working with grassland sites presented challenges, particularly with the N_{COSMIC} method. On the one hand it is a physics-based approach incorporating a comprehensive representation of the neutron counting process, but on the other hand, it relies on the detailed representation of the site characteristics in the hydrological model. This complexity could introduce additional uncertainties and limitations in the model, potentially affecting its performance, especially when the site is more complex than it has been modeled. The study suggests that the observed discrepancies between model and observations may be attributed to the representation of dynamic biomass, snow, surface ponding, and shallow groundwater dynamics, which are present at the grassland site, for instance. Addressing these features could further enhance the model's accuracy.

The evaluation with evapotranspiration from eddy covariance at *Hohes Holz* stations indicates deficiencies in mHM to deal with forest systems, but also great potential for CRNS measurements to improve the water partitioning as a whole. Especially in the growing season (March-August), deviations of the modeled and observed ETa indicate room for better representation of mixed soils and dynamic vegetation modules at the local scale within mHM. The calibration on neutron counts not only improved the soil moisture performance of the model but also helped to set the modeled evapotranspiration straight.

In conclusion, the incorporation of neutron counts estimation into mHM by accounting for vertical soil moisture profiles improves the model's accuracy and provides a more realistic representation of soil moisture dynamics at all four study sites and evapotranspiration at *Hohes Holz* site. This research presents a direction for future studies to explore. The next step in this research is to evaluate the ability of this CRNS module in mHM for estimating soil moisture through a large-scale soil moisture monitoring initiative, e.g. by utilizing more stationary CRNS networks or the novel rail-based CRNS data from Altdorff et al. (2023). To optimize accuracy and understanding, we recommend integrating both CRNS and satellite remote sensing data into mHM. Improving the model predictions will contribute to reducing the uncertainties associated with drought and flood management strategies and informed agricultural decisions.

Code availability. Simulation data is attached as supplemental material. The mesoscale Hydrological Model mHM (version 5.12) is open-source and can be freely accessed from GitLab: https://git.ufz.de/mhm/mhm/-/tree/v5.12.0?ref_type=tags.

Data availability. We kindly acknowledge the German Weather Service (DWD) for providing the meteorological datasets. The terrain elevation data was collected from USGS EROS Archive - Digital Elevation - Global Multi-resolution Terrain Elevation Data 2010 (GMTED2010), available at <https://www.usgs.gov/centers/eros/science/usgs-eros-archive-digital-elevation-global-multi-resolution-terrain-elevation>. Gridded soil characteristics are based on the BUEK200 database obtained from the German Federal Institute for Geosciences and Natural Resources (BGR, see online at <https://geoportal.bgr.de/mapapps/resources/apps/geoportal/index.html?lang=en#/datasets/portal/154997F4-3C14-4A53-B217-8A7C7509E05F>). The geological dataset was downloaded from Institute for Biogeochemistry and Marine

Chemistry, KlimaCampus, Universitt Hamburg (<https://www.geo.uni-hamburg.de/en/geologie/forschung/aquatische-geochemie/glim.html>).
Leaf Area Index (LAI) dataset was downloaded from the Global Land Cover Facility (GLCF), available at <http://iridl.ldeo.columbia.edu/SOURCES/UMD/GLCF/GIMMS/NDVIg/global/index.html>. The land cover dataset was downloaded from the European Space Agency
545 (ESA), available at http://due.esrin.esa.int/page_globcover.php. The ET data were obtained from <https://zenodo.org/record/7561854>.

Competing interests. RK and LS are members of the editorial board of Hydrology and Earth System Sciences.

Acknowledgements. The authors thank all the site owners for maintaining the local sensors, particularly to F. Böttcher (DWD) and E. Thiel (SKWP). The study has been made possible by the Terrestrial Environmental Observatories (TERENO), an infrastructural fund of the Helmholtz Association. The High-Performance Computing Cluster EVE has contributed to the computation of the scientific findings. Eshrat
550 Fatima is grateful for the financial support of the German Academic Exchange Service (DAAD) through the Graduate School Scholarship Program under Reference Number 91788160.

References

- Albergel, C., Calvet, J.-C., Rosnay, P. d., Balsamo, G., Wagner, W., Hasenauer, S., Naeimi, V., Martin, E., Bazile, E., Bouyssel, F., et al.: Cross-evaluation of modelled and remotely sensed surface soil moisture with in situ data in southwestern France, *Hydrology and Earth System Sciences*, 14, 2177–2191, 2010.
- 555 Altdorff, D., Oswald, S. E., Zacharias, S., Zengerle, C., Dietrich, P., Mollenhauer, H., Attinger, S., and Schrön, M.: Toward Large-Scale Soil Moisture Monitoring Using Rail-Based Cosmic Ray Neutron Sensing, *Water Resources Research*, 59, e2022WR033 514, 2023.
- Andreasen, M., Jensen, K. H., Desilets, D., Franz, T. E., Zreda, M., Bogaena, H. R., and Looms, M. C.: Status and perspectives on the cosmic-ray neutron method for soil moisture estimation and other environmental science applications, *Vadose Zone Journal*, 16, 1–11, 2017.
- 560 Avery, W. A., Finkenbiner, C., Franz, T. E., Wang, T., Nguy-Robertson, A. L., Suyker, A., Arkebauer, T., and Muñoz Arriola, F.: Incorporation of globally available datasets into the roving cosmic-ray neutron probe method for estimating field-scale soil water content, *Hydrology and Earth System Sciences*, 20, 3859–3872, <https://doi.org/10.5194/hess-20-3859-2016>, 2016.
- Baatz, R., Bogaena, H., Franssen, H.-J. H., Huisman, J., Qu, W., Montzka, C., and Vereecken, H.: Calibration of a catchment scale cosmic-ray probe network: A comparison of three parameterization methods, *Journal of Hydrology*, 516, 231–244, 2014.
- 565 Baatz, R., Bogaena, H., Hendricks Franssen, H.-J., Huisman, J., Montzka, C., and Vereecken, H.: An empirical vegetation correction for soil water content quantification using cosmic ray probes, *Water Resources Research*, 51, 2030–2046, 2015.
- Baatz, R., Hendricks Franssen, H.-J., Han, X., Hoar, T., Bogaena, H. R., and Vereecken, H.: Evaluation of a cosmic-ray neutron sensor network for improved land surface model prediction, *Hydrology and Earth System Sciences*, 21, 2509–2530, 2017.
- 570 Barbosa, L. R., Coelho, V. H. R., Scheffele, L. M., Baroni, G., Ramos Filho, G. M., Montenegro, S. M., Almeida, C. d. N., and Oswald, S. E.: Dynamic groundwater recharge simulations based on cosmic-ray neutron sensing in a tropical wet experimental basin, *Vadose Zone Journal*, 20, e20 145, 2021.
- Baroni, G. and Oswald, S.: A scaling approach for the assessment of biomass changes and rainfall interception using cosmic-ray neutron sensing, *Journal of Hydrology*, 525, 264–276, 2015.
- 575 Beck, H. E., Pan, M., Miralles, D. G., Reichle, R. H., Dorigo, W. A., Hahn, S., Sheffield, J., Karthikeyan, L., Balsamo, G., Parinussa, R. M., et al.: Evaluation of 18 satellite-and model-based soil moisture products using in situ measurements from 826 sensors, *Hydrology and Earth System Sciences*, 25, 17–40, 2021.
- BGR: Digital soil map of Germany 1 : 200,000 (BUEK 200) v0.5, https://www.bgr.bund.de/DE/Themen/Boden/Informationsgrundlagen/Bodenkundliche_Karten_Datenbanken/BUEK200/buek200_node.html, [Accessed: October 7, 2022], 2020.
- 580 Boeing, F., Rakovec, O., Kumar, R., Samaniego, L., Schrön, M., Hildebrandt, A., Reibmann, C., Thober, S., Müller, S., Zacharias, S., Bogaena, H., Schneider, K., Kiese, R., Attinger, S., and Marx, A.: High-resolution drought simulations and comparison to soil moisture observations in Germany, *Hydrology and Earth System Sciences*, 26, 5137–5161, <https://doi.org/10.5194/hess-26-5137-2022>, 2022.
- Bogaena, H., Huisman, J., Baatz, R., Hendricks Franssen, H.-J., and Vereecken, H.: Accuracy of the cosmic-ray soil water content probe in humid forest ecosystems: The worst case scenario, *Water Resources Research*, 49, 5778–5791, 2013.
- 585 Bogaena, H. R., Schrön, M., Jakobi, J., Ney, P., Zacharias, S., Andreasen, M., Baatz, R., Boorman, D., Duygu, M. B., Eguibar-Galán, M. A., et al.: COSMOS-Europe: a European network of cosmic-ray neutron soil moisture sensors, *Earth System Science Data*, 14, 1125–1151, 2022.

- Brunetti, G., Bogen, H., Baatz, R., Huisman, J. A., Dahlke, H., and Vereecken, H.: On the information content of cosmic-ray neutron data in the inverse estimation of soil hydraulic properties, *Vadose Zone Journal*, 18, 1–24, 2019.
- 590 Chan, S. K., Bindlish, R., O’Neill, P., Jackson, T., Njoku, E., Dunbar, S., Chaubell, J., Piepmeier, J., Yueh, S., Entekhabi, D., et al.: Development and assessment of the SMAP enhanced passive soil moisture product, *Remote sensing of environment*, 204, 931–941, 2018.
- Chen, F., Crow, W. T., Starks, P. J., and Moriasi, D. N.: Improving hydrologic predictions of a catchment model via assimilation of surface soil moisture, *Advances in Water Resources*, 34, 526–536, 2011.
- Desilets, D., Zreda, M., and Ferré, T. P.: Nature’s neutron probe: Land surface hydrology at an elusive scale with cosmic rays, *Water Resources*
595 *Research*, 46, 2010.
- Dimitrova-Petrova, K., Geris, J., Wilkinson, M. E., Rosolem, R., Verrot, L., Lilly, A., and Soulsby, C.: Opportunities and challenges in using catchment-scale storage estimates from cosmic ray neutron sensors for rainfall-runoff modelling, *Journal of Hydrology*, 586, 124–137, 2020.
- Dong, J. and Ochsner, T. E.: Soil texture often exerts a stronger influence than precipitation on mesoscale soil moisture patterns, *Water*
600 *Resources Research*, 54, 2199–2211, 2018.
- Dong, J., Ochsner, T. E., Zreda, M., Cosh, M. H., and Zou, C. B.: Calibration and validation of the COSMOS rover for surface soil moisture measurement, *Vadose Zone Journal*, 13, 1–8, 2014.
- Döpfer, V., Rocha, A. D., Berger, K., Gränzig, T., Verrelst, J., Kleinschmit, B., and Förster, M.: Estimating soil moisture content under grassland with hyperspectral data using radiative transfer modelling and machine learning, *International Journal of Applied Earth Observation*
605 *and Geoinformation*, 110, 102817, 2022.
- ESA: Global Land Cover Map for 2009, http://due.esrin.esa.int/files/Globcover2009_V2.3_Global_.zip, last accessed on 1 June 2021, 2009.
- Franz, T., Zreda, M., Rosolem, R., and Ferre, T.: A universal calibration function for determination of soil moisture with cosmic-ray neutrons, *Hydrology and Earth System Sciences*, 17, 453–460, 2013.
- Franz, T. E., Zreda, M., Ferre, T., Rosolem, R., Zweck, C., Stillman, S., Zeng, X., and Shuttleworth, W.: Measurement depth of the cosmic
610 ray soil moisture probe affected by hydrogen from various sources, *Water Resources Research*, 48, 2012.
- Fuamba, M., Branger, F., Braud, I., Batchabani, E., Sanzana, P., Sarrazin, B., and Jankowsky, S.: Value of distributed water level and soil moisture data in the evaluation of a distributed hydrological model: Application to the PUMMA model in the Mercier catchment (6.6 km²) in France, *Journal of Hydrology*, 569, 753–770, 2019.
- Greacen, E. L.: Soil water assessment by the neutron method, CSIRO, Australia, 1981.
- 615 Gupta, H. V., Sorooshian, S., and Yapo, P. O.: Status of automatic calibration for hydrologic models: Comparison with multilevel expert calibration, *Journal of hydrologic engineering*, 4, 135–143, 1999.
- Gupta, H. V., Kling, H., Yilmaz, K. K., and Martinez, G. F.: Decomposition of the mean squared error and NSE performance criteria: Implications for improving hydrological modelling, *Journal of Hydrology*, 377, 80–91, 2009.
- Hargreaves, G. H. and Samani, Z. A.: Reference Crop Evapotranspiration from Temperature, *Applied Engineering in Agriculture*, 1, 96–99,
620 <https://doi.org/10.13031/2013.26773>, 1985.
- Hawdon, A., McJannet, D., and Wallace, J.: Calibration and correction procedures for cosmic-ray neutron soil moisture probes located across Australia, *Water Resources Research*, 50, 5029–5043, 2014.
- Hermanns, F., Pohl, F., Rebmann, C., Schulz, G., Werban, U., and Lausch, A.: Inferring grassland drought stress with unsupervised learning from airborne hyperspectral VNIR imagery, *Remote Sensing*, 13, 1885, 2021.

- 625 Iman, R. L. and Helton, J. C.: An investigation of uncertainty and sensitivity analysis techniques for computer models, *Risk analysis*, 8, 71–90, 1988.
- Iwema, J., Rosolem, R., Baatz, R., Wagener, T., and Bogena, H.: Investigating temporal field sampling strategies for site-specific calibration of three soil moisture–neutron intensity parameterisation methods, *Hydrology and Earth System Sciences*, 19, 3203–3216, 2015.
- Iwema, J., Rosolem, R., Rahman, M., Blyth, E., and Wagener, T.: Land surface model performance using cosmic-ray and point-scale soil
630 moisture measurements for calibration, *Hydrology and Earth System Sciences*, 21, 2843–2861, 2017.
- Iwema, J., Schrön, M., Koltermann Da Silva, J., Schweiser De Paiva Lopes, R., and Rosolem, R.: Accuracy and precision of the cosmic-ray neutron sensor for soil moisture estimation at humid environments, *Hydrological Processes*, 35, e14419, 2021.
- Jablonowski, C.: Adaptive grids in weather and climate modeling, University of Michigan, 2004.
- James, L. D.: Selection, calibration, and testing of hydrologic models, *Hydrologic modeling of small watersheds*, pp. 437–472, 1982.
- 635 Kasner, M., Zacharias, S., and Schrön, M.: On soil bulk density and its influence to soil moisture estimation with cosmic-ray neutrons, *Hydrology and Earth System Sciences Discussions*, pp. 1–24, 2022.
- Köhli, M., Schrön, M., Zreda, M., Schmidt, U., Dietrich, P., and Zacharias, S.: Footprint characteristics revised for field-scale soil moisture monitoring with cosmic-ray neutrons, *Water Resources Research*, 51, 5772–5790, 2015.
- Kumar, R., Livneh, B., and Samaniego, L.: Toward computationally efficient large-scale hydrologic predictions with a multiscale regional-
640 ization scheme, *Water Resources Research*, 49, 5700–5714, 2013a.
- Kumar, R., Samaniego, L., and Attinger, S.: Implications of distributed hydrologic model parameterization on water fluxes at multiple scales and locations, *Water Resources Research*, 49, 360–379, 2013b.
- Kvålseth, T. O.: Cautionary note about R 2, *The American Statistician*, 39, 279–285, 1985.
- Köhli, M., Weimar, J., Schrön, M., Baatz, R., and Schmidt, U.: Soil Moisture and Air Humidity Dependence of the Above-Ground Cosmic-
645 Ray Neutron Intensity, *Frontiers in Water*, 2, 544 847, <https://doi.org/10.3389/frwa.2020.544847>, 2021.
- Liu, Y., Fernández-Ortega, J., Mudarra, M., and Hartmann, A.: Pitfalls and a feasible solution for using KGE as an informal likelihood function in MCMC methods: DREAM (ZS) as an example, *Hydrology and Earth System Sciences*, 26, 5341–5355, 2022.
- Livneh, B., Kumar, R., and Samaniego, L.: Influence of soil textural properties on hydrologic fluxes in the Mississippi river basin, *Hydrological Processes*, 29, 4638–4655, 2015.
- 650 Mai, J.: Ten strategies towards successful calibration of environmental models, *Journal of Hydrology*, p. 129414, <https://doi.org/10.1016/j.jhydrol.2023.129414>, 2023.
- Martinez, G. F. and Gupta, H. V.: Toward improved identification of hydrological models: A diagnostic evaluation of the “abcd” monthly water balance model for the conterminous United States, *Water Resources Research*, 46, 2010.
- Massoud, E. C., Xu, C., Fisher, R. A., Knox, R. G., Walker, A. P., Serbin, S. P., Christoffersen, B. O., Holm, J. A., Kueppers, L. M., Ricciuto,
655 D. M., et al.: Identification of key parameters controlling demographically structured vegetation dynamics in a land surface model: CLM4.5 (FATES), *Geoscientific Model Development*, 12, 4133–4164, 2019.
- McKay, M. and Conover, W.: RJ Beckman A comparison of three methods for selecting values of input variables in the analysis of output from a computer code, *Technometrics*, 21, 239–245, 1979.
- Nash, J. E. and Sutcliffe, J. V.: River flow forecasting through conceptual models part I—A discussion of principles, *Journal of Hydrology*,
660 10, 282–290, 1970.
- OpenStreetMap contributors: Planet dump retrieved from <https://planet.osm.org> (last access: 26 July 2021), available at: <https://www.openstreetmap.org> (last access: 26 July 2021), a, b, c, 2020.

- Orth, R., Staudinger, M., Seneviratne, S. I., Seibert, J., and Zappa, M.: Does model performance improve with complexity? A case study with three hydrological models, *Journal of Hydrology*, 523, 147–159, 2015.
- 665 Patil, A., Fersch, B., Hendricks Franssen, H.-J., and Kunstmann, H.: Assimilation of cosmogenic neutron counts for improved soil moisture prediction in a distributed land surface model, *Frontiers in Water*, p. 115, 2021.
- Pohl, F., Rakovec, O., Rebmann, C., Hildebrandt, A., Boeing, F., Hermanns, F., Attinger, S., Samaniego, L., and Kumar, R.: Long-term daily hydrometeorological drought indices, soil moisture, and evapotranspiration for ICOS sites, *Scientific Data*, 10, 281, 2023.
- Rakovec, O., Kumar, R., Attinger, S., and Samaniego, L.: Improving the realism of hydrologic model functioning through multivariate
670 parameter estimation, *Water Resources Research*, 52, 7779–7792, 2016.
- Rivera Villarreyes, C., Baroni, G., and Oswald, S. E.: Integral quantification of seasonal soil moisture changes in farmland by cosmic-ray neutrons, *Hydrology and Earth System Sciences*, 15, 3843–3859, 2011.
- Rivera Villarreyes, C. A., Baroni, G., and Oswald, S. E.: Inverse modelling of cosmic-ray soil moisture for field-scale soil hydraulic parameters, *European Journal of Soil Science*, 65, 876–886, <https://doi.org/10.1111/ejss.12162>, 2014.
- 675 Rosolem, R., Hoar, T., Arellano, A., Anderson, J. L., Shuttleworth, W. J., Zeng, X., and Franz, T. E.: Translating aboveground cosmic-ray neutron intensity to high-frequency soil moisture profiles at sub-kilometer scale, *Hydrology and Earth System Sciences*, 18, 4363–4379, 2014.
- Samaniego, L., Bárdossy, A., and Kumar, R.: Streamflow prediction in ungauged catchments using copula-based dissimilarity measures, *Water Resources Research*, 46, 2010a.
- 680 Samaniego, L., Kumar, R., and Attinger, S.: Multiscale parameter regionalization of a grid-based hydrologic model at the mesoscale, *Water Resources Research*, 46, 2010b.
- Samaniego, L., Kumar, R., Thober, S., Rakovec, O., Zink, M., Wanders, N., Eisner, S., Müller Schmied, H., Sutanudjaja, E. H., Warrach-Sagi, K., et al.: Toward seamless hydrologic predictions across spatial scales, *Hydrology and Earth System Sciences*, 21, 4323–4346, 2017.
- Samaniego, L., Thober, S., Wanders, N., Pan, M., Rakovec, O., Sheffield, J., Wood, E. F., Prudhomme, C., Rees, G., Houghton-Carr, H.,
685 Fry, M., Smith, K., Watts, G., Hisdal, H., Estrela, T., Buontempo, C., Marx, A., and Kumar, R.: Hydrological Forecasts and Projections for Improved Decision-Making in the Water Sector in Europe, *Bulletin of the American Meteorological Society*, 100, 2451–2472, <https://doi.org/10.1175/bams-d-17-0274.1>, 2019.
- Samaniego, L., Kumar, R., Zink, M., Cuntz, M., Mai, J., Thober, S., Schneider, C., Dalmasso, G., Musuuza, J., Rakovec, O., Craven, J., Schäfer, D., Prykhodko, V., Schrön, M., Spieler, D., Brenner, J., Langenberg, B., Schüler, L., Stisen, S., Demirel, C. M., Jing,
690 M., Kaluza, M., Schweppe, R., Shrestha, P. K., Döring, N., and Müller, S.: mesoscale Hydrologic Model - mHM v5.13.0, Zenodo, <https://doi.org/10.5281/zenodo.7997974>, 2023.
- Santanello Jr, J. A., Peters-Lidard, C. D., and Kumar, S. V.: Diagnosing the sensitivity of local land–atmosphere coupling via the soil moisture–boundary layer interaction, *Journal of Hydrometeorology*, 12, 766–786, 2011.
- Schreiner-McGraw, A. P., Vivoni, E. R., Mascaró, G., and Franz, T. E.: Closing the water balance with cosmic-ray soil moisture measurements
695 and assessing their relation to evapotranspiration in two semiarid watersheds, *Hydrology and Earth System Sciences*, 20, 329–345, 2016.
- Schrön, M.: Cosmic-ray Neutron Sensing and Its Applications to Soil and Land Surface Hydrology: On Neutron Physics, Method Development, and Soil Moisture Estimation Across Scales, Ph.D. thesis, Universität Potsdam, https://publishup.uni-potsdam.de/files/39543/schroen_diss.pdf, 2017.

- 700 Schrön, M., Köhli, M., Scheffele, L., Iwema, J., Bogena, H. R., Lv, L., Martini, E., Baroni, G., Rosolem, R., Weimar, J., et al.: Improving calibration and validation of cosmic-ray neutron sensors in the light of spatial sensitivity, *Hydrology and Earth System Sciences*, 21, 5009–5030, 2017.
- Schrön, M., Oswald, S. E., Zacharias, S., Kasner, M., Dietrich, P., and Attinger, S.: Neutrons on rails: Transregional monitoring of soil moisture and snow water equivalent, *Geophysical research letters*, 48, e2021GL093924, 2021.
- 705 Seneviratne, S. I., Lüthi, D., Litschi, M., and Schär, C.: Land–atmosphere coupling and climate change in Europe, *Nature*, 443, 205–209, 2006.
- Shuttleworth, J., Rosolem, R., Zreda, M., and Franz, T.: The COsmic-ray Soil Moisture Interaction Code (COSMIC) for use in data assimilation, *Hydrology and Earth System Sciences*, 17, 3205–3217, 2013.
- Smith, K. A., Barker, L. J., Tanguy, M., Parry, S., Harrigan, S., Legg, T. P., Prudhomme, C., and Hannaford, J.: A multi-objective ensemble approach to hydrological modelling in the UK: an application to historic drought reconstruction, *Hydrology and Earth System Sciences*, 710 23, 3247–3268, 2019.
- Van Steenberghe, N. and Willems, P.: Increasing river flood preparedness by real-time warning based on wetness state conditions, *Journal of Hydrology*, 489, 227–237, 2013.
- Vather, T., Everson, C., and Franz, T. E.: Calibration and validation of the cosmic ray neutron rover for soil water mapping within two South African land classes, *Hydrology*, 6, 65, 2019.
- 715 Viviroli, D., Zappa, M., Gurtz, J., and Weingartner, R.: An introduction to the hydrological modelling system PREVAH and its pre-and post-processing-tools, *Environmental Modelling & Software*, 24, 1209–1222, 2009.
- Wahbi, A., Heng, L., and Dercon, G.: Cosmic ray neutron sensing: estimation of agricultural crop biomass water equivalent, Springer Nature, 2018.
- Warm Winter, W.: Warm Winter 2020 Team and ICOS Ecosystem Thematic Centre: Warm Winter 2020 ecosystem eddy covariance flux product for 73 stations in FLUXNET-Archive format–release 2022-1 (Version 1.0), ICOS Carbon Portal [data set], 720 <https://doi.org/10.18160/2G60-ZHAK>, 2022.
- Wollschläger, U., Attinger, S., Borchardt, D., Brauns, M., Cuntz, M., Dietrich, P., Fleckenstein, J. H., Friese, K., Friesen, J., Harpke, A., et al.: The Bode hydrological observatory: a platform for integrated, interdisciplinary hydro-ecological research within the TERENO Harz/Central German Lowland Observatory, *Environmental Earth Sciences*, 76, 1–25, 2017.
- 725 Zacharias, S. and Wessolek, G.: Excluding organic matter content from pedotransfer predictors of soil water retention, *Soil Science Society of America Journal*, 71, 43–50, 2007.
- Zacharias, S., Bogena, H., Samaniego, L., Mauder, M., Fuß, R., Pu^ˆtz, T., Frenzel, M., Schwank, M., Baessler, C., Butterbach-Bahl, K., et al.: A network of terrestrial environmental observatories in Germany, *Vadose zone journal*, 10, 955–973, 2011.
- Zhao, H., Montzka, C., Baatz, R., Vereecken, H., and Franssen, H.-J. H.: The Importance of Subsurface Processes in Land Surface Modeling over a Temperate Region: An Analysis with SMAP, Cosmic Ray Neutron Sensing and Triple Collocation Analysis, *Remote Sensing*, 730 13, 3068, 2021.
- Zhuo, L., Dai, Q., Zhao, B., and Han, D.: Soil moisture sensor network design for hydrological applications, *Hydrology and Earth System Sciences*, 24, 2577–2591, 2020.
- Zink, M., Kumar, R., Cuntz, M., and Samaniego, L.: A high-resolution dataset of water fluxes and states for Germany accounting for 735 parametric uncertainty, *Hydrology and Earth System Sciences*, 21, 1769–1790, 2017.

- Zreda, M.: Land-surface hydrology with cosmic-ray neutrons: Principles and applications, *Journal of the Japanese Society of Soil Physics*, 132, 25–30, 2016.
- Zreda, M., Desilets, D., Ferré, T., and Scott, R. L.: Measuring soil moisture content non-invasively at intermediate spatial scale using cosmic-ray neutrons, *Geophysical research letters*, 35, 2008.
- 740 Zreda, M., Shuttleworth, W., Zeng, X., Zweck, C., Desilets, D., Franz, T., and Rosolem, R.: COSMOS: The cosmic-ray soil moisture observing system, *Hydrology and Earth System Sciences*, 16, 4079–4099, 2012a.
- Zreda, M., Shuttleworth, W. J., Zeng, X., Zweck, C., Desilets, D., Franz, T., and Rosolem, R.: COSMOS: the COsmic-ray Soil Moisture Observing System, *Hydrology and Earth System Sciences*, 16, 4079–4099, <https://doi.org/10.5194/hess-16-4079-2012>, 2012b.