

Answers to the comments of Referee 1

1. Page 2, line 54: *The authors use Bowers (2011) to support a statement about bacteria playing a significant role in the composition and dynamics of bioaerosols amongst all the different airborne microorganisms. The cited paper, though, focuses on the effect of weather and land-use type over diversity and abundance of airborne bacteria in the Colorado Front Range, it doesn't make any statements about other airborne bioaerosols. The reviewer suggests to revise this reference.*

The referee is right. We will change the reference with Gong et al., 2020

Gong, J., Qi, J., E, B., Yin, Y., Gao, D.: Concentration, viability and size distribution of bacteria in atmospheric bioaerosols under different types of pollution. *Environmental Pollution* 257, 113485, <https://doi.org/10.1016/j.envpol.2019.113485>, 2020.

2. Page 2, line 57: *The authors use Bauer et al. (2002) as a citation to substantiate the atmospheric concentration over land of bacteria, but the cited paper is about the carbon content of fungal spores. The reviewer suggests to use a more fitting reference or explicitate the connection between the number of bacteria in the atmosphere and the cited paper.*

The referee is right. We will change the reference by citing Burrows et al. , 2009.

Burrows, S. M., Butler, T., Jöckel, P., Tost, H., Kerkweg, A., Pöschl, U., Lawrence, M.G.: Bacteria in the global atmosphere – Part 2: Modeling of emissions and transport between different ecosystems, *Atmospheric Chem. Phys.* 9, 9281–9297. <https://doi.org/10.5194/acp-9-9281-2009>, 2009.

3. Page 3, line 65: *rather than Lighthart (2006), the reviewer suggests using “Lighthart B. Shaffer B.T. Marthi B. Ganio L.M. (1993) Artificial wind gust liberation of microbial bioaerosols previously deposited on plants. *Aerobiology*9, 189–196.” The latter is better suited to reference a statement about airborne bacterial agglomerates, as also cited by Lighthart (2006) itself.*

The referee is right. In the revised manuscript the reference will be changed as suggested.

4. Page 3, line 67: *Tong and Lighthart (1999) does not contain detailed information about particle size. The reviewer suggests to remove or change this reference.*

We will remove the Tong and Lighthart (1999) reference in the revised manuscript.

5. Page 3, line 77: *Bauer et al. (2003) it's not present in the reference section. The only Bauer paper in the reference section is from 2002 and is about carbon content of fungal spores. Said paper does not detail any interaction between bacteria and the atmosphere. The reviewer suggests adjustment to this reference.*

In the revised manuscript, the reference Bauer et al. (2003) will be added.

6. Page 4, line 96-97: *“where transdisciplinary studies gathering [...] issues are possible”. “Gathering issues” is an odd phrasing, the reviewer suggests substituting “gathering” with “addressing” or to better explicitate the meaning of the sentence.*

In the revised manuscript we will substitute the word “gathering” with “addressing” as suggested.

7. Page 6, line 167-168: *what sensor does the PID use to measure concentration?*

The PID (Proportional-Integral-Derivative) controller is a control loop mechanism employing feedback aimed at keeping the value of a certain parameter equal to a pre-defined setpoint value within a defined error. A PID controller is a mathematical construct that is then converted to an algorithm by the use of a software tool (developed by LabView in this case). It continuously calculates an error value as the difference between a desired setpoint (SP) and a measured process variable (PV) and applies correction based on proportional, integral, derivative terms in order to minimize the error on each loop cycle.

PV is the process variable read by a sensor and sampled by the hardware platform connected to the sensor (NI-cRIO, based on Labview program language).

In the first case the PVs are the CO₂ and SO₂ concentration values read as output of the corresponding gas analysers (sensors) connected to the cRIO platform. In the second case the PV is the ChAMBRé internal pressure as output of the pressure gauge (sensor) mentioned in the text.


For the reason above, In the revised manuscript, we will rephrase the sentence as follows:

“Two 5-lpm MFCs are designed for injection of CO₂ and other gases (i.e. SO₂, CO, NO and NO₂), respectively, whose concentration in the chamber can be selected by the operator (ppm or ppb units); a PID (Proportional-Integral-Derivative) controller, using the gas concentration values read from the corresponding gas analyzer, keeps the gas concentration in ChAMBRé constant during the experiment.

A 30-lpm MFC regulates the injection of dry air inside the chamber. In this case, the PID controller (using the ChAMBRé pressure values measured by pressure sensor mentioned above) allows to maintain a pre-defined pressure gap between inside and outside the chamber.”

8. Page 8, Figure 2: *The reviewer has some issues with the presented irradiance spectra. First of all, given that the data are presented over multiple wavelengths, shouldn't the correct unit for the y-axis be $\mu W cm^{-2} nm^{-1}$ as it actually represents a spectral, rather than an integrated, irradiance? The irradiance in the UV-C*

range seems unusually high also for the Sun measured on the terrace of the Genoa Physics Department. Even if it's not clear due to the coarse resolution of the y-axis scale, it seems that UV-C at 200 nm is well above $10 \mu\text{W cm}^{-2} \text{ nm}^{-1}$. That value seems unusually high. As a reference, the reviewer reports here a plot of average spectral solar irradiance measured by the TSIS-1 instrument on the International Space Station between the 1st of January and 06 March 2023. The instrument measures the solar irradiance at the top of atmosphere and normalizes it to 1 AU (i.e.: Sun-Earth distance). This measurement happens before any interaction between solar radiation and ozone happens in the Earth's atmosphere and, therefore, before reduction of any UV radiation. Data were originally in $\text{W m}^{-2} \text{ nm}^{-1}$ but were converted to $\mu\text{W cm}^{-2} \text{ nm}^{-1}$ by multiplying the original data by 100 in order to have units consistent with those presented in the paper. The plot shows the spectrum between 200 and 300 nanometers for better visualization of the range of interest and it shows that even above the Earth's atmosphere the irradiance up to (roughly) 250 nm is well below $10 \mu\text{W cm}^{-2} \text{ nm}^{-1}$. These values are almost constant with time: the plot contains also a shaded area which details the temporal standard deviation (SD), but the shading is not visible as the SD values are close to zero (maximum SD = $0.1263 \mu\text{W cm}^{-2} \text{ nm}^{-1}$).




Data used in this plot are accessible through the following API link: [https://lasp.colorado.edu/lisird/latis/dap/tsis_ssi_24hr.csv?time,wavelength,irradiance&time>=2023-01-01T00:00:00Z&time<=2023-03-07T23:59:59Z&formatTime\(yyyy-MM-dd'T'HH:mm:ss\)](https://lasp.colorado.edu/lisird/latis/dap/tsis_ssi_24hr.csv?time,wavelength,irradiance&time>=2023-01-01T00:00:00Z&time<=2023-03-07T23:59:59Z&formatTime(yyyy-MM-dd'T'HH:mm:ss)) and their description can be found at: https://lasp.colorado.edu/lisird/data/tsis_ssi_24hr

Again the presented data are top of atmosphere, so at the surface of the Earth these UV values should be even lower.

Given that UV-C are the most harmful for biological particles, any excess in this spectral range in the Solar Simulator could severely impact bioaerosols' viability. Due to this reasons, the reviewer kindly asks the author to better explain the Solar Simulator spectra and the measured sun irradiance.

The referee is right. We will change the Y axis unit using $\mu\text{W cm}^{-2} \text{ nm}^{-1}$ in the revised manuscript. Thanks to the detailed comment of the Referee, we found an error in figure 2 reported in the manuscript: not only in the original manuscript the “No filter” irradiance in the UV range is lower than IR filter, AM1.5G filter and sun irradiance (and this is not possible) but also the absolute values are wrong. We revised data from spectrometer, and we found an error in the subtraction of background noise collected in “dark condition” in the UV range (with the optic fiber covered by a black hood). Below the corrected figure will be replaced in the revised manuscript. The irradiance MDL of the spectrometer varies between 3 and 5 $\mu\text{W cm}^{-2} \text{ nm}^{-1}$ and, in the case of measure spectra with Solar Simulator and AM 1.5G filter and sun, it has met below 300 nm.

In conclusion, with the AM1.5G filer and Sun, we don't observe any sizeable irradiance below 300 nm.



9. Page 12, figure 5: What does the error bars represent? Standard deviation? Standard error? Where does this uncertainty comes from? Repeated WBS measurements at the same time?

The bar error is the uncertainty of bacteria concentrations calculated following the Poisson statistics i.e. the standard deviation. We will add this sentence in the revised manuscript.

10. Page 13, line 319: the acronym MISG is explained here, but was already used at page 7, line 173. Please move the acronym explanation at the first occurrence of it in the text.

We will move the acronym explanation at the first occurrence of MISG in the revised manuscript.

11. Page 16, line 420: the acronym OD (which I assume stands for “Optical Depth”) is used but never explained. Please explain it.

We will add the acronym of OD (Optical Density) to the revised manuscript.

12. Pages 17, lines 437-438: it is stated “Data, obtained by CFU counting on agar plates, are averaged and used to figure out the uncertainty”. What’s the uncertainty metric? Standard deviation?

The referee is right, the sentence is not clear. We will modify the sentence in the revised manuscript as: “Data, obtained by CFU counting on agar plates, are weighted averaged and used to figure out the uncertainty (standard error of the weighted mean) of the bacterial concentration in the solution. The weights are the relative uncertainties of CFU number on agar plates following the Poisson statistics.”

13. Page 18, figure 6: What are the error bars? Standard deviation? Standard Error? Also, while for CFU ml⁻¹ the uncertainty should come from the fact that bacteria are spread in duplicates (page 17, line 434), where does the uncertainty of the QTx TOT come from? Was it also measured in duplicates?

The referee is right. We never explained how calculate the uncertainty of QTx TOT. We will add this sentence in the revised manuscript at the Quantum TX section: “To evaluate the uncertainty on the bacteria count (QTx TOT), we repeated the measurement on the same sample 10 times, and we found an instrument repeatability of 5%. This uncertainty is much higher of the statistical error of total counting (assuming the Poisson statistic), and, for this reason, we adopted a 5% uncertainty to all Quantum Tx counts.”

In addition, we will add a sentence under the figure: “Error bars have, in most cases, the same size of the data points and they are calculated as previously described.”

14. Page 18, line 463: “OD600nm and QTx TOT have the same value of the b parameter”. Looking at Table 1 that’s not the case: for OD600nm the value is $(3.3 \pm 0.1) \times 10^{-2}$, while for QTx TOT the value is $(3.4 \pm 0.5) \times 10^{-2}$. If the meaning of the sentence was that the two values are not significantly different, please add a statistical test confirming it.

We will change the phrase in: The b values of OD600nm and QTx TOT are compatible within their uncertainties, and this result is expected since the OD600nm is an indirect measurement of the total concentration of cells in suspension.

15. Page 19, lines 476-477: “the number of the counted colonies are averaged, to retrieve the bacterial concentration in the solution and its statistical uncertainty”. Again, what’s the metric for this uncertainty?

We will revise the part from line 469 to 481 to explain better the preparation and the metric used to calculate the uncertainty of counted colonies. In the revised manuscript this part will be replaced by:

“To prepare the inoculum for the chamber experiments, the *E. coli* is grown in 30 ml of fresh TSB nonselective medium, in a shaking incubator at 37 °C and 200 rpm and its growth is followed by checking the OD600nm value until the mid-exponential phase. When OD600nm ~ 0.5, 20 ml of this liquid preparation is centrifuged at 3000 rpm for 10 min. Afterward, the bacteria pellet, separated by supernatant, is resuspended in 20 ml of sterile physiological solution (NaCl 0.9 % w/v) to prepare a suspension of approximately 10^8 CFU mL⁻¹, as verified by standard dilution plating. To retrieve the bacterial concentration, the average of CFU counting on agar plates and the uncertainty are calculated following the same metric described in 2.3.1 paragraph.

For the experiments performed at ChAMBRé, the typical bacterial concentration in the inoculum is 10^7 CFU ml⁻¹: to reach this concentration, a further dilution step is needed (i.e., typically 1:10 or 1:5) before the injection (see Massabò et al, 2018 for details).”

16. Page 20, line 499-501: The coefficient of variation of the experiments should be moved to the results section and then possibly discussed in terms of reproducibility of ChAMBRé experiments (see also comment #24).

We will move the coefficient of variation to the results section. See the answer of comment 24 about the discussion in terms of reproducibility.

17. Page 20: “Dark” conditions are never explained. Intuitively, it just means with the Solar Simulator turned off, but please explicitate that.

We will clarify the dark condition in the revised manuscript as follows: “solar simulator off” and “solar simulator on” to dark and light condition, respectively.

18. Page 22, Figure 7: in the caption the colours are swapped and wrong: red is indicated for total bacteria and green for viable, while it should be red for viable and blue for total.

The Referee is right: In the caption of figure 7, we will indicate as blue the total bacteria and red viable bacteria trend in the revised manuscript.

*19. Page 22, lines 563-564: “particles, in the size range of 1-2 μm ($\tau=2-3$ hours), the same of *E. Coli*”. Commas make the sentence awkward to read. The reviewer suggests rephrasing to “particles in the same size range of *E. Coli* (1-2 μm) and τ of 2-3 hours”.*

In the revised manuscript, we will rephrase the sentence as suggested by the referee.

20. Page 23, line 573: *The usage of “Then” at the beginning of the sentence is confusing, it seems to imply that dead bacteria after ozone treatment were exposed to NOx. The reviewer suggests to rephrase it starting with “In another experiment ...”.*

We will replace “Then” with “In another experiments” in the revised manuscript.

21. Page 24, Paragraph “Experiments with E.Coli and the Solar Simulator”: *all the results reported here must be discussed in view of the UV-C intensity in the Solar Simulator (see comment #8).*

By considering the answer to comment n. 8, we believe that no additional discussion is needed.

22. Page 24, lines 593-594: *This sentence ending with “and data are here gathered” is not very readable. Please clarify its meaning.*

The referee is right; in the revised manuscript, we will revise the sentence to: “No significant differences in results appeared changing the intensity of the Solar Simulator operated with the AM1.5G filter and the data with a Solar Simulator intensity of 100% are here reported.”

23. Page 24, lines 598-599: *“V:T (...) reaches zero after an hour”. In figure 9, though, V/T appears to be 0 already at the 30 minute mark. What’s the correct result? Also, please keep consistent the format of V:T in both text and figures (either V:T or V/T everywhere).*

The referee is right; in the revised manuscript we will modify the sentence to: “The viable concentration collapses quickly, reaching zero after 30 minutes. The comparison between V:T ratio obtained for dark and light baseline is shown in Figure 9.”

We will correct in the revised manuscript the Y axis of figure 8 and 9 reporting V:T instead of V/T.

24. Page 25, lines 615-617: *The reviewer thinks that reproducibility would need a more in depth discussion. First of all, in the text it is never explicitated the amount of performed replicates, with the exception of the caption of figure 7 where it is stated that eight repetitions were done for the baseline experiments. This point is, instead, extremely important as the reproducibility of the baseline experiments is crucial for the application of ChAMBRe itself. The reviewer encourages the authors to clearly state how the baseline experiments are performed in terms of replicates and how uncertainty and error propagation (if any) is estimated. Furthermore, the sentence here is not very clear, it suggests that the reproducibility (20% in terms of what? How was this reproducibility calculated?) is in line with the experimental sensitivity. What does this mean? That the*

experimental uncertainty is as big as the reproducibility? If so that would imply that any variation of viability could be inputtable to an error in reproducibility. The reviewer strongly suggest to clarify and clearly quantify all the parameters tied to the reproducibility and experimental uncertainty so that the reader can have a clear idea of the uncertainties tied to the ChAMBRé experimental protocol presented in this paper.

The Referee is right: some points are missing (for example the number of experiments as suggested) and the discussion in terms of reproducibility and uncertainties is not always clear and not well described. In the first version of manuscript, we have discussed the results in terms of average and std error of C_0 and τ and these values have been retrieved starting from the averaged results on number of experiments. Thanks to the Referee comment, we have thought that a revised following approach should be a better choice. For this reason, in the revised manuscript, we will add a column to table 3 reporting the number of experiments and we will review the “Results” and “Discussion, conclusion and perspectives” sections describing how we have calculated the uncertainties of C_0 and τ for total and viable *E.coli* concentration and V:T ratio. All data have been reviewed following this procedure:

- For each experiment, C_0 and τ have been retrieved fitting the data with the function $C(t) = C_0 e^{-\frac{t}{\tau}}$;
- In the tables 2 and 3 average and standard deviation of C_0 and τ of all experiments are reported.

The results have been discussed in terms of τ of V:T ratio and its coefficient of variability of baseline experiments compared to the values obtained with gases experiments.

Here the sections we would insert in the revised manuscript:

3.1 Baseline experiments with E. coli in dark conditions.

E. coli behaviour in a set of eight replicated experiments, led from separate cultures, was first determined in dark conditions. The average total concentration and standard deviation of *E. coli* inside the chamber at $t = 0$ (three minutes after the conclusion of the injection to allow proper mixing/homogenization inside the ChAMBRé volume) was (0.34 ± 0.08) cells cm^{-3} , as measured by the WIBS; the average viable concentration and standard deviation, determined by the Andersen impactor sampling at $t = 0$, was (0.04 ± 0.02) cells cm^{-3} . The viable concentration at $t = 0$ was obtained by measuring the CFUs on three petri consecutively sampled; the coefficient of variation on the CFUs collected on the three petri, resulted equal to 12%.

The average ratio and standard deviation of viable:total (V:T in the following) bacteria concentration inside ChAMBRé, at $t = 0$ turned out to be $V:T = (0.13 \pm 0.07)$. The total and viable bacteria concentration values, measured inside ChAMBRé, depended the V:T ratio in the inoculum to be injected (biological effects between each bacteria culture) and on the aerosolization process affecting the bacteria viability. The bacteria viable concentration in the inoculum was determined via standard dilution plating while the bacteria total concentration was calculated by the Quantum Tx. During baseline experiments, the V:T ratio of the inoculum ranged between 0.25 ± 0.03 and 0.50 ± 0.06 . Time-trends of the averaged total and viable concentration of the bacteria, nebulized inside ChAMBRé, are shown in Figure 7. Bacteria lifetime in ChAMBRé can be calculated by fitting the data of each experiment with an exponential function as:

$$C(t) = C_0 e^{-\frac{t}{\tau}} \quad (2)$$

where C_0 is the total or viable concentration of *E. coli* just after the injection ($t = 0$) and τ is the total or viable bacteria lifetime, respectively. In table 2, the average and standard deviation of C_0 and τ for the *E. coli* total and viable concentration of eight experiments are reported.




Figure 7: Time-trend of *E. coli* average bacteria total (blue) and viable (red) concentration inside ChAMBRé obtained by eight repetitions of baseline experiments.

Table 2: C_0 and τ (average \pm std deviation) of the exponential fit for total and viable concentration of *E. coli*.

Exponential function	Total <i>E. coli</i>	Viable <i>E. coli</i>
C_0	(0.33 ± 0.08) cells cm ⁻³	(0.04 ± 0.02) CFU cm ⁻³
τ (min)	153 ± 22	32 ± 5

The total *E. coli* averaged lifetime is about 150 minutes; this value agrees with data reported in (Massabò et al., 2018) for generic aerosols: particles in the same size range of *E. coli* (1-2 μm) and $\tau = 2$ -3 hours. The viable *E. coli* averaged lifetime is about 32 minutes, lower than the aerodynamic lifetime, this indicating the difficulty of this microorganism to survive in the atmospheric medium.

3.2 Experiments with *E. coli* and NO_x in dark conditions.

A preliminary check was performed exposing the *E. coli* to O₃, which is recognized to be a strong antimicrobial agent (Kim et al., 1999; Giuliani et al., 2018; Thanomsub et al., 2022), hence the expected result was a complete viability loss. The exposure of bacteria to O₃ (concentration > 1000 ppb) resulted in a complete cell mortality, as expected. The initial condition immediately after the injection was V:T = (0.03 ± 0.01) and no CFUs were collected in any of the following samplings (starting 30 minutes after the injection).

In another experiments, bacteria were exposed to NO₂ and NO concentrations, 900 and 1200 ppb for both the pollutants. The exposure of bacteria to such pollutants showed a V:T reduction. The average results, obtained in a set of total eight experiments, are shown in Figure 8.




Figure 8: Time-trend of the V:T ratio for *E. coli* in baseline (black) and in the experiments with ChAMBR_e maintained at a constant concentration of: NO₂ (900 ppb red and 1200 ppb blue) and NO (900 ppb dark red and 1200 ppb gray).

The quantitative reduction in the *E. coli* lifetime, due to the exposure to pollutants, can be evaluated considering the V:T ratio and fitting the data with an exponential curve, as previously described; the results are shown in Table 3.

Table 3: Initial values and τ (average and std deviation) of the exponential fit for V:T ratio of *E. coli* at different pollutants concentrations.

Exponential function	(V:T t = 0)	τ (min)	experiments #
----------------------	-------------	--------------	---------------

Baseline	0.13 ± 0.07	40 ± 5	8
NO ₂ 900 ppb	0.11 ± 0.02	25 ± 2	2
NO ₂ 1200 ppb	0.06 ± 0.02	11 ± 2	2
NO 900 ppb	0.05 ± 0.01	26 ± 3	2
NO 1200 ppb	0.10 ± 0.02	25 ± 4	2

E. coli averaged lifetime in baseline experiments, calculated on the V:T ratio, turned out to be about 40 min. The exposure of *E. coli* to NO₂ reduced the lifetime to about 25 and 11 min with a concentration of 900 ppb and 1200 ppb respectively. The exposure to 900 ppb and 1200 ppb of NO decreased bacteria lifetime to 26 and 25 min, respectively and the values are similar to the value obtained with the lowest NO₂ concentration. The increase of the NO concentration did not correspond to a decrease in the *E. coli* viability, as observed with NO₂: these results suggest a greater toxic effect of NO₂ than of NO on *E. coli*. The literature of a comparison of the toxic effects of NO and NO₂ on *E. coli* is poor. Some research articles have demonstrated negative effects of these two gases on bacterial strains: Kosaka et al. 1986 found a decrease in *E. coli* viability with increasing NO₂ concentration. Janvier et al, 2020 highlighted a significant adverse effect of NO₂ on some commensal skin bacterial strains. Mancinelli and McKay, 1983 found that a low concentration of NO is bacteriostatic for some organisms but not for others. It is worth noting that NO has a strong antimicrobial property, being an endogenously produced molecule that is critical for critical infection defence (Fang, 1997), although some bacteria are able to escape this NO action (Privett et al., 2012).

4 Discussion, conclusion and perspectives

The main result presented in this work is the assessment of a multi-step and well controlled protocol to perform experiments on the impact of air quality on bacteria viability by an atmospheric simulation chamber, ChAMBRé in this case. Even if the chamber configuration is still in progress and several new equipment will be deployed at ChAMBRé in the near future, the present set-up opens the possibility of systematic studies. The average τ of the V:T ratio of eight baseline experiments was 40 min with a standard deviation of 5 min; the coefficient of variation of 13% corresponds to the experimental sensitivity to changes in *E. coli* viability due to exposure to pollutants and/or other relevant parameters. The baseline reference must be experimentally determined for each bacteria strain and efforts are planned for repeating the observation with *Bacillus subtilis*, *Bacillus spizizenii* and *Pseudomonas fluorescens* in the near future. It is worthy to note that the experimental protocol returns the lifetime of total and viable bacteria injected in the chamber. The figure for total bacteria corresponds to the aerodynamic behavior of aerosol of diameter around 1 μm , already reported in (Massabò et al., 2008) while the lifetime of viable bacteria is much shorter (about half an hour) due to the difficulty of this microorganism to survive in the atmospheric medium. Such shorter lifetime posed clear constraints on the first experiments with exposure of *E. coli* to NO_x inside ChAMBRé. A time window of two hours after the bacteria injection was considered to observe the behaviour of *E. coli* viability and it was possible to quantify a lifetime reduction, in dark conditions, clearly related to NO and NO₂ concentration inside ChAMBRé. These findings

pave the road to systematic studies including other bacteria strains and pollutant species. In the near future With the *E. coli* exposed to the light produced by the Solar Simulator operated with the AM1.5 filter, the viability resulted very short even in the baseline conditions and therefore no further experiment with pollutants was performed. With other bacterial strains, the impact of light on viability will have to be reinvestigated. It is well known in the literature that the viable but non-culturable condition (VBNC) is a survival strategy of many bacteria in the environment in response to adverse environmental conditions (e.g., solar radiation). There is a growing scientific interest in studying VBNC cells, including to understand novel public health implications of VBNC cells. In our simulated experiments, we are investigating alternative methods to detect bacterial viability and VBNC state, such as “live and dead staining” by fluorescence microscopy. This assay can be used to monitor the viability of bacterial populations as a function of cell membrane integrity using different fluorescent dyes.

Further experiments with “flow cytometry” could certainly be more beneficial not only to enumerate live and dead bacteria, but also to evaluate the health and viability of bacterial cells by determining the activity of bacterial oxidases and reductases.

References

- Fang, F. C.: Perspectives series: host/pathogen interactions. Mechanisms of nitric oxide-related antimicrobial activity, *J Clin Invest.* 99(12): 2818–2825, <https://doi.org/10.1172%2FJCI119473>, 1997.
- Janvier, X., Alexandre, S., Boukerb, A. M., Souak, D., Maillot, O., Barreau, M., Gouriou, F., Grillon, C., Feuilloley, M. G. J., Groboillot, A.: Deleterious Effects of an Air Pollutant (NO₂) on a Selection of Commensal Skin Bacterial Strains, Potential Contributor to Dysbiosis?, *Frontiers of in Microbiology*, Volume 11, Article 591839, <https://doi.org/10.3389/fmicb.2020.591839>, 2020.
- Kosaka, H., Yamamoto, H., Oda, Y., Uozumi, M: Induction of SOS functions by nitrogen dioxide in *Escherichia coli* with different DNA-repair capacities, *Mutat Res. Aug;* 162(1):1-5, doi: 10.1016/0027-5107(86)90065-5, 1986.
- Mancinelli, R. L. and McKay, C.P.: Effects of Nitric Oxide and Nitrogen Dioxide on Bacterial Growth, *Applied and Environmental Microbiology*, 198-202, <https://doi.org/10.1128/aem.46.1.198-202.1983>, 1983.
- Privett, B. J., Broadnax, A. D., Bauman, S. J., Riccio, D. A., Schoenfisch, M. H.: Examination of bacterial resistance to exogenous nitric oxide, *Nitric Oxide*, Volume 26, Issue 3, Pages 169-173, <https://doi.org/10.1016/j.niox.2012.02.002>, 2012.

Answers to the comments of Referee 2

1. *Abstract: I think that it would be useful to indicate the mean residence/life times estimated for E. coli (baseline conditions).*

The Referee is right. We will add this information in the abstract of the revised manuscript.

2. *The manuscript is likely aimed at non-microbiologists, and so very basic protocols for culturing bacteria and characterizing their growth and numbers are given in great detail. I leave these to authors' choice, but these can be found in any handbook of basic microbiology that could eventually be referred to in order to shorten the manuscript. Examples include lines 401-419 regarding bacterial growth (which appeared in addition quite out-of-subject here), or lines 422-438 regarding preparation of cell suspension and estimation of viable cell numbers by dilution-plating. I thought that Figure 6 as well is unnecessary in the context of the study and could be put in supplement. Section 2.3.1 could therefore ultimately be greatly shortened and merged with another section (e.g. 2.3.2).*

The Referee is right, the manuscript is aimed at non-microbiologists, therefore we have decided to describe the protocols for culturing bacteria in detail, considering useful this part to the potential reader. We would prefer to keep this part in the main text; however, we leave the decision to the Editor (to maintain this section in the manuscript or move to supplement).

3. *Lines 28-29 (abstract): the sentence "This figure quantifies the protocol sensitivity as well to changes in viability when bacteria are exposed in other (e.g., polluted) conditions" is not clear (which figure? Conditions other than what?). In addition, "exposed in" should be "exposed to".*

The Referee is right, the sentence is not clear. We will modify the sentence in the revised manuscript as follows:

"The whole procedure showed an estimated life time of total (T) and viable (V) E. coli of about 153 and 32 minutes, respectively, and a V:T ratio lifetime of 40 ± 5 minutes when ChAMBRé is held in a reference "baseline" condition. The coefficient of variation of 13% shows how sensitive the protocol is also to changes in viability when the bacteria are exposed to other (e.g., polluted) conditions."

4. *Lines 43-45: DMS and other VOCs are given as examples of SBAs, but these are not aerosols! Literally, SBA would correspond to fragments of larger biological particles, material released from cells (disruption, excretion...), nucleated biogenic gases, or cells "born" in the air from microbial multiplication.*

We will modify the sentence in the revised manuscript as follows: “SBA are fragments of larger biological particles, material released from cells (disruption, excretion...), nucleated biogenic gases, or cells “born” in the air from microbial multiplication (Morris et al., 2014, Ervens et Amato, 2020).”

5. *Lines 46 and 47: the two “can” are not necessary. Moreover, do they really “vary” in size (at the individual scale)? Or do they rather fall within a range of sizes?*

We will modify the sentence in the revised manuscript as follows: “The PBAs vary in size depending on the specific biological material being aerosolized; they range from several nanometers (e.g., viruses, cell fragments) to a few hundred micrometers in aerodynamic diameter (e.g., pollen, plant debris) (Pöschl, 2005).”

6. *Line 57: It is said that bacteria are at concentration “usually greater than 10⁴ cell/m³” over land. The reference Bauer et al 2002 is not appropriate here for such statement. Rather cite the review by (Burrows et al., 2009) where Table 1 summarizes well literature data as a function of land use. In addition, the statement that concentrations in air over sea tend to be lower needs a reference, and it is not fully supported by literature (see for instance (Mayol et al., 2014), Table 1). This whole sentence thus needs revision.*

The Referee is right. In the revised manuscript, we will replace Bauer et al 2002 with Burrows et al., 2009. The sentence will be revised and replaced with: “They are ubiquitous in the atmosphere, and their presence and abundance can vary depending on factors such as location, season, and local environmental conditions: usually, over the land, the concentration in atmosphere is greater than 10⁴ cells m⁻³ (Burrows et al., 2009) while our understanding of airborne microbes over oceans, is indeed limited compared to the knowledge we have about microbes in terrestrial and aquatic environments. In a recent work (Mayol et al., 2014), the airborne prokaryotic abundance over the North Atlantic ocean ranged from about 3000 to 20000 prokaryotes m⁻³ (average about 8000 cells m⁻³).”

7. *Lines 61-62: “Bacteria have a relatively long atmospheric residence time, of the order of several days or more, compared to larger particles and can be transported over long distances, up to thousands of km (Després et al., 2012)” is barely a plagiarism of the cited reference, that says: “Due to their small size, bacteria have a relatively long atmospheric residence time (on the order of several days or more) compared to larger particles and can be transported over long distances (up to thousands of kilometres).” This may need rephrasing.*

We will rephrase the sentence in the revised manuscript as follows: “Bacteria, as small airborne particles or aerosols, can have relatively long atmospheric residence times compared to larger particles. This is due to their small size and low settling velocity, which allows them to remain suspended in the air for prolonged periods. (Després et al., 2012).”

8. *Lines 69-72: Is this statement about the lack of data on abundance and diversity of microbes in the air really relevant here? It would be more helpful to develop a bit on the subject of the paper: viability, and the impacts of environmental conditions, for which even more data is lacking.*

In the revised manuscript, we will add a sentence in line 78 as follows: “In particular, bacterial viability, the proportion of viable to total bacteria concentration, can act as Cloud Condensation Nuclei (CCN) thanks to the hygroscopic properties of their surfaces (Delort et al., 2010). Additionally, the near-surface atmosphere's viable bacteria can have a significant impact on human health, including allergies, acute toxic effects, and infections (Bolashikov and Melikov 2009).”

9. *Line 79-80: I would suggest to use more recent references here about the potential impacts on cloud chemistry, such as (Khaled et al., 2021; Jaber et al., 2021, 2020; Fankhauser et al., 2019).*

In the revised manuscript, we will replace the original references with those suggested by the Referee.

10. *Lines 86-88: what is “direct deposition”, and how does this differ from precipitation? Do you rather mean “wet or dry deposition”. Also, can you develop what “multifaceted” means here when referring to the mechanisms that govern transport, survival and activity. Eventually cite (Amato et al., 2023) which discusses the fact that biologically- and physically-driven processes are indeed intertwined.*

In the revised manuscript, we will rephrase the sentence as follows:

“Once in the air, they are carried upwards by air currents and may remain in the atmosphere for many days before being removed by wet or dry deposition onto surfaces. Indeed, the mechanisms that govern the transport, survival, and activity of bacteria in the atmosphere are complex and multifaceted. Understanding these mechanisms is crucial for various scientific disciplines, including microbiology, atmospheric science, and public health. This complexity is related to some key factors such as aerosolization, transport and dispersion, survival, hygroscopicity, interactions with other particles, droplet nucleation, deposition, activation of ice nucleation, impacts on cloud formation and chemistry and all these processes are indeed intertwined (Amato et al., 2023).”

11. *Line 88-90: what kind of impacts are in question here?*

We will add the words climate and health in the revised manuscript.

12. *Line 98-104: In addition of the references cited, I think that a number of other (and early) studies that have investigated the survival and activity of bacteria using simulation chambers should be referenced: (Ehrlich et al., 1970; Krumins et al., 2014; Wright et al., 1969).*

In the revised manuscript, we will add the sentence as follows: “Few studies have investigated bacterial survival and activity using simulation chambers, and some of them are old (Wright et al., 1969; Ehrlich et al., 1970; Krumins et al., 2014).”

13. *Line 177: provide the reference of the nebulizer inlet please.*

In the revised manuscript, we will add the reference at the paragraph 2.2.

14. *Line 180: I do not see the point of mentioning “(or in the laboratory)”. Please specify (or remove).*

In the revised manuscript, we will remove “in the laboratory”.

15. *Line 342: quartz filters can be used for offline analysis. Please consider that this is also the case with other sampling methods (impingers and cultures cannot be considered “online” methods).*

The referee is right: in the revised manuscript, at the line 327 we will add the following sentence: “All the methods described below allow to perform offline analyses”. We will remove “allowing subsequent offline laboratory analysis” from line 338 and “for offline analysis” from line 342.

16. *Lines 351-352: “The hood is equipped with HEPA filter and UV-light laminar is created inside the cabinet” probably needs rephrasing (what is UV-light laminar?).*

In the revised manuscript, we will change the sentence as follows: “The hood is equipped with HEPA filter and an UV-light lamp allows the sterilization of the illuminated surfaces inside the hood.”

17. *Line 358: specify that the spectrophotometer is dedicated to liquid samples, to avoid confusion.*

In the revised document, we will add the sentence: “designed for liquid samples.”

18. Line 364: *same as previous comment, regarding the shaker: it might not be obvious for non-microbiologists that these are intended for liquid cultures.*

In the revised document, we will add the sentence: “designed for liquid samples.”

19. Line 446: *“the results of the logistic fit are only shown” might rather read “the results of the logistic fit only are shown”.*

We will apply the correction to the revised manuscript.

20. Line 472: *“centrifugated” should be centrifuged.*

We will apply the correction to the revised manuscript.

21. Line 468-481: *this is quite useless as presented as many details are missing: what is the volume of culture used? And that volume of liquid were cells resuspended into after centrifugation? Dilution step from what (line 480)?*

We will revise the part from line 469 to 481 to explain better the preparation and providing more details. In the revised manuscript this part will be replaced by:

“To prepare the inoculum for the chamber experiments, the *E. coli* is grown in 30 ml of fresh TSB nonselective medium, in a shaking incubator at 37 °C and 200 rpm and its growth is followed by checking the OD_{600nm} value until the mid-exponential phase. When OD_{600nm} ~ 0.5, 20 ml of this liquid preparation is centrifuged at 3000 rpm for 10 min. Afterward, the bacteria pellet, separated by supernatant, is resuspended in 20 ml of sterile physiological solution (NaCl 0.9 % w/v) to prepare a suspension of approximately 10⁸ CFU mL⁻¹, as verified by standard dilution plating. To retrieve the bacterial concentration, the average of CFU counting on agar plates and the uncertainty are calculated following the same metric described in 2.3.1 paragraph.

For the experiments performed at ChAMBRé, the typical bacterial concentration in the inoculum is 10⁷ CFU ml⁻¹: to reach this concentration, a further dilution step is needed (i.e., typically 1:10 or 1:5) before the injection (see Massabò et al, 2018 for details).”

22. Line 489 *“relative standard deviation”* and Lines 500-501 *“the ratio between standard deviation and mean”*: *aren't these actually coefficients of variation?*

In the revised manuscript we will replace “relative standard deviation” and “the ratio between standard deviation and mean” with coefficient of variation.

23. Line 546: *nebulized*

The correction will be done in the revised manuscript.

24. Line 547: *Indicate if these are true independent replicates of experiments (i.e. led from separate cultures? So somehow accounting for biological variability), or just consecutive repeats from same cultures.*

Thanks for the comment: we will add the sentence “led from separate cultures” in the Results section in the revised manuscript.

25. Line 562-565: *check language (“the same of”, “this indicating”).*

In the revised manuscript we will rephrase the sentence as follows: “particles in the same size range of E.Coli (1-2 μm) and τ of 2-3 hours”.

26. Line 570: *what is a “roughly complete cell mortality”? It is either complete, or incomplete...*

In the revised manuscript we will delete “roughly” word since we had a complete cell mortality.

27. Table 3?

We don't understand the request of referee.

28. Lines 588-589: *NO is less toxic than NO2. It would be nice to develop on this (in the discussion section): why could this be? what are the processes of toxicity of these compounds? What does this imply for natural situations? What is known about it?*

In the revised manuscript, we plan to go deeply into the discussion of toxicity effect of NO and NO₂ on *E.coli*. Below, the phrases we would like to add:

“The increase of the NO concentration did not correspond to a decrease in the *E. coli* viability, as observed with NO₂: these results suggest a greater toxic effect of NO₂ than of NO on *E. coli*. The literature of a comparison of the toxic effects of NO and NO₂ on *E coli* is poor. Some research articles have demonstrated negative effects of these two gases on bacterial strains: Kosaka et al. 1986 found a decrease in *E.coli* viability with increasing NO₂ concentration. Janvier et al, 2020 highlighted a significant adverse effect of NO₂ on some commensal skin bacterial strains. Mancinelli and McKay, 1983 found that a low concentration

of NO is bacteriostatic for some organisms but not for others. It is worth noting that NO has a strong antimicrobial property, being an endogenously produced molecule that is critical for critical infection defence (Fang, 1997), although some bacteria are able to escape this NO action (Privett et al., 2012).

29. *Line 598: “30 min”, or less, given the data shown. This is difficult to tell from Figure 9, maybe showing it on log-scale would be more appropriate? (using the 1/10 detection limit to represent “0” for instance).*

The Referee is right. In the revised manuscript we will rephrase the sentence as follows: “The viable concentration collapses quickly, reaching zero after 30 minutes. The comparison between V:T ratio obtained for dark and light baseline is shown in Figure 9.”

30. *Line 600: the reference should be “Figure 9 “here.*

In the revised manuscript we will change the reference.

31. *Lines 614 and 621: “next future” probably should read “near future”.*

We will modify “next future” in “near future” in the revised manuscript.

32. *The perspectives section could be developed based on other sampling possibility. Notably the possibility of using methods other than cultures to evaluate viability would be relevant if this is the case (live/dead staining, RT-qPCR, and others), in particular as non-culturable state is common in viable bacteria (as mentioned line 608). Moreover, neither SMPS data nor their potential use are exploited. It could be mentioned for instance that investigations about the relationship between particle size and survival could be performed, using cascade impactors as samplers for instance.*

In the revised manuscript, we will add the following sentence:

“It is well known in the literature that the viable but non-culturable condition (VBNC) is a survival strategy of many bacteria in the environment in response to adverse environmental conditions (e.g., solar radiation). There is a growing scientific interest in studying VBNC cells, including to understand novel public health implications of VBNC cells. In our simulated experiments, we are investigating alternative methods to detect bacterial viability and VBNC state, such as “live and dead staining” by fluorescence microscopy. This assay can be used to monitor the viability of bacterial populations as a function of cell membrane integrity using different fluorescent dyes.

Further experiments with “flow cytometry” could certainly be more beneficial not only to enumerate live and dead bacteria, but also to evaluate the health and viability of bacterial cells by determining the activity of bacterial oxidases and reductases.”

References cited and added in the revised manuscript:

Amato, P., Mathonat, F., Nuñez Lopez, L., Péguilhan, R., Bourhane, Z., Rossi, F., Vyskocil, J., Joly, M., and Ervens, B.: The aeromicrobiome: the selective and dynamic outer-layer of the Earth’s microbiome, *Frontiers in Microbiology*, 14, <https://doi.org/10.3389/fmicb.2023.1186847>, 2023

Bauer, H., Giebl, H., Hitzenberger, R., Kasper-Giebl, A., Reischl, G., Zibuschka, F., Puxbaum, H.: Airborne bacteria as cloud condensation nuclei, *Journal of Geophysical Research*, vol. 108, no. D21, 4658, [doi:10.1029/2003JD003545](https://doi.org/10.1029/2003JD003545), 2003.

Delort, A. M., Vaïtilingom, M., Amato, P., Sancelme, M., Parazols, M., Mailhot, G., Laj, P., Deguillaume, L.: A short overview of the microbial population in clouds: Potential roles in atmospheric chemistry and nucleation processes, *Atmospheric Research*, Volume 98, Issues 2–4, Pages 249-260, <https://doi.org/10.1016/j.atmosres.2010.07.004>, 2010

Ehrlich, R., Miller, S., and Walker, R. L.: Relationship Between Atmospheric Temperature and Survival of Airborne Bacteria, *Appl Microbiol*, 19, 245–249, 1970.

Ervens, B. and Amato, P.: The global impact of bacterial processes on carbon mass. *Atmos. Chem. Phys.*, 20, 1777–1794, <https://doi.org/10.5194/acp-20-1777-2020>, 2020.

Fang, F. C.: Perspectives series: host/pathogen interactions. Mechanisms of nitric oxide-related antimicrobial activity, *J Clin Invest*. 99(12): 2818–2825, <https://doi.org/10.1172%2FJCI119473>, 1997.

Fankhauser, A. M., Antonio, D. D., Krell, A. M., Alston, S. J., Banta, S., and McNeill, V. F.: Constraining the impact of bacteria on the aqueous atmospheric chemistry of small organic compounds, *ACS Earth Space Chem.*, <https://doi.org/10.1021/acsearthspacechem.9b00054>, 2019.

Jaber, S., Lallement, A., Sancelme, M., Lereboure, M., Mailhot, G., Ervens, B., and Delort, A.-M.: Biodegradation of phenol and catechol in cloud water: comparison to chemical oxidation in the atmospheric multiphase system, *Atmospheric Chemistry and Physics*, 20, 4987–4997, <https://doi.org/10.5194/acp-20-4987-2020>, 2020.

Jaber, S., Joly, M., Brissy, M., Lereboure, M., Khaled, A., Ervens, B., and Delort, A.-M.: Biotic and abiotic transformation of amino acids in cloud water: experimental studies and atmospheric implications, *Biogeosciences*, 18, 1067–1080, <https://doi.org/10.5194/bg-18-1067-2021>, 2021.

Janvier, X., Alexandre, S., Boukerb, A. M., Souak, D., Maillot, O., Barreau, M., Gouriou, F., Grillon, C., Feuilloley, M. G. J., Groboillot, A.: Deleterious Effects of an Air Pollutant (NO₂) on a Selection of Commensal Skin Bacterial Strains, Potential Contributor to Dysbiosis?, *Frontiers of in Microbiology*, Volume 11, Article 591839, <https://doi.org/10.3389/fmicb.2020.591839>, 2020.

Khaled, A., Zhang, M., Amato, P., Delort, A.-M., and Ervens, B.: Biodegradation by bacteria in clouds: an underestimated sink for some organics in the atmospheric multiphase system, *Atmospheric Chemistry and Physics*, 21, 3123–3141, <https://doi.org/10.5194/acp-21-3123-2021>, 2021.

Kosaka, H., Yamamoto, H., Oda, Y., Uozumi, M.: Induction of SOS functions by nitrogen dioxide in *Escherichia coli* with different DNA-repair capacities, *Mutat Res. Aug*; 162(1):1-5, doi: 10.1016/0027-5107(86)90065-5, 1986.

Krumins, V., Mainelis, G., Kerkhof, L. J., and Fennell, D. E.: Substrate-Dependent rRNA Production in an Airborne Bacterium, *Environ. Sci. Technol. Lett.*, 1, 376–381, <https://doi.org/10.1021/ez500245y>, 2014.

Mancinelli, R. L. and McKay, C.P.: Effects of Nitric Oxide and Nitrogen Dioxide on Bacterial Growth, *Applied and Environmental Microbiology*, 198-202, <https://doi.org/10.1128/aem.46.1.198-202.1983>, 1983.

Mayol, E., Jiménez, M. A., Herndl, G. J., Duarte, C. M., and Arrieta, J. M.: Resolving the abundance and air-sea fluxes of airborne microorganisms in the North Atlantic Ocean. *Frontiers in Microbiology*, 5, <https://doi.org/10.3389/fmicb.2014.00557>, 2014.

Privett, B. J., Broadnax, A. D., Bauman, S. J., Riccio, D. A., Schoenfisch, M. H.: Examination of bacterial resistance to exogenous nitric oxide, *Nitric Oxide*, Volume 26, Issue 3, Pages 169-173, <https://doi.org/10.1016/j.niox.2012.02.002>, 2012.

Wright, D. N., Bailey, G. D., and Goldberg, L. J.: Effect of Temperature on Survival of Airborne *Mycoplasma pneumoniae*, *J Bacteriol*, 99, 491–495, 1969.