

Atmospheric teleconnections between the Arctic and the Baltic Sea region as simulated by CESM1-LE

Erko Jakobson¹ and Liisi Jakobson¹

¹Tartu Observatory of University of Tartu, Tartu, Estonia

5 *Correspondence to:* Erko Jakobson (erko.jakobson@ut.ee)

Abstract. This paper examines teleconnections between the Arctic and the Baltic Sea region and is based on two cases of CESM-LE climate model simulations¹: the stationary case with pre-industrial radiative forcing and the climate change case with ~~measured and~~ RCP8.5 radiative forcing.

10 Stationary control simulation 1800-year long time-series were used for stationary teleconnection and 40-member ensemble from the period 1920–2100 for teleconnections during ongoing climate change. We ~~analyzed~~analysed seasonal temperature at a 2-meter level, sea-level pressure, sea ice concentration, precipitations, geopotential height and 10-meter level wind speed. The Arctic was divided into seven areas.

~~the~~The Baltic Sea region climate has strong teleconnections with the Arctic climate; the strongest connections are with Svalbard and Greenland region. There is high seasonality in the teleconnections, with the strongest correlations in winter and 15 the lowest correlations in summer, when the local meteorological factors are stronger. ~~The majority of NAO and AO climate indices can explain most~~ teleconnections in winter and spring ~~can be explained by climate indexes NAO and AO.~~ During ongoing climate change, the teleconnection patterns did not show remarkable developments~~changes~~ by the end of the 21st century. Minor pattern changes are between the Baltic Sea region temperature and the sea ice concentration.

~~To~~We calculated the correlation between the parameter and its Ridge regression estimation to estimate different Arctic regions' 20 collective statistical connections with the Baltic Sea region, ~~we calculated the correlation between the parameter and its Ridge regression estimation.~~ Seasonal coefficient of determination, R^2 , were highest for winter: for temperature~~T2m~~ $R^2 = 0.64$, for surface pressure~~SLP~~ $R^2 = 0.44$ and for precipitation~~PREC~~ $R^2 = 0.35$. When doing the same for the seasons' previous month values in the Arctic, the relations are considerably weaker, with the highest $R^2 = 0.09$ for temperature in the spring. Hence, ~~the Arctic climate data~~ forecasting capacity ~~of Arctic climate data~~ for the Baltic Sea region is weak.

25 Although there are statistically significant teleconnections between the Arctic and Baltic Sea region, the Arctic impacts are regional and mostly connected with climate indexes. There are no simple cause-and-effect pathways. By the end of the 21st century, the Arctic ice concentration has significantly decreased. Still, the general ~~teleconnections~~ pattern~~teleconnection patterns~~ between the Arctic and the Baltic Sea region will not change considerably by the end of the 21st century.

1 Introduction

30 The Arctic region is warming at least twice ~~as fast as the whole planet~~ (IPCC, 2021; Nakamura ~~&~~ Sato, 2022; Overland et al., 2018; Meleshko et al., 2020); ~~), some authors showed that nearly four times (Rantanen et al., 2022) as fast as the whole planet~~; also, the Baltic Sea region is warming faster than the global average (BACC, 2015). The question of the faster warming in the Arctic region affecting mid-latitudes has been under debate for a long time. We want to address the Baltic Sea region and find out if it is affected by the changing Arctic and if this information can be used in long-term weather forecasts.

35 The faster warming in the Arctic compared to the global mean, a phenomenon known as Arctic amplification (AA), ~~results is a result~~ from ~~a number of positive feedbacks that are relatively large in the Arctic, including the interacting processes: sea ice loss and surface~~ albedo feedback, ~~lapse rate feedback and Planck feedback~~ (Lainé et al., 2016; Yoshimori et al., 2014; Serreze et al., 2009; Screen and Simmonds 2010), ~~changes in longwave and/or temperature feedbacks (Dai and Jenkins, 2023; Jenkins and Dai, 2021; Duan et al., 2019; Pithan and Mauritzen 2014), 2014; Lu and Cai, 2009), cloud changes (Taylor et al., 2022;~~
40 ~~Boeke and Taylor, 2018; Taylor et al., 2015; Taylor et al., 2013; Francis and Hunter, 2006; Vavrus, 2004), intraseasonal cycling of heat (Clark et al., 2021; Bintanja and Krikken, 2016; Bintanja and Linden, 2013), and poleward energy transport (Sang et al., 2022; Spielhagen et al., 2011).~~ However, the relative weight of these different factors is still under debate (Taylor et al., 2022; Dai et al., 2019).

The assessment of the potential for AA to influence broader hemispheric weather (referred to as linkages) is complex and
45 controversial (Dai and Song, 2020; Francis and Vavrus, 2015; Barnes and ~~Sereens~~Screen, 2015; Sun et al., 2016) and many details of linkage mechanisms remain elusive (Sun et al., 2018).

~~For a long time, it has been recognised that ice conditions in the Greenland region might be connected to several variables in Europe (Hildebrandsson, 1914; Wiese, 1924; Schell, 1956; etc.). Zhuo et al. (2023) found that the Greenland region is the most influential of all Arctic regions on teleconnections to the Baltic region. Deng et al. (2018) showed that the Greenland region~~
50 ~~was the most important of Arctic regions considering heat waves in Eastern Europe.~~

AA is expected to be related to further changes that affect mid-latitudes and the rest of the world (Jung et al., 2015; Vihma et al., 2019). According to Overland et al. (2015), potential Arctic teleconnections with Europe are less clear than with North America and Asia. The linkages between the Arctic and mid-latitudes depend strongly on the season and geographical region
(Zhuo et al., 2023; Coumou et al., 2018; Jakobson et al., 2017; ~~Coumou et al., 2018).~~ Furthermore, it has been
55 ~~reeognized~~recognised that extratropical impacts depend highly on the regional structure of the anomalous Arctic climate state (Kug et al., 2015). It appears that Arctic impacts will be regional and intermittent, clouding the identification of cause-and-effect and raising the issue of how to effectively communicate potential Arctic impacts (Overland et al., 2021; Rudeva and Simmonds, 2021; Luo et al., 2019; Cohen et al., 2018).

The Baltic Sea region is very sensitive to climate change; it is a region with spatially varying climate and diverse ecosystems
60 (Christensen et al., 2022). Climate change may bring profound ecological changes in the region (Halkka, 2022). During the last half-century, the duration of ~~permanents~~seasonal snow cover and snow depth have decreased (Viru and Jaagus, 2020);

during the last decades, there has been a major increase in both extreme mild ice winters and severe ice winters; a minor increase in the intense precipitations, heat waves and cold spells (Rutgersson et al., 2022). Because of the closeness to the Arctic, the Baltic Sea region receives influences from the Arctic either remotely (teleconnections) or directly. The weather in the region depends highly on the position of the polar front: it can be located northward as well as southward of the area (Jakobson et al., 2017).

~~One of the reasons scientists are so interested in knowing the Arctic and its influence on mid-latitude weather is that there is a possibility to glance to the future. The pace of climate change is a crucial metric in risk assessment surrounding future climate predictions and directly impacts the efforts required to adapt (Rondeau-Genesse and Braun 2019; Lehner et al., 2017). The planning processes of decision makers are very limited if the pace of changes is unknown (Klein et al., 2014). However, for short timescales, this pace can be masked by internal variability. Over a few decades, this can cause climate change effects to exceed what would be expected from the greenhouse gas (GHG) emissions alone or, to the contrary, cause slowdowns or even hiatuses (Rondeau-Genesse and Braun, 2019). Slowdowns, temporary hiatuses, and negative trends can be expected, regardless of the spatial scale (Medhaug et al., 2017; Fyfe et al., 2016). However, cooling trends should become less common or even nonexistent (Risbey et al., 2017; Grenier et al., 2015; Kay et al., 2015). At the same time, extremely warm decades should become more likely to happen (Risbey et al., 2017; Chavaillaz et al., 2016).~~

Furthermore, some direct impacts are influenced by remote processes in the Arctic. For example, possibly the Barents and Kara Seas warming associated with the sea ice loss affects the Ural blocking (Peings et al., 2023; Yao et al., 2017; Luo et al., 2017), which has been identified as precursors of sudden stratospheric warmings (Statnaia et al., 2020; Lee et al., 2019; Martius et al., 2009), and extreme temperature/precipitation anomalies over Europe (Yang et al., 2022; Peings, 2019; Cattiaux et al., 2010).

Widely used climate model simulations, such as those from the Coupled Model Intercomparison Project (~~CMIP5~~CMIP6), combines internal and inter-model variability caused by differing physics, dynamical cores, and resolutions, making it almost impossible to assess the portion of uncertainty caused by internal variability alone (Kay et al., 2015). To enable quantification of internal variability in the midst of transient climate change, large ensembles with individual models have been performed.

Comparison across ensemble members ~~that are~~ simulated with the same model and ~~the same~~ external forcing provides a measure of simulated internal variability. Here, we use the Community Earth System Model version 1 large ensemble (CESM-LE; Kay et al., 2015) to diagnose connections between the Arctic and Baltic regions. In the last few years, CESM-LE has aimed at better understanding internal variability (Rondeau-Genesse and Braun, 2019). The CESM-LE has been used in multiple studies of Arctic sea ice cover, performing well overall (Smith and Jahn, 2019; Labe et al., 2018; Jahn, 2018; Massonnet et al., 2018; Barnhart et al., 2016; Jahn et al., 2016; Swart et al., 2015). The CESM-LE also produces credible NAO interannual aspects, given the length of the observational record available for assessment (93 years) (Deser et al., 2017).

Our previous research studied Arctic-Baltic teleconnections and physical mechanisms behind Arctic-Baltic teleconnections (Jakobson et al., 2017), using ERA-Interim and NCEP-CFSR ~~reanalyzes~~reanalyses. In this paper, we used the CESM-LE model time-series to verify previous results, ~~search~~examine climate ~~change~~change's influence on teleconnections and study

different Arctic regions' collective forecasting capabilities. CESM-LE differs from ~~reanalyzesreanalyses~~ by a much longer time-series and the aspect that it is not constrained by observations, allowing projections. The purpose of the study is to understand which Arctic factors influence the Baltic Sea, how strong these connections are, how is AA affecting these relationships, and if the knowledge can be used in long-term weather forecasting ~~for next some months~~ in the Baltic Sea region.

The paper is ~~organizedorganised~~ as follows: Section 2 describes the used datasets and methodology. Section 3 explains the results of the spatial correlations of climatic variables (stationary, 20-year periods up to the year 2100 and lagged correlations), whereas Section 4 provides a discussion of the results and ~~completes with~~ conclusions.

2 Data and methodology

We used the CESM Large Ensemble Project (CESM-LE) set of climate model simulations on a $1^\circ \times 1^\circ$ horizontal grid (Kay et al., 2015). The CESM1 is a fully coupled model ~~as~~ described by Hurrell et al. (2013). A control simulation under pre-industrial (1850) radiative forcing conditions was run for 1800 years. A single ensemble member was branched from this control and ran from 1850 to 1920 with transient forcing. A 40-member ensemble was then performed for the period 1920-2100. All 40 CESM-LE ensemble members use the same model and the same external forcing. Each ensemble member has a unique climate trajectory because of small round-off level differences in their initial atmospheric conditions. Simply put, the CESM-LE ensemble spread results from internally generated climate variability alone (Kay et al., 2015). Each member is subject to the same radiative forcing scenario (historical up to 2005 and RCP8.5 thereafter).

We used Pearson's correlation coefficient to measure ~~of the~~ dependence between two variables. To measure the strength of a relationship between two variables without the possible controlling effect of a third variable, we used partial correlation:

$$R(X, Y|Z) = \frac{R(X, Y) - R(X, Z) \cdot R(Y, Z)}{\sqrt{(1 - R^2(X, Z)) \cdot (1 - R^2(Y, Z))}} \quad (1)$$

where $R(A, B)$ is the regular Pearson correlation. Partial correlation difference from Pearson correlation reveals the controlling factor Z effect on input variables X and Y .

The following parameters were ~~analyzedanalysed~~: the temperature at 2-meter level (T2m); sea-level pressure (~~PSL~~SLP); sea ice concentration (~~ICE~~ESIC); precipitations (PREC) as the sum of large-scale and convective precipitation; geopotential height at 500 mb (Z500); and wind speed at 10-meter level (U10). The following teleconnection indices – North Atlantic Oscillation (NAO), Arctic Oscillation (AO) and Barents Oscillation (BO) were calculated from the model data using `eofs.xarray` module in Python. NAO is defined as EOF-1 of seasonal SLP anomalies for 20–80N, 80W–40E, BO as EOF-2 of seasonal SLP anomalies for 30–90N, 90W–90E, and AO as EOF-1 of seasonal geopotential anomalies for 20–90N.

Correlations with and without the effect of teleconnection indices were ~~analyzedanalysed~~. For that purpose, partial correlations between atmospheric variables with the controlling impact of the teleconnection index were calculated.

For the testing area (TA), we chose a region around the Baltic Sea (50–65N, 10–40E). For the Arctic, we looked at the area ~~above 50N~~north of 60N. We chose ~~7~~seven important areas (IA) in the Arctic, ~~where the correlations with TA were stronger,~~

130 to ~~analyze~~analyze the relationships between regions in the Arctic and TA (~~Figure~~Fig. 1, Table 1). Initially, we also looked ~~also~~ at the Chukchi Sea and Canada Basin regions, but ~~all~~the correlations between climate parameters between TA and these regions' correlations with TA regions were clearly weaker ~~than with the remaining regions.~~



Figure 1: Important areas (IA, coloured areas) in the Arctic and test area (TA, black rectangle).

135 **Table 1. Important Areas (IA)**

	Area	Lat 1	Lat 2	Lon1	Lon 2	Mask
1	Central Arctic	85	90	0	360	No
2	Greenland	60	85	297	340	Land
3	West-Greenland	60	77	295	315	Sea

140	4	East-Greenland	60	75	315	340	Sea
	5	Svalbard	75	82	10	30	No
	6	Kara Sea	66	80	60	100	Sea
	7	Laptev Sea	66	80	100	170	Sea

145 We assessed the control simulation to reveal statistically significant seasonal correlations between TA and the Arctic region. For control simulation 1800-year long time-series, all correlations stronger than ± 0.046 are statistically significant at the confidence level of 95%. To ~~analyze~~analyse the strength and shape changes in teleconnections ~~transformations~~ during climate change, we looked at 20-year periods of the ensemble simulation from 1980 to 2100. For every ensemble simulation, 20-year long periods with 40 ensemble members (in a total of 800 events), all correlations stronger than ± 0.069 are statistically

150 significant at the confidence level of 95%. ~~This paper only looks at correlations stronger than ± 0.1 , as %.~~ In order to make the connections clearer, we do not show weaker correlations ~~are supposedly not important than ± 0.1 , though they were still statistically significant.~~ For control simulation, this gives a confidence level of 99.998%, while for ensemble simulations of 20-year-long series, the confidence level is 99.5%. Still, the relationship shown by the correlations may not be causal; there can be indirect connections, and we cannot rule out model internal feedback-generated correlations.

155 For understanding correlation maps, the first correlation parameter marks always the TA areal average, and the second one is from the Arctic. We also calculated correlations between TA and IA areal seasonal averages. We computed correlations with IAIA's previous month's averages to investigate the Arctic region's capability to forecast TA seasonal averages. Ridge regression (Saleh et al., 2019) is a multiple-regression method developed for cases with a strong correlation between input parameters. We used the Ridge regression method to estimate IA-s collective forecasting capability, using the module

160 `sklearn.linear_model` in Python with complexity parameter $\alpha = 1.0$.

3 Results


3.1 Stationary spatial correlations of climatic variables

The climatic variables of separate areas are usually dependent, but ~~correlations in space depend highly~~ the strength of the correlation depends on the distance and ~~climatic~~concrete variables. In addition to natural spatial correlations between climatic

165 variables within near distances, also correlations emerge within longer distances. To reveal stationary connections between remote regions, we used CESM-LE 1800 year-long stationary control run. This model run is very stable without significant trends, as the radiative forcing conditions were constantly at the year 1850 level during the whole period.

We found significant correlations between several variables between the Baltic Sea testing area (TA; shown with the rectangle in our Figures) and different Arctic areas. The most robust results were found for ~~2-meter temperature (T2m)~~; the negative

170 correlation between TA and the Greenland area is $R < -0.6$ in winter and only slightly weaker in spring and autumn (FigureFig. 2, row 1).



175 **Figure 2: The 2-meter temperature (T2m) seasonal correlations between the testing area (black rectangle) and surrounding areas according to the CESM-LE 1800-year control run. The second row shows the partial correlation with the controlling factor NAO.**

180 The summer is more influenced by local circulations, and remote correlations are weak. Most of these correlations are related to the NAO index – the same partial correlation with the controlling factor NAO has values in the Greenland area in winter weaker than -0.2 (Figure Fig. 2, row 2) and in spring and autumn mostly weaker than -0.3 . Teleconnections with TA that are not controlled by the NAO index are strongest at the Atlantic Ocean east to Iceland, but the correlation strength exceeds 0.3

185 The volume of physics-based analysis of these correlations does not fit in this paper, some methods and ideas can be found in Luo et al. (2016). TA parameters correlations with Z500 or U10 were weaker than with T2m, SIC and PREC and were not included in the following analysis.

190

Table 2. Seasonal correlations of control run 1800 years-long data between areal averages of testing area (TA, rows) and important area (IA, columns) parameters. The first column is the correlation between the parameter in TA and its estimation using Ridge regression over all parameters in IA-s. All shown correlations are statistically significant at the 95% confidence level. For each TA parameter in each IA-s, the strongest correlation is highlighted in bold.

	Correl	Central Arctic			Greenland			West-Greenland			East-Greenland			Svalbard			Kara Sea			Laptev Sea		
		Ridge	PSL	T2m	PSL	T2m	ICECSI	PSL	T2m	ICECSI	PSL	T2m	ICECSI	PSL	T2m	ICECSI	PSL	T2m	ICECSI	PSL	T2m	ICECSI
		P	S	C	P	S	C	P	S	C	P	S	C	P	S	C	P	S	C	P	S	C
MA	0.68	-0.42		0.05	-0.49	0.55	0.36	-0.39	0.51	0.14	-0.52	0.49	0.33	-0.54	0.07	-0.07	-0.41	0.43	-0.15	-0.38	0.39	-0.27
JJA	0.54	-0.31		-0.13	-0.25	0.17	0.13	-0.18	0.21	0.17	-0.28	-0.1		-0.29	0.23	-0.29	-0.15	0.12	0.06	-0.17	0.19	-0.16
SON	0.59	-0.21	0.09		-0.28	0.47	0.16	-0.15	0.38	0.14	-0.34	0.41	0.08	-0.42	0.09	-0.13	-0.13	0.19			0.06	
DJF	0.8	-0.47	-0.2	0.29	-0.58	0.65	0.28	-0.46	0.61	0.21	-0.61	0.57	0.23	-0.68	0.07		-0.52	0.47	-0.28	-0.37	0.2	-0.31
MA	0.51		0.11		-0.09	0.16	-0.07	-0.11		0.08	-0.07	0.28	-0.13	0.08	0.31	-0.22	0.21	-0.08		0.17		
JJA	0.47	-0.19	0.06	-0.06	-0.24	0.24	0.06	-0.16	0.19		-0.27	0.17		-0.28	0.09	-0.07		0.05		0.09	0.13	-0.05
SON	0.6	-0.08			-0.31	0.08	0.08	-0.3	0.14	0.12	-0.31	0.28			0.25	-0.1	0.08	-0.12				-0.07
DJF	0.66	0.1	0.13	-0.23	0.12	0.39	-0.12	0.07	0.21		0.15	0.47	-0.15	0.28	0.29	-0.17	0.31	-0.19	0.07	0.18		
MA	0.47	0.12			0.28	0.05		0.28	0.22	-0.12	0.27	0.11	0.06	0.13	-0.23	0.18					0.08	0.08
JJA	0.53	0.22			0.35	0.26	-0.12	0.28	0.27	-0.13	0.37	0.16		0.37	0.12	0.12	0.22	0.09			0.17	0.07
SON	0.56	0.16			0.39	0.02	-0.1	0.37	0.2	-0.15	0.38	-0.16		0.19	-0.15	0.07	0.11	0.07		0.1		0.06
DJF	0.59	-0.06	0.07	0.17		0.31	0.09		0.14	-0.05	-0.07	0.41	0.13	-0.19	0.22	0.15	-0.19	0.2		-0.12		0.06

195

To ~~generalize~~generalise the results, we divided the Arctic region into important areas (IA, [Figure Fig. 1](#)) and calculated correlations between IA and TA seasonal averages (Table 2).

200

The strongest correlations for IA-s are between T2m in TA and [PSL](#) in Svalbard, and seasonally, the correlation was strongest in winter ($R = -0.68$). Meanwhile, correlations with Svalbard T2m and [ICECSI](#) are much weaker with $|R| < 0.31$ in all seasons. TA T2m correlations with the Greenland region IA-s [PSL](#) are comparable to Svalbard, but correlations with T2m are much stronger. The correlations between T2m values in the Greenland region in winter and spring are $-0.65 \leq R \leq -0.49$ and in autumn $-0.47 \leq R \leq -0.38$.

205

Correlations between TA and IA-s parameters are the weakest in summer, where the strongest correlations are $R = 0.37$ between PREC in TA and [PSL](#) in both East Greenland and Svalbard. Correlations in summer between T2m in TA and [PSL](#) are even weaker, with the strongest correlation $R = -0.31$ in the Central Arctic. [ICECSI](#) in IA-s has generally weaker correlations with TA parameters than [PSL](#) or T2m.

210

Central Arctic parameters correlation with TA is stronger than 0.4 only between TA T2m and Central Arctic [PSL](#) in winter and spring. Laptev Sea parameters correlations with TA parameters were weaker than in other IA-s, with a correlation stronger than $|R| > 0.3$ occurring with TA T2m only in spring and winter.

To estimate IA-s collective forecasting capability, we calculated the correlation between the parameter in TA and its Ridge regression estimation (Table 2, first column). All IA-s seasonal [PSL](#), T2m and [ICECSI](#) values were used for the Ridge regression estimation. The correlation between T2m in TA and its Ridge regression estimation varies from 0.54 in summer to 0.8 in winter. The correlation between [PSL](#) in TA and its Ridge regression estimation is from 0.47 in summer to 0.66 in winter. The correlation between PREC in TA and its Ridge regression estimation is from 0.47 in spring to 0.59 in winter. ~~When~~

examining-As correlation square is the coefficient measure of determination R^2 the proportion of variance explained, then $0.82 = 64\%$ of T2m variability in TA in winter can be explained by PSLSLP, T2m and ICECSIC variability in Arctic IA-s. The

215 weakest connection with the Arctic is TA PSLSLP in summer and PREC in spring, with $R^2 = 0.472 = 22\%$.

Local correlations (at the same spot, no figures shown) are necessary to better understand teleconnections between different parameters. The abovementioned strong correlation between T2m in the Greenland region and TA can be connected through local relationships with other parameters. There is a robust local connection between T2m and ICECSIC, especially in the areas of ice margin ($R \sim 0.9$). ICEC The SIC correlation between TA and the Greenland region reaches $R = -0.4$; the correlation

220 between T2m at TA and ICECSIC in the Greenland region reaches $R = 0.5$.

The local correlation between ICECSIC and U10 is mostly strongly negative (strength up to -0.8 , not shown), especially in the regions where ICECSIC is lower than 0.8. No significant local correlation exists between ICECSIC and PSLSLP in the Greenland Sea (not shown).

3.2 Spatial correlations of climatic variables during 2020—2100

225 To analyze how teleconnections are modified by climate change, we investigated the differences between 20-year periods and the control run. Depending on the variable, the correlation might change its spatial pattern and value between different 20-year periods. The correlations between the following variables (~~the temperature at 2-meter level were analysed: T2m; sea-level pressure PSL and sea-ice concentration ICEC; precipitations, SLP, SIC, PREC; geopotential height at 500-mb, and Z500) were analyzed.~~ To get statistically reliable significant results, we used for every 20-year period all 40 ensemble

230 members, so we had in a total of 800 values for each period. Most of the correlations did not show significant changes in 20-year periods from the control run, including the most emphasized correlation between T2m in TA and the Arctic region.

However, there are some statistically significant changes in correlations between T2m in the TA and ICECSIC in the Arctic. Positive correlations between T2m in the TA and ICECSIC in the North Pole region winter (DJF) show simultaneously a

235 remarkable weakening in winter (DJF) after 2020. At the same time, the positive correlation strengthens significantly the North Pole region but a significant strengthening in the Davis Strait and Hudson Bay region- (Fig. 3).




Figure 3: The correlation in DJF between 2-meter temperature (T2m) in the testing area (black box) and sea ice concentration (SIC) in the control run and 20-year periods from ensemble simulations at the period 2020–2100.

The strong positive correlation in the Davis Strait is also remarkable in spring (MAM), but it does not strengthen as much as in winter (not shown). The correlation in winter in the region between Greenland and Iceland weakens. The negative correlation between T2m and ICECSIC in the control run in the coastal areas of Siberia fades away Russia becomes negligible after 2040, except in the Barents Sea, where it strengthens. A positive correlation is found in the East Siberian Sea in 2080–2100 (Fig. 3), supposedly connected with the decrease in the SIC (Fig. 4).

In regions of changing correlations, the average ICECSIC in winter will be lessening in the Barents Sea, Hudson Bay and between Greenland and Iceland (FigureFig. 4). ICECSIC in the North Pole region and coastal areas of Siberia eastward of the Barents Sea do not decrease in winter.





Figure 4: The difference of sea ice concentration (SIC) 20-year average in DJF from the control run average (first column) at the 20-year long periods of 2020–2100.

255

3.3 Lagged correlations

We are interested in factors driving variations in the Baltic Sea region and whether prior conditions may provide predictive capability. Given this, we studied the connections between earlier months (November for DJF, etc) average-~~month~~ values of different parameters in the Arctic region and seasonal values of TA conditions. The strongest correlations were between spring (MAM) T2m in TA and the previous month's (February) average T2m and PSL_{SLP} (Fig. 5). The correlation values were – 0.22 in Svalbard and Greenland regions; correlations between other parameters showed a weaker correlation ($|R| < 0.2$). Analogous correlations during other seasons were weaker. We used Ridge regression to determine the predictive capability of all previous months' average PSL_{SLP}, T2m and ICECSIC in all IA-s to TA next seasonal condition. Using Ridge regression did not improve the predictability much – the strongest correlation was $R = 0.30$ for T2m in MAM. Thus, even for the best case, the previous month's average values over the Arctic describe less than 10% of the variance of next season's TA average climate state variance.

265



270 **Figure 5: Lagged correlation between spring (MAM) mean 2-meter temperature (T2m) in TA and IA-s monthly means of 1) T2m (left); 2) sea level pressure (SLP, middle); and 3) sea ice concentration (SIC, right). On the x-axis are the monthly means in IA-s. All correlations stronger than ± 0.046 are statistically significant at the confidence level of 95 %.**

We focused on the testing area and searched for information at which rate the Arctic region climate parameters are statistically connected with incoming parameters in the testing area. It turned out that quite common are for several variables, the connection also occasionally worked contrariwise, where the values of the testing area variables can give information about the value of the Arctic region variables. For example, the spring average T2m in TA has $R > 0.35$ with ICECSIC in June at Greenland and East-Greenland. Stronger lagged correlations from TA to the Arctic can be explained by different averaging intervals – the monthly average in the Arctic has a lower influence on the next seasonal average in the TA than the TA seasonal average to the following month's average in the Arctic.

280 **4 Conclusion and Discussion**

The advantage of this study is the length of the stationary 1800-year long CESM-LE-self-consistent control database, which also reveals relations with a weaker strength (correlations stronger than ± 0.046 are statistically significant at the confidence level of 95 %). CESM-LE 40-member ensemble forecast until 2100 allows us to investigate how relationships may change in the changing climate. The most important teleconnections for the testing area are T2m, PSLSLP and ICECSIC in regions around Greenland and Svalbard (Figure Fig. 1, Table 1). For a long time, it has been recognized our results confirmed that ice conditions in the Greenland region might be connected is the most influential of all Arctic regions on teleconnection to several variables in Europe (Hildebrandsson 1914, Wiese 1924, Schell 1956, etc.). Our results did not confirm the old hypothesis by Hildebrandsson (1914) that the mean winter conditions over Europe depend on the summer sea ice extent in the Greenland Sea. Lagged (Hildebrandsson 1914). The lagged correlation

290 between summer ~~ICECSIC~~ in the Greenland Sea did not significantly correlate with any primary TA parameter in the following winter. It has to be considered that our testing area is only part of the area Hildebrandsson investigated.

As far as we know, the first attempt to reveal the teleconnections between the Baltic Sea region and the Arctic was made by our workgroup in 2017 (Jakobson et al., 2017). It was based on NCEP-CFSR_r and ERA-Interim reanalyses models for ~~the period~~ 1979—2015. Differences between the model parameters and different periods from ~~the~~ CESM-LE ensemble
295 complicated the comparison with the present study. The ~~comparablecomparison of~~ T2m in the present study and temperature at the 1000 hPa level in the study made in 2017 showed a different extent in the Greenland region but similar negative correlation strength in winter. In the present study, spring and autumn showed a much stronger correlation in much wider regions around Greenland. Summer showed a very weak correlation in both studies, probably due to the more
~~emphasizedemphasised~~ local circulation.

300 In investigating the influence of climatic parameters of the Arctic region on the testing area, we have to consider also the local correlations. Our results from the 1800-year long CESM-LE ensemble confirmed a strong ($R \sim 0.9$) local connection between T2m and ~~ICECSIC~~, as found in many other studies (e.g. Olonscheck et al., 2019; Vihma et al., 2014; Outten and Esau, 2012). Our results also confirm the strong negative local correlation between ~~ICECSIC~~ and U10 in the ice margin regions (strength up to -0.8), as shown in Jakobson et al. (2019). The negative correlation between SIC and W10 originates from the reduction
305 of both stratification and aerodynamic surface roughness with a reduction of SIC (Vavrus and Alkama, 2022; Jakobson et al., 2019). Many scientists have found a lower ~~PSLSLP~~ over the shrunk ice areas (Cassano et al., 2013; Alexander et al., 2004; Deser et al., 2000; Agnew, 1993), suggesting increased surface heating as a possible cause. Although Deser et al. (2000) found that mean ~~PSLSLP~~ has decreased over the retracted ice margin in the Greenland Sea (according to 1958–97 reanalysis products), our results did not show a significant correlation between ~~ICECSIC~~ and ~~PSLSLP~~ in the Greenland Sea region during
310 any season. According to Agnew (1993), the reason why the correlation is not present in the Greenland Sea may be due to the important role that ice export through Fram Strait and ocean currents play in determining ice extent in this region.

~~According to Chen et al. (2013),~~ Barents Oscillation (BO) is ~~through meridional flow and zonal wind anomalies~~ related to natural variability (the variation that humans do not cause) of Arctic surface air temperature (SAT) ~~variability through meridional flow and zonal wind anomalies (Chen et al., 2013).~~ Our T2m–T2m correlation pattern in winter (~~FigureFig.~~ 2) was
315 similar to the BO winter pattern. We tested ~~the~~ BO index influence on correlations between Arctic and TA, using partial correlation. The BO index effect was insignificant or more negligible than ~~the~~ NAO and AO index influence on all parameters we checked in all seasons. NAO and AO index had the largest impact in winter. In summer, the local effects are more dominant, and climate ~~indexesindices~~ influence is weaker.

We also aimed to reveal the ongoing climate change, especially AA's influence on teleconnections between TA and the Arctic
320 until ~~the year~~ 2100. Most of the correlations of 20-year periods did not show remarkable differences from the control run. Changes in the TA T2m correlations with ~~ICECSIC~~ in the Arctic ~~in the areas with decreasing ICEC~~ concur with negative trends in ~~ICECSIC~~ and positive trends in T2m in TA. Changes in the correlations in winter with regions with high and stable ~~ICECSIC~~ values are hard to suspect to have any direct physical basis. The strongest correlation between TA and Arctic region

parameters was T2m in TA and T2m in the Greenland region (Table 2). This correlation is constant up to 2100, ~~opposite to~~
325 ~~(not shown)~~. Sun et al. (2016), ~~who~~ declared that the "Warm Arctic, Cold Continents" regime is transient and becoming
increasingly unlikely as the climate continues to warm. There seems to be a discrepancy between these two results, but it is
not necessary. A strong negative correlation in winter means that warmer than average Arctic concur with colder than average
Baltic Sea region; it does not exclude that both regions' climate can continue to warm simultaneously.

To ~~generalize~~generalise separate Arctic regions' statistical connections with the TA, we used The Ridge regression. PSL_SLP,
330 T2m, and ICECSIC variability in Arctic IA-s can explain from 22_% of spring PREC to 64_% of winter T2m variability in TA.
A Climate indices can explain a substantial amount of it ~~can be explained by climate indices~~. The previous month's IA-s
averages forecasting capacity for TA seasonal average is much weaker – TA spring T2m has the highest coefficient of
determination $R^2 = 9\%$ with the Ridge regression estimation. Thus, we have to conclude that ~~the use of using~~ Arctic climate
data could not improve the Baltic Sea region's weather forecasting.

335 In conclusion – the Baltic Sea region climate has strong teleconnections with the Arctic climate; the strongest connections are
with the Svalbard and Greenland region. There is high seasonality in the teleconnections, with the strongest correlations in
winter and the weakest correlations in summer, when the local meteorological factors are stronger. The majority of
teleconnections in winter and spring can be explained by climate indexes NAO and AO.

By the end of the 21st century, the Arctic ice concentration has will significantly ~~decreased~~decrease. There will also be slight
340 changes in the teleconnection locations and strength. Still, the general teleconnections pattern between the Arctic and the Baltic
Sea region will not change during the 21st-century climate change.

The most important Arctic factors influencing the Baltic Sea are T2m and PSL_SLP, and the most important Arctic regions are
Greenland and Svalbard, but the mechanisms for these teleconnections ~~remained~~remain unknown. We have to agree with
Overland et al. (2016) that there are no simple cause-and-effect pathways in the Arctic and mid-latitude weather and climate
345 linkages.

Data availability

The CESM1-LE data are available online at <https://www.cesm.ucar.edu/community-projects/lens/data-sets>

Author contributions

EJ and LJ designed the experiments, ~~analyzed~~analysed the results and wrote the paper. EJ performed the experiments and
350 generated figures.

Competing interests

The contact author has declared that none of the authors has any competing interests.

Acknowledgements

We thank Marika Holland, Frederic Castruccio and the whole NCAR Oceanography Section for hosting our visit, for access
355 and support to use NCAR supercomputers and for the helpful discussions.

Financial support

We thank Baltic-American Freedom Foundation for financing EJ visit to NCAR via BAFF Research Scholar Program.

References

- 360 Agnew, T., 1993.: Simultaneous winter sea-ice and atmospheric circulation anomaly patterns, *Atmosphere-Atmos.-Ocean*,
31:2, 259–280, DOI:10.1080/07055900.1993.9649471, 1993.
- Alexander, M. A., Bhatt, U. S., Walsh, J. E., Timlin, M. S., Miller, J. S., Scott, J. D., 2004.: The atmospheric response to
realistic Arctic sea ice anomalies in an AGCM during winter. *J. ClimateClim.*, 17, 890–905, [https://doi.org/10.1175/1520-0442\(2004\)017<0890:TARTRA>2.0.CO;2](https://doi.org/10.1175/1520-0442(2004)017<0890:TARTRA>2.0.CO;2), 2004.
- BACC II Author Team: Second Assessment of Climate Change for the Baltic Sea Basin. Springer Open. 501 p. ISBN: 978-3-
365 319-16005-4 (Print) 978-3-319-16006-1 (Online), 2015.
- Barnes, E. A., and J. Screen, 2015:J.: The impact of Arctic warming on the midlatitude jet-stream: Can it? Has it? Will it?
Wiley Interdiscip. Rev.: *Climate. Clim. Change*, 6, 277–286, doi:10.1002/wcc.337, 2015.
- Barnhart, K. R., C. R. Miller, I. C. R., Overeem, I. and J. E. Kay, 2016:J. E.: Mapping the future expansion of Arctic open
water. *Nat. ClimateClim. Change*, 6, 280–285, <https://doi.org/10.1038/nclimate2848>, 2016.
- 370 Blackport, R., J. A. Screen, K. van der Wiel, and R. Bintanja, 2019: Minimal influenceR. and Krikken, F.: Magnitude and
pattern of reduced Arctic sea ice on coincident cold winters in mid-latitudes. *Nat. Climate Change*, 9, 697–704warming
governed by the seasonality of radiative forcing. *Sci. Rep.*, 6, 38287, 10.1038/srep38287, 2016.
- Bintanja, R. and van der Linden, E. C.: The changing seasonal climate in the Arctic. *Sci. Rep.*, 3, srep01556,
<https://doi.org/10.1038/s41558-019-0551-4srep01556>, 2013.
- 375 Boeke, R. C., and Taylor, P. C.: Seasonal Energy Exchange in Sea Ice Retreat Regions Contributes to Differences in Projected
Arctic Warming, *Nat. Commun*, 9, 5017. doi:10.1038/s41467-018-07061-9, 2018.

- Cassano, E. N., ~~J. J. Cassano, M. E. J. J., Higgins, M. E. and M. C. Serreze, 2013: M. C.:~~ Atmospheric impacts of an Arctic sea ice minimum as seen in the Community Atmosphere Model. *Int. J. Climatol.*, 34(3), 766–779. doi: 10.1002/joc.3723. ~~Chavaillaz et al., 2016, 2013.~~
- 380 ~~Cattiaux, J., Vautard, R., Cassou, C., Yiou, P., Masson-Delmotte, V., & Codron, F.: Winter 2010 in Europe: A cold extreme in a warming climate, *Geophys. Res. Lett.*, 37, L20704. <https://doi.org/10.1029/2010GL044613>, 2010.~~
- ~~Chen, H. W., Q. Zhang, H. Q., Körnich, and D. H. Chen (2013): D.: A robust mode of climate variability in the Arctic: The Barents Oscillation. *Geophysical Research Letters*, *Geophys. Res. Lett.*, 40, 2856–2861, <https://doi.org/10.1002/grl.50551>, 2013.~~
- 385 ~~Christensen, O. B., Kjellström, E., Dieterich, C., Gröger, M., and Meier, H. E. M.: Atmospheric regional climate projections for the Baltic Sea region until 2100, *Earth Syst. Dynam.*, 13, 133–157, <https://doi.org/10.5194/esd-13-133-2022>, 2022.~~
- ~~Clark, J. P. & Shenoy, V., Feldstein, S., Lee, S. 2019., Goss, M.: The Role of the Tropically Excited Arctic Warming Mechanism on the Warm Arctic Cold Continent Surface Air Horizontal Temperature Trend Pattern, *Geophysical Research Letters*, Volume 46, Issue 14 Advection in Arctic Amplification, *J. Clim.*, 34, 8, 2957–2976, <https://doi.org/10.1029/2019GL082714> 1175/JCLI-D-19-0937.1, 2021.~~
- 390 ~~Cohen, J., Zhang, X., Francis, J., Jung, T., et al., 2018.: Arctic change and possible influence on mid-latitude climate and weather: A US CLIVAR White Paper, US CLIVAR Rep. 2018 Mar; n/a., doi: 10.5065/D6TH8KGW, 2018.~~
- ~~Cohen, J., Agel, L., Barlow, M., Garfinkel, C.I., White, I. 2021, Linking Arctic variability and change with extreme winter weather in the United States, *Science*, Vol 373 Issue 6559, DOI: 10.1126/science.abi9167~~
- 395 ~~Coumou, D., Di Capua, G., Vavrus, S. et al., 2018.: The influence of Arctic amplification on mid-latitude summer circulation., *Nat. Commun.*, 9, 2959, <https://doi.org/10.1038/s41467-018-05256-8>, 2018.~~
- ~~Dai, A., Song, M., 2020.: Little influence of Arctic amplification on mid-latitude climate. *Nat. Clim. Chang.*, 10, 231–237, <https://doi.org/10.1038/s41558-020-0694-3>, 2020.~~
- ~~Dai, A., Jenkins, M.: Relationships among Arctic warming, sea-ice loss, stability, lapse rate feedback, and Arctic amplification, *Clim. Dyn.*, 61, 5217–5232, [10.1007/s00382-023-06848-x](https://doi.org/10.1007/s00382-023-06848-x), 2023.~~
- 400 ~~Dai, A., Luo, D., Song, M., Liu, J., 2019.: Arctic amplification is caused by sea-ice loss under increasing CO 2. *Nat. Nature Communications Commun.*, 10, 121., <https://doi.org/10.1038/s41467-018-07954-9>, 2019.~~
- ~~Deng, K., Yang, S., Ting, M., Lin, A., and Wang, Z.: An intensified mode of variability modulating the summer heatwaves in eastern Europe and Northern China, *Geophys. Res. Lett.*, 45, 11, 361–369, <https://doi.org/10.1029/2018GL079836>, 2018.~~
- 405 ~~Deser, C., Hurrell, J.W. & Phillips, A.S. The role of the North Atlantic Oscillation in European climate projections. *Clim Dyn* 49, 3141–3157 (2017). <https://doi.org/10.1007/s00382-016-3502-z>~~
- ~~Deser, C., Walsh, J.E., Timlin, M.S., 2000.: Arctic Sea Ice Variability in the Context of Recent Atmospheric Circulation Trends, *Journal of Climate*, Volume J. *Clim.*, 13—Issue, 3, [https://doi.org/10.1175/1520-0442\(2000\)013<0617:ASIVIT>2.0.CO;2](https://doi.org/10.1175/1520-0442(2000)013<0617:ASIVIT>2.0.CO;2), 2000.~~

- 410 [Deser, C., Hurrell, J.W. and Phillips, A.S.: The role of the North Atlantic Oscillation in European climate projections. *Clim. Dyn.*, 49, 3141–3157, <https://doi.org/10.1007/s00382-016-3502-z>, 2017.](#)
- [Duan, L., Cao, L., and Caldeira, K.: Estimating Contributions of Sea Ice and Land Snow to Climate Feedback. *J. Geophys. Res. Atmos.* 124 \(1\), 199–208, doi:10.1029/2018JD029093, 2019.](#)
- [Francis, J. A. and Hunter, E.: New insight into the disappearing Arctic sea ice. *EOS*, 87, 509–511, <https://doi.org/10.1029/2006EO460001>, 2006.](#)
- 415 ~~Francis, J. A. and Vavrus, S. J.: Francis, J. A. and Vavrus, S. J. 2015. Evidence for a wavier jet stream in response to rapid Arctic warming, *Environmental Research Letters*, Volume 10, Number 1, 10Environ. Res. Lett., 10, 014005~~
- ~~Francis, J. A., Vavrus, S. J. and Cohen, J. 2017. Amplified Arctic warming and mid latitude weather: new perspectives on emerging connections. *WIRES CLIM CHANGE*, 8, e474, doi: 10.1002/wcc.474.~~
- 420 ~~Fyfe, J. C., and et al., 2016: Making sense of the early 2000s warming slowdown. *Nat. Climate Change*, 6, 224–228, doi:10.1038/nclimate2938.~~
- ~~Grenier, P., Elia, R., Chaumont, D., 2015. Chances of Short Term Cooling Estimated from a Selection of CMIP5 Based Climate Scenarios during 2006–35 over Canada, *Journal of Climate* 28 (8): 3232–3249, DOI: 10.1175/JCLI-D-14-00224.1088/1748-9326/10/1/014005, 2015.~~
- 425 ~~Guan, W., Jiang, X., Ren, X., Chen, G., Ding, Q., 2020. Role of Atmospheric Variability in Driving the "Warm Arctic, Cold Continent" Pattern Over the North America Sector and Sea Ice Variability Over the Chukchi Bering Sea, *Geophysical Research Letters*, Volume 47, Issue 13, <https://doi.org/10.1029/2020GL088599>~~
- Halkka, A.: Changing climate and the Baltic region biota, Doctoral dissertation, Faculty of Biological and Environmental Sciences, University of Helsinki, Helsinki, Finland, 52 pp., <http://urn.fi/URN:ISBN:978-951-51-6021-8> (last access: 16
- 430 February 2022), 2020.
- ~~Hildebrandsson, H. H. (1914): Quelques recherches sur les centres d'action de l'atmosphère. *Kungl. Svenska vetenskapsakademiens handlingar*, 51, 3–16, 1914.~~
- [Hurrell, J.W. et al.: The Community Earth System Model: A Framework for Collaborative Research, *Bull. Am. Meteorol. Soc.*, 94, 1339–1360, <https://doi.org/10.1175/BAMS-D-12-00121.1>, 2013.](#)
- 435 IPCC, 2021: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press. In Press, 2021.
- ~~Jahn, A.: Reduced probability of ice-free summers for 1.5 °C compared to 2 °C warming. *Nature Nat. Clim. Change*, 8, 409–413 (2018), <https://doi.org/10.1038/s41558-018-0127-8>, 2018.~~
- 440 ~~Jahn, A., J. E. Kay, M. M. J. E., Holland, M. M. and D. M. Hall, 2016: D. M.: How predictable is the timing of a summer ice-free Arctic? *Geophys. Res. Lett.*, 43, 9113–9120, <https://doi.org/10.1002/2016GL070067>, 2016.~~

- Jakobson, L., Jakobson, E., Post, P., Jaagus, J., ~~2017.~~: Atmospheric teleconnections between the Arctic and the eastern Baltic Sea regions. *Earth Systems Dynamics, Syst. Dyn.*, 8, 1019–1030, <https://doi.org/10.5194/esd-8-1019-2017>, ~~2017~~.
- 445 Jakobson, L., Vihma, T., Jakobson, E., ~~2019.~~: Relationships between Sea Ice Concentration and Wind Speed over the Arctic Ocean during 1979–2015, *Journal of Climate, Volume J. Clim.*, 32, Issue-22, <https://doi.org/10.1175/JCLI-D-19-0271.1>, ~~2019~~.
- ~~Jung, T., Jenkins, M. and Dai, A.: The Impact of Sea-Ice Loss on Arctic Climate Feedbacks and Their Role for Arctic Amplification, Geophys. Res. Lett., 48, 15, https://doi.org/10.1029/2021GL094599, 2021.~~ Jung, T. et al.: Polar lower-latitude linkages and their role in weather and climate prediction, *Bull. Am. Meteorol. Soc.*, 96, ES197–ES200, 2015.
- 450 Kay, J. E., ~~and Coauthors, 2015.~~ et al.: The Community Earth System Model (CESM) large ensemble ~~project-project~~: A community resource for studying climate change in the presence of internal climate variability. *Bull. Amer. Meteor. Soc.*, 96, 1333–1349, doi:<https://doi.org/10.1175/BAMS-D-13-00255.1>, ~~2015~~.
- ~~Klein, R.J.T., G.F. Midgley, B.L. Preston, M. Alam, F.G.H. Berkhout, K. Dow, and M.R. Shaw, 2014. Adaptation opportunities, constraints, and limits. In: Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Field, C.B., V.R. Barros, D.J. Dokken, K.J. Mach, M.D. Mastrandrea, T.E. Bilir, M. Chatterjee, K.L. Ebi, Y.O. Estrada, R.C. Genova, B. Girma, E.S. Kissel, A.N. Levy, S. MacCracken, P.R. Mastrandrea, and L.L. White (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 899–943.~~
- 455 ~~Field, C.B., V.R. Barros, D.J. Dokken, K.J. Mach, M.D. Mastrandrea, T.E. Bilir, M. Chatterjee, K.L. Ebi, Y.O. Estrada, R.C. Genova, B. Girma, E.S. Kissel, A.N. Levy, S. MacCracken, P.R. Mastrandrea, and L.L. White (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 899–943.~~
- Kug, JSJ. S., Jeong, JH., Jang, YS. et al., ~~2015.~~: Two distinct influences of Arctic warming on cold winters over North America and East Asia. *Nature, Nat. Geosci.*, 8, 759–762. <https://doi.org/10.1038/ngeo2517>, ~~2015~~.
- 460 Labe, Z., ~~G.~~ Magnúsdóttir, G. and ~~H.~~ Stern, ~~2018:~~H.: Variability of Arctic sea ice thickness using PIOMAS and the CESM Large Ensemble. *J. Climate, Clim.*, 31, 3233–3247, <https://doi.org/10.1175/JCLI-D-17-0436.1>, ~~2018~~.
- ~~Lehner, F., Laine, A., Yoshimori, M. and Deser, Abe-Ouchi, A.: Surface Arctic amplification factors in CMIP5 models: land~~
- ~~€., Terray, L. 2017. Toward a New Estimate of "Time of Emergence" of Anthropogenic Warming: Insights from Dynamical~~
- 465 ~~Adjustment and a Large Initial Condition Model Ensemble. Journal of Climate, October 2017 (7739–7756), oceanic surfaces and seasonality. J. Clim., 29, 3297–3316, https://doi.org/10.1175/JCLI-D-15-0497.1, 2016.~~
- ~~Lee J. and Cai, M.: Seasonality of polar surface warming amplification in climate simulations, Geophys. Res. Lett., 36, L16704, 10.1029/2009GL040133, 2009.~~
- ~~Lee, S. H., Charlton-Perez, A. J., Furtado, J. C., and Woolnough, S. J.: Abrupt stratospheric vortex weakening associated with~~
- 470 ~~North Atlantic anticyclonic wave breaking, J. Geophys. Res.-Atmos., 124, 8563–8575, https://doi.org/10.1029/2019JD030940, 2019.~~
- ~~Luo, D., Y. Xiao, Y., Diao, Y., Dai, A., Franzke, C. And Simmonds, I.: Impact of Ural Blocking on winter Warm Arctic–Cold Eurasian anomalies. Part II: The link to the North Atlantic Oscillation, J. Clim., 29, 3949–3971, doi: 10.1175/JCLI-D-15-0612.1, 2016.~~

- 475 [Luo, D. et al.: Increased quasi-stationarity and persistence of winter Ural Blocking and Eurasian extreme cold events in response to Arctic warming. Part II: A theoretical explanation. *J. Clim.*, 30, 3569–3587, doi: 10.1175/JCLI-D-16-07920262.1, 2017.](#)
- [Luo, D., Chen, X., Overland, J., Simmonds, I., Wu, Y., Zhang, P.: Weakened potential vorticity barrier linked to recent winter Arctic sea ice loss and midlatitude cold extremes. *J. Clim.*, 32, 4235-4261, doi: 10.1175/JCLI-D-18-0449.1, 2019.](#)
- 480 [Martius, O., Polvani, L. M., and Davies, H. C.: Blocking precursors to stratospheric sudden warming events. *Geophys. Res. Lett.*, 36, 1–5, <https://doi.org/10.1029/2009GL038776>, 2009.](#)
- Massonnet, F., Vancoppenolle, M., Goosse, H., Docquier, D., Fifechet, T., and Blanchard-Wrigglesworth, E.: Arctic sea-ice change tied to its mean state through thermodynamic processes, *Nat. Clim. Change*, 8, 599–603, [10.1038/s41558-018-0204-z](https://doi.org/10.1038/s41558-018-0204-z), 2018.
- 485 ~~[Medhaug, I., M. B. Stolpe, E. M. Fischer, and R. Knutti, 2017: Reconciling controversies about the 'global warming hiatus.' *Nature*, 545, 41–47, doi:10.1038/nature22315.](#)~~
- Meleshko, V. P., Pavlova, T., Bobylev, L. P., Golubkin, P., ~~2020.~~: Current and Projected Sea Ice in the Arctic in the Twenty-First Century, in: Johannessen, O.M., Bobylev, L.P., Shalina, E.V., Sandven, S. (Eds.), *Sea Ice in the Arctic: Past, Present and Future*, Springer Polar Sciences, Springer International Publishing, Cham, pp.399–463, https://doi.org/10.1007/978-3-030-21301-5_10, 2020.
- 490 Nakamura, T. ~~&and~~ Sato, T., ~~2022.~~: A possible linkage of Eurasian heat wave and East Asian heavy rainfall in Relation to the Rapid Arctic warming, ~~*Environmental Research, Volume 209, 2022*~~*Environ. Res.*, 209, 112881, ISSN 0013-9351, <https://doi.org/10.1016/j.envres.2022.112881>, 2022.
- Olonscheck, D., Mauritsen, T. ~~&and~~ Notz, D.: Arctic sea-ice variability is primarily driven by atmospheric temperature fluctuations, *Nat. Geosci.*, 12, 430–434 (~~2019~~), <https://doi.org/10.1038/s41561-019-0363-1>, 2019.
- 495 Outten, S., and Esau, I., ~~2012.~~: A link between Arctic sea ice and recent cooling trends over Eurasia, ~~*Climate Clim. Change*, 110(3), 1069-1075, DOI: 10.1007/s10584-011-0334-z, 2012.~~
- ~~[Overland, J.E., Hanna, E., Hanssen-Bauer, I., Kim, S.-J., Walsh, J.E., Wang, M., Bhatt, U.S., Thoman, R.L., 2018. Surface air temperature. *Arctic Report Card*.](#)~~
- 500 ~~[Overland, J.E., Francis, J., Hall, R., Hanna, E., Kim, S.J., Vihma, T., 2015.: The melting Arctic and mid-latitude weather patterns: Are they connected? *J. Climate*, 28\(20\), 7917–7932, doi: 10.1175/JCLI-D-14-00822.1-, 2015.](#)~~
- ~~[Overland, J. E., Hanna, E., Hanssen-Bauer, I., Kim, S.-J., Walsh, J. Wood, K.R E., Wang, M., Bhatt, U. S., 2011. WarmThoman, R. L.: Surface air temperature. *Arctic Report Card*, 2018.](#)~~
- ~~[Overland, J. E., –cold continents: climate impacts of Ballinger, T. J., Cohen, J., Francis, J. A., Hanna, E., Jaiser, R., Kim, B.-](#)~~
- 505 ~~[M., Kim, S.-J., Ukita, J., Vihma, T., Wang, M. and Zhang, X.: How do intermittency and simultaneous processes obfuscate the newly open-Arctic influence on midlatitude winter extreme weather events?, *Env. Res. Lett.*, 16, 043002, doi: 10.1088/1748-9326/abdb5d, 2021.](#)~~

- Peings, Y.: Ural Blocking as a Driver of Early-Winter Stratospheric Warming, *Geophys. Res. Lett.*, 46, 5460-5468, <https://doi.org/10.1029/2019GL082097>, 2019.
- 510 Peings, Y., Davini, P., Magnusdottir, G.: Impact of Ural Blocking on Early Winter Climate Variability Under Different Barents-Kara Sea, *Polar Research*, Vol 30 (2011), Ice Conditions, *JGR Atmos.*, 128, e2022JD036994, <https://doi.org/10.3402/polar.v30i0.157871029/2022JD036994>, 2023.
- Pithan, F., Mauritsen, T.: Arctic amplification dominated by temperature feedbacks in contemporary climate models. *Nature, Nat. Geosci.*, 7, 181–184 (2014). <https://doi.org/10.1038/ngeo2071>, 2014.
- 515 Risbey, J.S., Grose, M.R., Moselesan, D., O’Kane T.J., Lewandowsky, Rantanen, M., Karpechko, A. Y., Lipponen, A., Nordling, K., Hyvärinen, O., Ruosteenoja, K., Vihma, T., and Laaksonen, A.: The Arctic has warmed nearly four times faster than the globe since 1979, *Commun. Earth Environ.*, 3, 168, doi: 10.1038/s43247-022-00498-3, 2022.
- ~~S., 2017. Transient response of the global mean warming rate and its spatial variation, *Weather and Climate Extremes* 18, DOI: 10.1016/j.wace.2017.11.002~~
- 520 Rondeau-Genesse, G., Braun, M., 2019.: Impact of internal variability on climate change for the upcoming decades: analysis of the CanESM2-LE and CESM-LE large ensembles. *Climatic, Clim. Change*, 156, 299–314 (2019). <https://doi.org/10.1007/s10584-019-02550-2>, 2019.
- Rudeva, I. and Simmonds, I.: Midlatitude winter extreme temperature events and connections with anomalies in the Arctic and tropics, *J. Clim.*, 34, 3733-3749, doi: 10.1175/JCLI-D-20-0371.1, 2021.
- 525 Rutgersson, A., Kjellström, E., Haapala, J., Stendel, M., Danilovich, I., Drews, M., Jylhä, K., Kujala, P., Larsén, X. G., Halsnæs, K., Lehtonen, I., Luomaranta, A., Nilsson, E., Olsson, T., Särkkä, J., Tuomi, L., and Wasmund, N.: Natural hazards and extreme events in the Baltic Sea region, *Earth Syst. DynamDyn.*, 13, 251–301, <https://doi.org/10.5194/esd-13-251-2022>, 2022.
- Schell, I. I., 1956.: Interrelations of Arctic ice with the atmosphere and the ocean in the North Atlantic-Arctic and adjacent areas, *Journal of the Atmospheric Sciences, Volume 13, Issue J. Atmos. Sci.*, 13, 1, [https://doi.org/10.1175/1520-0469\(1956\)013<0046:IOAIWT>2.0.CO;2](https://doi.org/10.1175/1520-0469(1956)013<0046:IOAIWT>2.0.CO;2), 1956.
- 530 Saleh, A. K., Arashi, M., Kibria Golam, B. M.: *Theory of Ridge Regression Estimation with Applications*. New York: John Wiley & Sons. ISBN 978-1-118-64461-4, 2019.
- Sang, X., Yang, XQ., Tao, L. *et al.*: Decadal changes of wintertime poleward heat and moisture transport associated with the amplified Arctic warming. *Clim. Dyn.*, 58, 137–159, <https://doi.org/10.1007/s00382-021-05894-7>, 2022.
- 535 Screen, J. A. and Simmonds, I.: Increasing fall-winter energy loss from the Arctic Ocean and its role in Arctic temperature amplification, *Geophys. Res. Lett.*, 37, L16707, <https://doi.org/10.1029/2010GL044136>, 2010.
- Serreze, M. C., Barrett, A. P., Stroeve, J. C., Kindig, D. N. and Holland, M. M.: The emergence of surface-based Arctic amplification, *TC*, 3, 11–19, <https://doi.org/10.5194/tc-3-11-2009>, 2009.
- 540 Smith, A., and Jahn, A., 2019.: Definition differences and internal variability affect the simulated Arctic sea ice melt season, *The CryosphereTC*, 13, 1-20, <https://doi.org/10.5194/tc-13-1-2019>, 2019.

- Spielhagen, R. F. et al.: Enhanced modern heat transfer to ~~Sun, L., Alexander, M., Deser, C., 2016. Evolution of the Global Coupled Climate Response to Arctic Sea Ice Loss during 1990–2090 by warm Atlantic water. *Sci.*, 331, 450–453, 10.1126/science.1197397, 2011.~~
- 545 ~~Statnaia, I. A., Karpechko, A. Y., and Its Contribution to Climate Change, *Journal of Climate*, Volume 31, Issue 19 Järvinen, H. J.: Mechanisms and predictability of sudden stratospheric warming in winter 2018, *Weather Clim. Dynam.*, 1, 657–674, <https://doi.org/10.1175/JCLI-D-18-0134.1-5194/wcd-1-657-2020>, 2020.~~
- ~~Sun, L., Perlwitz, J., and Hoerling, M.: What caused the recent "Warm Arctic, Cold Continents" trend pattern in winter temperatures? *Geophys. Res. Lett.*, 43, 5345–5352, doi: 10.1002/2016gl069024, 2016.~~
- 550 Sun, L., Alexander, M. A., & Deser, C. (2018).: Evolution of the global coupled climate response to Arctic sea ice loss during 1990–2090 and its contribution to climate change. ~~*Journal of Climate*, *J. Clim.*, 31(19), 7823–7843, <https://doi.org/10.1175/JCLI-D-18-0134.1>, 2018.~~
- ~~Swart, N. C., J.-C. Fyfe, E. J. C., Hawkins, E., Kay, J. E. Kay, and A. Jahn, 2015: A.: Influence of internal variability on Arctic sea-ice trends. *Nat. Climate Change*, 5, 86–89, <https://doi.org/10.1038/nclimate2483>, 2015.~~
- 555 ~~Taylor, P., Boeke, R., Boisvert, L., Feldl, N., Henry, M., Huang, Y., Langen, P., Liu, W., Pithan, F., Sejas, S., Tan, I.: Process Drivers, Inter-Model Spread, and the Path Forward: A Review of Amplified Arctic Warming, *Front. Earth Sci.*, 9, 10.3389/feart.2021.758361, 2022.~~
- ~~Taylor, P. C., Kato, S., Xu, K.-M. and Cai, M.: Covariance between Arctic sea ice and clouds within atmospheric state regimes at the satellite footprint level. *J. Geophys. Res. Atmos.*, 120, 12656–12678, <https://doi.org/10.1002/2015JD023520>, 2015.~~
- 560 ~~Taylor, P. C., Cai, M., Hu, A., Meehl, J., Washington, W., and Zhang, G. J.: A Decomposition of Feedback Contributions to Polar Warming Amplification. *J. Clim.*, 26 (18), 7023–7043. doi:10.1175/JCLI-D-12-00696.1, 2013.~~
- ~~Vavrus, S.: The impact of cloud feedbacks on Arctic climate under greenhouse forcing. *J. Clim.*, 17, 603–615, [https://doi.org/10.1175/1520-0442\(2004\)017<0603:TIOCF0>2.0.CO;2](https://doi.org/10.1175/1520-0442(2004)017<0603:TIOCF0>2.0.CO;2), 2004.~~
- 565 ~~Vavrus, S. J. and Alkama, R.: Future trends of arctic surface wind speeds and their relationship with sea ice in CMIP5 climate model simulations, *Clim. Dyn.*, 59, 1833–1848, doi: 10.1007/s00382-021-06071-6, 2022.~~
- ~~Vihma, T.: Effects of Arctic Sea Ice Decline on Weather and Climate: A Review, *Surv. Geophys.*, 35, 1175–1214, <https://doi.org/10.1007/s10712-014-9284-0>, 2014.~~
- ~~Vihma, T., Ruhe, G., Chen, L. et al., 2019.: Effects of the tropospheric large-scale circulation on European winter temperatures during the period of amplified Arctic warming *Int. J. Climatology*, 40, 509–529, <https://doi.org/10.1002/joc.6225>, 2019.~~
- ~~Vihma, T. Effects of Arctic Sea Ice Decline on Weather and Climate: A Review. *Surv Geophys* 35, 1175–1214 (2014). <https://doi.org/10.1007/s10712-014-9284-0>~~
- ~~Viru, B. and Jaagus, J.: Spatio-temporal variability and seasonal dynamics of snow cover regime in Estonia, *Theor. Appl. Climatol.*, 139, 759–771, <https://doi.org/10.1007/s00704-019-03013-5>, 2020.~~
- 575 ~~Wiese, W. (1924).: Polareis und atmosphärische Schwankungen. *Geograf. Ann.*, 6, 273–299, 1924.~~




Figure 1. Important areas (IA, coloured areas) in the Arctic and test area (TA, black rectangle).





Figure 2. The 2 meter temperature (T2m) seasonal correlations between the testing area (black rectangle) and surrounding areas according to CESM-LE 1800-year control run. Yang, S., Wen, X., Gao, M.: Relation of Mid-High-Latitude Eurasian ISO to Ural Blocking Frequency and Their Co-Effect on Extreme Hot Events during Boreal Summer, Atmosphere, Volume 13, Issue 12, 10.3390/atmos13122041, 2022.


Yao, Y., Luo, D., Dai, A. and Simmonds, I.: Increased quasi stationarity and persistence of Ural blocking and Eurasian extreme cold events in response to Arctic warming. Part I: Insights from observational analyses. *J. Clim.*, 30, 3549–3568, doi:10.1175/JCLI-D-16-0261.1, 2017.

Yoshimori, M., Watanabe, M., Abe-Ouchi, A., Shiogama, H. and Ogura, T.: Relative contribution of feedback processes to Arctic amplification of temperature change in MIROC GCM, *Clim. Dyn.*, 42, 1613–1630, <https://doi.org/10.1007/s00382-013-1875-9>, 2014.

Zhuo, W., Yao, Y., Luo, D., Simmonds, I., Huang, F.: The key atmospheric drivers linking regional Arctic amplification with East Asian cold extremes, *Atmos. Res.*, 283, 106557, ISSN 0169-8095, <https://doi.org/10.1016/j.atmosres.2022.106557>, 2023. The second row shows the partial correlation with the controlling factor NAO.



600 **Figure 3.** The correlation in DJF between 2-meter temperature (T2m) in the testing area (black box) and ice concentration (ICEC) in the control run and 20-year periods from ensemble simulations at the period 2020—2100.



605 **Figure 4.** The difference of ice concentration (ICEC) 20-year average in DJF from the control run average (first column) at the 20-year long periods of 2020—2100.




Figure 5. Lagged correlation between spring (MAM) mean 2-meter temperature (T2m) in TA and IA-s monthly means of 1) T2m (left); 2) pressure at sea level (PSL, middle); and 3) ice fraction (ICEC, right). On the x-axis are the monthly means in IA-s. All correlations stronger than ± 0.046 are statistically significant at the confidence level of 95%.