

# Zonal variability of methane trends derived from satellite data

Jonas Hachmeister<sup>1</sup>, Oliver Schneising<sup>1</sup>, Michael Buchwitz<sup>1</sup>, John P. Burrows<sup>1</sup>, Justus Notholt<sup>1</sup>, and Matthias Buschmann<sup>1</sup>

<sup>1</sup>Institute of Environmental Physics (IUP), University of Bremen FB1, Bremen, Germany

**Correspondence:** J. Hachmeister (jonas\_h@iup.physik.uni-bremen.de)

**Abstract.** The Tropospheric Monitoring Instrument (TROPOMI) on-board the satellite Sentinel-5 Precursor (S5P) is part of the latest generation of trace gas monitoring satellites and provides a new level of spatio-temporal information with daily global coverage, which enable the calculation of daily globally averaged CH<sub>4</sub> concentrations. To investigate changes of atmospheric methane, the background CH<sub>4</sub> level (i.e. the CH<sub>4</sub> concentration without seasonal and short-term variations) has to be determined. CH<sub>4</sub> growth rates vary in a complex manner and high-latitude zonal averages may have gaps in the time series, thus simple fitting methods don't produce reliable results. In this manuscript we present an approach based on fitting an ensemble of Dynamic Linear Models (DLMs) to TROPOMI data, from which the best model is chosen with the help of cross-validation to prevent overfitting. [This method is computationally fast and not dependent additional inputs, allowing for the fast and continuous analysis of the most recent time series data.](#) We present results of global annual methane increases (AMIs) for the first 4.5 years of S5P/TROPOMI data which show good agreement with AMIs from other sources. Additionally, we investigated what information can be derived from zonal bands. Due to the fast meridional mixing within hemispheres we use zonal growth rates instead of AMIs, since they provide a [daily-higher](#) temporal resolution. Clear differences can be observed between Northern and Southern Hemisphere growth rates, especially during 2019 and 2022. The growth rates show similar patterns within the hemispheres and show no short-term variations during the years, indicating that air masses within a hemisphere are well-mixed during a year. Additionally, the growth rates derived from S5P/TROPOMI data are largely consistent with growth rates derived from CAMS global inversion-optimized (CAMS/INV) data. In 2019 a reduction in growth rates can be observed for the Southern Hemisphere, while growth rates in the Northern Hemisphere stay stable or increase. During 2020 a strong increase in Southern Hemisphere growth rates can be observed, which is in accordance with recently reported increases in Southern Hemisphere wetland emissions. In 2022 the reduction of the global AMI can be attributed to decreased growth rates in the Northern Hemisphere, while growth rates in the Southern Hemisphere remain high. Investigations of fluxes from CAMS/INV data support these observations and suggest that the Northern Hemisphere decrease is mainly due to the decrease in anthropogenic fluxes while in the Southern Hemisphere wetland fluxes continued to rise.

## 1 Introduction

Methane (CH<sub>4</sub>) is one of the most important drivers of climate change with an effective radiative forcing of  $1.19 \text{ Wm}^{-2}$  [Arias *et al.*, 2021] and an atmospheric lifetime of 9.1 years [Szopa *et al.*, 2021]. The short lifespan of CH<sub>4</sub> compared to other greenhouse gases and the large fraction of anthropogenic emissions, makes CH<sub>4</sub> emission reduction [a straight-forward-way-to-tackle](#)

~~an attractive strategy to slow down or possibly reduce man-made~~ climate change in the short- to midterm. ~~A close monitoring of Accurate knowledge of the~~ atmospheric CH<sub>4</sub> concentrations ~~is therefore vital, both for scientists and policymakers, and~~ ~~dry column mixing ratios are therefore essential to improve our knowledge of the sources and sinks of CH<sub>4</sub> for science and~~ ~~international environmental policy.~~ The globally averaged surface concentration of CH<sub>4</sub> has increased by 156% between 1750 and 2019 reaching 1866 ± 3.3 ppb in 2019 (Gulev et al., 2021) and ~~1919.97~~ 1917.11 ppb in ~~February-June~~ 2023 [Lan et al., 2023].

While the concentrations have risen in total, the trend, i.e. the rate of change in the background level without seasonal or short-term variations, has evolved non-linearly. ~~An increase in global methane concentrations can be observed between the 1980s-2000~~ Global methane concentrations have been observed to increase in the period from the 1980s to 2000 and from 2007 until the present, ~~with~~. ~~However,~~ a plateau between 2000-2007 ~~which was observed. This~~ is referred to as “stabilization“. Whether to define the stabilization period or the period of renewed growth (2007-present) as anomalous has been the subject of debate. There have been a variety of explanations for the observed behavior in the literature (~~Turner et al., 2019~~) [Turner et al., 2019]. Recent publications suggest that the period of renewed growth can be attributed to the rise in microbial emissions (~~Lan et al., 2021; Basu et al., 2022~~) [Lan et al., 2021; Basu et al., 2022] and that tropical methane emissions explain a majority of recent changes in the atmospheric methane growth rate (~~Feng et al., 2022~~) [Feng et al., 2022]. In 2020 and 2021 record methane increases were observed by the Global Monitoring Laboratory of the ‘National Oceanic and Atmospheric Administration (NOAA-GML) [Lan et al., 2023] and the Copernicus Climate Change Service (C3S) [~~Copernicus, 2023e~~ Copernicus, 2023a]. The reasons for these increases are still debated, with studies attributing them to increases in wetland emissions and changes in the atmospheric methane sink to varying degrees. The main sink of methane is through reaction with the hydroxyl radical (OH) in the troposphere, ~~which itself is affected by various other atmospheric constituents. The rate of this reaction depends on the concentration of OH, which is determined by its photochemical sources and sinks.~~ Recent studies suggest that the steep decline of nitrogen dioxide (NO<sub>2</sub>) (~~Cooper et al., 2022~~) [Cooper et al., 2022], carbon monoxide (CO) and non-methane volatile organic compound emissions as a result of the measures introduced to control and limit the spread of the COVID-19 pandemic, lowered the levels of OH, and thus led to part of the increase in CH<sub>4</sub> concentrations in 2020 and 2021 (~~Stevenson et al., 2022; Laughner et al., 2021; Peng et al., 2022; Qu et al., 2022; Feng et al., 2023~~) [Stevenson et al., 2022; Laughner et al., 2022; Peng et al., 2022; Qu et al., 2022; Feng et al., 2023]. Additionally, enhanced wetland emissions, especially from tropical wetlands, contributed to the record increases of atmospheric CH<sub>4</sub> in 2020/21 (~~Peng et al., 2022; Feng et al., 2023, 2022; Qu et al., 2022~~) ~~—A potential uncertainty in studies using ground-based data originates in the high-latitude regions, where measurements are sparse and satellites can help with identifying methane emissions.~~ [Peng et al., 2022; Feng et al., 2023, 2022; Qu et al., 2022].

The Arctic contains large amounts of soil organic carbon (SOC) which is stored in the permafrost regions (ca. 1300 Pg) of which roughly 800 Pg is perennially frozen (~~Hugelius et al., 2014~~) [Hugelius et al., 2014]. The comparatively high temperature increase in the Arctic, compared to the rest of the world, also called “Arctic amplification“ (~~Serreze and Barry, 2011; Wendisch et al., 2017~~) [Serreze and Barry, 2011; Wendisch et al., 2017] may lead to increased permafrost degradation and rapid SOC loss (~~Plaza et al., 2019~~)

[[Plaza et al., 2019](#)] by the release of carbon dioxide (CO<sub>2</sub>) and/or methane. Latitudinally resolved growth rates are especially interesting in this regard and provided the initial motivation for this study.

65 In this paper we present methane growth rates and annual methane increases (AMIs) derived from Sentinel-5P/TROPOMI [XCH<sub>4</sub>](#) data using a Dynamic Linear Model (DLM) approach. In the second section we present the data used. Next, we describe our method for calculating [the these](#) growth rates, which is divided into four parts: (i) we discuss the preparation of the data; (ii) we provide a brief introduction into DLMs; (iii) we discuss our ensemble approach which utilizes cross validation to find the optimal DLM configuration for a given time series; (iv) we provide a method to calculate a bias related to the satellite  
70 sampling. In the fourth section we present global annual methane increases (AMIs) for the first 4.5 years of S5P/TROPOMI data and compare these to AMIs from other sources. In the fifth section we investigate zonal growth rates derived from 20° latitudinal bands to provide spatial information to the global AMIs. Additionally, we compare the growth rates to growth rates derived from CAMS global inversion-optimized methane data (CAMS/INV). In the sixth section we investigate CAMS/INV fluxes to help with the interpretation of our previous results. Finally, we summarize our results and discuss potential future uses  
75 of this method and suggestions for further research. [In the Appendix we provide additional information about our method and further results which are not included in the main text.](#)

## 2 Data

### 2.1 Sentinel-5P/TROPOMI WFMD product

The Sentinel-5 Precursor (S5P) satellite was launched on 13 October 2017 and has since delivered high quality data from  
80 its only scientific instrument, TROPOMI, which is a nadir viewing passive grating imaging spectrometer. Combined with a near-polar, sun-synchronous orbit, the swath width of 2600 km provides daily global coverage. Due to the orbit geometry and swath overlap multiple observations per day are possible in the polar regions. The spatial resolution depends on the bands and is  $5.5 \times 7$  km<sup>2</sup> for the short-wave infrared (SWIR) band ( $7 \times 7$  km<sup>2</sup> before August 2019) [[Ludewig, 2021](#)]. Methane is retrieved from TROPOMI measurements of sunlight reflected by the Earth's surface and atmosphere in the SWIR wavelengths.  
85 We use the latest release of the WFMD product (v1.8) [[Schneising et al., 2023](#)], which includes processing improvements such as an increased polynomial degree (cubic instead of quadratic) and an updated digital elevation model to account for various localized topography related biases [[Hachmeister et al., 2022](#)]. Furthermore, the machine-learning based quality filter in the post-processing is improved to further reduce scenes with residual clouds. We use data with a quality flag  $qf = 0$  (good) and don't include data with  $qf = 1$  (potentially bad). The WFMD product includes measurements for solar zenith angles up to 75°.  
90 ~~Data is currently available from 11.2017~~ [We performed this analysis using data from 05.2018](#) to 02.2023, ~~however, we exclude~~ [excluding](#) data from the commissioning phase (11.2017 to 04.2018).

## 2.2 CAMS global inversion-optimised greenhouse gas fluxes and concentrations (CAMS/INV)

The Copernicus Atmospheric Monitoring Service (CAMS) global inversion-optimised greenhouse gas fluxes and concentrations dataset (CAMS/INV) provides ~~model~~ data for carbon dioxide, nitrous oxide and methane. The methane data is produced using the CAMS CH<sub>4</sub> Flux Inversion system [Segers *et al.*, 2022], which is based on the TM5-4DVar inverse modeling system [Bergamaschi *et al.*, 2010, 2013]. We use release v22r1, where only ground-based observations from the NOAA network are used in the inversion (CAMS/INV-SRF) and release v21r1s, which includes satellite observations from the Greenhouse Gases Observing Satellite (GOSAT) in addition to ground-based observations (CAMS/INV-SRF-SAT). In our analysis we use the total column dry-air mole fractions and surface fluxes of methane from this dataset. The data is provided on a 2° × 3° grid from 1990 to 2022. We only apply our DLM approach to the methane concentrations from this data and use the corresponding fluxes directly to help with interpretation.

## 2.3 NOAA CH<sub>4</sub> Marine Boundary Layer Reference

The Marine Boundary Layer Reference (MBLR) is a 2-dimensional matrix (time vs. latitude) created from weekly air samples from the Cooperative Air Sampling Network [Dlugokencky *et al.*, 2021], which is created for various long-lived trace gases by NOAA-GML. The MBLR is created by first fitting the weekly data whereby the CH<sub>4</sub> level, seasonal component and short-term variations are separated. For each time step (48 evenly distributed per year) the different stations give a latitudinal distribution of CH<sub>4</sub> which is then smoothed. The global mean is calculated by averaging the smoothed latitudinal distribution for each time step. A detailed explanation can be found on the NOAA website [NOAA, 2022].

## 2.4 Univ. Bremen C3S/CAMS satellite data (UB-C3S-CAMS)

Annual methane increases are published by the Copernicus Climate Change Service (C3S, [Copernicus \[2023a\]](#)[Copernicus \[2023b\]](#)) in the context of the European State of the Climate (ESOTC) assessment. Here we use data from the ESOTC 2022 [[Copernicus, 2023b](#)[Coper climate indicator section \[Copernicus, 2023e\]](#)[Copernicus, 2023a](#)]. The methane data as shown on that website are (i) time series of monthly values of the column-averaged mole fraction of atmospheric methane, XCH<sub>4</sub>, as derived from satellite data, and (ii) annual mean methane growth rates including uncertainty estimates as derived from this time series.

115

The XCH<sub>4</sub> time series corresponds to averaged satellite data over land in the latitude band 60° S – 60° N and covers the period January 2003 to December 2022. The underlying satellite XCH<sub>4</sub> data product for 2002–2021 is XCH4\_OBS4MIPS version 4.4 available from the Copernicus Climate Data Store (CDS, [Copernicus \[2023d\]](#)) website [[Copernicus, 2018](#)]. The data product is derived from the satellite instruments SCIAMACHY/ENVISAT, TANSO-FTS/GOSAT and TANSO-FTS-2/GOSAT-2. A previous version of this data product is described in [Reuter \*et al.\* \[2020\]](#). This data set is extended using a year 2022 satellite-derived XCH<sub>4</sub> data product, generated for the Copernicus Atmosphere Monitoring Service (CAMS, [CAMS \[2023\]](#)) (see [Copernicus \[2023e\]](#)[Copernicus \[2023a\]](#) for details).

120

The combined C3S/CAMS XCH<sub>4</sub> time series has been generated by the University of Bremen (UB) and is [in the following referred to as UB-C3S-CAMS data set](#). This data set is also used to derive annual mean methane growth rates for 2003–2022 using the method as described in *Buchwitz et al.* [2017], for XCO<sub>2</sub>, which has later also been applied to XCH<sub>4</sub> [*Reuter et al.*, 2020]. This ~~data set is in the following referred to as UB-C3S-CAMS data set~~ method provides a new time series from the monthly XCH<sub>4</sub> time series described above. The new time series is generated by computing the difference in XCH<sub>4</sub> for a given calendar month between two consecutive years (e.g., January 2019 and 2020). The time assigned to this difference is the mean time between the two months (e.g., mid-July 2019). The annual mean growth rate for a give year is the weighted average of all monthly difference values of that year.

### 3 Method

The method section is split into four parts. First we describe how the data is prepared for the DLM, meaning how we get from single observations to a time-series we can fit our model to. Next, we shortly introduce DLMs and provide information on the specific types of models we are using. In the third subsection we explain how we use an ensemble of DLMs and cross-validation to select the best model. Lastly, we describe how we estimate the bias related to imperfect satellite sampling.

#### 3.1 Data preparation

The XCH<sub>4</sub> data to be used in the DLM fitting are preprocessed onto a latitude-longitude grid with sufficiently homogeneous sampling in space and time. Initially, the WFMD XCH<sub>4</sub> data product is gridded onto a 2° × 2° grid. For this we assign each measurement to a single grid cell and calculate the weighted average of all measurements per cell. The measurements are weighted using the inverse measurement uncertainty to disadvantage measurements with high uncertainty. ~~To account for inhomogeneities in spatial and temporal sampling of the data due to seasonal gaps and missing data (e.g. due to clouds)~~ For example, reported uncertainties are higher for low albedo scenes. Thus, these scenes contribute less to the average. The coverage of the WFMD data is roughly 25% in all regions and mostly constant, except for a few days with lower coverage and the seasonal data gaps in high latitudes (see Fig. D1). ~~To account for inhomogeneities in spatial and temporal sampling~~ we apply the method described by ~~Sofieva et al. [2014]~~ Sofieva et al. (2014). This method quantifies the sampling distribution's inhomogeneity using a measure denoted as  $0 \leq H \leq 1$ , which is defined as a linear combination of the asymmetry  $A$  and entropy  $E$  of the data:

$$H = \frac{1}{2}(A + (1 - E))$$

$$H = \frac{1}{2}(A + (1 - E)) \quad (1)$$

$$A = 2 \frac{|\bar{x} - x_0|}{\Delta x} \quad (2)$$

$$E = \frac{-1}{\log_e(N)} \sum_i \frac{n(i)}{n_0} \log_e \left( \frac{n(i)}{n_0} \right) \quad (3)$$

155 **Values**—In equation 2, the mean location of measurements is given by  $\bar{x}$  (e.g. mean spatial position or mean time),  $x_0$  is the central point and  $\Delta x$  the width of the region. Equation 3 represents the normalized entropy with  $N$  as the number of bins (e.g. grid cells or time steps),  $n(i)$  the number of observations in bin  $i$  and  $n_0$  the sample size.

$A$  can be intuitively understood as the asymmetry of the sampling distribution. For example,  $A$  would be high if only measurements in the eastern hemisphere are present for a given day. In contrast,  $A$ , would be zero if the measurements are 160 symmetrically distributed around the central point. The normalized entropy is  $E = 1$  for perfectly homogeneous sampling patterns and gets lower for each missing measurements. The entropy does however not capture the distribution of the sampling pattern. Hence, a combination of both measures is used to quantify the homogeneity of the sampling distribution. Values of  $H$  close to zero indicate a homogeneous sampling distribution while values close to one indicate a very inhomogeneous distribution. Inaccurate estimates and spurious features can arise without accounting for this inhomogeneous sampling [Sofieva et al., 165 2014]. **We**—The inhomogeneity can be calculated in the temporal domain (for each grid cell) and in the spatial domain (for each time step).


We first calculate the temporal inhomogeneity ( $H_T$ ) for each grid cell, which quantifies how even and symmetric the data for each grid cell is distributed in the temporal domain.  $H_T$  tends to be higher in cells with sparse data coverage, which are 170 often found over the oceans and tropical rain forests due to high cloud coverage (see Fig. 1). We then filter cells with  $H_T > 0.5$ . This threshold value was chosen empirically to exclude cells in ~~regions with known coverage problems, such as areas heavily affected by clouds (see Fig. 1)~~ these regions with limited cloud-free coverage. Next, we calculate the spatial inhomogeneity ( $H_S$ ) within the designated sub-grid, such as a zonal band. This allows to identify days with very inhomogeneous coverage. The spatial inhomogeneity can be calculated along both spatial dimensions. Hence, we define  $H_S$  as the equally weighted 175 linear combination of the latitudinal and longitudinal spatial inhomogeneity:

$$H_S = 0.5 \cdot H_S^{lat} + 0.5 \cdot H_S^{lon} \quad (4)$$

We determine a limit,  $H_S^{lim}$ , as the median of  $H_S$  plus two standard deviations

$$H_S^{lim} = \tilde{H}_S + 2\sigma_{H_S} \quad (5)$$

and filter out days with  $H_S > H_S^{lim}$ . ~~This filtering process removes days with highly inhomogeneous coverage.~~ The equation 180 for  $H_S^{lim}$  was empirically chosen and yields reliable limits for different sub-grids. Figure 1 illustrates the spatial and temporal



**Figure 1.** (a) Temporal inhomogeneity ( $H_T$ ) for global XCH<sub>4</sub> WFMDv1.8 data between May 2018 and February 2023. Grid cells with  $H_T > 0.5$  are omitted during analysis (b) Spatial inhomogeneity ( $H_S$ ) for global XCH<sub>4</sub> WFMDv1.8 data. Days above the  $H_S$  threshold (black line) are omitted from analysis, the threshold is set by Eq. 5 which was empirically chosen.

inhomogeneity for global WFMDv1.8 data.

Finally, we compute the area-weighted average of the chosen sub-grid, generating a time series for further the analysis. To further mitigate sampling bias in the global average, we first average over longitudes and subsequently over latitudes. This approach assumes a faster mixing of background methane levels within zonal bands, while acknowledging greater latitudinal disparities. A more detailed description is given in Sec. 3.4. CAMS/INV XCH<sub>4</sub> data is already provided on a grid with complete coverage and no inhomogeneity treatment is necessary. The time series are hence-just-therefore calculated using the area-weighted average of the sub-grid. Since the DLM approach is based on the assumption that errors are present and normally distributed, we add a Gaussian noise with  $\sigma = 0.5\sigma = 0.2$  ppb to the CAMS/INV XCH<sub>4</sub> time series.

### 190 3.2 Dynamic Linear Model fit

To extract information about the methane growth rate from the time-series we first need to calculate the underlying XCH<sub>4</sub> level, that is the smoothly changing background concentration without seasonal or short-term variation. While a simple approach, such as fitting a polynomial plus a trigonometric function to model the seasonality, may be considered, it is insufficient due to the complex change in methane-XCH<sub>4</sub> levels observed in historical records [*Lan et al., 2023; Copernicus, 2023eCopernicus, 2023a*].

195 The use of a moving average is not suitable due to possible data gaps, especially for high latitude bands. Therefore, we employ dynamic linear models to fit the XCH<sub>4</sub> data, which allow for the trend (i.e. the slope of the level, the growth rate) to change over time and can deal with missing data. For the analysis of global methane growth we use AMIs, since they are already used by other sources similar to other relevant studies [*Dlugokencky, 2022; Copernicus, 2023eCopernicus, 2023a; Schneising et al., 2023*]and therefore allow easy comparison of the results. This enables the methane growth of different studies to be readily

200 compared. AMIs are defined as the difference in methane level between the January 1st of two consecutive years, which is equal to. This is a measure of the integrated growth rate over the same time span. For zonal bands we directly investigate the



growth rate instead of AMIs (see Sec. 5 for a more detailed description).


205 A dynamic linear model is a regression model that can handle observations of varying accuracy, missing data, non-uniform sampling and non-stationary processes. It allows some of its parameters to change over time and directly models the observed variability using unobserved state variables [Laine, 2020]. These DLM properties allow the analysis of not only global but also zonal methane data, which can have higher uncertainties and more gaps, especially in the higher latitudes. Additionally, the direct modeling of the data allows the partition of the signal into different components, like such as an underlying level and seasonal component, which can prove advantageous beyond the scope of this paper. ~~DLMs have been previously used to successfully model stratospheric ozone Laine et al., 2014, methane from different GOSAT retrievals to investigate the seasonal cycle and trend Kivimäki et al., 2019 and methane from ground-based remote sensing Karppinen et al., 2020. For a detailed description of DLMs, including their formulation as a special case of a space-state model, we refer readers to Durbin and Koopman [2012] and Harvey [1990]. For a more concise introduction to DLMs, we refer to Laine [2020].~~

215 A DLM can be formulated as a special case of a space-state-state-space model, that is a model which consists of some unobserved components (represented by a state vector) and the observation vector. The evolution of the state vector and the relation between observation- and state-vector are modeled by a set of equations. If these equations are linear we have a so-called dynamic linear model. The DLM we use consists of three main components. First, a slowly changing background level, which captures the long-term trend of the methane concentration. Second, a seasonal component is included to model variations arising from seasonal cycles. This component enables variations in the phase and amplitude of the seasonal cycle to be accounted for. Third, an autoregressive component is incorporated to model noise and residual correlations in the data, accounting for short-term effects. Additionally, Gaussian noise can be included to model part of the errors. The ability of DLMs to capture changing components over time is achieved by modeling these changes as Gaussian random walks, allowing for smooth transitions and adjustments. The variances of these Gaussian random walks determine the overall variability of a certain parameter 220 (e.g. trend). A detailed description of the model setup and the different DLM components can be found in Appendix A. ~~Once the structure of the model is decided on, it can be brought into matrix form and the DLM consisting of the observation vector and the set of equations can then be solved using a Kalman Filter Durbin and Koopman, 2012; Harvey, 1990; Laine, 2020. The Kalman Filter is a recursive algorithm that estimates the unobserved states of a system using observed data and model equations Kalman, 1960.~~

230

In general, the model parameters (e.g. variances, see Tab. A1 for a complete list) are not known beforehand and have to be determined. For this purpose, Maximum Likelihood Estimation (MLE) [Durbin and Koopman, 2012; Harvey, 1990] can be used. MLE is a statistical method that estimates the parameters of a model by maximizing the likelihood of the observed data given the model's assumptions. Note that the data uncertainties are not used in the MLE but are indirectly included during the data preparation (gridding) as described above. For the end-user various software packages exist which provide the implementation 235 of this procedure, leaving only the model configuration open for the user. In our study we use the *UnobservedComponents*





**Figure 2.** DLM fit for daily area-weighted global WFMDv1.8 data. (a) Shows the daily area-weighted global XCH<sub>4</sub> together with the level and level+seasonal components from the DLM fit (b) The trend or growth rate is the slope of the level, the dashed lines show the  $\pm 1\sigma$  uncertainty, (c) Shows the seasonal component which captures the seasonal cycle, (d) The AR(1) component captures residual correlations in the data, (e) The residual shows the difference between the fit and the data, (f) A histogram of the residual shows that it is roughly normally distributed.

class of the python *statsmodel* package (Version [0.13.20.14.0](#), [Perktold et al. \[2023\]](#)), [which provides the means to define a DLM and to fit it using MLE \(see \*statsmodels-developers \[2023\]\* for documentation\)](#). An overview of a DLM fit for globally averaged WFMD data can be seen in Fig. 2.

240

[DLMs have been previously used to successfully model stratospheric ozone \[Laine et al., 2014\], methane from different GOSAT retrievals to investigate the seasonal cycle and trend \[Kivimäki et al., 2019\] and methane from ground-based remote sensing \[Karppinen et al., 2020\]. For a detailed description of DLMs, including their formulation as a special case of a state-space model, we refer readers to \*Durbin and Koopman \[2012\]\* and \*Harvey \[1990\]\*. For a more concise introduction to DLMs, we refer to \*Laine \[2020\]\*.](#)

245

### 3.3 Ensemble approach and Cross Validation

The choice of model configuration is a non-trivial problem, which is impacted by prior knowledge, empirical testing and different quality measures. [Prior knowledge may motivate From prior knowledge the inclusion of a seasonal component ,since is](#)

250 inferred, because the existence of a seasonality in atmospheric methane concentrations is known. Empirical testing can show that the inclusion of an autoregressive component is necessary, since because the data contains residual short-term variations. ~~And by quality measures we refer~~ The term quality measures refers to measures that facilitate model selection, such as the mean squared error (MSE), which is defined as the mean of squared differences between model and data. Additionally, the DLM provides variances (squared standard deviations) for each component which can be used to compare models; models  
255 with lower ~~variances-uncertainty~~ in the level and seasonality (~~i.e. higher certainty & less variability~~) are preferable to models with high ~~variances-uncertainties~~ for these terms.

To avoid the need for manual model selection, we employ an ensemble approach, fitting a range of DLMs to the data and automatically selecting the best model. The ensemble consists of different DLM configurations, with varying components,  
260 as described in Appendix A. Additionally, we perform k-fold cross-validation (CV) ~~five~~ with  $k = 5$  folds for each DLM to calculate an average mean squared error (AMSE). During CV, the DLM is ~~trained~~ fitted on a portion of the data while leaving out another portion (the fold) for testing. The difference between the model fit and the fold is used to calculate the MSE, and the average MSE across all five folds per DLM provides the AMSE. Low AMSE values indicate a better model fit and help in selecting the best model for a given time-series. The final model selection is based on an aggregated score, defined as the sum  
265 of the AMSE, the variance of the level and the variance of the seasonal term:

$$S_{agg} = AMSE + \sigma_{level}^2 + \sigma_{seas}^2 \quad (6)$$

The inclusion of the variances ensures that the uncertainty of the level and seasonal components is considered in the selection criterion. This approach aims to select DLMs that provide good estimates of the underlying methane signal while avoiding overfitting and reliance on expert knowledge.

270

Different methods and measure can be used for model selection and may yield different results. We want to emphasize that the problem of model selection is non-trivial and different approaches may be suitable for different data and use-cases. Here we select the model which yields the highest certainty fit of the level and seasonal component (i.e. the ~~methane~~ XCH<sub>4</sub> signal without noise) while avoiding overfitting and manual selection. Furthermore, we want to mention that in most cases the differences  
275 between all models in an ensemble are rather small with the best models producing approximately the same results. However, the use of a single DLM configuration for all zonal bands is not feasible due to the inherent differences in the seasonal signal for each zonal band. Additionally, an over-specified model can lead to high uncertainties in the resulting fit.

To quantify the impact of model selection, we calculate a model selection bias  $\sigma_{model}^2$ , which is included in the error budget  
280 of all AMIs and growth rates. For ~~this we calculate AMIs or growth rates~~ global data we calculate the AMIs for all models in

an ensemble and determine the weighted variance  $\div$ :

$$\sigma_{Model/AMI}^2 = \frac{\sum_i (AMI_{avg} - AMI_i)^2 \tilde{\sigma}_i^{-2}}{\sum_i \tilde{\sigma}_i^{-2}}; \quad \tilde{\sigma}_i^2 = \sigma_{avg}^2 + \sigma_i^2$$

$$\sigma_{Model/trend}^2 = \frac{\sum_i (\nu_{avg} - \nu_i)^2 \tilde{\sigma}_i^{-2}}{\sum_i \tilde{\sigma}_i^{-2}}; \quad \tilde{\sigma}_i^2 = \sigma_{avg}^2 + \sigma_i^2$$

for each year of interest:

$$\sigma_{Model/AMI,j}^2 = \frac{\sum_i (AMI_{avg,j} - AMI_{i,j})^2 \tilde{\sigma}_{i,j}^{-2}}{\sum_i \tilde{\sigma}_{i,j}^{-2}}; \quad \tilde{\sigma}_{i,j}^2 = \sigma_{avg,j}^2 + \sigma_{i,j}^2 \quad (7)$$

where  $\nu_i$  is the  $AMI_{i,j}$  is the AMI for year  $j$  and model  $i$ ,  $AMI_{avg,j}$  the average AMI of all models in year  $j$ , and  $\sigma_{avg,j}$  and  $\sigma_{i,j}$  the corresponding uncertainties. For the case of growth rates we calculate a single model uncertainty which is averaged over all time steps:

$$\sigma_{Model/trend}^2 = \frac{1}{T} \sum_t \left[ \frac{\sum_i (\nu_{avg,t} - \nu_{i,t})^2 \tilde{\sigma}_{i,t}^{-2}}{\sum_i \tilde{\sigma}_{i,t}^{-2}} \right]; \quad \tilde{\sigma}_{i,t}^2 = \sigma_{avg,t}^2 + \sigma_{i,t}^2 \quad (8)$$

where  $T$  is the total number of time steps,  $\nu_{i,t}$  the growth rate for model  $i$  at time step  $t$ ,  $\nu_{avg,t}$  the average growth rate of all models at time step  $t$ ,  $\sigma_{avg,t}$  the uncertainty of the average growth rate at time step  $t$ , and  $\sigma_{i,t}$  the uncertainty of model  $i$  at time step  $t$ .

The contribution of the model selection bias to the error budget can be seen in Tab. 1 for global AMIs and Tab. 2 for zonal growth rates. In case of global data, the contribution is small for all years with  $\sigma_{Model} < 1$  ppb but can be higher in specific cases. For example, this is the case for global AMIs in except for 2018, when only an incomplete time series is available. For zonal data,  $\sigma_{Model}$  varies between the bands and is on in the range of 0.60-1.03-4.81-34 ppb/y.

### 3.4 Estimation of sampling bias

The spatio-temporal coverage of S5P XCH<sub>4</sub> data is limited mostly by cloud coverage, the polar nights and poorly reflective surfaces, while additional gaps may exist due to technical problems with the satellite platform. Additionally, the sampling distribution is not completely random but is influenced by the total land mass per latitudinal band and seasonal cloud coverage over the tropical and subtropical oceans. Therefore, we don't calculate the direct global average but perform initial longitudinal averaging to ensure equal weighting of all latitudes, which is crucial given the heterogeneous daily distribution of The daily gridded data for S5P XCH<sub>4</sub> is therefore always incomplete, meaning we have some grid cells without any measurements. Here we investigate the systematic sampling bias due to polar nights, the total effect due to sampling and the effect of our zonal first averaging approach the averaging method. For this, we use CAMS/INV XCH<sub>4</sub> data (see Sec. 2.2) onto which we apply different filtering masks and averaging methods. We then compare AMIs. Since CAMS/INV data has a complete coverage (due to being model data) we can investigate the effects of different sampling masks or averaging and compare it to the results gained from

the unmasked data.

310

We therefore compare AMIs (for global data) and growth rates (for zonal data) calculated using different sampling and/or methods. To simulate the spatio-temporal pattern of S5P sampling, we created a daily mask from gridded WFMDv1.8 data and applied it to the model data. To only simulate the systematic effect of the polar nights, we created a daily mask using the average solar zenith angle (SZA) per grid cell with a cut-off value of  $75^\circ$ . The AMIs and growth rates for CAMS/INV XCH<sub>4</sub> data were calculated using the same ensemble approach used for WFMD data.

315

First, we investigate the effect of ~~our zonal-first averaging approach~~ the averaging method. We compare standard averaging, which is defined in this study as the area-weighted mean of all grid cells in a region, with an approach we call zonal-first averaging. Zonal-first averaging takes into account the inhomogeneous sampling at each latitude, which is influenced by the distribution of land mass and seasonal coverage. Since zonal transport occurs within weeks [Jacob, 1999], we first average the grid cells zonally, assuming that the available data within a zonal band provides a good estimate of the mean zonal XCH<sub>4</sub>. Subsequently, we calculate the global average from the zonal averages, which leads to a consistent weighting of all latitudes regardless of their individual coverage. Figure 3 shows AMIs calculated from CAMS/INV-SRF XCH<sub>4</sub> data using ~~(i) no-a) No~~ mask and standard averaging ~~(ii) the-b) The~~ S5P XCH<sub>4</sub> mask and standard averaging and ~~(iii) the-c) The~~ S5P XCH<sub>4</sub> mask and ~~zonal-first-zonal-first~~ averaging. Using ~~zonal-first-zonal-first~~ averaging the AMIs are closer to the AMIs derived from the complete data. This is especially visible for 2020, where the AMI is overestimated by roughly 3 ppb when using the standard averaging. Therefore, we use ~~zonal-first-zonal-first~~ averaging for all calculations of globally averaged data. ~~For zonally-averaged data we use standard averaging.~~

320

325

330

Figure 4 shows zonal growth rates calculated from CAMS/INV-SRF XCH<sub>4</sub> data using (a) No mask and standard averaging (b) The S5P XCH<sub>4</sub> mask and standard averaging and (c) The S5P XCH<sub>4</sub> mask and zonal-first averaging. Growth rates calculated on the masked data show  $1\sigma$  agreement with the growth rates calculated on the complete data. Growth rates calculated using the zonal-first averaging show better agreement for the  $50^\circ$ – $70^\circ$ S band while for the  $10^\circ$ S– $10^\circ$ N band the growth rate is more variable. For all other zonal bands the results are nearly identical. Noticeably, the  $70^\circ$ – $90^\circ$ N band shows no variation when using the S5P XCH<sub>4</sub> mask, indicating that insufficient data coverage hinders the detection of growth rate variation. Since sampling within a  $20^\circ$  zonal band can still vary with latitude we follow the same reasoning as in the previous paragraph and also use zonal-first averaging for zonal growth rate calculation.

335

Next, we investigated the ~~remaining bias due to sampling effect of satellite sampling and the polar nights by applying~~ corresponding masks to CAMS/INV XCH<sub>4</sub> data. For global data the sampling biases are calculated by taking the squared difference between the AMI calculated on the complete data (without any sampling filtering) and the AMI calculated on the

340

**Table 1.** Error budget for global AMIs.  $\sigma_{DLM}$  is the uncertainty provided by the DLM fit,  $\sigma_{Model}$  is the uncertainty from model selection and  $\sigma_{Sampling}$  is the bias due to satellite sampling. All values show  $\pm 1\sigma$  uncertainties.

Year	2018 <sup>†</sup>	2019	2020	2021	2022
$\sigma_{DLM}$	1.70	0.40	0.39	0.39	0.49
$\sigma_{Model}$	3.22	0.71	0.32	0.48	0.35
$\sigma_{Sampling}$	2.96	0.25	0.19	0.26	0.25
$\sigma_{Sampling(SZA)}$	0.06	0.37	0.05	0.22	0.73
$\sigma_{Total}$	4.70	0.85	0.53	0.67	0.65

All values are in ppb, <sup>†</sup> 2018 only includes data starting 01.05.2018.

~~reduced-masked~~ CAMS/INV data. We calculate a separate bias for each year in our analysis.

$$\sigma_{Sampling}^2 = (AMI - AMI_{Sampling})^2 \quad (9)$$

$$\sigma_{SZA}^2 = (AMI - AMI_{SZA})^2 \quad (10)$$


345 Table 1 shows the resulting errors for global AMIs. The sampling bias is around 0.25 ppb for all fully available years and much higher for 2018 with ~~2.92~~ 2.96 ppb. The contribution of the SZA related bias varies from year to year with a maximum value of 0.73 ppb in the year 2022. This variability might be related to the varying difference between the high- and mid-latitudes during the different years. When the difference between both is bigger, the masking of high-latitude regions is expected to have a larger effect on AMIs. For the analysis of zonal bands we calculate ~~an a zonal~~ error by taking the average squared difference  
350 between growth rates derived from the complete data and growth rates derived from the reduced CAMS/INV ~~data~~ XCH<sub>4</sub> data for each band:

$$\sigma_{Sampling/Zonal}^2 = \frac{1}{N} \sum_i^N (\nu_i - \nu_i^{Sampling})^2 \quad (11)$$

where  $\nu_i$  is the growth rate at time step  $i$  and  $N$  is the total number of data points. The sampling errors for zonal growth rates are shown in Tab. 2. Higher sampling errors correlate clearly with regions of high temporal inhomogeneity which ~~visualizes~~ regions with further shows the challenging sampling conditions in these regions (see Fig. 1).  
355

#### 4 Comparison of different global annual methane increases

In this section, we discuss global AMIs calculated using WFMDv1.8 data from May 2018 to February 2023. The results are shown in Fig. 5. An overall, although non-linear, rise in methane level is observed between 2019 and 2022. The most significant change occurs from 2019 to 2020, with an increase from  $6.89 \pm 0.85$  ppb to  $14.40 \pm 0.53$  ppb. The highest AMI is observed




**Figure 3.** Global AMIs derived from CAMS/INV-SRF  $XCH_4$  data using ~~the complete data~~ (a) The non-masked dataset with standard averaging, ~~the (b) The~~ S5P sampled CAMS/INV-SRF data with standard averaging and ~~the (c) The~~ S5P sampled CAMS/INV-SRF data using our ~~zonal-averaging first~~ zonal-first averaging approach.

360 in 2021, reaching  $16.93 \pm 0.67$  ppb. The globally averaged  $XCH_4$  has risen from  $1817.32 \pm 2.81$  ppb at the beginning of 2018 to  $1878.14 \pm 0.16$  ppb at the end of 2022.

To validate our findings, we compared our results with AMIs determined by *Schneising et al.* [2023], the NOAA–GML [*Dlugokencky*, 2022], and data generated for the ~~Copernicus Climate Change Service C3S~~ [*Copernicus*, 2023e] Copernicus, 2023a].  
 365 Additionally, we include AMIs derived using our DLM approach for monthly WFMDv1.8 data, NOAA–GML MBLR data and the UB–C3S–CAMS dataset. Table 3 and Fig. 7 provide a comparison of AMIs between 2018 and 2022. While absolute values may differ due to variations in data and methods, all AMIs exhibit the same qualitative trend. Differences are expected ~~specifically for for various reasons~~. First, the difference in sampling. NOAA–GML AMIs, ~~which~~ are based on surface flask measurements rather than ~~the satellite~~ total column observations used in the other calculations. ~~To assess~~ Second, different  
 370 methods are used to derive AMIs from the data (see the corresponding sources for description the other methods). And lastly, depending on the time resolution of the data and method used, the AMIs represent either the difference between the 1st of January of two consecutive years or the difference between the monthly January average of two consecutive years. The former is the case for our DLM approach and the AMIs derived by the NOAA–GML. The latter is the case for the AMIs derived by Schneising et al. [2023] and for the C3S AMIs.

375



**Figure 4.** Zonal growth rates derived from CAMS/INV-SRF XCH<sub>4</sub> data using (a) The non-masked dataset with zonal-first averaging, (b) The S5P sampled CAMS/INV-SRF data with standard averaging and (c) The S5P sampled CAMS/INV-SRF data using our zonal-first averaging approach.

To discern the impact of data and methodology, AMIs for the UB-C3S-CAMS and NOAA-GML datasets created using different methods can be compared. Our DLM based AMIs agree well of these datasets agree within  $1\sigma$  with the calculations done for the C3S and by the NOAA-GML, indicating no significant differences due to the method used. However, we see a comparatively high AMI in 2022 for the combination of DLM and NOAA-GML data (see Fig. 7). This is probably related to the higher uncertainty for AMIs at the start/end of a time series, which can also be seen for UB-C3S-CAMS AMIs. A longer input time series will likely lead to a reduction in uncertainty and to a reduced deviation compared to the other 2022 AMIs. An application of our DLM approach for the complete UB-C3S-CAMS and NOAA-GML MBLR data can be seen in Appendix B.

Additionally, we used our DLM approach on monthly WFMDv1.8 data, to better compare our method to the method used by Schneising *et al.* [2023], which also shows good agreement. Our results are therefore in good agreement with existing results from Schneising *et al.* [2023], the NOAA-GML Lan *et al.*, 2023 and the C3S Copernicus, 2023e, when accounting for the difference in methodology or data used. No significant differences agreement within  $1\sigma$ . Only differences smaller than  $1\sigma$  are found when comparing AMIs based on the same data or method but different methods. We also applied our approach to CAMS/INV XCH<sub>4</sub> data (see: 2.2), for which global AMIs can be seen in Fig. 6. The AMIs are in qualitative agreement and show the same structure over the 5-year period, however significant differences in absolute values are observed for 2019




**Table 2.** Sampling and model error for zonal growth rates.  $\sigma_{DLM}$  is not included in this table since it varies with time but is shown in Fig. 8. All values show  $\pm 1\sigma$  uncertainties.

Band	$\sigma_{Model}$	$\sigma_{Sampling}$	$\sigma_{Model} \sigma_{Sampling(SZA)}$
70–90° N	2.84–1.44	0.78–2.70	0.14
50–70° N	0.57–1.92	0.97–1.30	0.61
30–50° N	1.35–1.87	2.06–1.46	0.00
10–30° N	2.02–3.63	4.81–1.52	0.00
-10–10° N	0.93–3.34	2.81–2.65	0.00
10–30° S	0.86–1.73	1.93–1.87	0.00
30–50° S	2.33–1.03	0.60–3.98	0.00
50–70° S	3.49–2.95	3.95–1.46	1.85
70–90° S	1.82–4.34	2.02–1.94	0.67


All values are given in ppb/yr.

2020 and 2022. For 2019 the CAMSAMI seems to be in agreement with 2020 the CAMS/INV AMI is significantly lower compared to most other sources (see Tab. 3), while for 2022 CAMS is the clear outlier. the CAMS/INV AMI is significantly higher compared to most other sources (see Tab. 3).

395 The AMIs for 2020 and 2021 are the largest observed since NOAA began systematic records in 1983. The drivers contributing to these record increases have been the subject of recent debate and can be attributed to a rise in emissions, a reduction of the CH<sub>4</sub> sink, or a combination of both effects. According to the International Energy Agency (IEA) methane emissions from the energy sector decreased by approximately 10 % in 2020 [IEA, 2021]. However, additional emissions due to reduced maintenance of landfills and oil and gas infrastructure can be expected according to Laughner et al. (2021), while McNorton et al. (2022) suggest that the effect of the global slowdown on anthropogenic CH<sub>4</sub> emissions is relatively small. Some studies propose that the reduction of the OH sink, caused by decreased emissions of nitrogen oxides during the COVID-19 pandemic, may explain part of the increase (Stevenson et al., 2022; Laughner et al., 2021; Peng et al., 2022; Qu et al., 2022; Feng et al., 2023) [Stevenson et al., 2022; Laughner et al., 2021; Peng et al., 2022; Qu et al., 2022; Feng et al., 2023]. Specifically, Stevenson et al. (2022); Peng et al. (2022) suggest that approximately half of the increase can be attributed to this effect. Conversely, multiple  
405 several other studies attribute the majority of the CH<sub>4</sub> increase in 2020 to the growth in wetland emissions (Qu et al., 2022; Feng et al., 2023, 2022; Zhang et al., 2023).



**Figure 5.** Global annual methane increases derived from Sentinel-5P/TROPOMI WFMDv1.8 data. The errorbars show the  $\pm 1\sigma$  uncertainty and include the DLM, sampling and model error.




**Figure 6.** Global annual methane increases derived from CAMS global inversion-optimised greenhouse gas concentrations including only surface observations. The errorbars show the  $\pm 1\sigma$  uncertainty and include the DLM and model error, which are also shown in brackets.

**Table 3.** Comparison of global AMIs using different data and methods. The All errors represent  $\pm 1\sigma$  uncertainties.

<u>Method@Dataset (time resolution)</u>	2018	2019	2020	2021	2022
DLM@WFMDv1.8 (daily)	$9.74 \pm 4.70$ †	$6.89 \pm 0.85$	$14.40 \pm 0.53$	$16.93 \pm 0.53$	$16.93 \pm 0.53$
DLM@WFMDv1.8 (monthly)	$6.56 \pm 4.23$ †	$7.85 \pm 0.98$	$14.39 \pm 0.93$	$16.55 \pm 0.93$	$16.55 \pm 0.93$
<i>Schneising et al.</i> [2023] @WFMDv1.8 (monthly)		$7.80 \pm 0.60$	$15.00 \pm 1.00$	$16.40 \pm 1.00$	$16.40 \pm 1.00$
NOAA–GML (Version <del>2023-07</del> <u>2023-10</u> ) @NOAA MBLR ( <u>daily</u> ) *	<del><math>8.70 \pm 0.52</math></del> <u><math>8.76 \pm 0.52</math></u>	<del><math>9.70 \pm 0.60</math></del> <u><math>9.68 \pm 0.60</math></u>	<del><math>15.20 \pm 0.41</math></del> <u><math>15.16 \pm 0.41</math></u>	<del><math>17.75 \pm 0.41</math></del> <u><math>17.71 \pm 0.41</math></u>	<del><math>17.75 \pm 0.41</math></del> <u><math>17.71 \pm 0.41</math></u>
DLM@NOAA MBLR ( <u>daily</u> ) *	<del><math>9.37 \pm 0.89</math></del> <u><math>9.35 \pm 0.89</math></u>	<del><math>8.64 \pm 0.77</math></del> <u><math>8.60 \pm 0.77</math></u>	<del><math>15.97 \pm 0.97</math></del> <u><math>15.99 \pm 0.97</math></u>	<del><math>17.49 \pm 0.97</math></del> <u><math>18.10 \pm 0.97</math></u>	<del><math>17.49 \pm 0.97</math></del> <u><math>18.10 \pm 0.97</math></u>
<i>Buchwitz et al.</i> [2017] @UB–C3S–CAMS ( <u>monthly</u> )	$10.19 \pm 1.96$	$9.00 \pm 2.01$	$15.19 \pm 2.09$	$17.09 \pm 2.09$	$17.09 \pm 2.09$
DLM@UB–C3S–CAMS ( <u>monthly</u> )	$10.15 \pm 1.13$	$8.92 \pm 1.30$	$15.77 \pm 1.20$	$17.04 \pm 1.20$	$17.04 \pm 1.20$
DLM@CAMS/INV–SURF ( <u>daily</u> )	$6.24 \pm 0.97$	$9.86 \pm 0.39$	$13.34 \pm 0.38$	$18.04 \pm 0.38$	$18.04 \pm 0.38$

All values are in ppb. Uncertainties reflect one standard deviation. † Larger error since only data starting with May 2018 was used. \* Input data has a weekly time resolution but AMIs are provided of January of two consecutive years



**Figure 7.** Comparison of global AMIs listed in Tab. 3. The labels are formatted as Method@Dataset. Colors indicate the type of dataset used and the markers denote the method. All errors represent  $\pm 1\sigma$  uncertainties.

## 5 Investigation of zonal methane growth rates

In addition to our global analysis, we ~~also~~ investigated  $20^\circ$  zonal bands. ~~While the~~The good spatio-temporal coverage of S5P XCH<sub>4</sub> might suggest that ~~this would allow for the identification of~~ the same approach of using AMIs could be applied to

410 ~~identify~~ zonal bands with anomalous methane increases ~~using zonal AMIs, the~~. ~~However, the impact of~~ atmospheric transport  
has to be considered. While longitudinal mixing occurs on timescales of a few weeks, meridional transport is slower, taking  
1-2 months between mid-latitudes and tropics or polar regions and around a year between hemispheres [Jacob, 1999; Warneck,  
1999]. The relatively longer atmospheric lifetime of 9.1 years [Szopa *et al.*, 2021] compared to the mixing times, therefore  
guarantees a relatively even latitudinal distribution of methane in the troposphere, where the main difference is driven by the  
415 uneven distribution of CH<sub>4</sub> sources [Warneck, 1999]. Thus, we need to sample at about 1 year or less to observe differences  
between hemispheres and 1 month or less to observe differences within a hemisphere. The daily sampling of S5P is hence  
faster than meridional transport, however part of the temporal information gets lost when using AMIs which are obtained by  
integrating the growth rate over one year. Thus, we ~~look at~~ ~~investigate~~ the growth rate, which is the trend component of our  
DLM fits, to obtain a better temporal resolution of the ~~potential~~-zonal signals.

420

The results are shown in Fig. 8 and include growth rates derived from CAMS/INV-SRF XCH<sub>4</sub> data for comparison. The  
shown errors include the uncertainty gained from the DLM fit  $\sigma_{DLM}$ , the model selection error  $\sigma_{Model}$  and the sampling error  
 $\sigma_{Sampling}$  (see Sec. 3.3 & 3.4). Growth rates are similar within a hemisphere, while differences between hemispheres are  
clearly visible. Additionally, no significant sub-annual variations in zonal growth rates are present. Both observations are in  
425 good agreement with the ~~know-known~~ atmospheric mixing times and indicate that our data currently allows for identification  
of inter-hemispheric differences while short-term variations between zonal bands are not detected. The high latitude band be-  
tween 70°–90° is included for completeness, but shows no inter-annual variability. This may be due to a lack of real change  
in growth rates in this region ~~and/or~~, high uncertainties present in the data ~~and/or the sparse data coverage~~. The corresponding  
CAMS/INV-SRF XCH<sub>4</sub> growth rate indicates that the variability in growth rate is relatively small in this band, which supports  
430 the first point of our explanation. We therefore exclude this band from our following discussion. Hence, we mean bands be-  
tween 10° S and 90° S when we speak of the Southern Hemisphere (SH) and bands between 10° S and 70° N when we speak  
of the Northern Hemisphere (NH). The band between 10° S and 10° N represents the boundary region and is close to the global  
background as can be seen in Fig. 9, which presents zonal growth rate anomalies. Zonal growth rate anomalies are defined as  
the difference between the zonal and the global growth rate for each band.

435

Overall growth rates derived from WFMD data are close to growth rates derived from CAMS/INV-SRF XCH<sub>4</sub> data, with  
~~the strongest differences in the 50S and 70S band, which is also one of the most challenging bands to fit, due to the high cloud  
coverage and low landmass. an overall agreement within 1 $\sigma$~~ . The growth rates and the growth rate anomalies can be used to  
interpret the changes in global AMIs and allow the identification of hemispheres or zonal bands with anomalous growth rates.  
440 Differences between the hemispheres can be especially well seen in the zonal growth rate anomalies. During 2019 a decrease  
in growth rates can be observed for the whole SH (~~except the southernmost band~~), while growth rates in the NH increased or  
stayed stable. For 2020 growth rates ~~for all SH bands~~ increase strongly from roughly 0 ppb/y to 20 ppb/y. The NH growth  
rates increase more slowly, except for the 50° N – 70° N band which exhibits a small decrease in growth rate. During 2021  
most zonal growth rates move towards or around the global mean, with the strongest anomaly visible in the 10° N – 30° N band


445 which shows some additional increase in growth rate, which peaks mid of the year. During 2022 a clear difference between the hemispheres can be seen again, with a decrease of growth rates in the NH and an increase or stabilization of growth rates in the SH. This difference is especially clear when looking at the zonal growth rate anomalies in Fig. 9.

Recent studies, which discuss the record methane increases in 2020 and 2021 can help with interpreting the structure of  
450 zonal growth rates. *Peng et al.* [2022] employ an atmospheric inversion using ground-based data. They attribute the increase from 2019 to 2020 roughly equally to changes in the OH sink and an increase in wetland emissions located mainly in the NH. In contrast, studies based on the inversion of satellite data from the Japanese Greenhouse gases Observing SATellite (GOSAT) state that the majority of increase from 2019 to 2020 can be attributed to the African continent [*Feng et al.*, 2023; *Qu et al.*, 2022] with additional increases in tropical South America in 2021 [*Feng et al.*, 2023]. Our findings can thus be seen as  
455 aligning with recent studies by *Feng et al.* [2023]; *Qu et al.* [2022]. The increase in SH growth rates from 2019 to 2020 ~~can be associated~~ is consistent with increased wetland emissions. The rise in the NH latitudinal bands during 2020 can be explained by the decreasing OH sink, primarily located in the NH [*Peng et al.*, 2022; *Feng et al.*, 2023]. However, the continued increase in 2021 cannot be solely explained by the OH sink, as OH levels mostly recovered in that year according to *Feng et al.* [2023] and [*Peng et al.*, 2022]. ~~It is likely that~~ Possible explanations for the ongoing increase ~~is related to~~ are persistent wetland  
460 emissions *Feng et al.* [2023], as well as the return to pre-pandemic methane emissions from the energy sector in 2021 [*IEA*, 2023]. Finally, the decrease of growth rates in the Northern NH and increase of growth rates in the SH during 2022 hasn't been discussed to our knowledge. Our results therefore indicate that the decrease in global AMI from 2021 to 2022 can be attributed to a reduced growth rate in the NH. We further investigate this in the next section.


## 6 CAMS/INV Fluxes

465 As mentioned before, zonal growth rates provide information about the change of methane concentration in a given zonal band, including changes in sources, sinks and transport patterns. These transport patterns would average out for global AMIs given a perfect coverage. ~~Comparison of our global AMIs with global AMIs from other sources indicate,~~ In Sec. 3.4 we applied a S5P XCH<sub>4</sub> mask to CAMS/INV XCH<sub>4</sub> data (see Fig. 3) and compared AMIs calculated from this masked data with AMIs calculated from the complete data. Since differences between these AMIs are small, we conclude that the effect of ~~transport~~  
470 related sampling biases ~~seems to be~~ is limited. Therefore, changes in global AMIs can be attributed to the total source-sink balance of methane and not to changes in transport patterns. Whether this is also true for zonal growth rates is less clear, since transport effects are expected to be stronger especially within hemispheres. ~~The similarity~~

The agreement within 1 $\sigma$  of zonal growth rates derived from WFMD and CAMS/INV XCH<sub>4</sub> data in Fig. 8, suggest that the  
475 structures observed in our zonal growth rates are not artifacts from sampling related biases. However, we cannot rule out transport effects from this comparison, meaning we can't clearly attribute changes in hemispheres or zonal bands over the years to a change in the source-sink balance. Hence, we also investigated the change of surface fluxes ~~from~~ between consecutive years




**Figure 8.** Zonal growth rates for  $20^\circ$  bands derived from (a) Sentinel-5P/TROPOMI WFMDv1.8 data and (b) CAMS/INV-SRF  $XCH_4$  data. The errors show the  $\pm 1\sigma$  uncertainty.



**Figure 9.** Zonal growth rate anomalies for  $20^\circ$  bands derived from Sentinel-5P/TROPOMI WFMDv1.8 data. The anomalies are defined as the difference between zonal and global growth rates.

which are readily available for the CAMS/INV data ~~between consecutive years~~. In Figure 10, 11 and 12 we present total, wet-




**Figure 10.** Difference between total surface fluxes from the CAMS/INV-SRF data.


land and other fluxes for non-wetland fluxes from CAMS/INV-SRF data respectively. ~~Additionally, rice and biomass burning~~  
 480 ~~fluxes exist, which are not shown here.~~ The category of other non-wetland fluxes includes all other anthropogenic emissions  
 as well as contributions from oceans, wild animals, the soil sink ~~and termites~~. ~~Overall a great change in fluxes can be observed~~  
~~between all investigated years~~, termites and biomass burning.

Large changes in fluxes are identified between all the years investigated. The wetland flux difference between 2019 and  
 485 2020 indicate a strong increase in the NH as indicated by Peng et al., 2022 Peng et al. [2022] as well as some increase in  
 the SH wetlands as reported by Feng et al., 2023 ~~and~~ Qu et al., 2022 Feng et al. [2023] ~~and~~ Qu et al. [2022]. We expect the SH  
 wetland fluxes to be underestimated as indicated by Feng et al. [2023] since because the CAMS/INV-SRF data is only-based on  
 ground-based measurements from the NOAA network, similar to the inversion performed by Peng et al. [2022], due to the poor  
 coverage in the tropics. Interestingly, wetland fluxes from CAMS/INV-SRF-SAT data including satellite measurements from  
 490 GOSAT show stronger SH wetland emissions between 2019 and 2020 as shown in Figure 13. Additionally, an increase in other  
non-wetland fluxes occurs between 2019–2020 and 2020–2021. In the first case these increases are mainly focused on China,  
 while in the second case additional increases over the Indian subcontinent can be seen. Between 2021 and 2022 a clear decrease  
 of total surface fluxes can be seen in large parts of the NH, while strong increases can be observed over the whole of south  
South America. The large decreases in the NH can be clearly attributed to changes in the other non-wetland fluxes, while the  
 495 increase over south-South America seems to involve a combination of wetland and other fluxes. ~~This investigation of~~ Therefore,  
 CAMS/INV fluxes indicates imply that the changes in zonal growth rates we observed both in WFMD and CAMS/INV are  
 not merely due to changes in transport patterns but correlate with changes in surface methane fluxes between the years. This  
conclusion is strengthened by the qualitative agreement of the flux changes between CAMS/INV-SRF and the aforementioned  
studies by Peng et al. [2022]; Feng et al. [2023]; Qu et al. [2022]. However, further research is needed to substantiate these  
 500 inferences.






**Figure 11.** Difference between wetland surface fluxes from the CAMS/INV-SRF data.



**Figure 12.** Difference between other surface fluxes from the CAMS/INV-SRF data.

## 7 Conclusions

In this study, we presented a DLM based approach to calculate methane growth rates and AMIs from S5P/TROPOMI data. We addressed sampling-related biases by comparing AMIs and growth rates derived from CAMS/INV XCH<sub>4</sub> data both with and without S5P XCH<sub>4</sub> sampling. Further, we included a bias related to the model selection in our error budget. Our calculations of global AMIs based on WFMDv1.8 data from 2018 to 2022 demonstrate good agreement with other AMIs. Additionally, we separated the influence of the fitting method and the underlying data by applying our DLM approach to other datasets. We show that using the same ~~method or the same~~ input data results in agreement within  $\pm 1\sigma$  between all AMIs (~~except for CAMS/INV AMIs~~). ~~Moreover,~~ Using the same method but different input data results in qualitative agreement but with differences larger than  $1\sigma$ . Nevertheless, the consistency of AMIs derived from diverse datasets, such as ground-based data from NOAA and ~~total-column-dry-air-mole-fraction~~ data from WFMDv1.8 and UB-C3S-CAMS, highlights the robustness of these various



**Figure 13.** Difference between total surface fluxes from the CAMS/INV-SRF-SAT data. No Flux difference is available for 2022 - 2021, since the dataset currently ends in 2021.

approaches. The record methane increase in 2020 and 2021 is therefore well ~~documented using different data and methods identified in the different data sets, which use different methods to assess AMIs.~~ The underlying factors driving these increases, as discussed in Section 4, remain however a subject of debate.

515 In addition to global AMIs we investigated growth rates for 20° zonal bands which provide spatial information to the global AMIs. We argue that this is possible due to (a) the faster zonal mixing in comparison to meridional mixing and (b) the faster satellite sampling in comparison to the meridional mixing times. ~~Comparisons Firstly, comparisons~~ of zonal growth rates from S5P/TROPOMI data with growth rates from CAMS/INV-SRF ~~data show good agreement and XCH<sub>4</sub> data show agreement within 1σ.~~ Additionally, we investigated growth rates calculated from CAMS/INV-SRF XCH<sub>4</sub> data filtered using a S5P XCH<sub>4</sub> ~~mask which~~ indicate that no significant sampling biases exist for ~~zonal bands~~the zonal band approach. Still we want to emphasize that meridional transport ~~affects can affect~~ the zonal growth rates, meaning they don't ~~only necessarily~~ indicate changes in the sources and sinks of methane but might also show systematic changes in transport patterns. The zonal growth rates exhibit clear differences between the hemispheres for 2019 and 2022, ~~while whereas~~ growth rates are more similar for 2020 and 2021. Differences within a hemisphere are mostly smaller and no additional short-term variations are visible, which might reflect the   
525 well-mixed state of the atmosphere within a hemisphere. The low growth rates in the SH in 2019 and subsequent increases suggest a rise in atmospheric methane in that region, possibly driven by tropical wetland emissions according to *Feng et al.* [2023]; *Qu et al.* [2022]; *Zhang et al.* [2023]. Other factors potentially contributing to these changes include variations in the OH sink due to pollutant reductions during the COVID-19 pandemic [*Feng et al.*, 2023; *Qu et al.*, 2022; *Peng et al.*, 2022] and the changes in global methane emissions due to the COVID-19 pandemic and the subsequent recovery [*IEA*, 2021].

530

We further investigated this inter-hemispheric differences by investigating the surface fluxes ~~available~~ from CAMS/INV data, ~~which are consistent with the derived zonal growth rates.~~ ~~We argue that this is possible since (a) growth rates derived~~

from WFMDv1.8 data are similar to growth rates from CAMS/INV data and (b) no significant sampling bias is present as we showed in Sec. 3.4. The total surface fluxes show clear changes between the years, ~~indicating that the changes in zonal growth rates are not only due to changes in transport patterns. The~~ and the partition into wetland and ~~other non-wetland~~ (mainly anthropogenic) fluxes allows further interpretation ~~and shows~~. ~~Furthermore, changes in fluxes show~~ reasonable qualitative agreement with findings reported by *Feng et al.* [2023]; *Qu et al.* [2022]; *Peng et al.* [2022].

~~In addition to the confirmation of known results, new conclusions are also drawn.~~ Most notably, the decrease ~~in~~ of the global AMI in 2022 ~~can be attributed to~~ is caused by reduced NH zonal growth rates. This is clearly visible in zonal growth rates derived from S5P XCH<sub>4</sub> and CAMS/INV XCH<sub>4</sub> data. Investigation of the corresponding model surface fluxes, indicates that ~~changes in zonal growth rates are consistent with~~ the decrease of ~~other fluxes (which include mainly anthropogenic fluxes)~~ ~~non-wetland fluxes~~ in the NH, ~~while wetland fluxes continue to rise and the continuing increase of wetland fluxes~~ in the SH. ~~However, more research is needed to substantiate this inferred connection between the change in NH growth rates and fluxes.~~

In summary, our DLM-based approach allows ~~for the calculation of~~ ~~calculating~~ growth rates or AMIs for global and zonal S5P/TROPOMI data. This approach is computationally inexpensive and ~~readily~~ allows for the constant integration of new data, enabling timely assessments of global methane concentration changes. Importantly, no ~~additional~~ prior information about the atmospheric state is required. We believe that our approach provides an additional valuable tool for investigating atmospheric methane concentrations, enabling the rapid identification of regions of interest, ~~like such as~~ the 2022 NH. Furthermore, our approach can be readily applied to other datasets facing similar challenges, such as inhomogeneous sampling, non-linear trends, and data gaps. For the 70°–90° N band our method failed to identify any changes in growth rate, however this result is in good agreement with the growth rates from CAMS/INV-SRF XCH<sub>4</sub> data which themselves only show small variations. This indicates that (a) ~~The the~~ small changes in growth rate could not be distinguished from the random variability in the data ~~and~~ ~~or~~ (b) ~~No no~~ anomalous increases in growth rate are visible for the northern high latitude regions ~~in the observed period from 2018 – 2022.~~

Future research could aim to improve this approach, especially for high latitude regions, to identify smaller changes in growth rates. ~~To quantify~~ ~~Better estimates of~~ the impact of meridional transport on zonal growth rates, ~~the could help to~~ ~~provide better error estimates for our method.~~ The 2022 decrease in NH growth rates could be investigated in more detail and this approach be extended to include data sets of other atmospheric constituents. Data from future satellite ~~mission~~missions, with lower uncertainties and increased data coverage, could ~~further allow for~~ ~~enable~~ the investigation of sub-annual changes in growth rates, which are presently not detectable. Finally, zonal growth rates of long-lived gases (e.g. HF) without any significant sources or sinks could possibly ~~allow for~~ ~~enable~~ the quantification of atmospheric transport patterns.

565 *Code and data availability.* CAMS global inversion-optimized greenhouse gas fluxes and concentrations are available from <https://ads.atmosphere.copernicus.eu/>. Sentinel-5P/TROPOMI WFMD data is available from [https://www.iup.uni-bremen.de/carbon\\_ghg/products/tropomi\\_wfmd/](https://www.iup.uni-bremen.de/carbon_ghg/products/tropomi_wfmd/). NOAA MBL data is available from <https://gml.noaa.gov/ccgg/mb/>. Example code to recreate Fig. 5 (Global annual methane increases), including the gridding and processing of the data, is available under <https://doi.org/10.5281/zenodo.8178927>.

## Appendix A: Model setup and ensemble size

570 The structure of our DLMS ~~is informed by the basic assumption~~ assumes that the measured methane signal can be separated into a slowly changing background level, a seasonal component and noise term. This section closely follows the more detailed description in *Durbin and Koopman* [2012] and *Harvey* [1990].

The level component  ~~$\mu_t$~~  can be described by the following formulas

$$575 \quad \mu_{t+1} = \mu_t + \nu_t + \epsilon_{\text{level}, t, \text{level}}, \quad \epsilon_{\text{level}, t, \text{level}} \sim N(0, \sigma_{\text{level}}^2) \quad (\text{A1})$$

$$\nu_{t+1} = \nu_t + \epsilon_{\text{trend}, t, \text{trend}}, \quad \epsilon_{\text{trend}, t, \text{trend}} \sim N(0, \sigma_{\text{trend}}^2) \quad (\text{A2})$$

where  $\mu_t$  is the level,  $\nu_t$  is the trend (i.e. the slope or ~~change of the level~~ growth rate) and  $\epsilon_t$  are steps in a random walk sampled from a Gaussian distribution. The random walks allow components to change over time. Since we ~~assume~~ want to allow for a smoothly changing level we allow the trend to change over time, ~~which is modeled by a random walk with variance  $\sigma_{\text{trend}}^2$ .~~

580 Additionally. Additionally, we enforce a constraint of zero variance for the level to ensure that short-term fluctuations in the background level are not allowed:

$$\sigma_{\text{level}}^2 = 0 \quad (\text{A3})$$

$$\sigma_{\text{trend}}^2 > 0 \quad (\text{A4})$$

The seasonal part of the signal is modeled by a truncated Fourier-series with  $h$  harmonics:

$$585 \quad \gamma_t = \sum_{j=1}^h \gamma_{jt} \quad (\text{A5})$$

with

$$\gamma_{j,t+1} = \gamma_{jt} \cos(\lambda_j) + \gamma_{jt}^* \sin(\lambda_j) + \epsilon_{\text{seas}}, \quad \epsilon_{\text{seas}} \sim N(0, \sigma_{\text{seas}}^2), \quad \lambda_j = \frac{2\pi j}{s} \quad (\text{A6})$$

$$\gamma_{j,t+1}^* = -\gamma_{jt} \sin(\lambda_j) + \gamma_{jt}^* \cos(\lambda_j) + \epsilon_{\text{seas}}^*, \quad \epsilon_{\text{seas}}^* \sim N(0, \sigma_{\text{seas}}^2), \quad j = 1, \dots, h \quad (\text{A7})$$

590 where  $s$  describes the seasonality of the data, e.g. for monthly data  $s = 12$  or for daily data  $s = 365$  when modeling yearly patterns. The value of  $s = 365.25$  can be used to account for leap years. We use  $s = 365.2$  which is equal to the average number of days per year between 2018 and 2022. For  $\sigma_{\text{seas}}^2 > 0$  the seasonal cycle is allowed to change over time. We allow values of  $h \in \{1, 2, 3, 4\}$ , to account for varying levels of complexity in the seasonal cycle. The motivation for this is twofold. Firstly,

we want to only model the basic structure of the seasonal cycle and not the whole signal. Secondly, the inclusion of more harmonics introduces further parameters which have to be estimated. This quickly leads to high uncertainties in the produced fit since not enough information is included in the data to account for the growing number of parameters.

The noise term accounts for residual correlations as well as random Gaussian noise in the signal. ~~This is achieved by including~~ Residual correlations can be modeled by an autoregressive component which includes a serial dependence between the observations. An autoregressive noise of order  $n$  ~~therefore~~ includes a memory of the last  $n$  measurements. For  $n = 1$  this  $AR(1)$  term is

$$\eta_{t+1} = \rho\eta_t + \epsilon_{AR(1)}, \quad \epsilon_{AR(1)} \sim N(0, \sigma_{AR(1)}^2), \quad \rho \in [0, 1] \quad (\text{A8})$$

~~which introduce~~ where  $\eta_t$  is the autoregressive component,  $\rho$  determines the strength of the autocorrelation (the memory of the previous time step) and  $\epsilon_{AR(1)}$  is again a step in a Gaussian random walk. This component introduces the parameters  $\rho$  and  $\sigma_{AR(1)}^2$  to the model. We confine our autoregressive component to the order of  $n = 1$ , which is enough to model the residual correlations in ~~the our~~ data. Higher orders would introduce further parameters to be estimated and lead to a harder interpretability of the results. However, exclusion of the  $AR(1)$  component leads to bad fits since the model fails to account for the data variability.

An additional Gaussian noise can be included

$$\epsilon_{irr}, \quad \epsilon_{irr} \sim N(0, \sigma_{irr}^2) \quad (\text{A9})$$

which we call *irregular* component.

The complete signal can be then written as the sum of these components

$$y_t = \mu_t + \gamma_t + \eta_t + \epsilon_{irr} \quad (\text{A10})$$

~~The remaining tunable parameters are the~~

The ensemble size is determined by the number of possible model configurations. In our case this is determined by whether to allow variability of the seasonal ~~term, the inclusion of cycle, whether to include~~ a Gaussian error term and the number of harmonics ~~which determine the size of the ensemble ( $N = 2 \cdot 2 \cdot 4 = 16$ ).~~ An overview of all parameters can be found in Tab. A1.


## Appendix B: Replication of complete NOAA GML & UB-C3S AMIs

To investigate the effect of the fitting method on AMIs we replicated AMIs calculated by the NOAA-GML and C3S in Fig. B1 and B2 respectively. Here we present the comparison for the whole available time range (subset of data can be seen in Tab. 3).

**Table A1.** DLM parameters.

Parameter	Description	Allowed range
$\sigma_{level}^2$	Variability of level	0
$\sigma_{trend}^2$	Variability of trend	$[0, \infty]^*$
$\sigma_{seas}^2$	Variability of seasonal cycle	0 or <sup>†</sup> $[0, \infty]^*$
$\rho$	Parameter of AR(1)	$[0, 1]^*$
$\sigma_{AR(1)}^2$	Variability of AR(1)	$[0, \infty]^*$
$\sigma_{irr}^2$	Variability of Gaussian error	0 or <sup>†</sup> $[0, \infty]^*$
$h$	Number of harmonics	$1 - 4^\dagger$

\* determined by maximum likelihood estimation during DLM fit, <sup>†</sup> Different settings are part of ensemble




**Figure B1.** Comparison of global annual methane increases derived from the NOAA–GML MBLR data using different methods.


### Appendix C: Global AMIs and zonal growth rates derived from CAMS global inversion-optimised greenhouse concentrations

625

Here we present global AMIs and zonal growth rates for CAMS/INV-SRF-SAT data which includes satellite measurements from GOSAT in its optimization (see Fig. C1 and C2).




**Figure B2.** Comparison of global annual methane increases derived from C3S XCH<sub>4</sub>\_OBS4MIPS v4.4 data which is extended by CAMS NRT data after 2021 using different methods.



**Figure C1.** Global annual methane increases derived from CAMS global inversion-optimised greenhouse concentrations including both surface and satellite observations.

**Appendix D:** [WFMDv1.8 data coverage](#)





**Figure C2.** Zonal growth rates for 20° bands derived from (a) Sentinel-5P/TROPOMI WFMDv1.8 data and (b) CAMS/INV-SRF-SAT data. The errors show the  $\pm 1\sigma$  uncertainty.

630


[Figure D1](#) shows the area-normalized coverage of S5P/TROPOMI WFMDv1.8 data. It can be seen that tropics, mid-latitudes and high latitudes have mostly an average coverage of about 25%, while the high latitude regions have seasonal gaps due to the polar night.

635

*Author contributions.* JH developed the methodology and conducted the formal analysis. OS provided information on the WFMDv1.8 data. MiB provided information on the UB-C3S-CAMS data. JPB provided valuable input on atmospheric transport effects. MaB provided supervision and helped with the conceptualization. JH wrote the initial draft of this manuscript. All authors contributed significantly to the conception of the analysis, jointly discussed the results and provided constructive comments to improve the manuscript.

*Competing interests.* The authors declare that they have no conflict of interest.

*Disclaimer.* Neither the European Commission nor ECMWF is responsible for any use that may be made of the information it contains.



**Figure D1.** Area-normalized coverage of S5P/TROPOMI WFMDv1.8 data for 30° zonal bands, representing the high latitude, mid-latitude and tropic regions. For the calculation data on a 2° × 2° grid was used. The coverage is given on a scale from 0 to 1, where 1 represents complete coverage within a band.

*Acknowledgements.* ~~Contains~~ This publication uses and contains modified Copernicus Atmosphere Monitoring Service information (2018-2022) ~~-This publication contains and~~ modified Copernicus Sentinel data (2018-2023) 2018-2023. Sentinel-5 Precursor is an ESA mission implemented on behalf of the European Commission. The TROPOMI payload is a joint development by the ESA and the Netherlands Space Office (NSO). The Sentinel-5 Precursor ground-segment development has been funded by the ESA and with national contributions from the Netherlands, Germany, and Belgium. The pre-operational TROPOMI data processing was carried out on the Dutch national e-infrastructure with the support of SURF Cooperative. Scientific color maps [Crameri, 2021] are used in this study to prevent visual distortion of data and exclusion of readers with color-vision deficiencies [Crameri et al., 2020].

*Financial support.* This project is funded by the State and University of Bremen. In particular the University of Bremen funding for the junior research group ‘Greenhouse gases in the Arctic’ is acknowledged by JH and MB. We gratefully acknowledge the funding by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Projektnummer 268020496 – TRR 172, within the Transregional Collaborative Research Center “Arctic Amplification: Climate Relevant Atmospheric and SurfaCe Processes, and Feedback Mechanisms (AC)<sup>3</sup>”. We also acknowledge the received funding from the European Space Agency (ESA) via the projects GHG-CCI+ and MethaneCAMP (ESA contract nos. 4000126450/19/I-NB and 4000137895/22/I-AG) and from the german ministry of education and research (BMBF) within its project ITMS via grant 01 LK2103A. The TROPOMI/WFMD version 1.8 data have been generated using funding from ESA (GHG-CCI and MethaneCAMP projects). The TROPOMI/WFMD retrievals presented here were performed on HPC facilities of the IUP, University of Bremen, funded under DFG/FUGG grant nos. INST 144/379-1 and INST 144/493-1. Part of the AMIs presented were generated

for Copernicus ESOTC 2022 (<https://climate.copernicus.eu/climate-indicators/greenhouse-gas-concentrations>) funded by EU Copernicus  
655 Climate Change Service via project C3S2\_312a\_Lot2.

## References

- Arias, P., Bellouin, N., Coppola, E., Jones, R., Krinner, G., Marotzke, J., Naik, V., Palmer, M., Plattner, G.-K., Rogelj, J., Rojas, M., Sillmann, J., Storelvmo, T., Thorne, P., Trewin, B., Achuta Rao, K., Adhikary, B., Allan, R., Armour, K., Bala, G., Barimalala, R., Berger, S., Canadell, J., Cassou, C., Cherchi, A., Collins, W., Collins, W., Connors, S., Corti, S., Cruz, F., Dentener, F., Dereczynski, C., Di Luca, A., Diongue Niang, A., Doblas-Reyes, F., Dosio, A., Douville, H., Engelbrecht, F., Eyring, V., Fischer, E., Forster, P., Fox-Kemper, B., Fuglested, J., Fyfe, J., Gillett, N., Goldfarb, L., Gorodetskaya, I., Gutierrez, J., Hamdi, R., Hawkins, E., Hewitt, H., Hope, P., Islam, A., Jones, C., Kaufman, D., Kopp, R., Kosaka, Y., Kossin, J., Krakovska, S., Lee, J.-Y., Li, J., Mauritsen, T., Maycock, T., Meinshausen, M., Min, S.-K., Monteiro, P., Ngo-Duc, T., Otto, F., Pinto, I., Pirani, A., Raghavan, K., Ranasinghe, R., Ruane, A., Ruiz, L., Sallée, J.-B., Samset, B., Sathyendranath, S., Seneviratne, S., Sörensson, A., Szopa, S., Takayabu, I., Tréguier, A.-M., van den Hurk, B., Vautard, R., von Schuckmann, K., Zaehle, S., Zhang, X., and Zickfeld, K.: Technical Summary, in: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J., Maycock, T., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., p. 33-144, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/9781009157896.002>, type: Book Section, 2021.
- Basu, S., Lan, X., Dlugokencky, E., Michel, S., Schwietzke, S., Miller, J. B., Bruhwiler, L., Oh, Y., Tans, P. P., Apadula, F., Gatti, L. V., Jordan, A., Necki, J., Sasakawa, M., Morimoto, S., Di Iorio, T., Lee, H., Arduini, J., and Manca, G.: Estimating Emissions of Methane Consistent with Atmospheric Measurements of Methane and  $\delta^{13}\text{C}$  of Methane, *Atmospheric Chemistry and Physics Discussions*, 2022, 1–38, <https://doi.org/10.5194/acp-2022-317>, 2022.
- Bergamaschi, P., Krol, M., Meirink, J. F., Dentener, F., Segers, A., van Aardenne, J., Monni, S., Vermeulen, A. T., Schmidt, M., Ramonet, M., Yver, C., Meinhardt, F., Nisbet, E. G., Fisher, R. E., O'Doherty, S., and Dlugokencky, E. J.: Inverse modeling of European CH<sub>4</sub> emissions 2001–2006, *Journal of Geophysical Research: Atmospheres*, 115, <https://doi.org/10.1029/2010JD014180>, <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2010JD014180>, 2010.
- Bergamaschi, P., Houweling, S., Segers, A., Krol, M., Frankenberg, C., Scheepmaker, R. A., Dlugokencky, E., Wofsy, S. C., Kort, E. A., Sweeney, C., Schuck, T., Brenninkmeijer, C., Chen, H., Beck, V., and Gerbig, C.: Atmospheric CH<sub>4</sub> in the first decade of the 21st century: Inverse modeling analysis using SCIAMACHY satellite retrievals and NOAA surface measurements, *Journal of Geophysical Research: Atmospheres*, 118, 7350–7369, <https://doi.org/10.1002/jgrd.50480>, [\\_eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1002/jgrd.50480](https://onlinelibrary.wiley.com/doi/pdf/10.1002/jgrd.50480), 2013.
- Buchwitz, M., Schneising, O., Reuter, M., Heymann, J., Krautwurst, S., Bovensmann, H., Burrows, J. P., Boesch, H., Parker, R. J., Somkuti, P., Detmers, R. G., Hasekamp, O. P., Aben, I., Butz, A., Frankenberg, C., and Turner, A. J.: Satellite-derived methane hotspot emission estimates using a fast data-driven method, *Atmospheric Chemistry and Physics*, 17, 5751–5774, <https://doi.org/10.5194/acp-17-5751-2017>, publisher: Copernicus GmbH, 2017.
- C3S: Methane data from 2002 to present derived from satellite observations, <https://doi.org/10.24381/CDS.B25419F8>, 2018.
- C3S: Climate Indicators - Greenhouse gas concentrations, <https://climate.copernicus.eu/climate-indicators/greenhouse-gas-concentrations>, 2023a.
- C3S: Climate Data Store, <https://cds.climate.copernicus.eu#!/home>, 2023b.
- C3S: Copernicus Climate Change Service, <https://climate.copernicus.eu/>, 2023c.
- C3S: European State of the Climate 2022 | Copernicus, <https://climate.copernicus.eu/esotc/2022>, 2023d.

- CAMS: Copernicus Atmosphere Monitoring Service, <https://atmosphere.copernicus.eu/>, 2023.
- Cooper, M. J., Martin, R. V., Hammer, M. S., Levelt, P. F., Veefkind, P., Lamsal, L. N., Krotkov, N. A., Brook, J. R., and McLinden, C. A.:  
695 Global fine-scale changes in ambient NO<sub>2</sub> during COVID-19 lockdowns, *Nature*, 601, 380–387, <https://doi.org/10.1038/s41586-021-04229-0>, number: 7893 Publisher: Nature Publishing Group, 2022.
- Crameri, F.: Scientific colour maps, <https://doi.org/10.5281/zenodo.5501399>, 2021.
- Crameri, F., Shephard, G. E., and Heron, P. J.: The misuse of colour in science communication, *Nature Communications*, 11, 5444,  
<https://doi.org/10.1038/s41467-020-19160-7>, 2020.
- 700 Dlugokencky, E.: Trends in Atmospheric Methan, [https://www.gml.noaa.gov/ccgg/trends\\_ch4/](https://www.gml.noaa.gov/ccgg/trends_ch4/), last access: 20.09.2022, 2022.
- Dlugokencky, E. J., Lan, X., Crotwell, A., Thoning, K., and Crotwell, M.: Atmospheric Methane Dry Air Mole Fractions from the NOAA  
ESRL Carbon Cycle Cooperative Global Air Sampling Network, [ftp://aftp.cmdl.noaa.gov/data/trace\\_gases/ch4/flask/surface/](ftp://aftp.cmdl.noaa.gov/data/trace_gases/ch4/flask/surface/), 2021.
- Durbin, J. and Koopman, S. J.: Time Series analysis by state space methods, Second Edition, Oxford University press, 2012.
- Feng, L., Palmer, P. I., Zhu, S., Parker, R. J., and Liu, Y.: Tropical methane emissions explain large fraction of recent changes in global  
705 atmospheric methane growth rate, *Nature Communications*, 13, 1378, <https://doi.org/10.1038/s41467-022-28989-z>, 2022.
- Feng, L., Palmer, P. I., Parker, R. J., Lunt, M. F., and Bösch, H.: Methane emissions are predominantly responsible for record-breaking  
atmospheric methane growth rates in 2020 and 2021, *Atmospheric Chemistry and Physics*, 23, 4863–4880, <https://doi.org/10.5194/acp-23-4863-2023>, 2023.
- Gulev, S., Thorne, P., Ahn, J., Dentener, F., Domingues, C., Gerland, S., Gong, D., Kaufman, D., Nnamchi, H., Quaas, J., Rivera, J., Sathyendranath, S., Smith, S., Trewin, B., von Schuckmann, K., and Vose, R.: Changing State of the Climate System, in: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J., Maycock, T., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., pp. 287–422, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/9781009157896.004>, type: Book  
710 Section, 2021.
- Hachmeister, J., Schneising, O., Buchwitz, M., Lorente, A., Borsdorff, T., Burrows, J. P., Notholt, J., and Buschmann, M.: On the influence  
of underlying elevation data on Sentinel-5 Precursor TROPOMI satellite methane retrievals over Greenland, *Atmospheric Measurement  
Techniques*, 15, 4063–4074, <https://doi.org/10.5194/amt-15-4063-2022>, 2022.
- Harvey, A. C.: *Forecasting, Structural Time Series Models and the Kalman Filter*, Cambridge University Press,  
720 <https://doi.org/10.1017/CBO9781107049994>, 1990.
- Hugelius, G., Strauss, J., Zubrzycki, S., Harden, J. W., Schuur, E. A. G., Ping, C.-L., Schirrmeyer, L., Grosse, G., Michaelson, G. J., Koven, C. D., O'Donnell, J. A., Elberling, B., Mishra, U., Camill, P., Yu, Z., Palmtag, J., and Kuhry, P.: Estimated stocks of circumpolar permafrost carbon with quantified uncertainty ranges and identified data gaps, *Biogeosciences*, 11, 6573–6593, <https://doi.org/10.5194/bg-11-6573-2014>, 2014.
- 725 IEA: Methane Tracker 2021, <https://www.iea.org/reports/methane-tracker-2021>, license: CC BY 4.0, 2021.
- IEA: Global Methane Tracker 2022 – Analysis, <https://www.iea.org/reports/global-methane-tracker-2022>, 2023.
- Jacob, D. J.: *Introduction to Atmospheric Chemistry*, Princeton University Press, <https://doi.org/10.1515/9781400841547>, publication Title: *Introduction to Atmospheric Chemistry*, 1999.
- Kalman, R. E.: A New Approach to Linear Filtering and Prediction Problems, *Journal of Basic Engineering*, 82, 35–45,  
730 <https://doi.org/10.1115/1.3662552>, 1960.

- Karppinen, T., Lamminpää, O., Tukiainen, S., Kivi, R., Heikkinen, P., Hatakka, J., Laine, M., Chen, H., Lindqvist, H., and Tamminen, J.: Vertical Distribution of Arctic Methane in 2009–2018 Using Ground-Based Remote Sensing, *Remote Sensing*, 12, 917, <https://doi.org/10.3390/rs12060917>, number: 6 Publisher: Multidisciplinary Digital Publishing Institute, 2020.
- 735 Kivimäki, E., Lindqvist, H., Hakkarainen, J., Laine, M., Sussmann, R., Tsuruta, A., Detmers, R., Deutscher, N. M., Dlugokencky, E. J., Hase, F., Hasekamp, O., Kivi, R., Morino, I., Notholt, J., Pollard, D. F., Roehl, C., Schneider, M., Sha, M. K., Velazco, V. A., Warneke, T., Wunch, D., Yoshida, Y., and Tamminen, J.: Evaluation and Analysis of the Seasonal Cycle and Variability of the Trend from GOSAT Methane Retrievals, *Remote Sensing*, 11, 882, <https://doi.org/10.3390/rs11070882>, number: 7 Publisher: Multidisciplinary Digital Publishing Institute, 2019.
- Laine, M.: Introduction to Dynamic Linear Models for Time Series Analysis, in: *Geodetic Time Series Analysis in Earth Sciences*, pp. 139–156, Springer International Publishing, Cham, [https://doi.org/10.1007/978-3-030-21718-1\\_4](https://doi.org/10.1007/978-3-030-21718-1_4), 2020.
- 740 Laine, M., Latva-Pukkila, N., and Kyrölä, E.: Analysing time-varying trends in stratospheric ozone time series using the state space approach, *Atmospheric Chemistry and Physics*, 14, 9707–9725, <https://doi.org/10.5194/acp-14-9707-2014>, 2014.
- Lan, X., Basu, S., Schwietzke, S., Bruhwiler, L. M. P., Dlugokencky, E. J., Michel, S. E., Sherwood, O. A., Tans, P. P., Thoning, K., Etiope, G., Zhuang, Q., Liu, L., Oh, Y., Miller, J. B., Pétron, G., Vaughn, B. H., and Crippa, M.: Improved Constraints on Global Methane Emissions and Sinks Using  $\delta^{13}\text{C}\text{-CH}_4$ , *Global Biogeochemical Cycles*, 35, e2021GB007000, <https://doi.org/https://doi.org/10.1029/2021GB007000>, [\\_eprint: https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2021GB007000](https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2021GB007000), 2021.
- 745 Lan, X., Thoning, K., and Dlugokencky, E. J.: Trends in globally-averaged  $\text{CH}_4$ ,  $\text{N}_2\text{O}$ , and  $\text{SF}_6$  determined from NOAA Global Monitoring Laboratory measurements. Version 2023-07, <https://doi.org/10.15138/P8XG-AA10>, 2023.
- 750 Laughner, J. L., Neu, J. L., Schimel, D., Wennberg, P. O., Barsanti, K., Bowman, K. W., Chatterjee, A., Croes, B. E., Fitzmaurice, H. L., Henze, D. K., Kim, J., Kort, E. A., Liu, Z., Miyazaki, K., Turner, A. J., Anenberg, S., Avise, J., Cao, H., Crisp, D., de Gouw, J., Eldering, A., Fyfe, J. C., Goldberg, D. L., Gurney, K. R., Hasheminassab, S., Hopkins, F., Ivey, C. E., Jones, D. B. A., Liu, J., Lovenduski, N. S., Martin, R. V., McKinley, G. A., Ott, L., Poulter, B., Ru, M., Sander, S. P., Swart, N., Yung, Y. L., and Zeng, Z.-C.: Societal shifts due to COVID-19 reveal large-scale complexities and feedbacks between atmospheric chemistry and climate change, *Proceedings of the National Academy of Sciences*, 118, e2109481 118, <https://doi.org/10.1073/pnas.2109481118>, publisher: Proceedings of the National Academy of Sciences, 2021.
- 755 Ludewig, A.: S5P Mission Performance Centre Level 1b Readme, <https://sentinels.copernicus.eu/documents/247904/3541451/Sentinel-5P-Level-1b-Product-Readme-File.pdf/a89d82ce-7414-43e6-ac77-0c371ed1b096>, 2021.
- McNorton, J., Bousserez, N., Agustí-Panareda, A., Balsamo, G., Cantarello, L., Engelen, R., Huijnen, V., Inness, A., Kipling, Z., Parrington, M., and Ribas, R.: Quantification of methane emissions from hotspots and during COVID-19 using a global atmospheric inversion, *Atmospheric Chemistry and Physics*, 22, 5961–5981, <https://doi.org/10.5194/acp-22-5961-2022>, 2022.
- 760 NOAA: NOAA/GML calculation of global means, <https://gml.noaa.gov/ccgg/mbl/mbl.html>, 2022.
- Peng, S., Lin, X., Thompson, R. L., Xi, Y., Liu, G., Hauglustaine, D., Lan, X., Poulter, B., Ramonet, M., Saunio, M., Yin, Y., Zhang, Z., Zheng, B., and Ciais, P.: Wetland emission and atmospheric sink changes explain methane growth in 2020, *Nature*, 612, 477–482, <https://doi.org/10.1038/s41586-022-05447-w>, number: 7940 Publisher: Nature Publishing Group, 2022.
- 765 Perktold, J., Skipper Seabold, Sheppard, K., ChadFulton, Kerby Shedden, Jbrockmendel, J-Grana6, Quackenbush, P., Arel-Bundock, V., McKinney, W., Langmore, I., Baker, B., Gommers, R., Yogabonito, S-Scherrer, Zhurko, E., Brett, M., Giampieri, E., Yichuan Liu, Mill-

- man, J., Hobson, P., Vincent, Roy, P., Augspurger, T., Tvanzyl, Alexbrc, Hartley, T., Perez, F., Tamiya, Y., and Halchenko, Y.: statsmodel-  
s/statsmodels: Release 0.14.0, <https://doi.org/10.5281/ZENODO.593847>, 2023.
- 770 Plaza, C., Pegoraro, E., Bracho, R., Celis, G., Crummer, K. G., Hutchings, J. A., Hicks Pries, C. E., Mauritz, M., Natali, S. M., Salmon,  
V. G., Schädel, C., Webb, E. E., and Schuur, E. A. G.: Direct observation of permafrost degradation and rapid soil carbon loss in tundra,  
Nature Geoscience, 12, 627–631, <https://doi.org/10.1038/s41561-019-0387-6>, 2019.
- Qu, Z., Jacob, D. J., Zhang, Y., Shen, L., Varon, D. J., Lu, X., Scarpelli, T., Bloom, A., Worden, J., and Parker, R. J.: Attribution of  
the 2020 surge in atmospheric methane by inverse analysis of GOSAT observations, Environmental Research Letters, 17, 094003,  
775 <https://doi.org/10.1088/1748-9326/ac8754>, publisher: IOP Publishing, 2022.
- Reuter, M., Buchwitz, M., Schneising, O., Noël, S., Bovensmann, H., Burrows, J. P., Boesch, H., Di Noia, A., Anand, J., Parker, R. J.,  
Somkuti, P., Wu, L., Hasekamp, O. P., Aben, I., Kuze, A., Suto, H., Shiomi, K., Yoshida, Y., Morino, I., Crisp, D., O'Dell, C. W., Notholt, J.,  
Petri, C., Warneke, T., Velazco, V. A., Deutscher, N. M., Griffith, D. W. T., Kivi, R., Pollard, D. F., Hase, F., Sussmann, R., Té, Y. V., Strong,  
K., Roche, S., Sha, M. K., De Mazière, M., Feist, D. G., Iraci, L. T., Roehl, C. M., Retscher, C., and Schepers, D.: Ensemble-based satellite-  
780 derived carbon dioxide and methane column-averaged dry-air mole fraction data sets (2003–2018) for carbon and climate applications,  
Atmospheric Measurement Techniques, 13, 789–819, <https://doi.org/10.5194/amt-13-789-2020>, publisher: Copernicus GmbH, 2020.
- Schneising, O., Buchwitz, M., Hachmeister, J., Vanselow, S., Reuter, M., Buschmann, M., Bovensmann, H., and Burrows, J. P.: Advances  
in retrieving XCH<sub>4</sub> and XCO from Sentinel-5 Precursor: improvements in the scientific TROPOMI/WFMD algorithm, Atmospheric  
Measurement Techniques, 16, 669–694, <https://doi.org/10.5194/amt-16-669-2023>, 2023.
- 785 Segers, A., Steinke, T., and Houweling, S.: Description of the CH<sub>4</sub> Inversion Production Chain, [https://atmosphere.copernicus.eu/sites/  
default/files/2022-10/CAMS255\\_2021SC1\\_D55.5.2.1-2021CH4\\_202206\\_production\\_chain\\_CH4\\_v1.pdf](https://atmosphere.copernicus.eu/sites/default/files/2022-10/CAMS255_2021SC1_D55.5.2.1-2021CH4_202206_production_chain_CH4_v1.pdf), 2022.
- Serreze, M. C. and Barry, R. G.: Processes and impacts of Arctic amplification: A research synthesis, Global and Planetary Change, 77,  
85–96, <https://doi.org/https://doi.org/10.1016/j.gloplacha.2011.03.004>, 2011.
- Sofieva, V. F., Kalakoski, N., Päivärinta, S.-M., Tamminen, J., Laine, M., and Froidevaux, L.: On sampling uncertainty of satellite ozone  
790 profile measurements, Atmospheric Measurement Techniques, 7, 1891–1900, <https://doi.org/10.5194/amt-7-1891-2014>, 2014.
- Stevenson, D. S., Derwent, R. G., Wild, O., and Collins, W. J.: COVID-19 lockdown emission reductions have the potential to ex-  
plain over half of the coincident increase in global atmospheric methane, Atmospheric Chemistry and Physics, 22, 14243–14252,  
<https://doi.org/10.5194/acp-22-14243-2022>, 2022.
- Szopa, S., Naik, V., Adhikary, B., Artaxo, P., Bernsten, T., Collins, W., Fuzzi, S., Gallardo, L., Kiendler-Scharr, A., Klimont, Z., Liao, H.,  
795 Unger, N., and Zanis, P.: Short-Lived Climate Forcers, in: Climate Change 2021: The Physical Science Basis. Contribution of Working  
Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, edited by Masson-Delmotte, V., Zhai, P.,  
Pirani, A., Connors, S., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M., Huang, M., Leitzell, K., Lonnoy, E., Matthews,  
J., Maycock, T., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., pp. 817–922, Cambridge University Press, Cambridge, United Kingdom  
and New York, NY, USA, <https://doi.org/10.1017/9781009157896.008>, type: Book Section, 2021.
- 800 Turner, A. J., Frankenberg, C., and Kort, E. A.: Interpreting contemporary trends in atmospheric methane, Proceedings of the National  
Academy of Sciences, 116, 2805–2813, <https://doi.org/10.1073/pnas.1814297116>, 2019.
- Warneck, P.: Chemistry of the Natural Atmosphere, vol. 71 of *INTERNATIONAL GEOPHYSICS SERIES*, Academic Press, 2nd edition edn.,  
1999.
- Wendisch, M., Brückner, M., Burrows, J. P., Crewell, S., Dethloff, K., Ebell, K., Lüpkes, C., Macke, A., Notholt, J., Quaas, J., Rinke, A., and  
805 Tegen, I.: Understanding Causes and Effects of Rapid Warming in the Arctic, Eos, 98, <https://doi.org/10.1029/2017EO064803>, 2017.



Zhang, Z., Poulter, B., Feldman, A. F., Ying, Q., Ciais, P., Peng, S., and Li, X.: Recent intensification of wetland methane feedback, *Nature Climate Change*, 13, 430–433, <https://doi.org/10.1038/s41558-023-01629-0>, number: 5 Publisher: Nature Publishing Group, 2023.

-