

Measurement report: A one-year study to estimate maritime contributions to PM₁₀ in a coastal area in Northern France

Frédéric Ledoux^{1,*}, Cloé Roche^{1,*}, Gilles Delmaire², Gilles Roussel², Olivier Favez³, Marc Fadel^{1,*},
Dominique Courcot¹

5 ¹Unité de Chimie Environnementale et Interactions sur le Vivant, UCEIV UR4492, FR CNRS 3417, Université du Littoral
Côte d'Opale (ULCO), Dunkerque, France.

²Laboratoire d'Informatique Signal et Image de la Côte d'Opale, LISIC UR4491, Université du Littoral Côte d'Opale
(ULCO), Calais, France.

10 ³Institut National de l'Environnement Industriel et des Risques, INERIS, Parc Technologique ALATA, Verneuil-en-Halatte,
France.

Correspondence to: Marc FADEL (marc.fadel@univ-littoral.fr)

***These authors contributed equally to this work.**

Abstract. This work is focused on filling the lack of knowledge associated with natural and anthropogenic marine emissions on PM₁₀ concentrations in Northern France. For this purpose, a one-year measurement and sampling campaign for PM₁₀ has
15 been performed at a French coastal site situated in front of the Straits of Dover. The characterization of PM₁₀ samples was performed considering major and trace elements, water-soluble ions, organic carbon (OC), elemental carbon (EC), and organic markers of biomass burning and primary biogenic emissions. Furthermore, the source apportionment of PM₁₀ was achieved using the constrained weighted-non-negative matrix factorization (CW-NMF) model. The annual average PM₁₀ was 24.3 µg/m³ with six species contributing to 69% of its mass (NO₃⁻, OC, SO₄²⁻, Cl⁻, Na⁺, and NH₄⁺). The source
20 apportionment of PM₁₀ led to the identification of ~~940~~ sources. Fresh and aged sea-salts contributed to 37% of PM₁₀, while secondary nitrate and sulfate contributed ~~424~~%, biomass burning ~~840~~%, and Heavy Fuel Oil (HFO) combustion from shipping emissions contributed almost 5%, on yearly averages. Additionally, monthly evolution of the sources' contribution evidenced different behaviors with high contributions of secondary nitrate and biomass burning during winter. In the
25 summer season, ~~159~~ times higher concentrations for HFO combustion (July compared to January) and the predominance of aged sea-salts versus fresh sea-salts were observed. Constant weighted trajectories showed that the sources contributing to more than 80% of PM₁₀ at Cape Gris-Nez are of regional and/or long-range origins with the North Sea and the English Channel as hotspots for natural and anthropogenic marine emissions and Belgium, the Netherlands, and the West of Germany as hotspots for secondary inorganic aerosols.

1 Introduction

30 Maritime transport is considered nowadays as a crucial transportation system used to ship goods and people over long distances. Due to globalization of production processes and the growth of global trade, this mean of transportation has been

increasing with over 80% of the volume of international trade of goods carried by sea in the last decades (Marmer et al., 2009; UNCTAD, 2021). In the meantime, shipping is known as a significant atmospheric source of pollutants especially near harbors and surrounding coastal areas (Contini and Merico, 2021; Jonson et al., 2020; Ausmeel et al., 2020). It is an important source of carbon monoxide (CO), nitrogen oxides (NO_x), sulfur oxides (SO_x), volatile organic compounds (VOCs), and particulate matter (Lv et al., 2018; Seppälä et al., 2021). In particular, several studies have highlighted the impact of shipping emissions on human health (Andersson et al., 2009; Corbett et al., 2007) since most of their emissions are estimated to occur within 400 km of land (Endresen et al., 2003; Jutterström et al., 2021). According to Zhang et al. (2021), 94 200 premature deaths worldwide were associated with particulate matter (PM) exposure due to maritime shipping in 2015.

Over the last decades, European countries have made large efforts to reduce emissions from several anthropogenic sources such as road traffic, industrial, and power generation, ~~etc.~~ which led to the indirect increase of the contribution of shipping to the overall anthropogenic emissions (Viana et al., 2014). In order to reduce shipping emissions, different coastal areas (such as ~~(The North Sea, the English Channel, and the Baltic Sea), etc.)~~ have been classified as Sulfur Emission Control Areas (SECA) (EEA, 2013). In the latter areas, the sulfur content in marine fuels was limited to 0.1% starting 2015, after it was 1% between 2010 and 2015 and 1.5% before 2010 (Tang et al., 2020). The European Union has set a limit of 0.1% sulfur content fuel on ships at berth at EU ports since 2010 (EEA, 2021).

Several papers in the literature have estimated the contributions from shipping emissions (combustion of HFO) to air quality across Europe and were mainly focused on coastal areas of Italy, Spain, and Ireland (Mazzei et al., 2008; Viana et al., 2009; Pandolfi et al., 2011; Becagli et al., 2012; Contini et al., 2011; Hellebust et al., 2010; Donateo et al., 2014; Viana et al., 2014; Cesari et al., 2014; Bove et al., 2016; Amato et al., 2009). The annual mean contributions to PM in these studies varied between 1 and 12%. However, there were no studies focusing on the maritime contribution to PM in coastal areas of France and more specifically in the southern North Sea region that encompasses large harbours in different areas such as in Dunkerque, Calais, Rotterdam, and Antwerp, ~~etc.~~ (EEA, 2013).

In Northern France, studies focusing on the characterization of PM₁₀ and the identification of its emission sources were conducted in urban and industrial areas (Crenn et al., 2015; Kfoury et al., 2016; Waked et al., 2014; Rimetz-Planchon et al., 2008; Ledoux et al., 2017). However, coastline sites can also be impacted by high particulate atmospheric background levels even without any direct influence from urban and/or industrial sources. On one hand, this might be due to the long-range transport influence as well as the gas-to-particle conversion (Waked et al., 2014; Matthias et al., 2010). On the other hand, there is still a lack of information regarding the impact of natural maritime emissions such as sea-salts (Manders et al., 2010) and anthropogenic ones such as marine traffic. It is worth noting that the southern North Sea is considered as one of the higher ship traffic density in the world (EEA, 2013). This is due to the fact that the Northern France region is bordered by the North-Sea and the English Channel that form together with the Straits of Dover a narrow corridor leading to one of the highest shipping concentration in the world (EEA, 2013).

According to Ledoux et al. (2018), the impact of shipping emissions on PM₁₀ concentration in Calais urban area may reach up to about 30 µg/m³ on average, when the winds are blowing from the whole harbor area and compared to background concentrations. ~~Punctually, ship emissions led to a marked increase of PM₁₀ mass of +78.9 µg/m³.~~ Punctually, ship emissions led to a marked increase of PM₁₀ mass of 78.9 µg/m³ for a one-minute time interval.

The main objective of this work is to understand the impact of emissions resulting from the maritime compartment in coastal areas in Northern France. Therefore, a one-year PM₁₀ sampling and measurement campaign was performed at the Cape Gris-Nez, a French coastal site situated in front of the Straits of Dover in 2013. The collected samples were chemically characterized for their carbonaceous, ionic, and elemental fractions, as well as some organic tracers. Additionally, PM₁₀ sources were apportioned and studied, specifically natural emissions such as sea sprays and anthropogenic emissions linked to maritime traffic. This paper also highlights the seasonal variations of these PM₁₀ sources and the study of the regional influence. Even though the sampling campaign was conducted in 2013 and the regulations in SECA areas have evolved since, this study is still relevant today due to the scarcity of literature studies in front of the Straits of Dover considered as one of the world's busiest shipping lanes. Additionally, despite the IMO regulation for global sulfur limit of 0.5% from ship's fuel oil applied starting January 2020, different countries are still adopting higher sulfur limits. It is worth noting that these limits were only set for sulfur content in marine fuels in order to reduce not only SO₂ emissions but also primary PM emissions (Zetterdahl et al., 2016; Shen and Li, 2020). Hence, this study aims at highlighting the contribution of the natural and anthropogenic marine emissions to the PM₁₀ concentrations in a rural coastal ~~background~~ French site.

2 Materials and methods

2.1 Sampling site

The PM₁₀ filter sampling was conducted at the Cape Gris-Nez located in Northern France. It is a rural coastal site (50°52'08'' N, 1°35'49'' E, altitude 50 m), located on the edge of a cliff, 200 m from the sea and therefore strongly subject to marine influence from sectors 210° to 50° via the north (Figure 1). The Strait of Dover, connecting the English Channel and the North Sea, is an intense navigation area with more than 500 boats passing every day (passage of oil tankers, merchant ships and fishing boats). It is considered as a strategic passage between Northern Europe and China, on one hand, and the Americas, on the other, through the ports of Antwerp, Rotterdam and Hamburg, representing nearly the quarter of the world's freight traffic.

The sampling site is far from major continental pollution sources, ~~and can be considered as a background site~~. The nearest town to the sampling site is Boulogne-sur-Mer, which is 16 km to the south, while Calais – a city that encompasses the fourth largest port in France – is about 21 km to the northeast. The A16 motorway is at a minimum distance of 10 km to the south-east. It is also worth noting that the study area is located in a SECA area where sulfur content was limited to 1% (between 2010 and 2015) for marine vessels including passenger ships and 0.1% for ships docked at the port since the sampling was conducted in 2013.

130 2.2 Samples collection

PM₁₀ sampling and measurement campaign was carried out on 24-hour basis ~~every three days~~ from 1st of January to 31st of December 2013. During the sampling period, PM₁₀ concentrations were monitored using a MP101 beta gauge analyzer (Environment SA®). The analyzer was calibrated at the beginning of the campaign and routinely checked by the regional air quality network atmo Hauts-de-France. Moreover, PM₁₀ filters were collected using an automated high-volume sampler (DA80, DIGITEL®, Switzerland) operating at 30 m³/h onto 150 mm Pall® Tissuquartz™ 2500 QAT-UP filters (no binder) filters. Filters were pre-heated for 4 hr at 450 °C before sampling to decrease the impurities. Over the sampling period, ~~122~~ 362 samples have been collected ~~and stored at 20 °C until analysis.~~ 122 samples representing a sampling of one day over three were chosen for the chemical analysis and were presented in this study. Field blanks (two per month) were also considered by placing a blank filter in sampling conditions but without pumping. Additionally, meteorological data (temperature, wind speed and direction) were recorded on site using the WMT 52 ultrasonic wind sensor (Vaisala Windcap) coupled to the DIGITEL® DA80.

2.3 PM₁₀ chemical characterization

PM₁₀ chemical characterization included the analysis of the carbonaceous subfractions (OC and EC), major and trace elements, water-soluble ions, and some organic tracers. The detailed methods can be found in the supplementary information and will be briefly presented hereafter.

The analysis of OC and EC was done using a thermo-optical technique implementing the EUSAAR-2 protocol (Cavalli et al., 2010). Major and trace elements were analyzed following the protocol described in Ledoux et al. (2006) and Kfoury et al. (2016). Major elements (Al, Ba, Fe, Mn, P, Sr, Ti, and Zn) were analyzed by Inductively Coupled Plasma-Atomic Emission Spectrometry (ICP-AES), while trace elements (V, Cr, Ni, Sc, Co, Cu, As, Rb, Nb, Ag, Cd, Sn, Sb, Te, La, Ce, Tl, Pb, and Bi) were analyzed by ICP coupled to a mass spectrometer (ICP-MS). Cl⁻, SO₄²⁻, NO₃⁻, Ca²⁺, Mg²⁺, K⁺, Na⁺, and NH₄⁺ were analyzed by liquid ion chromatography following the protocol detailed in Ledoux et al. (2006) and Fadel et al. (2022). Finally, the analysis of organic compounds included the characterization of anhydrosugars (levoglucosan, mannosan, and galactosan), sugar alcohols (arabitol and mannitol), and monosaccharides (glucose and mannose) by High Performance Liquid Chromatography (HPLC) coupled to a Pulsed Amperometric Detector (PAD) (Srivastava et al., 2018).

2.4 Data analysis

2.4.1 Constrained weighted-non-negative matrix factorization (CW-NMF)

160 A Constrained Weighted Non-Negative Factorization (CW-NMF) model was applied in this study in order to identify and
quantify the contribution of the sources to PM₁₀ concentrations. This model, developed by the LISIC (Laboratoire
d'Informatique Signal et Image de la Côte d'Opale) at the University of Littoral Côte d'Opale, has the same principles as the
USEPA PMF (Delmaire et al., 2010; Kfoury, 2013; Limem et al., 2014; Scerri et al., 2023; Scerri et al., 2019). Moreover, the
model was adjusted in order to guide it in its calculations by adding constraints. The constraints added to this model are of
two types: the “equality” constraints that defines the presence or the absence of an element in the profile and the “boundary”
165 constraints that impose a wide range of values in which the optimal solution can be found. These constraints were added in
order to consider the “a priori” knowledge of the chemical composition of the sources. For example, the concentration of
levoglucosan was set zero in all the profiles except for the biomass burning source. Further details regarding the model can
be found in Ledoux et al. (2017), and Kfoury et al. (2016).

170 The input data of the model consisted of the concentrations of 258 species in the 122 samples as well as their uncertainties.
The chosen species were OC, EC, water-soluble ions (Cl⁻, SO₄²⁻, NO₃⁻, Ca²⁺, Mg²⁺, K⁺, Na⁺, and NH₄⁺), elements (Al, ~~Cr~~, Fe,
~~Mn~~, P, Sr, Ti, Zn, V, Ni, Co, Cu, Cd, Sb, ~~La~~, and Pb), and organic tracers namely levoglucosan for biomass burning and the
sum of the concentrations of sugar alcohols and monosaccharides (named polyols) as tracers for primary biogenic emissions.
Concerning the uncertainties, it was calculated by summing errors related to the analytical and sampling procedures. For
trace elements, the calculated uncertainty was majored of 10% as it is usually reported (Prendes et al., 1999). In the case of
175 species concentration below the detection limit (D.L.), it was replaced by D.L./2 and given an uncertainty of 100%. As for
missing data, it was replaced by the mean value and was assigned an uncertainty of 400% (Polissar et al., 1998; Kim et al.,
2004). Additionally, the stability of the solution was evaluated by bootstrap analysis. It is based on the same principles of the
bootstrap analysis used in EPA PMF 5.0. The profile of a source was validated if more than 80% of the bootstrap profiles
were correlated (with $r^2 > 0.6$) to the reference one (the one obtained by considering the original dataset).

180

2.4.2 Conditional bivariate probability function (CBPF)

The conditional bivariate probability function (CBPF) combines wind speed with the conditional probability function (CPF).
The latter is calculated as the probability that the concentration of a species, in a certain wind direction, is greater than a
specific concentration value (which is the 75th percentile concentration of the different species in this study). CBPF will help
185 to further understand the dependencies of the sources to the wind speed (Uria-Tellaetxe and Carslaw, 2014). These
representations were drawn using the open source software R and the open air package (Carslaw, 2015). The dataset used
consisted of the species concentrations for each PM₁₀ sample for the 24h period distributed in front of the 48 corresponding

wind speeds and directions (one measurement every 30 min). By that, the full dataset used to draw these representations count 122 x 48 lines.

220

2.4.3 Concentration weighted trajectories (CWT)

The Concentration Weighted Trajectories (CWT) approach investigates potential transport of pollutants and/or sources over large geographical scales (Polissar et al., 2001). It consists of combining species concentrations and/or source contributions with back-trajectories and use residence time information to identify air parcels that might be responsible of high concentrations observed at the receptor site (Petit et al., 2017).

Air mass back-trajectories were calculated using the HYSPLIT model considering the Gridded Meteorological Data Archives (GDAS 1 degrees). For each sample, eight 72h backward trajectories were assigned and the source contribution values for the 24 hours period were combined with the corresponding trajectories. These representations were achieved through the Zefir 4.0 software which is an Igor based package (Petit et al., 2017).

230 3. Results and discussions

3.1 PM₁₀ concentration and composition

The concentrations of PM₁₀ as well as the concentrations of OC, EC, water-soluble ions, and elements are reported in [Table 1](#) and the concentrations of the analyzed organic tracers are reported in [Table 2](#). The yearly-mean PM₁₀ concentration obtained for the set of sampling days was of 24.3 µg/m³ which is higher than the WHO PM₁₀ annual guideline value of 20 µg/m³ that was applicable in 2013 (WHO, 2006) and the newly annual guideline value of 15 µg/m³ (WHO, 2021). Even though it is a rural ~~background~~-site, the PM₁₀ average concentration found at Cape Gris-Nez (CGN) was similar to the ones reported for urban and industrial sites in the Northern region of France in 2013 (24 µg/m³) (Atmo, 2013), suggesting that PM₁₀ sources were mainly of regional influence rather than a local one (Ledoux et al., 2018).

The predominant chemical species were OC, Cl⁻, Na⁺, and secondary inorganic ions (NO₃⁻, SO₄²⁻, and NH₄⁺), accounting for 69% of the average PM₁₀ concentrations ([Table 1](#)). Secondary inorganic ions are found in the atmosphere due to the gas to particle conversion of their corresponding precursors (NO_x, SO₂, and NH₃) emitted by different anthropogenic activities. The neutralization ratio between NH₄⁺ and the sum of SO₄²⁻ and NO₃⁻ close to 1 shows that ammonium is predominately found in the atmosphere as ammonium sulfate and ammonium nitrate. The evaluation of the concentration ratios between SO₄²⁻ and NH₄⁺ on one hand and SO₄²⁻ + NO₃⁻ and NH₄⁺ on the other hand shows that ammonium nitrate (67%) is approximately as twice as abundant compared to ammonium sulfate (33%).

Na⁺ and Cl⁻ are typical seawater components and highlight the importance of the marine influence on air quality at the investigated site. Most of the other analyzed species contribute to less than 0.1% of PM₁₀ average concentration with the

exception of EC (1.3%), Mg²⁺ (1%), Ca²⁺ (0.9%), K⁺ (0.6%), Al (0.3%), Fe (0.4%), and levoglucosan (0.2%) (**Table 1** and **Table 2**).

OC and EC showed very strong correlation during the sampling period ($r=0.89$, $p<0.05$) meaning that these species were mainly emitted from the same sources. According to the literature, OC-to-EC concentration ratios between 0.3 and 1 were reported for vehicles running on diesel, higher values were reported for biomass burning (3.4-14), while emissions from heavy fuel oil vessels show OC-to-EC ratios higher than 10 (Zhang et al., 2020; Moldanová et al., 2009; Fadel et al., 2022; Khan et al., 2021). The OC-to-EC ratio obtained in this study varied between 2.7 and 26.3 with an average ratio of 7.6. The high variability in the OC-to-EC ratio shows that these species are emitted from several sources and the interval found highlights that the traffic exhaust emissions might not highly contribute to the emissions of carbonaceous matter at CGN.

The polar plot representations of OC and EC showed that the highest concentrations of carbonaceous matter were observed for winds blowing from the northeast (NE) and southeast (SE) sectors consisting mainly of continental winds (**Figures S1-a and S1-b**). Furthermore, a pollution rose of the OC-to-EC ratio was done in order to understand if different sources of carbonaceous matter exist in the different wind sectors (**Figures S1-c**). Without considering the wind speed parameter, the highest OC-to-EC ratios were observed in the southwest sector (values higher than 10) followed by the northeast (NE) and southeast (SE) sectors where values were between 6 and 10.

The evaluation of the polar plot representation of the OC-to-EC ratio (**Figures S1-d**) shows that ratios higher than 20 are observed in the southwest sector when wind speed is higher than 10 m/s (**Figure S1-d**). When examining the concentrations of OC and EC in these samples, the high OC-to-EC ratios was mainly due to the low EC concentrations ($< 200 \text{ ng/m}^3$) found in these samples under maritime influence which increases the obtained values of the considered ratio. When removing these samples, OC-to-EC ratios were observed higher than 6 in the sectors northeast to southwest emphasizing on the influence of shipping emissions, biomass burning, and primary biogenic emissions (**Figure S1-e**).

Anhydrosugars are mainly considered as tracers of biomass burning (Vincenti et al., 2022). These compounds show higher concentrations during the cold periods of the year. The average concentration of these compounds was 185 ng/m^3 between January and March 2013, 45 ng/m^3 between November and December, and 22 ng/m^3 for the rest of the year (warmer months). According to the literature, high levoglucosan-to-mannosan concentration ratio (close to 15) is indicative of hardwood combustion while low values (between 2 and 6) are mainly attributed to softwood combustion (Schmidl et al., 2008). The average ratio in this study is 7.3 which may be considered as indicative of a mix between the two types of wood.

Finally, the sugar alcohols and monosaccharides identified in this study are mainly of primary biogenic emissions. These compounds show at least three times higher concentrations during summer (June-August 2013) compared to the rest of the sampling days in 2013 ($64 \text{ vs. } 19 \text{ ng/m}^3$, respectively). Similar observations were reported by Samaké et al. (2019) in 28 French sites where the maximum concentrations were observed in the summer season. This might be linked to higher temperatures and humidity conditions in summer that leads to the growth and sporulation of fungal and prokaryotic cell activities (Samaké et al., 2019).

315

3.2 Source profiles

A progressive approach has been adopted in order to find the best solution by CW-NMF, consisting on increasing the number of factors and evaluating the profiles.

350 The best results were obtained for the ~~940~~ factors solution. The stability of the results was examined via bootstrap analysis and the different source profiles satisfied the validation criterion ([Table S1](#)). Additionally, the reconstructed and observed PM₁₀ concentrations were very strongly correlated (slope of 0.9~~65~~ and $r^2=0.99$) ([Figure S32](#)). Furthermore, the different species considered in the CW-NMF model were well reconstructed with slopes close to 1 ([varying between 0.8 and 1.1](#)) and r^2 higher than 0.~~875~~ ([Figure S3](#)).

355 The normalized profiles of the ~~940~~ identified sources at CGN are presented in [Figure 2](#) ~~Figure 2~~ and the profiles along with the percentiles 25th and 75th calculated via the bootstrap analysis can be found in [Figure S2](#). The time series of the different profiles ~~are~~ presented in the supplementary data ([Figure S43](#)). Additionally, the distribution of the chemical species in the ~~940~~ identified sources is presented in [Figure S54](#).

Two profiles related to sea-salts emissions were identified. The first profile shows the highest loading of Na⁺ and Cl⁻ between the profiles ([Figure S54](#)) with an average Cl⁻-to-Na⁺ ratio of 1.~~78~~ which is commonly observed for fresh sea-salts (Seinfeld and Pandis, 2016). The fresh sea-salts profile also included other ionic species such as K⁺, Mg²⁺, Ca²⁺, and SO₄²⁻. The second source profile was dominated by high contributions of Na⁺ but no Cl⁻ was found. This might be due to the Cl⁻ depletion resulting from the reaction between sea-salts and NO_x and SO₂ gas (Seinfeld and Pandis, 2006). Cl⁻ was compensated with much higher contributions of NO₃⁻ and SO₄²⁻ in this profile compared to the fresh sea-salts as well as some elements from anthropogenic origins leading to the attribution of this profile to aged sea-salts.

360 The third profile contained a high proportion of Al, Fe, Ca²⁺, and K⁺ which are mainly the signature of crustal dust resuspension (Moreno et al., 2013). Two profiles show high loadings of secondary inorganic ions: the first one shows high loadings of NO₃⁻ and NH₄⁺ and was identified as “secondary nitrate” while the second one shows an abundance of SO₄²⁻ and NH₄⁺ and was ascribable to “secondary sulfate”. The secondary nitrate profile also shows a considerable contribution of OC as well as some elements from anthropogenic origins. The presence of these species in the profile is mainly linked to the aging of the secondary inorganic aerosols and/or the effect of mixing with particles emitted from combustion sources (Koçak et al., 2015; Kfoury et al., 2016).

370 The following profile was attributed to biomass burning due to the presence of levoglucosan, OC, EC, as well as K⁺, SO₄²⁻, and NO₃⁻ (Fadel et al., 2022). The ~~levoglucosan to OC~~OC-to-levoglucosan concentration ratio found in this profile (which is equal to ~~139.6~~) is within the range found in different studies for wood burning (between 7 and 17) (Fine et al., 2002). Additionally, the OC-to-EC ratio, which is found equal to ~~7.85-3~~, is also indicative of biomass burning emissions (Sonwani et al., 2021).

A profile attributed to road traffic was identified due to high loading of species emitted from the exhaust emissions (such as OC and EC) as well as elements emitted from non-exhaust emissions (such as Fe, Al, Zn, Cu, Sb). The OC-to-EC ratio of 0.67 is within the range of diesel traffic exhaust (0.3-1) (Amato et al., 2011; Waked et al., 2014).

Heavy Fuel Oil (HFO) combustion profile was composed of high loadings of carbonaceous matter (OC and EC) as well as high contributions of V, Ni, NO₃⁻, and SO₄²⁻. The concentration of OC in this profile is 4.57 times higher than EC, which is consistent with other studies (Zhang et al., 2016). The evaluation of the V-to-Ni concentration ratio show a value of 1.3 in this study which is lower than the ones usually found (close to 3) from shipping emissions (Pandolfi et al., 2011; Becagli et al., 2012). Similar observations (V-to-Ni ratio of 1.6) were previously reported for other sites in the Northern region of France (Ledoux et al., 2017). This is mainly due to the position of the sampling sites reported in the latter study as well as the site of this study that are located in a SECA where sulfur content in marine fuels is limited to 1% during the sampling period in 2013, whereas the sites reported by Pandolfi et al. (2011) and Becagli et al. (2012) were not. The quality of the fuel with a low sulfur content used for shipping was shown to contain as well low contents of V and Ni and by that change their ratio (Zhang et al., 2016; Streibel et al., 2017). Gregoris et al. (2016) also found V/Ni concentration ratios that were less than 3 in Venice and the species were mainly attributed to HFO combustion from shipping.

Finally, the primary biogenic emissions factor was identified due to the high contribution of carbonaceous matter as well as the sum of sugar alcohols and monosaccharides, considered as tracers of the source (Bauer et al., 2008). Additionally, we can observe a ~~considerable~~ contribution of P in the profile. It is known that one of the sources of atmospheric P might be the primary biogenic ones (Shi et al., 2019).

~~Finally, the last profile shows high loadings of Al, Cr, Fe, Mn, and other elements such as Ni, Cu, Pb, La, V, Co, and Cd with neither ionic nor carbonaceous species. The profile was then named "metals rich" and might be due to the industrial activities in the region (Ledoux et al., 2017). This factor explains more than 50% of Cr concentrations (Figure S4).~~

400 3.3 PM₁₀ source contributions

The average contribution of the ~~940~~ sources to the PM₁₀ concentration at CGN during 2013 is presented in Figure 3. The secondary inorganic aerosols sources, namely secondary nitrate and secondary sulfate, contribute the most to the PM₁₀ concentration with a cumulative contribution of ~~421%~~ (8.210 µg/m³). The concentrations of secondary inorganic aerosols found in our study are higher than those found in Lens, an urban background site in Northern France, for a study in 2011-2012 (28% of PM₁₀, 5.9 µg/m³) and in Nogent-sur-Oise, an urban site in Northern France in 2013 (27% of PM₁₀, 7.1 µg/m³) (Oliveira, 2017; Waked et al., 2014).

The sea-salts (fresh and aged) contribute together to 37% of PM₁₀ concentration (7.29 µg/m³) (Table S2) with higher contribution found to the fresh sea-salts compared to the aged ones. These contributions are found higher than those found for non-coastal sites in the Northern region such as in Rouen between 2010 and 2011 (21% of PM₁₀, 4.6 µg/m³) and in Lens

between 2011 and 2012 (8% of PM₁₀, 1.6 µg/m³) (Waked et al., 2014; Favez et al., 2011) showing the high influence of sea spray emissions at the CGN site.

445 Biomass burning is also considered as an important source of PM₁₀ at CGN, contributing ~~84%~~ (1.52-4 µg/m³) (Table S2). This phenomenon is mainly observed in rural ~~background~~-sites where ~~biomass, specifically~~ wood; burning ~~for residential heating episodes are is~~ remarkably high (Golly et al., 2019). The contribution found in this study is similar to the yearly-average concentration found in France for biomass burning estimated at 2.5 ± 1.2 µg/m³ (Favez et al., 2021). This source is considered as the largest contributor to organic aerosols influencing the overall urban air quality in the country (Favez et al.,
450 2021).

The ~~five four~~ remaining sources contribute together to 12.83% of PM₁₀ (Figure 3Figure 3). Due to the rural typology of the site, the road traffic contribution to PM₁₀ (2.82%, 0.5442 µg/m³) is lower than the ones found in Lens (6%, 1.2 µg/m³) and in other urban sites in France (15 ± 7 % of PM₁₀) (Waked et al., 2014; Weber et al., 2021). It is important to mention that in CGN, the contribution of HFO combustion that might be mainly related to shipping emissions (4.55-3%) is higher than the
455 road traffic contribution. This percentage is in the range of the contribution from shipping emissions (1-7%) to PM₁₀ in different European coastal areas (Barcelona, Venice, Melilla, Algeciras, Lampedusa) (Pandolfi et al., 2011; Becagli et al., 2012; Viana et al., 2014; Amato et al., 2009; Viana et al., 2009). In addition to the contribution of shipping emissions to the PM₁₀ mass concentration, Viana et al. (2014) reported that this source may also influence new particle formation and thus further contributing to the air quality degradation. It is worth noting that the contribution to PM₁₀ presented in this work
460 corresponds to the direct emissions from the source. However, considerable amounts of SO₂ and NO_x can be emitted from shipping activities and can be transformed into secondary compounds by gas-to-particle conversion that largely contribute to PM mass. Ledoux et al. (2018) have reported that the impact of shipping in the harbor of Calais in Northern France was estimated to 35% of NO, 51% of SO₂, and 15% of NO₂ average concentrations. ~~showing that the contribution of shipping emissions to PM₁₀ at CGN could be underestimated.~~ Nevertheless, the transformation reaction of SO₂ emitted from the
465 ~~shipping emissions to SO₄²⁻ is probably not at the equilibrium yet and maybe SO₂ emissions contribute to the formation of more SO₄²⁻ than the quantity sampled close to the sea.~~

3.4 Monthly variations of the major PM₁₀ sources

470 The monthly variations of the contributions of the ~~940~~ identified sources at CGN during 2013 are presented in Figure S65. Secondary inorganic aerosols (secondary nitrate and sulfate) show a similar trend with higher contributions recorded during winter period. In this study, the highest contribution for ammonium nitrate was recorded in March (14 µg/m³ on average), when PM pollution episodes in Western Europe are dominated by secondary aerosols (Petit et al., 2019). Secondary nitrate shows higher contributions during spring (March-May) and lower contributions during summer (June-
475 August) compared to ammonium sulfate. This might be partially explained by the meteorological conditions and the semi-

volatility of nitrate while ammonium sulfate is considered more thermally stable (Petit et al., 2019). Similar seasonal trends were observed by Waked et al. (2014) and Weber et al. (2019). Biomass burning shows a strong seasonality with the highest contributions during the cold period (January to March) due to wood burning for residential heating. The average contribution of biomass burning to PM₁₀ during these three months is ~~15.4~~~~19.2~~% which is twice higher than the average contribution during the whole year (**Figure S6**).

Fresh and aged sea-salts as well as HFO combustion from shipping emissions, which contribute together to 42% of PM₁₀ during the total sampling period also show some temporal differences and were presented in **Figure 4**~~Figure 4~~. The figure also presents their conditional bivariate probability function (CBPF) plots in order to further interpret the highest contributions observed for these sources. Different trends of contributions were observed for the fresh and aged sea-salts. The higher contributions of fresh sea-salts source were recorded in November (~~6.97~~ $\mu\text{g}/\text{m}^3$) and in December ($7.5 \mu\text{g}/\text{m}^3$) while those of aged sea-salts in ~~April-May~~ (~~5.64-5~~ $\mu\text{g}/\text{m}^3$) and ~~June-May~~ ($5.4 \mu\text{g}/\text{m}^3$). Additionally, the aged sea-salts contribution is found lower than the fresh-salts throughout the sampling period except for summer months where the average contribution concentrations of aged sea-salts were ~~2.73~~ and ~~2.08~~ $\mu\text{g}/\text{m}^3$ compared to ~~2.04~~ and ~~0.78~~ $\mu\text{g}/\text{m}^3$ for fresh sea-salts in July and August, respectively. The main reason for these differences might be the meteorological conditions. Indeed, the CBPF representation of fresh sea-salts clearly evidenced that the maximum concentrations were observed for winds blowing from the southwest (SW) and northeast (NE) wind sectors corresponding to the English Channel and the North Sea, respectively and for medium to high wind speeds ($> 10 \text{ m/s}$) (**Figure 1**~~Figure 1~~). These wind directions were predominant during all months of the year except for the summer season which could explain the higher concentrations of fresh compared to aged sea-salts. On the other hand, the maximum concentrations of aged sea-salts were obtained when the wind blew from the northeast wind sector with wind speeds higher than 10 m/s (**Figure 4**~~Figure 4~~). This might be explained by the reaction of the fresh sea-salts with SO₂ and NO₂ that also show the highest concentrations in the northeast wind sector (**Figure S7**~~6~~) to yield aged sea-salts in the Strait of Dover and the North Sea area (**Figure 1**~~Figure 1~~). The high concentrations of these gases in this wind sector are predominantly emitted by ships crossing the English Channel as suggested by Ledoux et al. (2018).

As for the HFO combustion from shipping emissions, the highest contribution was observed during July with ~~ten-15~~ times higher average concentration during this month (~~2.5365~~ $\mu\text{g}/\text{m}^3$) compared to January (~~0.1725~~ $\mu\text{g}/\text{m}^3$). This source contributes to ~~184~~% of PM₁₀ concentration during July which is approximately ~~three-four~~ times higher than the yearly average contribution (**Figure 3**~~Figure 3~~). The CBPF representation shows that the high contribution concentrations of HFO combustion were observed for winds blowing from the northeast and southwest wind sectors with low wind speed ($< 5 \text{ m/s}$). According to the Marine Management Organization (MMO), the number of ferries was 1.5 times higher during this specific period while the traffic in the English Channel – North Sea remains constant which could account for explaining the higher contribution (MMO, 2014). In addition to that, meteorological conditions characterized by low wet deposition and low wind speed favor the accumulation of pollutants (Aulinger et al., 2015).

3.5 Regional influence

The representations of the source contributions with the air-mass back trajectories using the CWT method are useful to study the impact of regional emissions on PM₁₀ concentrations. The focus will be on the sources identified by CW-NMF that largely contribute to PM₁₀ concentrations and that can be of regional origins. For this purpose, we present in **Figure 5**, the CWT representations for fresh and aged sea-salts, secondary nitrate and sulfate, as well as HFO combustion that correspond altogether to **84%** of PM₁₀ during the sampling period at CGN (**Figure 3**). The CWT analysis exhibits specific hotspots for sea-salts factors over the North Sea, and the English Channel. Additionally, the coastal part of the Atlantic Ocean can be considered as an additional hotspot for fresh sea-salts (**Figure 5**). This can be also observed in the CBPF representations that show high contributions of the fresh sea-salts when the air blows from the southwest sector with high wind speeds (higher than 10 m/s) (**Figure 4**). The CWT model highlights once again the influence of HFO combustion corresponding to shipping emissions and shows that the North Sea, the English Channel, and the Strait of Dover can be considered as hotspots for this source.

On the other hand, secondary nitrate and secondary sulfate show similar geographical origins with main hotspots located over Belgium, the ~~southern part of the~~ Netherlands, and Western Germany.

These observations were consistent with other studies that highlighted that these sources are mainly inland and associated with continental air masses (Moufarrej et al., 2020; Petit et al., 2019). These regions were considered as important SO₂ emitters produced by power generation and transformation industries and NO₂ emitters from road transport, power plants, and other fuel converters (2012; Pay et al., 2010). Nevertheless, the North Sea can be also seen as a hotspot for these sources.

As for PM₁₀, the highest concentrations were traced back to marine areas such as the North Sea and the English Channel as well as some European countries such as Northern France, Belgium, Netherlands, and Western Germany. Waked et al. (2018) investigated the geographical origins of PM₁₀ impacting the North of France during the period between 2009 and 2013 and have also found similar results where air masses crossing Belgium, the Netherlands, and the North Sea were associated with intense anthropogenic activities and were considered as the highest potential source areas.

4 Conclusions

The main objective of this work was to determine the contribution of natural and anthropogenic maritime sources to PM₁₀ levels in a coastal site in the North of France. In order to do that, a PM₁₀ sampling and measurement campaign was performed in 2013 at a coastal site in front of the Straits of Dover, named Cape Gris-Nez. PM₁₀ was chemically characterized for its carbonaceous, elemental, and ionic fractions as well as some organic tracers of biomass burning and primary biogenic emissions. This exhaustive characterization was essential in order to explain the PM₁₀ concentration levels observed in the

North of France and to estimate the sources' contributions to PM₁₀ using a receptor model which was Constrained Weighted
605 Non-Negative Matrix Factorization.

In 2013, at Cape Gris-Nez, PM₁₀ mean value was 24.3 µg/m³ and is very similar to those observed in several other sites in Northern France. Six species explains about 70% of the total mass of PM₁₀: NO₃⁻, OC, SO₄²⁻, Cl⁻, Na⁺, and NH₄⁺. Using the CW-NMF model, ~~94~~ source profiles were identified and were associated to natural and anthropogenic emissions. During the period of study, the mean contributions of the different sources were 37% for fresh and aged sea-salts, ~~42~~ 42% for the
610 secondary inorganic aerosols, ~~84~~ 84% for biomass combustion, and almost 5% for marine traffic. Additionally, the study of the monthly evolution of the sources' contribution shows that secondary nitrate and biomass burning were predominant during the cold season. As for the summer season, the impact of marine traffic and the predominance of aged sea-salts versus fresh sea-salts were mainly evidenced. The contribution of HFO combustion from shipping in July was found to be ~~ten~~ 15 times
higher than in January.

615 Finally, CWT analysis showed that the North Sea, the English Channel, and the coastal part of the Atlantic Ocean are important hotspots for maritime emissions whereas Belgium, Netherlands, and the West of Germany are hotspots for secondary sources, emphasizing on the role of long-range transport on the air quality in the Cape Gris-Nez in particular and in the North of France in general.

Financial support

620 This project corresponds to a scientific contribution included in the Atmosphere Protection Plan funded by the Environment, Planning and Housing Regional Agency (Hauts-de-France) as well as by the French ministry of environment.

The "Unité de Chimie Environnementale et Interactions sur le Vivant", UCEIV UR4492, participates in the CLIMIBIO project, which is financially supported by the Hauts-de-France Region Council, the Ministry of Higher Education and Research, the European Regional Development Funds. Cloé Roche is grateful to the "Pôle Métropolitain Côte d'Opale"
625 (PMCO) for the funding of her PhD.

Acknowledgment

The authors would like to thank the "Centre Commun de Mesures, ULCO" and specifically Fabrice Cazier and Dorothee Dewaele, as well as the regional air quality monitoring network Atmo Haut-de-France, for their contribution to this project. The authors would also like to thank Amaury Kasprowiak for his help in the ionic chromatography analysis.

630 **Author contribution**

Frédéric Ledoux: data curation, conceptualization, supervision, investigation, review & editing, data analysis, and formal analysis.

Cloé Roche: sampling, chemical analysis, investigation, data curation, visualization, formal analysis, investigation, and writing.

635 Gilles Delmaire: Software, and formal analysis.

Gilles Roussel: Software, and formal analysis.

Olivier Favez: supervision of chemical analysis, review & editing.

Marc Fadel: data curation, visualization, writing original draft, review & editing.

640 Dominique Courcot: funding acquisition, conceptualization, supervision, investigation, formal analysis, project administration, and review & editing.

Competing interests

The authors declare no conflict of interest or competing interests.

Data availability

645 Data used for this study can be found at <https://doi.org/10.5281/zenodo.7664528>, <https://doi.org/10.5281/zenodo.8022729>

References

- Amato, F., Pandolfi, M., Escrig, A., Querol, X., Alastuey, A., Pey, J., Perez, N., and Hopke, P. K.: Quantifying road dust resuspension in urban environment by Multilinear Engine: A comparison with PMF2, *Atmos. Environ.*, 43, 2770-2780, <https://doi.org/10.1016/j.atmosenv.2009.02.039>, 2009.
- 650 Amato, F., Viana, M., Richard, A., Furger, M., Prévôt, A. S. H., Nava, S., Lucarelli, F., Bukowiecki, N., Alastuey, A., Reche, C., Moreno, T., Pandolfi, M., Pey, J., and Querol, X.: Size and time-resolved roadside enrichment of atmospheric particulate pollutants, *Atmos. Chem. Phys.*, 11, 2917-2931, 10.5194/acp-11-2917-2011, 2011.
- Andersson, C., Bergström, R., and Johansson, C.: Population exposure and mortality due to regional background PM in Europe – Long-term simulations of source region and shipping contributions, *Atmos. Environ.*, 43, 3614-3620, <https://doi.org/10.1016/j.atmosenv.2009.03.040>, 2009.
- 655 Atmo: Bilan territorial de la qualité de l'air, 2013.

- Aulinger, A., Matthias, V., Zeretzke, M., Bieser, J., Quante, M., and Backes, A.: The impact of shipping emissions on air pollution in the Greater North Sea region – Part 1: Current emissions and concentrations, *Atmos. Chem. Phys. Discuss.*, 15, 11277-11323, <https://doi.org/10.5194/acpd-15-11277-2015>, 2015.
- 660 Ausmeel, S., Eriksson, A., Ahlberg, E., Sporre, M. K., Spanne, M., and Kristensson, A.: Ship plumes in the Baltic Sea Sulfur Emission Control Area: chemical characterization and contribution to coastal aerosol concentrations, *Atmos. Chem. Phys.*, 20, 9135-9151, <https://doi.org/10.5194/acp-20-9135-2020>, 2020.
- Bauer, H., Claeys, M., Vermeylen, R., Schüller, E., Weinke, G., Berger, A., and Puxbaum, H.: Arabitol and mannitol as tracers for a quantification of airborne fungal spores, *Atmos. Environ.*, 42, 588-593, 665 <https://doi.org/10.1016/j.atmosenv.2007.10.013>, 2008.
- Becagli, S., Sferlazzo, D. M., Pace, G., di Sarra, A., Bommarito, C., Calzolari, G., Ghedini, C., Lucarelli, F., Meloni, D., Monteleone, F., Severi, M., Traversi, R., and Udisti, R.: Evidence for heavy fuel oil combustion aerosols from chemical analyses at the island of Lampedusa: a possible large role of ships emissions in the Mediterranean, *Atmos. Chem. Phys.*, 12, 3479-3492, <https://doi.org/10.5194/acp-12-3479-2012>, 2012.
- 670 Bove, M. C., Brotto, P., Calzolari, G., Cassola, F., Cavalli, F., Fermo, P., Hjorth, J., Massabò, D., Nava, S., Piazzalunga, A., Schembari, C., and Prati, P.: PM₁₀ source apportionment applying PMF and chemical tracer analysis to ship-borne measurements in the Western Mediterranean, *Atmos. Environ.*, 125, 140-151, <https://doi.org/10.1016/j.atmosenv.2015.11.009>, 2016.
- Carslaw, D. C.: The openair manual — open-source tools for analysing air pollution data, Manual for version 1.1-4, King's 675 College London, 2015.
- Cavalli, F., Viana, M., Yttri, K. E., Genberg, J., and Putaud, J. P.: Toward a standardised thermal-optical protocol for measuring atmospheric organic and elemental carbon: the EUSAAR protocol, *Atmos. Meas. Tech.*, 3, 79-89, <https://doi.org/10.5194/amt-3-79-2010>, 2010.
- Cesari, D., Genga, A., Ielpo, P., Siciliano, M., Mascolo, G., Grasso, F. M., and Contini, D.: Source apportionment of PM_{2.5} 680 in the harbour-industrial area of Brindisi (Italy): identification and estimation of the contribution of in-port ship emissions, *Sci Total Environ*, 497-498, 392-400, <https://doi.org/10.1016/j.scitotenv.2014.08.007>, 2014.
- Contini, D. and Merico, E.: Recent Advances in Studying Air Quality and Health Effects of Shipping Emissions, *Atmosphere*, 12, 92, <https://doi.org/10.3390/atmos12010092>, 2021.
- Contini, D., Gambaro, A., Belosi, F., De Pieri, S., Cairns, W. R. L., Donateo, A., Zanotto, E., and Citron, M.: The direct 685 influence of ship traffic on atmospheric PM_{2.5}, PM₁₀ and PAH in Venice, *Journal of Environmental Management*, 92, 2119-2129, <https://doi.org/10.1016/j.jenvman.2011.01.016>, 2011.
- Corbett, J. J., Winebrake, J. J., Green, E. H., Kasibhatla, P., Eyring, V., and Lauer, A.: Mortality from Ship Emissions: A Global Assessment, *Environ. Sci. Tech.*, 41, 8512-8518, <https://doi.org/10.1021/es071686z>, 2007.
- Crenn, V., Fronval, I., Petitprez, D., and Riffault, V.: Fine particles sampled at an urban background site and an 690 industrialized coastal site in Northern France — Part 1: Seasonal variations and chemical characterization, *Sci. Total Environ.*, 578, <https://doi.org/10.1016/j.scitotenv.2015.11.165>, 2015.
- Delmaire, G., Roussel, G., Hleis, D., and Ledoux, F.: Une version pondérée de la Factorisation Matricielle Non négative pour l'identification de sources de particules atmosphériques. Application au littoral de la Mer du Nord, *J. Eur. Syst. Autom.*, 44, <https://doi.org/10.3166/jesa.44.547-566>, 2010.

- 695 Donateo, A., Gregoris, E., Gambaro, A., Merico, E., Giua, R., Nocioni, A., and Contini, D.: Contribution of harbour activities and ship traffic to PM_{2.5}, particle number concentrations and PAHs in a port city of the Mediterranean Sea (Italy), *Environ Sci Pollut Res Int*, 21, 9415-9429, <https://doi.org/10.1007/s11356-014-2849-0>, 2014.
- EEA: The impact of international shipping on European air quality and climate forcing. European Environmental agency. Technical report No 4/2013, 2013.
- 700 EEA: European Maritime Transport Environmental Report 2021, 2021.
- Endresen, Ø., Sjørgård, E., Sundet, J. K., Dalsøren, S. B., Isaksen, I. S. A., Berglen, T. F., and Gravir, G.: Emission from international sea transportation and environmental impact, *Journal of Geophysical Research: Atmospheres*, 108, <https://doi.org/10.1029/2002JD002898>, 2003.
- 705 Fadel, M., Ledoux, F., Seigneur, M., Oikonomou, K., Sciare, J., Courcot, D., and Afif, C.: Chemical profiles of PM_{2.5} emitted from various anthropogenic sources of the Eastern Mediterranean: Cooking, wood burning, and diesel generators, *Environmental Research*, 211, 113032, <https://doi.org/10.1016/j.envres.2022.113032>, 2022.
- Favez, O., Lemeur, S., and Petit, J.-E.: Suivi de la composition chimique journalière des PM_{2.5} et PM₁₀ sur la station Petit Quevilly (agglomération de Rouen) d'Air Normand entre octobre 2010 et octobre 2011. NOTE DU LCSQA - Métrologie des particules, 2011.
- 710 Favez, O., Weber, S., Petit, J.-E., Alleman, L. Y., Albinet, A., Riffault, V., Chazeau, B., Amodeo, T., Salameh, D., Zhang, Y., Srivastava, D., Samaké, A., Aujay-Plouzeau, R., Papin, A., Bonnaire, N., Boullanger, C., Chatain, M., Chevrier, F., Detournay, A., Dominik-Sègue, M., Falhun, R., Garbin, C., Gherzi, V., Grignon, G., Levigoureux, G., Pontet, S., Rangognio, J., Zhang, S., Besombes, J.-L., Conil, S., Uzu, G., Savarino, J., Marchand, N., Gros, V., Marchand, C., Jaffrezo, J.-L., and Leoz-Garziandia, E.: Overview of the French Operational Network for In Situ Observation of PM Chemical
- 715 *Composition and Sources in Urban Environments (CARA Program)*, *Atmosphere*, 12, 207, <https://doi.org/10.3390/atmos12020207>, 2021.
- Fine, P. M., Cass, G. R., and Simoneit, B. R. T.: Chemical characterization of fine particle emissions from the fireplace combustion of woods grown in the southern United States, *Environ. Sci. Tech.*, 36, 1442-1451, <https://doi.org/10.1021/es0108988>, 2002.
- 720 Golly, B., Waked, A., Weber, S., Samake, A., Jacob, V., Conil, S., Rangognio, J., Chrétien, E., Vagnet, M. P., Robic, P. Y., Besombes, J. L., and Jaffrezo, J. L.: Organic markers and OC source apportionment for seasonal variations of PM_{2.5} at 5 rural sites in France, *Atmos. Environ.*, 198, 142-157, <https://doi.org/10.1016/j.atmosenv.2018.10.027>, 2019.
- Gregoris, E., Barbaro, E., Morabito, E., Toscano, G., Donateo, A., Cesari, D., Contini, D., and Gambaro, A.: Impact of maritime traffic on polycyclic aromatic hydrocarbons, metals and particulate matter in Venice air, *Environmental Science and Pollution Research*, 23, 6951-6959, <https://doi.org/10.1007/s11356-015-5811-x>, 2016.
- 725 Hellebust, S., Allanic, A., O'Connor, I. P., Jourdan, C., Healy, D., and Sodeau, J. R.: Sources of ambient concentrations and chemical composition of PM_{2.5-0.1} in Cork Harbour, Ireland, *Atmos. Res.*, 95, 136-149, <https://doi.org/10.1016/j.atmosres.2009.09.006>, 2010.
- Jonson, J. E., Gauss, M., Schulz, M., Jalkanen, J. P., and Fagerli, H.: Effects of global ship emissions on European air pollution levels, *Atmos. Chem. Phys.*, 20, 11399-11422, <https://doi.org/10.5194/acp-20-11399-2020>, 2020.
- 730

- Jutterström, S., Moldan, F., Moldanová, J., Karl, M., Matthias, V., and Posch, M.: The impact of nitrogen and sulfur emissions from shipping on the exceedance of critical loads in the Baltic Sea region, *Atmos. Chem. Phys.*, 21, 15827-15845, <https://doi.org/10.5194/acp-21-15827-2021>, 2021.
- 735 Kfoury, A.: Origin and physicochemical behaviour of atmospheric PM in cities located in the area of the Nord-Pas-de-Calais region, France.,
- Kfoury, A., Ledoux, F., Roche, C., Delmaire, G., Roussel, G., and Courcot, D.: PM_{2.5} source apportionment in a French urban coastal site under steelworks emission influences using constrained non-negative matrix factorization receptor model, *J. Environ. Sci.*, 40, 114-128, <https://doi.org/10.1016/j.jes.2015.10.025>, 2016.
- 740 Khan, J. Z., Sun, L., Tian, Y., Shi, G., and Feng, Y.: Chemical characterization and source apportionment of PM₁ and PM_{2.5} in Tianjin, China: Impacts of biomass burning and primary biogenic sources, *J. Environ. Sci.*, 99, 196-209, <https://doi.org/10.1016/j.jes.2020.06.027>, 2021.
- Kim, E., Hopke, P. K., and Edgerton, E. S.: Improving source identification of Atlanta aerosol using temperature resolved carbon fractions in positive matrix factorization, *Atmos. Environ.*, 38, 3349-3362, <https://doi.org/10.1016/j.atmosenv.2004.03.012>, 2004.
- 745 Koçak, M., Mihalopoulos, N., Tutsak, E., Theodosi, C., Zampas, P., and Kalegeri, P.: PM₁₀ and PM_{2.5} composition over the Central Black Sea: origin and seasonal variability, *Environmental Science and Pollution Research*, 22, 18076-18092, <https://doi.org/10.1007/s11356-015-4928-2>, 2015.
- Ledoux, F., Roche, C., Cazier, F., Beaugard, C., and Courcot, D.: Influence of ship emissions on NO_x, SO₂, O₃ and PM concentrations in a North-Sea harbor in France, *J. Environ. Sci.*, 71, 56-66, <https://doi.org/10.1016/j.jes.2018.03.030>, 2018.
- 750 Ledoux, F., Kfoury, A., Delmaire, G., Roussel, G., El Zein, A., and Courcot, D.: Contributions of local and regional anthropogenic sources of metals in PM_{2.5} at an urban site in northern France, *Chemosphere*, 181, 713-724, <https://doi.org/10.1016/j.chemosphere.2017.04.128>, 2017.
- Ledoux, F., Laversin, H., Courcot, D., Courcot, L., Zhilinskaya, E. A., Puskaric, E., and Aboukais, A.: Characterization of iron and manganese species in atmospheric aerosols from anthropogenic sources, *Atmos. Res.*, 82, 622-632, <https://doi.org/10.1016/j.atmosres.2006.02.018>, 2006.
- 755 Limem, A., Delmaire, G., Puigt, M., Roussel, G., and Courcot, D.: Non-negative Matrix Factorization under equality constraints—a study of industrial source identification, *Appl. Numer. Math.*, 85, 1-15, <https://doi.org/10.1016/j.apnum.2014.05.009>, 2014.
- 760 Lv, Z., Liu, H., Ying, Q., Fu, M., Meng, Z., Wang, Y., Wei, W., Gong, H., and He, K.: Impacts of shipping emissions on PM_{2.5} pollution in China, *Atmos. Chem. Phys.*, 18, 15811-15824, <https://doi.org/10.5194/acp-18-15811-2018>, 2018.
- Manders, A., Schaap, M., Querol, X., Albert, M. F. M. A., Vercauteren, J., Kuhlbusch, T. A. J., and Hoogerbrugge, R.: Sea salt concentrations across the European continent, *Atmos. Environ.*, 44, 2434-2442, <https://doi.org/10.1016/j.atmosenv.2010.03.028>, 2010.
- 765 Marmer, E., Dentener, F., Aardenne, J. v., Cavalli, F., Vignati, E., Velchev, K., Hjorth, J., Boersma, F., Vinken, G., Mihalopoulos, N., and Raes, F.: What can we learn about ship emission inventories from measurements of air pollutants over the Mediterranean Sea?, *Atmos. Chem. Phys.*, 9, 6815-6831, <https://doi.org/10.5194/acp-9-6815-2009>, 2009.

- Matthias, V., Bewersdorff, I., Aulinger, A., and Quante, M.: The contribution of ship emissions to air pollution in the North Sea regions, *Environ. Pollut.*, 158, 2241-2250, <https://doi.org/10.1016/j.envpol.2010.02.013>, 2010.
- 770 Mazzei, F., D'Alessandro, A., Lucarelli, F., Nava, S., Prati, P., Valli, G., and Vecchi, R.: Characterization of particulate matter sources in an urban environment, *Sci. Total Environ.*, 401, 81-89, <https://doi.org/10.1016/j.scitotenv.2008.03.008>, 2008.
- MMO: Mapping UK shipping density and routes from AIS. Marine Management Organisation. https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/317770/1066.pdf, 2014.
- 775 Moldanová, J., Fridell, E., Popovicheva, O., Demirdjian, B., Tishkova, V., Faccineto, A., and Focsa, C.: Characterisation of particulate matter and gaseous emissions from a large ship diesel engine, *Atmos. Environ.*, 43, 2632-2641, <https://doi.org/10.1016/j.atmosenv.2009.02.008>, 2009.
- 780 Moreno, T., Karanasiou, A., Amato, F., Lucarelli, F., Nava, S., Calzolari, G., Chiari, M., Coz, E., Artíñano, B., Lumberras, J., Borge, R., Boldo, E., Linares, C., Alastuey, A., Querol, X., and Gibbons, W.: Daily and hourly sourcing of metallic and mineral dust in urban air contaminated by traffic and coal-burning emissions, *Atmos. Environ.*, 68, 33-44, <https://doi.org/10.1016/j.atmosenv.2012.11.037>, 2013.
- Moufarrej, L., Courcot, D., and Ledoux, F.: Assessment of the PM_{2.5} oxidative potential in a coastal industrial city in Northern France: Relationships with chemical composition, local emissions and long range sources, *Sci. Total Environ.*, 748, 141448, <https://doi.org/10.1016/j.scitotenv.2020.141448>, 2020.
- Oliveira, D.: Identification of the main sources and geographical origins of PM₁₀ in the northern part of France, 2017.
- 785 Pandolfi, M., Gonzalez-Castanedo, Y., Alastuey, A., de la Rosa, J. D., Mantilla, E., de la Campa, A. S., Querol, X., Pey, J., Amato, F., and Moreno, T.: Source apportionment of PM₁₀ and PM_{2.5} at multiple sites in the strait of Gibraltar by PMF: impact of shipping emissions, *Environ Sci Pollut Res Int*, 18, 260-269, <https://doi.org/10.1007/s11356-010-0373-4>, 2011.
- 790 Pay, M. T., Jiménez-Guerrero, P., and Baldasano, J. M.: Assessing sensitivity regimes of secondary inorganic aerosol formation in Europe with the CALIOPE-EU modeling system, *Atmos. Environ.*, 51, 146-164, <https://doi.org/10.1016/j.atmosenv.2012.01.027>, 2012.
- Pay, M. T., Piot, M., Jorba, O., Gassó, S., Gonçalves, M., Basart, S., Dabdub, D., Jiménez-Guerrero, P., and Baldasano, J. M.: A full year evaluation of the CALIOPE-EU air quality modeling system over Europe for 2004, *Atmos. Environ.*, 44, 3322-3342, <https://doi.org/10.1016/j.atmosenv.2010.05.040>, 2010.
- 795 Petit, J.-E., Favez, O., Albinet, A., and Canonaco, F.: A user-friendly tool for comprehensive evaluation of the geographical origins of atmospheric pollution: Wind and trajectory analyses, *Environmental Modelling & Software*, 88, 183-187, <https://doi.org/10.1016/j.envsoft.2016.11.022>, 2017.
- Petit, J.-E., Pallarès, C., Favez, O., Alleman, L. Y., Bonnaire, N., and Rivière, E.: Sources and Geographical Origins of PM₁₀ in Metz (France) Using Oxalate as a Marker of Secondary Organic Aerosols by Positive Matrix Factorization Analysis, *Atmosphere*, 10, 370, <https://doi.org/10.3390/atmos10070370>, 2019.
- 800 Polissar, A. V., Hopke, P. K., and Poirot, R. L.: Atmospheric Aerosol over Vermont: Chemical Composition and Sources, *Environ. Sci. Tech.*, 35, 4604-4621, <https://doi.org/10.1021/es0105865>, 2001.

- Polissar, A. V., Hopke, P. K., Paatero, P., Malm, W. C., and Sisler, J. F.: Atmospheric aerosol over Alaska: 2. Elemental composition and sources, *Journal of Geophysical Research: Atmospheres*, 103, 19045-19057, <https://doi.org/10.1029/98JD01212>, 1998.
- 805 Prendes, P., Andrade, J. M., López-Mahía, P., and Prada, D.: Source apportionment of inorganic ions in airborne urban particles from Coruña city (N.W. of Spain) using positive matrix factorization, *Talanta*, 49, 165-178, [https://doi.org/10.1016/s0039-9140\(98\)00356-7](https://doi.org/10.1016/s0039-9140(98)00356-7), 1999.
- Rimetz-Planchon, J., Perdrix, E., Sobanska, S., and Brémard, C.: PM₁₀ air quality variations in an urbanized and industrialized harbor, *Atmos. Environ.*, 42, 7274-7283, <https://doi.org/10.1016/j.atmosenv.2008.07.005>, 2008.
- 810 Samaké, A., Jaffrezo, J.-L., Favez, O., Weber, S., Jacob, V., Canete, T., Albinet, A., Charron, A., Riffault, V., Perdrix, E., Waked, A., Golly, B., Salameh, D., Chevrier, F., Oliveira, D. M., Besombes, J.-L., Martins, J. M. F., Bonnaire, N., Conil, S., Guillaud, G., Mesbah, B., Rocq, B., Robic, P.-Y., Hulin, A., Le Meur, S., Descheemaeker, M., Chretien, E., Marchand, N., and Uzu, G.: Arabitol, mannitol, and glucose as tracers of primary biogenic organic aerosol: the influence of environmental factors on ambient air concentrations and spatial distribution over France, *Atmos. Chem. Phys.*, 19, 11013-11030, 815 <https://doi.org/10.5194/acp-19-11013-2019>, 2019.
- Scerri, M., Genga, A., Iacobellis, S., Delmaire, G., Giove, A., Siciliano, M., Siciliano, T., and Weinbruch, S.: Investigating the plausibility of a PMF source apportionment solution derived using a smaller dataset: A case study from a receptor in a rural receptor in Apulia - South East Italy, *Chemosphere*, 236, 124376, <https://doi.org/10.1016/j.chemosphere.2019.124376>, 2019.
- 820 Scerri, M. M., Weinbruch, S., Delmaire, G., Mercieca, N., Nolle, M., Prati, P., and Massabò, D.: Exhaust and non-exhaust contributions from road transport to PM₁₀ at a Southern European traffic site, *Environ. Pollut.*, 316, 120569, <https://doi.org/10.1016/j.envpol.2022.120569>, 2023.
- Schmidl, C., Bauer, H., Dattler, A., Hitztenberger, R., Weissenboeck, G., Marr, I. L., and Puxbaum, H.: Chemical characterisation of particle emissions from burning leaves, *Atmos. Environ.*, 42, 9070-9079, 825 <https://doi.org/10.1016/j.atmosenv.2008.09.010>, 2008.
- Seinfeld, J. H. and Pandis, S. N.: *Atmospheric chemistry and physics : from air pollution to climatic change*, John Wiley & Sons, Hoboken, N.J.2006.
- Seinfeld, J. H. and Pandis, S. N.: *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change*, 3r, John Wiley & Sons2016.
- 830 Seppälä, S. D., Kuula, J., Hyvärinen, A. P., Saarikoski, S., Rönkkö, T., Keskinen, J., Jalkanen, J. P., and Timonen, H.: Effects of marine fuel sulfur restrictions on particle number concentrations and size distributions in ship plumes in the Baltic Sea, *Atmos. Chem. Phys.*, 21, 3215-3234, <https://doi.org/10.5194/acp-21-3215-2021>, 2021.
- Shen, F. and Li, X.: Effects of fuel types and fuel sulfur content on the characteristics of particulate emissions in marine low-speed diesel engine, *Environmental Science and Pollution Research*, 27, 10.1007/s11356-019-07168-6, 2020.
- 835 Shi, J., Wang, N., Gao, H., Baker, A. R., Yao, X., and Zhang, D.: Phosphorus solubility in aerosol particles related to particle sources and atmospheric acidification in Asian continental outflow, *Atmos. Chem. Phys.*, 19, 847-860, <https://doi.org/10.5194/acp-19-847-2019>, 2019.

- 840 Sonwani, S., Saxena, P., and Shukla, A.: Carbonaceous Aerosol Characterization and Their Relationship With Meteorological Parameters During Summer Monsoon and Winter Monsoon at an Industrial Region in Delhi, India, *Earth and Space Science*, 8, e2020EA001303, <https://doi.org/10.1029/2020EA001303>, 2021.
- Srivastava, D., Favez, O., Bonnaire, N., Lucarelli, F., Haeffelin, M., Perraudin, E., Gros, V., Villenave, E., and Albinet, A.: Speciation of organic fractions does matter for aerosol source apportionment. Part 2: Intensive short-term campaign in the Paris area (France), *Sci Total Environ*, 634, 267-278, <https://doi.org/10.1016/j.scitotenv.2018.03.296>, 2018.
- 845 Streibel, T., Schnelle-Kreis, J., Czech, H., Harndorf, H., Jakobi, G., Jokiniemi, J., Karg, E., Lintelmann, J., Matuschek, G., Michalke, B., Müller, L., Orasche, J., Passig, J., Radischat, C., Rabe, R., Reda, A., Rüger, C., Schwemer, T., Sippula, O., Stengel, B., Sklorz, M., Torvela, T., Weggler, B., and Zimmermann, R.: Aerosol emissions of a ship diesel engine operated with diesel fuel or heavy fuel oil, *Environmental Science and Pollution Research*, 24, 10976-10991, 10.1007/s11356-016-6724-z, 2017.
- 850 Tang, L., Ramacher, M. O. P., Moldanová, J., Matthias, V., Karl, M., Johansson, L., Jalkanen, J. P., Yaramenka, K., Aulinger, A., and Gustafsson, M.: The impact of ship emissions on air quality and human health in the Gothenburg area – Part 1: 2012 emissions, *Atmos. Chem. Phys.*, 20, 7509-7530, <https://doi.org/10.5194/acp-20-7509-2020>, 2020.
- UNCTAD: Review of Maritime Transport 2021. United Nations, 2021.
- Uria-Tellaetxe, I. and Carslaw, D. C.: Conditional bivariate probability function for source identification, *Environmental Modelling & Software*, 59, 1-9, <https://doi.org/10.1016/j.envsoft.2014.05.002>, 2014.
- 855 Viana, M., Hammingh, P., Colette, A., Querol, X., Degraeuwe, B., Vliieger, I. d., and van Aardenne, J.: Impact of maritime transport emissions on coastal air quality in Europe, *Atmos. Environ.*, 90, 96-105, <https://doi.org/10.1016/j.atmosenv.2014.03.046>, 2014.
- 860 Viana, M., Amato, F., Alastuey, A., Querol, X., Moreno, T., García Dos Santos, S., Hecce, M. D., and Fernández-Patier, R.: Chemical Tracers of Particulate Emissions from Commercial Shipping, *Environ. Sci. Tech.*, 43, 7472-7477, <https://doi.org/10.1021/es901558t>, 2009.
- Vincenti, B., Paris, E., Carnevale, M., Palma, A., Guerriero, E., Borello, D., Paolini, V., and Gallucci, F.: Saccharides as Particulate Matter Tracers of Biomass Burning: A Review, *International Journal of Environmental Research and Public Health*, 19, 4387, <https://doi.org/10.3390/ijerph19074387>, 2022.
- 865 Waked, A., Bourin, A., Michoud, V., Perdrix, E., Alleman, L. Y., Sauvage, S., Delaunay, T., Vermeesch, S., Petit, J.-E., and Riffault, V.: Investigation of the geographical origins of PM₁₀ based on long, medium and short-range air mass back-trajectories impacting Northern France during the period 2009–2013, *Atmos. Environ.*, 193, 143-152, <https://doi.org/10.1016/j.atmosenv.2018.08.015>, 2018.
- 870 Waked, A., Favez, O., Alleman, L. Y., Piot, C., Petit, J. E., Delaunay, T., Verlinden, E., Golly, B., Besombes, J. L., Jaffrezo, J. L., and Leoz-Garziandia, E.: Source apportionment of PM₁₀ in a north-western Europe regional urban background site (Lens, France) using positive matrix factorization and including primary biogenic emissions, *Atmos. Chem. Phys.*, 14, 3325-3346, <https://doi.org/10.5194/acp-14-3325-2014>, 2014.
- 875 Weber, S., Salameh, D., Albinet, A., Alleman, L., Waked, A., Besombes, J.-L., Jacob, V., Guillaud, G., Meshbah, B., Rocq, B., Hulin, A., Dominik-Sègue, M., Chrétien, E., Jaffrezo, J.-L., and Favez, O.: Comparison of PM₁₀ Sources Profiles at 15 French Sites Using a Harmonized Constrained Positive Matrix Factorization Approach, *Atmosphere*, 10, <https://doi.org/10.3390/atmos10060310>, 2019.

Weber, S., Uzu, G., Favez, O., Borlaza, L. J. S., Calas, A., Salameh, D., Chevrier, F., Allard, J., Besombes, J.-L., Albinet, A., Pontet, S., Mesbah, B., Gille, G., Zhang, S., Pallares, C., Leoz-Garziandia, E., and Jaffrezo, J.-L.: Source apportionment of atmospheric PM₁₀ oxidative potential: synthesis of 15 year-round urban datasets in France, *Atmos. Chem. Phys.*, 21, 11353-11378, <https://doi.org/10.5194/acp-21-11353-2021>, 2021.

880 WHO: WHO air quality guidelines for particulate matter, ozone, nitrogen dioxide and sulfur dioxide : global assessment 2005 : summary of risk assessment, 2006.

WHO: WHO air quality guidelines. Particulate matter (PM_{2.5} and PM₁₀), ozone, nitrogen dioxide, sulfur dioxide and carbon monoxide. Geneva: World Health Organization Switzerland. Licence: CC BY-NC-SA 3.0 IGO., 2021.

885 Zetterdahl, M., Moldanová, J., Pei, X., Pathak, R. K., and Demirdjian, B.: Impact of the 0.1% fuel sulfur content limit in SECA on particle and gaseous emissions from marine vessels, *Atmos. Environ.*, 145, 338-345, <https://doi.org/10.1016/j.atmosenv.2016.09.022>, 2016.

Zhang, F., Chen, Y., Tian, C., Lou, D., Li, J., Zhang, G., and Matthias, V.: Emission factors for gaseous and particulate pollutants from offshore diesel engine vessels in China, *Atmos. Chem. Phys.*, 16, 6319-6334, <https://doi.org/10.5194/acp-16-6319-2016>, 2016.

890 Zhang, F., Guo, H., Chen, Y., Matthias, V., Zhang, Y., Yang, X., and Chen, J.: Size-segregated characteristics of organic carbon (OC), elemental carbon (EC) and organic matter in particulate matter (PM) emitted from different types of ships in China, *Atmos. Chem. Phys.*, 20, 1549-1564, <https://doi.org/10.5194/acp-20-1549-2020>, 2020.

Zhang, Y., Eastham, S. D., Lau, A. K. H., Fung, J. C. H., and Selin, N. E.: Global air quality and health impacts of domestic and international shipping, *Environ. Res. Lett.*, 16, 084055, <https://doi.org/10.1088/1748-9326/ac146b>, 2021.

895

900

905

Table 1: Average, standard deviations (S.D.), minimum (Min.), and maximum (Max.) concentrations of PM₁₀ and its chemical components (OC, EC, water-soluble ions in $\mu\text{g}/\text{m}^3$ and elements in ng/m^3) at Cape Gris-Nez (CGN) during the sampling period in 2013.

Concentrations		Average	S.D.	Min.	Max.
PM ₁₀ ($\mu\text{g}/\text{m}^3$)		24.3	13.6	5.0	74.0
Carbonaceous fraction ($\mu\text{g}/\text{m}^3$)	OC	2.14	2.05	0.33	13.7
	EC	0.32	0.28	0.02	1.69
Total carbon (TC) ($\mu\text{g}/\text{m}^3$)		2.45	2.33	0.35	15.4
Water-soluble ions ($\mu\text{g}/\text{m}^3$)	NO ₃ ⁻	5.16	5.92	0.21	33.2
	SO ₄ ²⁻	3.01	2.47	0.54	13.9
	Cl ⁻	2.48	2.56	0.003	9.3
	Na ⁺	2.11	1.52	0.10	7.88
	NH ₄ ⁺	1.94	2.54	0.03	12.9
	Mg ²⁺	0.25	0.18	0.02	0.89
	Ca ²⁺	0.21	0.26	0.05	2.26
	K ⁺	0.14	0.09	0.02	0.64
Total water-soluble ions ($\mu\text{g}/\text{m}^3$)		15.3	10.2	4.25	56.4
Elements (ng/m^3)	Fe	104	130	0.42	19.7
	Al	74.6	85.8	0.68	584
	P	30.8	43.5	1.37	253
	Zn	14.6	21.6	0.06	168
	V	5.68	6.19	0.31	33.8
	Ni	4.69	5.12	0.07	26.9
	Pb	4.67	5.62	0.12	38.4
	Ti	4.61	6.23	0.04	32.5
	Mn	4.31	7.03	0.01	40.3
	Sc	2.77	2.76	0.21	15.9
	Cu	2.22	3.08	0.03	22.3
	Ba	1.85	3.13	0.06	19.7
	Sr	1.75	1.05	0.15	5.98
	Cr	1.06	1.06	0.21	6.01
	Sn	0.69	0.88	0.01	5.49
	Sb	0.59	0.62	0.01	3.84
	As	0.29	0.38	<D.L.	2.18
	Rb	0.29	0.34	0.02	1.90
	Bi	0.22	0.44	<D.L.	2.21
	Te	0.21	0.30	<D.L.	2.01
	Co	0.18	0.20	0.01	1.25
	La	0.17	0.13	0.002	0.60
	Ce	0.12	0.13	0.005	0.72
	Cd	0.11	0.15	0.001	0.81
	Ag	0.04	0.06	0.001	0.31
	Tl	0.02	0.05	<D.L.	0.31
Nb	0.01	0.02	<D.L.	0.11	
Total elements (ng/m^3)		260	246	21.7	1434

Table 2: Average, standard deviations (S.D.), minimum (Min.), and maximum (Max.) concentrations of organic tracer species (in ng/m³) in PM₁₀ at Cape Gris-Nez (CGN) during the sampling period in 2013.

915

Concentrations in ng/m ³		Average	S.D.	Min.	Max.
Anhydrosugars	levoglucosan	55.2	110	0.01	853
	mannosan	7.60	12.2	0.01	90.6
	galactosan	3.18	7.16	0.01	51.4
Sugar alcohols	arabitol	10.7	25.1	0.01	232
	mannitol	10.7	20.0	0.01	184
Monosaccharides	glucose	6.71	9.89	0.01	60.0
	mannose	1.90	2.76	0.01	14.6

920

925






Figure 1: Location of the sampling site (© Google Earth)



930 Figure 2: PM₁₀ source profiles at Cape Gris-Nez (CGN) identified using the CW-NMF model









935

Figure 3: Mean source contributions to PM₁₀ collected at Cape Gris-Nez (CGN) during 2013


940

945



950

Figure 4: Monthly average, median and 25th and 75th percentiles of the contributions (ng/m^3) of the PM_{10} maritime sources at CGN as well the conditional bivariate probability function (CBPF) plots



955 **Figure 5: CWT results for PM₁₀ and some NMF factors (fresh sea-salts, aged sea-salts, secondary nitrate, secondary sulfate, and HFO combustion). Red colors highlight potential emission zones. Contribution scales are in $\mu\text{g}/\text{m}^3$.**