

Author's response

We are grateful to the editor and two reviewers for their time and expertise in reviewing our manuscript *Separating and Quantifying Facility-Level Methane Emissions with Overlapping Plumes for Spaceborne Methane Monitoring*.

After careful thought, we have decided to remove the LBPM part, as it is complicated and doesn't significantly improve the result. Another major modification is that we add a supplementary file with plumes of different observation pixel sizes and retrieval uncertainties, as well as comparable results for Exp2 with different observation settings. We also rephrase the text for correctness, conciseness and rigorousness.

Please see the point-by-point response as follows.

Response to Reviewer #1

- *“The overlapping of plumes is a real scenario that can occur worldwide.*

However, it is not understood the impact of these cases at a global level. Thus, it could be a very specific topic that is only applicable in some examples.”

Response: Thanks for your advice. Our motivation for separating methane plumes for quantification has arisen from real-world cases, e.g., the recently published research by EMIT (Thorpe et al., 2023). They manually inspected the L2 data and found plume overlapping can be common. The plume overlapping is especially common in O&G concentrated areas, which generally agrees with our analysis on VISTA-CA source inventory, as well as our Monte Carlo simulation for AVIRIS-NG observed sources.




Figure RC1.1 Overlapping plumes observed by (a) Worldview-3 (Sánchez-García et al., 2022); (b)

While a comprehensive global analysis of plume overlapping appears essential, it can be challenging due to its dependence on both instrumental and environmental factors, in addition to the source distribution. Thus, it seems beyond the scope of this paper. Future work could further investigate this phenomenon with real satellite observations for deeper understanding.

- *“Moreover, the proposed methodology is rather complex. Is it possible to just quantify the non-overlapped area of the plumes (of e.g. Fig 2 and 8) and apply (considering the caveats) the IME method over them?”*

Response: While quantifying the non-overlapped area of the plumes using direct morphological methods seems an interesting possibility, it presents challenges. This method requires predefined shape priors of each plume to eliminate the overlapping area and attribute plumes to sources, which can be challenging. It usually requires massive manual inspection and labeling work, and may also introduce extra systematic errors caused by the subjectivity of the operators.

Even if an algorithm were implemented to automatically eliminate the overlapping pixels and perform source attribution, obstacles may occur in specific cases. (1) When a source is located downwind of an interference source and its plume is fully covered by the interference plume, the method above is not applicable. (2) In cases where the isolated pixels are near the source (in most cases), the quantification can be unstable. As shown in Figure RC1.2, the downwind IME near the source exhibits instability, and this uncertainty will propagate proportionally into the quantification results.




Figure RC1.2 The integrated mass enhancement (IME) distribution downwind in our simulation.

The IME is calculated by integration across wind direction (i.e., over the y-axis).

- *“The results indicate a strong underestimation in the quantified flux rate when applying the separation methodology. The authors argue in L421 that this could be the result of some pixels not being attributed to any source. Thus, the results indicate that the methodology needs to be reviewed.”*

Response: We found an underestimation trend (see Figure 6), which exhibits a linear relationship with a correlation coefficient of 0.93 ($R=0.93$). Consequently, this underestimation is corrected using linear regression analysis.

We derived the correction coefficient from the simulation data and applied it to correct the quantification result of the EMIT observation. However, despite this correction, the separated quantification result on EMIT is still lower than the result of directly applying IME quantification, as shown in Table 2.

By superimposing plume pixels detected by the reference method (Varon et al., 2018; UNSEP) with those identified using our method (combination of separation and the reference method; SEP), we can see some plume pixels are excluded by our method (see Figure RC1.3). These excluded pixels can't be attributed to any source. This issue can be raised by irregular plume shapes, mislabeling of source(s), and interference from background noise. These challenges are common in this field. A detailed comprehensive

discussion of these issues seems beyond the scope of this article.




Figure RC1.3 Whole plume pixels (yellow) and separated & attributed plumes (Viridis coloring).

Response to Reviewer #2

Dear Referee,

Thank you for your insightful comment. We sincerely thank you for once again reviewing our manuscript carefully! I will present our point-by-point feedback as follows.

Point-by-Point Feedback

- *“The manuscript “Separating and Quantifying Facility-Level Methane Emissions with Overlapping Plumes for Spaceborne Methane Monitoring” by Y. Pang et al. documents an interesting topic and analysis regarding the quantification of pollutant or greenhouse gas emissions from industrial point sources based on satellite images. Such a study deserves a publication in AMT.”*

Response: Thank you sincerely for recognizing our efforts. Your insightful and practical advice has helped greatly in the improvement of our manuscript.

- *“1) However, the text of the manuscript needs a major improvement and clarifications. a) There are signs of a lack of proofreading or of rigor in the writing, such as typos (e.g. at lines 39, 257, 273, 317, 351-352, 420, 426...), clumsy formulas (l.33; “given the same emission rate” at l35; “generally” at l.38, “the high nonlinearity of the Gaussian plume model” at line 78, “accurate priors” at l211, l214-215, “the increases of multi-source Gaussian plume model” at l359-360...), misleading shortcuts (e.g. l41-42, l317-318...).”*

Response: We sincerely appreciate your pointing out our text shortcomings. These errors might have escaped our spelling and grammar checks, as well as our checks as authors for whom English is a second language. We will give it a multi-eyed and thorough check in the next revision, and try our best to avoid these grammatical errors.

- *“Some of the potential mistakes can be embarrassing: e.g. between mid-line 217 and line 218, I assume that the authors talk about chemistry rather than about diffusion, and I do not really understand what the link can be between "the low concentration of methane in the atmosphere" and the diffusion or the chemistry. The word at line 237 is "topography" rather than "topology" ? etc.”*

Response: Thank you for pointing out these potential mistakes. I will respond point-by-point as follows.

For, L217-L218: The original intent of the manuscript was that (on the simulation scale) tracer gas concentrations are low (compared to atmospheric background concentrations). For tracer gas transport, convective processes dominate the gas transport and the molecular diffusion term can be neglected. In the next revision, we will cite the opinion of Nottrott et al. (2014) directly here without a rationale.

For, L237: Thank you, it's a linguistic mistake. We will correct it in the next revision.

- *“b) In a general way, the concepts, plans and methods need to be better introduced and discussed.”*

Response: Thank you for the suggestion. The point-by-point responses are as follows.

- *“From line 68, the introduction loses the clear distinction between the plume detection and the subsequent emission quantification when processing the CH₄ plume images in two steps, or the distinction between such a two-step approach, and 1-step approaches such as Gaussian plume fitting (when using the emission rate from this fitting), while such distinctions are critical to follow the manuscript correctly.”*

Response: Thank you for the suggestion. Although in some practices Gaussian plume fitting also requires manual specification of plume pixels and background pixels, explicit pixel detection algorithms are generally not emphasized in the related studies.

If we perform fitting globally (e.g., the pixels in a surrounding rectangle), it can be considered as a “one-step” method. In comparison, other cases require a "two-step" detection-quantization approach.

However, we propose a different approach for the quantification of overlapping plumes. In this context, our proposed method can be considered as a "three-step" approach, i.e., separation-detection-quantification. We solve the overlapping plume separation by fitting Gaussian plumes with a "one-step" method and use a specific "two-step" method to achieve more robust quantification compared to the "one-step" method.

We will make this part of the introduction more concise and clearer in the next revision.

- *“Lines 72-83 (which have been extracted from section 2) should be improved and better merged in the introduction.”*

Response: Thank you for the suggestion. We will improve the text in the next revision.

- *“The abstract hardly manages to characterize the type of method that has been developed and tested in this manuscript: lines 3 to 7 seem to speak about a full quantification method and line 8 seems to introduce IME as a distinct benchmarking quantification method (and the rest of the abstract does not solve for this potential confusion).”*

Response: Thank you for the comments. As we mentioned above, we proposed a separation-detection-quantification schema. The detection-quantification method in the manuscript was basically modified from Varon’s method (Varon et al., 2018), and is considered as representative detection-quantification method. However, theoretically, the detection-quantification method here can be replaced by other methods of the same purpose, e.g., a similar IME-based method (Duren et al., 2019), angular width method (Jongaramrungruang et al., 2019), even deep learning method (Jongaramrungruang et al., 2022).

For the coherence of the text, we will move this part to the discussion and make the text more clear in the abstract and introduction.

- *“I thank the authors for having added the results of the quantification of the emissions based on the Gaussian plume model fitting in the result section. However,*

now, this benchmarking needs to be announced in the introduction or early in section 2, this quantification method needs to be detailed in section 2, and the abstract should probably highlight the comparison between this quantification method and IME.”

Response: Thanks for the suggestion for comparing the results of our method and the direct Gaussian plume fitting. We will add these parts in the next revision.

- *“This is a significant result, which tend to confirm that the Gaussian plume model inversion does not behave as well (compared to IME) when tackling turbulent plumes at fine resolution than when tackling mesoscale plumes with > 1 km resolution images.”*

Response: Our current findings show that the IME method outperforms Gaussian plume fitting in quantifying fine-scale and **small-length** plumes.

Limited by the computing constraints, we faced a resolution-scale trade-off in the WRF-LES simulation. Our preference for fine resolution limited the size of the simulation domain, preventing us from testing Gaussian plume fitting on larger scales. So, in Exp1, the number of plume pixels decreases with a sparser resolution, and the theoretical performance increase of the Gaussian plume with scale is not observed in the experiments.

- *“c) see the detailed comments below, which provide other illustrations of the general need to improve the text, its quality and its clarity, which applies to all sections.”*

Response: Thanks for the detailed suggestions. We endeavor to improve the text in the next revision. Please see our point-by-point responses as follows.

- *“2) Section 4 makes an effort to synthesize the results from section 3, to provide some explanations for the behaviors of the methods as a function of the "experimental parameters". However, it could be extended and strengthened to better characterize the problematic and successful cases depending on the methods and to provide more interpretation.”*

Response: Thanks for the suggestions and we will extend the discussion in the next revision.

- *“Furthermore, some items are probably missing such as a discussion on the impact of using the LBPM, and a discussion on the typical range of accuracy of the emissions estimates that can be expected as a function of the observation conditions.”*

Response: Thanks for your suggestion, we will extend the discussion in the next discussion.

- *“Regarding the LBPM, could this approach be hampered by the retrieval noise, or by the regular transition to the RMS metric in the optimization iterative process? See also some of the following comments that connect to potential gaps in this discussion section. Detailed comments:- is not the abstract misleading regarding the assessment of using the LBPM? the results show that it does not impact much the results, while the abstract seems to say it is successful. Section 3.3 does not really highlight the fact that this impact is relatively small, and section 4 does not discuss it at all.”*

Response: By definition (L196-197), the LBPM may be affected by strong noise ($\Delta C > 1\sigma$). The transition of LBPM to RMS primarily relies on empirical methods, as the stochastic nature of the differential evolution algorithm, makes it challenging to quantitatively analyze this iteration. In our preliminary trials, the total number of iterations generally ranges in the tens, with LBPM exponentially decaying to the same level as RMS within the first ~10 iterations.

The initial purpose of LBPM was to improve the sensitivity on shapes (gradient signs) for the fitting algorithm during the early iterations, thereby improving the fitting for weak point sources and avoiding local optima. LBPM performed exceptionally well in our preliminary tests on Gaussian plume synthetic observations. As illustrated in Figure RC2. 1, for fitting the synthetic observations (a), two estimates (b) and (c) were generated within an iteration. Despite a significant difference between its stronger source and the ground truth, (b) identified the weak source and had a lower LBPM metric from the observation. Therefore, it will be preserved for further iterated optimizations. Figure RC2. 2 shows that the LBPM outperforms the RMS in terms of convergence rate, and fitting accuracy (IoU, PWIoU) in a Monte Carlo stochastic test.

In the next revision, we will highlight LBPM's enhanced performance in detecting relatively weaker sources and extend the discussion, particularly focusing on potential areas for improvement. Additionally, we will refrain from making somewhat overly optimistic statements.




Figure RC2. 1 Demonstration of LBPM's outperformance over RMS in capturing relatively small sources in the iteration.




Figure RC2. 2 Preliminary results of plume separation of RMS (left) and RMS (right). IOU and PWIoU are two separation accuracy indicators.

- “- the abstract could mention the test on real data which is significant.”

Response: Thanks for the suggestion. We will mention it in the next revision.

- “- 111: MAPE has not been defined yet; you should rather speak about the average relative error in the estimates.”

Response: We will add it in the next revision. MAPE represents “mean absolute percentage error”.

- “- 163: how do you derive a "median interval distances of potential methane sources in California" ? I'm curious about it since the result (<200m) is surprisingly small (it is probably a matter of definition for such a term)”

Response: There are 234060 targets given in the VISTA-CA inventory with their locations in longitude and latitude. We project these locations into ESPG: 26941 (a Cartesian 2D axes for California, US, with an accuracy of ~2m). The exact intervals, Euclidean distance between one source and its nearest neighboring source, are then calculated. Over 90% of the source intervals are less than 200 m (Figure 1); if sources with intervals less than 30 m are merged, ~50% of the spacings are less than 200 m.

However, sources of type “oil and gas well” account for the vast majority (96.5%) of the inventory. If they are excluded, the median and mean interval become 495.9 m and 1247.1 m, respectively. The median is much lower than the mean, revealing the characteristic of spatial clustering distribution of emission sources. We will clarify this in the next revision.




Figure RC2. 3 Distribution of source intervals in VISTA-CA, with (left) and without (right) "Oil and Gas Wells".

- “- 168-70: unclear; you mean Nassar et al. fix the secondary plume to a specific concentration level corresponding to the secondary source rate given by the emission inventory ? ”

Response: Yes, we confirmed this point with Dr. Nassar in an Email. They fix the emission rates of the secondary sources. They also adjust them by $\pm 20\%$ to assess the uncertainty.

Some potential reasons why they cannot estimate multiple sources are listed here. Firstly, OCO-2 observations are too sparse to provide enough constraints to estimate multiple sources (Nassar’s opinion). Secondly, their estimation method LS methods are prone to local optima and are thus inferior to heuristic optimization algorithms in

finding global optima. Thirdly, Gaussian plume fitting is less robust than the combination of Gaussian plume fitting and IME method, as shown in our experiment (Exp1, Exp2).

- *“=> then you should also discuss other papers using multiple Gaussian plume model inversions to tackle multiple sources with observations similar to satellite images, i.e. something similar to the "Multi-source Gaussian plume" method tested in section 3 (e.g, Krings et al., 2011, which is already cited in the manuscript). I think that the introduction is a bit misleading regarding this: using multiple Gaussian plume model fitting to handle multiple sources is not a novelty. However, by developing an alternative approach with a multi-objective heuristic optimization for the Gaussian plume fitting combined with IME (which is often assumed to behave better than Gaussian plume model inversions when tackling fine resolution images of turbulent plumes) for the quantification, the authors bring new ideas and insights to improve the process of overlapping plumes.”*

Response: Thank you for pointing out our oversight. We have carefully reviewed the paper by Krings et al. (2011) and confirmed that they indeed employed a multi-point Gaussian plume model. We will organize and merge L68-L83 in the introduction as related work, with a specific emphasis on the contributions of Krings et al. (2011) and Nassar et al., (2017). We sincerely appreciate your summary of our innovation and we will add it to the text in the next revision!

- *“- 196; what does "the improvements of the separation algorithm in missed detection" mean ?”*

Response: Thank you for pointing out this oversight. In our latest revision, we have eliminated the pixel connectivity criterion and shifted our focus primarily to the Mean Absolute Percentage Error (MAPE) as the evaluation indicator for quantification, to make the content more concise. This sentence was overlooked during the inspection. We will scrutinize such issues in the next revision.

- *“- 1124: what could be the difference between σ_x and σ_y when using Pasquill stability classes to set-up such parameters? where could σ_x (and where does σ_x) stand in eq (2)? furthermore, the formula for the function*

sigma_y(x) needs to be given or explained”

Response: Thank you for pointing out this slip of the pen, where σ_x is redundant. cross-wind dispersion coefficient (σ_y) is a function of down-wind distance x and Pasquill stability class (which is a function to **wind speed u** , sun illumination, etc), and surface type. σ_y is given by a series of functions, which can be directly found in

Table 3. Brigg's parameterization for rural sites

Stability class	σ_y (m)	σ_z (m)
A	$0.22x(1 + 0.0001x)^{-1/2}$	$0.20x$
B	$0.16x(1 + 0.0001x)^{-1/2}$	$0.12x$
C	$0.11x(1 + 0.0001x)^{-1/2}$	$0.08x(1 + 0.0002x)^{-1/2}$
D	$0.08x(1 + 0.0001x)^{-1/2}$	$0.06x(1 + 0.0015x)^{-1/2}$
E	$0.06x(1 + 0.0001x)^{-1/2}$	$0.03x(1 + 0.0003x)^{-1}$
F	$0.04x(1 + 0.0001x)^{-1/2}$	$0.016x(1 + 0.0003x)^{-1}$

Table 4. Brigg's parameterization for urban sites

Stability class	σ_y (m)	σ_z (m)
A-B	$0.32x(1 + 0.0004x)^{-1/2}$	$0.24x(1 + 0.001x)^{1/2}$
C	$0.22x(1 + 0.0004x)^{-1/2}$	$0.20x$
D	$0.16x(1 + 0.0004x)^{-1/2}$	$0.14x(1 + 0.0003x)^{-1/2}$
E-F	$0.11x(1 + 0.0004x)^{-1/2}$	$0.08x(1 + 0.0015x)^{-1/2}$

Figure RC2. 4 Brigg's definition of dispersion coefficients.

the reference (Griffiths, 1994), as shown in the middle column in Figure RC2. 4. In some research, different definition of dispersion coefficients might be used. For instance, Masters (1998) proposed an alternative formulation, which has been adopted by studies such as those by Bovensmann et al. (2010) and Nassar et al. (2017); more localized formulations such as China's national standard [GB/T 39499-2020](#) may also be employed. Due to its diversity, we have chosen not to present them explicitly in the text.

- “- equations 3 and 4: we should have $C(x'_n, y'_n)$ rather than $C_n(x', y')$ ”

Response: The transferred coordinates (x', y') can be replaced by (x'_n, y'_n) , as they are dependent on the parameters of source n , including emission rates (Q_n) and source location (x_n, y_n) . However, by this convention, the suffix n of C_n shouldn't be omitted.

- “- 1146 The specific Gaussian plume fitting algorithm used here is questionable. Adjusting both u and Q to fit the observations can be problematic since these 2 parameters impact the plume amplitude in a similar way (there is no source of information to discriminate them in the fitting process). Furthermore, the lack of

adjustment of sigma_y in the plume fitting may limit the skill of the approach.”

Response: Thank you for inquiring about the fundamental principles of our method. As previously mentioned, the diffusion coefficient determines σ_y is a function of distance and atmospheric stability, which is a function mainly of wind speed. Thus, wind speed will indirectly determine the plume shape. As presented by Jongaramrungruang et al. (2019; Figure 9), wind speed determines the magnitude of the "half-mass angle" of the plume. Therefore, when the global optima is achieved in the plume fitting, it is possible to decouple the emission rate (Q) and the wind speed (u).

It is worth mentioning that our LPBM as an optimization objective is to some extent precisely to fit the plume shape, to avoid falling into a false local optimum.

- *“- 1170-1175 are very difficult to understand”*

Response: We will combine Figure 1 to provide a detailed explanation of the input, output, and implementation process. We will follow the sequence of parameter estimation, Gaussian plume, Gaussian blur, and weighted allocation for the implementation process.

- *“- 1281: the question is not whether these sources can be quantified, but rather whether their plumes can perturb the quantification of other sources (?)”*

Response: In the early experiments, the plume concentrations of these weak sources are generally similar to the noise level. Therefore, we exclude their analysis to focus on the larger targets, as well as saving to reduce the difficulty of program development.

- *“- 1282: I do not really understand this sentence, and in particular the term "aggressive". It seems to connect to the assumption that the emissions are constant, which makes the quantification problem and the process of the image easier.”*

Response: Assuming continuous emissions rates in our simulations serves to simplify the problem and increase the probability of plume overlap. We make this “aggressive” simplification as the inventory maker manually removes the overlapping plumes in its quality control phase (Duren et al., 2019). We will briefly state this simplification and its reasons in the next revision.

- *“- there is still a lack of explicit introduction to the fact that the whole study relies*

on scenes driven by homogeneous winds, except when tackling EMIT observations. It's implicitly guessed from the use of theta rather than theta_n in the equations, and by the quick mention to the wind speed at lines 254 (with the clumsy formula: "unified wind") and 285. There is also a lack of discussion on the fact that the use of homogeneous winds to generate the pseudo observations artificially inflates the skill of Gaussian plume models to support the plume separation: there is a piece of sentence about it at lines 461-462 but it is not very clear and it seems to be associated to very specific observation cases."

Response: Thank you for the suggestion. We employed the homogeneous winds assumption at two key stages:

1. We used homogeneous winds assumption to load LES plumes with the given 10-m winds load from ERA5 to synthesize the pseudo-observations. We will stress this point after Eq. 12.

2. Additionally, we leveraged the assumption of homogeneous winds to constrain the search range, thereby accelerating the convergence. We will emphasize this point after Eq. 4.

- *"- it would be useful to show images with different levels of observation noise, in suppl. material if not in main text (the presentation of fig 2 would be misleading if it does not include such a noise: does it ?) to provide an idea of the challenge associated to the plume separation when using noisy images."*

Response: Thanks for the suggestion, we will consider the possibility of including these images in the Supplement. In Figure 2, a noise of 1% (18 ppb) to the background concentration has been added. The source emission rates in Figure range from 0.2 to 2 t/h. It's worth noticing that the color bar is truncated, suppressing the noise visually.

- *"Actually, one would expect charts with the APE as a function of the level of noise since this level could be one of the drivers of the relative success of SEP vs. UNSEP. In a more general way, the retrieval uncertainty is a critical topic for the processing of plume images and the manuscript should bring more insights about it."*

Response: We tested on single source scenarios in EXP1, where the APE as a function of noise in bar graphs is shown in Figure 1. In EXP2, our focus shifted to quantification

comparisons under various overlapping scenarios, where factors directly influencing the overlapping index (OImass) include wind speed, wind direction, distance between two sources, and emission ratios between two sources (shown in Figure 2). The inversion noise in EXP2 was considered as a secondary factor, hence we treated it as fixed. We didn't expect there to be a large difference between SEP & UNSEP in Exp2 compared to Exp1. We will perform a new experiment of different noise levels in Exp2 to finally decide whether it is necessary to add it to the content.

- *“In particular, it should provide the values of the level of noise in a more visible way than at lines 266 and 272, and with some justification for such values.”*

Response: We will conduct a new experiment to determine the necessity of adding these plume images of different noise levels. The noise levels are mainly from Varon et al. (2018), where the typical retrieval error for GHGSat ranges from 1%-5%.

- *“What is the value of the retrieval uncertainty in EXP-3 ? Is the % applied to the CH4 background + plume signal? If yes, what is the corresponding background value ? If not, why (note that the full EMIT image is noisy in fig 8), and would not the values 1 to 3% be very low ? What are the typical relative uncertainties associated to EMIT observations ? Could the observation noise explain the relative failure of the use of the LBPM metric ?”*

Response: In EXP-3, the retrieval uncertainty is set at 1%, expressed as the fraction of the background concentration. The corresponding background is 1.8 ppm & 1 atm (with a concentration $\sim 10.3\text{g/m}^2$), as shown in L230-L231. We focused on the next generation of fine-spatial resolution and hyperspectral satellites, such as GHGSat, capable of obtaining XCH4 with high precision. The relatively limited performance of LBPM may primarily stem from the limited performance of Gaussian plume for such a small scale.

- *“- lines 290-300 poorly fit in section 2.2.3; having separate sections dedicated to the detection / quantification method(s) and to the data made available for the detection / quantification would make the presentation of the study much clearer.”*

Response: Thank you for the suggestion. In the next revision, we will introduce a new subsection to provide a more detailed implementation of both quantification schemas

(UNSEP & SEP) for the observations.

- “- L294-296: *this derivation of the wind driving the plume from the wind at 10 m height does not really make sense for the general process of satellite images. This is likely inherited from studies focusing on specific sites where dedicated local measurement of the wind at 10 m are available. In the general case, the wind should be derived from other source of knowledge (typically meteorological analysis) with a better vertical coverage (but less precision).*”

Response: Utilizing 10-m wind speed as the near-surface wind speed for the quantification is a common practice in recent studies within this field, as the direct availability of the 10-m wind components from meteorological reanalysis databases.

For our simulation, the decision to use 10-m wind speed is also driven by the need for simplification, as elaborated in L239-L243. While simulating specific scenarios with real driving wind profiles and other real boundary conditions would undoubtedly offer greater precision, the computational cost can be huge, especially when considering various factors such as wind speed and source location. Therefore, we simplify to simulate plumes of various wind speeds and use the 10-m wind speed from ERA5 as an index to load plumes from the WRF-LES database to synthesize the pseudo-observations.

- “- L297-300: *clarify how it is combined with the SEP approach*”

Response: In the next revision, we will introduce a new subsection to provide a more detailed implementation of both quantification schemas (UNSEP & SEP) for the observations.

- “- *there is no discussion on the CH₄ background mixing ratio fields (from sources outside the images, or from small point sources and diffuse area sources within the images), on how it is dealt with in the derivation of the images or when processing the images. Such a background is set to 0 in EXP 1 to EXP 3. How could it impact the theoretical results from these 3 experiments ? Do we see some residual pattern of background variations in the EMIT "enhancement data" ? could it explain part of the biases seen in section 3.4 ?*”

Response: The complexity of the real background, decided by factors such as surface

BRDF, instrument characteristics, as well as such “background” of interference concentration, has been stressed previously (Gorroño et al., 2022; Jongaramrungruang et al., 2022). Analysis considering such noise may require a specified instrument and scenario, and it is probably beyond the scope of this article. Therefore, currently, in our experiment, we only considered additive random noise for the background.

Although such a “background” wasn’t presented in the Exp1-3, it’s important to note that the unexpected enhancements in the “background” might slightly inflate the plume concentration. However, it’s still premature to conclude whether it will finally inflate the IME and Q, as the “background” may also influence pixel detection, introducing non-linearity and complicating this issue.

The monotonous surface type (desert) in Figure 8 and the relatively large pixel size of EMIT (~60 m) might help alleviate the interferences from unexpected surface features. However, as shown in Figure RC2. 5, there is a large portion of pixels identified as plume pixels by the UNSEP and rejected by the SEP. It is difficult to tell them the continuity of an up-wind plume or a new plume. This explanation could partially explain why the quantification results of the whole cluster by UNSEP are slightly higher than the result by SEP.

- *“- isn’t line 322 at odd with equation 14 ?”*

Response: Thank you for noticing this error in the text. Eq.14 is correct and the text in L322 should be “the ratio of plume mass integration from interference sources to that from the primary source”. We will correct it in the next revision.

- *“- line 328: I do not understand why "the quantification of methane source is considered as solving a regression problem". Is not the target of such a quantification the most precise estimate for a given source at a given time ?”*

Response: Thank you for the suggestion. It should be solving a parameter estimation problem.

- *“- 1367: so far, the metric for interference should be the OImass, not "the interference" ”*

Response: Thank you for pointing out this glitch. We will correct it in the next revision.

- *“- 1393: Fig 6 rather gives the feeling that UNSEP tends to overestimate the*

sources ? you meant SEP ? what do the person's R and p values correspond to in this line ?”

Response: We apologize for this slip of the pen. It should be SEP in L393. Pearson's R is the correlation coefficient between the ground truths and the predictions of source emission rates. we will correct it in the next revision.

- *“- I do not understand the correction of the SEP estimations in section 3.4 (l393-395, legend of fig 6): what is the rationale, what is done in practice ? Because of this correction, it is difficult to check whether SEP behaves better than UNSEP in section 3.4.”*

Response: As shown by Figure 6 (b), there was a systematic underestimation and we found this underestimation is linear ($R=0.93$, $p<0.01$). After a linear regression, there is $y = 0.51x - 1.31$, where y represents the quantification result of the SEP; and x represents the ground truth emission rates. So, the linear corrected quantification result can be simply expressed as $y' = 1.96y + 2.57$. The distribution of SEP is more concentrated, especially for source targets with low emission rates.

- *“- l413 vs l422: does the detection and "connectivity verification" underlying the application of the IME method really encompass the full extent of the whole set of plumes ?*

Response: In non-overlapping scenarios, the main structure is typically covered. Connectivity verification, a straightforward attribution strategy, forms the foundation of the applying IME method to scenarios with plume overlapping. However, there can be plume confusion for unseparated overlapping plumes. Moreover, this strategy may omit significant portions, as illustrated in Figure RC2. 5, where a substantial number of pixels identified as plume pixels by UNSEP are excluded by SEP.




Figure RC2. 5 Whole plume pixels (yellow, identified by SEP) and separated & attributed plumes (Viridis coloring; by UNSEP) in the EMIT observation.

- *l413 goes too fast so it is difficult to understand what it corresponds to.*”

Response: In L413, in comparison to our separated quantification approach, we also employed the traditional IME method by Varon et al. (2018) to quantify the entire plume cluster, denoted as “whole”. We will make it clearer for this point by elaborating on a more detailed implementation of two quantification schemas for the observations in a new subsection in the next revision.

- *“- l447-448: I do not understand the point which is made here, while I believe that the comparison between the Gaussian plume model inversion and IME for the emission quantification as a function of the spatial resolution and scale of the source quantification problem is an important topic”*

Response: We agree with your opinion and we tried to provide this comparison in the first experiment in 3.1. Our experiments demonstrate that for the quantification of small-scale plumes (within the 6×6 km² simulation domain), the IME method generally outperforms the Gaussian plume fitting (GPF). Limited by the computation and the scale of the simulation, a discussion for the comparison on a large scale seems beyond the scope of our current experiments.

However, it's worth noting, as shown by Varon et al. (2018), that the performance of Gaussian plume fitting increases with pixel size on a larger scale. L447-L448 tried to explain why we have not yet found the turning point of performance increase of the GPF. One explanation is that our simulation domain is too small, and the number of pixel samples begins to decrease as we increase the pixel size, which counteracts the benefit of the averaging of eddies by large pixels for performance improvement of the GPF. We will make this point clearer in the next revision.

- *“- l450 I do not understand "statistically correct", and the link made at lines 450-451 between the precision of the emission quantification using Gaussian plume fitting and the convolution kernel”*

Response: The LES simulated plume demonstrates Gaussian plume morphology after temporal averaging (Figure RC2. 6), and the LES simulation shows that a transient plume may deviate from the Gaussian behavior (Figure RC2. 7). In our initial

experiments, we found that the fitted Gaussian plume could be “spiky” near the source, resulting in missed capturing of some near-source-parts of the transient plume in some cases. To address this, we employ Gaussian blurring (i.e., convolving the image with a Gaussian kernel.), a subtle computer vision technique, to better capture these transient plumes when using Gaussian plume results as weights for the separation. Figure RC2. 8 illustrates how the Gaussian-blurred plume demonstrates more “diffusive” than the original Gaussian plume, particularly near the emission source. We will refine the text in the next revision.




Figure RC2. 6 Time-averaged plumes by WRF-LES. From left to right: 500, 800, and 1100 m of inversion height. From top to bottom: 1, 3, 5, 7, 9 m/s of wind speed.




Figure RC2. 7 Snapshots of transient plumes by WRF-LES.




Figure RC2. 8 Snapshot of LES-simulated transient plume (left), fitted Gaussian plume (middle), blurred Gaussian plume (right).

- “- I do not understand line 471”

Response: As we mentioned above, as well as shown in Figure 1, the separated plumes don’t rely on a specific plume detection and quantification method. So, a new plume pixel detection method and new quantification method can be introduced to quantify the separated plumes. We will make this point clearer in the next revision.

Reference

- Bovensmann, H., Buchwitz, M., Burrows, J. P., Reuter, M., Krings, T., Gerilowski, K., Schneising, O., Heymann, J., Tretner, A., & Erzinger, J. (2010). A remote sensing technique for global monitoring of power plant CO₂ emissions from space and related applications. *Atmospheric Measurement Techniques*, 3(4), 781–811. <https://doi.org/10.5194/amt-3-781-2010>
- Duren, R. M., Thorpe, A. K., Foster, K. T., Rafiq, T., Hopkins, F. M., Yadav, V., Bue, B. D., Thompson, D. R., Conley, S., Colombi, N. K., Frankenberg, C., McCubbin, I. B., Eastwood, M. L., Falk, M., Herner, J. D., Croes, B. E., Green, R. O., & Miller, C. E. (2019). California’s methane super-emitters. *Nature*, 575(7781), Article 7781. <https://doi.org/10.1038/s41586-019-1720-3>
- Green, R., Thorpe, A., Brodrick, P., Chadwick, D., Elder, C., Villanueva-Weeks, C., Fahlen, J., Coleman, R. W., Jensen, D., Olsen-Duvall, W., Lundeen, S., Lopez, A., & Thompson, D. (2023). EMIT L2B Methane Enhancement Data 60 m V001 [dataset]. NASA EOSDIS Land Processes Distributed Active Archive Center. <https://doi.org/10.5067/EMIT/EMITL2BCH4ENH.001>
- Griffiths, R. F. (1994). Errors in the use of the Briggs parameterization for atmospheric dispersion coefficients. *Atmospheric Environment*, 28(17), 2861–2865. [https://doi.org/10.1016/1352-2310\(94\)90086-8](https://doi.org/10.1016/1352-2310(94)90086-8)
- Jongaramrungruang, S., Frankenberg, C., Matheou, G., Thorpe, A. K., Thompson, D. R., Kuai, L., & Duren, R. M. (2019). Towards accurate methane point-source

quantification from high-resolution 2-D plume imagery. *Atmospheric Measurement Techniques*, 12(12), 6667–6681. <https://doi.org/10.5194/amt-12-6667-2019>

- Jongaramrungruang, S., Matheou, G., Thorpe, A. K., Zeng, Z.-C., & Frankenberg, C. (2021). Remote sensing of methane plumes: Instrument tradeoff analysis for detecting and quantifying local sources at global scale. *Atmospheric Measurement Techniques*, 14(12), 7999–8017. <https://doi.org/10.5194/amt-14-7999-2021>
- Masters, G. M. (1998). In *Introduction to environmental engineering and science* (2nd edn, p. 413). Prentice-Hall, Inc.
- Nassar, R., Hill, T. G., McLinden, C. A., Wunch, D., Jones, D. B. A., & Crisp, D. (2017). Quantifying CO₂ Emissions From Individual Power Plants From Space. *Geophysical Research Letters*, 44(19), Article 19. <https://doi.org/10.1002/2017GL074702>
- Nottrott, A., Kleissl, J., & Keeling, R. (2014). Modeling passive scalar dispersion in the atmospheric boundary layer with WRF large-eddy simulation. *Atmospheric Environment*, 82, 172–182. <https://doi.org/10.1016/j.atmosenv.2013.10.026>
- Sánchez-García, E., Gorroño, J., Irakulis-Loitxate, I., Varon, D. J., & Guanter, L. (2022). Mapping methane plumes at very high spatial resolution with the WorldView-3 satellite. *Atmospheric Measurement Techniques*, 15(6), Article 6. <https://doi.org/10.5194/amt-15-1657-2022>
- Thorpe, A. K., Green, R. O., Thompson, D. R., Brodrick, P. G., Chapman, J. W., Elder, C. D., Irakulis-Loitxate, I., Cusworth, D. H., Ayasse, A. K., Duren, R. M., Frankenberg, C., Guanter, L., Worden, J. R., Dennison, P. E., Roberts, D. A., Chadwick, K. D., Eastwood, M. L., Fahlen, J. E., & Miller, C. E. (2023). Attribution of individual methane and carbon dioxide emission sources using EMIT observations from space. *Science Advances*, 9(46), eadh2391. <https://doi.org/10.1126/sciadv.adh2391>
- Varon, D. J., Jacob, D. J., McKeever, J., Jervis, D., Durak, B. O. A., Xia, Y., & Huang, Y. (2018). Quantifying methane point sources from fine-scale satellite observations of atmospheric methane plumes. *Atmospheric Measurement Techniques*, 11(10), 5673–5686. <https://doi.org/10.5194/amt-11-5673-2018>