

# Uncertainties in cloud-radiative heating within an idealized extratropical cyclone

Behrooz Keshtgar<sup>1</sup>, Aiko Voigt<sup>2</sup>, Bernhard Mayer<sup>3</sup>, and Corinna Hoose<sup>1</sup>

<sup>1</sup>Institute of Meteorology and Climate Research Troposphere Research, Karlsruhe Institute of Technology, Karlsruhe, Germany

<sup>2</sup>Department of Meteorology and Geophysics, University of Vienna, Vienna, Austria

<sup>3</sup>Meteorological Institute, Ludwig Maximilian University of Munich, Munich, Germany

**Correspondence:** Behrooz Keshtgar (behrooz.keshtgar@kit.edu)

**Abstract.** Cloud radiative heating (CRH) within the atmosphere affects the dynamics and predictability of extratropical cyclones. However, CRH is uncertain in numerical weather prediction and climate models, and this could affect model predictions of extratropical cyclones. In this paper, we present a systematic quantification of CRH uncertainties. To this end, we study an idealized extratropical cyclone simulated at a convection-permitting resolution of 2.5 km, and combine large-eddy [model](#) simulations at 300 m resolution with offline radiative transfer calculations. We quantify four factors contributing to the CRH uncertainty in different regions of the cyclone: 3D cloud radiative effects, parameterization of ice-optical properties, cloud horizontal heterogeneity, and cloud vertical overlap. The two last factors can be considered to be essentially resolved at 300 m but need to be parametrized at 2.5 km resolution. Our results indicate that the parameterization of ice-optical properties and the cloud horizontal heterogeneity are the two factors contributing most to the [mean](#) uncertainty in CRH [at larger spatial scales and can be more relevant for the large-scale dynamics of the cyclone](#). On the other hand, 3D cloud radiative effects are much smaller [on average](#), especially for stratiform clouds within the warm conveyor belt of the cyclone. Our analysis in particular highlights the potential to improve the simulation of CRH by better representing ice-optical properties. Future work should in particular address how uncertainty in ice-optical properties affects the dynamics and predictability of extratropical cyclones.

## 1 Introduction

Our ability to predict extratropical cyclones is crucial for midlatitude weather as they are often associated with strong winds, heavy rain, and snow. Cloud diabatic processes, such as latent heat release due to phase changes of water, play an important role in shaping the dynamics of extratropical cyclones (Ahmadi-Givi et al., 2004; Booth et al., 2013; Binder et al., 2016). Yet, recent studies have shown that the impact of cloud radiative heating and cooling (CRH) on the dynamics of extratropical cyclones can also be substantial (Schäfer and Voigt, 2018; Keshtgar et al., 2023; Voigt et al., 2023). Our study is motivated by the CRH impact.

~~The most common effects of cloud-radiation interaction are cooling at the cloud top and warming at the cloud base, which promote convective instabilities.~~ [Within the atmosphere CRH results from the interaction of clouds with radiation in different parts of the electromagnetic spectrum. In the shortwave spectrum, clouds absorb the incoming shortwave radiation, which](#)

warms clouds and contributes to their stabilization. In the longwave spectrum, clouds absorb outgoing longwave radiation at their base and re-emit it at colder temperatures at their top, leading to substantial cooling. This pattern of cloud top cooling and the modest warming from below promotes convective instability within the cloud (Webster and Stephens, 1980). By doing so, CRH can alter cloud microphysical heating and precipitation (Klinger et al., 2017) and the intensity and predictability of an idealized extratropical cyclone (Keshtgar et al., 2023). Voigt et al. (2023) showed that the impact of CRH on extratropical cyclones differs between low-level and high-level clouds, implying that cyclones may respond differently to CRH depending on the vertical distribution of clouds. On the longer time scales of climate, modeling studies showed that CRH impacts the planetary-scale circulation of the atmosphere and its response to surface warming (Li et al., 2015; Voigt et al., 2020).

Small errors in the parameterization of subgrid-scale physical processes can quickly grow to the synoptic scale, leading to forecast errors of significant magnitude (e.g., Baumgart et al., 2019). The interaction between radiation and clouds is known to be uncertain due to many factors, such as simplified radiation schemes or poor representations of clouds in models. Keshtgar et al. (2023) showed that the modulation of latent heating by CRH within the ascending region of the cyclone changes the vertical motion and the divergent flow near the tropopause. After the divergent flow has changed, changes in potential vorticity are amplified by the rotational flow near the tropopause during the highly nonlinear stage of cyclone evolution. Keshtgar et al. (2023) further demonstrated that CRH continuously changes the latent heating during the growth phase of the cyclone. This result indicates that CRH uncertainties might affect the magnitude of latent heating and be relevant for error growth near the tropopause.

The correct simulation of CRH is challenging. Several previous studies have addressed the question of how the simulation of CRH depends on model assumptions, the representation of clouds in models, and the parameterization of radiative processes. Compared to models with coarse horizontal resolutions of 10-100 km and parameterized convection, clouds can be better simulated in kilometer-scale models and even better in large eddy models (Griewank et al., 2020; Stevens et al., 2020). In this paper, we make use of these modeling advances to present the first systematic assessment of CRH uncertainty within an extratropical cyclone by simulating the same idealized cyclone as in Keshtgar et al. (2023) at a convection-permitting resolution of 2.5 km. We further run large eddy model simulations at a horizontal resolution of 300 m for different regions of the cyclone and use their output for offline radiative transfer calculations. We assume that we have perfect knowledge of the clouds from the large eddy model simulations and this leads us to prioritize the following four factors as the potential sources of uncertainty in CRH in our study:

- 3D cloud radiative effects:

3D cloud radiative effects arise from horizontal photon propagation-transport that is not taken into account in 1D radiation schemes, as currently operational in weather and climate models. The main 3D cloud radiative effects are shortwave cloud-side illumination by shortwave radiation especially for (Jakub and Mayer, 2015, 2016), shortwave cloud-side radiation leakage (Hogan and Shonk, 2013), and longwave cloud-side absorption and emission (Klinger et al., 2017). At high solar zenith angles (Jakub and Mayer, 2015, 2016), cloud-side shortwave radiation-leakage-for cloud-side illumination increases the shortwave absorption at the cloud sides facing the sun. At low solar zenith angles (Hogan and Shonk, 2013)

~~-, and cloud-side longwave absorption and emission (Klinger et al., 2017)-,~~ however, photons can escape through cloud sides and lead to the reduction of shortwave cloud absorption. In the longwave, the horizontal transport of photons through cloud sides increases cloud radiative cooling. Thus, the largest differences between 3D and 1D radiative transfer calculations are expected to occur ~~at the interface~~ between cloudy and clear ~~areas~~ model grid boxes, where the gradient of cloud optical properties is large. Strong horizontal variability of in-cloud ~~optical properties might and subgrid cloud optical properties can~~ also lead to ~~noticeable horizontal radiative exchanges that are~~ horizontal radiative transfer that is neglected in 1D radiative transfer schemes. ~~Since we can assume that the clouds from the large eddy model simulations are perfectly known and no further subgrid cloud variability exists, 3D cloud radiative effects here are only attributed to the horizontal gradient of radiative fluxes between model columns and not within model grid boxes. For model grids with coarser horizontal resolution, the horizontal radiative exchange caused by the horizontal subgrid variability of cloud optical properties needs to be parameterized as part of the 3D cloud radiative effects.~~

– Ice-optical parameterization:

The representation of ice-optical properties in models is challenging due to the complexity of ice ~~particles and insufficient observations. This~~ crystals, especially with assumptions regarding their shape and surface roughness. ~~The lack of a consolidated understanding of the ice crystal shapes and how they should be represented in models~~ creates an important source of uncertainty for simulating CRH (e.g., Zhao et al., 2018; Yi, 2022). Several parameterizations for the ice-optical properties have been developed based on different assumptions about ice crystal shapes or habits, size distributions, and surface roughness (Yang et al., 2013; Baran et al., 2014; Baum et al., 2014). It is therefore important to examine how the assumed ice-optical properties affect CRH.

– Cloud horizontal heterogeneity and vertical overlap ~~assumption~~:

Clouds are not homogeneous over their horizontal and vertical extent, and this spatial variability significantly affects their interaction with radiation. Cloud horizontal heterogeneity refers to the horizontal variation of cloud optical properties within a grid box. The vertical overlap determines how cloud layers are stacked on top of each other. The ability of models to account for cloud horizontal heterogeneity and vertical overlap depends primarily on their horizontal resolution. ~~In large eddy model simulations, we can assume that there are no further subgrid cloud variabilities, and clouds are entirely characterized by the grid-box values. Thus, we quantify these sources of uncertainty at the convective-permitting resolution of 2.5 km for the numerical weather prediction model.~~ Neglecting cloud horizontal heterogeneity increases longwave emissivity and shortwave absorption of clouds, an effect known as the plane-parallel problem of radiative transfer calculations (e.g., Črnivec and Mayer, 2019). In current models with resolutions on the order of 1-10 km, parameterizations are used to represent both cloud horizontal heterogeneity within grid boxes and cloud vertical overlap between adjacent grid boxes. As a result, assumptions in the parameterizations of these two subgrid effects can lead to errors in the simulation of CRH (e.g., Wang et al., 2021).

90 We organize the manuscript as follows. The model simulation setup and radiative transfer calculations are described in Sect. 2. In Sect. 3 we quantify the CRH uncertainty due to the four factors discussed above, and we compare their importance in Sect. 4. The manuscript closes with a discussion and conclusion in Sect. 5.

## 2 Methods

This section describes the model setup for the idealized baroclinic life cycle simulation, the large eddy model simulations, 95 and the offline radiative transfer calculations that we use in combination to investigate the uncertainty in CRH within an extratropical cyclone.

### 2.1 Baroclinic life cycle simulation

We use the ICOSahedral Nonhydrostatic atmosphere model (ICON, version 2.6.2.2; Zängl et al. (2015)) to simulate a baroclinic life cycle. The simulation provides us with an idealized extratropical cyclone for which we can test the uncertainty in CRH. We 100 use the same model setup developed and described by Keshtgar et al. (2023) and only briefly describe the model setup here. The planar channel grid extends 4000 km and 9000 km in the zonal and meridional directions, respectively. The horizontal resolution is 2.5 km. Periodic boundary conditions are applied in the zonal direction, and the Coriolis parameter is set to a uniform value valid at 45°N. ~~In the vertical direction, ICON applies a terrain-following hybrid vertical coordinate, and we use~~ 75 model levels ~~are used.~~ The in the vertical direction. The layer's thickness increases from 20 m near the surface to 400 m at 105 10 km. Above 10 km, the layer thickness increases from 400 m to 1200 m at 30 km. The initial and lateral boundary conditions follow the life cycle type 1 of Polvani and Esler (2007). The model surface is configured as an ice-free ocean with a prescribed sea surface temperature that is 0.5 K lower than the initial temperature of the lowest model level. To trigger the development of the cyclone, we add a temperature perturbation with a wavelength equal to the domain length and an amplitude of 1 K at all levels. We run the ICON model with the numerical weather prediction (NWP) physics package, including the shallow 110 convection scheme of Bechtold et al. (2008), the two-moment microphysics of Seifert and Beheng (2006), and the turbulence scheme of Raschendorfer (2001). Due to the high horizontal resolution of the model, deep convection is treated explicitly.

We use the modeling technique developed by Keshtgar et al. (2023) to simulate the cyclone with only the contribution of CRH; the clear-sky radiative heating is not used for stepping the temperature forward in time. The CRH in the model is given by

$$115 \quad \left. \frac{\partial T}{\partial t} \right|_{\text{radiation}}^{\text{cloud}} = \left. \frac{\partial T}{\partial t} \right|_{\text{radiation}}^{\text{all-sky}} - \left. \frac{\partial T}{\partial t} \right|_{\text{radiation}}^{\text{clear-sky}} = \frac{1}{\rho c_v} \cdot \frac{\partial}{\partial z} (F^{\text{all-sky}} - F^{\text{clear-sky}}), \quad (1)$$


where  $\rho$  is the air density and  $F$  is the radiative flux in all-sky and clear-sky conditions defined as positive downward. Since ICON is based on height levels, the conversion of radiative fluxes to heating rate requires the heat capacity of air at constant volume,  $c_v$ . As described in Keshtgar et al. (2023), the technique eliminates the impact of strong clear-sky radiative cooling early in the simulation.

120 Different from Keshtgar et al. (2023), we use ecRad (Hogan and Bozzo, 2018), the new operational radiation scheme in ICON (Rieger et al., 2019) instead of the Rapid Radiative Transfer Model (Mlawer et al., 1997). The advantage of ecRad is that we can configure the offline radiative transfer setup in a manner that is very close to the radiative transfer setup used in ICON (Sect. 2.3). From the various radiation solvers available in ecRad, we use the homogeneous solver. The solver homogenizes the cloud properties over the entire grid box, i.e., it ignores the cloud fraction and does not require any assumption about cloud vertical overlap. Thus, for the purpose of radiative transfer calculation, clouds are treated as grid-box clouds. This treatment is the same as in the large eddy model simulations described later. For the cloud optical parameterization, we use the scheme from the Suite Of Community RAdiative Transfer codes (SOCRATES) based on Edwards and Slingo (1996) for liquid water droplets. For ice crystals, we use the scheme of Fu (Fu, 1996; Fu et al., 1998).

Overall, this model configuration results in a typical wintertime extratropical cyclone (Fig. 1 a). Note that for the latitude-longitude plots, we use the geographic coordinates assigned during the generation of the planar channel grid, but the model simulation is based on the Cartesian coordinate of the grid. The baroclinic simulation provides the initial and lateral boundary conditions for the large eddy model simulations described in the following section.

## 2.2 Large eddy model simulations

We target four different regions of the cyclone by the large eddy model (LEM) simulations, shown by black rectangles in Fig. 1. To characterize the 3D structure of clouds, panel b of Fig. 1 shows cloud classes. We derive the cloud classes from the clouds' vertical extent (Sullivan et al., 2023) using a 50% threshold for low-, mid-, and high-level cloud cover that are diagnosed within ICON during the model simulation. For example, if cloud cover exceeds the 50% threshold at low, middle, and high altitudes, the cloud class is High-Middle-Low.



**Figure 1.** (a) Surface pressure (black contours, hPa), cloud cover, and precipitation at day 4.5 in the baroclinic life cycle simulation with ICON-NWP. Panel (b) shows cloud classes. The rectangles in both panels indicate the location of the domains for the LEM simulations.

Domain 1 targets low-level shallow cumulus and stratocumulus clouds southwest of the cyclone center within the cold  
140 sector. These clouds are formed by the passage of cold, dry air over the warm ocean behind the cold front. Domain 2 targets the  
extensive vertical clouds within the warm conveyor belt (WCB) of the cyclone, where strong latent heating and precipitation  
occur. Domain 3 captures mid- and low-level clouds in the cyclonic branch of the WCB outflow, and Domain 4 captures mostly  
mid- and high-level clouds in the anticyclonic branch of the WCB outflow. We will refer to domains 1, 2, 3 and 4 as shallow  
cumulus, WCB ascent, WCB cyclonic outflow and WCB anticyclonic outflow, respectively.

145 For the LEM simulations, we use the ICON model in a limited area setup. The LEM simulations use a planar grid with  
perfectly uniform triangular grid-cell areas. Unlike the planar channel grid used in the NWP simulation, the planar grid does  
not apply zonal periodic and fixed meridional boundary conditions. Instead, hourly lateral boundary conditions are provided  
by the baroclinic life cycle simulation. The planar grid extends 471 km and 667 km in the zonal and meridional directions,  
respectively. This is approximately equivalent to a  $6^\circ \times 6^\circ$  domain at  $45^\circ\text{N}$ . The horizontal resolution is 300 m. We use 150  
150 model levels with layer thicknesses increasing from 20 m near the surface to 570 m at 30 km.

As in the NWP simulation, we run the LEM simulations with the two-moment microphysics scheme, the homogeneous  
radiation solver of ecRad, the same cloud optical parameterization, and with only the CRH contribution from the radiation  
scheme. In contrast to the NWP simulation, we use the 3D Smagorinsky diffusion model (Smagorinsky, 1963) for the turbu-  
lence scheme and switch off the shallow convection scheme. We configure the cloud cover scheme to treat clouds as grid-scale  
155 quantities, i.e. grid boxes are either fully cloudy or clear.


We run LEM simulations for 12 hours starting on day 4.25 of the baroclinic life cycle simulation when the cyclone is in its  
main growth phase. We exclude the first 3 hours to remove spin-up effects. Keshtgar et al. (2023) showed that between days 4  
and 5, the modulation of latent heating by the CRH is strong enough to affect the dynamics of the cyclone.

The offline radiative transfer calculations described in Sect. 2.3 require as input the 3D fields of cloud ice and liquid water  
160 content and effective particle radii simulated by the large eddy model. We briefly characterize the clouds from the LEM  
simulations in Fig. 2, which shows domain and time averages of cloud ice and liquid water contents and cloud fractions for the  
four LEM domains. We calculate the time average over nine snapshots that are each separated by a 30-minute time interval  
between ~~hours 10 and 14~~ domain local hours 12:30 and 16:30. We use the same snapshots for the offline radiative transfer  
calculation. During this time interval, the solar zenith angle changes from low to high values in all regions, which is an  
165 important factor for the shortwave 3D cloud radiative effects (Sect. 3.2).

In the shallow cumulus region, low-level liquid clouds prevail in the boundary layer between 0-2 km, with cloud fraction  
peaking at 40% at an altitude of 1.5 around 1 km (Fig. 2 a, b). In the WCB ascent region, extensive vertical clouds contain both  
ice and liquid water. The liquid water content in the lower and middle levels is much higher and more variable in time than the  
ice water content in the upper levels (Fig. 2 c). The decomposition of cloud fraction shows that liquid clouds are concentrated in  
170 the lower levels between 0-4 km, where their cloud fraction reaches a maximum of 60% at around 1 km. Mixed-phase clouds  
are located at intermediate levels between 2-7 km. Above the boundary layer, the fraction of ice clouds increases rapidly,  
reaching a maximum of nearly 60% at about 8 km (Fig. 2 d).

Clouds in the WCB cyclonic outflow region also contain both ice and liquid water but are located at lower altitudes compared to the WCB ascent region (Fig. 2 e). The total cloud fraction is dominated by ice clouds and reaches nearly 90% at around 6 km. Mixed-phase clouds exist only in the boundary layer (Fig. 2 f). In the WCB anticyclonic outflow, mid- and high-level clouds contain both ice and liquid water, with ice water content dominating over liquid water content (Fig. 2 g). The total cloud fraction is dominated by ice clouds (Fig. 2 h).

The WCB ascent and outflow regions are much cloudier than the shallow cumulus region. In the WCB outflow regions, stratiform ice clouds cover almost the entire domain.



**Figure 2.** Profiles of domain-averaged cloud hydrometeor contents and total cloud fraction, decomposed into the contributions from liquid, ice, and mixed-phase clouds for all four LEM domains. Thin lines show profiles for nine snapshots between domain local hours 12:30 to 16:30. The thick lines show time-averaged profiles. The threshold used to determine cloudy boxes for both ice and liquid water contents is  $10^{-8}$  kg kg<sup>-1</sup> (Costa-Surós et al., 2020). Note the different y-axes for panels (a) and (b).

### 180 2.3 Offline radiative transfer calculations

For the offline radiative transfer calculations, we use the [LibRadtran-libRadtran](#) software (Mayer and Kylling, 2005; Emde et al., 2016), which includes several radiation solvers and different cloud optical parameterizations. In order to achieve the best possible match between the offline calculations and the CRH calculated with ICON, we configure the radiative transfer setup of [LibRadtran-libRadtran](#) as similar as possible to ICON.

185 For gas absorption, we use the parameterization of Fu (Fu and Liou, 1992, 1993), which employs the correlated-k method with 6 and 12 spectral bands in the solar and thermal spectrums, respectively. This parameterization is optimized for use in climate models. The optical properties of cloud droplets are prescribed according to the parameterization of Hu and Stamnes (1993), which assumes spherical droplets. The optical properties of ice crystals are prescribed according to the parameterization of Fu (Fu, 1996; Fu et al., 1998), which is the same scheme used in the ICON simulations. The scheme of Fu uses an  
190 approximate phase function and assumes a pristine hexagonal column habit for the ice crystals. For the radiative transfer solver, we use the classical Delta-Eddington two-stream solver (Zdunkowski, 2007), which is suitable for horizontally homogeneous clouds. This solver is similar to the ecRad homogeneous solver used in the ICON simulations.

In [LibRadtran](#)~~libRadtran~~, inputs are required to have normal Cartesian rectangular coordinates. Therefore, we remap the LEM simulation output from the unstructured triangular grid to a regular Cartesian grid with the same resolution using nearest  
195 neighbor interpolation. For the clear-sky background, we use pressure, temperature, density, specific humidity, and trace gas concentration from the LEM simulations. We set other inputs such as surface type, albedo, and emissivity to the same values used in the ICON simulations. We use the solar zenith angles used in the LEM simulations for the offline solar radiative transfer calculations.

The offline radiative transfer calculations are very time-consuming, especially for the 3D Monte Carlo radiative transfer  
200 calculation (Mayer, B., 2009). Ideally, one would run the calculations over the entire LEM domain using the 3D cloud and clear-sky fields. However, this is not feasible because [LibRadtran](#)~~libRadtran~~ cannot be run in parallel and the wall-clock time of the computing nodes of the Levante DKRZ supercomputer that we are using for our study is limited to 8 hours. To overcome the computational challenge, we divide the LEM domains into  $36\ 1^\circ \times 1^\circ$  subdomains for which we run the radiative transfer calculations individually. For each subdomain, we use a 1D clear-sky background profile that we derive by horizontally  
205 averaging the clear-sky fields. After the radiative transfer calculations for all subdomains are finished, we merge the results to obtain the radiative heating rate over the entire LEM domain. While this means that we neglect horizontal variability of the clear-sky background within the subdomains, we show in Sect. 3.1 that this has a negligible impact and we are able to achieve a very good agreement between the CRH calculated offline with [LibRadtran](#)~~libRadtran~~ in the reference setup using the 1D radiation calculation and online in ICON-LEM.

210 The partitioning of the LEM domains into subdomains allows us to perform expensive 3D Monte Carlo radiative transfer calculation, but can introduce errors in the radiative transfer calculation due to the discontinuity of the horizontal photon [transport](#)~~propagation~~ at the lateral boundaries of the subdomains. As an example, consider a shortwave radiative transfer calculation with a cumulus cloud near the northern boundary of a subdomain. The shadow of this cloud should be present at the southern boundary of the subdomain to the north of it, yet the periodic boundary conditions applied in the 3D solver mean  
215 that the shadow is placed at the southern boundary of the cloud's own subdomain. The gravity of such an error depends on the solar zenith angle, cloud top height, and cloud cover. We minimize this issue by making the subdomains overlap in all directions by approximately 12 km. The overlap is sufficient for shallow cumulus clouds with a maximum height of 2 km. The overlap is too small for clouds at higher altitudes in the WCB, but since WCB clouds are mostly stratiform the resulting errors tend to be small and do not affect the generality of our results.



220 To investigate the uncertainty in CRH, we perform several radiative transfer calculations that are listed in Table. 1. For  
3D cloud radiative effects, we use the "Monte Carlo code for physically correct tracing of photons in cloudy atmospheres"  
(MYSTIC, Mayer, B. (2009)). MYSTIC can also be run in the independent column approximation mode (MYSTIC-ICA),  
which neglects horizontal photon transport between model columns and is equivalent to a 1D radiation scheme. We use the  
standard forward photon tracing method. The azimuth angle is set to a constant value of ~~zero~~180 degrees, which directs the  
225 solar radiation from south to north. To ~~obtain low~~reduce the Monte Carlo noise, we run MYSTIC ~~with and~~MYSTIC-ICA with  
72 million photons for each subdomain in the LEM domains at each time step and repeat the calculations 10 times, resulting in  
a total of 720 million photons traced per subdomain (nearly 5000 photons per LEM column)~~for each subdomain in the LEM~~  
~~domains.~~We then average over these 10 calculations to derive the radiative heating in each LEM domain. Overall, the total  
computational time required to perform the entire set of 3D radiative transfer calculations amounts to about 1500 hours on a  
230 single node of the Levante DKRZ supercomputer.

To estimate the Monte Carlo noise of the MYSTIC solver in our setup, we split 10 MYSTIC calculations for the shallow  
cumulus domain at a single time step into two sets of 5 calculations and average the heating rates over these two sets of 5  
calculations. We then calculate the relative standard deviation of the radiative heating between these two sets at each grid box.  
The relative standard deviation represents the relative variability of the radiative heating to the mean values calculated from  
235 the two sets of MYSTIC calculations. The median relative standard deviations in the shortwave, longwave, and net are less  
than 10 percent for almost all grid boxes (not shown). This estimate is an upper bound, as the true Monte Carlo noise in our  
calculations with 720 million photons can be expected to be smaller. The low Monte Carlo noise of our radiation calculation  
allows us to calculate the CRH uncertainty due to 3D cloud radiative effects locally at the scale of the horizontal grid resolution  
of the LEM domains (Sect. 4).


240 To assess the impact of the ice-optical parameterization, we use the more complex scheme of Baum-v36 (Yang et al., 2013;  
Baum et al., 2014) in addition to the scheme of Fu. The scheme of Baum includes the full phase function and three different  
ice habits: general habit mixture, solid column, and rough-aggregated. Ice crystal surfaces are treated as severely roughened in  
the Baum scheme. Zhao et al. (2018) compared the radiative fluxes calculated with the schemes of Fu and Baum and showed  
that in the longwave spectrum, the mass absorption coefficient is smaller in the Baum scheme than in Fu, resulting in weaker  
245 longwave radiative cooling at the cloud top and weaker warming below (Fig.1 of Zhao et al. (2018)). In the shortwave spectrum,  
the backward scattering of radiation by ice crystals is higher in the Baum scheme due to the reduced parameterized asymmetry  
factor and forward peak. This reduces the radiative flux gradient between cloudy and clear sky layers, resulting in a weaker  
CRH with the ice scheme of Baum.

To study the impact of cloud horizontal heterogeneity on CRH, it would be misleading to directly compare the CRH between  
250 the NWP and LEM setups since the two setups simulate different clouds. Therefore, we instead create homogeneous clouds  
at the 2.5 km NWP resolution by coarse-graining the LEM clouds. The method is illustrated in Fig. 3. We derive two types  
of homogeneous NWP clouds. For the first type, we average the cloud fields over all LEM grid boxes that fall within the  
corresponding NWP grid box with a resolution of 2.5 km. We call this type "homogeneous grid-box clouds". The type assumes  
that clouds are resolved at the NWP resolution so that grid boxes are either entirely cloudy or clear and no vertical overlap

**Table 1.** List of radiative transfer calculations to quantify the uncertainty in CRH. The first column shows the purpose of the calculations, the second column the radiation solvers, and the third column the ice-optical parameterization. The last column lists the sections in which the calculations are discussed.

Purpose	Radiation solvers	Ice-optical parameterization	Section
Reference	$\delta$ -Eddington two-stream	Fu	3.1, 3.3
3D cloud radiative effects	MYSTIC	Fu	3.2
	MYSTIC-ICA	Fu	3.2
Ice-optical parameterization	$\delta$ -Eddington two-stream	Baum-v36, general habit mixture	3.3
	$\delta$ -Eddington two-stream	Baum-v36, solid column	3.3
	$\delta$ -Eddington two-stream	Baum-v36, rough-aggregated	3.3
Cloud horizontal heterogeneity and overlap at 2.5 km NWP resolution (Fig. 3)	$\delta$ -Eddington two-stream	Fu	3.4
	$\delta$ -Eddington two-stream with maximum-random vertical overlap	Fu	3.4

255 assumption is needed. For the second type, we average the cloud fields only over those LEM boxes that are cloudy, and we use the number of clear-sky boxes to determine the cloud fraction. We refer to this type as "homogeneous clouds with fractional cloud cover". The type requires an overlap assumption, for which we use the Delta-Eddington two-stream solver that applies maximum-random overlap (Črnivec and Mayer, 2019).



**Figure 3.** Illustration of the method to derive homogeneous NWP clouds from LEM clouds. The plot shows a layer of grid boxes. (a) Clouds in the LEM simulation. (b) Homogeneous grid-box cloud at a resolution of 2.5 km. (c) Homogeneous cloud with fractional cloud cover at a resolution of 2.5 km.


### 3 Results

260 In this section, we assess CRH in the four regions of the cyclone. We derive CRH from all-sky and clear-sky radiative heating in the shortwave and longwave spectrum for each of the radiative transfer settings listed in Table. 1. To quantify the impact of the four factors described in Sects. 1 and 2, we compare the time and domain averages of the CRH from the different radiative transfer calculations.

#### 3.1 Average profiles of CRH

265 Fig. 4 shows domain and time averages of longwave, shortwave, and net CRH in the four regions of the cyclone for the ICON and the reference offline radiative transfer calculation. Despite the differences in the radiative transfer setup between ICON and [LibRadtran](#), the CRH profiles agree very well (cf. dashed and solid lines in Fig. 4). In all four regions, longwave radiation leads to cooling at the cloud tops and weak warming at the cloud bases (Fig. 4, blue), resulting in radiative destabilization of the cloud. In contrast, shortwave radiation warms the cloud tops and slightly cools the lower parts of the clouds, resulting in radiative stabilization of the cloud (Fig. 4, red). The longwave CRH is about twice as large as the shortwave CRH. Thus, the net CRH profiles are dominated by the stronger longwave CRH (Fig. 4, black).

270



**Figure 4.** Domain and time averages of shortwave, longwave, and net CRH. Profiles are shown for the (a) shallow cumulus, (b) WCB ascent, (c) WCB cyclonic outflow, and (d) WCB anticyclonic outflow regions. The ICON CRH is shown in dashed lines, and the CRH derived from the offline reference 1D radiative transfer calculation is shown in solid lines. Note the different x- and y-axes between panels (a) and (b-d).


The CRH in the shallow cumulus region is almost twice as strong as in the WCB regions (Fig. 4 a). Grise et al. (2019) and Voigt et al. (2023) showed that the strong cloud-radiative cooling of the boundary layer has a weakening effect on the intensity of extratropical cyclones. The CRH in WCB regions is strongly influenced by the radiative heating of the ice clouds (Fig. 4 b, c and d). Although the CRH in the WCB regions is weaker than the CRH in shallow cumulus region, previous work showed it to be important for near-tropopause dynamics of the cyclone (Li et al., 2015; Keshtgar et al., 2023; Voigt et al., 2023). Thus, it is important to quantify CRH uncertainties in all four regions of the extratropical cyclone.

275

### 3.2 3D cloud radiative effects

We quantify 3D cloud radiative effects as the difference in CRH calculated with the MYSTIC solver (3D, Sect. 2.3) and the  
 280 MYSTIC solver run in the independent column approximation mode (1D). The latter is a 1D radiation scheme and neglects  
 horizontal ~~radiative transfer~~ photon transport between model columns. To visualize the 3D cloud radiative effects, Fig. 5 shows  
 cross-sections of 3D and 1D all-sky radiative heating and their differences for shallow cumulus clouds southwest of the cyclone  
 center at ~~hour 14~~ domain local hour 16:30. At this location and time, the solar zenith angle is approximately  $65^\circ$ .

In the shortwave spectrum, neglecting horizontal ~~radiative transfer~~ photon transport leads to incorrect positions of illuminated  
 285 and shadowed areas in the atmosphere and on the surface (Fig. 5 a, b and c). Črnivec and Mayer (2019) showed that this can  
 have a substantial impact on surface radiative fluxes and hence the surface energy balance. ~~Our focus here, however,~~ However,  
~~our focus~~ our focus is on the 3D radiative effects of clouds within the atmosphere. In Fig. 5 a, the southern sides of the clouds (cf. the  
shallow cumulus domain in Fig. 1) facing the sun receive more radiative energy in the 3D calculation than the 1D calculation  
 (red colors around 1.5 km altitude in Fig. 5 c). In some cases, the northern sides of the clouds are shadowed and receive less  
 290 energy (blue colors around 1.5 km altitude in Fig. 5 c). The shortwave cloud side leakage is small in this example due to the  
 low position of the sun.




**Figure 5.** Cross-sections of shortwave, longwave and net all-sky radiative heating visualized by a logarithmic color scale for shallow cumulus clouds southwest of the cyclone center. The upper row shows 3D calculations, the middle row shows 1D calculations, and the lower row shows the differences between the 3D and 1D calculations. The cross sections are at domain local hour 16:30 and  $-11.5^\circ$  longitude and between  $37.5^\circ$  and  $38^\circ$  north. The solar zenith angle is  $65^\circ$ . Note that the impression of a lower solar zenith angle in the figure is due to the aspect ratio of the figures.

In the longwave spectrum, radiative cooling from the cloud tops and modest warming from the bottom are clearly visible in both the 3D and 1D calculations (cloud "shadows", visible as white areas below clouds (panels d and e of Fig. 5 d, e)), are weaker in the 3D calculation than in the 1D calculation due to the horizontal photon transport between model columns. However, the radiative cooling is stronger. Largest differences between the 3D and 1D calculations occur at the cloud-clear sky boundaries, where horizontal emission of longwave radiation from the cloud tops and cloud sides leads to stronger radiative cooling in the 3D calculation (blue colors around 1.5 km altitude height in Fig. 5 f).

In the net, most features of cloud radiative heating and cooling within the atmosphere are present in both 3D and 1D calculations (Fig. 5 g, and h). However, due to the emission of longwave radiation from the cloud sides, which is neglected in the 1D calculations, shortwave cloud-side illumination and horizontal longwave cloud absorption and emission, large differences exist at the interface of clouds and clear sky regions around 1.5 km height in Fig. 5 i and in the position of cloud shadows.

Fig. 6 shows the average profiles of 3D cloud radiative effects for the four regions of the cyclone. In all regions, shortwave and longwave CRH are stronger in the 3D radiation calculation due to shortwave cloud-side illumination and longwave cloud-side cooling, respectively. There is a direct relationship between cloud-side illumination and solar zenith angle (Fig. 6 a, thin lines). In the 3D calculation, clouds receive more radiative energy from their sides at higher solar zenith angles, which increases shortwave cloud radiative warming. Črnivec and Mayer (2019) also showed this direct relationship between cloud-side illumination and solar zenith angle. The magnitude of the cloud-side cooling depends on the magnitude of the longwave CRH, such that the stronger the longwave CRH, the stronger the longwave cooling from the cloud sides.

The 3D cloud radiative effects are much weaker for WCB regions which are dominated by stratiform clouds, than the shallow cumulus region, which contains a lot of broken clouds (Fig. 6). This is expected due to the weak horizontal gradients of cloud optical properties and the small aspect ratio of the stratiform clouds in the WCB regions. For shallow cumulus clouds, the net 3D cloud radiative effect is dominated by cloud-side longwave cooling (Fig. 6 a). This could lead to a stronger radiative destabilization of clouds during both day and night. In contrast, for stratiform clouds in the WCB, shortwave cloud-side illumination dominates the daytime 3D cloud radiative effects, except for shallow liquid clouds in the boundary layer of the WCB cyclonic outflow region, where longwave cloud-side cooling becomes dominant (Fig. 6 b, c and d). The stronger shortwave cloud-side illumination in the WCB regions is most likely due to the higher solar zenith angle at higher latitudes compared to the shallow cumulus region at lower latitudes. Although the incoming shortwave radiative fluxes are weaker at higher latitudes, shortwave cloud-side warming becomes stronger at higher zenith angles.




**Figure 6.** Average profiles of shortwave, longwave, and net 3D cloud radiative effects for the four cyclone regions. 3D cloud radiative effects are calculated as the difference in domain mean CRH between 3D (MYSTIC) and 1D (MYSTIC-ICA) radiative transfer calculations. The thin lines in panel (a) show the 3D radiative effects for nine snapshots between [domain local](#) hours ~~10~~12:30 to ~~14~~16:30, with the legend indicating the time and domain mean solar zenith angles for all snapshots. Note the different x-axes in panels.

### 3.3 Impact of ice-optical parametrization

320 We now quantify the impact of ice-optical parameterization by comparing the CRH calculated with the ice-optical parameterization of Fu and the more complex ice-optical parameterization of Baum. The ice scheme of Baum includes three ice habits, which allows us to also assess the effect of different ice habits. We only consider clouds in the three WCB domains, where ice crystals are abundant.


325 The top row in Fig. 7 shows cross-sections of shortwave, longwave, and net CRH in the WCB ascent region for the ice scheme of Fu. The bottom row in Fig. 7 shows the CRH differences between the ice schemes of Fu and Baum with the general habit mixture. Fu results in stronger longwave and shortwave CRH than Baum (Fig. 7 d, e), with CRH differences reaching up to 20% of the absolute values. As a result, Fu leads to stronger radiative destabilization in the longwave but stronger stabilization in the shortwave.



**Figure 7.** Cross-sections of (a) shortwave, (b) longwave, and (c) net CRH calculated with the ice-optical parameterization of Fu in the WCB ascent region. Panels (d-f) show CRH differences between the ice schemes of Fu and the ice scheme of Baum with the general habit mixture (Fu - Baum). Cross-sections are shown at domain local hour 14:30 and at 3° longitude.

Fig. 8 shows CRH differences between Fu and Baum. In all three WCB regions, longwave and shortwave CRH are amplified  
 330 in Fu compared to Baum. The magnitude of the CRH differences depends on which ice crystal habit is chosen in the Baum  
 scheme, with different impacts in the longwave and shortwave. In the longwave, the differences between Fu and Baum are the  
 same for the general mixture and rough-aggregated habits, but smaller by a factor of two when the solid column habit is used  
 in Baum. In the shortwave, the impact of the ice habit is much smaller. The stronger impact of the ice habit on the longwave  
 compared to the shortwave CRH is an interesting aspect of our results.

335 Compared to Baum, Fu leads to stronger shortwave as well as longwave cloud-radiative warming in the middle and lower  
 parts of the clouds, and therefore to net cloud warming (Fig. 8, lower row). Although Fu leads to some increase in cloud-  
 radiative cooling of the upper cloud parts, this suggests that the radiative destabilization of clouds within the WCB regions is  
 smaller in Fu than Baum.



**Figure 8.** Impact of ice-optical parameterization on time- and domain-averaged CRH in the three WCB regions. The CRH differences are calculated between the radiative transfer calculations with the ice schemes of Fu and Baum (Fu - Baum). The differences between Fu and Baum with general habit mixture (ghm), solid column (sc), and rough-aggregated (ra) habits are shown with solid lines, dashed lines, and dotted lines, as is indicated in the legend.

### 3.4 Cloud horizontal heterogeneity and vertical overlap

340 We continue by quantifying the impact of cloud horizontal heterogeneity and vertical overlap. Both factors are not resolved at 2.5 km and therefore need to be parameterized. We calculate the CRH differences between the radiation calculation using the two types of homogeneous NWP clouds and the LEM clouds (Sect. 2.3, Fig. 3). For homogeneous grid-box clouds, an overlap assumption is not needed. For the homogeneous clouds with fractional cloud cover, we use a maximum-random overlap assumption.


345 The solid lines in the first row of Fig. 9 show CRH differences between homogeneous grid-box clouds and the LEM clouds, decomposed into longwave and shortwave components. In all regions, the magnitude of the longwave and shortwave CRH profiles are overestimated when homogeneous grid-box clouds are used, with the longwave CRH being more strongly affected than the shortwave CRH and the largest differences occurring for shallow cumulus clouds.

The second row of Fig. 9 compares the impact of the two types of homogeneous clouds. When the overlap scheme is used for the homogeneous clouds with fractional cloud cover, the differences in CRH between the NWP and LEM clouds are strongly reduced in the shallow cumulus region, from  $-2 \text{ K day}^{-1}$  to  $0.5 \text{ K day}^{-1}$ . This illustrates a clear positive impact of taking



into account cloud fraction and partial overlap for these clouds, and that treating these clouds as grid-box clouds at 2.5 km resolution is problematic.

In the three WCB regions, cloud vertical overlap assumption has a smaller impact on CRH because the cloud fraction is higher. However, in contrast to the shallow cumulus region, taking into account cloud overlap assumption has a detrimental effect and in fact increases the CRH differences for shortwave, longwave and net (Fig. 9 c to h). This supports the idea that clouds in the WCB regions can be treated as grid-box clouds at 2.5 km resolution for radiative purposes because there is no obvious benefit by taking into account cloud fraction and vertical overlap assumption.



**Figure 9.** Impact of cloud horizontal heterogeneity and vertical overlap on CRH that are not resolved at 2.5 km horizontal resolution. The CRH differences are calculated between the LEM reference calculation and the radiative transfer calculations for the homogeneous NWP clouds (NWP - LEM), with the solid lines for the homogeneous grid-box clouds (without overlap assumption) and the dotted lines for the homogeneous clouds with fractional cloud cover (with overlap assumption). Note the different x and y-axes for panels (a) and (b).

#### 4 Relative importance of uncertainties

In the previous [sections](#), we have assessed the individual impacts of the four factors on CRH: 3D cloud radiative effects, ice-optical parametrization, cloud horizontal heterogeneity and cloud vertical overlap. In this section, we characterize the relative importance of these four factors [for extratropical cyclones](#), with the hope that this can guide efforts to improve CRH in models. As part of this characterization, we distinguish between mean uncertainties on spatial scales of around 500

km, which is approximately the size of the LEM domain, and local uncertainties, which arise on the scale of the horizontal  
 365 grid resolutions ~~from~~. For the uncertainties due to 3D cloud radiative effects and ice-optical parameterization, the scale of  
 the horizontal grid resolution is 300 m to, and for the uncertainties due to cloud horizontal heterogeneity and cloud vertical  
overlap, the scale of the horizontal grid resolution is 2.5 km.

For the mean uncertainties, we compute the *absolute mean* difference,

$$\delta CRH_{mean} = |\overline{CRH}_a - \overline{CRH}_b|, \quad (2)$$

370 where the bars represent the average over time and domain and the subscripts  $a$  and  $b$  indicate different radiative transfer  
 calculations. This means that the mean uncertainties are obtained from the absolute values of the figures shown in Sect. 3.

For the local uncertainties, we compute the *mean absolute* difference by first computing the absolute CRH differences at all  
 grid points and then averaging over time and domain,

$$\delta CRH_{local} = \frac{1}{n \cdot t} \sum_{j=1}^t \sum_{i=1}^n |CRH_a(i, j) - CRH_b(i, j)|. \quad (3)$$


375  $n$  is the number of horizontal grid points  $i$  at each vertical layer, and  $t$  is the number of time steps  $j$ . ~~We do not compute  
 the local uncertainty due to 3D cloud radiative effects. The latter would require to first quantify the Monte Carlo noise of the  
 MYSTIC solver, which if substantial can on its own create strong local uncertainties.~~ To derive the local uncertainties due to  
 horizontal cloud heterogeneity and vertical overlap, we coarse-grain the LEM CRH to the same horizontal resolution as the  
 NWP CRH.

380 Fig. 10 shows mean uncertainties by means of stacked bar plots in altitude intervals of 2 km calculated based on Eq.2. For  
 shallow cumulus clouds in the boundary layer between 0-2 km, the largest source of mean uncertainty is cloud horizontal  
 heterogeneity without overlap assumption (panels a, b and c; blue bars). Allowing for partial overlap for homogeneous clouds  
 with cloud fraction significantly reduces the mean uncertainty (cf. orange-green and blue bars in panels a, b and c). The mean  
 uncertainty due to 3D cloud radiative effects is about half of the uncertainty due to cloud horizontal heterogeneity when vertical  
 385 overlap is considered.

For clouds in the WCB ascent region, cloud horizontal heterogeneity has a significant impact on the mean uncertainty at all  
 levels, and dominates the CRH uncertainty in the boundary layer (panels d, e and f; blue bars). Apart from the boundary layer  
 and in contrast to shallow cumulus clouds, considering vertical overlap assumption increases the mean uncertainty (cf., orange  
 green and blue bars in panels d, e and f). Ice-optical parameterization dominates the mean uncertainty of net CRH at middle  
 390 and upper levels above 6 km (panel f). 3D cloud radiative effects contribute relatively little, except at altitudes between 2-6 km  
 in the shortwave spectrum and the net.

The relative importance of the four factors for the mean uncertainties in the WCB cyclonic outflow region is similar to that  
 in the WCB ascent region (panels g, h and i). The ice-optical parameterization contributes most significantly to the mean un-  
 certainty in the middle and upper levels, while cloud horizontal heterogeneity dominates the mean uncertainty in the boundary  
 395 layer. The vertical overlap assumption increases the mean uncertainty, and the impact of the 3D cloud radiative effects is small.  
 The mean uncertainty in the WCB anticyclonic outflow region is dominated by the ice-optical parameterization, and the impact


of cloud horizontal heterogeneity and vertical overlap is much weaker compared to the other regions of the cyclone, except between 4-6 km (panels j, k and l).



**Figure 10.** Mean uncertainties in CRH diagnosed as the absolute difference of domain and time mean CRH between different radiative transfer calculations. Uncertainties are decomposed into shortwave, longwave, and net. Uncertainties are computed as mass-weighted averages over 2 km altitude intervals. For the uncertainty due to the ice-optical parameterization, the difference between the ice schemes of Fu and Baum with the general habit mixture is used. The contribution of each factor is given by the horizontal length of its colored bar. Note the different x-axes in the panels.

Now we turn to local uncertainties in CRH. These are illustrated by means of Fig.11 for net CRH. **In all four regions of the cyclone** For comparison of mean and local uncertainties, Fig.11 includes the mean net CRH uncertainties from Fig. 10 only for values larger than  $0.05 \text{ K day}^{-1}$  as gray hatched bars. In contrast to the mean uncertainties, the impact of 3D cloud radiative effects is much stronger at the scale of the horizontal grid resolution of 300 m. Except for the WCB anticyclonic domain, cloud horizontal heterogeneity dominates local uncertainties at the boundary layer between 0-2 km and mid-levels between 2-8 km in all regions of the cyclone. As for the mean uncertainties, taking into account the vertical overlap assumption reduces the local uncertainties for shallow cumulus clouds, but slightly increases them for stratiform clouds in the WCB regions. For comparison, Fig. 11 includes the mean uncertainties from Fig. 10 as hatched bars. Local uncertainties due to 3D cloud radiative effects, cloud horizontal heterogeneity, and vertical overlap are much larger compared to their mean uncertainties in all four regions of the cyclone. However, the ice-optical parameterization has similar impacts on local and mean uncertainties. This

shows that ~~cloud-3D cloud radiative effects and cloud~~ horizontal heterogeneity and vertical overlap have a much stronger  
 410 impact on CRH locally than on the domain mean. ~~In contrast, the-~~



**Figure 11.** Same as Fig.10 but for local uncertainties of net CRH. For comparison, the mean net CRH uncertainties from Fig.10 are superimposed as gray hatched bars. The mean uncertainties smaller than 0.05 K day<sup>-1</sup> are not shown as they would not be visible in the plot. Note the different x-axes in the panels.


To understand the relative importance of CRH uncertainties at different horizontal spatial scales and for the dynamics of extratropical cyclones, we coarse-grain CRH from different radiative transfer calculations from their original horizontal resolution to horizontal resolutions equivalent to 2.5, 5, 10, 50, 100, and 500 km, which is approximately the spatial extent of the LEM domains. We calculate the CRH uncertainty at different spatial scales  $x$  by computing the *mean absolute* difference  
 415 of net CRH between different radiative transfer calculations from different sets of coarse-grained CRH and average over time and domain.

$$\delta CRH_x = \frac{1}{n \cdot t} \sum_{j=1}^t \sum_{i=1}^n |CRH_{x,a}(i,j) - CRH_{x,b}(i,j)|. \quad (4)$$

Here,  $x$  is the horizontal resolution of the coarse-graining, the subscripts a and b indicate different radiative transfer calculations,  $n$  is the number of horizontal grid points  $i$  at each vertical layer for different resolutions, and  $t$  is the number of time steps  $j$ .  
 420 The CRH uncertainty we calculate from Eq. 4 is an intermediate between the mean and the local CRH uncertainties described earlier such that at the horizontal resolutions of 300 m and 2.5 km this equation is equivalent to Eq. 3 and at the resolution of 500 km, the equation is equivalent to Eq. 2. For the shallow cumulus domain, where clouds are present only in the boundary layer, we apply a mass-weighted vertical average to the CRH uncertainties between 0-2 km altitude interval but for the WCB domains, we apply the averaging between 0-12 km altitude interval (Fig. 12).

Fig. 12 shows that in all regions of the cyclone, the net CRH uncertainty due to 3D cloud radiative effects and cloud horizontal heterogeneity with overlap assumption decreases with increasing the horizontal scale. In the WCB regions, these uncertainties decrease more rapidly than the uncertainty due to the ice-optical parameterization ~~has similar impacts on local and mean uncertainties. This suggests that~~ (cf. green and dark blue lines with red lines in panels b, c, and d of Fig. 12). This analysis indicates that while the CRH uncertainty due to ~~the~~ 3D cloud radiative effects is large at horizontal resolutions of hundreds of  
 430 meters, its spatial extent is limited and becomes less relevant at larger spatial extents. The uncertainty due to cloud horizontal heterogeneity shows a similar pattern but is larger than the uncertainty due to 3D cloud radiative effects. In contrast, the

435 uncertainty due to the ice-optical parameterization is less important for km-scale circulations and more strongly imprints on circulations on larger-scales of hundreds of kilometers. On the other hand, uncertainty from more or less constant as a function of horizontal scale in the WCB regions and dominates the uncertainty at spatial scales of 100 km and larger. This is due to the large-scale stratiform ice clouds that cover the entire domains in the WCB region of the cyclone, and therefore nearly the same level of uncertainty occurs over the entire domains.



**Figure 12.** Net CRH uncertainties as a function of horizontal scale from 300 m to approximately 500 km for all LEM domains. Uncertainties are computed as mass-weighted vertical averages between 0-2 km and 0-12 km altitude intervals for shallow cumulus and WCB regions, respectively. For the uncertainty due to the ice-optical parameterization, the CRH difference between the ice schemes of Fu and the ice scheme of Baum with the general habit mixture is used. Note the different y-axes in the panels.

440 Our analysis suggests that the large-scale changes in the dynamics of the cyclone are more susceptible to CRH uncertainties due to cloud horizontal heterogeneity and vertical overlap should also affect circulations on smaller as well as larger scales (assuming resolved clouds at the horizontal resolution of the NWP model) and ice-optical parameterization than to 3D cloud radiative effects.

## 5 Discussion and conclusion

Recent studies have demonstrated that cloud-radiative heating and cooling of within the atmosphere (hereafter referred to as CRH) is important for the dynamics and predictability of extratropical cyclones. However, CRH is uncertain in numerical weather prediction (NWP) and climate models, and these uncertainties may have systematic effects on model predictions of extratropical cyclones. This motivates us to here study CRH and its uncertainty within extratropical cyclones. To this end, we simulate an idealized extratropical cyclone with the ICON atmosphere model at a horizontal resolution of 2.5 km and focus on CRH within four regions of the cyclone with distinct cloud patterns: shallow cumulus clouds southwest of the cyclone center, deep clouds in the ascending region of the warm conveyor belt (WCB), stratiform low- and mid-level clouds in the WCB cyclonic outflow, and high stratiform ice clouds in the WCB anticyclonic outflow. For these four regions, we perform

450 large eddy model (LEM) simulations at a horizontal resolution of 300 m, which we use to drive offline radiative transfer calculations. This provides a framework to for the first time assess and compare uncertainty in CRH due to four factors within an extratropical cyclone: 3D cloud radiative effects, ice-optical parameterization, cloud horizontal heterogeneity, and cloud vertical overlap. Since we can assume that clouds from the LEM simulations are perfectly known for the purpose of radiative transfer calculation, we quantify the last two factors by coarse-graining the LEM clouds to the horizontal resolution of 2.5  
455 km of the NWP model. We create two sets of NWP homogeneous clouds with and without cloud fraction. By doing so, we quantify to what extent ignorance of the cloud subgrid variability at scales below 2.5 km affects CRH.

We find that 3D cloud radiative effects are ~~overall small and have a noticeable impact only~~ large at the scale of the horizontal grid resolution of 300 m but negligible on larger spatial scales of hundreds of kilometers. 3D cloud radiative effects are also more important for shallow cumulus clouds ~~-Horizontal radiative transfer-~~ than for clouds within the WCB. Horizontal photon  
460 transport within the cloud and from the cloud sides is small in the three WCB regions, where more uniform stratiform clouds prevail. These results are consistent with previous findings that 3D cloud radiative effects are small for stratiform clouds (Črnivec and Mayer, 2021), but important for the dynamics of shallow cumulus clouds (Jakub and Mayer, 2016). ~~Thus, while 3D cloud radiative effects have been shown to affect the organization of subtropical low-level clouds, our study indicates that they do not strongly affect extratropical cyclones.~~

465 The uncertainty in CRH due to different ice-optical parameterizations is substantial in the WCB of the cyclone. We show that using the more complex ice-optical parameterization of Baum leads to weaker shortwave and longwave CRH. Our analysis shows that the ice-optical parameterization is the largest source of uncertainty on spatial scales of 500 km in the upper troposphere where ice clouds prevail. The uncertainty due to ice-optical parameterization is also rather uniform in space, which suggests that it can affect the cyclone by modulating the large-scale radiative heating. Our findings are consistent with the  
470 results of Zhao et al. (2018), ~~who showed that the Baum ice scheme~~ that the ice scheme of Baum has a weaker effect on shortwave cloud radiative effects compared to the ice scheme of Fu. Zhao et al. (2018) showed that the Baum scheme decreases static stability and increases vertical motion in the midlatitudes. An important potential implication of this result that should be tested in future work is to what extent this might enhance latent heating in the WCB ascent region and alter the dynamics of the cyclone (Keshtgar et al., 2023). Our expectation is in line with studies showing that perturbations on a larger spatial  
475 scale are more effective for baroclinic error growth (e.g., Sun and Zhang, 2016). Recently, Lloveras et al. (2023) showed that small-scale perturbations, even with large amplitudes, have a negligible impact on the dynamics of the cyclone and the error growth near the tropopause than larger-scale perturbations with smaller amplitudes.

~~Cloud horizontal heterogeneity and cloud vertical overlap are not resolved at 2.5 resolution, which leads to cloud subgrid variability that can affect CRH. We show that this variability~~ Cloud horizontal heterogeneity contributes significantly to CRH  
480 uncertainty in all regions of the cyclone. This is consistent with previous findings that when cloud subgrid variability is neglected, both longwave and shortwave CRH are overestimated (e.g., Črnivec and Mayer, 2019). The impact is strong for shallow cumulus clouds, while it is much smaller for the stratiform clouds in the WCB. Including the vertical overlap assumption significantly improves the simulation of CRH for shallow cumulus clouds, but in fact slightly degrades CRH for clouds in the WCB since the maximum-random overlap assumption misrepresents the vertical arrangement of cloud layers in

485 sheared flows and a more complex form of the overlap assumption would be needed (e.g., Giuseppe and Tompkins, 2015). The comparison between the impact of cloud horizontal heterogeneity and cloud vertical overlap shows that for shallow cumulus clouds, vertical overlap has a stronger impact on CRH than cloud horizontal heterogeneity. In contrast, for stratiform clouds in the WCB, vertical overlap has a weaker effect on the CRH compared to cloud horizontal heterogeneity. An interesting corollary of our work is that for extratropical cyclones, treating clouds as grid-box quantities in km-scale models with resolutions of  
490 around 2 km appears a reasonable choice, because schemes for cloud cover and vertical overlap themselves can act as sources of uncertainty that are rarely quantified. This supports the approach taken by Hohenegger et al. (2023) for the ICON-Sapphire km-scale model, who decided to remove the cloud cover scheme and treat grid boxes as either entirely cloudy or entirely cloud free.

Our study indicates that ~~for extratropical cyclones,~~ improving the representation of ice-optical properties is particularly  
495 ~~important for reducing CRH uncertainty~~ relevant for the dynamics of extratropical cyclones. This is ~~because i) ice-optical properties represent an important source of uncertainty, ii)~~ due to several reasons. First, for the baroclinic error growth the spatial scale of the uncertainty is more important than the amplitude of the uncertainty (e.g., Lloveras et al., 2023). Although 3D cloud radiative effects are small, and iii) large at the scales of LEM model grid resolution and have been shown to affect the organization of subtropical low-level clouds, their spatial extent is limited. Second, cloud subgrid variability and  
500 3D cloud radiative effects can be taken into account in existing state-of-the-art radiation schemes such as TripleClouds (Črnivec and Mayer, 2021) ~~and SPARTACUS (Hogan et al., 2016).~~ Third, the parametrization of ice-optical properties represents an important source of uncertainty and dominates the CRH uncertainty at larger scales.

Although advances in radiation solvers can account for both 3D radiative effects and cloud subgrid variability, the lack of  
505 knowledge about the shape and surface roughness of ice particles continues to introduce large variability in the CRH and remains a critical challenge to better represent ice-optical properties in models. Future work in the direction of ice-optical properties should also address the problem that in many current models including ICON, the ice crystal effective radii are not treated consistently between the microphysics and radiation schemes (Sullivan and Voigt, 2021), and longwave cloud scattering is neglected (~~Fan et al., 2022~~) (Fan et al., 2023). In view of our results, we believe that future studies should focus on the impact of uncertainties in the ice-optical properties on the dynamics and predictability of extratropical cyclones.

510 *Code and data availability.* The code repository containing scripts for ICON simulations, offline radiative transfer calculations, and the analysis is available in the Gitlab repository (<https://gitlab.phaidra.org/climate/keshtgar-et-al-2024-cyclone-crh-uncertainties>) hosted by the University of Vienna. The simulation's raw outputs are archived on the High-Performance Storage System at the German Climate Computing Center (DKRZ). The post-processed data for producing the figures in the manuscript along with a copy of the Git repository will be published at the LMU open data server.

515 *Author contributions.* The study was designed by AV. The ICON simulations and offline radiative transfer calculations were carried out by BK. The data analysis was done by BK with inputs from AV, BM, and CH. BK led the writing process of the manuscript, with editing from AV and input from all authors.

*Competing interests.* At least one of the (co-)authors is a member of the editorial board of Atmospheric Chemistry and Physics, and the authors have also no other competing interests to declare.

520 *Acknowledgements.* This research has been performed within project B4 of the Transregional Collaborative Research Center SFB/TRR 165 “Waves to Weather” funded by the German Research Foundation (DFG). AV received partial support from the German Ministry of Education and Research (BMBF) and FONA: Research for Sustainable Development ([www.fona.de](http://www.fona.de)) under grant agreement 01LK1509A. We thank the German Climate Computing Center (DKRZ, Hamburg) for providing computing and storage resources as part of project bb1135. We are also thankful to the developers and maintainers of the open-source Python packages NumPy (Harris et al., 2020), Xarray (Hoyer and Hamman, 525 2017), and Matplotlib (Hunter, 2007) that were used for the data analysis.



## References

- Ahmadi-Givi, F., Graig, G. C., and Plant, R. S.: The dynamics of a midlatitude cyclone with very strong latent-heat release, *Quarterly Journal of the Royal Meteorological Society*, 130, 295–323, <https://doi.org/10.1256/qj.02.226>, 2004.
- Baran, A. J., Hill, P., Furtado, K., Field, P., and Manners, J.: A Coupled Cloud Physics–Radiation Parameterization of the Bulk Optical Properties of Cirrus and Its Impact on the Met Office Unified Model Global Atmosphere 5.0 Configuration, *Journal of Climate*, 27, 7725–7752, <https://doi.org/10.1175/JCLI-D-13-00700.1>, 2014.
- Baum, B. A., Yang, P., Heymsfield, A. J., Bansemer, A., Cole, B. H., Merrelli, A., Schmitt, C., and Wang, C.: Ice cloud single-scattering property models with the full phase matrix at wavelengths from 0.2 to 100 $\mu\text{m}$ , *Journal of Quantitative Spectroscopy and Radiative Transfer*, 146, 123–139, <https://doi.org/10.1016/j.jqsrt.2014.02.029>, electromagnetic and Light Scattering by Nonspherical Particles XIV, 2014.
- 535 Baumgart, M., Ghinassi, P., Wirth, V., Selz, T., Craig, G. C., and Riemer, M.: Quantitative view on the processes governing the up-scale error growth up to the planetary scale using a stochastic convection scheme, *Monthly Weather Review*, 147, 1713–1731, <https://doi.org/10.1175/MWR-D-18-0292.1>, 2019.
- Bechtold, P., Köhler, M., Jung, T., Doblas-Reyes, F., Leutbecher, M., Rodwell, M. J., Vitart, F., and Balsamo, G.: Advances in simulating atmospheric variability with the ECMWF model: From synoptic to decadal time-scales, *Quarterly Journal of the Royal Meteorological Society*, 134, 1337–1351, <https://doi.org/10.1002/qj.289>, 2008.
- 540 Binder, H., Boettcher, M., Joos, H., and Wernli, H.: The Role of Warm Conveyor Belts for the Intensification of Extratropical Cyclones in Northern Hemisphere Winter, *Journal of the Atmospheric Sciences*, 73, 3997 – 4020, <https://doi.org/10.1175/JAS-D-15-0302.1>, 2016.
- Booth, J. F., Wang, S., and Polvani, L.: Midlatitude storms in a moister world: Lessons from idealized baroclinic life cycle experiments, *Climate Dynamics*, 41, 787–802, <https://doi.org/10.1007/s00382-012-1472-3>, 2013.
- 545 Costa-Surós, M., Sourdeval, O., Acquistapace, C., Baars, H., Carbajal Henken, C., Genz, C., Hesemann, J., Jimenez, C., König, M., Kretzschmar, J., Madenach, N., Meyer, C. I., Schrödner, R., Seifert, P., Senf, F., Brueck, M., Cioni, G., Engels, J. F., Fieg, K., Gorges, K., Heinze, R., Siligam, P. K., Burkhardt, U., Crewell, S., Hoose, C., Seifert, A., Tegen, I., and Quaas, J.: Detection and attribution of aerosol–cloud interactions in large-domain large-eddy simulations with the ICOSahedral Non-hydrostatic model, *Atmospheric Chemistry and Physics*, 20, 5657–5678, <https://doi.org/10.5194/acp-20-5657-2020>, 2020.
- 550 Črnivec, N. and Mayer, B.: Quantifying the bias of radiative heating rates in numerical weather prediction models for shallow cumulus clouds, *Atmospheric Chemistry and Physics*, 19, 8083–8100, <https://doi.org/10.5194/acp-19-8083-2019>, 2019.
- Črnivec, N. and Mayer, B.: Towards an improved treatment of cloud–radiation interaction in weather and climate models: exploring the potential of the Tripleclouds method for various cloud types using libRadtran 2.0.4, *Geoscientific Model Development*, 14, 3663–3682, <https://doi.org/10.5194/gmd-14-3663-2021>, 2021.
- 555 Edwards, J. M. and Slingo, A.: Studies with a flexible new radiation code. I: Choosing a configuration for a large-scale model, *Quarterly Journal of the Royal Meteorological Society*, 122, 689–719, <https://doi.org/10.1002/qj.49712253107>, 1996.
- Emde, C., Buras-Schnell, R., Kylling, A., Mayer, B., Gasteiger, J., Hamann, U., Kylling, J., Richter, B., Pause, C., Dowling, T., and Bugliaro, L.: The libRadtran software package for radiative transfer calculations (version 2.0.1), *Geoscientific Model Development*, 9, 1647–1672, <https://doi.org/10.5194/gmd-9-1647-2016>, 2016.

- Fan, C., Chen, Y.-H., Chen, X., Lin, W., Yang, P., and Huang, X.: A Refined Understanding of the [Ice Cloud Long-wave Scattering Effects in Climate Model](#), ~~2022~~, [Journal of Advances in Modeling Earth Systems](#), *15*, e2023MS003810, <https://doi.org/https://doi.org/10.1029/2023MS003810>, 2023.
- 565 Fu, Q.: An Accurate Parameterization of the Solar Radiative Properties of Cirrus Clouds for Climate Models, *Journal of Climate*, *9*, 2058 – 2082, [https://doi.org/10.1175/1520-0442\(1996\)009<2058:AAPOTS>2.0.CO;2](https://doi.org/10.1175/1520-0442(1996)009<2058:AAPOTS>2.0.CO;2), 1996.
- Fu, Q. and Liou, K. N.: On the Correlated k-Distribution Method for Radiative Transfer in Nonhomogeneous Atmospheres, *Journal of Atmospheric Sciences*, *49*, 2139 – 2156, [https://doi.org/10.1175/1520-0469\(1992\)049<2139:OTCDMF>2.0.CO;2](https://doi.org/10.1175/1520-0469(1992)049<2139:OTCDMF>2.0.CO;2), 1992.
- Fu, Q. and Liou, K. N.: Parameterization of the Radiative Properties of Cirrus Clouds, *Journal of Atmospheric Sciences*, *50*, 2008 – 2025, [https://doi.org/10.1175/1520-0469\(1993\)050<2008:POTRPO>2.0.CO;2](https://doi.org/10.1175/1520-0469(1993)050<2008:POTRPO>2.0.CO;2), 1993.
- 570 Fu, Q., Yang, P., and Sun, W. B.: An Accurate Parameterization of the Infrared Radiative Properties of Cirrus Clouds for Climate Models, *Journal of Climate*, *11*, 2223 – 2237, [https://doi.org/10.1175/1520-0442\(1998\)011<2223:AAPOTI>2.0.CO;2](https://doi.org/10.1175/1520-0442(1998)011<2223:AAPOTI>2.0.CO;2), 1998.
- Giuseppe, F. D. and Tompkins, A. M.: Generalizing Cloud Overlap Treatment to Include the Effect of Wind Shear, *Journal of the Atmospheric Sciences*, *72*, 2865 – 2876, <https://doi.org/10.1175/JAS-D-14-0277.1>, 2015.
- Griewank, P. J., Heus, T., Lareau, N. P., and Neggers, R. A. J.: Size dependence in chord characteristics from simulated and observed  
575 continental shallow cumulus, *Atmospheric Chemistry and Physics*, *20*, 10 211–10 230, <https://doi.org/10.5194/acp-20-10211-2020>, 2020.
- Grise, K. M., Medeiros, B., Benedict, J. J., and Olson, J. G.: Investigating the Influence of Cloud Radiative Effects on the Extratropical Storm Tracks, *Geophysical Research Letters*, *46*, <https://doi.org/10.1029/2019GL083542>, 2019.
- Harris, C. R., Millman, K. J., van der Walt, S. J., Gommers, R., Virtanen, P., Cournapeau, D., Wieser, E., Taylor, J., Berg, S., Smith, N. J., Kern, R., Picus, M., Hoyer, S., van Kerkwijk, M. H., Brett, M., Haldane, A., del Río, J. F., Wiebe, M., Peterson, P., Gérard-Marchant,  
580 P., Sheppard, K., Reddy, T., Weckesser, W., Abbasi, H., Gohlke, C., and Oliphant, T. E.: Array programming with NumPy, *Nature*, *585*, 357–362, <https://doi.org/10.1038/s41586-020-2649-2>, 2020.
- Hogan, R. J. and Bozzo, A.: A Flexible and Efficient Radiation Scheme for the ECMWF Model, *Journal of Advances in Modeling Earth Systems*, *10*, 1990–2008, <https://doi.org/10.1029/2018MS001364>, 2018.
- Hogan, R. J. and Shonk, J. K. P.: Incorporating the Effects of 3D Radiative Transfer in the Presence of Clouds into Two-Stream Multilayer  
585 Radiation Schemes, *Journal of the Atmospheric Sciences*, *70*, 708 – 724, <https://doi.org/10.1175/JAS-D-12-041.1>, 2013.
- [Hogan, R. J., Schäfer, S. A. K., Klinger, C., Chiu, J. C., and Mayer, B.: Representing 3-D cloud radiation effects in two-stream schemes: 2. Matrix formulation and broadband evaluation, \*Journal of Geophysical Research: Atmospheres\*, \*121\*, 8583–8599, <https://doi.org/https://doi.org/10.1002/2016JD024875>, 2016.](#)
- Hohenegger, C., Korn, P., Linardakis, L., Redler, R., Schnur, R., Adamidis, P., Bao, J., Bastin, S., Behraves, M., Bergemann, M., Biercamp,  
590 J., Bockelmann, H., Brokopf, R., Brüggemann, N., Casaroli, L., Chegini, F., Datseris, G., Esch, M., George, G., Giorgetta, M., Gutjahr, O., Haak, H., Hanke, M., Ilyina, T., Jahns, T., Jungclaus, J., Kern, M., Klocke, D., Kluft, L., Kölling, T., Kornblueh, L., Kosukhin, S., Kroll, C., Lee, J., Mauritsen, T., Mehlmann, C., Mieslinger, T., Naumann, A. K., Paccini, L., Peinado, A., Praturi, D. S., Putrasahan, D., Rast, S., Riddick, T., Roeber, N., Schmidt, H., Schulzweida, U., Schütte, F., Segura, H., Shevchenko, R., Singh, V., Specht, M., Stephan, C. C., von Storch, J.-S., Vogel, R., Wengel, C., Winkler, M., Ziemann, F., Marotzke, J., and Stevens, B.: ICON-Sapphire: simulating  
595 the components of the Earth system and their interactions at kilometer and subkilometer scales, *Geoscientific Model Development*, *16*, 779–811, <https://doi.org/10.5194/gmd-16-779-2023>, 2023.
- Hoyer, S. and Hamman, J.: xarray: N-D labeled arrays and datasets in Python, *Journal of Open Research Software*, *5*, <https://doi.org/10.5334/jors.148>, 2017.

- Hu, Y. X. and Stamnes, K.: An Accurate Parameterization of the Radiative Properties of Water Clouds Suitable for Use in Climate Models, *Journal of Climate*, 6, 728 – 742, [https://doi.org/10.1175/1520-0442\(1993\)006<0728:AAPOTR>2.0.CO;2](https://doi.org/10.1175/1520-0442(1993)006<0728:AAPOTR>2.0.CO;2), 1993.
- 600 Hunter, J. D.: Matplotlib: A 2D graphics environment, *Computing in Science & Engineering*, 9, 90–95, <https://doi.org/10.1109/MCSE.2007.55>, 2007.
- Jakub, F. and Mayer, B.: A three-dimensional parallel radiative transfer model for atmospheric heating rates for use in cloud resolving models—The TenStream solver, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 163, 63–71, <https://doi.org/10.1016/j.jqsrt.2015.05.003>, 2015.
- 605 Jakub, F. and Mayer, B.: 3-D radiative transfer in large-eddy simulations - experiences coupling the TenStream solver to the UCLA-LES, *Geoscientific Model Development*, 9, 1413–1422, <https://doi.org/10.5194/gmd-9-1413-2016>, 2016.
- Keshtgar, B., Voigt, A., Hoose, C., Riemer, M., and Mayer, B.: Cloud-radiative impact on the dynamics and predictability of an idealized extratropical cyclone, *Weather and Climate Dynamics*, 4, 115–132, <https://doi.org/10.5194/wcd-4-115-2023>, 2023.
- 610 Klinger, C., Mayer, B., Jakub, F., Zinner, T., Park, S. B., and Gentine, P.: Effects of 3-D thermal radiation on the development of a shallow cumulus cloud field, *Atmospheric Chemistry and Physics*, 17, 5477–5500, <https://doi.org/10.5194/acp-17-5477-2017>, 2017.
- Li, Y., Thompson, D. W. J., and Bony, S.: The Influence of Atmospheric Cloud Radiative Effects on the Large-Scale Atmospheric Circulation, *Journal of Climate*, 28, 7263 – 7278, <https://doi.org/10.1175/JCLI-D-14-00825.1>, 2015.
- [Lloveras, D. J., Durran, D. R., and Doyle, J. D.: The Two- to Four-Day Predictability of Midlatitude Cyclones: Don't Sweat the Small Stuff. \*Journal of the Atmospheric Sciences\*, 80, 2613 – 2633, <https://doi.org/10.1175/JAS-D-22-0232.1>, 2023.](https://doi.org/10.1175/JAS-D-22-0232.1)
- 615 Mayer, B. and Kylling, A.: Technical note: The libRadtran software package for radiative transfer calculations - description and examples of use, *Atmospheric Chemistry and Physics*, 5, 1855–1877, <https://doi.org/10.5194/acp-5-1855-2005>, 2005.
- Mayer, B.: Radiative transfer in the cloudy atmosphere, *EPJ Web of Conferences*, 1, 75–99, <https://doi.org/10.1140/epjconf/e2009-00912-1>, 2009.
- 620 Mlawer, E. J., Taubman, S. J., Brown, P. D., Iacono, M. J., and Clough, S. A.: Radiative transfer for inhomogeneous atmospheres: RRTM, a validated correlated-k model for the longwave, *Journal of Geophysical Research: Atmospheres*, 102, 16 663–16 682, <https://doi.org/10.1029/97JD00237>, 1997.
- Polvani, L. M. and Esler, J. G.: Transport and mixing of chemical air masses in idealized baroclinic life cycles, *Journal of Geophysical Research: Atmospheres*, 112, <https://doi.org/10.1029/2007JD008555>, 2007.
- 625 Raschendorfer, M.: The new turbulence parameterization of LM.COSMO Newsletter,1, 89–97, 2001.
- [Rieger, D., Köhler, M., Hogan, R. J., Schäfer, S. A. K., Seifert, A., de Lozar, A., and Zängl, G.: ecRad in ICON - Details on the Implementation and First Results \(Technical report\), DWD, issue 004, \[https://www.dwd.de/DE/leistungen/reports\\\_on\\\_icon/pdf\\\_einzelbaende/2019\\\_04.html\]\(https://www.dwd.de/DE/leistungen/reports\_on\_icon/pdf\_einzelbaende/2019\_04.html\), 2019.](https://www.dwd.de/DE/leistungen/reports_on_icon/pdf_einzelbaende/2019_04.html)
- Schäfer, S. A. K. and Voigt, A.: Radiation Weakens Idealized Midlatitude Cyclones, *Geophysical Research Letters*, 45, <https://doi.org/10.1002/2017GL076726>, 2018.
- 630 Seifert, A. and Beheng, K. D.: A two-moment cloud microphysics parameterization for mixed-phase clouds. Part 1: Model description, *Meteorology and Atmospheric Physics*, 92, 45–66, <https://doi.org/10.1007/s00703-005-0112-4>, 2006.
- Smagorinsky, J.: General circulation experiments with the primitive equations: i. the basic experiment, *Monthly Weather Review*, 91, 99 – 164, [https://doi.org/10.1175/1520-0493\(1963\)091<0099:GCEWTP>2.3.CO;2](https://doi.org/10.1175/1520-0493(1963)091<0099:GCEWTP>2.3.CO;2), 1963.
- 635 Stevens, B., Acquistapace, C., Hansen, A., Heinze, R., Klinger, C., Klocke, D., Rybka, H., Schubotz, W., Windmiller, J., Adamidis, P., Arka, I., Barlakas, V., Biercamp, J., Brueck, M., Brune, S., a. Buehler, S., Burkhardt, U., Cioni, G., Costa-Surós, M., Crewell, S., Crüger, T.,

- Deneke, H., Friederichs, P., Henken, C. C., Hohenegger, C., Jacob, M., Jakub, F., Kalthoff, N., Köhler, M., w. Van Laar, T., Li, P., Löhnert, U., Macke, A., Madenach, N., Mayer, B., Nam, C., Naumann, A. K., Peters, K., Poll, S., Quaas, J., Röber, N., Rochetin, N., Scheck, L., Schemann, V., Schnitt, S., Seifert, A., Senf, F., Shapkalijevski, M., Simmer, C., Singh, S., Sourdeval, O., Spickermann, D., Strandgren, J., Tessiot, O., Vercauteren, N., vial, J., Voigt, A., and Zängl, G.: The Added Value of Large-eddy and Storm-resolving Models for Simulating Clouds and Precipitation, *Journal of the Meteorological Society of Japan. Ser. II*, 98, 395–435, <https://doi.org/10.2151/jmsj.2020-021>, 2020.
- 640 Sullivan, S. and Voigt, A.: Ice microphysical processes exert a strong control on the simulated radiative energy budget in the tropics, *Comms. Earth and Env.*, 2, <https://doi.org/10.1038/s43247-021-00206-7>, 2021.
- 645 Sullivan, S., Keshtgar, B., Albern, N., Bala, E., Braun, C., Choudhary, A., Hörner, J., Lentink, H., Papavasileiou, G., and Voigt, A.: How does cloud-radiative heating over the North Atlantic change with grid spacing, convective parameterization, and microphysics scheme in ICON version 2.1.00?, *Geoscientific Model Development*, 16, 3535–3551, <https://doi.org/10.5194/gmd-16-3535-2023>, 2023.
- [Sun, Y. Q. and Zhang, F.: Intrinsic versus Practical Limits of Atmospheric Predictability and the Significance of the Butterfly Effect, \*Journal of the Atmospheric Sciences\*, 73, 1419 – 1438, <https://doi.org/https://doi.org/10.1175/JAS-D-15-0142.1>, 2016.](https://doi.org/https://doi.org/10.1175/JAS-D-15-0142.1)
- 650 Voigt, A., Albern, N., Ceppi, P., Grise, K., Li, Y., and Medeiros, B.: Clouds, radiation, and atmospheric circulation in the present-day climate and under climate change, *WIREs Clim. Change*, 12, <https://doi.org/10.1002/wcc.694>, 2020.
- Voigt, A., Keshtgar, B., and Butz, K.: Tug-Of-War on Idealized Midlatitude Cyclones Between Radiative Heating From Low-Level and High-Level Clouds, *Geophysical Research Letters*, 50, e2023GL103188, <https://doi.org/10.1029/2023GL103188>, 2023.
- Wang, X., Miao, H., Liu, Y., and Bao, Q.: Dependence of cloud radiation on cloud overlap, horizontal inhomogeneity, and vertical alignment in stratiform and convective regions, *Atmospheric Research*, 249, 105–138, <https://doi.org/10.1016/j.atmosres.2020.105358>, 2021.
- 655 Webster, P. J. and Stephens, G. L.: Tropical Upper-Tropospheric Extended Clouds: Inferences from Winter MONEX, *Journal of Atmospheric Sciences*, 37, 1521 – 1541, [https://doi.org/10.1175/1520-0469\(1980\)037<1521:TUTECI>2.0.CO;2](https://doi.org/10.1175/1520-0469(1980)037<1521:TUTECI>2.0.CO;2), 1980.
- Yang, P., Bi, L., Baum, B. A., Liou, K.-N., Kattawar, G. W., Mishchenko, M. I., and Cole, B.: Spectrally Consistent Scattering, Absorption, and Polarization Properties of Atmospheric Ice Crystals at Wavelengths from 0.2 to 100  $\mu\text{m}$ , *Journal of the Atmospheric Sciences*, 70, 330 – 347, <https://doi.org/10.1175/JAS-D-12-039.1>, 2013.
- 660 Yi, B.: Diverse cloud radiative effects and global surface temperature simulations induced by different ice cloud optical property parameterizations, *Scientific Reports*, 12, 10539, <https://doi.org/10.1038/s41598-022-14608-w>, 2022.
- Zängl, G., Reinert, D., Rípodas, P., and Baldauf, M.: The ICON (ICOsahedral Non-hydrostatic) modelling framework of DWD and MPI-M: Description of the non-hydrostatic dynamical core, *Quarterly Journal of the Royal Meteorological Society*, 141, 563–579, <https://doi.org/10.1002/qj.2378>, 2015.
- 665 Zdunkowski, W. Trautmann, T. B. A.: *Radiation in the atmosphere – A course in theoretical meteorology*, Cambridge University Press, New York, 2007.
- Zhao, W., Peng, Y., Wang, B., Yi, B., Lin, Y., and Li, J.: Comparison of three ice cloud optical schemes in climate simulations with community atmospheric model version 5, *Atmospheric Research*, 204, 37–53, <https://doi.org/10.1016/j.atmosres.2018.01.004>, 2018.