

Importance of microphysical settings for climate forcing by stratospheric SO₂ injections as modelled by SOCOL-AERv2

Sandro Vattioni^{1,*}, Andrea Stenke^{1,2,3,*}, Beiping Luo¹, Gabriel Chiodo¹, Timofei Sukhodolov⁴, Elia Wunderlin¹, and Thomas Peter¹

¹Institute for Atmospheric and Climate Science, ETH Zurich, Switzerland

²Institute of Biogeochemistry and Pollutant Dynamics, ETH Zurich, Switzerland

³Eawag, Swiss Federal Institute of Aquatic Science and Technology, Dübendorf, Switzerland

⁴Physikalisch-Meteorologisches Observatorium Davos and World Radiation Center, Davos, Switzerland

*These authors contributed equally to this work.

Correspondence: Sandro Vattioni (sandro.vattioni@env.ethz.ch), Andrea Stenke (andrea.stenke@env.ethz.ch)

Abstract. Solar radiation modification as a sustained deliberate source of SO₂ into the stratosphere (strat-SRM) has been proposed as an option for climate intervention. Global interactive aerosol-chemistry-climate models are often used to investigate the potential cooling efficiencies and associated side effects of hypothesised strat-SRM scenarios. A recent model intercomparison study for composition-climate models with interactive stratospheric aerosol suggests that the modelled climate response to a particular assumed injection strategy, depends on the type of aerosol microphysical scheme used (e.g., modal or sectional representation), alongside also host model resolution and transport. Compared to short-duration volcanic SO₂ emission, the continuous SO₂ injections in strat-SRM scenarios may pose a greater challenge to the numerical implementation of microphysical processes such as nucleation, condensation, and coagulation. This study explores how changing the timesteps and sequencing of microphysical processes in the sectional aerosol-chemistry-climate model SOCOL-AERv2 (40 mass bins) affect model predicted climate and ozone layer impacts considering strat-SRM by SO₂ injections of 5 and 25 Tg(S) yr⁻¹ at 20 km altitude between 30°S and 30°N. The model experiments consider year 2040 boundary conditions for ozone depleting substances and greenhouse gases. We focus on the length of the microphysical timestep and the call sequence of nucleation and condensation, the two competing sink processes for gaseous H₂SO₄. Under stratospheric background conditions, we find no effect of the microphysical setup on the simulated aerosol properties. However, at the high sulfur loadings reached in the scenarios injecting 25 Tg(S) yr⁻¹ of sulfur with a default microphysical timestep of 6 min, changing the call sequence from the default "condensation first" to "nucleation first" leads to a massive increase in the number densities of particles in the nucleation mode ($R < 0.01 \mu\text{m}$) and a small decrease in coarse mode particles ($R > 1 \mu\text{m}$). As expected, the influence of the call sequence becomes negligible when the microphysical timestep is reduced to a few seconds, with the model solutions converging to a size distribution with a pronounced nucleation mode. While the main features and spatial patterns of climate forcing by SO₂ injections are not strongly affected by the microphysical configuration, the absolute numbers vary considerably. For the extreme injection with 25 Tg(S) yr⁻¹, the simulated net global radiative forcing ranges from -2.3 W m^{-2} to -5.3 W m^{-2} , depending on the microphysical configuration. "Nucleation first" shifts the size distribution towards radii better suited for solar scattering ($0.3 \mu\text{m} < R < 0.4 \mu\text{m}$), enhancing the intervention efficiency. The size-distribution shift however generates more

25 ultra-fine aerosol particles, increasing the surface area density, resulting in 10 DU less ozone (about 3% of total column) in the northern midlatitudes and 20 DU less ozone (6%) over the polar caps, compared to the "condensation first" approach. Our results suggest that a reasonably short microphysical time step of 2 minutes or less must be applied to accurately capture the magnitude of the H_2SO_4 supersaturation resulting from SO_2 injection scenarios or volcanic eruptions. Taken together these results underscore how structural aspects of model representation of aerosol microphysical processes become important under conditions of elevated stratospheric sulfur in determining atmospheric chemistry and climate impacts.

30 1 Introduction

The idea of increasing the Earth's albedo by injecting sulfur containing gases into the stratosphere to reduce some of the adverse effects of greenhouse gas (GHG) induced global warming dates back to the 1970s (Budyko, 1974), and was 30 years later further elaborated by Crutzen (2006). The arguments presented by Crutzen called for active scientific research of the kind of activity, which became known under the somewhat misleading term "geoengineering". Therefore, in this study we are using the term "stratospheric solar radiation modification" (strat-SRM). Crutzen's idea is based on the fact that sulfur containing 35 gases, such as SO_2 , H_2S or OCS , injected into the lower stratosphere will form aqueous sulfuric acid aerosol particles via a chain of chemical and microphysical processes (Thomason and Peter, 2006; Kremser et al., 2016). The resulting binary H_2SO_4 - H_2O solution droplets scatter solar radiation back to space, causing a cooling at the Earth's surface. At the same time, however, they heat the stratosphere due to absorption of upwelling long-wave radiation. Moreover, sulfate aerosols play an important role in stratospheric ozone chemistry by providing surfaces for heterogeneous reactions (Solomon, 1999). While 40 the infrared absorptivity is determined to good approximation by the total aerosol volume, the efficiency of scattering solar radiation depends strongly on the detailed aerosol size distribution: Many small particles are more efficient than a few large particles, but they also provide a larger surface area density (SAD) accelerating heterogeneous chemistry (Heckendorn et al., 2009).

45 In the stratosphere, the total aerosol number density and size distribution are governed by the microphysical processes of nucleation, coagulation, condensation, evaporation, and gravitational settling (Kremser et al., 2016, and references therein). The formation of new sulfate aerosol particles occurs via binary homogeneous nucleation of H_2SO_4 and H_2O molecules, or, via heterogeneous nucleation in the presence of appropriate condensation nuclei like meteoritic dust or ions, which requires lower saturation ratios than homogeneous nucleation. The freshly formed particles can grow further through coagulation as well as condensation of H_2SO_4 (together with H_2O). As stratospheric temperatures increase with altitude, the sulfate aerosol 50 particles eventually evaporate above 32 to 35 km, releasing H_2SO_4 back to the gas phase.

The effectiveness of strat-SRM by SO_2 injection has been intensively investigated by using models of different complexity and assuming different injection scenarios (e.g., Heckendorn et al., 2009; Pierce et al., 2010; Niemeier et al., 2011; English et al., 2011; Niemeier and Timmreck, 2015; Tilmes et al., 2018; Vattioni et al., 2019; Weisenstein et al., 2022; Laakso et al., 55 2022; Tilmes et al., 2022). Such modelling studies have advanced our understanding of stratospheric aerosols, but they also highlighted uncertainties regarding the transport, chemistry, and microphysics of the aerosol size distribution. In a recent study,

Weisenstein et al. (2022) presented a model intercomparison exploring the impacts of stratospheric injections of SO₂ gas as well as accumulation-mode sulfuric acid aerosol (AM-H₂SO₄) on atmospheric chemistry and climate. Three general circulation models (GCMs) with interactive aerosol microphysics conducted strictly coordinated model experiments within the framework of the Geoengineering Model Intercomparison Project (GeoMIP, Kravitz et al., 2011), namely the Community Earth System Model (CESM2) with the Whole Atmosphere Community Climate Model (WACCM) atmospheric configuration (Danabasoglu et al., 2020), the middle atmosphere version of ECHAM5 with the HAM microphysical module (MA-ECHAM5-HAM; Stier et al., 2005), and the SOLar Climate Ozone Links model with AER microphysics (SOCOL-AERv2) version 2 (Feinberg et al., 2019). The model experiments included injections of 5 and 25 Tg(S) yr⁻¹ in form of SO₂ gas or AM-H₂SO₄, emitted either as two "point injections" at 30°N and 30°S or as "regional injection" between 30°N and 30°S. Two of the participating models, CESM2 and MA-ECHAM5-HAM, assume the aerosol size distribution can be described by superimposed lognormal size distributions (modal scheme), while Socol-AERv2 uses a size-resolving (sectional) scheme.

The analysis of the simulated particle size distributions for the SO₂ injection scenarios revealed substantial differences between each pair of the three models. CESM2 generates new particles and adds them directly to the Aitken mode ($R \gtrsim 10$ nm), so that there are no nm-sized particles. In contrast, Socol-AERv2 treats these tiny particles down to 0.4 nm. Compared to MA-EACHM-HAM, Socol shows substantially fewer nucleation mode particles, suggesting different roles of nucleation and condensation in both models: the microphysical scheme in Socol-AERv2 appears to prefer condensational growth of existing particles by uptake of H₂SO₄ over the formation of new particles, while the opposite seems to be the case for MA-ECHAM5-HAM. The description of the results of the microphysical processes by means of lognormal functions in modal models, such as CESM2 and MA-ECHAM5-HAM, further complicates the interpretation.

Nucleation and condensation are competing sink processes for gas-phase H₂SO₄, which occur simultaneously in the atmosphere, but typically with different speeds. The characteristic time scale τ for removal of H₂SO₄ molecules by condensation is given by the following Equation:

$$\tau_{\text{cond}} = \frac{4}{Av}, \quad (1)$$

with A being the aerosol surface area density and v the mean thermal velocity of H₂SO₄ molecules. For background conditions with typical SAD values of 5 to 10 $\mu\text{m}^2\text{cm}^{-3}$ in nucleation regions, the equilibrium time scale for condensation is around 0.5–1 h. This value decreases inversely with increasing SAD. Under volcanic or strat-SRM conditions with SAD $\sim 80 \mu\text{m}^2\text{cm}^{-3}$, typical for the 25 Tg(S) yr⁻¹ injection scenario, the equilibrium time scale is less than 5 minutes. As the nucleation rate strongly depends on the gas-phase H₂SO₄ supersaturation, the model timestep used for condensation and nucleation must be significantly smaller than the time required to approach gas-phase equilibrium in order to avoid one process erroneously dominating the gas-to-particle transfer of H₂SO₄. Furthermore, coagulation is also affected by the competition between nucleation and condensation, as it is most efficient at (initially) high number densities and between particles of different size. Small particles move fast, but have only small cross-sections for collision, while large particles exhibit slower Brownian motion, but provide good collision targets for smaller particles (Seinfeld and Pandis, 1997). The correct numerical representation of these simultaneously occurring processes is challenging, especially under sulfur-rich conditions, when characteristic time

scales become extremely short. This motivated us to critically question the microphysical scheme of the sectional SOCOL-AERv2 model and to systematically test the impact of the call sequence of the subroutines for condensation and nucleation, as well as the microphysical timestep on the simulated aerosol properties and the modeled climate response to stratospheric SO₂ injection.

95 The paper is organised as follows: Section 2 presents a brief description of the SOCOL-AERv2 model and details of the experimental setup. Section 3 discusses the impact of the microphysical settings on the aerosol size distribution under stratospheric background conditions as well as under stratospheric injections of SO₂ gas (Section 3.1), on the global mean particle size, aerosol burden and radiative forcing (Section 3.2), and on the meridional distributions of aerosol burden, radiative forcing, and ozone (Section 3.3) resulting from the SO₂ injections. The influence of the microphysical settings on profiles of various
100 quantities is briefly mentioned (Section 3.4) and detailed in the Supplement. To evaluate the changes in SOCOL aerosol microphysics against observations we also tested different settings for the 1991 eruption of Mt. Pinatubo (Section 3.5). Section 4 includes a summary and discussion.

2 Model description and experimental setup

2.1 SOCOL-AERv2

105 A first version of the aerosol-chemistry-climate model SOCOL-AER had been introduced by Sheng et al. (2015), who integrated the size-resolving (sectional) sulfate aerosol module AER (Weisenstein et al., 1997) into the three dimensional grid of the chemistry-climate model (CCM) SOCOLv3 (Stenke et al., 2013), which consists of the middle atmosphere version of the spectral general circulation model MA-ECHAM5 (Roeckner et al., 2003, 2006) and the chemistry-transport model MEZON (Rozanov et al., 1999; Egorova et al., 2003). Since then, the model's tropospheric and stratospheric sulfur cycle have undergone
110 several improvements, resulting in the publication of SOCOL-AERv2 (Feinberg et al., 2019).

SOCOL-AERv2 resolves the sulfate aerosol particles in 40 mass bins, ranging from 0.39 nm to 3.2 μm in radius. Since the mass bins refer to dry aerosol radius, they can also be interpreted as aerosol H₂SO₄ mass bins, ranging from about 2.8 molecules to 1.6×10^{12} molecules of H₂SO₄ per aerosol particle. Neighboring mass bins differ by molecule number doubling.

Detailed descriptions of the original AER microphysics and their adaptations for the coupled model are provided in Weisenstein et al. (1997, 2007) and Sheng et al. (2015), respectively. Aerosol composition, i.e. the sulfuric acid weight percent in the
115 particles, is calculated as function of ambient temperature and H₂O partial pressure using the parameterization of Tabazadeh et al. (1997), which is also used for the calculation of the wet aerosol radius of each mass bin. For the formation of new particles by binary homogeneous nucleation the scheme of Vehkamäki et al. (2002) is used. The scheme calculates the nucleation rate as well as the radius and composition of new particles, meaning that the nucleated mass is added to a single mass bin. The
120 particles can grow through H₂SO₄ condensation and shrink through evaporation, both processes depending on the equilibrium concentration of H₂SO₄ above the particle surface (Ayers et al., 1980; Kulmala and Laaksonen, 1990). Condensational growth leads to an increase of mass in the aerosol phase and a shift of particles to larger mass bins, while evaporation does the opposite. Changes in the net number density occur only upon evaporation from the smallest mass bin or condensational growth

of the largest mass bin. Finally, coagulation reduces number densities and shifts aerosol mass to larger bins. Coagulation is
125 solved by a semi-implicit method (Jacobson and Seinfeld, 2004), whereas at most 90% of the available mass in one mass
bin is allowed to be lost by coagulation within one microphysical time step. Otherwise, the coagulation timestep is reduced.
The coagulation kernel, which defines the collision probability of two particles, depends on the particle radius and the diffu-
sion coefficient (Fuchs, 1964). Finally, sedimentation, which affects the vertical distribution of aerosol particles and reduces
their residence time in the stratosphere, is parameterised following the numerical scheme of Walcek (2000). The gravitational
130 settling velocities of aerosol particles are calculated following Kasten (1968).

The CCM SOCOLv3 and the aerosol module AER are interactively coupled via the chemistry and radiation routines. Sulfur
chemical reactions (Sheng et al., 2015, see Table 1) are fully integrated into the model's chemical solver, which is based
on the implicit iterative Newton-Raphson scheme (Stott and Harwood, 1993). In addition to gas-phase chemistry, the model
includes aqueous-phase oxidation of S(IV) to S(VI) by ozone (O_3) and hydrogen peroxide (H_2O_2) in cloud water (Jacob, 1986).
135 The modeled sulfate aerosol is fed directly into the heterogeneous chemistry and radiation schemes. The aerosol radiative
properties (extinction coefficients, single scattering albedos, and asymmetry factors as functions of wavelength) required to
drive the model dynamics are calculated online from the aerosol size distribution using Mie theory with a temperature- and
humidity-dependent look-up table, which accounts for the aerosol H_2SO_4 weight percent.

The model uses operator splitting. The dynamics module is called every 15 min, whereas the chemistry, aerosol micro-
140 physics, and radiation schemes are called every 2 h. For the microphysical processes, especially for nucleation with its highly
nonlinear dependence on gaseous H_2SO_4 concentration, sub-timesteps are used within the 2-h chemistry loop, to avoid that the
process called first mistakenly dominates the H_2SO_4 -to-particle exchange rate. The default procedure is to use $N_{\text{micro}} = 20$
sub-loops within the chemical timestep, which results in a microphysical timestep of 6 minutes ($2 \text{ h}/N_{\text{micro}} = 2 \text{ h}/20 = 6 \text{ min}$).
The parameter N_{micro} can be easily adjusted between runs. By default, the call sequence for the microphysical processes is
145 condensation first, followed by nucleation (see "CN" sequence in Fig. 1), and finally coagulation. At each chemical timestep,
the model first calculates the new H_2SO_4 gas-phase concentration resulting from chemical production and transport. In the mi-
crophysical loop, the H_2SO_4 concentration is then consecutively updated by condensation and nucleation. As we will see later,
it is also important to distribute the gaseous H_2SO_4 molecules produced during the 2-h chemical timestep homogeneously over
the N_{micro} sub-timesteps (see ΔH_2SO_4 in Fig. 1), rather than passing them as a total amount at the beginning of the micro-
150 physical loop. Otherwise, under conditions of SO_2 injections the 2-hourly call frequency of the chemistry scheme would lead
to initially unrealistically high H_2SO_4 supersaturations in the microphysical loop, which then causes artefacts in the aerosol
microphysics. The aerosol composition as well as the coagulation kernel are calculated only once every 2 h and are kept con-
stant for the microphysical calculations. Finally, sedimentation is calculated after the microphysical sub-loop, again once every
2 h. To test the implications of the aerosol microphysics on the simulated aerosol size distribution under various stratospheric
155 sulfur loadings, we performed several sensitivity simulations, for which we changed the call sequence for condensation and
nucleation or increased the number of microphysical sub-timesteps $N_{\text{micro}} > 20$.




Figure 1. Schematic representation of the call sequence for the microphysical processes in SOCOL-AERv2. The scheme shows the setup with the microphysical sub-loop with $N_{\text{micro}} = 20$ steps by default, or an increased number of steps (60, 80, or 200) in the test runs. By default, the condensation routine is called first and nucleation second ("CN"), which was reversed ("NC") for the tests. Furthermore, the amount of H_2SO_2 gas molecules produced by the chemistry scheme is uniformly distributed over the N_{micro} time steps, instead of providing the total amount at the first microphysical time step, as done in the original set-up.

2.2 SOCOLv4

160 Additionally, we performed simulations with the fully coupled Earth System Model (ESM) SOCOLv4 (Sukhodolov et al., 2021), which is a further development of SOCOL-AERv2. SOCOLv4 is based on ECHAM6, but incorporates the same aerosol
 165 module, AER, as SOCOL-AERv2 (Section 2.1). The major differences between the model versions is that SOCOLv4 is based on the MPI-ESM1.2 (Mauritsen et al., 2019), which incorporates the fully coupled interactive ocean module MPIOM (Jungclaus et al., 2013). SOCOLv4 has a finer horizontal and vertical atmospheric resolution with T63 truncation ($1.9^\circ \times 1.9^\circ$) and 47 vertical pressure levels also reaching up to 0.01 hPa. Compared to SOCOL-AERv2 the default dynamical timestep was halved to 7.5 min, while the default chemical and microphysical time steps are the same as for SOCOL-AERv2 (2h and 6 min, respectively). The interactive ocean as well as the finer spatial resolution make SOCOLv4 computationally much more expensive than SOCOL-AERv2. Therefore, we performed most sensitivity simulations with SOCOL-AERv2 using fixed sea surface temperatures (SST) and sea ice coverage (SIC, see Section 2.3), while SOCOLv4 was primarily used to look at the impact on surface temperature anomalies.

2.3 Experimental setup

170 For the present study, we employed SOCOL-AERv2 with a resolution of 39 hybrid sigma-pressure levels in the vertical and a horizontal truncation of T42 ($\sim 2.8^\circ \times 2.8^\circ$ in latitude and longitude). The simulations for this study include a reference scenario for stratospheric background conditions as well as two perturbation scenarios including stratospheric sulfur injections. The boundary conditions are identical to the GeoMIP test-bed experiment "accumH2SO4"¹ with GHGs and ozone-depleting substances taken from the projections for 2040 from the SSP5-8.5 scenario (see also Weisenstein et al., 2022). SST and
175 SIC are prescribed using an average of the years 1988–2007 of the CMIP5 PCMDI-AMIP-1.1.0 SST/Sea Ice dataset (Taylor et al., 2000). As SOCOL-AERv2 with 39 vertical levels does not generate a quasi-biennial oscillation (QBO) internally, the simulated wind in the equatorial stratosphere is nudged towards observed wind profiles (Stenke et al., 2013). We ran 20 model years for each scenario. The first 5 years are considered as spin-up period (sufficient for the present application), and we use the subsequent 15 years for our analysis.

180 Consistent with Weisenstein et al. (2022), the intervention scenarios examined here apply gaseous SO₂ injections of 5 and 25 Tg(S) yr⁻¹ emitted uniformly in a 2 km thick layer centred around 20 km altitude in the region between 30°S and 30°N over all longitudes. These so-called "regional injections" are complemented by an example of a "point injection" performed with SOCOLv4 (see section 2.2) injecting 5 Tg(S) yr⁻¹ in the form of SO₂ at the same vertical extent but constrained to a region from 10°N to 10°S at the equator only emitting at the 0° meridian. These "point injection" scenarios with constant
185 injection rates were motivated by the G4 GeoMIP experiment described in Kravitz et al. (2011). However, instead of RCP4.5 GHG and injections of 5 Tg(SO₂) yr⁻¹ as specified in Kravitz et al. (2011), we used SSP5-8.5 GHG and injected 5 Tg(S) yr⁻¹ in the form of SO₂, which is consistent with Wunderlin et al. (2024). The transient SSP5-8.5 boundary conditions allow us to explore the sensitivity of surface temperature to the call sequence in a fully coupled ESM.

To determine the effects of the setup of the microphysical scheme (see Fig. 1) on the computed size distribution and aerosol
190 burden, we performed several model simulations for background conditions as well as conditions of strat-SRM. The different simulations vary by reversing the call sequence of the condensation and nucleation routines, or by increasing the number of microphysical timesteps N_{micro} . The model simulations are summarized in Table 1. The experiment BG_CN_20 represents the default setup of the microphysical scheme in SOCOL-AERv2 and is used as the reference simulation.

In the absence of observational data of the stratospheric aerosol layer under strat-SRM conditions, we also tested the effect
195 of different microphysical settings in the modeling of the 1991 Mt. Pinatubo eruption. The 1991 Mt. Pinatubo eruption was specified as 5 Tg S emitted in the form of SO₂ at 21-23 km altitude (2 model levels) above the Mt. Pinatubo geographical location (i.e., two model grid boxes) during one day. This set up corresponds to the "HErSEA_Pin_EI_Ism" scenario proposed by the Interactive Stratospheric Aerosol Model Intercomparison Project (ISA-MIP, Timmreck et al., 2018), which has been shown to have better agreement with observations for some variables compared to scenarios with larger emission amounts and
200 different emission altitudes (Quaglia et al., 2023).

¹Details of the experiment protocol: <http://climate.envsci.rutgers.edu/geomip/testbed.html>

Table 1. Overview of model simulations performed with SOCOL-AERv2 (except for S5p, which was performed with SOCOLv4). Simulation names refer to the following naming convention: "SO₂ injection scenario"_"Call sequence"_" N_{micro} ". BG: background; S5: 5 Tg(S)yr⁻¹, "regional injection"; S5p: 5 Tg(S)yr⁻¹, "point injection" simulated with SOCOLv4; S25: 25 Tg(S)yr⁻¹, "regional injection"; PIN: Pinatubo eruption ("shallow injection" scenario (HErSEA_Pin_El_Ism) of ISA-MIP, Timmreck et al., 2018); CN: condensation first; NC: nucleation first; N_{micro} : number of microphysical timesteps.

Simulation name	SO ₂ injection scenario	Microphysical call sequence	Microphysical timesteps
BG_CN_20	background	Cond-Nuc	20
BG_NC_20		Nuc-Cond	20
S5_CN_20	5 Tg(S) yr ⁻¹ (regional)	Cond-Nuc	20
S5_CN_200		Cond-Nuc	200
S5_NC_20		Nuc-Cond	20
S5_NC_200		Nuc-Cond	200
S5p_CN_20	5 Tg(S) yr ⁻¹ (point)	Cond-Nuc	20
S5p_NC_20		Nuc-Cond	20
S25_CN_20	25 Tg(S) yr ⁻¹ (regional)	Cond-Nuc	20
S25_CN_200		Cond-Nuc	200
S25_NC_20		Nuc-Cond	20
S25_NC_60		Nuc-Cond	60
S25_NC_200		Nuc-Cond	200
PIN_CN_20	Pinatubo 5 Tg(S) (single event, point)	Cond-Nuc	20
PIN_NC_20		Nuc-Cond	20
PIN_NC_200		Nuc-Cond	200

3 Results

In this section, we first analyze how the microphysical settings in SOCOL-AERv2 affect the calculated aerosol size distributions under stratospheric background conditions and under scenarios with SO₂ injection. Next, we examine how the changes in size distributions affect global aerosol properties, such as aerosol loading and net radiative forcing. Finally, we show that microphysical settings directly affect stratospheric chemistry and thus the ozone layer via aerosol surface area density under conditions with strat-SRM.

3.1 Influence of microphysical settings on aerosol size distribution

The upper row of panels in Fig. 2 shows particle size distributions at 55 hPa in the low latitudes (30°S-30°N) for unperturbed conditions and for conditions with strat-SRM. As obvious from panel (a), changing the call sequence of the nucleation and condensation subroutines does not influence the simulated aerosol size distribution under background conditions. Since
210 maximum nucleation rates occur about 2-3 km below the tropical tropopause (Thomason and Peter, 2006), we also examined the size distributions at 115 hPa (not shown), and again find that the call sequence has no impact on the model results. This indicates that the default microphysical timestep of 6 min is sufficiently shorter than the characteristic times of nucleation and condensation under background conditions, so that none of the two processes inappropriately dominates the H₂SO₄-to-particle
215 conversion.

In contrast to background conditions, the SO₂ injections scenarios are highly sensitive to the microphysical settings. Initially, we kept the microphysical timestep constant ($N_{\text{micro}} = 20$), but reversed the call sequence from the default "condensation first" (CN) to "nucleation first" (NC). This modification leads to a massive increase of nucleation mode particles ($R < 0.01 \mu\text{m}$) (Fig. 2c and f, yellow and blue dotted lines).

220 To highlight differences in the coarse mode ($R > 1 \mu\text{m}$), we calculated the fifth moment of the corresponding size distributions (lower row in Fig. 2). This provides an estimate of the downward mass flux due to aerosol sedimentation, which is determined by the product of particle volume (proportional to the third moment) and sedimentation speed (roughly proportional to the second moment). Swapping from CN to NC leads to a significant decrease of coarse mode particles (by one order of magnitude) for the S25 scenario (inset in Fig. 2f).

225 These significant differences in the size distributions demonstrate the dominating role of the first-called process as H₂SO₄ sink, either condensation or nucleation, indicating that the default timestep ($2 \text{ h} / N_{\text{micro}} = 2 \text{ h} / 20 = 6 \text{ min}$) is too long to properly handle elevated stratospheric sulfur loadings. Therefore, we increased the number of microphysical substeps until the resulting particle size distributions of the CN and NC simulations converge. For a sufficiently short microphysical timestep (0.6 min with $N_{\text{micro}} = 200$), the simulations develop a pronounced peak of nucleation mode particles (Fig. 2c, e, orange and blue solid
230 lines) similar to the CN_20 simulations, but with somewhat lower particle number densities.

As expected, the computational costs of the model increase with a shorter microphysical timestep. Increasing the number of microphysical substeps from 20 to 200 almost doubles the required wall-clock time per model year from 4.6 h to 9 h, using parallel computing on 64 CPUs. To assess possibilities to reduce the computational costs, we tested the efficiency of $N_{\text{micro}} = 60$. Since coagulation has the largest timescale and is computationally the most expensive process within the microphysical sub-
235 loop, we also tested a scenario with 80 sub-steps for nucleation and condensation, but only 40 for coagulation. Both sensitivity simulations resulted in similar results (not shown). The red lines in Fig. 2e,f show the results for S25_CN_60, demonstrating excellent agreement with $N_{\text{micro}} = 200$, which gives us confidence in the accuracy of the model solution. Furthermore, the computational demand increased only moderately by about 33% (60 min) per model year (relative to $N_{\text{micro}} = 20$). In conclusion, in SOCOL-AERv2 nucleation first with $N_{\text{micro}} = 60$ provides a very good description of strat-SRM scenarios, even
240 when the loading is extremely high.

We also explored the effects of the distribution of gaseous H_2SO_4 molecules produced during the 2-hourly call of the chemistry routine, either homogeneously across the N_{micro} sub-timesteps or as a total amount at the beginning of the microphysical loop. As Fig. S1 in the Supplement shows, proper partitioning of the H_2SO_4 molecules among the N_{micro} sub-timesteps is critical to avoid an excessive formation of nucleation mode particles due to artificially high H_2SO_4 supersaturations at the beginning of the microphysical substepping. More details can be found in the supplementary material (see Section S2).




Figure 2. Upper row: Size distributions ($dN/d \ln R$, particles cm^{-3}) averaged between 30°S and 30°N at 55 hPa for the model simulations with (a) stratospheric background conditions, (b) "regional" SO_2 injections of 5 Tg(S) yr^{-1} , and (c) 25 Tg(S) yr^{-1} . Lower row: The fifth moment ($dNR^5/d \ln R$, $\mu\text{m}^5 \text{cm}^{-3}$) of the aerosol size distributions as an estimate for aerosol sedimentation mass flux (particle volume ($\propto R^3$) times sedimentation velocity ($\propto R^2$)). Blue lines: simulations with condensation first; orange and red lines: nucleation first. Dashed lines: $N_{\text{micro}} = 20$ microphysical timesteps; solid orange and blue lines: $N_{\text{micro}} = 200$; solid red lines in (c) and (f): $N_{\text{micro}} = 60$. Insert in (f) highlights the differences for coarse particles.

3.2 Influence of microphysical settings on global means of particle size, aerosol burden and radiative forcing

The large differences in the simulated size distribution have wide implications for other key metrics of stratospheric aerosols, namely the average size of the aerosol particles, burden and radiative forcing: these are collectively shown in Fig. 3 in the three sets of experiments. Figure 3a shows the globally averaged effective radius (R_{eff}) at 55 hPa. For background conditions both microphysical settings, CN and NC, result in an average R_{eff} of $0.15 \mu\text{m}$. For the SO_2 injections scenarios, most of the additional sulfur condenses onto existing particles or is consumed in nucleation of new particles, which coagulate preferentially onto the larger background particles. This increases the simulated R_{eff} compared to the background case, moving towards and




Figure 3. Effect of microphysical settings on the global averages of various calculated aerosol and radiative quantities. **(a)** Global mean effective radius (μm) at 55 hPa. **(b)** Global mean sulfuric acid aerosol burden increase (Tg(S)). **(c)** Global mean change in net top-of-atmosphere (shortwave + longwave) radiative forcing (Wm^{-2}). Injection scenarios are: BG = background conditions (no injection); S5 = 5 Tg(S) yr^{-1} "regional" SO_2 injection; S5p = 5 Tg(S) yr^{-1} "point" SO_2 injection; and S25 = 25 Tg(S) yr^{-1} regional SO_2 injection. Blue symbols: condensation first. Orange and red symbols: nucleation first. Open or filled circles: $N_{\text{micro}} = 20$ (or 60). Crosses: $N_{\text{micro}} = 200$. Light blue shading in **(a)**: optimal effective radii for scattering of solar radiation from Dykema et al. (2016) and Figure 4 in Weisenstein et al. (2022).

into the range of optimal effective radius for scattering of solar radiation between 0.3 and 0.4 μm (Weisenstein et al., 2022, see their Fig. 4). The standard microphysical setup (CN, $N_{\text{micro}} = 20$, solid blue circles) result in the largest simulated R_{eff} , as
 255 condensation partly suppresses the subsequent formation of smaller particles via nucleation. Conversely, nucleation first with

long microphysical timesteps (NC, $N_{\text{micro}} = 20$, solid orange circles) exaggerates the formation of small particles, resulting in an underestimation of R_{eff} . Given a sufficiently short timestep ($N_{\text{micro}} = 200$), CN and NC converge to R_{eff} of $0.38 \mu\text{m}$ for the S5 scenario, and $0.48 \mu\text{m}$ for the S25 scenario (blue and orange crosses). Compared to the modal models MA-ECHAM5-HAM and CESM2, the sectional SOCOL-AERv2 in general produces smaller R_{eff} for the "regional injections". Hence, improving
260 the SOCOL-AERv2 aerosol microphysics by swapping the sequence to nucleation first and increasing N_{micro} leads to a slight reduction in the spread in R_{eff} among these models.

Figure 3b shows the impact of microphysical settings on the total (troposphere and stratosphere) aerosol burden increase in the intervention scenarios compared to background conditions. For background conditions, CN and NC with $N_{\text{micro}} = 20$ show an almost identical aerosol burden (see also Table S1 in the Supplement). For the SO_2 injection scenarios, the original
265 setup CN_20 reveals the smallest aerosol burden. The largest aerosol burden is simulated by the simulation with CN_20 setting, since this setup shifts the size distribution towards small particles, which have a longer stratospheric residence time. For the S5 scenario the dependence on call sequence is small, but for S25 the simulated aerosol burdens differ by more than 30% (Table S1). Despite this large spread in the simulated burden increase, SOCOL-AERv2 still falls between the CESM2 and the MA-ECHAM5-HAM models, which showed for most of the simulated injection scenarios the largest and smallest burden
270 increase, respectively (Weisenstein et al., 2022, their Fig. 1).

Figure 3c displays globally averaged changes in the net top-of-atmosphere short- and longwave radiative forcing (RF) due to SO_2 injections. Since SOCOL-AERv2 uses prescribed SST and SIC, the strat-SRM runs remain in non-equilibrium and the perturbation in radiative fluxes at TOA directly quantify the Effective RF (Forster et al., 2016). All S5 simulations show a rather consistent RF change of around -1.5 W m^{-2} . For the S25 simulations, however, we find a large spread in RF,
275 ranging from -2.3 W m^{-2} for the original microphysical setup (CN_20) to -5.4 W m^{-2} for the simulation with reversed call sequence (NC_20). As already mentioned in Weisenstein et al. (2022), the differences in RF between the various SOCOL-AERv2 simulations, but also between different models, are mainly related to the respective burden increases (Fig. 3b). The simulations with the largest burden increase also show the smallest R_{eff} , which efficiently scatters the incoming solar radiation and enhances the negative RF.

As discussed in previous studies (Heckendorn et al., 2009; Kleinschmitt et al., 2018), the efficacy of the SO_2 injection, i.e. the RF per Tg of sulfur injected annually, decreases with increasing injection rate, since the aerosol particles grow larger, which increases sedimentation and decreases scattering efficiency. However, the model intercomparison by Weisenstein et al. (2022) revealed that not only the radiative efficacy itself, but also its decrease with increasing injection rates is strongly model dependent. For SOCOL in Fig. 3c, the radiative efficacy of the various S5 simulations ranges moderately between -
285 0.29 and $-0.32 \text{ W m}^{-2} (\text{Tg}(\text{S}) \text{ yr}^{-1})^{-1}$. For the S25 simulations, the simulations with highest and lowest efficacy differ by more than a factor of 2. The applied microphysical improvements lead to a significantly stronger radiative efficacy ($-0.13 \text{ W m}^{-2} (\text{Tg}(\text{S}) \text{ yr}^{-1})^{-1}$ for S25_CN_60) compared to the default setup ($-0.09 \text{ W m}^{-2} (\text{Tg}(\text{S}) \text{ yr}^{-1})^{-1}$ for S25_CN_20).

As SOCOL-AERv2 does not include an interactive ocean model, but prescribed SSTs, it is unfeasible to test the impact of the call sequence on surface temperature anomalies. To overcome this limitation, we performed a $5 \text{ Tg}(\text{S}) \text{ yr}^{-1}$ "point
290 injection" scenario with the CN setup (S5p_CN_20, see section 2.3) using the ESM SOCOLv4, a coupled model which shares

the same exact aerosol module as SOCOL-AERv2 (see methods). The simulation shows an increase of 25% in stratospheric aerosol burden compared to the conventional S5p_CN_20 scenario (see Fig. 3b, left). The corresponding global averaged surface cooling is 0.65 K and 1.02 K for S5p_CN_20 and S5p_NC_20, respectively, which is an increase of 57%, whereas no significant differences in global stratospheric aerosol burden and RF were found among "regional" S5 scenarios performed
295 with SOCOL-AERv2 (see Fig. 3b, middle). This underlines the sensitivity of our results to the chosen injection scenario (point vs. regional) as well as to the model resolution (Section 2.2). Both the model resolution and the regional confinement of the injection scenario can lead to locally very different H₂SO₄ supersaturation, which influences the sensitivity to the microphysical settings due to the strong non-linearity of nucleation to the H₂SO₄ supersaturation.

3.3 Influence of settings on meridional distributions of aerosol burden, radiative forcing, and ozone

300 Figure 4a,b show the influence of microphysical settings on the modeled latitudinal variation of the sulfate aerosol column burden (stratosphere plus troposphere) for the strat-SRM scenarios simulated with SOCOL-AERv2 (S5 and S25). In contrast, background simulations (not shown) have almost no dependence on the call sequence (see Table S1 in the Supplement). The SO₂ injection scenarios show similar latitudinal patterns, with aerosol column burdens peaking over the inner tropics, confined by the tropical leaky pipe. After overcoming the subtropical jet, the burden again maximizes around 45°N/S in the
305 stratospheric surf zone, whereas the polar regions are isolated by the polar jets. As discussed before (Fig. 3b), the original setting CN_20 results in the smallest aerosol burden, whereas NC_20 with "nucleation first" shifts the size distribution towards smaller particles with less gravitational settling (see also Table S1).

The latitudinal variations of the burden in Fig. 4a,b are reflected in the changes in radiative forcing (RF) in Fig. 4c,d, with reduced irradiance at high aerosol loading, and illustrate the direct radiative effects of the aerosol. However, in contrast
310 to the smooth distributions of aerosol loading, RF exhibits a much higher degree of small fluctuations due to tropospheric cloud variability. The latitudinal variations in RF are very similar for all S5 simulations and the S25 simulations also show a consistent geographic pattern. The negative RF covers more than 80% of the globe, with the exception of the polar caps where absorption of outwelling infrared radiation by the aerosol predominates and the RF becomes positive. The differences between the individual simulations become largest in the tropics, reflecting the sensitivity of the aerosol loading to the microphysical
315 setup.

Figure 4e,f shows the impact of the simulated SO₂ injections on zonally averaged total column ozone as difference to the reference simulation BG_CN_20. As already discussed by Weisenstein et al. (2022), the SO₂ injections lead to a massive reduction of the ozone column. This is caused by accelerated ClO_x-induced and HO_x-induced ozone destruction cycles, which in turn are due to heterogeneous N₂O₅ hydrolysis on the aerosol particles (leaving less NO₂ required for ClO_x and HO_x
320 deactivation). The N₂O₅ hydrolysis rate is proportional to the SAD (see next Section and Figs. S2 and S3 in the Supplement). Both injection scenarios, S5 and S25, show similar patterns with the most pronounced changes in mid- to high latitudes. In the polar regions, the ozone loss is mainly caused by enhanced heterogeneous ClONO₂ activation on the additional aerosol SAD. Furthermore, in agreement with the CESM2 model, SOCOL-AERv2 simulates a decrease of the ozone column in the tropics, where the accelerated Brewer-Dobson circulation results in faster transport of ozone towards higher latitudes. In the tropics,

325 the presented microphysical modifications do not show any significant impact on the simulated ozone decrease (Fig. 4e and f),
despite clear differences in the simulated SAD for the same sulfur injection (Figs. S2 and S3). This result indicates that above
a certain threshold a further SAD increase does not affect the NO_x cycle and its coupling to the ClO_x and HO_x cycles anymore.
The fact that the S25 simulations result in a more pronounced total column ozone change than the S5 simulations is related to a
330 to the increased stratospheric H_2O entry, which enhances ozone loss by the HO_x cycle (Tilmes et al., 2018).

In mid-to high latitudes both injection scenarios, S5 and S25, reveal substantial differences in the total ozone loss simulated,
depending on the microphysical settings used in the simulations. For the S5 simulations (Fig. 4e), the total ozone losses over
Antarctica range between 23 and 26 DU. For the Northern Hemisphere, the spread in simulated polar ozone losses is with 13
to 21 DU even larger. For the S25 simulations (Fig. 4f), the simulated polar ozone loss range between 60 and 80 DU over the
335 Southern Hemisphere, and between 35 and 60 DU over the Northern Hemisphere. It should be noted that the microphysical
setup with the smallest ozone change in one hemisphere, does not necessarily also show the smallest ozone change on the other
hemisphere, which might be related to the dynamical variability.

It should be emphasized that the discussed changes in total column ozone caused by stratospheric SO_2 injections refer to
stratospheric concentrations of ozone depleting substances and GHGs projected for the year 2040. With further decreasing
340 stratospheric chlorine loadings in the future, the impact of the enhanced aerosol SAD under SO_2 injections on total column
ozone might change as the coupling between the ClO_x and NO_x cycle becomes less important.

3.4 Influence of settings on SAD and stratospheric temperature

Strat-SRM by stratospheric SO_2 injection yields an increase in aerosol surface area density (SAD), which enables heteroge-
neous chemical reactions such as N_2O_5 hydrolysis, but which is also an approximate measure of the extinction and, hence,
345 the backscatter of shortwave radiation. Moreover, the aqueous H_2SO_4 aerosol absorbs outwelling longwave radiation, which
increases the air temperature, with repercussions for stratospheric dynamics.

Both quantities, SAD and temperature, are also affected by the microphysical settings CN versus NC and N_{micro} . In brief,
the NC setting with $N_{\text{micro}} = 200$ yields larger SAD than CN with $N_{\text{micro}} = 20$, roughly by 20%. This is due to the smaller
particles with larger SAD and larger burden (see Figs. S2 and S3 in the Supplement). The larger burden, in turn, leads to
350 more longwave radiative heating, which increases stratospheric temperatures. This is a marginal effect in the S5 scenario, but
corresponds to about 1 K higher temperatures under S25 conditions (see Fig. S4 in the Supplement). A strongly temperature
dependent reaction such as $\text{O}_3 + \text{O} \rightarrow 2 \text{O}_2$ changes by about 4% for $\Delta T = 1 \text{ K}$, so that the impact of microphysical settings on
ozone via SAD-changes is by far more important than the impact via T -changes. Also differences from dynamical feedbacks
between the different settings are likely small since the absolute temperature increase from the S25 scenarios is up to 15 K and
355 thus much larger.




Figure 4. Effect of microphysical settings on the zonal averages of various calculated aerosol, radiative and chemical quantities. Left column: regional SO₂ injections of 5 Tg(S) yr⁻¹. Right column: regional SO₂ injections of 25 Tg(S) yr⁻¹. **(a,b)** Sulfuric acid aerosol burden per square meter (mg(S) m⁻²). **(c,d)** Zonal mean net top-of-atmosphere (shortwave + longwave) radiative forcing (Wm⁻²). **(e,f)** Change in zonal average column ozone (Dobson units). Blue lines: simulations with condensation first; orange and red lines: nucleation first. Dashed lines: $N_{\text{micro}} = 20$ microphysical timesteps; solid orange and blue lines: $N_{\text{micro}} = 200$; solid red lines: $N_{\text{micro}} = 60$. All panels use the simulation BG_CN_20 as reference.

3.5 Influence of settings on modeling the eruption of Mt. Pinatubo

So far, our study has highlighted the impacts of the microphysical settings for continuous injections in strat-SRM scenarios. Here, we expand this analysis, by evaluating the effects under conditions of volcanic eruptions on the 1991 eruption of Mt. Pinatubo by using the PIN_CN_20, PIN_NC_20 and the PIN_CN_200 settings (Table 1). We compared the evolution of the computed global stratospheric aerosol burden with SAGE and HIRS satellite data and the evolution of the computed mean effective particle radius with balloon measurements over Laramie (Wyoming, see Fig. 5). Details on the observational data sets and their uncertainties as well as model and inter-model uncertainties can be found in Sukhodolov et al. (2018) and Quaglia et al. (2023). All model settings show a very similar peak in the stratospheric aerosol burden, but distinctly different declines during the years 1992/93. "Nucleation first" shifts the size distribution towards smaller particles, which have a longer stratospheric residence time. The slower decline is in better agreement with observations, although it should be mentioned that the agreement with observations strongly depends on the assumed SO₂ emissions profile (Quaglia et al., 2023). Regarding the mean R_{eff} , PIN_NC_20 simulates smaller values than PIN_CN_20 for the first couple of months after the eruption and larger values later on, as PIN_CN_20 returns faster towards background conditions due to faster sedimentation of larger particles. Overall, the microphysical modifications do not overly influence the discrepancy between modeled and observed R_{eff} (Fig. 5b).

However, other than under strat-SRM conditions the evolution of the aerosol burden and R_{eff} in the PIN_CN_200 scenario are much closer to PIN_CN_20 than to PIN_NC_20. The volcanic eruption is a "point" event in time and space, whereas the strat-SRM scenarios have continuous injections across all longitude and 30°N and 30°S in latitude, which establish a steady-state situation. This leads to H₂SO₄ production rates, which locally are about $10^4 - 10^5$ times larger in the Mt. Pinatubo case compared to S5 and S25. Since nucleation is exponentially dependant on the H₂SO₄ supersaturation this leads to erroneously large nucleation rates in the PIN_NC_20 scenario. Coagulation is not efficient enough to remove the large amount of nucleation mode particles in that scenario. When increasing N_{micro} to 200 (PIN_NC_200) the burden and the R_{eff} of the plume evolve following the PIN_CN_20 scenario since local supersaturation are smaller now and coagulation can keep up with efficiently removing the nucleation mode particles.

Therefore, for volcanic eruptions, where H₂SO₄ supersaturations are locally much larger compared to climate engineering scenarios, the correct solution is much closer to CN_20, since otherwise nucleation would erroneously dominate over condensation. This is a good example of how the very different distributions of H₂SO₄ supersaturation in space and time when simulating volcanic eruptions and strat-SRM scenarios lead to different challenges within aerosol microphysics schemes (Heckendorn et al., 2009; Vattioni et al., 2019).

4 Comparison with other work and conclusions

In this study, we have shown the importance of properly setting the length of the microphysical timestep and the call sequence of nucleation and condensation for modeling the global stratospheric sulfuric acid aerosol under conditions of SO₂ injections for climate engineering. In the aerosol-chemistry-climate model SOCOL-AERv2, the evolution of the H₂SO₄ concentration in the gas-phase is determined by sequential operator splitting using a sub-stepping approach for aerosol microphysics with




Figure 5. (a) Evolution of the simulated global stratospheric aerosol burden (Tg(S)) for PIN_CN_20 and PIN_NC_20 compared with HIRS- and SAGE-II-derived data (SAGE-3,4λ and GLOSSACv2.2, Arfeuille et al., 2013; Thomason et al., 2018; Kovilakam et al., 2020). HIRS-derived total (troposphere and stratosphere) aerosol sulfur burden assumes 75% sulfuric acid by weight (Baran and Foot, 1994). Light blue shaded area: uncertainties of HIRS. (b) Effective particle radius (μm) averaged over the altitude range from 14 to 30 km compared to in-situ measurements taken at Laramie, Wyoming (OPC UWv2.0, Deshler et al., 2019). Thin blue whiskers reflect the measurement uncertainty (adapted from Quaglia et al., 2023). For comparison the steady-state values for S5_CN_20 and S5_NC_20 from this work are shown as thin horizontal red and gray lines in both panels.

a default timestep of 6 min, i.e. the H_2SO_4 gas-phase concentration is consecutively updated by H_2SO_4 production from chemistry, condensation and nucleation. We found the following:

- Under stratospheric background conditions, the call sequence does not affect the model results, indicating that the default number of microphysical sub-timesteps is sufficient to prevent the first called process from spuriously dominating the size distribution.
- Under elevated H_2SO_4 supersaturations in the stratosphere the characteristic times for nucleation and condensation may become shorter than the default microphysical timestep. In such cases, the competition between the two H_2SO_4 sinks affects the simulated aerosol size distribution and the microphysical time step must be reduced.
- The default setting "condensation first" can massively underestimate the fraction of nucleation mode particles, whereas "nucleation first" tends to underestimate the number of coarse mode particles. Tests of numerical convergence with very short timesteps indicate that "nucleation first" yields smaller numerical errors for continuous "regional" SO_2 injections, whereas "condensation first" yields smaller numerical errors for the simulation of volcanic eruptions with locally and temporally more confined emissions resulting in extremely high H_2SO_4 supersaturations.
- Despite significant shifts in simulated aerosol size distributions, the main response patterns of atmospheric chemistry and climate to stratospheric SO_2 injections as simulated with SOCOL-AERv2 are robust to microphysical time integration adjustments, but the strength of the response can differ substantially in the case of high injection rates (such as

405 25 Tg(S) yr⁻¹) or locally confined injections (such as 5 Tg(S) yr⁻¹ "point" injections, S5p), which both lead to large H₂SO₄ supersaturations.

– Depending on the microphysical setting the globally averaged surface temperature response from "point injections" in SOCOLv4 ranges from -0.65 K to -1.02 K cooling when injecting 5 Tg(S) yr⁻¹. In SOCOL-AERv2, the radiative forcing found for the 25 Tg(S) yr⁻¹ "regional injection" scenario varies by more than a factor of 2 between the different microphysical settings. Nevertheless, this model-internal uncertainty in SOCOL-AERv2 is still smaller than the scatter between the three GCMs with interactive aerosol microphysics – CESM2-WACCM6, MA-ECHAM5-HAM, and SOCOL-AERv2 – compared by Weisenstein et al. (2022) in strictly coordinated strat-SRM modeling.

410

The first part of our conclusions confirms the study by Wan et al. (2013), who investigated different time integration methods to solve the H₂SO₄ continuity equation using different versions of the ECHAM-HAM model: HAM1 with an Euler forward scheme with sequential operator splitting similar to SOCOL-AERv2, but without microphysical substeps; HAM2 with a two-step time integration scheme implemented by Kokkola et al. (2009) as well as a version which implicitly solves production, condensation and nucleation simultaneously via linearization of nucleation. They identified sequential operator splitting with too long timesteps as major source of numerical error in HAM1, and found that the implicit two time stepping scheme with simultaneous processing of H₂SO₄ production, condensation and nucleation most accurately represents the competition between these processes. The microphysical sub-stepping technique as applied in SOCOL-AERv2 improves the results of the operator splitting approach, but requires a sufficiently large number of substeps. Instead of using a fixed number of substeps, a dynamical timestep adjustment could be beneficial, but we have not tested this here. However, the best way forward would be to directly implement an implicit solver of H₂SO₄ production, condensation and nucleation into the next version of SOCOL as it is presented in (Wan et al., 2013). This will likely generate more accurate results while avoiding the need for sub-loops, which are computationally more expensive. However, these implicit solvers have not yet been tested for numerical stability under conditions of continuously larger SO₂ injections as would occur in strat-SRM scenarios.

415

420

425

The importance of aerosol microphysics and the competition between nucleation and condensation on the simulated aerosol size distribution and the radiative efficiency of stratospheric sulfur injections was also shown by Laakso et al. (2022), who investigated different injection strategies using the ECHAM-HAMMOZ model with two different aerosol schemes, the sectional SALSA scheme as well as the modal M7 scheme. SALSA describes the aerosol size distribution in 10 mass bins, while M7 uses 7 lognormal modes. The authors found that nucleation of new particles dominates over condensational particle growth in the sectional SALSA scheme, while the opposite is the case in the modal M7 module. In addition, the use of M7 with lognormal modes can result in a minimum in the particle size distribution in the optimal size range for solar scattering due to the accumulation mode reaching its largest size, which adds mass to the coarse mode in the injection scenario applied in Laakso et al. (2022). The resulting gap between the two modes tends to underestimate gravitational settling. These differences resulted in smaller particles in SALSA and, therefore, a stronger radiative forcing. For an injection scenario of 20 Tg(S) yr⁻¹, SALSA revealed a global net ToA radiative forcing of around -8 W m⁻², M7 resulted in -3 W m⁻². This spread is even larger than what we found for the S25 simulations S25_CN_20 and S25_NC_20. Laakso et al. (2022) further investigated the impact

430

435

of the competition between nucleation and condensation by performing simulations with the nucleation being switched off in
440 both aerosol modules by emitting 25% of the sulfur directly as 3 nm particles. The results of these sensitivity studies indicate
that the different treatment of nucleation and condensation explains the differences in radiative forcing between SALSA and
M7 only partly: The difference in radiative forcing was weakened from -5 W m^{-2} to about -3 W m^{-2} .

Apart from time integration or representation of the aerosol size distribution, numerical parameterizations of individual
processes are another source of uncertainty. The binary-homogeneous nucleation scheme by Vehkamäki et al. (2002), for
445 example, is widely used in models, including SOCOL-AERv2 or the above mentioned aerosol schemes SALSA and M7.
The latter two include an extension of the scheme for high sulfate concentrations implemented by Kokkola et al. (2009),
using the collision rate as maximum possible nucleation rate. In a very recent study, Yu et al. (2023) evaluated simulated
nucleation rates in the lowermost stratosphere by CLOUD laboratory measurements under stratospheric temperatures. They
found that the Vehkamäki scheme overestimates observed nucleation rates by 3 to 4 orders of magnitude. As the particle size
450 distribution is not only determined by nucleation, but also by particle growth through condensation and coagulation, Yu et al.
(2023) compared the simulated size distributions to in-situ measurements of the particle number densities down to a diameter
of 3 nm obtained during the NASA Atmospheric Tomography Mission (ATom) between 2016 and 2018. In the size range
between 3 to 10 nm, the number densities simulated with the GEOS-Chem model using the Vehkamäki et al. scheme were
1-2 orders of magnitude larger than observed. The same holds true for SOCOL-AERv2: under background conditions in the
455 southern hemisphere lowermost stratosphere (70° S , 12 km altitude), modeled number densities for particles smaller than
10 nm in diameter range between 10^3 and 10^4 cm^{-3} , while the ATom observations indicate values between slightly below
 10^1 to 10^2 cm^{-3} . Using the kinetic scheme for ion-mediated and binary homogeneous nucleation (Yu et al., 2020) calculated
nucleation rates, but also the size distributions simulated by GEOS-Chem were closer to ATom. Furthermore, the results by Yu
et al. (2023) suggest that under low stratospheric background H_2SO_4 concentrations nucleation on ions, which is usually not
460 represented in global models, dominates over binary homogeneous H_2SO_4 - H_2O nucleation. However, the importance of binary
homogeneous nucleation is expected to increase under high H_2SO_4 concentrations. Unfortunately, CLOUD measurements of
nucleation rates refer to stratospheric background conditions only and do not cover strongly elevated H_2SO_4 concentrations
under SO_2 injection scenarios or after volcanic eruptions, but based on the findings of Yu et al. (2023) it may be that all
models using the Vehkamäki scheme overestimate the role of nucleation. This might explain the low bias in the simulated
465 mean effective radius compared to in-situ measurements following the eruption of Mt. Pinatubo. Furthermore, this might
have substantial repercussions on the simulated aerosol size distribution, aerosol burdens and radiative forcing under strat-
SRM conditions, most likely resulting in a decreased efficiency of SO_2 injections. The reported weaknesses of the Vehkamäki
scheme were addressed by Määttänen et al. (2018), who presented a new parameterisation for sulfuric acid aerosol formation
including homogeneous and ion-induced nucleation pathways validated by CLOUD laboratory measurements. In future work,
470 the Määttänen et al. nucleation scheme, which is reported to be valid for the whole range of atmospheric conditions including
high stratospheric sulfuric acid concentrations during SO_2 injection scenarios, will be used in SOCOL-AER instead of the
Vehkamäki scheme (Määttänen et al., 2018).

This work adds to a series of recent publications that highlight the crucial role of aerosol microphysics for simulated aerosol properties and modeled estimates of strat-SRM effects on atmospheric chemistry and climate. Our results clearly demonstrate that there is considerable uncertainty when numerical schemes like the aerosol microphysics in SOCOL-AERv2 are applied under unprecedented conditions, such as strat-SRM with continuously large SO₂ injections, even if these models have been thoroughly evaluated and are well capable of reproducing observations under background or moderately perturbed conditions like volcanic eruptions. It should be emphasized that our conclusions are mainly based on simulations of "regional" SO₂ injections, which are supported by "point injection" scenarios and simulations of the 1991 Mt. Pinatubo eruption. As the nucleation rate strongly depends on the gas-phase H₂SO₄ concentration, ambient temperatures and relative humidities, the optimal number of microphysical (sub-)timesteps will depend on the assumed SO₂ injection rates, but also on the injection scenario and spatial confinement of the injections. "Point" injections of SO₂, for example result in very high, but locally confined H₂SO₄ supersaturations, which makes the results more sensitive to the details of the microphysical approach. This effect is shown with the "point injection" scenarios (S5p), which are much more sensitive to the microphysical settings compared to the "regional injection" scenarios (S5, see Fig. 3b). The intention of this paper is to raise awareness within the (aerosol) modelling community for potential numerical problems within conventional aerosol microphysics modules when applying them to unprecedented extreme conditions such as high H₂SO₄ supersaturations from SO₂ injection for strat-SRM. To increase confidence in the different aerosol microphysics modules included in the various aerosol-chemistry climate models, we recommend conducting a model intercomparison study including to explore differences among the process-split sub-stepping methods of aerosol microphysics schemes under conditions of continuous large H₂SO₄ supersaturation, such as would occur under stratospheric solar strat-SRM scenarios.

While this study focused on the importance of a proper temporal resolution of aerosol microphysics, it did not address effects of spatial model resolution, which also affects model results (e.g., Niemeier et al., 2020). Properly resolving the various temporal and spatial scales, ranging from nanometers and seconds for microphysical processes to kilometers and decades for global climate, remains a significant challenge for aerosol-chemistry-climate models (Vattioni et al., 2019; Weisenstein et al., 2022). Continuous model development, such as embedded SO₂ injection plume modelling (Sun et al., 2022), is indispensable to close the spatial and temporal gap between aircraft injection plumes and large-scale model grids, and to effectively reduce existing model uncertainties with respect to the effectiveness of strat-SRM by stratospheric sulfur injections. Furthermore, additional laboratory and co-ordinated model intercomparison studies of aerosol formation, growth and dispersion under various stratospheric conditions could also be beneficial to evaluate and improve existing numerical models or to develop new explicit aerosol schemes, which potentially will be directly integrated in chemical solvers.

Supplement. The supplement related to this article is available online at <https://doi.org/10.5194/amt-16-xxx-2023-supplement>.

Code and data availability. The original SOCOI-AERv2 code is available at <https://doi.org/10.5281/zenodo.5733121> (Brodowsky et al., 2018). The simulation data using that model code, which does not account for the interpolation of H₂SO₄ production within the
505 microphysical sub-loop are available at <https://doi.org/10.3929/ethz-b-000610854> (Stenke et al., 2023). The modified source code of SOCOL-AERv2 handling the microphysical sub-loop by taking into account the interpolation of H₂SO₄ production within the microphysical sub-loop can be found here: <https://doi.org/10.5281/zenodo.10791230> (Vattioni et al., 2024) and the data from these simulations are available at <http://hdl.handle.net/20.500.11850/622193> (Vattioni et al., 2023).

Author contributions. SV proposed the study and implemented the modifications to the microphysical scheme. SV and AS performed the
510 strat-SRM simulations with SOCOL-AERv2 and analysed the model results. EW performed and analyzed the simulations with SOCOLv4. TS conducted and analyzed the Pinatubo simulations. Significant scientific guidance on the overall project was provided by TP and BL. AS drafted a first version of the manuscript. All co-authors contributed to the discussion and the text.

Competing interests. At least one of the (co-)authors is a member of the editorial board of Geoscientific Model Development. Other than this, the authors declare that they have no conflict of interest.

515 *Acknowledgements.* Our special thanks go to Debra Weisenstein for discussions about the original AER code. We also thank Ilaria Quaglia for providing the processed OPC data. Furthermore, we appreciate the reviews from Jim Haywood, Daniele Visioni, Ulrike Niemeier and Anton Laakso, which significantly improved the manuscript. Support for Gabriel Chiodo and Andrea Stenke was provided by the Swiss Science Foundation within the Ambizione grant no. PZ00P2_180043. Support for Sandro Vattioni was provided by the ETH Research grant no. ETH-1719-2 as well as by the Harvard Geoenvironment Research Program. Gabriel Chiodo and Sandro Vattioni received also funding
520 from the Simons Foundation (grant no. SFI-MPS-SRM-00005217). Timofei Sukhodolov acknowledges the support from the Swiss National Science Foundation (grant no. 200020-182239), the Karbacher Fonds, Graubünden, Switzerland and the Simons Foundation (grant no. SFI-MPS-SRM-00005217).

References

- Arfeuille, F., Luo, B. P., Heckendorn, P., Weisenstein, D., Sheng, J. X., Rozanov, E., Schraner, M., Brönnimann, S., Thomason, L. W., and
525 Peter, T.: Modeling the stratospheric warming following the Mt. Pinatubo eruption: uncertainties in aerosol extinctions, *Atmos. Chem. Phys.*, 13, 11 221–11 234, <https://doi.org/10.5194/acp-13-11221-2013>, 2013.
- Ayers, G., Gillett, R., and Gras, J.: On the vapor pressure of sulfuric acid, *Geophys. Res. Lett.*, 7, 433–436,
<https://doi.org/10.1029/GL007i006p00433>, 1980.
- Baran, A. J. and Foot, J. S.: New application of the operational sounder HIRS in determining a climatology of sulphuric acid aerosol from
530 the Pinatubo eruption, *J. Geophys. Res.*, 99, 25 673–25 679, <https://doi.org/10.1029/94JD02044>, 1994.
- Brodowsky, C. et al.: SOCOL-AERv2 model code, Zenodo, <https://doi.org/10.5281/zenodo.5733121>, 2018.
- Budyko, M. I.: *Climate and Life*, Academic Press, New York, USA, 1974.
- Crutzen, P.: Albedo Enhancement by Stratospheric Sulfur Injections: A Contribution to Resolve a Policy Dilemma?, *Climatic Change*, 77,
211–220, <https://doi.org/10.1007/s10584-006-9101-y>, 2006.
- 535 Danabasoglu, G., Lamarque, J.-F., Bacmeister, J., Bailey, D. A., DuVivier, A. K., Edwards, J., Emmons, L. K., Fasullo, J., Garcia, R.,
Gettelman, A., Hannay, C., Holland, M. M., Large, W. G., Lauritzen, P. H., Lawrence, D. M., Lenaerts, J. T. M., Lindsay, K., Lipscomb,
W. H., Mills, M. J., Neale, R., Oleson, K. W., Otto-Bliessner, B., Phillips, A. S., Sacks, W., Tilmes, S., van Kampenhout, L., Vertenstein,
M., Bertini, A., Dennis, J., Deser, C., Fischer, C., Fox-Kemper, B., Kay, J. E., Kinnison, D., Kushner, P. J., Larson, V. E., Long, M. C.,
Mickelson, S., Moore, J. K., Nienhouse, E., Polvani, L., Rasch, P. J., and Strand, W. G.: The Community Earth System Model Version 2
540 (CESM2), *Journal of Advances in Modeling Earth Systems*, 12, e2019MS001 916, <https://doi.org/10.1029/2019MS001916>, 2020.
- Deshler, T., Luo, B., Kovilakam, M., Peter, T., and Kalnajs, L. E.: Retrieval of Aerosol Size Distributions From In Situ Particle Counter
Measurements: Instrument Counting Efficiency and Comparisons With Satellite Measurements, *Journal of Geophysical Research:*
Atmospheres, 124, 5058–5087, <https://doi.org/10.1029/2018JD029558>, 2019.
- Dykema, J. A., Keith, D. W., and Keutsch, F. N.: Improved aerosol radiative properties as a foundation for solar geoengineering risk
545 assessment, *Geophysical Research Letters*, 43, 7758–7766, <https://doi.org/10.1002/2016GL069258>, 2016.
- Egorova, T., Rozanov, E., Zubov, V., and Karol, I. L.: Model for Investigating Ozone Trends (MEZON), *Izvestiya, Atmos. Oceanic Phys.*,
39, 277–292, 2003.
- English, J. M., Toon, O. B., Mills, M. J., and Yu, F.: Microphysical simulations of new particle formation in the upper troposphere and
lower stratosphere, *Atmos. Chem. Phys.*, 11, 9303–9322, <https://doi.org/10.5194/acp-11-9303-2011>, 2011.
- 550 Feinberg, A., Sukhodolov, T., Luo, B.-P., Rozanov, E., Winkel, L. H. E., Peter, T., and Stenke, A.: Improved tropospheric and stratospheric
sulfur cycle in the aerosol–chemistry–climate model SOCOL-AERv2, *Geosci. Model Dev.*, 12, 3863–3887,
<https://doi.org/10.5194/gmd-12-3863-2019>, 2019.
- Forster, P. M., Richardson, T., Maycock, A. C., Smith, C. J., Samset, B. H., Myhre, G., Andrews, T., Pincus, R., and Schulz, M.:
Recommendations for diagnosing effective radiative forcing from climate models for CMIP6, *Journal of Geophysical Research:*
555 *Atmospheres*, 121, 12,460–12,475, <https://doi.org/10.1002/2016JD025320>, 2016.
- Fuchs, N.: *The Mechanics of Aerosols*, Pergamon Press, Oxford, 1964.
- Heckendorn, P., Weisenstein, D., Fueglistaler, S., Luo, B. P., Rozanov, E., Schraner, M., Thomason, L. W., and Peter, T.: The impact of
geoengineering aerosols on stratospheric temperature and ozone, *Environ. Res. Lett.*, 4, 045 108,
<https://doi.org/10.1088/1748-9326/4/4/045108>, 2009.

- 560 Jacob, D. J.: Chemistry of OH in remote clouds and its role in the production of formic acid and peroxymonosulfate, *J. Geophys. Res.*, 91, 9807–9826, <https://doi.org/10.1029/JD091iD09p09807>, 1986.
- Jacobson, M. Z. and Seinfeld, J. H.: Evolution of nanoparticle size and mixing state near the point of emission, *Atmos. Environ.*, 38, 1839–1850, <https://doi.org/10.1016/j.atmosenv.2004.01.014>, 2004.
- Jungclaus, J. H., Fischer, N., Haak, H., Lohmann, K., Marotzke, J., Matei, D., Mikolajewicz, U., Notz, D., and von Storch, J. S.:
565 Characteristics of the ocean simulations in the Max Planck Institute Ocean Model (MPIOM) the ocean component of the MPI-Earth system model, *Journal of Advances in Modeling Earth Systems*, 5, 422–446, <https://doi.org/10.1002/jame.20023>, 2013.
- Kasten, F.: Falling speed of aerosol particles, *J. Appl. Meteorol.*, 7, 944–947, [https://doi.org/10.1175/1520-0450\(1968\)007<0944:FSOAP>2.0.CO;2](https://doi.org/10.1175/1520-0450(1968)007<0944:FSOAP>2.0.CO;2), 1968.
- Kleinschmitt, C., Boucher, O., and Platt, U.: Sensitivity of the radiative forcing by stratospheric sulfur geoengineering to the amount and
570 strategy of the SO₂ injection studied with the LMDZ-S3A model, *Atmos. Chem. Phys.*, 18, 2769–2786, <https://doi.org/10.5194/acp-18-2769-2018>, 2018.
- Kokkola, H., Hommel, R., Kazil, J., Niemeier, U., Partanen, A.-I., Feichter, J., and Timmreck, C.: Aerosol microphysics modules in the framework of the ECHAM5 climate model – intercomparison under stratospheric conditions, *Geosci. Model Dev.*, 2, 97–112, <https://doi.org/10.5194/gmd-2-97-2009>, 2009.
- 575 Kovilakam, M., Thomason, L. W., Ernest, N., Rieger, L., Bourassa, A., and Millán, L.: The Global Space-based Stratospheric Aerosol Climatology (version 2.0): 1979–2018, *Earth. Syst. Sci. Data*, 12, 2607–2634, <https://doi.org/10.5194/essd-12-2607-2020>, 2020.
- Kravitz, B., Robock, A., Boucher, O., Schmidt, H., Taylor, K. E., Stenchikov, G., and Schulz, M.: The Geoengineering Model Intercomparison Project (GeoMIP), *Atmospheric Science Letters*, 12, 162–167, <https://doi.org/10.1002/asl.316>, 2011.
- Kremser, S., Thomason, L. W., von Hobe, M., Hermann, M., Deshler, T., Timmreck, C., Toohey, M., Stenke, A., Schwarz, J. P., Weigel, R.,
580 Fueglistaler, S., Prata, F. J., Vernier, J.-P., Schlager, H., Barnes, J. E., Antuña-Marrero, J.-C., Fairlie, D., Palm, M., Mahieu, E., Notholt, J., Rex, M., Bingen, C., Vanhellemont, F., Bourassa, A., Plane, J. M. C., Klocke, D., Carn, S. A., Clarisse, L., Trickl, T., Neely, R., James, A. D., Rieger, L., Wilson, J. C., and Meland, B.: Stratospheric aerosol – Observations, processes, and impact on climate, *Rev. Geophys.*, 54, 278–335, <https://doi.org/10.1002/2015RG000511>, 2016.
- Kulmala, M. and Laaksonen, A.: Binary nucleation of water–sulfuric acid system: Comparison of classical theories with different H₂SO₄
585 saturation vapor pressures, *J. Chem. Phys.*, 93, 696–701, <https://doi.org/10.1063/1.459519>, 1990.
- Laakso, A., Niemeier, U., Vioni, D., Tilmes, S., and Kokkola, H.: Dependency of the impacts of geoengineering on the stratospheric sulfur injection strategy – Part I: Intercomparison of modal and sectional aerosol modules, *Atmos. Chem. Phys.*, 22, 93–118, <https://doi.org/10.5194/acp-22-93-2022>, 2022.
- Mauritsen, T., Bader, J., Becker, T., Behrens, J., Bittner, M., Brokopf, R., Brovkin, V., Claussen, M., Crueger, T., Esch, M., Fast, I., Fiedler, S., Fläschner, D., Gayler, V., Giorgetta, M., Goll, D. S., Haak, H., Hagemann, S., Hedemann, C., Hohenegger, C., Ilyina, T., Jahns, T., Jimenez-de-la Cuesta, D., Jungclaus, J., Kleinen, T., Kloster, S., Kracher, D., Kinne, S., Kleberg, D., Lasslop, G., Kornblueh, L., Marotzke, J., Matei, D., Meraner, K., Mikolajewicz, U., Modali, K., Möbis, B., Müller, W. A., Nabel, J. E. M. S., Nam, C. C. W., Notz, D., Nyawira, S.-S., Paulsen, H., Peters, K., Pincus, R., Pohlmann, H., Pongratz, J., Popp, M., Raddatz, T. J., Rast, S., Redler, R., Reick, C. H., Rohrschneider, T., Schemann, V., Schmidt, H., Schnur, R., Schulzweida, U., Six, K. D., Stein, L., Stemmler, I., Stevens, B., von
595 Storch, J.-S., Tian, F., Voigt, A., Vrese, P., Wieners, K.-H., Wilkenskjaeld, S., Winkler, A., and Roeckner, E.: Developments in the MPI-M Earth System Model version 1.2 (MPI-ESM1.2) and Its Response to Increasing CO₂, *Journal of Advances in Modeling Earth Systems*, 11, 998–1038, <https://doi.org/10.1029/2018MS001400>, 2019.

- Määttänen, A., Merikanto, J., Henschel, H., Duplissy, J., Makkonen, R., Ortega, I. K., and Vehkamäki, H.: New Parameterizations for Neutral and Ion-Induced Sulfuric Acid-Water Particle Formation in Nucleation and Kinetic Regimes, *Journal of Geophysical Research: Atmospheres*, 123, 1269–1296, <https://doi.org/10.1002/2017JD027429>, 2018.
- 600 Niemeier, U. and Timmreck, C.: What is the limit of climate engineering by stratospheric injection of SO₂?, *Atmos. Chem. Phys.*, 15, 9129–9141, <https://doi.org/10.5194/acp-15-9129-2015>, 2015.
- Niemeier, U., Schmidt, H., and Timmreck, C.: The dependency of geoengineered sulfate aerosol on the emission strategy, *Atmospheric Science Letters*, 12, 189–194, <https://doi.org/10.1002/asl.304>, 2011.
- 605 Niemeier, U., Richter, J. H., and Tilmes, S.: Differing responses of the quasi-biennial oscillation to artificial SO₂ injections in two global models, *Atmospheric Chemistry and Physics*, 20, 8975–8987, <https://doi.org/10.5194/acp-20-8975-2020>, 2020.
- Pierce, J. R., Weisenstein, D. K., Heckendorn, P., Peter, T., and Keith, D. W.: Efficient formation of stratospheric aerosol for climate engineering by emission of condensable vapor from aircraft, *Geophys. Res. Lett.*, 37, <https://doi.org/10.1029/2010GL043975>, 2010.
- Quaglia, I., Timmreck, C., Niemeier, U., Visionsi, D., Pitari, G., Brodowsky, C., Brühl, C., Dhomse, S. S., Franke, H., Laakso, A., Mann, G. W., Rozanov, E., and Sukhodolov, T.: Interactive stratospheric aerosol models’ response to different amounts and altitudes of SO₂ injection during the 1991 Pinatubo eruption, *Atmospheric Chemistry and Physics*, 23, 921–948, <https://doi.org/10.5194/acp-23-921-2023>, 2023.
- 610 Roeckner, E., Bäuml, G., Bonaventura, L., Brokopf, R., Esch, M., Giorgetta, M., Hagemann, S., Kirchner, I., Kornblüeh, L., Manzini, E., Rhodin, A., Schlese, U., Schulzweida, U., and Tompkins, A.: The atmospheric general circulation model ECHAM 5. PART I: Model description, MPI-Report No. 349, Max-Planck-Institut für Meteorologie, Hamburg, http://www.mpimet.mpg.de/fileadmin/publikationen/Reports/max_scirep_349.pdf, 2003.
- Roeckner, E., Brokopf, R., Esch, M., Giorgetta, M., Hagemann, S., Kornblüeh, L., Manzini, E., Schlese, U., and Schulzweida, U.: Sensitivity of simulated climate to horizontal and vertical resolution in the ECHAM5 atmosphere model, *J. Climate*, 19, 3771–3791, <https://doi.org/10.1175/JCLI3824.1>, 2006.
- 620 Rozanov, E. V., Zubov, V. A., Schlesinger, M. E., Yang, F., and Andronova, N. G.: The UIUC three-dimensional stratospheric chemical transport model: Description and evaluation of the simulated source gases and ozone, *J. Geophys. Res.*, 104, 11 755–11 781, <https://doi.org/10.1029/1999JD900138>, 1999.
- Seinfeld, J. H. and Pandis, S. N.: *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change*, John Wiley & Sons Inc., 1997.
- 625 Sheng, J., Weisenstein, D. K., Luo, B., Rozanov, E., Stenke, A., Anet, J., Bingemer, H., and Peter, T.: Global atmospheric sulfur budget under volcanically quiescent conditions: Aerosol-chemistry-climate model predictions and validation, *J. Geophys. Res.- Atmos.*, 120, 256–276, <https://doi.org/10.1002/2014JD021985>, 2015.
- Solomon, S.: Stratospheric ozone depletion: A review of concepts and history, *Rev. Geophys.*, 37, 275–316, <https://doi.org/doi:10.1029/1999RG900008>, 1999.
- 630 Stenke, A., Schraner, M., Rozanov, E., Egorova, T., Luo, B., and Peter, T.: The SOCOL version 3.0 chemistry–climate model: description, evaluation, and implications from an advanced transport algorithm, *Geosci. Model Dev.*, 6, 1407–1427, <https://doi.org/10.5194/gmd-6-1407-2013>, 2013.
- Stenke, A., Vattioni, S., Chiodo, G., Luo, B., Sukhodolov, T., and Peter, T.: Simulation data for testing the aerosol microphysical scheme of SOCOL-AERv2, ETH Zurich, <https://doi.org/10.3929/ethz-b-000610854>, 2023.

- 635 Stier, P., Feichter, J., Kinne, S., Kloster, S., Vignati, E., Wilson, J., Ganzeveld, L., Tegen, I., Werner, M., Balkanski, Y., Schulz, M., Boucher, O., Minikin, A., and Petzold, A.: The aerosol-climate model ECHAM5-HAM, *Atmos. Chem. Phys.*, 5, 1125–1156, <https://doi.org/10.5194/acp-5-1125-2005>, 2005.
- Stott, P. A. and Harwood, R. S.: An implicit time-stepping scheme for chemical species in a global atmospheric circulation model, *Ann. Geophys.*, 11, 377–388, 1993.
- 640 Sukhodolov, T., Sheng, J.-X., Feinberg, A., Luo, B.-P., Peter, T., Revell, L., Stenke, A., Weisenstein, D. K., and Rozanov, E.: Stratospheric aerosol evolution after Pinatubo simulated with a coupled size-resolved aerosol–chemistry–climate model, *SOCOL-AERv1.0*, *Geosci. Model Dev.*, 11, 2633–2647, <https://doi.org/10.5194/gmd-11-2633-2018>, 2018.
- Sukhodolov, T., Egorova, T., Stenke, A., Ball, W. T., Brodowsky, C., Chiodo, G., Feinberg, A., Friedel, M., Karagodin-Doyennel, A., Peter, T., Sedlacek, J., Vattioni, S., and Rozanov, E.: Atmosphere–ocean–aerosol–chemistry–climate model *SOCOLv4.0*: description and
- 645 evaluation, *Geosci. Model Dev.*, 14, 5525–5560, <https://doi.org/10.5194/gmd-14-5525-2021>, 2021.
- Sun, H., Eastham, S., and Keith, D.: Developing a Plume-in-Grid Model for Plume Evolution in the Stratosphere, *J. Adv. Model. Earth Syst.*, 14, e2021MS002816, <https://doi.org/10.1029/2021MS002816>, 2022.
- Tabazadeh, A., Toon, O. B., Clegg, S. L., and Hamill, P.: A new parameterization of H₂SO₄/H₂O aerosol composition: Atmospheric implications, *Geophys. Res. Lett.*, 24, 1931–1934, <https://doi.org/10.1029/97GL01879>, 1997.
- 650 Taylor, K. E., Williamson, D., and Zwiers, F.: The sea surface temperature and sea ice concentration boundary conditions for AMIP II simulations, Program for Climate Model Diagnosis and Intercomparison, PCMDI report 60, Lawrence Livermore National Laboratory, <https://pcmdi.llnl.gov/report/pdf/60.pdf?id=32>, (last access: 3 January 2023), 2000.
- Thomason, L. and Peter, T.: SPARC Assessment of Stratospheric Aerosol Properties (ASAP), SPARC Report No. 4, WCRP-124, WMO/TD-No. 1295, Stratospheric Processes And their Role in Climate (SPARC),
- 655 <http://www.sparc-climate.org/publications/sparc-reports/>, (last access: 18 January 2023), 2006.
- Thomason, L. W., Ernest, N., Millán, L., Rieger, L., Bourassa, A., Vernier, J.-P., Manney, G., Luo, B., Arfeuille, F., and Peter, T.: A global space-based stratospheric aerosol climatology: 1979–2016, *Earth Syst. Sci. Data*, 10, 469–492, <https://doi.org/10.5194/essd-10-469-2018>, 2018.
- Tilmes, S., Richter, J. H., Mills, M. J., Kravitz, B., MacMartin, D. G., Garcia, R. R., Kinnison, D. E., Lamarque, J.-F., Tribbia, J., and Vitt, F.: Effects of Different Stratospheric SO₂ Injection Altitudes on Stratospheric Chemistry and Dynamics, *J. Geophys. Res.*, 123, 4654–4673, <https://doi.org/10.1002/2017JD028146>, 2018.
- 660 Tilmes, S., Visioni, D., Jones, A., Haywood, J., Séférian, R., Nabat, P., Boucher, O., Bednarz, E. M., and Niemeier, U.: Stratospheric ozone response to sulfate aerosol and solar dimming climate interventions based on the G6 Geoengineering Model Intercomparison Project (GeoMIP) simulations, *Atmospheric Chemistry and Physics*, 22, 4557–4579, <https://doi.org/10.5194/acp-22-4557-2022>, 2022.
- 665 Timmreck, C., Mann, G. W., Aquila, V., Hommel, R., Lee, L. A., Schmidt, A., Brühl, C., Carn, S., Chin, M., Dhomse, S. S., Diehl, T., English, J. M., Mills, M. J., Neely, R., Sheng, J., Toohey, M., and Weisenstein, D.: The Interactive Stratospheric Aerosol Model Intercomparison Project (ISA-MIP): motivation and experimental design, *Geosci. Model Dev.*, 11, 2581–2608, <https://doi.org/10.5194/gmd-11-2581-2018>, 2018.
- Vattioni, S., Weisenstein, D., Keith, D., Feinberg, A., Peter, T., and Stenke, A.: Exploring accumulation-mode H₂SO₄ versus SO₂
- 670 stratospheric sulfate geoengineering in a sectional aerosol–chemistry–climate model, *Atmos. Chem. Phys.*, 19, 4877–4897, <https://doi.org/10.5194/acp-19-4877-2019>, 2019.

- Vattioni, S., Stenke, A., Chiodo, G., Luo, B., Sukhodolov, T., and Peter, T.: Simulation data for testing the aerosol microphysical scheme of SOCOL-AERv2, ETH Zurich, <https://doi.org/10.3929/ethz-b-000622193>, 2023.
- Vattioni, S., Stenke, A., and Chiodo, G.: SOCOL-AERv2 [Software], <https://doi.org/10.5281/zenodo.10791230>, 2024.
- 675 Vehkamäki, H., Kulmala, M., Napari, I., Lehtinen, K. E. J., Timmreck, C., Noppel, M., and Laaksonen, A.: An improved parameterization for sulfuric acid–water nucleation rates for tropospheric and stratospheric conditions, *J. Geophys. Res.*, 107, AAC 3–1–AAC 3–10, <https://doi.org/10.1029/2002JD002184>, 2002.
- Walcek, C. J.: Minor flux adjustment near mixing ratio extremes for simplified yet highly accurate monotonic calculation of tracer advection, *J. Geophys. Res.*, 105, 9335–9348, <https://doi.org/10.1029/1999JD901142>, 2000.
- 680 Wan, H., Rasch, P. J., Zhang, K., Kazil, J., and Leung, L. R.: Numerical issues associated with compensating and competing processes in climate models: an example from ECHAM-HAM, *Geosci. Model Dev.*, 6, 861–874, <https://doi.org/10.5194/gmd-6-861-2013>, 2013.
- Weisenstein, D. K., Yue, G. K., Ko, M. K., Sze, N., Rodriguez, J. M., and Scott, C. J.: A two-dimensional model of sulfur species and aerosols, *J. Geophys. Res.*, 102, 13 019–13 035, <https://doi.org/10.1029/97JD00901>, 1997.
- Weisenstein, D. K., Penner, J. E., Herzog, M., and Liu, X.: Global 2-d intercomparison of sectional and modal aerosol modules, *Atmos. Chem. Phys.*, 7, 2339–2355, <https://doi.org/10.5194/acp-7-2339-2007>, 2007.
- 685 Weisenstein, D. K., Visionsi, D., Franke, H., Niemeier, U., Vattioni, S., Chiodo, G., Peter, T., and Keith, D. W.: An interactive stratospheric aerosol model intercomparison of solar geoengineering by stratospheric injection of SO₂ or accumulation-mode sulfuric acid aerosols, *Atmos. Chem. Phys.*, 22, 2955–2973, <https://doi.org/10.5194/acp-22-2955-2022>, 2022.
- Wunderlin, E., Chiodo, G., Sukhodolov, T., Vattioni, S., Visionsi, D., and Tilmes, S.: Side Effects of Sulfur-Based Geoengineering Due To Absorptivity of Sulfate Aerosols, *Geophysical Research Letters*, 51, e2023GL107 285, <https://doi.org/10.1029/2023GL107285>, 2024.
- 690 Yu, F., Nadykto, A. B., Luo, G., and Herb, J.: H₂SO₄–H₂O binary and H₂SO₄–H₂O–NH₃ ternary homogeneous and ion-mediated nucleation: lookup tables version 1.0 for 3-D modeling application, *Geosci. Model Dev.*, 13, 2663–2670, <https://doi.org/10.5194/gmd-13-2663-2020>, 2020.
- Yu, F., Luo, G., Nair, A. A., Eastham, S., Williamson, C. J., Kupc, A., and Brock, C. A.: Particle number concentrations and size distributions in the stratosphere: implications of nucleation mechanisms and particle microphysics, *Atmos. Chem. Phys.*, 23, 1863–1877, <https://doi.org/10.5194/acp-23-1863-2023>, 2023.
- 695