

1 **The Regional Climate-Chemistry-Ecology**

2 **Coupling Model RegCM-Chem (v4.6)-YIBs (v1.0):**

3 **Development and Application**

4 Nanhong Xie¹, Tijian Wang^{1*}, Xiaodong Xie², Xu Yue², Filippo Giorgi³, Qian Zhang¹,
5 Danyang Ma¹, Rong Song¹, Beiyao Xu¹, Shu Li¹, Bingliang Zhuang¹, Mengmeng Li¹, Min
6 Xie¹, Natalya Andreeva Kilifarska⁴, Georgi Gadzhev⁵, Reneta Dimitrova⁶

7 ¹School of Atmospheric Sciences, Nanjing University, Nanjing, 210023, China

8 ²School of Environmental Sciences and Engineering, Nanjing University of Information Science and Technology,
9 Nanjing, 210023, China

10 ³Earth System Physics Section, the Abdus Salam International Centre for Theoretical Physics, Trieste, 34100, Italy

11 ⁴Climate, Atmosphere and Waters Research Institute, Bulgarian Academy of Sciences, Sofia, 1113, Bulgaria

12 ⁵National Institute of Geophysics, Geodesy and Geography, Bulgarian Academy of Sciences, Sofia, 1113,
13 Bulgaria

14 ⁶Department of Meteorology and Geophysics, Faculty of Physics, Sofia University, Sofia, 1113, Bulgaria

15 *Corresponding to:* Tijian Wang (tjwang@nju.edu.cn)

16 **Abstract.** The interactions between the terrestrial biosphere, atmospheric chemistry, and climate involve complex
17 feedbacks that have traditionally been modeled separately. We present a new framework that couples the Yale
18 Interactive terrestrial Biosphere (YIBs), a dynamic plant-chemistry model, with the RegCM-Chem model.
19 RegCM-Chem-YIBs integrates meteorological variables and atmospheric chemical composition from
20 RegCM-Chem with land surface parameters from YIBs. The terrestrial carbon flux calculated by YIBs, are fed
21 back into RegCM-Chem interactively, thereby representing the interactions between fine particulate matter
22 (PM_{2.5}), ozone (O₃), and carbon dioxide (CO₂). For testing purposes, we carry out a one-year simulation (2016) at
23 a 30 km horizontal resolution over East Asia with RegCM-Chem-YIBs. The model accurately captures the spa-
24 tio-temporal distribution of climate, chemical composition, and ecological parameters. In particular, the estimated
25 O₃ and PM_{2.5} are consistent with ground observations, with correlation coefficients (R) of 0.74 and 0.65, respec-
26 tively. The simulated CO₂ concentration is consistent with observations from six sites (R ranged from 0.89 to 0.97)
27 and exhibits a similar spatial pattern when compared to carbon assimilation products. RegCM-Chem-YIBs pro-
28 duces reasonably good gross primary productivity (GPP) and net primary productivity (NPP), showing seasonal
29 and spatial distributions consistent with satellite observations, and mean biases (MBs) of 0.13 and 0.05 kg C m⁻²
30 year⁻¹. This study illustrates that the RegCM-Chem-YIBs is a valuable tool to investigate coupled interactions
31 between the terrestrial carbon cycle, atmospheric chemistry, and climate change at a higher resolution in regional
32 scale.

33 **1 Introduction**

34 Air pollution and climate change are major focal points in atmospheric and environmental science (Hong et
35 al., 2019; Kan et al., 2012). In this respect, China exhibits both high air pollution levels and large greenhouse
36 gas emissions (Zheng et al., 2018; Li et al., 2016a). The consequences of China's air pollution on global, region-
37 al, and urban climate are significant (Liu et al., 2022; Lu et al., 2020). Conversely, global warming impacts the
38 dynamics, physics, and chemical mechanisms underlying atmospheric pollutant formation, underscoring a ro-
39 bust link between atmospheric chemistry and climate change (Baklanov et al., 2016; Fiore et al., 2015; Fiore et
40 al., 2012).

41 $PM_{2.5}$, O_3 , and CO_2 are important for regional air pollution and climate. O_3 , a potent pollutant, is harmful
42 for human health and can also harm chloroplasts in plant cells, consequently influencing the carbon assimilation
43 efficiency of land ecosystems (Xie et al., 2019; Ainsworth et al., 2012). Similarly, $PM_{2.5}$ is not only one of the
44 most dangerous pollutants for human health (Kim et al., 2015), but also affects atmospheric radiation mechanics,
45 modulates radiation fluxes reaching vegetation canopies, and hence impacts plant physiological processes and
46 terrestrial carbon fluxes (Lu et al., 2017; Strada and Unger, 2016). Terrestrial ecosystems, absorbing nearly 30%
47 of anthropogenic CO_2 emissions, play an essential role in the global carbon cycle, for which even minor altera-
48 tions can trigger significant oscillations in atmospheric CO_2 concentrations, potentially destabilizing the global
49 climate (Forkel et al., 2016; Ahlstrom et al., 2015). As a result, $PM_{2.5}$, O_3 , and CO_2 exhibit intricate interplays.

50 Models that couple climate and chemistry are vital tools for investigating the interplay between environ-
51 mental pollution and climate warming (Dunne et al., 2020; Yahya et al., 2017), and in particular the direct and
52 indirect influences of aerosols, O_3 , and greenhouse gases on climates at different scales (Chutia et al., 2019; Pu
53 et al., 2017; Li et al., 2017a). For example, the Atmospheric Chemistry and Climate Model Intercomparison
54 Project (ACCMIP) addresses this issue through the use of a range of global coupled climate-chemistry models
55 (Young et al., 2013; Shindell et al., 2013; Lamarque et al., 2013). In fact, China has achieved significant ad-
56 vancements in atmospheric chemistry and coupled climate models during recent years, both at the global and
57 regional scale. Representative models encompass BCC_AGCM2.0_CAM, BCC-AGCM_CUACE2.0,
58 RIEMS-Chem, and RegCCMS.

59 BCC_AGCM2.0_CAM was coupled by the China Meteorological Administration through direct integra-
60 tion of the National Climate Center's atmospheric circulation model (BCC-AGCM) with the Canadian aerosol
61 model (CAM) (Zhang et al., 2012). Atmospheric model BCC-AGCM2.0 was developed by the National Climate

62 Center. For example, at the regional scale the Institute of Atmospheric Physics of the Chinese Academy of Sci-
63 ences, has constructed the Regional Integrated Environmental Modeling System (RIEMS), which is widely used
64 in studies on East Asian regional climate change and severe weather systems (Scheuch et al., 2015; Xiong et al.,
65 2009). It incorporates atmospheric chemistry and aerosol dynamics into the Regional Integrated Environment
66 Modeling System and produces online simulations of meteorological parameters, aerosol chemical composition,
67 optical characteristics, radiation forcing, and aerosol-induced climate feedback (Li et al., 2014; Li et al., 2013a;
68 Han et al., 2012).

69 The Nanjing University developed the Regional Climate Chemistry Modeling System (RegCCMS), a syn-
70 thesis of the regional climate model RegCM2 and the tropospheric atmospheric chemistry model TACM, pri-
71 marily oriented toward investigating the spatio-temporal distribution, radiation forcing, and climatic effects of
72 tropospheric O₃ and sulfate aerosols. Subsequently, RegCM3 was coupled with TACM, integrating modules for
73 aerosols into RegCCMS (Zhang et al., 2014; Li et al., 2009). The system incorporates parameterization schemes
74 facilitating the simulation of aerosols' direct, indirect, and semidirect climatic effects. Extensive evaluations
75 have been carried out regarding major aerosol impacts on the meteorology and regional climate within East Asia
76 (Zhuang et al., 2013; Zhuang et al., 2011; Wang et al., 2010). Subsequently, Shalaby et al. (2012) developed the
77 regional climate-chemistry model RegCM-Chem, by coupling the CBM-Z gas phase chemistry module to ver-
78 sion 4 of the RegCM system, RegCM4 (Giorgi et al., 2012). RegCM-Chem also includes a simplified aerosol
79 scheme including radiatively interactive sulfates, carbonaceous aerosols, sea salt, and desert dust (Zakey et al.,
80 2006; Solmon et al., 2006), and it has been used for a variety of applications in different domains.

81 By developing the regional climate-chemistry-ecology model RegCM-Chem-YIBs, in which the interactive
82 biosphere model YIBs is coupled to RegCM-Chem. The model can produce multi-process simulations of re-
83 gional climate, atmospheric chemistry, and ecology, especially PM_{2.5}, O₃, and CO₂, and their interactions with
84 atmospheric variables (Xu et al., 2023; Ma et al., 2023b; Ma et al., 2023a; Xu et al., 2022; Gao et al., 2022; Xie
85 et al., 2020). Here we expand on these previous studies. We carry out a one-year simulation (2016) at a 30 km
86 horizontal resolution over East Asia with RegCM-Chem-YIBs and conduct a comprehensive assessment. We
87 validate the simulation not only in terms of atmospheric variables but also in terms of atmospheric composition
88 and ecological parameters, by comparison with a range of observations available for this period.


89 The paper is organized as follows. In section 2 we first describe the RegCM-Chem-YIBs system, focusing
90 in particular on the newly implemented coupling with the ecological component. We also describe the observa-

91 tion datasets used in the model assessment. The simulations are then analyzed in section 3, while section 4 pre-
 92 sents our conclusions and a general discussion of our results and future developments.

93 **2 Model and Methods**

94 **2.1 Overall Framework**

95 In RegCM-Chem-YIBs, the atmospheric variables produced by RegCM (temperature, humidity, precipita-
 96 tion, radiation, etc.) and atmospheric chemical compounds, such as O₃ and PM_{2.5}, produced by the chemis-
 97 try/aerosol module are input into YIBs, which simulates the physiological processes of vegetation (such as pho-
 98 tosynthesis, respiration, etc.), and calculates land process variables such as CO₂ fluxes, BVOC emissions, and
 99 stomatal conductance. The output from YIBs is then fed back to RegCM-Chem, which adjusts the CO₂, O₃, and
 100 PM_{2.5} concentrations and their radiative and microphysical effects on the meteorological fields in the lower at-
 101 mosphere, thereby achieving a full coupling between climate, chemistry, and ecology. Figure 1 shows the basic
 102 framework of the RegCM-Chem-YIBs coupled model.



103
 104 **Figure 1.** RegCM-Chem-YIBs Coupling Model Framework

105 **2.2 Descriptions of the RegCM-Chem model**

106 The inception of the RegCM system traces back to the late 1980s and early 1990s, when NCAR's (U.S. Na-
 107 tional Center for Atmospheric Research) RegCM 1 was first developed for climate downscaling (Giorgi, 1990;

108 Giorgi and Bates, 1989; Dickinson et al., 1989). After a series of developments, subsequent versions were in-
109 troduced, such as RegCM2 (Giorgi et al., 1993), RegCM2.5 (Giorgi and Mearns, 1999), RegCM3 (Pal et al.,
110 2007), RegCM4 (Giorgi et al., 2012). The RegCM system presently managed, maintained, and expanded by the
111 Earth System Physics (ESP) section of the Abdus Salam International Center for Theoretical Physics (ICTP), is
112 open-source and extensively employed in regional climate studies, contributing to the establishment of a com-
113 prehensive Regional Climate Research Network (RegCNET) (Giorgi et al., 2006). The model can be applied to
114 all regions of the globe (Giorgi et al., 2012) and is moving into a fully-coupled regional Earth system model
115 framework through coupling with the ocean (Turuncoglu et al., 2013; Artale et al., 2010), lake (Small et al.,
116 1999), aerosol (Solmon et al., 2006), dust (Zakey et al., 2006), chemistry (Shalaby et al., 2012), hydrology
117 (Coppola et al., 2003), land surface processes (Oleson et al., 2008). Of specific interest for our study, Shalaby et
118 al. (2012) added a radiatively interactive gas-phase chemical module (CBM-Z) to RegCM4, generating
119 RegCM-Chem, in which atmosphere physics and chemistry are fully coupled.

120 **2.2.1 Aerosol Mechanisms**

121 The RegCM model integrates a simplified aerosol framework, enabling the simulation of sulfate, black
122 carbon (BC), organic carbon (OC), sea salt, and desert dust. The model specifies an external mix of aerosols and
123 accounts for the influence of horizontal advection, turbulent diffusion, vertical transport, emissions, dry and wet
124 deposition, and gas-liquid transition on aerosol concentration (Solmon et al., 2012; Giorgi et al., 2012; Zakey et
125 al., 2006). The secondary organic aerosol scheme VBS (volatile basis set) has also been introduced into the
126 model to further improve RegCM-Chem's simulation of tropospheric aerosols (Yin et al., 2015). The model in-
127 corporates the ISORROPIA thermodynamic equilibrium scheme to describe the formation process of secondary
128 inorganic salts, thus enhancing the model's capability to simulate secondary inorganic aerosols (Li et al., 2016b).
129 The further addition of bioaerosols was carried out by Liu (Liu et al., 2016).

130 **2.2.2 Gas phase chemical mechanism**

131 RegCM4-Chem includes the CBM-Z (Carbon Bond Mechanism-Z) atmospheric chemistry mechanism
132 (Zaveri and Peters, 1999). The CBM-IV mechanism, recognized for its widespread use, serves as the basis for
133 CBM-Z (Gery et al., 1989) and was developed to balance simulation accuracy and computational speed. Both
134 CBM-IV and CBM-Z categorize volatile organic compounds (VOCs) into groups dependent on their carbon

135 bond formation and use lumped species to represent each group. However, CBM-Z includes additional species
136 and reactions compared to CBM-IV, which are crucial for simulating typical urban environments and long-term
137 simulations at regional to global scales. Enhancements in CBM-Z include (1) specific representation of stable
138 alkanes; (2) updated parameters for higher alkanes; (3) separation of olefins into two categories based on differ-
139 ing reactions; (4) addition of peroxy alkane self-reactions significant in low-NO_x, such as remote regions; (5)
140 incorporation of reactions among alkanes, peroxyacyl radicals, and NO₃, which are crucial nocturnally; (6) in-
141 clusion of long-lived organic nitrates and peroxides; and (7) refinement of isoprene and its peroxy radical chem-
142 istry. Collectively, these updates to the CBM-Z chemistry mechanism enhance the model's ability to more accu-
143 rately simulate long-lived VOCs and address the atmospheric chemistry transition from urban to rural settings.

144 **2.2.3 Radiation scheme**

145 RegCM4 adopts the CCM3 radiation scheme, which uses the delta-Eddington approximation for solar
146 spectral radiation and accounts for the attenuation effect of atmospheric components such as O₃, H₂O, CO₂, O₂
147 on solar radiation (Kiehl et al., 1996). The CCM3 radiation scheme, implemented in RegCM4, extends from 0.2
148 to 5 μm, and is segmented into 18 bands. It uses the cloud scattering and absorption parameter scheme, and
149 cloud optical characteristics. As cumulus clouds form, the cloud optical characteristics stretch from the cloud
150 base up to the cloud top, and the radiation calculations assume random overlap. It is assumed in the model that
151 the cloud thickness is equivalent to that of the model's vertical layers, with distinctive cloud water and ice
152 contents assigned to high, middle, and low clouds (Slingo, 1989).

153 **2.2.4 Photolysis rate**

154 Meteorological conditions and chemical input fields determine the photolysis rate, with most variables
155 dynamically produced by the RegCM's modules and updated every 3-30 minutes. SO₂ and NO_x, inverted from
156 the US standard atmosphere's vertical profile, are model-defined. Owing to the computational demands of
157 precise photolysis rates from the Tropospheric Ultraviolet-Visible Model (TUV) method (Madronich and
158 Flocke, 1998) and eight data stream spherical harmonics discretization, a look-up table and interpolation method
159 are adopted. Considering the significant impact of clouds on the photolysis rate, it becomes crucial to adjust the
160 cloud amount. Here we use the cloud optical depth information for each grid cell within the model. As the
161 absorption and scattering of ultraviolet radiation by clouds reduce the photolysis rate inside and below the cloud

162 while enhancing it above the cloud, the correction value for the photolysis rate under clear sky conditions de-
163 pends on the position to the cloud layer. Hence, cloud height and optical depth are necessary for the photolysis
164 rate computation (Chang et al., 1987).

165 **2.2.5 Deposition Processes**

166 In the model, dry deposition serves as the principal removal process for trace gases, with the deposition
167 velocity being determined by three categories of resistance: aerodynamic, quasi-laminar sublayer, and surface
168 resistance, encompassing soil and vegetation absorption. The latter is inclusive of both stomatal and nonstomatal
169 absorption. The dry deposition module, taken from the CLM4 surface scheme, covers 29 gas-phase species and
170 comprises 11 types of land cover. To enhance the accuracy of the daily variation in dry deposition simulation,
171 both stomatal and nonstomatal resistances are accounted for in the dry deposition scheme. The calculation of all
172 deposition resistances is performed within the CLM land surface model (Wesely, 1989). Wet deposition uses the
173 MOZART global model's wet deposition parameterization scheme (Emmons et al., 2010; Horowitz et al., 2003),
174 including 26 gas-phase species in CBM-Z, and the wet deposition amount is based on the simulated precipita-
175 tion.

176 **2.3 Descriptions of the YIBs model**

177 The YIBs model, pioneered by Yale University, integrates plant physiological mechanisms to simulate how
178 photosynthesis, respiration, and other physiological processes respond to environmental drivers such as radia-
179 tion, temperature, and moisture. Moreover, YIBs simulates the carbon cycle both regionally and globally (Yue
180 and Unger, 2015). For example, its simulation of terrestrial carbon flux closely matches ground flux observa-
181 tions and satellite-derived data in diverse geographical areas such as the United States and China (Yue and
182 Unger, 2017; Yue et al., 2017).

183 **2.3.1 The main processes in YIBs**

184 In the YIBs model, eight distinct Plant Functional Types (PFTs) are incorporated, encompassing evergreen
185 coniferous forest, evergreen broad-leaved forest, deciduous broad-leaved forest, shrub forest, tundra, C3 grass-
186 land, C4 grasslands, and crops. The model employs the Michaelis–Menten enzyme-kinetics scheme for simulat-
187 ing plant photosynthesis (Farquhar et al., 1980), and the total photosynthesis (A_{tot}) of leaves is affected by Ru-

188 bisco enzyme activity (J_e), electron transfer rate (J_e), and photosynthetic product (triose phosphate) transport
189 capacity (J_s) limitation.

190 2.3.2 Canopy Radiation Scheme

191 A multilayer canopy radiation transmission scheme is adopted in YIBs for canopy radiation transmission
192 (Spitters et al., 1986), consisting of a radiation transfer model based on the total leaf area index, extinction coef-
193 ficient, and vegetation height. The entire vegetation canopy is usually divided into 2 to 16 layers, and the spe-
194 cific number of layers can be automatically adjusted according to the height of the canopy.

195 2.3.3 Biogenic Volatile Organic Compound Emission Scheme

196 Differently from the traditional MEGAN scheme, the YIBs model applies a biogenic volatile organic com-
197 pound (BVOC) emission scheme on a leaf scale, which is better suited to describe the photosynthesis process in
198 vegetation (Guenther et al., 1995). This introduces an effect of plant photosynthesis on BVOC emissions which
199 is more closely related to the real physiological process of vegetation. BVOC emissions from leaves to the can-
200 opy are integrated to obtain total canopy emissions. The intensity of leaf BVOC emission depends on the rate of
201 photosynthesis J_e under electron transfer rate limitation, leaf surface temperature, and intracellular CO_2 con-
202 centration (Yue and Unger, 2015):

$$203 I = J_e \cdot \beta \cdot \kappa \cdot \tau \cdot \varepsilon \quad (1)$$

204 where I is the intensity of leaf BVOC emission in units of $\mu\text{mol m}^{-2}[\text{leaf}] \text{ s}^{-1}$, J_e is the electron
205 transport-limited photosynthesis rate, the calculation formula is as follows:

$$206 J_e = a_{leaf} \cdot PAR \cdot \alpha_{qe} \cdot \frac{C_i - \Gamma^*}{C_i - 2\Gamma^*} \quad (2)$$

207 where a_{leaf} is the leaf-specific light absorbance, PAR is photosynthetically active radiation, α_{qe} is the intrin-
208 sic quantum efficiency for photosynthetic CO_2 uptake in the chlorophyll reaction system, C_i is the internal leaf
209 CO_2 concentration, Γ^* is the CO_2 concentration compensation point in the absence of non-photorespiratory
210 respiration (Collatz et al., 1991).

211 β is the coefficient for converting electron transfer flux into BVOC emissions (Niinemets et al., 1999;
212 Pacifico et al., 2011):

$$213 \beta = \frac{C_i - \Gamma^*}{\delta(4.67C_i + 9.33\Gamma^*)} \quad (3)$$

214 where κ is related to the internal leaf CO_2 concentration:

Formatted: Font: Times New Roman

Formatted

Formatted: Font: Times New Roman, Not Italic

Formatted: Indent: First line: 0 ch

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted

Formatted

Formatted

Formatted

Formatted: Font: 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: 10 pt

Formatted

Formatted

Formatted

215

$$\kappa = \frac{C_{i_standard}}{C_a} \quad (4)$$

216 where $C_{i_standard}$ is the internal leaf CO_2 concentration under standard conditions (when atmospheric CO_2 is
 217 370 ppm). The τ term reflects the response of BVOC emission intensity to temperature:

$$\tau = \exp[0.1(T - T_{ref})] \quad (5)$$

218 where T is the blade surface temperature. T_{ref} is the standard temperature (30 °C). When the blade temperature
 219 is 40 °C, the BVOC emission intensity is maximum. As the temperature further rises, the BVOC emission grad-
 220 ually weakens. In reality, such high temperatures are relatively rare and may only occur under extremely dry
 221 climate conditions.

223 2.3.4 Ozone Damage Protocol

224 When tropospheric ozone enters plants through stomata, it can directly damage plant cell tissues, thereby
 225 slowing the photosynthesis rate and further weakening the carbon sequestration capacity of vegetation. The
 226 YIBs model incorporates the semi-mechanistic parameterization scheme to delineate ozone's effect on plants
 227 (Sitch et al., 2007):

$$A = A_{tot} \cdot F \quad (6)$$

229 where A is photosynthesis minus the influence of ozone, A_{tot} is the total photosynthesis of leaves, F is the
 230 proportion of photosynthesis minus the influence of ozone, which depends on the ozone flux from the stomata
 231 into the vegetation that exceeds the threshold.

$$F = 1 - a \cdot \max[(F_{ozn} - F_{ozncrit})_0] \quad (7)$$

233 where a is the sensitivity parameter of vegetation to ozone obtained based on observation data. $F_{ozncrit}$ repre-
 234 sents the threshold corresponding to the damage caused by ozone to vegetation, F_{ozn} represents the flux of
 235 ozone entering the page through the stomata:

$$F_{ozn} = \frac{[O_3]}{r_b + \frac{\kappa_{O_3}}{r_s}} \quad (8)$$

237 where $[O_3]$ is the ozone concentration at the top of the canopy, r_b is the boundary layer resistance, κ_{O_3} is the
 238 ratio of O_3 leaf resistance to water vapor blade resistance, r_s is the stomatal resistance considering the influence
 239 of ozone:

$$r_s = g_s \cdot F \quad (9)$$

241 g_s is the leaf conductance without O_3 effects. The set of equations (7), (8) and (9) yields a quadratic term in F

- Formatted: Font: 10 pt
- Formatted: Font: 10 pt
- Formatted: Font: 10 pt
- Formatted: Font: 10 pt
- Formatted: Font: 10 pt
- Formatted: Font: 10 pt
- Formatted: Font: (Default) Times New Roman, 10 pt, Not Italic
- Formatted: Font: 10 pt
- Formatted: Font: (Default) Times New Roman, 10 pt, Italic
- Formatted: Font: (Default) Times New Roman, 10 pt
- Formatted: Font: 10 pt
- Formatted: Font: 10 pt
- Formatted: Font: (Default) Times New Roman, 10 pt
- Formatted: Font: (Default) Times New Roman, 10 pt, Subscript
- Formatted: Font: (Default) Times New Roman, 10 pt
- Formatted: Font: (Default) Times New Roman, 10 pt, Subscript
- Formatted: Font: (Default) Times New Roman, 10 pt
- Formatted: Font: 10 pt
- Formatted: Font: (Default) Times New Roman, 10 pt
- Formatted: Font: 10 pt
- Formatted: Font: 10 pt
- Formatted: Font: 10 pt
- Formatted: Font: 10 pt
- Formatted: Font: (Default) Times New Roman, 10 pt, Not Italic
- Formatted: Font: 10 pt
- Formatted: Font: (Default) Times New Roman, 10 pt
- Formatted: Font: 10 pt
- Formatted: Font: (Default) Times New Roman, 10 pt
- Formatted: Font: Times New Roman, Not Italic
- Formatted: Font: Times New Roman, Not Italic
- Formatted: Font: Times New Roman, Not Italic
- Formatted: Font: Times New Roman, Not Italic
- Formatted: Subscript
- Formatted: Subscript
- Formatted: Font: Times New Roman, Not Italic
- Formatted: Indent: First line: 0 ch
- Formatted: Subscript
- Formatted: Font: (Default) Times New Roman, 10 pt

242 that can be solved analytically.

243 **2.4 Descriptions of the RegCM-Chem-YIBs model**

244 **2.4.1 Coupling between RegCM-Chem and YIBs**

245 The integrated RegCM-Chem-YIBs model, an enhancement to the original RegCM-Chem, introduces CO₂
246 as an atmospheric constituent, incorporating its source-sink dynamics, transport, and diffusion processes. At-
247 mospheric CO₂ concentration is primarily influenced by atmosphere-ocean CO₂ exchange flux, biomass com-
248 bustion emissions, fossil fuel emissions, and terrestrial ecosystem CO₂ flux. The model prescribes fossil fuel
249 emissions, biomass combustion emissions, and atmosphere-ocean CO₂ fluxes, while the terrestrial ecosystem
250 CO₂ fluxes are computed in real time via the coupled YIBs terrestrial ecosystem model.

251 Within the coupled model system, meteorological variables (including temperature, humidity, precipitation,
252 radiation, etc.) and atmospheric pollutant concentrations (O₃ and PM_{2.5}) generated by RegCM-Chem are incor-
253 porated into the YIBs model every six-minute intervals. This integration step is to be consistent with the integra-
254 tion time step of the chemistry module, thus maintaining synchronization between modules. Considering the
255 complexity of chemical reactions and ecological processes, dynamic adjustments at short intervals enable the
256 model to better capture transient interactions between ecology and the atmosphere. The choice of this adjust-
257 ment frequency balances the representation of actual processes with computational efficiency, ensuring that
258 simulation results are both accurate and efficient. YIBs then simulates vegetation physiological processes such
259 as photosynthesis and respiration, computing land surface parameters including CO₂ flux, BVOC, and stomatal
260 conductance. These outputs from the YIBs are subsequently integrated back into the RegCM-Chem model every
261 six-minute intervals, the intricacies of this integration process lead to significant changes in various environ-
262 mental parameters. The major direct changes, prominently influencing the model's behavior, arise from altera-
263 tions in CO₂ concentration. These changes are directly attributed to intricate physiological processes within the
264 vegetation, including photosynthesis and respiration. The fluxes of CO₂ through these biological processes play
265 a pivotal role in shaping the atmospheric composition. On the indirect front, the integration of YIBs outputs in-
266 duces intricate variations in PM_{2.5} and O₃ concentrations. These indirect changes are primarily orchestrated by
267 shifts in BVOC emissions. The dynamic nature of these emissions contributes to the complexity of atmospheric
268 chemistry, influencing the levels of PM_{2.5} and O₃. Simultaneously, the integration process plays a crucial role in
269 shaping the temporal variations of atmospheric temperature, humidity, and circulation. These changes over time

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

270 are intricately linked to variations in land surface parameters. The interplay of these variables illustrates the dy-
271 namic feedback loops between climate, chemical composition, and ecological processes within the integrated
272 model system modulating atmospheric composition (CO_2 , O_3 , and $\text{PM}_{2.5}$) and atmospheric variables (atmos-
273 pheric temperature, humidity, and circulation), thereby describing the interplay of climate, chemical, and eco-
274 logical processes.

275 2.4.2 Model input data

276 The input data of RegCM-Chem-YIBs mainly includes four categories: surface data, initial boundary data,
277 anthropogenic emission data and CO_2 surface flux data, which are detailed below.

278 (1) Surface data include surface vegetation cover type, terrain, and leaf area index. Land cover type infor-
279 mation is obtained from the MODIS and AVHRR satellites, employing the classification scheme suggested by
280 Lawrence and Chase (Lawrence and Chase, 2007), which uses MODIS data to preliminarily distinguish forest,
281 grassland, bare soil, etc., and combine this with AVHRR data to make a detailed forest classification. The dataset
282 contains a total of 16 different vegetation functional types. To align with the classification conventions of the
283 YIBs model, the original 16 vegetation functional types were converted into the corresponding 8 types recog-
284 nized by the YIBs model. The results are shown in Figure S1.

285 (2) Initial and boundary data include initial and boundary conditions of meteorological variables and at-
286 mospheric chemical composition. Here we use ERA-Interim reanalysis meteorological data, a product from the
287 European Center for Medium-Range Weather Forecasts (ECMWF) created through four-dimensional variational
288 assimilation. The data is on 37 vertical levels, with a horizontal resolution of $0.125^\circ \times 0.125^\circ$, and time resolution
289 of 6 hours. Data for Sea Surface Temperature (SST) is provided by the weekly averaged Optimum Interpolation
290 SST product (OI_WK) of the National Oceanic and Atmospheric Administration (NOAA) (Reynolds et al.,
291 2002). The initial and boundary conditions of atmospheric chemical components (e.g. O_3), come from simula-
292 tions carried out with the global chemistry model MOZART (Emmons et al., 2010; Horowitz et al., 2003). In
293 addition, the initial and boundary conditions for CO_2 species come from the CarbonTracker global carbon as-
294 similation system (Peters et al., 2007) developed by NOAA Earth System Research Laboratory ESRL (Earth
295 System Research Laboratory), which uses the ensemble Kalman filter algorithm to assimilate ESRL greenhouse
296 gas observations and CO_2 observation data provided by the network of collaborating institutions worldwide. The
297 assimilated data includes not only conventional fixed-site observations but also mobile monitoring data such as

298 aircraft and ships. Since 2007, yearly updated carbon assimilation products are provided by CarbonTracker, de-
299 livering global CO₂ three-dimensional concentration data products every three hours. In this study, we utilized
300 the CT2019 product, updated in 2019, spanning a period from January 1, 2000 to March 29, 2019.

301 (3) Anthropogenic emission data include precursors of ozone and particulate matter such as NO_x, VOC,
302 BC, OC, etc. The MIX Asian anthropogenic emission inventory developed by the Tsinghua University is used
303 (Li et al., 2017b), which integrates the results of the emission inventories of various regions in Asia. The emis-
304 sions in China come from China's multi-scale emission inventory MEIC (Multi-resolution Emission Inventory
305 for China) and the high-resolution NH₃ emission inventory developed by Peking University. The anthropogenic
306 emissions in India come from the Indian local emission inventory developed by ANL (Argonne National Labor-
307 atory), while the anthropogenic emissions in South Korea come from the CAPSS (The Korean local emission
308 inventory developed by the Policy Support System), and the man-made emissions in other regions are provided
309 by the REAS (Regional Emission inventory in Asia) emission inventory version 2.1. The anthropogenic emis-
310 sions of major pollutants in the simulated area are shown in Figure S2.

311 (4) Data pertaining to fossil fuel CO₂ emissions are sourced from the MIX Asian anthropogenic emission
312 inventory with a monthly time resolution. CO₂ emissions resulting from biomass burning are derived from the
313 FINN (Fire Inventory from NCAR) inventory (Wiedinmyer et al., 2011) developed by the National Center for
314 Atmospheric Research. The FINN inventory has a daily time resolution. The model's ocean-atmosphere CO₂
315 exchange flux is obtained from the carbon flux product of the CarbonTracker assimilation system, constructed
316 with the global atmospheric transport model TM5 and assimilating CO₂ observation data via an ensemble Kal-
317 man filter algorithm. This provides global 1°×1° resolution CO₂ exchange flux data between the ocean and the
318 atmosphere updated every three hours. The emissions are detailed in Figure S3.

319 **3 Model Application**

320 **3.1 Model setup**

321 To evaluate the performance of RegCM-Chem-YIBs we carried out a one-year simulation starting from
322 December 1st, 2015, through December 31st, 2016. The initial month is used as spin-up period, and thus it is not
323 included in the analysis. The simulation domain is centered at 36°N, 107°E, and covers a considerable part of
324 East Asia, including China, Japan, the Korean Peninsula, and Mongolia, along with significant parts of India and

325 Southeast Asia (Figure S4). The horizontal grid spacing is 30 km and we use 14 levels in the vertical, reaching
 326 up to 50 hPa. Section 2.4.2 provides a comprehensive description of the model input data.

327 3.2 Climate simulations in East Asian

328 Given the importance of the climate for the East Asia region, we first present an assessment of the simula-
 329 tion for the climate 2016 by comparison with the ERA-Interim data. The simulated temperature, specific
 330 humidity, and wind fields at varying altitudes and seasons compared well with the reanalyzed data (Figure S5~
 331 Figure S9), especially temperature and specific humidity, while a tendency to overestimate wind speed is
 332 observed at the near surface and 850 hPa levels. The fields at 500 hPa show very close agreement with
 333 reanalysis data, indicating a strong mid-atmosphere forcing by the boundary conditions, while the simulated
 334 circulation patterns near the surface and at 850 hPa in summer tend to deviate more from the driving reanalysis.
 335 The simulated circulation patterns in the other seasons are basically consistent with the reanalysis data.

336 We first calculated the daily average of the meteorological variables, such as temperature, wind speed, and
 337 specific humidity, from the model simulation and reanalysis data, respectively. Then we calculate the
 338 corresponding statistical indicator correlation coefficient (R), mean deviation (MB), and root mean square error
 339 (RMSE) based on the daily averages. Table 1 reports a number of statistical metrics of comparison between
 340 simulated and reanalysis meteorological variables at different heights. Correlation coefficients (R) range from
 341 0.95 to 0.98 for temperature, 0.71 to 0.97 for longitudinal wind, 0.81 to 0.92 for latitudinal wind, and 0.91-0.92
 342 for specific humidity, indicating a general good consistency between model and driving data, in line with previ-
 343 ous studies (Zhuang et al., 2018; Zhou et al., 2014; Wang et al., 2010).

344 **Table 1.** Statistical indicators for comparison between model simulation results and reanalysis data

Heights	Statistical index	Air Temperature(K)	Longitudinal wind (m/s)	Latitudinal wind (m/s)	Specific humidity (kg kg ⁻¹)
500 hpa	R	0.98	0.97	0.92	0.91
	MB	0.15	0.35	-0.03	0.00015
	RMSE	0.93	0.75	0.51	0.00019
850 hpa	R	0.96	0.77	0.85	0.94
	MB	-0.98	0.38	0.15	-0.00066
	RMSE	1.1	1.08	0.59	0.00077

Near sur- face	R	0.95	0.71	0.81	0.92
	MB	-1.21	0.33	0.23	-0.00098
	RMSE	1.35	0.59	0.54	0.00112

345 (Correlation coefficients (R), mean biases (MB), and root mean square error (RMSE))


346 The magnitude of surface radiation flux directly determines the rates of photosynthesis in vegetation. For
 347 verification purposes, model surface solar fluxes were compared with data on solar energy at the surface
 348 retrieved from the Clouds and the Earth's Radiant Energy System (CERES) satellite, which has a $1^\circ \times 1^\circ$
 349 horizontal and monthly temporal resolution. Figure S10 shows the simulated surface net shortwave radiation in
 350 different seasons and comparison with observational data. The model tends to overestimate surface net
 351 shortwave radiation in spring and winter over India and summer over North China (Yin et al., 2014). Overall,
 352 the simulated surface net shortwave radiation agrees well with the CERES satellite retrieval results, capturing
 353 the spatial distribution and seasonal fluctuation patterns of surface shortwave radiation. The simulation findings
 354 from our study are consistent with earlier research regarding surface net shortwave radiation (Han et al., 2016).

355 In conclusion, RegCM-Chem-YIBs demonstrates a good performance in simulating the climatological
 356 features of the East Asia atmospheric circulations, effectively reproducing the spatial distribution and seasonal
 357 variations of temperature, specific humidity, and radiation.

358 3.3 Simulations of PM_{2.5}, O₃ and CO₂

359 In this section, we compare simulated PM_{2.5} and O₃ concentrations against observational data from 366
 360 stations provided by the China National Environmental Monitoring Center. The geographical distribution of the
 361 simulated annual mean near-surface daily PM_{2.5} and maximum daily 8-hour average (MDA8) O₃ concentration,
 362 along with the observed values, are shown in Figure 2. Supplementary Figure S11 then compares in a scat-
 363 ter-plot format the observation and simulation results. Both figures demonstrate that the model reproduces the
 364 spatial distribution patterns of PM_{2.5} and O₃, with a significant agreement between modeled and measured
 365 values across all stations. The statistical indicators of simulated and measured surface PM_{2.5} and O₃ levels are
 366 shown in Table S1, showing a correlation between simulation and observations of O₃ and PM_{2.5} of 0.74 and 0.65,
 367 respectively. The simulated O₃ concentrations are generally lower than observed in the Fenwei Plain of China, a
 368 discrepancy possibly attributable to uncertainties in the emission inventory for this region. In summary, the
 369 RegCM-Chem-YIBs model demonstrates a good ability to capture the spatial distribution of observed

370 near-surface ozone and particulate matter, especially in highly polluted areas.



371
372 **Figure 2.** Simulation and observation comparison of (a, b) O₃ and (c, d) PM_{2.5} and their differences (e, f) in China.
373 The differences are simulation minus observation. The colored circles in the figure represent station observations.
374 Units: µg m⁻³.


375 Measured and calculated monthly mean CO₂ concentrations at six observation stations in East Asia from
376 the World data Center for Greenhouse Gases are shown in Figure 3. Information on the six sites is listed in Table
377 2. The simulated CO₂ concentration agrees well with observations, with correlation coefficients ranging from
378 0.89 to 0.97. However, in urban and coastal areas, the model performance deteriorates likely due to local emis-
379 sion fluctuations and errors in biogenic fluxes. Nevertheless, the model overall captures the seasonal variations
380 in CO₂ concentrations (Figure 3). This result likely stems from the complex relationship between biogenic and

381 fossil fuel emissions, which are known contributors to observed seasonal CO₂ patterns (Kou et al., 2015). A high
 382 CO₂ mixing ratio (412.3 ppm) is observed at the TAP site, which is associated with strong local emissions. Fur-
 383 ther analysis into the specific sources contributing to elevated CO₂ levels would provide valuable insights into
 384 localized patterns of emissions and their effects on regional carbon cycle processes. The model's ability to re-
 385 produce the geographical and seasonal CO₂ patterns serves as an illustration of its ability to capture the main
 386 processes driving CO₂ dynamics. In summary, while discrepancies in urban or coastal areas highlight the chal-
 387 lenges associated with capturing localized CO₂ dynamics, the model's overall performance and ability to repro-
 388 duce geographical and seasonal CO₂ patterns demonstrates its usefulness in studying CO₂ dynamics at a regional
 389 scale.

390 **Table 2.** Information on six CO₂ stations in East Asia and statistical indicators of observed and modeled CO₂.

Sites	Latitude	Longitude	Elevation	Observations	Simulations	R	RMSE
				(ppm)	(ppm)		
WLG	36.29	100.90	3810	404.3	402.9	0.94	1.75
TAP	36.72	126.12	20	412.3	414.8	0.97	2.70
UUM	44.45	111.08	992	405.7	403.7	0.96	2.66
LLN	23.46	120.86	2867	406.0	407.2	0.93	1.63
YON	24.47	123.02	30	407.1	407.4	0.89	2.80
HK	22.31	114.17	65	407.9	409.7	0.92	15.67

391 (Correlation coefficients (R) and root mean square error (RMSE))




392
 393 **Figure 3.** Modeled (blue) and observed (black) monthly mean CO₂ concentrations validated at six sites in East
 394 Asia. Units: ppm.

395 The limitations of ground-based CO₂ observation stations, particularly their sparse spatial distribution, pose
 396 challenges in obtaining high-resolution CO₂ data. To offset this limitation, data assimilation methods have been
 397 implemented to ensure a coherent global distribution of atmospheric CO₂, effectively filling the void left by
 398 sparse ground-based observations. Here we utilize the Carbon Tracker global carbon assimilation system
 399 developed by the NOAA Earth System Research Laboratory (ESRL) to validate the simulated CO₂
 400 concentrations (Peters et al., 2007). This comparison for the year 2016 is shown in Figure 4. The simulated CO₂
 401 concentrations tend to be lower than observed in Northeastern India and Northeastern China, while they show a
 402 better agreement with observations in other regions. These discrepancies can be traced back to factors such as
 403 the underestimation of localized CO₂ emissions along with the effects of complex topography and circulation
 404 patterns. However, the closer agreement in other regions suggests that the model effectively captures the
 405 primary processes driving CO₂ concentrations.

406 Seasonal variations in the spatial distribution of CO₂ concentrations for 2016 are illustrated in supplemen-
 407 tary Figure S12. The simulations show marked seasonal variations, with elevated concentrations in spring,
 408 autumn, and lower values during summer. In northern regions, including Russia, Mongolia, and Northeast China,
 409 the lowest near-surface CO₂ concentrations occur in summer. This pattern can be attributed to the enhanced

410 photosynthetic activity of terrestrial vegetation in summer, leading to enhanced atmospheric CO₂ sequestration.
 411 Conversely, winter months are characterized by lower solar radiation fluxes and reduced vegetation
 412 photosynthesis, resulting in relatively higher CO₂ concentrations. In specific regions, notably the eastern coastal
 413 zones of China and South Korea, the seasonal pattern of CO₂ concentration is reduced, likely because of the
 414 high levels of urbanization, dense population, and intense anthropogenic emissions in these areas. In contrast,
 415 regions such as Yunnan, the southern side of the Qinghai-Tibet Plateau, and Southeast Asia exhibit consistently
 416 low CO₂ concentrations during summer because of significant vegetation sinks in these densely vegetated areas.
 417 An increase in CO₂ concentrations can be observed over these regions during spring due to local forest fires and
 418 straw-burning processes, which release substantial amounts of CO₂ into the atmosphere (Chuang et al., 2014).



419 **Figure 4.** Evaluation of simulated CO₂ (a) using Carbon Tracker products (b) and their difference (c) in 2016. The
 420 differences are simulation minus observation. Units: ppm.
 421

422 3.4 Simulations of carbon fluxes in terrestrial systems

423 Our assessment of GPP and NPP uses the MOD17A3 Collection 6, a global product originating from
 424 MODIS satellite observations. GPP data include 8-day values with a resolution of 500 meters, as produced in

425 MOD17A2H Version 6 based on radiation use efficiency theory. Such data can be used as input to computations
426 of terrestrial carbon and energy flows, water cycling processes, and vegetation biogeochemistry. Moreover, the
427 MOD17A3H Version 6 product provides information on annual NPP, also on a resolution of 500 meters. All
428 8-day Net Photosynthesis (PSN) products (MOD17A2H) from a particular year are combined to derive annual
429 NPP values (He et al., 2018; Madani et al., 2014; Running, 2012).


430 Figure 5 (a, b, e) shows the geographical distribution of the mean GPP in 2016 from the model simulations
431 and MODIS products. RegCM-Chem-YIBs effectively captures the observed spatial GPP features, with high
432 values mostly over Southwest, Central, and Southeastern China, areas characterized by deciduous broad-leaf
433 and evergreen coniferous forests (Figure S1). The annual average GPP simulated by RegCM-Chem-YIBs is
434 higher than observed over Southwest and Central China by 6.8% and 12.7%, respectively. The annual average
435 simulated GPP over China is 6.18 Pg C yr⁻¹, which is about 7.56% higher than the GPP in MODIS.

436 Figure 6 (a) and Table S2 show the scatter plots of the simulated annual average GPP on each model grid
437 point compared with MODIS. A correlation coefficient of 0.91 and root mean square error of 0.4 kg C m⁻² yr⁻¹ is
438 found, reflecting an overall good simulation by the model. Compared with the results obtained from the global
439 model NASA ModelE2-YIBs (Yue and Unger, 2017), the GPP value estimated here compares better with the
440 MODIS product, which may also be attributed to the higher spatial resolution of the regional system. Moreover,
441 our GPP results are also in line with earlier findings, such as from Li (Li et al., 2013b) who estimated an annual
442 average GPP over China of 6.04 Pg C yr⁻¹ based on the light energy utilization model EC-LUE.


443 Figure 5 (c, d, f) shows the spatial distribution of mean NPP for both the simulations and MODIS
444 products in 2016. NPP, similarly to GPP, exhibits a gradual reduction from southeast to northwest China. The
445 scatter plot comparing the simulated and MODIS annual average NPP across the model grid is illustrated in
446 Figure 6 (b). According to Table S2, a correlation coefficient of 0.87 is found between the simulated and
447 MODIS NPP, with a root mean square error of 0.22 kg C m⁻² yr⁻¹. Notably, the simulated NPP shows a distinct
448 underestimation over regions with higher NPP values. Compared with the MODIS NPP data products, the
449 annual average NPP simulated for the entire China region in 2016 is overestimated by approximately 8.64%,
450 mostly because of the model overestimate in Central China (16.6%).

451 Part of the reason for this result is the relatively simple treatment of the nitrogen deposition process in YIBs
452 (Yue and Unger, 2015). On the other hand, some studies have noted that due to the limitations of driving data
453 and algorithm parameters, the MODIS NPP products have some problems in China (Li et al., 2013b).

454 Furthermore, the NPP value estimated by the model over China is $3.21 \text{ Pg C yr}^{-1}$, in line with the mean value
455 ($2.92 \pm 0.12 \text{ Pg C yr}^{-1}$) found in previous 37 studies (Wang et al., 2017).




456
457 **Figure 5.** Spatial distribution of modeled (a, c) and MODIS (b, d), annual mean GPP (a, b) and NPP (c, d), and
458 their differences (e, f). The differences are simulation minus observation. Units: $\text{kg C m}^{-2} \text{ year}^{-1}$.




459
 460 **Figure 6.** Density scatter plots of (a) GPP and (b) NPP for model simulations and inversion-based products for
 461 2016. Units: kg C m⁻² year⁻¹.

462 Figure 7 and Figure 8 illustrate the seasonal fluctuations in GPP and NPP, as simulated for 2016 in East
 463 Asia. Both GPP and NPP present pronounced seasonal variations, with negligible values during winter, and a
 464 strong peak in summer. The winter minimum is attributable to limiting environmental factors such as reduced
 465 solar radiation, lower temperatures, and suppressed photosynthetic activity by vegetation. Conversely, summer
 466 shows the highest GPP and NPP values due to extended daylight hours, increased solar radiation, and
 467 temperatures facilitating increased photosynthetic activity and vegetation metabolism.



468
 469 **Figure 7.** Spatial distribution of GPP simulated by model of spring(a), summer(b), autumn(c) and winter(d) in
 470 2016. Units: g C m⁻²



471
 472 **Figure 8.** Spatial distribution of NPP simulated by model of spring(a), summer(b), autumn(c) and winter(d) in
 473 2016. Units: g C m^{-2}

474 **3.5 Simulations of other carbon-bearing species**

475 The analysis of additional carbonaceous compounds such as BC, OC and carbon monoxide (CO), is crucial
 476 due to their considerable influence on climate and the carbon cycle. The spatial distribution of simulated BC for
 477 each season of 2016 is shown in Figure S13. BC concentrations are mainly centered in North China, Central
 478 China, the Sichuan Basin, Chongqing, and Northeast India, regions with a higher concentration of industrial and
 479 residential emission sources. BC displays a marked seasonal variation, with elevated levels in winter, possibly
 480 attributed to residential heating, more stagnant conditions, and reduced removal by precipitation.

481 Figure S14 then shows the spatial corresponding distribution of seasonal OC, which is also higher over
 482 North China, Central China, Sichuan and Chongqing, and Northeast India. Finally, Figure S15 reports the
 483 annual mean near-surface CO concentrations for observations and simulation data across the monitoring sites in
 484 China. While simulated CO concentrations agree well spatially with observations, the simulations produce
 485 higher values than observed in Central China, likely linked to uncertainties in emission inventories. Figure S16
 486 presents the seasonal spatial distributions of CO, with simulated high values mostly localized in
 487 Sichuan-Chongqing and Central China, and a peak in winter.

488 **4 Conclusions**

489 Regional climate-chemical coupled models can be used to study the characteristics of regional-scale cli-
490 mate and pollutants, and is an important means to investigate the behavior of atmospheric pollutants and their
491 radiative climate effects. However, current coupled regional climate models describe the physiological process
492 of terrestrial vegetation relatively simply and do not consider the interaction between atmospheric pollutants
493 (such as PM_{2.5} and O₃) and CO₂, as well as their impacts on terrestrial ecosystems.

494 To overcome this problem, in this work we coupled the YIBs biogeochemical model to the RegCM-CHEM
495 regional climate-chemistry model, and tested this coupled modeling system over a domain covering East Asia at
496 a 30 km horizontal grid spacing for the year 2016. The model output was validated against reanalysis data, ob-
497 servational data, and satellite remote sensing data, both for the atmosphere and the carbon cycle.

498 Our simulations show that the coupled RegCM-Chem-YIBs system can effectively reproduce the spa-
499 tio-temporal distribution of meteorological variables, atmospheric composition (PM_{2.5}, O₃, and CO₂) and terres-
500 trial carbon fluxes (GPP and NPP). Comparisons of the simulated temperature, longitudinal wind, latitudinal
501 wind, and specific humidity for different seasons with the driving ERA-Interim reanalysis data showed correla-
502 tion coefficients of 0.95-0.98, 0.71-0.97, 0.81-0.92, and 0.91-0.92, respectively. The correlation coefficients
503 between observed and simulated O₃ and PM_{2.5} levels in China were 0.74 and 0.65, respectively, while the corre-
504 sponding correlations for CO₂ were in the range of 0.89 to 0.97. Comparison of the ecological parameters GPP
505 and NPP simulated in East Asia with the observed data showed correlation coefficients of 0.91 and 0.87, respec-
506 tively. In addition, in all cases, the seasonal variation of the different variables was captured by the model.
507 Therefore, we conclude that, overall, the RegCM-Chem-YIBs model demonstrates a good performance in simu-
508 lating the spatio-temporal distribution characteristics of regional meteorological characteristics, atmospheric
509 composition, and ecological parameters over East Asia.

510 In the future, we will continue to improve RegCM-Chem-YIBs in the following aspects. First, we will in-
511 vestigate the impact of CO₂ and O₃ inhomogeneity on radiation calculations by integrating temporally and spa-
512 tially varying concentrations derived from YIBs and Chem into the RegCM radiation module. This will enable
513 additional accurate computation of longwave radiation flux, improving the representation of the regional radia-
514 tion balance. Second, we intend to assimilate a module representing various chemical transformations happening
515 on the surfaces of aerosol particles. Finally, we will include the wet removal process of O₃. These advancements
516 will contribute to the refinement of RegCM-Chem-YIBs, enhancing our ability to investigate the interactions

517 between regional atmosphere, carbon cycle, and vegetation processes.

518 **Code and data availability**

519 The RegCM-Chem source code can be obtained from <https://github.com/ICTP/RegCM> (last access: 10 July
520 2023). The YIBs model code is available at https://github.com/YIBS01/YIBs_site (last access: 10 July 2023).
521 The input data and source code for RegCM-Chem-YIBs have been archived on Zenodo at
522 <https://doi.org/10.5281/zenodo.8186164> (Xie and Wang, 2023). The CarbonTracker data are provided at
523 (<https://gml.noaa.gov/ccgg/carbontracker/>). The CERES surface radiation data are available at
524 (<https://ceres.larc.nasa.gov/>). WDCGG data are available at (<https://gaw.kishou.go.jp/>). CNEMC data are pro-
525 vided at (<http://www.cnemc.cn/>). MODIS data are available at (<https://ladsweb.modaps.eosdis.nasa.gov/>).

526 **Author contributions**

527 TW led the development of RegCM-Chem-YIBs with significant contributions from NX and XX. NX per-
528 formed the evaluation. NX, TW drafted the manuscript and all authors contributed to review and editing of the
529 manuscript.

530 **Competing interests**

531 The corresponding author has stated that all the authors have no conflicts of interest.

532 **Disclaimer**

533 Publisher's note: Copernicus Publications remains neutral about jurisdictional claims in published maps and
534 institutional affiliations.

535 **Acknowledgments**

536 This work was supported by the National Natural Science Foundation of China (42077192), the National Key
537 Basic Research & Development Program of China (2020YFA0607802), the Creative talent exchange program
538 for foreign experts in the Belt and Road countries, and the Emory University-Nanjing University Collaborative
539 Research Grant.

540 **References**

- 541 Ahlstrom, A., Raupach, M. R., Schurgers, G., Smith, B., Arneth, A., Jung, M., Reichstein, M., Canadell, J. G., *et*
542 *al.*: The dominant role of semi-arid ecosystems in the trend and variability of the land CO₂ sink, *Science*,
543 348, 895-899, <https://doi.org/10.1126/science.aaa1668>, 2015.
- 544 Ainsworth, E. A., Yendrek, C. R., Sitch, S., Collins, W. J., and Emberson, L. D.: The Effects of Tropospheric
545 Ozone on Net Primary Productivity and Implications for Climate Change, *Annu. Rev. Plant Biol.*, 63,
546 637-661, <https://doi.org/10.1146/annurev-arplant-042110-103829>, 2012.
- 547 Artale, V., Calmanti, S., Carillo, A., Dell'Aquila, A., Herrmann, M., Pisacane, G., Ruti, P. M., Sannino, G., *et al.*:
548 An atmosphere-ocean regional climate model for the Mediterranean area: assessment of a present climate
549 simulation, *Clim. Dyn.*, 35, 721-740, <https://doi.org/10.1007/s00382-009-0691-8>, 2010.
- 550 Baklanov, A., Molina, L. T., and Gauss, M.: Megacities, air quality and climate, *Atmos. Environ.*, 126, 235-249,
551 <https://doi.org/10.1016/j.atmosenv.2015.11.059>, 2016.
- 552 Chang, J. S., Brost, R. A., Isaksen, I. S. A., Madronich, S., Middleton, P., Stockwell, W. R., and Walcek, C. J.: A
553 3-DIMENSIONAL EULERIAN ACID DEPOSITION MODEL - PHYSICAL CONCEPTS AND
554 FORMULATION, *J. Geophys. Res.: Atmos.*, 92, 14681-14700, <https://doi.org/10.1029/JD092iD12p14681>,
555 1987.
- 556 Chuang, M. T., Lee, C. T., Chou, C. C. K., Lin, N. H., Sheu, G. R., Wang, J. L., Chang, S. C., Wang, S. H., *et al.*:
557 Carbonaceous aerosols in the air masses transported from Indochina to Taiwan: Long-term observation at
558 Mt. Lulin, *Atmos. Environ.*, 89, 507-516, <https://doi.org/10.1016/j.atmosenv.2013.11.066>, 2014.
- 559 Chutia, L., Ojha, N., Girach, I. A., Sahu, L. K., Alvarado, L. M. A., Burrows, J. P., Pathak, B., and Bhuyan, P. K.:
560 Distribution of volatile organic compounds over Indian subcontinent during winter: WRF-chem simulation
561 versus observations, *Environ. Pollut.*, 252, 256-269, <https://doi.org/10.1016/j.envpol.2019.05.097>, 2019.
- 562 [Collatz, G. J., Ball, J. T., Grivet, C., and Berry, J. A.: Physio-logical and Environmental-Regulation of Stomatal](#)
563 [Conductance, Photosynthesis and Transpiration – a Model That Includes a Laminar Boundary-Layer, *Agric.*](#)
564 [For. Meteorol.](#), 54, 107-136, [https://doi.org/10.1016/0168-1923\(91\)90002-8](https://doi.org/10.1016/0168-1923(91)90002-8), 1991.
- 565 Coppola, E., Poulton, M., Charles, E., Dustman, J., and Szidarovszky, F.: Application of Artificial Neural
566 Networks to Complex Groundwater Management Problems, *Nat. Resour. Res.*, 12, 303-320,
567 <https://doi.org/10.1023/B:NARR.0000007808.11860.7e>, 2003.
- 568 Dickinson, R. E., Errico, R. M., Giorgi, F., and Bates, G. T.: A Regional Climate Model for the Western
569 United-States, *Clim. Change*, 15, 383-422, <https://doi.org/10.1007/BF00240465>, 1989.
- 570 Dunne, J. P., Horowitz, L. W., Adcroft, A. J., Ginoux, P., Held, I. M., John, J. G., Krasting, J. P., Malyshev, S., *et*
571 *al.*: The GFDL Earth System Model Version 4.1 (GFDL-ESM 4.1): Overall Coupled Model Description
572 and Simulation Characteristics, *J. Adv. Model. Earth Syst.*, 12,
573 <https://doi.org/ARTNe2019MS00201510.1029/2019MS002015>, 2020.
- 574 Emmons, L. K., Walters, S., Hess, P. G., Lamarque, J. F., Pfister, G. G., Fillmore, D., Granier, C., Guenther, A.,
575 *et al.*: Description and evaluation of the Model for Ozone and Related chemical Tracers, version 4
576 (MOZART-4), *Geosci. Model Dev.*, 3, 43-67, <https://doi.org/10.5194/gmd-3-43-2010>, 2010.
- 577 Farquhar, G. D., Caemmerer, S. V., and Berry, J. A.: A biochemical model of photosynthetic CO₂ assimilation in
578 leaves of C₃ species, *Planta*, 149, 78-90, <https://doi.org/10.1007/BF00386231>, 1980.
- 579 Fiore, A. M., Naik, V., and Leibensperger, E. M.: Air Quality and Climate Connections, *J. Air Waste Manage.*
580 *Assoc.*, 65, 645-685, <https://doi.org/10.1080/10962247.2015.1040526>, 2015.
- 581 Fiore, A. M., Naik, V., Spracklen, D. V., Steiner, A., Unger, N., Prather, M., Bergmann, D., Cameron-Smith, P. J.,
582 *et al.*: Global air quality and climate, *Chem. Soc. Rev.*, 41, 6663-6683, <https://doi.org/10.1039/c2cs35095e>,

583 2012.

584 Forkel, M., Carvathais, N., Rodenbeck, C., Keeling, R., Heimann, M., Thonicke, K., Zaehle, S., and Reichstein,
585 M.: Enhanced seasonal CO₂ exchange caused by amplified plant productivity in northern ecosystems,
586 *Science*, 351, 696-699, <https://doi.org/10.1126/science.aac4971>, 2016.

587 Gao, Y. M., Zhuang, B. L., Wang, T. J., Chen, H. M., Li, S., Wei, W., Lin, H. J., and Li, M. M.:
588 Climatic-Environmental Effects of Aerosols and Their Sensitivity to Aerosol Mixing States in East Asia in
589 Winter, *Remote Sens.*, 14, <https://doi.org/10.3390/rs14153539>, 2022.

590 Gery, M. W., Whitten, G. Z., Killus, J. P., and Dodge, M. C.: A Photochemical Kinetics Mechanism for Urban
591 and Regional Scale Computer Modeling, *J. Geophys. Res.: Atmos.*, 94, 12925-12956,
592 <https://doi.org/10.1029/JD094iD10p12925>, 1989.

593 Giorgi, F.: Simulation of Regional Climate Using a Limited Area Model Nested in a General-Circulation Model,
594 *J. Clim.*, 3, 941-963, [https://doi.org/10.1175/1520-0442\(1990\)003<0941:Sorcu>2.0.Co;2](https://doi.org/10.1175/1520-0442(1990)003<0941:Sorcu>2.0.Co;2), 1990.

595 Giorgi, F. and Bates, G. T.: The Climatological Skill of a Regional Model over Complex Terrain, *Mon. Weather*
596 *Rev.*, 117, 2325-2347, [https://doi.org/10.1175/1520-0493\(1989\)117<2325:TCSOAR>2.0.CO;2](https://doi.org/10.1175/1520-0493(1989)117<2325:TCSOAR>2.0.CO;2), 1989.

597 Giorgi, F. and Mearns, L. O.: Introduction to special section: Regional climate modeling revisited, *J. Geophys.*
598 *Res.: Atmos.*, 104, 6335-6352, <https://doi.org/10.1029/98jd02072>, 1999.

599 Giorgi, F., Marinucci, M. R., Bates, G. T., and Decanio, G.: DEVELOPMENT OF A 2ND-GENERATION
600 REGIONAL CLIMATE MODEL (REGCM2) .2. CONVECTIVE PROCESSES AND ASSIMILATION OF
601 LATERAL BOUNDARY-CONDITIONS, *Mon. Weather Rev.*, 121, 2814-2832,
602 [https://doi.org/10.1175/1520-0493\(1993\)121<2814:Doasgr>2.0.Co;2](https://doi.org/10.1175/1520-0493(1993)121<2814:Doasgr>2.0.Co;2), 1993.

603 Giorgi, F., Pal, J. S., Bi, X., Sloan, L., Elguindi, N., and Solmon, F.: Introduction to the TAC special issue: The
604 RegCNET network, *Theor. Appl. Climatol.*, 86, 1-4, <https://doi.org/10.1007/s00704-005-0199-z>, 2006.

605 Giorgi, F., Coppola, E., Solmon, F., Mariotti, L., Sylla, M. B., Bi, X., Elguindi, N., Diro, G. T., *et al.*: RegCM4:
606 model description and preliminary tests over multiple CORDEX domains, *Clim. Res.*, 52, 7-29,
607 <https://doi.org/10.3354/cr01018>, 2012.

608 Guenther, A., Hewitt, C. N., Erickson, D., Fall, R., Geron, C., Graedel, T., Harley, P., Klinger, L., *et al.*: A
609 Global-Model of Natural Volatile Organic-Compound Emissions, *J. Geophys. Res.: Atmos.*, 100,
610 8873-8892, <https://doi.org/10.1029/94jd02950>, 1995.

611 Han, Z. W., Li, J. W., Xia, X. G., and Zhang, R. J.: Investigation of direct radiative effects of aerosols in dust
612 storm season over East Asia with an online coupled regional climate-chemistry-aerosol model, *Atmos.*
613 *Environ.*, 54, 688-699, <https://doi.org/10.1016/j.atmosenv.2012.01.041>, 2012.

614 Han, Z. W., Xie, Z. X., Wang, G. H., Zhang, R. J., and Tao, J.: Modeling organic aerosols over east China using
615 a volatility basis-set approach with aging mechanism in a regional air quality model, *Atmos. Environ.*, 124,
616 186-198, <https://doi.org/10.1016/j.atmosenv.2015.05.045>, 2016.

617 He, M. Z., Kimball, J. S., Maneta, M. P., Maxwell, B. D., Moreno, A., Begueria, S., and Wu, X. C.: Regional
618 Crop Gross Primary Productivity and Yield Estimation Using Fused Landsat-MODIS Data, *Remote Sens.*,
619 10, <https://doi.org/10.3390/rs10030372>, 2018.

620 Hong, C. P., Zhang, Q., Zhang, Y., Davis, S. J., Tong, D., Zheng, Y. X., Liu, Z., Guan, D. B., *et al.*: Impacts of
621 climate change on future air quality and human health in China, *Proc. Natl. Acad. Sci. U.S.A.*, 116,
622 17193-17200, <https://doi.org/10.1073/pnas.1812881116>, 2019.

623 Horowitz, L. W., Walters, S., Mauzerall, D. L., Emmons, L. K., Rasch, P. J., Granier, C., Tie, X. X., Lamarque, J.
624 F., *et al.*: A global simulation of tropospheric ozone and related tracers: Description and evaluation of
625 MOZART, version 2, *J. Geophys. Res.: Atmos.*, 108, <https://doi.org/10.1029/2002jd002853>, 2003.

626 Kan, H. D., Chen, R. J., and Tong, S. L.: Ambient air pollution, climate change, and population health in China,

627 Environ. Int., 42, 10-19, <https://doi.org/10.1016/j.envint.2011.03.003>, 2012.

628 Kiehl, J. T., Hack, J. J., Bonan, G. B., Boville, B. A., and Briegleb, B. P.: Description of the NCAR community
629 climate model (CCM3), National Center for Atmospheric Research, Boulder, CO (United States). Climate
630 and Global Dynamics Div., Technical Report PB-97-131528/XAB;NCAR/TN-420-STR TRN: 70341499,
631 1996.

632 Kim, K. H., Kabir, E., and Kabir, S.: A review on the human health impact of airborne particulate matter,
633 Environ. Int., 74, 136-143, <https://doi.org/10.1016/j.envint.2014.10.005>, 2015.

634 Kou, X. X., Zhang, M. G., Peng, Z., and Wang, Y. H.: Assessment of the biospheric contribution to surface
635 atmospheric CO₂ concentrations over East Asia with a regional chemical transport model, Adv. Atmos. Sci.,
636 32, 287-300, <https://doi.org/10.1007/s00376-014-4059-6>, 2015.

637 Lamarque, J. F., Dentener, F., McConnell, J., Ro, C. U., Shaw, M., Vet, R., Bergmann, D., Cameron-Smith, P., *et*
638 *al.*: Multi-model mean nitrogen and sulfur deposition from the Atmospheric Chemistry and Climate Model
639 Intercomparison Project (ACCMIP): evaluation of historical and projected future changes, Atmos. Chem.
640 Phys., 13, 7997-8018, <https://doi.org/10.5194/acp-13-7997-2013>, 2013.

641 Lawrence, P. J. and Chase, T. N.: Representing a new MODIS consistent land surface in the Community Land
642 Model (CLM 3.0), J. Geophys. Res.: Biogeosci., 112, <https://doi.org/10.1029/2006jg000168>, 2007.

643 Li, B., Gasser, T., Ciais, P., Piao, S., Tao, S., Balkanski, Y., Hauglustaine, D., Boisier, J.-P., *et al.*: The
644 contribution of China's emissions to global climate forcing, Nature, 531, 357-361,
645 <https://doi.org/10.1038/nature17165>, 2016a.

646 Li, J. W., Han, Z. W., and Xie, Z. X.: Model analysis of long-term trends of aerosol concentrations and direct
647 radiative forcings over East Asia, Tellus B: Chem. Phys. Meteorol., 65,
648 <https://doi.org/10.3402/tellusb.v65i0.20410>, 2013a.

649 Li, J. W., Han, Z. W., and Zhang, R. J.: Influence of aerosol hygroscopic growth parameterization on aerosol
650 optical depth and direct radiative forcing over East Asia, Atmos. Res., 140, 14-27,
651 <https://doi.org/10.1016/j.atmosres.2014.01.013>, 2014.

652 Li, M., Wang, T., Han, Y., Xie, M., Li, S., Zhuang, B., and Chen, P.: Modeling of a severe dust event and its
653 impacts on ozone photochemistry over the downstream Nanjing megacity of eastern China, Atmos.
654 Environ., 160, 107-123, <https://doi.org/10.1016/j.atmosenv.2017.04.010>, 2017a.

655 Li, M., Zhang, Q., Kurokawa, J., Woo, J. H., He, K. B., Lu, Z. F., Ohara, T., Song, Y., *et al.*: MIX: a mosaic
656 Asian anthropogenic emission inventory under the international collaboration framework of the MICS-Asia
657 and HTAP, Atmos. Chem. Phys., 17, 935-963, <https://doi.org/10.5194/acp-17-935-2017>, 2017b.

658 Li, S., Wang, T. J., Zhuang, B. L., and Han, Y.: Indirect radiative forcing and climatic effect of the anthropogenic
659 nitrate aerosol on regional climate of China, Adv. Atmos. Sci., 26, 543-552,
660 <https://doi.org/10.1007/s00376-009-0543-9>, 2009.

661 Li, S., Wang, T. J., Solmon, F., Zhuang, B. L., Wu, H., Xie, M., Han, Y., and Wang, X. M.: Impact of aerosols on
662 regional climate in southern and northern China during strong/weak East Asian summer monsoon years, J.
663 Geophys. Res.: Atmos., 121, 4069-4081, <https://doi.org/10.1002/2015jd023892>, 2016b.

664 Li, X. L., Liang, S. L., Yu, G. R., Yuan, W. P., Cheng, X., Xia, J. Z., Zhao, T. B., Feng, J. M., *et al.*: Estimation
665 of gross primary production over the terrestrial ecosystems in China, Ecol. Model., 261, 80-92,
666 <https://doi.org/10.1016/j.ecolmodel.2013.03.024>, 2013b.

667 Liu, L., Solmon, F., Vautard, R., Hamaoui-Laguel, L., Torma, C. Z., and Giorgi, F.: Ragweed pollen production
668 and dispersion modelling within a regional climate system, calibration and application over Europe,
669 Biogeosciences, 13, 2769-2786, <https://doi.org/10.5194/bg-13-2769-2016>, 2016.

670 Liu, Z., Deng, Z., He, G., Wang, H. L., Zhang, X., Lin, J., Qi, Y., and Liang, X.: Challenges and opportunities

671 for carbon neutrality in China, *Nat Rev Earth Env*, 3, 141-155, <https://doi.org/10.1038/s43017-021-00244-x>,
672 2022.

673 Lu, X., Zhang, S. J., Xing, J., Wang, Y. J., Chen, W. H., Ding, D., Wu, Y., Wang, S. X., *et al.*: Progress of Air
674 Pollution Control in China and Its Challenges and Opportunities in the Ecological Civilization Era,
675 *Engineering*, 6, 1423-1431, <https://doi.org/10.1016/j.eng.2020.03.014>, 2020.

676 Lu, X. L., Chen, M., Liu, Y. L., Miralles, D. G., and Wang, F. M.: Enhanced water use efficiency in global
677 terrestrial ecosystems under increasing aerosol loadings, *Agric. For. Meteorol.*, 237, 39-49,
678 <https://doi.org/10.1016/j.agrformet.2017.02.002>, 2017.

679 Ma, D., Wang, T., Wu, H., Qu, Y., Liu, J., Liu, J., Li, S., Zhuang, B., *et al.*: The effect of anthropogenic emission,
680 meteorological factors, and carbon dioxide on the surface ozone increase in China from 2008 to 2018
681 during the East Asia summer monsoon season, *Atmos. Chem. Phys.*, 23, 6525-6544,
682 <https://doi.org/10.5194/acp-23-6525-2023>, 2023a.

683 Ma, D. Y., Wang, T. J., Xu, B. Y., Song, R., Gao, L. B., Chen, H. M., Ren, X. J., Li, S., *et al.*: The mutual
684 interactions among ozone, fine particulate matter, and carbon dioxide on summer monsoon climate in East
685 Asia, *Atmos. Environ.*, 299, <https://doi.org/10.1016/j.atmosenv.2023.119668>, 2023b.

686 Madani, N., Kimball, J. S., Affleck, D. L. R., Kattge, J., Graham, J., van Bodegom, P. M., Reich, P. B., and
687 Running, S. W.: Improving ecosystem productivity modeling through spatially explicit estimation of
688 optimal light use efficiency, *J. Geophys. Res.: Biogeosci.*, 119, 1755-1769,
689 <https://doi.org/10.1002/2014jg002709>, 2014.

690 Madronich, S. and Flocke, S.: The role of solar radiation in atmospheric chemistry, *Environ. Chem.*, 1-26,
691 https://doi.org/10.1007/978-3-540-69044-3_1, 1998.

692 [Niinemets, Ü., Tenhunen, J. D., Harley, P. C., and Steinbrecher, R.: A model of isoprene emission based on](#)
693 [energetic requirements for isoprene synthesis and leaf photosynthetic properties for Liquidambar and](#)
694 [Quercus. *Plant, Cell & Environment*, 22, 1319-1335. <https://doi.org/10.1046/j.1365-3040.1999.00505.x>.](#)
695 [2002.](#)

696 Oleson, K. W., Niu, G. Y., Yang, Z. L., Lawrence, D. M., Thornton, P. E., Lawrence, P. J., Stockli, R., Dickinson,
697 R. E., *et al.*: Improvements to the Community Land Model and their impact on the hydrological cycle, *J.*
698 *Geophys. Res.: Biogeosci.*, 113, <https://doi.org/10.1029/2007jg000563>, 2008.

699 [Pacífico, F., Harrison, S. P., Jones, C. D., Arneth, A., Sitch, S., Weedon, G. P., Barkley, M. P., Palmer, P. I., et al.:](#)
700 [Evaluation of a photosynthesis-based biogenic isoprene emission scheme in JULES and simulation of](#)
701 [isoprene emissions under present-day climate conditions. *Atmos. Chem. Phys.*, 11, 4371-4389.](#)
702 <https://doi.org/10.5194/acp-11-4371-2011>.

703 Pal, J. S., Giorgi, F., Bi, X. Q., Elguindi, N., Solmon, F., Gao, X. J., Rauscher, S. A., Francisco, R., *et al.*:
704 Regional climate modeling for the developing world - The ICTP RegCM3 and RegCNET, *Bull. Am.*
705 *Meteorol. Soc.*, 88, 1395-+, <https://doi.org/10.1175/Bams-88-9-1395>, 2007.

706 Peters, W., Jacobson, A. R., Sweeney, C., Andrews, A. E., Conway, T. J., Masarie, K., Miller, J. B., Bruhwiler, L.
707 M. P., *et al.*: An atmospheric perspective on North American carbon dioxide exchange: CarbonTracker,
708 *Proc. Natl. Acad. Sci. U.S.A.*, 104, 18925-18930, <https://doi.org/10.1073/pnas.0708986104>, 2007.

709 Pu, X., Wang, T. J., Huang, X., Melas, D., Zanis, P., Papanastasiou, D. K., and Poupkou, A.: Enhanced surface
710 ozone during the heat wave of 2013 in Yangtze River Delta region, China, *Sci. Total Environ.*, 603,
711 807-816, <https://doi.org/10.1016/j.scitotenv.2017.03.056>, 2017.

712 Reynolds, R. W., Rayner, N. A., Smith, T. M., Stokes, D. C., and Wang, W. Q.: An improved in situ and satellite
713 SST analysis for climate, *J. Clim.*, 15, 1609-1625,
714 [https://doi.org/10.1175/1520-0442\(2002\)015<1609:Aiisas>2.0.Co;2](https://doi.org/10.1175/1520-0442(2002)015<1609:Aiisas>2.0.Co;2), 2002.

715 Running, S. W.: A Measurable Planetary Boundary for the Biosphere, *Science*, 337, 1458-1459,
716 <https://doi.org/10.1126/science.1227620>, 2012.

717 Scheuch, M., Hoper, D., and Beer, M.: RIEMS: a software pipeline for sensitive and comprehensive taxonomic
718 classification of reads from metagenomics datasets, *BMC Bioinf.*, 16,
719 <https://doi.org/10.1186/s12859-015-0503-6>, 2015.

720 Shalaby, A., Zaakey, A. S., Tawfik, A. B., Solmon, F., Giorgi, F., Stordal, F., Sillman, S., Zaveri, R. A., *et al.*:
721 Implementation and evaluation of online gas-phase chemistry within a regional climate model
722 (RegCM-CHEM4), *Geosci. Model Dev.*, 5, 741-760, <https://doi.org/10.5194/gmd-5-741-2012>, 2012.

723 Shindell, D. T., Lamarque, J. F., Schulz, M., Flanner, M., Jiao, C., Chin, M., Young, P. J., Lee, Y. H., *et al.*:
724 Radiative forcing in the ACCMIP historical and future climate simulations, *Atmos. Chem. Phys.*, 13,
725 2939-2974, <https://doi.org/10.5194/acp-13-2939-2013>, 2013.

726 Sitch, S., Cox, P. M., Collins, W. J., and Huntingford, C.: Indirect radiative forcing of climate change through
727 ozone effects on the land-carbon sink, *Nature*, 448, 791-U794, <https://doi.org/10.1038/nature06059>, 2007.

728 Slingo, A.: A Gcm Parameterization for the Shortwave Radiative Properties of Water Clouds, *J. Atmos. Sci.*, 46,
729 1419-1427, [https://doi.org/10.1175/1520-0469\(1989\)046<1419:Agpfts>2.0.Co;2](https://doi.org/10.1175/1520-0469(1989)046<1419:Agpfts>2.0.Co;2), 1989.

730 Small, E. E., Sloan, L. C., Hostetler, S., and Giorgi, F.: Simulating the water balance of the Aral Sea with a
731 coupled regional climate-lake model, *J. Geophys. Res.: Atmos.*, 104, 6583-6602, <https://doi.org/10.1029/98jd02348>, 1999.

732 Solmon, F., Elguindi, N., and Mallet, M.: Radiative and climatic effects of dust over West Africa, as simulated
733 by a regional climate model, *Clim. Res.*, 52, 97-113, <https://doi.org/10.3354/cr01039>, 2012.

734 Solmon, F., Giorgi, F., and Lioussé, C.: Aerosol modelling for regional climate studies: application to
735 anthropogenic particles and evaluation over a European/African domain, *Tellus B: Chem. Phys. Meteorol.*,
736 58, 51-72, <https://doi.org/10.1111/j.1600-0889.2005.00155.x>, 2006.

737 Spitters, C. J. T., Toussaint, H., and Goudriaan, J.: SEPARATING THE DIFFUSE AND DIRECT
738 COMPONENT OF GLOBAL RADIATION AND ITS IMPLICATIONS FOR MODELING CANOPY
739 PHOTOSYNTHESIS .1. COMPONENTS OF INCOMING RADIATION, *Agric. For. Meteorol.*, 38,
740 217-229, [https://doi.org/10.1016/0168-1923\(86\)90060-2](https://doi.org/10.1016/0168-1923(86)90060-2), 1986.

741 Strada, S. and Unger, N.: Potential sensitivity of photosynthesis and isoprene emission to direct radiative effects
742 of atmospheric aerosol pollution, *Atmos. Chem. Phys.*, 16, 4213-4234,
743 <https://doi.org/10.5194/acp-16-4213-2016>, 2016.

744 Turuncoglu, U. U., Dalfes, N., Murphy, S., and DeLuca, C.: Toward self-describing and workflow integrated
745 Earth system models: A coupled atmosphere-ocean modeling system application, *Environ. Model. Software*,
746 39, 247-262, <https://doi.org/10.1016/j.envsoft.2012.02.013>, 2013.

747 Wang, T., Xue, L. K., Brimblecombe, P., Lam, Y. F., Li, L., and Zhang, L.: Ozone pollution in China: A review
748 of concentrations, meteorological influences, chemical precursors, and effects, *Sci. Total Environ.*, 575,
749 1582-1596, <https://doi.org/10.1016/j.scitotenv.2016.10.081>, 2017.

750 Wang, T. J., Li, S., Shen, Y., Deng, J. J., and Xie, M.: Investigations on direct and indirect effect of nitrate on
751 temperature and precipitation in China using a regional climate chemistry modeling system, *J. Geophys.*
752 *Res.: Atmos.*, 115, <https://doi.org/10.1029/2009jd013264>, 2010.

753 Wesely, M. L.: Parameterization of Surface Resistances to Gaseous Dry Deposition in Regional-Scale
754 Numerical-Models, *Atmos. Environ.*, 23, 1293-1304, [https://doi.org/10.1016/0004-6981\(89\)90153-4](https://doi.org/10.1016/0004-6981(89)90153-4), 1989.

755 Wiedinmyer, C., Akagi, S. K., Yokelson, R. J., Emmons, L. K., Al-Saadi, J. A., Orlando, J. J., and Soja, A. J.:
756 The Fire INventory from NCAR (FINN): a high resolution global model to estimate the emissions from
757 open burning, *Geosci. Model Dev.*, 4, 625-641, <https://doi.org/10.5194/gmd-4-625-2011>, 2011.

759 Xie, X. D., Wang, T. J., Yue, X., Li, S., Zhuang, B. L., and Wang, M. H.: Effects of atmospheric aerosols on
760 terrestrial carbon fluxes and CO₂ concentrations in China, *Atmos. Res.*, 237,
761 <https://doi.org/10.1016/j.atmosres.2020.104859>, 2020.

762 Xie, X. D., Wang, T. J., Yue, X., Li, S., Zhuang, B. L., Wang, M. H., and Yang, X. Q.: Numerical modeling of
763 ozone damage to plants and its effects on atmospheric CO₂ in China, *Atmos. Environ.*, 217,
764 <https://doi.org/10.1016/j.atmosenv.2019.116970>, 2019.

765 Xiong, Z., Fu, C. B., and Yan, X. D.: Regional integrated environmental model system and its simulation of East
766 Asia summer monsoon, *Chin. Sci. Bull.*, 54, 4253-4261, <https://doi.org/10.1007/s11434-009-0669-2>, 2009.

767 Xu, B. Y., Wang, T. J., Ma, D. Y., Song, R., Zhang, M., Gao, L. B., Li, S., Zhuang, B. L., *et al.*: Impacts of
768 regional emission reduction and global climate change on air quality and temperature to attain carbon
769 neutrality in China, *Atmos. Res.*, 279, <https://doi.org/10.1016/j.atmosres.2022.106384>, 2022.

770 Xu, B. Y., Wang, T. J., Gao, L. B., Ma, D. Y., Song, R., Zhao, J., Yang, X. G., Li, S., *et al.*: Impacts of
771 meteorological factors and ozone variation on crop yields in China concerning carbon neutrality objectives
772 in 2060, *Environ. Pollut.*, 317, <https://doi.org/10.1016/j.envpol.2022.120715>, 2023.

773 Yahya, K., Wang, K., Campbell, P., Chen, Y., Glotfelty, T., He, J., Pirhalla, M., and Zhang, Y.: Decadal
774 application of WRF/Chem for regional air quality and climate modeling over the U.S. under the
775 representative concentration pathways scenarios. Part 1: Model evaluation and impact of downscaling,
776 *Atmos. Environ.*, 152, 562-583, <https://doi.org/10.1016/j.atmosenv.2016.12.029>, 2017.

777 Yin, C. Q., Wang, T. J., Solmon, F., Mallet, M., Jiang, F., Li, S., and Zhuang, B. L.: Assessment of direct
778 radiative forcing due to secondary organic aerosol over China with a regional climate model, *Tellus Series*
779 *B-Chemical and Physical Meteorology*, 67, <https://doi.org/10.3402/tellusb.v67.24634>, 2015.

780 Yin, J. F., Wang, D. H., Zhai, G. Q., and Xu, H. B.: An investigation into the relationship between liquid water
781 content and cloud number concentration in the stratiform clouds over north China, *Atmos. Res.*, 139,
782 137-143, <https://doi.org/10.1016/j.atmosres.2013.12.004>, 2014.

783 Young, P. J., Archibald, A. T., Bowman, K. W., Lamarque, J. F., Naik, V., Stevenson, D. S., Tilmes, S.,
784 Voulgarakis, A., *et al.*: Pre-industrial to end 21st century projections of tropospheric ozone from the
785 Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP), *Atmos. Chem. Phys.*, 13,
786 2063-2090, <https://doi.org/10.5194/acp-13-2063-2013>, 2013.

787 Yue, X. and Unger, N.: The Yale Interactive terrestrial Biosphere model version 1.0: description, evaluation and
788 implementation into NASA GISS ModelE2, *Geosci. Model Dev.*, 8, 2399-2417,
789 <https://doi.org/10.5194/gmd-8-2399-2015>, 2015.

790 Yue, X. and Unger, N.: Aerosol optical depth thresholds as a tool to assess diffuse radiation fertilization of the
791 land carbon uptake in China, *Atmos. Chem. Phys.*, 17, 1329-1342,
792 <https://doi.org/10.5194/acp-17-1329-2017>, 2017.

793 Yue, X., Strada, S., Unger, N., and Wang, A. H.: Future inhibition of ecosystem productivity by increasing
794 wildfire pollution over boreal North America, *Atmos. Chem. Phys.*, 17, 13699-13719,
795 <https://doi.org/10.5194/acp-17-13699-2017>, 2017.

796 Zakey, A. S., Solmon, F., and Giorgi, F.: Implementation and testing of a desert dust module in a regional
797 climate model, *Atmos. Chem. Phys.*, 6, 4687-4704, <https://doi.org/10.5194/acp-6-4687-2006>, 2006.

798 Zaveri, R. A. and Peters, L. K.: A new lumped structure photochemical mechanism for large-scale applications, *J.*
799 *Geophys. Res.: Atmos.*, 104, 30387-30415, <https://doi.org/10.1029/1999jd900876>, 1999.

800 Zhang, H., Jing, X., and Li, J.: Application and evaluation of a new radiation code under McICA scheme in
801 BCC_AGCM2.0.1, *Geosci. Model Dev.*, 7, 737-754, <https://doi.org/10.5194/gmd-7-737-2014>, 2014.

802 Zhang, H., Wang, Z. L., Wang, Z. Z., Liu, Q. X., Gong, S. L., Zhang, X. Y., Shen, Z. P., Lu, P., *et al.*: Simulation

803 of direct radiative forcing of aerosols and their effects on East Asian climate using an interactive
804 AGCM-aerosol coupled system, *Clim. Dyn.*, 38, 1675-1693, <https://doi.org/10.1007/s00382-011-1131-0>,
805 2012.

806 Zheng, B., Tong, D., Li, M., Liu, F., Hong, C. P., Geng, G. N., Li, H. Y., Li, X., *et al.*: Trends in China's
807 anthropogenic emissions since 2010 as the consequence of clean air actions, *Atmos. Chem. Phys.*, 18,
808 14095-14111, <https://doi.org/10.5194/acp-18-14095-2018>, 2018.

809 Zhou, Y., Huang, A. N., Jiang, J., and La, M. K.: Modeled interaction between the subseasonal evolving of the
810 East Asian summer monsoon and the direct effect of anthropogenic sulfate, *J. Geophys. Res.: Atmos.*, 119,
811 1993-2016, <https://doi.org/10.1002/2013jd020612>, 2014.

812 Zhuang, B. L., Jiang, F., Wang, T. J., Li, S., and Zhu, B.: Investigation on the direct radiative effect of fossil fuel
813 black-carbon aerosol over China, *Theor. Appl. Climatol.*, 104, 301-312,
814 <https://doi.org/10.1007/s00704-010-0341-4>, 2011.

815 Zhuang, B. L., Li, S., Wang, T. J., Deng, J. J., Xie, M., Yin, C. Q., and Zhu, J. L.: Direct radiative forcing and
816 climate effects of anthropogenic aerosols with different mixing states over China, *Atmos. Environ.*, 79,
817 349-361, <https://doi.org/10.1016/j.atmosenv.2013.07.004>, 2013.

818 Zhuang, B. L., Li, S., Wang, T. J., Liu, J., Chen, H. M., Chen, P. L., Li, M. M., and Xie, M.: Interaction between
819 the Black Carbon Aerosol Warming Effect and East Asian Monsoon Using RegCM4, *J. Clim.*, 31,
820 9367-9388, <https://doi.org/10.1175/Jcli-D-17-0767.1>, 2018.

821