

Sharp increase of Saharan dust intrusions over the western Euro-Mediterranean in February-March 2020-2022 and associated atmospheric circulation

5 Emilio Cuevas-Agulló^{1,*}, David Barriopedro², Rosa Delia García^{1,3}, Silvia Alonso-Pérez^{4,5}, Juan Jesús González-Alemán⁶, Ernest Werner⁷, David Suárez⁸, Juan José Bustos¹, Gerardo García-Castrillo⁷, Omaira García¹, África Barreto¹, Sara Basart^{9,10}

¹Izaña Atmospheric Research Center (IARC), State Meteorological Agency (AEMET), Santa Cruz de Tenerife, 38001 Spain

10 ²Instituto de Geociencias (IGEO), Consejo Superior de Investigaciones Científicas - Universidad Complutense de Madrid (CSIC-UCM), Madrid, 28040, Spain

³TRAGSATEC, Madrid, Spain

⁴Departamento de Ingeniería Industrial, Universidad de La Laguna (ULL), La Laguna (Tenerife), 38002, Spain

15 ⁵Instituto Universitario de Estudios de las Mujeres, Universidad de La Laguna (ULL), La Laguna (Tenerife), 38002, Spain

⁶Modeling Area-Department of Development and Applications, State Meteorological Agency (AEMET), Madrid, 28040, Spain

⁷Territorial Delegation of AEMET in Catalonia, State Meteorological Agency (AEMET), Barcelona, 08071, Spain

20 ⁸Territorial Delegation of AEMET in Canary Islands, State Meteorological Agency of Spain (AEMET), Las Palmas, 35017, Spain

⁹Barcelona Supercomputing Center (BSC), Barcelona, 08034, Spain

¹⁰Now at the World Meteorological Organization (WMO), Geneva, 1201, Switzerland

*Now, retired

Correspondence to: Sara Basart (sbasart@wmo.int) and Emilio Cuevas-Agulló (emilio.cuevas@uva.es)

Abstract

During the months of February-March (FM) of the 2020-2022 period, several intense dust intrusions from northern Africa affected Europe. The frequency of dust events was exceptional, considering that wintertime is the season with minimum dust activity in the Mediterranean, and some episodes displayed a duration and/or intensity never recorded before, affecting large areas of the western Euro-Mediterranean region (WEM). The main objective of this work is to construct a catalogue of FM dust events over WEM for the 2003-2022 period based on satellite aerosol retrievals and analyse their atmospheric drivers at the synoptic and large scales, paying special attention to the recent 2020-2022 period of high dust activity. Overall, our results indicate large intra-seasonal and inter-annual variability in the occurrence of wintertime dust events over WEM. Dust events of FM 2020-2022 were characterised by enhanced dust concentration and high maximum altitudes in comparison with those of previous years (2003-2019). WEM dust events are associated with enhanced activity of high-pressure systems over the Euro-Atlantic sector, which favour the obstruction of the westerlies and the occurrence of cut-off lows at subtropical latitudes. However, these high-pressure systems can exhibit a large variety of configurations, including meridional dipole blocking patterns with poleward shifted jets, or Mediterranean subtropical ridges with an intensified mid-latitude jet. The former is the dominant favourable pattern for WEM dust occurrence, but the latter was relatively common during the 2020-2022 period.

1 Introduction

Mineral dust, when considering its mass, is the most abundant type of atmospheric aerosol found over the ocean and continents, representing about 75% of the global aerosols present in the atmosphere (Kinne et al., 2006; Wu et al., 2020). Dust affects the climate system by interacting with longwave and shortwave radiation, as well as contributing to the formation of cloud condensation nuclei and ice nucleating particles (e.g. Szopa et al., 2021). Recent studies revealed that the magnitude of radiative forcing due to mineral dust is small (Kok et al., 2017; Ryder et al., 2018) because dust particles seem to be larger than previously thought (Kok et al., 2017; Szopa et al., 2021). This new finding, along with the fact that global models underestimate the amount of coarse dust in the atmosphere (Adebiyi and Kok, 2020) raises the possibility that dust emissions may even be warming the climate system (Szopa et al., 2021). Primary emissions of mineral dust from deserts are the most important contributor to the aerosol budget in the Middle East, Eurasia, and Africa, accounting for 40–70 % of the annual average atmospheric

55 concentrations (Szopa et al., 2021). In large regions of the Earth, mineral dust exerts a great impact on ecosystems (Middleton and Kappelle, 2020), human health (Aghababaeian et al., 2021; Kinni et al., 2021) and other socio-economic sectors such as energy and transportation (Cuevas et al., 2021; ESCAP, 2021; Monteiro et al., 2022).

The Sahara and surrounding regions contribute to more than half of global dust emissions (Kok et al., 2021). Dust mobilization and transport in northern Africa shows high variability at different time scales: diurnal (Knippertz, 2008; Cuesta et al., 2009; Fiedler et al., 2014), intraseasonal (Ashpole and Washington, 2013; Cuevas et al., 2017), 60 seasonal (Barnaba and Gobbi, 2004; Israelevich et al., 2012), interannual (Rodríguez et al., 2015; Tegen et al., 2013; Wagner et al., 2016) and multidecadal (Wang et al., 2015) ones. African dust can travel around the globe to parts of Europe, South America, Central America, the Caribbean, and the United States (e.g., Shepherd et al., 2016). Mineral dust mobilized in the Sahara and North Africa often reaches the Mediterranean basin and the Canary Islands (Moulin et al., 1998; Basart et al., 2009, Varga et al., 2014, Gkikas et al., 2015; Cuevas et al., 2021; Monteiro 65 et al., 2022), affecting air quality and causing exceedances of the thresholds established by the World Health Organisation (WHO) Air Quality Guidelines (WHO, 2021). Specifically, in southern European countries, African dust intrusions cause the exceedance of the European Union Air Quality Directive (Kuula et al., 2022) for particulate matter (PM) (Gerasopoulos et al., 2006; Escudero et al., 2007; Querol et al., 2009; Basart et al., 2012; Pey et al., 2013; Marconi et al., 2014). Furthermore, extreme dust events in Europe lead to large impacts on socio-economic activities such as energy and transportation with associated economic losses (e.g., Monteiro et al., 2022). 70 Finally, Saharan dust deposition fertilizes the Mediterranean Sea by triggering marine phytoplankton growth, but also harmful cyanobacterial blooms (e.g., Guerzoni et al., 1999; Gallisai et al., 2014; Lagaria et al., 2017).

The challenge of mitigating the impacts of sand and dust storms is recognised globally. The United Nations (UN) agencies are promoting measures to confront the problem and their inclusion in national policies through the UN 75 Coalition for Combating SDS (Pitkanen-Brunnsberg, 2019). Aligned with the UN Coalition's objectives, the World Meteorological Organization (WMO) Sand and Dust Storm Warning Advisory and Assessment System (SDS-WAS; WMO, 2015) aims to enhance the ability of countries to deliver timely and good quality SDS forecasts, observations, and knowledge to users through an international hub of research and operational communities (Terradellas et al., 2015, Basart et al., 2019). Monitoring and forecasting are pillars for the deployment of Early 80 Warning Systems that are the most effective tool to mitigate the impacts of natural hazards such as SDS.

Winters from 2020 to 2022 were characterized by an unusually high frequency of intense and long-lasting dust events over the western Mediterranean impacting the Euro-Atlantic region, which merited preliminary and widely disseminated analyses by international institutions such as WMO, Copernicus and the European Organisation for

the Exploitation of Meteorological Satellites (EUMETSAT). These institutions described some of these dust events
85 (see Table S1) as "unusual", "historic" or "exceptional". In February 2020, two strong Saharan dust events affected
the Canary Islands. The first one, on 4-5 February 2020, registered a daily peak of $1000 \mu\text{g}\cdot\text{m}^{-3}$ over the Canary
Islands, and reached Iceland and Scandinavia (WMO, 2021). The second event, on 22-24 February 2020, stands as
the most intense dust event since records began in the Canary Islands (WMO, 2021; Cuevas et al., 2021). This dust
intrusion produced extremely high hourly concentrations of PM with diameters below $10 \mu\text{m}$ (PM₁₀) in the Canary
90 Islands, which exceeded $3000 \mu\text{g}\cdot\text{m}^{-3}$, and significantly impacted aviation (closure of the eight regional airports
and cancellation of ~ 1000 flights), agriculture and solar energy (Cuevas et al., 2021). This dust event also affected
the western half of the Iberian Peninsula. On 18 February 2021, a dust intrusion from the Sahara swept across much
of southern and central Europe, turning the snow-covered mountains of the Pyrenees and Alps orange because of
dust deposition. The exceptional Saharan dust transport to Europe during this episode was associated with
95 atmospheric rivers (Francis et al. 2022), carrying dust away as far as Scandinavia at the end of the month. On 15-
17 March 2022, a plume of Saharan dust travelled from northern Africa and across the Mediterranean into western
Europe, reaching the United Kingdom (WMO, 2023). The dust intrusion caused record PM₁₀ concentrations over
the Spanish air quality network, red- and orange-coloured skies and dust deposition on the ground across Europe,
especially visible on snow in the Pyrenees and Alps. The media disseminated shocking images of the dramatic
100 reduction in visibility in southern European cities and of people skiing on reddish-coloured snow in the Pyrenees
and the Alps. These dust events caused great social alarm, being sometimes associated with ongoing climate
change.

The most frequent time of the year for Saharan dust transport to the Mediterranean basin varies strongly from one
region to another. Overall, we can identify three large regions (Moulin et al., 1998; Basart et al., 2009; Pey et al.,
105 2013; Varga et al., 2014; Gkikas et al., 2013): eastern, central and western Mediterranean. In the eastern
Mediterranean, dust intrusions typically occur in spring (Israelevich et al., 2012), while in the western
Mediterranean the maximum Saharan dust transport occurs later, in summer (Salvador et al., 2014). The central
Mediterranean shows a bimodal distribution with two secondary dust peaks in spring and summer, respectively
(Barnaba and Gobbi, 2004; Marconi et al., 2014).

110 Thunderstorms and cyclones can produce high-speed winds that lift the dust and transport it thousands of miles
away through the air. Fiedler et al. (2014) reported that cyclones are related to up to 20 % of the total dust events
over the Mediterranean, although this contribution may increase during extreme dust events. Dust-related cyclones
are more frequent in spring and summer (see Moulin et al., 1998; Varga et al., 2014; Gkikas et al., 2016), coinciding

with the maximum of dust activity in the Mediterranean. As for winter, cyclones on the leeward side of the Atlas Mountains, also known as “Sharav cyclones” (Winstanley, 1972), and more recently as North African cyclones or Saharan cyclones have been associated with dust storms (Flaounas et al., 2022; and references herein). However, the release of dust from western Sahara, which is the primary source of dust events reaching the western Mediterranean (e.g. Gkikas et al., 2016), takes place at a considerable distance from recognized cyclogenesis areas. This implies that the transport of dust may be more closely linked to intense North Atlantic cyclones (Flaounas et al., 2022) or upper-level cut-off lows in the western Mediterranean (Portmann et al., 2021).

Both North Atlantic cyclones and western Mediterranean cut-off lows have been linked to large-scale atmospheric blocking (e.g. Trigo et al., 2004; Nieto et al., 2007). Blocking is a quasi-stationary equivalent-barotropic high-pressure system that can persist for several days or even weeks (Woollings et al., 2018; Kautz et al., 2022, and references herein). Atmospheric blocking interrupts the westerly flow in mid-latitudes and diverts the eastward path of synoptic cyclones (Rex, 1950). The region with the highest frequency of winter blocking is the eastern North Atlantic (Barriopedro et al., 2006; Wazneh et al., 2021, and references herein). In this region, blocking induces an equatorward displacement of the storm track towards the Mediterranean basin and North Africa region (Pfahl, 2014; Kautz et al., 2022). The meridional circulation and wavy jet induced by blocks can also result in deep troughs and cut-off lows upstream, downstream or in the equatorial flank of the block (Barriopedro et al., 2006; Nieto et al., 2007). This favourable large-scale environment for the development of cyclones and cut-off lows suggests an active role of blocking in the occurrence of dust intrusions in western Mediterranean.

Winter dust intrusions over the western Mediterranean have received little attention and existing studies on the associated atmospheric circulation have mainly focused on synoptic systems during individual case studies. For example, deep North Atlantic cyclones entering the western Mediterranean through North Morocco and Algeria were associated with an extreme dust intrusion in February 2017 over western Europe (Fernández et al., 2019; Oduber et al., 2019), whereas Bou Karam et al. (2009) described a strong cyclogenesis over the southern side of the Atlas Mountains during the African dust intrusion of February 2007 over western Mediterranean. However, a comprehensive climatological study of winter dust intrusions over the western Mediterranean, including the associated large-scale and synoptic atmospheric conditions is still lacking. This might be a result of 1) a relatively low frequency of occurrence of cyclone-related dust events in this region and season compared with those in spring and summer; 2) hampered assessment of the dust emission and transport in winter and spring due to cyclone-induced clouds (Schepanski et al., 2009), 3) less dust mobilization as a result of greater soil moisture during the winter season in North Africa (Gherboudj et al., 2015), and 4) the high wind speed thresholds needed to generate

dust in winter, which would require severe weather systems, such as deep upper-level troughs/cut-offs (Cowie et al., 2014). All this leads to poor knowledge of the drivers of dust transport to Europe in wintertime.

The main goal of the present study is two-fold. Firstly, to describe the winter anomaly of dust events that affected the Euro-Atlantic region (from now on the western Euro-Mediterranean region, WEM, for brevity) during the 2020-2022 period, and to assess to what extent they were exceptional compared to those recorded in the previous period of 2003-2019 (see Figure S1). Secondly, to characterise the atmospheric circulation systems associated with these winter dust events, emphasising the role of the large-scale weather patterns and blocking, which have been largely unexplored so far.

The paper is structured as follows. Section 2 describes the datasets and methods used to generate the results. Section 3 analyses the results of the dust-related datasets focusing on the identification of winter dust events over WEM for the 2003-2022 period, the associated meteorological systems and the comparison of the recent anomalous period of 2020-2022 with 2003-2019. In Section 4 are discussed the main results and identified those issues that need further investigation. Finally, the main conclusions are presented in Section 5.

2. Materials and methods

2.1 Catalogue and characterisation of dust events over the western Euro-Mediterranean

The identification and characterisation of dust events relies on satellite-based aerosol products. We have used aerosol optical depth at 550 nm (AOD) from the Moderate Resolution Imaging Spectroradiometer (MODIS; Kaufman et al., 1997) daily aerosol joint product (Collection 6.1), specifically, the Combined Land and Ocean product (Sayer et al., 2013) from Aqua satellite available between 2003 and 2022 with a $1^\circ \times 1^\circ$ spatial horizontal resolution. For the assessment of the meridional transport of dust, monthly mean AOD values have been computed for each month of the extended winters (from December to March) of 2003-2022 and over a large domain [27-60°N, 30°W-36°E], which encompasses northern Africa, the Mediterranean basin, Europe and the eastern North Atlantic. The identification of winter dust events in WEM, defined as [35-50°N, 20°W-5°E] is based on the daily AOD value averaged over the WEM (AOD_{avg}) and considers three categories (Moderate, Strong and Extreme events), following the thresholds proposed by Gkikas et al. (2016).

For an event to occur these conditions must persist for at least three consecutive days (Table 1). For example, a moderate dust event occurs when the daily AOD_{avg} is higher than the climatological mean (AOD_{Clim}) plus one standard deviation (SD), the two latter being defined from all daily AOD_{avg} values of the winter months (January-to-March) of the 2003-2022 period. This yields a threshold of 0.18 for AOD_{avg} , which must be exceeded for at least

three consecutive days. Similar values ($AOD > 0.15$) have also been associated with dust conditions in previous studies (Dubovic et al., 2002; Basart et al., 2009).

175 A drawback of the MODIS dataset might be the presence of clouds within some atmospheric systems. This could be particularly inconvenient in winter, when the strongest cyclones are expected over the Mediterranean Sea. However, we have verified that our MODIS-based approach, already used in previous studies (e.g., Moulin et al., 1998; Barnaba and Gobbi, 2004; Gkikas et al., 2016), is robust and reliable. Following Gkikas et al. (2016), we have also compared the MODIS AOD_{avg} with ground-based observations from Aerosol Robotic Network (AERONET) (Giles et al., 2019; see details in Section S2.1). In the comparison with five European stations (see 180 Figure S2), the correlation for dust days of the January-March period of 2003-2022 is in the 0.41-0.67 range (see Table S2).

Table 1. Aerosol optical depth (AOD) thresholds used to identify and classify the intensity of dust events. The imposed criteria must be satisfied for at least three consecutive days.

Type of dust event	Criteria	AOD thresholds
Moderate	$AOD_{Clim} + SD < AOD_{avg} \leq AOD_{Clim} + 2SD$	$0.18 < AOD_{avg} \leq 0.23$
Strong	$AOD_{Clim} + 2SD < AOD_{avg} \leq AOD_{Clim} + 4SD$	$0.23 < AOD_{avg} \leq 0.33$
Extreme	$AOD_{avg} > AOD_{Clim} + 4SD$	$AOD_{avg} > 0.33$

185 We have also used dust profiles based on the Vertical Feature Mask (VFM) aerosol product obtained from the National Aeronautics and Space Administration (NASA) Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observations (CALIPSO; Stephens et al., 2002; Winker et al., 2009), which is an active sensor measuring the backscatter signal at 532 and 1064 nm and the polarization at 532 nm (Winker et al., 2009). The identification of cloud and aerosol layers within the atmosphere (Vaughan et al., 2009) is made through the cloud aerosol discrimination algorithm (Liu et al., 2009; Kim et al., 2018). The VFM product aerosol subtyping algorithm distinguishes between tropospheric and stratospheric aerosols. It considers seven primary aerosol types: clean marine, dust, polluted continental, clean continental, polluted dust, smoke, and dusty marine (Liu et al., 2019). Here, we use the available CALIPSO Lidar Level 2 VFM product from NASA Atmospheric Science Data Center 190

195 (<https://asdc.larc.nasa.gov/project/CALIPSO>; last access 15 June 2023), which includes Version 4.2 (2007-2021) and Version 3.1 (for year 2022). Please, note that this CALIPSO dataset is the available products at by the time the results were processed. Despite the use of two different processing algorithms, the comparison between the two versions shows dust typing remains consistent (https://www-calipso.larc.nasa.gov/resources/calipso_users_guide/qs/cal_lid_l2_all_v4-20.php, last access 15 September 2023).

200 The aerosol profile products are generated at a horizontal resolution of 5 km, and the vertical resolution varies from 60 to 180 m depending on the altitude range and the parameter. For the present analysis, the VFM aerosol profile from 2007 to 2022 has been reduced to seven vertical ranges, namely, [0-1km), [1-2km), [2-4km), [4-6km), [6-8km), [8-10km) and [>10km).

2.2 Atmospheric circulation

205 To analyse the atmospheric circulation patterns associated with winter dust events over WEM we used data from the National Centre for Environmental Prediction/National Centre for Atmospheric Research (NCEP/NCAR) Reanalysis (Kalnay et al., 1996). The NCEP/NCAR Reanalysis is one of the state-of-the-art atmospheric reanalysis systems, providing consistent information from 1948 to present. For this study, six-hourly and daily mean fields at different pressure levels and $2.5^{\circ} \times 2.5^{\circ}$ spatial resolution for the 1991-2022 period have been used to characterise
210 the large-scale atmospheric conditions associated with the dust events identified in Section 2.1. Following international benchmark standards to assess changes in the meteorological conditions (WMO, 2007), we use 1991-2020 as the climatological normal period of analysis and the baseline to compute the anomalies of the meteorological fields. Although this reference period is not entirely covered by the MODIS observations employed for the characterisation of dust events (Section 2.1), the results of the atmospheric circulation analyses are robust
215 to the study period (e.g. the 2003-2022 dust period) and the atmospheric reanalysis employed (e.g. ERA5; Hersbach et al., 2020).

For the analysis of the atmospheric circulation, we employed several tools and algorithms, which allowed us to classify the atmospheric configurations associated with dust events, diagnose specific weather systems and circulation components (e.g. blocking and the jet stream), and establish linkages between these atmospheric
220 circulation patterns and dust intrusions. First, we have carried out a characterisation of the synoptic patterns associated with dust events in order to discriminate different flow configurations leading to dust events. The analysis of favourable patterns for dust intrusions is based on the k-means clustering algorithm (e.g., Wilks, 2011), which partitions data into a predefined number of groups, so that the inter-cluster (intra-cluster) Euclidean distance

is maximised (minimised). To obtain the typical patterns associated with dust events we have followed the methodology of Alonso-Pérez et al. (2011). The clustering has been applied to the daily anomalies of geopotential height at 500 hPa (Z500) of all dust days of the wintertime catalogue over the Euro-Atlantic sector, [20-70° N, 65° W-25° E]), and for the periods 2003-2019 and 2020-2022, separately. The selected number of clusters was limited to two, considering the relatively low number of dust days, particularly for the recent period of 2020-2022. The method assigns each dust day to one of the two clusters, allowing us to explore the two main Z500 patterns associated with WEM dust intrusions.

Second, we have explored if dust events tend to occur preferentially under specific recurrent patterns of the Euro-Atlantic sector. To categorise the large-scale atmospheric circulation in a limited number of recurrent weather regimes (WRs) we have followed Grams et al. (2017), which uses an extended year-round classification of the Euro-Atlantic atmospheric circulation in seven WRs. These WRs are derived from a k-means clustering in the phase space spanned by the seven leading empirical orthogonal functions of 10-day low-pass filtered Z500* fields, with Z500* denoting the normalized Z500 anomaly over the Euro-Atlantic sector. Following Michel and Rivière (2011), for each day and WR we compute a WR index as the spatial projection of the daily unfiltered Z500* onto the cluster centroid (i.e. the mean Z500* for all days in the cluster). The resulting indices are normalised (zero mean and one SD). A day is assigned to a given WR if the respective index is greater than 1 and higher than that of all other WRs. The seven WRs are defined as follows: the Zonal regime (ZO) and its variant Atlantic Trough (AT); the Scandinavian Trough (ScTr) and Atlantic Ridge (AR); and the blocking types, which include European Blocking (EuBL), Scandinavian Blocking (ScBL) and Greenland Blocking (GL). For further information on WR characteristics see Beerli and Grams (2019).

Finally, we analysed the occurrence of specific large-scale phenomena during dust days, including the North Atlantic jet stream and blocking events. The localisation of the North Atlantic eddy-driven jet stream and blocking patterns is performed on a daily basis, following the detection algorithms of Barriopedro et al. (2023) and Woollings et al. (2018), respectively. The retrieved fields (daily maps of binary fields) allow us to quantify the local frequency of jet and blocking occurrences during a given time interval or subsample of days. They are herein used to identify the favourable conditions (jet configurations and blocking activity) for the occurrence of dust events during the period of 2003-2019 and the recent period of 2020-2022. We note that the jet stream is detected every day of the analysed period, whereas blocking may or not be present on a given day. Therefore, the blocking catalogue is also employed to distinguish dust days with and without blocking, which allows us to assess the importance of blocking on dust.

255 The jet stream is diagnosed from 10-day low-pass filtered fields of the 925-700 hPa averaged zonal wind (see more details in Barriopedro et al., 2023). The latitudinal position of the jet is inferred from the daily meridional profile of the zonal wind at each longitude, which has been previously smoothed by averaging the zonal wind over a longitudinal sector of 60° width centred at that longitude. The latitudinal position of the jet is the latitude of the zonal wind maximum in the meridional profile. The poleward and equatorward boundaries of the jet correspond to the latitudes at each side of the jet latitude where the zonal wind maximum decreases by a given amount (defined as half the difference between the zonal wind peak and the meridional mean of the zonal wind over 15-75° N). All latitudes between the poleward and equatorward flanks of the jet identify the jet structure at that longitude. When applied to all longitudes of the Euro-Atlantic sector, the method provides a 2-D daily identification of the North Atlantic jet.

265 Blocking patterns are identified from Z500 daily fields, following the meridional gradient reversal method described in Woollings et al. (2018). For each day d and longitude λ , meridional Z500 gradients are computed to the north and south of a given latitude ϕ as follows:

$$GHGN(\lambda, \phi, d) = \frac{Z500(\lambda, \phi + \Delta, d) - Z500(\lambda, \phi, d)}{\Delta} \quad (1a)$$

$$GHGS(\lambda, \phi, d) = \frac{Z500(\lambda, \phi, d) - Z500(\lambda, \phi - \Delta, d)}{\Delta} \quad (1b)$$

$$GHGS2(\lambda, \phi, d) = \frac{Z500(\lambda, \phi - \Delta, d) - Z500(\lambda, \phi - 2\Delta, d)}{\Delta} \quad (1c)$$

270 where $45^\circ < \phi < 70^\circ$ N and $\Delta = 15^\circ$ latitude. GHGS is employed to detect flow reversals, whereas GHGN and GHGS2 aim to filter out other aspects unrelated to block. The meridional gradients in Eq. (1) are averaged over $\Delta/2$ in longitude, denoted by the overbars in Eq. (2):

$$\overline{GHGN}(\lambda, \phi, d) < -10 \text{ m}/^\circ \quad (2a)$$

$$\overline{GHGS}(\lambda, \phi, d) > 0 \text{ m}/^\circ \quad (2b)$$

275
$$\overline{GHGS2}(\lambda, \phi, d) < -5 \text{ m}/^\circ \quad (2c)$$

Daily blocking-like structures are identified as spatial clusters of contiguous grid points satisfying the conditions in Eq. (2) and covering a minimum areal extension of 500,000 km². Finally, the blocked areas detected in consecutive days are required to overlap and persist for at least five days to be considered a blocking event.

3. Results

280 The results are structured in two sections, which describe the climatological characteristics of wintertime dust events in WEM, paying special attention to the anomalous-dust 2020-2022 period (Section 3.1) and the atmospheric circulation associated with these events (Section 3.2).

3.1 Climatological assessment of winter dust events over WEM

285 We start assessing the spatial patterns of MODIS monthly mean AOD averaged for the 2020-2022 period in order to find footprints of the strong wintertime Saharan dust intrusions over the Mediterranean basin, and specifically over WEM at that time. The analysis is performed for each month of the extended winter (December-March) separately and compared with the corresponding maps of the previous period with available MODIS observations (2003-2019) (Figure 1).




Figure 1. MODIS/Aqua AOD at 550nm monthly averages for: (a, e) December, (b, f) January, (c, g) February, and (d, h) March of the 2002-2019 period (a-d) and the 2020-2022 period (e-h). The black box delimits the WEM spatial domain (i.e. [35°-50°N, 20°W-5°E]).

The climatological assessment (2003-2019 period) shows that wintertime Saharan dust intrusions towards the Mediterranean basin take place preferentially over the eastern Mediterranean through the coasts of Libya and western Egypt, affecting mainly Greece and southern Italy (Figures 1a-d). Moreover, dust intrusions increase in intensity and areal extent as winter progresses, reaching a maximum in March (Figure 1d). However, the impact on AOD over WEM is very small, which agrees with the results reported in previous studies (e.g., Varga et al., 2014; Gkikas et al., 2015; Gavrouzou et al., 2021). During 2020-2022, no significant changes were found in the AOD patterns of December and January, which showed very low AOD values (Figures 1e-f), like in the 2003-2019 period. However, in February 2020-2022 the AOD spatial pattern changed drastically: dust transport from western

Sahara occurred through Morocco and Algeria, yielding high AOD values over WEM (Figure 1g). Moreover, dust transport through the usual corridors of the eastern Mediterranean almost disappeared, which represents an extraordinary situation. These results suggest an increase of dust transport towards the western Mediterranean in February-March months of the 2020-2022 period compared to those recorded in the previous observational (2003-2019) period.

Based on these results, we have selected an appropriate study area (black rectangle in Figure 1) to detect dust intrusions over WEM. The southern boundary of the domain has been chosen far enough from northern Africa to avoid the detection of short-range weak dust intrusions associated with mesoscale processes. The northernmost boundary does not extend too far north to get non-negligible mean dust content over the geographical domain. We have analysed the time series (2003-2022) of the monthly mean AOD averaged over WEM to assess the interannual variability of Saharan dust intrusions for each month of the extended winter (Figure 2). The historical AOD series of December and January show relatively low values and few interannual variations, which contrasts with the higher and more variable AOD values at interannual scales for February and March. Notably, the February series of AOD displayed anomalously high values in our study region during three consecutive years (2020-2022). Similarly, the two highest AOD values of the March series were registered in 2021 and 2022. Although the climatological mean AOD over WEM increases in spring, the AOD values in March 2020-2022 is larger than in April 2020-2022, indicating that the anomalous character of late winter 2020-2022 did not extend towards the spring.





Figure 2. Time series of monthly averages of MODIS/Aqua AOD at 550 nm of the extended winters months (i.e., December, January, February, and March) of the 2003-2022 period and for WEM (black box in Figure 1).

As the frequency of dust intrusions in WEM is almost negligible in December and January (Figures 1a-b, e-f), from now on the analyses will focus on the months of February and March only. Our choice of the wintertime, defined as February-March (FM), is supported by both climatological arguments (higher AOD in late than in early winter; Figure 2) and the degree of extremeness of the 2020-2022 period as compared to the 2003-2019 period (see Section S2). Because of the coverage limitations associated to satellite-based products to capture the daily cycle (i.e. satellite overpasses or cloud contamination), the same dust event analysis is being done using the NASA MERRA-2 global reanalysis (Gelaro et al., 2017) considering the same period (i.e. 2003-2022) and study WEM domain. Atmospheric-chemistry reanalysis such as MERRA-2 provides representative and complete dust fields in space and time. The MERRA-2 confirmed the results obtained by MODIS (Section S2.2) showing the exceptional dust 2020-2022 period in comparison with 2003-2019 (see Figures S3 and S4) but also, previous decades (see Figures S5 and S6).

According to the AOD thresholds specified in Table 1, during FM 2003-2022 we have identified a total of 30 dust events (1.5 dust-events/year) over WEM. Recall that dust events lasting less than 3 days, or with daily mean AOD_{avg} lower than 0.18 are not considered in this study. The main characteristics of the dust events are summarised in the time series of Figure 3, including the number of dust days and its average AOD_{avg} for February and March of each year of the 2003-2022 period. This information is completed with Table 2, which shows the number of dust days and events for each calendar month and analysed period. In February the occurrence of dust events is quite irregular over the analysed period, including a period of six consecutive years (2010-2015) without any event (Figure 3). The intensity of February events tends to be moderate (the only one labelled as extreme occurred in 2017). The 2020-2022 period displayed an outstanding increase in the frequency of February dust intrusions, both in terms of events and days, which were approximately five and three times higher than those registered in the 2003-2019 period (Table 2). Indeed, five out of the 11 dust events recorded in February since 2003 have occurred in the 2020-2022 period, including one strong event (Figure 3). As expected, March, halfway between late winter and early spring, is the analysed month with the highest frequency of dust events in WEM. After a quiet period of four years (2016-2019), the comparatively shorter 2020-2022 period registered five dust events, including two extreme and two strong episodes. When FM is considered as a whole, the recent 2020-2022 period experienced an outstanding (eight-fold increase) in both the mean frequency of dust events and days as compared to 2003-2019.



350

Figure 3. Monthly time series (2003-2022) with the number of dust days (blue dots and lines; left y-axis) and the AOD at 550 nm averaged over WEM and for all dust days in the month (bars; right y-axis) for: (a) February; (b) March. The colour of the bars indicates the type of dust event: moderate ($0.18 < AOD_{avg} \leq 0.23$) in white, strong ($0.23 < AOD_{avg} \leq 0.33$) in grey, and extreme ($AOD_{avg} > 0.33$) in black.

355

In summary, every February and March of the 2020-2022 period reported dust intrusions, with an average of more than 15 dust days per FM period and at least one strong/extreme event. Overall, the analyses (Figure 3 and Table 2) confirm that the 2020-2022 period was the most active one in terms of frequency, duration, and intensity of dust events in WEM since observational records began. Compared to previous years of the 2003-2019 period, many of those dust events were long-lasting (>8 days) and intense ($AOD > 0.25$). The occurrence of intense dust events over WEM during 2020-2022 is significantly higher than during the 2003-2019 period (p-value of $\sim 10^{-6}$) as derived from the Wilcoxon rank sum test. In particular, the 14-day dust event of March 2022, with an AOD mean of 0.4 constitutes a milestone in dust transport from northern Africa to WEM. This event broke all records in terms of duration, intensity, and areal extent, and affected air quality in western and central Europe, with high PM₁₀ values ($50-100 \mu\text{g}\cdot\text{m}^{-3}$) (see Figure S14) even on high mountains such as the Pyrenees and the Alps (see Figure S15).

365

Table 2. Mean frequency of dust events ($\# \text{ year}^{-1}$) and dust days (days year^{-1}) for February and March of the 2003-2019 and 2020-2022 periods.

	2003-2019		2020-2022	
	Feb	Mar	Feb	Mar
Dust events ($\# \text{ year}^{-1}$)	0.35	0.82	1.66	1.66
Dust days ($\# \text{ year}^{-1}$)	1.6	6.8	5.0	12.3

370

According to the dust climatology of Gkikas et al. (2016), dust layers in the Mediterranean are normally observed between 2 and 6 km height during summer, while in winter they are mainly detected below 3 km and over the central and eastern Mediterranean Sea. Here, we have calculated the mean percentage of pure dust in seven tropospheric layers (between 0 and 10 km) of our geographical region based on the aerosol vertical profiles from CALIPSO (see Section 2.1), which are available for 2020-2022 and an earlier 2007-2019 period. For the 2007-2019 period, the results show slightly higher percentages of pure dust at higher altitudes (2-4 km) than those reported by Gkikas et al. (2016) (Table 3). The percentages of pure dust increased significantly in FM 2020-2022. In March 2020-2022, a non-negligible percentage of pure mineral dust is detected in an upper layer, between 4 and 6 km of altitude. These results suggest that the enhanced intensity of dust intrusions during 2020-2022 could have been mediated by atmospheric circulation processes lifting mineral dust from the desert to heights higher than usual. Other factors could have contributed to exacerbate the frequency, duration and/or intensity of recent dust events. During FM 2021-2022, the severe drought conditions over Europe (Garrido-Pérez et al., 2024) affected the arable areas of northern Morocco and Algeria (see Figure S7). Indeed, during the winter months of 2022, the monthly mean series of the Normalized Difference Vegetation Index (NDVI), which accounts for vegetation cover and activity, recorded some of the lowest values of 2003-2022 over the northernmost strip of the Maghreb (see Figure S8).

380

385

390 **Table 3.** Mean fraction of pure dust (with respect to the total available aerosol observations) in different vertical layers for the FM dust days of the 2007-2019 and 2020-2022 periods, as inferred from the CALIPSO Vertical Feature Mask (VFM) product.

	2007-2019			2020-2022		
	Jan	Feb	Mar	Jan	Feb	Mar
0-1 km	0.02	0.03	0.03	0.02	0.05	0.04
1-2 km	0.02	0.04	0.04	0.03	0.08	0.06
2-4 km	0.02	0.05	0.05	0.04	0.12	0.14
4-6 km	0.02	0.03	0.03	0.02	0.03	0.09
6-8 km	0.02	0.02	0.03	0.02	0.02	0.04
8-10 km	0.02	0.01	0.02	0.02	0.02	0.02
> 10 km	0.02	0.01	0.02	0.02	0.02	0.01


3.2 Atmospheric drivers of winter dust events over WEM

395 This section describes the favourable patterns and preferred WRs for the occurrence of dust events (Section 3.2.1), as well as their associated meteorological systems at synoptic and large scales (Section 3.2.2). The diversity of atmospheric configurations will be assessed through statistical tools (clusters and WRs) and automatic algorithms tracking specific phenomena (blocking and jet stream), as described in Section 2.2. The meteorological systems associated with dust events at synoptic scales (cyclones, cut-off lows, etc.) have been identified qualitatively (by visual inspection), which has also allowed us to confirm the results of the objective analyses.

400

3.2.1. Dust-related patterns and weather regimes

Figure 4 shows the two most recurrent Z500 anomaly patterns obtained from the k-means clustering of FM dust days for the 2003-2019 and 2020-2022 periods. As explained in Section 2.2, we have only retained two clusters, which yields a robust partitioning of the data while still retaining the major features of the dominant patterns associated with dust events. The dominant pattern in both periods (Cluster C#1; Figures 4a, c) is characterised by a meridional pressure dipole with positive Z500 anomalies between the British Isles and Scandinavia, and negative Z500 anomalies to the south. This favourable configuration corresponds to a European Rex-like block (Rex, 1950; Sousa et al., 2021). Comparatively, the second most frequent pattern during FM dust days (Cluster C#2; Figures 4b, d) shows a more zonally-oriented dipole, but with noticeable differences between the two periods. In the 2003-2019 period, the C#2 pattern (45% of dust days of that period) featured a pronounced trough over the eastern Atlantic, and positive Z500 anomalies over the mid-latitudes of eastern Europe. In the 2020-2022 period, C#2 (30%) displayed a cut-off low signature in the Z500 anomalous field over the subtropical eastern Atlantic, accompanied by positive Z500 anomalies over the Euro-Mediterranean region.



415

Figure 4. Two clusters (C#1 and C#2) of Z500 anomalies (in m) for the FM dust days of the 2003-2019 (a-b) and the 2020-2022 (c-d) period. The relative frequency of each cluster (in % with respect to the total number of dust days of each period) is indicated on the top.

420 These results show that blocking (C#1) is a recurrent driver of dust intrusions, being present in both periods, albeit with different frequencies of occurrence. There are also differences in the C#2 spatial patterns of the two periods, suggesting that favourable configurations other than high-latitude blocks occurred in the 2020-2022 period. Likewise, the occurrence of cut-off lows was a common signature of dust intrusions during the 2020-2022 period. They were accompanied by either high-latitude blocks over Europe (C#1; Figure 4c) or high-pressure systems at

425 at lower latitudes (C#2; Figure 4d). There are also signatures of subtropical cut-off lows during dust days of the 2003-2019 period (mainly in C#1; Figure 4a). Another distinctive signature of the recent 2020-2022 period was the prevalence of positive Z500 anomalies over southwestern Europe, which is absent in the cluster patterns of the 2003-2019 period (cf. Figures 4a, b, d). These results suggest that cut-off lows are actively involved in dust intrusions. At larger spatial scales, the enhanced dust activity of the recent 2020-2022 period could partially be

430 explained by a high frequency of favourable configurations, including both recurrent (high-latitude blocking) and uncommon (e.g. subtropical high-pressure systems) dust-related patterns of the 2003-2019 period.




Figure 5. Probability of occurrence of weather regimes (WRs) during dust days of 2003-2019 (black bars) and 2020-2022 (red bars). The following acronyms are used. No: no regime; AT: Atlantic trough; AR: Atlantic ridge; GL: Greenland blocking; EuBL: European blocking; ScBL: Scandinavian blocking; ZO: zonal regime and ScTr: Scandinavian trough.

The dust-related configurations identified in Figure 4 do not necessarily match with the dominant large-scale patterns responsible for the day-to-day variability of the extratropical circulation (WRs). To investigate whether the occurrence of dust days over WEM is favoured by specific recurrent patterns, we used the WR classification described in Section 2.2. Figure 5 shows the frequency density distribution of the seven WRs for the FM dust days of each period. We note that none of the WRs coincide exactly with the dust-related patterns of Figure 4, which is reflected in the dispersion of dust days across different WRs. As such, dust intrusions can occur under different WRs, stressing again the multiplicity of large-scale patterns compatible with dust events. In spite of this, a more detailed analysis of WRs and their correspondence with the clusters of Figure 4 reveal the following results:

- The WR with a higher probability of occurrence during dust days is the EuBL WR (Figure 5), which is consistent with the high frequency of occurrence of the C#1 clusters in the two periods (Figures 4a, c).
- Several WRs featuring high-pressure systems at subtropical latitudes of the Euro-Mediterranean (e.g. AR and AT) show a small increase in probability of occurrence during dust days of the 2020-2022 period, which is consistent with the distinctive positive Z500 anomalies over the Euro-Mediterranean region of C#2 (Figure 4d).
- The frequency of ZO also increased (by a factor of two) during dust days of the 2020-2022 period with respect to 2003-2019. ZO does not have a direct correspondence with any of the clusters of Figure 4 but its occurrence is consistent with the presence of European subtropical ridges, which are associated with an intensification of the mid-latitude zonal flow (Sousa et al., 2021). Therefore, the high frequency of ZO in the 2020-2022 period could be interpreted as an increase in subtropical ridges (readily apparent in C#2; Figure 4d)

Therefore, dust intrusions can occur during WRs featuring high-pressure systems at different latitudes (mainly European Blocking, EuBl, and Atlantic Ridge, AR), which is consistent with the clustering analysis. The relative influence of blocking and other high-pressure systems is analysed in more detail in the next section.

3.2.2. Meteorological systems

In this section, we analyse the weather systems associated with WEM dust events following the methods described in Section 2.2. First, we carried out a preliminary qualitatively analysis of the Z fields at different pressure levels (from 925 to 100 hPa) for each day of the 30 dust events of 2003-2022. The results revealed the following common signatures: 1) synoptic systems: negative Z anomalies (typically < -80 m) at all levels from 925 to 200 hPa located in the subtropical region of the eastern North Atlantic or in north-western Africa, which are identified as cut-off lows; and 2) large-scale systems: relatively extensive regions of positive Z anomalies (typically > 100 m at 500 hPa). The latter can occur under a variety of flow configurations and locations (high pressure systems from subtropical to high latitudes), as confirmed by Figures 4 and 5. A substantial number of cases resemble the C#1 patterns (Figures 4a, e) or EuBL WR, with positive Z anomalies located to the north of a cut-off low, therefore forming a dipole-like block (Rex, 1950; Barriopedro et al., 2010). A second group, like AR, reveals positive Z anomalies at comparatively lower latitudes, which are more characteristic of subtropical ridges (Sousa et al., 2021) or low-latitude blocks (Davini et al., 2012) over the eastern North Atlantic or western Mediterranean. The next sections describe the synoptic (cut-off lows) and large-scale (high-pressure) systems associated with dust days. We also assess the jet stream, since these weather systems are often associated with substantial departures of the westerlies and the storm track activity (Trigo et al., 2004). The analysis relies on composites anomalies for the 2003-2019 and 2020-2022 period, separately, to emphasise distinctive features of the recent anomalous period.

Section S4 provides a more detailed analysis of the three dust episodes labelled as “extreme” in the 2003-2022 period, which occurred on 20-24 February 2017 (Section S4.1), 27-31 March 2021 (Section S4.2), and 15-31 March 2022 (Section 4.3). Furthermore, for the three case studies analysed, the geographical location as well as the activation time of each dust hotspot has also been identified manually (Schepanski et al., 2007, 2009, 2012) by using the 15-min EUMETSAT RGB dust animations (MetOffice; EUMETSAT, 2022; see timelapse videos at <https://repositorio.aemet.es/handle/20.500.11765/15054>). Although the results should not be generalised to all dust events, the inspection of case studies confirms that cut-off lows are commonly involved during WEM dust events. Moreover, the three cases featured large-scale high-pressure systems with different configurations and at different latitudes, ranging from high-latitude blocks to subtropical ridges (Sousa et al., 2021). The synoptic and large-scale patterns favourable for the occurrence of dust events will be addressed in the next sections.


3.2.2.1. Synoptic scales

490 For all dust events, the vertical cross-section shows negative Z anomalies between 925 and 200 hPa (not shown). These Z anomalies are in all cases prominent at 200 hPa and weaken towards the surface, consistent with the typical signatures of upper-level cut-off lows (Nieto et al., 2005, and references herein). Indeed, more than half (65%) of the FM dust events of the 2003-2019 period concurred with a cut-off low. This relationship strengthened during the 2020-2022 period, particularly in March, when all dust events were accompanied by cut-off lows. In few cases, 495 a near-surface cyclone initiated during the later phases of the cut-off low. For dust days, cut-off lows are normally located between the Canary Islands and the Iberian Peninsula, activating the dust sources over Morocco and Algeria and transporting dust northwards (see Section S4). Noticeable negative Z anomalies were also found in the lower stratosphere (100 hPa) during at least 50 % of the 2003-2019 cases and all dust events of the 2020-2022 period. Although the negative Z anomalies at the lowest (925 hPa) level are weaker than at higher levels (not shown), they 500 are intense enough to generate surface winds exceeding the wind speed threshold for dust mobilization in western Sahara, which is between 5 and 12.5 m·s⁻¹ (Helgren and Prospero, 1987). This is supported by 6-hourly surface wind data from NCEP/NCAR reanalysis over Morocco, Algeria, western Sahara and Mauritania (see Section S5).

For some dust events the centres of these cut-off lows remained stationary over the Atlantic coast of Morocco during ~3-4 days, mobilizing dust on its eastern flank (see some examples in Figure S16). However, stationary cut-off lows were relatively uncommon; in most cases, cut-off lows moved eastward from the subtropical North 505 Atlantic to the western Mediterranean (see e.g. the cut-off low to the west of the Atlas Mountains during the events of 20-24 February 2017 in Section S4.1 and 27-31 March 2021 in Section S4.2). The same event can comprise two or more successive cut-off lows. This situation was identified in at least one third of the 2003-2019 dust events, but in almost all March dust events of the 2020-2022 period (see e.g. the sequence of cut-off lows over the coasts of 510 Morocco and Algeria in the 15-31 March 2022 event; Section S4.3). The maximum number of cut-off lows in a dust event was four, registered in the historical episode of 15-31 March 2022 (see case #3 in Section S4.3). Finally, the cut-off lows detected during dust events tend to be long-lasting, in agreement with the characteristic time scale (~1 week) of the accompanying high-pressure systems (Barriopedro et al., 2006), which act to weaken the mean zonal flow and slow down the reabsorption of cut-off lows by the westerlies. Apart from this, we do not identify 515 distinctive features of the cut-off lows associated with dust events. Indeed, their characteristics are consistent with those reported in previous climatological studies (e.g. Nieto et al., 2005, 2007): 1) winter cut-off low occurrence is mostly related to blocking events; 2) winter cut-off lows of the European sector tend to occur in northern Africa and southern Europe; 3) more than 80% of cut-off lows are highly transient during their life cycle.

520 3.2.2.2. Large scale

To assess the influence of large-scale blocking and jet stream configurations, we computed the local frequency of blocking and jet stream occurrence for all FM dust days and compared them with their climatological frequencies (regardless of dust occurrence). Figures 6a and c show blocking and jet frequency exceedances above the climatology during dust days, for the 2003-2019 and the 2020-2022 periods. Overall, dust days are associated with enhanced blocking activity over a latitudinal band centred at mid latitudes (~50-60° N) and extending from the eastern Atlantic to Scandinavia (Figure 6a). Therein, the probability of blocking occurrence increases up to ~12% (red shading in Figure 6a), which more than doubles the expected values from the climatology (contours in Figure 6a). The North Atlantic jet is shifted poleward, with preferred locations over high latitudes and reduced frequencies over typical jet latitudes (Figure 6c), consistent with the obstruction of the westerlies by the blocking action and its diverting effect on the jet (Trigo et al., 2004; Barriopedro et al., 2010; Sousa et al., 2018). Therefore, European blocking with poleward shifted jets represent a favourable configuration for dust occurrence.



535 **Figure 6.** Blocking (a,b) and jet stream (c,d) frequency anomaly during dust days (in our study region) in FM of the 2003-
 2019 (a,c) and 2020-2022 (b,d) period. Frequencies are expressed in percentage of dust days of the FM period and anomalies
 (colour shading) are defined with respect to the expected frequency of occurrence (contours). Dots denote significant positive
 anomalies at $p < 0.05$ (i.e. grid points where local blocking and jet frequency is significantly higher than expected), as inferred
 from a 500-trial bootstrap. Significance is assessed with a bootstrapping of 500 members, each one derived from a random
 choice of FM days of the same size as the number of dust days. The expected frequency of occurrence is defined as the mean
 540 of the 500 random members, all with the same sample size as the number of considered days but random dates of occurrence.

545 Figures 6b and d show the anomalies in blocking and jet frequency for FM 2020-2022. Overall, blocking activity
 was not significantly higher than the climatology, suggesting that the anomalous frequency of dust days in 2020-
 2022 cannot be fully explained by a corresponding blocking increase over the favourable region for dust intrusions.
 However, the lack of statistical significance in the blocking and jet frequency of FM 2020-2022 partially results
 from contrasting signatures between the corresponding patterns for February and March 2020-2022 (Figure S17).
 March 2020-2022 was characterized by poleward jets and enhanced blocking over the climatological region of

occurrence, like in Figure 6(a) and the canonical pattern associated with dust (cluster C#1 of Figure 4). Differently, during February 2020-2022, blocking activity was almost suppressed over central Europe, but it was recurrent at lower latitudes, resulting in a significant increase over the Mediterranean (also apparent in Figure 4d). A high frequency of low-latitude blocks is commonly associated with subtropical ridges, which can temporarily exhibit meridional gradient reversals (Sousa et al., 2021). These high-pressure low-latitude structures reinforce the zonal wind at their poleward flanks, favouring a mid-latitude intensification of the jet (consistent with the zonal flow configuration of Figure 6d), rather than its poleward migration (see e.g. Sousa et al., 2018; Woollings et al., 2018; Barriopedro et al., 2023). The enhanced occurrence of high-pressure systems at subtropical latitudes during the 2020-2022 period is also consistent with the high frequency of AR, AT and ZO WRs (Figure 5).

These results indicate that dust events can be favoured by high-pressure systems at very different latitudes, ranging from subtropical ridges with mid-latitude jets to high-latitude blocks with poleward-shifted jets. The occurrence of subtropical ridges was particularly enhanced during dust days of FM 2020-2022, weakening the strong block-dust linkage reported over the historical period. To address this question, we quantified the association between blocking and dust days as the percentage of dust days when a block was detected at any latitude of the [25° W-30° E] domain. During the 2003-2019 period, more than 65% of the FM dust days coincided with block, which is significantly higher than that expected by random chance (binomial test, $p < 0.01$). This conditional probability decreased to ~50% during FM 2020-2022 ($p > 0.1$). Therefore, although blocking activity was favourable for the anomalously high frequency of FM dust days in 2020-2022, they explained less exceedances than those expected from the historical record.

To stress the key role of subtropical (non-blocking) high-pressure systems in driving dust events of the 2020-2022 period, we computed the composites of Z500 anomalies for dust days with and without simultaneous blocking over the [25°W-30°E] domain (Figure 7). The analysis is performed for the 2003-2019 and 2020-2022 periods separately, aiming to uncover the type of weather systems (other than blocking) that contributed to the anomalous frequency of dust intrusions in 2020-2022. As expected, block-dust concurrence during the 2003-2019 period shows the typical configuration of a high-low pressure dipole, with low pressures to the south of a blocking high that is fully detached from the subtropical belt (Figure 7a). During the 2020-2022 period, dust-related blocks were located at comparatively more easterly longitudes, and displayed higher Z500 anomalies, which extended towards the Mediterranean (Figure 7b). This equatorial elongation of Z500 anomalies denotes an Omega-like blocking pattern (i.e., without complete detachment of the blocking high; e.g. Barriopedro et al., 2010). The distinctive structure of the 2020-2022 blocked flows becomes clear in Figure 7c, which shows the Z500 difference between

the blocking-like patterns associated with dust days in 2020-2022 and 2003-2019. The pattern evidences an eastern shift and equatorial elongation of 2020-2022 blocking (positive Z500 differences over continental Europe), with a deeper trough upstream of the blocking high (negative Z500 differences over the eastern Atlantic).
580




Figure 7. Composites of Z500 anomalies (in m) for dust days of: (a), (d), and (c), the 2003-2019 period; (b), (e), and (h), the 2020-2022 period; (c), (f), and (i), the difference of composites between the 2020-2022 and 2003-2019 periods. Top and middle panels show the composites for dust days ((a), (b) and (c)) with and without blocking ((d), (e), and (f)) panels show the composites for dust days with and without blocking over the 25° W-30° E domain, respectively.
585

On the other hand, non-blocking dust days are favoured by positive Z500 anomalies over central Europe and the Mediterranean (Figures 7d, e). These situations denote high-pressure systems with no meridional reversals of the Z500 gradient (i.e. no block), which is a distinctive feature of pure subtropical ridges (Sousa et al., 2021). This non-blocking pattern was recurrent during 2020-2022 (~44% of dust days) and it actually concurred with more dust days than in the historical period, explaining the reduced intervention of blocking. Moreover, the difference between the non-blocking dust days composites of Z500 for 2020-2022 and 2003-2019 indicates an anomalous occurrence and/or location of cut-off lows over the Canary Islands and eastern northern Africa during 2020-2022 (Figure 7f). The results are consistent with those of Section 3.2.2.1, which report an increased percentage of dust
590
595

events accompanied by cut-off lows during the 2020-2022 period. They also agree with the emergence of a dust-related pattern with positive Z500 anomalies at subtropical latitudes (C#2; Figure 4d) and the increased frequency of WRs featuring high-pressure signatures over the Mediterranean (Figure 5). Therefore, while blocking was still the dominant pattern associated with dust days in 2020-2022, an enhanced occurrence of subtropical ridges and cut-off lows at low latitudes also contributed to the outstanding frequency of dust intrusions. Indeed, the high frequency of dust days associated with high-pressure systems at low-latitudes (different to blocking) suggests that these systems were particularly relevant for the anomalous frequency of dust events in 2020-2022. Dedicated studies are required to address the causes of the anomalous activity of cut-off lows and associated subtropical ridges (blocking) in February (March) 2020-2022.

600

605

4. Discussion

In this study, we analyse winter dust events over WEM for the 2003-2022 period, paying special attention to the recent period of exceptional dust activity (FM 2020-2022). This section discusses additional pertinent questions not addressed in the results. For the summer season, Cuevas et al. (2017) found that Saharan dust pulses towards the Mediterranean and the North Atlantic were the result of the coupling between mesoscale dust uplift processes and longitudinal shifts of the Sahara heat low governed by the North African Dipole and Rossby waves. Similarly, winter dust events in WEM may constitute a good example of atmospheric processes interacting at very different scales: microscale to mesoscale processes, necessary for dust lifting and mobilization (Knippertz and Todd, 2012) from small dust sources located in Morocco and Algeria (Ginoux et al., 2012), synoptic-scale systems such as cut-off lows driving its transport (Muñoz et al., 2020), and a large-scale environment favouring these synoptic disturbances and the persistence of high AOD values. The latter can occur under a variety of large-scale high-pressure systems, including European blocking with poleward shifted jets (as in the 2003-2019 period; Figures 6a, c) or Mediterranean subtropical ridges with an intensified mid-latitude jet (e.g., as in the 2020-2022 period; Figures 6b, d). Previous studies have reported a strong relationship between the occurrence of cut-off lows and blocking (e.g. Nieto et al., 2007), and our analyses suggest a similar correspondence between cut-off lows and subtropical ridges, particularly strong during the 2020-2022 period. However, it is unclear whether subtropical ridges and cut-off lows represent independent phenomena (and hence their unusually high concurrence in 2020-2022 was by chance) or they can be dynamically linked, as in the case of blocking and cut-off lows. Additional studies are encouraged to address connections between subtropical ridges and cut-off lows. It is also unclear if the influence

610

615

620

625 of cut-off lows on dust events depends on specific characteristics of the weather system. In this sense, dust-related cut-off lows of the 2020-2022 period were deeper and extended to higher pressure levels than those of the 2003-2019 period. This seems to be in agreement with the enhanced percentage of pure dust in the lower troposphere (0-4 km) and the non-negligible concentration of pure dust in upper layers (4-6 km) recorded during recent dust events.

To the author's knowledge, this work represents the first comprehensive study showing a close relationship between Saharan dust events and different types of large-scale high-pressure systems, namely blocking and subtropical ridges. Reliable forecasts of these weather systems might enhance the mean-range predictability of dust events, which is highly valuable since these episodes can lead to severe pollution episodes locally and in remote regions. The relationship between blocking, and air quality is also supported by recent studies describing atmospheric blocking influences on air stagnation and PM10 concentrations in Europe (Garrido-Pérez et al., 2018). High surface ozone levels in Europe have also been related to blocking (Otero et al., 2022), as well as to the location of subtropical ridges (Ordóñez et al., 2017). Therefore, blocks and ridges represent potential predictors of pollution events in Europe, as also shown by Maddison et al. (2021) using statistical models.

Atmospheric rivers have also been pointed out as an important triggering factor of Saharan dust intrusions in Europe (Francis et al., 2022). Preliminary analyses based on a visual identification of atmospheric rivers -herein defined as long and narrow structures with vertically integrated water vapour higher than 20 Kg m^{-2} (Gimeno et al., 2014)- indicate that ~10% of the FM dust days of our catalogue concurred with atmospheric rivers. The linkage strengthens during the 2020-2022 period (~17% of the FM dust days with atmospheric rivers), which is comparable to the values reported by Francis et al. (2022), but much lower than that obtained by conditioning on cut-off lows or blocks (Section 3.2.2). Moreover, atmospheric rivers might not be fully independent of blocking and cut-off lows, as suggested in previous studies (Benedict et al., 2019). Indeed, case studies (Francis et al., 2022) and composite analyses (Lorente-Plazas et al., 2020) of Mediterranean atmospheric rivers reveal high-low pressure configurations that are consistent with those reported herein for WEM dust events (Figure 7). In any case, atmospheric rivers might play an important role in the wet deposition of dust, as during some of the extreme events analysed herein. Additional studies are encouraged to address the similarities and differences between dry and wet dust events in terms of driving processes and associated impacts.

The shocking images of European ski slopes covered by Saharan dust that were published by the media during the winters of 2021-2022 raised the question of whether these events are becoming more likely because of climate change. Several reasons prevent making attribution statements at this stage. First, our catalogue of dust events is short and shows large temporal variations (such as the sharp increase of 2020-2022) and relatively long periods

655 with no Saharan intrusions, which hamper the detection of long-term trends. Secondly, the atmospheric weather
systems linked to dust events do not exhibit robust trends and/or our confidence in their regional climate change
responses is low. For example, winter blocking trends over the last decades are seasonally and spatially
heterogeneous, with no robust evidence of generalised significant changes, partially due to discrepancies among
660 the detection indices and their large variability on interannual-to-multidecadal time scales (e.g. Woollings et al.,
2018), which makes trends sensitive to the analysed period (Sousa et al., 2021). Although several studies point to
future decreases in blocking frequency, the spatial and seasonal patterns of change can be complex or depend on
the characteristic addressed. For example, recent studies have reported both significant decreases (Davini and
D'Andrea, 2020) and non-significant changes (Bacer et al., 2022) in the frequency and duration of Northern
Hemisphere blocking during the 21st century, while other attributes such as blocking size may increase in response
665 to climate change (Nabizadeh et al., 2019). On the other hand, subtropical ridges have been less addressed than
blocking, and hence their observed and projected changes remain uncertain (Sousa et al., 2021). Finally, the
detection of trends in cut-off lows is complicated by the diversity of detection approaches (which can be sensitive
to methodological aspects such as the spatial resolution or the vertical level), the large interannual variability at
regional scales and the influence of internal modes of variability. Nieto et al. (2007) did not report significant trends
670 in the frequency of winter cut-of lows over the European sector for the 1958-1998 period. Although this period
does not inform on the last decades (with stronger anthropogenic forcing) or our study period, it suggests weak cut-
off low responses to long-term climate change. However, the key role of Euro-Atlantic cut-off lows in WEM dust
activity calls for updated analyses of their trends and variability.

In addition to atmospheric dynamics, the sources and atmospheric loading of dust are sensitive to changes in other
675 climate components and forcings, including vegetation and land use (IPCC, 2022). In this sense, the 2020-2022
period coincided with dry conditions and the lowest winter NDVI monthly values (Figure S8). Warm Sea Surface
Temperature conditions, as those reported in the easternmost subtropical North Atlantic and the westernmost
Mediterranean in the period 2003-2023 (Figure S20) seem to play a crucial role in the development and evolution
of Mediterranean cyclones, by fuelling them with energy and water vapour (Stathopoulos et al., 2020), which may
680 also favour dust intrusions (Flaounas et al., 2015). Finally, Thomas and Nigam (2018) reported a modest northward
advance of the Sahara Desert during the boreal winter. Therefore, desertification processes and their complex
interactions with the ocean, orography and soil conditions over northern Africa could also affect the frequency
and/or severity of winter dust intrusions.

685 5. Summary and conclusions

This work provides a catalogue of dust events over the western Euro-Mediterranean region (WEM) for 2003-2022, focusing on their characteristics during the unusual 2020-2022 period of exceptional activity. Dust events are detected based on satellite aerosol products from MODIS/Aqua. This study also contributes to the understanding of WEM dust events and their synoptic and large-scale environments, which have been poorly addressed in the literature. It shows that wintertime WEM dust events are linked to meridional flows typically induced by subtropical cut-off lows and/or large-scale high-pressure systems at mid or low latitudes. The main results can be summarized as follows:

- Winter dust intrusions over WEM show irregular activity and large differences among the winter months. December and January are characterised by very low frequencies, and the winter months of highest activity (February and March, FM) display large multi-year variations characterised by calm (e.g. 2018-2019) and active (2020-2022) periods (see Figure 3). It is unclear if these intra-seasonal differences are meaningful or just reflect the effects of limited sampling (a short record with a low number of events).
- More than half of all FM dust events over WEM were accompanied by cut-off lows between the Canary Islands and the Iberian Peninsula, which is substantially higher than the frequencies for central and eastern Mediterranean events reported in literature. These results indicate that cut-off lows are a key synoptic driver of WEM dust intrusions in winter. This association increased during the 2020-2022 period, and all March dust events of that period were concurrent with cut-off lows, suggesting an unusual frequency and/or efficiency of cut-off lows.
- Concerning the large-scale circulation, WEM dust days are associated with enhanced blocking activity over the mid-latitudes of the Euro-Atlantic sector. This meridional pressure dipole favours the obstruction of the westerlies, which are shifted poleward by the blocking high, and the occurrence of cut-off lows at subtropical latitudes. Although European blocking is not a necessary condition for the occurrence of dust, it represents the most recurrent pattern associated with FM dust days. Consistently, WEM dust days tend to occur under anticyclonic weather regimes.
- Enhanced mid-latitude blocking and poleward jet configurations were favourable for the outstanding occurrence of WEM dust days, particularly during March 2020-2022. However, a substantial fraction of dust days (mainly those in February 2020-2022) concurred with high-pressure structures at low latitudes or subtropical ridges, which favour a mid-latitude intensification of the jet, rather than its

715 poleward migration. Although subtropical ridges do not necessarily develop secluded troughs, they
were frequently accompanied by cut-off lows enhancing the southerly winds over northern Africa.

To assess whether the anomalous conditions persisted during the winter of 2023 we have evaluated the already
available data for February and March 2023 (see Section 7). No dust event was recorded in March 2023, however,
a six-day strong dust event occurred on 18-23 February 2023 (Figure S19). Like for the dust events of the 2020-
720 2022 period, dust uplift and transport were associated with two consecutive cut-off lows in the vicinity of the
Canary Islands. This event was also accompanied by positive Z500 anomalies over central Europe and the
Mediterranean, like the characteristic pattern of subtropical ridges (Figure 4d). Further research is required to
address the ultimate causes of the anomalous activity of dust events and their atmospheric drivers.

725 **Data availability**

The MODIS and MERRA-2 data used in this work can be freely obtained from
<https://giovanni.gsfc.nasa.gov/giovanni>. CALIPSO Level 2 Vertical Feature Mask (VFM) Version 3.1 and 4.2
product available from January to March 2007-2022 download from NASA's Atmospheric Science Data Center
(ASDC, <https://asdc.larc.nasa.gov/project/CALIPSO>). NCEP/NCAR Reanalysis data used in this work were
730 obtained the NOAA PSL, Boulder, Colorado, USA, from their website at <https://psl.noaa.gov/> (Kalnay et al.,
1996). In the supplementary material is also considered: AOD direct-sun Version 3 Level 1.5 data can be download
from the AERONET database (<https://aeronet.gsfc.nasa.gov>; Holben et al., 1998; Giles et al., 2019). Daily mean
PM10 data used was obtained from the European Environment Agency's air quality database (AirBase;
<https://www.eea.europa.eu/data-and-maps/data/aqereporting-9>). EUMESAT RGB dust images (and processed
735 with the Man computer Interactive Data Access System, McIDAS) as well as dust forecasts from the MONARCH
(Multiscale Online Non-hydrostatic Atmosphere Chemistry) are available through the WMO Barcelona Dust
Regional Center (<https://dust.aemet.es/products/daily-dust-products>).

Author contributions

740 EC designed the study and coordinated the different contributions of the analysis. EC, DB and SB discussed the
main results and wrote the draft, with contributions from all co-authors. RDG, JJB, OG and AB collected and
prepared MODIS, MERRA-2 and AERONET datasets for the elaboration of the dust catalogue. SB collected and

745 prepared the CALIPSO datasets used in the dust catalogue. EC performed the qualitatively analyses of the atmospheric circulation patterns associated with winter dust events. SAP provided the main atmospheric patterns
750 favour dust intrusions using the k-means clustering algorithm. SAP and JJGA prepared the results on weather regimes. DB provided the localisation of the jet stream and blocking patterns using the detection algorithms of Barriopedro et al. (2023) and Woollings et al. (2018), respectively. EC and SB led the discussion on the identification of the dust events. DB prepared and lead the discussion on meteorological drivers of dust events. EW, DS and GGC collected and prepared the modelling and monitoring datasets included in the study cases considered in the Supplement. All co-authors reviewed and contributed to the final editing of the paper.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

755 The authors thank all the principal investigators and their staff, for establishing and maintaining the NASA and PHOTONS AERONET sites, the NASA MODIS and CALIPSO mission, the NASA MERRA scientists and associated NASA personnel, to produce the data used in this study. Also, EUMESAT and WMO are sincerely acknowledge for the provision of monitoring and forecasting products. Ongoing support for GrADS is provided by
760 an omnibus grant jointly funded by the NSF, NOAA and NASA that forms the core support for all research at COLA. NCEP/NCAR data is provided by the NOAA/ESRL PSL, Boulder, Colorado, USA, from their Website at <https://psl.noaa.gov/>. E. Cuevas, E. Werner, G. García-Castrillo, and S. Basart acknowledge the WMO Barcelona Dust Regional Center. S. Basart acknowledges CAMS-84 and CAMS2-82 (part of the Copernicus Atmospheric Monitoring Services, CAMS). Also, Dr. Christian Grams, Karlsruher Institut für Technologie (KIT), is
765 acknowledged for providing the weather regimes data in ERA5 reanalysis used in this work. The authors thank Ms Elena Morato, head of the AEMET Library, for hosting the timelapses of the three case studies in the AEMET ARCIMIS repository. Finally, the authors thank the editor and two anonymous referees who kindly reviewed the earlier version of this manuscript and provided valuable suggestions and comments.

770 **References**

- Adebiyi, A.A. and Kok, J.F.: Climate models miss most of the coarse dust in the atmosphere, *Science Advances*, 6(15), eaaz9507, doi:10.1126/sciadv.aaz9507, 2020.
- Aghababaeian, H., Ostadtaghizadeh, A., Ardalan, A., Asgary, A., Akbary, M., Yekaninejad, M.S., Stephens, C.: Global Health Impacts of Dust Storms: A Systematic Review, *Environmental Health Insights*, V15, <https://doi.org/10.1177/11786302211018390>, 2021.
- Alonso-Pérez, S., Cuevas, E., Querol, X.: Objective identification of synoptic meteorological patterns favouring African dust intrusions into the marine boundary layer of the subtropical eastern north Atlantic region, *Meteorol. Atmos. Phys.*, 113, 109-124, <https://doi.org/10.1007/s00703-011-0150-z>, 2011.
- Ashpole, I., and Washington, R.: Intraseasonal variability and atmospheric controls on daily dust occurrence frequency over the central and Western Sahara during the boreal summer, *J. Geophys. Res.-Atmos* 118, 12, 12915-12926, <http://dx.doi.org/10.1002/2013JD020267>, 2013.
- Bacer, S., Jomaa, F., Beaumet, J., Gallée, H., Le Bouëdec, E., Ménégoz, M., and Staquet, C.: Impact of climate change on wintertime European atmospheric blocking, *Weather Clim. Dynam.*, 3, 377–389, <https://doi.org/10.5194/wcd-3-377-2022>, 2022.
- Barnaba, F., and Gobbi, G.P.: Aerosol seasonal variability over the Mediterranean region and relative impact of maritime, continental and Saharan dust particles over the basin from MODIS data in the year 2001, *Atmos. Chem. Phys.*, 4, 2367–2391, <https://doi.org/10.5194/acp-4-2367-2004>, 2004.
- Barriopedro, D., García-Herrera, R., Lupo, A., and Hernández, E. R.: A climatology of northern hemisphere blocking, *J. Climate*, 19, 1042–1063, <https://doi.org/10.1175/JCLI3678.1>, 2006.
- Barriopedro D, García-Herrera R and Trigo RM.: Application of blocking diagnosis methods to general circulation models. Part I: a novel detection scheme, *Clim. Dyn.*, 35(7–8),1373–91, <https://doi.org/10.1007/s00382-010-0767-5>, 2010.
- Barriopedro, D., Ayarzagüena, B., Garcia-Burgos, M., and García-Herrera, R.: A multi-parametric perspective of the North Atlantic eddy-driven jet, *Climate Dynamics*, 61, 375–397, <https://doi.org/10.1007/s00382-022-06574-w>, 2023.

- Basart, S., Pérez C., Cuevas E., Baldasano J.M., Gobbi G.P.: Aerosol characterization in Northern Africa, Northeastern Atlantic, Mediterranean Basin and Middle East from direct-sun AERONET observations. *Atmos. Chem. Phys.*, 9, 8265-8282, <https://doi.org/10.5194/acp-9-8265-2009>, 2009.
- 800 Basart, S., Pay, M. T., Jorba, O., Pérez, C., Jiménez-Guerrero, P., Schulz, M., & Baldasano, J. M: Aerosols in the CALIOPE air quality modelling system: evaluation and analysis of PM levels, optical depths and chemical composition over Europe. *Atmos. Chem. Phys.*, 12(7), 3363-3392, 2012.
- Basart, S., Nickovic, S., Terradellas, E., Cuevas, E., Pérez García-Pando, C., García-Castrillo, G., Werner, E. and Benincasa, F.: The WMO SDS-WAS Regional Center for Northern Africa, Middle East and Europe, *Web of Conferences*, 99, 04008, <https://doi.org/10.1051/e3sconf/20199904008>, 2019.
- 805 Beerli, R., and Grams, C.M.: Stratospheric modulation of the large-scale circulation in the Atlantic–European region and its implications for surface weather events, *Q. J. R. Meteorol. Soc.*, 145, 3732– 3750, <https://doi.org/10.1002/qj.3653>, 2019.
- Benedict, J. J., Clement, A. C. and Medeiros, B.: Atmospheric blocking and other large-scale precursor patterns of landfalling atmospheric rivers in the North Pacific: A CESM2 study. *Journal of Geophysical Research: Atmospheres*, 124, <https://doi.org/10.1029/2019JD030790>, 2019.
- 810 Bou Karam, D., Flamant, C., Tulet, P., Todd, M.C., Pelon, J., and Williams, E.: Dry cyclogenesis and dust mobilization in the intertropical discontinuity of the West African Monsoon: A case study, *J. Geophys. Res.-Atmos.*, 114, D05115, <https://doi.org/10.1029/2008JD010952>, 2009.
- Cowie, S. M., Knippertz, P., and Marsham, J. H.: A climatology of dust emission events from northern Africa using long-term surface observations, *Atmos. Chem. Phys.*, 14, 16, 8579–8597, <https://doi.org/10.5194/acp-14-8579-2014>, 2014.
- 815 Cuesta, J., Marsham, J.H., Parker, D.J. and Flamant, C.: Dynamical mechanisms controlling the vertical redistribution of dust and the thermodynamic structure of the West Saharan atmospheric boundary layer during summer, *Atmos. Sci. Lett.*, 10, 34-42. <http://dx.doi.org/10.1002/asl.207>, 2009.
- 820 Cuevas, E., Gómez-Peláez, A.J., Rodríguez, S., Terradellas, E., Basart, S., García, R.D, García, O.E., Alonso-Pérez, S.: The pulsating nature of large-scale Saharan dust transport as a result of interplays between mid-latitude Rossby waves and the North African Dipole Intensity, *Atmospheric Environment*, 167, 586-602, <https://doi.org/10.1016/j.atmosenv.2017.08.059>, 2017.

- Cuevas, E., Milford, C., Barreto, A., Bustos, J. J., García, R. D., Marrero, C. L., Prats, N., Bayo, C., Ramos, R.,
825 Terradellas, E., Suárez, D., Rodríguez, S., de la Rosa, J., Vilches, J., Basart, S., Werner, E., López-Villarrubia,
E., Rodríguez-Mireles, S., Pita Toledo, M. L., González, O., Belmonte, J., Puigdemunt, R., Lorenzo, J.A., Oromí,
P., and del Campo-Hernández, R.: Desert Dust Outbreak in the Canary Islands (February 2020): Assessment and
Impacts. (Eds. Cuevas, E., Milford, C. and Basart, S.), State Meteorological Agency (AEMET), Madrid, Spain
and World Meteorological Organization, Geneva, Switzerland, WMO Global Atmosphere Watch (GAW) Report
830 No. 259, WWRP 2021-1, 2021.
- Davini, P., Cagnazzo, C., Gualdi, S., and Navarra, A.: Bidimensional Diagnostics, Variability, and Trends of
Northern Hemisphere Blocking, *J. Climate*, 25, 6496–6509, <https://doi.org/10.1175/JCLI-D-12-00032.1>, 2012.
- Davini, P., and D'Andrea, F.: From CMIP3 to CMIP6: Northern Hemisphere Atmospheric Blocking Simulation in
Present and Future Climate. *J. Climate*, 33, 10021–10038, <https://doi.org/10.1175/JCLI-D-19-0862.1>, 2020.
- 835 Dubovik, O., Holben, B. N., Eck, T. F., Smirnov, A., Kaufman, Y. J., King, M. D., Tanré, D., and Slutsker, I.:
Variability of Absorption and Optical Properties of Key Aerosol Types Observed in Worldwide Locations,
Atmos. Sci., 59, 590–608, [https://doi.org/10.1175/1520-0469\(2002\)059<0590:VOAAOP>2.0.CO;2](https://doi.org/10.1175/1520-0469(2002)059<0590:VOAAOP>2.0.CO;2), 2002.
- ESCAP/APDIM. Sand and dust storms risk assessment in Asia and the Pacific. Retrieved from:
<https://hdl.handle.net/20.500.12870/4452>. (last access on 13 November 2023), 2021.
- 840 Fernández, A.J., Sicard, M., Costa, M.J., Guerrero-Rascado, J.L., Gómez-Amo, J.L., Molero, F., Barragán, R.,
Basart, S. Bortoli, D., Bedoya-Velásquez, A.E., Utrillas, M.P., Salvador, P., Granados-Muñoz, M.J., Potes, M.,
Ortiz-Amezcu, P., Martínez-Lozano, J.A., Artíñano, B., Muñoz-Porcar, C., Salgado, R., Román, R.,
Rocadenbosch, F., Salgueiro, V., Benavent-Oltra, A., Rodríguez-Gómez, A., Alados-Arboledas, L., Comerón,
A. and Pujadas, M.: Extreme, wintertime Saharan dust intrusion in the Iberian Peninsula: Lidar monitoring and
845 evaluation of dust forecast models during the February 2017 event, *Atmospheric Research*, 228, 223-241, ISSN
0169-8095, <https://doi.org/10.1016/j.atmosres.2019.06.007>, 2019.
- Escudero, M., Querol, X., Ávila A., and Cuevas, E.: Origin of the exceedances of the European daily PM limit
value in regional background areas of Spain, *Atmos. Environ.*, 41, 730–744,
<https://doi.org/10.1016/j.atmosenv.2006.09.014>, 2007.

- 850 Fiedler, S., Schepanski, K., Knippertz, P., Heinold, B., and Tegen, I.: How important are atmospheric depressions and mobile cyclones for emitting mineral dust aerosol in North Africa?, *Atmos. Chem. Phys.*, 14, 8983–9000, <https://doi.org/10.5194/acp-14-8983-2014>, 2014.
- Flaounas, E., Davolio, S., Raveh-Rubin, S., Pantillon, F., Miglietta, M. M., Gaertner, M. A., Hatzaki, M., Homar, V., Khodayar, S., Korres, G., Kotroni, V., Kushta, J., Reale, M., and Ricard, D.: Mediterranean cyclones: current
855 knowledge and open questions on dynamics, prediction, climatology and impacts, *Weather Clim. Dynam.*, 3, 173–208, <https://doi.org/10.5194/wcd-3-173-2022>, 2022.
- Flaounas, E., Kotroni, V., Lagouvardos, K., Kazadzis, S., Gkikas, A. and Hatzianastassiou, N.: Cyclone contribution to dust transport over the Mediterranean region. *Atmos. Sci. Lett.*, 16, 473-478, <https://doi.org/10.1002/asl.584>, 2015.
- 860 Francis, D., Fonseca, R., Nelli, N., Bozkurt, D., Picard, G., and Guan, B. Atmospheric rivers drive exceptional Saharan dust transport towards Europe. *Atmospheric Research*, 266, 105959, <https://doi.org/10.1016/j.atmosres.2021.105959>, 2022.
- Gallisai, R., Peters, F., Volpe, G., Basart, S., and Baldasano, J., M.: Saharan dust deposition may affect phytoplankton growth in the Mediterranean Sea at ecological time scales, *PLoS ONE* 9(10): e110762.,
865 <https://doi.org/10.1371/journal.pone.0110762>, 2014.
- Garrido-Pérez, J.M., Ordóñez, C., García-Herrera, R. and Barriopedro, D.: Air stagnation in Europe: Spatiotemporal variability and impact on air quality, *Science of The Total Environment*, 645, 2018, 1238-1252, ISSN 0048-9697, <https://doi.org/10.1016/j.scitotenv.2018.07.238>, 2018.
- Garrido-Pérez, J.M., Vicente-Serrano, S.M., Barriopedro, D., García-Herrera, R., Trigo, R. and Beguería, S.:
870 Examining the outstanding Euro-Mediterranean drought of 2021-2022 and its historical context. *Journal of Hydrometeorology*, 130653, 10.1016/j.jhydrol.2024.130653, 2024
- Gavrouzou, M., Hatzianastassiou, N., Gkikas, A., Lolis, C.J. and Mihalopoulos, N. A.: Climatological Assessment of Intense Desert Dust Episodes over the Broader Mediterranean Basin Based on Satellite Data. *Remote Sens.*, 13, 2895, <https://doi.org/10.3390/rs13152895>, 2021.
- 875 Gelaro, R., McCarty, W., Suarez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C., Darmenov, A., Bosilovich, M. G., Reichle, R., Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V., Conaty, A., da Silva, A., Gu, W., Kim, G. K., Koster, R., Lucchesi, R., Merkova, D., Nielsen, J. E., Partyka, G., Pawson, S., Putman,

- 880 W., Rienecker, M., Schubert, S. D., Sienkiewicz, M., and Zhao, B.: The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2), *J. Climate*, 30, 5419–5454, <https://doi.org/10.1175/JCLI-D-16-0758.1>, 2017.
- Gerasopoulos, E., Kouvarakis, G., Babasakalis, P., Vrekoussis, M., Putaud, J. P., and Mihalopoulos, N.: Origin and variability of particulate matter (PM₁₀) mass concentrations over the Eastern Mediterranean, *Atmos. Environ.*, 40, 4679–4690, <https://doi.org/10.1016/j.atmosenv.2006.04.020>, 2006.
- 885 Gherboudj, I., Beegum, S.N, Marticorena, B., and Ghedira, H.: Dust emission parameterizationscheme over the MENA region: Sensitivity analysis to soil moisture andsoil texture,*J. Geophys. Res. Atmos.*,120,10,915–10,938, <https://doi.org/10.1002/2015JD023338>, 2015.
- Giles, D. M., Sinyuk, A., Sorokin, M. G., Schafer, J. S., Smirnov, A., Slutsker, I., Eck, T. F., Holben, B. N., Lewis, J. R., Campbell, J. R., Welton, E. J., Korokin, S. V., and Lyapustin, A. I.: Advancements in the Aerosol Robotic Network (AERONET) Version 3 database – automated near-real-time quality control algorithm with improved cloud screening for Sun photometer aerosol optical depth (AOD) measurements, *Atmos. Meas. Tech.*, 12, 169-209, <https://doi.org/10.5194/amt-12-169-2019>, 2019.
- 890 Gimeno, L., Nieto, R., Vázquez, M. and Lavers, D.: Atmospheric rivers: a mini-review. *Frontiers in Earth Science*, 2, <https://doi.org/10.3389/feart.2014.00002>, 2014
- Ginoux, P., Prospero, J. M., Gill, T. E., Hsu, N. C., and Zhao, M.: Global-scale attribution of anthropogenic and natural dust sources and their emission rates based on MODIS Deep Blue aerosol products, *Rev. Geophys.*, 50, RG3005, <https://doi.org/10.1029/2012RG000388>, 2012.
- 895 Gkikas, A., Hatzianastassiou, N., Mihalopoulos, N., Katsoulis, V., Kazadzis, S., Pey, J., Querol, X., and Torres, O.: The regime of intense desert dust episodes in the Mediterranean based on contemporary satellite observations and ground measurements, *Atmos. Chem. Phys.*, 13, 12135–12154, <https://doi.org/10.5194/acp-13-12135-2013>, 2013.
- 900 Gkikas, A., Houssos, E.E., Lolis, C.J., Bartzokas, A., Mihalopoulos, N. and Hatzianastassiou, N.: Atmospheric circulation evolution related to desert dust episodes over the Mediterranean, *Quarterly Journal of the Royal Meteorological Society*, 141, 690, 1634-1645, <https://doi.org/10.1002/qj.2466>, 2015.

- 905 Gkikas, A., Basart, S., Hatzianastassiou, N., Marinou, E., Amiridis, V., Kazadzis, S., Pey, J., Querol, X., Jorba, O.,
Gassó, S., and Baldasano, J. M.: Mediterranean intense desert dust outbreaks and their vertical structure based on
remote sensing data, *Atmos. Chem. Phys.*, 16, 8609–8642, <https://doi.org/10.5194/acp-16-8609-2016>, 2016.
- Grams, C.M., Beerli, R., Pfenninger, S., Stafell, I., and Wernli, H.: Balancing Europe’s wind-power output through
spatial deployment informed by weather regimes, *Nature Clim Change*, 7, 557–562,
<https://doi.org/10.1038/nclimate3338>, 2017.
- 910 Guerzoni, S., Chester, R., Dulac, F., Herut, B., Loye-Pilot, M.-D., Measures, C., Migon, C., Molinaroli, E., Moulin,
C., Rossini, P., Saydam, C., Soudine, A., and Ziveri, P.: The role of atmospheric deposition in the biogeochemistry
of the Mediterranean Sea, *Prog. Oceanogr.*, 44, 147–190, [https://doi.org/10.1016/S0079-6611\(99\)00024-5](https://doi.org/10.1016/S0079-6611(99)00024-5), 1999.
- Helgren, D. M., and Prospero, J. M.: Wind Velocities Associated with Dust Deflation Events in the Western
Sahara, *Journal of Climate and Applied Meteorology*, 26, 9, 1147–1151. <https://www.jstor.org/stable/26183511>,
915 1987.
- Hersbach, H, Bell, B, Berrisford, P, Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu,
R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot,
J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R.,
Fuentes M., Geer, A., Haimberger, L., Healy, S., Hogan, R.J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P.,
920 Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S. and Thépaut, J-N.: The
ERA5 global reanalysis. *Q J R Meteorol Soc.*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- Holben, B. N., Eck, T. F., Slutsker, I., Tanre, D., Buis, J. P., Setzer, A., Vermote, E., Reagan, J. A., Kaufman, Y.,
Nakajima, T., Lavenue, F., Jankowiak, I., and Smirnov, A.: AERONET – A federated instrument network and
data archive for aerosol characterization, *Remote Sens. Environ.*, 66, 1–16, [https://doi.org/10.1016/S0034-4257\(98\)00031-5](https://doi.org/10.1016/S0034-4257(98)00031-5), 1998.
925
- IPCC: Climate Change 2022: Impacts, Adaptation, and Vulnerability. Contribution of Working Group II to the
Sixth Assessment Report of the Intergovernmental Panel on Climate Change [H.-O. Pörtner, D.C. Roberts, M.
Tignor, E.S. Poloczanska, K. Mintenbeck, A. Alegría, M. Craig, S. Langsdorf, S. Löschke, V. Möller, A. Okem,
B. Rama (eds.)]. Cambridge University Press. Cambridge University Press, Cambridge, UK and New York, NY,
930 USA, 3056 pp., [doi:10.1017/9781009325844](https://doi.org/10.1017/9781009325844), 2022.

- Israelevich, P., Ganor, E., Alpert, P., Kishcha, P. and Stupp, A.: Predominant transport paths of Saharan dust over the mediterranean sea to Europe, *J. Geophys. Res.*, 117, D02205, <http://dx.doi.org/10.1029/2011JD016482>, 2012.
- 935 Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, I. S., White, G., Woollen, J., Zhu, Y., Leetmaa, A., Reynolds, R., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J., Jenne, R., and Joseph, D.: The NCEP/NCAR 40-Year Reanalysis Project, *B. Am. Meteorol. Soc.*, 77, 437–471, [https://doi.org/10.1175/1520-0477\(1996\)077<0437:TNYRP>2.0.CO;2](https://doi.org/10.1175/1520-0477(1996)077<0437:TNYRP>2.0.CO;2), 1996.
- Kaufman, Y.J., Tanré, D., Remer, L.A., Vermote, E.F., Chu, A., Holben, B.N.: Operational remote sensing of tropospheric aerosol over land from EOS moderate resolution imaging spectroradiometer, *J. Geophys. Res.*, 102, 17-51, <https://doi.org/10.1029/96JD03988>, 1997.
- 940 Kautz, L.-A., Martius, O., Pfahl, S., Pinto, J. G., Ramos, A. M., Sousa, P. M., and Woollings, T.: Atmospheric blocking and weather extremes over the Euro-Atlantic sector – a review, *Weather Clim. Dynam.*, 3, 305–336, <https://doi.org/10.5194/wcd-3-305-2022>, 2022.
- Kim, M.-H., Omar, A. H., Tackett, J. L., Vaughan, M. A., Winker, D. M., Trepte, C. R., Hu, Y., Liu, Z., Poole, L. R., Pitts, M. C., Kar, J., and Magill, B. E.: The CALIPSO version 4 automated aerosol classification and lidar ratio selection algorithm, *Atmos. Meas. Tech.*, 11, 6107–6135, <https://doi.org/10.5194/amt-11-6107-2018>, 2018.
- 945 Kinne, S., Schulz, M., Textor, C., Guibert, S., Balkanski, Y., Bauer, S. E., Berntsen, T., Berglen, T. F., Boucher, O., Chin, M., Collins, W., Dentener, F., Diehl, T., Easter, R., Feichter, J., Fillmore, D., Ghan, S., Ginoux, P., Gong, S., Grini, A., Hendricks, J., Herzog, M., Horowitz, L., Isaksen, I., Iversen, T., Kirkevåg, A., Kloster, S., Koch, D., Kristjansson, J. E., Krol, M., Lauer, A., Lamarque, J. F., Lesins, G., Liu, X., Lohmann, U., Montanaro, V., Myhre, G., Penner, J., Pitari, G., Reddy, S., Seland, O., Stier, P., Takemura, T., and Tie, X.: An AeroCom initial assessment – optical properties in aerosol component modules of global models, *Atmos. Chem. Phys.*, 6, 1815–1834, <https://doi.org/10.5194/acp-6-1815-2006>, 2006.
- Kinni, P., Kouis, P., Dimitriou, H., Yarza, S., Papatheodorou, S., Kampriani, E., Charalambous, M., Middleton, N., Novack, V., Galanakis, E., Yiallourous, P. K.: Health effects of desert dust storm events in the south-eastern Mediterranean: perceptions and practices of local stakeholders, *East Mediterr Health – Vol. 27 No. 11*, 1092-1101, <https://doi.org/10.26719/emhj.21.037>, 2021
- 955 Knippertz, P.: Dust mobilization in the West African heat trough—The role of the diurnal cycle and of extratropical synoptic disturbances, *Meteorol. Z.*, 17, 553–563, <https://doi.org/10.1127/0941-2948/2008/0315>, 2008.

- Knippertz, P. and Todd, M. C.: Mineral dust aerosols over the Sahara: Meteorological controls on emission and transport and implications for modeling, *Rev. Geophys.*, 50, RG1007, <https://doi.org/10.1029/2011RG000362>, 2012.
- Kok, J.F., Ridley, D.A., Zhou, Q., Miller, R.L., Zhao, C., Heald, C.L., Ward, D.S., Albani S. and Haustein, K.: Smaller desert dust cooling effect estimated from analysis of dust size and abundance, *Nature Geoscience*, 10, 274–278, <https://doi.org/10.1038/ngeo2912>, 2017.
- 965 Kok, J. F., Adebisi, A. A., Albani, S., Balkanski, Y., Checa-Garcia, R., Chin, M., Colarco, P. R., Hamilton, D. S., Huang, Y., Ito, A., Klose, M., Li, L., Mahowald, N. M., Miller, R. L., Obiso, V., Pérez García-Pando, C., Rocha-Lima, A., and Wan, J. S.: Contribution of the world’s main dust source regions to the global cycle of desert dust, *Atmos. Chem. Phys.*, 21, 8169–8193, <https://doi.org/10.5194/acp-21-8169-2021>, 2021.
- Kuula, J., Timonen, H., Niemi, J. V., Manninen, H. E., Rönkkö, T., Hussein, T., Fung, P. L., Tarkoma, S., Laakso, 970 M., Saukko, E., Ovaska, A., Kulmala, M., Karppinen, A., Johansson, L., and Petäjä, T.: Opinion: Insights into updating Ambient Air Quality Directive 2008/50/EC, *Atmos. Chem. Phys.*, 22, 4801–4808, <https://doi.org/10.5194/acp-22-4801-2022>, 2022.
- Lagaria, A., Mandalakis, M., Mara, P., Papageorgiou, N., Pitta, P., Tsiola, A., Kagiorgi, M., and Psarra, S.: Phytoplankton response to Saharan dust depositions in the Eastern Mediterranean Sea: A mesocosm study, *Front. 975 Mar. Sci.*, 3, 287, <https://doi.org/10.3389/fmars.2016.00287>, 2017.
- Liu, Z., Vaughan, M., Winker, D., Kittaka, C., Getzewich, B., Kuehn, R., Omar, A., Powell, K., Trepte, C., and Hostetler, C.: The CALIPSO lidar cloud and aerosol discrimination: Version 2 algorithm and initial assessment of performance, *J. Atmos. Ocean. Tech.*, 26, 1198–1213, <https://doi.org/10.1175/2009JTECHA1229.1>, 2009.
- Liu, Z., Kar, J., Zeng, S., Tackett, J., Vaughan, M., Avery, M., Pelon, J., Getzewich, B., Lee, K.-P., Magill, B., 980 Omar, A., Lucker, P., Trepte, C., and Winker, D.: Discriminating between clouds and aerosols in the CALIOP version 4.1 data products, *Atmos. Meas. Tech.*, 12, 703–734, <https://doi.org/10.5194/amt-12-703-2019>, 2019.
- Lorente-Plazas, R., Montavez, J. P., Ramos, A. M., Jerez, S., Trigo, R. M. and Jimenez-Guerrero, P.: Unusual atmospheric-river-like structures coming from Africa induce extreme precipitation over the western Mediterranean Sea. *Journal of Geophysical Research: Atmospheres*, 125, e2019JD031280, 985 <https://doi.org/10.1029/2019JD031280>, 2020.

Maddison, J. W., Abalos, M., Barriopedro, D., García-Herrera, R., Garrido-Pérez, J. M., and Ordóñez, C.: Linking air stagnation in Europe with the synoptic- to large-scale atmospheric circulation. *Weather Clim. Dynam.*, 2, 675–694, <https://doi.org/10.5194/wcd-2-675-2021>, 2021.

990 Marconi, M., Sferlazzo, D. M., Becagli, S., Bommarito, C., Calzolari, G., Chiari, M., di Sarra, A., Ghedini, C., Gómez-Amo, J. L., Lucarelli, F., Meloni, D., Monteleone, F., Nava, S., Pace, G., Piacentino, S., Rugi, F., Severi, M., Traversi, R., and Udisti, R.: Saharan dust aerosol over the central Mediterranean Sea: PM10 chemical composition and concentration versus optical columnar measurements, *Atmos. Chem. Phys.*, 14, 2039–2054, <https://doi.org/10.5194/acp-14-2039-2014>, 2014.

Met Office; EUMETSAT (2022): MSG: Dust imagery in the RGB channels over the full disc at 41.5 degrees East (LEDF41, up to 0900 UTC 1st June 2022). NERC EDS Centre for Environmental Data Analysis, date of citation. 995 <https://catalogue.ceda.ac.uk/uuid/b1dacc09b42f4d8ab492c5d5c751efa9>.

Michel, C., and Rivière, G.: The link between Rossby wave breakings and weather regime transitions, *J. Atmos. Sci.*, 68, 1730-1748, <https://doi.org/10.1175/2011JAS3635.1>, 2011.

Middleton N., and Kappelle M.: Impacts of sand and dust storms on oceans: a scientific environmental assessment for policy makers. United Nations Environment Programme, Nairobi, Kenya. 1000 <https://www.unep.org/resources/report/impacts-sand-and-dust-storms-oceans>, 2020

Monteiro, A., Basart, S., Kazadzis, S., Votsis, A., Gkikas, A., Vandenbussche, S., Tobias, A., Gama, C., Pérez García-Pando, C., Terradellas, E., Notas, G., Middleton, N., Kushta, J., Amiridis, V., Lagouvardos, K., Kosmopoulos, P., Kotroni, V., Kanakidou, M., Mihalopoulos, N., Kalivitis, N., Dagsson-Waldhauserová, P., El-Askary, H., Sievers, K., Giannaros, T., Mona, L., Hirtl, M., Skomorowski, P., Virtanen, T.H., Christoudias, T., Di Mauro, B., Trippetta, S., Kutuzov, S., Meinander, O., and Nickovic, S.: Multi-sectoral impact assessment of an extreme African dust episode in the Eastern Mediterranean in March 2018. *Science of The Total Environment*, 843, 156861, <https://doi.org/10.1016/j.scitotenv.2022.156861>, 2022. 1005

Moulin, C., Lambert, C. E., Dayan, U., Masson, V., Ramonet, M., Bousquet, P., Legrand, M., Balkanski, Y. J., Guelle, W., Marticorena, B., Bergametti, G., and Dulac, F.: Satellite climatology of African dust transport in the Mediterranean atmosphere, *J. Geophys. Res.*, 103, 13137–13144, <https://doi.org/10.1029/98JD00171>, 1998. 1010

Muñoz, C., Schultz, D. and Vaughan, G.: A Midlatitude Climatology and Interannual Variability of 200- and 500-hPa Cut-Off Lows, *Journal of Climate*, 33, 6, 2201-2222, <https://doi.org/10.1175/JCLI-D-19-0497.12020>, 2020.

- 1015 Nabizadeh, E., Hassanzadeh, P., Yang, D., and Barnes, E. A.: Size of the atmospheric blocking events: Scaling law and response to climate change, *Geophys. Res. Lett.*, 46, 13,488–13,499, <https://doi.org/10.1029/2019GL084863>, 2019.
- Nieto, R., Gimeno, L., de la Torre, L., Ribera, P., Gallego, D., García-Herrera, R., García, J.A., Nuñez, M., Redaño, A., Lorente, J.: Climatological features of Cut-off low systems in the Northern Hemisphere, *J. Climate*, 18, 16, 3085–3103, <https://doi.org/10.1175/JCLI3386.1>, 2005.
- 1020 Nieto, R., L. Gimeno, L. de la Torre, Ribera, P., Barriopedro, D., García-Herrera, R., Serrano, A., Gordillo, A., Redaño, A. and Lorente, J.: Interannual variability of cut-off low systems over the European sector: The role of blocking and the Northern Hemisphere circulation modes, *Meteorol. Atmos. Phys.*, 96, 85–101, <https://doi.org/10.1007/s00703-006-0222-7>, 2007.
- 1025 Oduber, F., Calvo, A.I., Blanco-Alegre, C., Castro, A., Nunes, T., Alves, C., Sorribas, M., Fernández-González, D., Vega-Maray, A.M., Valencia-Barrera, R.M., Lucarelli, F., Nava, S., Calzolari, G., Alonso-Blanco, E., Fraile, B., Fialho, P., Coz, E., Prevo, A.S.H., Pont, V. and Fraile, R.: Unusual winter Saharan dust intrusions at Northwest Spain: Air quality, radiative and health impacts, *Sci. Total Environ.*;669,213-228, <https://doi.org/10.1016/j.scitotenv.2019.02.305>, 2019.
- 1030 Ordóñez, C., Barriopedro, D., García-Herrera, R., Sousa, P. M., and Schnell, J. L.: Regional responses of surface ozone in Europe to the location of high-latitude blocks and subtropical ridges, *Atmos. Chem. Phys.*, 17, 3111–3131, <https://doi.org/10.5194/acp-17-3111-2017>, 2017.
- Otero, N., Jurado, O. E., Butler, T., and Rust, H. W.: The impact of atmospheric blocking on the compounding effect of ozone pollution and temperature: a copula-based approach, *Atmos. Chem. Phys.*, 22, 1905–1919, <https://doi.org/10.5194/acp-22-1905-2022>, 2022.
- 1035 Pey, J., Querol, X., Alastuey, A., Forastiere, F., and Stafoggia, M.: African dust outbreaks over the Mediterranean Basin during 2001–2011: PM10 concentrations, phenomenology and trends, and its relation with synoptic and mesoscale meteorology, *Atmos. Chem. Phys.*, 13, 1395–1410, <https://doi.org/10.5194/acp-13-1395-2013>, 2013.
- Pfahl, S.: Characterising the relationship between weather extremes in Europe and synoptic circulation features, *Nat. Hazards Earth Syst. Sci.*, 14, 1461–1475, <https://doi.org/10.5194/nhess-14-1461-2014>, 2014.
- 1040 Pitkanen-Brunnsberg, J.: Background Note on the UN Sand and Dust Storm (SDS) Coalition, United Nations Convention to Combat Desertification. France. Retrieved from

<https://policycommons.net/artifacts/2227328/background-note-on-the-un-sand-and-dust-storm-sds-coalition/2984760/> (last access on 17 June 2023), 2019.

1045 Portmann, R., Sprenger, M., and Wernli, H.: The three-dimensional life cycles of potential vorticity cutoffs: a global and selected regional climatologies in ERA-Interim (1979–2018), *Weather Clim. Dynam.*, 2, 507–534, <https://doi.org/10.5194/wcd-2-507-2021>, 2021.

Querol, X., Pey, J., Pandolfi, M., Alastuey, A., Cusack, M., Pérez, N., Moreno, N., Viana, M., Mihalopoulos, N., Kallos, G., and Kleanthous, S.: African dust contributions to mean ambient PM₁₀ mass-levels across the Mediterranean basin, *Atmos. Environ.*, 43, 4266–4277, <https://doi.org/10.1016/j.atmosenv.2009.06.013>, 2009.

1050 Rex, D.: Blocking Action in the Middle Troposphere and its Effect upon Regional Climate. *Tellus*, 2: 275-301, <https://doi.org/10.1111/j.2153-3490.1950.tb00339.x>, 1950.

Rodríguez, S., Cuevas, E., Prospero, J. M., Alastuey, A., Querol, X., López-Solano, J., García, M. I., and Alonso-Pérez, S.: Modulation of Saharan dust export by the North African dipole, *Atmos. Chem. Phys.*, 15, 7471–7486, <https://doi.org/10.5194/acp-15-7471-2015>, 2015.

1055 Ryder, C. L., Marengo, F., Brooke, J. K., Estelles, V., Cotton, R., Formenti, P., McQuaid, J. B., Price, H. C., Liu, D., Ausset, P., Rosenberg, P. D., Taylor, J. W., Choularton, T., Bower, K., Coe, H., Gallagher, M., Crosier, J., Lloyd, G., Highwood, E. J., and Murray, B. J.: Coarse-mode mineral dust size distributions, composition and optical properties from AER-D aircraft measurements over the tropical eastern Atlantic, *Atmos. Chem. Phys.*, 18, 17225–17257, <https://doi.org/10.5194/acp-18-17225-2018>, 2018.

1060 Salvador, P., Alonso-Pérez, S., Pey, J., Artíñano, B., de Bustos, J. J., Alastuey, A., and Querol, X.: African dust outbreaks over the western Mediterranean Basin: 11-year characterization of atmospheric circulation patterns and dust source areas, *Atmos. Chem. Phys.*, 14, 6759–6775, <https://doi.org/10.5194/acp-14-6759-2014>, 2014.

1065 Sayer, A.M., Hsu, N.C., Bettenhausen, C., Jeong, M.-J.: Validation and uncertainty estimates for MODIS Collection 6 “Deep Blue” aerosol data, *J. Geophys. Res. Atmos.*, 118, 7864–7872, <http://dx.doi.org/10.1002/jgrd.50600>, 2013.

Schepanski, K., Tegen, I., Laurent, B., Heinold, B., and Macke, A.: A new Saharan dust source activation frequency map derived from MSG-SEVIRI IR-channels, *Geophys. Res. Lett.*, 34, L18803, <https://doi.org/10.1029/2007GL030168>, 2007.

- 1070 Schepanski, K., Tegen, I., Todd, M. C., Heinold, B., Bönisch, G., Laurent, B., and Macke, A.: Meteorological processes forcing Saharan dust emission inferred from MSGSEVIRI observations of subdaily dust source activation and numerical models, *J. Geophys. Res.*, 114, D10201, <https://doi.org/10.1029/2008JD010325>, 2009.
- Schepanski, K., Tegen, I., and Macke, A.: Comparison of satellite based observations of Saharan dust source areas, *Remote Sens. Environ.*, 123, 90–97, ISSN 0034-4257, <https://doi.org/10.1016/j.rse.2012.03.019> , 2012.
- 1075 Shepherd, G., Terradellas, E., Baklanov, A., Kang, U., Sprigg, W.A., Nickovic, S., Bolorani, A.D., Al-Dousari, A., Basart, S., Benedetti, A., Sealy, A., Tong, D., Zhang, X., Shumake-Guillemot, J., Kebin, Z., Knippertz, P., Mohammed, A.A.A., Al-Dabbas, M., Cheng, L., Otani, S., Wang, F., Zhang, C., Ryoo, SB., Cha, J.: Global assessment of sand and dust storms, United Nations Environment Programme, Nairobi, 139 pp, ISBN: 978-92-807-3551-2, 2016.
- 1080 Sousa P.M., Trigo R. M., Barriopedro D., Soares P. M. M., Santos J. A. European temperature responses to blocking and ridge regional patterns. *Climate Dynamics*, 50, 1-2, 457-477, <https://doi.org/10.1007/s00382-017-3620-2>, 2018.
- Sousa, P. M., Barriopedro, D., García-Herrera, R., Woollings, T., and Trigo, R.M.: A New Combined Detection Algorithm for Blocking and Subtropical Ridges, *J. Climate*, 34, 7735–7758, <https://doi.org/10.1175/JCLI-D-20-0658.1>, 2021.
- 1085 Stathopoulos, C.; Patlakas, P.; Tsalis, C. and Kallos, G.: The Role of Sea Surface Temperature Forcing in the Life-Cycle of Mediterranean Cyclones, *Remote Sens.*, 12, 825. <https://doi.org/10.3390/rs12050825>, 2020.
- Stephens, G.L., D.G. Vane, R.J. Boain, G.G. Mace, K. Sassen, Z. Wang, A.J. Illingworth, E.J. O'Connor, W.B. Rossow, S.L. Durden, S.D. Miller, R.T. Austin, A. Benedetti, C. Mitrescu, and CloudSat Science Team: The CloudSat mission and the A-Train: A new dimension of space-based observations of clouds and precipitation. *Bull. Amer. Meteorol. Soc.*, 83, 1771-1790, <https://doi.org/10.1175/BAMS-83-12-1771>, 2002.
- 1090 Szopa, S., V. Naik, B. Adhikary, P. Artaxo, T. Berntsen, W.D. Collins, S. Fuzzi, L. Gallardo, A. Kiendler-Scharr, Z. Klimont, H. Liao, N. Unger, and P. Zanis, 2021: Short-Lived Climate Forcers. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K.
- 1095

Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 817–922, doi:10.1017/9781009157896.008, 2021.

1100 Tegen, I., Schepanski, K., and Heinold, B.: Comparing two years of Saharan dust source activation obtained by regional modelling and satellite observations, *Atmos. Chem. Phys.*, 13, 2381–2390, <https://doi.org/10.5194/acp-13-2381-2013>, 2013.

Terradellas, E., Nickovic, S., and Zhang, X. Y.: Airborne dust: a hazard to human health, environment and society, *WMO Bull*, 64, 42–46, <https://sds-was.aemet.es/materials/hazard.pdf> (last access on 15 February 2024), 2015.

Thomas, N., and Nigam, S.: Twentieth-Century Climate Change over Africa: Seasonal Hydroclimate Trends and Sahara Desert Expansion, *J. Climate*, 31, 3349–3370, <https://doi.org/10.1175/JCLI-D-17-0187.1>, 2018.

1105 Trigo, R. M., Trigo, I. F., DaCamara, C.C., and Osborn, T.J.: Winter blocking episodes in the European–Atlantic sector: Climate impacts and associated physical mechanisms in the Reanalysis, *Climate Dynamics* (2004) 23: 17–28, DOI 10.1007/s00382-004-0410-4, 2004.

Varga, G., Újvári, G, Kovács, J.: Spatiotemporal patterns of Saharan dust outbreaks in the Mediterranean Basin, *Aeolian Res.*, 15, 151–160, <https://doi.org/10.1016/j.aeolia.2014.06.005>, 2014.

1110 Vaughan, M. A., Powell, K. A., Kuehn, R. E., Young, S. A., Winker, D. M., Hostetler, C. A., Hunt, W. H., Liu, Z., McGill, M. J., and Getzewich, B. Z.: Fully automated detection of cloud and aerosol layers in the CALIPSO lidar measurements, *J. Atmos. Ocean. Tech.*, 26, 2034–2050, <https://doi.org/10.1175/2009JTECHA1228.1>, 2009.

Wagner R., Schepanski, K., Heinold, B, and Tegen, I.: Interannual variability in the Saharan dust source activation—Toward differences between 2007 and 2008, *J. Geophys. Res. Atmos.*, 121, 4538–4562, 1115 <https://doi.org/10.1002/2015JD024302>, 2016.

Wang, W., A. Evan, C. Flamant, and C. Lavaysse,: On the decadal scale correlation between African dust and Sahel rainfall: The role of Saharan heat low-forced winds, *Sci. Adv.*, 1(9), e1500646, doi:10.1126/sciadv.1500646, 2015.

1120 Wazneh, H., Gachon, P., Laprise, R., de Vernal, A., Tremblay, B.: Atmospheric blocking events in the North Atlantic: trends and links to climate anomalies and teleconnections, *Clim. Dyn.*, 56, 2199–2221, <https://doi.org/10.1007/s00382-020-05583-x>, 2021.

WHO (World Health Organization). WHO global air quality guidelines: particulate matter (PM_{2.5} and PM₁₀), ozone, nitrogen dioxide, sulfur dioxide and carbon monoxide. World Health Organization. <https://iris.who.int/handle/10665/345329>. Licencia: CC BY-NC-SA 3.0 IGO, 2021.

1125 Wilks, D. S.: Statistical Methods in the Atmospheric Sciences, Cambridge, MA, USA, ISBN: 9780123850232, 2011.

Winker, D. M., Vaughan, M. A., Omar, A. H., Hu, Y., Powell, K. A., Liu, Z., Hunt, W. H., and Young, S. A.: Overview of the CALIPSO mission and CALIOP data processing algorithms, *J. Atmos. Ocean. Tech.*, 26, 2310–2323, <https://doi.org/10.1175/2009JTECHA1281.1>, 2009.

1130 Winstanley, D.: Sharav, *Weather*, 27, 146–160, <https://doi.org/10.1002/j.1477-8696.1972.tb04279.x>, 1972.

WMO: The Role of Climatological Normals in a Changing Climate, WCDMP-No.61, WMO-TD No.1377, Edited by: Omar Baddour and Hama Kontongomde, 2007

WMO: Sand and Dust Storm Warning Advisory and Assessment System (SDS-WAS), Science and Implementation Plan: 2015-2020, WWRP 2015–5, 30pp, <https://repositorio.aemet.es/handle/20.500.11765/2852> (last access on 15 February 2024), 2015.

1135 WMO: Airborne dust bulletin No. 5, permalink: <https://library.wmo.int/idurl/4/57560>, 2021.

WMO: Airborne dust bulletin No. 7, permalink: <https://library.wmo.int/idurl/4/57560>, 2023.

Woollings, T., Barriopedro, D., Methven, J., Son, S-W, Martius, O., Harvey, B., Sillmann, J., Lupo, A. R., and Seneviratne, S.: Blocking and its response to climate change. *Current Climate Change Reports*, <https://doi.org/10.1007/s40641-018-0108-z>, E-ISSN: 2198-6061, 2018.

1140 Wu, M., Liu, X., Yu, H., Wang, H., Shi, Y., Yang, K., Darmenov, A., Wu, C., Wang, Z., Luo, T., Feng, Y., and Ke, Z.: Understanding processes that control dust spatial distributions with global climate models and satellite observations, *Atmos. Chem. Phys.*, 20, 13835–13855, <https://doi.org/10.5194/acp-20-13835-2020>, 2020.

1145