

# Cretaceous-Paleocene extension at the southwest continental margin of India and opening of the Laccadive Basin: Constraints from geophysical data

Mathews George Gilbert<sup>1</sup>, Parakkal Unnikrishnan<sup>1</sup>, and Munukutla Radhakrishna<sup>1</sup>

<sup>1</sup>Department of Earth Sciences, Indian Institute of Technology Bombay, Powai, Mumbai 400076, India.


**Correspondence:** Mathews George Gilbert (georgegilbertm@gmail.com)

**Abstract.** Previous geophysical investigations of the Western Continental Margin of India (WCMI) confirm the ~~double two-phase~~ breakup history of the margin with the first breakup taking place between India and Madagascar in Late Cretaceous and the second breakup event in early Paleocene with Seychelles ~~separation~~ ~~separating~~ from India. Despite numerous geoscientific studies along the WCMI, the opening of the Laccadive Basin, lying along the southern part of the margin, ~~is not discussed~~ ~~remains~~ ~~poorly constrained~~. In this study, we evaluate the multi-channel seismic reflection and gravity anomalies at the margin to identify the early rift signatures in conjunction with the magnetic anomaly identifications in the Mascarene Basin. The analysis led to the identification of two extensional directions, a ENE-WSW oriented extension over the Laccadive Ridge north of Tellicherry Arch, and the NW-SE extension in the Laccadive Basin region towards ~~south~~. ~~Plate the south~~. ~~Previous plate~~ reconstruction models of the Mascarene Basin ~~suggest using marine magnetic lineations suggest that~~ the ENE-WSW extension observed over the Laccadive Ridge could be related to the India-Madagascar separation. ~~However, the sediment deposition pattern~~ ~~We associate the pattern of sediment deposition~~ and the presence of ~~Paleocene trap~~ ~~a Paleocene trap~~, associated with the NW-SE extensional grabens observed in the Laccadive Basin region ~~has been attributed to~~, ~~to the~~ extension between the Laccadive Ridge and the West coast of India after the separation of Madagascar from India. We further propose that the anti-clockwise rotation of India and the passage of the Réunion plume have facilitated the opening of the Laccadive Basin.

15 *Copyright statement.* TEXT

## 1 Introduction


The ~~Western~~ ~~breakup of Gondwanaland into the Eastern and Western Gondwana during the Early Jurassic period has initiated the formation of western Indian Ocean (fig. 1). The subsequent breakup of Madagascar and the Seychelles from India, and the seafloor spreading along the Carlsberg and Central Indian ridges culminated into the present northwest Indian Ocean. The~~ ~~Laccadive Ridge (LR), the Maldives ridge (MR), the Chagos Bank (CB) and the Saya-de-Malha Bank (SMB), the Nazareth~~ ~~Bank (NB), the Mauritius Island (MI), the Reunion Island (RI) are the major topographic features present in the northwest~~ ~~Indian Ocean (fig. 1A), and some of these are believed to be the micro-continents (Torsvik et al., 2013).~~



**Figure 1.** A) Regional tectonic map of the northwestern Indian Ocean with satellite-derived seafloor topography (Smith and Sandwell, 1997). The area shown in the dashed black rectangle is shown in figure 2A & 2B. B) The map shows the position of India relative to Madagascar and Deccan Volcanic Province in late Paleozoic fit (Lovecchio et al., 2020, modified after). The area of interest is marked in black rectangle. SB: Seychelles Bank; SMB: Saya-de Malha Bank; NB: Nazarat Bank; MI: Mauritius Island; RI: Reunion Island; RTJ: Rodrigues Triple Junction; LR: Laccadive Ridge; MA: Maldives; CB: Chagos Bank; MR: Murray Ridge.

The Western Continental Margin of India (WCMI) formed through the breakup and separation of India and Madagascar in the late-Cretaceous (Storey et al., 1995; Pande et al., 2001). This breakup event resulted in the formation the Mascarene Basin (fig. 1A & 2B), the details of which are recorded in the magnetic anomalies in the basin (fig. 2B). The northern part of the margin then experienced another breakup event when the Seychelles block separated from the combined Laxmi Ridge and India in the early Paleocene time. The This second breakup event is well studied with the pre-drift juxtaposition of the continental blocks fairly well established from the magnetic anomaly identifications and geochronology (Collier et al., 2008; Chaubey et al., 2002; Ganerød et al., 2011; Shellnutt et al., 2015, 2017). Earlier-reconstruction-studies-place-the-southwestern-margin of India close to the southeastern margin of Madagascar (Katz and Premoli, 1979), whereas, more recent studies incorporate The southern part of the margin is considered to be conjugate with the Madagascar (Katz and Premoli, 1979) based on the matching of the major shear zones and reconstructed to 2000 m isobath. However, recent close-fit reconstruction models have incorporated the continental fragments like Laccadive Ridge (Bhattacharya and Yatheesh, 2015) or Mauritia, consisting comprising of Mauritius, the Southern Mascarene Plateau, the Laccadive Plateau and the Chagos Bank (Torsvik et al., 2013) between India and Madagascar in the India-Madagascar pre-drift scenario, and suggest a breakup timing of around 83 Ma.

Since the Laccadive Basin and the adjoining southwest margin lying south of the Tellicherry basement Archis the conjugate segment of the Madagascar, the margin is important in terms of examining the early breakup history. Also, the basin Laccadive



**Figure 2.** A) Regional tectonic map of the western continental margin of India (Smith and Sandwell, 1997). Black solid lines in the offshore region represent shear zones and faults. B) Tectonic map of Madagascar and Mascarene basin showing mapped seafloor spreading type magnetic lineations in solid green coloured lines (Bhattacharya and Yatheesh, 2015, and references therein). Solid white lines represent the mapped fracture zones or pseudo-faults. SB: Seychelles Bank; SMB: Saya-de Malha Bank; NB: Nazarat Bank; MI: Mauritius Island; RI: Reunion Island; NMR: Northern Madagascar Ridge; DC: Dharwar Craton; SGT: Southern Granulite Terrain; LB: Laccadive Basin; AP: Alleppey Platform; TT Trivandrum Terrace; CKE: Chain-Kairali Escarpment; COB: Continent-Ocean transition.

Ridge (Plateau) is the bathymetric high feature in the southwest offshore margin and lies parallel to the west coast of India. The Laccadive Basin lies between the Laccadive Ridge and the continental shelf (fig. 2A) south of Tellicherry Arch. It is known well that the southwest margin was affected by Réunion plume volcanism towards the end of Cretaceous (Singh and


Lal, 1993) ~~Further, due to as revealed by~~ the presence of wide-spread trap layer below the thin Tertiary sediment cover at the margin (Singh et al., 2007; Singh and Lal, 1993), ~~as a result~~, the vintage seismic data was not of much help to decipher the basin architecture. ~~However, the recent high-resolution seismic data could image the deeper parts of the basin more clearly and provided some new insights on the pre-Tertiary history of the margin.~~

45 ~~Most of the drilled-wells along this margin neither penetrated the Paleocene trap layer nor encountered the crystalline basement, except the well CH-1-1 well located within the shelf, which penetrated through the trap and encountered Santonian formations (Singh and Lal, 1993).~~ One of the key ~~questions that remain to be question that was not~~ resolved is the absence of Late Cretaceous sediments in the ~~basin, that is the long-time Laccadive basin as a whole and the long time~~ gap of more than 20 ~~m.y.r Ma~~ between the India-Madagascar breakup at 83 Ma and the oldest sediments of ~~65 Ma encountered in almost all wells~~ ~~except at CH-1-1 well which could penetrate through Santonian formations (Singh and Lal, 1993)~~ Paleocene age. This long-time gap indicates either the presence of older sediments below the Paleocene trap layer or the opening of the Laccadive basin subsequent to the ~~Madagascar India-Madagascar~~ separation. Further, the accommodation of the continental fragments between India and Madagascar (Bhattacharya and Yatheesh, 2015; Torsvik et al., 2013) ~~brings new complexity to the geodynamics of this area makes a complex geodynamic setting~~, as how this separation took place, and therefore provides some ~~insight into the inheritance of the lithosphere that existed before insights into the pre-existing lithospheric inheritance~~. Hence, examining the ~~Laccadive Basin formation development of the Laccadive Basin~~ will provide important constraints on the early breakup evolution of the margin. In this study, we ~~present made a correlative analysis of the multi-channel seismic data with the residual gravity anomalies which provided~~ evidence of a major extensional event that occurred at the southwest margin that ~~cannot be correlated with is not related to~~ the India-Madagascar ~~separation or the India-Seychelles separation. This helped us to present an evolutionary model that explains the formation of extensional features and the opening of the Laccadive Basin. Understanding and breakup. The~~ time-stamping of this major extensional event ~~will provide provides~~ important constraints on the evolution of the ~~western continental margin of India and help in tight-fit reconstruction studies WCMI and help to build improved tectonic reconstruction models.~~

## 2 Description of tectonic elements ~~The of the study area under investigation falls towards~~


65 ~~The study area lies~~ south of Tellicherry Arch and contains the southern part of the Laccadive Ridge and the Laccadive Basin ~~in the offshore. (fig. 2A)~~. The major geomorphic features present in the study area from west to east are the Laccadive Ridge, the Laccadive Basin, the Alleppey Platform and Trivandrum Terrace together called the Alleppey-Trivandrum Terrace Complex (ATTC) ~~and the continental shelf (fig. 12A)~~. The ATTC is ~~bounded to characterized by horst-graben structures and bounded in~~ the west by the Chain-Kairali Escarpment (CKE) ~~and shows the horst-graben structures within it feature~~ (Yatheesh et al., 70 2006, 2013; Nathaniel, 2013) as revealed in the seismic sections (See fig. S1). ~~There are numerous Numerous~~ seamounts/guyots/knolls ~~are~~ present in the ~~Laccadive Basin which lies region~~ between the Laccadive Ridge and the continental shelf ~~within the Laccadive Basin (See fig. S2)~~ (Bijesh et al., 2018). In addition to this, the entire region is characterized by several intrusive

Regional tectonic map of the western continental margin of India (Smith and Sandwell, 1997) on the left. Solid green coloured lines represent the mapped seafloor spreading type magnetic lineations (Bhattacharya and Yatheesh, 2015, and references therein). Solid white lines represent the mapped fracture zones or pseudo-faults. Black solid lines in the offshore region represent shear zones and faults. The larger study area is marked in a black rectangle. The map on the right shows the position of India relative to Madagascar and Deccan Volcanic Province in late Paleozoic fit (Lovecchio et al., 2020, modified after). The area of interest is marked in red rectangle. SVP: Saurashtra volcanic province; DVP: Deccan Volcanic Province; DC: Dharwar Craton; SGT: Southern Granulite Terrain; LB: Laccadive Basin; AP: Alleppey Platform; TT: Trivandrum Terrace.




**Figure 3.** A) Satellite-derived free-air anomaly map B) Crustal Bouguer anomaly map. The location and orientation of identified extensional features, grabens and intrusives are marked. Black solid lines represent the location of the profiles. Interpreted seismic sections are shown in fig. 4. The thick broken reddish line in the centre of the basin represents the identified volcanic ridge. (refer to fig. 4 & S1 for seismic sections). CRS represents the Cannanore Rift System as identified by DGH. CKE: Chain-Kairali Escarpment; AP: Alleppey Platform; TT: Trivandrum Terrace.

structures within the Tertiary sediments (Unnikrishnan et al., 2023). The southern part of the Cannanore Rift system (CRS) identified over the Laccadive Ridge (fig. 3) by the Director General of Hydrocarbon (source: DGH) lies within the study area.



**Figure 4.** Figure showing interpreted seismic sections. The faults are marked and the intrusives are shown in dotted circles. The inset free-air anomaly map on the right bottom corner shows that location of seismic profiles. The time is given two-way travel times (TWT). TWT is the elapsed time for a seismic wave to travel from its source to a given reflector and return to a receiver at the Earth's surface.



**Figure 5.** A) Crustal Bouguer anomaly, B) Band-pass filtered crustal Bouguer anomaly, C) First vertical derivative (FVD) of band-pass filtered crustal Bouguer anomaly, D) Total horizontal derivative (THD) of band-pass filtered crustal Bouguer anomaly. The black lines show the structural highs and the blue dotted lines show the continuity of rift basins identified. The chartreuse coloured solid ticked line represents the Chain-Kairali escarpment (CKE). The brown line represents the identified volcanic ridge. The thin black lines close to the coast represents major faults identified on the continental shelf Singh and Lal (1993). TA: Tellicherry Arch.

### 75 3 Data and Methods

In this study, we used the satellite-derived free-air gravity (Sandwell et al., 2014) and bathymetry from General Bathymetric Chart of the Oceans (GEBCO, 2020) for comparative analysis with the long-offset multichannel multi-channel seismic reflection profiles. The large volume of industry seismic reflection data (see supplementary fig S3) at this margin provided information on the sediment thickness above the Paleocene trap layer and the intermediate horizons and various horizons

80 ~~within the post-Paleocene sediments~~ (Unnikrishnan et al., 2023). We also ~~compiled-gathered~~ a few published seismic sections (Nathaniel, 2013; Yatheesh et al., 2013) within the study area ~~-(see supplementary fig S1).~~


The ~~crustal Bouguer anomaly is calculated by removing the~~ gravity effects of ~~bathymetry and sediments~~ ~~the water layer and the sediments are removed~~ from the satellite-derived free-air anomaly ~~data which gave rise to crustal Bouguer anomalies~~. High-resolution sediment thickness derived from ~~FWT-two-way travel time (TWT)~~ maps is used to calculate the gravity effect of the sediments. The ~~two-way travel time (FWT)-TWT~~ maps are available for three different times: the early Paleocene, early Eocene, early Miocene ~~-(Unnikrishnan et al., 2023)~~. These maps were converted to depth with respective interval velocities ~~(after Unnikrishnan et al., 2023)~~ ~~(after Unnikrishnan et al., 2023)~~ and the total sediment layer is used to calculate the gravity effect of sediments ~~by assigning the~~. The densities of 2.3 g/cc for sediments and 1.03 g/cc for the water column ~~respectively are used~~. For the crustal rocks, an average density of 2.8 g/cc was considered as the study area lies within the extended continental crust ~~(Unnikrishnan et al., 2023)~~. A ~~band-pass filter of~~ ~~(Unnikrishnan et al., 2023)~~. A 10-200 km ~~wavelengths-wavelength band-pass filter~~ was applied to the crustal Bouguer anomaly map to highlight the crustal heterogeneities. ~~This map's~~, and the first vertical derivative (FVD) ~~was also as well as the total horizontal derivative (THD) of the band-pass filtered map was~~ prepared to identify shallow structural features. The ~~identified features~~ ~~location of the identified features such as rifts and volcanic intrusives~~ in the seismic ~~section-sections~~ are then transferred ~~on~~ to these gravity anomaly maps ~~and their continuity is mapped~~. ~~in order to understand their continuity~~. Scientifically derived colour maps are used to prepare maps (Cramer et al., 2020).

95 ~~Satellite-derived free-air anomaly map of the study area showing the location and orientation of identified extensional features, grabens and intrusives. Black solid lines represent the location of the profiles. The grabens marked in white are given in fig. S2. Interpreted seismic sections are shown on the right. The faults are marked and the intrusives are shown in circles. The locations of seismic sections is marked in the free-air anomaly diagram. The broken brown line in the centre of the basin represents the identified volcanic ridge. (refer to fig. S2 & S3 for more seismic sections). CRS represents the Cannanore Rift System as identified by DGH. CKE: Chain-Kairali Escarpment; AP: Alleppey Platform; TT: Trivandrum Terrace~~

#### 4 Results

~~We present twelve~~ ~~We present thirteen~~ interpreted seismic sections ~~which reveal~~ ~~(fig. 4)~~ ~~which reveal several rift-related horst-graben structures and extensional features in the Laccadive Ridge area (see fig. 2 & fig. S2-S3). In the area north of the Tellicherry Arch, the~~ ~~(extensional features) at the margin. We correlated these structures with the gravity anomaly trends and noticed that the~~ grabens are oriented in the NNW-SSE direction ~~interpreted as a ENE-WSW extension and towards the south, in the area north of the Tellicherry Arch, whereas,~~ the grabens are oriented in NNE-SSW direction ~~which is interpreted as extension is in the NNW-SSE direction.~~ ~~south of Tellicherry Arch (see fig. 3 & fig. 4)~~. These identified extensional structures show low gravity anomalies ~~and~~ the continuity of which can be traced as gravity lows surrounded by ~~gravity~~ highs in the anomaly maps (fig. ~~2-3A-B~~ & fig. ~~3A-C5A-D~~). This is particularly ~~true towards the south of the Tellicherry Arch. A volcanic intrusive is identified~~ ~~prominent within the study area (south of Tellicherry Arch). The seismic sections also reveal series of~~





**Figure 6.** Isochron maps prepared from the TWT for selected time intervals: A) Early Paleocene to Early Eocene; B) Early Eocene to Early Miocene; C) Early Miocene to Present day; and D) Total sedimentary layers. The brown line in B represents trend of the identified volcanic ridge. (Refer to text for detailed explanation and interpretation). TA: Tellicherry Arch; CKE: Chain-Kairali Escarpment.

volcanic intrusives in the centre of the Laccadive Basin which is traced almost parallel to the identified extensional trend in the south and to CKE (fig. 2). This is seen in the gravity anomaly map also as a broken chain of highs (fig. 2 & 3).-

- 115 A) Crustal Bouguer anomaly, B) Band-pass filtered crustal Bouguer anomaly, C) First vertical derivative (FVD) of band-pass filtered crustal Bouguer anomaly, D) Depth to the basement map with all identifications, E) Proposed Tectonic map of the study area, F)  $\beta$ -value map (Adapted from Unnikrishnan et al. (2023)). The black lines show the structural highs and the red lines show

the continuity of rift basins identified. The green solid line represents the Chain-Kairali escarpment (CKE). The broken brown line represents the identified volcanic ridge. Shelfal tectonic elements shown are from Singh and Lal (1993). TA: Tellicherry Arch.

120 , 4 & 5).

The isochron ~~map for the maps were prepared for three time periods in the study area (fig. 6). These maps show that for~~ the early Paleocene to early Eocene time interval ~~shows significant deposition of sediments~~, significant sediment deposition occurred parallel to the coast with very less sedimentation in the Laccadive Basin. During this period, maximum deposition ~~occurred took place~~ in the area between Tellicherry Arch and off Kochi (marked sediment patch in fig. ~~4A6A~~) with a minor sediment channel extending into the basin (~~fig. 4A marked sediment channel in fig. 6A~~). During the Early Eocene to Early Miocene time interval, the deposition within the sediment patch is almost absent, whereas, significant deposition is observed in the northern part of the Laccadive Basin on either side of the identified volcanic ridge (fig ~~4B6B~~). During Early Miocene to recent time, the sedimentation is uniform in the Laccadive Basin (fig. ~~4C6C~~). The sediment deposition ~~in the coast parallel grabens along the coast~~ remained high throughout the time intervals (fig ~~4A-C6A-C~~). The total sediment deposition pattern 130 shows that most of the sediments were accommodated parallel to the coast and towards the south there is an axis of high sediment deposition into the Laccadive Basin (Figure ~~4D6D~~).

Isochron maps prepared from the TWT for selected time intervals: A) Early Paleocene to Early Eocene; B) Early Eocene to Early Miocene; C) Early Miocene to Present day; and D) Total sedimentary layers. The brown line in B represents trend of the identified volcanic Ridge. (Refer to text for detailed explanation and interpretation). TA: Tellicherry Arch

## 135 5 DiscussionsDiscussion

### 5.1 Development of Cannanore Rift System (CRS)

The above analysis indicates that subsequent to the India-Madagascar breakup, the southwest margin of India encountered a passive extensional event which resulted in the separation of Laccadive ridge from the mainland. However, the change in the extensional direction from ENE-WSW in the northern part This study identifies two major extensional events in the southern 140 part of WCMI, one, the NNW-SSE oriented rift grabens over the Laccadive Ridge north of Tellicherry Arch to NW-SE in south which facilitated the opening of the Laccadive Basin needs to be considered in the generation of tight-fit reconstruction models for this part of the margin. The seafloor spreading magnetic lineations in the Mascarene Basin, and the second, an NNE-SSW graben system in the Laccadive Basin area south of Tellicherry Arch (fig. S1) provide some insights on the Indian-Madagascar separation. The identified anomalies clearly reveal that during the initial stages, the separation was E-W (83-79 Ma) and then 145 it changed to NE-SW (73 Ma) (Shuhail et al., 2018). During this period, the Cannanore Rift System (CRS) had developed due to continued extensional activity in the Eastern Basin of western 3). The NNW-SSE rifts conforms with the orientation of the onshore Dharwar structural trend which suggests that the pre-existing structural weakness have important influence on localization of deformation and thinning (Zwaan et al., 2021). Detailed analysis of multichannel seismic data in the southwest margin of India It is to be noted that the CRS extends southward till the Tellicherry Arch and ceases to exist further south (fig.

150 2). Normally one would expect this trend to continue southwards but a change in the direction of extension in the southern part signifies that this region may have a different evolutionary history compared to the northern part.

### 5.1 Distribution of Bathymetry highs and intrusives


A striking feature along this margin is the presence of many intrusives and bathymetric highs observed in the seismic and bathymetry data, respectively (Unnikrishnan et al., 2023; Bijesh et al., 2018). These features have very clear expressions on the gravity image of the area (figs. 2 & 3A-C). The intrusives and bathymetric highs in the study, south of Mangaluru by the Directorate General of Hydrocarbons (DGH) helped to identify a complex block-like basement structures comprising of grabens, half-grabens and faults over the Laccadive Ridge (source: DGH) north of Tellicherry Arch, and named as the Cannanore Rift System (CRS). These rifts were mostly observed along the eastern flank of the Laccadive Ridge and extends between 17°N-9.5°N parallel to the west coast of India. The NNW-SSE rifts identified in this study are part of the CRS (fig. 1) appear to be elongated roughly parallel to the trend of the Laccadive Basin. In the centre of the Laccadive Basin, we noticed a series of volcanic mounds with a trend almost parallel to the CKE (fig. 2), which are clearly expressed in the seismic sections. The observed trend correlates well with the crustal Bouguer anomaly map 3A-B). As mentioned earlier, India-Madagascar breakup took place at 83 Ma and resulted in the opening of the Mascarene Basin. The analysis of magnetic anomaly identifications together with plate-tectonic reconstruction studies (Shuhail et al., 2018) revealed that during the initial period (83-79 Ma), the spreading in the Mascarene Basin was E-W (fig. 3A) as well as the depth to basement map prepared by removing the sediment thickness from sea bottom (fig. 3D). This trend divides the Laccadive Basin into eastern and western basins. The composite tectonic map of the study area is shown in fig. 3E2B). This spreading direction matches well with the NNW-SSE oriented rift system observed over the Laccadive Ridge which had resulted from the E-W extension during initial stages of India-Madagascar separation.

### 5.1 Opening of the Laccadive Basin

Since the entire Laccadive Ridge area was in extensional regime during India-Madagascar (as it was sandwiched between India and Madagascar), normally one would expect this trend to continue further southwards in the Laccadive basin area. Further, the  $\beta$ -value (crustal stretching factor) map (fig. 3F) calculated clearly reveals the extensional structures. However, the observed NNE-SSW orientation of the identified grabens in the study area (Unnikrishnan et al., 2023). The high  $\beta$ -values in the centre of south of Tellicherry Arch suggests that the Laccadive Basin indicate maximum thinning, confirming our observation in this study.

### 5.2 Timing of opening

had formed through a different evolutionary history that is not related to India-Madagascar separation. The sediment deposition in the basin is interpreted from the perspective of the creation of accommodation space and sediment supply. The available high-resolution time-structure maps (fig. 4A-C6A-D) provide insights on the timing of opening of the Laccadive Basin. These



**Figure 7.** A) Depth to the basement map with all structures identified in the study, B)  $\beta$ -value map redrawn from Unnikrishnan et al. (2023), C) Proposed Tectonic map of the study area. Bottom panel shows a seismic section (Seis1), showing the general characteristic of the Laccadive Basin area. The seismic section is from Unnikrishnan (2018). The black lines show the structural highs and the blue dotted lines show the continuity of rift basins identified. The chartreuse coloured solid ticked line represents the Chain-Kairali escarpment (CKE). The solid brown line represents the identified volcanic ridge. The thin black lines close to the coast in A & B represents major faults identified on the continental shelf (Singh and Lal, 1993). TA: Tellicherry Arch; AP: Alleppy platform; TT: Trivandrum Terrace; CKE: Chain-Kairali Escarpment. AP: Alleppy platform; VR: Volcanic Ridge.

maps clearly reveal significant sediment deposition along the coast parallel grabens within the shelfal part of the margin in all time periods. Further, during the Paleocene-Eocene period, sediment deposition was very significant on the northern fringe of the Laccadive Basin (sediment patch in fig. 4A-6A) with negligible sediments elsewhere in the basin. During Eocene to Miocene the sediment deposition shifted further offshore into the Laccadive Basin (Fig. 4B-fig. 6B). The development of the median-high volcanic ridge within the basin is also noticed. This occurred during this time since it is observed that the sediments are deposited on either side of the location of the ridge. Further the seismic sections show sediments onlapping to the identified volcanic ridge (fig. 4 sections NN' & PP'). The comparative analysis of fig. 6A & 6B indicates that the basin opened sometime after some-time after the early Eocene as a result of which accommodation space was created and all the incoming sediments migrated southward into the basin. Note that the sediment deposition occurred on either side of the identified volcanic ridge.

185  
190 A small channel of sediment deposition into the Laccadive Basin towards the southwest of the sediment patch (marked as sediment channel in Fig. 4A-6A) may represent the initial stage of opening of the basin. During the Miocene to recent period

(Fig. 4C6C), the sediment deposition is more or less uniform throughout the basin. The western edge of the western basin has relatively less deposition which may be due to the area's location far from any sediment supply. ~~By this time, the basin attained the present-day configuration.~~

195 Unnikrishnan et al. (2018) identified the Alleppy platform as a continental fragment and inferred its development during the Oligocene-Miocene period, ~~which agrees with this study's observations.~~ Alleppy platform is located adjacent to the Laccadive Basin and hence the development of the basin and the platform is related. The timing of the development of the Alleppy platform given by Unnikrishnan et al. (2018) closely agrees with the inferred timing of the opening of the Laccadive Basin from this study.

200 ~~Map showing the evolution of the region in four stages. Stage I: The pre-rift juxtaposition of the continental fragments. Note that the Laccadive Ridge is larger since it incorporates all the fragments that are littered between India and Madagascar. Stage II: The formation of the faults system or the rifts system due to the influence of spreading in the Mascarene Basin. Stage III: The opening of the basin with CKE + ATTC and India moving away from the Laccadive Ridge. See how the orientation of the extensional feature's changes with the anticlockwise movement of India. Stage IV: The present-day configuration of the margin with all extensional features and the volcanic ridge. LaX: Laxmi Ridge; SEY: Seychelles; MAD: Madagascar; SL: Sri Lanka; LR: Laccadive Ridge~~

## 5.2 Opening Distribution of the Laccadive Basin Bathymetry highs and intrusives

A striking feature along this margin is the presence of many intrusives and bathymetric highs (see fig. S2) observed in the seismic and bathymetry data, respectively (Unnikrishnan et al., 2023; Bijesh et al., 2018). These features have very clear expressions on the gravity image of the area (figs. 3 & 5A-D). The intrusives and bathymetric highs in the study area, south of Tellicherry Arch (fig. S2) appear to be elongated roughly parallel to the trend of the Laccadive Basin. In the centre of the Laccadive Basin, we noticed a series of volcanic mounds with a trend almost parallel to the CKE (fig. 3), which are clearly expressed in the seismic sections (fig. 4). The observed trend correlates well with the crustal Bouguer anomaly map (fig. 5A) as well as the depth to basement map prepared by removing the sediment thickness from sea bottom (fig. 7A). This trend divides

215 the Laccadive Basin into eastern and western basins. The composite tectonic map of the study area is shown in fig. 7C.

Further, the  $\beta$ -value (crustal stretching factor) map (fig. 7B) calculated (Unnikrishnan et al., 2023) clearly reveals the extensional trend in the study area. The high  $\beta$ -values in the centre of the Laccadive Basin indicate maximum thinning, confirming our observation in this study.

~~During the India-Madagascar separation, the Laccadive area was near the fracture zones in the Mascarene Basin. It has been proposed that the CKE was connected to the spreading in the Mascarene Basin through a long transform fault (Shuhail et al., 2018). There are~~

220

## 5.3 Evolutionary model

Existing plate tectonic reconstruction models (Bhattacharya and Yatheesh, 2015; Shuhail et al., 2018), the pre-existing structural trends along WCMI, India and Madagascar (Subrahmanyam et al., 1994; Kolla and Coumes, 1990; Bhattacharya and Yatheesh, 2015)

225 , and the results from this study have been used to build a schematic evolutionary model for the opening of the Laccadive Basin at the southwest margin of India. Additionally, we also looked at analog and numerical modelling studies that explain extension along passive continental margin (Péron-Pinvidic and Manatschal, 2010; Zwaan et al., 2021; Bonini et al., 1997; Henza et al., 2010). The locations of India and Madagascar in various stages (I-IV) were adopted from the reconstruction model of Shuhail et al. (2018). The salient aspects of each stage is given below:

### 230 5.3.1 Stage I

Stage I shows India and Madagascar along with all the continental fragments in between them and a large number of suture zones ~~on land in onshore for~~ both India and Madagascar (Bhattacharya and Yatheesh, 2015, and references therein) (~~fig-1 & fig-5 Stage I~~) which immediately after the spreading started at 83 Ma. These suture zones were earlier used as piercing points to find the relative position of India and Madagascar in ~~plate reconstruction studies (Katz and Premoli, 1979; Subrahmanyam and Chand, 2006)~~.  
235 ~~Earlier studies along WCMI has the matching of coastlines (Katz and Premoli, 1979; Subrahmanyam and Chand, 2006).~~ Besides, studies along the WCMI have shown the extension of onshore structural trends into the offshore region (Subrahmanyam et al., 1994; Kolla and Coumes, 1990) and studies (~~Péron-Pinvidic and Manatschal, 2010~~) (Péron-Pinvidic and Manatschal, 2010; show that structural inheritance plays a role during rifting and breakup. Dating of volcanics in Madagascar yields an age of 88 Ma (Storey et al., 1995; Pande et al., 2001) and magnetic anomalies indicate spreading started around 83 Ma (Shuhail et al., 2018).  
240 Hence, we infer that ~~these structural trends would have continued into the continental fragments located in between them. Due to this~~ during this extensional span, CRS formed on the Laccadive Ridge following the Precambrian Dharwarian trend which is dominant north of Tellicherry Arch.

### 5.3.2 Stage II

Stage II shows the proximity of Laccadive Ridge to the spreading centre. As discussed in stage I, the lithosphere between  
245 India and Madagascar was weak and as a result, when the area was proximal to the spreading centre ~~in the Mascarene Basin,~~ a number of parallel trans-tensional faults may have formed on the Laccadive Ridge south of Tellicherry Arch (~~fig-5 stage II~~).  
~~Subsequently.~~ Even though the spreading was happening along the entire basin, the spreading was very active in the southern part of the Mascarene basin, and also characterized by long transform faults (Shuhail et al., 2018). Also, Shuhail et al. (2018) proposed that the CKE was connected to the spreading in the Mascarene Basin through a long transform fault. As discussed in  
250 section 5.1 distribution of bathymetric highs and intrusives south of Tellicherry Arch provide some evidence for this. It is very likely that, later when the Reunion passed over the area, the magma may have migrated through the faults formed during this stage and hence intrusives show this orientation. Similar arguments can also explain the orientation of bathymetric features in this area.

### 5.3.3 Stage III

255 Stage III shows the initial stages of opening of the Laccadive basin. By this time, the entire region was flooded by Deccan volcanics during the passage of Réunion plume in the Paleocene time. Studies by Patriat and Achache (1984) and Dewey (1989) showed that the Indian plate rotated anticlockwise about  $40^{\circ}$  since 84 Ma, out of which, it underwent about  $25^{\circ}$  after the soft collision at 50 Ma (Treloar and Coward, 1991). The ~~plume's proximity coupled with India's~~ passage of the Réunion plume over the area may have weakened the overlying lithosphere and together with India's anti-clockwise rotation and the  
260 presence of inherited structural weakness may have led to the reactivation and further extension in the Laccadive Basin. The sediment thickness data during the Paleocene-Eocene period shows a sediment channel (fig. 5-stage III). The ATTC remained 6A) extending into the Laccadive basin and an area of high sediment deposition (Sediment patch in fig. 6A) which indicate the initial stages of opening of the basin. Fig. 6B shows sediments being deposited on either side of the volcanic ridge in the Laccadive basin. By this time the basin opened and sediments were accommodated in the basin.

#### 265 5.3.4 Stage IV

Stage IV shows the present-day configuration of the Laccadive Basin. The ATTC remain attached to the Indian continent with CKE forming its western boundary. The centre of the Laccadive Basin experienced maximum crustal thinning and a series of intrusives got emplaced in the crust (~~fig. 5 stage IV~~). The change in the orientation of horst-graben structures in the ATTC from that in the southern part of Laccadive Ridge can be noticed.

## 270 6 Conclusions

The seismic and gravity data analysis along southern part of WCMI (~~the area south of Tellicherry Arch) has a different evolutionary history than the northern segment. The study provided evidences of pre-Deccan ENE-WSW extensional tectonics revealed two significantly different orientations of the extensional graben system, one a NNW-SSE oriented rifts over the Laccadive Ridge, north of Tellicherry Arch which were masked by volcanic traps. In the~~, and the other NNE-SSW oriented  
275 rifts in the Laccadive Basin region south of Tellicherry Arch, ~~a significant change is observed in the extensional direction to ENE-WSW in the post-Deccan period. This is evidenced by tilted intrusive features within grabens bounded by normal faults in the seismic section (Fig. 2 seismic section DD').~~ The change in extensional direction in the Laccadive basin along with the isochron maps for different times provided evidence of opening of the Laccadive Basin. The Laccadive Basin opened within the post-Eocene period with maximum extension along the centre of the basin where the volcanic intrusives are emplaced.  
280 The lithosphere that existed had zones of weakness and this along with the ~~plume-plume proximity to Réunion plume and anti-clockwise rotation of India~~ led to complex rifting and ~~evolution of evolutionary history for~~ the southern part of the margin.

*Data availability.* The authors do not have permission to share data.

*Author contributions.* MGG – conceptualisation, methodology, validation, formal analysis, writing (original draft and editing), visualisation. MR – conceptualisation, validation, resources, writing ( review and editing), supervision. PU – conceptualisation, writing (review and editing).  
285

*Competing interests.* The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.


*Acknowledgements.* This study forms a part of PhD work of MGG at IIT Bombay. MGG thanks the Council of Scientific and Industrial Research (CSIR), New Delhi, India, for the Financial support under Junior Research Fellowship (Grant No. 09/087(1005)/2019-EMR-I).  
290 The second author gratefully acknowledges the permission of the Oil and Natural Gas Corporation Limited (ONGC) to use the geoscientific data pertaining to the Kerala-Konkan Basin. The authors acknowledge the support from Science & Engineering Research Board (SERB/CRG/2021/006505).



## References

- Bhattacharya, G. C. and Yatheesh, V.: Plate-tectonic evolution of the deep ocean basins adjoining the western continental margin of India—a proposed model for the early opening scenario, *Petroleum geosciences: Indian contexts*, pp. 1–61, 2015.
- 295 Bijesh, C., Kurian, P. J., Yatheesh, V., Tyagi, A., and Twinkle, D.: Morphotectonic characteristics, distribution and probable genesis of bathymetric highs off southwest coast of India, *Geomorphology*, 315, 33–44, 2018.
- Bonini, M., Souriot, T., Boccaletti, M., and Brun, J. P.: Successive orthogonal and oblique extension episodes in a rift zone: Laboratory experiments with application to the Ethiopian Rift, *Tectonics*, 16, 347–362, 1997.
- 300 Chaubey, A., Dymant, J., Bhattacharya, G. C., Royer, J.-Y., Srinivas, K., and Yatheesh, V.: Paleogene magnetic isochrons and palaeo-propagators in the Arabian and Eastern Somali basins, NW Indian Ocean, Geological Society, London, Special Publications, 195, 71–85, 2002.
- Collier, J., Sansom, V., Ishizuka, O., Taylor, R., Minshull, T., and Whitmarsh, R.: Age of Seychelles–India break-up, *Earth and Planetary Science Letters*, 272, 264–277, 2008.
- 305 Crameri, F., Shephard, G. E., and Heron, P. J.: The misuse of colour in science communication, *Nature communications*, 11, 5444, 2020.
- Dewey, J.: Tectonic evolution of the India/Eurasia collision zone, *Eclogae Geologicae Helveticae*, 82, 717–734, 1989.
- DGH: Directorate General of Hydrocarbons (DGH), Noida, India web page: Kerala Konkan Basin, <https://www.dghindia.gov.in/index.php?page?pageId=67&name=Indian%20Geology>, last access 23 June 2023.
- Ganerød, M., Torsvik, T., Van Hinsbergen, D., Gaina, C., Corfu, F., Werner, S., Owen-Smith, T., Ashwal, L., Webb, S., and Hendriks, B.: Palaeoposition of the Seychelles microcontinent in relation to the Deccan Traps and the Plume Generation Zone in Late Cretaceous–Early Palaeogene time, Geological Society, London, Special Publications, 357, 229–252, 2011.
- 310 GEBCO: General Bathymetric Chart of the Oceans, <https://doi.org/10.5285/a29c5465-b138-234d-e053-6c86abc040b9>, 2020.
- Henza, A. A., Withjack, M. O., and Schlische, R. W.: Normal-fault development during two phases of non-coaxial extension: An experimental study, *Journal of Structural Geology*, 32, 1656–1667, 2010.
- 315 Katz, M. and Premoli, C.: India and Madagascar in Gondwanaland based on matching Precambrian lineaments, *Nature*, 279, 312–315, 1979.
- Kolla, V. and Coumes, F.: Extension of structural and tectonic trends from the Indian subcontinent into the eastern Arabian Sea, *Marine and petroleum Geology*, 7, 188–196, 1990.
- Lovecchio, J. P., Rohais, S., Joseph, P., Bolatti, N. D., and Ramos, V. A.: Mesozoic rifting evolution of SW Gondwana: a poly-phased, subduction-related, extensional history responsible for basin formation along the Argentinean Atlantic margin, *Earth-Science Reviews*, 320, 203, 103 138, 2020.
- Nathaniel, D.: Hydrocarbon potential of sub-basalt Mesozoics of deepwater Kerala Basin, India, in: 10th Biennial International Conference & Exposition Kochi, pp. 1–7, 2013.
- Pande, K., Sheth, H. C., and Bhutani, R.: 40Ar–39Ar age of the St. Mary’s Islands volcanics, southern India: record of India–Madagascar break-up on the Indian subcontinent, *Earth and Planetary Science Letters*, 193, 39–46, 2001.
- 325 Patriat, P. and Achache, J.: India–Eurasia collision chronology has implications for crustal shortening and driving mechanism of plates, *Nature*, 311, 615–621, 1984.
- Péron-Pinvidic, G. and Manatschal, G.: From microcontinents to extensional allochthons: witnesses of how continents rift and break apart?, *Petroleum Geoscience*, 16, 189–197, 2010.

- Sandwell, D. T., Müller, R. D., Smith, W. H., Garcia, E., and Francis, R.: New global marine gravity model from CryoSat-2 and Jason-1  
330 reveals buried tectonic structure, *Science*, 346, 65–67, 2014.
- Shellnutt, J., Lee, T.-Y., Chiu, H.-Y., Lee, Y.-H., and Wong, J.: Evidence of Middle Jurassic magmatism within the Seychelles microcontinent: Implications for the breakup of Gondwana, *Geophysical Research Letters*, 42, 10–207, 2015.
- Shellnutt, J. G., Yeh, M.-W., Suga, K., Lee, T.-Y., Lee, H.-Y., and Lin, T.-H.: Temporal and structural evolution of the Early Palaeogene rocks of the Seychelles microcontinent, *Scientific reports*, 7, 179, 2017.
- 335 Shuhail, M., Yatheesh, V., Bhattacharya, G. C., Müller, R. D., Raju, K. K., and Mahender, K.: Formation and evolution of the Chain-Kairali Escarpment and the Vishnu Fracture Zone in the western Indian Ocean, *Journal of Asian Earth Sciences*, 164, 307–321, 2018.
- Singh, K., Radhakrishna, M., and Pant, A.: Geophysical structure of western offshore basins of India and its Implications to the evolution of the Western Ghats, *Geological Society of India*, 70, 445–458, 2007.
- Singh, N. K. and Lal, N. K.: Geology and Petroleum Prospects of Konkan-Kerala Basin, in: *In: Proc. 2nd Seminar on Petroleum Basins of India*, KDMIPE and ONGC, vol. 2, pp. 461–469, 1993.
- 340 Smith, W. H. and Sandwell, D. T.: Global sea floor topography from satellite altimetry and ship depth soundings, *Science*, 277, 1956–1962, 1997.
- Storey, M., Mahoney, J. J., Saunders, A. D., Duncan, R. A., Kelley, S. P., and Coffin, M. F.: Timing of hot spot—related volcanism and the breakup of Madagascar and India, *Science*, 267, 852–855, 1995.
- 345 Subrahmanyam, C. and Chand, S.: Evolution of the passive continental margins of India—a geophysical appraisal, *Gondwana Research*, 10, 167–178, 2006.
- Subrahmanyam, V., Krishna, K., Murthy, G., Rao, D. G., Ramana, M., and Rao, M. G.: Structural interpretation of the Konkan basin, southwestern continental margin of India, based on magnetic and bathymetric data, *Geo-marine letters*, 14, 10–18, 1994.
- Torsvik, T. H., Amundsen, H., Hartz, E. H., Corfu, F., Kusznir, N., Gaina, C., Doubrovine, P. V., Steinberger, B., Ashwal, L. D., and Jamtveit, B.: A Precambrian microcontinent in the Indian Ocean, *Nature geoscience*, 6, 223–227, 2013.
- 350 Treloar, P. J. and Coward, M. P.: Indian Plate motion and shape: constraints on the geometry of the Himalayan orogen, *Tectonophysics*, 191, 189–198, 1991.
- Unnikrishnan, P.: Crustal structure, tectonic and sedimentation history along the Kerala-Konkan, basin, western continental margin of India based on integrated geophysical studies, Ph.D. thesis, IIT Bombay, Mumbai, India, 2018.
- 355 Unnikrishnan, P., Radhakrishna, M., and Prasad, G. K.: Crustal structure and sedimentation history over the Alleppey platform, southwest continental margin of India: Constraints from multichannel seismic and gravity data, *Geoscience Frontiers*, 9, 549–558, 2018.
- Unnikrishnan, P., Gilbert, M., and Radhakrishna, M.: Crustal structure along the Kerala-Konkan Basin, southwest continental margin of India, using multi-channel seismic and gravity modelling: Implications on India-Madagascar rifting and basin evolution, *Journal of Asian Earth Sciences*, 242, 105 504, 2023.
- 360 Yatheesh, V., Bhattacharya, G. C., and Mahender, K.: The terrace like feature in the mid-continental slope region off Trivandrum and a plausible model for India–Madagascar juxtaposition in immediate pre-drift scenario, *Gondwana Research*, 10, 179–185, 2006.
- Yatheesh, V., Kurian, P. J., Bhattacharya, G. C., and Rajan, S.: Morphotectonic architecture of an India–Madagascar breakup related anomalous submarine terrace complex on the southwest continental margin of India, *Marine and Petroleum geology*, 46, 304–318, 2013.
- Zwaan, F., Chenin, P., Erratt, D., Manatschal, G., and Schreurs, G.: Complex rift patterns, a result of interacting crustal and mantle weaknesses, or multiphase rifting? Insights from analogue models, *Solid Earth*, 12, 1473–1495, 2021.
- 365



**Figure 8.** Map showing the evolution of the region in four stages. Stage I: The pre-rift juxtaposition of the continental fragments. Note that the Laccadive Ridge is larger since it incorporates all the fragments that are littered between India and Madagascar. Stage II: The formation of the faults system or the rifts system due to the influence of spreading in the Mascarene Basin. Stage III: The opening of the basin with CKE + ATTC and India moving away from the Laccadive Ridge. See how the orientation of the extensional feature's changes with the anticlockwise movement of India. Stage IV: The present-day configuration of the margin with all extensional features and the volcanic ridge. LaX: Laxmi Ridge; SEY: Seychelles; MAD: Madagascar; SL: Sri Lanka; LR: Laccadive Ridge; ATTC: Alleppey-Trivandrum Terrace Complex; CKE: Chain-Kairali Escarpment.