

Climatology of Large Hail in Europe: Characteristics of the European Severe Weather Database

Faye Hulton^{1,a}, David M. Schultz^{1,2}

¹ Centre for Atmospheric Science, Department of Earth and Environmental Sciences, University of Manchester, Manchester, M13 9PL, United Kingdom

² Centre for Crisis Studies and Mitigation, University of Manchester, Manchester, M13 9PL, United Kingdom

^a Now at: MetDesk, Aylesbury, HP22 6NJ, United Kingdom

Correspondence to: Faye Hulton, faye.hulton@gmail.com

Submitted as an Article to *Natural Hazards and Earth System Science*, 4 February 2023, Revised 24 May 2023

Abstract. Large hail (greater than 2 cm in diameter) can cause devastating damage to crops and property, and can even cause loss of life. Because hail reports are often collected by individual countries, constructing a European-wide large-hail climatology has been challenging to date. However, the European Severe Storm Laboratory's European Severe Weather Database provides the only pan-European dataset for severe convective storm reports. The database is comprised of 62,053 large-hail reports from 40 C.E. to September 2020, yet its characteristics have not been evaluated. Thus, the purpose of this study is to evaluate hail reports from this database for the purposes of constructing a climatology of large hail. For the period 2000–2020, large-hail reports are most prominent in June, whereas large-hail days are most common in July. Large hail is mostly reported between 1300–1900 local time, a consistent pattern since 2010. The intensity, as measured by maximum hail size, shows decreasing frequency with increasing hailstone diameter, and little change over the 20-year period. The quality of reports by country varies, with the most complete reporting being from central European countries. These results suggest that despite its short record, many indications are that the dataset represents some reliable aspects of European large-hail climatology, albeit with some limitations.

1 Introduction

Hail, with a diameter of at least 2 cm in the longest direction is called *large hail*, and it can cause damage to crops, property, or even loss of life. Several recent studies have documented the occurrence and variability of large hail, with special emphasis on the United States and Europe where large hail is common (e.g., Allen and Tippett 2015; Punge and Kunz 2016; Brooks et al. 2019; Púčik et al. 2019; Tang et al. 2019; Taszarek et al. 2020; Raupach et al. 2021). The strongest severe convective storms in Europe are often perceived to be less intense than the strongest storms in the United States, although they can be just as damaging. For example, one of the most devastating large-hail events took place over Germany on 28 July 2013 when two supercells formed almost simultaneously, producing hailstones of up to 10 cm in diameter and more than EUR 1 billion in insurance payouts (Kunz et al. 2018). Other similar events occurred over southern Germany on 10–12 June 2019, with one storm producing 6-cm hailstones and causing EUR 1 billion in damages (Wilhelm et al. 2021). More recently, several large-hail events were reported during summer 2021 in Poland, the Czech Republic, Germany, and Italy, with reported maximum hail sizes in excess of 7 cm (Associated Press 2021; Space 2021a,b,c). Although these extreme events are widely reported by the media, meteorological research on these storms may be hindered by the lack of ground-truth hail data, such as onset and ending times, duration, and hailstone size.

Deleted: For s

Deleted: ssion

Deleted: 2

Deleted: s

Deleted: no

Deleted: in Europe

Deleted: Another

Deleted: intensity,

52 Such hail data in Europe is generally collected on a national scale, and hence most climatologies are produced
53 on a country-by-country basis (e.g., Brooks et al. 2009). Given the relatively small sizes of many European
54 countries, each country has a low probability of large hail occurring at any given time (e.g., Brooks et al. 2019).
55 A summary table of past European hail climatologies can be found in Tuovinen et al. (2009), and a review was
56 published by Punge and Kunz (2016). Because countries that have a similar spatial extent as Europe have produced
57 their own climatologies—such as the United States (Tang et al. 2019), Canada (Etkin and Brun 2001), and China
58 (Zhang et al. 2008)—a pan-European large-hail climatology would be highly desired.

59 Climatologies of European convective storms and their impacts have been constructed using a number of
60 datasets. For example, some studies have examined the climatology of convective storms using remote-sensed
61 data such as lightning, radar, and satellite (e.g., Punge et al. 2017). Others have examined the environments that
62 favor such storms, such as through reanalyses or soundings (Rädler et al. 2018; Taszarek et al. 2017, 2018, 2019)
63 or reanalyses coupled with hailpad data (Sanchez et al. 2017).

64 To create a pan-European dataset of surface reports from severe convective storms (including large hail,
65 tornadoes, and severe wind gusts), the European Severe Storms Laboratory formed the European Severe Weather
66 Database (ESWD) in 2006 (Dotzek et al. 2009; Groenemeijer et al. 2017). In addition to collecting contemporary
67 data, the ESWD has an ongoing objective of synthesizing historical large-hail data which helps produce a longer
68 and more complete climatology. Despite the tremendous potential value of the ESWD being the only pan-
69 European large-hail dataset, its characteristics need to be examined to understand its suitability for answering
70 certain scientific questions about large hail. For example, Taszarek et al. (2019) found substantial variability
71 across Europe in the frequency of ESWD reports and the frequency of favorable environments for convective
72 storms.

73 To this effect, Púčik et al. (2019) constructed a climatology of large hail from the ESWD. They examined
74 hail size, occurrence, annual cycle, diurnal cycle, and societal impacts (e.g., damages, injuries) for 39,537 reports
75 during the 13-yr period 2006–2018. Although their work shed the first light on the pan-European distribution and
76 characteristics of large hail and large-hail days from surface reports, they concluded by foreseeing “an update to
77 this study as the reporting homogeneity improves in future.” In the present article, we explore whether increasing
78 the size of the dataset through lowering the quality-control levels of the reports and extending the period of
79 analysis yields comparable results, increasing the generality of Púčik et al.’s (2019) results. In doing so, we also
80 document the reporting characteristics of the database as a function of time both throughout the 20th century and
81 within the last 20 years. In particular, we seek the possible existence of a relatively homogeneous period of time
82 in the database that could be used as a baseline for climatologies and climate-change studies.

83 This article consists of nine sections. Section 2 describes the data from the ESWD used in the present study.
84 Section 3 discusses the frequency of large-hail reports and days on decadal, annual, and diurnal time scales.
85 Section 4 investigates the intensity distribution of large hail, as segregated into 1-cm diameter bins, and discusses
86 how the frequency of large-hail size has changed over the past 20 years. Section 5 looks at the time accuracy of
87 these reports, how it has changed over the past 20 years, and how it varies by individual countries. Section 6
88 investigates the spatial distribution of reports by country. Because of the large number of reports from Poland
89 during the 1930s to 1950s, section 7 focuses on the data from Poland, comparing the historical frequency of reports
90 during this period to that from the period 2000–2020. Section 8 offers a discussion comparing our work to previous

Deleted:

Formatted: Font color: Text 1

Formatted: Font: Times New Roman, 10 pt, Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font: Times New Roman, 10 pt, Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font: Times New Roman, 10 pt, Font color: Text 1

Deleted: ¶

Deleted: will

Deleted: ¶

95 hail climatologies and reflects on the prospects of using the ESWD as a baseline for climate-change research.
96 Section 9 summarizes the findings of this paper.

97

98 2 Data and methods

99 The climatology of European large hail in this present article is produced from the ESWD (Dotzek et al. 2009;
100 Groenemeijer et al. 2017). Large hail in the ESWD is defined as hail with a diameter of at least 2 cm in the longest
101 direction (Groenemeijer and Liang 2020), comparable to the severe-hail criterion of 0.75 inch (1.9 cm) in the
102 United States. The current ESWD data on hail is a mixture of historical entries, insurance data information, reports
103 provided by storm-spotters, national European meteorological organizations, and public entries via the ESWD
104 website at www.eswd.eu (Dotzek et al. 2009). Since December 2015, reports have also been collected via ESSL's
105 European Weather Observer app (Groenemeijer et al. 2017).

Deleted: organization

106 At the time this study commenced, the ESWD consisted of 62,053 large-hail reports from 59 countries dating
107 from 40 C.E. to 26 September 2020. All reports with hail sizes less than 2 cm were removed. Of the 59 countries
108 included with the initial dataset received from the European Severe Storms Laboratory, only 41 were in Europe.
109 Of those removed, the highest reporting countries were Turkey, Armenia, and Azerbaijan. Reports from other
110 countries that were removed included Morocco, Turkmenistan, Egypt, and Jordan. The Russian Federation was
111 included in the present study, even though a small number of reports were from the Asian part of the country. A
112 small part of Turkey is geographically in Europe, but their data was not included in this study.

Deleted: T

113 We also examined two periods of time from the ESWD. The first period is the nearly 121-yr period from 1
114 January 1900 to 26 September 2020 (when work on this research commenced). We hereafter refer to this period
115 as 1900–2020, recognizing the omission of data from the last three months and four days of 2020. The second
116 period is more focused on the most recent large-hail data for the nearly 21-yr period 1 January 2000 to 26
117 September 2020, hereafter referred to as 2000–2020.

118 All data is imputed in a standard format and is given a single quality-control level by the maintenance team
119 (Dotzek et al. 2009). There are four quality-control levels given to these entries (Groenemeijer and Kühne 2014):

- 120 • Q0: “as received”, any report straight from the public,
- 121 • QC0+: “plausibility checked”, any report checked by staff at the European Severe Storms Laboratory or a
122 partner organization,
- 123 • QC1: “report confirmed”, any report confirmed by a reliable source such as a national meteorological
124 organization or storm-spotter network, and
- 125 • QC2: “event fully verified”, any report from an event that has been subject of a scientific case study.

126 As mentioned in section 1, Půčik et al. (2019) used only QC1 and QC2 events. However, to see if the quality-
127 control level affects the interpretation of the results, this present study uses QC0+, QC1, and QC2. For the period
128 1900–2020, there were 9173 QC0+, 45,805 QC1, and 2391 QC2 reports, producing a total of 57,369 large-hail
129 reports. For the period 2000–2020, there were 6330 QC0+, 20,585 QC1, and 1310 QC2 reports, producing a total
130 of 28,225 large-hail reports. Thus, the addition of the QC0+ reports increased the size of the 1900–2020 dataset
131 by 19% and the 2000–2020 dataset by 29%.

Deleted: plausibly checked

132 With these two datasets constructed, we can then look at their characteristics. In particular, we are interested
133 in the number of large-hail days, size of the large-hail reports, and time accuracy of the reports. The annual
134 number of large-hail days was derived from the annual number of large-hail reports by removing duplicate dates.

138 The size of the hail in each hail report was defined as the maximum hail diameter recorded in cm. Although the
139 ESWD contains fields for the fall speed and density of the hailstones, these were infrequently reported and were
140 not considered as part of the present article. To represent the size distribution of the reports, the reports were
141 classified into 1-cm bins based on their maximum hail diameter, starting at the minimum threshold of large hail
142 of 2 cm. The *time accuracy* of reports is a field in the ESWD that allows the user to know how reliable the
143 reporting time of the large-hail report is. The time accuracy represents the total time window that a given report
144 was recorded in. For example, a 30-min time accuracy would indicate that the hail fell in the window of 15 min
145 before the recorded time to a maximum of 15 min after the recorded time. The existing ESWD dataset is a result
146 of both meteorological variations in hail and reporting issues, much as other severe-weather datasets have (e.g.,
147 Groenemeijer and Kühne 2014; Punge and Kunz 2016; Antonescu et al. 2017; Púčik et al. 2019). Indeed,
148 underreporting from rural areas and nighttime storms may influence this dataset. ~~These and other characteristics~~
149 of the large-hail dataset will be explored in subsequent sections.

Moved down [1]: These and other characteristics of the large-hail dataset will be explored in subsequent sections.

Moved (insertion) [1]

Formatted: Not Strikethrough

150

151

3 Frequency of large hail across Europe: 1900–2020

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171


172

173

To understand the number of large-hail reports as a function of time, the annual number of large-hail reports and annual number of large-hail days were plotted versus year from 1900 to 2020 (Fig. 1). Throughout much of this period, the annual number of reports was quite small, with peaks during the 1930s, 1940s–1950s, and early 1980s before a steady increase starting around 2000. These two peaks in the 1930s and 1940s–1950s were associated with a large number of reports from Poland and are investigated further in section 8. The lesser peak during the 1980s was associated with a number of reports from Italy, but is not considered further.

Figure 1 also shows the annual number of hail days from 1900 to 2020. The peaks in large-hail days during the 1930s and 1940s–1950s suggest that there were many large-hail events, not just many reports. Moreover, these periods illustrate that, while some periods and some locations may be well represented in the database, reporting of large hail throughout much of the 20th century in the ESWD is far from complete.

Focusing on the last 30 years, the number of reports increased starting around 2000 and continued to rise until 2020. (Recall that the 2020 data was only available until 26 September, which may explain the fewer number reports, although most large-hailfall in Europe is reported between April and September.) In contrast, the number of large-hail days began rising a few years earlier in the late 1990s before reaching a plateau during the 2010s with around 175 annual large-hail days per year, similar to Taszarek et al. (2020, their Fig. 2a). This result suggests that the database grew around this time by first obtaining data from a larger number of days on which hail fell, followed by the database growing with a larger number of reports within the same day. The inconsistency in reports over time is also seen in other convective-storm research, such as for tornadoes as described by Antonescu et al. (2017), and may be a reflection in scientific interest in severe convective storms, or due to economic or political changes.




176

177 **Figure 1. Time series of annual numbers of large-hail reports (orange line) and large-hail days (green**


178 **line) across Europe 1900–2020.**

179 To show these data in a slightly different way, a scatterplot is created of the number of hail days versus
 180 number of hail reports for each year in the dataset, with different colors for the period before and after 2000 (Fig.
 181 2). The dataset from 1900 onwards suggests a positive linear relationship between large-hail reports and large-
 182 hail days; however, the spread is sometimes large. The high number of large-hail reports during 1949–1955
 183 (mostly from Poland, section 8) and early 1950s all congregate in one region of the graph and 2010–2020 also
 184 congregate in one region. As fewer reports are needed for a greater quantity of large-hail days, either areal extent
 185 of spotters has improved, the number of reporters has decreased in hail-prone regions, or the ESWD maintenance
 186 team have improved their ability to detect reports linked to the same event, and hence have removed duplicate
 187 events from the dataset. Thus, the 1950s are a time when reports mostly came from Poland (section 8) and captured
 188 a large number of large-hail days, indicating that certain periods of time can be fruitful for hail research using the
 189 ESWD. The spatial distribution of these reports is discussed in section 7.

190
191




192
 193 **Figure 2. Scatterplot of the annual number of large-hail days versus annual number of large-hail reports**
 194 **across Europe: 1900–2000 (green dots) and 2000–2020 (orange dots), with corresponding linear regression**
 195 **lines. These quantities are not divided by the number of years because of the incomplete data for the year**
 196 **2020.**



Deleted:

198 The average monthly distribution of the number of large-hail reports and large-hail days from 2000 to 2020
 199 is plotted in Fig. 3. The warm-season months of May, June, and July have the highest number of large-hail reports,
 200 and the cool-season months from October to March have the lowest. Whereas the month with the highest number
 201 of large-hail reports is June, the month with the highest number of large-hail days is July. Figure 3 can be compared
 202 to Půček et al. (2019, their Fig. 4) who break down the annual cycle into the frequency of reports for the continental
 203 regions of Europe north of 46°N and the more Mediterranean-influenced regions south of 46°N. Despite these
 204 differences, these two distributions look similar, with the added information coming from the distribution of large-
 205 hail days in the present study. The distribution of large-hail days in Fig. 3 is more similar to the shape of the
 206 distribution of north of 46°N in Půček et al. (2019, their Fig. 4), meaning that fewer reports occur later in the
 207 season although the number of large-hail days remains relatively high. These distributions are also similar to
 208 those from Kunz et al. (2020, their Fig. 2a) for hailstorms in central Europe using radar-derived hail streaks
 209 combined with all quality levels from the ESWD, indicating that this larger dataset of QC0+ events derived using
 210 different methods is a reliable source of large-hail data.

Deleted: larger



213
 214 **Figure 3. Combined line graph and bar chart of the total monthly numbers of large-hail reports (orange**
 215 **line) and large-hail days (green bars) across Europe: 2000–2020. These quantities are not divided by the**
 216 **number of years because of the incomplete data for the year 2020.**

Deleted: Histogram

219 The percentage of hail days reported by month per country for the period 2000–2020 was investigated in Fig.
220 4. Greece is the only country to not have over 50% of its reports being within the months of May, June, and July,
221 having a more consistent number of hail days throughout the year. Many countries do not have any reports before
222 April or after September. Spain, Italy, France, and Croatia have similar distributions of hail days throughout the
223 year, which may be linked to their Mediterranean setting, although Slovenia, Bosnia and Herzegovina, and
224 Bulgaria do not share the same characteristics, despite also being situated along the Mediterranean. Previous
225 studies such as Tzarek et al. (2020) have investigated hail distribution in Europe by linking events to
226 meteorological and climatological factors, which may help explain some of the differences seen in Fig. 4.
227 Furthermore, Sanchez et al. (2017) investigated hail events in southern Europe, concluding that even small
228 geographical and climatological differences can have a large impact on the number of hail days reported, but also
229 with the peak month of hailfall, which may also explain some of the differences in Fig. 4.

Moved (insertion) [2]

Deleted: also

Formatted: Justified, Indent: First line: 0.25", Line spacing: 1.5 lines

Deleted: f

Deleted: ure

Deleted: (???)

Deleted: ,

Deleted: appear to

Deleted: f

Deleted: ure

Deleted: (?)


Deleted: seen here

Deleted: f

Deleted: ure

Deleted: (?)

Deleted: ¶



244
245
246


Figure 4. Horizontal bar charts of the monthly distribution for countries with 100 or more reports: 2000–2020.

Commented [DMS1]: Confirm this is correct or revise.
Formatted: Justified, Space Before: 0 pt, Tab stops: 0.25", Left

247 The average diurnal cycle for the number of large-hail reports between 2000 and 2020 is shown in Fig. 5.
 248 The hour 1500–1559 UTC (labelled 1500 UTC) was the most common time for large hail to be reported with a
 249 gentle rise and a slightly more rapid decline. When corrected for local time (LT), this peak shifts to 1700–1759
 250 LT because most of Europe is east of the Prime Meridian. Figure 5 can be compared to Půček et al. (2019, their
 251 Fig. 5), who also found a peak during the 1500-UTC hour. These distributions are also similar to those from Kunz
 252 et al. (2020, their Fig. 2b) who found a peak during 1500–1800 LT for hailstorms in central Europe using all
 253 quality levels from the ESWD. Thus, the QC0+ data over a longer period of time used in this study produces a
 254 similar climatology and is consistent with previously published research using a shorter period and more selective
 255 quality-control levels, indicating that this larger dataset is a reliable source of large-hail data.

Deleted: 4

Deleted: 4




258
 259 **Figure 5.** Distribution of the hourly time of large-hail reports across Europe in UTC (green line) and local
 260 time (orange line): 2000–2020. Reports are associated with the starting hour (i.e., a report at 1515 UTC
 261 would be placed in the 1500-UTC bin).

Deleted: 4

265 To examine the year-by-year consistency of the diurnal cycle, the distribution of large-hail reports as a
 266 function of local time for each year during the period 2000–2020 is plotted in Fig. 6. Each year mostly reproduces
 267 the diurnal cycle seen in Fig. 5. The exception is some years, particularly early during this period, that have
 268 unusual peaks at 1000–1200 UTC. These reports are associated with hail events in the early part of the database
 269 that occurred at an unknown time during the night or day and were placed in 0000 UTC or 1200 UTC, respectively
 270 (Půčik et al. 2019, p. 3906). However, by 2010, the diurnal distributions seemed to have settled down to look like
 271 that in Fig. 5. The consistency after 2010 suggests the possibility that the dataset becomes more consistent in
 272 reporting events and could represent a stable period for documenting the present large-hail climate of Europe.

273
 274



275

276 **Figure 6. Hourly percentage of large hail in local time across Europe in local time for each year 2000 to**
 277 **2020.**

Deleted: 5

Deleted: 4


Deleted: 4

Deleted: 5

282 **4 Intensity of large hail: 2000–2020**

283 It is not just the frequency of events that determines their impact on society, but also the intensity of the
284 events, here represented by the maximum diameter of the hail associated with each report. Maximum hail size
285 can be difficult to measure for several reasons as highlighted by Pilorz (2015). For example, as hail is often
286 irregular in shape, the maximum diameter is actually the longest axis of the stone. Therefore, if a stone were more
287 spherical, then its maximum diameter would be smaller than an oblate stone, even though it would have a larger
288 volume. Furthermore, there is always the possibility that the largest hailstone from any given event has not been
289 found or that it has partially melted before discovery.

290 For the 28,225 large-hail reports in the present study between 2000 and 2020, 18,132 (64%) had data for the
291 maximum diameter. These reports were organized into 1-cm bins, ranging from 2.0–2.9 cm to 10+ cm. Frequency
292 of hail reports decreased with increasing hail size (Fig. 7). The maximum hail size in the database from 2000 to
293 2020 was 15 cm and was reported in Romania on 26 May 2016. This report was rated QC1, so has been confirmed.
294 The second largest hail size was 14.1 cm and was reported in Germany on 6 August 2013. This particular hailstone
295 set the record for the largest hailstone in Germany (ESKP 2013). This report is recorded as QC2 and includes
296 additional information in the ESWD database, such as the average hailstone size being 8 cm.
297



298 **Figure 7. Bar chart of the number of large-hail reports across Europe by maximum diameter in 1-cm bins:**
299 **2000–2020.**

Deleted: an
Deleted: its

Deleted: 6
Deleted: plausibly check


Deleted: 6
Deleted: Histogram

307 To investigate the distribution of large-hail size over time, Fig. 8 presents the percentage of each hail-size bin
 308 per year from 2000 to 2020. During this 21-yr period, the percentage of each bin size does not change dramatically.
 309 This distribution is similar to the 1989–2018 average from Púčik et al. (2019, their Fig. 7), with about 40% of
 310 large-hail reports being smaller than 3 cm, about 70% being smaller than 4 cm, and about 84% being smaller than
 311 5 cm. Therefore, the large-hail size distribution during 2000–2020 may represent a period of stability in reporting
 312 with little detectable change in large-hail size distributions in the ESWD dataset. For determining the present
 313 large-hail climate, the stability in the large-hail size distribution after 2000 represents a slightly longer period of
 314 record compared to that of the diurnal cycle, which stabilized after 2010 (Fig. 6).

Deleted: 7

Deleted: 5

315



316


317 **Figure 8.** Time series of bar charts of the annual distributions of large-hail size across Europe in 1-cm
 318 diameter bins: 2000–2020.

Deleted: 7

322 The ESWD has information on average hail size, although only 12% (2237 out of 18,132) of reports contain
 323 this information for 2000–2020. There is, however, a strong positive linear relationship between the average and
 324 maximum hail size recorded (Fig. 9). There were two outliers that are most likely data-entry errors, such as events
 325 with a 2-cm maximum size and 5-cm or 3-cm average size. Both were QC1. The linear relationship ($R^2 = 0.76$)
 326 between maximum and average hail size suggests that the average hail size is about 60% of the maximum hail
 327 size, although there is considerable spread around this line.

Deleted: 8


Deleted: that the entries



328

329

330 **Figure 9. Scatterplot representing 2237 hail reports of the maximum large-hail size versus average large-**
 331 **hail size across Europe during 2000–2020, with corresponding linear regression line (green dotted line).**
 332 **The 1:1 line is plotted as a blue line. Two pink dots represent likely data-entry errors where the average**
 333 **diameter is greater than the maximum diameter.**



Deleted:


Deleted: 8

Deleted: red

339 **5 Time accuracy of reports: 2000–2020**

340 The ESWD includes a quantity called time accuracy, defined as the time interval over which the report could
341 have occurred. For example, a time accuracy of 5 min would mean that the large hail fell within 2.5 min on either
342 side of the time recorded in the ESWD. Groenemeijer and Liang (2020) specify ten categories of time accuracy:
343 1 min, 5 min, 15 min, 30 min, 1 h, 3 h, 6 h, 12 h, 1 day, and greater than 1 day. The time accuracy of large hail
344 in the ESWD has improved over time, with over 50% of reports having a time accuracy of 30 min by 2012,
345 followed by 50% having a time accuracy of 15 min by 2017 (Fig. 10). Moreover, between 2009 and 2010, reports
346 with a time accuracy of 30 min became more common, replacing some of the reports with time accuracy of 1 h,
347 and time accuracy of 12 h and greater become negligible. Viewing the ESWD from 2000–2020 as a whole, these
348 improvements in time accuracy means that the ESWD is becoming a more reliable source of data, with more
349 highly temporally resolved data on hail occurrence.
350

Deleted: 9




351

352 **Figure 10. Time series of bar charts of the annual distributions of time accuracy of reports across Europe:**
353 **2000–2020.**

Deleted: 9

356 On the scale of individual countries, however, work remains to improve the quality of the ESWD. The
 357 average time accuracy for each country with 100 or more reports during 2000–2020 is shown in Fig. 11. The
 358 distribution of time accuracy varies considerably among these 24 countries. Germany, Finland, and the Czech
 359 Republic have more than 40% of their reports with time accuracy of 5 min, whereas Bulgaria, Russian Federation,
 360 and Moldova have the lowest (1% or less). Figure 11 also indicates the countries for which there is opportunity
 361 to improve engagement in severe-weather reporting.

Deleted: 0
 Deleted: from
 Deleted: 0
 Deleted: perhaps requiring greater support and mentorship from ESSL and other organizations
 Deleted:



362
 363 Figure 11. Horizontal bar charts of the time accuracy for countries with 100 or more reports: 2000–2020.

Deleted: 0

371 **6 Spatial distribution by country: 2000–2020**

372 Hail reports across Europe are heterogenous, not just in time, but also in space. Countries such as Germany,
373 Russian Federation, and Italy reported 4956, 4182, and 2447 large-hail events between 2000 and 2020, compared
374 to others such as Switzerland, the UK, and Denmark only reporting 266, 85 and 31 cases, respectively (Table 1).
375 Central and western European countries reported more large hail with 5 out of the top 10 countries located there
376 (Table 1). Germany has more large-hail reports than the Russian Federation for fewer large-hail days, similarly to
377 Poland having more reports than Italy, and Austria more reports than Greece.


378 Besides meteorological reasons for the variability, other reasons that may explain these reporting differences
379 include the existence, size, and enthusiasm of spotter networks within each country; variations in the ability or
380 enthusiasm of citizens to input into the ESWD; and the availability of information to quality-control reports. In
381 fact, many central European countries have larger and more enthusiastic spotter networks [e.g., Poland, as
382 discussed in Pacey et al. (2021) and section 7 of the present article] and are more likely to enter their reports into
383 the ESWD. Population density and area of the country were considered as possible explanations for the number
384 of hail reports varying by country, although neither had a statistically significant relationship with the number of
385 hail reports (not shown). As with the time-accuracy data (section 5), greater engagement with some countries to
386 encourage entering their reports into the ESWD would lead to a larger and more complete dataset.

387 **Table 1. Number of large-hail days and large-hail reports by country: 2000–2020.**

| Country | Number of large-hail reports | Number of large-hail days |
|------------------------|------------------------------|---------------------------|
| Germany | 4956 | 692 |
| Russian Federation | 4182 | 1012 |
| Poland | 3226 | 471 |
| Italy | 2447 | 555 |
| France | 1707 | 440 |
| Austria | 1502 | 353 |
| Spain | 1027 | 295 |
| Ukraine | 1021 | 319 |
| Romania | 983 | 267 |
| Greece | 975 | 395 |
| Hungary | 903 | 226 |
| Bulgaria | 820 | 238 |
| Serbia and Kosovo | 490 | 146 |
| Czech Republic | 490 | 174 |
| Moldova | 451 | 117 |
| Croatia | 399 | 181 |
| Finland | 382 | 139 |
| Slovenia | 332 | 116 |
| Switzerland | 266 | 87 |
| Belarus | 261 | 103 |
| Slovakia | 234 | 104 |
| Bosnia and Herzegovina | 169 | 65 |
| Netherlands | 165 | 76 |
| Belgium | 121 | 49 |
| Latvia | 86 | 50 |
| United Kingdom | 85 | 41 |
| Estonia | 79 | 38 |
| Portugal | 77 | 34 |
| Sweden | 74 | 50 |
| Cyprus | 68 | 45 |
| Lithuania | 42 | 23 |
| Luxembourg | 39 | 6 |
| Denmark | 31 | 18 |
| Albania | 22 | 12 |
| Montenegro | 21 | 3 |
| North Macedonia | 21 | 13 |
| Norway | 21 | 15 |
| Malta | 11 | 9 |
| Andorra | 6 | 4 |
| Iceland | 4 | 4 |
| Ireland | 2 | 2 |

388

389 Similar to Fig. 2 where the number of large-hail reports was plotted versus the number of large-hail days by
 390 year, Fig. 12 shows a scatterplot between the number of large-hail reports versus the number of large-hail days by
 391 country from Table 1. There is a positive linear relationship ($R^2 = 0.88$) between large-hail reports and large-hail
 392 days by country (Fig. 12), suggesting that large-hail reports are proportional to large-hail days. This relationship
 393 would therefore imply that reporting frequency is similar across all hail frequencies and countries, except for
 394 Germany and Poland which have a much greater number of reports proportional to the number of days.




395
 396 **Figure 12. Scatterplot of the total number of large-hail reports versus large-hail days by country: 2000–**
 397 **2020.**

398 We further investigated the hail size distribution by country for the period 2000–2020 (Fig. 13). Only one
 399 report of each size diameter was taken per country per day to minimize some of the reporting biases. Finland has
 400 the greatest proportion of the lowest hail bin size, whereas Slovenia has the lowest. For sizes 5 cm in diameter
 401 and greater, the proportion of hail sizes recorded starts to diminish drastically, which would be expected as larger
 402 hailstones are rarer. Although Slovenia has the greatest proportion of hail sizes above 5 cm, these reports came
 403 from a sample of 116 hail reports, one of the smallest of the countries analyzed. For hail days with a report above
 404 10 cm, Russia has the greatest quantity with 10 reports over this period, whereas Italy came second with 9 reports
 405 and France with 8. Slovenia, although having a greater proportion, had 5 days with a hail report above 10 cm for
 406 this period.
 407

Deleted: 1
 Deleted: 1
 Deleted: implies
 Deleted: s

Deleted: ¶ ... [1]
 Deleted: 1
 Deleted: ure
 Deleted: ?
 Formatted: Indent: First line: 0.25", Line spacing: 1.5 lines
 Deleted: ile
 Deleted: Above
 Deleted: once more
 Deleted: but was only taken from
 Deleted: se
 Deleted: In terms of
 Deleted: ile
 Deleted: recorded



424

425 **Figure 13. Horizontal bar charts of the size distribution of large hail for countries with 100 or more**
 426 **reports: 2000–2020.**

- Formatted: Caption
- Commented [DMS2]: Is this correct? Revise accordingly.
- Deleted: ¶
- Formatted: Font: 10 pt, Bold, Kern at 14 pt

428 **7 Poland: 1900–2020**

429 As noted in association with Fig. 1, nearly all large-hail reports and large-hail days during the 1930s and
430 1940s–1950s originated in Poland (Figs. 14a,b). Very few hail days were recorded between 1956 and 2000, before
431 the general increase along with the rest of Europe for the last 20 years (Fig. 14). There appears to be far fewer
432 large-hail days over the past 20 years in Poland (30–40 days a year) compared to the 1940s–1950s (100–120 days
433 a year). With an overall increase in reporting numbers and accuracy, it would be unlikely that the current Polish
434 reports are missing many events, and therefore the difference in annual numbers of large-hail days seems unlikely.

435 The addition of this data in the ESWD was due to Igor Laskowski who reports:

436 “those reports were based on annual records collected by a Polish National Institute of
437 Meteorology founded in 1919, now Institute of Meteorology and Hydrology - National
438 Research Institute (<https://imgw.pl/instytut/historia>). The data was collected via hail
439 questionnaires, which provided information on the size of the hail (vetch-sized, pea-sized, broad
440 bean-sized, hazelnut-sized, walnut-sized, pigeon egg-sized, hen egg-sized and goose egg-sized)
441 and also details about time of its occurrence, storm direction and the size of the expected yield
442 decrease (in percent). The questionnaires were filled in both by agricultural correspondents of
443 the Polish Central Statistical Office (whose number was growing larger, especially in the
444 [19]50s) and existing insurance companies which provided hail insurance at this time. Those
445 records also contain observations of hail reported by observers at meteorological stations.”

446 At the time of this study, data from yearbooks from 1930–1937 and 1946–1955 had been added.

447 Suwała (2011) investigated Polish hail based on data from 23 meteorological stations recorded in the
448 Meteorological Yearbooks published by the Institute of Meteorology and Water Management for the years 1973–
449 1980 and the Polish National Climatic Data Centre for the years 1981–2009. They found that over the 37-year
450 period, March was the month with the highest hail frequency across the country, followed by February and
451 January. For individual stations, December and January recorded the highest hailfall, with the two stations along
452 the Baltic coasts having a mean of 8 days. Although these results may indicate a cool-season preference for hail,
453 there is the possibility that ice pellets or graupel might have been classified as hail (e.g., Punge and Kunz 2018).
454 Overall, the Baltic coast showed the highest annual mean, whereas central Poland showed the lowest. This result
455 contradicts the findings of Pilorz (2015) who investigated large hail in Poland for 2007–2015, concluding that
456 southeast Poland had the greatest number of storms and associated large hail events.

457 Furthermore, the warm months of June to September had the lowest mean hail frequency for all stations. This
458 contradicts the results found in this present study and those by Púčík et al. (2019), that hail is most frequent in the
459 warm season, but also contradicts those by Taszarek and Suwała (2014) who investigated large hail in Poland in
460 2012. In addition, there appeared to be some cyclicity in frequency over the 37-yr period, although this cyclicity
461 varied greatly when investigating individual stations, and no trends were observed. These results may explain why
462 Poland possesses a different annual distribution to other locations.

463 Suwała (2011) mentioned previous hail studies in Poland, such Schmuck (1949), Koźmiński (1964), and
464 Zinkiewicz and Michna (1955), which may offer an explanation on the high number of hail reports during the
465 1930s and 1950s. Unfortunately, these are not currently available to read. Access to these historical studies may
466 help explain the quantity of Polish entries in the ESWD during the 1930s, 1940s, and 1950s. Moreover, an effort
467 to retrieve and input the data from 1973 to 2009 into the ESWD would greatly help with the homogeneity of the

Moved up [2]: The percentage of hail days reported by month per country for the period 2000–2020 was also investigated in figure (??). Greece is the only country to not have over 50% of its reports being within the months of May, June, and July, having a more consistent number of hail days throughout the year. Many countries do not have any reports before April, or after September. Spain, Italy, France, and Croatia appear to have similar distributions of hail days throughout the year, which may be linked to their Mediterranean setting, although Slovenia, Bosnia and Herzegovina, and Bulgaria do not share the same characteristics, despite also being situated along the Mediterranean. Previous studies such as Tazarek have investigated hail distribution in Europe by linking events to meteorological and climatological factors, which may help explain some of the differences seen in figure (?). Furthermore, Sanchez et al. (2017) investigated hail events in southern Europe, concluding that even small geographical and climatological differences can have a large impact on the number of hail days reported, but also with the peak month of hailfall, which may also explain some of the differences seen here in figure (?).

Deleted: Figure (?) shows the proportion of hail days per country from 2000–2020, which shows that there is much variation year to year. However, much of this variation is likely due to an increased involvement and establishment of the ESWD as a reporting vessel, and not a depiction of any changing trends in hail days. Belarus and Bosnia and Herzegovinian were the last to start entering data into the ESWD, with the first hail days reported being in 2006, when the database was founded. Germany appears to have the most consistent reporting over this period, closely followed by the Netherlands and Russia, with similar proportions reported each year. Furthermore, this figure helps to picture the change in report patterns on a country by country basis, from which figure 1 showed a more stable time period. [2]

Deleted: 2

Deleted: 2

Formatted: Indent: Left: 0.5", Right: 0.52"

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt


Formatted: Font: (Default) Times New Roman, 10 pt

Deleted: amount


Deleted: ,

Deleted: s

509 Polish dataset. There remains the possibility that this data does not exist as the country suffered major economic
 510 difficulties during this period.



511



512

513 **Figure 14.** Time series of annual numbers of (a) large-hail reports for Europe (green line) and Poland (red
 514 line), and (b) large-hail days for Europe (green line) and Poland (red line): 1900–2020.

Deleted: 2

516 As in Fig. 10, the time accuracy of large-hail reports can be plotted as a function of time during 1930–2020
 517 in Fig. 15. The time accuracy of reporting in Poland has improved over the past 20 years, with over half the
 518 reports having a time accuracy of 15 min by 2015 (Fig. 15). During the 1930s and 1950s, the time accuracy was
 519 much lower, around 3 h (Fig. 15). Although this result may suggest that reports were less reliable during this
 520 period, the consistency in time accuracy (especially during the 1950s) may also suggest that the data-collection
 521 methods were more consistent. These reports were later found to be based upon the Meteorological Yearbooks
 522 from the Polish National Institute of Meteorology (I. Laskowski 2022, personal communication). The yearbooks
 523 contained information on hail size, time of occurrence and storm direction based upon questionnaires posed to
 524 insurance companies, agricultural correspondents of the Polish Central Statistical Office alongside observations
 525 from meteorological stations. Laskowski also mentioned that yearbooks from the 1960s and 1970s also existed,
 526 but was currently unable to find any existing copies. Hence, such data – when it is found – remains to be entered
 527 into the ESWD.

528 In addition, the reported location accuracy was also investigated, with the most common distances being 1
 529 and 3 km, similar to those found in the broader 2000–2020 dataset. This result reiterates the importance of these
 530 earlier reports in constructing a reliable hail climatology, and gives credit to the data-collection method.


531 The historical Polish datasets offer an insight into past hail frequency and reporting accuracies. Results by
 532 Suwała (2011) for the period 1973–2009 contradict those found for more recent time periods in terms of peak
 533 annual frequency and spatial distribution of large hail. The potential implications of these discrepancies may
 534 suggest that distributions of hail size, frequency, and location have changed over time and have not yet been
 535 established or studied due to the lack of historical pan-European data, highlighting the importance of building the
 536 ESWD further. Moreover, the existence of Meteorological Yearbooks in Poland could also suggest that other
 537 nations might hold similar records that remain to be analyzed and could contribute toward building a more
 538 complete climatology.
 539

Deleted: 9

Deleted: 3

Deleted: 3

Deleted: 3



Deleted: hail

Deleted: trends

540
 541 **Figure 15. Time series of bar charts the annual distributions of time accuracy of reports for Poland: 1930–**
 542 **2020.**

Deleted: 3

550 **8 Comparison to previous hail climatologies and prospects for a baseline for climate-change research**

551 The ultimate goal of severe-storm climatologies is to create a consistent and complete database in space and
552 time. Consistent data acquisition methods throughout the study area and through time would assist in achieving
553 this goal; however, consistency is not achievable across Europe. Punge and Kunz (2016) synthesized all European
554 hail studies in their review, not just large hail. They concluded that not all regions have the same threat of hail,
555 and efforts to report and record these events vary by country. They further concluded that there was insufficient
556 evidence to determine any trends in hail events, both in terms of spatial and temporal extent, highlighting the need
557 for the continuation of the ESWD to form a reliable climatology. Previous studies have provided pan-European
558 climatologies of hail based using other methods such as Punge et al. (2014, 2017), who used overshooting cloud
559 tops, Rädler et al. (2018) who used reanalysis data, or Taszarek et al. (2018) who used a combination of data
560 sources. Some studies are projecting increases in hailstorms with climate change in Italy (Piani et al. 2005),
561 Netherlands (Botzen et al. 2010), and Germany (Mohr et al. 2015), as well as across much of Europe (Taszarek
562 et al. 2021). Other studies have also concluded that there were no positive trends in the frequency of hail in hailpad
563 data in northern Italy and France (e.g., Eccel et al. 2012; Dessens et al. 2015; Raupach et al. 2021; Manzato et al.
564 2023). Tazarek et al. (2019) argue that a combination of datasets is important to construct a robust climatology,
565 particularly as the spatial and temporal resolutions would often differ between methods. Furthermore, studies such
566 as Rädler et al. (2018) compared their reanalysis results to surface observed reports from the ESWD to strengthen
567 their arguments. Therefore, understanding the characteristics of the current surface observations via the ESWD
568 helps not only build a climatology of large hail in Europe, but can also be used in association with other research
569 methods to identify the underlying factors which lead to such events.

570 Examining the evidence presented in the present article, we seek a stable time period during 2000–2020.
571 Based on the number of large-hail reports, no stable time period exists (Fig. 1). Based on the number of large-
572 hail days, the time period starts around 2012 (Fig. 1). Based on the diurnal cycle of large-hail reports, the time
573 period starts around 2010 (Fig. 6). Based on the large-hail size distributions, the time period starts around 2004
574 (Fig. 8). Based on the time accuracy of reports, the time period possibly starts around 2018 (Fig. 10). However,
575 if one is prepared to accept an accuracy of 3 h or less, then the time period starts around 2010 (Fig. 10).

576
577 **9 Conclusion**

578 The ESWD provides the only pan-European dataset for large-hail reports. The frequency of reports is sporadic
579 pre-2000, and hence the focus of this study is for the period 2000 to 2020. Hail reports have continuously increased
580 since 2000. The annual number of large-hail days have remained steady after 2010 at around 175 days per year,
581 although some interannual variability is still observed. Increased large-hail reports for similar large-hail days
582 suggests that a greater spotter network is in operation, and that the engagement with the ESWD is increasing.

583 The warm season of May to August shows the highest number of large-hail reports and large-hail days, with
584 June showing the highest large-hail reports and July the highest large-hail days. The number of large-hail reports
585 decrease faster than large-hail days from June to September. The diurnal cycle shows that the peak hailfall time
586 is 1500 UTC and 1700 local time.

587 The number of large-hail reports decreases with increasing diameter, and the percentage distribution of each
588 large-hail size by year does not appear to have changed over the past 20 years. The possibility that hail-size
589 distribution is changing remains, as smaller, less damaging hail size events are being recorded more regularly.

Deleted:

Deleted: and Punge et al. 2017

Deleted: based theirs on

Deleted: with climate change

Deleted: in order

Deleted: v

Deleted: Given that some studies are projecting increases in hailstorms in Italy (Piani et al. 2005), Netherlands (Botzen et al. 2010), and Germany (Mohr et al. 2015), as well as across much of Europe (Taszarek et al. 2021) with climate change, creating a robust climatology with a consistent record is needed.

Deleted: 5

Deleted: 7

Deleted: 9

Deleted: 9

606 The diurnal cycle by year shows that for the past 10 years, a consistent pattern has emerged, with a rise in the
607 early afternoon and a decline in the evening. Furthermore, the time accuracy of reports has improved with over
608 50% of reports being reported to within a 30-minute window by 2012, followed by 50% being reported to within
609 a 15-minute window by 2017. Not all countries display improved time accuracies. Germany, Finland, and the
610 Czech Republic have the greatest proportions of 5-minute time-accuracy reports, whereas Russia, Moldova, and
611 Bulgaria have the highest proportions of 1-h or greater time-accuracy reports. Efforts to improve monitoring and
612 reporting in these regions is therefore suggested to improve the completeness of the ESWD.

613 Poland possessed anomalously large numbers of large-hail reports during the 1930s, 1940s, and 1950s. The
614 reason is linked to scientific interest in severe convective storms during these periods alongside a nationwide
615 effort by the Polish National Institute of Meteorology to record hail events via questionnaires. Yearbooks also
616 exist for the 1960s and 1970s; however, copies are yet to be retrieved and entered into the database.

617 Even though the dataset remains too short to extract any trends in large-hail pattern distribution, the
618 climatology presented here provides insight into which countries and geographical regions to target for
619 improvements in data acquisition. This climatology also helps advance the idea that some time series are starting
620 to show consistent behavior, suggesting their utility as climate-change baselines. Furthermore, the differences in
621 both spatial and annual frequencies of hail in Poland over different time periods may suggest that hail trends have
622 been changing, highlighting the importance of building and maintaining such climatologies. Therefore, the
623 usefulness of the ESWD will only continue to expand and offer avenues for future severe convective storm
624 research.

625
626 *Data availability.* The data were obtained from the European Severe Storms Laboratory European Severe Weather
627 Database, in accordance with their data policies: <http://www.eswd.eu>.

628
629 *Author contributions.* FH performed the analyses and wrote the paper. DMS supervised the research, and helped
630 write and edit the paper.

631
632 *Competing interests.* The authors declare that they have no conflicts of interest.

633
634 *Acknowledgments.* This article is derived from FH's BSc dissertation at the University of Manchester. We thank
635 the European Severe Storms Laboratory for their kind access to the ESWD that made this work available. We
636 thank Neil Mitchell for his comments on the dissertation. We thank Igor Laskowski for explaining the Polish hail
637 dataset. We thank the anonymous reviewers for their comments that have improved this article.

638
639 *Financial support.* Schultz is partially supported by the Natural Environment Research Council grant numbers
640 NE/N003918/1 and NE/W000997/1.

641 642 **References**

643
644 Allen, J. T., and M. K. Tippett: The characteristics of United States hail reports: 1955–2014. *Electron. J. Severe*
645 *Storms Meteor.*, **10** (3), <http://www.ejssm.org/ojs/index.php/ejssm/article/viewArticle/149>, 2015.

Deleted: displayed

Deleted: quantities

Deleted: temporally incomplete

649 Antonescu, B., D. M. Schultz, A. Holzer, and P. Groenemeijer: Tornadoes in Europe: An underestimated
650 threat. *Bull. Amer. Meteorol. Soc.*, **98**, 713–728, 2017.

651 Associated Press, The: Storm floods German vaccine center, 5 injured by heavy hail. ABC News. Available at:
652 [https://abcnews.go.com/International/wireStory/german-vaccine-center-flooded-injured-heavy-hail-](https://abcnews.go.com/International/wireStory/german-vaccine-center-flooded-injured-heavy-hail-78459538)
653 [78459538](https://abcnews.go.com/International/wireStory/german-vaccine-center-flooded-injured-heavy-hail-78459538) (Accessed: 16 July 2021), 2021.

654 Brooks, H. E., Doswell, C.A. III, X. Zhang, A. A. Chernokulsky, E. Tochimoto, B. Hanstrum, E. de Lima
655 Nascimento, D. M. Sills, B. Antonescu, and B. Barrett: A century of progress in severe convective storm
656 research and forecasting. *Meteorological Monographs*, **59**, 18.1–18.41, 2019.

657 Botzen, W. J. W., L. M. Bouwer, and J. C. J. M. Van den Bergh: Climate change and hailstorm damage: Empirical
658 evidence and implications for agriculture and insurance. *Resource and Energy Economics*, **32**, 341–362,
659 2010.

660 *Dessens, J., C. Berthet, and J. L. Sanchez: Change in hailstone size distributions with an increase in the melting*
661 *level height. Atmos. Res.*, **158–159**, 245–253, 2015.

662 Dotzek, N., P. Groenemeijer, B. Feuerstein, and A. M. Holzer: Overview of ESSL's severe convective storms
663 research using the European Severe Weather Database ESWD. *Atmos. Res.*, **93**, 575–586, doi:
664 10.1016/j.atmosres.2008.10.020, 2009.

665 *Eccel, E., P. Cau, K. Riemann-Campe, and F. Biasioli: Quantitative hail monitoring in an alpine area: 35-year*
666 *climatology and links with atmospheric variables. Intl. J. Climatol.*, **32**, 503–517, 2012.

667 ESKP: <https://www.eskp.de/en/natural-hazards/largest-hailstone-found-over-swabian-jura-935341/>. (Accessed
668 25 March 2022), 2013.

669 Fluck, E., M. Kunz, P. Geissbuehler, and S. P. Ritz: Radar-based assessment of hail frequency in Europe. *Natural*
670 *Hazards and Earth System Sciences*, **21**, 683–701, 2021.

671 Groenemeijer, P., M. Kuehne, Z. Liang, and N. Dotzek: New capabilities of the European Severe Weather
672 Database. *5th European Conference on Severe Storms*, Landshut, Germany, 311–312, 2009.

673 Groenemeijer, P., and T. Kühne: A climatology of tornadoes in Europe: Results from the European Severe
674 Weather Database. *Mon. Wea. Rev.*, **142**, 4775–4790, 2014.

675 Groenemeijer, P., T. Púčík, A. M. Holzer, B. Antonescu, K. Riemann-Campe, D. M. Schultz, T. Kühne, B.
676 Feuerstein, H. E. Brooks, C. A. Doswell, H.-J. Koppert, and R. Sausen: Severe convective storms in Europe:
677 Ten years of research and education at the European Severe Storms Laboratory. *Bull. Amer. Meteor. Soc.*, **98**,
678 2641–2651, doi: 10.1175/BAMS-D-16-0067.1, 2017.

679 Groenemeijer, P., and Z. Liang: ESWD data format specification Version 1.60 and 1.60-csv. Revision 1. Tech.
680 Report 2020-11, European Severe Storms Laboratory, 66 pp. Available at: [https://www.essl.org/cms/wp-](https://www.essl.org/cms/wp-content/uploads/ESSL-rep-2020-11.pdf)
681 [content/uploads/ESSL-rep-2020-11.pdf](https://www.essl.org/cms/wp-content/uploads/ESSL-rep-2020-11.pdf), 2020.

682 Hand, W. H., and G. Cappelluti: A global hail climatology using the UK Met Office convection diagnosis
683 procedure (CDP) and model analyses. *Meteorol. Appl.*, **18**, 446–4589, 2011.

684 Hawkins, E., D. Frame, L. Harrington, M. Joshi, A. King, M. Rojas, and R. Sutton: Observed emergence of the
685 climate change signal: From the familiar to the unknown. *Geophys. Res. Lett.*, **47**, e2019GL086259, 2020.

686 Istrate, V., R. V. Dobri, F. Bărcăcianu, R. A. Ciobanu, and L. Apostol: A ten years hail climatology based on
687 ESWD hail reports in Romania, 2007-2016. *Geographia Technica*, **12**, 110–118, 2017.

Formatted: Font: Italic

Formatted: Font: Bold

Formatted: Font: Italic

Formatted: Font: Bold

688 Kaonga, G.: BBC Weather: Europe braces for hail storms as heavy downpours to cover continent, Express.co.uk.
689 Available at: [https://www.express.co.uk/news/weather/1454518/BBC-Weather-Europe-forecast-hail-](https://www.express.co.uk/news/weather/1454518/BBC-Weather-Europe-forecast-hail-storms-latest-June-2021-vn)
690 [storms-latest-June-2021-vn](https://www.express.co.uk/news/weather/1454518/BBC-Weather-Europe-forecast-hail-storms-latest-June-2021-vn) (Accessed: 16 July 2021), 2021.

691 Koźmiński, C.: Geograficzne rozmieszczenie większych burz gradowych zanotowanych na obszarze Polski w
692 latach 1946–1956. [Geographical distribution of major hailstorms recorded in Poland in 1946–1956.] *Przegl.*
693 *Geogr.*, **36** (1), 87–102, 1964.

694 Kunz, M., U. Blahak, J. Handwerker, M. Schmidberger, H. J. Punge, S. Mohr, E. Fluck, and K. M. Bedka: The
695 severe hailstorm in southwest Germany on 28 July 2013: Characteristics, impacts and meteorological
696 conditions. *Quart. J. Roy. Meteor. Soc.*, **144**, 231–250, 2018.

697 Kunz, M., J. Sander, and C. Kottmeier: Recent trends of thunderstorm and hailstorm frequency and their relation
698 to atmospheric characteristics in southwest Germany. *Intl. J. Climatol.*, **29**, 2283–2297, 2009.

699 Kunz, M., J. Wandel, E. Fluck, S. Baumstark, S. Mohr, and S. Schemm: Ambient conditions prevailing during
700 hail events in central Europe. *Natural Hazards and Earth System Sciences*, **20**, 1867–1887, 2020.

701 [Manzato, A., Fasano, G., Cicogna, A., Sioni, F., and Pucillo, A.: Trends of sounding-derived indices and](#)
702 [observations in NE Italy in the last 30 years. 11th European Conference on Severe Storms, Bucharest,](#)
703 [Romania, 8–12 May 2023, ECSS2023-3, <https://doi.org/10.5194/ecss2023-3>, 2023.](#)

704 Mohr, S., M. Kunz, and K. Keuler: Development and application of a logistic model to estimate the past and future
705 hail potential in Germany. *J. Geophys. Res. Atmos.*, **120**, 3939–3956, 2015.

706 Pacey, G. P., D. M. Schultz, and L. Garcia-Carreras: Severe convective windstorms in Europe: Climatology,
707 preconvective environments, and convective mode. *Wea. Forecasting*, **36**, 237–252, doi: [10.1175/WAF-D-](https://doi.org/10.1175/WAF-D-20-0075.1)
708 [20-0075.1](https://doi.org/10.1175/WAF-D-20-0075.1), 2021.

709 Piani, F., A. Crisci, G. De Chiara, G. Maracchi, and F. Meneguzzo: Recent trends and climatic perspectives of
710 hailstorms frequency and intensity in Tuscany and Central Italy. *Nat. Hazards Earth Syst. Sci.*, **5**, 217–224,
711 2005.

712 Pilorz, W.: Very large hail occurrence in Poland from 2007 to 2015. *Contemporary Trends in Geoscience*, **4**, 45–
713 55, 2015.

714 Püčik, T., C. Castellano, P. Groenemeijer, T. Kühne, A. T. Rädler, B. Antonescu, and E. Faust: Large hail
715 incidence and its economic and societal impacts across Europe. *Mon. Wea. Rev.*, **147**, 3901–3916, 2019.

716 Punge, H. J., K. M. Bedka, M. Kunz, and A. Werner: A new physically based stochastic event catalog for hail in
717 Europe. *Natural Hazards*, **73**, 1625–1645, 2014.

718 [Punge, H. J., K. M. Bedka, M. Kunz, and A. Reinbold: Hail frequency estimation across Europe based on a](#)
719 [combination of overshooting top detections and the ERA-INTERIM reanalysis. *Atmos. Res.*, **198**, 34–43,](#)
720 [2017.](#)

721 Punge, H. J., and M. Kunz: Hail observations and hailstorm characteristics in Europe: A review. *Atmos. Res.*, **176**,
722 159–184, 2016.

723 [Rädler, A. T., P. Groenemeijer, E. Faust, and R. Sausen: Detecting severe weather trends using an additive](#)
724 [Regressive Convective Hazard Model \(AR-CHaMo\). *J. Appl. Meteor. Climatol.*, **57**, 569–587, 2018.](#)

725 Raupach, T. H., O. Martius, J. T. Allen, M. Kunz, S. Lasher-Trapp, S. Mohr, K. L. Rasmussen, R. J. Trapp, and
726 Q. Zhang: The effects of climate change on hailstorms. *Nature Reviews Earth & Environment*, **2**, 213–226,
727 2021.

Formatted: Font: 10 pt

Formatted: Normal

Formatted: Font: 10 pt

Formatted: Justified, Indent: Left: 0", Hanging: 0.25", Line spacing: 1.5 lines

Formatted: Default Paragraph Font, Font: 10 pt

Formatted: Font: 10 pt

728 [Sanchez, J. L., A. Merino, P. Melcón, E. García-Ortega, S. Fernández-González, C. Berthet, and H. Dessens: Are](#)
729 [meteorological conditions favoring hail precipitation change in southern Europe? Analysis of the period](#)
730 [1948–2015. *Atmos. Res.*, **198**, 1–10, 2017.](#)

731 Space: Weather Armageddon hits Poland: Huge hail and a powerful tomado. Earth Chronicles News. Available
732 at: [https://earth-chronicles.com/natural-catastrophe/weather-armageddon-hits-poland-huge-hail-and-a-](https://earth-chronicles.com/natural-catastrophe/weather-armageddon-hits-poland-huge-hail-and-a-powerful-tornado.html)
733 [powerful-tornado.html](https://earth-chronicles.com/natural-catastrophe/weather-armageddon-hits-poland-huge-hail-and-a-powerful-tornado.html) (Accessed: 16 July 2021), 2021a.

734 Space: Giant hail hits highway in Italy damaging hundreds of cars. Earth Chronicles News. Available at:
735 [https://earth-chronicles.com/natural-catastrophe/giant-hail-hits-highway-in-italy-damaging-hundreds-of-](https://earth-chronicles.com/natural-catastrophe/giant-hail-hits-highway-in-italy-damaging-hundreds-of-cars.html)
736 [cars.html](https://earth-chronicles.com/natural-catastrophe/giant-hail-hits-highway-in-italy-damaging-hundreds-of-cars.html) (Accessed: 31 July 2021), 2021b.

737 Space: Russia: Hail the size of a quail’s egg fell in the Kemerovo region. Earth Chronicles News. Available at:
738 [https://earth-chronicles.com/natural-catastrophe/russia-hail-the-size-of-a-quails-egg-fell-in-the-kemerovo-](https://earth-chronicles.com/natural-catastrophe/russia-hail-the-size-of-a-quails-egg-fell-in-the-kemerovo-region.html)
739 [region.html](https://earth-chronicles.com/natural-catastrophe/russia-hail-the-size-of-a-quails-egg-fell-in-the-kemerovo-region.html) (Accessed: 31 July 2021), 2021c.

740 Schmuck, A.: Burze gradowe. *Czas. Geogr.*, **20**, 260–267, 1949.

741 Suwała, K.: Hail occurrence in Poland. *Quaestiones Geographicae*, **30**, 115–126, 2011.

742 Suwała, K., and E. Bednorz: Climatology of hail in Central Europe. *Quaestiones Geographicae*, **32**, 99–110, 2013.

743 Tang, B. H., V. A. Gensini, and C. R. Homeyer: Trends in United States large hail environments and
744 observations. *npj Climate and Atmospheric Science*, **2**, 1–7, 2019.

745 [Taszarek, M., H. E. Brooks, B. Czernecki, P. Szuster, and K. Fortuniak: Climatological aspects of convective](#)
746 [parameters over Europe: A comparison of ERA-Interim and sounding data. *J. Climate*, **31**, 4281–4308, 2018.](#)

747 Taszarek, M., J. Allen, T. Púčik, P. Groenemeijer, B. Czernecki, L. Kolendowicz, K. Lagouvardos, V. Kotroni,
748 and W. Schulz: A climatology of thunderstorms across Europe from a synthesis of multiple data sources. *J.*
749 *Climate*, **32**, 1813–1837, 2019.

750 Taszarek, M., J. T. Allen, P. Groenemeijer, R. Edwards, H. E. Brooks, V. Chmielewski, and S. Enno: Severe
751 convective storms across Europe and the United States. Part I: Climatology of lightning, large hail, severe
752 wind, and tornadoes. *J. Climate*, **33**, 10239–10261, 2020.

753 Taszarek, M., J. T. Allen, H. E. Brooks, N. Pilguy, and B. Czernecki: Differing trends in United States and
754 European severe thunderstorm environments in a warming climate. *Bull. Amer. Meteor. Soc.*, **102**, E296–
755 E322, 2021.

756 Tuovinen, J. P., A. J. Punkka, J. Rauhala, H. Hohti, and D. M. Schultz: Climatology of severe hail in Finland:
757 1930–2006. *Mon. Wea. Rev.*, **137**, 2238–2249, 2009.

758 Wilhelm, J., S. Mohr, H. J. Punge, B. Mühr, M. Schmidberger, J. E. Daniell, K. M. Bedka, and M. Kunz: Severe
759 thunderstorms with large hail across Germany in June 2019. *Weather*, **76**, 228–237, 2021.

760 Zhang, C., Q. Zhang, and Y. Wang: Climatology of hail in China: 1961–2005. *J. Appl. Meteor. Climatol.*, **43**,
761 795–804, 2008.

762 Zinkiewicz, W., and E. Michna: Częstotliwość występowania gradów w województwie lubelskim w zależności
763 od warunków fizjograficznych. [The frequency of hailstorms in the Lubelskie Province depending on the
764 physiographic conditions.] *Annales UMCS B*, **10**, 223–300, 1995.

Formatted: Font: Italic

Formatted: Font: Bold

Formatted: Font: 10 pt

Formatted: Justified, Indent: Left: 0", Hanging: 0.25", Line spacing: 1.5 lines

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Page 19: [1] Deleted

Faye Hulton

5/22/23 8:44:00 PM

Page 21: [2] Deleted

David Schultz

5/22/23 7:27:00 AM