

# 1 Climatology of Large Hail in Europe: Characteristics of 2 the European Severe Weather Database

3 Faye Hulton<sup>1,a</sup>, David M. Schultz<sup>1,2</sup>

4 <sup>1</sup> Centre for Atmospheric Science, Department of Earth and Environmental Sciences, University of Manchester,  
5 Manchester, M13 9PL, United Kingdom

6 <sup>2</sup> Centre for Crisis Studies and Mitigation, University of Manchester, Manchester, M13 9PL, United Kingdom

7 <sup>a</sup> Now at: MetDesk, Aylesbury, HP22 6NJ, United Kingdom

8 Correspondence to: Faye Hulton, [faye.hulton@gmail.com](mailto:faye.hulton@gmail.com)

9  
10 Submitted as an Article to *Natural Hazards and Earth System Science*, 4 February 2023, Revised 12 February,  
11 2024

12 **Deleted: 12**

13 **Deleted: January**

14  
15 **Abstract.** Large hail (greater than 2 cm in diameter) can cause devastating damage to crops and property, and can  
16 even cause loss of life. Because hail reports are often collected by individual countries, constructing a European-  
17 wide large-hail climatology has been challenging to date. However, the European Severe Storm Laboratory's  
18 European Severe Weather Database provides the only pan-European dataset for severe convective storm reports.  
19 The database is comprised of 62,053 large-hail reports from 40 C.E. to September 2020, yet its characteristics  
20 have not been evaluated. Thus, the purpose of this study is to evaluate hail reports from this database for the  
21 purposes of constructing a climatology of large hail. For the period 2000–2020, large-hail reports are most  
22 prominent in June, whereas large-hail days are most common in July. Large hail is mostly reported between 1300–  
23 1900 local time, a consistent pattern since 2010. The intensity, as measured by maximum hail size, shows  
24 decreasing frequency with increasing hailstone diameter, and little change over the 20-year period. The quality of  
25 reports by country varies, with the most complete reporting being from central European countries. These results  
26 suggest that despite its short record, many indications are that the dataset represents some reliable aspects of  
27 European large-hail climatology, albeit with some limitations.

## 28 29 **1 Introduction**

30  
31 Hail with a diameter of at least 2 cm in the longest direction is called *large hail*, and it can cause damage to  
32 crops, property, or even loss of life. Several recent studies have documented the occurrence and variability of  
33 large hail, with special emphasis on the United States and Europe where large hail is common (e.g., Allen and  
34 Tippett 2015; Punge and Kunz 2016; Brooks et al. 2019; Púčik et al. 2019; Tang et al. 2019; Taszarek et al. 2020;  
35 Raupach et al. 2021). The strongest severe convective storms in Europe are often perceived to be less intense  
36 than the strongest storms in the United States, although they can be just as damaging. For example, one of the  
37 most devastating large-hail events took place over Germany on 28 July 2013 when two supercells formed almost  
38 simultaneously, producing hailstones of up to 10 cm in diameter and more than EUR 1 billion in insurance payouts  
39 (Kunz et al. 2018). Other similar events occurred over southern Germany on 10–12 June 2019, with one storm  
40 producing 6-cm hailstones and causing EUR 1 billion in damages (Wilhelm et al. 2021). More recently, several  
41 large-hail events were reported during summer 2021 in Poland, the Czech Republic, Germany, and Italy, with  
42 reported maximum hail sizes in excess of 7 cm (Associated Press 2021; Space 2021a,b,c). Although these extreme  
43 events are widely reported by the media, meteorological research on these storms may be hindered by the lack of  
44 ground-truth hail data, such as onset and ending times, duration, and hailstone size.

47 Such hail data in Europe is generally collected on a national scale, and hence most climatologies are produced  
48 on a country-by-country basis (e.g., Brooks et al. 2009). Given the relatively small sizes of many European  
49 countries, each country has a low probability of large hail occurring at any given time (e.g., Brooks et al. 2019).  
50 A summary table of past European hail climatologies can be found in Tuovinen et al. (2009), and an updated  
51 review was published by Punge and Kunz (2016). Because countries that have a similar spatial extent as Europe  
52 have produced their own climatologies—such as the United States (Tang et al. 2019), Canada (Etkin and Brun  
53 2001), and China (Zhang et al. 2008)—a pan-European large-hail climatology would be highly desired.

54 Climatologies of European convective storms and their impacts have been constructed using a number of  
55 datasets. For example, some studies have examined the climatology of convective storms using remote-sensed  
56 data such as lightning, radar, and satellite (e.g., Punge et al. 2017). Others have examined the environments that  
57 favor such storms, such as through reanalyses or soundings (Rädler et al. 2018; Taszarek et al. 2017, 2018, 2019)  
58 or reanalyses coupled with hailpad data (Sanchez et al. 2017).

59 To create a pan-European dataset of in situ surface reports from severe convective storms (including large  
60 hail, tornadoes, and severe wind gusts), the European Severe Storms Laboratory formed the European Severe  
61 Weather Database (ESWD) in 2006 (Dotzek et al. 2009; Groenemeijer et al. 2017). In addition to collecting  
62 contemporary data, the ESWD has an ongoing objective of synthesizing historical large-hail data which helps  
63 produce a longer and more complete climatology. Despite the tremendous potential value of the ESWD being the  
64 only pan-European large-hail dataset, its characteristics need to be examined to understand its suitability for  
65 answering certain scientific questions about large hail. For example, Taszarek et al. (2019) found substantial  
66 variability across Europe in the frequency of ESWD reports and the frequency of favorable environments for  
67 convective storms.

68 To this effect, Púćik et al. (2019) constructed a climatology of large hail from the ESWD. They examined  
69 hail size, occurrence, annual cycle, diurnal cycle, and societal impacts (e.g., damages, injuries) for 39,537 reports  
70 during the 13-yr period 2006–2018. Although their work shed the first light on the pan-European distribution and  
71 characteristics of large hail and large-hail days from surface reports, they concluded by foreseeing “an update to  
72 this study as the reporting homogeneity improves in future.” In the present article, we explore whether increasing  
73 the size of the dataset through lowering the quality-control levels of the reports and extending the period of  
74 analysis yields comparable results, increasing the generality of Púćik et al.’s (2019) results. In doing so, we also  
75 document the reporting characteristics of the database as a function of time both throughout the 20th century and  
76 within the last 20 years. In particular, we seek the possible existence of a relatively homogeneous period of time  
77 in the database that could be used as a baseline for climatologies and climate-change studies.

78 This article consists of nine sections. Section 2 describes the data from the ESWD used in the present study.  
79 Section 3 discusses the frequency of large-hail reports and days on decadal, annual, and diurnal time scales.  
80 Section 4 investigates the intensity distribution of large hail, as segregated into 1-cm diameter bins, and discusses  
81 how the frequency of large-hail size has changed over the past 20 years. Section 5 looks at the time accuracy of  
82 these reports, how it has changed over the past 20 years, and how it varies by individual countries. Section 6  
83 investigates the spatial distribution of reports by country. Because of the large number of reports from Poland  
84 during the 1930s to 1950s, section 7 focuses on the data from Poland, comparing the historical frequency of reports  
85 during this period to that from the period 2000–2020. Section 8 offers a discussion comparing our work to previous

86 hail climatologies and reflects on the prospects of using the ESWD as a baseline for climate-change research.  
87 Section 9 summarizes the findings of this paper.

88

## 89 **2 Data and methods**

90 The climatology of European large hail in this present article is produced from the ESWD (Dotzek et al. 2009;  
91 Groenemeijer et al. 2017). Large hail in the ESWD is defined as hail with a diameter of at least 2 cm in the longest  
92 direction (Groenemeijer and Liang 2020), comparable to the severe-hail criterion of 0.75 inch (1.9 cm) in the  
93 United States. The current ESWD data on hail is a mixture of historical entries, insurance data information, reports  
94 provided by storm-spotters, national European meteorological organizations, and public entries via the ESWD  
95 website at [www.eswd.eu](http://www.eswd.eu) (Dotzek et al. 2009). Since December 2015, reports have also been collected via ESSL's  
96 European Weather Observer app (Groenemeijer et al. 2017).

97 At the time this study commenced, the ESWD consisted of 62,053 large-hail reports from 59 countries dating  
98 from 40 C.E. to 26 September 2020. All reports with hail sizes less than 2 cm were removed. Of the 59 countries  
99 included with the initial dataset received from the European Severe Storms Laboratory, only 41 were in Europe.  
100 Of those removed, the highest reporting countries were Turkey, Armenia, and Azerbaijan. Reports from other  
101 countries that were removed included Morocco, Turkmenistan, Egypt, and Jordan. The Russian Federation was  
102 included in the present study, even though a small number of reports were from the Asian part of the country. A  
103 small part of Turkey is geographically in Europe, but their data was not included in this study.

104 We also examined two periods of time from the ESWD. The first period is the nearly 121-yr period from 1  
105 January 1900 to 26 September 2020 (when work on this research commenced). We hereafter refer to this period  
106 as 1900–2020, recognizing the omission of data from the last three months and four days of 2020. The second  
107 period is more focused on the most recent large-hail data for the nearly 21-yr period 1 January 2000 to 26  
108 September 2020, hereafter referred to as 2000–2020.

109 All data is imputed in a standard format and is given a single quality-control level by the maintenance team  
110 (Dotzek et al. 2009). There are four quality-control levels given to these entries (Groenemeijer and Kühne 2014):

- 111 • Q0: “as received”, any report straight from the public,
- 112 • QC0+: “plausibility checked”, any report checked by staff at the European Severe Storms Laboratory or a  
113 partner organization,
- 114 • QC1: “report confirmed”, any report confirmed by a reliable source such as a national meteorological  
115 organization or storm-spotters network, and
- 116 • QC2: “event fully verified”, any report from an event that has been subject of a scientific case study.

117 As mentioned in section 1, Púčik et al. (2019) used only QC1 and QC2 events. However, to see if the quality-  
118 control level affects the interpretation of the results, this present study uses QC0+, QC1, and QC2. For the period  
119 1900–2020, there were 9173 QC0+, 45,805 QC1, and 2391 QC2 reports, producing a total of 57,369 large-hail  
120 reports. For the period 2000–2020, there were 6330 QC0+, 20,585 QC1, and 1310 QC2 reports, producing a total  
121 of 28,225 large-hail reports. Thus, the addition of the QC0+ reports increased the size of the 1900–2020 dataset  
122 by 19% and the 2000–2020 dataset by 29%.

123 With these two datasets constructed, we can then look at their characteristics. In particular, we are  
124 interested in the number of large-hail days, size of the large-hail reports, and time accuracy of the reports. The  
125 annual number of large-hail days was derived from the annual number of large-hail reports by removing duplicate

126 dates. We analyzed not only the number of hail reports, but the number of hail days, as well. Hail days are a more  
127 robust measure of hail occurrence and helps minimize variability due to variability in hail reporting across  
128 different countries. Hail days are also useful for certain purposes. For example, Punge and Kunz (2016) wrote that  
129 the insurance industry measures hail damage per region per day instead of measuring damage per individual  
130 hailstorm. Therefore, a pan-European overview of hail days may be of use given that these insurance portfolios  
131 cover large parts of Europe, often including data from multiple countries. However, an awareness of the spatial  
132 distribution of these reports is necessary to identify the most at-risk regions. ▾

133 The size of the hail in each hail report was defined as the maximum hail diameter recorded in cm. Although  
134 the ESWD contains fields for the fall speed and density of the hailstones, these were infrequently reported and  
135 were not considered as part of the present article. To represent the size distribution of the reports, the reports were  
136 classified into 1-cm bins based on their maximum hail diameter, starting at the minimum threshold of large hail  
137 of 2 cm. The *time accuracy* of reports is a field in the ESWD that allows the user to know how reliable the  
138 reporting time of the large-hail report is. The time accuracy represents the total time window that a given report  
139 was recorded in. For example, a 30-min time accuracy would indicate that the hail fell in the window of 15 min  
140 before the recorded time to a maximum of 15 min after the recorded time. The existing ESWD dataset is a result  
141 of both meteorological variations in hail and reporting issues, much as other severe-weather datasets have (e.g.,  
142 Groenemeijer and Kühne 2014; Punge and Kunz 2016; Antonescu et al. 2017; Púčik et al. 2019). Indeed,  
143 underreporting from rural areas and nighttime storms may influence this dataset. These and other characteristics  
144 of the large-hail dataset will be explored in subsequent sections.

145  
146

### 147 **3 Frequency of large hail across Europe: 1900–2020**

148 To understand the number of large-hail reports as a function of time, the annual number of large-hail reports  
149 and annual number of large-hail days were plotted versus year from 1900 to 2020 (Fig. 1). Throughout much of  
150 this period, the annual number of reports was quite small, with peaks during the 1930s, 1940s–1950s, and early  
151 1980s before a steady increase starting around 2000. These two peaks in the 1930s and 1940s–1950s were  
152 associated with a large number of reports from Poland and are investigated further in section 8. The lesser peak  
153 during the 1980s was associated with a number of reports from Italy, but is not considered further.


154 Figure 1 also shows the annual number of hail days from 1900 to 2020. The peaks in large-hail days during  
155 the 1930s and 1940s–1950s suggest that there were many large-hail events, not just many reports. Moreover, these  
156 periods illustrate that, while some periods and some locations may be well represented in the database, reporting  
157 of large hail throughout much of the 20th century in the ESWD is far from complete.

158 Focusing on the last 30 years, the number of reports increased starting around 2000 and continued to rise until  
159 2020. (Recall that the 2020 data was only available until 26 September, which may explain the fewer number  
160 reports, although most large-hailfall in Europe is reported between April and September.) In contrast, the number  
161 of large-hail days began rising a few years earlier in the late 1990s before reaching a plateau during the 2010s  
162 with around 175 annual large-hail days per year, similar to Taszarek et al. (2020, their Fig. 2a). This result suggests  
163 that the database grew around this time by first obtaining data from a larger number of days on which hail fell,  
164 followed by the database growing with a larger number of reports within the same day. The inconsistency in  
165 reports over time is also seen in other convective-storm research, such as for tornadoes as described by Antonescu

**Deleted:** hail days are also aligned with information that the insurance industry uses, as their portfolios cover regions larger than countries and hailstorm outbreaks may cover more than one country. ↗


170 et al. (2017), and may be a reflection in scientific interest in severe convective storms, or due to economic or  
171 political changes.

172



173  
174 **Figure 1.** Time series of annual numbers of large-hail reports (orange line) and large-hail days (green  
175 line) across Europe 1900–2020.


176 To show these data in a slightly different way, a scatterplot is created of the number of hail days versus  
 177 number of hail reports for each year in the dataset, with different colors for the period before and after 2000 (Fig.  
 178 2). The dataset from 1900 onwards suggests a positive linear relationship between large-hail reports and large-  
 179 hail days; however, the spread is sometimes large. The high number of large-hail reports during 1949–1955  
 180 (mostly from Poland, section 8) and early 1950s all congregate in one region of the graph and 2010–2020 also  
 181 congregate in one region. As fewer reports are needed for a greater quantity of large-hail days, either areal extent  
 182 of spotters has improved, the number of reporters has decreased in hail-prone regions, or the ESWD maintenance  
 183 team have improved their ability to detect reports linked to the same event. Thus, the 1950s are a time when  
 184 reports mostly came from Poland (section 8) and captured a large number of large-hail days, indicating that certain  
 185 periods of time can be fruitful for hail research using the ESWD. The spatial distribution of these reports is  
 186 discussed in section 7.  
 187  
 188



189  
 190 **Figure 2. Scatterplot of the annual number of large-hail days versus annual number of large-hail reports**  
 191 **across Europe: 1900–2000 (green dots) and 2000–2020 (orange dots), with corresponding linear regression**  
 192 **lines. These quantities are not divided by the number of years because of the incomplete data for the year**  
 193 **2020.**


194 The average monthly distribution of the number of large-hail reports and large-hail days from 2000 to 2020  
 195 is plotted in Fig. 3. The warm-season months of May, June, and July have the highest number of large-hail reports,  
 196 and the cool-season months from October to March have the lowest. Whereas the month with the highest number  
 197 of large-hail reports is June, the month with the highest number of large-hail days is July. Figure 3 can be compared  
 198 to Púćik et al. (2019, their Fig. 4) who break down the annual cycle into the frequency of reports for the continental  
 199 regions of Europe north of 46°N and the more Mediterranean-influenced regions south of 46°N. Despite these  
 200 differences, these two distributions look similar, with the added information coming from the distribution of large-  
 201 hail days in the present study. The distribution of large-hail days in Fig. 3 is more similar to the shape of the  
 202 distribution of north of 46°N in Púćik et al. (2019, their Fig. 4), meaning that fewer reports occur later in the  
 203 season although the number of large-hail days remains relatively high. These distributions are also similar to  
 204 those from Kunz et al. (2020, their Fig. 2a) for hailstorms in central Europe using radar-derived hail streaks  
 205 combined with all quality levels from the ESWD, indicating that this larger dataset including QC0+ events derived  
 206 using different methods is a reliable source of large-hail data.

207  
 208



209  
 210 **Figure 3. Combined line graph and bar chart of the total monthly numbers of large-hail reports (orange**  
 211 **line) and large-hail days (green bars) across Europe: 2000–2020. These quantities are not divided by the**  
 212 **number of years because of the incomplete data for the year 2020.**

213 The percentage of hail days by month per country (for countries with 100 or more reports) for the period  
214 2000–2020 is shown in Fig. 4. Greece is the only country to not have over 50% of its reports being within the  
215 months of May, June, and July, having a more consistent number of hail days throughout the year. Many countries  
216 do not have any reports before April or after September. Spain, Italy, France, and Croatia have similar distributions  
217 of hail days throughout the year, which may be linked to their Mediterranean setting, although Slovenia, Bosnia  
218 and Herzegovina, and Bulgaria do not share the same characteristics, despite also being situated along the  
219 Mediterranean. Previous studies such as Tazarek et al. (2020) have investigated hail distribution in Europe by  
220 linking events to meteorological and climatological factors, which may help explain some of the differences seen  
221 in Fig. 4. Furthermore, Sanchez et al. (2017) investigated hail events in southern Europe, concluding that even  
222 small geographical and climatological differences can have a large impact on the number of hail days reported,  
223 which may also explain some of the differences in Fig. 4.



224

225


226

Figure 4. Horizontal bar charts of the monthly distributions of large-hail reports (%) for countries with 100 or more reports: 2000–2020.

227 The average diurnal cycle for the number of large-hail reports between 2000 and 2020 is shown in Fig. 5.  
 228 The hour 1500–1559 UTC (labelled 1500 UTC) was the most common time for large hail to be reported with a  
 229 gentle rise and a slightly more rapid decline. When corrected for local legal time (LT) based on each country's  
 230 official time zone, this peak shifts to 1700–1759 LT because most of Europe is east of the Prime Meridian. Figure  
 231 5 can be compared to Púčik et al. (2019, their Fig. 5), who also found a peak during the 1500-UTC hour. These  
 232 distributions are also similar to those from Kunz et al. (2020, their Fig. 2b) who found a peak during 1500–1800  
 233 LT for hailstorms in central Europe using all quality levels from the ESWD, although small differences (e.g.,  
 234 relatively more hail during 1200–1500 LT in Kunz et al. (2020) compared to Fig. 5) may be due to the different  
 235 study areas between these two studies. Thus, the QC0+ data over a longer period of time used in this study  
 236 produces a similar climatology and is consistent with previously published research using a shorter period and  
 237 more selective quality-control levels, indicating that this larger dataset is a reliable source of large-hail data.

238

239




240

241 **Figure 5. Distribution of the hourly time of large-hail reports across Europe in UTC (green line) and local**  
 242 **time (orange line): 2000–2020. Reports are associated with the starting hour (i.e., a report at 1515 UTC**  
 243 **would be placed in the 1500-UTC bin).**

244 To examine the year-by-year consistency of the diurnal cycle, the distribution of large-hail reports as a  
 245 function of local time for each year during the period 2000–2020 is plotted in Fig. 6. Each year mostly reproduces  
 246 the diurnal cycle seen in Fig. 5. The exception is some years, particularly early during this period, that have  
 247 unusual peaks at 1000–1200 UTC. These reports are associated with hail events in the early part of the database  
 248 that occurred at an unknown time during the night or day and were placed in 0000 UTC or 1200 UTC, respectively  
 249 (Púćik et al. 2019, p. 3906). However, by 2010, the diurnal distributions seemed to have settled down to look like  
 250 that in Fig. 5. The consistency after 2010 suggests the possibility that the dataset becomes more consistent in  
 251 reporting events and could represent a stable period for documenting the present large-hail climate of Europe.


252  
 253



255 **Figure 6. Hourly percentage of large hail in local time across Europe in local time for each year 2000 to**  
 256 **2020.**

257  
 258 The diurnal distribution by country was also investigated for countries with 100 or more reports (Fig. 7). For  
 259 most countries, the time period with the most hail reports is between 1400 and 1800 LT, with little variation  
 260 between east and west, and north and south Europe. Belgium seems to be the exception with a larger spread of  
 261 times, but has the lowest number of reports out of these countries, with only 121 reports for 49 hail days (Table  
 262 1), which is likely not representative of the meteorological conditions that would favor large-hail production.

Percentage of hail reports per hour by country (100+) in local time



263


264 **Figure 7. Horizontal bar charts of the hourly distributions of large-hail reports (percentage divided by 265 100) for countries with 100 or more reports: 2000–2020.**

266 **4 Intensity of large hail: 2000–2020**

267 It is not just the frequency of events that determines their impact on society, but also the intensity of the  
268 events, here represented by the maximum diameter of the hail associated with each report. Maximum hail size  
269 can be difficult to measure for several reasons as highlighted by Pilorz (2015). For example, as hail is often  
270 irregular in shape, the maximum diameter is actually the longest axis of the stone. Therefore, if a stone were more  
271 spherical, then its maximum diameter would be smaller than an oblate stone, even though it would have a larger  
272 volume. Furthermore, there is always the possibility that the largest hailstone from any given event has not been  
273 found or that it has partially melted before discovery.

274 For the 28,225 large-hail reports in the present study between 2000 and 2020, 18,132 (64%) had data for the  
275 maximum diameter. These reports were organized into 1-cm bins, ranging from 2.0–2.9 cm to 10+ cm. Frequency  
276 of hail reports decreased with increasing hail size (Fig. 8). The maximum hail size in the database from 2000 to  
277 2020 was 15 cm and was reported in Romania on 26 May 2016. This report was rated QC1, so has been confirmed.  
278 The second largest hail size was 14.1 cm and was reported in Germany on 6 August 2013. This particular hailstone  
279 set the record for the largest hailstone in Germany (ESKP 2013). This report is recorded as QC2 and includes  
280 additional information in the ESWD database, such as the average hailstone size being 8 cm.


281



283 **Figure 8. Bar chart of the number of large-hail reports across Europe by maximum diameter in 1-cm bins:**  
284 **2000–2020.**


285 To investigate the distribution of large-hail size over time, Fig. 9 presents the percentage of each hail-size bin  
 286 per year from 2000 to 2020. During this 21-yr period, the percentage of each bin size does not change dramatically.  
 287 This distribution is similar to the 1989–2018 average from Púćik et al. (2019, their Fig. 7), with about 40% of  
 288 large-hail reports being smaller than 3 cm, about 70% being smaller than 4 cm, and about 84% being smaller than  
 289 5 cm. Therefore, the large-hail size distribution during 2000–2020 may represent a period of stability in reporting  
 290 with little detectable change in large-hail size distributions in the ESWD dataset. For determining the present  
 291 large-hail climate, the stability in the large-hail size distribution after 2000 represents a slightly longer period of  
 292 record compared to that of the diurnal cycle, which stabilized after 2010 (Fig. 6).

293



294  
 295 **Figure 9.** Time series of bar charts of the annual distributions of large-hail size across Europe in 1-cm  
 296 diameter bins (%): 2000–2020.

297 The ESWD has information on average hail size, although only 12% (2237 out of 18,132) of reports contain  
298 this information for 2000–2020. There is, however, a strong positive linear relationship between the average and  
299 maximum hail size recorded (Fig. 10). There were two outliers that are most likely data-entry errors, such as  
300 events with a 2-cm maximum size and 5-cm or 3-cm average size. Both were QC1. The linear relationship ( $R^2 =$   
301 0.76) between maximum and average hail size suggests that the average hail size is about 60% of the maximum  
302 hail size, although there is considerable spread around this line.




303  
304  
305 **Figure 10.** Scatterplot representing 2237 hail reports of the maximum large-hail size versus average large-  
306 hail size across Europe during 2000–2020, with corresponding linear regression line (green dotted line).  
307 The 1:1 line is plotted as a blue line. Two pink dots represent likely data-entry errors where the average  
308 diameter is greater than the maximum diameter.

309 **5 Time accuracy of reports: 2000–2020**


310 The ESWD includes a quantity called time accuracy, defined as the time interval over which the report could  
311 have occurred. For example, a time accuracy of 5 min would mean that the large hail fell within 2.5 min on either  
312 side of the time recorded in the ESWD. Groenemeijer and Liang (2020) specify ten categories of time accuracy:  
313 1 min, 5 min, 15 min, 30 min, 1 h, 3 h, 6 h, 12 h, 1 day, and greater than 1 day. The time accuracy of large hail  
314 in the ESWD has improved over time, with over 50% of reports having a time accuracy of 30 min by 2012,  
315 followed by 50% having a time accuracy of 15 min by 2017 (Fig. 11). Moreover, between 2009 and 2010, reports  
316 with a time accuracy of 30 min became more common, replacing some of the reports with time accuracy of 1 h,  
317 and time accuracy of 12 h and greater become negligible. Viewing the ESWD from 2000–2020 as a whole, these  
318 improvements in time accuracy means that the ESWD is becoming a more reliable source of data, with more  
319 highly temporally resolved data on hail occurrence.

320



321 **Figure 11. Time series of bar charts of the annual distributions of time accuracy of reports across Europe**  
322 **(%): 2000–2020.**

324 On the scale of individual countries, however, work remains to improve the quality of the ESWD. The  
 325 average time accuracy for each country with 100 or more reports during 2000–2020 is shown in Fig. 12. The  
 326 distribution of time accuracy varies considerably among these 24 countries. Germany, Finland, and the Czech  
 327 Republic have more than 40% of their reports with time accuracy of 5 min, whereas Bulgaria, Russian Federation,  
 328 and Moldova have the lowest (1% or less). Figure 12 also indicates the countries for which there is opportunity  
 329 to improve engagement in severe-weather reporting.



330  
 331 **Figure 12. Horizontal bar charts of the time accuracy for countries with 100 or more reports (%): 2000–  
 332 2020.**

333 **6 Spatial distribution by country: 2000–2020**


334 Hail reports across Europe are heterogenous, not just in time, but also in space. Countries such as Germany,  
335 Russian Federation, and Italy reported 4956, 4182, and 2447 large-hail events between 2000 and 2020, compared  
336 to others such as Switzerland, the UK, and Denmark only reporting 266, 85 and 31 cases, respectively (Table 1).  
337 Central and western European countries reported more large hail with 5 out of the top 10 countries located there  
338 (Table 1). Germany has more large-hail reports than the Russian Federation for fewer large-hail days, similarly to  
339 Poland having more reports than Italy, and Austria more reports than Greece. The ESWD grew out of other data-  
340 collecting efforts such as TorDACH (i.e., a tornado dataset collection effort from Germany, Austria, and  
341 Switzerland), which may partially explain why there are more reports for a similar amount of days in Germany,  
342 and Poland has a long history of hail reports (section 7).

343 Besides meteorological reasons for the variability, other reasons that may explain these reporting differences  
344 include the existence, size, and enthusiasm of spotter networks within each country; variations in the ability or  
345 enthusiasm of citizens to input into the ESWD; and the availability of information to quality-control reports. In  
346 fact, many central European countries have larger and more enthusiastic spotter networks [e.g., Poland, as  
347 discussed in Pacey et al. (2021) and section 7 of the present article] and are more likely to enter their reports into  
348 the ESWD. KERAUNOS, based in France, or the MeteoSwiss app based in Switzerland, for example, also  
349 encourage citizen involvement in reporting of extreme events, which are imputed into the ESWD database.  
350 Population density and area of the country were considered as possible explanations for the number of hail reports  
351 varying by country, although neither had a statistically significant relationship with the number of hail reports  
352 (not shown). As with the time-accuracy data (section 5), greater engagement with some countries to encourage  
353 entering their reports into the ESWD would lead to a larger and more complete dataset.

**Table 1. Number of large-hail days and large-hail reports by country: 2000–2020.**

Country	Number of large-hail reports	Number of large-hail days
Germany	4956	692
Russian Federation	4182	1012
Poland	3226	471
Italy	2447	555
France	1707	440
Austria	1502	353
Spain	1027	295
Ukraine	1021	319
Romania	983	267
Greece	975	395
Hungary	903	226
Bulgaria	820	238
Serbia and Kosovo	490	146
Czech Republic	490	174
Moldova	451	117
Croatia	399	181
Finland	382	139
Slovenia	332	116
Switzerland	266	87
Belarus	261	103
Slovakia	234	104
Bosnia and Herzegovina	169	65
Netherlands	165	76
Belgium	121	49
Latvia	86	50
United Kingdom	85	41
Estonia	79	38
Portugal	77	34
Sweden	74	50
Cyprus	68	45
Lithuania	42	23
Luxembourg	39	6
Denmark	31	18
Albania	22	12
Montenegro	21	3
North Macedonia	21	13
Norway	21	15
Malta	11	9
Andorra	6	4
Iceland	4	4
Ireland	2	2

356      Similar to Fig. 2 where the number of large-hail reports was plotted versus the number of large-hail days by  
 357      year, Fig. 13 shows a scatterplot between the number of large-hail reports versus the number of large-hail days by  
 358      country from Table 1. There is a positive linear relationship ( $R^2 = 0.88$ ) between large-hail reports and large-hail  
 359      days by country (Fig. 13), suggesting that large-hail reports are proportional to large-hail days. This relationship  
 360      would therefore imply that reporting frequency is similar across all hail frequencies and countries, except for  
 361      Germany and Poland which have a much greater number of reports proportional to the number of days.



362  
 363 **Figure 13. Scatterplot of the total number of large-hail reports versus large-hail days by country: 2000–  
 364 2020.**

365  
 366      We also investigated the number of large-hail days for each country with 100+ reports for the period  
 367      2000–2020 (Fig 14 a, b, c, and d). We separated these countries into 4 groups based on their total number of  
 368      large-hail days for ease of visualization. We do note that 2020 may show slightly fewer large-hail days than other  
 369      years since the last 3 months of the year are omitted from this dataset.

370      Figure 14 (a) shows the number of large-hail days per country for the top 6 countries with 100+ reports  
 371      for the period 2000–2020. In this subset, Greece displays the fewest large-hail days with 395 days, and Russia the  
 372      greatest, with 1012 days. Germany appears to have the most stable number of annual large-hail days over this  
 373      period, notably from 2003 onwards. However, there remains some year-to-year variation, with 2003–2009  
 374      showing the most stable period. Russia also shows a consistently high number of annual large-hail days throughout  
 375      this period, and although there is a lot of variation up until 2013, the number of large-hail days appears to stabilise  
 376      after this. Italy shows a steady increase in large-hail days up until 2014, after which a slight decline is seen before  
 377      rising again from 2016 onwards. France, Poland, and Greece all appear to see a rise in large-hail days from around



378 2005 onwards, with Poland showing a consistent number of large-hail days from then on, while large variability  
379 is still seen in France and Greece.

380 Figure 14 (b) shows the number of large-hail days per country for the upper middle 6 countries with  
381 100+ reports for the period 2000–2020. In this subset, Hungary displays the fewest large-hail days with 226 days,  
382 and Austria the greatest, with 353 days. Out of the 4 groups, this one shows the most consistent and significant  
383 rise in large-hail days over this period, although there remains much annual variation for each country. Ukraine  
384 displays an anomalous spike of 40 large-hail days in 2002, a total which is not again reached over this period,  
385 with the second highest large-hail year being 2019 with 36 days. Bulgaria and Hungary have a similar number of  
386 large-hail days throughout this period, these gradually increasing until 2016, after which they start to decline.  
387 Additionally, with the exception of 2012, Austria shows a consistent number of large-hail days for the 2010–2018  
388 period.

389 Figure 14 (c) shows the number of large-hail days per country for the lower middle 6 countries with 100+  
390 reports for the period 2000–2020. In this subset, Slovenia displays the fewest hail days with 116 days, and Croatia  
391 the greatest, with 181 days. This group shows significant year-to-year variation in large-hail days, notably for  
392 Slovenia, Moldova, and Finland. Finland has the largest variation in annual large-hail days in this group, with 22  
393 reports in 2008, and none in 2012 and 2015. Serbia and Kosovo, The Czech Republic, and Croatia have a similar  
394 number of large-hail days over this period, although between 2006–2008 and then again from 2019, they display  
395 a greater difference. Slovenia has seen several peaks in large-hail days, the first being in 2005, followed by the  
396 greatest peak in 2009 with 18 large-hail days, which was then followed by a quick decline, before increasing again  
397 from 2015 onwards. Moldova initially demonstrated a steady increase in large-hail days, followed by a peak  
398 between 2013 and 2016, with 2014 seeing the greatest number of large-hail days here with 24 days.

399 Figure 14 (d) shows the number of large-hail days per country for the bottom 6 countries with 100+  
400 reports for the period 2000–2020. In this subset, Belgium displays the fewest hail days with 49 days, and Slovakia  
401 the greatest, with 104 days. There appears to be a rising trend in the number of large-hail days reported for each  
402 country. However, as these countries have few annual large-hail days, it is difficult to determine whether this rise  
403 is due to increased reporting or an increase in large-hail events. Furthermore, although all countries exhibit annual  
404 variation, Belarus shows the greatest variation in this group.

405 Although there generally is a rise in the number of large-hail days for each country throughout the period  
406 2000–2020, there remains much annual variation. The top 50% of countries with 100+ reports for this period are  
407 mostly showing more consistency in the number of annual large-hail days than the bottom 50%. However, the  
408 bottom 25% of countries are generally showing an increase in annual hail-days for this period, although it is  
409 difficult to assess any real trends in large-hail days as these may be due to a better reporting, and not more large-  
410 hail events.




411

412 **Figure 14. Line graph of large hail days per country for countries with 100+ reports: 2000–2020. (a) top 6 countries, (b) upper middle 6 countries, (c) lower middle 6 countries, and (d) bottom 6 countries.**

413

414 We further investigated the hail-size distributions by country for the period 2000–2020 (Fig. 15). Only one  
415 report of each size diameter was taken per country per day to minimize some of the reporting biases. Finland has  
416 the greatest proportion of the lowest hail bin size, whereas Slovenia has the lowest. For sizes 5 cm in diameter  
417 and greater, the proportion of hail sizes recorded starts to diminish drastically, which would be expected as larger  
418 hailstones are rarer. Although Slovenia has the greatest proportion of hail sizes above 5 cm, these reports came  
419 from a sample of 116 hail reports, one of the smallest of the countries analyzed. For hail days with a report above  
420 10 cm, Russia has the greatest quantity with 10 reports over this period, whereas Italy came second with 9 reports  
421 and France with 8. Slovenia, although having a greater proportion, had 5 days with a hail report above 10 cm for  
422 this period.

Distribution of large-hail size by country (2000–2020)



423

Figure 15. Horizontal bar charts of the size distribution of large hail for countries with 100 or more reports (%): 2000–2020.

424

425

426

427 **7 Poland: 1900–2020**

428 As noted in association with Fig. 1, nearly all large-hail reports and large-hail days during the 1930s and  
429 1940s–1950s originated in Poland (Figs. 16a,b). Very few hail days were recorded between 1956 and 2000, before  
430 the general increase along with the rest of Europe for the last 20 years (Fig. 15). There appears to be far fewer  
431 large-hail days over the past 20 years in Poland (30–40 days a year) compared to the 1940s–1950s (100–120 days  
432 a year). With an overall increase in reporting numbers and accuracy, it would be unlikely that the current Polish  
433 reports are missing many events, and therefore the difference in annual numbers of large-hail days seems unlikely.

434 The addition of this data in the ESWD was due to Igor Laskowski who reports:

435 “those reports were based on annual records collected by a Polish National Institute of  
436 Meteorology founded in 1919, now Institute of Meteorology and Hydrology - National  
437 Research Institute (<https://imgw.pl/institut/historia>). The data was collected via hail  
438 questionnaires, which provided information on the size of the hail (vetch-sized, pea-sized, broad  
439 bean-sized, hazelnut-sized, walnut-sized, pigeon egg-sized, hen egg-sized and goose egg-sized)  
440 and also details about time of its occurrence, storm direction and the size of the expected yield  
441 decrease (in percent). The questionnaires were filled in both by agricultural correspondents of  
442 the Polish Central Statistical Office (whose number was growing larger, especially in the  
443 [19]50s) and existing insurance companies which provided hail insurance at this time. Those  
444 records also contain observations of hail reported by observers at meteorological stations.”


445 At the time of this study, data from yearbooks from 1930–1937 and 1946–1955 had been added.

446 Suwala (2011) investigated Polish hail based on data from 23 meteorological stations recorded in the  
447 Meteorological Yearbooks published by the Institute of Meteorology and Water Management for the years 1973–  
448 1980 and the Polish National Climatic Data Centre for the years 1981–2009. They found that over the 37-year  
449 period, March was the month with the highest hail frequency across the country, followed by February and  
450 January. For individual stations, December and January recorded the highest hailfall, with the two stations along  
451 the Baltic coasts having a mean of 8 days. Although these results may indicate a cool-season preference for hail,  
452 there is the possibility that ice pellets or graupel might have been classified as hail (e.g., Punge and Kunz 2018).  
453 Overall, the Baltic coast showed the highest annual mean, whereas central Poland showed the lowest. This result  
454 contradicts the findings of Pilarz (2015) who investigated large hail in Poland for 2007–2015, concluding that  
455 southeast Poland had the greatest number of storms and associated large hail events.

456 Furthermore, the warm months of June to September had the lowest mean hail frequency for all stations. This  
457 contradicts the results found in this present study and those by Púčik et al. (2019) that hail is most frequent in the  
458 warm season, but also contradicts those by Taszarek and Suwala (2014) who investigated large hail in Poland in  
459 2012. In addition, there appeared to be some cyclicity in frequency over the 37-yr period, although this cyclicity  
460 varied greatly when investigating individual stations, and no trends were observed. These results may explain why  
461 Poland possesses a different annual distribution to other locations.

462 Suwala (2011) mentioned previous hail studies in Poland, such Schmuck (1949), Koźmiński (1964), and  
463 Zinkiewicz and Michna (1955), which may offer an explanation on the high number of hail reports during the  
464 1930s and 1950s. Unfortunately, these are not currently available to read. Access to these historical studies may  
465 help explain the quantity of Polish entries in the ESWD during the 1930s, 1940s, and 1950s. Moreover, an effort  
466 to retrieve and input the data from 1973 to 2009 into the ESWD would greatly help with the homogeneity of the

467 Polish dataset. There remains the possibility that this data does not exist as the country suffered major economic  
468 difficulties during this period.



470  
471 **Figure 16. Time series of annual numbers of (a) large-hail reports for Europe (green line) and Poland (red**  
472 **line), and (b) large-hail days for Europe (green line) and Poland (red line): 1900–2020.**

473 As in Fig. 11, the time accuracy of large-hail reports can be plotted as a function of time during 1930–2020  
 474 in Fig. 17. The time accuracy of reporting in Poland has improved over the past 20 years, with over half the  
 475 reports having a time accuracy of 15 min by 2015 (Fig. 16). During the 1930s and 1950s, the time accuracy was  
 476 much lower, around 3 h (Fig. 16). Although this result may suggest that reports were less reliable during this  
 477 period, the consistency in time accuracy (especially during the 1950s) may also suggest that the data-collection  
 478 methods were more consistent. These reports were later found to be based upon the Meteorological Yearbooks  
 479 from the Polish National Institute of Meteorology (I. Laskowski 2022, personal communication). The yearbooks  
 480 contained information on hail size, time of occurrence and storm direction based upon questionnaires posed to  
 481 insurance companies, agricultural correspondents of the Polish Central Statistical Office alongside observations  
 482 from meteorological stations. Laskowski also mentioned that yearbooks from the 1960s and 1970s also existed,  
 483 but was currently unable to find any existing copies. Hence, such data – when it is found – remains to be entered  
 484 into the ESWD.

485 In addition, the reported location accuracy was also investigated, with the most common distances being 1  
 486 and 3 km, similar to those found in the broader 2000–2020 dataset. This result reiterates the importance of these  
 487 earlier reports in constructing a reliable hail climatology, and gives credit to the data-collection method.

488 The historical Polish datasets offer an insight into past hail frequency and reporting accuracies. Results by  
 489 Suwala (2011) for the period 1973–2009 contradict those found for more recent time periods in terms of peak  
 490 annual frequency and spatial distribution of large hail. The potential implications of these discrepancies may  
 491 suggest that distributions of hail size, frequency, and location have changed over time and have not yet been  
 492 established or studied due to the lack of historical pan-European data, highlighting the importance of building the  
 493 ESWD further. Moreover, the existence of Meteorological Yearbooks in Poland could also suggest that other  
 494 nations might hold similar records that remain to be analyzed and could contribute toward building a more  
 495 complete climatology.

496



497  
 498 **Figure 17. Time series of bar charts the annual distributions of time accuracy of reports for Poland (%):**  
 499 **1930–2020.**

500 **8 Comparison to previous hail climatologies and prospects for a baseline for climate-change research**

501 The ultimate goal of severe-storm climatologies is to create a consistent and complete database in space and  
502 time. Consistent data acquisition methods throughout the study area and through time would assist in achieving  
503 this goal; however, consistency is not achievable across Europe. Punge and Kunz (2016) synthesized all European  
504 hail studies in their review, not just large hail. They concluded that not all regions have the same threat of hail,  
505 and they found that efforts to report and record these events vary by country. They further concluded that there  
506 was insufficient evidence to determine any trends in hail events, both in terms of spatial and temporal extent,  
507 highlighting the need for the continuation of the ESWD to form a reliable climatology. Previous studies have  
508 provided pan-European climatologies of hail based using other methods such as Punge et al. (2014, 2017) who  
509 used overshooting cloud tops, Rädler et al. (2018) who used reanalysis data, or Taszarek et al. (2018) who used a  
510 combination of data sources. Some studies projected increases in hailstorms with climate change in Italy (Piani et  
511 al. 2005), Netherlands (Botzen et al. 2010), and Germany (Mohr et al. 2015), as well as across much of Europe  
512 (Taszarek et al. 2021). Other studies have also concluded that there were no positive trends in the frequency of  
513 hail in hailpad data in northern Italy and France (e.g., Eccel et al. 2012; Dessens et al. 2015; Raupach et al. 2021;  
514 Manzato et al. 2023). Tazarek et al. (2019) argued that a combination of datasets is important to construct a robust  
515 climatology, particularly as the spatial and temporal resolutions would often differ between methods. Furthermore,  
516 studies such as Rädler et al. (2018) compared their reanalysis results to surface observed reports from the ESWD  
517 to strengthen their arguments. Therefore, understanding the characteristics of the current surface observations via  
518 the ESWD helps not only build a climatology of large hail in Europe, but can also be used in association with  
519 other research methods to identify the underlying factors which lead to such events.

520 Examining the evidence presented in the present article, we seek a stable time period during 2000–2020.  
521 Based on the number of large-hail reports, no stable time period exists (Fig. 1). Based on the number of large-  
522 hail days, the time period starts around 2012 (Fig. 1). Based on the diurnal cycle of large-hail reports, the time  
523 period starts around 2010 (Fig. 6). Based on the large-hail size distributions, the time period starts around 2004  
524 (Fig. 9). Based on the time accuracy of reports, the time period possibly starts around 2018 (Fig. 11). However,  
525 if one is prepared to accept an accuracy of 3 h or less, then the time period starts around 2010 (Fig. 11).

526 **9 Conclusion**

528 The ESWD provides the only pan-European dataset for large-hail reports. The frequency of reports is sporadic  
529 pre-2000, and hence the focus of this study is for the period 2000 to 2020. Hail reports have continuously increased  
530 since 2000. The annual number of large-hail days have remained steady after 2010 at around 175 days per year,  
531 although some interannual variability is still observed. Increased large-hail reports for similar large-hail days  
532 suggests that a greater spotter network is in operation, and that the engagement with the ESWD is increasing.  
533 When considering the annual number of large-hail days per country, there does appear to be an overall increase  
534 in the quantity observed for the countries which previously reported fewer hail-days, while those which observed  
535 greater numbers throughout this period seem to be stabilizing.

536 The warm season of May to August shows the highest number of large-hail reports and large-hail days, with  
537 June showing the highest large-hail reports and July the highest large-hail days. The number of large-hail reports  
538 decrease faster than large-hail days from June to September. The diurnal cycle shows that the peak hailfall time  
539 is 1500 UTC and 1700 local time.

540 The number of large-hail reports decreases with increasing diameter, and the percentage distribution of each  
541 large-hail size by year does not appear to have changed over the past 20 years. The possibility that hail-size  
542 distribution is changing remains, as smaller, less damaging hail size events are being recorded more regularly.

543 The diurnal cycle by year shows that for the past 10 years, a consistent pattern has emerged, with a rise in the  
544 early afternoon and a decline in the evening. Furthermore, the time accuracy of reports has improved with over  
545 50% of reports being reported to within a 30-minute window by 2012, followed by 50% being reported to within  
546 a 15-minute window by 2017. Not all countries display improved time accuracies. Germany, Finland, and the  
547 Czech Republic have the greatest proportions of 5-minute time-accuracy reports, whereas Russia, Moldova, and  
548 Bulgaria have the highest proportions of 1-h or greater time-accuracy reports. Efforts to improve monitoring and  
549 reporting in these regions is therefore suggested to improve the completeness of the ESWD.

550 Poland possessed anomalously large numbers of large-hail reports during the 1930s, 1940s, and 1950s. The  
551 reason is linked to scientific interest in severe convective storms during these periods alongside a nationwide  
552 effort by the Polish National Institute of Meteorology to record hail events via questionnaires. Yearbooks also  
553 exist for the 1960s and 1970s; however, copies are yet to be retrieved and entered into the database.

554 Even though the dataset remains too short to extract any trends in large-hail pattern distribution, the  
555 climatology presented here provides insight into which countries and geographical regions to target for  
556 improvements in data acquisition. This climatology also helps advance the idea that some time series are starting  
557 to show consistent behavior, suggesting their utility as climate-change baselines. Furthermore, the differences in  
558 both spatial and annual frequencies of hail in Poland over different time periods may suggest that hail trends have  
559 been changing, highlighting the importance of building and maintaining such climatologies. Therefore, the  
560 usefulness of the ESWD will only continue to expand and offer avenues for future severe convective storm  
561 research.

562 *Data availability.* The data were obtained from the European Severe Storms Laboratory European Severe Weather  
563 Database, in accordance with their data policies: <http://www.eswd.eu>.

565  
566 *Author contributions.* FH performed the analyses and wrote the paper. DMS supervised the research, and helped  
567 write and edit the paper.

568  
569 *Competing interests.* The authors declare that they have no conflicts of interest.

570  
571 *Acknowledgments.* This article is derived from FH's BSc dissertation at the University of Manchester. We thank  
572 the European Severe Storms Laboratory for their kind access to the ESWD that made this work available. We  
573 thank Neil Mitchell for his comments on the dissertation. We thank Igor Laskowski for explaining the Polish hail  
574 dataset. We thank the anonymous reviewers for their comments that have improved this article.

575  
576 *Financial support.* Schultz is partially supported by the Natural Environment Research Council grant numbers  
577 NE/N003918/1 and NE/W000997/1.

578  
579 **References**

580

581 Allen, J. T., and M. K. Tippett: The characteristics of United States hail reports: 1955–2014. *Electron. J. Severe*  
582 *Storms Meteorol.*, **10** (3), <http://www.ejssm.org/ojs/index.php/ejssm/article/viewArticle/149>, 2015.

583 Antonescu, B., D. M. Schultz, A. Holzer, and P. Groenemeijer: Tornadoes in Europe: An underestimated  
584 threat. *Bull. Amer. Meteorol. Soc.*, **98**, 713–728, 2017.

585 Associated Press, The: Storm floods German vaccine center, 5 injured by heavy hail. ABC News. Available at:  
586 <https://abcnews.go.com/International/wireStory/german-vaccine-center-flooded-injured-heavy-hail-78459538> (Accessed: 16 July 2021), 2021.

587

588 Brooks, H. E., Doswell, C.A. III, X. Zhang, A. A. Chernokulsky, E. Tochimoto, B. Hanstrum, E. de Lima  
589 Nascimento, D. M. Sills, B. Antonescu, and B. Barrett: A century of progress in severe convective storm  
590 research and forecasting. *Meteorological Monographs*, **59**, 18.1–18.41, 2019.

591 Botzen, W. J. W., L. M. Bouwer, and J. C. J. M. Van den Bergh: Climate change and hailstorm damage: Empirical  
592 evidence and implications for agriculture and insurance. *Resource and Energy Economics*, **32**, 341–362,  
593 2010.

594 Dessens, J., C. Berthet, and J. L. Sanchez: Change in hailstone size distributions with an increase in the melting  
595 level height. *Atmos. Res.*, **158–159**, 245–253, 2015.

596 Dotzek, N., P. Groenemeijer, B. Feuerstein, and A. M. Holzer: Overview of ESSL's severe convective storms  
597 research using the European Severe Weather Database ESWD. *Atmos. Res.*, **93**, 575–586, doi:  
598 10.1016/j.atmosres.2008.10.020, 2009.

599 Eccel, E., P. Cau, K. Riemann-Campe, and F. Biasioli: Quantitative hail monitoring in an alpine area: 35-year  
600 climatology and links with atmospheric variables. *Intl. J. Climatol.*, **32**, 503–517, 2012.

601 ESKP: <https://www.eskp.de/en/natural-hazards/largest-hailstone-found-over-swabian-jura-935341/>. (Accessed  
602 25 March 2022), 2013.

603 Fluck, E., M. Kunz, P. Geissbuehler, and S. P. Ritz: Radar-based assessment of hail frequency in Europe. *Natural*  
604 *Hazards and Earth System Sciences*, **21**, 683–701, 2021.

605 Groenemeijer, P., M. Kuehne, Z. Liang, and N. Dotzek: New capabilities of the European Severe Weather  
606 Database. *5th European Conference on Severe Storms*, Landshut, Germany, 311–312, 2009.

607 Groenemeijer, P., and T. Kühne: A climatology of tornadoes in Europe: Results from the European Severe  
608 Weather Database. *Mon. Wea. Rev.*, **142**, 4775–4790, 2014.

609 Groenemeijer, P., T. Púčik, A. M. Holzer, B. Antonescu, K. Riemann-Campe, D. M. Schultz, T. Kühne, B.  
610 Feuerstein, H. E. Brooks, C. A. Doswell, H.-J. Koppert, and R. Sausen: Severe convective storms in Europe:  
611 Ten years of research and education at the European Severe Storms Laboratory. *Bull. Amer. Meteor. Soc.*, **98**,  
612 2641–2651, doi: 10.1175/BAMS-D-16-0067.1, 2017.

613 Groenemeijer, P., and Z. Liang: ESWD data format specification Version 1.60 and 1.60-csv. Revision 1. Tech.  
614 Report 2020-11, European Severe Storms Laboratory, 66 pp. Available at: <https://www.essl.org/cms/wp-content/uploads/ESSL-rep-2020-11.pdf>, 2020.

615

616 Hand, W. H., and G. Cappelluti: A global hail climatology using the UK Met Office convection diagnosis  
617 procedure (CDP) and model analyses. *Meteorol. Appl.*, **18**, 446–4589, 2011.

618 Hawkins, E., D. Frame, L. Harrington, M. Joshi, A. King, M. Rojas, and R. Sutton: Observed emergence of the  
619 climate change signal: From the familiar to the unknown. *Geophys. Res. Lett.*, **47**, e2019GL086259, 2020.

620 Istrate, V., R. V. Dobri, F. Bărcăianu, R. A. Ciobanu, and L. Apostol: A ten years hail climatology based on  
621 ESWD hail reports in Romania, 2007-2016. *Geographia Technica*, **12**, 110–118, 2017.

622 Kaonga, G.: BBC Weather: Europe braces for hail storms as heavy downpours to cover continent, Express.co.uk.  
623 Available at: [https://www.express.co.uk/news/weather/1454518/BBC-Weather-Europe-forecast-hail-  
624 storms-latest-June-2021-vn](https://www.express.co.uk/news/weather/1454518/BBC-Weather-Europe-forecast-hail-storms-latest-June-2021-vn) (Accessed: 16 July 2021), 2021.

625 Koźmiński, C.: Geograficzne rozmieszczenie największych burz gradowych zanotowanych na obszarze Polski w  
626 latach 1946–1956. [Geographical distribution of major hailstorms recorded in Poland in 1946–1956.] *Przegl.  
627 Geogr.*, **36** (1), 87–102, 1964.

628 Kunz, M., U. Blahak, J. Handwerker, M. Schmidberger, H. J. Punge, S. Mohr, E. Fluck, and K. M. Bedka: The  
629 severe hailstorm in southwest Germany on 28 July 2013: Characteristics, impacts and meteorological  
630 conditions. *Quart. J. Roy. Meteor. Soc.*, **144**, 231–250, 2018.

631 Kunz, M., J. Sander, and C. Kottmeier: Recent trends of thunderstorm and hailstorm frequency and their relation  
632 to atmospheric characteristics in southwest Germany. *Intl. J. Climatol.*, **29**, 2283–2297, 2009.

633 Kunz, M., J. Wandel, E. Fluck, S. Baumstark, S. Mohr, and S. Schemm: Ambient conditions prevailing during  
634 hail events in central Europe. *Natural Hazards and Earth System Sciences*, **20**, 1867–1887, 2020.

635 Manzato, A., Fasano, G., Cicogna, A., Sioni, F., and Pucillo, A.: Trends of sounding-derived indices and  
636 observations in NE Italy in the last 30 years, 11th European Conference on Severe Storms, Bucharest,  
637 Romania, 8–12 May 2023, ECSS2023-3, <https://doi.org/10.5194/eccs2023-3>, 2023.

638 Mohr, S., M. Kunz, and K. Keuler: Development and application of a logistic model to estimate the past and future  
639 hail potential in Germany. *J. Geophys. Res. Atmos.*, **120**, 3939–3956, 2015.

640 Pacey, G. P., D. M. Schultz, and L. Garcia-Carreras: Severe convective windstorms in Europe: Climatology,  
641 preconvective environments, and convective mode. *Wea. Forecasting*, **36**, 237–252, doi: [10.1175/WAF-D-  
642 20-0075.1](https://doi.org/10.1175/WAF-D-20-0075.1), 2021.

643 Piani, F., A. Crisci, G. De Chiara, G. Maracchi, and F. Meneguzzo: Recent trends and climatic perspectives of  
644 hailstorms frequency and intensity in Tuscany and Central Italy. *Nat. Hazards Earth Syst. Sci.*, **5**, 217–224,  
645 2005.

646 Pilorz, W.: Very large hail occurrence in Poland from 2007 to 2015. *Contemporary Trends in Geoscience*, **4**, 45–  
647 55, 2015.

648 Púčik, T., C. Castellano, P. Groenemeijer, T. Kühne, A. T. Rädler, B. Antonescu, and E. Faust: Large hail  
649 incidence and its economic and societal impacts across Europe. *Mon. Wea. Rev.*, **147**, 3901–3916, 2019.

650 Punge, H. J., K. M. Bedka, M. Kunz, and A. Werner: A new physically based stochastic event catalog for hail in  
651 Europe. *Natural Hazards*, **73**, 1625–1645, 2014.

652 Punge, H. J., K. M. Bedka, M. Kunz, and A. Reinbold: Hail frequency estimation across Europe based on a  
653 combination of overshooting top detections and the ERA-INTERIM reanalysis. *Atmos. Res.*, **198**, 34–43,  
654 2017.

655 Punge, H. J., and M. Kunz: Hail observations and hailstorm characteristics in Europe: A review. *Atmos. Res.*, **176**,  
656 159–184, 2016.

657 Rädler, A. T., P. Groenemeijer, E. Faust, and R. Sausen: Detecting severe weather trends using an additive  
658 Regressive Convective Hazard Model (AR-CHaMo). *J. Appl. Meteor. Climatol.*, **57**, 569–587, 2018.

659 Raupach, T. H., O. Martius, J. T. Allen, M. Kunz, S. Lasher-Trapp, S. Mohr, K. L. Rasmussen, R. J. Trapp, and  
660 Q. Zhang: The effects of climate change on hailstorms. *Nature Reviews Earth & Environment*, **2**, 213–226,  
661 2021.

662 Sanchez, J. L., A. Merino, P. Melcón, E., García-Ortega, S. Fernández-González, C. Berthet, and H. Dessens: Are  
663 meteorological conditions favoring hail precipitation change in southern Europe? Analysis of the period  
664 1948–2015. *Atmos. Res.*, **198**, 1–10, 2017.

665 Space: Weather Armageddon hits Poland: Huge hail and a powerful tornado. Earth Chronicles News. Available  
666 at: <https://earth-chronicles.com/natural-catastrophe/weather-armageddon-hits-poland-huge-hail-and-a-powerful-tornado.html> (Accessed: 16 July 2021), 2021a.

667 Space: Giant hail hits highway in Italy damaging hundreds of cars. Earth Chronicles News. Available at:  
668 <https://earth-chronicles.com/natural-catastrophe/giant-hail-hits-highway-in-italy-damaging-hundreds-of-cars.html> (Accessed: 31 July 2021), 2021b.

669 Space: Russia: Hail the size of a quail's egg fell in the Kemerovo region. Earth Chronicles News. Available at:  
670 <https://earth-chronicles.com/natural-catastrophe/russia-hail-the-size-of-a-quails-egg-fell-in-the-kemerovo-region.html> (Accessed: 31 July 2021), 2021c.

671 Schmuck, A.: Burze gradowe. *Czas. Geogr.*, **20**, 260–267, 1949.

672 Suwala, K.: Hail occurrence in Poland. *Quaestiones Geographicae*, **30**, 115–126, 2011.

673 Suwala, K., and E. Bednorz: Climatology of hail in Central Europe. *Quaestiones Geographicae*, **32**, 99–110, 2013.

674 Tang, B. H., V. A. Gensini, and C. R. Homeyer: Trends in United States large hail environments and  
675 observations. *npj Climate and Atmospheric Science*, **2**, 1–7, 2019.

676 Taszarek, M., H. E. Brooks, B. Czernecki, P. Szuster, and K. Fortuniak: Climatological aspects of convective  
677 parameters over Europe: A comparison of ERA-Interim and sounding data. *J. Climate*, **31**, 4281–4308, 2018.

678 Taszarek, M., J. Allen, T. Púčik, P. Groenemeijer, B. Czernecki, L. Kolendowicz, K. Lagouvardos, V. Kotroni,  
679 and W. Schulz: A climatology of thunderstorms across Europe from a synthesis of multiple data sources. *J. Climate*, **32**, 1813–1837, 2019.

680 Taszarek, M., J. T. Allen, P. Groenemeijer, R. Edwards, H. E. Brooks, V. Chmielewski, and S. Enno: Severe  
681 convective storms across Europe and the United States. Part I: Climatology of lightning, large hail, severe  
682 wind, and tornadoes. *J. Climate*, **33**, 10239–10261, 2020.

683 Taszarek, M., J. T. Allen, H. E. Brooks, N. Pilguy, and B. Czernecki: Differing trends in United States and  
684 European severe thunderstorm environments in a warming climate. *Bull. Amer. Meteor. Soc.*, **102**, E296–  
685 E322, 2021.

686 Tuovinen, J. P., A. J. Punkka, J. Rauhala, H. Hohti, and D. M. Schultz: Climatology of severe hail in Finland:  
687 1930–2006. *Mon. Wea. Rev.*, **137**, 2238–2249, 2009.

688 Wilhelm, J., S. Mohr, H. J. Punge, B. Mühr, M. Schmidberger, J. E. Daniell, K. M. Bedka, and M. Kunz: Severe  
689 thunderstorms with large hail across Germany in June 2019. *Weather*, **76**, 228–237, 2021.

690 Zhang, C., Q. Zhang, and Y. Wang: Climatology of hail in China: 1961–2005. *J. Appl. Meteor. Climatol.*, **43**,  
691 795–804, 2008.

692 Zinkiewicz, W., and E. Michna: Częstotliwość występowania gradów w województwie lubelskim w zależności  
693 od warunków fizjograficznych. [The frequency of hailstorms in the Lubelskie Province depending on the  
694 physiographic conditions.] *Annales UMCS B*, **10**, 223–300, 1995.