

# Thermal stratification and meromixis in four dilute temperate zone lakes

Elizabeth D. Swanner<sup>1\*</sup>, Chris Harding<sup>1</sup>, Sajjad A. Akam<sup>1</sup>, Ioan Lascu<sup>2</sup>, Gabrielle Ledesma<sup>1, 3</sup>, Pratik Poudel<sup>1, 4</sup>, Heeyeon Sun<sup>1, 4</sup>, Samuel Duncanson<sup>1, 4</sup>, Karly Bandy<sup>4</sup>, Alex Branham<sup>4</sup>, Liza Bryant-Tapper<sup>4</sup>, Tanner Conwell<sup>4</sup>, Omri Jamison<sup>4</sup>, and Lauren Netz<sup>4</sup>.

<sup>1</sup>Department of Geological & Atmospheric Sciences, Iowa State University, Ames, IA, 50010, USA

<sup>2</sup>Department of Mineral Sciences, National Museum of Natural History, Smithsonian Institution, Washington, DC, 20560, USA

<sup>3</sup>Department of Earth Sciences, Memorial University of Newfoundland, St. John's, Canada

<sup>4</sup>Students in GEOL 406/506, Spring 2022, Iowa State University, Ames, IA, 50010, USA

10

*Correspondence to:* Elizabeth D. Swanner ([eswanner@iastate.edu](mailto:eswanner@iastate.edu))

**Abstract.** Four adjacent lakes (Arco, Budd, Deming, and Josephine) within Itasca State Park in Minnesota, USA are reported to be meromictic in the scientific literature. However, seasonally persistent chemoclines have never been documented. We collected seasonal profiles of temperature and specific conductance and placed temperature sensor chains in two lakes for ~ 1 year to explore whether these lakes remain stratified through seasonal mixing events, and what factors contribute to their stability. The results indicate that all lakes are predominantly thermally stratified and are prone to mixing in isothermal periods during spring and fall. Despite brief, semi-annual erosion of thermal stratification, Deming Lake showed no signs of complete mixing from 2006 to 2009 and 2019-2022 and is likely meromictic. However, the other lakes are not convincingly meromictic. Geochemical data indicate that water in Budd Lake, ~~the-which the most dilute lakewater~~, is predominantly sourced from precipitation. The water in the other three lakes is calcium-magnesium bicarbonate type, reflecting a source of water that has interacted with the ~~deglaciated~~ landscape.  $\delta^{18}\text{O}_{\text{H}_2\text{O}}$  and  $\delta^2\text{H}_{\text{H}_2\text{O}}$  measurements indicate the lakes are supplied by precipitation modified by evaporation. ~~Josephine, Arco, and Deming lakes sit in a valley with likely permeable sediments and may be hydrologically connected through wetlands, and recharged with shallow groundwater, as no streams are present.~~ The water residence time in meromictic Deming Lake is short (100 days), yet it maintains a large reservoir of dissolved iron, ~~indicating that shallow groundwater may be an additional source of water and dissolved ions.~~ ~~Josephine, Arco, and Deming lakes sit in a valley with likely permeable sediments and may be hydrologically connected through wetlands, and recharged with shallow groundwater, as no streams are present.~~ All four lakes develop subsurface chlorophyll maxima layers during the summer. All lakes also develop subsurface oxygen maxima that may result from oxygen trapping in the spring by rapidly developed ~~summer~~ thermoclines. Documenting the mixing status and general chemistry of these lakes enhances their utility and accessibility for future biogeochemical studies, which is important as lake stratification and anoxia become more prevalent due to climate and land use changes.

## 1 Introduction

Four lakes in Itasca State Park, Minnesota, USA - Arco, Budd, Deming, and Josephine (Figure 1) - have been described as meromictic in the scientific literature since the 1960s (Baker and Brook, 1971; Anderson et al., 1985; Stewart et al., 2009). However, there has never been physical and chemical data from all lakes during all seasons - especially spring and fall when mixing would be expected - to document through the maintenance of a chemocline that they are meromictic. The identification of meromictic lakes is important as they are critical analogues for understanding of the biogeochemistry of past oxygen-stratified oceans (Degens and Stoffers, 1976). In that regard, the four lakes in Itasca are particularly well suited, if they are meromictic, as they have low amounts of sulfur and abundant iron, ~~producing~~ resulting in anoxic and “ferruginous” conditions under stratification that are chemically similar to oceans during the Precambrian Era (Swanner et al., 2020).

Meromictic lakes are characterized by a sharp increase in electrical conductivity at a chemocline separating a cold and/or dense monimolimnion from the epilimnion that persists year-round (Boehrer and Schultze, 2008). Thermal stratification is typically not the main mechanism stabilizing meromixis in lakes of the temperate zone as it is in the tropics due to the erosion of the thermocline in autumn (Boehrer et al., 2017). Meromictic lakes are present on all continents but are thought to be rare (Hall and Northcote, 2012; Stewart et al., 2009). However, seasonally persistent stratification, and even meromixis, are becoming more common as a result of temperature increases driven by climate change (Ficker et al., 2017). Furthermore, increased monimolimnion water density can also result from increased chemical constituents that enrich the water density of monimolimnion, such as from runoff, groundwater inputs, remineralization processes associated with organic carbon loading (Hakala, 2004), and ~~road de-icing~~ salt contamination (Koretsky et al., 2012).

While anoxia and hypoxia are common features of temperate lakes under thermal stratification (Nürnberg, 1995), meromictic lakes have a permanently anoxic monimolimnion. These conditions restrict the habitat of aerobic organisms, including zooplankton and fish, can amplify microbial or geochemical cycling of elements across the redoxcline (Busigny et al., 2016), and can impact the emission of greenhouse gases such as methane (Lambrecht et al., 2020). The investigation of meromictic lakes can help to understand the alterations to global biogeochemical cycles that may result from climate change and land-use changes that are driving increases in hypoxia in lakes (Jenny et al., 2016).

Finally, the sediments of meromictic lakes are valuable as archives of past climatic transitions, vegetation changes, changes in sediment transport, and atmospheric deposition patterns (O’Sullivan, 1983). For instance, meromictic Crawford Lake in Ontario, Canada was recently promoted as a “golden spike” for the proposed Anthropocene Epoch based on radionuclide patterns recorded in its laminated sediments (McCarthy et al., 2023). Laminated sediments that form in many meromictic lakes are also useful for studying the formation, deposition, and diagenetic transformation of chemically precipitated minerals (Ledesma et al., 2023). Such minerals can potentially be used as proxies for paleoredox conditions, which have application for paleo marine and lacustrine studies (Swanner et al., 2020).

The goals of this study are to 1) determine whether Arco, Budd, Deming, and Josephine lakes are meromictic and the factors that drive stratification, 2) investigate the water type, sources, and reasons for meromixis, and 3) describe the unique

biological features of these lakes. To achieve these goals, we conducted fieldwork from 2006-2009 and 2019-2022, and integrated data from University of Minnesota students working at the Itasca Biological Station and Laboratories since the 1950s. The description of the lakes, their mixing status, and the factors that control it, and key biological characteristics will provide an expanded understanding why lakes resist mixing ~~and~~ ~~It will also~~ highlight the utility of these lakes for further scientific investigations.

## 2 Materials and Methods

Formation of the four study lakes occurred during the late-Wisconsin glaciation ~12,000 years ago (marine isotope stage 2; Jennings and Johnson, 2011). They occupy a tunnel valley that was formed beneath the Wadena lobe of the Laurentide Ice Sheet. Following glacial retreat, the melting of stagnant ice blocks within the tunnel valley left depressions in the landscape now occupied by lakes and wetlands (Wright Jr., 1993).

Today, the four lakes investigated in this study sit in the HUC-12 watershed that sources the headwaters of the Mississippi River (Watershed Boundary Dataset, 2022). Budd (478.6 meters above mean sea level; MAMSL) is the highest elevation, while Arco (465.8 MAMSL) and Josephine (465.4 MAMSL) lie at similar elevations (Figure 1). Deming is the lowest elevation of the lakes (464.8 MAMSL).

Lake depth measurements were collected using a Garmin Striker 4 dual-beam transducer (sonar) attached to a rowboat or canoe. Depth and GPS measurements were taken every six seconds while the boat was in motion. A Garmin GLO 2 GPS receiver and ArcGIS Collector app was used to navigate, track the boat's course, and ensure even coverage. The shoreline of the lakes was obtained by walking along accessible areas of the shore with the Garmin GLO 2 GPS receiver, or from Lidar-derived digital elevation models. Bathymetry rasters (1 m resolution) were generated from the depth measurements in ArcGIS Pro 3.0 using a 3rd-degree Local Polynomial Interpolation. These rasters were used to calculate lake volumes and contour maps. Rasters and volume data have been deposited with the Environmental Data Initiative (Swanner et al., 2023).

Morphometric data were used to calculate relative depth ( $Z_r$ ; %), which is the ratio of the maximum lake depth ( $Z_m$ ; m) to the diameter of a circle of area equal to that of the lake ( $A_0$ ; m<sup>2</sup>), expressed as a percentage in eq. 1 (Wetzel, 2001):

$$Z_r = \frac{Z_m}{2} \sqrt{\frac{\pi}{A_0}} \times 100 \quad (1)$$

Chemical, physical, and biological parameters measured on the four lakes included depth, temperature, specific conductance, salinity, turbidity, pH, oxidation-reduction potential, dissolved oxygen, photosynthetically active radiation, chlorophyll-a, and phycocyanin. Major cations, anions, and isotopes of water ( $\delta^2\text{H-H}_2\text{O}$  and  $\delta^{18}\text{O-H}_2\text{O}$ ) were determined on lake water retrieved from different depths within the four lakes. Taxon-specific chlorophyll-a fluorescence was collected with a Fluoroprobe (BBE Moldaenke). The data and description of methods are available in the Environmental Data Initiative (Swanner et al., 2023). Measurements were made and samples were collected from a boat anchored within the deepest basin of each lake.

100 A string of temperature loggers (HOBO Water Temp Pro v2) placed at different depths were deployed into the deep areas of Deming, Arco, and Budd Lakes for one year. A conductance logger (HOBO Conductivity Logger) was added near the bottom of the strings in Arco and Budd after six months. These sensors measured temperature every thirty minutes and specific conductivity every 2 hours. The sensor string was not retrieved from Budd, as it could not be located in May 2022. Conductance measurements with a Yellow Spring Instruments ProDSS temperature/conductivity sensor on deployment and removal were used to check for drift in the HOBO conductance logger. Hourly wind speed data for the duration of sensor deployment utilized the ITCM5 (47.2400, -95.1900, 1480 feet elevation) weather station in Itasca State Park. Data was downloaded from MesoWest (<https://mesowest.utah.edu/>). Plots and analyses were produced in Python or R Studio (2022.07.2).

105 The density, Brunt-Väisälä or buoyancy frequency ( $N^2$ ,  $s^{-2}$ ), and the Schmidt stability were calculated using data from seasonal profiles or the loggers in the RLakeAnalyzer package v.1.11.4.1 with equations described in that package's documentation (Winslow et al., 2019).

110 Major anions ( $CO_3^{2-}$ ,  $HCO_3^-$ ,  $Cl^-$ , and  $SO_4^{2-}$ ) and cations ( $Na^+$ ,  $K^+$ ,  $Ca^{2+}$ ,  $Mg^{2+}$ ) were used to produce a Piper diagram in Geochemist's Workbench 15.0. The concentration of cations and anions was calculated as the percentage of total cations and anions in millequivalents (meq)  $L^{-1}$ .

115 The isotopes of water ( $\delta^2H_{H_2O}$  and  $\delta^{18}O_{H_2O}$ ) were measured on spring or seep water that had been filtered with 0.45 micron nylon syringe filters and stored at 4 °C with minimal headspace until analysis. Samples were analyzed with a Picarro L1102-i Isotopic Liquid Water Analyzer at the Stable Isotope Laboratory at Iowa State University. The analytical uncertainty and average correction factor for  $\delta^{18}O_{H_2O}$  are  $\pm 0.05$  ‰ and  $\pm 0.30$  ‰ for  $\delta^2H_{H_2O}$  relative to V-SMOW.

120 Samples for microscopy and water color were collected into amber bottles with a Van Dorn sampler from three different depths in each lake, including the subsurface chlorophyll maximum layer (SCML) SCML, if present, as determined with the YSI ProDSS. Water color was determined on water filtered through a GF-75 (Advantec) glass fiber filter (Cuthbert and del Giorgio, 1992). Absorbance was measured at 440 nm and 750 nm. The absorption coefficient ( $g; m^{-1}$ ) was calculated by subtracting the absorbance at 750 nm from the absorbance at 440 nm and dividing by the path length (m):

$$g_{440} = (2.303 * A_{440} - A_{750}) / (\text{path length}) \quad (2)$$

A conversion was necessary to determine the color (mg Pt  $L^{-1}$ ) of the lake water:

$$\text{Color} = 18.216 * (g_{440}) - 0.209 \quad (3)$$

125 Water sampled from the SCML was preserved with 1% Lugol's solution upon returning to the laboratory. Fixed samples were settled in the dark for three to seven days.

Student reports from courses taking place over several decades at the Itasca Biological Station and Laboratories (IBSL) (Knoll and Cotner, 2018), formerly the Itasca Biological Station, were acquired from the library at the University of Minnesota, Twin Cities. Morphometric data, temperature and conductance profiles, observations of water column mixing, and inferences from biological experiments in these reports are referenced in the current study.



### 130 3 Results and Discussion

135 To ascertain if the study lakes are meromictic, morphometric data were acquired to assess whether the basin shape was prone to stratification (sec. 3.1). Specific conductance profiles collected in this study and from historical records are used to determine if seasonal mixing occurred. Where those records are lacking or incomplete, temperature profiles or high frequency sensor data are used to assess the stability of stratification. Because of the importance of specific conductance measurements in indicating mixing, the major ions that contribution to the lakes' conductance were determined (sec. 3.2). This data along with watershed characteristics is also used to assess potential water and ion sources to the lakes. Finally, because meromixis can result in distinctive biological zones and resulting biogeochemical reactions, an assessment of these major features was made and compared to historical biological records (sec. 3.3).

#### 3.1 Morphometry and Mixing Status

140 All four study lakes exceed a  $Z_r$  of 4% and have surface areas of less than 500,000 m<sup>2</sup>, typical of meromictic lakes in the temperate zone (Swanner et al., 2020). Deming Lake is the largest and deepest of the four study lakes with a 54,325 m<sup>2</sup> surface area and a maximum depth of 20.8 meters (Table 1), with 4.3% of the surface area encompassing water depths at or below 17 m. The maximum depth of Deming Lake has previously been reported as 16.5 m (Hooper, 1951), 17 m (Baker and Brook, 1971), and 17.6 m (Lascu and Plank, 2013). Josephine Lake has the longest fetch and lowest  $Z_r$  of all study lakes. The maximum depth of Josephine Lake (14.8 m) has previously been reported as 10.3 m (Baker and Brook, 1971), 12-13 m (Callis et al., 1976), and 13 m (Gage and Gorham, 1985). The surface area with depths at or below 13 m in Josephine Lake is 5.1%. Budd Lake has the highest  $Z_r$  (8.8%). There was the greatest variation between the previously reported maximum depth of Budd Lake (10.8 m; Baker and Brook, 1971) and measurements presented here (16.1 m). Arco Lake has the smallest surface area of 24,180 m<sup>2</sup> and a maximum depth of 12.8 m. Arco and Budd Lakes have steep banks along their eastern sides (Figure 1).

150 In all lakes, steep-sided deep holes were detected that were deeper than previously reported (Figure 2; Supplementary Figure 1). Differences in technology and the small surface areas encompassing the deepest parts of the basins could account for the variation – these basins may have been difficult to locate and map with manual techniques. Alternately, the temperature, pressure, and/or salinity gradients present at the tops of these holes could deflect sound beams, causing errors in our depth measurements (Boehrer and Schultze, 2008). Maximum depths should be verified with measurements by multiple techniques in the future.

160 Deming Lake has the longest record of seasonal profiles of temperature and specific conductance (2006-2009 and 2019-2022), as its sediments have previously been used to investigate Holocene climate variations (Lascu et al., 2012; McLauchlan et al., 2013). A chemocline, or sharp increase in specific conductance, persists in Deming Lake at all sampled times (Figure 3), which is expected for a meromictic lake. The chemocline occurs at 11-13 m (Figure 3). The chemocline was below 10 m in 1989 (Church et al., 1989), and from 11-13 m a decade later (Reiter et al., 1998) based on temperature-

compensated conductance measurements from IBSL student reports, indicating that the chemocline has been at a consistent depth for decades.

165 A systematic increase in the magnitude of specific conductance readings below the chemocline in Deming Lake was observed from the 2006-2009 to the 2019-2022 datasets (Figure 3). This could result from variations in the amount of in-lake primary productivity over time (Campbell, 1977), or could increase with time since a mixing event (Katsev et al., 2010). Deming Lake mixed after a beaver dam broke on the western side of the lake in 1997 (Frane and Walberg, 1997). If the lake has not mixed since aside from this catastrophic event, the specific conductance should have increased from 1997 to 2006-2009 and from 2006-2009 to 2019-2022 at similar rates. The 1997 conductivity data was converted to specific conductance  
170 using the reported temperature values (Frane and Walberg, 1997), and the value at 14 m in July 1997 was compared to the values at 14 m in July 2006 and July 2021. The specific conductance values have increased between both intervals, but the rate has changed from  $31 \mu\text{Siemens cm}^{-1} \text{ yr}^{-1}$  from July 1997 to July 2006 to  $11 \mu\text{Siemens cm}^{-1} \text{ yr}^{-1}$  from July 2006 to July 2021. This could imply a partial mixing event in the years between 2006 and 2021, or some change to the supply of water and dissolved ions to the lake between the two time intervals being compared.

175 Changes in precipitation could have affected the rate of the increase of specific conductance in Deming Lake over this period. Historical drought records are available for the region encompassing Itasca State Park from the year 2000 (National Integrated Drought Information System; Supplementary Figure 2). Drought over this period could have modulated the rate of the lake's specific conductance increase through increased evaporation (Jellison and Melack, 1993). Alternatively, or in addition, the magnitude of water sources of differing specific conductance supplying the lake may have been altered (Ludlam  
180 and Duval, 2001). In the second case, drought would decrease the magnitude of low specific conductance precipitation and increase the proportion of higher specific conductance groundwater to the annual water budget. As Deming Lake does not have a [permanent](#) surface inlet, changes in streamflow can be neglected. The amount of recharge to the groundwater system may have also decreased due to drought, effectively decreasing groundwater inputs to the lake, and counteracting the effects of evaporation on specific conductance in Deming Lake.

185 The record of seasonal specific conductance profiles, which is necessary to document the seasonally persistent chemocline that characterizes meromictic lakes, is only available for the period of 2019-2022 in the other three lakes. We therefore take a comparative approach to establish the likely mixing status of the other lakes. Budd Lake had the lowest range of specific conductivity values ( $23\text{-}90 \mu\text{S cm}^{-1}$ ). A weakly demarcated chemocline was observed around 4-5.5 m water depths for May 2021, July 2021, October 2021, and May 2022. The January 2021 profile shows a deeper chemocline (Figure 3).  
190 Specific conductance values were uniform down to 8 m in Josephine Lake during the current study period (Figure 3). A chemocline was only detected below 9 m in October 2021, while depths below 11 m were not assessed on other dates during the study period due to ~~the~~ difficulty of finding the deepest area before mapping in 2022. Autumnal circulation down to 10 m was reported in Josephine Lake in 1975 (Gage and Gorham, 1985). A chemocline was also present at all time points for Arco Lake between 7-10 m except for May 2022 (Figure 3). It was poorly developed in May 2021, and its deepest occurrence was  
195 in January 2021. Arco Lake has a maximum depth of 12.6 m, but no specific conductance data was collected below 12 m as

part of the profiles acquired during the study period due to ~~the difficulty of~~ finding the small deep spot (representing 1.7% of the surface area). The chemocline is shallowest in mid-summer, consistent with the 7 m chemocline observed in temperature-equilibrated conductivity readings in 1975 and 1976 (Evans and Bjerklie, 1975; Barnes et al., 1976), and 2011 (Harren et al., 2011).

200 Arco Lake has the next best documentation of chemocline dynamics after Deming Lake due to the placement of temperature sensors (1.5, 3.5, 5.5, 7.5, 9.1, 9.5 m) and a conductance sensor (9.1 m) from May 2021 to May 2022 (Figure 4). In the summer of 2021, Arco Lake developed a summer thermocline. The temperatures at different depths started to converge in late fall until the lake became isothermal and continued cooling from 6 to 4°C in mid-November 2021, representing the erosion of thermal stratification. The Brunt-Väisälä or buoyancy frequency ( $N$ ), commonly expressed as  $N^2$  in units  $s^{-2}$ ,  
205 calculated from the time series temperature data is near zero at that time, indicating an erosion of stability, which persisted until the sensors were removed on May 16, 2022, over a week after the ice-off on May 8 (Supplementary Figure 3). A drop in the conductance value at 9.1 m in early November 2021 preceded isothermal conditions and indicated mixing down to at least 9.1 m (Figure 4). The dimensionless lake number (~~Imberger and Patterson, 1989~~) was calculated according to Imberger and Patterson (1989) using the time series and hourly wind data from Itasca State Park (Supplementary Figure 4). It started to  
210 consistently rise on May 15, 2022, as the epilimnion warmed. In combination with the seasonally variable chemocline depth (Figure 3), these analyses indicate that Arco Lake mixes during isothermal periods and is not a meromictic lake. The data presented is most consistent with Arco being a holomictic lake.

Data from the temperature sensors (0.5, 2.5, 5.5, 8.5, 11.5, and 14.5 m) in Deming Lake from June 2019 to May 2020 are presented in Supplementary Figure 5. The lake became isothermal in early November 2019. The Brunt-Väisälä frequency  
215 calculated from this data dropped to near zero and rebounded slightly due to under-ice thermal stratification, a phenomenon not observed in Arco Lake (Supplementary Figure 3). The specific conductance profiles in Deming consistently show the same trend, although the exact depth and magnitude of the chemocline vary seasonally and on the decadal scale (Figure 3). This indicates partial seasonal mixing in Deming, likely during isothermal periods, but it may be insufficient or of too short a duration to fully mix the lake. Deming was ice-free on April 25, 2020 (Lake Ice Out Dates, 2022), and the lake number started  
220 rising immediately due to the onset of thermal stratification (Supplementary Figure 4), limiting the opportunity for mixing. Long-term seasonal observation of a chemocline is consistent with Deming being a meromictic lake (Zadereev et al., 2017).

Arco therefore presents an end-member of a holomictic lake and Deming an end-member of a meromictic lake among the four lakes studied here. In the absence of high-frequency sensor profiles for Budd and Josephine Lakes, the Schmidt stability (Supplementary Figure 6) and the Brunt-Väisälä frequency (Supplementary Figures 7-10) were calculated from  
225 seasonal profiles of temperature (Supplementary Figure 11) and salinity for all four lakes. The similar volumes, depths, and surface areas of the lakes (Table 1) permit comparison of the Schmidt stability results. The Schmidt stability indicates that Deming and Budd Lakes are the most strongly stratified, followed by Josephine and Arco Lakes. It is unknown when ice came off Budd Lake in 2022, but Deming Lake was ice-free on May 7, and Arco and Josephine Lakes were ice-free on May 8 (Lake Ice Out Dates, 2022) following strong winds on the evening of May 7. Likely, Budd Lake was also ice-free by May 8 given

230 its similar morphometry and location. A chemocline consistent with other summer observations was present in Budd Lake on  
May 17, 2022, which could indicate a lack of spring mixing. However, the January 2021 specific conductance profile for Budd  
Lake shows a very weak and deep chemocline, which would be expected following an autumn mixing event. Without full  
profiles of specific conductance in Josephine Lake, it is impossible to evaluate if the lake mixed. However, based on existing  
235 profiles, if a chemocline persists through spring turnover in Josephine Lake as suggested by the sharp and deep chemocline  
observed in May 2022, then the monimolimnion must be limited to depths below 11 m, representing only 3.6% of total lake  
volume (11,404.5 m<sup>3</sup> of 313,295.5 m<sup>3</sup>).

Water colour is related to the abundance of dissolved organic carbon (DOC) (Pace and Cole, 2002). Of the four study  
lakes, Budd Lake had the most coloured water, followed by Deming Lake (Supplementary Table 1). Arco and Josephine Lakes  
had very little colour. Visually, Budd and Deming Lakes appear brown, while Arco and Josephine Lakes appear green.  
240 Enhanced water colour could lead to a shallower thermocline and stronger stratification in Budd and Deming Lakes due to  
greater light absorption by compounds conferring colour (Houser, 2006). The Brunt-Väisälä frequencies, calculated from  
temperature and salinity profiles, were generally highest in the epilimnia during late summer (July 2021; Supplementary  
Figures 7-10). This indicates that temperature is more important than salinity to the stability of stratification in these lakes. In  
August 2022, when Deming Lake had the highest N<sup>2</sup> values observed from the 2019-2022 dataset (data not plotted in  
245 Supplementary Figure 9), the meromictic stability (S') calculated as described by Walker (1974) was 7.81 J m<sup>-2</sup>, within the  
range observed for the meromictic and ferruginous Lake 120 in Canada (Campbell, 1977) and Lake Nordbytjernet in Norway  
(Hongve, 1999). These observations suggest that meromixis in Deming Lake is primarily due to thermal stratification, with a  
small contribution from salinity that limits-prevents mixing in this lake but maybe not in the others -nearby. This phenomenon  
could be common to dilute meromictic lakes. The mixing event at Deming Lake in the summer of 1997 due to a breached  
250 beaver dam on the western side of the lake (Frane and Walberg, 1997) indicates that mixing can occur due to catastrophic  
events.

### 3.2 Chemical Characteristics and Water Sources

A Piper diagram was used to define the water types of the four lakes based on their major cations and anions. The  
predominant cation in all four study lakes was calcium (Figure 5). Budd Lake had a greater proportion of milliequivalents from  
255 sodium and potassium than the other three lakes, with the charge balance mostly from magnesium. Bicarbonate and carbonate  
ions represented nearly all anion milliequivalents in Deming, Arco, and Josephine Lakes whereas chloride and sulfate  
contributed to the anion balance in Budd Lake. The water type for Deming, Arco, and Josephine Lakes is a calcium-bicarbonate  
type, whereas Budd Lake does not have a dominant water type. The low specific conductance and lack of water type in Budd  
Lake indicate the water source is predominantly precipitation. Calcium, magnesium, and carbonate/bicarbonate ions source  
260 from the calcareous Itasca moraine and are typical for the Itasca region (Megard et al., 1993). This indicates that water  
supplying Deming, Arco, and Josephine Lakes has undergone more water-rock reaction than the water supplying Budd Lake,  
such as might be expected from an increasing proportion of groundwater seepage to a lake's water inputs. Deming, Arco, and

Josephine Lakes lie in a tunnel valley channel containing coarser sands and gravel while Budd Lake is poorly integrated hydrologically with the other three lakes, and the net seepage in Budd Lake is probably the lowest among all the lakes. Arco Lake (465.8 MAMSL) is located 51 meters north of Josephine Lake (465.4 MAMSL) and Deming Lake (464.8 MAMSL) is located 284 meters north of Arco Lake. The similar lake levels and [the occurrence of](#) wetlands between Arco and Deming Lakes (Figures 1 and Supplementary Figure 1) indicate these lakes are likely to be hydrologically connected. If the water table mimics the surface topography, groundwater potentially flows from Arco-Josephine to Deming Lake. However, Budd Lake (478.6 MAMSL) is perched highest in the watershed, located off-axis of Josephine-Arco-Deming Lakes, to the west of a ridge (Figure 1). Because Budd Lake is perched, the ridge is likely composed of less permeable material, possibly till, than the valley sediments. The presence of wetlands on the north end of Budd Lake that follow a small valley towards Deming Lake suggests that there could be a [periodic](#) surface or near surface hydrological connection from Budd Lake to Deming Lake. Lakes highest in their watershed are more likely to receive a greater proportion of water from precipitation (Kratz et al., 1997), consistent with the dilute nature of Budd Lake water.

The isotopic composition of water (i.e.,  $\delta^2\text{H}_{\text{H}_2\text{O}}$  and  $\delta^{18}\text{O}_{\text{H}_2\text{O}}$  in ‰) in May 2021 varied little with depth in each lake (Supplementary Figure 12). In Arco, Budd, and Deming Lakes sampled in June 2019, the epilimnion waters were depleted relative to the deeper water. Additional measurements at Deming Lake in July and October 2021 show enriched values in the epilimnion relative to the deeper water. Preferential evaporation of light isotopes during the summer and early fall drives the epilimnia to more enriched values. In Figure 6, all lakes data points diverge from a local meteoric water line (LMWL; Stelling et al., 2021), falling on a lake evaporation line (LEL) with equation  $\delta^2\text{H}_{\text{H}_2\text{O}} (\text{‰}) = 4.5 * \delta^{18}\text{O}_{\text{H}_2\text{O}} (\text{‰}) - 29.9$ . The intersection of the LMWL and the LEL is the composition of regional groundwater, representing isotopically depleted snowmelt as the predominant source of recharge to the local aquifers (Krabbenhoft et al., 1994).

To ascertain the isotopic composition of groundwater in the area, historical  $\delta^2\text{H}_{\text{H}_2\text{O}}$  and  $\delta^{18}\text{O}_{\text{H}_2\text{O}}$  data from springs at Elk Lake in Itasca State Park were retrieved from the Minnesota Spring Inventory (Minnesota Department of Natural Resources, 2022). The Elk Lake springs were sampled again in May 2022, and samples were collected from spring along Nicollet Creek and the bog on the southeast side of Deming Lake (Supplementary Table 2). These sites had abundant marsh marigolds (Supplementary Figure 13), which bloom in May at groundwater discharge sites in northern Minnesota (Rosenberry et al., 2000). The sites at Elk Lake and Nicollet Creek had visible iron mineralization, indicating reducing, iron-bearing water discharge at the surface (Supplementary Figure 13). The Nicollet Creek spring sample lies closest to the intersection of the LEL and LMWL, whereas the Deming bog sample lies closest to the lakes but is more enriched than the lakes (Figure 6).

The seasonal amplitude of a lake's oxygen isotopes can be used to calculate a lake's water residence time (Engel and Magner, 2019). A time series of temperature data from weather station ITCM5 was used to determine average daily temperatures, and seasonal maximum and minimum temperature values were used to calculate the amplitude in  $\delta^{18}\text{O}_{\text{H}_2\text{O}}$  inputs of precipitation and water vapor  $\delta^{18}\text{O}_{\text{H}_2\text{O}}$  (Supplementary Figure 14). These two values were used to estimate a mean water residence time of 100 days for Deming Lake. The Deming Lake  $\delta^{18}\text{O}_{\text{H}_2\text{O}}$  values encompass early spring, mid-summer, and late

summer data, but data from the other three lakes was only from May and June time points (Supplementary Figure 12), and so the mean water residence time could not be calculated by this method.

Potential water sources to Deming Lake are precipitation, outflow from the boggy areas to the south and west, and groundwater. In addition to higher specific conductance values than the other lakes, Deming also has higher concentrations of iron in the monimolimnion than the other lakes, exceeding 1 mM, and has sediments enriched in biogenic magnetite (Lascu et al., 2012). As ~~springs-groundwater~~ in Itasca State Park ~~has-have~~ visible iron mineralization (Supplementary Figure 13), it is possible that groundwater supplies iron into Deming Lake (Swanner et al., 2020). Due to active iron redox cycling, which scavenges iron into the monimolimnion in meromictic lakes, dissolved iron can accumulate to high concentrations in its ferrous form (Busigny et al., 2016; Campbell, 1977). The low abundance of sulfur (Figure 5) means very little sulfide can be produced in the monimolimnion. The lack of anion for precipitation of ferrous iron as sulfide mineral allows for the maintenance of these high ferrous iron concentrations and “ferruginous” conditions.

### 3.3 Biological Characteristics

A ~~subsurface-chlorophyll-maximum-layer (SCML)~~ often develops within the metalimnion, the middle seasonally mixed layer of meromictic lakes (Baker and Brook, 1971; Camacho, 2006). Stable stratification, the availability of both light and nutrients, and ~~avoidance of~~ zooplankton grazing ~~in the epilimnion~~ drive the formation and location of the SCML in meromictic lakes (Klausmeier and Litchman, 2001).

Prior work on these four lakes indicated the presence of a lake-wide turbidity maximum layer that was persistently ~~present~~ just below 5 m in Deming Lake (Baker and Brook, 1971). This peak was observed between late May and August from 1968-1970 but was absent in the winter. It was reported in that study that bacteria and phytoplankton both contributed to the turbidity maximum based on microscopic observations. This work indicated the layer was populated by the cyanobacteria *Oscillatoria agardhii* var. *isothrix* (this genus is now called *Planktothrix*).

From 2019 to 2022, chlorophyll-a measured via multi-wavelength fluorescence acquired with a Fluoroprobe allowed for deconvolution of the fluorescence signal so chlorophyll-a could be attributed to one of four taxonomic groups: Cyanobacteria, Chlorophyta, Diatoms and Dinophyta, or Cryptophyta. Taxon-specific concentrations of chlorophyll-a in each lake inform phytoplankton community structures with depth (Figure 7). Deming Lake has a persistent SCML around 5-6 m attributable to Cyanobacteria. The SCML was less distinct in May 2022, one and a half weeks after ice off. It was also less distinct in May 2021. This suggests that the SCML in Deming Lake is a summer phenomenon that develops at or below the thermocline (Supplementary Figure 11), consistent with the observations from Baker and Brook (1971). In June 2019, a profile of photosynthetically active radiation was also recorded (Supplementary Figure 15), and showed an inflection point at 5.5 m, corresponding to the depth of the SCML at that time. These observations are consistent with cyanobacteria forming a dense accumulation at that depth, attenuating the light flux into deeper waters.

Samples from the epilimnia or the SCML were taken from the four lakes in May 2022: Arco (5 m), Budd (4.5 m), Deming (3.5 m), and Josephine (4.5 m). The most common phytoplankton of the four lakes were the cyanobacteria *Planktothrix*



sp. and *Aphanocapsa* sp., and the green alga *Monoraphidium* sp. (Supplementary Figure 16). Some other phytoplankton species identified were the red alga *Cryptomonas* sp., the cyanobacteria *Planktolyngbya* sp. and *Dolichospermum* sp., as well as other colonial green algae and cyanobacteria species.

All four lakes have subsurface oxygen maxima exceeding air saturation during the summer but lack this feature in the fall (Figure 8). The subsurface oxygen maximum in Deming Lake usually occurs at 4 m, near the top of the thermocline (Bieter et al., 1991; Church et al., 1989; Balk et al., 2007). Metalimnetic oxygen maxima can result because dissolved oxygen is more soluble in cold water after spring mixing, but as the water warms the gas solubility decreases, causing oxygen to become supersaturated. Stratification induced by the thermocline then limits the discharge of supersaturated oxygen to the atmosphere (Wilkinson et al., 2015). Deming Lake rapidly develops a thermocline after ice-off (Supplementary Figure 5), providing a mechanism for gas trapping. Enhanced biological productivity can also contribute to the oxygen maximum, but is generally a smaller contributor (Wilkinson et al., 2015; Craig et al., 1992). The subsurface oxygen maximum is above the SCML in Deming Lake, suggesting that the entrainment of spring oxygen is a larger contributor than photoautotrophy to the oxygen maximum. However, there is also a pH maximum at ~~that~~ 4 m during the summer (Bieter et al., 1991; Supplementary Figure 18; Church et al., 1989; Barkow and Habedank, 1990), as is expected for oxygenic photosynthesis. Itasca Biological Station and Laboratories student experiments that quantified photosynthesis and respiration via dissolved oxygen measurements in bottle experiments generally found that net oxygen release was minimal in the SCML due to vigorous respiration (Barkow and Habedank, 1990; Engstrom et al., 1974). However, primary productivity is often highest within the SCML of other meromictic lakes (Camacho, 2006). Measurements of the O<sub>2</sub>/Ar ratio would help to quantify the contribution of oxygen trapping vs. oxygenic photosynthesis to the Deming Lake subsurface oxygen maximum (Craig et al., 1992).

The occurrence of a summer SCML in Arco, Budd, and Josephine Lakes was more variable than in Deming Lake (Figure 7). Arco Lake had a spring bloom of cyanobacteria in May 2021 and in May 2022, at which time the highest chlorophyll-a of this study was recorded. At both times, the SCML occurred below the oxygen maximum, in a hypoxic zone of the water column. The May 2022 SCML at 5 m in Arco Lake also ~~enhanced light attenuation~~ attenuated light below that depth (Supplementary Figure 15), as was observed in Deming Lake in June 2019.

Arco, Budd, and Josephine Lakes also have subsurface oxygen maxima during summer, although not as consistently as Deming. When the depth of the thermocline is plotted against the depth of the oxygen maximum for all four lakes, there is a significant correlation if October 2021 samples are excluded (Supplementary Figure 19). In the fall, the subsurface oxygen maximum is absent (the highest dissolved oxygen values are at the surface), and the thermocline is deepest (Figure 8). This may reflect that in fall the thermocline has deepened into hypoxic or anoxic waters.

#### 4 Conclusions

Although reported to be meromictic, based on seasonal temperature and specific conductance profiles, Arco, Budd, and Josephine Lakes may be holomictic. Higher frequency specific conductance profiles in multiple seasons during turnover



events would better establish their frequency of mixing. Deming Lake has been meromictic in the periods of data collection presented here (2006-2009 and 2019-2022) but experienced a mixing event in 1997 when an adjacent beaver dam broke. Although a seasonally persistent chemocline occurs in Deming Lake, stratification is predominantly a thermal phenomenon, with stability mostly conferred by a thermocline that develops rapidly after isothermal spring and fall periods. Thermal stratification is not typical of meromictic lakes in the temperate zone, but this weak stratification may be common in dilute boreal lakes (Meriläinen, 1970; Campbell, 1977) and may be enhanced by water color. However, such meromixis may be easily perturbed by hydrographic or land-use changes (Hongve, 2002), which is consistent with Deming Lake mixing in 1997.

The chemical composition of the study lakes reflects their position in the watershed and water sources. The highest-elevation lake, Budd Lake, was very dilute and had no distinguishable water chemical type and likely received most of its water from precipitation. Arco, Deming, and Josephine Lakes are all Ca-Mg-bicarbonate type waters (Figure 5). Deming Lake had the highest ionic load, consistent with a greater contribution of water that had undergone some water-rock interaction, likely groundwater. These results further support the conclusion that due to the dilute nature of the water in these lakes, stratification is predominantly a thermal rather than chemical phenomenon. Further work is necessary to characterize groundwater composition and quantify and localize groundwater input to determine if Deming Lake could be classified as crenogenic meromictic. The short water residence time in Deming Lake implied by the analysis of  $\delta^2\text{H}_{\text{H}_2\text{O}}$  and  $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ , along with the minimal surface water inputs and likely high permeability of tunnel valley deposits suggest the shallow groundwater system should be considered as a major water source.

All four lakes have SCML during summer, as was reported over decades of observations, consistent with all four lakes being thermally stratified in the summer months. The depth of the SCML varies in Arco, Budd, and Josephine and is sometimes absent, but persists at around 5 m in Deming, corresponding to the base of the photic zone. Subsurface oxygen maxima above air saturation likely develop by trapping gas in cold water that then warms with rapid thermocline development in the spring. Corresponding pH maxima in Arco and Deming indicate that oxygenic photosynthesis could also contribute to the subsurface oxygen maxima, and indicate that zones of maximum oxygen production, primary productivity, and biomass may be spatially decoupled in these lakes due to the strong gradients resulting from stratification.

**Code availability** Codes for data analysis are available at <https://github.com/eswanner/Itasca>.

**Data availability** Water properties of Arco Lake, Budd Lake, Deming Lake, and Josephine Lake in Itasca State Park from 2006-2009 and 2019-2021, v. 2. Environmental Data Initiative. <https://doi.org/10.6073/pasta/6f4292da3adcc029934805fdb9a314d4>

**Supplement link:** the link to the supplement will be included by Copernicus, if applicable.

**Author contribution:** Individuals in parenthesis contributed substantially to the study's conception (EDS), data acquisition (EDS, CH, SAA, IL, GL, PP, HS, SD, KB, AB, LB-T, TC, OJ, LN), or analysis (EDS, CH, SAA, GL, PP, HS, SD, KB, AB, LB-T, TC, OJ, LN); contributed substantially to drafting the manuscript (EDS, CH, SAA, GL, PP, HS, SD, KB, AB, LB-T,

395 TC, OJ, LN); and approved the final submitted manuscript (EDS, CH, SAA, IL, GL, PP, HS, SD, KB, AB, LB-T, TC, OJ, LN).

**Competing interests:** The authors declare that they have no conflict of interest.

**Acknowledgements:** This work was supported by a National Science Foundation CAREER award to EDS (1944946).

The GeoAllies program, funded by the National Science Foundation, supported student participation in the GEOL 406/506  
400 field course, as did funds from the Department of Geological & Atmospheric Sciences at Iowa State University. Support for  
the field course was also provided by Dr. Jonathan Schilling, Director of the Itasca Biological Station and Laboratories, and  
Dr. Emily Schilling. Phytoplankton was identified at the genus level with the help of Dr. Lesley Knoll, Associate Director of  
the Itasca Biological Station and Laboratories, who also provided assistance with field operations for the duration of the 2019-  
2022 field seasons. Chad Wittkop and Moses Lado (Minnesota State University) assisted with fieldwork in 2019. Jazlyn Beeck  
405 and Elise Bauer (Iowa State University) assisted with fieldwork in 2021. Julia Kelly and Shannon Farrell of the University of  
Minnesota Library assisted with accessing Itasca Biological Station and Laboratories student reports.

## References

- Anderson, R. Y., Dean, W. E., Bradbury, P., and Love, D.: Meromictic Lakes and Varved Lake Sediments in North America,  
410 19 pp., 1985.
- Baker, A. and Brook, A.: Optical density profiles as an aid to the study of microstratified phytoplankton populations in lakes,  
Arch. für Hydrobiol., 69, 214–233, 1971.
- Balk, D., Graetz, A., Morantes, A., and Olson, D.: Itasca Biological Station Student Report 2485: Comparative investigation  
of zooplankton diel vertical migration patterns in Arco and Deming Lake: the effect of predation, University of Minnesota, 24  
415 pp., 2007.
- Barkow, M. and Habedank, T.: Itasca Biological Station Student Report 1906: A Limnological study of Demming Lake, 1990.
- Barnes, P., Halpern, T., and Meier, T.: Itasca Biological Station Student Report 1157: A limnological investigation of Arco  
Lake, University of Minnesota, Itasca Biological Station, 47 pp., 1976.
- Bieter, J., Romig, G., and Waugh, J.: Itasca Biological Station Student Report 1905: An ecological study of the meromictic  
420 Deming Lake, University of Minnesota, 23 pp., 1991.
- Boehrer, B. and Schultze, M.: Stratification of lakes, Rev. Geophys., 46, <https://doi.org/10.1029/2006RG000210>, 2008.
- Boehrer, B., von Rohden, C., and Schultze, M.: Physical Features of Meromictic Lakes: Stratification and Circulation, in:  
Ecology of Meromictic Lakes, edited by: Gulati, R. D., Zadereev, E. S., and Degermendzhi, A. G., Springer International  
Publishing, Cham, 15–34, [https://doi.org/10.1007/978-3-319-49143-1\\_2](https://doi.org/10.1007/978-3-319-49143-1_2), 2017.
- 425 Busigny, V., Jezequel, D., Cosmidis, J., Viollier, E., Benzerara, K., Planavsky, N. J., Alberic, P., Lebeau, O., Sarazin, G., and  
Michard, G.: The Iron Wheel in Lac Pavin: Interaction with Phosphorus Cycle, in: Lake Pavin: History, Geology,


- Biogeochemistry, and Sedimentology of a Deep Meromictic Maar Lake, edited by: Sime-Ngando, T., Boivin, P., Chapron, E., Jézéquel, D., and Meybeck, M., 1–421, <https://doi.org/10.1007/978-3-319-39961-4>, 2016.
- 430 Callis, J., Hummon, M. R., and Skerrett, P.: Itasca Biological Station Student Report 1158: Lake Josephine, Summer 1976, University of Minnesota, 34 pp., 1976.
- Camacho, A.: On the occurrence and ecological features of deep chlorophyll maxima (DCM) in Spanish stratified lakes, *Limnetica*, 25, 453–478, 2006.
- Campbell, P.: Descriptive limnology of Lake 120, a meromictic lake on the Precambrian Shield in northwestern Ontario, University of Manitoba, 1977.
- 435 Church, T., Tillotson, M., and Wendler, L.: Itasca Biological Station Student Report 1830: Survey of the meromictic Demming Lake, Summer 1989, University of Minnesota, 27 pp., 1989.
- Craig, A. H., Wharton, R. A., McKay, C. P., Science, S., Series, N., and Jan, N.: Oxygen Supersaturation in Ice-Covered Antarctic Lakes : Biological Versus Physical Contributions, *Science* (80-. ), 255, 318–321, 1992.
- Cuthbert, I. D. and del Giorgio, P.: Toward a standard method of measuring color in freshwater, *Limnol. Oceanogr.*, 37, 1319–  
440 1326, <https://doi.org/10.4319/lo.1992.37.6.1319>, 1992.
- Degens, E. T. and Stoffers, P.: Stratified waters as a key to the past, *Nature*, 263, 22–27, <https://doi.org/10.1038/263022a0>, 1976.
- Engel, L. and Magner, J.: Estimating the direction of lake hydraulic residence in Minnesota’s sentinel lakes: implications for management, *Int. J. Hydrol.*, 3, 362–367, <https://doi.org/10.15406/ijh.2019.03.00201>, 2019.
- 445 Engstrom, G., Dahl, B., Nelson, D., and Netko, D.: Itasca Biological Station Student Report 777: An ecological study of Deming Lake, University of Minnesota, 39 pp., 1974.
- Evans, J. and Bjerklie, D.: Itasca Biological Station Student Report 842: Some limnological investigations of Arco Lake, Itasca State Park, Minnesota, University of Minnesota, 40 pp., 1975.
- Ficker, H., Luger, M., and Gassner, H.: From dimictic to monomictic: Empirical evidence of thermal regime transitions in  
450 three deep alpine lakes in Austria induced by climate change, *Freshw. Biol.*, 62, 1335–1345, <https://doi.org/10.1111/fwb.12946>, 2017.
- Frane, M. A. and Walberg, T. J.: Itasca Biological Station Student Report 2260: Arco and Demming: Two Worlds Divided, Demming: When the Levee Breaks, 1997.
- Gage, M. A. and Gorham, E.: Alkaline phosphatase activity and cellular phosphorus as an index of the phosphorus status of  
455 phytoplankton in Minnesota lakes, *Freshw. Biol.*, 15, 227–233, <https://doi.org/10.1111/j.1365-2427.1985.tb00195.x>, 1985.
- Hakala, A.: Meromixis as a part of lake evolution - observations and a revised classification of true meromictic lakes in Finland, *Boreal Environ. Res.*, 9, 37–53, 2004.
- Hall, K. and Northcote, T.: Meromictic Lakes, in: *Encyclopedia of Lakes and Reservoirs*, edited by: Bengtsson, L., Herschy, R., and Fairbridge, R., 519–524, 2012.
- 460 Harren, S., Kartak, J., Knight, J., and Lehman, J.: The effects of fish presence and mixing patterns on water clarity in lakes:

- Arco, Deming, and Itasca, Field Ecology, University of Minnesota, Itasca Biological Station and Laboratories, 17 pp., 2011.
- Hongve, D.: Long-term Variation in the Stability of the Meromictic Lake Nordbytjernet Caused by Groundwater Fluctuations, *Nord. Hydrol.*, 30, 21–38, 1999.
- Hongve, D.: Seasonal Mixing and Genesis of Endogenic Meromixis in Small Lakes in Southeast Norway, *Hydrol. Res.*, 33, 189 LP – 206, 2002.
- 465 Hooper, F. F.: Limnological Features of a Minnesota Seepage Lake, *Am. Midl. Nat.*, 46, 462–481, <https://doi.org/10.2307/2421989>, 1951.
- Houser, J. N.: Water color affects the stratification, surface temperature, heat content, and mean epilimnetic irradiance of small lakes, *Can. J. Fish. Aquat. Sci.*, 63, 2447–2455, <https://doi.org/10.1139/F06-131>, 2006.
- 470 Imberger, J. and Patterson, J. C.: Physical Limnology, vol. 27, edited by: Hutchinson, J. W. and Wu, T. Y. B. T.-A. in A. M., Elsevier, 303–475, [https://doi.org/10.1016/S0065-2156\(08\)70199-6](https://doi.org/10.1016/S0065-2156(08)70199-6), 1989.
- Jellison, R. and Melack, J. M.: Meromixis in hypersaline Mono Lake, California. 1. Stratification and vertical mixing during the onset, persistence, and breakdown of meromixis, *Limnol. Oceanogr.*, 38, 1008–1019, <https://doi.org/10.4319/lo.1993.38.5.1008>, 1993.
- 475 Jennings, C. E. and Johnson, M. D.: The Quaternary of Minnesota, in: Quaternary Glaciations Extent and Chronology, Part IV - a closer look, edited by: Ehlers, J., Gibbard, P. L., and Hughes, P. D., 2011.
- Jenny, J.-P., Francus, P., Normandeau, A., Lapointe, F., Perga, M.-E., Ojala, A., Schimmelmann, A., and Zolitschka, B.: Global spread of hypoxia in freshwater ecosystems during the last three centuries is caused by rising local human pressure, *Glob. Chang. Biol.*, 22, 1481–1489, <https://doi.org/10.1111/gcb.13193>, 2016.
- 480 Katsev, S., Crowe, S. A., Mucci, A., Sundby, B., Nomosatryo, S., Haffner, G. D., and Fowle, D. A.: Mixing and its effects on biogeochemistry in persistently stratified, deep, tropical Lake Matano, Indonesia, *Limnol. Ocean.*, 55, 763–776, 2010.
- Klausmeier, C. A. and Litchman, E.: Algal games: The vertical distribution of phytoplankton in poorly mixed water columns, *Limnol. Oceanogr.*, 46, 1998–2007, <https://doi.org/10.4319/lo.2001.46.8.1998>, 2001.
- Knoll, L. B. and Cotner, J. B.: University of Minnesota Itasca Biological Station and Laboratories: Over 100 Years of Field-  
485 based Education and Research, *Limnol. Oceanogr. Bull.*, 27, 42–44, 2018.
- Koretsky, C. M., MacLeod, A., Sibert, R. J., and Snyder, C.: Redox Stratification and Salinization of Three Kettle Lakes in Southwest Michigan, USA, *Water, Air, Soil Pollut.*, 223, 1415–1427, <https://doi.org/10.1007/s11270-011-0954-y>, 2012.
- Krabbenhoft, D. P., Bowser, C. J., Kendall, C., and Gat, J. R.: Use of Oxygen-18 and Deuterium To Assess the Hydrology of Groundwater-Lake Systems, in: Environmental Chemistry of Lakes and Reservoirs, edited by: Baker, L. A., American  
490 Chemical Society, Washington, D.C., 67–90, <https://doi.org/10.1021/ba-1994-0237.ch003>, 1994.
- Kratz, T. K., Webster, K. E., Bowser, C. J., Magnuson, J. J., and Benson, B. J.: The influence of landscape position on lakes in northern Wisconsin, *Freshw. Biol.*, 37, 209–217, <https://doi.org/10.1046/j.1365-2427.1997.00149.x>, 1997.
- Lambrecht, N., Katsev, S., Wittkop, C., Hall, S. J., Sheik, C. S., Picard, A., Fakhraee, M., and Swanner, E. D.: Biogeochemical and physical controls on methane fluxes from two meromictic ferruginous lakes, *Geobiology*, 18, 54–69,

- 495 <https://doi.org/10.1111/gbi.12365>, 2020.
- Lascu, I. and Plank, C.: A new dimension to sediment magnetism: Charting the spatial variability of magnetic properties across lake basins, *Glob. Planet. Change*, 110, 340–349, <https://doi.org/10.1016/j.gloplacha.2013.03.013>, 2013.
- Lascu, I., McLauchlan, K. K., Myrbo, A., Leavitt, P. R., and Banerjee, S. K.: Sediment-magnetic evidence for last millennium drought conditions at the prairie–forest ecotone of northern United States, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 337–338, 500 99–107, <https://doi.org/https://doi.org/10.1016/j.palaeo.2012.04.001>, 2012.
- Ledesma, G., Islam, R., and Swanner, E. D.: Evaluation of preservation protocols for oxygen-sensitive minerals within laminated aquatic sediments, *Limnol. Oceanogr. Methods*, 21, 127–140, 2023.
- Ludlam, S. D. and Duval, B.: Natural and management-induced reduction in monimolimnetic volume and stability in a Coastal, Meromictic Lake, *Lake Reserv. Manag.*, 17, 71–81, <https://doi.org/10.1080/07438140109353975>, 2001.
- 505 McCarthy, F. M. G., Patterson, T., Head, M. J., Riddick, N. L., Cumming, B. F., Hamilton, P. B., Pisaric, M. F. J., Gushulak, C., Leavitt, P. R., Lafond, K. M., Llew-Williams, B., Marshall, M., Heyde, A., Pilkington, P. M., Moraal, J., Boyce, J. I., Nasser, N. A., Walsh, C., Garvie, M., Roberts, S., Rose, N. L., Cundy, A. B., Gaca, P., Milton, A., Hajdas, I., Crann, C. A., Boom, A., Finkelstein, S. A., and McAndrews, J. H.: The varved succession of Crawford Lake, Milton, Ontario, Canada as a candidate Global boundary Stratotype Section and Point for the Anthropocene series, *Anthr. Rev.*, 510 <https://doi.org/10.1177/20530196221149281>, 2023.
- McLauchlan, K. K., Myrbo, A., Lascu, I., and Leavitt, P. R.: Variable ecosystem response to climate change during the Holocene in northern Minnesota, USA, *GSA Bull.*, 125, 445–452, <https://doi.org/10.1130/B30737.1>, 2013.
- Megard, R. O., Bradbury, J. P., and Dean, W. E.: Climatic and limnologic setting of Elk Lake, in: *Elk Lake, Minnesota: Evidence for Rapid Climate Change in the North-Central United States*, edited by: Bradbury, J. P. and Dean, W. E., The Geological Society of America, 1993.
- 515 Meriläinen, J.: On the limnology of the meromictic Lake Valkiajärvi, in the Finnish Lake District, *Ann. Bot. Fenn.*, 7, 29–51, 1970.
- Lake Ice Out Dates: [https://www.dnr.state.mn.us/ice\\_out/index.html](https://www.dnr.state.mn.us/ice_out/index.html), last access: 25 August 2022.
- Minnesota Department of Natural Resources: Minnesota Spring Inventory, 520 [https://www.dnr.state.mn.us/waters/groundwater\\_section/springs/msi.html](https://www.dnr.state.mn.us/waters/groundwater_section/springs/msi.html), 2022.
- National Integrated Drought Information System:
- Nürnberg, G. K.: Quantifying anoxia in lakes, *Limnol. Oceanogr.*, 40, 1100–1111, <https://doi.org/10.4319/lo.1995.40.6.1100>, 1995.
- O’Sullivan, P. E.: Annually-laminated lake sediments and the study of Quaternary environmental changes — a review, *Quat. Sci. Rev.*, 1, 245–313, [https://doi.org/https://doi.org/10.1016/0277-3791\(83\)90008-2](https://doi.org/https://doi.org/10.1016/0277-3791(83)90008-2), 1983.
- 525 Pace, M. L. and Cole, J. J.: Synchronous variation of dissolved organic carbon and color in lakes, *Limnol. Oceanogr.*, 47, 333–342, <https://doi.org/10.4319/lo.2002.47.2.0333>, 2002.
- Reiter, C., Schmitz, J., and Helfman, S.: Itasca Biological Station Student Report 2318: A limnological study of Deming Lake,





University of Minnesota, 1998.

- 530 Rosenberry, D. O., Streigl, R. G., and Hudson, D. C.: Plants as indicators of focused ground water discharge to a northern Minnesota lake, *Ground Water*, 2000.
- Stelling, J. M., Sebestyen, S. D., Griffiths, N. A., Mitchell, C. P. J., and Green, M. B.: The stable isotopes of natural waters at the Marcell Experimental Forest, *Hydrol. Process.*, 35, 1–9, <https://doi.org/10.1002/hyp.14336>, 2021.
- Stewart, K. M., Walker, K. F., and Likens, G. E.: Meromictic Lakes, in: *Encyclopedia of Inland Waters*, edited by: Likens, G. E. B. T.-E. of I. W., Academic Press, Oxford, 589–602, <https://doi.org/10.1016/B978-012370626-3.00027-2>, 2009.
- Swanner, E. D., Lambrecht, N., Wittkop, C., Harding, C., Katsev, S., Torgeson, J., and Poulton, S. W.: The biogeochemistry of ferruginous lakes and past ferruginous oceans, *Earth-Science Rev.*, 211, 103430, <https://doi.org/10.1016/j.earscirev.2020.103430>, 2020.
- 540 Swanner, E. D., Lascu, I., Harding, C., Ledesma, G. M., Leung, T., and Akam, S. A.: Water properties of Arco Lake, Budd Lake, Deming Lake, and Josephine Lake in Itasca State Park from 2006-2009 and 2019-2023, v. 4, <https://doi.org/10.6073/pasta/eb8757dedf22ce40e79ca410c2ab9a34>, 2023.
- Watershed Boundary Dataset: <https://www.usgs.gov/national-hydrography/watershed-boundary-dataset>, last access: 20 May 2022.
- 545 Walker, K. F.: The stability of meromictic lakes in central Washington, *Limnol. Oceanogr.*, 19, 209–222, <https://doi.org/10.4319/lo.1974.19.2.0209>, 1974.
- Wetzel, R. G.: *Limnology: Lake and River Ecosystems*, Third., Academic Press, San Diego, 1006 pp., 2001.
- Wilkinson, G. M., Cole, J. J., Pace, M. L., Johnson, R. A., and Kleinmans, M. J.: Physical and biological contributions to metalimnetic oxygen maxima in lakes, *Limnol. Oceanogr.*, 60, 242–251, <https://doi.org/10.1002/lno.10022>, 2015.
- 550 Winslow, L. A., Read, J. S., Woolway, R., Brentrup, J. A., Leach, T. H., Zwart, J. A., Albers, S., and Collinge, D.: *rLakeAnalyzer*, 2019.
- Wright Jr., H. E.: History of the landscape in the Itasca region, in: *Elk Lake, Minnesota: Evidence for Rapid Climate Change in the North-Central United States*, edited by: Bradbury, J. P. and Dean, W. E., United States Geological Survey, Boulder, CO, 7–17, 1993.
- 555 Zadereev, E. S., Bohrer, B., and Gulati, R. D.: Introduction: Meromictic Lakes, Their Terminology and Geographic Distribution, in: *Ecology of Meromictic Lakes*, edited by: Gulati, R. D., Zadereev, E. S., and Degermendzhi, A. G., Springer International Publishing, Cham, 1–11, [https://doi.org/10.1007/978-3-319-49143-1\\_1](https://doi.org/10.1007/978-3-319-49143-1_1), 2017.




560 **Figure 1: Map of the four study lakes within Itasca State Park, Minnesota. Basemaps: USGS, ESRI.**






**Figure 2. Bathymetric maps of the study lakes overlain on the digital elevation models. Depth is in meters. The fence diagrams correspond to the cross-sections in Supplementary Figure 1.**

565




**Figure 3. Specific conductance profiles of a) Arco Lake, b) Budd Lake, c) Josephine Lake, and d) Deming Lake. For**

Deming Lake, seasons were classified according to solstice and equinox dates. Note the variability in the x and y axis scales.




570

**Figure 4. Hobo sensor data for temperature and conductance from Arco Lake, collected from May 2021 to May 2022 and from October 2021 to May 2022, respectively (top). Isotherm plot of the temperature data from the Hobo sensor (bottom).**




575

Figure 5. Piper Diagram of major cations and anions collected in May 2021.




580

Figure 6. Isotope data (points) and fit (dashed line) of an evaporation line ( $\delta^2\text{H}_{\text{H}_2\text{O}} = 4.5 \cdot \delta^{18}\text{O}_{\text{H}_2\text{O}} - 29.9$ ,  $r\text{-squared} = 0.93$ ) from the four study lakes plotted with a Local Meteoric Water Line (solid; Stelling et al., 2021). Crosses are spring or bog water data from Itasca State Park (Supplementary Table 2). The cross closest to the lake data points is the Deming bog sample, and the cross closest to the intersection of the LEL and LMWL is the Nicollet Creek sample. Analytical precision is within the symbol sizes.



585 **Figure 7. Multi-wavelength chlorophyll fluorescence (i.e., Fluoroprobe) measurements at a) Arco, b) Budd, c) Josephine, and d) Deming Lakes, showing seasonal and depth trends in major taxonomic groups. Note that the x scales vary between plots.**



590 **Figure 8. Dissolved oxygen measurements at a) Arco, b) Budd, c) Josephine, and d) Deming Lakes, showing seasonal and depth trends.**

**Table 1. Morphometric parameters for the four study lakes.**

Lake	Volume ( $m^3$ )	Surface Area ( $m^2$ )	Maximum Depth (m)	Relative Depth (%)	Fetch Distance (m)	Fetch Heading (degrees)
Arco	142,628.73 $m^3$	24180 $m^2$	12.6 m	7.3%	210.4 m	175 deg.
Budd	183,441.22 $m^3$	28184.5 $m^2$	16.1 m	8.8%	310.3 m	169 deg.
Deming	242,701.12 $m^3$	54325 $m^2$	20.8 m	7.3%	346.7 m	188 deg.
Josephine	315,765.05 $m^3$	52711 $m^2$	14.5 m	5.7%	432.4 m	136 deg.