

Uplift and denudation history of the Ellsworth Mountains: insights from low temperature thermochronology

Joaquín Bastías-Silva^{1,2,3}, David Chew¹, Fernando Poblete⁴, Paula Castillo⁵, William Guenther⁶, Anne Grunow⁷, Ian W.D. Dalziel⁸, Airton N. C. Dias⁹, Cristóbal Ramírez de Arellano¹⁰, Rodrigo Fernandez⁷

5

¹ Department of Geology, Trinity College Dublin, College Green, Dublin 2, Ireland.

² Escuela de Geología, Facultad de Ingeniería, Universidad Santo Tomás, Santiago, Chile.

³ Institute of Geochemistry and Petrology, Department of Earth Sciences, ETH Zürich, Switzerland.

⁴ Departamento de Geología, Universidad de Chile, Santiago.

10 ⁵ Institut für Geologie und Paläontologie, Westfälische Wilhelms-Universität, Münster, Germany.

⁶ Department of Geology, University of Illinois Urbana-Champaign, Urbana, IL, USA.

⁷ Byrd Polar and Climate Research Center, Ohio State University, Columbus, USA.

⁸ Institute for Geophysics, Jackson School of Geosciences, The University of Texas, USA.

⁹ Departamento de Física, Química e Matemática, CCTS, UFSCar - Campus Sorocaba, Brazil.

15 ¹⁰ Carrera Geología, Facultad de Ingeniería, Universidad Andres Bello, Chile.

Correspondence to: Joaquín Bastías-Silva (j.bastias.silva@gmail.com)

Abstract. While thermochronological studies have constrained the landscape evolution of several of the crustal blocks of West and East Antarctica, the tectono-thermal evolution of the Ellsworth Mountains remains relatively poorly constrained. These mountains are among the crustal blocks that comprise West Antarctica and exhibit an exceptionally well-preserved Palaeozoic sedimentary sequence. Despite the seminal contribution of Fitzgerald and Stump (1991), who suggested an Early Cretaceous uplift event for the Ellsworth Mountains, further thermochronological studies are required to improve the current understanding of the landscape evolution of this mountain chain. We present new zircon (U-Th)/He (ZHe) ages, which provide insights into the landscape evolution of the Ellsworth Mountains. The ZHe ages collected from near the base and the top of the sequence suggest that these rocks underwent burial reheating after deposition. A cooling event is recorded during the Jurassic–Early Cretaceous, which we interpret as representing exhumation in response to rock uplift of the Ellsworth Mountains. Moreover, our results show that, while ZHe ages at the base of the sequence are fully reset, towards the top ZHe are partially reset. Uplift and exhumation of the Ellsworth Mountains during the Jurassic–Early Cretaceous was contemporaneous with the rotation and translation of this crustal block with respect to East Antarctica and possibly the Antarctic Peninsula. Furthermore, this period is characterised by widespread extension associated with the disassembly and breakup of Gondwana, with the Ellsworth Mountains playing a key role in the opening of the far South Atlantic. Based on these results, we suggest that uplift of the Ellsworth Mountains during the disassembly of Gondwana provides additional evidence for major rearrangement of the crustal blocks between the South American, African, Australian and Antarctic plates. Finally, uplift of the Ellsworth Mountains commenced during the Jurassic, which predates the Early Cretaceous uplift of the Transantarctic Mountains. We suggest that the rift-related exhumation of the Ellsworth Mountains occurred throughout two events: (i) a Jurassic uplift associated with

35 the disassembly of Gondwana and (ii) an Early Cretaceous uplift related with the separation between Antarctica and Australia, which is also recorded in the Transantarctic Mountains.

1 Introduction

The Ellsworth Mountains extend for ~350 km between the Transantarctic Mountains and the Antarctic Peninsula and are ~50 km wide (Fig. 1). They are located within West Antarctica, which is composed of crustal blocks that amalgamated along the Pacific margin of Gondwana during the latest Precambrian to middle Phanerozoic (e.g. Dalziel and Elliot, 1982; Dalziel and Lawver 2001; Jordan et al., 2020; Riley et al., 2023). Furthermore, the Ellsworth Mountains form part of the most isolated and enigmatic crustal block (the Ellsworth-Whitmore Mountains block) in West Antarctica (Schopf, 1969; Dalziel and Elliot, 1982). Their exposures are comprised of a series of nunataks and mountains, which are dominated by the Heritage and Sentinel ranges (Fig. 1). Although the Antarctic icecap is intensively developed at these latitudes, the Ellsworth Mountains yield an extensive sedimentary record that extends from the Neoproterozoic to the Permian (Fig. 2; Craddock, 1969; Webers et al., 1992a; Castillo et al., 2017). Several studies have assessed the geological evolution of the Ellsworth Mountains along with its paleogeographic significance (Curtis et al., 1999; Curtis, 2001; Flowerdew et al., 2007; Dalziel, 2007; 2014, Castillo et al., 2017, Craddock et al., 2017). However, its tectonothermal evolution is relatively poorly constrained with the exception of the study of Fitzgerald and Stump (1991), who reported an Early Cretaceous uplift event based on apatite fission-track analyses. Most studies concerning the tectonothermal history of West Antarctica have instead been conducted on the Antarctic Peninsula (e.g. Guenther et al., 2010; Twinn et al., 2022; Bastias et al., 2022) and Thurston Island (e.g. Zundel et al., 2019). Further efforts to constrain the regional landscape evolution have been undertaken in the Transantarctic Mountains (e.g. Fitzgerald, 1994; Fitzgerald and Stump, 1997), which extends for ~3,000 km and divide East from West Antarctica (e.g. Goodge, 2020). To better understand the thermal evolution of the Ellsworth Mountains, we present herein new zircon (U-Th)/He data to constrain its thermal history and hence the formation of its present-day landscape. Furthermore, the (U-Th)/He is a thermochronometric system that is sensitive to low temperatures (Wolf et al., 1996) and has the potential to provide robust constraints on the thermal evolution of basins along with their subsequent exhumation histories (e.g. Ault et al., 2019; Dai et al., 2019).

2 Ellsworth Mountains

60 The Ellsworth Mountains hosts a stratigraphic sequence that spans the Palaeozoic era and is up to ~13 km thick (Fig. 2; e.g. Webers et al., 1992a; Castillo et al., 2017). At the base of the sequence is the lower Palaeozoic Heritage Group (Webers et al., 1992b), which consist of ~7.5 km of strata that are almost exclusively present in the Heritage Range (Fig. 2). They consist of sedimentary and volcanic rocks that were deposited in a rapidly subsiding basin (e.g. Curtis and Lomas, 1999). The group is composed, from base to top, by the Union Glacier, Hyde Glacier, Drake Icefall, Conglomerate Ridge, Liberty Hills, Springer

65 Peak, Frazier Ridge and Minaret formations (Fig. 2). The Union Glacier Formation includes continental volcanic and
volcaniclastic rocks (~3 km thick) and its age is constrained by a U-Pb zircon age from a Cambrian hyaloclastite (512 ± 14
Ma; personal communication, Rees et al., 1998). However, two metavolcaniclastic rocks from this formation yield U-Pb zircon
ages of ~675 Ma, raising questions as to the depositional age of this unit (Castillo et al 2017). The Hyde Glacier Formation
70 locally overlies the Union Glacier Formation and is composed of fluvial to shallow-marine deltaic deposits (Webers et al.,
1992b). Overlying these formations is the Drake Icefall Formation, comprised of black shales interbedded with limestones
deposited in a shallow-marine environment (Jago and Webers, 1992). Conglomeratic quartzite and polymictic conglomerates
of the Conglomerate Ridge Formation (Webers et al., 1992b) structurally overlie the Drake Icefall Formation with their contact
defined by a reverse fault. Three laterally equivalent formations overlie the Conglomerate Ridge Formation. Termed the
Springer Peak, Liberty Hills and Frazier Ridge formations. They are mostly clastic in composition and are comprised of
75 argillite, graywacke and quartzite (Webers et al., 1992b). Locally overlying these deposits is the Minaret Formation (Curtis
and Lomas, 1999), which is dominated by marble and carbonate rocks deposited during the Late Cambrian (e.g. Buggisch and
Webers, 1992; Jago and Webers, 1992). The Transitions Beds represent the uppermost unit of the Heritage Group and are
comprised of by a thin succession of sandstone interbedded with argillite (Spörli, 1992).

Overlying the Heritage Group is the Crashsite Group, a ~3 km-thick sequence dominated by quartzite, argillite conglomerate,
80 limestone and basic volcanic rocks (Fig. 2; Goldstrand et al., 1994; Spörli, 1992). This sequence is comprised from bottom to
top by the Howard Nunataks, Mount Liptak and Mount Wyatt Earp formations, which were deposited in a shallow-marine to
fluvial environment (Curtis and Lomas, 1999; Spörli, 1992). The age of the Crashsite Group has been constrained to the late
Cambrian–Devonian by trilobite faunas, sedimentation rates and detrital zircon ages (Shergold and Webers, 1992; Spörli,
1992; Webers et al., 1992b; Flowerdew et al., 2007).

85 The Crashsite Group is conformably overlain by the Whiteout Conglomerate (Fig. 2). These rocks are 1 km thick and
dominated by late Carboniferous to early Permian grey to black diamictites, which are associated with the Permo–
Carboniferous Gondwanan glaciation (Matsch and Ojakangas, 1992). Overlying the Whiteout Conglomerate is a 1 km thick
sequence of argillites, siltstone, sandstone and coal of the Polarstar Formation (Fig. 2; Collinson et al., 1992). Detrital zircons
geochronology implies this unit was deposited during the Permian (Elliot et al., 2016).

90 **3 Methods**

3.1 (U-Th)/He zircon thermochronology

Low-temperature thermochronometry is a robust method to constrain the time–temperature histories of rocks (Bargnesi et al.,
2016). The zircon (U-Th)/He system has a closure temperature to He diffusion of ~195-175 °C (Dodson, 1973), which provides
cooling ages that can be associated with shallow processes in the crust. Additionally, the robustness of zircon to weathering
95 and alteration during transport and diagenesis is particularly useful in clastic systems, such as the rocks exposed in the
Ellsworth Mountains. Therefore zircon (U-Th)/He dating is a powerful tool to constrain the thermal evolution of a given rock.

Zircon separates were previously prepared for the U-Pb geochronology and Hf isotope studies presented in Castillo et al. (2017), which applied standard separation procedures. Two to three single-grain aliquots from each sample were selected for (U-Th)/He analysis (Table 1). Zircon (U-Th)/He analytical methods followed those described in Guenther et al. (2016). The details of the data reduction is shared in the Supplementary Files.

Helium extraction and analysis consisted of in vacuo diode laser heating, cryogenic purification and quadrupole mass-spectrometry on a Pfeiffer Prisma Plus at the University of Illinois. Zircon dissolution was followed by U and Th analysis via isotope-dilution inductively coupled plasma-mass spectrometry on a Thermo Element2 at the University of Arizona. Dimension measurements for zircon were collected for both the alpha ejection correction and to calculate eU concentrations. The alpha ejection correction employed the equations of Hourigan et al. (2005) and Farley (2002), with U and Th specific ejection values as listed in Farley (2002, Table 1). Depending on the degree of abrasion, one of two equations was used: tetragonal prism with pyramidal terminations (when terminations are present and measurable) or prolate spheroid (when terminations are absent). Further detail on the results of the samples and standards analysed are presented in the Table 1.

4 Results

4.1 Heritage Group

We selected three samples from the Heritage Group (Fig. 2). The samples (13EG01, 13EG05 and EHD0801A) are located in the Heritage Range in the southern sector of the Ellsworth Mountains, with an altitude that ranges from ~1450 to 990m (Table 1). Sample 13EG01 is a sandstone collected from the base of the early to middle Cambrian Union Glacier Formation and yielded ZHe ages of 179, 156 and 140 Ma (Fig. 3). The middle to late Cambrian Springer Peak Formation (Jago and Webers, 1992; Shergold and Webers, 1992; Randall et al., 2000) is part of the upper section of the Heritage Group, from which we analysed the sandstone sample 13EG05 which yielded ZHe ages of 184, 170 and 158 Ma (Fig. 3). The Springer Peak Formation is in lateral contact with the middle to late Cambrian Liberty Hill Formation. ZHe ages from a sandstone (sample EHD0801A) from Liberty Hill Formation yield ages of 150, 149 and 103 Ma (Fig. 3). While most of the ZHe ages from the Heritage Group are Jurassic (from 184 to 149 Ma) two zircon grains yielded Early Cretaceous ages (140 and 103 Ma). The two younger ages are found in both the base and the upper parts of the Heritage Group, in the Union Glacier and Liberty Hills formations, respectively (Fig. 3).

4.2 Whiteout Conglomerate

Two samples of matrix from conglomeratic sandstones were analysed for ZHe ages from the Whiteout Conglomerate, samples 13EG10 and 13EG15, which are located in the upper and base of this sequence, respectively. This unit was deposited during the Permian-Carboniferous (Collison et al., 1992; Matsch and Ojakangas, 1992) and is part of the upper section of the stratigraphic succession exposed in the Ellsworth Mountains. Sample 13EG10 was collected from the Whiteout Nunatak in the Sentinel Range in the northern section of the Ellsworth Mountains (Fig. 2) and yielded a ZHe age of 182 Ma. A second

rock was analysed (13EG15) from this unit to the south in the Heritage Range and to the east of the samples analysed from the Heritage Group (Fig. 2). ZHe ages from sample 13EG15 are 791, 468 and 159 Ma (Table 1). While two grains yielded ages older than the depositional age of this unit (~360 to 300 Ma), they agree with the provenance studies collected from the same sample and presented by Castillo et al. (2017), who showed the presence of Palaeozoic, Neoproterozoic and Mesoproterozoic sources in the Whiteout Conglomerate. Therefore, these results suggest that the ZHe system in the Whiteout Conglomerate is partially reset. There is a significant distance (~250 km) between the two samples of the Whiteout Conglomerate (Fig. 2); their altitude is 1520 and 580 m for 13EG10 and 13EG15, respectively (Table 1).

135 **5 Discussion**

5.1 Landscape evolution

Our results predominantly indicate Jurassic–Early Cretaceous ZHe ages from the Ellsworth Mountains. These results help to constrain the landscape evolution of the Ellsworth Mountains, which is thought to have experienced an uplift event during the Early Cretaceous (Fitzgerald and Stump, 1991) based on apatite fission-track analyses. The Jurassic–Early Cretaceous ZHe ages are significantly younger than the age of the host sedimentary rocks, indicating the ZHe ages have been reset. While this indicates a post-depositional thermal event, the heat source and the causative mechanism is not clear. The Palaeozoic sedimentary sequence of the Ellsworth Mountains, although faulted and deformed (e.g. Curtis, 2001), has not experienced significant regional metamorphism. Furthermore, sedimentary analysis shows that significant stratigraphic repetition related to tight folding or thrust faults is not likely (e.g. Collison et al., 1992; Matsch and Ojakangas, 1992; Spörli, 1992; Webers et al., 1992a). The sedimentary sequence exposed in the Ellsworth Mountains has a thickness of ~13 km which can account for the resetting of the ZHe ages by burial alone, assuming a geothermal gradient of 30°C km⁻¹ and a ZHe partial retention zone in the range of ~200–130°C (Wolfe and Stockli, 2010). Therefore, burial heating associated with deposition of the sequence exposed in the Ellsworth Mountains may account for the resetting of the ZHe ages. It is noteworthy that the heat flux distribution in West Antarctica is relatively complex (e.g. Martos et al., 2017). The samples are distributed along a distance of ~300 km (Fig. 2), which allows for extrapolation to a regional scope, but the distance is insufficient to consider a significant change between the geothermal gradient value of each sample. We nevertheless acknowledge that the chosen value may vary as there is a better understanding of the paleo-geothermal gradient.

While all the ZHe ages in the Heritage Group range from the Jurassic to the Early Cretaceous, only two of the four grains from the Whiteout Conglomerate are within that age range. Furthermore, the two older grains from the Whiteout Conglomerate yield ZHe dates that are concordant with their detrital U-Pb ages, which were presented in Castillo et al. (2017), who dated the same samples (13EG10 and 13EG15). This suggest only partial ZHe resetting occurred in the Whiteout Conglomerate and burial heating did not reset all ZHe ages. A lesser degree of burial heating in the Whiteout Conglomerate compared to the Heritage Group is in agreement with their respective positions in the stratigraphic sequence, in which they are towards the top and the base of the section, respectively (Fig. 2). Zircon grains may not be fully reset in a given rock if they experience

160 reheating below the range of temperatures of the ZHe partial retention zone (e.g. Schneider and Issler, 2019; Malusà and
Fitzgerald, 2019) as in the Whiteout Conglomerate. Conversely, zircon grains of the Heritage Group were fully reset because
they experienced temperatures above the ZHe partial retention zone. Taking into the account that (i) the partial retention zone
for low to moderately damaged zircon grains for the (U-Th)/He system is in the range of ~200-130°C (Wolfe and Stockli,
2010) and (ii) assuming a geothermal gradient of 30°C km⁻¹; the Whiteout Conglomerate may have experienced between 7 –
165 4 km of burial. The Heritage Group experienced temperatures above ~200°C and burial by at least 7 – 6 km.

5.2 Gondwana fragmentation

Several tectono-magmatic events preceded the formation of oceanic lithosphere that led to the fragmentation of Gondwana
and one key tectono-magmatic event is recorded in the Ellsworth Mountains. Dalziel et al. (2013 and references therein) argued
that the key to understanding Gondwana's initial fragmentation in the South Atlantic-Weddell Sea region is the opposed sense
170 of rotations of the Falklands/Malvinas Plateau and the Ellsworth-Whitmore Mountains block. These rotations were termed by
Martin (2007) as 'double-saloon-door' tectonics and ascribed by them to seafloor spreading above a curved and retreating
subduction zone. This event has been interpreted as an extensional episode that followed the emplacement of the Karoo and
Ferrar LIPs at ~184-182 Ma (e.g. Svensen et al., 2012; Burgess et al., 2015; Greber et al., 2020) and coincides with the early
development of the silicic magmatism of the Chon Aike province in the Antarctic Peninsula and Patagonia (Pankhurst et al.,
175 2000; Bastias et al., 2021).

Grunow et al. (1987) presented a thorough revision of the paleogeographic evolution of the Ellsworth-Whitmore Mountains
block based on paleomagnetic constraints. They suggested that the Ellsworth-Whitmore Mountains block and the Antarctic
Peninsula have undergone little relative movement since the Middle Jurassic. Furthermore, they suggested that along with
Thurston Island, the Ellsworth-Whitmore Mountains block and the Antarctic Peninsula define a single entity termed
180 'Weddellia'. Between the Middle Jurassic and Early Cretaceous, these crustal blocks remained attached to West Gondwana,
while East Antarctica moved southward (dextrally) relative to Weddellia. The present-day position of the Ellsworth-Whitmore
Mountains block was attained during the Early and mid-Cretaceous by clockwise rotation of Weddellia along with a sinistral
movement relative to East Antarctica. Randall and MacNiocaill (2004) investigated the paleopositions of the Ellsworth-
Whitmore Mountains block prior to the break-up of Gondwana, again based on paleomagnetic studies. Their findings suggest
185 that the Ellsworth-Whitmore Mountains block was located near the junction of East Antarctica and Africa. However, Castillo
et al. (2017), employing provenance studies, suggested a closer affinity of the Ellsworth-Whitmore Mountains block to the
Australo-Antarctic plate and located this crustal block further east than that proposed by Randall and MacNiocaill (2004).
Nevertheless, prior to the disassembly of Gondwana, all these studies place the Ellsworth-Whitmore Mountains block in the
vicinity of the margin of the junction between East and West Gondwana, as proposed by Schopf (1969).

190 The sequence exposed in the Ellsworth Mountains experienced burial reheating after Palaeozoic deposition. The ZHe ages
presented here yield predominantly Jurassic and Early Cretaceous dates. This suggests that these rocks cooled through the
~200-130°C ZHe partial retention zone during the Jurassic and Early Cretaceous, which we interpret as exhumation related to

a specific rock uplift event. Although our dataset does not contradict the seminal work of Fitzgerald and Stump (1991), who reported apatite fission-track ages ranging from ~141–117 Ma, with the exception of two zircons yielding Early Cretaceous, most of the ZHe presented herein predates 141 Ma. This suggests that our results provide an older age (Jurassic-Early Cretaceous) for the uplift of the Ellsworth Mountains than what was previously reported (Early Cretaceous; Fitzgerald and Stump, 1991). While this may be simply explained by the relatively higher thermal sensitivity of ZHe (~200-130°C; Wolfe and Stockli, 2010) compared to that of apatite fission-tracks (~120-60°C; Fleischer et al., 1965; Green et al., 1985), and therefore ZHe ages may potentially yield older ages than those of apatite fission-tracks, it also suggests that the Ellsworth Mountains may have uplifted earlier than previously considered, during the Jurassic. Nevertheless, this Jurassic-Early Cretaceous episode is recorded in both the Sentinel and Heritage ranges (Fig. 2), implying the presence of a regional event that affected the Ellsworth Mountains. Although the structures that uplifted these rocks are poorly constrained, the Jurassic and Early Cretaceous is dominated by widespread extension and magmatism associated with the disassembly of Gondwana (e.g. Dalziel et al., 2013; Jordan et al., 2017; Pankhurst et al., 2000; Bastias et al., 2021). We suggest that the uplift event responsible for the exhumation of the Ellsworth Mountains sequences is also part of the major plate reconfiguration associated with the break-up of Gondwana. An extensional setting prevailed in this sector of Gondwana during the Jurassic and Cretaceous (e.g. Dalziel et al., 2013) and therefore, we suggest that the Ellsworth Mountains were uplifted through this deformative event. However, we acknowledge that likely the Ellsworth Mountains were still being accommodated along West Antarctica during the Jurassic–Late Cretaceous (Grunow et al., 1987) and the uplift may have been also caused by the transtensional movement. Nevertheless, the widespread evidence of extensional tectonics in this sector of Gondwana during the Jurassic-Late Cretaceous favours an extensional mechanism.

5.3 Connection with the Transantarctic Mountains

The Transantarctic Mountains extend for ~3,200 km from northern Victoria Land area of the Australian-New Zealand sector in Antarctica, to the Pensacola Mountains near the Ronne Ice Shelf (Fig. 1). They rise to elevations of >4,500 m directly from sea level along the Ross Sea coastline (e.g. Goodge, 2020). This mountain chain is a major feature of the Earth landscape, as it is the longest intraplate mountain belt. It defines the limit between East and West Antarctica, a thick stable craton and a large accretionary province, respectively (e.g. Goodge, 2020; Jordan et al., 2020). Furthermore, the Transantarctic Mountains are the world’s largest rift mountain system (e.g. Goodge, 2020). Although the Ellsworth Mountains are geographically separated from the Transantarctic Mountains, both contain Precambrian and Palaeozoic rocks of similar provenance affinity (e.g. Schopf, 1969; Bradshaw, 2013) and thus they have been often correlated (e.g. Goodge, 2020). The dataset presented herein suggests that while the Transantarctic Mountains and the Ellsworth Mountains have similar rocks, their uplift may be associated with a different tectonic event. Furthermore, the thermal-tectonic history of the Transantarctic Mountains shows three major exhumation events, which occurred during the Early Cretaceous, Late Cretaceous and Cenozoic (Fitzgerald et al., 2002). These events have been associated with regional tectonic events, which are (i) the initial separation between Antarctica and Australia during the Early Cretaceous, (ii) Late Cretaceous extension (main phase) between West and East Antarctica and (iii) the Cenozoic southward seafloor propagation of the Adare Trough into the Ross Sea (e.g. Fitzgerald and Gleadow, 1988; Fitzgerald, 1992, 1994, 2002; Balestrieri et al., 1997; Miller et al., 2010; Goodge, 2020). The ZHe results presented herein consistently yield Jurassic ages (nine out of eleven; Fig. 3), which predate the first uplift of the Antarctic Mountains during the Early Cretaceous (e.g. Goodge, 2020). Hence, although there is some overlap in the timing of exhumation

235 between these two mountain chains, it also suggests that the uplift of the Ellsworth Mountains may be older. Therefore,
the historical correlation between the Ellsworth and Transantarctic Mountains may not be correct, at least on their
uplift histories. While the uplift of the Ellsworth Mountains and the Transantarctic Mountains may be part of a
diachronous exhumation event whereby uplift commenced in the Ellsworth Mountains during the Jurassic and
advanced progressively towards the south, propagating into the Transantarctic Mountains during the Early
240 Cretaceous, such a tectonic event remains unlikely considering the Gondwana breakup clockwise rotations during this
period in Africa, India, Australia and New Zealand (e.g. Jokat et al., 2003). Alternatively, the Jurassic-Early Cretaceous
uplift of the Ellsworth Mountains may be associated with two different tectonic episodes. First, a Jurassic uplift event
occurred associated with the disassembly of southwestern Gondwana, which is exclusively recorded in the Ellsworth
Mountains. A second exhumation event occurred during the Early Cretaceous in both the Ellsworth Mountains and
the Transantarctic Mountains. This event corresponds to the first uplift of the Transantarctic Mountains, which is
associated with the separation of Antarctica and Australia (e.g. Goode, 2020).6 Conclusions

The Palaeozoic stratigraphic sequence exposed in the Ellsworth Mountains consistently yield Jurassic and Early Cretaceous
ZHe ages, which we interpret as exhumation of the Ellsworth Mountains as a direct response to rock uplift. The stratigraphic
245 succession reached temperatures within or above the ZHe partial retention zone (~200-130°C; Wolfe and Stockli, 2010) by
burial reheating associated with the ~13 km thick stratigraphic column, assuming a geothermal gradient of 30°C km⁻¹.
However, while all zircon grains from the Heritage Group yield reset ZHe ages, the Whiteout Conglomerate also yielded non-
reset ZHe ages. Additionally, these non-reset ZHe ages are broadly contemporaneous with the ages of the detrital material of
this unit presented by Castillo et al. (2017) on the same samples. This suggests that the temperature associated with burial
250 reheat was, as would be expected, progressively higher towards the base of the sequence exposed in the Ellsworth Mountains.
The Jurassic–Early Cretaceous uplift of the Ellsworth Mountains is older than the Early Cretaceous exhumation of the
Transantarctic Mountains. We suggest that the widespread extension that dominated this sector of Gondwana and related to
its fragmentation during the Jurassic and Cretaceous, is also responsible of the uplift of the Ellsworth Mountains. While the
first uplift event recorded in the Transantarctic Mountains, which occurred during the Early Cretaceous, is also present in the
255 Ellsworth Mountains, the latter also yield evidence of an older and independent Jurassic uplift episode, which we tentatively
associate with the disassembly of Gondwana during this period.

Acknowledgements

This study was financed by project RT-4418 funded by INACH (Chilean Antarctic Survey). JB was funded by the Swiss
National Science Foundation (project P5R5PN_217947). DC acknowledges support from Science Foundation Ireland (SFI)
260 under Grant Number 13/RC/2092 and 13/RC/2092_P2 (SFI Research Centre in Applied Geosciences, iCRAG). Kei Ogata is
thanked for editorial handling and the authors are grateful to tow anonymous reviewers for providing constructive criticism
that improved the paper.

References

- Ault, A.K., Gautheron, C., and King, G.E. 2019. Innovations in (U–Th)/He, fission track, and trapped charge
265 thermochronometry with applications to earthquakes, weathering, surface-mantle connections, and the growth and decay
of mountains: *Tectonics*, v. 38, p. 3705–3739, <https://doi.org/10.1029/2018TC005312>.
- Balestrieri, M.L., Bigazzi, G., Ghezzi, C. 1997. Uplift-denudation of the Transantarctic Mountains between the David and the
Mariner glaciers, northern Victoria Land (Antarctica); constraints by apatite fission-track analysis, in VII international
symposium on Antarctic earth sciences, Siena. Italy 547–554.
- 270 Bargnesi, E. A., Stockli, D. F., Hourigan, J. K., Hager, C. 2016. Improved accuracy of zircon (U–Th)/He ages by rectifying
parent nuclide zonation with practical methods. *Chemical Geology*, v. 426, p. 158–169.
<https://doi.org/10.1016/j.chemgeo.2016.01.017>.
- Bastías, J., Spikings, R., Riley, T., Ulianov, A., Grunow, A., Chiaradia, M., & Hervé, F. 2021. A revised interpretation of the
Chon Aike magmatic province: active margin origin and implications for the opening of the Weddell Sea. *Lithos*, 386,
275 106013.
- Bastias, J., Spikings, R., Riley, T., Chew, D., Grunow, A., Ulianov, A., Chiaradia, M., & Burton-Johnson, A. 2023. Cretaceous
magmatism in the Antarctic Peninsula and its tectonic implications. *Journal of the Geological Society*, 180(1).Bradshaw,
M.A. 2013. The Taylor Group (Beacon Supergroup): the Devonian sediments of Antarctica, in Hambrey, M. J. et al., eds.,
Antarctic Palaeoenvironments and Earth-Surface Processes: Geological Society, London. Special Publications 381, 67–97.
- 280 Buggisch, W., and Webers, G.F. 1992. Facies of Cambrian carbonate rocks, Ellsworth Mountains, West Antarctica, in Webers,
G.F., Craddock, C., and Spletstoesser, J.F., eds., *Geology and Paleontology of the Ellsworth Mountains, West Antarctica:*
Geological Society of America Memoir 170, p. 81–100.
- Burgess, S.D., Bowring, S.A., Fleming, T.H., Elliot, D.H. 2015. High-precision geochronology links the Ferrar large igneous
province with early-Jurassic Ocean anoxia and biotic crisis. *Earth Planet. Sci. Lett.* 415, 90–99.
- 285 Castillo, P., Fanning, C. M., Fernandez, R., Poblete, F., & Hervé, F. 2017. Provenance and age constraints of Paleozoic
siliciclastic rocks from the Ellsworth Mountains in West Antarctica, as determined by detrital zircon geochronology. *GSA*
Bulletin, 129(11-12), 1568-1584.
- Collinson, J.W., Vavra, C.L., and Zawiskie, J.M. 1992. Sedimentology of the Polarstar Formation (Permian), Ellsworth
Mountains, West Antarctica, in Webers, G.F., Craddock, C., and Spletstoesser, J.F., eds., *Geology and Paleontology of*
290 *the Ellsworth Mountains, West Antarctica: Geological Society of America Memoir 170*, p. 63–79.
- Craddock, C. 1969. Geological Map of Antarctica. Sheet 4. Ellsworth Mountains 1: 1,000,000. American Geographical
Society.
- Craddock, J.P., Fitzgerald, P., Konstantinou, A., Nereson, A., Thomas, R.J. 2017. Detrital zircon provenance of upper
Cambrian-Permian strata and tectonic evolution of the Ellsworth Mountains, West Antarctica: *Gondwana Research*, v. 45,
295 p. 191–207.

- Curtis, M.L., 2001, Tectonic history of the Ellsworth Mountains, West Antarctica: Reconciling a Gondwana enigma: Geological Society of America Bulletin, v. 113, no. 7, p. 939–958.
- Curtis, M.L., and Lomas, S.A. 1999. Late Cambrian stratigraphy of the Heritage Range, Ellsworth Mountains: Implications for basin evolution: Antarctic Science, v. 11, no. 1, p. 63–77.
- 300 Curtis, M.L., Leat, P.T., Riley, T.R., Storey, B.C., Millar, I.L., and Randall, D.E. 1999. Middle Cambrian rift-related volcanism in the Ellsworth Mountains, Antarctica: Tectonic implications for the palaeo-Pacific margin of Gondwana: Tectonophysics, v. 304, no. 4, p. 275–299.
- Dai, J. G., Fox, M., Shuster, D. L., Hourigan, J., Han, X., Li, Y. L., & Wang, C. S. 2020. Burial and exhumation of the Hoh Xil Basin, northern Tibetan Plateau: Constraints from detrital (U-Th)/He ages. Basin Research, 32(5), 894-915.
- 305 Dalziel, I. W., & Elliot, D. H. 1982. West Antarctica: problem child of Gondwanaland. Tectonics, 1(1), 3-19.
- Dalziel, I. W. D., & Lawver, L. A. 2001. The lithospheric setting of the West Antarctic ice sheet. The West Antarctic ice sheet: behavior and environment, 77, 29-44.
- Dalziel, I. W., Lawver, L. A., Norton, I. O., & Gahagan, L. M. 2013. The Scotia Arc: genesis, evolution, global significance. Annual Review of Earth and Planetary Sciences, 41, 767-793.
- 310 Dalziel, I.W.D. 2007. The Ellsworth Mountains: Critical and enduringly enigmatic, in Cooper, A.K., and Raymond, C.R., eds., Antarctica: A Keystone in a Changing World: U.S. Geological Survey Open-File Report 2007-1047, Short Research Paper 004, 5 p., doi:10.3133/of2007-1047.srp004.
- Dalziel, I.W.D. 2014. Cambrian transgression and radiation linked to an Iapetus-Pacific oceanic connection?: Geology, v. 42, no. 11, p. 979–982, doi:10.1130/G35886.1.
- 315 Dodson, M.H. 1973. Closure temperature in cooling geochronological and petrological systems. Contrib Mineral Petrol 40: 259-274.
- Elliot, D.H., Fanning, C.M., and Laudon, T.S. 2016. The Gondwana plate margin in the Weddell Sea sector: Zircon geochronology of Upper Paleozoic (mainly Permian) strata from the Ellsworth Mountains and eastern Ellsworth Land, Antarctica: Gondwana Research, v. 29, no. 1, p. 234–247.
- 320 Farley, K. A. 2002. (U-Th)/He dating: Techniques, calibrations, and applications. Reviews in Mineralogy and Geochemistry, 47(1), 819-844.
- Fitzgerald, P.G. 1992. The Transantarctic Mountains of southern Victoria Land: the application of apatite fission track analysis to a rift shoulder uplift. Tectonics 11, 634–662.
- Fitzgerald, P.G. 1994. Thermochronologic constraints on post-Paleozoic tectonic evolution of the central Transantarctic Mountains, Antarctica. Tectonics 13, 818–836.
- 325 Fitzgerald, P.G. 2002. Tectonics and landscape evolution of the Antarctic plate since the breakup of Gondwana, with an emphasis on the West Antarctic Rift System and the Transantarctic Mountains. In: Gamble, J., Skinner, D.A., Henrys, S. (Eds.), Antarctica at the Close of a Millennium, Proceedings of the 8th International Symposium on Antarctic Earth Science: Wellington. 35, pp. 453–469 Royal Society of New Zealand Bulletin.

- 330 Fitzgerald, P.G., Gleadow, A.J.W. 1988. Fission-track geochronology, tectonics and structure of the Transantarctic Mountains in Northern Victoria Land, Antarctica. *Isotope Geoscience* 73, 169–198.
- Fitzgerald, P. G., & Stump, E. 1991. Early cretaceous uplift in the Ellsworth Mountains of West Antarctica. *Science*, 254(5028), 92-94.
- Fitzgerald, P.G., Stump, E. 1997. Cretaceous and Cenozoic episodic denudation of the Transantarctic Mountains, Antarctica:
335 New constraints from apatite fission track thermochronology in the Scott Glacier region. *Journal of Geophysical Research* 102 (B4), 7747–7765.
- Fleischer, R. L., Price, P. B., & Walker, R. M. 1965. Tracks of Charged Particles in Solids: These tracks are becoming useful tools in many fields of science and engineering. *Science*, 149(3682), 383-393.
- Flowerdew, M.J., Millar, I.L., Curtis, M.L., Vaughan, A.P.M., Horstwood, M.S.A., Whitehouse, M.J., and Fanning, C.M.
340 2007. Combined U-Pb geochronology and Hf isotope geochemistry of detrital zircons from early Paleozoic sedimentary rocks, Ellsworth-Whitmore Mountains block, Antarctica: *Geological Society of America Bulletin*, v. 119, no. 3-4, p. 275–288, doi:10.1130/B25891.1.
- Ghidella, M. E., Yáñez, G., & LaBrecque, J. L. 2002. Revised tectonic implications for the magnetic anomalies of the western Weddell Sea. *Tectonophysics*, 347(1-3), 65-86.
- 345 Goldstrand, P.M., Fitzgerald, P.G., Redfield, T.F., Stump, E., and Hobbs, C. 1994. Stratigraphic evidence for the Ross orogeny in the Ellsworth Mountains, West Antarctica: Implication for the evolution of the paleo-Pacific margin of Gondwana: *Geology*, v. 22, p. 427–430, doi:10.1130/0091-7613(1994)022<0427:SEFTRO>2.3.CO;2.
- Goodge, J. W. 2020. Geological and tectonic evolution of the Transantarctic Mountains, from ancient craton to recent enigma. *Gondwana Research*, 80, 50-122.
- 350 Greber, N.D., Davies, J.H., Gaynor, S., Jourdan, F., Bertrand, H., Schaltegger, U. 2020. New high precision U-Pb ages and Hf isotope data from the Karoo large igneous province; implications for pulsed magmatism and early Toarcian environmental perturbations. *Results Geochem.*, 100005. <https://doi.org/10.1016/j.ringeo.2020.100005>.
- Green, P. F., Duddy, I. R., Gleadow, A. J. W., Tingate, P. R., & Laslett, G. M. 1985. Fission-track annealing in apatite: track length measurements and the form of the Arrhenius plot. *Nuclear Tracks and Radiation Measurements* (1982), 10(3), 323-
355 328.
- Grunow, A.M., Kent, D.V., and Dalziel, I.W.D. 1987. Mesozoic evolution of West Antarctica and the Weddell Sea Basin: New paleomagnetic constraints: *Earth and Planetary Science Letters*, v. 86, no. 1, p. 16–26, doi:10.1016/0012-821X(87)90184-1.
- Guenther, W. R., Barbeau Jr, D. L., Reiners, P. W., & Thomson, S. N. 2010. Slab window migration and terrane accretion
360 preserved by low-temperature thermochronology of a magmatic arc, northern Antarctic Peninsula. *Geochemistry, Geophysics, Geosystems*, 11(3).
- Guenther, W. R., Reiners, P. W., & Chowdhury, U. 2016. Isotope dilution analysis of Ca and Zr in apatite and zircon (U-Th)/He chronometry. *Geochemistry, Geophysics, Geosystems*, 17(5), 1623-1640.

- Hourigan, J. K., Reiners, P. W., & Brandon, M. T. 2005. U-Th zonation-dependent alpha-ejection in (U-Th)/He chronometry. *Geochimica et Cosmochimica Acta*, 69(13), 3349-3365.
- 365
- Jago, J.B., and Webers, G.F. 1992. Middle Cambrian trilobites from the Ellsworth Mountains, West Antarctica, in Webers, G.F., Craddock, C., and Spletstoeser, J.F., eds., *Geology and Paleontology of the Ellsworth Mountains, West Antarctica: Geological Society of America Memoir 170*, p. 101–124, doi:10.1130/MEM170-p101
- Jokat, W., Boebel, T., König, M., & Meyer, U. 2003. Timing and geometry of early Gondwana breakup. *Journal of Geophysical Research: Solid Earth*, 108(B9).
- 370
- Jordan, T. A., Ferraccioli, F., & Leat, P. T. 2017. New geophysical compilations link crustal block motion to Jurassic extension and strike-slip faulting in the Weddell Sea Rift System of West Antarctica. *Gondwana Research*, 42, 29-48.
- Jordan, T. A., Riley, T. R., & Siddoway, C. S. 2020. The geological history and evolution of West Antarctica. *Nature Reviews Earth & Environment*, 1(2), 117-133.
- 375
- König, M., & Jokat, W. 2006. The Mesozoic breakup of the Weddell Sea. *Journal of Geophysical Research: Solid Earth*, 111(B12).
- Malusà, M. G., & Fitzgerald, P. G. (Eds.). 2019. *Fission-track thermochronology and its application to geology*. Switzerland: Springer International Publishing.
- Martin AK. 2007. Gondwana breakup via double-saloon-door rifting and seafloor spreading in a backarc basin during subduction rollback. *Tectonophysics* 445:245–72.
- 380
- Martos, Y. M., Catalán, M., Jordan, T. A., Golynsky, A., Golynsky, D., Eagles, G., & Vaughan, D. G. 2017. Heat flux distribution of Antarctica unveiled. *Geophysical Research Letters*, 44, 11,417–11,426. <https://doi.org/10.1002/2017>.
- Matsch, C. L., & Ojakangas, R. W. 1992. Stratigraphy and sedimentology of the Whiteout Conglomerate; an upper Paleozoic glacial unit, Ellsworth Mountains, West Antarctica.
- 385
- Miller, S.R., Fitzgerald, P.G., Baldwin, S.L. 2010. Cenozoic range-front faulting and development of the Transantarctic Mountains near Cape Surprise, Antarctica; thermochronologic and geomorphologic constraints. *Tectonics* 29, TC1003 (21 pp). <https://doi.org/10.1029/2009TC002457>.
- Pankhurst, R.J., Riley, T.R., Fanning, C.M., Kelley, S.P. 2000. Episodic silicic volcanism in Patagonia and the Antarctic Peninsula: chronology of magmatism associated with the breakup of Gondwana. *J. Petrol.* 41, 603–625.
- 390
- Randall, D.E., and Mac Niocaill, C. 2004. Cambrian palaeomagnetic data confirm a Natal Embayment location for the Ellsworth-Whitmore Mountains, Antarctica, in *Gondwana reconstructions: Geophysical Journal International*, v. 157, no. 1, p. 105–116, doi:10.1111/j.1365-246X.2004.02192.x.
- Randall, D.E., Curtis, M.L., and Millar, I.L. 2000. A new late middle Cambrian paleomagnetic pole for the Ellsworth Mountains, Antarctica: *The Journal of Geology*, v. 108, p. 403–425, doi:10.1086/314408.
- 395
- Rees, M.N., Smith, E.I., Duebendorfer, E.M., and Keenan, D. 1998. Cambrian marginal basin rifting and subduction recorded in the Ellsworth-Whitmore Mountains terrane, West Antarctica, in Almond, J., et al., eds., *Gondwana 10: Event Stratigraphy of Gondwana: Journal of African Earth Sciences*, v. 27, no. 1, supplement 1, p. 151–153.

- Reiners, P. W. (2005). Zircon (U-Th)/He thermochronometry. *Reviews in mineralogy and geochemistry*, 58(1), 151-179.
- Riley, T. R., Burton-Johnson, A., Flowerdew, M. J., Poblete, F., Castillo, P., Hervé, F., Leat, P., Millar, I., Bastias, J., &
400 Whitehouse, M. J. (2023). Palaeozoic–Early Mesozoic geological history of the Antarctic Peninsula and correlations with Patagonia: Kinematic reconstructions of the proto-Pacific margin of Gondwana. *Earth-Science Reviews*, 236, 104265.
- Schneider, D. A., & Issler, D. R. 2019. Application of low-temperature thermochronology to hydrocarbon exploration. *Fission-Track Thermochronology and its Application to Geology*, 315-333.
- Schopf, J. M. 1969. Ellsworth Mountains: position in West Antarctica due to sea-floor spreading. *Science*, 164(3875), 63-66.
- 405 Shergold, J.H., and Webers, G.F. 1992. Late Dresbachian (Idamean) and other trilobite faunas from the Heritage Range, Ellsworth Mountains, West Antarctica, in Webers, G.F., Craddock, C., and Spletstoesser, J.F., eds., *Geology and Paleontology of the Ellsworth Mountains, West Antarctica: Geological Society of America Memoir 170*, p. 125–168, doi:10.1130/MEM170-p125.
- Spörli, K.B. 1992. Stratigraphy of the Crashsite Group, Ellsworth Mountains, West Antarctica, in Webers, G.F., Craddock, C.,
410 and Spletstoesser, J.F., eds., *Geology and Paleontology of the Ellsworth Mountains, West Antarctica: Geological Society of America Memoir 170*, p. 21–36, doi:10.1130/MEM170-p21.
- Svensen, H., Corfu, F., Polteau, S., Hammer, O., Planke, S. 2012. Rapid magma emplacement in the Karoo large igneous province. *Earth Planet. Sci. Lett.* 325-326, 1–9. <https://doi.org/10.1016/j.epsl.2012.01.015>.
- Twinn, G., Riley, T., Fox, M., & Carter, A. 2022. Thermal history of the southern Antarctic Peninsula during Cenozoic oblique
415 subduction. *Journal of the Geological Society*, 179(6), jgs2022-008.
- Webers, G. F., Craddock, C., & Spletstoesser, J. F. 1992a. *Geology and paleontology of the Ellsworth Mountains. West Antarctica: Geological Society of America Memoir*, 170, 459.
- Webers, G.F., Bauer, R.L., Anderson, J.M., Buggisch, W., Ojakangas, R.W., and Spörli, K.B. 1992b. The Heritage Group of
420 the Ellsworth Mountains, West Antarctica, in Webers, G.F., Craddock, C., and Spletstoesser, J.F., eds., *Geology and Paleontology of the Ellsworth Mountains, West Antarctica: Geological Society of America Memoir 170*, p. 9–20, doi:10.1130/MEM170-p9.
- Wolf, R.A., Farley, K.A., Silver, L.T. 1996. Helium diffusion and low-temperature thermochronology of apatite. *Geochim. Cosmochim. Acta* 60, 4231–4240.
- Wolfe, M. R., & Stockli, D. F. 2010. Zircon (U–Th)/He thermochronometry in the KTB drill hole, Germany, and its
425 implications for bulk He diffusion kinetics in zircon. *Earth and Planetary Science Letters*, 295(1-2), 69-82.
- Zundel, M., Spiegel, C., Mehling, A., Lisker, F., Hillenbrand, C. D., Monien, P., & Klügel, A. 2019. Thurston Island (West Antarctica) between Gondwana subduction and continental separation: A multistage evolution revealed by apatite thermochronology. *Tectonics*, 38(3), 878-897.






Figure 1. Antarctica and crustal blocks of West Antarctica: AP—Antarctic Peninsula; EWM—Ellsworth-Whitmore Mountains block; MBL—Marie Byrd Land; TI—Thurston Island; N—Nunatak.



435 **Figure 2. (A) Simplified geological map of the Ellsworth Mountains from Craddock (1969) with the sample locations (circled numbers): 1—13EG-01; 2—13EG-05; 3—EHD0801A; 4—13EG10; 5—13EG15. (B) Stratigraphic column of the Ellsworth Mountains succession after Curtis (2001). Sample locations are placed within the column according to their stratigraphic position.**



440 **Figure 3. Depositional ages compared with ZHe ages of the grains analysed in this study. The major uplift events that formed the Transantarctic Mountains are indicated; (i) the separation between Antarctica and Australia during the Early Cretaceous, (ii) Late Cretaceous extension (main phase) between West and East Antarctica and (iii) the Cenozoic southward seafloor propagation of the Adare Trough into the Ross Sea (e.g. Fitzgerald and Gleadow, 1988; Fitzgerald, 1992, 1994, 2002; Balestrieri et al., 1997; Miller et al., 2010; Goode, 2020) along with the timing of the Weddell Sea opening (Ghidella et al., 2002; König and Jokat, 2006).**

Table 1. Single grain zircon (U-Th)/He dating results from the Heritage Group and Whiteout Conglomerate.

Sample and aliquots	South	West	Altitude	Lithology	Group	Unit	Stratigraphical Age*	⁴ He	±	U	±	Th	±	Rs ^b	Mass	⁴ He	±	U	±	Th	±	eU	Uncorr date	±2σ	F _T	Corrected date
								(pmol)		(ng)		(ng)		(μm)	(g)	(nmol/g)		(ppm)		(ppm)		(ppm)	(Ma)	(Ma)		(Ma)
13EG01 z01 z02 z03	-79.80	- 83.65	987	Sandstone	Heritage Group	Union Glacier Formation	early Cambrian (~540-520 Ma)																			
								0.54	0.0033	0.88	0.013	0.30	0.004	49	0.00271	129	0.8	210	3.0	72	1.0	227	104.6	3.0	0.8	139.3
								0.33	0.0021	0.39	0.006	0.34	0.005	45	0.00213	101	0.6	118	1.7	103	1.5	142	130.5	3.5	0.7	178.2
								0.58	0.0035	0.76	0.011	0.45	0.006	58	0.00392	86	0.5	113	1.6	67	1.0	129	123.0	3.4	0.8	155.8
13EG05 z01 z02 z03	-79.61	- 84.45	1443	Sandstone	Heritage Group	Springer Peak	middle- to late Cambrian (~520-480 Ma)																			
								0.62	0.0038	0.72	0.010	0.42	0.006	49	0.00309	133	0.8	155	2.2	91	1.3	177	138.2	3.8	0.8	184.0
								0.21	0.0009	0.29	0.004	0.15	0.002	41	0.00165	84	0.3	115	1.6	58	0.8	128	119.9	3.2	0.7	169.2
								0.56	0.0023	0.78	0.011	0.42	0.006	48	0.00230	144	0.6	200	2.8	107	1.5	225	117.6	3.1	0.7	157.5
EHD0801A z01 z02 z03	-79.97	- 82.94	1240		Heritage Group	Liberty Hills Formation	middle- to late Cambrian (~520-480 Ma)																			
								0.14	0.0006	0.22	0.003	0.12	0.002	41	0.00137	64	0.3	99	1.4	57	0.8	112	105.5	2.8	0.7	149.7
								0.02	0.0001	0.05	0.001	0.01	0.000	40	0.00004	8	0.0	20	0.3	5	0.1	21	72.2	2.1	0.7	102.9
								0.11	0.0005	0.18	0.003	0.08	0.001	37	0.00090	71	0.3	118	1.7	54	0.8	130	100.6	2.7	0.7	148.3
13EG10 z03	-77.60	- 86.32	1810	Conglomeratic sandstone (matrix)		Whiteout Conglomerate	Permian - Carbonifeours (~360-260 Ma)																			
								1.13	0.0047	1.61	0.023	0.14	0.002	39	0.00201	466	1.9	662	9.4	57	0.8	676	126.6	3.6	0.7	181.7
13EG15 z01	-79.77	- 81.30	600	Conglomeratic sandstone (matrix)		Whiteout Conglomerate	Permian - Carbonifeours (~360-260 Ma)																			
								2.65	0.0109	3.80	0.054	0.39	0.006	58	0.00508	324	1.3	465	6.6	47	0.7	476	124.9	3.5	0.8	158.2

z02								0.77	0.0035	0.37	0.005	0.05	0.001	50	0.00029	156	0.7	76	1.1	11	0.2	78	357.4	10.5	0.8	467.2	
z03								1.16	0.0053	0.33	0.005	0.00	0.000	51	0.00001	222	1.0	63	0.9	1	0.0	63	614.2	18.8	0.8	790.4	
Standard																											
FCTjm_Zr14								0.14	0.0007	1.04	0.015	0.88	0.013	41	0.00158	63	0.3	455	6.6	387	5.6	546	21.4	0.5	0.7	30.3	
FCTjm_Zr13								0.19	0.0012	1.42	0.020	0.82	0.012	43	0.00224	67	0.4	509	7.3	295	4.2	578	21.5	0.6	0.7	30.0	

* Stratigraphical ages have been estimated from the column presented in Curtis (2001)

Code/Data availability

All raw unprocessed data related to this study can be requested by contacting the corresponding author.

Author Contribution

- 450 Joaquín Bastías-Silva: conceptualisation, leading manuscript development, leading data interpretation, visualisation.
David Chew: manuscript development, visualisation.
Fernando Poblete: funding acquisition, conceptualisation, sample collection, manuscript development.
Paula Castillo: conceptualisation, sample collection, manuscript development.
William Guenther: data collection, manuscript development.
- 455 Anne Grunow: manuscript development.
Ian W.D. Dalziel: manuscript development.
Airtón N. C. Dias: manuscript development.
Cristóbal Ramírez de Arellano: sample collection.
Rodrigo Fernandez: conceptualisation, sample collection.

460 **Competing interest**

The (co-)authors are not member of the editorial board of Solid Earth and/or a guest member of the editorial board of Solid Earth. The peer-review process will be guided by an independent editor, and the authors also have no other competing interests to declare.