

High-resolution Mapping of Nitrogen Oxide Emissions in Large US Cities from TROPOMI Retrievals of Tropospheric Nitrogen Dioxide Columns

5 Fei Liu^{1,2}, Steffen Beirle³, Joanna Joiner², Sungeon Choi^{2,4}, Zhining Tao^{1,2}, K. Emma Knowland^{1,2},
Steven J. Smith⁵, Daniel Q. Tong^{6,7}, Siqi Ma^{6,7}, Zachary T. Fasnacht^{2,4}, Thomas Wagner³

¹Goddard Earth Sciences Technology and Research (GESTAR) II, Morgan State University, Baltimore, MD 21251, USA

²NASA Goddard Space Flight Center, Greenbelt, MD, 20771, USA

³Max-Planck-Institut für Chemie, Mainz, 55128, Germany

10 ⁴Science Systems and Applications Inc., Lanham, MD, 20706, USA

⁵Joint Global Change Research Institute, Pacific Northwest National Laboratory, College Park, MD, 20740, USA

⁶Department of Atmospheric, Oceanic and Earth Sciences, George Mason University, Fairfax, 22030, Virginia, USA

⁷Center for Spatial Information Science and Systems, George Mason University, Fairfax, 22030, Virginia, USA

15 *Correspondence to:* Fei Liu (fei.liu@nasa.gov)

Abstract. Satellite-derived spatiotemporal patterns of nitrogen oxide (NO_x) emissions can improve accuracy of emission inventories to better support air quality and climate research and policy studies. In this study, we develop a new method by coupling the chemical transport Model-Independent SATellite-derived Emission estimation Algorithm for Mixed-sources (MISATEAM) with a divergence method to map high-resolution NO_x emissions across US cities using TROPOspheric
20 Monitoring Instrument (TROPOMI) tropospheric nitrogen dioxide (NO₂) retrievals. The accuracy of the coupled method is validated through application to synthetic NO₂ observations from the NASA-Unified Weather Research and Forecasting (NU-WRF) model, with a horizontal spatial resolution of 4 km × 4 km for 33 large and mid-size US cities. Validation reveals excellent agreement between inferred and NU-WRF-provided emission magnitudes ($R = 0.99$, Normalized Mean Bias, NMB = -0.01) and a consistent spatial pattern when comparing emissions for individual grid cells ($R = 0.88 \pm 0.06$). We then develop
25 a TROPOMI-based database reporting annual emissions for 39 US cities at a horizontal spatial resolution of 0.05°×0.05° from 2018 to 2021. This database demonstrates a strong correlation ($R = 0.90$) with the national emission inventory (NEI) but reveals some bias (NMB = -0.24). There are noticeable differences in the spatial patterns of emissions in some cities. Our analysis suggests that uncertainties in TROPOMI-based emissions and potential misallocation of emissions and/or missing sources in bottom-up emission inventories both contribute to these differences.

30 1 Introduction

Nitrogen oxides (NO_x), including nitrogen dioxide (NO₂) and nitric oxide (NO), play a vital role as trace gases in the atmosphere. They substantially contribute to the formation of secondary aerosols and tropospheric ozone (Seinfeld and Pandis, 2006), which in turn impact climate and human health. The primary source of NO_x emissions is fossil fuel combustion from mobile and industrial sources, often concentrated in urban areas (Crippa et al., 2018). Traditionally, NO_x emissions are compiled using "bottom-up" methods that rely on total fuel usage and average emission factors. However, determining urban emissions is challenging due to the difficulty of obtaining comprehensive data on operating conditions of emitters and their rapidly changing emission factors (Liu et al., 2016b) at the city level (Butler et al., 2008). Spatial distribution of emissions is commonly estimated by applying proxies, such as road network maps and/or population density maps, to allocate total emissions onto a grid. This process may introduce significant biases due to spatial mismatches between emission locations and spatial proxies (Woodard et al., 2014; Hogue et al., 2016).

Tropospheric NO₂ vertical column densities (VCDs) retrieved from satellite observations offer valuable insights into the magnitude and location of global NO_x emissions (Martin et al., 2003; Lamsal et al., 2011). Techniques leveraging Chemical Transport Models (CTMs) have been developed to relate NO₂ VCDs to NO_x emissions. For example, the extended (Ding et al., 2017) and ensemble (Miyazaki et al., 2017) Kalman filter, the four-dimensional variational (4D-Var) method (Henze et al., 2007, 2009), and the hybrid mass balance/4D-Var (Qu et al., 2019).

Alternative methods, which are independent of CTMs, have also been proposed to characterize NO_x plumes from major sources (e.g., Beirle et al., 2011; Liu et al., 2016a; Laughner and Cohen, 2019). Early studies employed one dimensional (1D) empirical plume dispersion functions to fit NO₂ VCDs surrounding isolated sources. Liu et al. (2016a, 2022) refined these functions to accommodate the description of NO₂ plumes from sources in polluted background. The derived fitting parameters yield the magnitude of NO_x emissions for point sources, such as power plants (de Foy et al., 2014), and cities, by assuming these sources as equivalent point sources (Lu et al., 2015; Liu et al., 2017; Goldberg et al., 2019). A recent study (Beirle et al., 2019) introduced a two-dimensional (2D) divergence approach, enabling the identification of finer details in NO_x distributions and thus facilitating the detection of smaller sources. Subsequent studies have further refined this approach to enhance divergence calculation (de Foy and Schauer, 2022) and to optimize its performance over mountainous regions (Sun, 2022). Current applications of this divergence method primarily focus on inferring emissions from point sources, e.g., power plants in South Asia (de Foy and Schauer, 2022), oil and gas production areas in the United States (Dix et al., 2022), and a global catalog of point sources (Beirle et al., 2021, 2023). However, the robustness of applying the approach for area sources such as cities has seldom been investigated.

The main goal of this study is to map NO_x emissions across major cities in the US based on the TROPOspheric Monitoring Instrument (TROPOMI; Veefkind et al., 2012) retrievals of NO₂ VCDs (Ialongo et al., 2020). We will couple the 1D (Liu et al., 2022) and 2D (Beirle et al., 2019) CTM-independent approaches to infer gridded NO_x emissions. Due to the absence of established "true emissions" to serve as a standard for comparison, gauging the precision of the coupled method becomes

challenging. We will employ synthetic NO₂ observations, generated by a model simulation, to appraise the accuracy of the approach. Section 2 offers a summary of the satellite data used to map urban NO_x emissions, and the synthetic NO₂ observations used for validation. In Section 3.1, we assess the coupled approach by juxtaposing the emissions inferred from synthetic NO₂ observations with the true emissions used by the model for generating the synthetic observations. In Section 3.2, we compare the emissions derived from satellite data with the National Emission Inventory (NEI) developed by the United States Environmental Protection Agency (US EPA) to shed light on the uncertainties of both and bottom-up emissions. Section 3.3 summarizes the uncertainties of the method and the TROPOMI-derived emissions. We discuss the robustness of the derived emissions and outline the plan work in Section 4.

2 Data and Methods

2.1 TROPOMI NO₂ dataset

TROPOMI is a UV-VIS-NIR-SWIR nadir-viewing imaging spectrometer (Veefkind et al., 2012) on board the Sentinel 5 Precursor (S5P) satellite that was launched in 2017. It has a ground pixel size at nadir of 7.5 km×3.5 km before August 6, 2019, and improved to 5.5 km ×3.5 km afterwards. It provides daily global coverage with a local equator crossing time of approximately 13:30 h. The instrument's radiance and irradiance measurements are utilized to obtain slant NO₂ columns using the Differential Optical Absorption Spectroscopy (DOAS) algorithm (Platt and Stutz, 2008). The slant columns are subsequently differentiated into stratospheric and tropospheric parts, with the tropospheric slant columns being further converted to vertical columns based on air mass factors.

We selected TROPOMI NO₂ retrieved by NASA Goddard Space Flight Center (GSFC), TROPOMI Multi-Decadal Nitrogen Dioxide and Derived Products from Satellites (MINDS) NO₂ product (Lamsal et al., 2022), in this study due to its utility in investigating emission trends. The official NO₂ product available at the commencement of this study switched the processor version in December 2020, which introduced a discontinuity in the time series (van Geffen et al., 2022). The GSFC product (Lamsal et al., 2022) utilizes slant columns from the official product to retrieve a full time series of tropospheric NO₂ VCDs from spring to autumn months of 2018 to 2021 based on the algorithm developed for the Ozone Monitoring Instrument (OMI) Aura NO₂ standard product version 4.0 (Lamsal et al., 2021). TROPOMI NO₂ data has been reported to have an overall low bias (Judd et al., 2020; Tack et al., 2021; Verhoelst et al., 2021; Wang et al., 2020), which will be propagated into emissions inferred from TROPOMI NO₂.

We further aggregate TROPOMI's individual measurements at spatial resolution of 0.05°×0.05° by averaging the original pixels weighted by the fraction of the overlapping surface area. Only high-quality pixels with a quality assurance value (qa_value) above 0.75 are considered for averaging, which excludes cloud-covered scenes with cloud radiance fraction > 0.5. Following the approach of earlier research (e.g., Liu et al., 2017), our analysis is confined to the data from May through September. This

decision is made to omit winter data, characterized by longer NO_x lifetimes, which in turn lead to greater uncertainties to the method.

95 2.2 Emission mapping algorithm

We couple our 1D CTM-Independent SATellite-derived Emission estimation Algorithm for Mixed-sources (MISATEAM; Liu et al., 2022) with the 2D divergency method of Beirle et al. (2019). We couple our 1D CTM-Independent SATellite-derived Emission estimation Algorithm for Mixed-sources (MISATEAM; Liu et al., 2022) with the 2D divergence method of Beirle et al. (2019). 1D MISATEAM quantifies the magnitude of city-level NO_x lifetime and emissions by conceptualizing
 100 urban areas as point sources and thus does not capture the spatial variability within the urban areas. Conversely, the 2D divergence method allows for the resolution of finer spatial details in NO_x distributions but relies on additional, often external, sources for determining NO_x lifetimes, which can be a significant limitation. The coupled algorithm (hereafter referred to as 2D MISATEAM for simplicity) leverages the strengths of both: it maps NO_x emissions over urban areas with enhanced spatial detail and does so independently by deriving lifetimes directly from NO₂ measurements, thereby overcoming the need for
 105 prescribed or externally sourced lifetime constraints. 1D and 2D MISATEAM methods deliver consistent estimates for total emissions, displaying a small relative difference of $-5\% \pm 9\%$. While this result demonstrates the internal consistency of the MISATEAM methods, it's important to note that such coherence is due to the shared fundamental principle of mass balance underlying both methodologies. Therefore, the similarity in emissions estimations should not be viewed as an independent validation metric.

110 We average both NO₂ VCDs and reanalysis wind data from May to September each year. We then use those averaged data to infer NO_x emissions E by summing the divergence of the NO_x flux D with the NO_x sink S based on the continuity equation for steady state following:

$$E = D + S \quad (1)$$

$$D = R_{NO_x:NO_2} \times \nabla \cdot \vec{F} = R_{NO_x:NO_2} \times \nabla \cdot \vec{v}(\Omega - b) \quad (2)$$

$$115 \quad S = R_{NO_x:NO_2} \times (\Omega - b)/\tau, \quad (3)$$

Where \vec{F} is the NO_x flux. It is calculated from the horizontal fluxes of NO₂ VCDs Ω . $R_{NO_x:NO_2}$ is the ratio of NO_x to NO₂ columns. Following previous studies (Beirle et al., 2019), we use an $R_{NO_x:NO_2}$ value of 1.32 to represent “typical urban conditions and noontime sun” (Seinfeld and Pandis, 2006). We interpolate the Goddard Earth Observing System Forward Processing for Instrument Teams (GEOS FP-IT) reanalysis wind vectors (Lucchesi, 2015) to the TROPOMI overpass time
 120 and average layers from surface to 1000 m altitude to derive \vec{v} used in Eq. (2). Since the NO_x sinks are dominated by the chemical loss through the reaction of NO₂ with OH at TROPOMI’s local overpass time (13:30 local time), it can be characterized by a first order effective NO_x lifetime τ . Consequently, it bears a proportionate relationship to the NO₂ VCD itself in Eq. (3). Note that we subtract the NO₂ background b from Ω in the calculation of the divergences and the sinks,

because we aim to remove the natural and non-local contributions from the total emissions for each urban area in order to infer urban emissions. For each city, we infer b and τ by applying 1D MISATEAM (Liu et al., 2022) to NO₂ VCDs averaged from May through September, 2018–2021, assuming b and τ are constant over years. Additional technical details for deriving b and τ are given in Fig. S1 and Text S1 of the Supplement.

The city of New York serves as a case study to showcase our approach. With its substantial size and numerous point and area sources, this city is an ideal illustration of the capability of 2D MISATEAM to map emissions from various sources. Figure 1a–c illustrate maps of the derived sink S , the divergence D , and the resulting NO_x emissions E , respectively. The divergence of the NO_x flux (Fig. 1b) presents enhancement throughout the urban area due to emissions from traffic and industrial sources. The divergence is negative outside the urban areas, because the change of the NO_x flux is dominated by chemical loss here. The addition of sinks (Fig. 1a) compensates for such negative divergences, resulting in the emission pattern E (Fig. 1c). Two point sources emissions on Long Island stand out: Glenwood Landing power station (labelled as power plant PP) and John F. Kennedy (JFK) airport, respectively.

We apply 2D MISATEAM to US cities with populations exceeding 200,000, a categorization that corresponds to medium to large urban areas as designated in Organization for Economic Co-operation and Development (OECD) countries. Adjacent cities, those situated within 50 km of the most populous city in an urban conglomeration, are treated as a single city cluster. Cities producing significantly weaker NO₂ signals than the surroundings are excluded (see criteria in Text S2 of Supplement). The specified criteria yield a combined count of 52 cities and urban conglomerations (refer to Table S1) that are suitable for the application of 2D MISATEAM. We obtained valid results from 39 of these cities shown in Fig. S2. Cities with invalid results are associated with large fitting errors (see Text S2 of the Supplement).

2.3 National Emission Inventory (NEI)

We compare emissions derived in this study with those of the bottom-up NEI 2019 processed by George Mason University (Ma and Tong, 2022). NEI is a comprehensive estimate of emissions of criteria air pollutants and their precursors from point, mobile and area sources, which has been widely used to support urban air quality model simulations. The large point sources in NEI are compiled based on direct stack emissions measurements by continuous emissions monitoring systems (CEMS). For non-point sources, the gridded data is generated using spatial proxies such as roadway-level traffic data for distributing aggregate emissions to grid cells. Figure 1d displays a map of NEI estimates E_{NEI} at a spatial resolution of 12 km for 2019. NEI has been reported to be biased high by 30–70% in the early years of the 2010s (Choi and Souri, 2015; Dickerson et al., 2019). For example, NEI NO_x estimates for 2011 were potentially overestimated by 51–70% over the Baltimore-Washington region (Anderson et al., 2014), 30–60% in the Southeastern US (Travis et al., 2016), and 30–60% over urban areas of Texas (Souri et al., 2016). NEI NO_x emissions for mobile sources alone were reported to be biased high by 28% (McDonald et al., 2018). We use the lower bound of the reported bias (30%) as the uncertainty for NEI emissions in this study. The spatial distribution of our estimated emissions E (Fig. 1c) is generally in good agreement with that of NEI emissions (Fig. 1d), with a correlation coefficient of 0.71. More comparisons between these two inventories will be discussed in Section 3.2.

2.4 NU-WRF simulations

We use a regional modeling system, the NASA-Unified Weather Research and Forecasting (NU-WRF; Tao et al., 2013; Peters-Lidard et al., 2015), to provide synthetic tropospheric NO₂ VCDs Ω_{NU-WRF} and wind fields $\overrightarrow{v_{NU-WRF}}$ over the continental US. The domain of the simulation is illustrated in Fig. 2 of Liu et al. (2022). We perform the 2016 NU-WRF simulation at a high horizontal spatial resolution of 4 km × 4 km, comparable to the TROPOMI footprint. The meteorological and chemical initial conditions and their lateral boundaries are obtained from NASA’s Modern Era Retrospective-Analysis for Research and Applications version 2 (MERRA-2; Gelaro et al., 2017) and the Community Atmosphere Model with chemistry (CAM-chem; Lamarque et al., 2012), respectively. Additional model set-up information, including the chemical mechanism, aerosol module, and emissions are detailed in Tao et al. (2020) and Liu et al. (2022). We integrate simulated NO₂ concentrations from the surface to the tropopause to provide Ω_{NU-WRF} . We average wind fields used by NU-WRF from the surface to an altitude of 1000 m to compute $\overrightarrow{v_{NU-WRF}}$.

We apply 2D MISATEAM to the synthetic NO₂ VCDs Ω_{NU-WRF} and wind fields $\overrightarrow{v_{NU-WRF}}$ to map NO_x emissions E_{NU-WRF} for cities in Table S1. We derive valid results for 33 cities (see Text S2 of Supplement). We have valid results for a smaller number of cities compared to results derived from TROPOMI data, as cities at the edge of the NU-WRF domain, such as Seattle and San Francisco, are omitted. Since the model output partially lacks the data for their inflow/outflow plumes, it fails to satisfy the requirement for the application of 2D MISATEAM. We further compare E_{NU-WRF} with “true emissions” directly given by the model E'_{NU-WRF} (hereafter referred to as “given emissions”), which are used to drive NU-WRF simulations, to assess the accuracy of 2D MISATEAM.

2.5 Performance evaluation

We sum up NO_x emissions from each grid cell within the city’s domain to provide a total emission for an individual city. The city domain is defined as 70 km × 70 km around city center for most cities, which is large enough to include all urban areas. We use a larger domain of 100 km × 100 km for New York, Chicago, Los Angeles and Houston due to their larger expanse. The model performance metrics of the Normalized Mean Bias (NMB) and the Root Mean Squared Error (RMSE) for the evaluation are defined as

$$NMB = \frac{\sum_{i=1}^n (Emis_i - Emis'_i)}{\sum_{i=1}^n Emis'_i} \quad (5)$$

and

$$RMSE = \sqrt{\frac{\sum_{i=1}^n (Emis_i - Emis'_i)^2}{n}}, \quad (6)$$

respectively, where i denotes an individual city, while n refers to the total count of cities utilized for the evaluation. $Emis$ denotes the total emission from the MISATEAM-derived datasets (i.e., E or E_{NU-WRF}), and $Emis'$ denotes the total emission

from benchmark emission datasets (i.e., E_{NEI} or E'_{NU-WRF}). We also assess the intracity spatial correlation $R_{intracity}$ by calculating the correlation coefficient of emissions at grid level over the city domain between E and E_{NEI} or between E_{NU-WRF} and E'_{NU-WRF} .

3 Results and discussions

3.1 Validation using NU-WRF simulations

190 We compare MISATEAM-derived NO_x emissions E_{NU-WRF} with given emissions E'_{NU-WRF} to validate 2D MISATEAM. The validation indicates the uncertainty of MISATEAM assuming a best-case scenario with perfect knowledge of the winds and errorless satellite NO_2 retrievals. Figure 2 compares the total emissions from the two datasets for the 33 cities (Table S1). The correlation coefficient is 0.99, which indicates an excellent agreement between the two datasets. The overall bias computed over all valid cities is generally low (NMB = -0.01) and the RMSE is also low (0.21 kg s^{-1}).

195 We assess 2D MISATEAM's performance to infer the spatial distribution of emissions. Figure 3 displays the emission maps around the city of Jacksonville, Florida. E_{NU-WRF} (Fig 3c) and E'_{NU-WRF} (Fig. 3d) show a high intracity spatial correlation $R_{intracity}$ of 0.92. This good consistency is substantially better than the comparison of E'_{NU-WRF} with the tropospheric NO_2 VCD Ω_{NU-WRF} (Fig. 3a; $R_{intracity} = 0.75$), which is often considered to be a reasonable spatial proxy of emissions locations. We further compare the correlation with the comparison of E'_{NU-WRF} with Ω_{NU-WRF} under calm wind conditions (Fig. 3b). We
200 use the threshold of 2 m s^{-1} as the criterion for calm wind, allowing for the attainment of an adequate sample size. This threshold reduces the error associated with transport by increasing $R_{intracity}$ from 0.75 (Fig. 3a) to 0.80 (Fig. 3b). However, this correlation is still smaller than that between E_{NU-WRF} and E'_{NU-WRF} , suggesting that 2D MISATEAM is successful in allocating emissions by accounting for NO_x transport. Similar enhanced correlations are observed for all 33 cities. Figure 2 illustrates $R_{intracity}$ of E_{NU-WRF} and E'_{NU-WRF} for individual cities (0.88 ± 0.06 , mean \pm standard deviation). These correlations are larger than the
205 comparison of Ω_{NU-WRF} (0.78 ± 0.09) or Ω_{NU-WRF} under calm wind conditions (0.80 ± 0.08) against E'_{NU-WRF} .

3.2 TROPOMI-based NO_x emissions

We compare TROPOMI-based NO_x emissions E with NEI estimates E_{NEI} for 2019 in Fig. 4. The total emission estimates for individual cities in the two datasets generally agree well with each other, with a correlation R of 0.90. This level of correlation is comparable to the validation using NU-WRF simulations. The relative difference of the total emission between E and E_{NEI}
210 is within the uncertainty range of E (47%; see Section 3.3) for 31 out of 39 cities. The comparison for all cities shows a bias with NMB of -0.24. The bias is likely associated with uncertainties in the TROPOMI NO_2 retrievals, which have been reported to be biased low by 23% on average (van Geffen et al., 2022). The bias may also arise from the uncertainties in NEI, which has been reported to be biased high by over 30% (Section 2.3).

Figure 5 compares the NO_x emission patterns from TROPOMI NO_2 with those reported in the NEI, using Dallas and Tucson
215 as case studies. Consistent with observations in New York (as shown in Fig. 1), TROPOMI-derived emission maps reveal

several more pronounced point sources as compared to NEI. Notable emissions from the Dallas/Fort Worth International Airport, Perot Field Fort Worth Alliance Airport, and three major cement factories—TXI, Holcim, and Ash Grove—are distinctly evident (Fig. 5a), whereas these details appear diffused in NEI (Fig. 5b). Similarly, emissions from the Asarco Mission copper mine are clearly discernible in the TROPOMI data (Fig. 5c) but are not as apparent in the NEI data (Fig. 5d).

220 We use $R_{intracity}$ to compare the intracity spatial distribution of emissions for more cities in Fig. 4. We upscale E to the same spatial resolution of E_{NEI} to calculate their $R_{intracity}$ (Fig. S3). $R_{intracity}$ between E and E_{NEI} is 0.57 ± 0.16 , which is smaller than that between E_{NU-WRF} and E'_{NU-WRF} in the evaluation using model data (0.88 ± 0.06 ; Fig. 2). The generally smaller values of $R_{intracity}$ are likely caused by the uncertainties of both TROPOMI-based and NEI emissions. Compared to E_{NU-WRF} inferred from perfect NO_2 columns and wind fields, the uncertainties of TROPOMI NO_2 retrievals (25%; van Geffen et al., 2022) and GEOS

225 FP-IT wind reanalysis (30%; Liu et al., 2022) are propagated into the uncertainties of TROPOMI-based emissions E and may result in incorrect spatial patterns. More details about the uncertainties are discussed in Section 3.3. Uncertainties in E_{NEI} also contribute to the disagreement. NEI uses spatial-distribution proxies, such as maps of population densities or road networks, to allocate country-level emissions from non-point sources onto a grid. This procedure may be associated with biases due to either a spatial mismatch between the locations of emissions and spatial proxies or incorrect emission magnitudes. Some

230 hotspots shown in E are missing from E_{NEI} , for instance, JFK airport (Fig.1) and Asarco Mission mine (Fig.5), indicating missing sources or misallocation of sources.

Figure 6 presents the trends of derived NO_x emissions across US cities from 2018 to 2021. A significant dip in emissions was observed in 2020, primarily attributed to the diminished NO_x emissions in response to COVID-related lockdown policies (e.g., Liu et al., 2020). The 2020 emissions saw an average reduction of approximately 25% from 2019, in agreement with the

235 findings of other studies (Goldberg et al., 2020; Miyazaki et al., 2021). Emissions in 2021 experienced a rebound compared to those in 2020; however, they are still lower than the pre-pandemic levels. The slopes of the linear regression lines in Fig. 6 decrease from 0.91 in 2019 to 0.85 in 2021. This decline aligns with the long-term trend of decreasing emissions in the US, primarily driven by the downturn trend in vehicular NO_x emissions (McDonald et al., 2018). The reduced slope in 2021 (0.85) relative to 2019 (0.91) may also encapsulate lingering impacts of diminished traffic from the COVID-19 pandemic, since the

240 traffic and commercial flight metrics at the end of 2020 were still substantially lower than their January 2020 levels (Laughner et al., 2021).

3.3 Uncertainty analysis

We follow the method proposed by Liu et al. (2022) to evaluate the uncertainties associated with the derived results. We attribute an uncertainty of 20% to the inferred emissions, based on the relative differences between E_{NU-WRF} and E'_{NU-WRF} for

245 all 33 cities using the NU-WRF synthetic data ($2 \pm 24\%$). Since the differences are less than 20% for most cities (73%), this estimate may be conservative.

Instead of using data spanning multiple years (2018-2021), we apply 1D MISATEAM to annual data to investigate the uncertainty introduced by presuming a consistent NO_x lifetime over several years. For each city, we infer τ by applying 1D

MISATEAM to TROPOMI NO₂ VCDs, averaged from May to September for each individual year from 2018 to 2021. Among
250 the cities analyzed, 14 cities (listed in Table S1) have valid NO_x lifetimes available for all four years. The ratio of the standard
deviation of the fitted τ for these individual years to the average τ , as derived from data from 2018 to 2021, stands at 16%. This
indicates an uncertainty of 16% in relation to the assumption of a constant NO_x lifetime.

We identify additional uncertainties that may arise when implementing 2D MISATEAM to TROPOMI and GEOS-IT data as
opposed to synthetic data, as follows:

255 • The uncertainty of TROPOMI NO₂ observations carries over into the uncertainty of the inferred emissions. We
consider an overall uncertainty of 25% for TROPOMI tropospheric NO₂ VCDs, following the recommendation from
a recent validation using ground-based measurements (van Geffen et al., 2022). The uncertainty originates from
various factors, including the spectral fitting process during the retrieval, the separation of stratospheric and
tropospheric columns, and the tropospheric air mass factor (AMF).

260 AMF accounts for the variable sensitivity of satellite observations to NO₂ at different atmospheric altitudes, which is
informed by a priori knowledge of NO₂ vertical distribution as provided by chemical transport models. Laughner et
al. (2016) demonstrated that urban NO_x emissions estimated via NO₂ VCDs with daily, high-resolution a priori
profiles are considerably higher than those derived from retrievals using coarser resolution profiles. This presents a
relevant challenge for our study. To refine the assessment of AMF influences on 2D MISATEAM-derived emissions,
265 future work could include a sensitivity analysis where TROPOMI's AMFs are applied to NU-WRF profiles. This
would generate NO₂ columns with AMF biases, which, when used to calculate emissions, can be contrasted with
those derived from idealized columns. Such a comparison would reveal the extent to which AMF biases systematically
affect the emissions determined by 2D MISATEAM.

The potential bias arising from the separation of stratospheric and tropospheric columns is eliminated by the
270 employment of the background terms b and b_{calm} within the model functions of MISATEAM. Since the random
uncertainty of the tropospheric NO₂ observations could be suppressed due to the consideration of long-term means,
this estimate may be conservative.

• The presence of clouds is an additional source of uncertainties. We exclude TROPOMI NO₂ data with cloud radiance
fraction of 0.5 or greater in our analysis. This exclusion may lead to a bias in the NO₂ VCD averages, a consequence
275 of eliminating data with changing NO_x lifetime and NO_x/NO₂ ratio during cloudy conditions (Geddes et al., 2012).
We attribute an uncertainty of 10% to cloud selection criteria based on an assessment carried out at urban locations
(Geddes et al., 2012). More sensitivity analysis regarding the impact of clouds has been discussed by Liu et al. (2022).

• The precision of wind fields plays an important role in determining the total uncertainty, as it limits the model
functions' ability to describe NO₂ transport. We estimate the corresponding uncertainties to be 30% based on an
280 assessment of reanalysis wind products using sounding measurements (refer to Table S3 in Liu et al., 2016a).

We define the overall uncertainty of the inferred emissions as the root of the quadratic sum of the above-mentioned uncertainties, which are assumed to be independent. We thus calculate that the total uncertainty of MISATEAM-derived NO_x emissions for a mid-size US city is 47%.

4 Conclusions

285 In this study, we developed a new method by coupling the 1D CTM-independent methodology (Liu et al. 2022), MISATEAM, with the 2D divergence approach (Beirle et al., 2019) to generate maps of NO_x emissions across US cities, using TROPOMI NO₂ observations. This coupled method, 2D MISATEAM, is suitable for sources within polluted backgrounds. Our initial application of the approach used synthetic NU-WRF-generated tropospheric NO₂ VCDs over the continental US to evaluate the method. The evaluation yielded robust agreement with the NU-WRF given values, presenting a high correlation coefficient
290 (R = 0.99) and a minimal bias (NMB = -0.01).

Subsequently, we applied 2D MISATEAM to TROPOMI NO₂ retrievals across the US cities from 2018 to 2021. We estimated NO_x emissions for 39 US cities. Our resulting total city emissions estimates align strongly with NEI (R = 0.90), albeit with a moderate bias (NMB = -0.24). Our derived emissions delineate differences in spatial patterns across certain cities, implying potential inaccuracies in emission allocation and/or missing sources in NEI. Our current estimates suggest that uncertainties
295 in NO_x emissions arising from 2D MISATEAM method itself are approximately 20% for a large and mid-size US city. Additional uncertainties stem primarily from errors in the reanalysis wind dataset as well as the TROPOMI NO₂ retrieval, increasing the overall uncertainties of resulting emissions to about 47%.

2D MISATEAM is contingent on 1D MISATEAM for NO_x lifetime estimation. 1D MISATEAM relies on NO₂ observations under calm wind conditions to infer lifetimes, which in turn influences the temporal resolution of the emissions data we can
300 confidently derive. Our investigation indicates that typically 3 to 6 months of TROPOMI data are required to ensure comprehensive data coverage of calm-wind NO₂ observations for urban emissions analysis. Therefore, we advise caution when considering the use of 2D MISATEAM for emission estimations over periods shorter than three months, unless we want to use prescribed or externally sourced lifetimes.

In our future research, we plan to extend the application of 2D MISATEAM to observations from geostationary satellites to
305 estimate diurnal variations in urban emissions. Geostationary sensors include the Korean Geostationary Environmental Monitoring Spectrometer (GEMS; Kim et al., 2012), NASA's Tropospheric Emissions: Monitoring of Pollution (TEMPO; Chance et al., 2012), and ESA's Sentinel-4 (Ingmann et al., 2012). These instruments possess spatial resolutions similar to TROPOMI and the validation of NU-WRF simulation (4 km) utilized in this study. For applications based on geostationary satellites with local observation time extending beyond the early afternoon time frame of TROPOMI in this study, further
310 exploration into the impact of the diurnal cycle of NO_x lifetime will be necessary. When applying the 2D MISATEAM to cities globally, particularly in non-OECD countries, it may be necessary to adjust population thresholds to reflect local demographics and urban profiles. We advise removing cities with too weak emissions signals following this study, i.e., $b_{calm}/\text{mean VCDs} >$

50% (Text S2 of the Supplement), as such a high ratio can introduce significant uncertainties associated with determined background b_{calm} .

315 In the next phase of our work, we will strive to harmonize bottom-up and satellite-derived urban emissions estimates to produce a fused emission inventory (Liu et al., 2018). This will enable the provision of timely NO_x emissions estimates that will be of value to both air quality and climate modelling communities.

Data availability. The NU-WRF model outputs are available upon request from Zhining Tao (zhining.tao@nasa.gov).

320 Additional data related to this paper may be requested from the corresponding author.

Author contributions. Conceptualization and methodology: F.L., J.J., and S.B.; Model simulation: Z.T.; Satellite data processing: S.C.; Wind data processing: Z.F.; NEI data processing: D.T. and S.M.; Formal analysis: F.L.; Visualization: F.L.; Writing—original draft: F.L., review and editing: All authors; Funding acquisition: F.L. and J.J.

Competing interests. The authors declare that they have no competing interests.

325 *Acknowledgements.* This work was funded by NASA through the Aura project data analysis program and through the Atmospheric Composition Modeling and Analysis Program (ACMAP) program (grant no. 80NSSC19K0980). We thank Prof. Russell R. Dickerson for helpful discussion on uncertainties in NEI. We thank Dr. Laughner and one anonymous reviewer for helpful comments.

330 **References**

Anderson, D. C., Laughner, C. P., Diskin, G., Weinheimer, A., Canty, T. P., Salawitch, R. J., Worden, H. M., Fried, A., Mikoviny, T., Wisthaler, A., and Dickerson, R. R.: Measured and modeled CO and NO_y in DISCOVER-AQ: An evaluation of emissions and chemistry over the eastern US, *Atmos. Environ.*, 96, 78–87, <https://doi.org/10.1016/j.atmosenv.2014.07.004>, 2014.

335 Beirle, S., Borger, C., Dörner, S., Li, A., Hu, Z., Liu, F., Wang, Y., and Wagner, T.: Pinpointing nitrogen oxide emissions from space, *Sci. Adv.*, 5, eaax9800, <https://doi.org/10.1126/sciadv.aax9800>, 2019.

Beirle, S., Borger, C., Dörner, S., Eskes, H., Kumar, V., de Laat, A., and Wagner, T.: Catalog of NO_x emissions from point sources as derived from the divergence of the NO₂ flux for TROPOMI, *Earth Syst. Sci. Data*, 13, 2995–3012, <https://doi.org/10.5194/essd-13-2995-2021>, 2021.

340 Beirle, S., Borger, C., Jost, A., and Wagner, T.: Improved catalog of NO_x point source emissions (version 2), *Earth Syst. Sci. Data Discuss.*, 1–37, <https://doi.org/10.5194/essd-2023-44>, 2023.


- Butler, T. M., Lawrence, M. G., Gurjar, B. R., van Aardenne, J., Schultz, M., and Lelieveld, J.: The representation of emissions from megacities in global emission inventories, *Atmos. Environ.*, 42, 703–719, <https://doi.org/10.1016/j.atmosenv.2007.09.060>, 2008.
- 345 Chance, K., Lui, X., Suleiman, R. M., Flittner, D. E., and Janz, S. J.: Tropospheric Emissions: monitoring of Pollution (TEMPO), presented at the 2012 AGU Fall Meeting, San Francisco, USA, 3–7 December 2012, A31B-0020, 2012.
- Choi, Y. and Souri, A. H.: Chemical condition and surface ozone in large cities of Texas during the last decade: Observational evidence from OMI, CAMS, and model analysis, *Remote Sens. Environ.*, 168, 90–101, <https://doi.org/10.1016/j.rse.2015.06.026>, 2015.
- 350 Crippa, M., Guizzardi, D., Muntean, M., Schaaf, E., Dentener, F., van Aardenne, J. A., Monni, S., Doering, U., Olivier, J. G. J., Pagliari, V., and Janssens-Maenhout, G.: Gridded emissions of air pollutants for the period 1970–2012 within EDGAR v4.3.2, *Earth Syst Sci Data*, 10, 1987–2013, <https://doi.org/10.5194/essd-10-1987-2018>, 2018.
- Dickerson, R. R., Anderson, D. C., and Ren, X.: On the use of data from commercial NO_x analyzers for air pollution studies, *Atmos. Environ.*, 214, 116873, <https://doi.org/10.1016/j.atmosenv.2019.116873>, 2019.
- 355 Ding, J., van der A, R. J., Mijling, B., and Levelt, P. F.: Space-based NO_x emission estimates over remote regions improved in DECSO, *Atmospheric Meas. Tech.*, 10, 925–938, <https://doi.org/doi:10.5194/amt-10-925-2017>, 2017.
- Dix, B., Francoeur, C., Li, M., Serrano-Calvo, R., Levelt, P. F., Veeffkind, J. P., McDonald, B. C., and de Gouw, J.: Quantifying NO_x Emissions from U.S. Oil and Gas Production Regions Using TROPOMI NO₂, *ACS Earth Space Chem.*, 6, 403–414, <https://doi.org/10.1021/acsearthspacechem.1c00387>, 2022.
- 360 de Foy, B. and Schauer, J. J.: An improved understanding of NO_x emissions in South Asian megacities using TROPOMI NO₂ retrievals, *Environ. Res. Lett.*, 17, 024006, <https://doi.org/10.1088/1748-9326/ac48b4>, 2022.
- de Foy, B., Wilkins, J. L., Lu, Z., Streets, D. G., and Duncan, B. N.: Model evaluation of methods for estimating surface emissions and chemical lifetimes from satellite data, *Atmos. Environ.*, 98, 66–77, <https://doi.org/10.1016/j.atmosenv.2014.08.051>, 2014.
- 365 van Geffen, J., Eskes, H., Compernelle, S., Pinardi, G., Verhoelst, T., Lambert, J.-C., Sneep, M., ter Linden, M., Ludewig, A., Boersma, K. F., and Veeffkind, J. P.: Sentinel-5P TROPOMI NO₂ retrieval: impact of version v2.2 improvements and comparisons with OMI and ground-based data, *Atmospheric Meas. Tech.*, 15, 2037–2060, <https://doi.org/10.5194/amt-15-2037-2022>, 2022.
- Geddes, J. A., Murphy, J. G., O'Brien, J. M., and Celarier, E. A.: Biases in long-term NO₂ averages inferred from satellite observations due to cloud selection criteria, *Remote Sens. Environ.*, 124, 210–216, <http://dx.doi.org/10.1016/j.rse.2012.05.008>, 2012.
- 370 Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G., Reichle, R., Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V., Conaty, A., Silva, A. M. da, Gu, W., Kim, G.-K., Koster, R., Lucchesi, R., Merkova, D., Nielsen, J. E., Partyka, G., Pawson, S., Putman, W., Rienecker, M., Schubert, S.

- 375 D., Sienkiewicz, M., and Zhao, B.: The Modern-Era Retrospective analysis for Research and Applications, version 2 (MERRA-2), *J. Clim.*, 30, 5419–5454, <https://doi.org/10.1175/JCLI-D-16-0758.1>, 2017.
- Goldberg, D. L., Lu, Z., Streets, D. G., de Foy, B., Griffin, D., McLinden, C. A., Lamsal, L. N., Krotkov, N. A., and Eskes, H.: Enhanced capabilities of TROPOMI NO₂: Estimating NO_x from North American cities and power plants, *Environ. Sci. Technol.*, 53, 12594–12601, <https://doi.org/10.1021/acs.est.9b04488>, 2019.
- 380 Goldberg, D. L., Anenberg, S. C., Griffin, D., McLinden, C. A., Lu, Z., and Streets, D. G.: Disentangling the Impact of the COVID-19 Lockdowns on Urban NO₂ From Natural Variability, *Geophys. Res. Lett.*, 47, e2020GL089269, <https://doi.org/10.1029/2020GL089269>, 2020.
- Henze, D. K., Seinfeld, J. H., and Shindell, D. T.: Inverse modeling and mapping US air quality influences of inorganic PM_{2.5} precursor emissions using the adjoint of GEOS-Chem, *Atmospheric Chem. Phys.*, 9, 5877–5903, [https://doi.org/10.5194/acp-](https://doi.org/10.5194/acp-9-5877-2009)
- 385 9-5877-2009, 2009.
- Hogue, S., Marland, E., Andres, R. J., Marland, G., and Woodard, D.: Uncertainty in gridded CO₂ emissions estimates, *Earths Future*, 4, 225–239, <https://doi.org/10.1002/2015EF000343>, 2016.
- Ialongo, I., Virta, H., Eskes, H., Hovila, J., and Douros, J.: Comparison of TROPOMI/Sentinel-5 Precursor NO₂ observations with ground-based measurements in Helsinki, *Atmospheric Meas. Tech.*, 13, 205–218, [https://doi.org/10.5194/amt-13-205-](https://doi.org/10.5194/amt-13-205-2020)
- 390 2020, 2020.
- Ingmann, P., Veihelmann, B., Langen, J., Lamarre, D., Stark, H., and Courrèges-Lacoste, G. B.: Requirements for the GMES atmosphere service and ESA's implementation concept: Sentinels-4/-5 and -5p, *Remote Sens. Environ.*, 120, 58–69, <https://doi.org/10.1016/j.rse.2012.01.023>, 2012.
- Judd, L. M., Al-Saadi, J. A., Szykman, J. J., Valin, L. C., Janz, S. J., Kowalewski, M. G., Eskes, H. J., Veeffkind, J. P., Cede,
- 395 A., Mueller, M., Gebetsberger, M., Swap, R., Pierce, R. B., Nowlan, C. R., Abad, G. G., Nehrir, A., and Williams, D.: Evaluating Sentinel-5P TROPOMI tropospheric NO₂ column densities with airborne and Pandora spectrometers near New York City and Long Island Sound, *Atmos Meas Tech*, 13, 6113–6140, <https://doi.org/10.5194/amt-13-6113-2020>, 2020.
- Kim, J.: GEMS (Geostationary Environment Monitoring Spectrometer) onboard the GeoKOMPSAT to monitor air quality in high temporal and spatial resolution over Asia-Pacific Region, *EGU General Assembly Conference Abstracts*, 4051, 2012.
- 400 Lamarque, J.-F., Emmons, L. K., Hess, P. G., Kinnison, D. E., Tilmes, S., Vitt, F., Heald, C. L., Holland, E. A., Lauritzen, P. H., Neu, J., Orlando, J. J., Rasch, P. J., and Tyndall, G. K.: CAM-chem: description and evaluation of interactive atmospheric chemistry in the Community Earth System Model, *Geosci. Model Dev.*, 5, 369–411, <https://doi.org/10.5194/gmd-5-369-2012>, 2012.
- Lamsal, L. N., Martin, R. V., Padmanabhan, A., van Donkelaar, A., Zhang, Q., Sioris, C. E., Chance, K., Kurosu, T. P., and
- 405 Newchurch, M. J.: Application of satellite observations for timely updates to global anthropogenic NO_x emission inventories, *Geophys. Res. Lett.*, 38, L05810, <https://doi.org/doi:10.1029/2010gl046476>, 2011.

- Lamsal, L. N., Krotkov, N. A., Marchenko, S. V., Joiner, J., Oman, L., Vasilkov, A., Fisher, B., Qin, W., Yang, E.-S., Fasnacht, Z., Choi, S., Leonard, P., and Haffner, D.: TROPOMI/S5P NO₂ tropospheric, stratospheric and total columns MINDS 1-Orbit L2 Swath 5.5 km x 3.5 km, last accessed May, 2023, doi: 10.5067/MEASURES/MINDS/DATA203, 2022.
- 410 Laughner, J. L., Zare, A., and Cohen, R. C.: Effects of daily meteorology on the interpretation of space-based remote sensing of NO₂, *Atmos. Chem. Phys.*, 16, 15247–15264, <https://doi.org/10.5194/acp-16-15247-2016>, 2016.
- Laughner, J. L. and Cohen, R. C.: Direct observation of changing NO_x lifetime in North American cities, *Science*, 366, 723–727, <https://doi.org/10.1126/science.aax6832>, 2019.
- Liu, F., Beirle, S., Zhang, Q., Dörner, S., He, K., and Wagner, T.: NO_x lifetimes and emissions of cities and power plants in
415 polluted background estimated by satellite observations, *Atmospheric Chem. Phys.*, 16, 5283–5298, <https://doi.org/doi:10.5194/acp-16-5283-2016>, 2016a.
- Liu, F., Zhang, Q., A, R. J. van der, Zheng, B., Tong, D., Yan, L., Zheng, Y., and He, K.: Recent reduction in NO_x emissions over China: synthesis of satellite observations and emission inventories, *Environ. Res. Lett.*, 11, 114002, <https://doi.org/10.1088/1748-9326/11/11/114002>, 2016b.
- 420 Liu, F., Beirle, S., Zhang, Q., van der A, R. J., Zheng, B., Tong, D., and He, K.: NO_x emission trends over Chinese cities estimated from OMI observations during 2005 to 2015, *Atmospheric Chem. Phys.*, 17, 9261–9275, <https://doi.org/doi:10.5194/acp-17-9261-2017>, 2017.
- Liu, F., Choi, S., Li, C., Fioletov, V. E., McLinden, C. A., Joiner, J., Krotkov, N. A., Bian, H., Janssens-Maenhout, G., Darmenov, A. S., and da Silva, A. M.: A new global anthropogenic SO₂ emission inventory for the last decade: a mosaic of
425 satellite-derived and bottom-up emissions, *Atmospheric Chem. Phys.*, 18, 16571–16586, <https://doi.org/doi:10.5194/acp-18-16571-2018>, 2018.
- Liu, F., Page, A., Strode, S. A., Yoshida, Y., Choi, S., Zheng, B., Lamsal, L. N., Li, C., Krotkov, N. A., Eskes, H., van der A, R., Veefkind, P., Levelt, P. F., Hauser, O. P., and Joiner, J.: Abrupt decline in tropospheric nitrogen dioxide over China after the outbreak of COVID-19, *Sci. Adv.*, 6, eabc2992, <https://doi.org/10.1126/sciadv.abc2992>, 2020.
- 430 Liu, F., Tao, Z., Beirle, S., Joiner, J., Yoshida, Y., Smith, S. J., Knowland, K. E., and Wagner, T.: A new method for inferring city emissions and lifetimes of nitrogen oxides from high-resolution nitrogen dioxide observations: a model study, *Atmospheric Chem. Phys.*, 22, 1333–1349, <https://doi.org/10.5194/acp-22-1333-2022>, 2022.
- Lu, Z., Streets, D. G., de Foy, B., Lamsal, L. N., Duncan, B. N., and King, J.: Emissions of nitrogen oxides from US urban areas: estimation from Ozone Monitoring Instrument retrievals for 2005–2014, *Atmospheric Chem. Phys.*, 15, 10367–10383,
435 <https://doi.org/doi:10.5194/acp-15-10367-2015>, 2015.
- Lucchesi, R.: File Specification for GEOS-5 FP-IT. GMAO Office Note No. 2 (Version 1.3), 60 pp, available from http://gmao.gsfc.nasa.gov/pubs/office_notes, 2015.
- Ma, S. and Tong, D. Q.: Neighborhood Emission Mapping Operation (NEMO): A 1-km anthropogenic emission dataset in the United States, *Sci. Data*, 9, 680, <https://doi.org/10.1038/s41597-022-01790-9>, 2022.

- 440 Martin, R. V., Jacob, D. J., Chance, K., Kurosu, T. P., Palmer, P. I., and Evans, M. J.: Global inventory of nitrogen oxide emissions constrained by space-based observations of NO₂ columns, *J. Geophys. Res.*, 108, 4537, <https://doi.org/doi:10.1029/2003jd003453>, 2003.
- McDonald, B. C., McKeen, S. A., Cui, Y. Y., Ahmadov, R., Kim, S.-W., Frost, G. J., Pollack, I. B., Peischl, J., Ryerson, T. B., Holloway, J. S., Graus, M., Warneke, C., Gilman, J. B., de Gouw, J. A., Kaiser, J., Keutsch, F. N., Hanisco, T. F., Wolfe,
445 G. M., and Trainer, M.: Modeling ozone in the eastern U.S. using a fuel-based mobile source emissions inventory, *Environ. Sci. Technol.*, 52, 7360–7370, <https://doi.org/10.1021/acs.est.8b00778>, 2018.
- Miyazaki, K., Eskes, H., Sudo, K., Boersma, K. F., Bowman, K., and Kanaya, Y.: Decadal changes in global surface NO_x emissions from multi-constituent satellite data assimilation, *Atmospheric Chem. Phys.*, 17, 807–837, <https://doi.org/doi:10.5194/acp-17-807-2017>, 2017.
- 450 Miyazaki, K., Bowman, K., Sekiya, T., Takigawa, M., Neu, J. L., Sudo, K., Osterman, G., and Eskes, H.: Global tropospheric ozone responses to reduced NO_x emissions linked to the COVID-19 worldwide lockdowns, *Sci. Adv.*, 7, eabf7460, <https://doi.org/10.1126/sciadv.abf7460>, 2021.
- Peters-Lidard, C. D., Kemp, E. M., Matsui, T., Santanello, J. A., Kumar, S. V., Jacob, J. P., Clune, T., Tao, W.-K., Chin, M., Hou, A., Case, J. L., Kim, D., Kim, K.-M., Lau, W., Liu, Y., Shi, J., Starr, D., Tan, Q., Tao, Z., Zaitchik, B. F., Zavadsky, B.,
455 Zhang, S. Q., and Zupanski, M.: Integrated modeling of aerosol, cloud, precipitation and land processes at satellite-resolved scales, *Environ. Model. Softw.*, 67, 149–159, <https://doi.org/10.1016/j.envsoft.2015.01.007>, 2015.
- Platt, U. and Stutz, J.: *Differential absorption spectroscopy*, 91–99, 135–159, Berlin and Heidelberg, Germany, Springer, 2008.
- Qu, Z., Henze, D. K., Theys, N., Wang, J., and Wang, W.: Hybrid mass balance/4D-Var joint inversion of NO_x and SO₂ emissions in East Asia, *J. Geophys. Res.*, 124, 8203–8224, <https://doi.org/10.1029/2018JD030240>, 2019.
- 460 Seinfeld, J. H. and Pandis, S. N.: *Atmospheric chemistry and physics: From air pollution to climate change*, 2nd ed., John Wiley and Sons, New York, 204–275 pp., 2006.
- Souri, A. H., Choi, Y., Jeon, W., Li, X., Pan, S., Diao, L., and Westenbarger, D. A.: Constraining NO_x emissions using satellite NO₂ measurements during 2013 DISCOVER-AQ Texas campaign, *Atmos. Environ.*, 131, 371–381, <https://doi.org/10.1016/j.atmosenv.2016.02.020>, 2016.
- 465 Sun, K.: Derivation of emissions from satellite-observed column amounts and its application to TROPOMI NO₂ and CO observations, *Geophys. Res. Lett.*, 49, e2022GL101102, <https://doi.org/10.1029/2022GL101102>, 2022.
- Tack, F., Merlaud, A., Iordache, M.-D., Pinardi, G., Dimitropoulou, E., Eskes, H., Bomans, B., Veeffkind, P., and Van Roozendaal, M.: Assessment of the TROPOMI tropospheric NO₂ product based on airborne APEX observations, *Atmospheric Meas. Tech.*, 14, 615–646, <https://doi.org/10.5194/amt-14-615-2021>, 2021.
- 470 Tao, Z., Santanello, J. A., Chin, M., Zhou, S., Tan, Q., Kemp, E. M., and Peters-Lidard, C. D.: Effect of land cover on atmospheric processes and air quality over the continental United States – a NASA Unified WRF (NU-WRF) model study, *Atmospheric Chem. Phys.*, 13, 6207–6226, <https://doi.org/10.5194/acp-13-6207-2013>, 2013.


- Tao, Z., Kawa, S. R., Jacob, J. P., Liu, D. Y., Collatz, G. J., Wang, J. S., Ott, L. E., and Chin, M.: Application of NASA- Unified WRF model to carbon dioxide simulation- model development and evaluation, *Environ. Model. Softw.*, 132, 104785, 475 <https://doi.org/10.1016/j.envsoft.2020.104785>, 2020.
- Travis, K. R., Jacob, D. J., Fisher, J. A., Kim, P. S., Marais, E. A., Zhu, L., Yu, K., Miller, C. C., Yantosca, R. M., Sulprizio, M. P., Thompson, A. M., Wennberg, P. O., Crounse, J. D., St. Clair, J. M., Cohen, R. C., Laughner, J. L., Dibb, J. E., Hall, S. R., Ullmann, K., Wolfe, G. M., Pollack, I. B., Peischl, J., Neuman, J. A., and Zhou, X.: Why do models overestimate surface ozone in the Southeast United States?, *Atmos Chem Phys*, 16, 13561–13577, <https://doi.org/10.5194/acp-16-13561-2016>, 2016.
- 480 Veefkind, J. P., Aben, I., McMullan, K., Förster, H., de Vries, J., Otter, G., Claas, J., Eskes, H. J., de Haan, J. F., Kleipool, Q., van Weele, M., Hasekamp, O., Hoogeveen, R., Landgraf, J., Snel, R., Tol, P., Ingmann, P., Voors, R., Kruizinga, B., Vink, R., Visser, H., and Levelt, P. F.: TROPOMI on the ESA Sentinel-5 Precursor: A GMES mission for global observations of the atmospheric composition for climate, air quality and ozone layer applications, *Remote Sens. Environ.*, 120, 70–83, 2012.
- Verhoelst, T., Compernelle, S., Pinardi, G., Lambert, J.-C., Eskes, H. J., Eichmann, K.-U., Fjæraa, A. M., Granville, J., 485 Niemeijer, S., Cede, A., Tiefengraber, M., Hendrick, F., Pazmiño, A., Bais, A., Bazureau, A., Boersma, K. F., Bognar, K., Dehn, A., Donner, S., Elokhov, A., Gebetsberger, M., Goutail, F., Grutter de la Mora, M., Gruzdev, A., Gratsea, M., Hansen, G. H., Irie, H., Jepsen, N., Kanaya, Y., Karagkiozidis, D., Kivi, R., Kreher, K., Levelt, P. F., Liu, C., Müller, M., Navarro Comas, M., PETERS, A. J. M., Pommereau, J.-P., Portafaix, T., Prados-Roman, C., Puentedura, O., Querel, R., Remmers, J., Richter, A., Rimmer, J., Rivera Cárdenas, C., Saavedra de Miguel, L., Sinyakov, V. P., Stremme, W., Strong, K., Van 490 Roozendaal, M., Veefkind, J. P., Wagner, T., Wittrock, F., Yela González, M., and Zehner, C.: Ground-based validation of the Copernicus Sentinel-5P TROPOMI NO₂ measurements with the NDACC ZSL-DOAS, MAX-DOAS and Pandonia global networks, *Atmospheric Meas. Tech.*, 14, 481–510, <https://doi.org/10.5194/amt-14-481-2021>, 2021.
- Wang, P., PETERS, A., van Geffen, J., Tuinder, O., Stammes, P., and Kinne, S.: Shipborne MAX-DOAS measurements for validation of TROPOMI NO₂ products, *Atmospheric Meas. Tech.*, 13, 1413–1426, <https://doi.org/10.5194/amt-13-1413-2020>, 495 2020.
- Woodard, D., Branham, M., Buckingham, G., Hogue, S., Hutchins, M., Gosky, R., Marland, G., and Marland, E.: A spatial uncertainty metric for anthropogenic CO₂ emissions, *Greenh. Gas Meas. Manag.*, 4, 139–160, <https://doi.org/10.1080/20430779.2014.1000793>, 2014.



500

Figure 1: NO_x budget inferred from TROPOMI NO_2 observations around New York City from May through September, 2019. (a) sinks S , (b) divergence D , (c) derived NO_x emissions $E=S+D$, (d) NEI NO_x emissions E_{NEI} . The locations of Glenwood Landing power station, JFK airport, and the city center of Newark are labeled as PP, JFK, and Newark, respectively. Wind barbs at TROPOMI overpass time from May to September of 2019 are averaged and shown in (b). Wind speed is given in the units of knots, which is a nautical miles per hour (1.9 km per hour). Each short and long barb represents 5 knots (9.3 km/h) and 10 knots (18.5 km/h), respectively.


505




510

Figure 2: Scatterplot of the derived NO_x total emissions for the investigated cities based on the NO_2 tropospheric VCDs simulated by NU-WRF (y axis) as compared to the given emissions used to drive the NU-WRF simulation (x axis). NO_x emissions from all grid cells within the domain of $70 \text{ km} \times 70 \text{ km}$ around city center are summed up to derive the total emission for most cities; a $100 \text{ km} \times 100 \text{ km}$ domain is used for New York, Chicago, Los Angeles and Houston. Error bars show the standard error of the


515 derived emissions for all wind directions with derived NO_x lifetime τ . Standard error is defined as standard deviation divided by \sqrt{n} , with n being the number of wind directions with derived NO_x lifetime τ . The intracity spatial correlation $R_{intracity}$ between the derived and given emissions for individual cities are color coded. The dashed line represents the 1:1 line. Statistics are provided in the inset table.




520 **Figure 3: Improved spatial correlation of derived and given NO_x emissions compared to that of derived NO_x emissions and NO_2 columns. (a) Mean NU-WRF tropospheric NO_2 VCDs $\Omega_{\text{NU-WRF}}$; (b) Mean NU-WRF tropospheric NO_2 VCDs under calm wind conditions only; (c) Mean NO_x emission rates $E_{\text{NU-WRF}}$ derived from (a); (d) Mean NO_x emission rates used to drive the NU-WRF simulation $E'_{\text{NU-WRF}}$. Hourly mean data at 14:00 LT are averaged from May through September 2016. The city of Jacksonville, Florida is in the centre of the domain shown.**



525 **Figure 4: Similar to Figure 2, but for the comparison between the derived NO_x total emissions based on TROPOMI tropospheric NO_2 VCDs (y axis) with NEI total emissions (x axis) for 2019. Error bars show the uncertainties of NEI (30%) and fitted (47%) emissions.**



530 **Figure 5: Comparison of NO_x emissions inferred from TROPOMI NO_2 observations with NEI emissions from May through September, 2019. Dallas: (a) TROPOMI-derived NO_x emissions E , (b) NEI NO_x emissions E_{NEI} . The locations of Dallas/Fort Worth International Airport, Perot Field Fort Worth Alliance Airport and the city center of Fort Worth are labeled as DFW, AFW, and Fort Worth, respectively. The locations of three large cement factories (TXI, Holcim, and Ash Grove) are labeled as dots. Tucson: (c) TROPOMI-derived NO_x emissions E , (d) NEI NO_x emissions E_{NEI} . The location of a large copper mine Asarco Mission is labeled as M.**



535 **Figure 6: Comparison of TROPOMI-derived NO_x emission estimates for (a) 2019, (b) 2020, (c) 2021 with those for 2018. The dashed line represents the 1:1 line. The slope of the least-squares linear regression line is provided in the figure.**