

Changing effects of external forcing on Atlantic-Pacific interactions

Soufiane Karmouche^{1,2}, Evgenia Galytska^{1,2}, Gerald A. Meehl³, Jakob Runge^{4,5}, Katja Weigel^{1,2}, and Veronika Eyring^{2,1}

¹University of Bremen, Institute of Environmental Physics (IUP), Bremen, Germany

²Deutsches Zentrum für Luft- und Raumfahrt e.V. (DLR), Institut für Physik der Atmosphäre, Oberpfaffenhofen, Germany

³Climate and Global Dynamics Laboratory, National Center for Atmospheric Research (NCAR), Boulder, CO, USA

⁴Deutsches Zentrum für Luft- und Raumfahrt e.V. (DLR), Institut für Datenwissenschaften, Jena, Germany

⁵Fachgebiet Klimainformatik, Technische Universität Berlin, Berlin, Germany

Correspondence: Soufiane Karmouche (sou_kar@uni-bremen.de)

1 Supplementary Material

Table S1. Directing unoriented contemporaneous links based on most detected direction within a specific run among the 10 simulations of the CESM2 pacemaker ensemble (Fig. 7).

Unoriented Links	Instances	Direction	Ruling
PNA—NAO	Fig. 5d	NAO→PNA	Most detected direction for "1985-2014 minus MEM" regime
Niño3.4—PNA	Fig. 5d	Niño3.4→PNA	Based on the 1-season lagged instances in all regimes
Niño3.4—PWC _u	P1 and P2 in Fig. 8b	PWC_u→Niño3.4	Most detected direction in all regimes

Atlantic-driven period (1986-2014)




Figure S1. Atlantic-driven regime starting in 1986 (instead of 1985) shows vanishing TNA → PWC_u link when MEM is subtracted.
 a) Similar to Fig. 5b but for 1986-2014. b) Similar to Fig. 5d but for 1986-2014 period