

# Influence of Lower Tropospheric Moisture on Local Soil Moisture-Precipitation Feedback over the U.S. Southern Great Plains

Gaoyun Wang<sup>1,2,3</sup>, Rong Fu<sup>2</sup>, Yizhou Zhuang<sup>2</sup>, Paul A. Dirmeyer<sup>4</sup>, Joseph A. Santanello<sup>5</sup>, Guiling Wang<sup>6</sup>, Kun Yang<sup>7</sup>, Kaighin McColl<sup>8</sup>

<sup>1</sup>Department of Atmospheric and Oceanic Sciences, School of Physics, Peking University, Beijing, China

<sup>2</sup>Department of Atmospheric and Oceanic Sciences, University of California, Los Angeles, CA, USA

<sup>3</sup>The High School Affiliated to Southern University of Science and Technology, Shenzhen, China

<sup>4</sup>Department of Atmospheric, Oceanic and Earth Sciences, George Mason University, Fairfax, VA, USA

<sup>5</sup>NASA Godard Space Flight Center, Greenbelt, MD, USA

<sup>6</sup>Department of Civil and Environmental Engineering, University of Connecticut, Storrs, CT, USA

<sup>7</sup>Department of Earth System Science, Tsinghua University, Beijing, China

<sup>8</sup>Department of Earth and Planetary Sciences, Harvard University, Cambridge, MA, USA

Correspondence to: Yizhou Zhuang ([zhuangyz@atmos.ucla.edu](mailto:zhuangyz@atmos.ucla.edu))

**Abstract.** Land-atmosphere coupling (LAC) has long been studied focusing on land surface and atmospheric boundary layer processes. However, the influence of ~~lower tropospheric (LT)~~ humidity in the lower troposphere (LT), especially that above the planetary boundary layer (PBL), on LAC remains largely unexplored. In this study, we use radiosonde observations from the U.S. Southern Great Plains (SGP) site and an entrained parcel buoyancy model to investigate the impact of LT humidity on LAC there during the warm season (May-September). We quantify the effect of LT humidity on convective buoyancy by measuring the difference between the 2-4 km vertically integrated buoyancy with and without the influence of background LT humidity. Our results show that, under dry soil conditions, anomalously high LT humidity is necessary to produce the buoyancy profiles required for afternoon precipitation events (APEs). These APEs under dry soil moisture cannot be explained by commonly used local land-atmosphere coupling indices such as the convective triggering potential/low-level humidity index (CTP/HI<sub>Low</sub>), which do not account for the influence of the LT humidity. On the other hand, consideration of LT humidity is unnecessary to explain APEs under wet soil moisture conditions, suggesting the boundary layer moisture alone could be sufficient to generate the required buoyancy profiles. These findings highlight the need to consider the impact of LT humidity, which is often decoupled from the humidity near the surface and largely controlled by moisture transport, in understanding land-atmospheric feedbacks over dry soil conditions, especially during droughts or dry spells over the SGP.

## 1 Introduction

Land-atmosphere coupling (LAC) plays an important role in determining local and regional climate variability, including surface temperature, humidity, cloud, precipitation, and climate extremes such as drought and floods, especially during the warm season over interior continents (e.g., Fernando et al., 2016; Koster et al., 2004, 2006; Guo

et al., 2006; Wang et al., 2007; Roundy and Santanello, 2017; Santanello et al., 2009; Konings et al., 2010; Song et al., 2016; Roundy et al., 2013). To provide a consistent characterization of land-atmosphere coupling, the International Global Energy and Water Exchanges Project (GEWEX) developed the Local Land-Atmosphere Coupling (LoCo) initiative to coordinate and promote process-level metrics that quantify and characterize LAC (Santanello et al., 2018).

40 The LoCo initiative develops a suite of integrative metrics to quantify the complex relationships and feedback between the land surface and atmosphere. For example, the mixing diagram approach (Santanello et al., 2009) relates the daytime coevolution of 2-m potential temperature and humidity to the energy and water budgets and growth of the planetary boundary layer (PBL). The convective triggering potential/low-level humidity index (CTP/ $HI_{Low}$ ; Findell & Eltahir, 2003) characterizes the lower tropospheric lapse rate and dewpoint depression of the PBL for convection.

45 The heated condensation framework (HCF, Tawfik et al., 2015a, 2015b) diagnoses the contribution of surface fluxes to convective initiation based on temperature and humidity profiles. The soil moisture-precipitation (SM-P) feedback is one of the most extensively studied land-atmospheric feedbacks in the literature (e.g., Koster et al., 2004; Ferguson & Wood, 2010; Roundy & Santanello, 2017; Santanello et al., 2018), particularly regarding its effects on the frequency and intensity of convective precipitation (e.g. Taylor, 2015; Tuttle & Salvucci, 2016; Yin et al., 2015).

50 LoCo investigates the links in the chain coupling soil moisture with the PBL, which connect through surface fluxes, 2-meter temperature and humidity, PBL growth and entrainment, cloud, and precipitation. However, the humidity in the lower troposphere (LT) above the PBL, i.e., ~2-4 km above ground level (AGL), is not explicitly included in previous research. Recent research indicates that specific humidity in the LT ( $q_{LT}$ ) plays a central role in triggering (or the development of) deep convection in the tropics, subtropics, and mid-latitudes during the warm season (Bretherton et al., 2004; Holloway and Neelin, 2009; Zhang and Klein, 2010; Zhuang et al., 2018) and in the convective initiation driven by land surface heating (Tawfik et al., 2015b, a). The lateral entrainment of  $q_{LT}$  dominates buoyancy above the PBL, which is crucial for deep convection development, while entrainment of air at the cloud base has a stronger influence on shallow convection (Holloway and Neelin, 2009; Mapes et al., 2006).

60 A moist LT can enhance convection by entraining moist air plumes, while low LT humidity can dilute moist plumes originating from the PBL, thereby interfering with the surface influence on convection and precipitation. Thus,  $q_{LT}$  determines whether shallow convection can develop into deep convection locally (Schiro et al., 2016; Zhang and Klein, 2010; Zhuang et al., 2017, 2018) and the occurrence and intensity of mesoscale convection (Schiro et al., 2018).  $q_{LT}$  is influenced by moisture transport from the PBL (which is largely influenced by land surface), horizontal moisture advection, and subsidence that mixes dry air from aloft. Therefore, it is important to study the relative influences of

65 land surface versus large-scale atmospheric circulation on rainfall and clouds.

The Department of Energy's Atmospheric Radiation Measurement (DOE ARM) project has been pivotal in providing comprehensive datasets for investigating land-atmospheric interactions over the past two decades (e.g., Zhang and Klein, 2010; Santanello et al., 2018). Among the various ARM sites, the Southern Great Plains (SGP) site stands out as the project's inaugural site and one of the most heavily instrumented sites. The SGP region is also widely known as a hotspot of land-atmosphere interactions, as evidenced by numerous past research (e.g., Wakefield et al., 2019; Santanello et al., 2018; Dimmeyer et al., 2006; Koster et al., 2004; Koster et al., 2006; Guo et al., 2006). In this study, we explore the effect-impact of LT humidity on the SM-P relationship and local

70

LAC at the SGP site by utilizing an entrained parcel buoyancy model (Zhuang et al. 2018) and the correlation between LT humidity and near-surface humidity to determine whether it can significantly influence the local LAC. The dataset and methods are described in section 2. The results are reported in section 3. Discussion and conclusions are provided in section 4.

## 2 Data and method

### 2.1 Dataset

This study focuses on the local warm season (May-September) when thermodynamically driven convection is most prevalent and land-surface feedback plays an important role in determining precipitation (Myoung and Nielsen-Gammon, 2010). Unless stated otherwise, all measurements are taken at the U.S. Department of Energy's Atmospheric Radiation Measurement (DOE ARM) Southern Great Plain (SGP) central facility (CF) in north-central Oklahoma (36.60°N, 97.48°W), and the region within a 50-km radius of the CF for the period of 2001-2018. Below are the specific details about the datasets used in this study.

#### 2.1.1 Sounding profiles

Sounding profile data at the SGP CF were obtained through balloon sonde observation. This data is available four times daily at 05:30, 11:30, 17:30, and 23:30 local standard time (LST). We only use the 11:30 LST sounding data immediately before noon, specifically at 11:30 local standard time (LST), were used in this study as it they best represents the preconditions of afternoon convection. Because the vertical levels vary with each sounding, data were re-gridded into a uniform vertical resolution of 20 m to facilitate composite analysis. Profiles of the dry-bulb temperature ( $T$ ), dew point temperature ( $T_d$ ), and atmospheric pressure ( $p$ ) were used to calculate the mixing ratio ( $r$ ), specific humidity ( $q$ ), and buoyancy ( $b$ ) using the entrained parcel model (described in section 2.2.1). The data used are available online at <https://www.arm.gov/capabilities/instruments/sonde>.

#### 2.1.2 Soil moisture

Fractional water index (FWI) is a normalized measurement specifically developed for the Campbell 229-L soil moisture sensor and ranges from 0 for very dry soil to 1 for saturated soil (Schneider et al., 2003). FWI can capture soil wetness independent of soil texture, so it standardizes the observation and allows for intercomparison among different sites with different soil types. Most root biomass in the SGP region and its vicinity is within the top 30 cm of the soil profile (e.g., Raz-Yaseef et al., 2015). Because evapotranspiration, a vital link in the SM-P relationship, is heavily influenced by plant and root zone soil moisture, we used FWI at 25 cm measurement depth provided by from a surface station near the SGP CF. The 30-min averaged Oklahoma Mesonet Soil Moisture (OKMSOIL) value-added product (VAP) was used to represent soil wetness (available at <https://www.arm.gov/capabilities/vaps/okmsoil>). This data has a 30-min resolution, and we use the average FWI during 06:00-12:00 LST to represent soil moisture condition before afternoon precipitation at daily scale. Wet soils

105 are defined as those with FWI greater than 0.7, which is considered optimal for the plant, and dry soils are defined as  
FWI smaller than 0.4, which could result in water stress and plant wilting (Illston et al., 2008; Wakefield et al., 2019).

### 2.1.3 Precipitation

110 ~~We utilize hourly gridded precipitation products from The~~ Arkansas-Red Basin River Forecast Center (ABRFC)  
~~precipitation data is~~ based on WSR-88D Nexrad radar precipitation estimates ~~combined with~~ and rain gauge reports  
with extensive quality control (Fulton et al., 1998). ~~This is an hourly gridded data product and~~ is available ~~online~~ at  
<https://www.arm.gov/capabilities/vaps/abrfc>. We used spatially averaged data over the region within a 50 km radius  
of the SGP CF for this study.

### 2.1.4 PBL height

115 ~~We applied the boundary layer regime identified by the criteria defined in Liu and Liang (2010) to estimate the PBL~~  
~~height by using gradients of potential temperature profiles from the sounding observations that are available four times~~  
~~per day.~~ PBL height data are obtained from the ARM's Planetary Boundary Layer Height (PBLHT) value-added  
~~products derived from radiosonde data using the algorithm developed by Liu and Liang (2010).~~ This data is available  
at <https://www.arm.gov/capabilities/vaps/pblht>.

## 2.2 Quantifying contributions of surface and LT humidity to convective buoyancy

120 Previous research on local land-atmosphere interaction mainly focused on the influence of surface flux and moisture  
in the PBL on convection initiation and precipitation, such as those related to HCF and mixing diagram metrics. The  
CTP/ $H_{Low}$  metric considers the effect of lapse rate 100-300 hPa (or about 2-4 km) above ground level (AGL) on  
vertically integrated buoyancy in the LT (i.e., CTP) and humidity of the PBL (i.e.,  $H_{Low}$ ), but it does not account for  
the impact of LT moisture.

125 Isolating the local influence from other factors in observation ~~presents a significant is-challenging.~~ ~~Understanding~~  
~~the relationship between near-surface and upper-level information could be crucial to address this.~~ In this study, we  
~~use first examine~~ the correlation between specific humidity ( $q$ ) profile in the LT and ~~the mixed-layer humidity ( $q_m$ ),~~  
~~defined as average  $q$  in the 0-1 km AGL mixed layer, that at the near-surface layer to assess infer~~ the potential influence  
of the land surface on the LT moisture (Figure 1). ~~As expected, the correlation between the  $q$  near the surface and~~  
130 ~~that in the PBL is significant and larger than 0.95.~~ We choose mixed layer humidity over humidity directly above  
surface to represent land surface moisture condition because: 1) radiosonde measurements near the surface are often  
more susceptible to errors and local disturbances, which could skew the representation of actual surface moisture  
condition; 2) at noon, 0-1 km mixed layer offers a more representative snapshot of the land surface moisture by  
capturing the integrated effect of surface evaporation and convective mixing process; 3) we observe strong  
135 correlations, exceeding 0.95 ( $p < 0.05$ ), between the  $q$  near the surface and  $q_m$ . Above the PBL However, these  
correlation between  $q$  of LT and  $q$  in the near-surface layer decreases rapidly with height ~~diminishes with increasing~~  
height above the PBL. Notably, ~~the~~ LT humidity above 2 km maintains a ~~is still significantly~~ correlated to near-  
surface moisture with  $q_m$  above 2km, ~~suggesting a which indicates~~ the potential influence from the surface. To separate

Formatted: Font: Italic, Not Highlight

Formatted: Font: Italic, Not Highlight

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Subscript

isolate the effect of the land surface on LT humidity, we establish define a “land-coupled LT humidity profile  $q_{LC}$ ” from for 2–km to 4 km AGL, which is linked related to the land surface moisture condition. This profile which is derived reconstructed from the using a linear regression between  $q(h,t)$  profile of within this layer ( $2\text{ km} \leq h \leq 4\text{ km}$ ) and  $q_m(t)$  the averaged  $q$  within the mixed layer. In our regression model, represented by the equation  $y = a \times x + b$ ,  $y$  is  $q$  at a given height and time  $q(h,t)$ , and  $x$  is  $q_m(t)$ , with  $a(h)$  and  $b(h)$  being the linear coefficients at each height level. By solving  $a(h)$  and  $b(h)$  for each height level in the LT, we can then calculate the “land-coupled LT humidity” as the fitted LT humidity, i.e.,  $q_{LC}(h,t) = \hat{q}(h,t) = a(h) * q_m(t) + b(h)$ .

To quantify the direct influence of LT moisture on convective buoyancy, we adopt an entraining parcel model used in Zhuang et al. (2018). In this model, the air parcel is lifted with the initial condition of average value within the mixed layer. The ascending air parcel then goes through three processes at each vertical level: dry adiabatic process (parcel ascends without interacting with the environment, entropy conservation), entrainment process (interacts with ambient air, enthalpy conservation), precipitation process (releases condensate, temperature conservation). We apply the deep inflow-A entrainment (DIA) scheme which has been shown to more realistically represent the buoyancy profile required for deep convection compared to the other assumptions of the lateral entrainment rates such as the constant fractional entrainment rate scheme (e.g., Holloway & Neelin, 2009; Schiro et al., 2016; Siebesma et al., 2007). The DIA scheme uses an entrainment rate inversely proportional to the altitude ( $\alpha z^{-1}$ ) for simplicity. All condensates formed from the previous two processes are set to fall out in the precipitation process (pseudo-adiabatic process). And finally, buoyancy is calculated by  $b = g \frac{T_{pv} - T_{ev}}{T_{ev}}$ , where  $T_{pv}$  and  $T_{ev}$  are the virtual temperature of the parcel and the environment, respectively. Details of this model are provided in Zhuang et al. (2018).

To quantify the influence of LT humidity on the lateral entrainment of the convection and, consequently, the buoyancy of the convective air parcel, we consider two  $q$  profiles for the lateral entrainment process: 1) the observed humidity profile ( $q_R$ ) and 2) the land-coupled humidity profile ( $q_{LC}$ ). The  $q_{LC}$  below 2 km AGL equals to the observed  $q$ , and  $q_{LC}$  between 2 km and 4 km AGL is a coupled LT humidity profile constructed from the regression between  $q$  and averaged  $q$  in the mixed layer (as described in 2.2). Since the effect of entrainment accumulates continuously after the parcel is lifted, we calculate the buoyancy profile with  $q_R$  as humidity profile ( $b_R$ ) and that with  $q_{LC}$  as humidity profile ( $b_{LC}$ ), and then use the vertical integral of their difference in the LT (2-4 km AGL),  $B_{LT} = \int_{2\text{ km AGL}}^{4\text{ km AGL}} (b_R - b_{LC}) dz$ , to quantify the additional effect of LT moisture variation on the parcel buoyancy that is not coupled with the PBL. We also calculate the integral of buoyancy based on  $q_{LC}$ , i.e.,  $B_{LC} = \int_{2\text{ km AGL}}^{4\text{ km AGL}} b_{LC} dz$ , to assess the land-coupled effect on convection. To make the results comparable, we apply a normal percentile transform (Wilks, 2011) to obtain standardized scores of  $B_{LT}$ , which we use for further analysis.

### 2.3 Identify dry/wet soil regime and coupled afternoon precipitation events

The CTP/ $HI_{Low}$  framework developed by Findell & Eltahir (2003) is commonly used to identify atmospheric preference of LAC state. CTP is calculated by integrating the difference between moist adiabat temperature and the ambient temperature profile from 100 to 300 hPa AGL. It is a measure of the energy available for convection and the 100–300 hPa AGL is a critical level for the development of the daytime boundary layer.  $HI_{Low}$ , on the other hand,

Formatted: Font: Italic, Not Highlight

Formatted: Subscript, Not Highlight

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: Italic, Not Highlight

Formatted: Font: Not Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Subscript

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

175 indicates the pre-existing moisture of the very lower atmosphere, and is defined as  $HI_{Low} = (T - T_d)_{150hPa AGL} + (T - T_d)_{50hPa AGL}$ . In this framework, wet soil advantage regime occurs when the atmospheric state is closer to the wet adiabatic rate, resulting in a low CTP and triggering convection through a strong increase in moist static energy from the land surface moisture large latent heat flux (small  $HI_{Low}$ ). Conversely, dry soil advantage regime occurs when the temperature profile is close to the dry adiabatic lapse rate with weak thermal stability (high CTP), and the soil provides less water vapor but more heat. This condition favors convection lifted by the boundary layer growth due to high sensible heat fluxes at the surface (Ek and Holtlag, 2004; Huang and Margulis, 2011; Gentine et al., 2013).  
180 We adopt a modified CTP/ $HI_{Low}$  framework proposed by Wakefield et al. (2019) using the standardized score of CTP/ $HI_{Low}$ . We first calculate CTP and  $HI_{Low}$  using sounding data at 11:30 LSTAs in Wakefield et al. (2019), and average FWI during 06:00-12:00 LST. Then dry-coupling cases are defined by as days with anomalously high CTP (higher than climatological CTP for our analysis period) over dry soils (FWI < 0.4), which is similar to the dry soil advantage regime in Findell & Eltahir (2003); ~~W~~wet-coupling cases, on the other hand, are characterized by anomalously low  $HI_{Low}$  over wet soil (FWI > 0.7), which corresponds to a moisture-abundant, energy-limited regime where the atmospheric profile is likely near moist adiabatic (Findell and Eltahir, 2003). The wet soil condition is expected to promote precipitation recycling through the addition of moist static energy via evapotranspiration and provides a continuous supply of low-level moisture.  
190 Since LAC would mostly affect the thermodynamically driven afternoon convection, we focus on the morning (0600-1300 LST), afternoon (1400-2000 LST), and evening (2100-2400 LST) precipitation events in our analysis. Afternoon precipitation events (APEs) are identified as daily samples that meet the following two criteria: 1) daily precipitation peaks during the afternoon hours defined above; and 2) the afternoon precipitation is at least twice as large as the morning precipitation, and also greater than the evening precipitation (filter out organized precipitation). The cases  
195 not categorized as APEs are referred to as non-APEs afterward. We obtain a total of 368 APEs from the 2172 sounding. We further select APEs associated with either dry-coupling or wet-coupling condition, resulting in 94 dry-coupling APEs and 79 wet-coupling APEs. These which account for 24.2% of the total 388 dry-coupling cases and 20.3 % of the total 389 wet-coupling cases, respectively. The comparable number of APEs for both dry- and wet-coupling conditions aligns with the finding of Findell and Eltahir (2003b) that the SGP is in the transitional region where negative and positive feedback days occurred with similar frequency. In addition, our analysis also shows that (Figure S1), within all 368 APEs, 16 instances exhibit a  $HI_{Low}$  lower than 5°C — a threshold established in Findell and Eltahir (2003a, 2003b). Among these, 8 are wet-coupling APEs and have significantly higher FWI compared to other groups. This suggests that the low  $HI_{Low}$  values observed before noon in these cases are likely influenced by soil moisture evaporation, rather than being purely controlled by atmospheric factors. Furthermore, one of these cases is categorized as dry-coupling APE, and seven as "other APEs", which are APEs not categorized as either dry-coupling or wet-coupling APEs. These cases likely represent "atmospherically controlled days", as per the CTP-  $HI_{Low}$  framework, and only account for a small fraction (~2.2%) of all APEs we identified.  
200  
205

Formatted: Subscript

### 3 Results

#### 3.1 Thermodynamic pre-conditions for APEs under the dry and wet-coupling regimes

210 To identify favorable atmospheric conditions for coupling APEs, we evaluate the differences in temperature ( $T$ ),  
specific humidity ( $q$ ), and relative humidity (RH) at 11:30 LST between averaged local coupling APEs and non-APE  
cases in the warm seasons (May to September), as shown in Figure 2. Regardless of soil moisture conditions and  
coupling regimes, APEs are always associated with a wetter lower troposphere (0-4 km) than non-APEs. For dry-  
coupling regimes, the increases of  $q$  and RH in the PBL and the LT associated with APEs are also stronger than those  
215 with the non-APEs, especially between 0.5 km and 3.5 km AGL, and the contrast between APEs and non-APEs in  
dry-coupling regime is larger than that for wet-coupling regimes. These humidity differences are expected, as more  
humid pre-conditions favor the occurrence of APEs. Notice that the greatest contrast between RH of APEs and that  
of non-APEs occurs between 1 km and 3.5 km AGL (above the PBL), which is the combined result of high  $q$  and  
decreasing  $T$  over this layer, highlighting the possible strong influence of LT humidity on deep convection.

220 In contrast, the sign of the temperature difference between APEs and non-APEs below 1.7 km AGL varies between  
the dry-coupling and wet-coupling regimes. For the dry-coupling regimes, the average temperature of APEs is lower  
than that of the non-APEs. This is presumably due to the stronger surface sensible flux and less stable atmosphere (a  
steep lapse rate or faster decrease of temperature with height) associated with APEs than the non-APEs under the dry-  
coupling regime. For the wet-coupling regime, the average temperature of the APEs is warmer than that of the non-  
225 APEs below 1.7 km AGL. This is consistent with a weaker lapse rate associated with a more humid environment,  
presumably due to vertical mixing of shallow convection, in the APEs than in the non-APEs cases. Above 1.7 km  
AGL, temperature of the APEs cases is lower than that of the non-APEs for both dry and wet-coupling regimes, as  
expected from less stable thermodynamic conditions in the APEs than in the non-APEs.

To investigate the difference in atmospheric conditions that favor APEs under the dry- versus wet-coupling regime,  
230 we compare the composite differential profiles of RH,  $q$ , and  $T$  between dry- and wet-coupling APEs, as shown in  
Figure 3. In general,  $T$  is higher for APEs under dry-coupling than under wet-coupling regime (Figure 3a), especially  
in the PBL (below 2 km). This can be attributed to stronger sensible heat flux and temperature mixing over a dry  
surface. Notably, there is a significant difference in LT specific humidity between dry- and wet-coupling regimes  
(Figure 3b), with  $q$  associated with APEs being slightly lower below 1 km AGL under dry-coupling than under wet-  
235 coupling regime, but higher above 1 km, especially between 2 km and 3 km AGL. This suggests that APEs require  
entrainment of higher LT moisture under dry-coupling than under wet-coupling regimes. Both Figures 2b and 3b  
suggest that higher LT specific humidity is needed for APEs under the dry-coupling than under the wet-coupling  
regimes. Moreover, RH associated with dry-coupling regimes is less than that of wet-coupling regimes in the PBL  
(Figure 3c), as expected from a drier PBL over a dry surface. However, such an RH difference becomes smaller and  
240 eventually disappears in the LT (2-4 km). This is because the lower RH in the dry-coupling regimes is mainly due to  
warmer  $T$  below 2 km AGL (Figure 3b), whereas above 2 km AGL, the higher  $q$  and slightly warmer  $T$  in the dry-  
coupling regimes balance each other out and lead to a similar RH as in the wet-coupling regimes.

Recent studies on the SM-P relationship have highlighted the greater impact of soil-moisture anomalies on boundary-  
layer stability and precipitation formation than on the ambient moisture (e.g., Seneviratne et al., 2010; Santanello et

245 al., 2018). To investigate how humidity affects the preconditioning of the convective environment and how it impacts instability in dry- and wet-coupling regimes, we use an entraining parcel model to calculate buoyancy profiles for  $b_R$  and  $b_{LC}$ , respectively, with parcels originating in the mixed layer. We then compute differences in the integral buoyancy between  $b_R$  and  $b_{LC}$  profiles for the 2-4 km AGL range to explore the influence of LT humidity-related convective thermodynamic instability.

250 To evaluate the atmospheric thermodynamic structure associated with the  $B_{LT}$ , we evaluate the composite average sounding profiles based on three tertile of  $B_{LT}$  in the warm season as shown in Figure 4. The  $B_{LT}$  values for the three tertiles range from -55.8 to -8.8 J/kg, -8.8 to 10.20J/kg, and 10.0 to 72.8 J/kg, respectively. It is noteworthy that the temperature and dew point are similar among these three tertiles of  $B_{LT}$  near the surface (below 900hPa) but clearly different from above 900 hPa up to at least 400 hPa AGL, which indicates the importance of LT humidity.

255 For the lower tertile (0% - 33%) of  $B_{LT}$  (Figure 4a), dew point values are substantially lower than those for the middle and upper tertile of the  $B_{LT}$  (Figures 4b and 4c). The sharp decrease of dew point values with height, near-constant temperature, and large gap between the temperature and dew point profiles at 700-900 hPa suggest strong dry shallow convection. For the middle tertile (Figure 4b), dewpoint and temperature decrease gradually with a height between 900 hPa and 700 hPa, and the gap between the dew point and temperature profiles is smaller than those for the lower tertile of the  $B_{LT}$  (Figure 4a). These features suggest a mixture of dry and moist shallow convection. For the upper tertile of  $B_{LT}$  (Figure 4c, 67% - 100%), dew point values are nearly constant from the surface to 700 hPa, and the humidity of free troposphere, as indicated by the gap between temperature and dew point profiles, is substantially wetter (implying higher RH) than for the middle and lower tertile of the  $B_{LT}$ . These features suggest that moist shallow convection dominates LT. Higher dew point values between 700 hPa and 500 hPa also suggest a more humid middle troposphere associated with the upper  $B_{LT}$  tertile (Figure 4c) than with the middle and lower  $B_{LT}$  tertile (Figure 4a and 4b). Thus, Figure 4 suggests that  $B_{LT}$  variations are strongly influenced by the humidity in the lower and middle troposphere.

### 3.2 The influence of LT humidity on afternoon precipitation under different LAC regimes

To investigate the effect of the LT humidity versus land surface air humidity on APEs, we evaluate the probability distribution of APEs as a function of  $B_{LT}$  and  $B_{LC}$  for dry- and wet-coupling APEs, respectively, in Figure 5.  $B_{LC}$  is usually more negative over dry soil (Figure 5b) than over wet soil (Figure 5a), and therefore usually more negative over dry PBL than over wet PBL. Figure 5b shows that dry-coupling APEs occur more frequently with negative  $B_{LC}$  (69%) and positive  $B_{LT}$  (77%), indicating that the influence of a more humid LT can override the influence of surface air aridity. For land surface, favoring local precipitation ( $B_{LC} > 0$ ), dry-coupling APEs occur more frequently with positive  $B_{LT}$  (26%) than with negative  $B_{LT}$  (5%). For the wet-coupling conditions, 57% of APEs occur with positive  $B_{LT}$ , while 31% occur with humid PBL pre-condition (Figure 5a). Overall, for dry-coupling cases, more APEs are associated with humid LT (positive  $B_{LT}$ ) than with humid surface air (positive  $B_{LC}$ ), but this trend is not evident for wet-coupling cases.

To evaluate the influence of LT humidity on LAC under the CTP/HI<sub>Low</sub> framework, we compare the joint distribution of all APEs, wet-coupling APEs, and dry-coupling APEs, respectively, as a function of  $B_{LT}$  scores and HI<sub>Low</sub> scores



(Figure 6S1) using a normal percentile transform (Wilks 2011). Larger than normal  $B_{LT}$  (humid LT) values were associated with 61% of all APEs, regardless of the PBL humidity. Smaller than normal  $H_{Low}$  (humid PBL) occurred in 63% of all APEs, regardless of the LT humidity. About 40% of all APEs occur under both humid PBL and humid LT. Thus, the probabilities for APEs to occur under either humid PBL or humid LT are similar, with a preference for APEs to occur under both humid PBL and humid LT conditions. For dry-coupling condition, 76% of the APEs occur under humid LT versus 59% under humid PBL. Therefore, APEs appear to prefer humid LT more than humid PBL under dry-coupling conditions, but this preference is not found in wet-coupling conditions. This result is consistent with the findings in Figure 5.

To further investigate how LT humidity or  $B_{LT}$  can affect the probability of APEs under the dry-coupling and wet-coupling regimes, respectively, we present three statistical measures of APEs as a function of  $B_{LT}$  in Figure 76. Figure 76a shows that, for the dry-coupling cases, the fractional occurrence of APEs (defined as the proportion of APEs relative to all dry-coupling cases) in each  $B_{LT}$  bin increases with  $B_{LT}$  up to its 70<sup>th</sup> percentile, with a significant correlation ( $R = 0.65$ ,  $p < 0.05$ ). For the wet-coupling cases, the fractional occurrence of the APEs peaks when  $B_{LT}$  is between the 30<sup>th</sup> and 70<sup>th</sup> percentile. Thus, APEs appear to prefer higher LT humidity under dry coupling than under wet coupling.

Next, we explore how  $B_{LT}$  affects the partition of dry-coupling versus wet-coupling APEs. Figure 76b shows that the proportion of the dry-coupling APEs relative to all APEs increases with  $B_{LT}$ , with a strong correlation ( $R = 0.89$ ,  $p < 0.05$ ). The proportion ranges from 0.04 at the bottom 10% to 0.47 at the top 10% of  $B_{LT}$ . However, the proportion of wet-coupling APEs per  $B_{LT}$  bin peaks at lower to medium  $B_{LT}$  percentiles (30%-50%) and decreases almost monotonically with increasing  $B_{LT}$  from 50% to 100%.

We also investigate how  $B_{LT}$  affects rain rates associated with dry-coupling and wet-coupling APEs, respectively. Figure 76c shows a clear increase in rain rate with  $B_{LT}$  for the dry-coupling APEs, except for the 90-100<sup>th</sup> percentile of  $B_{LT}$ , where there are few APEs samples. In contrast, we find no clear dependence of rain rate on  $B_{LT}$  for the wet-coupling APEs. Thus, our findings suggest that a high  $B_{LT}$  tends to increase the frequency and intensity of the dry-coupling APEs, as well as the relative frequency of dry-coupling APEs compared to wet-coupling APEs.

In addition, we evaluate the variations of deep- (cloud top height (CTH)  $> 8$  km), shallow- (CTH  $< 3$  km), and convective congestus (CTH between 3 km and 8 km) associated with APEs based on hourly precipitation and cloud fraction as in following Zhuang et al. (2017, Figure S2). In general, APEs associated with all three convective types increase with  $B_{LT}$  under dry-coupling conditions (Figure S2). Under wet-coupling condition, APEs associated with deep convection does not exhibit a clear dependence on  $B_{LT}$ . However, APEs associated with shallow convection decreases with increasing  $B_{LT}$ , while those associated with congestus increase with increasing  $B_{LT}$ . These results imply that the increase in  $B_{LT}$  can lead to a deepening of shallow convection into congestus due to reduced buoyancy dilution caused by entraining wetter LT air for wet-coupling convection.

#### 4 Conclusions

315 Land-atmosphere interactions occur when local land surface and subsurface conditions influence the moisture and energy budgets of the overlying atmosphere. The relative impacts of soil moisture on convective precipitation can vary depending on the atmospheric conditions. In this study, we compared the difference in RH,  $q$ , and  $T$  profiles between APEs and non-APEs under both dry- and wet-coupling conditions.

Our initial analysis revealed that APEs had an overall wetter PBL and LT (0-4km AGL) than non-APEs, especially 320 under dry-coupling regimes. The RH difference between APEs and non-APEs in the LT was driven by differences in both  $q$  and  $T$ , with dry-coupling APEs exhibiting lower humidity in the PBL than wet-coupling APEs. However, as the altitude increases, the difference in RH between dry- and wet-coupling APEs decreases due to the increasing difference in  $q$  and decreasing difference in  $T$ . Above 4 km AGL, the difference in  $q$  becomes zero. Therefore, we could infer the importance of LT humidity in the SM-P relationship, and the APEs under dry-coupling conditions 325 necessitate more LT humidity than that under wet-coupling conditions.

To further investigate the influence of LT humidity on the SM-P relationship, we employ an entraining parcel model and a new metric  $B_{LT}$ , which measures the 2-4 km vertical integral of the difference between buoyancy calculated from the observed humidity profile and that correlated to (regressed against) the average specific humidity in the PBL. Statistical analysis reveals that the wetter LT and normal PBL were associated with larger  $B_{LT}$  values, whereas drier 330 LT was linked to smaller  $B_{LT}$  values. Moreover, there is a higher likelihood of APEs occurring with positive  $B_{LT}$  percentile under dry-coupling conditions, while this relationship is not apparent for the probability distribution of  $B_{LT}$  percentile for wet-coupling APEs. Additionally, as the  $B_{LT}$  percentile increases, the frequency of dry-coupling APEs also increases, whereas the opposite tendency was observed for wet-coupling APEs. In the meantime, the ratio of dry-coupling APEs to all APEs increases with the  $B_{LT}$  percentile, while this tendency is the opposite for the ratio of wet-coupling APEs to all APEs. Regarding precipitation, the average rain rate tends to rise with increasing  $B_{LT}$  percentile 335 under dry-coupling conditions, but this trend is not significant for wet-coupling APEs. Overall, our results indicate that the impact of LT humidity differs between dry- and wet-coupling APEs, with dry-coupling APEs being more influenced by LT humidity compared to wet-coupling APEs.

The Great Plain Low-Level Jet (GPLLJ) is widely acknowledged as a primary mechanism responsible for the regional-scale water vapor transport from the Gulf of Mexico during May-September. The GPLLJ creates a thermodynamic environment that facilitates convection and precipitation, making it a key factor in initiating and sustaining mesoscale weather phenomena (e.g., Higgins et al., 1997; Hodges & Pu, 2019; Mo et al., 1997; Pu et al., 2016; Pu & Dickinson, 2014; Weaver & Nigam, 2008). It is well established that the GPLLJ can enhance the occurrence of nocturnal convective precipitation in the SGP (Pu and Dickinson, 2014). However, our findings imply that the moisture carried 345 by the GPLLJ could also play an important role in generating local diurnal afternoon precipitation when it reaches the region during the daytime, particularly over dry soil conditions. This result is consistent with Ford et al. (2015) that suggested-soil moisture feedback to precipitation could potentially manifest itself over wetter- and drier-than-normal soils, depending on the overall synoptic and dynamic conditions, and precipitation favors dry soil when the low-level jet is present. Therefore, these results collectively suggest that the GPLLJ plays a significant role in alleviating drought 350 conditions in the SGP by influencing both diurnal and nocturnal precipitation.

Our study not only presents new insights into the role of LT humidity on the SM-P relationship but also serves as a quantitative elucidation of the negative feedback behavior discussed in Findell and Eltahir (2003b) using the CTP-HI<sub>low</sub> framework. Specifically, they highlighted the topographic and dynamical circumstances that commonly result in a moist air layer originating from the elevated Mexican plateau, typically with its base around 850 hPa. This moist layer, while not captured by the HI<sub>low</sub> metric, can be detected by our approach. This work reveals the varying impact of LT humidity on the SM-P relationship, highlighting the importance of incorporating land-atmosphere interaction into model parameterizations. However, there are still limitations in our work. One key concern is the potential uncertainties introduced by constructing the land-coupled LT humidity profile via linear regression. Such uncertainties arise mainly from the linear model's inherent assumptions, including the constancy of relationships under varying conditions and the potential oversight of non-linearity. A more thorough investigation into the model's residuals and additional sensitivity analyses could provide deeper insights into these uncertainties. Furthermore, our categorization of APEs may not be always associated with convective precipitation, given that it relies solely on the region-average precipitation data. Improving the classification of APEs, possibly by integrating convection classification results from radar observations, could lead to more precise interpretations. This study only focuses on a single location, i.e., SGP, thus expanding research to include a variety of climatic zones would be crucial in assessing the broader applicability of our methods and conclusions. Our future work will also involve investigating the primary source of LT humidity and employing both  $B_{LT}$  and CTP/HI<sub>low</sub> as atmospheric indicators to identify the global regions exhibiting various with diverse LT humidity-SM-P relationships, thereby advancing our knowledge-understanding of LAC on a broader scale.

### 370 Competing Interests

The contact author has declared that none of the authors has any competing interests.

### Acknowledgment

G.W. was funded by the China Scholarship Council (CSC; 201806010052). Y.Z. and R.F. were supported by the National Oceanic and Atmospheric Administration-Climate Program Office (NOAA-CPO) Modelling, Analysis, Predictions, and Projections (MAPP) Program (NA20OAR4310426), and the National Science Foundation (NSF) Physical and Dynamic Meteorology (PDM) Program (AGS-2214697). P.A.D. was supported by the National Aeronautics and Space Administration (NASA) (80NSSC21K1801). J.S. was supported by funding from NASA HQ and PM Jared Entin.

### Reference

380 Bretherton, C. S., Peters, M. E., and Back, L. E.: Relationships between water vapor path and precipitation over the tropical oceans, *J Clim*, [https://doi.org/10.1175/1520-0442\(2004\)017<1517:RBWVPA>2.0.CO;2](https://doi.org/10.1175/1520-0442(2004)017<1517:RBWVPA>2.0.CO;2), 2004.

[Dirmeyer, P. A., Koster, R. D., and Guo, Z. Guo: Do Global Models Properly Represent the Feedback between Land and Atmosphere?. J Hydrometeorol, 7, 1177–1198, <https://doi.org/10.1175/JHM532.1>, 2006.](#)

385 Ek, M. B. and Holtslag, A. A. M.: Influence of soil moisture on boundary layer cloud development, J Hydrometeorol, 5, 86–99, [https://doi.org/10.1175/1525-7541\(2004\)005<0086:IOSMOB>2.0.CO;2](https://doi.org/10.1175/1525-7541(2004)005<0086:IOSMOB>2.0.CO;2), 2004.

Ferguson, C. R. and Wood, E. F.: An evaluation of satellite remote sensing data products for land surface hydrology: Atmospheric infrared sounder, J Hydrometeorol, 11, 1234–1262, <https://doi.org/10.1175/2010JHM1217.1>, 2010.

Fernando, D. N., Mo, K. C., Fu, R., Pu, B., Bowerman, A., Scanlon, B. R., Solis, R. S., Yin, L., Mace, R. E., Mioduszewski, J. R., Ren, T., and Zhang, K.: What caused the spring intensification and winter demise of the 2011 drought over Texas?, Clim Dyn, 47, 3077–3090, <https://doi.org/10.1007/s00382-016-3014-x>, 2016.

390 Findell, K. L. and Eltahir, E. A. B.: Atmospheric controls on soil moisture-boundary layer interactions. Part I: Framework development, J Hydrometeorol, 4, 552–569, [https://doi.org/10.1175/1525-7541\(2003\)004<0552:ACOSML>2.0.CO;2](https://doi.org/10.1175/1525-7541(2003)004<0552:ACOSML>2.0.CO;2), 2003a.

[Findell, K. L. and Eltahir, E. A. B.: Atmospheric Controls on Soil Moisture-Boundary Layer Interactions. Part II: Feedbacks within the Continental United States, 570–583 pp., 2003b.](#)

395 Ford, T. W., Rapp, A. D., and Quiring, S. M.: Does afternoon precipitation occur preferentially over dry or wet soils in Oklahoma?, J Hydrometeorol, <https://doi.org/10.1175/JHM-D-14-0005.1>, 2015.

Fulton, R. A., Breidenbach, J. P., Seo, D. J., Miller, D. A., and O'Bannon, T.: The WSR-88D rainfall algorithm, Weather Forecast, [https://doi.org/10.1175/1520-0434\(1998\)013<0377:TWRA>2.0.CO;2](https://doi.org/10.1175/1520-0434(1998)013<0377:TWRA>2.0.CO;2), 1998.

400 Gentine, P., Holtslag, A. A. M., D'Andrea, F., and Ek, M.: Surface and atmospheric controls on the onset of moist convection over land, J Hydrometeorol, 14, 1443–1462, <https://doi.org/10.1175/JHM-D-12-0137.1>, 2013.

Guo, Z., Dirmeyer, P. A., Koster, R. D., Bonan, G., Chan, E., Cox, P., Gordon, C. T., Kanoe, S., Kowalczyk, E., Lawrence, D., Liu, P., Lu, C. H., Malyshev, S., McAvaney, B., McGregor, J. L., Mitchell, K., Mocko, D., Oki, T., Oleson, K. W., Pitman, A., Sud, Y. C., Taylor, C. M., Verseghy, D., Vasic, R., Xue, Y., and Yamada, T.: GLACE: The Global Land-Atmosphere Coupling Experiment. Part II: Analysis, J Hydrometeorol, 7, 611–625, <https://doi.org/10.1175/JHM511.1>, 2006.

405 Higgins, R. W., Yao, Y., Yarosh, E. S., Janowiak, J. E., and Mo, K. C.: Influence of the Great Plains Low-Level Jet on Summertime Precipitation and Moisture Transport over the Central United States, Journal of Climate, American Meteorological Society, 481–507 pp., [https://doi.org/10.1175/1520-0442\(1997\)010<0481:IOTGPL>2.0.CO;2](https://doi.org/10.1175/1520-0442(1997)010<0481:IOTGPL>2.0.CO;2), 1997.

410 Hodges, D. and Pu, Z.: Characteristics and variations of low-level jets in the contrasting warm season precipitation extremes of 2006 and 2007 over the Southern Great Plains, Theor Appl Climatol, 136, 753–771, <https://doi.org/10.1007/S00704-018-2492-7/FIGURES/19>, 2019.

Holloway, C. E. and Neelin, D. J.: Moisture vertical structure, column water vapor, and tropical deep convection, J Atmos Sci, 66, 1665–1683, <https://doi.org/10.1175/2008JAS2806.1>, 2009.

415 Huang, H. Y. and Margulis, S. A.: Investigating the impact of soil moisture and atmospheric stability on cloud development and distribution using a coupled large-eddy simulation and land surface model, J Hydrometeorol, <https://doi.org/10.1175/2011JHM1315.1>, 2011.

- Illston, B. G., Basara, J. B., Fisher, D. K., Elliott, R., Fiebrich, C. A., Crawford, K. C., Humes, K., and Hunt, E.: Mesoscale monitoring of soil moisture across a statewide network, *J Atmos Ocean Technol*, <https://doi.org/10.1175/2007JTECHA993.1>, 2008.
- 420 Konings, A. G., Katul, G. G., and Porporato, A.: The rainfall-no rainfall transition in a coupled land-convective atmosphere system, *Geophys Res Lett*, <https://doi.org/10.1029/2010GL043967>, 2010.
- [Koster, R. D., Dirmeyer, P. A., Guo, Z., Bonan, G., Chan, E., Cox, P., Gordon, C. T., Kanae, S., Kowalczyk, E., Lawrence, D. and Liu, P.: Regions of strong coupling between soil moisture and precipitation, \*Science\*, 305, 1138-1140, <https://doi.org/10.1126/science.1100217>, 2004.](https://doi.org/10.1126/science.1100217)
- 425 Koster, R. D., Guo, Z., Dirmeyer, P. A., Bonan, G., Chan, E., Cox, P., Davies, H., Gordon, C. T., Kanae, S., Kowalczyk, E., Lawrence, D., Liu, P., Lu, C. H., Malyshev, S., McAvaney, B., Mitchell, K., Mocko, D., Oki, T., Oleson, K. W., Pitman, A., Sud, Y. C., Taylor, C. M., Verseghy, D., Vasic, R., Xue, Y., and Yamada, T.: GLACE: The Global Land-Atmosphere Coupling Experiment. Part I: Overview, <https://doi.org/10.1175/JHM510.1>, 2006.
- 430 Liu, S. and Liang, X. Z.: Observed diurnal cycle climatology of planetary boundary layer height, *J Clim*, <https://doi.org/10.1175/2010JCLI3552.1>, 2010.
- Mapes, B., Tulich, S., Lin, J., and Zuidema, P.: The mesoscale convection life cycle: Building block or prototype for large-scale tropical waves?, *Dynamics of Atmospheres and Oceans*, <https://doi.org/10.1016/j.dynatmoce.2006.03.003>, 2006.
- 435 Mo, K. C., Paegle, J. N., and Higgins, R. W.: Atmospheric processes associated with summer floods and droughts in the central United States, *J Clim*, [https://doi.org/10.1175/1520-0442\(1997\)010<3028:APAWSF>2.0.CO;2](https://doi.org/10.1175/1520-0442(1997)010<3028:APAWSF>2.0.CO;2), 1997.
- Myoung, B. and Nielsen-Gammon, J. W.: The convective instability pathway to warm season drought in Texas. Part I: The role of convective inhibition and its modulation by soil moisture, *J Clim*, <https://doi.org/10.1175/2010JCLI2946.1>, 2010.
- 440 Pu, B. and Dickinson, R. E.: Diurnal Spatial Variability of Great Plains Summer Precipitation Related to the Dynamics of the Low-Level Jet, *J Atmos Sci*, 71, 1807–1817, <https://doi.org/10.1175/JAS-D-13-0243.1>, 2014.
- Pu, B., Dickinson, R. E., and Fu, R.: Dynamical connection between Great Plains low-level winds and variability of central Gulf States precipitation, *Journal of Geophysical Research: Atmospheres*, 121, 3421–3434, <https://doi.org/10.1002/2015JD024045>, 2016.
- 445 Raz-Yaseef, N., Billesbach, D. P., Fischer, M. L., Biraud, S. C., Gunter, S. A., Bradford, J. A., and Torn, M. S.: Vulnerability of crops and native grasses to summer drying in the U.S. Southern Great Plains, *Agric Ecosyst Environ*, <https://doi.org/10.1016/j.agee.2015.07.021>, 2015.
- Roundy, J. K. and Santanello, J. A.: Utility of Satellite Remote Sensing for Land–Atmosphere Coupling and Drought Metrics, *J Hydrometeorol*, 18, 863–877, <https://doi.org/10.1175/jhm-d-16-0171.1>, 2017.
- 450 Roundy, J. K., Ferguson, C. R., and Wood, E. F.: Temporal Variability of Land-Atmosphere Coupling and Its Implications for Drought over the Southeast United States, *J Hydrometeorol*, 14, 622–635, <https://doi.org/10.1175/JHM-D-12-090.1>, 2013.

- Santanello, J. A., And, J. R., Peters-Lidard, C. D., Kumar, S. v, Alonge, C., and Tao, W.-K.: A Modeling and Observational Framework for Diagnosing Local Land-Atmosphere Coupling on Diurnal Time Scales, *J. Hydrometeorol*, 10, 577–599, <https://doi.org/10.1175/2009JHM1066.1>, 2009.
- 455 Santanello, J. A., Dirmeyer, P. A., Ferguson, C. R., Findell, K. L., Tawfik, A. B., Berg, A., Ek, M., Gentine, P., Guillod, B. P., van Heerwaarden, C., and Roundy, J.: LAND-ATMOSPHERE INTERACTIONS The LoCo Perspective, <https://doi.org/10.1175/BAMS-D-17-0001.1>, 2017.
- Santanello, J. A., Dirmeyer, P. A., Ferguson, C. R., Findell, K. L., Tawfik, A. B., Berg, A., Ek, M., Gentine, P., Guillod, B. P., Van Heerwaarden, C., Roundy, J., and Wulfmeyer, V.: Land-Atmosphere Interactions: The LoCo Perspective, *Bull Am Meteorol Soc*, 99, 1253–1272, <https://doi.org/10.1175/BAMS-D-17-0001.1>, 2018.
- 460 Schiro, K. A., Neelin, J. D., Adams, D. K., and Lintner, B. R.: Deep convection and column water vapor over tropical land versus tropical ocean: A comparison between the amazon and the tropical Western Pacific, *J Atmos Sci*, 73, 4043–4063, <https://doi.org/10.1175/JAS-D-16-0119.1>, 2016.
- 465 Schiro, K. A., Ahmed, F., Giangrande, S. E., and David Neelin, J.: GoAmazon2014/5 campaign points to deep-inflow approach to deep convection across scales, 115, 4577–4582, <https://doi.org/10.1073/pnas.1719842115>, 2018.
- Schneider, J. M., Flsher, D. K., Elliott, R. L., Brown, G. O., and Bahmann, C. P.: Spatiotemporal variations in soil water: First results from the ARM SGP CART network, *J Hydrometeorol*, [https://doi.org/10.1175/1525-7541\(2003\)004<0106:SVISWF>2.0.CO;2](https://doi.org/10.1175/1525-7541(2003)004<0106:SVISWF>2.0.CO;2), 2003.
- 470 Seneviratne, S. I., Corti, T., Davin, E. L., Hirschi, M., Jaeger, E. B., Lehner, I., Orlowsky, B., and Teuling, A. J.: Investigating soil moisture–climate interactions in a changing climate: A review, *Earth Sci Rev*, 99, 125–161, <https://doi.org/10.1016/j.earscirev.2010.02.004>, 2010.
- Siebesma, A. P., Soares, P. M. M., and Teixeira, J.: A Combined Eddy-Diffusivity Mass-Flux Approach for the Convective Boundary Layer, <https://doi.org/10.1175/JAS3888.1>, 2007.
- 475 Song, H. J., Ferguson, C. R., and Roundy, J. K.: Land-atmosphere coupling at the southern great plains atmospheric radiation measurement (ARM) field site and its role in anomalous afternoon peak precipitation, *J Hydrometeorol*, 17, 541–556, <https://doi.org/10.1175/JHM-D-15-0045.1>, 2016.
- Tawfik, A. B., Dirmeyer, P. A., and Santanello, J. A.: The heated condensation framework. Part I: Description and southern great plains case study, *J Hydrometeorol*, 16, 1929–1945, <https://doi.org/10.1175/JHM-D-14-0117.1>, 2015a.
- 480 Tawfik, A. B., Dirmeyer, P. A., and Santanello, J. A.: The heated condensation framework. Part II: Climatological behavior of convective initiation and land-atmosphere coupling over the conterminous United States, *J Hydrometeorol*, 16, 1946–1961, <https://doi.org/10.1175/JHM-D-14-0118.1>, 2015b.
- Taylor, C. M.: Detecting soil moisture impacts on convective initiation in Europe, *Geophys Res Lett*, 42, 4631–4638, <https://doi.org/10.1002/2015GL064030>, 2015.
- 485 Tuttle, S. and Salvucci, G.: Atmospheric science: Empirical evidence of contrasting soil moisture-precipitation feedbacks across the United States, *Science* (1979), 352, 825–828, <https://doi.org/10.1126/science.aaa7185>, 2016.
- Wakefield, R. A., Basara, J. B., Furtado, J. C., Illston, B. G., Ferguson, C. R., and Klein, P. M.: A modified framework for quantifying land–atmosphere covariability during hydrometeorological and soil wetness extremes in oklahoma, *J Appl Meteorol Climatol*, 58, 1465–1483, <https://doi.org/10.1175/JAMC-D-18-0230.1>, 2019.

490 Weaver, S. J. and Nigam, S.: Variability of the Great Plains Low-Level Jet: Large-Scale Circulation Context and Hydroclimate Impacts, *J Clim*, 21, 1532–1551, <https://doi.org/10.1175/2007JCLI1586.1>, 2008.

Wilks, D. S.: *Statistical methods in the atmospheric sciences*, 676, 2011.

Yin, J., Albertson, J. D., Rigby, J. R., and Porporato, A.: Land and atmospheric controls on initiation and intensity of moist convection: CAPE dynamics and LCL crossings, *Water Resour Res*, 51, 8476–8493, <https://doi.org/10.1002/2015WR017286>, 2015.

495 Zhang, Y. and Klein, S. A.: Mechanisms Affecting the Transition from Shallow to Deep Convection over Land: Inferences from Observations of the Diurnal Cycle Collected at the ARM Southern Great Plains Site, *J Atmos Sci*, 67, 2943–2959, <https://doi.org/10.1175/2010jas3366.1>, 2010.

Zhuang, Y., Fu, R., Marengo, J. A., and Wang, H.: Seasonal variation of shallow-to-deep convection transition and its link to the environmental conditions over the Central Amazon, *J Geophys Res*, 122, 2649–2666, <https://doi.org/10.1002/2016JD025993>, 2017.

500 Zhuang, Y., Fu, R., and Wang, H.: How do environmental conditions influence vertical buoyancy structure and shallow-to-deep convection transition across different climate regimes?, *J Atmos Sci*, 75, 1909–1932, <https://doi.org/10.1175/JAS-D-17-0284.1>, 2018.

505




Figure 1: Correlation coefficient profiles between specific humidity ( $q$ ) at each vertical level from 0 to 4 km AGL and mean  $q$  in the mixed layer (0-1km AGL). The correlation coefficients are calculated for the warm season of each year. The black line indicates the average value of 18 years, and the green shade shows the standard error. All correlation coefficients at 0-4 km for all 18 years are significant at 0.05 level.

510





Figure 2: Composite difference of a) temperature ( $T_{diff}$ ), b) specific humidity ( $q_{diff}$ ), and c) relative humidity ( $RH_{diff}$ ) profiles between APEs and non-APEs for dry- (red lines) and wet- (black lines) coupling cases. The thicker portions of the lines indicate where the differences are statistically significant at 0.05 level.

515








Figure 3: Composite difference of a) temperature ( $T_{diff}$ ), b) specific humidity( $q_{diff}$ ), and c) relative humidity ( $RH_{diff}$ ) between dry- and wet-coupling APEs (Dry minus Wet). The thicker portions of the lines indicate where the differences are statistically significant at 0.05 level.

520






Figure 4: Composite temperature ( $T$ ; solid line) and dewpoint temperature ( $T_d$ ; dash line) profiles at the ARM SGP site for all days during May–September from 2001 to 2018, above the 950hPa level, based on  $B_{LT}$  tercile: 0-33% (lower  $B_{LT}$ , black); 33%-67% (medium  $B_{LT}$ , green); 67%-100% bins (higher  $B_{LT}$ , red).

525




530

Figure 5: Joint frequency distributions of a) wet-coupling APEs and b) dry-coupling APEs as a function of  $B_{LT}$  (representing contribution of LT humidity to convective buoyancy) and  $B_{LC}$  (representing contribution of surface humidity) with white shades representing no APEs and darkest shades representing more than 5 APEs occurs of each  $B_{LC} - B_{LT}$  bin. The number indicates the fraction in each quadrant.



535 **Figure 6:** Joint distribution of (a) all APEs, (b) wet-coupling APEs, and (c) dry-coupling APEs as functions of  $B_{LT}$  score and  $H_{Low}$  score. The number indicates the fraction in each quadrant.

Formatted: Normal



540 **Figure 7:** (a) The fraction of wet- (dry-) coupling APEs over wet- (dry-) coupling cases ( $\frac{N_{wet-coupling-APEs}}{N_{wet-coupling-cases}}$  or  $\frac{N_{dry-coupling-APEs}}{N_{dry-coupling-cases}}$ ) in each  $B_{LT}$  bin for the dry- and wet-coupling cases, respectively. (b) Same as (a), but for the percentage of wet- or dry-coupling APEs relative to all APEs ( $\frac{N_{wet-coupling-APEs}}{N_{APEs}}$  and  $\frac{N_{dry-coupling-APEs}}{N_{APEs}}$ ). (c) Same as (a), but for the mean afternoon precipitation rate (PR). The correlation coefficients between  $B_{LT}$  percentiles and the fraction/PR are listed for the dry- and wet-coupling cases, respectively; correlation coefficients between  $B_{LT}$  percentiles and y-axis value significant at 0.05 level are marked with two asterisks.