



On the time scale of meteorological, soil moisture, and snow drought indices to assess streamflow drought over catchments with different hydrological regime: a case study using a hundred Chilean catchments

Oscar M. Baez-Villanueva^{1,2}, Mauricio Zambrano-Bigiarini^{3,4}, Diego G. Miralles¹, Hylke E. Beck⁵, Jonatan F. Siegmund⁶, Camila Alvarez-Garretón⁴, Koen Verbist⁷, René Garreaud^{8,4}, Juan Pablo Boisier^{8,4}, and Mauricio Galleguillos^{9,4}

¹Hydro-Climate Extremes Lab (H-CEL), Ghent University, Ghent, Belgium

²Institute for Technology and Resources Management in the Tropics and Subtropics (ITT), TH Köln, Cologne, Germany

³Department of Civil Engineering, Universidad de la Frontera, Temuco, Chile

⁴Center for Climate and Resilience Research, Universidad de Chile, Santiago, Chile

⁵Climate and Livability Initiative, Physical Sciences and Engineering, King Abdullah University of Science and Technology, Thuwal, Saudi Arabia

⁶Ernst Young GmbH, Wirtschaftsprüfungsgesellschaft, Stuttgart, Germany

⁷UNESCO International Hydrological Programme, Paris, France

⁸Department of Geophysics, Universidad de Chile, Santiago, Chile

⁹Faculty of Engineering and Sciences, Universidad Adolfo Ibáñez, Santiago, Chile

Correspondence: Mauricio Zambrano-Bigiarini (mauricio.zambrano@ufrontera.cl)

Abstract. A wide variety of drought indices exist today without consensus on suitable indices and temporal scales for monitoring streamflow drought across diverse hydrological settings. Considering the growing interest in spatially-distributed indices for ungauged areas, this study addresses the following questions: *i*) what temporal scales of precipitation-based indices are most adequate to assess streamflow drought in catchments with different hydrological regimes?, *ii*) do soil moisture indices outperform meteorological indices as proxies for streamflow drought?, *iii*) are snow indices more effective than meteorological indices for assessing streamflow drought in snow-influenced catchments? To answer these questions, we used one hundred near-natural catchments with four main types of hydrological regimes. The Standardised Precipitation Index (SPI), Standardised Precipitation and Evapotranspiration Index (SPEI), Empirical Standardised Soil Moisture Index (ESSMI), and standardised Snow Water Equivalent Index (SWEI) were computed across various time scales over the catchments for 1979–2020. Cross-correlation and event coincidence analysis were applied between these indices and the Standardised Streamflow Index at a temporal scale of one month (SSI-1), as representative of streamflow drought events. Finally, the linear correlation values and precursor coincidence rates were analysed for all catchments simultaneously, and separated by the hydrological regime. Our results indicate that *i*) there is no single meteorological, soil moisture, or snow drought index and temporal scale that could be used to characterise all streamflow droughts across Chile, and *ii*) the greater the snow influence in a catchment, the larger the temporal scale of the drought index to be used as proxy of streamflow drought. Finally, to avoid considering the influence of non-drought periods when analysing time series of drought indices, we suggest that future studies use the event



coincidence analysis to evaluate which meteorological, soil moisture, and/or snow drought indices can be used as proxies of streamflow drought events.

20 1 Introduction

A wide variety of drought indices coexists today (e.g., Svoboda et al., 2002; Hayes et al., 2012; Cammalleri et al., 2021) without consensus on which are more appropriate for monitoring streamflow drought, and over which temporal scales (Vicente-Serrano and López-Moreno, 2005; Lorenzo-Lacruz et al., 2013; Blauhut et al., 2015; Jain et al., 2015; Bachmair et al., 2016; Barker et al., 2016b; Zhao et al., 2017; Zhu et al., 2016; Slette et al., 2019; Hoffmann et al., 2020; Vorobeuskii et al., 2022). Primarily because just a few of them have truly been tested against drought impact data (Blauhut et al., 2015). Drought indices are often used alone or in combination to operationally monitor the onset, duration, and severity of drought events. Examples include the U.S. Drought Monitor (Svoboda et al., 2002; Leeper et al., 2022), the European Drought Observatory (Vogt, 2011; Cammalleri et al., 2021), the German Drought Monitor (Zink et al., 2016), the South Asia Drought Monitor (Saha et al., 2021), the Australian Drought Monitor (Cobon et al., n.d.), and the Global Drought Monitor Portal (Heim Jr. and Brewer, 2012; Hao et al., 2017). The low agreement among these indices, as well as the limited knowledge about their critical thresholds related to tangible impacts on society, economy, and ecosystems (Steinemann, 2003; Lac; Bachmair et al., 2015; Stagge et al., 2015a; Bachmair et al., 2016; Torelló-Sentelles and Franzke, 2022), hinders their effective use for decision-making purposes, drought mitigation, and adaptation (Steinemann and Cavalcanti, 2006; Steinemann, 2014; Steinemann et al., 2015). Before selecting a drought index, proper case-specific understanding is essential. This understanding enables effective monitoring, management, and planning, and can lead to developing mitigation strategies that ensure the resilience of our environment and communities (Bachmair et al., 2016).

Drought events are frequently classified as meteorological, soil moisture, and hydrological (Haile et al., 2020), each with different characteristics and impacts, but also with different indices used to monitor their development. Well-known meteorological drought indices include percent of normal (Zargar et al., 2011), deciles (Gibbs and Maher, 1967), Z-score (ZSI; Palmer, 1965a; Zargar et al., 2011), Rainfall Anomaly Index (RAI; van Rooy, 1965; Keyantash and Dracup, 2002), Palmer Drought Severity Index (PDSI; Palmer, 1965b), Standardised Precipitation Index (SPI; McKee et al., 1993), Effective Drought Index (EDI; Byun and Wilhite, 1999), and Standardised Precipitation Evapotranspiration Index (SPEI; Vicente-Serrano et al., 2010).

Although meteorological and soil moisture drought indices are relatively easy to calculate due to either their reliance on frequently-measured precipitation (P) or their reduced number of input variables, most of the socio-ecological impacts of droughts are related to surface water and groundwater deficits (altogether referred to as hydrological droughts), i.e., water shortages in rivers, lakes, and groundwater (Van Loon, 2015). Likewise, numerous indices have been used to monitor soil moisture drought, including the Palmer Moisture Anomaly Index (Z-index; Karl, 1986; Keyantash and Dracup, 2002), the



Soil Moisture Deficit Index (SMDI; Narasimhan and Srinivasan, 2005), and the Empirical Standardised Soil Moisture Index (ESSMI; Carrão et al., 2016). Drought indices describing hydrological droughts use a variety of variables related to surface and groundwater amounts. Examples are: the Standardised Streamflow Index (SSI; Vicente Serrano et al., 2012) and the Standardised Runoff Index (SRI; Shukla and Wood, 2008)) both based on streamflow (Q); the Surface Water Supply Index (SWSI; Shafer and Dezman, 1982) based on snow water equivalent (SWE) as representative of the snowpack, Q , reservoir storage, and P ; the Standardised Groundwater level Index (SGI; Bloomfield and Marchant, 2013; Van Loon et al., 2017) based on groundwater levels or terrestrial water storage; the Standardised Snow Melt and Rain Index (SMRI; Staudinger et al., 2014) based on snow storage and P ; and the Standardised Snow Water Equivalent Index (SWEI; Huning and AghaKouchak, 2020) based on SWE . As these variables are frequently less easily accessible, streamflow drought indices are often used to characterise hydrological droughts (Barker et al., 2016a; Wu et al., 2017a; Peña-Gallardo et al., 2019; Bhardwaj et al., 2020).

The propagation of meteorological drought into streamflow drought is a complex process modulated by the land component of the hydrological cycle. This propagation may lead to clustered, attenuated, delayed and prolonged events, which are mediated by catchments characteristics, climate factors, and soil moisture dynamics (Van Loon, 2015). Various approaches have been used to evaluate how P deficits propagate through the hydrological cycle into Q deficits (Lorenzo-Lacruz et al., 2013; Rahiz and New, 2013; Haslinger et al., 2014; Zhu et al., 2016; Apurv et al., 2017; Huang et al., 2017; Wu et al., 2017b; Gevaert et al., 2018; Jehanzaib et al., 2020), and studies suggest that combining drought indices based on different variables could provide a stronger agreement with ground-based hydrological impacts (Niemeyer et al., 2008; Sivakumar et al., 2011; Zargar et al., 2011). However, there is still a need to identify which indices and time scales are most appropriate for assessing streamflow droughts, and how this choice varies as a function of the specific hydrological regime. This is particularly important in catchments with a sparse monitoring network, where the identification and assessment of streamflow droughts can be challenging.

Therefore, this study aims to answer the following questions: *i*) what temporal scales of (the frequently used and easy to calculate) SPI and SPEI meteorological indices can be used as proxies for streamflow drought in catchments with different hydrological regimes? *ii*) considering that the soil acts as a natural reservoir to maintain streamflow during periods of reduced P , can a soil moisture drought index be used to assess streamflow droughts instead of the SPI and SPEI meteorological indices?; and *iii*) considering that snowmelt is an important moisture contribution to streamflow and surface water availability during spring and summer months in catchments with a pronounced snow influence, can a SWE index be used to assess streamflow droughts instead of the SPI and SPEI in those catchments? Answering these questions can help us to design effective monitoring and early warning systems, to elaborate better drought mitigation strategies (Huang et al., 2017) and management practices to reduce societal vulnerability to drought (Svoboda et al., 2002), and to better understand how future changes in meteorological variables will impact on streamflow droughts.

2 Study area and selected catchments

Continental Chile spans 4,300 km of latitudinal extension (17.5° – 56.0° S) but only 200 km of longitudinal extension on average (76.0°–66.0°W). It is bounded to the north by Peru, to the west by the Pacific Ocean, and to the east by Bolivia and Argentina,



with elevations ranging from 0 to 6,892 m a.s.l. in the Andes cordillera. Figure 1 shows the elevation (Jarvis et al., 2008), the land cover classification (Zhao et al., 2016), the Köppen-Geiger climate classification (Beck et al., 2018), and the 100 near-natural catchments with different hydrological regimes that are used in this study (same as in Baez-Villanueva et al., 2021). The climate transitions from arid in the north to temperate and humid climates in the Central Chile and South regions. In general, P increases with latitude (in the southern direction) and elevation. Likewise, Q also tends to increase from north to south (Alvarez-Garreton et al., 2018; Vásquez et al., 2021). The country is home of about 18 million inhabitants, two thirds of them concentrated in central Chile (30°–38°S), region where intensive agriculture and other economic activities prevail.

The El Niño-Southern Oscillation (ENSO) influences winter P in Central Chile, producing positive and negative anomalies during El Niño and La Niña events, respectively (Montecinos and Aceituno, 2003; Verbist et al., 2010; Robertson et al., 2014). Recently, Central Chile experienced a prolonged period of dry conditions from 2010 onward, known as the Central Chile *megadrought* (Garreaud et al., 2017), with P deficits of 25–80% combined with an unprecedented increase in evaporative demand, which affected approximately 70% of the Chilean population. These conditions had adverse effects on snowpack, groundwater, and surface water levels, with a decrease in mean river discharge up to 90% (Garreaud et al., 2020) and up to 60% increase in areas affected by forest fires (González et al., 2018).

To exclude antropogenic influences in our study, we selected a subset of 100 near-natural catchments as presented in Baez-Villanueva et al. (2021), which are included in the Catchment Attributes and Meteorology for Large-sample Studies dataset in Chile (CAMELS-CL; Alvarez-Garreton et al., 2018), and summarised in the Supplementary Material. CAMELS-CL dataset provides hydroclimatic indices and geomorphological characteristics for 516 catchments in continental Chile, including their location, topography, geology, soil types, land cover, hydrological signatures, degree of human intervention, among others.

100 3 Data and methods

To address the research questions outlined in Section 1, we followed a three-step procedure, as illustrated in Figure 2: *i*) computing meteorological (SPI, SPEI), soil moisture (ESSMI), and snow (SWEI) drought indices; *ii*) applying cross-correlation and event coincidence analyses among the previous drought indices and the SSI at temporal scale of 1 month as representative of streamflow deficit conditions, within the study catchments; and finally *iii*) evaluating the cross-correlation and event coincidence analysis results considering the hydrological regime of the selected catchments.

3.1 Gridded datasets

Table 1 summarises the gridded datasets used in this study and their related variables. CR2METv2.5 (Boisier et al., 2018) provides daily gridded estimates of P and maximum and minimum temperature (T) over continental Chile at a spatial resolution of 0.05° from 1979 to near present. P estimates are computed by combining ground-based observations with reanalysis data from ERA5, while T is estimated using multivariate regression from Moderate Resolution Imaging Spectroradiometer (MODIS) land surface temperature (LST) and ERA5 as covariates (Alvarez-Garreton et al., 2018; Boisier et al., 2018). The Hargreaves-Samani equation (Hargreaves and Samani, 1985) was used to obtain daily potential evaporation (PE) from CR2METv2.5




Figure 1. Study area adapted from Baez-Villanueva et al. (2021): *a*) elevation (SRTMv4.1; Jarvis et al., 2008) and major macroclimatic zones of Chile: *i*) Far North (17.50–26.00°S), *ii*) Near North (26.00–32.18°S), *iii*) Central Chile (32.18–36.40°S), *iv*) South (36.40–43.70°S), and *v*) Austral/Far South (43.70–56.00°S); *b*) land cover classification (Zhao et al., 2016); *c*) Köppen-Geiger climate classification (Beck et al., 2018); and *d*) 100 near-natural catchments with different hydrological regimes (Baez-Villanueva et al., 2021).

maximum and minimum daily T at the same 0.05° spatial resolution. We selected CR2METv2.5 due to its strong performance compared to other products in previous studies (Baez-Villanueva et al., 2020, 2021).

115 ERA5-Land (Muñoz-Sabater et al., 2021) is an enhanced hourly global dataset for the land component of ERA5 that describes the evolution of energy and water cycles consistently over land. This product was developed through numerical integrations of the ECMWF land surface model driven by the downscaled meteorological forcing from ERA5, which included an elevation correction for the thermodynamic near-surface state (Muñoz-Sabater et al., 2021). ERA5-Land has a spatial resolution of 0.10° and spans from 1950 to the present. The volumetric soil water in this product is associated with soil texture, soil
 120 depth, and the underlying groundwater level. Additionally, ERA5-Land also provides snow depth as SWE .




Figure 2. Schematic summarising the three-step approach to answer the research questions presented in Section 1.

Other soil moisture and *SWE* datasets were also evaluated and deemed less adequate for the study (see Section 1 of the Supplementary Material).

Table 1. Gridded hydrometeorological products used in this study.

Variable	Product	Period	Spatio-temporal resolution	References
<i>P</i>	CR2METv2.5	1979–present	0.05°; daily	Boisier et al. (2018)
<i>T_{max}</i> and <i>T_{min}</i>	CR2METv2.5	1979–present	0.05°; daily	Boisier et al. (2018)
Soil moisture	ERA5-Land	1940–present	0.10°; hourly	Muñoz-Sabater et al. (2021)
<i>SWE</i>	ERA5-Land	1950–present	0.10°; hourly	Muñoz-Sabater et al. (2021)



3.2 Drought Indices

3.2.1 Standardised Precipitation Index

125 The Standardised Precipitation Index (SPI; McKee et al., 1993) is considered the standard for monitoring and characterising
meteorological drought at different time scales (Hayes et al., 2011; WMO, 2012) and has been widely used for decades (e.g.,
Wu et al., 2001; Zhang et al., 2013; Meroni et al., 2017; Liu et al., 2021) including Chile (Zambrano et al., 2017; Zambrano-
Bigiarini and Baez-Villanueva, 2019). The SPI compares the accumulated P for a specific temporal scale at a given location
with its long-term (e.g., 30 years) distribution, making it useful for comparing P deficits or excesses across regions with
130 different climates. The SPI follows a three-step approach: *i*) selecting a probability density function (PDF) and a method for
estimating its parameters (e.g., Hosking and Wallis, 1995) to represent the long-term distribution of accumulated P over the
desired period; *ii*) obtaining the cumulative distribution function (CDF) from the fitted distribution; and *iii*) transforming
accumulated P into a normal distribution with a mean of zero and standard deviation of one, using an equi-percentile inverse
transformation to derive the SPI values. The SPI represent the number of standard deviations that cumulative P deviates from
135 its long-term average (JRC, 2011).

In this study, the SPI was calculated at different scales (1, 3, 6, 9, 12, 18, and 24 months) using CR2METv2.5. The selected P
accumulation periods were adjusted through the two-parameter gamma distribution, as recommended by Stagge et al. (2015b),
in combination with the unbiased Probability Weighted Moments (Hosking and Wallis, 1995) to estimate the parameters of the
gamma distribution.

140 3.2.2 Standardised Precipitation and Evapotranspiration Index

The Standardised Precipitation and Evapotranspiration Index (SPEI; Vicente-Serrano et al., 2010; Beguería et al., 2014) is a
multi-scalar index similar to the SPI. However, instead of using only P as input, the SPEI uses the climatic water balance
given by P minus potential evaporation ($P - PE$). The SPEI is suited to identifying, tracking, and exploring how drought
conditions are affected by global warming, e.g., it allows to robustly consider the influence of recent positive trends in the
145 evaporative demand on the temporal evolution of streamflow (Vicente-Serrano et al., 2014, 2010; Liu et al., 2021). To calculate
this index, the often highly skewed distribution of the climatic water balance is transformed to the standard normal distribution
using the univariate log-logistic probability distribution (Vicente-Serrano et al., 2014, 2010). Temporal scales used to calculate
this index are similar to those of SPI, and share the same relation with monitoring water availability. In this study, the SPEI
was calculated at different scales (1, 3, 6, 9, 12, 18, and 24 months). The selected accumulation periods were adjusted through
150 the log-logistic distribution as recommended by Vicente-Serrano et al. (2010), in combination with the unbiased Probability
Weighted Moments method (Hosking and Wallis, 1995) to estimate the parameters of the log-logistic distribution.



3.2.3 Empirical Standardised Soil Moisture Index

Soil moisture modulates both the land surface water and energy cycles, and influences plant productivity and yields, playing a key role in agriculture and water supply. Among the many indices available for monitoring soil moisture deficits (e.g. Zargar et al., 2011; Sims et al., 2002; Wang et al., 2015), we chose the Empirical Standardised Soil Moisture Index (ESSMI; Carrão et al., 2016), a spatially-invariant probabilistic index designed to detect soil moisture anomalies. The ESSMI was chosen because its non-parametric nature has proven to be more accurate than previously proposed parametric indices (Carrão et al., 2016). The computation of the ESSMI was carried out by fitting a non-parametric empirical probability density function (ePDF) to historical time-series of SM, using a kernel density estimation (Silverman, 1986) optimised for each grid-cell, and then transforming it into a normal distribution with a mean of zero and standard deviation of one. As for the SPI and SPEI, negative standard normal values indicate dry conditions, whereas positive values indicate wet conditions. Typical accumulation periods used for ESSMI are 1, 3, 6, or 12 months. We used soil moisture data of ERA5-Land after evaluating other products such as ERA5, SMAP-L3E and SMAP-L4SM (see Section 1 of the Supplementary Material).

3.2.4 Standardised Snow Water Equivalent Index

The Standardised Snow Water Equivalent Index (SWEI; Huning and AghaKouchak, 2020) aims at quantifying snow drought conditions. To compute this index we followed Huning and AghaKouchak (2020) who used a non-parametric approach (Farahmand and AghaKouchak, 2015) to standardise SWE time series. In general, instead of fitting a specific distribution function to the data, the SWEI computes the probabilities associated with the SWE time series using the empirical Gringorten plotting position (Gringorten, 1963):

$$p(A_{m,i}) = (i - 0.44)/(N + 0.12) \quad (1)$$

where i is the rank of the non-zero variable (in increasing order), and N is the sample size. These ranks are determined using the 3-month SWE integration for any given month ($A_{m,i} = SWE_m + SWE_{m-1} + SWE_{m-2}$). In this sense, $A_{m,i}$ provides an integrated measure of the SWE amount and its persistence over each value. Afterwards, $A_{m,i}$ is standardised by transforming the empirical probability p to the standard normal distribution:

$$SWEI(A_{m,i}) = \phi^{-1}[p(A_{m,i})] \quad (2)$$

While this index has the potential to be used across different time scales, its interpretation differs from other drought-related variables because areas with seasonal snowpack frequently experience several months without SWE during the warm season. Therefore, Huning and AghaKouchak (2020) recommend the use of the SWEI with scales lower than six months to assess multi-year droughts. As recommended by Huning and AghaKouchak (2020) *i*) the SWEI index was only calculated for accumulation periods of 1, 3, and 6 months, and *ii*) the warm season months (November–April) were discarded to avoid introducing biases in the analysis of the index, as in regions with seasonal snowpack there are multiple months without SWE . We assumed that the selected SWE product provides useful information on the spatial distribution of snow and, therefore, discarded grid-cells that had SWE values < 1 mm.



3.2.5 Standardised Streamflow Index

185 Streamflow drought indices directly explain the consequences of climatic anomalies on current hydrological conditions, in contrast to climate-based indices that describe climate anomalies in isolation from their hydrological context (Shukla and Wood, 2008). The SSI follows the same idea of the SPI: *i*) selecting a PDF and estimating its parameters to model the long-term distribution of Q for a given period, and *ii*) deriving the corresponding CDF to transform Q into a standard normal distribution. Here, we used the SSI with an accumulation period of one month (SSI-1) to characterise streamflow drought
190 because *i*) it integrates catchment-scale hydrogeological processes (Barker et al., 2016a), *ii*) similarly to the soil moisture, it already integrates the hydrometeorology of the past months (Stahl et al., 2020), and *iii*) has been widely used in previous studies of streamflow droughts (Barker et al., 2016a; Stahl et al., 2020; Li et al., 2020; Wang et al., 2020; Bevacqua et al., 2021). Due to the important variability in the statistical properties of monthly series of climatic water balance, several probability distributions can be used to compute a reliable SSI (Vicente Serrano et al., 2012). Here, we use the GEV distribution to
195 calculate the SSI, as it was one of the recommended distributions by Vicente Serrano et al. (2012), besides the log-logistic distribution. Q data were obtained for each catchment from the Center for Climate and Resilience Research (CR2; <http://www.cr2.cl/datos-de-caudales/>).

3.3 Evaluation of proxies of streamflow drought

For evaluating the ability of the selected drought indices to be used as proxies for streamflow drought, we employed two
200 analyses in each one of the 100 near-natural catchments: *i*) a cross-correlation analysis, to measure the similarity between the time series of the selected drought indices (SPI, SPEI, ESSMI, SWEI) at different temporal scales and the SSI-1, and *ii*) an event coincidence analysis, to evaluate the statistical interdependence between specific events from the selected indices and SSI-1. Both analysis were performed for the period 1979–2020, using lag times from 0 to 12 months between the analysed drought index and SSI-1.

205 Results of the cross-correlation and event coincidence analysis for the selected 100 near-natural catchments were evaluated both for all the catchments at once, and for the catchments grouped by hydrological regime. The analysed hydrological regimes are: *i*) nival (16 catchments), *ii*) nivo-pluvial (nival catchments with a rain component: 25 catchments), *iii*) pluvio-nival, (pluvial catchments with a snow component: 40 catchments), and *iv*) pluvial (19 catchments), as shown in Figure 1d. These hydrological regimes were determined according to the contribution of solid and liquid P to the mean monthly Q values as
210 described in (Baez-Villanueva et al., 2021). Figure S1 of the Supplementary Material, adapted from Baez-Villanueva et al. (2021), displays conceptual hydrographs for each one of these hydrological regimes.

3.3.1 Cross-correlation analysis

A cross-correlation analysis (Equation 3) measures the similarity of two time series as a function of the relative displacement of one relative to the other. Two time series (X_t and Y_t) may be related at different lags between each other (e.g., the time
215 series Y_t may be related to past or future lags of X_t). The cross-correlation function is defined as a set of correlations between



X_{t+h} and Y_t for different lags ($h = 0, \pm 1, \pm 2, \dots, \pm n$). In this study, X_t represents the time series of the SSI-1 for each one of the 100 selected catchments, while Y_t represents the time series of SPI, SPEI, ESSMI, and SWEI, alternatively. This analysis is useful for identifying whether Y_t could be useful to predict X_t . Therefore, we used positive lag values from zero (cross-correlation analysis at the same period for both variables) to 12 months assuming that the variable X lags the variable Y . In this sense, X lags Y means that changes in the selected drought indices (the SPI, SPEI, ESSMI, and SWEI) precede changes in the SSI-1; and therefore, variations in P , PE , soil moisture, and SWE might have a delayed impact on Q . The cross-correlation analysis have been previously applied to analyse the propagation from meteorological to streamflow droughts and investigate the influence of climate and catchment properties on streamflow drought characteristics (Barker et al., 2016a; Peña-Gallardo et al., 2019).

$$r_h = \frac{\sum_{t=1}^n [(Y_t - \bar{Y})(X_{t+h} - \bar{X})]}{\sqrt{\sum_{t=1}^n (Y_t - \bar{Y})^2} \sqrt{\sum_{t=1}^n (X_{t+h} - \bar{X})^2}} \quad (3)$$

where

3.3.2 Event coincidence analysis

The cross-correlation analysis uses the whole time series of the selected drought indices and, therefore, takes into account the influence of non-drought periods. To overcome the previous limitation of the cross-correlation analysis, we also use the event coincidence analysis in this work (Donges et al., 2016; Siegmund et al., 2017), a statistical technique specifically designed to assess the inter-dependency between two event time series; in this case, those of the meteorological, soil moisture or snow drought index, and the SSI-1. This method examines the degree of temporal association between events by assessing the frequency of simultaneous events, considering delayed responses (specified by a lag term τ) and uncertain timing (specified by a coincidence window ΔT) between binary time series (Siegmund et al., 2017). The binary series are computed using a threshold, where entries with a value of one correspond to time steps with an event, and entries with zero correspond to time steps without an event (Donges et al., 2016). This method is able to quantify the strength of statistical interrelationships between two event series A and B, addressing B-type events as precursors (the event B precedes event A) and triggers (the event B triggers the event A). We analysed only precursor coincidence rates in this study, and not trigger rates, because we were interested in determining the number of streamflow drought events preceded by or coinciding with abnormal meteorological, soil moisture, and snow drought events, and not the number of streamflow droughts that were triggered by meteorological, soil moisture, or snow droughts. The precursor coincidence rate is calculated as follows:

$$r_p(\Delta T, \tau) = \frac{1}{N_A} \sum_{i=1}^{N_A} \Theta \left[\sum_{j=1}^{N_B} 1_{[0, \Delta T]}((t_i^A - \tau) - t_j^B) \right] \quad (4)$$

Equation 4 quantifies the proportion of A-type events that are preceded by at least one B-type event. It considers multiple B-type events within the coincidence interval as a single occurrence. The Heaviside function $\Theta(\cdot)$ is used in the equation,



245 defined as $\Theta(x) = 0$ for $x \leq 0$ and $\Theta(x) = 1$, otherwise. Additionally, the indicator function $1_I(\cdot)$ is employed to represent the presence of values within a specific interval I , defined as $1_I(x) = 1$ for x within I and $1_I(x) = 0$ otherwise. The optimal value of the precursor coincidence rate is one and represents the ratio of events A that coincide with events B. For example, if a value of 0.6 is obtained in this case, it means that 60% of streamflow droughts align with a drought event identified using the chosen index.

250 This method, has been widely used to analyse *i*) floods as triggers of epidemic outbreaks (Donges et al., 2016); *ii*) meteorological extremes (Siegmund et al., 2016); *iii*) the timing of droughts and floods (Siegmund et al., 2017); and *iv*) droughts and wet spells (He and Sheffield, 2020), and to calculate precursor and trigger coincidence rates (Donges et al., 2016). Here, we use two different thresholds to define an event (Lloyd-Hughes and Saunders, 2002; Barker et al., 2016a): *i*) a value ≤ -1.0 to evaluate moderate droughts, and *ii*) a value ≤ -1.5 to evaluate severe droughts. We excluded extreme droughts (values
255 ≤ -2.0) as not all indices presented standardised values below -2.0 . In this sense, values lower than or equal to the selected threshold are considered as events (value = 1), while values greater than the threshold represent periods without an event (value = 0).

Additionally, to assess the strength of the statistical relationship between these drought types, we performed a significance test assuming a Poisson process, which quantifies the robustness of the relationship between them. This test evaluates the
260 hypothesis that the occurrence of these signals is distributed randomly and not related to each other. The methodology for this approach is explained in detail in Donges et al. (2016) and has been used in previous studies (Donges et al., 2016; Siegmund et al., 2017; He and Sheffield, 2020). A schematic representation of the event coincidence analysis is presented in Figure 3.




Figure 3. Schematic of the event coincidence analysis between meteorological, soil moisture, or snow drought indices and the SSI-1 considering a time lag τ with a certain window ΔT .



4 Results and discussion

4.1 Proxies of streamflow drought: all catchments

265 This section analyses the cross-correlation values and precursor coincidence rates between the selected drought indices (SPI, SPEI, ESSMI, SWEI) and the SSI-1 when all the 100 near-natural catchments are analysed at once. An arbitrary lower threshold of 0.4 was used for cross-correlation values and precursor coincidence rates to separate drought indices and temporal scales that could be considered a good proxy of streamflow droughts, from those that could not. While the analysis was done for diverse lags, the cross-correlation values and precursor coincidence rates, decreased gradually from zero lag (lag = zero months) to 270 a lag of 12 months. Therefore, in this section, all the results are analysed only for a lag of zero months, i.e., when the SSI-1 has no time delay respect to the analysed drought indices. The remainder of the analysis can be found in the Supplementary Material (Figures S2–S4 and S5–S7).

Figure 4 displays the linear correlation values (zero lag) between the selected drought indices and the SSI-1. The highest median correlation values are obtained for SPI-6 (~ 0.58), followed closely by SPI-3 and SPI-9 (~ 0.57 both of them). A similar 275 behaviour is obtained for SPEI, which yields the maximum median correlation for SPEI-3 and SPEI-6 (~ 0.52). Interestingly, the values of SPEI exhibit a lower median correlation than those of the SPI, which suggests that, in the selected Chilean catchments, the variations in the climatological water balance ($P - PE$) are of less importance than the variations in P alone. This happens mainly in arid regions, where the PE is very high compared to the actual evaporation. Our results are in line with those of Barker et al. (2016a), who found the highest correlation values between SSI-1 and SPI at temporal scales ranging 280 from 1 to 9 over most of the 121 evaluated UK near-natural catchments; and those of Peña-Gallardo et al. (2019), who found that SPEI at temporal scales from two to four months are the most correlated with SSI-1.

The soil moisture drought index ESSMI reaches its highest median correlation value at a temporal scale of 1 month (ESSMI-1 ~ 0.50), followed by ESSMI-3 (~ 0.47). This indicates that shorter accumulation periods of soil moisture (from the antecedent 1 to 3 months) align better with the SSI-1 signal over these catchments. Finally, the median correlation values for SWEI are all 285 lower than 0.4, with the highest values obtained for SWEI-1 at lags from zero to five months, suggesting that snow accumulated during the cold season influences the Q observed in the next hot season. This results indicate that nival catchments have a higher streamflow predictability as concluded by Pechlivanidis et al. (2020). In this sense, SWEI-1 might provide useful information related to deficits and surplus of streamflow values up to five months in advance as the correlation analysis also includes periods with near-normal and surplus conditions. However, the relatively low correlation values of the SWEI could also be influenced 290 by the ability of ERA5-Land in representing appropriately the SWE over the evaluated catchments.

Figure 5 displays the spatial distribution of the linear correlation values (zero lag) between the selected drought indices and the SSI-1 at different temporal scales. The spatial patterns observed at different temporal scales are similar, therefore, here we focus on the spatial patterns obtained at zero lag and the scales that yielded the highest cross-correlation values. Figures S8–S11 in the Supplementary Material show the spatial distribution of the cross-correlation values of the selected drought 295 indices at different temporal scales and lag values ranging from 0 to 12 months.




Figure 4. Linear correlation (zero lag) between the selected drought indices and the SSI-1 over all catchments. The horizontal blue line indicates the optimal correlation value. The solid line within each box represents the median value, the edges of the boxes represent the first and third quartiles, and the whiskers extend to the most extreme data point which is no more than 1.5 times the interquartile range from the box.

The spatial distribution of the median correlation values of SPI-6 and SPEI-3 is similar. For both indices, the highest values (~ 0.4 – 0.8) are obtained in Central and Southern Chile (30 – 43°S), with low values (~ 0.4 – 0.2) in the Far and Near North (17.5 – 30°S) and Far South (43 – 56.6°S , ~ 0.4 – 0.0), which are related to the reduced capability of CR2METv2.5 (and other P products) in representing accurately the P patterns of these areas (Baez-Villanueva et al., 2020). It is worth noting that the highest values in Central Chile (32 – 36°S) are obtained with SPI-12 (~ 0.6 – 0.8) and SPEI-12 (~ 0.4 – 0.8), indicating that streamflow droughts are related to longer P deficits over those catchments, which could be attributed to higher baseflow coefficients (Alvarez-Garreton et al., 2018, their Figure 10e). The ESSMI-1 shows the highest median correlation values (~ 0.4 – 0.7) in Central and Southern Chile (34 – 43°S), with low values (~ 0.2 – 0.4) in the Far and Near North (17.5 – 34°S) and Far South (43 – 56.5°S , ~ 0.5 – 0.0). Similarly to SPI and SPEI, the highest values of correlation between ESSMI and SSI-1 in the Near North (32 – 36°S) are obtained with ESSMI-6 (~ 0.5 – 0.8). Finally, SWEI-1 shows its highest median correlation values (~ 0.3 – 0.6) in Central Chile (30 – 34°S) yielding relatively low values (≤ 0.3) elsewhere.

We used two thresholds to define a drought event in the event coincidence analysis: *i*) -1.0 for *moderate* drought events, and *ii*) -1.5 for *severe* drought events as done in previous studies (Andreadis et al., 2005; Sheffield et al., 2009). Figure 6 shows boxplots with the precursor coincidence rates (hereafter referred to as coincidence rates) for all evaluated indices for moderate




Figure 5. Spatial distribution of the linear correlation (lag zero) between the selected drought indices (SPI, SPEI, ESSMI, SWEI) and the SSI-1 over the 100 near-natural catchments.

310 drought events ($thr = -1.0$). Coloured boxplots indicate the temporal scales at which coincidence rates are significant at a confidence level of 95% for at least 75% of the evaluated catchments (75 in this case). In contrast, white boxplots show the temporal scales where less than 75% of the catchments had statistically significant coincidence rates.

The highest median coincidence rates are obtained for SPI-6 (~ 0.53), followed closely by SPI-3 (~ 0.52) and SPI-9 (~ 0.50). A similar behaviour is obtained for SPEI, which achieves the maximum coincidence rates for SPEI-6 (~ 0.51), followed



315 closely by SPEI-3 and SPEI-9 (~ 0.50 both of them). Our results indicate that, as expected, meteorological droughts strongly
control streamflow droughts and that P influences meteorological droughts the most compared to the evaporative demand as
suggested by the small differences found between the SPI and SPEI. On the other hand, the median coincidence rates for
ESSMI are all lower than 0.4, and they reach their highest values at a temporal scale of 3 months (ESSMI-3 ~ 0.31), followed
by ESSMI-6 (~ 0.29) and ESSMI-1 (~ 0.28). Similarly to the SWE , the lower performance of the ESSMI could be related,
320 to some extent, to the ability of the product in representing the soil moisture patterns across Chile.

While the results of the cross-correlation analysis and the event coincidence analysis cannot be directly compared, it is worth
noting the substantial differences between both analyses regarding the ESSMI at the evaluated lags. Although soil moisture
and Q signals are related, as evidenced by the relatively high cross-correlation results between the ESSMI and the SSI-1, the
substantially lower event coincidence rates between these indices suggest that a high correlation does not mean that a specific
325 drought index could be used to assess streamflow droughts as the correlation also includes periods with near-normal and surplus
conditions (in this case of soil moisture and Q). These differences in the case of the ESSMI could be related to the fact that
soil moisture has a more immediate influence on the runoff coefficient via infiltration during rainy periods compared to its
influence on evaporation. Therefore, as correlation evaluates also periods with surplus of the evaluated variables, it provides
a better agreement between the standardised soil moisture and Q signals. On the other hand, our implementation of the event
330 coincidence analysis focuses only on dry conditions making it a better method to relate the selected indices to streamflow
droughts.

The SWEI scored the lowest results (median values lower than 0.4 and not statistically significant) among the evaluated
indices, which is expected since the snow influence is confined to a limited number of catchments. Finally, the median coinci-
dence rates for SWEI are all also lower than 0.4 and not statistically significant. This can be explained by the fact that the SWEI
335 and ESSMI have fewer events below the -1.5 threshold used to define severe events compared to the SSI-1, which affects the
characterisation of drought events. In other words, different indices might have different thresholds related to the occurrence
of specific drought events. This result suggests that the threshold used to define moderate or severe droughts for the case of the
event coincidence analysis could vary according to the component of the hydrological cycle that is being evaluated; however,
the evaluation of the optimal threshold to characterise droughts is out of the scope of this study.

340 Figure 7 shows the spatial distribution of the coincidence rates at lag zero between the selected drought indices and the
SSI-1 at different temporal scales for moderate droughts, while Figures S12–S20 and S16–S19 in the Supplementary Material
show coincidence rates for moderate and severe droughts, respectively, considering lag values ranging from 0 to 12 months.
The spatial distribution of the median coincidence rates of SPI-6 and SPEI-6 are similar (Figure 7).

For both indices, the highest coincidence rates (~ 0.4 – 0.8) are obtained in Central and Southern Chile (30 – 43°S), with
345 low values (~ 0.0 – 0.3) in the Far and Near North (17.5 – 30°S) and Far South (43 – 56.6°S). It is worth noting that the highest
coincidence rates in the Near North (30 – 34°S) are obtained with SPI-24 (~ 0.8 – 0.5) and SPEI-24 (~ 0.8 – 0.3), closely followed
by SPI-12 (~ 0.8 – 0.3) and SPEI-12 (~ 0.8 – 0.4). This finding is consistent with Liu et al. (2019), who found weaker relationships
between the SPEI and the Standardised Runoff Index (SRI) in the arid region of the Tibetan Plateau. This could be attributed
to the fact that in periods when P is very low there is a point where the evaporative demand fluctuations could be deemed as




Figure 6. Precursor coincidence rates between the selected drought indices (SPI, SPEI, ESSMI, SWEI) and the SSI-1 at a lag of zero months for moderate droughts ($thr = -1.0$). Coloured boxplots indicate that -at least- 75% of the catchments presented significant coincidence rates at a 95% confidence interval, while white boxplots indicate no statistically significant rates. The horizontal blue line indicates the maximum possible coincidence rate. The solid line within each box represents the median value, the edges of the boxes represent the first and third quartiles, and the whiskers extend to the most extreme data point which is no more than 1.5 times the interquartile range from the box.

350 irrelevant, which is consistent with periods of more than 3 to 6 months without P , a normal situation in northern Chile. The high accumulation periods for the SPI and SPEI (up to 24 months) is likely due to the occurrence of multi-annual droughts, which have particularly affected the central-northern region of Chile (Garreaud et al., 2020). Regarding the spatial distribution of the soil moisture index ESSMI-3, it shows its highest median coincidence rates ($\sim 0.5-0.3$) in Southern Chile ($36-43^\circ\text{S}$), with low values lower than 0.4 elsewhere. Finally, SWEI-1 shows its highest median linear correlation values ($\sim 0.6-0.3$) in
 355 Central Chile ($30-34^\circ\text{S}$), and lower values (≤ 0.3) elsewhere.

The highest cross-correlation and event coincidence rates were obtained at a lag zero, which is in agreement with Peña-Gallardo et al. (2019). Considering the results of all catchments in the cross-correlation and event coincidence analyses, it can be observed that —in general— the SPI-6, followed by SPI-3, and SPI-9 are the best indices to assess streamflow drought, with median correlation values ranging from 0.57 to 0.58, and precursory coincidence rates ranging from 0.52 to 0.53 for moderate
 360 drought events (and from 0.45 to 0.53 for severe drought events, see Table S2 of the Supplementary Material). However, in the Far and Near North, the SPI-12, SPI-24, and SPEI-12 are the best meteorological proxies of streamflow droughts, which highlights that these indices provide complementary information and that a single SPI or SPEI cannot describe the complex




Figure 7. Spatial distribution of the event coincidence analysis results for moderate droughts between the selected indices and the SSI-1 over 100 near-natural catchments and at a lag of zero months.

P-Q interaction over Chile. Therefore, a subregional approach should be used considering the climatic gradient of Chile and the hydrological response of the catchments.

365 On the other hand, the median correlation values and coincidence rates for ESSMI at different temporal scales were all lower than 0.4, indicating that ESSMI is not a better proxy for streamflow droughts than SPI and SPEI. Similarly, Figures 4, 5, 6, and 7 show that when all the catchments are analysed at once, SWEI is not a better proxy than SPI or SPEI at any temporal scale.



Overall, the results suggest that various hydrological processes are likely to influence the response of the 100 near-natural catchments situated within Chile's edapho-topo-climatologically diverse territory. Although the ESSMI and the SWEI did not yield higher values compared to the SPI and SPEI, these indices could be combined into a sequential drought index. This index could consider P deficits first, followed by soil moisture deficits and/or snow deficits to mitigate the occurrence of false drought signals.

4.2 Influence of hydrological regime

The influence of the hydrological regime of the catchments on the relationship between different scales of meteorological, soil moisture, and snow drought indices and streamflow droughts, although expected and intuitive, is almost never considered in large-scale drought analyses and characterisations. Although evaluating all catchments together provided general recommendations for which indices could be used to assess streamflow droughts, analysing the results for catchments with similar hydrological regimes provided additional insights into the relationship between the selected drought indices and how they relate to streamflow droughts. Therefore, this section analyses the cross-correlation values and precursor coincidence rates between the selected drought indices and the SSI-1 considering the hydrological regimes of the 100 near-natural catchments: *i*) nival (16 catchments); *ii*) nivo-pluvial (25 catchments); *iii*) pluvio-nival (40 catchments); and *iv*) pluvial (19 catchments; see Figure1).

In this sense, only cross-correlation values and precursor coincidence rates higher than 0.4 are considered as indicative that the analysed drought index and time scale is a good proxy of streamflow droughts. Figures S21–S24 and S25–S33 of the Supplementary Material show that the cross-correlation and event coincidence analysis values decreased gradually from zero lag to a lag of 12 months for SPI, SPEI and ESSMI at all temporal scales, with the only exception of SPI and SPEI for nival catchments, where high correlation values and coincidence rates extend to lags of 3-6 months. Therefore, all indices are analysed hereafter only for a lag of zero months, i.e., when the SSI-1 has no delay in time with the analysed drought indices.

Figure 8 displays the correlation values between the selected drought indices and the SSI-1 at a lag of zero months. For nival catchments the highest median correlation values are obtained for SPI-12 (~ 0.75), followed by SPEI-12 (~ 0.51), ESSMI-6 (~ 0.51) and SWEI-6 (~ 0.51). Nivo-pluvial catchments show a slightly different behaviour, with the highest median correlation values obtained for SPI-12 (~ 0.52), followed by ESSMI-3 (~ 0.49), SPEI-3 (~ 0.48), and SWEI-1 being a poor proxy of streamflow drought (~ 0.38). The highest median correlation values for pluvio-nival catchments are obtained for SPI-3 (~ 0.73), followed by SPEI-3 (~ 0.68), ESSMI-3 (~ 0.55) and SWEI-1 (~ 0.25). Finally, for pluvial catchments the highest median correlation values are obtained for SPEI-3 (~ 0.63), followed by SPI-3 (~ 0.60), ESSMI-1 (~ 0.52) and, as expected, SWEI-3 not being a good proxy of streamflow drought (~ 0.1). The SWEI showed the highest values over nival catchments, followed by nivo-pluvial and pluvio-nival catchments, respectively. These findings are consistent with Tijdeman et al. (2018), who obtained high correlation values over cold climates in the US, and with Staudinger et al. (2014), who found similar results over alpine catchments in Switzerland. The larger scales of the SPI and SPEI over nival catchments are related to the fact that Q is mostly controlled by snow melt that occurs after snow accumulation; contrastingly, the pluvial catchments react rapidly to P and therefore, P deficits are more quickly propagated into Q deficits.




Figure 8. Linear correlation values between the selected drought indices (SPI, SPEI, ESSMI, SWEI) and the SSI-1, using a lag of zero months for SPI, SPEI and ESSMI and a lag of 6 months for SWEI. From left to right, the four columns corresponds to *i*) nival; *ii*) nivo-pluvial; *iii*) pluvio-nival; and *iv*) pluvial catchments. The horizontal blue line indicates the optimal cross-correlation. The solid line within each boxplot represents the median value, the edges of the boxes represent the first and third quartiles, and the whiskers extend to the most extreme data point which is no more than 1.5 times the interquartile range from the box. A detailed summary of the highest results can be found in Table S3 of the Supplementary Material.



Figure 9 shows the coincidence rates between the selected drought indices and the SSI-1 for moderate drought events ($thr = -1$) with zero lag. For all hydrological regimes, ESSMI and SWEI presented median coincidence rates lower than 0.4 (i.e., less than 40% of streamflow droughts preceded by soil moisture and snow droughts) for all time scales and, therefore, are not considered good proxies of streamflow drought events. For nival catchments, the highest coincidence rates are obtained for SPI-24 (~ 0.74), followed by SPEI-18 (~ 0.52). Nivo-pluvial catchments show a different behaviour, with the highest median correlation values obtained for SPEI-9 (~ 0.50), followed by SPI-6 (~ 0.48). The highest median correlation values for pluvio-nival catchments are obtained for SPI-6 (~ 0.68), followed by SPEI-3 (~ 0.60). Finally, for pluvial catchments the highest median correlation values are obtained for SPI-3 (~ 0.55), followed by SPEI-3 (~ 0.49). The results of the event coincidence analysis per hydrological regime for severe drought events ($thr = -1.5$) show a behaviour similar to that of moderate droughts, but with lower values, except for SPI and SPEI in pluvio-nival catchments, where coincidence rates for severe events are higher than those of moderate ones (Figure S25).

Our results show that the SPI scored the highest cross-correlation and event coincidence values among all evaluated indices, which can be attributed to the fact that P is the most important variable that drives the Q response. Similarly, Barker et al. (2016a) found high cross-correlation values between the SPI-1 and SPI-3, and the SSI-1 over the majority of 121 near natural catchments in the UK. The event coincidence analysis showed relatively low values of the ESSMI compared to those obtained with the cross-correlation analysis, indicating that even if an index has a high correlation with the SSI-1, it may not be able to accurately detect specific streamflow drought events. This result highlights the fact that a high correlation between two indices does not imply that they can be interchangeably used to assess drought conditions. The SWEI showed the highest values over pluvio-nival catchments, which could be related to T increases (Garreaud et al., 2020) and the loss of snow cover over these catchments (Notarnicola, 2022). However, these values were substantially low compared of those of the cross-correlation analysis, which could be attributed to the fact that nival catchments present 5-6 months of delay between the rainfall season (June–August) and the months with maximum discharges (October–December), reinforcing the idea that a correlation analysis does not necessarily indicate that an index could be used to assess streamflow droughts.

Figure 10 depicts the spatial distribution of the temporal scale with the highest correlation (zero lag) and precursor coincidence rates for all indices. These results show that, although the cross-correlation analysis and event coincidence analysis are in close agreement concerning the spatial distribution of the recommended scales for each index, the event coincidence analysis suggests slightly higher scales compared to the cross-correlation analysis. For example, over the South region, the SPI provides the highest results between accumulation periods of 1–9 months, while for the event coincidence analysis for moderate droughts, scales up to 12 months are visible. We conducted a Spearman correlation analysis to examine the relationship between the scale that yielded the highest values for each index and the hydrological regime of the catchments. Our findings show that only the SPEI exhibited a significant correlation for all analyses at a 95% significance level (the cross-correlation analysis and the event coincidence analysis for both moderate and severe droughts). The SPI yielded significant results for the cross-correlation analysis and the event coincidence analysis for severe droughts, while the ESSMI yielded significant results only for the cross-correlation analysis. Finally, the SWEI provided significant results for the event coincidence analysis when both the moderate and severe droughts were analysed despite their values being relatively low. These results suggest that the




Figure 9. Event coincidence analysis results of the selected drought indices and the SSI-1 for moderate droughts at a lag of zero months for *i*) nival; *ii*) nivo-pluvial; *iii*) pluvio-nival; and *iv*) pluvial catchments. Coloured boxplots indicate lags where at least 75% of the catchments presented significant results at the 95% confidence interval, while white boxplots indicate the opposite. The blue line indicates the optimal cross-correlation. The solid line within each box represents the median value, the edges of the boxes represent the first and third quartiles, and the whiskers extend to the most extreme data point which is no more than 1.5 times the interquartile range from the box. A detailed summary of the highest results can be found in Table S4 of the Supplementary Material.



SWEI alone struggles in representing the surpluses and deficits of Q over snow-influenced catchments, which can be related to the ability of ERA5-Land in representing the SWE over these catchments. However, its inclusion in combination with other indices could be beneficial, which is in line with previous studies highlighting the importance of incorporating snow-related information in drought analyses and drought prediction studies (Huning and AghaKouchak, 2020; Gottlieb and Mankin, 2021).

For the specific case of Chile, the drought indices presented in Table 2 could be used to assess streamflow droughts based on the evaluated hydrological regimes (Figures 8–10). These results suggest that the greater the snow influence in the catchments, the higher the accumulation period required for the variables in the evaluated meteorological, soil moisture, and snow drought indices. These results are in line with Fluixá-Sanmartín et al. (2018), who recommended the use of SPI-6 to assess droughts using a catalogue of historical droughts over the snow-influenced Jinsha River Basin in China, and with Vicente-Serrano and López-Moreno (2005) who determined that Q respond to short SPI time scales in the snow-influenced high basin of the Aragon River in Spain.

Table 2. Drought indices that, for the particular case of Chile, could be used to assess streamflow droughts based on the evaluated hydrological regimes.

Hydrological Regime	Scales			
	SPI	SPEI	ESSMI	SWEI
Nival	12–24	12–18	6–12	-
Nivo-pluvial	3–12	3–6	1–3	-
Pluvio-nival	3–6	3–6	1–3	-
Pluvial	3–6	1–6	1–3	-

This evaluation provides a new perspective on the importance of the hydrological regime of catchments in the relationship between meteorological, soil moisture, and snow indices and streamflow droughts, with specific focus on the complex topography and climate of Chile. Previous studies have focused primarily on the influence of geomorphological catchment characteristics on these relationships. For example, Peña-Gallardo et al. (2019) found that elevation is an important driver in the response of SSI to SPEI, while (Van Loon and Laaha, 2015) concluded that drought deficit over Austrian catchments is influenced by catchment average wetness and elevation. Similarly, our findings show that nival catchments, which are located at higher elevations, exhibit different behaviour in the response of the evaluated indices to SSI-1 compared to catchments with other hydrological regimes. Additionally, it is worth also highlighting the different behaviour of central-northern catchments that exhibit longer accumulation periods compared to the more southern, wetter catchments.

5 Conclusion

Assessing streamflow droughts in data-scarce regions can be challenging due to the unavailability of Q data (Peña-Gallardo et al., 2019), which can cause mismatches between the streamflow drought hazards and their related impacts. Publicly available gridded datasets of P , air temperature, evaporation, soil moisture, and SWE , among others, can be used to compute drought




Figure 10. Spatial distribution of the scale that scored the highest correlation (left panel) and precursor coincidence rates (middle and right panel for moderate and severe droughts, respectively) at lag zero over 100 near-natural catchments for each index.

indices in data-scarce areas and serve as a proxy to assess streamflow droughts. In this study, the CR2MET Chilean dataset was used to provide P and PE data, while soil moisture and SWE were collected from ERA5-Land for 1979–2020. We used 100



near-natural catchments in Chile to evaluate which drought indices and temporal scales can be used as proxies of streamflow droughts in catchments with four main types of hydrological regimes (nival, nivo-pluvial, pluvio-nival, pluvial). First, we
465 computed the SPI, SPEI, ESSMI, and SWEI at different temporal scales in the selected catchments. Second, we applied a cross-correlation and event coincidence analysis between the previous indices and the SSI-1, as representative of streamflow drought events. Finally, linear correlation and precursor coincidence rates were analysed for all catchments simultaneously and separated by the hydrological regime. Our results indicate that there is no single best index and temporal scale that could be used as a proxy for streamflow droughts, and that this relationship depends, to some extent, on the hydrological regime of the
470 catchments, and therefore, on their hydrological behaviour. In the following lines we summarise our key findings:

1. There is no single meteorological, soil moisture, or snow drought index and temporal scale that could be used to characterise streamflow droughts in the diverse climate regions of Chile. The best index and temporal scale depend on the hydrological regime of each catchment.
2. The greater the snow influence in a catchment, the larger the temporal scale of the drought index to be used as proxy of streamflow drought, and the more valuable SWEI becomes.
475
3. The nival catchments show a less pronounced decrease in cross-correlation and event coincidence values at increasing lags, and therefore, the evaluated indices could be used to assess streamflow droughts up to five or six months in advance. Contrastingly, pluvio-nival and pluvial catchments show a rapid decay in cross-correlation and event coincidence values at increasing lags, and therefore low predictability of streamflow droughts.
4. For the Chilean case study in particular, the following drought indices are advised (individually or jointly) to monitor streamflow droughts depending on the hydrological regime of the study area: *i*) for nival catchments: SPI-12–24, SPEI-18; *ii*) for nivo-pluvial catchments: SPI-6–12, SPEI-3–9; *iii*) for pluvio-nival catchments: SPI-3–6, SPEI-3; and *iv*) for pluvial catchments: SPI-3 and SPEI-3.
480
5. Notwithstanding soil moisture is a key variable modulating the propagation of meteorological deficits into hydrological ones, meteorological indices are better proxies of streamflow droughts than soil moisture ones.
485
6. Meteorological drought indices are better proxies of streamflow drought events than SWEI index over snow-influenced catchments.
7. Interestingly, the cross-correlation values and event coincidence rates of the SPI were generally higher than those of the SPEI, indicating that for the selected Chilean catchments, the variations in the climatological water balance ($P - PE$) are of less importance than the variations in P alone. This was more pronounced over arid regions, where the PE is very
490 high compared to the actual evaporation.
8. As the cross-correlation analysis includes the full signal of the evaluated indices and the SSI-1 (periods with surplus and deficits), a high cross-correlation value does not necessarily imply that the evaluated drought indices could be used to



495 assess streamflow drought conditions. Therefore, future studies evaluating the use of meteorological, soil moisture, and
snow drought indices as proxies of streamflow drought should consider methods that evaluate solely drought conditions,
such as the event coincidence analysis.

The approach demonstrated in this paper highlights the effectiveness of using individual drought indices to characterise
streamflow drought, while also highlighting the varying behaviour of these indices across catchments with diverse hydro-
logical regimes. This suggests that combining different drought indices based on catchment-specific responses could provide
500 further avenues for strengthening the relationship between meteorological, soil moisture, and snow drought indices as proxies
of streamflow drought. Although catchment characteristics are important in understanding the relationship between meteorolo-
gical, soil moisture, snow drought and streamflow droughts, this article provides a complementary perspective on how this
relationship can also be related to the hydrological regime of the catchments. Our findings indicate that hydrological regimes
offer valuable insights into the time lag at which these indices could be used to assess streamflow drought. For instance, nival
505 catchments exhibited relatively high rates of event coincidence at lags of up to 5 months in the case of SPI and SPEI. These
results suggest that current SPI and SPEI values could hold predictive value to assess future streamflow droughts. In contrast,
the rapid decline in coincidence rates at increasing lags over pluvial catchments makes it challenging to employ these indices
for projecting future streamflow droughts over these catchments. The relatively low values obtained by the ESSMI and SWEI
could be partially influenced by the ability of ERA5-Land in representing the soil moisture and *SWE* patterns. Future studies
510 could focus in analysing the relative changes in performance of these indices using adjusted soil moisture and *SWE* datasets.
We hope this information would help to improve current drought monitoring and shift towards more effective and proactive
management strategies.

<https://doi.org/10.5194/egusphere-2023-1911>

Preprint. Discussion started: 24 August 2023

© Author(s) 2023. CC BY 4.0 License.



Author contributions. OMBV and MZB conceptualised the research. OMBV, MZB, and DGM wrote the manuscript with help of all authors. OMBV designed the computational framework and analysed the data. DGM and JFS verified the analytical methods. DGM, HEB, JFS, CAG, 515 KV, RG, and MG contributed to the interpretation of the results. All authors provided critical feedback and helped shape the research, analysis and manuscript.

Competing interests. No competing interests

Acknowledgements. The authors thank the projects ANID-Sequía FSEQ210001, ANID Fondecyt Regular 1212071, ANID-PCI 190018, and ANID-Fondap 1522A0001.



520 References

Tech. rep.

- Alvarez-Garretón, C., Mendoza, P. A., Boisier, J. P., Addor, N., Galleguillos, M., Zambrano-Bigiarini, M., Lara, A., Puelma, C., Cortes, G., Garreaud, R., McPhee, J., and Ayala, A.: The CAMELS-CL dataset: catchment attributes and meteorology for large sample studies – Chile dataset, *Hydrology and Earth System Sciences*, 22, 5817–5846, <https://doi.org/10.5194/hess-22-5817-2018>, 2018.
- 525 Andreadis, K. M., Clark, E. A., Wood, A. W., Hamlet, A. F., and Lettenmaier, D. P.: Twentieth-century drought in the conterminous United States, *Journal of Hydrometeorology*, 6, 985–1001, 2005.
- Apurv, T., Sivapalan, M., and Cai, X.: Understanding the Role of Climate Characteristics in Drought Propagation, *Water Resources Research*, 53, 9304–9329, <https://doi.org/10.1002/2017WR021445>, 2017.
- Bachmair, S., Kohn, I., and Stahl, K.: Exploring the link between drought indicators and impacts, *Natural Hazards and Earth System Science*, 15, 1381–1397, <https://doi.org/10.5194/nhess-15-1381-2015>, 2015.
- 530 Bachmair, S., Stahl, K., Collins, K., Hannaford, J., Acreman, M., Svoboda, M., Knutson, C., Smith, K. H., Wall, N., Fuchs, B., Crossman, N. D., and Overton, I. C.: Drought indicators revisited: the need for a wider consideration of environment and society, *WIREs Water*, 3, 516–536, <https://doi.org/10.1002/wat2.1154>, 2016.
- Baez-Villanueva, O. M., Zambrano-Bigiarini, M., Beck, H. E., McNamara, I., Ribbe, L., Nauditt, A., Birkel, C., Verbist, K., Giraldo-Osorio, J. D., and Think, N. X.: RF-MEP: A novel Random Forest method for merging gridded precipitation products and ground-based measurements, *Remote Sensing of Environment*, 239, 111 606, 2020.
- 535 Baez-Villanueva, O. M., Zambrano-Bigiarini, M., Mendoza, P. A., McNamara, I., Beck, H. E., Thurner, J., Nauditt, A., Ribbe, L., and Think, N. X.: On the selection of precipitation products for the regionalisation of hydrological model parameters, *Hydrology and Earth System Sciences*, 25, 5805–5837, 2021.
- 540 Barker, L. J., Hannaford, J., Chiverton, A., and Svensson, C.: From meteorological to hydrological drought using standardised indicators, *Hydrology and Earth System Sciences*, 20, 2483–2505, 2016a.
- Barker, L. J., Hannaford, J., Chiverton, A., and Svensson, C.: From meteorological to hydrological drought using standardised indicators, *Hydrology and Earth System Sciences*, 20, 2483–2505, 2016b.
- Beck, H. E., Zimmermann, N. E., McVicar, T. R., Vergopolan, N., Berg, A., and Wood, E. F.: Present and future Köppen-Geiger climate classification maps at 1-km resolution, *Scientific data*, 5, 180 214, 2018.
- 545 Beguería, S., Vicente-Serrano, S. M., Reig, F., and Latorre, B.: Standardized precipitation evapotranspiration index (SPEI) revisited: parameter fitting, evapotranspiration models, tools, datasets and drought monitoring, *International journal of climatology*, 34, 3001–3023, 2014.
- Bevacqua, A. G., Chaffe, P. L., Chagas, V. B., and AghaKouchak, A.: Spatial and temporal patterns of propagation from meteorological to hydrological droughts in Brazil, *Journal of Hydrology*, 603, 126 902, 2021.
- 550 Bhardwaj, K., Shah, D., Aadhar, S., and Mishra, V.: Propagation of meteorological to hydrological droughts in India, *Journal of Geophysical Research: Atmospheres*, 125, e2020JD033 455, 2020.
- Blauhut, V., Gudmundsson, L., and Stahl, K.: Towards pan-European drought risk maps: quantifying the link between drought indices and reported drought impacts, *Environmental Research Letters*, 10, 014 008, <https://doi.org/10.1088/1748-9326/10/1/014008>, 2015.
- 555 Bloomfield, J. P. and Marchant, B. P.: Analysis of groundwater drought building on the standardised precipitation index approach, *Hydrology and Earth System Sciences*, 17, 4769–4787, <https://doi.org/10.5194/hess-17-4769-2013>, 2013.



- Boisier, J. P., Alvarez-Garretón, C., Cepeda, J., Osses, A., Vásquez, N., and Rondanelli, R.: CR2MET: A high-resolution precipitation and temperature dataset for hydroclimatic research in Chile, in: EGU General Assembly Conference Abstracts, p. 19739, 2018.
- Byun, H.-R. and Wilhite, D. A.: Objective quantification of drought severity and duration, *Journal of climate*, 12, 2747–2756, 1999.
- 560 Cammalleri, C., Arias-Muñoz, C., Barbosa, P., de Jager, A., Magni, D., Masante, D., Mazzeschi, M., McCormick, N., Naumann, G., Spinoni, J., et al.: A revision of the Combined Drought Indicator (CDI) used in the European Drought Observatory (EDO), *Natural Hazards and Earth System Sciences*, 21, 481–495, 2021.
- Carrão, H., Russo, S., Sepulcre-Canto, G., and Barbosa, P.: An empirical standardized soil moisture index for agricultural drought assessment from remotely sensed data, *International journal of applied earth observation and geoinformation*, 48, 74–84, 2016.
- 565 Cobon, D., Gacenga, F., An-Vo, D.-A., Pudmenzky, C., Nguyen-Huy, T., Stone, R., Guillory, L., Jackson, J., Svoboda, M., Swigart, J., et al.: Australian Drought Monitor, <https://doi.org/https://doi.org/10.26192/dmek-v625>, n.d.
- Donges, J. F., Schleussner, C.-F., Siegmund, J. F., and Donner, R. V.: Event coincidence analysis for quantifying statistical interrelationships between event time series, *The European Physical Journal Special Topics*, 225, 471–487, 2016.
- Farahmand, A. and AghaKouchak, A.: A generalized framework for deriving nonparametric standardized drought indicators, *Advances in*
570 *Water Resources*, 76, 140–145, 2015.
- Fluixá-Sanmartín, J., Pan, D., Fischer, L., Orlowsky, B., García-Hernández, J., Jordan, F., Haemmig, C., Zhang, F., and Xu, J.: Searching for the optimal drought index and timescale combination to detect drought: a case study from the lower Jinsha River basin, China, *Hydrology and Earth System Sciences*, 22, 889–910, <https://doi.org/10.5194/hess-22-889-2018>, 2018.
- Garreaud, R. D., Alvarez-Garretón, C., Barichivich, J., Pablo Boisier, J., Christie, D., Galleguillos, M., LeQuesne, C., McPhee, J., and
575 Zambrano-Bigiarini, M.: The 2010–2015 megadrought in central Chile: impacts on regional hydroclimate and vegetation, 2017.
- Garreaud, R. D., Boisier, J. P., Rondanelli, R., Montecinos, A., Sepúlveda, H. H., and Veloso-Aguila, D.: The Central Chile Mega Drought (2010–2018): A climate dynamics perspective, *International Journal of Climatology*, 40, 421–439, 2020.
- Gevaert, A. I., Veldkamp, T. I. E., and Ward, P. J.: The effect of climate type on timescales of drought propagation in an ensemble of global hydrological models, *Hydrology and Earth System Sciences*, 22, 4649–4665, <https://doi.org/10.5194/hess-22-4649-2018>, 2018.
- 580 Gibbs, W. J. and Maher, J. V.: Rainfall deciles as drought indicators, *Meteorology Bull.* 48, 48, Bureau of Meteorology. Commonwealth of Australia, Melbourne, Australia, 1967.
- González, M. E., Gómez-González, S., Lara, A., Garreaud, R., and Díaz-Hormazábal, I.: The 2010–2015 Megadrought and its influence on the fire regime in central and south-central Chile, *Ecosphere*, 9, e02300, 2018.
- Gottlieb, A. R. and Mankin, J. S.: Observing, measuring, and assessing the consequences of snow drought, *Bulletin of the American Meteorological Society*, 2021.
585
- Gringorten, I. I.: A plotting rule for extreme probability paper, *Journal of Geophysical Research*, 68, 813–814, 1963.
- Haile, G. G., Tang, Q., Li, W., Liu, X., and Zhang, X.: Drought: Progress in broadening its understanding, *Wiley Interdisciplinary Reviews: Water*, 7, e1407, <https://doi.org/https://doi.org/10.1002/wat2.1407>, 2020.
- Hao, Z., Yuan, X., Xia, Y., Hao, F., and Singh, V. P.: An overview of drought monitoring and prediction systems at regional and global scales,
590 *Bulletin of the American Meteorological Society*, 98, 1879–1896, 2017.
- Hargreaves, G. H. and Samani, Z. A.: Reference crop evapotranspiration from ambient air temperature, *American Society of Agricultural Engineers*, (fiche no. 85-2517), (Microfiche collection)(USA). no. fiche no. 85-2517., 1985.
- Haslinger, K., Koffler, D., Schöner, W., and Laaha, G.: Exploring the link between meteorological drought and streamflow: Effects of climate-catchment interaction, *Water Resources Research*, 50, 2468–2487, <https://doi.org/10.1002/2013WR015051>, 2014.



- 595 Hayes, M., Svoboda, M., Wall, N., and Widhalm, M.: The Lincoln Declaration on Drought Indices: Universal Meteorological Drought Index Recommended, *Bulletin of the American Meteorological Society*, 92, 485–488, <https://doi.org/10.1175/2010BAMS3103.1>, 2011.
- Hayes, M. J., Svoboda, M. D., Wardlow, B. D., Anderson, M. C., and Kogan, F.: Drought monitoring: Historical and current perspectives, in: *Remote Sensing of Drought: Innovative Monitoring Approaches*, edited by Wardlow, B. D., Anderson, M. C., and Verdin, J. P., CRC Press/Taylor & Francis, 2012.
- 600 He, X. and Sheffield, J.: Lagged compound occurrence of droughts and pluvials globally over the past seven decades, *Geophysical Research Letters*, 47, e2020GL087924, 2020.
- Heim Jr., R. R. and Brewer, M. J.: The global drought monitor portal: The foundation for a global drought information system, *Earth interactions*, 16, 1–28, <https://doi.org/10.1175/2012EI000446.1>, 2012.
- Hoffmann, D., Gallant, A. J. E., and Arblaster, J. M.: Uncertainties in Drought From Index and Data Selection, *Journal of Geophysical Research: Atmospheres*, 125, e31946, <https://doi.org/10.1029/2019JD031946>, 2020.
- 605 Hosking, J. R. M. and Wallis, J. R.: A Comparison of Unbiased and Plotting-Position Estimators of L Moments, *Water Resources Research*, 31, 2019–2025, 1995.
- Huang, S., Li, P., Huang, Q., Leng, G., Hou, B., and Ma, L.: The propagation from meteorological to hydrological drought and its potential influence factors, *Journal of Hydrology*, 547, 184–195, <https://doi.org/10.1016/j.jhydrol.2017.01.041>, 2017.
- 610 Huning, L. S. and AghaKouchak, A.: Global snow drought hot spots and characteristics, *Proceedings of the National Academy of Sciences*, 117, 19753–19759, 2020.
- Jain, V. K., Pandey, R. P., Jain, M. K., and Byun, H.-R.: Comparison of drought indices for appraisal of drought characteristics in the Ken River Basin, *Weather and Climate Extremes*, 8, 1, <https://doi.org/10.1016/j.wace.2015.05.002>, 2015.
- Jarvis, A., Reuter, H. I., Nelson, A., Guevara, E., et al.: Hole-filled SRTM for the globe Version 4, available from the CGIAR-CSI SRTM 90m Database (<http://srtm.csi.cgiar.org>), 15, 25–54, 2008.
- 615 Jehanzaib, M., Sattar, M. N., Lee, J.-H., and Kim, T.-W.: Investigating effect of climate change on drought propagation from meteorological to hydrological drought using multi-model ensemble projections, *Stochastic Environmental Research and Risk Assessment*, 34, 7–21, <https://doi.org/10.1007/s00477-019-01760-5>, 2020.
- JRC: Product Fact Sheet: SPI - Europe, Tech. rep., European Commission, Joint Research Centre, DESERT Action, LMNH Unit, available on http://edo.jrc.ec.europa.eu/documents/factsheets/factsheet_spi.pdf. [Last visited: October 2016], 2011.
- 620 Karl, T. R.: The Sensitivity of the Palmer Drought Severity Index and Palmer’s Z-Index to their Calibration Coefficients Including Potential Evapotranspiration, *Journal of Applied Meteorology*, 25, 77–86, [https://doi.org/10.1175/1520-0450\(1986\)025<0077:TSOTPD>2.0.CO;2](https://doi.org/10.1175/1520-0450(1986)025<0077:TSOTPD>2.0.CO;2), 1986.
- Keyantash, J. and Dracup, J. A.: The quantification of drought: an evaluation of drought indices, *Bulletin of the American Meteorological Society*, 83, 1167–1180, 2002.
- 625 Leeper, R. D., Bilotta, R., Petersen, B., Stiles, C. J., Heim, R., Fuchs, B., Prat, O. P., Palecki, M., and Ansari, S.: Characterizing US drought over the past 20 years using the US Drought Monitor, *International Journal of Climatology*, 42, 6616–6630, <https://doi.org/10.1002/joc.7653>, 2022.
- Li, Q., He, P., He, Y., Han, X., Zeng, T., Lu, G., and Wang, H.: Investigation to the relation between meteorological drought and hydrological drought in the upper Shaying River Basin using wavelet analysis, *Atmospheric Research*, 234, 104743, 2020.
- 630 Liu, C., Yang, C., Yang, Q., and Wang, J.: Spatiotemporal drought analysis by the standardized precipitation index (SPI) and standardized precipitation evapotranspiration index (SPEI) in Sichuan Province, China, *Scientific Reports*, 11, 1–14, 2021.



- Liu, Y., Zhu, Y., Ren, L., Singh, V. P., Yong, B., Jiang, S., Yuan, F., and Yang, X.: Understanding the spatiotemporal links between meteorological and hydrological droughts from a three-dimensional perspective, *Journal of Geophysical Research: Atmospheres*, 124, 3090–3109, 635 2019.
- Lloyd-Hughes, B. and Saunders, M. A.: A drought climatology for Europe, *International Journal of Climatology: A Journal of the Royal Meteorological Society*, 22, 1571–1592, 2002.
- Lorenzo-Lacruz, J., Vicente-Serrano, S. M., González-Hidalgo, J. C., López-Moreno, J. I., and Cortesi, N.: Hydrological drought response to meteorological drought in the Iberian Peninsula, *Climate Research*, 58, 117–131, 2013.
- 640 McKee, T. B., Doesken, N. J., Kleist, J., et al.: The relationship of drought frequency and duration to time scales, in: *Proceedings of the 8th Conference on Applied Climatology*, vol. 17, pp. 179–183, California, 1993.
- Meroni, M., Rembold, F., Fasbender, D., and Vrieling, A.: Evaluation of the Standardized Precipitation Index as an early predictor of seasonal vegetation production anomalies in the Sahel, *Remote Sensing Letters*, 8, 301–310, <https://doi.org/10.1080/2150704X.2016.1264020>, 2017.
- 645 Montecinos, A. and Aceituno, P.: Seasonality of the ENSO-related rainfall variability in central Chile and associated circulation anomalies, *Journal of climate*, 16, 281–296, 2003.
- Muñoz-Sabater, J., Dutra, E., Agustí-Panareda, A., Albergel, C., Arduini, G., Balsamo, G., Boussetta, S., Choulga, M., Harrigan, S., Hersbach, H., et al.: ERA5-Land: A state-of-the-art global reanalysis dataset for land applications, *Earth System Science Data*, 13, 4349–4383, 2021.
- 650 Narasimhan, B. and Srinivasan, R.: Development and Evaluation of Soil Moisture Deficit Index (SMDI) and Evapotranspiration Deficit Index (ETDI) for Agricultural Drought Monitoring, *Agricultural and Forest Meteorology*, 133, 69–88, <https://doi.org/10.1016/j.agrformet.2005.07.012>, 2005.
- Niemeyer, S. et al.: New drought indices, *Options Méditerranéennes. Série A: Séminaires Méditerranéens*, 80, 267–274, 2008.
- Notarnicola, C.: Overall negative trends for snow cover extent and duration in global mountain regions over 1982–2020, *Scientific Reports*, 655 12, 13 731, 2022.
- Palmer, W.: Meteorological drought, US Weather Bureau. Research Paper 45, 58 pp, 1965a.
- Palmer, W. C.: Meteorological drought, vol. 30, US Department of Commerce, Weather Bureau, 1965b.
- Pechlivanidis, I., Crochemore, L., Rosberg, J., and Bosshard, T.: What are the key drivers controlling the quality of seasonal streamflow forecasts?, *Water Resources Research*, 56, e2019WR026 987, 2020.
- 660 Peña-Gallardo, M., Vicente-Serrano, S. M., Hannaford, J., Lorenzo-Lacruz, J., Svoboda, M., Domínguez-Castro, F., Maneta, M., Tomas-Burguera, M., and El Kenawy, A.: Complex influences of meteorological drought time-scales on hydrological droughts in natural basins of the contiguous United States, *Journal of Hydrology*, 568, 611–625, 2019.
- Rahiz, M. and New, M.: Does a rainfall-based drought index simulate hydrological droughts?, *International Journal of Climatology*, p. n/a, <https://doi.org/10.1002/joc.3879>, 2013.
- 665 Robertson, A. W., Baethgen, W., Block, P., Lall, U., Sankarasubramanian, A., de Souza Filho, F. d. A., and Verbist, K. M.: Climate risk management for water in semi-arid regions, *Earth Perspectives*, 1, 12, 2014.
- Saha, T. R., Shrestha, P. K., Rakovec, O., Thober, S., and Samaniego, L.: A drought monitoring tool for South Asia, *Environmental Research Letters*, 16, 054 014, 2021.
- Shafer, B. A. and Dezman, L. E.: Development of a Surface Water Supply Index (SWSI) to assess the severity of drought conditions in 670 snowpack runoff areas, in: *Proceedings of the Western Snow Conference*, vol. 50, pp. 164–175, 1982.



- Sheffield, J., Andreadis, K., Wood, E. F., and Lettenmaier, D.: Global and continental drought in the second half of the twentieth century: Severity–area–duration analysis and temporal variability of large-scale events, *Journal of Climate*, 22, 1962–1981, 2009.
- Shukla, S. and Wood, A. W.: Use of a standardized runoff index for characterizing hydrologic drought, *Geophysical research letters*, 35, 2008.
- 675 Siegmund, J. F., Sanders, T. G., Heinrich, I., Van der Maaten, E., Simard, S., Helle, G., and Donner, R. V.: Meteorological drivers of extremes in daily stem radius variations of beech, oak, and pine in northeastern Germany: an event coincidence analysis, *Frontiers in plant science*, 7, 733, 2016.
- Siegmund, J. F., Siegmund, N., and Donner, R. V.: CoinCalc—A new R package for quantifying simultaneities of event series, *Computers & Geosciences*, 98, 64–72, 2017.
- 680 Silverman, B. W.: *Density estimation for statistics and data analysis*, Chapman Hall, 1986.
- Sims, A. P., Niyogi, D. d. S., and Raman, S.: Adopting drought indices for estimating soil moisture: A North Carolina case study, *Geophysical Research Letters*, 29, 24–1, <https://doi.org/10.1029/2001GL013343>, 2002.
- Sivakumar, M. V., Wilhite, D. A., Svoboda, M. D., Hayes, M., and Motha, R.: Drought risk and meteorological droughts, background paper prepared for the, 2011.
- 685 Slette, I. J., Post, A. K., Awad, M., Even, T., Punzalan, A., Williams, S., Smith, M. D., and Knapp, A. K.: How ecologists define drought, and why we should do better, *Global Change Biology*, 25, 3193–3200, <https://doi.org/10.1111/gcb.14747>, 2019.
- Stagge, J. H., Kohn, I., Tallaksen, L. M., and Stahl, K.: Modeling drought impact occurrence based on meteorological drought indices in Europe, *Journal of Hydrology*, 530, 37–50, <https://doi.org/10.1016/j.jhydrol.2015.09.039>, 2015a.
- Stagge, J. H., Tallaksen, L. M., Gudmundsson, L., Van Loon, A. F., and Stahl, K.: Candidate Distributions for Climatological Drought Indices (SPI and SPEI), *International Journal of Climatology*, 35, 4027–4040, <https://doi.org/10.1002/joc.4267>, 2015b.
- 690 Stahl, K., Vidal, J.-P., Hannaford, J., Tjeldeman, E., Laaha, G., Gauster, T., and Tallaksen, L. M.: The challenges of hydrological drought definition, quantification and communication: an interdisciplinary perspective, *Proceedings of the International Association of Hydrological Sciences*, 383, 291–295, 2020.
- Staudinger, M., Stahl, K., and Seibert, J.: A drought index accounting for snow, *Water Resources Research*, 50, 7861–7872, <https://doi.org/10.1002/2013WR015143>, 2014.
- 695 Steinemann, A.: Drought indicators and triggers: a stochastic approach to evaluation, *Journal of the American Water Resources Association (JAWRA)*, 39, 1217–1233, <https://doi.org/10.1111/j.1752-1688.2003.tb03704.x>, 2003.
- Steinemann, A.: Drought information for improving preparedness in the western states, *Bulletin of the American Meteorological Society*, 95, 843–847, <https://doi.org/10.1175/BAMS-D-13-00067.1>, 2014.
- 700 Steinemann, A., Iacobellis, S. F., and Cayan, D. R.: Developing and Evaluating Drought Indicators for Decision-Making, *Journal of Hydrometeorology*, 16, 1793–1803, <https://doi.org/10.1175/JHM-D-14-0234.1>, 2015.
- Steinemann, A. C. and Cavalcanti, L. F. N.: Developing Multiple Indicators and Triggers for Drought Plans, *Journal of Water Resources Planning and Management*, 132, 164–174, [https://doi.org/10.1061/\(ASCE\)0733-9496\(2006\)132:3\(164\)](https://doi.org/10.1061/(ASCE)0733-9496(2006)132:3(164)), 2006.
- Svoboda, M., LeComte, D., Hayes, M., Heim, R., Gleason, K., Angel, J., Rippey, B., Tinker, R., Palecki, M., Stooksbury, D., et al.: The drought monitor, *Bulletin of the American Meteorological Society*, 83, 1181–1190, 2002.
- 705 Tjeldeman, E., Barker, L., Svoboda, M., and Stahl, K.: Natural and human influences on the link between meteorological and hydrological drought indices for a large set of catchments in the contiguous United States, *Water Resources Research*, 54, 6005–6023, 2018.



- Torelló-Sentelles, H. and Franzke, C. L. E.: Drought impact links to meteorological drought indicators and predictability in Spain, *Hydrology and Earth System Sciences*, 26, 1821–1844, <https://doi.org/10.5194/hess-26-1821-2022>, 2022.
- 710 Van Loon, A. and Laaha, G.: Hydrological drought severity explained by climate and catchment characteristics, *Journal of hydrology*, 526, 3–14, 2015.
- Van Loon, A. F.: Hydrological drought explained, *Wiley Interdisciplinary Reviews: Water*, 2, 359–392, <https://doi.org/10.1002/wat2.1085>, 2015.
- Van Loon, A. F., Kumar, R., and Mishra, V.: Testing the use of standardised indices and GRACE satellite data to estimate the European 2015
715 groundwater drought in near-real time, *Hydrology and Earth System Sciences*, 21, 1947–1971, <https://doi.org/10.5194/hess-21-1947-2017>, 2017.
- van Rooy, M.: A Rainfall Anomaly Index independent of time and space, *Notos*, 14, 43–48, 1965.
- Vásquez, N., Cepeda, J., Gómez, T., Mendoza, P. A., Lagos, M., Boisier, J. P., Álvarez-Garretón, C., and Vargas, X.: Catchment-Scale Natural Water Balance in Chile, in: *Water Resources of Chile*, pp. 189–208, Springer, 2021.
- 720 Verbist, K., Robertson, A. W., Cornelis, W. M., and Gabriels, D.: Seasonal predictability of daily rainfall characteristics in central northern Chile for dry-land management, *Journal of Applied Meteorology and Climatology*, 49, 1938–1955, 2010.
- Vicente-Serrano, S. M. and López-Moreno, J. I.: Hydrological response to different time scales of climatological drought: an evaluation of the Standardized Precipitation Index in a mountainous Mediterranean basin, *Hydrology and Earth System Sciences*, 9, 523–533, <https://doi.org/10.5194/hess-9-523-2005>, 2005.
- 725 Vicente-Serrano, S. M., Beguería, S., and López-Moreno, J. I.: A multiscalar drought index sensitive to global warming: the standardized precipitation evapotranspiration index, *Journal of climate*, 23, 1696–1718, 2010.
- Vicente Serrano, S. M., López-Moreno, J. I., Beguería, S., Lorenzo-Lacruz, J., Azorin-Molina, C., and Morán-Tejeda, E.: Accurate computation of a streamflow drought index, *American Society of Civil Engineers*, 2012.
- Vicente-Serrano, S. M., Lopez-Moreno, J.-I., Beguería, S., Lorenzo-Lacruz, J., Sanchez-Lorenzo, A., García-Ruiz, J. M., Azorin-Molina, C.,
730 Morán-Tejeda, E., Revuelto, J., Trigo, R., et al.: Evidence of increasing drought severity caused by temperature rise in southern Europe, *Environmental Research Letters*, 9, 044 001, 2014.
- Vogt, J.: The European Drought Observatory, in: 2011 GEOSS Workshop XL-Managing Drought through Earth Observation, pp. 1–16, IEEE, <https://doi.org/10.1109/GEOSS.2011.5948940>, 2011.
- Vorobevskii, I., Kronenberg, R., and Bernhofer, C.: Linking different drought types in a small catchment from a statistical perspective –
735 Case study of the Wernersbach catchment, Germany, *Journal of Hydrology X*, 15, 100 122, <https://doi.org/10.1016/j.hydroa.2022.100122>, 2022.
- Wang, F., Wang, Z., Yang, H., Di, D., Zhao, Y., Liang, Q., and Hussain, Z.: Comprehensive evaluation of hydrological drought and its relationships with meteorological drought in the Yellow River basin, China, *Journal of Hydrology*, 584, 124 751, 2020.
- Wang, H., Rogers, J. C., and Munroe, D. K.: Commonly used drought indices as indicators of soil moisture in China, *Journal of Hydrometeorology*, 16, 1397–1408, <https://doi.org/10.1175/JHM-D-14-0076.1>, 2015.
- 740 WMO: Standardized Precipitation Index User Guide (M. Svoboda, M. Hayes and D. Wood), WMO-No. 1090, World Meteorological Organization, Geneva, Available online on http://www.wamis.org/agm/pubs/SPI/WMO_1090_EN.pdf. [Last accessed October-2016], 2012.
- Wu, H., Hayes, M. J., Weiss, A., and Hu, Q.: An evaluation of the Standardized Precipitation Index, the China-Z Index and the statistical Z-Score, *International Journal of Climatology: A Journal of the Royal Meteorological Society*, 21, 745–758, 2001.



- 745 Wu, J., Chen, X., Yao, H., Gao, L., Chen, Y., and Liu, M.: Non-linear relationship of hydrological drought responding to meteorological drought and impact of a large reservoir, *Journal of Hydrology*, 551, 495–507, 2017a.
- Wu, J., Chen, X., Yao, H., Gao, L., Chen, Y., and Liu, M.: Non-linear relationship of hydrological drought responding to meteorological drought and impact of a large reservoir, *Journal of Hydrology*, 551, 495–507, <https://doi.org/10.1016/j.jhydrol.2017.06.029>, 2017b.
- Zambrano, F., Wardlow, B., Tadesse, T., Lillo-Saavedra, M., and Lagos, O.: Evaluating satellite-derived long-term historical precipitation datasets for drought monitoring in Chile, *Atmospheric Research*, 186, 26–42, <https://doi.org/10.1016/j.atmosres.2016.11.006>, 2017.
- 750 Zambrano-Bigiarini, M. and Baez-Villanueva, O. M.: Characterizing meteorological droughts in data scarce regions using remote sensing estimates of precipitation, in: *Extreme Hydroclimatic Events and Multivariate Hazards in a Changing Environment*, pp. 221–246, Elsevier, 2019.
- Zargar, A., Sadiq, R., Naser, B., and Khan, F. I.: A review of drought indices, *Environmental Reviews*, 19, 333–349, 2011.
- 755 Zhang, F., Zhang, L.-w., Wang, X.-z., and Hung, J.-f.: Detecting Agro-Droughts in Southwest of China Using MODIS Satellite Data, *Journal of Integrative Agriculture*, 12, 159–168, [https://doi.org/10.1016/S2095-3119\(13\)60216-6](https://doi.org/10.1016/S2095-3119(13)60216-6), 2013.
- Zhao, H., Gao, G., An, W., Zou, X., Li, H., and Hou, M.: Timescale differences between SC-PDSI and SPEI for drought monitoring in China, *Physics and Chemistry of the Earth*, 102, 48–58, <https://doi.org/10.1016/j.pce.2015.10.022>, 2017.
- Zhao, Y., Feng, D., Yu, L., Wang, X., Chen, Y., Bai, Y., Hernández, H. J., Galleguillos, M., Estades, C., Biging, G. S., Radke, J. D., and Gong, P.: Detailed dynamic land cover mapping of Chile: Accuracy improvement by integrating multi-temporal data, *Remote Sensing of Environment*, 183, 170–185, <https://doi.org/10.1016/j.rse.2016.05.016>, 2016.
- 760 Zhu, Y., Wang, W., Singh, V. P., and Liu, Y.: Combined use of meteorological drought indices at multi-time scales for improving hydrological drought detection, *Science of the Total Environment*, 571, 1058–1068, 2016.
- Zink, M., Samaniego, L., Kumar, R., Thober, S., Mai, J., Schäfer, D., and Marx, A.: The German drought monitor, *Environmental Research Letters*, 11, 074002, 2016.