

1 **Seasonal characteristics of emission, distribution, and radiative effect of marine organic aerosols**  
2 **over the western Pacific Ocean: an investigation with a coupled regional climate-aerosol model**

3  
4 Jiawei Li<sup>1</sup>, Zhiwei Han<sup>1,2\*</sup>, Pingqing Fu<sup>3</sup>, Xiaohong Yao<sup>4</sup>, Mingjie Liang<sup>1,2</sup>

5  
6 <sup>1</sup>CAS Key Laboratory of Regional Climate-Environment for Temperate East Asia (RCE-TEA),  
7 Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing 100029, China

8 <sup>2</sup>University of Chinese Academy of Sciences, Beijing 100049, China

9 <sup>3</sup>School of Earth System Science, Tianjin University, Tianjin 300072, China

10 <sup>4</sup>Laboratory of Marine Environmental Science and Ecology, Ministry of Education, Ocean University  
11 of China, Qingdao 266100, China

12  
13  
14 Correspondence to: Zhiwei Han (hzw@mail.iap.ac.cn)

15  
16 **Abstract:** Organic aerosols from marine sources over the western Pacific Ocean of East Asia were  
17 investigated by using an online-coupled regional chemistry-climate model RIEMS-Chem for the  
18 entire year 2014. Model evaluation against a wide variety of observations from research cruises and  
19 in-situ measurements demonstrated a good skill of the model in simulating temporal variation and  
20 spatial distribution of particulate matter with aerodynamic diameter less than 2.5  $\mu\text{m}$  and 10  $\mu\text{m}$   
21 ( $\text{PM}_{2.5}$  and  $\text{PM}_{10}$ ), black carbon (BC), organic carbon (OC), sodium, and aerosol optical depth (AOD)  
22 in the marine atmosphere. The inclusion of marine organic aerosols improved model performance on  
23 OC concentration by reducing model biases of up to 20%. The regional and annual mean near surface  
24 marine organic aerosol (MOA) concentration was estimated to be 0.27  $\mu\text{g m}^{-3}$ , with the maximum in  
25 spring and the minimum in winter and contributed 26% of the total organic aerosol concentration on  
26 average over the western Pacific. Marine primary organic aerosol (MPOA) accounted for the majority  
27 of marine organic aerosol (MOA) mass and MPOA concentration exhibited the maximum in autumn  
28 and the minimum in summer, whereas marine secondary organic aerosol (MSOA) was approximately  
29 1~2 orders of magnitude lower than MPOA, having a distinct summer maximum and a winter  
30 minimum. MOA induced a direct radiative effect ( $\text{DRE}_{\text{MOA}}$ ) of  $-0.27 \text{ W m}^{-2}$ , and an indirect radiative

31 effect ( $IRE_{MOA}$ ) of  $-0.66 \text{ W m}^{-2}$  at TOA ( $IRE_{MOA}$ ) in terms of annual and oceanic average over the  
32 western Pacific, with the highest seasonal mean  $IRE_{MOA}$  up to  $-0.94 \text{ W m}^{-2}$  in spring.  $IRE_{MOA}$  was  
33 stronger than but in a similar magnitude to the IRE due to sea salt aerosol on average, and it was  
34 approximately 9% of the IRE due anthropogenic aerosols in terms of annual mean over the western  
35 Pacific, and this ratio increased to 19% in the northern parts of the western Pacific in autumn. This  
36 study reveals an important role of MOA in perturbing cloud properties and shortwave radiation fluxes  
37 in the western Pacific of East Asia.

38

## 39 **1 Introduction**

40 Atmospheric aerosol is one of the most important and uncertain factors in climate change issues  
41 (IPCC, 2013). Aerosols can alter radiation balance by scattering/absorbing solar/infrared radiation,  
42 and affect cloud microphysics and lifetime by activating as cloud condensation nuclei (CCN), exerting  
43 significant effects on climate system directly and indirectly. Aerosols are originated from  
44 anthropogenic and natural sources and of high spatial and temporal variability and short atmospheric  
45 lifetime relative to greenhouse gases. Consequently, aerosol radiative and climatic effects often have  
46 strong regional characteristics.

47 The western Pacific Ocean is frequently influenced by continental outflow of both anthropogenic  
48 and natural aerosols. Due to continuous growth of economy and energy consumption in the past  
49 decades, the aerosol level in China has been enhanced (Smith et al., 2011; Li M. et al., 2017) and may  
50 have potentially significant effects on radiation and cloud over not only the East Asian continent but  
51 also the wide downwind oceanic areas. Besides, East Asia is one of the major dust source regions on  
52 earth (Shao and Dong, 2006). Dust storms often occur in spring and dust particles can be transported  
53 eastward from the deserts and Gobi areas of north China and southern Mongolia to the western Pacific  
54 Ocean (Gong et al., 2003), providing nutrients (e.g., iron) for phytoplankton or even triggering the  
55 outbreak of algae bloom in oceans (Calil et al., 2011; Tan et al., 2017). In addition to anthropogenic  
56 and dust aerosols, marine aerosols also significantly affect aerosol chemical composition, radiation  
57 transfer, and cloud properties in marine atmosphere. The behaviors and climatic impacts of sea salt  
58 and non-sea-salt sulfate oxidized from dimethylsulphide (DMS) have been extensively investigated  
59 (Graf et al., 1997; Liao et al., 2004; Rap et al., 2013). In recent years, particular attentions have been  
60 paid on the sources and impacts of marine organic aerosols (O'Dowd et al., 2004; Meskhidze and

61 Nenes, 2006; Luo and Yu, 2010; Vignati et al., 2010; Gantt et al., 2011; Burrow et al., 2014; Quinn et  
62 al., 2017; Bertram et al., 2018; Huang et al., 2018), however, such studies were still very limited,  
63 especially for the western Pacific.

64 O'Dowd et al. (2004) found that organic matter dominated the chemical composition of marine  
65 aerosol during plankton bloom periods from spring to autumn over the North Atlantic Ocean,  
66 contributing 63% to sub-micron aerosol mass. Meshkidze and Nenes (2006) revealed a significant  
67 impact of phytoplankton bloom on cloud droplet number concentration and radiation balance in the  
68 Southern Ocean and proposed a major contribution of secondary organic aerosol (SOA) from  
69 phytoplankton produced isoprene. Some studies indicated that primary marine sources may dominate  
70 marine organic matter, whereas SOA oxidized from marine isoprene could only comprise a small  
71 fraction of the observed organic aerosol mass over marine environment (Facchini et al. 2008; Arnold  
72 et al., 2009; Myriokefalitakis et al., 2010). The estimated global emission amounts of primary marine  
73 organic matter varied largely among models. Using the global aerosol-climate model ECHAM5-HAM,  
74 Roelofs (2008) estimated a global production of marine organic aerosols to be 75 TgC yr<sup>-1</sup>. Spracklen  
75 et al. (2008) estimated the marine organic carbon emission to be approximately 8 TgC yr<sup>-1</sup> based on  
76 measured organic carbon mass and satellite retrieved chlorophyll-a (Chl-a) concentration. Vignati et al.  
77 (2010) derived a global emission of marine primary organic matter in the sub-micron size by sea spray  
78 process to be 5.8 TgC yr<sup>-1</sup> by using an off-line global Chemistry-Transport Model TM5 with a  
79 parameterization relating organic emission fraction to sea surface Chl-a concentration. Gantt et al.  
80 (2011) found that the combination of 10 m wind speed and sea surface Chl-a concentration were the  
81 most consistent predictors of organic mass fraction of sea spray aerosol based on observations from  
82 the Mace Head atmospheric research station on the Atlantic coast of Ireland and a site at the Point  
83 Reyes National Seashore on the Pacific coast of California. They developed a new MPOA emission  
84 function and estimated the global annual MPOA emission associated with sea spray to be from 15.9  
85 TgC yr<sup>-1</sup> to 18.7 TgC yr<sup>-1</sup> (2.8~5.6 TgC yr<sup>-1</sup> in the sub-micron size). However, Quinn et al. (2014)  
86 found that the organic carbon content of sea spray aerosol is weakly correlated with satellite retrieved  
87 chlorophyll-a concentration based on cruise measurements in the North Atlantic Ocean and the coastal  
88 waters of California. Bates et al (2020) reported that plankton bloom has little effect on the emission  
89 flux, organic fraction or cloud condensation nuclei of sea spray aerosol based on cruise experiment  
90 over the North Atlantic. Burrows et al. (2014) developed a novel physically based framework for

91 parameterizing the organic fractionation of sea spray aerosol by consideration of ocean  
92 biogeochemistry processes, and their predicted relationships between Chl-a and organic fraction are  
93 similar to existing empirical parameterizations associated with ocean Chl-a concentrations at high  
94 Chl-a levels, but the empirical relationships may not be adequate to predict OM fraction of sea spray  
95 aerosol outside of strong seasonal blooms.

96 Regarding the influence on climatic factors, such as cloud condensation nuclei (CCN),  
97 Ovadnevaite et al. (2011) revealed that MPOA was a dichotomy of low hygroscopicity and high CCN  
98 activity through analysis of ambient measurements of aerosol chemical compositions and size  
99 distributions at the Mace Head atmospheric research station, and highlighted the importance of MPOA  
100 in CCN activation over marine atmosphere. A later study of Westervelt et al. (2012) indicated that  
101 marine organic aerosols was able to increase CCN by up to 50% in the Southern Ocean and by 3.7%  
102 globally during the austral summer based on the model simulation of GISS GCM II'. Based on the  
103 measurements from seven research cruises over the Pacific, Southern, Arctic, and Atlantic oceans  
104 between 1993 and 2015, Quinn et al. (2017) indicated that sea spray aerosol generally makes a  
105 contribution of less than 30% to CCN population at supersaturation of 0.1 to 1.0% on a global basis.  
106 Burrows et al. (2022) pointed out that sea spray organic aerosol strengthened shortwave radiative  
107 cooling by clouds by  $-0.36 \text{ W m}^{-2}$  in the global annual mean, with the zonal mean contribution  
108 exceeding  $-3.5 \text{ W m}^{-2}$  in the Southern Ocean in summertime.

109 The above studies reveal the important role of marine organic aerosols in chemical composition,  
110 radiation budget, and cloud microphysics with focus on the global scale. However, there is very  
111 limited modeling research on this important and challenging issue for the western Pacific Ocean of  
112 East Asia. To our knowledge, only two of our previous studies explored the effects of MPOA on  
113 chemical composition, radiation, cloud and precipitation over the western Pacific in springtime with  
114 an online-coupled regional chemistry/aerosol-climate model RIEMS-Chem (Han et al., 2019; Li et al.,  
115 2019), whereas the seasonality and annual aspect of MPOA and MSOA produced by marine isoprene  
116 and terpene are still unknown. In this study, we conducted a one-year simulation with the developed  
117 RIEMS-Chem to further explore the characteristics and radiative impacts of marine organic aerosols  
118 over the western Pacific. The model simulated aerosol compositions were validated against a wide  
119 series of observations from ground and cruise measurements, and the simulated MSOA was evaluated  
120 by comparison with cruise measured secondary organic tracer in marine air masses. To our knowledge,

121 for the first time, the seasonality of emissions, concentrations, direct and indirect radiative effects of  
122 marine organic aerosols was characterized and the annual means were estimated specifically for the  
123 western Pacific and for the key oceanic regions of concern over East Asia. This study would provide  
124 new insights into properties and impacts of marine organic aerosols over the western Pacific and  
125 would be a necessary supplement to the global perspective of marine organic aerosols.

126

## 127 2 Model and data

### 128 2.1 Model description and key processes

129 An online-coupled regional atmospheric chemistry/aerosol-climate model RIEMS-Chem was  
130 used to investigate marine organic aerosols in this study. RIEMS-Chem composes of the host regional  
131 climate model RIEMS (Fu et al., 2005; Xiong et al., 2009; Wang S.Y. et al., 2015) and a  
132 comprehensive atmospheric chemistry/aerosol module. RIEMS was developed based on the dynamic  
133 structure of the fifth-generation Pennsylvania State University NCAR Mesoscale Model (MM5; Grell  
134 et al., 1995) with a series of parameterizations to represent major physical processes, such as a  
135 modified Biosphere-Atmosphere Transfer Scheme (BATS; Dickinson et al., 1993) for land-surface  
136 process, the Medium-Range Forecasts scheme (MRF; Hong and Pan, 1996) for planetary boundary  
137 layer process, the Grell cumulus convective parameterization scheme (Grell, 1993) for convective  
138 process, the Reisner explicit moisture scheme (Reisner et al., 1998) and a modified radiation package  
139 of the NCAR Community Climate Model (CCM3; Kiehl et al., 1996) for radiation transfer processes  
140 with aerosol effect. RIEMS has participated in the Regional Climate Model Intercomparison Project  
141 (RMIP) for Asia and it was one of the best models in predicting surface air temperature and  
142 precipitation over East Asia (Fu et al., 2005).

143 Atmospheric chemistry/aerosol modules have been incorporated into RIEMS in recent years,  
144 establishing the online-coupled model RIEMS-Chem, which can account for the interactions among  
145 chemistry, radiation, cloud, and meteorology (Han, 2010; Han et al., 2012). RIEMS-Chem has been  
146 successfully applied in previous modeling studies on anthropogenic aerosols, mineral dust and marine  
147 aerosols regarding spatial-temporal distributions, physical and chemical evolutions, and radiative and  
148 climatic effects over East Asia (Han et al., 2012; 2013; 2019; Li et al., 2014; 2016a; 2016b; 2019;  
149 2020). It is now participating in the international model comparison project MICS-Asia III (Model  
150 Inter Comparison Study for Asia phase III) and shows a good ability in predicting aerosol

151 concentrations and AOD over East Asia (Gao et al., 2018).

152 RIEMS-Chem includes atmospheric chemistry and aerosol processes, such as gas and aqueous  
153 phase chemistries which are represented by the CB-IV mechanism (Gery et al. 1989) and RADM  
154 scheme (Chang et al., 1987), respectively; Sulfate is mainly produced from the oxidation of SO<sub>2</sub> by  
155 OH radical in gas phase and the oxidation of dissolved SO<sub>2</sub> by H<sub>2</sub>O<sub>2</sub>, O<sub>3</sub>, and metal catalysis in  
156 aqueous phase (Chang et al., 1987). Nitrate and ammonium are produced through thermodynamic  
157 processes represented by the ISORROPIA II model (Fountoukis and Nenes, 2007). BC, POA, and  
158 anthropogenic primary PMs are considered chemically inert. SOA formation from anthropogenic and  
159 biogenic VOC precursors is treated by a bulk yield scheme from Lack et al. (2004), with SOA yield of  
160 424 μg m<sup>-3</sup> ppm<sup>-1</sup> for toluene, 342 μg m<sup>-3</sup> ppm<sup>-1</sup> for xylene, and 762 μg m<sup>-3</sup> ppm<sup>-1</sup> for monoterpene.  
161 For irreversible conversion of marine VOCs to SOA, a 28.6% mass yield is assumed for isoprene  
162 (Surratt et al., 2010, Meskhidze et al., 2011) and 30% for monoterpene (Lee et al., 2006).  
163 Heterogeneous reactions between gaseous precursors and aerosols are also taken into account (Li and  
164 Han, 2010; Li J. W. et al., 2018). Dry deposition velocity is represented by a size-dependent  
165 parameterization over different underlying surfaces (Han et al., 2004). Dry deposition velocity of  
166 particle is expressed as the inverse of the sum of resistant plus a gravitational settling term. Over sea  
167 or ocean surfaces, the quasi-laminar boundary layer (QBL) may be disrupted by bursting bubbles,  
168 resulting in an increase in downward movement of particles, which is parameterized by the approach  
169 of Van den Berg et al. (2000), in which quasi-laminar resistance  $r_b$  is determined by Brownian  
170 diffusion and impaction when QBL is intact, and by turbulence and washout velocity of particles by  
171 spray drops when QBL is broken down. Below-cloud scavenging (BCS) of particles between cloud  
172 base and ground surface represents capture processes of particle by falling hydrometeor through  
173 Brownian and turbulent shear diffusion, interception and inertial impaction, and is parameterized by a  
174 scavenging rate, which is a function of precipitation rate and collision efficiency of particle by  
175 hydrometeor (Slinn, 1984).

176 A physically based scheme (namely A-G scheme) developed based on classical Köhler theory by  
177 Abdul-Razzak and Ghan (1998, 2004) is incorporated into RIEMS-Chem to represent aerosol  
178 activation into cloud droplet processes. This scheme calculates cloud droplet number concentration  
179 ( $N_c$ ) with not only aerosol number concentration, but also aerosol size distribution and composition,  
180 updraft velocity and ambient supersaturation. Aerosols are activated if their critical supersaturation is

181 less than the maximum ambient supersaturation. The critical supersaturation for activating particles is  
182 determined by curvature effect and solute effect. The maximum ambient supersaturation is calculated  
183 by solving supersaturation balance equation (Abdul-Razzak and Ghan, 1998). The updraft velocity is  
184 represented by the sum of grid mean updraft velocity and sub-grid updraft velocity, which is  
185 diagnosed from vertical eddy diffusivity according to Ghan et al. (1997). The A-G scheme in  
186 RIEMS-Chem has been applied over the western Pacific Ocean in spring 2014 and its prediction for  
187 hourly CCN concentration at different supersaturations has been validated by cruise measurements  
188 from the marginal seas of China to remote oceans southeast of Japan, which demonstrates a good  
189 ability, with the correlation coefficient of 0.87 and normalized mean bias within 20%. More details on  
190 the treatment and evaluation of marine aerosol activation refer to Han et al. (2019). Once  $N_c$  is  
191 calculated by the A-G scheme, the cloud droplet effective radius  $r_e$  is calculated with the method of  
192 Martin et al. (1994). The activated aerosols (into cloud droplet) are removed from the air. The auto  
193 conversion rate from cloud water to rainwater is parameterized by the scheme of Beheng (1994),  
194 which depends on  $N_c$  diagnosed and cloud liquid water content. The effect of aerosols on ice nuclei  
195 and convective cloud is not treated in this model due to limited knowledge at present.

196

## 197 2.2 Aerosol physical and chemical properties

198 Ten aerosol types are simulated in RIEMS-Chem, which are sulfate ( $\text{SO}_4^{2-}$ ), nitrate ( $\text{NO}_3^-$ ),  
199 ammonium ( $\text{NH}_4^+$ ), black carbon (BC), primary organic aerosol (POA), secondary organic aerosol  
200 (SOA), anthropogenic primary PMs ( $\text{PM}_{2.5}$  and  $\text{PM}_{10}$ ), dust, and sea salt.

201 Based on the observational analysis of aerosol mixing state in eastern China (Ma et al., 2017; Wu  
202 et al., 2017), an internal mixing assumption is adopted for anthropogenic aerosols and they are  
203 externally mixed with natural aerosols. The geometric mean radius and standard deviation of the  
204 anthropogenic internal mixture are estimated to be 0.11  $\mu\text{m}$  and 1.65, respectively, based on field  
205 measurements (Ma et al., 2017). Mineral dust is represented by 5 size bins (0.1~1.0, 1.0~2.0, 2.0~4.0,  
206 4.0~8.0 and 8.0~20.0  $\mu\text{m}$ ), while sea salt is represented by 2 size bins, with the mean fine mode radius  
207 being 0.1 $\mu\text{m}$  and the coarse mode radius being 1  $\mu\text{m}$  according to measurements from Gong et al.  
208 (1997). Feng et al. (2017) indicated that the measured total organic carbon (TOC) in the western  
209 Pacific Ocean during the same period as this study was enriched in <0.3  $\mu\text{m}$  (volume median  
210 diameter), which was mainly contributed by MPOA and the super-micron TOC was generally below

211 the detection limit. Accordingly, the geometric mean diameter of marine organic aerosol number  
212 concentration was set to be 0.1  $\mu\text{m}$ , with a standard deviation of 1.6. The number concentration is  
213 calculated by mass concentration as the formula in Curci et al. (2015). MPOA can be mixed with sea  
214 salt both externally or internally. It is more likely to be externally mixed with sea salt for finer  
215 aerosols (<200 nm in diameter) (Gantt and Meskhidze, 2013) and the effect of externally mixed  
216 MPOA was found to be much more important than that of internally mixed MPOA (Gantt et al.,  
217 2012b). So an external mixture of MPOA and sea salt is assumed in this study, this means additional  
218 marine organic aerosols are produced to affect cloud properties and represents an upper limit of  
219 indirect effect. There is little information on physical and chemical properties of marine organic  
220 aerosols, some key parameters for calculation of aerosol activation, i.e. the number of ions the salt  
221 dissociates into water, the osmotic coefficient, the mass fraction of soluble material, the density, and  
222 molecular weight are set to be 3.0, 1, 0.1, 1.5  $\text{g m}^{-3}$ , and 90, respectively, according to a few previous  
223 studies (Abdul-Razzak and Ghan, 2004; Roelofs, 2008). The soluble mass fraction of MPOA is  
224 assumed to be 0.2, slightly higher than that of MPOA (Liu and Wang, 2010; Westervelt et al., 2012).  
225 An OM/OC ratio of 1.4 was applied to convert organic matter (OM) to OC (Gantt and Meskhidze,  
226 2013).

227 The hygroscopic growth of aerosol is parameterized by a  $\kappa$  parameterization (Petters and  
228 Kreidenweis, 2007). The hygroscopicity parameters ( $\kappa$ ) for inorganic aerosol components, BC, POA,  
229 SOA, dust, and sea salt are set to be 0.65, 0, 0.1, 0.2, 0.01 and 0.98, respectively (Riemer et al., 2010;  
230 Liu et al., 2010; Westervelt et al., 2012). The hygroscopicity for MPOA and MPOA are assumed to be  
231 the same as those for anthropogenic POA and SOA. Because there was limited information on the  
232 optical properties of marine organic aerosols, the refractive index of anthropogenic POA and SOA was  
233 used instead. The aerosol refractive index and hygroscopicity ( $\kappa$ ) of the internally mixed aerosol are  
234 calculated by volume-weighting of the parameters for each aerosol component. Aerosol optical  
235 parameters including extinction coefficient, single scattering albedo, and asymmetry factor are  
236 calculated by a Mie-theory based method developed by Ghan and Zaveri (2007), in which the aerosol  
237 optical parameters are pre-calculated by the Mie theory and then fitted by Chebyshev polynomials  
238 with a table of polynomial coefficients for looking up for aerosols with certain size and refractive  
239 index. More detailed description refers to Li et al. (2020). This approach is much faster than  
240 traditional Mie code with a similar level of accuracy and has been successfully used in estimating



241 aerosol optical properties over East Asia (Han et al. 2011).

242

### 243 2.3 Anthropogenic and natural emissions

244 Monthly mean anthropogenic emissions of sulfur dioxide (SO<sub>2</sub>), nitrogen (NO<sub>x</sub>), ammonia (NH<sub>3</sub>),  
245 non-methane volatile organic compounds (NMVOC), carbon monoxide (CO), BC, POA, and other  
246 anthropogenic primary PM<sub>2.5</sub> and PM<sub>10</sub> in China for the year 2014 are obtained from the MEIC  
247 inventory (Multi-resolution Emission Inventory for China) which was developed by Tsinghua  
248 University (<http://meicmodel.org>, last access: 2020/01/20). Anthropogenic emissions outside China  
249 are taken from the MIX inventory which was developed to support the Model Inter-Comparison Study  
250 for Asia phase III (MICS-Asia III) and the Hemispheric Transport of Air Pollution (HTAP) projects  
251 (Li M. et al., 2017). Both inventories of MEIC and MIX have the same resolution of 0.5 degree. Open  
252 biomass burning emissions of aerosols and gas precursors for the year 2014 with a spatial resolution  
253 of 0.5 degree are derived from the Global Fire Emissions Database, Version 4.0 (GFED4) on a daily  
254 basis (Giglio et al., 2013). The biogenic VOC emission is derived from the CAMS-BIO Global  
255 biogenic emissions dataset (CAMS-GLOB-BIO v3.1) (Granier et al., 2019; Sindelarova et al., 2014)  
256 distributed by ECCAD-GEIA (<https://permalink.aeris-data.fr/CAMS-GLOB-BIO>, last access:  
257 2020/02/10) and the monthly mean biogenic emission for the year 2014 with a horizontal resolution of  
258 0.25° is used. All the above emission data are bilinearly interpolated to the lambert projection of  
259 RIEMS-Chem. The deflation of mineral dust is represented by the scheme of Han et al. (2004), which  
260 is calculated online with RIEMS predicted meteorology.

261 The generation of sea salt aerosol through bubbles is represented by Gong (2003) which is  
262 developed for sea salt radius from 0.07μm to 20μm based on the scheme of Monahan et al. (1986) and  
263 it is modified by considering the influences of relative humidity (RH) (Zhang et al., 2005).

264 Considering the strong bloom seasonality in the western Pacific region and the availability of  
265 global satellite data for Chl-a concentration, and the lack of cruise measurements on the relationship  
266 between sea spray organic aerosol fluxes and Chl-a in this region, we adopted the scheme of Gantt et  
267 al (2011) for parameterizing marine primary organic aerosol emission in this study. The size-resolved  
268 marine primary organic aerosol (MPOA) emission is parameterized based on the method of Gantt et al.  
269 (2011; 2012a), in which the emission rate of MPOA is the product of sea spray emission rate and  
270 organic matter fraction of sea spray aerosol, which is expressed as a function of wind speed, surface

271 seawater Chl-a concentration, and aerosol size. The Chl-a data used in this study is the Level-3 daily  
272 mean Chl-a concentration ( $\text{mg m}^{-3}$ ) product with 9 km resolution retrieved from the VIIRS (Visible  
273 infrared Imaging Radiometer) sensor onboard the Suomi National Polar-orbiting Partnership (SNPP)  
274 satellite platform (OBPG, 2018) (<http://oceandata.sci.gsfc.nasa.gov>, last access: 2020/12/8). A brief  
275 description on this scheme with formulas is presented in the supplement.

276 Marine isoprene emission released by phytoplankton activities is parameterized using the scheme  
277 of Gantt et al. (2009) which considers light sensitivity of phytoplankton isoprene production and  
278 dynamic euphotic depth (see more details in the supplement). Marine emission of monoterpene is  
279 scaled by 0.2 to those of isoprene following the suggestion from Myriokefalitakis et al. (2010). The  
280 marine abiotic source of isoprene (due to photochemical production in the sea surface microlayer)  
281 may be important according to recent studies (Brüggemann et al., 2018; Conte et al., 2020). This  
282 mechanism is not considered in this study because the production mechanism for marine abiotic  
283 isoprene is poorly understood at present.

284

#### 285 2.4 Model setup and experiment design

286 This study focused on the western Pacific Ocean of East Asia. The model domain covered most  
287 areas of eastern China, the Korean Peninsula, Japan, parts of Southeast Asia, and a wide area of the  
288 western Pacific Ocean ( $11^{\circ}\text{N}\sim 54^{\circ}\text{N}$ ,  $89^{\circ}\text{E}\sim 173^{\circ}\text{E}$ ) (Figure 1). A Lambert conformal projection with 60  
289 km horizontal resolution was applied in the model. 16 vertical layers stretched unevenly from the  
290 surface to tropopause in a terrain-following sigma coordinate with the first 8 layers within planetary  
291 boundary layer. The time step of model integration is 180 seconds. The simulation period was from 1  
292 December 2013 to 31 December 2014 with the first month as model spin-up and the whole year of  
293 2014 was used for analysis. Year 2014 is a normal year (neither El Niño nor La Niña) which can be  
294 reflected by the ENSO index (<https://psl.noaa.gov/enso/mei/>, last access: 2023/12/15). The reason we  
295 choose 2014 as the study period is the availability of cruise campaigns (for model validation and  
296 analysis) over the East China Sea and the western Pacific from spring to early summer, when BC and  
297 OC concentrations were measured. Final reanalysis data with  $1^{\circ}\times 1^{\circ}$  resolution and 6-hour interval  
298 from the National Centers for Environmental Prediction (NOAA/NCEP, 2000) was used to provide  
299 initial and boundary conditions for meteorology. Chemical results derived from the MOZART-4  
300 (Model for Ozone and Related chemical Tracers, version 4; Emmons et al., 2010) simulation with

301 6-hour interval were used to provide lateral conditions for trace gases and aerosols. The full  
302 simulation (FULL) is designed by considering all anthropogenic and natural emissions, including  
303 marine emissions of primary organic aerosol and sea salt, and a series of model simulations are also  
304 conducted to estimate the direct and indirect radiative effect of MOA and their sensitivity to MOA  
305 properties, which are described in the following sections.

306

## 307 2.5 Observations

308 In-situ measurements of PM<sub>10</sub>, PM<sub>2.5</sub>, and gas precursors (O<sub>3</sub>, SO<sub>2</sub>, and NO<sub>x</sub>/NO<sub>2</sub>) at coastal and  
309 island sites in Japan and Republic of Korea were obtained from EANET (Acid Deposition Monitoring  
310 Network in East Asia, <http://www.eanet.asia>, last access: 2020/01/23) (Figure 1). Hourly  
311 concentrations of PM<sub>10</sub>, SO<sub>2</sub>, NO<sub>x</sub> in Japan, NO<sub>2</sub> in Korea, and O<sub>3</sub> were automatically monitored at  
312 six Japanese sites (Rishiri, Tappi, Sado-seki, Oki, Hedo, and Ogasawara) and three Korean sites (Jeju,  
313 Kanghwa, and Imsil), whereas hourly PM<sub>2.5</sub> concentrations were only available at three Japanese sites  
314 (Rishiri, Sado-seki, and Oki). Sodium (Na<sup>+</sup>) concentrations sampled on a bi-weekly basis at the 6  
315 coast/island EANET sites in Japan were also collected. Besides, hourly PM<sub>10</sub> and PM<sub>2.5</sub>  
316 concentrations monitored in the three coastal cities of China (Qingdao, Shanghai, and Fuzhou) were  
317 also obtained from the CNEMC (China National Environmental Monitoring Center,  
318 <http://www.cnemc.cn/>, last access: 2020/01/23) and used for model validation (Figure 1).

319 Carbonaceous aerosol (OC and BC) concentrations measured from two research cruise campaigns  
320 covering the western Pacific during the spring and summer of 2014 (Figure 1) were collected and used  
321 for model validation. The spring cruise campaign was carried out from 17 March to 22 April 2014  
322 onboard the research vessel R/V Dongfanghong II, which started from Qingdao, sailed to the western  
323 Pacific Ocean, and then returned (Figure 1) (Luo et al., 2016; Feng et al., 2017). OC and BC samples  
324 were collected by an 11-stage MOUDI (Models110-IITM) (0.054~18 μm) equipped with  
325 pre-combusted quartz filters onboard the vessel. Mass concentrations of total OC (primary and  
326 secondary) and BC were determined by the thermal/optical carbon analyzer (Sunset Laboratory Inc.,  
327 Forest Grove, OR). Totally 19 daily BC and OC samples were collected during the cruise. Detailed  
328 information about this campaign and the sampling and analysis techniques were documented in Feng  
329 et al (2017). The early summer campaign was carried out from 18 May to 12 June 2014 (Kang et al.,  
330 2018). Total suspended particles (TSP) were collected on pre-combusted quartz filters using a

331 high-volume air sampler (Kimoto, Japan) onboard the KEXUE-1 Research Vessel during a National  
332 Natural Science Foundation of China (NSFC) sharing cruise (Figure 1). This campaign covered low-  
333 to mid-latitudes of the western Pacific Ocean (over the Yellow Sea and the East China Sea). Totally 51  
334 half-day (daytime/nighttime) OC samples were obtained during this campaign. Detailed information  
335 about this campaign and samples were described in Kang et al. (2018).

336 Besides the in-situ observations and cruise campaigns mentioned above, long-term observations  
337 of OC and BC from previous publications were collected to help model comparison and analysis.  
338 Carbonaceous aerosol samples (OC and BC) in TSP were continuously collected on a weekly basis  
339 from 2001 to 2012 at Chichijima Island (the same place as Ogasawara in Figure 1), a remote island  
340 located in the western North Pacific. The monthly mean OC and BC concentrations of the 12-year  
341 average were reported by Boreddy et al. (2018) and used to verify the model performance over remote  
342 oceans. Measurements of seasonal mean OC and BC concentrations in TSP at Huaniao Island (a  
343 pristine island about 100 km southeast of Shanghai over the East China Sea, see Figure 1) from  
344 October 2011 to August 2012 (Wang F. W. et al., 2015) and at Okinawa island (the same place as  
345 Hedo in Figure 1) in the western Pacific Ocean from October 2009 to October 2010 (Kunwar and  
346 Kawamura, 2014) were collected and used in this study. BC observations were conducted at Fukue  
347 Island of western Japan using a continuous soot-monitoring system (COSMOS) (Figure 1) by Kanaya  
348 et al. (2016) from 2009 to 2015.

349 Ground observations of AOD were obtained from the Aerosol Robotic Network (AERONET,  
350 <https://aeronet.gsfc.nasa.gov/>, last access: 2020/06/03). Level 2 AOD observations for the year 2014  
351 were collected at 7 coastal sites shown in Figure 1. Hourly and monthly mean observations were  
352 derived from raw data and used for model comparison and statistics calculation. AOD at 550 nm was  
353 used to match the model output. The level-3, daily deep blue global AOD product (in  $1^{\circ}\times 1^{\circ}$  horizontal  
354 resolution and at 550nm) retrieved by VIIRS sensor onboard the SNPP satellite platform (Sayer et al.  
355 2018) were also collected to examine AOD spatial distribution.

356

### 357 3 Model validations

358 In this section, the model results for OC, BC, PM<sub>10</sub>, PM<sub>2.5</sub>, sodium concentrations, and AOD were  
359 compared with a variety of observations from research cruise and monitoring networks to help  
360 evaluate the model ability over wide areas from eastern China to the western Pacific Ocean. Because

361 the above comparison was for total OC mass concentration, we also compared the simulated SOA  
362 from marine sources to cruise measured SOA tracer to examine the model performance for marine  
363 organic aerosols. Model results are extracted from the model grid closest to the observational site for  
364 comparison with observations.

365

### 366 3.1 Particulate matters (PM<sub>10</sub> and PM<sub>2.5</sub>), sodium (Na<sup>+</sup>) and gas precursors

367 As particulate matter in remote marine atmosphere is mainly composed of sea salt, the model  
368 performance for PM<sub>10</sub> and PM<sub>2.5</sub> may partly reflect the model ability for sea salt simulation, which is  
369 crucial to the estimation of MPOA emission.

370 Because the focus of this study is seasonal variation, the hourly PM<sub>10</sub> and PM<sub>2.5</sub> observations and  
371 corresponding simulations were averaged to be monthly means and shown in Figure 2. In general,  
372 RIEMS-Chem performed quite well in simulating monthly variation of PM<sub>10</sub> concentrations at both  
373 the EANET sites (Figure 2a~2i) and CNEMC sites (Figure 2j~2l) for the year 2014, although model  
374 biases still occurred at some sites, such as the underprediction in winter and spring in Jeju (Figure 2g)  
375 and Imsil (Figure 2i) and the overprediction in May in Oki (Figure 2d) and Rishiri (Figure 2a). It was  
376 striking that PM<sub>10</sub> concentration peaked in May and was lowest in July-August at all Korean sites and  
377 Japanese sites over northeast Asia (Figure 2a~2i). The long-range transport of mineral dust from north  
378 China and Mongolia in spring could contribute to the PM<sub>10</sub> maximum in May. It was noteworthy that  
379 the model simulated seasonality and magnitude of PM<sub>10</sub> agreed quite well with observations at the  
380 four island sites of northern Japan (Rishiri, Tappi, Sado, and Oki) (Figure 2a~2d), where sea salt  
381 aerosol played a more important role than those sites in Korea, implying sea salt concentrations could  
382 also be well reproduced by the model. The PM<sub>10</sub> level at Ogasawara (Figure 2f) was much lower than  
383 those at the other sites and its seasonality was characterized by the minimum in summer (5  $\mu\text{g m}^{-3}$ )  
384 and the maximum in spring. The model reasonably reproduced the seasonality at Hedo (Figure 2e) and  
385 Ogasawara (Figure 2f) as well, although it generally predicted lower values at Hedo and higher values  
386 at Ogasawara. As for PM<sub>10</sub> concentrations at the CNEMC sites of eastern China, the model simulated  
387 PM<sub>10</sub> concentrations very well for Shanghai (Figure 2k) and Fuzhou (Figure 2l) in terms of both  
388 monthly variation and magnitude, showing higher values in spring and the maximum in winter in  
389 Shanghai, and an almost stable level around 60  $\mu\text{g m}^{-3}$  in Fuzhou throughout the year except for the  
390 elevated value in January. The PM<sub>10</sub> level in Qingdao (Figure 2j) was higher than those in Shanghai

391 and Fuzhou, and reached the maximum of  $170 \mu\text{g m}^{-3}$  in January due to anthropogenic sources and the  
392 peak in March was resulted from the effect of mineral dust.

393 The monthly variations of  $\text{PM}_{2.5}$  concentrations at Rishiri, Sado, and Oki (Figure 2m~2o) were  
394 similar to those of  $\text{PM}_{10}$ , but the peaks in May were not as evident as those of  $\text{PM}_{10}$ , because mineral  
395 dust comprises a small fraction of fine particles and has less effect on  $\text{PM}_{2.5}$  variation. The model  
396 reproduced  $\text{PM}_{2.5}$  concentrations very well at the three coastal sites of eastern China (Figure 2p~2r)  
397 and the monthly variation of  $\text{PM}_{2.5}$  concentrations resembled those of  $\text{PM}_{10}$ , because fine particle  
398 accounts for a large fraction of PM mass in these Chinese megacities due to the dominant effect of  
399 anthropogenic sources.

400 Aerosol chemical components including primary anthropogenic PM, sea salt, mineral dust, MOA,  
401 anthropogenic carbonaceous aerosols (BC+OC), and inorganic aerosols (sulfate+nitrate+ammonium)  
402 in  $\text{PM}_{10}$  and  $\text{PM}_{2.5}$  at all the EANET and CNEMC sites are also shown in Figure 2. It is found that sea  
403 salt dominated  $\text{PM}_{10}$  mass at the coastal and island sites of Japan and Korea in most months except  
404 May and June when dust aerosols were abundant (Figure 2a~2g). MOA accounted for a small fraction  
405 of  $\text{PM}_{10}$  mass at all sites, however, the relative contribution of MOA to  $\text{PM}_{2.5}$  mass appeared to be  
406 larger than that of sea salt at the coastal sites of Japan (Figure 2m~2o), indicating the potential  
407 importance of MOA in fine aerosol mode in marine atmosphere.

408 Table 1 shows that for all the 9 EANET sites, the overall mean  $\text{PM}_{10}$  concentration was  $30.0 \mu\text{g m}^{-3}$   
409  $\text{m}^{-3}$  from observation and  $28.5 \mu\text{g m}^{-3}$  from simulation, with the overall Pearson correlation coefficient  
410 (R) of 0.65 (0.48~0.64) and the normalized mean bias (NMB) of -5% (-27~36%). For  $\text{PM}_{2.5}$ , the mean  
411 concentrations averaged over the EANET sites were  $10.9 \mu\text{g m}^{-3}$  from observation and  $12.3 \mu\text{g m}^{-3}$   
412 from simulation, with R and NMB of 0.61 (0.53~0.64) and 12% (0~21%), respectively. The annual  
413 mean observed and simulated  $\text{PM}_{10}$  concentrations at the 3 CNEMC sites (Table 2) were  $81.6 \mu\text{g m}^{-3}$   
414 and  $80.7 \mu\text{g m}^{-3}$ , with R and NMBs of 0.65 (0.38~0.61) and -1% (-4~1%), respectively, while the  
415 annual mean observed and simulated  $\text{PM}_{2.5}$  concentrations, R, and NMB were  $46.6 \mu\text{g m}^{-3}$ ,  $43.4 \mu\text{g m}^{-3}$ ,  
416  $\text{m}^{-3}$ , 0.70 (0.44~0.72), and -7% (-12~0%), respectively. The good performance statistics shown in  
417 Table 1 and Table 2 suggest a good skill of RIEMS-Chem in reproducing PM levels from the coastal  
418 regions of east China to the remote western Pacific. Figure 2, Table 1, and Table 2 also illustrate that  
419 the spatial distribution of PM exhibited higher concentrations at the continental (coastal) sites  
420 (CNEMC sites, Jeju, Kanghwa, and Imsil) and lower concentrations at the remote island site

421 (Ogasawara) over the western Pacific, which were also reasonably reproduced by RIEMS-Chem.

422 Seasonal mean statistics of PM<sub>10</sub> and PM<sub>2.5</sub> concentrations at the EANET and CNEMC sites were  
423 also listed in Table 1 and Table 2. Statistics for spring (March, April, and May, MAM), summer (June,  
424 July, and August, JJA), autumn (September, October, and November, SON), and winter (December,  
425 January, and February, DJF) were calculated. PM<sub>10</sub> observations generally exhibited higher  
426 concentrations in MAM and DJF, moderate concentrations in SON, and lower concentrations in JJA at  
427 most sites covering coastal areas (CNEMC sites, Jeju, Kanghwa, and Imsil) and remote islands (e.g.,  
428 Oki, Hedo, and Ogasawara). The model reproduced such seasonal variation of PM<sub>10</sub> reasonably well  
429 although some underestimations occurred from winter to spring at Jeju and Imsil (Figure 2g, 2i),  
430 which could be attributed to the uncertainties in emissions (anthropogenic, biomass burning).

431 Comparison with observations of Sodium (Na<sup>+</sup>) concentration at 6 Japan coastal/island sites from  
432 EANET is conducted to further examine the model performance for sea salt. The modeled sodium is  
433 estimated to be 38.56% of sea salt mass (Kelly et al., 2010), and the agreement between observation  
434 and model simulation is generally satisfactory at all sites except at Oki in December, when the model  
435 largely underpredict Na<sup>+</sup>. The model well reproduces the seasonality of sodium concentration, with  
436 the maximum in winter and the minimum in summer (Figure 3). The model predicts sodium  
437 concentration best at Ogasawara, with the correlation coefficient of 0.85 and NMB of 5%. The overall  
438 correlation coefficient for all sites is 0.50, with NMB of -11% (Table S1).

439 In all, RIEMS-Chem was able to reasonably reproduce the spatial distribution and seasonal  
440 variation of PM<sub>10</sub>, PM<sub>2.5</sub>, and sodium concentrations in the marine environment of the western Pacific.  
441 The above good performances give us confidence in the estimation of marine sea salt emission.

442 In addition, the overall statistics were generally acceptable for gas precursors (O<sub>3</sub>, SO<sub>2</sub>, and  
443 NO<sub>x</sub>/NO<sub>2</sub>), indicating atmospheric chemistry processes could be reasonably represented by the model  
444 over the western Pacific (see statistics in Table S2).

445

### 446 3.2 Carbonaceous aerosols

447 Modeled BC and OC concentrations were compared with observations from research cruises and  
448 from previous publications at coastal/remote islands. BC is considered to be inert and chemical  
449 inactive, so it is governed solely by physical processes and a good indicator of long-range transport.  
450 The analysis of BC can help identify regions with large continental influence.

451

### 452 3.2.1 Comparison with research cruise measurements

453 Figure 4a shows the observed and simulated daily BC concentrations along the cruise track  
454 during the spring campaign. An obvious spatial gradient was found for BC concentration, which was  
455 characterized by apparent higher concentrations of 0.5~4.2  $\mu\text{g m}^{-3}$  over the marginal seas of China  
456 (the Yellow Sea and East China Sea, 18~19 March and 21~22 April) and very low concentrations of  
457  $<0.2 \mu\text{g m}^{-3}$  over open oceans (during most of the measurement days). It is interesting to note that an  
458 observed BC peak occurred on 21 March, which could be attributed to the long-range transport of  
459 biomass burning plumes from northeast Asia (Luo et al., 2016; 2018). The model generally  
460 reproduced the spatial and temporal variations of BC concentration during the campaign period;  
461 however, the BC peak on 21 March was missed by the model simulation. Uncertainties in biomass  
462 burning emission could be responsible for such model bias. On average, the measured and simulated  
463 BC concentrations during this campaign onboard the Dongfanghong II cruise were 0.49  $\mu\text{g m}^{-3}$  and  
464 0.55  $\mu\text{g m}^{-3}$ , respectively, with the R and NMB of 0.87 and 13% (Table 3).

465 Figure 4b shows the daily mean OC concentrations from observation and model simulation for  
466 the same cruise. In general, the observed OC exhibited a similar spatial distribution and temporal  
467 variation to that of BC, with higher concentrations over the marginal seas and relatively lower  
468 concentrations over open oceans. The model generally captured the spatial-temporal features along the  
469 cruise track. Like BC, the observed OC concentrations were high on 21 and 25~26 March mainly due  
470 to the continental outflow of biomass burning emissions from northeast Asia, and the model largely  
471 underpredict the high OC observation in these days. It is noteworthy that two OC peaks appeared on  
472 10 and 12 April when the ship was over the open ocean east of Japan (the ship location was around  
473 33.5°N, 146.0°E on 10 April and around 36.5°N, 145.0°E on 12 April, approximately 400~500 km to  
474 the east of Japan), whereas the elevation of BC concentration was not evident. Because BC and OC  
475 are often originated from the same anthropogenic and biomass sources, the inconsistency in daily  
476 variation between BC and OC in these areas implied a potential influence of marine sources rather  
477 than that from anthropogenic and biomass burning emissions. Coincidentally, during these days, daily  
478 Chl-a concentrations over the oceanic areas east of Japan (the region of 35°N to 43°N and 140.0°E to  
479 150.0°E, north to the ship location) reached as high as 45  $\text{mg m}^{-3}$ , as a comparison, the monthly mean  
480 Chl-a concentration in April over the same region was in a range of 2 to 14  $\text{mg m}^{-3}$ . The apparent



481 higher Chl-a concentration during these days could induce changes in marine primary organic  
482 emissions. On 10 April, the wind direction in the vicinity of the cruise was mainly southwesterly,  
483 backward trajectory (figure not shown) indicates that air parcels travelled over low Chl-A regions to  
484 the southwest of the cruise, implying a small effect of MOA. On this day, the fraction of land-OC (OC  
485 originated from continental sources) in total OC was 68%, which was larger than that of marine  
486 organic carbon (marine-OC) (32%) as shown in Figure 4b. On 12 April, northwesterly winds prevailed  
487 over the cruise region, with backward air trajectory traveling over strong Chl-A regions to the  
488 northeast of Japan (figure not shown), marine-OC aerosols produced from the bloom regions could be  
489 blown to the southeast where ship located, leading to the elevation of OC concentrations (Figure 4b).  
490 Marine-OC (percentage contribution of 74%) dominated over land-OC (26%) in the total OC  
491 concentration on this day. The model improves OC simulation on 10 and 12 April when considering  
492 marine organic aerosols (marine-OC in Figure 4b). However, it should be acknowledged that the  
493 model appears to generally overpredict OC concentrations during 5~16 April over the ocean southeast  
494 of Japan, especially on 5~6, 11 and 13 April. The high model biases could be due to potential  
495 overpredictions for either land-source OC or marine-source OC. The cruise campaign average OC  
496 concentration was  $1.2 \mu\text{g m}^{-3}$  from observation and  $1.1 \mu\text{g m}^{-3}$  from simulation, with the R and NMB  
497 of 0.66 and -5%, respectively (Table 3). For the coastal/marginal sea areas (cruising time on 18 to 20  
498 March and 20 to 22 April), the mean observed and simulated OC concentrations were  $3.1 \mu\text{g m}^{-3}$  and  
499  $3.0 \mu\text{g m}^{-3}$ , respectively. For open sea areas (cruising time from 21 March to 19 April), the observed  
500 and simulated OC concentrations were  $0.69 \mu\text{g m}^{-3}$  and  $0.65 \mu\text{g m}^{-3}$ , respectively. The inclusion of  
501 marine-OC (including both primary and secondary OC) reduced the model bias from -33% to -5%  
502 along the cruise. The average contribution of marine-OC to the total OC mass in the marine  
503 atmosphere was approximately 29% along the cruise, with lower contributions of 11~27% over the  
504 marginal seas of China (18~19 March and 21~22 April) and higher contributions of 32~74% over the  
505 open oceans (5~18 April) (Figure 4b), demonstrating an increasing importance of marine organic  
506 aerosols to total OC mass from the marginal seas to remote open oceans.

507 Shown in Figure 4c is OC samples collected onboard the KEXUE-1 Research Vessel over the  
508 East China Sea during the early summer campaign and the corresponding model results along the  
509 cruise track. There were four OC peaks observed during the campaign, with three occurring over the  
510 northern parts of the East China Sea (on 20 May, 26~29 May, and 1~5 June) and one over the southern

511 part of the East China Sea on 22 May. The model reproduced the OC variation quite well during most  
512 of the cruise track, capturing the three OC peaks over the northern parts of the East China Sea  
513 although low biases occurred for the first peak (over the area of 27.5°N to 30.0°N and 121.6°E to  
514 121.9°E). The model missed the second OC peak on 22 May over the southern part of the East China  
515 Sea (over the area of 22°N to 23°N and 121.5°E to 122.2°E). Kang et al. (2018) proposed that this  
516 peak was seriously affected by biogenic and biomass burning emissions from Southeast Asia  
517 (Philippines) because the OC concentrations from 21 to 25 May were characterized by high  
518 abundance of sesquiterpene-derived SOA which was mainly originated from terrestrial photosynthetic  
519 vegetation (e.g. trees and plants). Uncertainties in emission inventories, such as missing some  
520 biogenic sources (e.g. fungal spores, Fröhlich-Nowoisky et al. 2016) could be partly responsible to the  
521 model biases. In addition, some regions of Southeast Asia (e.g. Philippines) were not included in the  
522 study domain, instead, their influence on the study domain was represented by chemical boundary  
523 conditions from MOZART simulation, so, the uncertainties in chemical boundary conditions may also  
524 contributed to such biases. At the time of the third (25°N to 26°N and 118.8°E to 121.7°E) and fourth  
525 (28°N to 28.7°N and 119.6°E to 122.7°E) OC peaks, the ship was close to the shore and  
526 predominately affected by continental sources (such as anthropogenic and biomass burning emissions),  
527 the model captured the peaks quite well in terms of both temporal variation and magnitude. On  
528 average, the observed and simulated OC concentrations from the KEXUE-1 cruise were  $4.3 \mu\text{g m}^{-3}$   
529 and  $3.7 \mu\text{g m}^{-3}$ , respectively, with R and NMB of 0.75 and -13% (Table 3). The inclusion of  
530 marine-OC reduced the NMB from -19% to -13%. Along the cruise track, marine-OC was estimated  
531 to account for 6% (1~60%) of the total OC mass on average, with lower contribution over the seas  
532 close to the continent (1~9%) and higher contribution over the seas far from the continent (7~60%).  
533 During the KEXUE-1 cruise campaign, the contribution of marine-OC to total OC mass was  
534 obviously lower than that during the spring campaign conducted by the Dongfanghong II, because this  
535 cruise over the marginal seas of China was more affected by continental outflow of anthropogenic and  
536 biomass emissions compared with that mainly over the open oceans.

537

### 538 3.2.2 Comparison with measurements at island and coastal sites

539 Figure S1 shows the modeled BC is generally consistent with observations at island sites  
540 (Huaniao, Fukue, Okinawa, Chichijima) in terms of both spatial distribution and seasonal variation,

541 indicating a good skill of RIEMS-Chem in representing the physical processes and long-rang transport  
542 of carbonaceous aerosols over the western Pacific.

543 OC observations are limited in the western Pacific Ocean. We collected observations at islands  
544 from previous publications (Boreddy et al., 2018; Kunwar and Kawamura, 2014; Wang F. W. et al.,  
545 2015) for model comparison. Figure 5 shows the model simulated and observed seasonal/monthly  
546 mean OC concentrations at the three islands. It should be kept in mind that the observations are  
547 averages of different years. At Huaniao Island (Figure 5a), a distinct seasonality of OC observation  
548 was shown, with the highest OC concentration of  $4.7 \mu\text{g m}^{-3}$  in DJF, followed by  $3.7 \mu\text{g m}^{-3}$  in MAM  
549 and  $3.8 \mu\text{g m}^{-3}$  in SON, and the minimum of  $1.1 \mu\text{g m}^{-3}$  in JJA (Table 4). It was encouraging that  
550 RIEMS-Chem reproduced the OC seasonality at Huaniao Island quite well (Figure 5a), despite the  
551 different years between simulation and observation. The simulated OC was also divided into land-OC  
552 and marine-OC to quantify the relative contribution of these sources to total OC mass. The simulated  
553 annual mean OC concentration was  $3.2 \mu\text{g m}^{-3}$ , in which  $2.6 \mu\text{g m}^{-3}$  (81%) was contributed by  
554 land-OC and  $0.6 \mu\text{g m}^{-3}$  (19%) by marine-OC (Table 4). The simulation was very close to the  
555 observation of  $3.3 \mu\text{g m}^{-3}$  (Table 4). It was striking that the inclusion of marine-OC obviously  
556 improved the model performance, reducing the NMB from -21% to -3%, although the improvement of  
557 prediction for SOA from land source may also reduce the model bias at Huaniao island. It was  
558 noteworthy that marine-OC exhibited the maximum value in MAM and the minimum value in JJA.  
559 The higher Chl-a concentration over the East China Sea in MAM might be responsible for the  
560 maximum at Huaniao Island (Figure 7h and Table 7), whereas the lowest sea salt emission flux could  
561 result in the minimum in summer (Table 7). In terms of seasonal mean, marine-OC accounted for 12%,  
562 22%, 19%, and 23% of the total OC concentration in DJF, MAM, JJA, and SON, respectively, with an  
563 annual mean contribution of 19% at Huaniao Island. The lowest relative contribution (12%) of  
564 marine-OC in winter was attributed to the maximum anthropogenic OC emissions in eastern China in  
565 this season.

566 At Okinawa (Figure 5b), the observed total OC showed the maximum in MAM, followed by that  
567 in JJA, and the lower ones in DJF and SON during October 2009-2010. Figures 5a and 5b also show  
568 that the seasonal cycling of OC concentration at Okinawa (Figure 5b) differed a lot from that at  
569 Huaniao Island (Figure 5a). The high OC concentration in summer at Okinawa could be attributed to  
570 higher SOA produced by local biogenic VOC emissions (Kunwar and Kawamura, 2014). The model

571 generally reproduced the seasonal variation of OC except that it predicted lower OC level in summer,  
572 which could be due to the exclusion of local biogenic VOC emissions in the CAMS-GLOB-BIO  
573 emission inventory. In terms of annual average, the observed OC concentration was  $1.8 \mu\text{g m}^{-3}$ , larger  
574 than the simulations of  $1.3 \mu\text{g m}^{-3}$  from the FULL case including marine-OC and of  $1.1 \mu\text{g m}^{-3}$  from  
575 the case excluding marine organic emissions (Table 4). The inclusion of marine organic emissions  
576 improved OC simulation at Okinawa, reducing the NMB from -39% to -28%. It was estimated that  
577 marine-OC accounted for 18%, 17%, 10%, and 18% of total OC mass concentration at Okinawa in  
578 DJF, MAM, JJA, and SON, respectively, with an annual mean contribution of 17%. The relatively  
579 smaller contribution of marine-OC to the total OC mass at Okinawa than that at Huaniao Island (19%)  
580 could be attributed to the higher Chl-a concentration and MPOA emission flux in the marginal seas of  
581 China than those over remote western Pacific south of Japan (Figure 7).

582 Long-term average (2001-2012) of monthly mean OC concentrations at Chichijima Island  
583 reported by Boreddy et al. (2018) and the simulated monthly mean OC concentration in 2014 were  
584 shown in Figure 5c. The observations show higher OC levels from January to March mainly due to  
585 continental outflows. It was noticed that the simulated OC levels in April-May were apparently higher  
586 than observations, which could be associated with different time periods between observation and  
587 simulation, and with potentially stronger continental outflows and bloom in spring 2014 than those of  
588 ten-year averages. OC observations were relatively lower in summer and autumn due to the  
589 dominance of high-pressure system and pristine ocean air mass over the western Pacific (Figure 9d  
590 and 9e). The model tended to predict lower OC level in summer and autumn (Figure 5c). Boreddy et  
591 al. (2018) indicated that in summer and autumn, OC at Chichijima was often influenced by long-range  
592 transport of biomass burning plumes from Southeast Asia, which was not well represented in the  
593 model (using chemical boundary conditions from MOZART-4 instead) and led to low model bias. On  
594 average, the annual mean OC concentration was  $0.76 \mu\text{g m}^{-3}$  from observation, and  $0.78 \mu\text{g m}^{-3}$  from  
595 the FULL case, and  $0.65 \mu\text{g m}^{-3}$  without considering marine-OC (Table 5). The inclusion of marine  
596 organic emissions reduced the annual mean NMB from -13% to 3% and enhanced the correlation  
597 coefficient from 0.56 to 0.6 at this site. The apparent better simulation from the FULL case indicated  
598 the necessity of inclusion of marine organic emissions for simulating OC over the remote oceans of  
599 the western Pacific. Both observation and model simulation revealed higher seasonal mean OC  
600 concentrations in MAM (observed:  $0.83 \mu\text{g m}^{-3}$ , simulated:  $0.91 \mu\text{g m}^{-3}$ ) and DJF (observed:  $0.9 \mu\text{g m}^{-3}$ )

601  $\text{m}^{-3}$ , simulated:  $1.2 \mu\text{g m}^{-3}$ ) when the measurement site was frequently influenced by continental  
602 outflows, whereas lower concentrations in JJA (observed:  $0.65 \mu\text{g m}^{-3}$ , simulated:  $0.47 \mu\text{g m}^{-3}$ ) and  
603 SON (observed:  $0.66 \mu\text{g m}^{-3}$ , simulated:  $0.57 \mu\text{g m}^{-3}$ ) when clean maritime air masses or biomass  
604 burning plumes from Southeast Asia (e.g. Philippine) influenced this region. The highest marine-OC  
605 concentration was  $0.19 \mu\text{g m}^{-3}$  in MAM, followed by  $0.16 \mu\text{g m}^{-3}$  in DJF and  $0.11 \mu\text{g m}^{-3}$  in SON, and  
606 the lowest one of  $0.05 \mu\text{g m}^{-3}$  in JJA. However, the percentage contribution of marine-OC to the total  
607 OC mass was estimated to be largest in SON (20%), followed by 18% in DJF, 16% in MAM, and  
608 lowest in JJA (10%), with an annual mean contribution of 16% (Table 5). The largest contribution in  
609 SON was associated with the relatively lower total OC concentration as shown in Figure 5c. The  
610 relative contribution from marine-OC to total OC at Chichijima Island resembled that at Okinawa in  
611 terms of annual and season averages.

612 The above comparison against a variety of OC observations demonstrated a generally good skill  
613 of RIEMS-Chem in simulating OC over the western Pacific in terms of seasonal variation and  
614 magnitude. The model results from the FULL case indicated that including marine organic emissions  
615 improved OC simulation over the western Pacific Ocean.

616

### 617 3.2.3 SOA over the western Pacific

618 Recently, Guo et al. (2020) reported SOA observations in the marine atmosphere from the  
619 marginal seas of east China to the northwest Pacific Ocean. The measurements were conducted on  
620 three research cruises in the spring and early summer of 2014 and in the spring of 2017. Total  
621 suspended particulate (TSP) samples were collected from 19 March to 21 April 2014 over the  
622 northwestern Pacific Ocean (NWPO), from 30 April to 17 May 2014 over the Yellow and Bohai seas  
623 (YBS), and from 29 March to 4 May 2017 over the South China Sea (SCS). SOA concentration was  
624 derived by using a tracer-based method. The measured SOA concentrations were  $467 \pm 384 \text{ ng m}^{-3}$  over  
625 the YBS,  $617 \pm 649 \text{ ng m}^{-3}$  over the SCS, and  $155 \pm 236 \text{ ng m}^{-3}$  over the NWPO, respectively. The  
626 model simulated period and regional mean SOA concentrations were  $664 \text{ ng m}^{-3}$  over the YBS,  $466 \text{ ng}$   
627  $\text{m}^{-3}$  over the SCS, and  $157 \text{ ng m}^{-3}$  over the NWPO, which were generally consistent with the above  
628 observations, although the study periods are not exactly the same. Guo et al. (2020) also presents the  
629 tracer-based estimations of isoprene and monoterpene derived SOA in the air masses from ocean  
630 (assuming marine sources), which were  $1.7 \text{ ng m}^{-3}$  and  $0.3 \text{ ng m}^{-3}$ , respectively, over the western

631 Pacific to the southeast of Japan, whereas the modeled SOA concentrations produced from marine  
632 isoprene and monoterpene emissions along the cruise track were  $1.6 \text{ ng m}^{-3}$  and  $0.28 \text{ ng m}^{-3}$ ,  
633 respectively, generally agreeing with the tracer-estimation. However, it should be mentioned that there  
634 could be uncertainties in such comparison. First, the isoprene- and monoterpene-derived SOA tracers  
635 in the air masses categorized as marine sources by Guo et al (2020) might include SOA tracers from  
636 terrestrial isoprene and monoterpene under the prevailing northwesterly winds in spring, which could  
637 bias the estimation high; second, the measured tracer could just comprise a part of total SOA tracers,  
638 which might bias the estimation low. Despite these uncertainties, the cruise measured SOA  
639 concentration derived from marine isoprene and monoterpene was approximately several  $\text{ng m}^{-3}$  over  
640 the western Pacific, and it can reach approximately  $10 \text{ ng m}^{-3}$  even through dividing by a mass  
641 fraction of tracer compound to yield the concentration of total SOA tracers. It was noteworthy that  
642 both observation and model simulation exhibited a decreasing SOA concentration from marginal seas  
643 of China to remote oceanic areas. In all, the model reproduced the SOA levels in the marine  
644 atmosphere of the western Pacific Ocean reasonably well.

645 The comparison of the magnitudes between SOA and OA mass (1.4 times OC mass)  
646 concentrations shown above indicates that SOA concentration was approximately 1~2 orders of  
647 magnitude lower than OA over the western Pacific. Previous observation studies using the  
648 tracer-based approach also indicated that the percentage contribution of SOA to OA was quite low  
649 over some marine areas (Fu et al., 2011; Hu et al., 2013; Bikkina et al., 2014; Zhu et al., 2016). For  
650 example, at Okinawa island, even considering all biogenic sources (including isoprene, monoterpene,  
651 and sesquiterpene of both terrestrial and oceanic origins), the measured concentration of total  
652 biogenic-SOA tracers was still less than  $100 \text{ ng m}^{-3}$ , with majority of SOA tracers from local terrestrial  
653 biogenic emissions (Zhu et al., 2016). The above studies suggested that primary organic aerosols were  
654 more important in remote marine atmosphere.

655

### 656 3.3 Aerosol optical depth

657 Figure 6 shows the temporal variations of the observed and simulated monthly mean AOD at the  
658 7 AERONET sites. In general, RIEMS-Chem simulated the monthly mean AOD reasonably well in  
659 terms of magnitude and monthly variation at almost all sites, although some biases occurred during  
660 some months, such as the overpredictions in August at Fukuoka and in April at EPA-NCU, and the

661 underprediction in July at Yonsei University. For the sites in the northern oceanic areas (Ussuriysk,  
662 Yonsei\_University, Gwangju\_GIST, and Fukuoka, Figure 6a~6d), both observations and simulations  
663 generally exhibited higher AOD values in summer (JJA), moderately high AOD values from late  
664 winter (JF) to spring (MAM), and relatively lower AOD values in autumn (SON). The simulated  
665 higher inorganic aerosol concentrations in summer and late spring months could be responsible for the  
666 higher AOD values in these regions. Besides, the higher relative humidity in summer due to the  
667 predominant influence of maritime air masses also contributed to the maximum AOD values during  
668 summer months (JJA) at these sites. On the other hand, for the sites in the southern oceanic areas  
669 (EPA-NCU and Chen-Kung\_ Univ, Figure 6e and 6f), the monthly mean AOD was apparently higher  
670 from March to April and remained low levels during the rest months. The above AOD peaks in spring  
671 could be attributed to the continental outflows of biomass burning plumes originated from Southeast  
672 Asia, which were most active in springtime in those regions (Hsiao et al., 2017; Tao et al., 2020).  
673 Table 6 shows the performance statistics for hourly AOD at these AERONET sites. The overall annual  
674 mean AOD for the 7 sites was 0.34 from model simulation, which was very close to the observation of  
675 0.36, with the NMB of -6% and the overall correlation coefficient of 0.54 (0.40~0.67). The statistics  
676 indicate that the model was able to reproduce aerosol optical properties over the coastal regions and  
677 islands around the western Pacific Ocean. Contributions of MOA, sea salt, and non-oceanic aerosols  
678 (anthropogenic and other natural aerosols from the Asian continent) to total AOD are also shown in  
679 Figure 6. It clearly shows the predominant contribution to AOD from non-oceanic aerosols at all sites.  
680 It is also noticed that AOD induced by sea salt is generally larger than that by MOA, but they are  
681 comparable at the Chen-Kung Univ and EPA-NCU sites in spring (Figure 6f and 6g).

682 The model simulated annual mean AOD at 550 nm are also compared with the VIIRS retrievals  
683 (Figure S2), which indicates the model is generally capable of reproducing AOD distribution and  
684 magnitude in the study domain. The generally high model bias over the western Pacific could be  
685 attributed to potential overpredictions of inorganic aerosol concentration and relative humidity. AOD  
686 reflects the column integrated extinction coefficient due to all aerosols.

687 At the AERONET sites, the model simulated annual mean percentage contribution of MOA to  
688 AOD varied from 1.4% to 3.2% with an overall average of 1.9%. For the oceanic VIIRS region, the  
689 mean contribution of MOA to AOD was approximately 2%.

690

## 691 4 Model results

### 692 4.1 Marine primary organic and isoprene emissions

693 Figure 7 shows the estimated annual and seasonal mean MPOA emission rates over the western  
694 Pacific of East Asia. The MPOA emission mainly occurred over two hotspot regions: the marginal  
695 seas of China including the East China Sea, the Yellow Sea, and the Bohai Sea (EYB, 27°N~40°N,  
696 115°E~123°E, denoted in Figure 7a) and the northern parts of the western Pacific northeast of Japan  
697 (NWP, 35°N~54°N, 140°E~160°E, denoted in Figure 7a), with annual mean emission rates varying  
698 from  $0.9 \times 10^{-2} \mu\text{g m}^{-2} \text{ s}^{-1}$  to  $1.8 \times 10^{-2} \mu\text{g m}^{-2} \text{ s}^{-1}$ . In SON, high MPOA emission occurred in both the  
699 EYB and NWP regions, with the maximum up to  $3.5 \times 10^{-2} \mu\text{g m}^{-2} \text{ s}^{-1}$  in the NWP (Figure 7e),  
700 whereas MPOA emission was very low over the EYB in JJA (Figure 7d). The maximum seasonal  
701 mean emission rate of MPOA approached  $3.6 \times 10^{-2} \mu\text{g m}^{-2} \text{ s}^{-1}$  over the Yellow Sea in DJF (Figure 7b),  
702 which was approximately 1/10 of the annual mean anthropogenic POA emission rate in north China  
703 (on the order of  $1.0 \sim 3.0 \times 10^{-1} \mu\text{g m}^{-2} \text{ s}^{-1}$ ). Table 7 presents the seasonal and annual averages of MPOA  
704 emission averaged over the western Pacific and the EYB and NWP regions (note they are oceanic  
705 averages with land grids excluded). In terms of oceanic average of the western Pacific, the mean  
706 MPOA emission generally exhibited the largest emission rate in SON ( $0.2 \times 10^{-2} \mu\text{g m}^{-3} \text{ s}^{-1}$ ),  
707 moderately high emission rates in DJF ( $0.18 \times 10^{-2} \mu\text{g m}^{-2} \text{ s}^{-1}$ ) and MAM ( $0.17 \times 10^{-2} \mu\text{g m}^{-2} \text{ s}^{-1}$ ), and the  
708 lowest one in JJA ( $0.08 \times 10^{-2} \mu\text{g m}^{-2} \text{ s}^{-1}$ ), with an annual average of  $0.16 \times 10^{-2} \mu\text{g m}^{-2} \text{ s}^{-1}$  (Table 7). It is  
709 interesting to note that the seasonal variation of MPOA emission was not consistent with that of Chl-a  
710 concentration, which exhibited higher values in SON and JJA and the lowest one in DJF (Table 7).  
711 This is because MPOA emission rate is determined by the combined effect of Chl-a concentration and  
712 sea salt emission flux, and sea salt flux is mainly controlled by surface wind speed according to the  
713 scheme of Gong (2003). In terms of seasonal and domain average over the western Pacific, the  
714 maximum Chl-a concentration and the second largest sea salt emission flux in SON led to the largest  
715 MPOA emission in autumn (Table 7). However, although Chl-a concentration was also high in JJA  
716 ( $1.07 \text{ mg m}^{-3}$ , Table 7), the sea salt flux was minimum in JJA ( $0.14 \mu\text{g m}^{-2} \text{ s}^{-1}$ , Table 7) due to the  
717 weakest wind speed ( $3.0 \text{ m s}^{-1}$ , Table 9), resulting in the lowest MPOA emission in summer (Table 7).  
718 Although the sea salt emission flux reached the maximum in DJF (Table 7) due to the largest wind  
719 speed in this season (Table 9), the winter Chl-a concentration was lowest, leading to a moderate  
720 MPOA emission in winter (Table 7), in a similar magnitude to that in spring when moderately high



721 Chl-a concentration and relatively low sea salt flux occurred. In all, the MPOA emission rate over the  
722 western Pacific exhibited an apparent seasonality of SON > DJF  $\approx$  MAM > JJA.

723 For the EYB region, the maximum MPOA emission occurred in winter (DJF) (Figure 7b and  
724 Table 7) with a seasonal and domain average of  $1.2 \times 10^{-2} \mu\text{g m}^{-2} \text{s}^{-1}$ , which was 10 times larger than  
725 the minimum of  $0.12 \times 10^{-2} \mu\text{g m}^{-2} \text{s}^{-1}$  in summer (JJA) (Figure 7d and Table 7). Although Chl-a  
726 concentrations were similar between DJF and JJA, the sea salt flux in DJF was approximately 9 times  
727 that in JJA (Table 7). So, the seasonality of MPOA emission in the EYB region was mainly  
728 determined by that of sea salt emission flux due to the weak seasonal variation of Chl-a concentration.  
729 Differently, in the NWP region, MPOA emission exhibited the maximum value in SON, followed by  
730 those in MAM and DJF, and the lowest ones in JJA (Table 7). It is interesting to note that although  
731 both the Chl-a concentration and sea salt emission flux were slightly higher in MAM than those in  
732 SON, the MPOA emission (related to both Chl-a concentration and sea salt emission) was higher in  
733 SON, which could be due to the slight negative correlation between Chl-a and sea salt emission in MAM,  
734 but the slight positive one in SON. The MPOA emissions in winter and summer were in a similar level  
735 in the NWP region, about 40% lower than that in autumn.

736 The distribution pattern of MPOA emission in the western Pacific from this study is similar to  
737 those from previous model studies (Spracklen et al., 2008; Gantt et al., 2009; Huang et al., 2018), but  
738 the magnitude of the simulated MPOA emission flux is larger than previous estimates. For example,  
739 the annual mean MPOA emission rates over the western Pacific were estimated to vary from 0.1 to  
740 approximately  $12 \text{ ng m}^{-2} \text{s}^{-1}$  in previous studies (Spracklen et al., 2008; Vignati et al., 2010; Gantt et  
741 al., 2011; Long et al., 2011; Huang et al., 2018), whereas the estimates in this study ranged from 3 to  
742  $18 \text{ ng m}^{-2} \text{s}^{-1}$  (Figure 7a). The larger marine POA emission estimated in this study could be attributed  
743 to the application of the daily mean Chl-a concentration from satellite retrievals and of a finer model  
744 grid resolution (60 km) compared with those in global models. On average, the annual MPOA  
745 emission was estimated to be  $0.78 \text{ Tg yr}^{-1}$  over the western Pacific (with an ocean area of  $1.6 \times 10^7 \text{ km}^2$ )  
746 from this study. The regions of EYB and NWP comprised approximately 2% and 18% of the western  
747 Pacific in terms of area, respectively, but they contributed 8% and 46% of the MPOA annual emission  
748 ( $\text{Tg yr}^{-1}$ ). This study revealed that the EYB and NWP are important bloom regions, totally accounting  
749 for more than half of the total MPOA emission over the western Pacific.

750 Table S3 presents the simulated marine isoprene emission fluxes from this study and the estimates

751 based on cruise observations over the western Pacific of East Asia from previous studies. Over the  
752 western North Pacific, the observed marine isoprene emission flux showed larger values in May (140  
753  $\text{nmol m}^{-2} \text{ day}^{-1}$  from Matsunaga et al. (2002) and  $143.8 \text{ nmol m}^{-2} \text{ day}^{-1}$  from Ooki et al., (2015)), a  
754 moderate value in August ( $55.6 \text{ nmol m}^{-2} \text{ day}^{-1}$  from Ooki et al., (2015)), and the lowest one in winter  
755 ( $21.4 \text{ nmol m}^{-2} \text{ day}^{-1}$  from Ooki et al., (2015)). The simulation from this study generally agreed with  
756 observation in terms of both seasonality and magnitude except for the low bias in May ( $85\sim 89 \text{ nmol}$   
757  $\text{m}^{-2} \text{ day}^{-1}$  vs observation-based estimates of  $140\sim 143.8 \text{ nmol m}^{-2} \text{ day}^{-1}$ ), which could be associated  
758 with the different years. According to equations (2) and (3) (in the supplement), both Chl-a  
759 concentration and incoming solar radiation determine marine biogenic VOCs emission, the larger  
760 isoprene flux in May was mainly due to the maximum Chl-a concentration in spring over the NWP  
761 region (Table 7). Over the marginal seas of China, Li J. L. et al. (2017; 2018) observed higher marine  
762 isoprene emission flux in July-August ( $161.5 \text{ nmol m}^{-2} \text{ day}^{-1}$ ) than those in October-November ( $48.3$   
763  $\text{nmol m}^{-2} \text{ day}^{-1}$ ) and May-June ( $36.1 \text{ nmol m}^{-2} \text{ day}^{-1}$ ) during 2013-2014. The model results from this  
764 study show the similar seasonal variation and magnitude of isoprene flux, with corresponding mean  
765 values of  $130 \text{ nmol m}^{-2} \text{ day}^{-1}$ ,  $48 \text{ nmol m}^{-2} \text{ day}^{-1}$ , and  $35 \text{ nmol m}^{-2} \text{ day}^{-1}$  during the same periods of  
766 2014, respectively. The apparently higher isoprene flux in July-August was mainly resulted from the  
767 strongest solar radiation in summer, although the Chl-a concentration was not highest in this season in  
768 the EYB region (Table 7). The domain-wide annual marine isoprene emission estimated over the  
769 western Pacific was  $0.015 \text{ Tg yr}^{-1}$  in this study.

770

#### 771 4.2 Marine organic aerosols and their relative importance

772 Annual and seasonal mean near surface MOA concentrations, MSOA concentrations, and the  
773 percentage contributions of MOA to total OA mass in the study domain were shown in Figure 8. The  
774 spatial distributions of MOA concentrations (Figure 8a~8e) generally resembled those of MPOA  
775 emissions (Figure 7a~7e). It is remarkable that MPOA concentration (MOA minus MSOA) was  
776 approximately 1~2 orders of magnitude higher than MSOA concentration (with concentration of  
777 several  $\text{ng m}^{-3}$ ) in the western Pacific (Figure 8a~8e vs Figure 8f~8j), indicating that MPOA  
778 constituted a dominant fraction of MOA, which will be discussed below. Figure 8a shows that high  
779 MOA concentrations mainly occurred over the EYB and NWP regions, with the annual and regional  
780 averages being  $0.48 \mu\text{g m}^{-3}$  and  $0.59 \mu\text{g m}^{-3}$ , respectively (Table 8), accounting for 13% (6~30%) and

781 42% (30~60%) of total OA mass in these two regions, respectively (Figure 8k and Table 8). The larger  
782 MOA contribution over the NWP was attributed to the high MOA level and the relatively low total OA  
783 level there. It is noticed that MOA even influenced the coastal areas of eastern China. The annual  
784 mean MOA concentration decreased from approximately  $0.5 \mu\text{g m}^{-3}$  in coastal areas to  $0.1 \mu\text{g m}^{-3}$  in  
785 the inland areas approximately 600 km away from the coastline (Figure 8a), accounting for  
786 approximately 2% to 6% of the near-surface OA mass in the coastal regions (Figure 8k). The  
787 maximum seasonal mean MOA concentration over the coastal areas of eastern China could be up to  
788  $0.6 \mu\text{g m}^{-3}$  to  $0.8 \mu\text{g m}^{-3}$  in MAM (Figure 8c) and SON (Figure 8e). The domain and seasonal mean  
789 MOA concentration over the western Pacific exhibited the maximum value in MAM ( $0.37 \mu\text{g m}^{-3}$ ),  
790 follow by that in SON ( $0.26 \mu\text{g m}^{-3}$ ), and relatively lower concentrations in JJA ( $0.23 \mu\text{g m}^{-3}$ ) and DJF  
791 ( $0.21 \mu\text{g m}^{-3}$ ) (Table 8). It was noteworthy that the seasonality of MOA concentration was different  
792 from that of MPOA emission, which could be attributed to the influence of different meteorological  
793 conditions and physical processes. In the western Pacific, although MPOA emission peaked in SON  
794 (Table 7), MOA concentration peaked in MAM (Table 8). It is noticed that precipitation was lowest  
795 and wind speed was low in MAM (Figure 9c and 9h, Table 9), leading to a smaller dry deposition  
796 velocity (Zhang et al. 2001) and the weakest wet scavenging, both favored accumulation of MOA and  
797 thus resulted in the highest MOA level in spring. On the contrary, due to the maximum wind speed  
798 and relatively more precipitation in DJF (Figure 9b and 9g, Table 9), the mean MOA concentration  
799 was lowest in winter.

800 For the EYB region, northwesterly winds prevailed In DJF and SON and turned to northeasterly  
801 winds over marginal seas of southeast China (Figure 9b and 9e), which transported MOA from the  
802 major MPOA source region (EYB) to the northern part of the South China Sea (Figure 8b and 8e). As  
803 wind speed over the EYB was low in MAM and JJA (Figure 9c and 9d, Table 9), MOA was mainly  
804 restricted within this region (Figure 8c and 8d). In terms of seasonal average, MOA concentration  
805 experienced its maximum in MAM, followed by those in DJF and SON, and the minimum in JJA  
806 (Figure 8b~8e). The seasonal and regional mean MOA concentrations over the EYB were  $0.62 \mu\text{g m}^{-3}$ ,  
807  $0.54 \mu\text{g m}^{-3}$ ,  $0.52 \mu\text{g m}^{-3}$ , and  $0.22 \mu\text{g m}^{-3}$  for MAM, DJF, SON, and JJA, respectively (Table 8). The  
808 different seasonality between MOA concentration (Table 8) and MPOA emission (Table 7) in the EYB  
809 region could also be mainly attributed to meteorological conditions. The MPOA emission was  
810 relatively low in MAM (Table 7), but the second lowest wind speed and less precipitation (Table 9)

811 favored aerosol accumulation, resulting in the highest MOA concentration in spring (Table 8). The  
812 minimum MPOA emission and the maximum precipitation in JJA led to the minimum MOA  
813 concentration in summer. Although MPOA emission was largest in SON and DJF (Table 7), the  
814 maximum wind speeds (Table 9) led to stronger dry deposition of aerosols and thus a moderate MOA  
815 concentration in the two seasons (Table 8).

816 MOA concentration over the NWP region exhibited apparent higher concentrations in MAM and  
817 JJA than those in SON and DJF (Figure 8b~8e), with the regional and seasonal averages reaching 0.81  
818  $\mu\text{g m}^{-3}$  in MAM, 0.80  $\mu\text{g m}^{-3}$  in JJA, 0.52  $\mu\text{g m}^{-3}$  in SON, and 0.23  $\mu\text{g m}^{-3}$  in DJF, respectively (Table  
819 8). Using GEOS-Chem with a different marine organic aerosol emission scheme, Spracklen et al.  
820 (2008) also showed that in the North Atlantic along the similar latitude bands to the NWP ( $\sim 35^\circ\text{N}$  to  
821  $\sim 55^\circ\text{N}$ ), both observation and simulation exhibited higher OC concentrations in summer and spring  
822 than in the other seasons at the Azores Island and Mace Head Island. The strong seasonality of MOA  
823 over the NWP was also attributed to the combined effects of MPOA emission, wind speed, and  
824 precipitation. In MAM, the high MOA concentration over the NWP was mainly due to the large  
825 MPOA emission (Figure 7c and Table 7), which was just smaller than that in SON, and partly due to  
826 the relatively weak dry deposition and wet scavenging caused by moderate wind speed and  
827 precipitation in this season (Table 9). In JJA, although the MPOA emission was small, the lowest wind  
828 speed and precipitation in JJA over the NWP ( $2.5 \text{ m s}^{-1}$  and  $3.7 \text{ cm grid}^{-1} \text{ month}^{-1}$ , Table 9) led to the  
829 weakest dry deposition and wet scavenging of particles in summer, resulting in a long residence time  
830 of MOA and consequently the high MOA concentration in summer over the NWP. In SON, although  
831 the MPOA emission was largest over the NWP (Table 7), the mean wind speed was high over the  
832 northern part of the NWP (Figure 9e) where MPOA emission mainly occurred (Figure 7e), leading to  
833 strong dilution of MOA particles in autumn. Furthermore, SON had the secondly largest precipitation  
834 over the NWP (Table 9), which caused strong wet scavenging of particles and also contributed to the  
835 relatively low MOA level. In DJF, the wind speed was largest, about 2 times those in the other seasons,  
836 and the precipitation was also the maximum (Table 9, Figure 9b and 9g), leading to the lowest MOA  
837 concentration in winter over the NWP (Figure 8b and Table 8).

838 As shown in Figures 8k~8o, MOA generally accounted for approximately 30% to over 60% of  
839 total OA concentration over the remote oceans of high ( $>35^\circ\text{N}$ ) and low ( $<25^\circ\text{N}$ ) latitudes within the  
840 model domain. The large MOA/OA ratios over the remote oceans of high latitude (including NWP)

841 could be attributed to the high MOA concentration due to large marine emissions there; whereas, the  
842 large MOA/OA ratios over the subtropical oceans of low latitude were mainly due to the low total OA  
843 level (small denominator). Averaged over the NWP region, the annual mean MOA/OA ratio was 42%,  
844 with higher contributions in MAM (52%) and SON (48%) and lower ones in DJF (36%) and JJA  
845 (32%) (Table 8). Although MOA concentration over the NWP was secondly highest in JJA, its  
846 contribution was small because OA transported from land sources also subject to weak dry deposition  
847 and wet scavenging, which led to higher OA level and lower MOA/OA ratio. Over the EYB region,  
848 MOA accounted for approximately 6% to 30% of the total OA in terms of annual mean (Figures 8k).  
849 In terms of annual and regional average, the MOA/OA ratio was 13%, with higher ratios in SON (18%)  
850 and MAM (15%), a moderate one in DJF (11%), and the lowest one in JJA (6%) (Table 8), similar to  
851 the seasonality over the NWP. For the oceanic areas, the domain and annual mean MOA/OA ratio was  
852 26%, indicating an important contribution of MOA to airborne OA over the western Pacific. It was  
853 found that the importance of MOA in total OA increased as the distance to the East Asian continent  
854 increased over the western Pacific. It is also interesting to note that MOA even accounted for  
855 approximately 2~6% of the annual mean OA mass over portions of southeast China (Figures 8k), and  
856 such contribution could be as high as 8~10% in the coastal areas in SON (Figures 8o) and MAM  
857 (Figures 8m).

858 In all, both the MOA concentration and the MOA contribution to total OA were lowest in summer  
859 (JJA) in the EYB region, which was mainly due to the much smaller MPOA emission in this season.  
860 However, in the NWP region, although the MPOA emission was also lowest in summer, MOA  
861 concentration in summer was in a same level as that in spring, and larger than those in the other  
862 seasons, because dry deposition velocity and precipitation were lowest in summer, which favored  
863 aerosol accumulation and a high level of MOA.

864 SOA produced by marine biogenic VOCs (isoprene and terpene) was on the order of  $10^{-2}\sim 10^{-3}$   $\mu\text{g}$   
865  $\text{m}^{-3}$  (Figure 8f~8j), which was much lower than the MPOA concentration. The spatial distribution of  
866 MSOA exhibited high concentrations over the EYB and NWP regions in terms of annual mean, with  
867 values up to  $6 \text{ ng m}^{-3}$  (approximately 0.5% of MOA concentration) over these two regions (Figure 8f).  
868 MSOA concentration exhibited the maximum in JJA, with seasonal mean values of  $7 \text{ ng m}^{-3}$  to  $11 \text{ ng}$   
869  $\text{m}^{-3}$  extending from the marginal seas of China (EYB) to remote western North Pacific (NWP) (Figure  
870 8i). MSOA distribution in MAM was similar to that in JJA but with lower mean concentrations (4~7

871  $\text{ng m}^{-3}$ ) over the EYB and NWP regions (Figure 8h). In SON (Figure 8j), MSOA concentrations were  
872  $2\sim 4 \text{ ng m}^{-3}$  in the above two regions. In DJF (Figure 8g), MSOA concentration was lowest, with  
873 values of  $0.4\sim 2 \text{ ng m}^{-3}$  over the marginal seas of China and the southern parts of the model domain.  
874 The maximum seasonal mean MSOA concentration was up to  $14 \text{ ng m}^{-3}$  over oceanic areas of the  
875 EYB to NWP regions in JJA, and the maximum daily mean MSOA value exceeded  $28 \text{ ng m}^{-3}$  on some  
876 days, e.g. June 6~7 (figure not shown). Table 8 shows the domain and seasonal/annual averages of  
877 MSOA over the oceanic regions of concern. The annual mean MSOA concentrations were  $2.2 \text{ ng m}^{-3}$ ,  
878  $4.1 \text{ ng m}^{-3}$  and  $3.8 \text{ ng m}^{-3}$  averaged over the western Pacific, the EYB and NWP regions. It is striking  
879 that the domain average MSOA concentration consistently exhibited a distinct seasonality, with the  
880 maximum in summer and the minimum in winter throughout all the oceanic regions of the western  
881 Pacific, which was resulted from the combined effects of isoprene emission flux and meteorological  
882 conditions. The domain average MSOA concentrations reached the maximums of  $3.9 \text{ ng m}^{-3}$ ,  $7.5 \text{ ng m}^{-3}$ ,  
883  $8.3 \text{ ng m}^{-3}$ , respectively, over the western Pacific, the EYB and NWP regions in JJA. The  
884 seasonality of MSOA concentration over the western Pacific is similar to the simulation result from  
885 Myriokefalitakis et al. (2010). According to Table 8, the annual mean fraction of MSOA in MOA was  
886 estimated to be 0.8%, 0.9% and 0.6%, over the western Pacific, the EYB and NWP regions,  
887 respectively. The maximum and minimum fractions of MSOA in MOA averaged over the western  
888 Pacific occurred in JJA (1.7%) and DJF (0.3%), respectively, with the maximum regional and  
889 seasonal average MSOA fraction up to 3.4% in summer over the EYB region. Based on the  
890 GEOS-Chem model simulation, Arnold et al. (2009) indicated that SOA produced by marine isoprene  
891 contributed only a very small fraction (0.01~1.4%) of the observed organic aerosol mass at remote  
892 marine sites (Amsterdam Island in southern Indian Ocean, Azores and Mace Head islands in northern  
893 Atlantic Ocean). In a global model simulation from Myriokefalitakis et al. (2010), the annual mean  
894 marine isoprene and monoterpene derived SOA concentrations were approximately  $0.4\sim 1 \text{ ng m}^{-3}$   
895 (accounting for  $\sim 0.4\%$  of marine OA) over the western Pacific. Meskhidze et al. (2011) illustrated the  
896 marine SOA from phytoplankton-derived isoprene and monoterpenes contributed  $<10\%$  of surface  
897 OM concentration of marine source in most areas of the western Pacific.

898

### 899 4.3 Direct radiative effect due to MOA

900 In this section, the direct radiative effect (DRE) due to MOA ( $\text{DRE}_{\text{MOA}}$ ) over the western Pacific

901 of East Asia was estimated and analyzed. DRE is defined as the difference in net shortwave radiation  
 902 flux at TOA (or at the surface) induced by aerosols. The DRE of MOA is derived by the difference  
 903 between two radiation calls with aerosol optical properties (aerosol optical depth, single scattering  
 904 albedo, asymmetry factor) due to total aerosols including sea spray (sea salt+MOA), anthropogenic  
 905 aerosols and other natural aerosols (mineral dust, SOA from vegetation emitted VOCs) in this model  
 906 and with those due to total aerosols but MOA (i.e., call two times in the radiation module in the same  
 907 simulation), reflecting an instantaneous change in shortwave radiation fluxes induced by MOA. DRE  
 908 of other aerosol components is estimated by using the same method. The direct (indirect) radiative  
 909 effect is calculated with the following general equation in this study:

$$910 \quad D(I)RE_{ai} = (F\downarrow - F\uparrow)_{all\ aerosols} - (F\downarrow - F\uparrow)_{all\ aerosols\ but\ ai}$$

911 Here  $F\downarrow$  and  $F\uparrow$  represents incoming and outgoing shortwave radiation fluxes, respectively,  $a_i$  denotes  
 912 a specific aerosol component, e.g., MOA.

913 Figures 10a to 10e show the annual and seasonal mean  $DRE_{MOA}$  at TOA under all-sky condition.  
 914 MOA induced negative DRE over the entire western Pacific. Consistent with the spatial distribution of  
 915 MOA concentration, the maximum  $DRE_{MOA}$  up to  $-0.9\ W\ m^{-2}$  occurred over the NWP region (Figure  
 916 10a), with statistical significance at the 99% confidence level over limited areas of NWP (note the  
 917 areas of statistical significance at the 95% confidence level is larger). Over the EYB region, the other  
 918 hotspot of MOA mass concentration, the  $DRE_{MOA}$  was weaker, with an annual mean  $DRE_{MOA}$  of  $\sim$   
 919  $-0.4\ W\ m^{-2}$  (Figure 10a). In terms of domain average, the annual mean  $DRE_{MOA}$  was estimated to be  
 920  $-0.27\ W\ m^{-2}$  over the western Pacific, smaller than that over the NWP ( $-0.5\ W\ m^{-2}$ ) but similar to that  
 921 over the EYB ( $-0.33\ W\ m^{-2}$ ) (Table 10). The weaker  $DRE_{MOA}$  over the EYB than that over the NWP  
 922 could be attributed to both the lower MOA concentration (Table 8) and lower relative humidity (73%  
 923 vs 83%, Table 9) because low relative humidity is unfavorable for hygroscopic growth of MOA,  
 924 leading to a smaller light extinction coefficient and thus weaker DRE. The mean  $DRE_{MOA}$  over the  
 925 western Pacific was largest in spring ( $-0.38\ W\ m^{-2}$ ) and lowest in winter ( $-0.18\ W\ m^{-2}$ ) (Table 10),  
 926 consistent with the seasonality of MOA concentration.

927 In the NWP region, MOA induced the largest DRE up to  $-2.0\ W\ m^{-2}$  to the northeast of Japan in  
 928 MAM (Figure 10c) and followed by that in JJA (up to  $-1.5\ W\ m^{-2}$ ) (Figure 10d) with statistical  
 929 significance at the 99% confidence level, mainly due to higher MOA concentrations in the two  
 930 seasons. The  $DRE_{MOA}$  value was relatively low in SON ( $\sim -0.7\ W\ m^{-2}$ ) (Figure 10e), and it was lowest

931 in DJF, with the maximum of just  $-0.4 \text{ W m}^{-2}$  without statistical significance (Figure 10b) due to the  
932 lowest MOA concentration in winter (Table 8). The regional and seasonal means of  $\text{DRE}_{\text{MOA}}$  over the  
933 NWP were estimated to be  $-0.76 \text{ W m}^{-2}$ ,  $-0.72 \text{ W m}^{-2}$ ,  $-0.32 \text{ W m}^{-2}$ , and  $-0.21 \text{ W m}^{-2}$  in MAM, JJA,  
934 SON, and DJF, respectively (Table 10). On the contrary, in the EYB region,  $\text{DRE}_{\text{MOA}}$  exhibited a  
935 different seasonal trend from that over the NWP, exhibiting the largest DRE in SON (Figure 10e),  
936 moderate DREs in MAM (Figure 10c) and DJF (Figure 10b), and the lowest one in JJA (Figure 10d),  
937 with corresponding mean values of  $-0.38 \text{ W m}^{-2}$ ,  $-0.34 \text{ W m}^{-2}$ ,  $-0.32 \text{ W m}^{-2}$ , and  $-0.26 \text{ W m}^{-2}$ ,  
938 respectively, for the four seasons (Table 10). The different seasonal variation of DRE in EYB and  
939 NWP could be mainly attributed to the different seasonality of MOA concentrations in these two  
940 regions.

941 It is of interest to estimate the relative importance of MOA in directly perturbing solar radiation  
942 compared with other types of aerosols over the western Pacific. Table 10 lists the simulated annual  
943 and seasonal mean DREs due to sea salt and anthropogenic aerosols over the western Pacific and the  
944 regions of NWP and EYB, respectively. It is noted that the annual mean  $\text{DRE}_{\text{MOA}}$  was approximately  
945 one third of that due to sea salt and one order of magnitude lower than that due to anthropogenic  
946 aerosols on average over the western Pacific in this simulation. Over the EYB region,  $\text{DRE}_{\text{MOA}}$  was  
947 almost negligible compared with that by anthropogenic aerosols due to the predominance of  
948 anthropogenic emissions near the continent, however, it is noteworthy that  $\text{DRE}_{\text{MOA}}$  was comparable  
949 in magnitude to the DRE by sea salt, especially in springtime ( $-0.34$  vs  $-0.36 \text{ W m}^{-2}$  in MAM) when  
950 MOA concentration reached its maximum (Table 8). Over the remote oceans (NWP),  $\text{DRE}_{\text{MOA}}$  was  
951 approximately 19% of the DRE by anthropogenic aerosols due to weakened influence of continental  
952 anthropogenic emissions over open oceans, and it was comparable in magnitude to the DRE by sea  
953 salt, especially in summertime ( $0.72$  vs  $0.79 \text{ W m}^{-2}$ ) when sea salt emission flux was lowest (Table 7).  
954 The annual and regional mean all-sky  $\text{DRE}_{\text{MOA}}$  were approximately 10%, 5%, and 19% of the DREs  
955 due to anthropogenic aerosols over the western Pacific, the EYB, and NWP, respectively, and it was  
956 comparable in magnitude to the DREs by sea salt in the EYB and NWP regions.

957 It should be mentioned that due to the much smaller MSOA concentration than MPOA  
958 concentration, the above  $\text{DRE}_{\text{MOA}}$  was dominantly contributed by MPOA, similar to the findings from  
959 previous studies (Arnold et al., 2009; Booge et al., 2016; Li et al., 2019). MPOA also dominated over  
960 MSOA in the following IRE estimation.



961

#### 962 4.4 Indirect radiative effect due to MOA

963 The first indirect effect refers to that aerosol particles tend to increase cloud droplet number  
964 concentration, decrease cloud effective radius under stable cloud liquid water content, and thus  
965 modify cloud optical properties and radiation, which is also denoted as the indirect radiative effect  
966 (IRE) in this study. IRE due to marine organic aerosols ( $IRE_{MOA}$ ) over the western Pacific of East Asia  
967 was explored in this section. The model is able to generally reproduce the major distribution features  
968 of annual mean cloud fraction and cloud optical depth in comparison with MODIS retrievals, which  
969 generally shows higher values from southeast China to the NWP region (Figure not shown). The first  
970 indirect radiative effect or IRE of MOA is estimated by the instantaneous difference in net shortwave  
971 radiation flux at TOA (or at the surface) between two radiation calls with cloud properties (cloud  
972 optical depth, cloud single scattering albedo, cloud asymmetry factor) due to total aerosols and with  
973 those due to total aerosols but MOA in one simulation. The lower bound of cloud droplet number  
974 concentration ( $N_c$ ) is set to  $10/\text{cm}^3$ , which roughly represents  $N_c$  in liquid clouds in clean ocean  
975 conditions, according to satellite observations and global simulations. Hoose et al. (2009) pointed out  
976 that the choice of low bound of  $N_c$  may lead to uncertainties in IRE estimation.

977 Here, the calculation of IRE due to MOA (the sum of MPOA and MSOA) is called the base case  
978 (BASE), from which a series of sensitivity simulations are carried out below.

979  $N_c$  due to aerosol activation is diagnosed by the A-G scheme, then the cloud effective radius  $r_e$  is  
980 calculated as a function of  $N_c$  and cloud liquid water content following the approach of Martin et al.  
981 (1994), and the cloud optical parameters (liquid cloud extinction optical depth, single scatter albedo,  
982 asymmetry factor etc.) are calculated by the scheme of Slingo et al. (1989), finally, shortwave  
983 radiation fluxes are calculated by the CCM3 radiation scheme (Kiehl et al., 1996).

984 The annual and seasonal mean  $IRE_{MOA}$  at TOA are shown in Figure 10f to 10j.  $IRE_{MOA}$  was  
985 negative in the entire domain, resulting from a series of changes in cloud properties induced by MOA,  
986 i.e., an increase in cloud droplet number concentration, a decrease in cloud droplet effective radius, an  
987 increase in cloud optical depth and cloud water path, and consequently more reflection of solar  
988 radiation at TOA. The model simulated cloud properties have been compared against satellite  
989 retrievals in spring 2014 in our previous study (Han et al., 2019), which indicated the model was able  
990 to reasonably reproduce the major features of cloud distribution. It is remarkable that  $IRE_{MOA}$  was

991 stronger than  $DRE_{MOA}$  over the western Pacific, with the maximum annual mean of  $IRE_{MOA}$  being  
992 approximately three times the maximum of  $DRE_{MOA}$ , and the positions of their maximum values were  
993 different. The annual mean  $IRE_{MOA}$  of  $-0.9 \sim -2.5 \text{ W m}^{-2}$  distributed from southwest to northeast over  
994 wide areas of the western Pacific (Figure 10f) with statistical significance at the 99% level over most  
995 of the NWP region. It was evident that the strongest  $IRE_{MOA}$  occurred in spring (MAM), with the  
996 seasonal mean values of  $-1.2 \sim -3.0 \text{ W m}^{-2}$  over vast areas from the East China Sea to the oceans east  
997 of Japan (Figure 10h).  $IRE_{MOA}$  in SON was similar in distribution pattern to that in MAM, with lower  
998 values in the above regions (Figure 10j). The  $IRE_{MOA}$  was weakest in JJA, with the maximum up to  
999  $-1.5 \text{ W m}^{-2}$  over a portion of the western Pacific east of Japan (Figure 10i), whereas the  $IRE_{MOA}$  value  
1000 in DJF was between those in MAM and JJA with a similar distribution pattern.  $IRE_{MOA}$  is statistically  
1001 significant at the 99% level over specific areas of EYB and NWP in all seasons. The seasonal  
1002 variation of  $IRE_{MOA}$  was likely influenced by both the seasonal changes in cloud amount and MOA  
1003 concentration. In terms of domain average, the seasonal mean  $IRE_{MOA}$  was strongest ( $-0.94 \text{ W m}^{-2}$ ) in  
1004 MAM over the western Pacific (Table 10), which was mainly attributed to the maximum MOA  
1005 concentration in spring (Table 8).  $IRE_{MOA}$  was secondly strongest in SON ( $-0.7 \text{ W m}^{-2}$ ) because MOA  
1006 concentration and cloud fraction (Figure S3j) were both high in autumn. The weakest  $IRE_{MOA}$   
1007 occurred in JJA, which was mainly attributed to both the lower MOA concentration and cloud fraction  
1008 in summer (Table 8, Figure S3i). Figure S4 further presents the monthly mean distributions of Chl-a  
1009 concentration, MPOA emission, MOA concentration, and  $IRE_{MOA}$  in April, when Chl-a concentration  
1010 and MPOA emission resulting from phytoplankton were distinctly high in the EYB and NWP regions  
1011 (Figure S4a and S4b). It can be found that MOA was transported from the high Chl-a regions to the  
1012 southeast under northwesterly winds over the oceans (Figure S4c), resulting in an elevated  $IRE_{MOA}$  up  
1013 to  $-5 \text{ W m}^{-2}$  over the western Pacific east of Japan (Figure S4d). Previous studies were very limited to  
1014 compare with. Meskhidze and Nenes (2006) estimated based on satellite retrievals a reduction of  $15$   
1015  $\text{W m}^{-2}$  in shortwave radiation at TOA due to changes in cloud properties during a strong  
1016 phytoplankton bloom event near South Georgia Island in the Southern Ocean in summertime.

1017 In terms of annual and regional mean,  $IRE_{MOA}$  was estimated to be  $-0.66 \text{ W m}^{-2}$  for the western  
1018 Pacific,  $-0.23 \text{ W m}^{-2}$  over the EYB region, and  $-1.0 \text{ W m}^{-2}$  over the NWP region, respectively (Table  
1019 10). There was an apparent seasonality in the  $IRE_{MOA}$ , with the maximum of  $-0.94 \text{ W m}^{-2}$  in MAM  
1020 and the minimum of  $-0.36 \text{ W m}^{-2}$  in JJA over the western Pacific (Table 10). However, the seasonality

1021 of  $IRE_{MOA}$  in the EYB and NWP regions are different from that over the western Pacific. Over the  
1022 EYB, the estimated  $IRE_{MOA}$  reached its maximum ( $-0.38 \text{ W m}^{-2}$ ) in SON, which was due to the  
1023 combined effect of a moderately high MOA concentration (Table 8) and the maximum cloud fraction  
1024 (Figure S3j) in this region. Although MOA concentration reached the maximum in MAM, there was a  
1025 minimum total cloud fraction in spring among seasons (Figure S3h), leading to a moderate  $IRE_{MOA}$ .  
1026 Over the NWP region,  $IRE_{MOA}$  in DJF ( $-0.57 \text{ W m}^{-2}$ ) was smaller than those in other seasons ( $-0.78 \sim$   
1027  $-1.4 \text{ W m}^{-2}$ ), which was mainly attributed to the lowest MOA concentration in winter (Table 8),  
1028 although cloud fraction was highest among seasons in this region (Figure S3g).

1029 The relative importance of MOA in the aerosol indirect radiative effect over the western Pacific  
1030 was investigated by comparing the  $IRE_{MOA}$  with the IREs induced by sea salt and anthropogenic  
1031 aerosols. In terms of annual and oceanic average, the IREs by sea salt and anthropogenic aerosols  
1032 were estimated to be  $-0.41 \text{ W m}^{-2}$  and  $-7.7 \text{ W m}^{-2}$  (Table 10), respectively, indicating  $IRE_{MOA}$  ( $-0.66 \text{ W}$   
1033  $\text{m}^{-2}$ ) was larger than the IRE by sea salt and was approximately 9% of that by anthropogenic aerosols.  
1034 Although the hygroscopicity of sea salt is larger than that of MOA, because the mean radius of MOA  
1035 ( $0.05 \mu\text{m}$ ) is smaller than the radius of sea salt ( $0.1 \mu\text{m}$  for fine mode), the number concentration of  
1036 MOA is larger than that of sea salt, leading to larger activated cloud droplet number concentration and  
1037 IRE by MOA than those by sea salt. It is noted that the relative magnitude of  $IRE_{MOA}$  compared with  
1038 the IRE by anthropogenic aerosols was reduced (5%) in the EYB and enhanced over the NWP (12%),  
1039 because anthropogenic aerosols from the East Asian continent dominated aerosol magnitude in the  
1040 marginal seas of China. In terms of seasonal and domain average over the western Pacific,  $IRE_{MOA}$   
1041 was approximately 1.6 times the IRE by sea salt, and approximately 10% of the IRE by anthropogenic  
1042 aerosols in MAM. In summer (JJA) when both sea salt and anthropogenic aerosol concentrations were  
1043 lowest,  $IRE_{MOA}$  was similar in magnitude to the IRE by sea salt, and about 8% of the IRE by  
1044 anthropogenic aerosols. In the EYB region,  $IRE_{MOA}$  was just about 5% of the IRE by anthropogenic  
1045 aerosols, but approximately three times the IRE by sea salt. In the NWP,  $IRE_{MOA}$  was about 12% of  
1046 the IRE by anthropogenic aerosols in terms of annual mean, whereas in autumn when anthropogenic  
1047 aerosol level was relatively low,  $IRE_{MOA}$  was as high as the maximum in MAM and approximately 18%  
1048 of the IRE by anthropogenic aerosols. The above model estimation demonstrates that  $IRE_{MOA}$  was  
1049 generally stronger than the IRE due to sea salt over the western Pacific, and approximately 6~18% of  
1050 the IRE due to anthropogenic aerosols in the four seasons over the NWP, suggesting an important role

1051 of MOA in perturbing radiation transfer through modifying cloud properties over the western Pacific  
1052 Ocean of East Asia. The estimated IRE due to MSOA accounted for approximately 1% of the annual  
1053 mean  $IRE_{MOA}$  averaged over the western Pacific, consistent with the very low proportion of MSOA in  
1054 the MOA mass concentration (Table 8). Overall, MSOA plays a minor role in perturbing cloud  
1055 properties and shortwave radiation compared with MPOA.

1056 To address potential uncertainty in the estimated  $IRE_{MOA}$  due to limited knowledge on MOA  
1057 properties, three additional sensitivity simulations from the base case (results shown in Figure S5 and  
1058 Table S4) were carried out regarding particle size, solubility, and molecule weight, which are crucial  
1059 to aerosol activation (note we focus on MPOA due to its dominant fraction in MOA as shown above).  
1060 The first sensitivity simulation (SENS1) assumes a smaller geometric mean radius ( $0.02\mu\text{m}$  instead of  
1061  $0.05\mu\text{m}$  in the base case) for MPOA, resulting in a weaker domain-annual mean  $IRE_{MOA}$  ( $-0.53\text{ W m}^{-2}$ )  
1062 than that in the base case ( $-0.66\text{ W m}^{-2}$ ) over the oceanic region (Figure S5b, Table S4). The second  
1063 sensitivity simulation (SENS2) assigns a lower solubility (0.05) with relatively large molecule weight  
1064 ( $146\text{ g mol}^{-1}$ ) for MPOA (which is similar to the properties of adipic acid, Huff Hartz et al., 2006;  
1065 Miyazaki et al., 2010) instead of the slight solubility (0.1) with a smaller molecule weight ( $90\text{ g mol}^{-1}$ )  
1066 (which is similar to the properties of oxalic acid, Roelofs, 2008; Miyazaki et al., 2010) in the base case,  
1067 in this case, the  $IRE_{MOA}$  reduces to  $-0.2\text{ W m}^{-2}$  (Figure S5c, Table S4). The third simulation (SENS3)  
1068 combines the above two cases, assuming a smaller geometric mean radius as in SENS1 together with  
1069 the lower solubility and larger molecule weight as in SENS2, it produces a further reduced  $IRE_{MOA}$  of  
1070  $-0.14\text{ W m}^{-2}$  (Figure S5d, Table S4). The above sensitivity simulations exhibit a high sensitivity of  
1071  $IRE_{MOA}$  to the MPOA properties.

1072 It is interesting to note that Quinn et al. (2017) indicated that sea spray aerosol generally makes a  
1073 contribution of less than 30% to CCN population at supersaturation of 0.1 to 1.0% on a global basis  
1074 based on measurements onboard seven research cruises over the Pacific, Southern, Arctic and Atlantic  
1075 oceans. However, their cruise tracks did not cover the western Pacific Ocean. The results from this  
1076 study exhibits the annual and oceanic mean contribution of sea spray aerosol (the sum of MOA and  
1077 sea salt) to the IRE by total aerosols was approximately 12% in the base case over the western Pacific  
1078 region (Table 10). This percentage contribution increases to 19% in the NWP in autumn due to the  
1079 highest  $IRE_{MOA}$  and lower IRE by anthropogenic aerosols in autumn (Table 10).

1080 The western Pacific Ocean is just downwind of the East Asian continent, which have large

1081 amounts of anthropogenic aerosols, mineral dust, and nutrients inputs to the marginal seas of China  
1082 from the Yangtze and Yellow Rivers, could be very different from remote clean oceans in the world.  
1083 Although some cruise measurements have been carried out and a few knowledges on MOA properties  
1084 (e.g., size distribution) was gained, the cruise measurement and observational analysis for MOA  
1085 chemical properties are so far almost absent in the western Pacific of East Asia. Therefore, marine  
1086 biogeochemistry, marine aerosol sources and properties, as well as their potentials to be CCN and  
1087 impacts on radiation, cloud, and precipitation deserve further investigation in the future.

1088

#### 1089 4.5 Indirect radiative effect of MOA due to aerosol-radiation and aerosol-cloud interactions

1090 IPCC AR5 proposes the concept of effective radiative forcing (ERF), which is defined as the  
1091 change in net radiative flux at TOA after rapid adjustment of the atmosphere (including atmospheric  
1092 temperature, water vapor, cloud, circulation) to radiation perturbation with prescribed SST and sea ice,  
1093 which can better represent climate response to perturbation of forcing factors (Boucher et al., 2013).

1094 Similarly, this study also estimates the direct radiative effect of MOA due to aerosol-radiation  
1095 interaction (denoted as  $DRE_{ari}$  hereinafter) and the indirect radiative effect of MOA due to  
1096 aerosol-cloud interaction (denoted as  $IRE_{aci}$  hereinafter). Note the method for calculating  $DRE_{ari}$  and  
1097  $IRE_{aci}$  is different from that for  $DRE_{MOA}$  and  $IRE_{MOA}$ .  $DRE_{ari}$  is derived by the difference in shortwave  
1098 radiation flux at TOA induced by MOA scattering between two simulations with aerosol optical  
1099 properties due to total aerosols and with those due to total aerosols but MOA, while the perturbation  
1100 of MOA to cloud properties is turned off.  $IRE_{aci}$  is calculated by the difference in shortwave radiation  
1101 flux at TOA induced by MOA perturbation to cloud albedo between two simulations with cloud  
1102 properties due to total aerosols and with those due to total aerosols but MOA, while the direct  
1103 perturbation of MOA to radiation is closed. The calculation of  $DRE_{ari}$  and  $IRE_{aci}$  considers the  
1104 adjustment of atmospheric temperature, water vapor, and cloud (including cloud microphysical and  
1105 lifetime change, i.e., the second indirect effect) to the MOA-induced radiation perturbation in the two  
1106 simulations.

1107 Figure 11a shows the annual mean MOA  $DRE_{ari}$  in the study domain. The distribution of MOA  
1108  $DRE_{ari}$  was similar in distribution to that of  $DRE_{MOA}$  without statistical significance at the 99% level.  
1109 It is noted that  $DRE_{ari}$  was not consistently negative in the domain, the effect of atmospheric  
1110 adjustment on radiation can be seen in some locations over the Ocean and the continent, with a few

1111 positive values of 0.1~0.3 W m<sup>-2</sup>.

1112 Figure 11b shows the annual mean MOA IRE<sub>aci</sub> in the study domain, which was similar in  
1113 magnitude and distribution pattern to IRE<sub>MOA</sub> (Figure 10f), but it distributed unevenly in the domain  
1114 with some positive values exceeding 0.2 W m<sup>-2</sup> over both the Western Pacific and the continent. IRE<sub>aci</sub>  
1115 of MOA was statistically significant at the 99% confidence level over the NWP region. The  
1116 atmospheric response and adjustment induced by IRE<sub>MOA</sub> could be somewhat stronger than that by  
1117 DRE<sub>MOA</sub>, with respect to the positive values of IRE<sub>aci</sub> up to 0.3 W m<sup>-2</sup> over the continent. The small  
1118 positive values could be associated with the radiative feedback and atmospheric and cloud adjustment.

1119 The annual and domain average of DRE<sub>ari</sub> and IRE<sub>aci</sub> over the western Pacific are estimated to be  
1120 -0.25 W m<sup>-2</sup> and -0.61 W m<sup>-2</sup>, respectively, both are somewhat weaker than the DRE<sub>MOA</sub> (-0.27 W m<sup>-2</sup>)  
1121 and IRE<sub>MOA</sub> (-0.66 W m<sup>-2</sup>).

1122

## 1123 5. Conclusions.

1124 The organic aerosols of marine origin over the western Pacific Ocean of East Asia was  
1125 investigated by an online-coupled regional climate-chemistry model RIEMS-Chem for the year 2014.  
1126 Emissions and relevant processes of marine MPOA, isoprene and monoterpene were incorporated into  
1127 RIEMS-Chem. A wide variety of observational datasets from EANET, CNEMC and AERONET  
1128 networks, cruise measurements and previous publications were collected for model validation. The  
1129 modeled SOA from marine VOC sources was also compared with secondary organic tracers measured  
1130 by research cruise. The model performed well for PM<sub>2.5</sub> and PM<sub>10</sub> in marine environment, producing  
1131 overall correlation coefficients and NMBs of 0.61/0.70 and 12%/-7% for PM<sub>2.5</sub> concentration,  
1132 0.65/0.65 and -5%/-1% for PM<sub>10</sub> concentration at the EANET/CNEMC sites, respectively. The model  
1133 reasonably reproduced the spatial distribution and temporal variation of BC and OC concentrations  
1134 along cruise tracks and at islands over the west Pacific, with the correlation coefficients and NMBs  
1135 being 0.6~0.75 and -28%~3% for OC, respectively. The modeled OC concentration was apparently  
1136 improved while taking into account marine organic aerosols. The model results clearly showed an  
1137 increasing contribution of marine organic aerosols to total OC mass concentration from the marginal  
1138 seas of China to remote oceans. Organic aerosol mass of marine origin was dominated by MPOA  
1139 because MSOA produced by marine isoprene and monoterpene emissions was about 1~2 orders of  
1140 magnitude lower than MPOA. The model simulates AOD reasonably well at the 7 coastal or island

1141 AERONET sites, with an overall correlation coefficient of 0.54 and an NMB of -6%.

1142 High MPOA emission mainly occurred over the marginal seas of China (EYB) and the northern  
1143 parts of western Pacific northeast of Japan (NWP). For the western Pacific, MPOA emission reached  
1144 the maximum in SON, followed by those in DJF and MAM, and the minimum in JJA, with an annual  
1145 and domain average emission rate of  $0.16 \times 10^{-2} \mu\text{g m}^{-2} \text{ s}^{-1}$ . The combination of Chl-a concentration and  
1146 sea salt emission flux determined the seasonality of MPOA emission. The annual MPOA emission for  
1147 the year 2014 was estimated to be  $0.78 \text{ Tg yr}^{-1}$  over the western Pacific.

1148 Consistent with the distribution pattern MPOA emission, high MOA concentration mainly  
1149 distributed over the EYB and NWP, with an annual and domain mean concentration of  $0.27 \mu\text{g m}^{-3}$ ,  
1150  $0.48 \mu\text{g m}^{-3}$  and  $0.59 \mu\text{g m}^{-3}$ , over the western Pacific, the EYB and NWP regions, respectively. MOA  
1151 concentration was highest in MAM and lowest in DJF, with the seasonal and domain mean values of  
1152  $0.37 \mu\text{g m}^{-3}$  and  $0.21 \mu\text{g m}^{-3}$ , respectively, over the western Pacific. The seasonality of MOA  
1153 concentration was determined by the combined effect of MPOA emission, dry and wet depositions.

1154 On average, the annual mean percentage contribution of MOA to total OA mass was 26% over the  
1155 western Pacific, with the largest seasonal mean contribution of 32% in SON and the lower ones in  
1156 DJF (24%) and JJA (23%). Over the NWP, the domain average contribution of MOA to OA could be  
1157 as high as 42% in terms of annual mean and approaching 52% in MAM; however, over the EYB, the  
1158 annual mean contribution was just 13% and the percentage contribution was even reduced to 6% in  
1159 JJA. This indicated that the relative importance of MOA in total OA concentration increased with the  
1160 distance away from the East Asian continent. MSOA concentration was approximately 1~2 orders of  
1161 magnitude lower than MPOA, with the simulated annual and regional mean MSOA being  $2.2 \text{ ng m}^{-3}$   
1162 and the maximum daily mean value up to  $28 \text{ ng m}^{-3}$  in summer over the western Pacific.

1163 An annual/oceanic mean all-sky  $\text{DRE}_{\text{MOA}}$  of  $-0.27 \text{ W m}^{-2}$  at TOA was estimated over the western  
1164 Pacific, which was about 40% of the  $\text{IRE}_{\text{MOA}}$  ( $-0.66 \text{ W m}^{-2}$ ). The domain mean  $\text{IRE}_{\text{MOA}}$  was strongest  
1165 in spring ( $-0.94 \text{ W m}^{-2}$ ) and weakest in summer ( $-0.36 \text{ W m}^{-2}$ ) over the western Pacific, and the  
1166 monthly mean  $\text{IRE}_{\text{MOA}}$  can reach  $-5 \text{ W m}^{-2}$  in the NWP region east of Japan in April. The changes in  
1167 MOA concentration and cloud amount both contributed to the seasonality of  $\text{IRE}_{\text{MOA}}$ . In terms of  
1168 annual and oceanic mean over the western Pacific, MSOA just contributed approximately 1% of the  
1169  $\text{IRE}_{\text{MOA}}$ .  $\text{IRE}_{\text{MOA}}$  was generally larger than the IRE due to sea salt on average. The annual and oceanic  
1170 mean IRE due to sea spray aerosols (MOA + sea salt) was approximately 12% of that due to total

1171 aerosols over the western Pacific, but this ratio can increase up to 19% in autumn in the NWP region.  
1172 The estimation of  $IRE_{MOA}$  was sensitive to MOA properties, which decreased apparently while a  
1173 smaller geometric mean radius together with a lower solubility and a larger molecule weight were  
1174 assigned for MOA. Overall, the indirect radiative effect of MOA was larger than the direct radiative  
1175 effect, and had a nonnegligible impact on radiation budget and cloud over the western Pacific. The  
1176 direct and indirect radiative effect considering atmospheric feedback and adjustment were estimated  
1177 as well, which was similar in magnitude to the  $DRE_{MOA}$  and  $IRE_{MOA}$ , with a few positive changes in  
1178 shortwave radiation fluxes in some locations.

1179 While this study presents new insights into the seasonal variation and annual means of emissions,  
1180 concentrations, and radiative effects of MOA in the western Pacific, it is still subject to some  
1181 uncertainties as follows: 1.) the properties of marine organic aerosols, including size distribution,  
1182 molecular weight, solubility, surfactant amount etc. are still poorly characterized, which are crucial to  
1183 aerosol activation, dry deposition, and wet scavenging; 2.) the sources and chemical formation  
1184 processes of marine organic aerosols including secondary organics are highly complex, and poorly  
1185 understood and represented in the model; 3.) the indirect effects of MOA in this study is for warm  
1186 stratiform cloud. Further research on MOA sources, properties, chemical processes, and climatic  
1187 impacts will be conducted together with the advances in both field experiments (integrated cruise,  
1188 aircraft and satellite observations) and model development in the future.

1189

#### 1190 **Author Contributions.**

1191 ZH designed the study, JL and ZH developed the model, processed and analyzed the model results, JL  
1192 performed the model simulation, ZH and JL wrote the paper, PF and XY provided and analyzed the  
1193 cruise measurement data, ML analyzed the model results.

1194

#### 1195 **Data availability.**

1196 The observational data can be accessed through contacting the corresponding author.

1197

#### 1198 **Competing interests.**

1199 The authors declare that they have no conflict of interests.

1200



1201 **Acknowledgement.**

1202 This study was supported by the National Key R&D Program of China (No. 2019YFA0606802),  
1203 the National Natural Science Foundation of China (No. 42275118, No. 42375107), the Jiangsu  
1204 Collaborative Innovation Center for Climate Change. The authors appreciate the science teams of  
1205 EANET, CNEMC, AERONET, MODIS and VIIRS for their works in data maintenance.

1206

1207 **Reference**

1208 Abdul-Razzak, H., Ghan, S.J., and Rivera-Carpio, C.: A parameterization of aerosol activation: 1.  
1209 Single aerosol type, *J. Geophys. Res.*, 103, 6123–6131, 1998.

1210 Abdul-Razzak, H. and Ghan, S.J.: Parameterization of the influence of organic surfactants on aerosol  
1211 activation, *J. Geophys. Res.*, 109, D03205, doi:10.1029/2003JD004043, 2004.

1212 Arnold, S.R., Spracklen, D.V., Williams, J., Yassaa, N., Sciare, J., Bonsang, B., Gros, V., Peeken, I.,  
1213 Lewis, A.C., Alvaïn, S., and Moulin, C.: Evaluation of the global oceanic isoprene source and its  
1214 impacts on marine organic carbon aerosol, *Atmos. Chem. Phys.*, 9, 1253–1262.  
1215 <https://doi.org/10.5194/acp-9-1253-2009>, 2009.

1216 Bates, T. S., Quinn, P. K., Coffman, D. J., Johnson, J. E., Upchurch, L., Saliba, G., Lewis, S., Graff, J.,  
1217 Russell, L. M., and Behrenfeld, M. J.: Variability in Marine Plankton Ecosystems Are Not  
1218 Observed in Freshly Emitted Sea Spray Aerosol Over the North Atlantic Ocean, *Geophys. Res.  
1219 Lett.*, 47, e2019GL085938. <https://doi.org/10.1029/2019GL085938>, 2020.

1220 Beheng, K.D.: A parameterization of warm cloud microphysical conversion processes. *Atmos. Res.*,  
1221 33, 193–206. [https://doi.org/10.1016/0169-8095\(94\)90020-5](https://doi.org/10.1016/0169-8095(94)90020-5), 1994.

1222 Bertram, T.H., Cochran, R.E., Grassian, V.H., and Stone, E.A.: Sea spray aerosol chemical  
1223 composition: elemental and molecular mimics for laboratory studies of heterogeneous and  
1224 multiphase reactions, *Chem. Soc. Rev.*, 47(7), 2374-2400, DOI: 10.1039/c7cs00008a, 2018.

1225 Bennartz, R.: Global assessment of marine boundary layer cloud droplet number concentration from  
1226 satellite, *J. Geophys. Res.*, 112, D02201, doi:10.1029/2006jd007547, 2007.

1227 Bikkina, S., Kawamura, K., Miyazaki, Y., and Fu, P.: High abundances of oxalic, azelaic and glyoxylic  
1228 acids and methylglyoxal in the open ocean with high biological activity: Implication for  
1229 secondary OA formation from isoprene, *Geophys. Res. Lett.*, 41, 3649–3657, 2014.

1230 Booge, D., Marandino, C.A., Schlundt, C., Palmer, P.I., Schlundt, M., Atlas, E.L., Bracher, A.,

1231 Saltzman, E.S., and Wallace, D.W.R.: Can simple models predict large-scale surface ocean  
1232 isoprene concentrations? *Atmos. Chem. Phys.*, 16, 11807–11821,  
1233 <https://doi.org/10.5194/acp-16-11807-2016>, 2016.

1234 Boreddy, S.K.R., Haque, M.M., and Kawamura, K.: Long-term (2001–2012) trends of carbonaceous  
1235 aerosols from a remote island in the western North Pacific: an outflow region of Asian pollutants,  
1236 *Atmos. Chem. Phys.*, 18, 1291–1306, <https://doi.org/10.5194/acp-18-1291-2018>, 2018.

1237 Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., Kerminen, V.-M., Kondo,  
1238 Y., Liao, H., Lohmann, U., Rasch, P., Satheesh, S.K., Sherwood, S., Stevens, B., and Zhang, X.Y.:  
1239 Clouds and Aerosols. In: *Climate Change 2013: The Physical Science Basis. Contribution of*  
1240 *Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate*  
1241 *Change* (pp. 616–632) [Stocker, T.F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S.K., Boschung,  
1242 J., Nauels, A., Xia, Y., Bex, V., and Midgley, P.M. (eds.)]. Cambridge University Press,  
1243 Cambridge, United Kingdom and New York, NY, USA, 2013.

1244 Brüggemann, M., Hayeck, N. and George, C.: Interfacial photochemistry at the ocean surface is a  
1245 global source of organic vapors and aerosols, *Nat. Commun.*, 9, 2101.  
1246 <https://doi.org/10.1038/s41467-018-04528-7>, 2018.

1247 Burrows, S.M., Ogunro, O., Frossard, A.A., Russell, L.M., Rasch, P.J., and Elliott, S.M.: A physically  
1248 based framework for modeling the organic fractionation of sea spray aerosol from bubble film  
1249 Langmuir equilibria, *Atmos. Chem. Phys.*, 14, 13601–13629,  
1250 <https://doi.org/10.5194/acp-14-13601-2014>, 2014.

1251 Burrows, S.M., Easter, R.C., Liu, X., Ma, P.-L., Wang, H., Elliott, S.M., Singh, B., Zhang, K., and  
1252 Rasch, P.J.: OCEANFILMS (Organic Compounds from Ecosystems to Aerosols: Natural Films  
1253 and Interfaces via Langmuir Molecular Surfactants) sea spray organic aerosol emissions –  
1254 implementation in a global climate model and impacts on clouds, *Atmos. Chem. Phys.*, 22, 5223–  
1255 5251, <https://doi.org/10.5194/acp-22-5223-2022>, 2022.

1256 Calil, P.H.R., Doney, S.C., Yumimoto, K., Eguchi, K., and Takemura, T.: Episodic upwelling and dust  
1257 deposition as bloom triggers in low-nutrient, low-chlorophyll regions, *J. Geophys. Res.: Oceans*,  
1258 116, C06030, <http://dx.doi.org/10.1029/2010JC006704>, 2011.

1259 Chang, J.S., Brost, R.A., Isaksen, I.S.A., Madronich, S., Middleton, P., Stockwell, W.R., Walcek, C.J.:  
1260 A three-dimensional Eulerian acid deposition model: physical concepts and formulation, *J.*

- 1261 Geophys. Res., 92(D12), 14681-14700, doi:10.1029/JD092iD12p14681, 1987.
- 1262 Conte, L., Szopa, S., Aumont, O., Gros, V., and Bopp, L.: Sources and sinks of isoprene in the global  
1263 open ocean: Simulated patterns and emissions to the atmosphere, *J. Geophys. Res: Oceans*, 125,  
1264 e2019JC015946. <https://doi.org/10.1029/2019JC015946>, 2020.
- 1265 Curci, G., Hogrefe, C., Bianconi, R., Im, U., Balzarini, A., Baró, R., Brunner, D., Forkel, R., Giordano,  
1266 L., Hirtl, M., Honzak, L., Jiménez-Guerrero, P., Knote, C., Langer, M., Makar, P.A., Pirovano, G.,  
1267 Pérez, J.L., San José, R., Syrakov, D., Tuccella, P., Werhahn, J., Wolke, R., Žabkar, R., Zhang, J.,  
1268 and Galmarini, S.: Uncertainties of simulated aerosol optical properties induced by assumptions  
1269 on aerosol physical and chemical properties: An AQMEII-2 perspective, *Atmos. Environ.*, 115,  
1270 541–552, 2015.
- 1271 Dickinson, R.E., Henderson-Sellers, A., Kennedy, P.J.: Biosphere-Atmosphere Transfer Scheme  
1272 (BATS) Version 1e as coupled to NCAR Community Climate Model, NCAR Technical Note,  
1273 NCAR/TN-387+STR, p. 72, 1993.
- 1274 Emmons, L.K., Walters, S., Hess, P.G., Lamarque, J.-F., Pfister, G.G., Fillmore, D., Granier, C.,  
1275 Guenther, A., Kinnison, D., Laepple, T., Orlando, J., Tie, X., Tyndall, G., Wiedinmyer, C.,  
1276 Baughcum, S.L., and Kloster, S.: Description and evaluation of the Model for Ozone and Related  
1277 chemical Tracers, version 4 (MOZART-4), *Geosci. Model Dev.*, 3, 43-67,  
1278 doi:10.5194/gmd-3-43-2010, 2010.
- 1279 Facchini, M.C., Rinaldi, M., Decesari, S., Carbone, C., Finessi, E., Mircea, M., Fuzzi, S., Ceburnis, D.,  
1280 Flanagan, R., Nilsson, E.D., de Leeuw, G., Martino, M., Woeltjen, J., and O’Dowd, C.D.:  
1281 Primary submicron marine aerosol dominated by insoluble organic colloids and aggregates,  
1282 *Geophys. Res. Lett.*, 35, L17814, doi:10.1029/2008GL034210, 2008.
- 1283 Feng, L.M., Shen, H.Q., Zhu, Y.J., Gao, H.W., and Yao, X.H.: Insight into Generation and Evolution  
1284 of Sea-Salt Aerosols from Field Measurements in Diversified Marine and Coastal Atmospheres,  
1285 *Sci. Rep.*, 7, 41260; doi: 10.1038/srep41260, 2017.
- 1286 Fountoukis, C. and Nenes, A.: ISORROPIA II: a computationally efficient thermodynamic  
1287 equilibrium model for  $K^+Ca^{2+}Mg^{2+}NH_4^+Na^+SO_4^{2-}NO_3^-Cl^-H_2O$  aerosols, *Atmos. Chem.*  
1288 *Phys.*, 7, 4639-4659, 2007.
- 1289 Fröhlich-Nowoisky J., Kampf, C.J., Weber, B., Huffman, J.A., Pöhlker, C., Andreae, M.O., Lang-Yona,  
1290 N., Burrows, S.M., Gunthe, S.S., Elbert, W., Su, H., Hoor, P., Thines, E., Hoffmann, T., Després,

1291 V.R., and Pöschl, U.: Bioaerosols in the Earth system: Climate, health, and ecosystem  
1292 interactions, *Atmos. Res.*, 182, 346-376, 2016.

1293 Fu, C. B., Wang, S.Y., Xiong, Z., Gutowski, W. J., Lee, D., McGregor, J. L., Sato, Y., Kato, H., Kim, J.,  
1294 and Suh, M.: Regional climate model intercomparison project for Asia, *Bull. Amer. Meteor. Soc.*,  
1295 86, 257-266, 2005.

1296 Fu, P., Kawamura, K., and Miura, K.: Molecular characterization of marine organic aerosols collected  
1297 during a round-the-world cruise, *J. Geophys. Res.*, 116, D13302, doi:10.1029/2011JD015604,  
1298 2011.

1299 Gantt, B., Meskhidze, N., and Kamykowski, D.: A new physically-based quantification of marine  
1300 isoprene and primary organic aerosol emissions, *Atmos. Chem. Phys.*, 9, 4915-4927,  
1301 <https://doi.org/10.5194/acp-9-4915-2009>, 2009.

1302 Gantt, B., Meskhidze, N., Facchini, M.C., Rinaldi, M., Ceburnis, D., and O'Dowd, C.D.: Wind speed  
1303 dependent size-resolved parameterization for the organic mass fraction of sea spray aerosol,  
1304 *Atmos. Chem. Phys.*, 11, 8777-8790, <https://doi.org/10.5194/acp-11-8777-2011>, 2011.

1305 Gantt, B., Johnson, M.S., Meskhidze, N., Sciare, J., Ovadnevaite, J., Ceburnis, D., and O'Dowd, C.D.:  
1306 Model evaluation of marine primary organic aerosol emission schemes, *Atmos. Chem. Phys.*, 12,  
1307 8553-8566, <https://doi.org/10.5194/acp-12-8553-2012>, 2012a.

1308 Gantt, B., Xu, J., Meskhidze, N., Zhang, Y., Nenes, A., Ghan, S.J., Liu, X., Easter, R., and Zaveri, R.:  
1309 Global distribution and climate forcing of marine organic aerosol – Part 2: Effects on cloud  
1310 properties and radiative forcing, *Atmos. Chem. Phys.*, 12, 6555-6563,  
1311 <https://doi.org/10.5194/acp-12-6555-2012>, 2012b.

1312 Gantt, B. and Meskhidze, N.: The physical and chemical characteristics of marine primary organic  
1313 aerosol: a review, *Atmos. Chem. Phys.*, 13, 3979–3996, 2013.

1314 Gao, M., Han, Z., Liu, Z., Li, M., Xin, J., Tao, Z., Li, J., Kang, J.-E., Huang, K., Dong, X., Zhuang, B.,  
1315 Li, S., Ge, B., Wu, Q., Cheng, Y., Wang, Y., Lee, H.-J., Kim, C.-H., Fu, J.S., Wang, T., Chin, M.,  
1316 Woo, J.-H., Zhang, Q., Wang, Z., and Carmichael, G.R.: Air quality and climate change, Topic 3  
1317 of the Model Inter-Comparison Study for Asia Phase III (MICS-Asia III) – Part 1: Overview and  
1318 model evaluation, *Atmos. Chem. Phys.*, 18, 4859-4884, 2018.

1319 Gery, M.W., Whitten, G.Z., Killus, J.P., Dodge, M.C.: A photochemical kinetics mechanism for urban  
1320 and regional scale computer modeling, *J. Geophys. Res.*, 94, 12925-12956, 1989.

- 1321 Ghan, S.J., Leung, L.R., Easter, R.C., Abdul-Razzak, K.: Prediction of cloud droplet number in a  
1322 general circulation model, *J. Geophys. Res.*, 102 (D18), 21,777–21,794, 1997.
- 1323 Ghan, S. and Zaveri R.A.: Parameterization of optical properties for hydrated internally mixed aerosol,  
1324 *J. Geophys. Res.*, 112, D10201, doi:10.1029/2006JD007927, 2007.
- 1325 Giglio, L., Randerson, J.T., and van der Werf, G.R.: Analysis of daily, monthly, and annual burned  
1326 area using the fourth generation Global Fire Emissions Database (GFED4), *J. Geophys. Res.:*  
1327 *Biogeosciences*, doi:10.1002/jgrg.20042, 2013.
- 1328 Gong, S.L., Barrie, L.A., and Blanchet, J.-P.: Modeling sea-salt aerosols in the atmosphere 1. Model  
1329 development, *J. Geophys. Res.*, 102(D3), 3805-3818, doi:10.1029/96JD02953, 1997.
- 1330 Gong, S.L., Zhang, X.Y., Zhao, T., Mckendry, I.G., Jaffe, D.A., and Lu, N.: Characterization of soil  
1331 dust aerosol in China and its transport and distribution during 2001 ACE-Asia: 2. Model  
1332 simulation and validation, *J. Geophys. Res.*, 108(D9), 4262, doi:10.1029/2002JD002633, 2003.
- 1333 Gong, S.L.: A parameterization of sea-salt aerosol source function for sub- and super-micron particles,  
1334 *Global Biogeochem. Cy.*, 17(4), 1097, doi:10.1029/2003GB002079, 2003.
- 1335 Graf, H.-F., Feichter, J., and Langmann, B.: Volcanic sulfur emissions: Estimates of source strength  
1336 and its contribution to the global sulfate distribution. *J. Geophys. Res.*, 102(D9), 10727-10738,  
1337 doi: 10.1029/96JD03265, 1997.
- 1338 Granier, C., Darras, S., Denier van der Gon, H., Doubalova, J., Elguindi, N., Galle, B., Gauss, M.,  
1339 Guevara, M., Jalkanen, J.-P., Kuenen, J., Liousse, C., Quack, B., Simpson, D., and Sindelarova,  
1340 K.: The Copernicus Atmosphere Monitoring Service global and regional emissions (April 2019  
1341 version), Report April 2019 version, doi:10.24380/d0bn-kx16, 2019.
- 1342 Grell, G.A.: Prognostic evaluation of assumptions used by cumulus parameterizations, *Mon. Wea. Rev.*  
1343 121, 764-787, 1993.
- 1344 Grell, G.A., Dudhia, J., Stauffer, D.R.: A Description of the Fifth-Generation Penn State/NCAR  
1345 Mesoscale Model (MM5), NCAR Technical Note, NCAR/TN-398tSTR, p.117, 1995.
- 1346 Guo, T., Guo, Z., Wang, J., Feng, J., Gao, H., and Yao, X.: Tracer-based investigation of organic  
1347 aerosols in marine atmospheres from marginal seas of China to the northwest Pacific Ocean,  
1348 *Atmos. Chem. Phys.*, 20, 5055–5070, <https://doi.org/10.5194/acp-20-5055-2020>, 2020.
- 1349 Han, X., Zhang, M.G., Han, Z.W., Xin, J.Y., Liu, X.H.: Simulation of aerosol direct radiative forcing  
1350 with RAMS-CMAQ in East Asia, *Atmos. Environ.*, 45, 6576-6592, 2011.

- 1351 Han, Z.: Direct radiative effect of aerosols over East Asia with a regional coupled climate/chemistry  
1352 model, *Meteor. Zeit.*, 19, 287–298, 2010.
- 1353 Han, Z.W., Ueda, H., Matsuda, K., Zhang, R.J., Arao, K., Kanai, Y., Hasome, H.: Model study on  
1354 particle size segregation and deposition during Asian dust events in March 2002, *J. Geophys.*  
1355 *Res.*, 109, D19205, doi:10.1029/2004jd004920, 2004.
- 1356 Han, Z.W., Li, J.W., Xia, X.A., Zhang, R.J.: Investigation of direct radiative effects of aerosols in dust  
1357 storm season over East Asia with an online coupled regional climate-chemistry-aerosol model,  
1358 *Atmos. Environ.*, 54, 688-699, 2012.
- 1359 Han, Z.W., Li, J.W., Guo, W.D., Xiong, Z., Zhang, W.: A study of dust radiative feedback on dust  
1360 cycle and meteorology over East Asia by a coupled regional climate-chemistry-aerosol model,  
1361 *Atmos. Environ.*, 68, 54-63, 2013.
- 1362 Han, Z.W., Li, J.W., Yao, X.H., Tan, S.C.: A regional model study of the characteristics and indirect  
1363 effects of marine primary organic aerosol in springtime over East Asia, *Atmos. Environ.*, 197,  
1364 22–35, 2019.
- 1365 Hsiao, T.-C., Chen, W.-N., Ye, W.-C., Lin, N.-H., Tsay, S.-C., Lin, T.-H., Lee, C.-T., Chuang, M.-T.,  
1366 Pantina, P., Wang, S.-H.: Aerosol optical properties at the Lulin Atmospheric Background Station  
1367 in Taiwan and the influences of long-range transport of air pollutants, *Atmos. Environ.*, 150,  
1368 366-378, 2017.
- 1369 Hong, S.H. and Pan, H.L.: Nonlocal boundary layer vertical diffusion in a medium range forecast  
1370 model, *Mon. Wea. Rev.*, 124, 2322–2339, 1996.
- 1371 Hoose, C., Kristjánsson, J.E., Iversen T., A. Kirkevåg, Ø. Seland, and Gettelman A.: Constraining  
1372 cloud droplet number concentration in GCMs suppresses the aerosol indirect effect, *Geophys.*  
1373 *Res. Lett.*, 36, L12807, doi:10.1029/2009GL038568, 2009.
- 1374 Hu, Q.-H., Xie, Z.-Q., Wang, X.-M., Kang, H., He, Q.-F., and Zhang, P.: Secondary organic aerosols  
1375 over oceans via oxidation of isoprene and monoterpenes from Arctic to Antarctic, *Sci. Rep.*, 3,  
1376 2280, doi:10.1038/srep02280, 2013.
- 1377 Huang, W.T.K., Ickes, L., Tegen, I., Rinaldi, M., Ceburnis, D., and Lohmann, U.: Global relevance of  
1378 marine organic aerosol as ice nucleating particles, *Atmos. Chem. Phys.*, 18, 11423–11445,  
1379 <https://doi.org/10.5194/acp-18-11423-2018>, 2018.
- 1380 Huff Hartz, K.E., Tischuk, J.E., Chan, M.N., Chan, C.K., Donahue, N.M., and Pandis, S.N.: Cloud

1381 condensation nuclei activation of limited solubility organic aerosol, *Atmos. Environ.*, 40, 605–  
1382 617, 2006.

1383 IPCC: Intergovernmental Panel on Climate Change, *Climate Change 2013: The Physical Science*  
1384 *Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental*  
1385 *Panel on Climate Change, Report*, edited by: Stocker, T.F., Qin, D.H., Plattner, G.K., Tignor,  
1386 M.M.B., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P.M., Cambridge  
1387 University Press, New York, <http://www.ipcc.ch/report/ar5>, 2013 (last access: 2020/04/30).

1388 Kanaya, Y., Pan, X., Miyakawa, T., Komazaki, Y., Taketani, F., Uno, I., and Kondo, Y.: Long-term  
1389 observations of black carbon mass concentrations at Fukue Island, western Japan, during 2009–  
1390 2015: constraining wet removal rates and emission strengths from East Asia, *Atmos. Chem. Phys.*,  
1391 16, 10689–10705, <https://doi.org/10.5194/acp-16-10689-2016>, 2016.

1392 Kang, M., Fu, P., Kawamura, K., Yang, F., Zhang, H., Zang, Z., Ren, H., Ren, L., Zhao, Y., Sun, Y.,  
1393 and Wang, Z.: Characterization of biogenic primary and secondary organic aerosols in the marine  
1394 atmosphere over the East China Sea, *Atmos. Chem. Phys.*, 18, 13947–13967, 2018.

1395 Kelly, J.T., Bhave, P.V., Nolte, C.G., Shankar, U., and Foley, K.M.: Simulating emission and chemical  
1396 evolution of coarse sea-salt particles in the Community Multiscale Air Quality (CMAQ) model.  
1397 *Geosci. Model Dev.*, 3, 257–273, 2010.

1398 Kiehl, J.T., Hack, J.J., Bonan, G.B., Boville, B.A., Briegleb, B.P., Williamson, D.L., Rasch, P.J.:  
1399 Description of the NCAR Community Climate Model (CCM3), NCAR Technical Note,  
1400 NCAR/TN-420+STR, p.152, 1996.

1401 Kunwar, B. and Kawamura, K.: One-year observations of carbonaceous and nitrogenous components  
1402 and major ions in the aerosols from subtropical Okinawa Island, an outflow region of Asian dusts,  
1403 *Atmos. Chem. Phys.*, 14, 1819–1836, <https://doi.org/10.5194/acp-14-1819-2014>, 2014.

1404 Lack, D.A., Tie, X.X., Bofinger, N.D., Wiegand, A.N., and Madronich, S.: Seasonal variability of  
1405 secondary organic aerosol: A global modeling study, *J. Geophys. Res.: Atmosphere*, 109, D03203,  
1406 doi:10.1029/2003JD003418, 2004.

1407 Lee, A., Goldstein, A.H., Kroll, J.H., Ng, N.L., Varutbangkul, V., Flagan, R.C., and Seinfeld, J.H.:  
1408 Gas-phase products and secondary aerosol yields from the photooxidation of 16 different  
1409 terpenes, *J. Geophys. Res.*, 111, D17305, doi:10.1029/2006JD007050, 2006.

1410 Leibensperger, E.M., Mickley, L.J., Jacob, D.J., Chen, W.-T., Seinfeld, J.H., Nenes, A., Adams, P.J.,

1411 Streets, D.G., Kumar, N., and Rind, D.: Climatic effects of 1950–2050 changes in US  
1412 anthropogenic aerosols–Part 1: Aerosol trends and radiative forcing, *Atmos. Chem. Phys.*, 12,  
1413 3333–3348, 2012.

1414 Li, J.-L., Zhang, H.-H., and Yang, G.-P.: Distribution and sea-to-air flux of isoprene in the East China  
1415 Sea and the South Yellow Sea during summer, *Chemosphere*, 178, 291-300, 2017.

1416 Li, J.-L., Zhai, X., Zhang, H.-H., and Yang, G.-P.: Temporal variations in the distribution and  
1417 sea-to-air flux of marine isoprene in the East China Sea, *Atmos. Environ.*, 187, 131–143, 2018.

1418 Li, J.W. and Han, Z.W.: A modeling study of the impact of heterogeneous reactions on mineral aerosol  
1419 surfaces on tropospheric chemistry over East Asia, *Particuology*, 8, 433-441, 2010.

1420 Li, J.W., Han, Z.W., and Zhang, R.J.: Influence of aerosol hygroscopic growth parameterization on  
1421 aerosol optical depth and direct radiative forcing over East Asia, *Atmos. Res.*, 140-141, 14-27,  
1422 2014.

1423 Li, J.W. and Han, Z.W.: Aerosol vertical distribution over east china from RIEMS-Chem simulation in  
1424 comparison with CALIPSO measurements, *Atmos. Environ.*, 143, 177-189, 2016a.

1425 Li, J.W. and Han, Z.W.: Seasonal variation of nitrate concentration and its direct radiative forcing over  
1426 East Asia, *Atmosphere*, 7(8), 105, 2016b.

1427 Li, J.W., Han, Z.W., Yao, X.H.: A modeling study of the influence of sea salt on inorganic aerosol  
1428 concentration, size distribution, and deposition in the western Pacific Ocean, *Atmos. Environ.*,  
1429 188, 157-173, 2018.

1430 Li, J.W., Han, Z.W., Yao, X.H., Xie, Z.X., Tan, S.C.: The distributions and direct radiative effects of  
1431 marine aerosols over East Asia in springtime, *Sci. Tot. Environ.*, 651, 1913–1925, 2019.

1432 Li, J., Han, Z., Wu, Y., Xiong, Z., Xia, X., Li, J., Liang, L., and Zhang, R.: Aerosol radiative effects  
1433 and feedbacks on boundary layer meteorology and PM<sub>2.5</sub> chemical components during winter  
1434 haze events over the Beijing-Tianjin-Hebei region, *Atmos. Chem. Phys.*, 20, 8659–8690,  
1435 <https://doi.org/10.5194/acp-20-8659-2020>, 2020.

1436 Li, M., Zhang, Q., Kurokawa, J.-I., Woo, J.-H., He, K., Lu, Z., Ohara, T., Song, Y., Streets, D.G.,  
1437 Carmichael, G.R., Cheng, Y., Hong, C., Huo, H., Jiang, X., Kang, S., Liu, F., Su, H., and Zheng,  
1438 B.: MIX: a mosaic Asian anthropogenic emission inventory under the international collaboration  
1439 framework of the MICS-Asia and HTAP, *Atmos. Chem. Phys.*, 17, 935-963,  
1440 [doi:10.5194/acp-17-935-2017](https://doi.org/10.5194/acp-17-935-2017), 2017.



1441 Liao, H., Seinfeld, J.H., Adams, P.J., Mickley, L.J.: Global radiative forcing of coupled tropospheric  
1442 ozone and aerosols in a unified general circulation model, *J. Geophys. Res.: Atmosphere*, 109,  
1443 D16207. <https://doi.org/10.1029/2003JD004456>, 2004.

1444 Liu, X. and Wang, J.: How important is organic aerosol hygroscopicity to aerosol indirect forcing?  
1445 *Environ. Res. Lett.*, 5(4), 044010, <http://iopscience.iop.org/1748-9326/5/4/044010>, 2010.

1446 Lohmann, U. and Feichter J.: Global indirect aerosol effects: a review, *Atmos. Chem. Phys.*, 5, 715–  
1447 737, 2005.

1448 Long, M.S., Keene, W.C., Kieber, D.J., Erickson, D.J., and Maring, H.: A sea-state based source  
1449 function for size- and composition-resolved marine aerosol production, *Atmos. Chem. Phys.*, 11,  
1450 1203–1216, <https://doi.org/10.5194/acp-11-1203-2011>, 2011.

1451 Luo, G. and Yu, F.: A numerical evaluation of global oceanic emissions of  $\alpha$ -pinene and isoprene,  
1452 *Atmos. Chem. Phys.*, 10, 2007–2015, doi:10.5194/acp-10-2007-2010, 2010.

1453 Luo, L., Yao, X.H., Gao, H.W., Hsu, S.C., Li, J.W., and Kao, S.J.: Nitrogen speciation in various types  
1454 of aerosols in spring over the northwestern Pacific Ocean, *Atmos. Chem. Phys.*, 16, 325-341,  
1455 2016.

1456 Luo, L., Kao, S.-J., Bao, H., Xiao, H., Xiao, H., Yao, X., Gao, H., Li, J., and Lu, Y.: Sources of  
1457 reactive nitrogen in marine aerosol over the Northwest Pacific Ocean in spring, *Atmos. Chem.*  
1458 *Phys.*, 18, 6207–6222, 2018.

1459 Ma, Q.X., Wu, Y.F., Zhang, D.Z., Wang, X.J., Xia, Y.J., Liu, X.Y., Tian, P., Han, Z.W., Xia, X.A.,  
1460 Wang, Y., and Zhang, R.J.: Roles of regional transport and heterogeneous reactions in the PM<sub>2.5</sub>  
1461 increase during winter haze episodes in Beijing, *Sci. Total. Environ.*, 599/600, 246–253, 2017.

1462 Martin, G.M., Johnson, D.W., and Spice, A.: The Measurements and Parameterization of Effective  
1463 Radius of Droplets in Warm Stratocumulus Clouds, *J. Atmos. Sci.*, 51, 1823–1842, 1994.

1464 Matsunaga, S., Mochida, M., Saito, T., and Kawamura, K.: In situ measurement of isoprene in the  
1465 marine air and surface seawater from the western North Pacific, *Atmos. Environ.*, 36, 6051–6057,  
1466 doi:10.1016/s1352-2310(02)00657-x, 2002.

1467 Meskhidze, N. and Nenes, A.: Phytoplankton and cloudiness in the Southern Ocean, *Science*, 314,  
1468 1419–1423, 2006.

1469 Meskhidze, N., Xu, J., Gantt, B., Zhang, Y., Nenes, A., Ghan, S.J., Liu, X., Easter, R., Zaveri, R.:  
1470 Global distribution and climate forcing of marine organic aerosol: 1. Model improvements and

1471 evaluation, *Atmos. Chem. Phys.*, 11, 11689–11705, 2011.

1472 Miyazaki, Y., Kawamura, K., and Sawano, M.: Size distributions and chemical characterization of  
1473 water-soluble organic aerosols over the western North Pacific in summer, *J. Geophys. Res.*, 115,  
1474 D23210, doi:10.1029/2010JD014439, 2010.

1475 Monahan, E.C., Spiel, D.E., Davidson, K.L.: A model of marine aerosol generation via white caps and  
1476 wave disruption. In: *Oceanic Whitecaps*. D. Reidel, Norwell, Mass, pp. 167–174, 1986.

1477 Myriokefalitakis, S., Vignati, E., Tsigaridis, K., Papadimas, C., Sciare, J., Mihalopoulos, N., Facchini,  
1478 M.C., Rinaldi, M., Dentener, F.J., Ceburnis, D., Hatzianastasiou, N., O'Dowd, C.D., van Weele,  
1479 M., and Kanakidou, M.: Global modeling of the oceanic source of organic aerosols. *Adv. Meteor.*,  
1480 939171, doi:10.1155/2010/939171, 2010.

1481 NOAA/NCEP: NCEP FNL Operational Model Global Tropospheric Analyses, continuing from July  
1482 1999. Research Data Archive at the National Center for Atmospheric Research, Computational  
1483 and Information Systems Laboratory. Dataset. <https://doi.org/10.5065/D6M043C6>, 2000, Last  
1484 accessed, 2019/12/12.

1485 OBPB: NASA Goddard Space Flight Center, Ocean Ecology Laboratory, Ocean Biology Processing  
1486 Group: Visible and Infrared Imager/Radiometer Suite (VIIRS) Chlorophyll Data; NASA  
1487 OB.DAAC, Greenbelt, MD, USA. doi:10.5067/NPP/VIIRS/L3M/CHL/2018.

1488 O'Dowd, C.D., Facchini, M.C., Cavalli, F., Ceburnis, D., Mircea, M., Decesari, S., Fuzzi, S., Yoon,  
1489 Y.J., Putaud, J.P.: Biogenically driven organic contribution to marine aerosol, *Nature*, 431, 676–  
1490 680, 2004.

1491 Ooki, A., Nomura, D., Nishino, S., Kikuchi, T., and Yokouchi, Y.: A global-scale map of isoprene and  
1492 volatile organic iodine in surface seawater of the Arctic, Northwest Pacific, Indian, and Southern  
1493 Oceans, *J. Geophys. Res.: Oceans* 120, 4108–4128, doi:10.1002/2014JC010519, 2015.

1494 Ovadnevaite, J., Ceburnis, D., Martucci, G., Bialek, J., Monahan, C., Rinaldi, M., Facchini, M.C.,  
1495 Berresheim, H., Worsnop, D.R., O'Dowd, C.: Primary marine organic aerosol: a dichotomy of  
1496 low hygroscopicity and high CCN activity, *Geophys. Res. Lett.*, 38, L21806.  
1497 <https://doi.org/10.1029/2011GL048869>, 2011.

1498 Petters, M.D. and Kreidenweis, S.M.: A single parameter representation of hygroscopic growth and  
1499 cloud condensation nucleus activity, *Atmos. Chem. Phys.*, 7, 1961–1971, 2007.

1500 Quinn, P.K., Bates Timothy S., Schulz Kristen S., Coffman D.J., Frossard A.A., Russell L.M., Keene

1501 W.C. and Kieber D.J.: Contribution of sea surface carbon pool to organic matter enrichment in  
1502 sea spray aerosol. *Nat. Geosci.* 7, 228-232, 2014.

1503 Quinn, P.K., Coffman, D.J., Johnson, J.E., Upchurch, L.M., and Bates, T.S.: Small fraction of marine  
1504 cloud condensation nuclei made up of sea spray aerosol, *Nat. Geosci.*, 10, 674–679, 2017.

1505 Rap, A., Scott, C.E., Spracklen, D.V., Bellouin, N., Forster, P.M., Carslaw, K.S., Schmidt, A., and  
1506 Mann, G.: Natural aerosol direct and indirect radiative effects, *Geophys. Res. Lett.*, 40,  
1507 3297-3301, doi:10.1002/grl.50441, 2013.

1508 Reisner, J., Rasmussen, R.M., Bruintjes, R.T.: Explicit forecasting of super cooled liquid water in  
1509 winter storms using the MM5 mesoscale model, *Quart. J. Roy. Meteor. Soc.*, 124 (548), 1071–  
1510 1107, 1998.

1511 Riemer, N., West, M., Zaveri, R., Easter, R.: Estimating black carbon aging time-scales with a  
1512 particle-resolved aerosol model, *J. Aerosol Sci.*, 41, 143-158, 2010.

1513 Roelofs, G.J.: A GCM study of organic matter in marine aerosol and its potential contribution to cloud  
1514 drop activation, *Atmos. Chem. Phys.*, 8, 709–719 2008, 2008.

1515 Sayer, A.M., Hsu, N.C., Bettenhausen, C., Lee, J., Kim, W.V., and Smirnov, A.: Satellite Ocean  
1516 Aerosol Retrieval (SOAR) Algorithm Extension to S-NPP VIIRS as Part of the “Deep Blue”  
1517 Aerosol Project, *J. Geophys. Res. Atmos.*, 123, doi:10.1002/2017JD027412, 2018.

1518 Shao, Y. and Dong, C.H.: A review on East Asian dust storm climate, modeling and monitoring, *Glob.*  
1519 *Planet. Change*, 52, 1-22, 2006.

1520 Sindelarova, K., Granier, C., Bouarar, I., Guenther, A., Tilmes, S., Stavrou, T., Müller, J.-F., Kuhn,  
1521 U., Stefani, P., and Knorr, W.: Global data set of biogenic VOC emissions calculated by the  
1522 MEGAN model over the last 30 years. *Atmos. Chem. Phys.*, 14, 9317–9341.  
1523 <https://doi.org/10.5194/acp-14-9317-2014>, 2014.

1524 Slingo A.: A GCM Parameterization for the Shortwave Radiative Properties of Water Clouds, *J. Atmos.*  
1525 *Sci.*, 46(10), 1419-1427, 1989.

1526 Slinn, W.G.N.: Precipitation scavenging: in *Atmospheric Science and Power Production*. pp. 466–532,  
1527 Technical Information Center, Office of Science and Technology Information, Department of  
1528 Energy, Washington, D. C., 1984.

1529 Smith, S.J., van Aardenne, J., Klimont, Z., Andres, R.J., Volke, A., and Delgado Arias, S.:  
1530 Anthropogenic sulfur dioxide emissions: 1850-2005, *Atmos. Chem. Phys.*, 11, 1101-1116, 2011.

1531 Spracklen, D.V., Arnold, S.R., Carslaw, K.S., Sciare, J., and Pio, C.: Globally significant oceanic  
1532 source of organic carbon aerosol, *Geophys. Res. Lett.*, 35, L12811, doi:10.1029/2008GL033359,  
1533 2008.

1534 Surratt, J.D., Chan, A.W.H., Eddingsaas, N.C., Chan, M.N., Loza, C.L., Kwan, A.J., Hersey, S.P.,  
1535 Flagan, R.C., Wennberg, P.O., and Seinfeld, J.H.: Reactive intermediates revealed in secondary  
1536 organic aerosol formation from isoprene, *Proc. Natl. Acad. Sci. USA*, 107, 6640–6645,  
1537 doi:10.1073/pnas.0911114107, 2010.

1538 Tan, S.C., Li, J.W., Che, H.Z., Chen, B., and Wang, H.: Transport of East Asian dust storms to the  
1539 marginal seas of China and the southern North Pacific in spring 2010, *Atmos. Environ.*, 148,  
1540 316-328, 2017.

1541 Tao, J., Surapipith, V., Han, Z.W., Prapamontol, T., Kawichai, S., Zhang, L.M., Zhang, Z.S., Wu, Y.F.,  
1542 Li, J.W., Li J., Yang, Y.H., and Zhang, R.J.: High mass absorption efficiency of carbonaceous  
1543 aerosols during the biomass burning season in Chiang Mai of northern Thailand, *Atmos.*  
1544 *Environ.*, 240, 117821, 2020.

1545 Van den Berg, A., Dentener, F., and Lelieveld, J.: Modeling the chemistry of the marine boundary  
1546 layer: Sulphate formation and the role of sea-salt aerosol particles, *J. Geophys. Res.*, 105, 11671–  
1547 11698. <https://doi.org/10.1029/1999JD901073>, 2000.

1548 Vignati, E., Facchini, M.C., Rinaldi, M., Scannell, C., Ceburnis, D., Sciare, J., Kanakidou, M.,  
1549 Myriokefalitakis, S., Dentener, F., and O’Dowd, C.D.: Global scale emission and distribution of  
1550 seaspray aerosol: sea-salt and organic enrichment, *Atmos. Environ.*, 44, 670–677, 2010.

1551 Xiong, Z., Fu, C.B., and Yan, X.D.: Regional Integrated environmental model system and its  
1552 simulation of East Asia summer monsoon, *Chinese Sci. Bull.*, 54(22), 4253-4261, 2009.

1553 Wang, F.W., Guo, Z.G., Lin, T., Hu, L.M., Chen, Y.J., and Zhu, Y.F.: Characterization of carbonaceous  
1554 aerosols over the East China Sea: The impact of the East Asian continental outflow, *Atmos.*  
1555 *Environ.*, 110, 163-173, 2015.

1556 Wang, M., and Penner J.E.: Aerosol indirect forcing in a global model with particle nucleation, *Atmos.*  
1557 *Chem. Phys.*, 9, 239– 260, 2009.

1558 Wang, S.Y., Fu, C.B., Wei, H.L., Qian, Y., Xiong, Z., Feng, J.M., Zhao, D.M., Dan, L., Han, Z.W., Su,  
1559 B.K., Zhao, M., Zhang, Y.C., Tang, J.P., Liu, H.N., Wu, J., Zeng, X.M., Chen, M., Wang, L.Z.:  
1560 Regional integrated environmental modeling system: development and application, *Climatic*

1561 Change, 129, 499-510, 2015.

1562 Westervelt, D.M., Moore, R.H., Nenes, A., and Adams, P.J.: Effect of primary organic sea spray  
1563 emissions on cloud condensation nuclei concentrations, *Atmos. Chem. Phys.*, 12, 89–101.  
1564 <https://doi.org/10.5194/acp-12-89-2012>, 2012.

1565 Wu, Y.F., Wang, X.J., Tao, J., Huang, R.J., Tian, P., Cao, J.J., Zhang, L.M., Ho, K.F., Han, Z.W.,  
1566 Zhang, R.J.: Size distribution and source of black carbon aerosol in urban Beijing during winter  
1567 haze episodes, *Atmos. Chem. Phys.*, 17, 7965-7975, 2017.

1568 Zeng, S., Riedi, J., Trepte, C.R., Winker, D.M., and Hu, Y.X.: Study of global cloud droplet number  
1569 concentration with A-Train satellites. *Atmos. Chem. Phys.* 14, 7125–7134, 2014.

1570 Zhang, K.M., Knipping, E.M., Wexler, A.S., Bhave, P.V., and Tonnesen, G.S.: Size distribution of  
1571 sea-salt emissions as a function of relative humidity, *Atmos. Environ.*, 39, 3373-3379, 2005.

1572 Zhang, L.M., Gong, S.L., Padro, J., and Barrie, L.: A size-segregated particle dry deposition scheme  
1573 for an atmospheric aerosol module, *Atmos. Environ.*, 35(3), 549-560, 2001.

1574 Zhao, B., Liou, K.N., Gu, Y., Li, Q., Jiang, J.H., Su, H., He, C., Tseng, H.R., Wang, S., Liu, R., Qi, L.,  
1575 Lee, W.L., and Hao, J.: Enhanced PM<sub>2.5</sub> pollution in China due to aerosol-cloud interactions, *Sci.*  
1576 *Rep.*, 7, 4453, <https://doi.org/10.1038/s41598-017-04096-8>, 2017.

1577 Zhu, C., Kawamura, K., and Fu, P.: Seasonal variations of biogenic secondary organic aerosol tracers  
1578 in Cape Hedo, Okinawa, *Atmos. Environ.*, 130, 113-119, 2016.

1579

1580

1581

1582




Figure 1. Model domain, observational sites, and research cruise tracks. EANET sites are marked in light-blue. Observation sites of carbonaceous aerosols are marked in red (Chichijima Island: Boreddy et al., 2018; Fukue: Kanaya et al., 2016; Okinawa: Kunwar and Kawamura, 2014; Huaniao Island: Wang F. W. et al., 2014). Three CNEMC sites are marked in grey (Qingdao, Shanghai, and Fuzhou). Two research cruise tracks are represented by green line (Dongfanghong II from 17 March to 22 April 2014: Luo et al., 2016; Feng et al., 2017) and orange line (KEXUE-1 from 18 May to 12 June 2014: Kang et al., 2018), respectively. AERONET sites are represented by triangles with numbers (1-Ussuriysk, 2-Yonsei\_University, 3-Gwangju\_GIST, 4-EPA-NCU, 5-Chen-Kung\_Univ, 6-Fukuoka, and 7-Shirahama). Full names of abbreviations are given in the text.



Figure 2. The model simulated (Sim) and observed (Obs) monthly  $PM_{10}$  (a~l) and  $PM_{2.5}$  (m~r) concentrations at EANET and CNEMC sites for the year 2014. The monthly data were averaged from hourly observations and the simulations were sampled according to the observations. Simulated aerosol components (PPM: primary anthropogenic PM, Sea salt, Dust, MOA, Car: anthropogenic BC+OC, SNA: sulfate+nitrate+ammonium) are also shown.




Figure 3. Observed and model simulated monthly mean sodium ( $\text{Na}^+$ ) concentrations at 6 coastal and island EANET sites of Japan for the year 2014.






Figure 4. The model simulated (bars) and observed (dotted lines) daily BC and OC concentrations from the spring campaign (a, b) and half-day OC concentrations from the early summer campaign (c). The modeled total OC concentration was decomposed into those from marine (green bars) and land (blue bars) sources.




Figure 5. The model simulated (bars) and observed (dotted lines) OC concentrations at different sites. Seasonal mean concentrations were provided at (a) Huaniao Island (Wang et al., 2015) and (b) Okinawa (Kunwar and Kawamura, 2014) while monthly mean concentrations were provided at (c) Chichijima Island (Boreddy et al., 2018). Standard deviations were available at Okinawa and Chichijima. The modeled OC concentrations were decomposed to marine (green bars) and land (blue bars) sources. The simulation is for the year 2014.




Figure 6. The model simulated (Sim) and observed (Obs) monthly mean AOD at 7 AERONET sites for the year 2014. The monthly mean observations were calculated from hourly data and the simulations were sampled according to the observations. Contributions of aerosol components to total AOD by MOA, sea salt, and aerosols of non-oceanic sources (anthropogenic, dust etc.) are shown.




Figure 7. Model simulated annual and seasonal mean distributions of MPOA emissions (a~e), VIIRS retrieved surface sea water chlorophyll-a (Chl-a) concentrations (f~j), and model simulated sea salt (SS) emissions (k~o). Two hotspot regions are marked with red boxes: the region including the East China Sea, the Yellow Sea, and the Bohai Sea (EYB, 27~40°N, 115~123°E) and the region including the northern parts of the western Pacific to the northeast of Japan (NWP, 35~55°N, 140~160°E). Units are given in parentheses.




Figure 8. Model simulated annual and seasonal mean near surface MOA (primary+secondary) concentrations (a~e), near surface MSOA concentrations (f~j), and percentage contributions of MOA to total OA (k~o). The two regions of the EYB (27~40°N, 115~123°E) and the NWP (35~55°N, 140~160°E) are marked in 8a. Units are given in parentheses.




Figure 9. Model simulated annual and seasonal mean near surface temperatures (unite:  $^{\circ}\text{C}$ ) overlaid with wind vectors (unit:  $\text{m s}^{-1}$ ) (a~e) and mean monthly precipitations (unit:  $\text{cm month}^{-1}$ ) (f~j).




Figure 10. Model simulated annual and seasonal mean direct radiative effect due to MOA ( $DRE_{MOA}$ ) (a~e) and indirect radiative effect due to MOA ( $IRE_{MOA}$ ) (f~j) at the top of atmosphere (TOA) under all-sky condition (unit:  $W m^{-2}$ ). Red dot indicates areas where the difference is statistically significant at the 99% confidence level according to a two-tailed t-test.




Figure 11. Model simulated annual mean (a) direct radiative effect of MOA due to aerosol-radiation interaction ( $DRE_{ari}$ ), (b) indirect radiative effect of MOA due to aerosol-cloud interaction ( $IRE_{aci}$ ) at the top of atmosphere (TOA) (unit:  $W m^{-2}$ ). Red dot indicates areas where the difference is statistically significant at the 99% confidence level according to a two-tailed t-test.



Table 1. Annual and seasonal performance statistics for hourly PM<sub>10</sub> and PM<sub>2.5</sub> concentrations (unit:  $\mu\text{g m}^{-3}$ ) at EANET sites for the year 2014. Mean observation (Obs), mean simulation (Sim), correlation coefficient (R), and normalized mean bias (NMB in %) are listed. ANN=annual, DJF=December-January-February, MAM=March-April-May, JJA=June-July-August, and SON=September-October-November.

| Sites                   | Samples | ANN  |      |      | DJF |      |      | MAM  |     |      | JJA  |      |     | SON  |      |      |     |      |      |      |     |  |
|-------------------------|---------|------|------|------|-----|------|------|------|-----|------|------|------|-----|------|------|------|-----|------|------|------|-----|--|
|                         |         | Obs  | Sim  | R    | NMB | Obs  | Sim  | R    | NMB | Obs  | Sim  | R    | NMB | Obs  | Sim  | R    | NMB |      |      |      |     |  |
| <b>PM<sub>10</sub></b>  |         |      |      |      |     |      |      |      |     |      |      |      |     |      |      |      |     |      |      |      |     |  |
| Rishiri                 | 8381    | 18.0 | 19.9 | 0.53 | 11  | 13.6 | 18.4 | 0.65 | 35  | 20.1 | 23.0 | 0.56 | 15  | 15.2 | 19.4 | 0.42 | 28  | 23.0 | 18.8 | 0.51 | -18 |  |
| Tappi                   | 8584    | 20.1 | 23.2 | 0.49 | 15  | 17.4 | 22.3 | 0.54 | 28  | 29.0 | 28.0 | 0.59 | -4  | 16.1 | 23.8 | 0.18 | 48  | 17.6 | 18.5 | 0.39 | 5   |  |
| Sado                    | 8640    | 22.8 | 26.4 | 0.63 | 16  | 23.6 | 29.5 | 0.68 | 25  | 29.2 | 30.3 | 0.65 | 4   | 18.6 | 24.5 | 0.55 | 32  | 19.6 | 21.4 | 0.53 | 9   |  |
| Oki                     | 8424    | 31.3 | 29.2 | 0.68 | -7  | 33.2 | 31.1 | 0.65 | -7  | 40.2 | 37.5 | 0.71 | -7  | 26.7 | 26.6 | 0.61 | 0   | 25.8 | 22.3 | 0.66 | -14 |  |
| Hedo                    | 8008    | 27.7 | 21.7 | 0.56 | -22 | 28.8 | 22.4 | 0.66 | -22 | 30.5 | 28.3 | 0.58 | -7  | 20.7 | 14.8 | 0.54 | -28 | 30.9 | 20.8 | 0.34 | -33 |  |
| Ogasawara               | 8120    | 11.5 | 15.7 | 0.48 | 36  | 13.4 | 20.3 | 0.38 | 52  | 14.2 | 19.7 | 0.40 | 39  | 7.0  | 6.8  | 0.46 | -2  | 11.2 | 15.0 | 0.30 | 34  |  |
| Jeju                    | 7101    | 46.9 | 36.9 | 0.64 | -21 | 50.1 | 38.2 | 0.71 | -24 | 62.6 | 46.6 | 0.66 | -26 | 36.4 | 31.5 | 0.44 | -13 | 34.7 | 29.3 | 0.44 | -15 |  |
| Kanghwa                 | 8524    | 49.2 | 51.2 | 0.59 | 4   | 50.2 | 46.0 | 0.60 | -8  | 59.9 | 61.9 | 0.66 | 3   | 40.0 | 47.2 | 0.47 | 18  | 46.5 | 49.3 | 0.38 | 6   |  |
| Imsil                   | 8383    | 44.5 | 32.3 | 0.58 | -27 | 48.8 | 27.9 | 0.63 | -43 | 58.0 | 42.1 | 0.62 | -27 | 38.4 | 31.1 | 0.47 | -19 | 33.0 | 28.2 | 0.42 | -15 |  |
| Average                 | 74165   | 30.0 | 28.5 | 0.65 | -5  | 30.8 | 28.3 | 0.67 | -8  | 37.9 | 35.2 | 0.65 | -7  | 23.9 | 25.1 | 0.59 | 5   | 26.9 | 25.0 | 0.58 | -7  |  |
| <b>PM<sub>2.5</sub></b> |         |      |      |      |     |      |      |      |     |      |      |      |     |      |      |      |     |      |      |      |     |  |
| Rishiri                 | 8331    | 8.6  | 8.7  | 0.54 | 0   |      |      |      |     |      |      |      |     |      |      |      |     |      |      |      |     |  |
| Sado                    | 6517    | 11.0 | 13.4 | 0.53 | 21  | 8.1  | 7.4  | 0.78 | -8  | 9.2  | 9.2  | 0.56 | 0   | 7.2  | 11.5 | 0.54 | 59  | 10.0 | 6.7  | 0.31 | -33 |  |
| Oki                     | 8410    | 13.1 | 15.0 | 0.64 | 14  | 8.5  | 9.8  | 0.60 | 14  | 14.4 | 16.1 | 0.63 | 12  | 11.4 | 16.8 | 0.47 | 48  | 9.1  | 9.1  | 0.24 | 0   |  |
| Average                 | 23258   | 10.9 | 12.3 | 0.61 | 12  | 13.0 | 12.9 | 0.77 | -1  | 17.4 | 18.7 | 0.64 | 8   | 12.1 | 18.3 | 0.55 | 51  | 10.1 | 10.0 | 0.39 | -1  |  |

Table 2. Same as Table 1 but for CNEMC sites.

| Sites                   | Samples | ANN   |       |       | DJF |       |       | MAM  |     |       | JJA   |      |       | SON  |       |       |     |      |       |      |    |
|-------------------------|---------|-------|-------|-------|-----|-------|-------|------|-----|-------|-------|------|-------|------|-------|-------|-----|------|-------|------|----|
|                         |         | Obs   | Sim   | R NMB | Obs | Sim   | R NMB | Obs  | Sim | R NMB | Obs   | Sim  | R NMB | Obs  | Sim   | R NMB |     |      |       |      |    |
| <b>PM<sub>10</sub></b>  |         |       |       |       |     |       |       |      |     |       |       |      |       |      |       |       |     |      |       |      |    |
| Qingdao                 | 7622    | 107.0 | 108.6 | 0.61  | 1   | 131.0 | 124.3 | 0.76 | -5  | 117.3 | 109.9 | 0.49 | -6    | 83.6 | 108.4 | 0.64  | 30  | 97.1 | 101.4 | 0.59 | 4  |
| Shanghai                | 7581    | 73.4  | 70.5  | 0.55  | -4  | 93.9  | 81.1  | 0.72 | -14 | 83.2  | 80.2  | 0.58 | -4    | 59.6 | 64.0  | 0.37  | 7   | 64.8 | 60.6  | 0.43 | -7 |
| Fuzhou                  | 7610    | 63.7  | 63.8  | 0.38  | 0   | 69.9  | 72.9  | 0.30 | 4   | 69.8  | 72.6  | 0.32 | 4     | 55.3 | 45.8  | 0.28  | -17 | 60.5 | 64.5  | 0.30 | 7  |
| Average                 | 22813   | 81.6  | 80.7  | 0.65  | -1  | 98.4  | 92.6  | 0.74 | -6  | 89.9  | 86.7  | 0.58 | -4    | 66.0 | 72.0  | 0.61  | 9   | 74.1 | 75.4  | 0.51 | 2  |
| <b>PM<sub>2.5</sub></b> |         |       |       |       |     |       |       |      |     |       |       |      |       |      |       |       |     |      |       |      |    |
| Qingdao                 | 7627    | 55.2  | 48.7  | 0.72  | -12 | 75.1  | 67.8  | 0.83 | -10 | 56.3  | 43.7  | 0.61 | -22   | 40.5 | 42.8  | 0.60  | 6   | 48.2 | 43.9  | 0.74 | -9 |
| Shanghai                | 7724    | 51.9  | 51.8  | 0.62  | 0   | 68.0  | 59.6  | 0.80 | -12 | 57.2  | 57.5  | 0.60 | 0     | 42.6 | 49.8  | 0.46  | 17  | 42.6 | 42.9  | 0.51 | 1  |
| Fuzhou                  | 7641    | 32.3  | 30.0  | 0.44  | -7  | 40.3  | 40.2  | 0.25 | 0   | 35.8  | 36.8  | 0.37 | 3     | 24.0 | 15.8  | 0.38  | -34 | 29.2 | 27.3  | 0.29 | -7 |
| Average                 | 22992   | 46.6  | 43.4  | 0.70  | -7  | 61.1  | 55.8  | 0.78 | -9  | 49.7  | 45.6  | 0.63 | -8    | 35.6 | 35.5  | 0.62  | 0   | 39.9 | 38.0  | 0.62 | -5 |

Table 3. Performance statistics for BC and OC from the two research campaigns in 2014. BC and OC were measured on Dongfanghong II during the spring campaign whereas only OC were collected on KEXUE-1 during the early summer campaign. Mean observation (Obs), mean simulation (Sim), correlation coefficient (R), and normalized mean bias (NMB in %) are listed. The modeled concentrations of marine-OC (including MPOA and MSOA) and its contribution to total OC were estimated.

|                              | Dongfanghong II |      |                     | KEXUE-1 |                     |
|------------------------------|-----------------|------|---------------------|---------|---------------------|
|                              | BC              | OC   | Marine-OC (% in OC) | OC      | Marine-OC (% in OC) |
| Samples                      | 19              | 19   |                     | 51      |                     |
| Obs ( $\mu\text{g m}^{-3}$ ) | 0.49            | 1.2  |                     | 4.3     |                     |
| Sim ( $\mu\text{g m}^{-3}$ ) | 0.55            | 1.1  | 0.33 (29%)          | 3.7     | 0.23 (6%)           |
| R                            | 0.87            | 0.66 |                     | 0.75    |                     |
| NMB (%)                      | 13              | -5   |                     | -13     |                     |

Table 4. Comparison of model simulated and observed seasonal OC concentrations (unit:  $\mu\text{g m}^{-3}$ ) at Huaniao Island and Okinawa. The modeled concentrations of marine-OC and its contribution to total OC were estimated. ANN=annual, DJF=December-January-February, MAM=March-April-May, JJA=June-July-August, and SON=September-October-November.

|                |           | Time                  | ANN <sup>c</sup> | DJF   | MAM   | JJA   | SON   | Reference                    |
|----------------|-----------|-----------------------|------------------|-------|-------|-------|-------|------------------------------|
| <b>OC</b>      |           |                       |                  |       |       |       |       |                              |
| Huaniao Island | Obs       | Oct 2011~<br>Aug 2012 | 3.3              | 4.7   | 3.7   | 1.1   | 3.8   | Wang F. W. et al.,<br>2015   |
|                | Sim       | 2014                  | 3.2              | 4.5   | 3.9   | 1.7   | 2.9   |                              |
|                | Marine-OC |                       | 0.6              | 0.56  | 0.88  | 0.32  | 0.65  |                              |
|                | (% in OC) |                       | (19%)            | (12%) | (22%) | (19%) | (23%) |                              |
| Okinawa        | Obs       | Oct 2009~<br>Oct 2010 | 1.8              | 1.5   | 2.4   | 1.8   | 1.4   | Kunwar and<br>Kawamura, 2014 |
|                | Sim       | 2014                  | 1.3              | 1.4   | 1.9   | 0.6   | 1.2   |                              |
|                | Marine-OC |                       | 0.21             | 0.25  | 0.32  | 0.06  | 0.23  |                              |
|                | (% in OC) |                       | (17%)            | (18%) | (17%) | (10%) | (18%) |                              |

a: The location of Huaniao Island is 30.86°N, 122.67°E.

b: The location of Okinawa Island is 26.15°N, 128.03°E.

c: The annual means are averages of the four seasonal means.

Table 5. Comparison of model simulated and observed monthly mean OC concentrations (unit:  $\mu\text{g m}^{-3}$ ) at Chichijima Island. Marine-OC concentration and its contribution to total OC were estimated.

| Month            | Jan   | Feb   | Mar   | Apr   | May   | Jun  | Jul   | Aug  | Sep  | Oct   | Nov   | Dec   | Annual |
|------------------|-------|-------|-------|-------|-------|------|-------|------|------|-------|-------|-------|--------|
| Obs <sup>a</sup> | 0.80  | 0.95  | 1.1   | 0.77  | 0.80  | 0.74 | 0.58  | 0.63 | 0.60 | 0.62  | 0.75  | 0.73  | 0.76   |
| Sim <sup>b</sup> | 0.95  | 1.0   | 1.0   | 1.4   | 1.2   | 0.64 | 0.34  | 0.44 | 0.84 | 0.37  | 0.50  | 0.71  | 0.78   |
| Marine-OC        | 0.17  | 0.11  | 0.15  | 0.22  | 0.19  | 0.06 | 0.04  | 0.04 | 0.07 | 0.12  | 0.14  | 0.19  | 0.13   |
| (% in OC)        | (18%) | (11%) | (15%) | (16%) | (16%) | (9%) | (11%) | (9%) | (8%) | (33%) | (28%) | (27%) | (16%)  |

a: Observations at Chichijima Island (27.07°N, 142.22°E) were obtained from Boreddy et al. (2018) and are 12-yr averages (2001-2012).

b: Simulations are for the year 2014.

Table 6. Performance statistics for hourly AOD (unitless) at AERONET sites for the year 2014. Mean observation (Obs), mean simulation (Sim), correlation coefficient (R), and normalized mean bias (NMB in %) are listed. IDs are marked in Figure 1.

| ID | Site              | Obs  | Sim  | R    | NMB | Samples |
|----|-------------------|------|------|------|-----|---------|
| 1  | Ussuriysk         | 0.22 | 0.21 | 0.41 | -6  | 945     |
| 2  | Yonsei_University | 0.48 | 0.37 | 0.67 | -23 | 1629    |
| 3  | Gwangju_GIST      | 0.33 | 0.36 | 0.53 | 7   | 900     |
| 4  | EPA-NCU           | 0.38 | 0.39 | 0.43 | 4   | 685     |
| 5  | Chen-Kung_Univ    | 0.49 | 0.37 | 0.60 | -25 | 657     |
| 6  | Fukuoka           | 0.28 | 0.34 | 0.50 | 18  | 1144    |
| 7  | Shirahama         | 0.26 | 0.31 | 0.40 | 19  | 752     |
|    | Average           | 0.36 | 0.34 | 0.54 | -6  | 6712    |

Table 7. Modeled domain and annual/seasonal mean MPOA emission rates, surface sea water chlorophyll-a (Chl-a) concentrations, and sea salt emission fluxes over the western Pacific of East Asia (Mean), the region including the East China Sea, the Yellow Sea, and the Bohai Sea (EYB) and the region including northern parts of western Pacific to the northeast of Japan (NWP).

|     | MPOA emission<br>( $\times 10^{-2} \mu\text{g m}^{-2} \text{s}^{-1}$ ) |                  |                  |                  | Chl-a concentration<br>( $\text{mg m}^{-3}$ ) |                  |                  | Sea salt emission flux<br>( $\mu\text{g m}^{-2} \text{s}^{-1}$ ) |                  |                  |
|-----|--|------------------|------------------|------------------|---|------------------|------------------|--|------------------|------------------|
|     | Mean <sup>a</sup>  | Max <sup>b</sup> | EYB <sup>c</sup> | NWP <sup>d</sup> | Mean <sup>a</sup>                             | EYB <sup>c</sup> | NWP <sup>d</sup> | Mean <sup>a</sup>  | EYB <sup>c</sup> | NWP <sup>d</sup> |
| ANN | 0.16   | 1.8              | 0.65             | 0.40             | 1.2   | 3.5              | 0.96             | 0.36   | 0.18             | 0.59             |
| DJF | 0.18   | 3.6              | 1.2              | 0.33             | 0.67  | 3.2              | 0.37             | 0.63   | 0.35             | 1.1              |
| MAM | 0.17   | 2.5              | 0.41             | 0.43             | 0.97  | 4.0              | 1.1              | 0.30   | 0.11             | 0.61             |
| JJA | 0.08   | 1.9              | 0.12             | 0.29             | 1.1   | 3.1              | 0.90             | 0.14   | 0.04             | 0.15             |
| SON | 0.20   | 3.5              | 0.88             | 0.54             | 1.1   | 2.9              | 0.90             | 0.38   | 0.24             | 0.53             |

a: Mean over oceanic areas.

b: Maximums over oceanic areas.

c: Ocean areas within 27~40°N, 115~123°E.

d: Ocean areas within 35~55°N, 140~160°E.

Table 8. Modeled domain and annual/seasonal mean near surface MOA concentrations, MSOA concentrations, and MOA to total OA ratios over the western Pacific of East Asia (Mean), the EYB region, and the NWP region.

|     | MOA concentration<br>( $\mu\text{g m}^{-3}$ ) |                  |                  |                  | MSOA concentration<br>( $\times 10^{-3} \mu\text{g m}^{-3}$ ) |                  |                  |                  | MOA/OA<br>(%)     |                  |                  |                  |
|-----|---|------------------|------------------|------------------|---|------------------|------------------|------------------|-------------------|------------------|------------------|------------------|
|     | Mean <sup>a</sup>                             | Max <sup>b</sup> | EYB <sup>c</sup> | NWP <sup>d</sup> | Mean <sup>a</sup>   | Max <sup>b</sup> | EYB <sup>c</sup> | NWP <sup>d</sup> | Mean <sup>a</sup> | Max <sup>b</sup> | EYB <sup>c</sup> | NWP <sup>d</sup> |
| ANN | 0.27  | 1.2              | 0.48             | 0.59             | 2.2   | 6.9              | 4.1              | 3.8              | 26%               | 62%              | 13%              | 42%              |
| DJF | 0.21  | 0.8              | 0.54             | 0.23             | 0.7   | 3.2              | 1.0              | 0.4              | 24%               | 57%              | 11%              | 36%              |
| MAM | 0.37  | 1.9              | 0.62             | 0.81             | 2.7   | 10.5             | 5.3              | 4.1              | 26%               | 69%              | 15%              | 52%              |
| JJA | 0.23  | 2.3              | 0.22             | 0.8              | 3.9   | 13.6             | 7.5              | 8.3              | 23%               | 69%              | 6%               | 32%              |
| SON | 0.26  | 1.3              | 0.52             | 0.52             | 1.5   | 4.2              | 2.6              | 2.2              | 32%               | 73%              | 18%              | 48%              |

a: Mean over oceanic areas.

b: Maximums over oceanic areas.

c: Ocean areas within 27~40°N, 115~123°E.

d: Ocean areas within 35~55°N, 140~160°E.

Table 9. Modeled domain and annual/seasonal mean near surface wind speed, temperature, precipitation, and relative humidity (RH) over the western Pacific of East Asia (Mean), the EYB region, and the NWP region.

|     | Wind speed<br>( $\text{m s}^{-1}$ ) |                  |                  | Temperature<br>(°C) |                  |                  | Precipitation<br>( $\text{cm month}^{-1}$ ) |                  |                  | RH<br>(%)         |                  |                  |
|-----|-------------------------------------|------------------|------------------|---------------------|------------------|------------------|---|------------------|------------------|-------------------|------------------|------------------|
|     | Mean <sup>a</sup>                   | EYB <sup>b</sup> | NWP <sup>c</sup> | Mean <sup>a</sup>   | EYB <sup>b</sup> | NWP <sup>c</sup> | Mean <sup>a</sup>                           | EYB <sup>b</sup> | NWP <sup>c</sup> | Mean <sup>a</sup> | EYB <sup>b</sup> | NWP <sup>c</sup> |
| ANN | 4.3                                 | 2.9              | 4.0              | 19.2                | 15.1             | 8.5              | 6.1   | 2.7              | 8.0              | 78                | 73               | 83               |
| DJF | 6.4                                 | 4.5              | 6.9              | 14.0                | 4.5              | 1.0              | 7.0   | 1.8              | 12.4             | 75                | 67               | 77               |
| MAM | 3.8                                 | 2.0              | 3.7              | 16.9                | 13.4             | 5.1              | 4.3   | 2.1              | 7.0              | 79                | 75               | 84               |
| JJA | 3.0                                 | 1.9              | 2.5              | 24.0                | 23.2             | 15.8             | 5.1   | 3.5              | 3.7              | 83                | 80               | 94               |
| SON | 4.1                                 | 3.1              | 3.1              | 21.7                | 17.9             | 12.0             | 7.9   | 3.2              | 9.0              | 76                | 71               | 77               |

a: Mean over oceanic areas.

b: Ocean areas within 27~40°N, 115~123°E.

c: Ocean areas within 35~55°N, 140~160°E.

Table 10. Modeled regional and annual/seasonal mean all-sky TOA direct radiative effect (DRE) and indirect radiative effects (IRE) due to MOA, anthropogenic aerosols, and sea salt over oceanic areas of the western Pacific (WP), the EYB region, and the NWP region. The units are  $W m^{-2}$ .

|     | MOA             |                  |                  | Anthropogenic   |                  |                  | Sea salt        |                  |                  |
|-----|-----------------|------------------|------------------|-----------------|------------------|------------------|-----------------|------------------|------------------|
|     | WP <sup>a</sup> | EYB <sup>b</sup> | NWP <sup>c</sup> | WP <sup>a</sup> | EYB <sup>b</sup> | NWP <sup>c</sup> | WP <sup>a</sup> | EYB <sup>b</sup> | NWP <sup>c</sup> |
|     | DRE             |                  |                  |                 |                  |                  |                 |                  |                  |
| ANN | -0.27           | -0.33            | -0.50            | -2.8            | -6.6             | -2.7             | -0.86           | -0.56            | -0.89            |
| DJF | -0.18           | -0.32            | -0.21            | -1.2            | -3.5             | -1.2             | -0.93           | -0.48            | -0.95            |
| MAM | -0.38           | -0.34            | -0.76            | -2.8            | -6.2             | -3.0             | -0.79           | -0.36            | -1.1             |
| JJA | -0.28           | -0.26            | -0.72            | -4.9            | -11.0            | -5.0             | -0.77           | -0.85            | -0.79            |
| SON | -0.26           | -0.38            | -0.32            | -2.2            | -5.5             | -1.4             | -0.94           | -0.55            | -0.73            |
|     | IRE             |                  |                  |                 |                  |                  |                 |                  |                  |
| ANN | -0.66           | -0.23            | -1.0             | -7.7            | -4.6             | -8.7             | -0.41           | -0.08            | -0.43            |
| DJF | -0.64           | -0.28            | -0.57            | -10.8           | -4.8             | -9.2             | -0.43           | -0.06            | -0.31            |
| MAM | -0.94           | -0.21            | -1.4             | -9.9            | -4.9             | -10.7            | -0.47           | -0.07            | -0.58            |
| JJA | -0.36           | -0.07            | -0.78            | -4.5            | -3.7             | -7.3             | -0.30           | -0.07            | -0.46            |
| SON | -0.70           | -0.38            | -1.4             | -5.7            | -5.1             | -7.5             | -0.45           | -0.13            | -0.37            |

a: Mean over oceanic areas.

b: 27~40°N, 115~123°E.

c: 35~55°N, 140~160°E.