



Peatland evaporation across hemispheres: contrasting controls and sensitivity to climate warming driven by plant functional types

Leeza Speranskaya¹, David I. Campbell¹, Peter M. Lafleur², Elyn R. Humphreys³

¹ Environmental Research Institute, Faculty of Science and Engineering, University of Waikato, Hamilton, 3216, New Zealand

² School of Environment, Trent University, Peterborough, ON K9L 0G2, Canada

³ Geography and Environmental Studies, Carleton University, Ottawa, ON K1S 5B6, Canada

Correspondence to: David I. Campbell (david.campbell@waikato.ac.nz)

Abstract. Peatlands store disproportionately large amounts of carbon per unit area, a function that is dependent on maintaining high and stable water tables. Climate change is likely to negatively impact carbon storage in peatlands, in part due to increases in vapour pressure deficit (VPD) driving higher evaporation (E) rates. However, the response of E to increasing VPD depends on the dominant vegetation type within peatlands. In this study, we used multiple years of eddy covariance (EC) measurements to compare E regimes at two peatlands with contrasting vegetation types - Kopuatai bog in Aotearoa New Zealand, dominated by the vascular jointed wire rush *Empodisma robustum*, and Mer Bleue bog in Canada, a ‘typical’ shrub and moss-dominated Northern Hemisphere peatland. We examined seasonal variability in E and equilibrium E (E_{eq}), energy balance partitioning, and the response of E , evaporative fraction (EF), and canopy conductance (g_c) to VPD. Mean annual E was 45 % lower than mean annual E_{eq} at Kopuatai, but only 16 % lower at Mer Bleue, demonstrating much greater limitations on E at Kopuatai. In addition, the mean midday (10:00-14:30) dry canopy Bowen ratio (β) at Kopuatai was 1.96, compared to 0.77 at Mer Bleue; therefore, the sensible heat flux (H) dominated over the latent heat flux (LE) at Kopuatai, and vice versa at Mer Bleue. The responses of E , EF, and g_c to increasing VPD at Kopuatai differed from those at Mer Bleue in a way that resulted in stronger limitations on E above ~ 0.7 kPa at the former bog. The observed limitations at Kopuatai were attributed to strong stomatal control by *E. robustum* due to the rapid decrease in g_c with increasing VPD, however surface E could also be limited by its dense standing litter. At Mer Bleue, however, E was only weakly limited at VPD > 2 kPa, likely due to weak stomatal control over transpiration by the sparse shrub canopy and relatively large surface E from *Sphagnum* carpets. As such, the results of this study suggest that *E. robustum* drives a greater “hydrological resistance” to increasing VPD than the vegetation at Mer Bleue, leading to greater water retention at Kopuatai. This may enable greater resilience of the carbon sink function at Kopuatai to climatic warming and drying than at Mer Bleue.

1 Introduction

Global peatlands have accumulated carbon (C) stocks of 500 Pg or more over thousands of years, resulting in a net cooling effect on the climate (Yu et al., 2010; Frolking and Roulet, 2007). This C stock is estimated to make up a third of global soil C (based on a median estimate of 1460.5 Pg), despite only occupying ~ 3 % of Earth’s land surface area (Scharlemann et al.,



2014; Xu et al., 2018). Carbon accumulation as peat occurs in undisturbed peatlands when the rate of organic matter deposition exceeds decomposition, due to limitations imposed on decomposition by waterlogged conditions and the presence of decay-resistant plant litter (Joosten and Clarke, 2002; Holden, 2005). Therefore, long-term C sequestration in peatlands depends on maintaining a high and stable water table (Kim et al., 2021; Ma et al., 2022).

40

Evaporation (E)^a is an important indirect influence on peat accumulation, as it reduces available water and contributes to water table drawdown. Evaporative fluxes consist of three main components - transpiration, surface E (either from water, soil, or non-vascular plant surfaces), and interception loss from vegetation surfaces, and are influenced by a number of meteorological (solar radiation, vapour pressure deficit (VPD)) and surface factors (vegetation type and water table depth (WTD); Takagi et al., 1999; Shimoyama et al., 2003; Wu et al., 2010). In bog ecosystems, where precipitation is the sole water input (Holden, 2005), conservation of water is particularly important; this occurs through limitations imposed on E by bog vegetation (Campbell and Williamson, 1997), as well as limitations on lateral and vertical drainage by low hydraulic conductivity in the highly decomposed peat soils (Fraser et al., 2001). As such, vegetation type is often a major factor regulating E from bogs (Lafleur and Roulet, 1992; Takagi et al., 1999; Admiral et al., 2006).

50

The response of E to increasing atmospheric demand, i.e., increasing VPD, is of particular interest, as climate change-related warming has been predicted to elevate VPD (Ficklin and Novick, 2017; Fang et al., 2022). As VPD increases, contrasting E responses have been observed across different vegetation types (Takagi et al., 1999; Admiral et al., 2006; Massman et al., 2019) and ecosystems (Helbig et al., 2020). For example, using eddy covariance (EC) data from 95 boreal peatland and forest sites, Helbig et al. (2020) demonstrated that E rates were elevated by increasing VPD in both ecosystem types, however E rates were 30 % higher in boreal peatlands than in boreal forests at high VPD. This suggests that the stability of peatland hydrology and C stores may be at risk under future changes to the climate.

55

Kopuatai bog is a warm-temperate, ombrotrophic peatland on Te Ika-a-Māui North Island of Aotearoa New Zealand. The vegetation at Kopuatai is dominated by the vascular plant *Empodisma robustum*, which has been implicated in severely restricting E (Campbell and Williamson, 1997). The hypothesised mechanisms for this are stomatal control of transpiration and limitation of surface E due to restricted water vapour diffusion from the substrate through the dense standing litter layer (Campbell and Williamson, 1997; Thompson et al., 1999). Studies on ecosystem C balances have shown that C uptake at Kopuatai is greater than in analogous Northern Hemisphere peatlands, and exhibits resilience to drought (i.e., high annual C uptake rates despite drought disturbance in summer; Goodrich et al., 2017). This could be a result of the year-round growing conditions (Campbell et al., 2014), in contrast to the temperature-limited growing season lengths in Northern Hemisphere peatlands which constrain C uptake (Roehm and Roulet, 2003; Lafleur et al., 2001; Helfter et al., 2015). However, high C

65

^a 'Evaporation' refers to "the bulk flux of water, including transpiration". We did not use the term 'evapotranspiration', as both evaporation and transpiration involve transformation of water from a liquid to a vapour (Miralles et al., 2020).



uptake rates and drought resilience could also be indirectly attributed to the extremely low E , which maximises available water and maintains a high and stable water table. Therefore, the *resilience* of C uptake at Kopuatai could be due to
70 *resistance* of the hydrology to environmental stressors through limitations on E , thereby preventing major shifts in the hydrological regime (resistance and resilience definitions are from Nimmo et al., 2015). As such, E limitation could be a very important self-regulation mechanism in this peatland ecosystem type.

Peatlands have formed in warm climates, however the majority of the global peatland area occurs in boreal regions of the
75 Northern Hemisphere (Vitt, 2006). One such peatland is the Mer Bleue bog in Canada, a shrub and *Sphagnum*-dominated ecosystem located in a cool continental climate zone (Moore et al., 2002). As one of the most studied Northern Hemisphere peatlands, Mer Bleue has an extensive record of E measurements and knowledge of E processes. Daily E rates at Mer Bleue are comparable to many other Northern Hemisphere peatlands; E has been shown to be reduced by deep water tables (Lafleur et al., 2005), and regulated by available energy, which drives increases in E , and by VPD, the effect of which varies
80 depending on the partitioning of E between vascular and non-vascular sources (Admiral et al., 2006). The average net annual C uptake at Mer Bleue is much lower than at Kopuatai, primarily due to net C losses during the winter period, and a shorter growing season (Roulet et al., 2007; Goodrich et al., 2017).

Knowledge of E at Kopuatai bog is limited to the results of two short-term studies (Campbell and Williamson, 1997;
85 Thompson et al., 1999); therefore, long-term E datasets from Kopuatai have not yet been examined, so our understanding of the variability and mechanisms of E restriction by *E. robustum* is incomplete. In addition, it is not yet known how these limitations on E may affect the response of the ecosystem at Kopuatai to climate warming. Therefore, the aim of this study is to compare the E regime at Kopuatai to that of Mer Bleue, as an example of a 'typical' Northern Hemisphere peatland, to better understand the potential relative sensitivity of these systems to hydrological change under continued climate warming.
90 Compared with previous studies, our study benefits from extensive multi-annual data records at each site, thereby analysing E characteristics over a wide range of weather conditions. The objectives of this study are to (1) compare seasonal variability in actual E and equilibrium E (E_{eq}) at Kopuatai and Mer Bleue, (2) investigate energy balance partitioning and (3) examine the responses of E , evaporative fraction (EF), and canopy conductance (g_c) to VPD at each site. Our hypothesis is that differences in seasonal E limitations, energy balance partitioning, and responses of E to VPD, such that more water is
95 conserved at Kopuatai relative to Mer Bleue, can be attributed to differences in vegetation. If this hypothesis is correct, it is possible that the water-conserving traits of *E. robustum* will enable greater resistance of the hydrological regime, and hence greater resilience of the C stores, to climate warming compared to typical Northern Hemisphere peatlands.



2 Methodology

2.1 Site descriptions

100 2.1.1 Kopuatai bog

Kopuatai bog is a 96 km² ombrotrophic peatland located in a warm-temperate oceanic climate in the Waikato region of Aotearoa New Zealand (37.388° S, 175.554° E). Kopuatai is Aotearoa's largest undisturbed bog, representing a remnant of formerly widespread and diverse lowland wetlands, which have been reduced to 10 % of their previous extent due to widespread drainage, primarily for agriculture (McGlone, 2009; Dymond et al., 2021). Probing at the research site showed a
105 mean peat depth of 11 m (maximum depth = 14 m), which has accumulated over the last 11,700 years at an average rate of 0.9 mm year⁻¹ (Newnham et al., 1995; Shearer, 1997).

The dominant peat-forming vegetation at Kopuatai is *E. robustum* (Wagstaff and Clarkson, 2012), a vascular, evergreen jointed rush-like plant that is part of the family Restionaceae (known as 'restiads'; Wagstaff and Clarkson, 2012). At the
110 field site, *E. robustum* has a mean leaf area index (LAI) of 1.32 (Goodrich et al., 2015) and mean canopy height of 0.48 m. *E. robustum* forms a dense, negatively geotropic surface root mat approximately 50 mm deep which can hold up to 15 times its dry weight in water (Campbell, 1964; Agnew et al., 1993; Clarkson et al., 2009). In addition, the standing litter of *E. robustum* intercepts a considerable quantity of precipitation (Campbell and Williamson, 1997). It has been hypothesised that
115 this dense standing litter layer (litter biomass = 0.92 kg m⁻², where total canopy biomass = 1.80 kg m⁻²) can also restrict water vapour diffusion from the peat surface by limiting turbulent transport (Campbell and Williamson, 1997; Keyte-Beattie, 2014). However, *E. robustum* can also restrict water loss via transpiration through strong stomatal control (Campbell and Williamson, 1997). The roots of *E. robustum* are the main material from which peat is formed, however live mosses and decayed canopy litter are also bound by the roots for peat formation (Campbell, 1964; Agnew et al., 1993). Other vegetation
120 types found at this site include sedges (*Machaerina* spp. and *Schoenus brevifolius*), isolated shrubs of *Leptospermum scoparium* (manuka) and *Epacris pauciflora*, as well as small patches of *Sporadanthus ferrugineus*, another restiad plant.

2.1.2 Mer Bleue bog

Mer Bleue bog is a 28 km² ombrotrophic peatland (45.411° N, -75.481° E) located near Ottawa, Canada, in a cool continental climate. Mer Bleue has many features in common with boreal peatlands, but is located near the southern limit of the boreal climate zone (Hember et al., 2005). This peatland was formed 8400 years ago, initially as a fen, transitioning into a bog
125 around 7100–6800 years ago (Roulet et al., 2007). At the research site, Mer Bleue is characterised by a hummock-hollow microtopography, and has peat depths of 5–6 m (Lafleur et al., 2005). The dominant vegetation consists of evergreen ericaceous and deciduous shrubs (e.g., *Chamaedaphne calyculata*, *Rhododendron groenlandicum*, *Kalmia angustifolium*, *Kalmia polifolia*, *Vaccinium myrtilloides*), with *Sphagnum* moss species as the dominant ground cover, including *S.*



130 *capillifolium*, *S. papillosum*, and *S. magellanicum* (Bubier et al., 2006). The average shrub canopy height is 0.18 m (range = 0.10–0.30 m; Bubier et al., 2006); shrub vegetation makes up 61 % of the total biomass on average, while *Sphagnum* capitula make up 30 % (Moore et al., 2002). The total biomass ranges between 0.147 and 1.011 kg m⁻², while the LAI of dominant vascular species is 1.3 on average (Moore et al., 2002; Bubier et al., 2006).

2.2 Data collection and processing

2.2.1 Data collection

135 Data has been collected at both sites for many years - in this study, data collected between 1 January 2012 and 31 December 2022 at Kopuatai and 1 January 1999 and 31 December 2018 at Mer Bleue were used. At each site, 30-minute fluxes of latent heat (*LE*), sensible heat (*H*) and net radiation (*R_n*), along with data on environmental and weather variables, including air temperature (*T_{air}*), vapour pressure deficit (*VPD*), precipitation (*P*), and water table depth (*WTD*; measured relative to the hummock surface at Mer Bleue), were collected. Latent and sensible heat fluxes were measured using the eddy covariance
140 (EC) technique (Burba, 2022). The EC system at Kopuatai is an open path system (Appendix A) situated at 4.25 m height, with uninterrupted fetch greater than 500 m in all directions. At Mer Bleue, a closed path EC system is situated at 3.0 m height, with a fetch greater than 500 m in all directions except south, where it is 200–300 m. Full details of data processing, quality control, and gap-filling are provided in Goodrich et al. (2017) for Kopuatai. For gap-filling of *LE* and *H*, separate daytime and night-time neural network models used drivers of *R_n*, *T_{air}*, *VPD*, and modelled canopy wetness state (detailed
145 below). At Mer Bleue, gap-filling of *LE* was carried out by developing a linear relationship between available energy (*R_a*) and *LE* for summer, and calculating a multiplier to adjust the estimated *LE* to observed *LE* over a moving window period (the window width is 100 consecutive available half hours moved in increments of 20 half hours). *R_a* was calculated as *R_n* minus the rate of change in energy storage terms (*J*) and the ground heat flux (*G*), which was calculated using peat surface temperature profiles. Sensible heat fluxes were then calculated as the difference between *R_a* and *LE*.

150 2.2.2 Equilibrium evaporation

To provide a reference *E* rate at each site, the equilibrium evaporation (*E_{eq}*) was calculated using gap-filled 24-hour mean data via the following equation:

$$E_{eq} = \frac{s(R_n - G)}{L_v(s + \gamma)} \quad (1)$$

155 where *s* is the slope of the saturation vapour pressure versus air temperature curve, *L_v* is the latent heat of vaporisation (dependent on *T_{air}*), and γ is the psychrometric constant (0.066 kPa °C⁻¹). *G* was assumed to be 0 W m⁻² due to the use of 24-hour mean data.



The proportion E/E_{eq} , i.e. the Priestley–Taylor α , was then calculated in order to be able to compare E regimes at the two sites despite differing climates. Gap-filled LE data were converted to $E = LE/L_v$ for this analysis.

160

E_{eq} was selected as a measure of potential E as it provides a more conservative estimate compared to other methods, such as the Penman equation (Granger, 1989). In addition, E_{eq} is less dependent on energy partitioning compared to Penman open water E (Thompson et al., 1999). As explained by Thompson et al. (1999), due to the high ratio of sensible to latent heat flux (i.e., a high Bowen ratio (β)) at Kopuatai, VPD is potentially larger than it would be above an extensive water surface in the same region, driving Penman open water E to an unrealistic level. Due to the lack of a VPD term in the E_{eq} equation, this effect is mitigated.

165

2.2.3 Energy balance partitioning and VPD analysis

The analysis of energy balance partitioning and the relationships of E , EF , and g_c with VPD involved the use of non-gap-filled E , R_n , LE , H , and VPD data, pre-filtered through quality control processes. These data were then filtered further - firstly, data were filtered by growing season months (May–Oct at Mer Bleue and Sep–May at Kopuatai). In addition, all 30-minute data from both sites were filtered to select only “middle of day” (MoD) conditions (10:00–14:30 local standard time), and $R_n \geq 200 \text{ W m}^{-2}$. These 30-minute data were also filtered by dry or wet canopy conditions, which were identified using an antecedent precipitation index (API) at both Kopuatai and Mer Bleue (Appendix B). An $API \leq 0.2$ signifies a dry canopy, while $API \geq 1$ indicates fully wet canopy conditions. These filtered 30-minute data were then used to calculate MoD means of all variables. Days with less than three acceptable 30-minute data points for all variables were removed to ensure representative MoD means. Overall, this filtering resulted in 56 % and 65 % of MoD 30-minute data being rejected for Kopuatai and Mer Bleue, respectively.

170

175

To analyse energy balance partitioning, the filtered growing season MoD mean LE and H data were binned by R_n (bin width = 50 W m^{-2}) for each site, taking the mean value of each bin. This was done separately for dry and wet canopy conditions. Seasonal energy balance partitioning was also investigated by calculating monthly means of MoD dry canopy LE , H , and R_n .

180

To show the relationship of variables with VPD, filtered dry canopy growing season MoD means of E , EF , and g_c were binned by VPD (0.1 kPa intervals), following Helbig et al. (2020), taking the mean of each bin for all variables. EF was used to assess the change in energy balance partitioning into LE with increasing VPD, and was calculated as:

185

$$EF = \frac{LE}{LE + H} \quad (2)$$



The denominator $LE + H$ was used rather than R_n to prevent incomplete energy balance closure (Appendix C) from affecting our results. In addition, β was calculated as:

190

$$\beta = \frac{H}{LE} \quad (3)$$

Finally, g_c was calculated as in Campbell and Williamson (1997):

$$\frac{1}{g_c} = \frac{(1 + \beta) \rho c_p VPD}{\gamma R_n} + \frac{r_a s \beta}{\gamma} - r_a \quad (4)$$

195 where ρ is air density (1.2 kg m^{-3}), c_p is the specific heat of air ($1010 \text{ J kg}^{-1} \text{ }^\circ\text{C}^{-1}$), and r_a is aerodynamic resistance. Values of r_a were also calculated using the method from Campbell and Williamson (1997), which firstly required calculation of the aerodynamic resistance to momentum transfer from the measurement height to the canopy (r_{aM}):

$$r_{aM} = \frac{u}{u_*^2} \quad (5)$$

200 where u is the horizontal wind speed and u_* the friction velocity measured by the sonic anemometer.

Then, r_a was calculated as:

$$r_a = \frac{1.6}{k u_*} + r_{aM} \quad (6)$$

where k is von Karman's constant (0.4).

2.2.4 Energy balance closure

205 Energy balance closure was imperfect at both sites, with 85 % closure at Kopuatai and 89 % at Mer Bleue (Appendix C). This could be due to a number of factors, such as mismatch between footprint characteristics for the turbulent and non-turbulent energy balance terms, or increasing uncertainty of surface soil heat flux measurements due to peat accumulation, which increases the depth of soil heat flux plates over time. Energy imbalances could also be a result of a lack of sustained turbulence at these sites (Anderson and Wang, 2014), or due to the use of MoD means of energy balance components rather
210 than 24-hour means (Leuning et al., 2012).


All analyses were carried out using MATLAB R2021b.



3 Results

3.1 Climate and hydrology

215 Mean annual precipitation (P) was 1213 ± 147 mm (± 95 % confidence interval) at Kopuatai (2012–2022), compared to 879
 ± 58.1 mm at Mer Bleue (1999–2018). Mean monthly P ranged between 49–139 mm and 50–100 mm at Kopuatai and Mer
Bleue, respectively, with the minimum monthly P occurring during the warm season at Kopuatai and the cool season at Mer
Bleue (Fig. 1a). The mean monthly air temperature (T_{air}) range was much smaller at Kopuatai (9.3–19.0 °C) than at Mer
Bleue (-10.0–20.2 °C) (Fig. 1b), with mean annual temperatures of 14.1 ± 0.27 °C and 6.2 ± 0.31 °C at the two sites,
220 respectively. Mean annual VPD was 0.39 ± 0.013 kPa at Kopuatai and 0.38 ± 0.023 kPa at Mer Bleue, and monthly mean
VPD ranged between 0.17–0.68 kPa and 0.07–0.77 kPa (Fig. 1c). Mean annual WTD was -82.6 ± 14.4 mm at Kopuatai and
 -377 ± 19.8 mm at Mer Bleue. Monthly mean WTD had narrower ranges and was shallower at Kopuatai than at Mer Bleue
year-round; as a result, the WTD ranges at these sites did not overlap (-153 to -28 mm and -460 to -278 mm, respectively;
Fig. 1d).



225

Figure 1. Mean monthly (a) precipitation (b) air temperature (c) vapour pressure deficit, and (d) water table depth at Kopuatai (orange) and Mer Bleue (blue). These values were calculated using data between 1999–2018 (inclusive) at Mer Bleue and 2012–2022 (inclusive) at Kopuatai. Error bars are 95 % confidence intervals. Note that the Northern and Southern Hemisphere seasons have been aligned by using separate *x*-axes for each site (Jan-Dec for Mer Bleue and Jul-Jun for Kopuatai).

230 **3.2 Evaporation and equilibrium evaporation**

Mean annual E and E_{eq} were larger at Kopuatai compared to Mer Bleue, however mean annual E/E_{eq} was lower at Kopuatai (Table 1). In addition, growing season and year-round mean E/E_{eq} values were very similar at each respective site. At both sites, mean monthly E was below E_{eq} , except during winter (Fig. 2). Both variables followed an expected seasonal pattern of low values in winter and high values in summer, mostly driven by radiation receipts. The difference between E_{eq} and E was larger at Kopuatai than at Mer Bleue in non-winter months; the maximum difference between monthly E_{eq} and E was 71 mm in January at Kopuatai, but only 21 mm at Mer Bleue in May. Both E and E_{eq} were close to zero at Mer Bleue during winter




(Dec–Feb), when air temperature was below 0 °C and the bog was snow covered, and both values were similar during spring (March) and autumn (October) months.

240 **Table 1.** Mean annual ecosystem evaporation (E) and equilibrium evaporation (E_{eq}), and means and ranges of their ratios (i.e. Priestley–Taylor α) at each site ($n = 11$ and 20 years for Kopuatai and Mer Bleue, respectively). These values are given for both year-round data and for the growing season only (Sep–May at Kopuatai and May–Oct at Mer Bleue). Values in parentheses are 95 % confidence intervals.

Site	Annual				Growing season only			
	E (mm)	E_{eq} (mm)	E/E_{eq}	E/E_{eq} range	E (mm)	E_{eq} (mm)	E/E_{eq}	E/E_{eq} range
Kopuatai	578 (± 13)	910 (± 18)	0.64 (± 0.022)	0.57–0.70	482 (± 11)	816 (± 19)	0.59 (± 0.023)	0.52–0.65
Mer Bleue	450 (± 17)	530 (± 13)	0.85 (± 0.026)	0.76–0.95	388 (± 16)	466 (± 11)	0.83 (± 0.027)	0.70–0.94

245



250 **Figure 2.** Mean monthly E and E_{eq} at (a) Kopuatai and (b) Mer Bleue. Error bars represent 95 % confidence intervals. Note that the x-axis for Kopuatai in the Southern Hemisphere starts from July (a six-month offset from the graph for Mer Bleue in the Northern Hemisphere).



3.3 Dry canopy energy balance partitioning

Seasonal energy balance partitioning was markedly different at each site. At Kopuatai, H was greater than LE in most months, with LE only comprising 21 % of the annual R_n (Fig. 3). Between autumn and winter (April to August), however, the difference between H and LE was smaller. In contrast, LE was a much larger component of the energy balance at Mer
 255 Bleue, representing 43 % of the annual R_n due to LE exceeding H throughout most of the growing season. During the rest of the year, however, LE was either less than or similar to H . There was also a difference in the seasonality of LE , H , and R_n between sites - at Kopuatai, all components of the energy balance reached a maximum during summer; LE and R_n peaked in mid-summer (January), while H peaked a month earlier. At Mer Bleue, LE and R_n also peaked during summer (in June and July, respectively), however H peaked in the middle of spring (April).

260

Mean monthly LE increased substantially (range = 257 W m^{-2}) towards summer at Mer Bleue, while at Kopuatai, there was less monthly variation in LE (range = 79 W m^{-2}). Although the maximum monthly R_n ($\pm 95 \%$ confidence interval) at Kopuatai was higher than at Mer Bleue ($578 \pm 26.0 \text{ W m}^{-2}$ and $480 \pm 17.7 \text{ W m}^{-2}$, respectively), maximum monthly LE at Kopuatai ($124 \pm 6.45 \text{ W m}^{-2}$) was lower than at Mer Bleue ($264 \pm 17.0 \text{ W m}^{-2}$). Consequently, the maximum monthly H was
 265 higher at Kopuatai than at Mer Bleue ($304 \pm 30.5 \text{ W m}^{-2}$ and $191 \pm 7.57 \text{ W m}^{-2}$, respectively). In addition, R_n was particularly low in winter at Mer Bleue due to snow cover and the lower solar receipt. During spring, R_n increased steeply (March and April) due to a decrease in albedo resulting from snowmelt. In contrast, Kopuatai does not receive snow, and seasonal albedo does not vary substantially (not shown).

270




Figure 3. Mean monthly LE , H , and R_n at (a) Kopuatai and (b) Mer Bleue for middle-of-day, dry canopy conditions. Error bars are 95 % confidence intervals. Note that the x-axis for Kopuatai starts from July (a six-month offset from the graph for Mer Bleue).



275 Daily mean energy balance partitioning during dry canopy conditions was also distinctly different between Kopuatai and Mer Bleue (Fig. 4). At Kopuatai, H was a much larger component of the energy balance than LE (Fig. 4a); H was greater than LE at $R_n \geq 250 \text{ W m}^{-2}$, which was reflected in a mean Bowen ratio (β) of 1.96 (β range based on binned H and LE values was 0.37–2.91). At Mer Bleue, however, LE was greater than H at $R_n \geq 350 \text{ W m}^{-2}$ (Fig. 4b). As a result, mean β was 0.77, with a range of 0.61–1.05.

280




Figure 4. Relationship between binned middle-of-day mean LE , H , and R_n for dry canopy conditions at (a) Kopuatai and (b) Mer Bleue. Values are for the growing season only (Sep–May at Kopuatai and May–Oct at Mer Bleue). Error bars are standard deviations.

285 3.4 Wet canopy energy balance partitioning

During wet canopy conditions, the limitation on LE observed at Kopuatai under dry canopy conditions was not apparent (Fig. 5a); LE was greater than H at both low and high R_n in these conditions (β ranged between -0.01–0.93, with mean 0.51). Both LE and H increased at approximately the same rate with increasing R_n . At Mer Bleue, the relationship between LE , H , and R_n under wet canopy conditions was similar to dry canopy conditions, with a lower mean β of 0.57 and a range of -0.38–

290 0.83 (Fig. 5b).




Figure 5. Relationship between binned daily middle-of-day mean LE , H , and R_n for wet canopy conditions at (a) Kopuatai and (b) Mer Bleue. Values are for the growing season only (Sep–May at Kopuatai and May–Oct at Mer Bleue). Error bars are standard deviations.

295 3.5 Response of E , EF , and g_c to VPD

E increased with increasing VPD at both sites, i.e. greater water loss occurred at high VPD (Fig. 6). Below 2.0 kPa, the rate of increase in E with increasing VPD was 3.7 times larger at Mer Bleue (0.221 mm hr⁻¹ / kPa) than at Kopuatai (0.060 mm hr⁻¹ / kPa). This resulted in a higher E of 0.44 mm h⁻¹ at VPD = 2.0 kPa at Mer Bleue, compared to only 0.23 mm h⁻¹ at the same VPD at Kopuatai. Beyond VPD = 2.0 kPa, E showed little or no change with increasing VPD at both sites.

300





Figure 6. Middle-of-day mean growing season E for 0.1 kPa bins of VPD during dry canopy conditions at Kopuatai and Mer Bleue. Shaded areas represent 95 % confidence intervals.

305

At Kopuatai, EF decreased in response to increasing VPD, while the opposite trend was observed at Mer Bleue (Fig. 7). This contrast in responses of EF to increasing VPD was also evident in 30-minute data (Appendix D). At Kopuatai, mean daytime EF decreased from 0.44 at VPD = 0.5 kPa to 0.28 at 2.7 kPa, and increased from 0.31 at 0.4 kPa to 0.66 at 3.3 kPa at Mer Bleue bog. At both sites, the rate of increase or decrease in EF was reduced at VPD > 1.0 kPa.

310



315 **Figure 7. Middle-of-day mean growing season EF for 0.1 kPa bins of VPD during dry canopy conditions at Kopuatai and Mer Bleue. Shaded areas represent 95 % confidence intervals.**

There was a steep decline in g_c with increasing VPD at Kopuatai (Fig. 8); mean g_c decreased from a maximum of 10.6 mm s^{-1} at a VPD of 0.5 kPa to a minimum of 2.57 mm s^{-1} at 2.7 kPa . In contrast, the pattern of declining g_c with increasing VPD was weak at Mer Bleue, decreasing from 9.10 mm s^{-1} at 0.4 kPa to 6.23 mm s^{-1} at 3.3 kPa .

320




Figure 8. Middle-of-day mean growing season g_c for 0.1 kPa bins of VPD during dry canopy conditions at Kopuatai and Mer Bleue. Shaded areas represent 95 % confidence intervals.

325 4 Discussion

4.1 Evaporation regimes

Our study demonstrates the crucial importance of vegetation characteristics in controlling water loss from peatlands. Kopuatai and Mer Bleue, two ombrotrophic bogs with different vegetation communities, display strikingly different E regimes, energy balance partitioning, and responses of E , EF , and g_c to VPD. Despite being in a warmer climate zone with warm winters, the E regime at Kopuatai was more conservative than at Mer Bleue. Although mean annual E and E_{eq} were 25 % and 53 % higher at Kopuatai than at Mer Bleue, mean annual E/E_{eq} (i.e., Priestley–Taylor α) was 28 % lower at 330 Kopuatai. This occurred because the mean annual E was 45 % lower than mean annual E_{eq} at Kopuatai, while at Mer Bleue there was only a 16 % difference. The large difference between E and E_{eq} at Kopuatai occurred due to the relatively high E_{eq} , resulting from the higher R_n (due to latitudinal differences), warmer climate, and longer growing season compared to Mer 335 Bleue. These results indicate that greater surface limitations were imposed on E at Kopuatai relative to Mer Bleue. The difference in E/E_{eq} between the two sites is highly likely to be a result of differences in vegetation, as different vegetation



communities, even at the same peatland, have been shown to result in different E regimes (Takagi et al., 1999; Strilesky and Humphreys, 2012). The limitations on E at Kopuatai likely occurred due to the physiological and canopy structural features of the dominant vascular plant, *E. robustum*. However, some limitation on E was observed at Mer Bleue, as evidenced by the decrease in E/E_{eq} towards mid-summer. It is possible that this occurred due to stomatal closure in the shrub vegetation in response to high VPD (Admiral et al., 2006). However, this limitation was weaker than at Kopuatai, likely due to high surface E from *Sphagnum* carpets (Admiral et al., 2006).

At Kopuatai, the range of E/E_{eq} values tended to be lower than at most Northern Hemisphere peatlands, while the E/E_{eq} range at Mer Bleue overlapped with the lower end of most of these ranges (Table 2). However, there were some sites with E/E_{eq} ranges that were closer to those of Kopuatai. For example, a tree-dominated peatland in Canada had a lower E/E_{eq} than at Kopuatai, which could be attributed to stomatal limitation of transpiration (Brümmer et al., 2012), i.e., the same mechanism of E limitation as exhibited at Kopuatai, but to a greater extent. Of interest is also the Fäjemyr bog in Sweden, which had lower growing season E/E_{eq} values than most peatlands in Table 2 (but not Kopuatai), possibly due to a low water table and low surface conductance (Alekseychik et al., 2018). In addition, in an earlier study of E rates at Kopuatai, Thompson et al. (1999) reported lower E/E_{eq} values than those in this study; this could have occurred due to measurements being made only during summer, when limitations on E are most prominent. Apart from these outliers, the E/E_{eq} values in Table 2 are generally much greater than at Kopuatai bog, indicating that *E. robustum* may be more adept at restricting E than the vascular and non-vascular vegetation in most Northern Hemisphere peatlands.

4.2 Controls on evaporation

The steep decline in EF with increasing VPD at Kopuatai suggests strong surface controls on E at this site. A similar negative relationship between midsummer LE/R_n and VPD was shown by Takagi et al. (1999) at a section of a peatland invaded by vascular vegetation, which was attributed to stomatal closure. This is also likely to be the main factor limiting E from Kopuatai due to the non-linear decrease in canopy conductance (g_c) with increasing VPD, a trend that has also been observed in other peatlands (Humphreys et al., 2006; Peichl et al., 2013; Runkle et al., 2014; Alekseychik et al., 2018). However, it has also been hypothesised that the standing litter layer formed by *E. robustum* reduces E (Campbell and Williamson, 1997); this canopy feature has been observed to limit E in marsh, restored wetland, and dryland ecosystems (Goulden et al., 2007; Villegas et al., 2012; Eichelmann et al., 2018). Standing litter restricts E by decoupling the subcanopy environment from atmospheric turbulence, and by preventing solar radiation from reaching the surface, which minimises the variability of T_{air} and VPD below the litter layer (Goulden et al., 2007; Eichelmann et al., 2018). Although the results of this study cannot separate the effect of stomatal closure and standing litter on E , it is likely that both of these factors act in tandem to severely restrict E at Kopuatai.



370 **Table 2. Ranges of E/E_{eq} calculated for peatlands globally (including this study), with a description of the type of variability covered in the range of E/E_{eq} values, and the time period for which they were calculated.**

Location	Dominant vegetation types	E/E_{eq} (i.e., Priestley–Taylor α)	Range type	Time period	Reference
Kopuatai bog, Aotearoa	<i>Empodisma robustum</i>	0.57–0.70	Interannual variability	Year-round	This study
		0.52–0.65	Interannual variability	Growing season (Sep–May)	This study
	<i>Empodisma robustum</i>	0.34	N/A	Growing season (Nov–Mar)	Thompson et al. (1999)
	<i>Sporadanthus ferrugineus</i> ^a	0.58	N/A	Summer period (Jan–Mar)	Thompson et al. (1999)
Mer Bleue bog, Canada	Shrubs and <i>Sphagnum</i> moss	0.76–0.95	Interannual variability	Year-round	This study
		0.70–0.94	Interannual variability	Growing season (May–Oct)	This study
Plotnikovo, Russia	Sedges, shrubs, and <i>Sphagnum</i> moss	0.99–1.29 (1999) 0.96–1.07 (2000)	Seasonal variability	Growing season (Apr–Oct)	Shimoyama et al. (2004)
Sandhill fen, Canada	Trees, shrubs, and brown moss	0.79–1.04	Interannual variability	Snow-free periods (May–Nov)	Sonnentag et al. (2010)
Alberta, Canada	Trees, shrubs, and various mosses	0.55–0.57	Interannual variability	Year-round	Brümmer et al. (2012)
Degerö Stormyr, Sweden	Shrubs, grasses, and <i>Sphagnum</i> moss	0.86–1.17	Interannual variability	Year-round	Peichl et al. (2013)
Siikaneva-1 & Siikaneva-2, Finland	Shrubs, sedges, and <i>Sphagnum</i> moss	1.09–1.21 & 1.11–1.13	Interannual variability	Growing season (May–Oct)	Alekseychik et al. (2018)
Fäjemyr bog, Sweden	Shrubs, sedges, <i>Sphagnum</i> moss, and sparse trees	0.66–0.71	Interannual variability	Growing season (May–Oct)	Alekseychik et al. (2018)
Seven peatlands in Canada	Shrubs, sedges, mosses (all peatlands), trees, and herbs (some peatlands)	0.82–1.05	Spatial variability	Midsummer period (Jul–Aug)	Humphreys et al. (2006)

^a A tall restiad plant that dominates portions of Kopuatai bog, but is not present at the EC site



375 At Mer Bleue, E rates were almost double those of Kopuatai at high VPD, and there was a greater absolute increase in dry
canopy LE with increasing R_n at Mer Bleue than at Kopuatai (an increase of 254 W m^{-2} compared to 93 W m^{-2} , respectively),
indicating weaker water loss restrictions at Mer Bleue. Weak restrictions likely occurred because of the large contribution of
Sphagnum moss E during well-watered conditions, as moss E is enhanced substantially by increasing VPD (Admiral and
Lafleur, 2007). Since LE increased with increasing R_n and E and EF increased with increasing VPD, both of these factors
380 were important controls on E and energy partitioning at Mer Bleue. Similar positive correlations between LE and R_n (or
available energy, $R_n - G$) at daily and monthly scales have previously been observed at other peatlands (Lafleur and Roulet,
1992; Kurbatova et al., 2002; Brümmer et al., 2012), in addition to positive relationships between daily E and VPD (Wu et
al., 2010; Peichl et al., 2013; Wang et al., 2020).

385 There were also reduced rates of increase in E and EF with increasing VPD at Mer Bleue (i.e., a ‘saturating’ effect; also
observed by Peichl et al., 2013), which only became evident at high VPD (above $\sim 2.0 \text{ kPa}$ for E and $\sim 1.0 \text{ kPa}$ for EF ; Figs. 6
and 7). This relatively weak E limitation was likely caused by stomatal control of transpiration by vascular vegetation
(Admiral et al., 2006), as a slight decrease in g_c was observed with increasing VPD. A similar relationship between surface
conductance and VPD has previously been observed at Mer Bleue (Humphreys et al., 2006). It has also been suggested that
390 E may become limited at high VPD if the transport rate of water through mosses is insufficient to meet the atmospheric
demand (Liljedahl et al., 2011). While we cannot determine whether this occurred at Mer Bleue using the data available, it is
possible that this may have been a contributing factor to limitations on E .

While there was almost no difference in energy balance partitioning between dry and wet canopy conditions at Mer Bleue,
395 the canopy wetness state had a major effect at Kopuatai. During dry canopy conditions, H was the dominant convective flux,
as the rate of increase in LE with increasing R_n was suppressed. However, LE became dominant over H during wet canopy
conditions, with a much greater rate of increase in LE with increasing R_n compared to dry canopy conditions. A previous
study at Kopuatai also reported dominance of LE over H when the canopy was rain-wetted, and vice versa during dry canopy
conditions (Campbell and Williamson, 1997). This indicates that the restrictions on E imposed by *E. robustum* break down
400 in wet canopy conditions, likely due to high interception loss. It is possible that interception loss is much higher at Kopuatai
relative to Mer Bleue due to the greater canopy height (means 48 cm and 18 cm , respectively) and density.

The responses of E and EF to increasing VPD at Mer Bleue were consistent with previous findings by Helbig et al. (2020),
which synthesised data from 35 boreal peatlands, including Mer Bleue; however, the response at Kopuatai bog was closer to
405 that of the boreal forests included in their study. As such, the response of E to VPD at Kopuatai bog is distinctive compared
to boreal peatlands. This may be a critical adaptation that allows peat formation in a climate zone that is uncharacteristic for
ombrotrophic peatlands (McGlone, 2009) due to the combination of moderate precipitation and a higher mean annual



temperature than at Northern Hemisphere peatlands. Tropical peatlands also experience higher temperatures, however annual precipitation tends to be much higher than at Kopuatai (>2500 mm in Southeast Asia, for example; Page et al., 2006), which
410 maintains adequately wet conditions. Therefore, it is likely that the dominance of *E. robustum* at Kopuatai is crucial to the persistence of this peatland in its unusual climate zone due to restrictions imposed on transpiration and sub-canopy *E*, as these adaptations likely contribute to maintaining a shallow and stable water table.

4.3 Implications for future peatland water balances

Northern Hemisphere peatlands, such as Mer Bleue, may be at risk of increased evaporative water loss as VPD increases due
415 to climate warming (Helbig et al., 2020). This could lead to more frequent low water tables and water deficits, which in turn may reduce net C uptake (Zhong et al., 2020; Kwon et al., 2022). The effect of low water tables on the C balance has previously been observed at Mer Bleue, where decreases in water levels at a beaver pond near the EC site caused a decrease in C uptake; further modelling also predicted that a beaver pond water table below ~1.7 m would cause Mer Bleue to switch
420 from a C sink to a source (He et al., 2022). A lower mean water table at a drained *E. robustum* bog near Kopuatai also caused lower C uptake rates compared to Kopuatai, but likely remained a C sink due to the tolerance of *E. robustum* to a low and fluctuating water table (Ratcliffe et al., 2018). Therefore, the C uptake of both peatlands could be impacted by decreased water tables; however, under increased VPD due to climate warming, it appears that Kopuatai may be able to retain more water than Northern Hemisphere peatlands through strong constraints on *E*. This greater water retention at Kopuatai could enable current high and stable net annual C uptake rates (Goodrich et al., 2017) to be maintained despite warmer and
425 potentially drier conditions in the future (Lawrence et al., 2022). Therefore, we propose that the previously observed *resilience* of C uptake at Kopuatai to dry conditions (Goodrich et al., 2017) primarily results from “hydrological resistance” to increasing VPD driven by the properties of the *E. robustum* canopy.

5 Conclusions

We compared ecosystem *E* regimes, energy balance partitioning, and the response of *E* to VPD at two peatlands - Kopuatai
430 bog in Aotearoa New Zealand and Mer Bleue bog in Canada. Our motivation was to determine whether regulation of water losses in these peatlands with different vegetation communities enables hydrological resistance to climatic drying. Our results demonstrated that the *E* regime was much more conservative at Kopuatai than at Mer Bleue at high VPD because of greater limitations on *E* due to reduced g_c . At Mer Bleue, *E* was only weakly limited at high VPD, as the impacts of reduced shrub g_c were likely offset by water loss from non-vascular moss surfaces. The greater limitations on *E* observed at Kopuatai
435 indicate hydrological resistance of this vascular plant-dominated ecosystem to hydro-climatic change. Importantly, this resistance likely contributes to resilience of the net C sink at Kopuatai, despite being located in a warm-temperate climate zone, and thus could be a key self-regulation mechanism. As a result, it appears that Kopuatai may be better equipped to conserve water and maintain high C uptake rates than Mer Bleue, and perhaps other Northern Hemisphere peatlands, in the face of climate change-induced warming and drying. Further investigation of *E* partitioning into transpiration, interception



440 loss, and peat surface E , and the role of the dense standing litter layer for E regulation at Kopuatai may yield additional insights into this observed hydrological resistance.

Appendices

Appendix A

Table A1. Methods and instruments used for data collection at Kopuatai and Mer Bleue.

Variable	Kopuatai	Mer Bleue
Latent heat flux (LE)	4.25 m CSAT3 & CSAT3B (CSI); LI-7500 (LI-COR)	3.0 m 1012R3 & R3-50 (Gill); LI-6262 & LI7000 (LI-COR)
Sensible heat flux (H)	4.25 m CSAT3 & CSAT3B (CSI)	1012R3 & R3-50 (Gill)
Soil (peat) heat flux (G)	HFP01 (Hukseflux); TCAV (CSI)	N/A
Net radiation flux density (R_n)	2 m NR01 (Hukseflux)	CNR1 (Kipp & Zonen)
Air temperature (T_{air}) and vapour pressure deficit (VPD)	4.2 m HMP155 (Vaisala)	2 m HMP various models (Vaisala)
Precipitation (P)	0.6 m TB03 (Hydrological Services)	0.3 m TM525 tipping bucket gauge (Texas Instruments) Winter snowfall from Environment Canada weather station at Ottawa Airport
Water table depth (WTD)	Dipwell, WL1000 (Hydrological Services)	Float & potentiometer, and OTT PLS

445



Appendix B

The antecedent precipitation index (API) is an exponential function used to predict the duration of canopy wetness after a rainfall event, with inputs of rainfall depth and time since rainfall (Keyte Beattie, 2014). Initially conceived by Woods and
450 Rowe (1996) for predicting catchment moisture conditions prior to a rain event, the API function was then modified by Smith (2003) to estimate canopy wetness state:

$$API = \sum_{i=1}^j \frac{P_i}{1.104 \times 1.024^i} \quad (B1)$$

where P_i is precipitation (mm) measured during the i^{th} half-hour period before the current half hour, and j is the moving
455 window size in half hours (48 in this study). An example of modelled canopy drying using the API function is shown in Figure B1. An API value of 0.2 units or less approximately represents a dry canopy, meaning that E is predominantly sourced from peat surface E and transpiration. When the API value is greater than or equal to 1, the vegetation canopy is deemed wet and interception loss is the dominant source of E .

Predictions of canopy wetness using the API have been shown to correspond reasonably well with leaf wetness sensor
460 measurements at Kopuatai, using an API period of 18 hours for the upper canopy, 36 hours for the standing litter layer, and 60 hours for the lower canopy (Keyte Beattie, 2014). In this study, we defined wet canopy periods as times when the upper canopy was saturated, i.e., when interception loss likely made up the majority of E , so an 18-hour period would likely be appropriate for this study. However, in order to obtain a conservative identification of periods with a dry upper canopy, i.e.,
465 for greater certainty that the canopy is dry when $API \leq 0.2$, a slightly longer period of 24 hours was used. While this approach has only been tested in the field at Kopuatai, for consistency we used the same function for separating wet and dry canopy measurements at Mer Bleue. Given the relatively sparse shrub canopy and absence of standing litter at Mer Bleue, API should provide conservative estimates of canopy dryness. The use of MoD data in this study is beneficial as it would have reduced the impact of early morning wetting from dew, therefore reducing the number of instances where API
470 predicted canopy wetness incorrectly.





Figure B1. Examples of predicted canopy drying after rainfall using the antecedent precipitation index (API) with 24, 48, and 96 half hour time parameters. The rain event in (a) and (c) is from the Kupuatai dataset, while (b) and (d) consist of Mer Bleue data.




Appendix C



475


Figure C1. Energy balance closure at Kopuatai for 2012–2022 (inclusive), based on daily middle-of-day means of energy balance components. *LE*: latent heat flux; *H*: sensible heat flux; *G*: soil heat flux; *R_n*: net radiation; ΔS_{can} : canopy heat storage change. EBR is the energy balance ratio.




480 **Figure C2.** Energy balance closure at Mer Bleue for 1999–2018 (inclusive), based on daily middle-of-day means of energy balance components. For term definitions, see Figure C1. G is assumed to be 10 % of R_n , while ΔS_{can} is assumed to be 3 % of R_n . EBR is the energy balance ratio.



Appendix D



485 **Figure D1.** Relationship between the evaporative fraction (EF) and vapour pressure deficit (VPD) at Kopuatai bog. Light grey data points represent raw, unfiltered 30-minute eddy covariance data, while dark grey data points show growing season (Sep–May), dry canopy, middle-of-day mean data used to calculate the binned means (black line).



490 **Figure D2. Relationship between evaporative fraction (EF) and vapour pressure deficit (VPD) at Mer Bleue bog. Light grey data points represent raw, unfiltered 30-minute eddy covariance data, while dark grey data points show growing season (May–Oct), dry canopy, middle-of-day mean data used to calculate the binned means (black line).**

Data availability

Data sets used in this paper are available on Zenodo at <https://doi.org/10.5281/zenodo.8226863> (Speranskaya et al., 2023).

Author contribution

495 DIC and PML initiated the project and developed the methodology with LS. DIC collected and processed data from Kopuatai, and PML and ERH collected and processed data from Mer Bleue. LS analysed the data and wrote the manuscript draft. DIC, PML, and ERH contributed to review and editing of the manuscript.

Competing interests

The authors declare that they have no conflict of interest.



500 Acknowledgements

Kopuatai bog (Aotearoa New Zealand): Te Kupenga O Ngāti Hako and The Department of Conservation permitted us to access Kopuatai bog and supported our longstanding research presence there. Funding was from Manaaki Whenua Landcare Research under its wetland research programme and The University of Waikato, including a scholarship to LS. Access across farmland was facilitated by the Brewster family.

505

Mer Bleue bog (Canada): The Mer Bleue research site has been supported through research grants to PML and ERH (and others) from the Natural Science and Engineering Research Council of Canada. We are grateful to the National Capital Commission for access to the site and continued support of our research.

References

510 Admiral, S. W. and Lafleur, P. M.: Partitioning of latent heat flux at a northern peatland, *Aquat. Bot.*, 86, 107-116, <https://doi.org/10.1016/j.aquabot.2006.09.006>, 2007.

Admiral, S. W., Lafleur, P. M., and Roulet, N. T.: Controls on latent heat flux and energy partitioning at a peat bog in eastern Canada, *Agr. Forest Meteorol.*, 140, 308-321, <https://doi.org/10.1016/j.agrformet.2006.03.017>, 2006.

515

Agnew, A. D. Q., Rapson, G. L., Sykes, M. T., and Bastow Wilson, J.: The functional ecology of *Empodisma minus* (Hook, f.) Johnson & Cutler in New Zealand ombrotrophic mires, *New Phytol.*, 124, 703-710, 1993.

Alekseychik, P., Mammarella, I., Lindroth, A., Lohila, A., Aurela, M., Laurila, T., Kasurinen, V., Lund, M., Rinne, J., Nilsson, M. B., Peichl, M., Minkinen, K., Shurpali, N. J., Tuittila, E.-S., Martikainen, P. J., Tuovinen, J.-P., and Vesala, T.: Surface energy exchange in pristine and managed boreal peatlands, *Mires and Peat*, 21, 1-26, <https://doi.org/10.19189/MaP.2018.OMB.333>, 2018.

525 Anderson, R. G. and Wang, D.: Energy budget closure observed in paired Eddy Covariance towers with increased and continuous daily turbulence, *Agr. Forest Meteorol.*, 184, 204-209, <https://doi.org/10.1016/j.agrformet.2013.09.012>, 2014.

Brümmer, C., Black, T. A., Jassal, R. S., Grant, N. J., Spittlehouse, D. L., Chen, B., Nesic, Z., Amiro, B. D., Arain, M. A., Barr, A. G., Bourque, C. P. A., Coursolle, C., Dunn, A. L., Flanagan, L. B., Humphreys, E. R., Lafleur, P. M., Margolis, H. A., McCaughey, J. H., and Wofsy, S. C.: How climate and vegetation type influence evapotranspiration and water use efficiency in Canadian forest, peatland and grassland ecosystems, *Agr. Forest Meteorol.*, 153, 14-30, <https://doi.org/10.1016/j.agrformet.2011.04.008>, 2012.



- Bubier, J. L., Moore, T. R., and Crosby, G.: Fine-scale vegetation distribution in a cool temperate peatland, *Can. J. Botany*, 84, 910-923, <https://doi.org/10.1139/b06-044>, 2006.
- 535
- Burba, G.: *Eddy Covariance Method for Scientific, Regulatory, and Commercial Applications*, LI-COR Biosciences, Lincoln, Nebraska, USA, 702 pp., 2022.
- Campbell, E. O.: The restiad peat bogs at Motumaoho and Moanatuatua, *Transactions of the Royal Society of New Zealand : Botany*, 2, 1964.
- 540
- Campbell, D. I. and Williamson, J. L.: Evaporation from a raised peat bog, *J. Hydrol.*, 193, 142-160, [https://doi.org/10.1016/S0022-1694\(96\)03149-6](https://doi.org/10.1016/S0022-1694(96)03149-6), 1997.
- 545
- Campbell, D. I., Smith, J., Goodrich, J. P., Wall, A. M., and Schipper, L. A.: Year-round growing conditions explains large CO₂ sink strength in a New Zealand raised peat bog, *Agr. Forest Meteorol.*, 192-193, 59-68, <https://doi.org/10.1016/j.agrformet.2014.03.003>, 2014.
- Clarkson, B. R., Schipper, L. A., and Silvester, W. B.: Nutritional niche separation in coexisting bog species demonstrated by ¹⁵N-enriched simulated rainfall, *Austral Ecol.*, <https://doi.org/10.1111/j.1442-9993.2009.01939.x>, 2009.
- 550
- Dymond, J. R., Sabetizade, M., Newsome, P. F., Harmsworth, G. R., and Ausseil, A.: Revised extent of wetlands in New Zealand, *N. Z. J. Ecol.*, 45, 1-8, <https://doi.org/10.20423/nzjecol.45.32>, 2021.
- 555
- Eichelmann, E., Hemes, K. S., Knox, S. H., Oikawa, P. Y., Chamberlain, S. D., Sturtevant, C., Verfaillie, J., and Baldocchi, D. D.: The effect of land cover type and structure on evapotranspiration from agricultural and wetland sites in the Sacramento–San Joaquin River Delta, California, *Agr. Forest Meteorol.*, 256-257, 179-195, <https://doi.org/10.1016/j.agrformet.2018.03.007>, 2018.
- 560
- Fang, Z., Zhang, W., Brandt, M., Abdi, A. M., and Fensholt, R.: Globally Increasing Atmospheric Aridity Over the 21st Century, *Earth's Future*, 10, <https://doi.org/10.1029/2022ef003019>, 2022.
- Ficklin, D. L. and Novick, K. A.: Historic and projected changes in vapor pressure deficit suggest a continental-scale drying of the United States atmosphere, *J. Geophys. Res.: Atmos.*, 122, 2061-2079, <https://doi.org/10.1002/2016jd025855>, 2017.
- 565



- Fraser, C. D. J., Roulet, N. T., and Lafleur, P. M.: Groundwater flow patterns in a large peatland, *J. Hydrol.*, 246, 142-154, [https://doi.org/10.1016/S0022-1694\(01\)00362-6](https://doi.org/10.1016/S0022-1694(01)00362-6), 2001.
- 570 Frolking, S. and Roulet, N. T.: Holocene radiative forcing impact of northern peatland carbon accumulation and methane emissions, *Global Change Biol.*, 13, 1079-1088, <https://doi.org/10.1111/j.1365-2486.2007.01339.x>, 2007.
- Goodrich, J. P., Campbell, D. I., and Schipper, L. A.: Southern Hemisphere bog persists as a strong carbon sink during droughts, *Biogeosciences*, 14, 4563-4576, <https://doi.org/10.5194/bg-14-4563-2017>, 2017.
- 575 Goodrich, J. P., Campbell, D. I., Clearwater, M. J., Rutledge, S., and Schipper, L. A.: High vapor pressure deficit constrains GPP and the light response of NEE at a Southern Hemisphere bog, *Agr. Forest Meteorol.*, 203, 54-63, <https://doi.org/10.1016/j.agrformet.2015.01.001>, 2015.
- Goulden, M. L., Litvak, M., and Miller, S. D.: Factors that control *Typha* marsh evapotranspiration, *Aquat. Bot.*, 86, 97-580 106, <https://doi.org/10.1016/j.aquabot.2006.09.005>, 2007.
- Granger, R. J.: An examination of the concept of potential evaporation, *J. Hydrol.*, 111, 9-19, [https://doi.org/10.1016/0022-1694\(89\)90248-5](https://doi.org/10.1016/0022-1694(89)90248-5), 1989.
- 585 He, H., Moore, T., Humphreys, E. R., Lafleur, P. M., and Roulet, N. T.: Water level variation at a beaver pond significantly impacts net CO₂ uptake of a continental bog, *Hydrol. Earth Syst. Sc.*, 27, 213-227, <https://doi.org/10.5194/hess-27-213-2023>, 2023.
- 590 Helbig, M., Waddington, J. M., Alekseychik, P., Amiro, B. D., Aurela, M., Barr, A. G., Black, T. A., Blanken, P. D., Carey, S. K., Chen, J., Chi, J., Desai, A. R., Dunn, A., Euskirchen, E. S., Flanagan, L. B., Forbrich, I., Friborg, T., Grelle, A., Harder, S., Heliasz, M., Humphreys, E. R., Ikawa, H., Isabelle, P.-E., Iwata, H., Jassal, R., Korkiakoski, M., Kurbatova, J., Kutzbach, L., Lindroth, A., Löfvenius, M. O., Lohila, A., Mammarella, I., Marsh, P., Maximov, T., Melton, J. R., Moore, P. A., Nadeau, D. F., Nicholls, E. M., Nilsson, M. B., Ohta, T., Peichl, M., Petrone, R. M., Petrov, R., Prokushkin, A., Quinton, W. L., Reed, D. E., Roulet, N. T., Runkle, B. R. K., Sonnentag, O., Strachan, I. B., Taillardat, P., Tuittila, E.-S., Tuovinen, 595 J.-P., Turner, J., Ueyama, M., Varlagin, A., Wilmking, M., Wofsy, S. C., and Zyrianov, V.: Increasing contribution of peatlands to boreal evapotranspiration in a warming climate, *Nat. Clim. Change*, 10, 555-560, <https://doi.org/10.1038/s41558-020-0763-7>, 2020.



600 Helfter, C., Campbell, C., Dinsmore, K. J., Drewer, J., Coyle, M., Anderson, M., Skiba, U., Nemitz, E., Billett, M. F., and
Sutton, M. A.: Drivers of long-term variability in CO₂ net ecosystem exchange in a temperate peatland, *Biogeosciences*, 12,
1799-1811, <https://doi.org/10.5194/bg-12-1799-2015>, 2015.

605 Hember, R. A., Lafleur, P. M., and Cogley, J. G.: Synoptic controls on summer evapotranspiration from a bog peatland in
southern Canada, *Int. J. Climatol.*, 25, 793-809, <https://doi.org/10.1002/joc.1154>, 2005.

Holden, J.: Peatland hydrology and carbon release: why small-scale process matters, *Philosophical Transactions of the Royal
Society A: Mathematical Physical and Engineering Sciences*, 363, 2891-2913, <https://doi.org/10.1098/rsta.2005.1671>, 2005.

610 Humphreys, E. R., Lafleur, P. M., Flanagan, L. B., Hedstrom, N., Syed, K. H., Glenn, A. J., and Granger, R.: Summer
carbon dioxide and water vapor fluxes across a range of northern peatlands, *J. Geophys. Res.: Biogeo.*,
111, <https://doi.org/10.1029/2005jg000111>, 2006.

Joosten, H. and Clarke, D.: *Wise use of mires and peatlands*, Totnes, 2002.

615 Keyte Beattie, A. M.: *The role of *Empodisma robustum* litter in CO₂ exchange at Kopuatai bog*, University of Waikato,
Hamilton, New Zealand, 2014.

620 Kim, J., Rochefort, L., Hogue-Hugron, S., Alqulaiti, Z., Dunn, C., Pouliot, R., Jones, T. G., Freeman, C., and Kang, H.:
Water Table Fluctuation in Peatlands Facilitates Fungal Proliferation, Impedes *Sphagnum* Growth and Accelerates
Decomposition, *Front. Earth Sci.*, 8, <https://doi.org/10.3389/feart.2020.579329>, 2021.

625 Kurbatova, J., Arneth, A., Vygodskaya, N. N., Kolle, O., Varlargin, A. V., Milyukova, I. M., Tchebakova, N. M., Schulze, E.
D., and Lloyd, J.: Comparative ecosystem-atmosphere exchange of energy and mass in a European Russian and a central
Siberian bog I. Interseasonal and interannual variability of energy and latent heat fluxes during the snowfree period, *Tellus*
B, 54, <https://doi.org/10.3402/tellusb.v54i5.16683>, 2002.

630 Kwon, M. J., Ballantyne, A., Ciais, P., Qiu, C., Salmon, E., Raoult, N., Guenet, B., Gockede, M., Euskirchen, E. S.,
Nykanen, H., Schuur, E. A. G., Turetsky, M. R., Dieleman, C. M., Kane, E. S., and Zona, D.: Lowering water table reduces
carbon sink strength and carbon stocks in northern peatlands, *Global Change Biol.*, 28, 6752-
6770, <https://doi.org/10.1111/gcb.16394>, 2022.



- Lafleur, P. M. and Roulet, N. T.: A comparison of evaporation rates from two fens of the Hudson Bay Lowland, *Aquat. Bot.*, 44, 59-69, [https://doi.org/10.1016/0304-3770\(92\)90081-S](https://doi.org/10.1016/0304-3770(92)90081-S), 1992.
- 635 Lafleur, P. M., Roulet, N. T., and Admiral, S. W.: Annual cycle of CO₂ exchange at a bog peatland, *J. Geophys. Res.: Atmos.*, 106, 3071-3081, <https://doi.org/10.1029/2000jd900588>, 2001.
- Lafleur, P. M., Hember, R. A., Admiral, S. W., and Roulet, N. T.: Annual and seasonal variability in evapotranspiration and water table at a shrub-covered bog in southern Ontario, Canada, *Hydrol. Process.*, 19, 3533-640 3550, <https://doi.org/10.1002/hyp.5842>, 2005.
- Lawrence, J., Mackey, B., Chiew, F., Costello, M. J., Hennessy, K., Lansbury, N., Nidumolu, U. B., Pecl, G., Rickards, L., Tapper, N., A.Woodward, and A.Wreford: Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, in: *Climate Change 2022: Impacts, Adaptation and Vulnerability*, edited by: 645 Pörtner, H.-O., Roberts, D. C., Tignor, M., Poloczanska, E. S., Mintenbeck, K., Alegría, A., Craig, M., Langsdorf, S., S. Löschke, Möller, V., Okem, A., and Rama, B., Cambridge University Press, Cambridge, UK and New York, NY, USA, 1581–1688, <https://doi.org/10.1017/9781009325844.013>, 2022.
- Leuning, R., van Gorsel, E., Massman, W. J., and Isaac, P. R.: Reflections on the surface energy imbalance problem, *Agr. 650 Forest Meteorol.*, 156, 65-74, <https://doi.org/10.1016/j.agrformet.2011.12.002>, 2012.
- Liljedahl, A. K., Hinzman, L. D., Harazono, Y., Zona, D., Tweedie, C. E., Hollister, R. D., Engstrom, R., and Oechel, W. C.: Nonlinear controls on evapotranspiration in arctic coastal wetlands, *Biogeosciences*, 8, 3375-655 3389, <https://doi.org/10.5194/bg-8-3375-2011>, 2011.
- Ma, L., Zhu, G., Chen, B., Zhang, K., Niu, S., Wang, J., Ciais, P., and Zuo, H.: A globally robust relationship between water table decline, subsidence rate, and carbon release from peatlands, *Communications Earth & Environment*, 3, <https://doi.org/10.1038/s43247-022-00590-8>, 2022.
- 660 Massmann, A., Gentine, P., and Lin, C.: When Does Vapor Pressure Deficit Drive or Reduce Evapotranspiration?, *J. Adv. Model. Earth Sy.*, 11, 3305-3320, <https://doi.org/10.1029/2019MS001790>, 2019.
- McGlone, M. S.: Postglacial history of New Zealand wetlands and implications for their conservation, *N. Z. J. Ecol.*, 33, 1-665 23, 2009.



- Miralles, D. G., Brutsaert, W., Dolman, A. J., and Gash, J. H.: On the Use of the Term "Evapotranspiration", *Water Resour. Res.*, 56, e2020WR028055, <https://doi.org/10.1029/2020WR028055>, 2020.
- Moore, T. R., Bubier, J. L., Frohling, S. E., Lafleur, P. M., and Roulet, N. T.: Plant biomass and production and
670 CO₂ exchange in an ombrotrophic bog, *J. Ecol.*, 90, 25-36, <https://doi.org/10.1046/j.0022-0477.2001.00633.x>, 2002.
- Newnham, R. N., de Lange, P. J., and Lowe, D. J.: Holocene vegetation, climate and history of a raised bog complex, northern New Zealand based on palynology, plant macrofossils and tephrochronology, *The Holocene*, 5, 267-282, 1995.
- 675 Nimmo, D. G., Mac Nally, R., Cunningham, S. C., Haslem, A., and Bennett, A. F.: Vive la resistance: reviving resistance for 21st century conservation, *Trends Ecol. Evol.*, 30, 516-523, <https://doi.org/10.1016/j.tree.2015.07.008>, 2015.
- Page, S. E., Rieley, J. O., and Wüst, R.: Lowland tropical peatlands of Southeast Asia, in: *Peatlands - Evolution and Records of Environmental and Climate Changes*, edited by: Martini, I. P., Martinez Cortizas, A., and Chesworth, W., Elsevier,
680 Amsterdam, The Netherlands, 145-172, [https://doi.org/10.1016/s0928-2025\(06\)09007-9](https://doi.org/10.1016/s0928-2025(06)09007-9), 2006.
- Peichl, M., Sagerfors, J., Lindroth, A., Buffam, I., Grelle, A., Klemetsson, L., Laudon, H., and Nilsson, M. B.: Energy exchange and water budget partitioning in a boreal minerogenic mire, *J. Geophys. Res.: Biogeo.*, 118, 1-13, <https://doi.org/10.1029/2012jg002073>, 2013.
- 685 Ratcliffe, J. L., Campbell, D. I., Clarkson, B. R., Wall, A. M., and Schipper, L. A.: Water table fluctuations control CO₂ exchange in wet and dry bogs through different mechanisms, *Sci. Total Environ.*, 655, 1037-1046, <https://doi.org/10.1016/j.scitotenv.2018.11.151>, 2019.
- 690 Roehm, C. L. and Roulet, N. T.: Seasonal contribution of CO₂ fluxes in the annual C budget of a northern bog, *Global Biogeochem. Cy.*, 17, <https://doi.org/10.1029/2002gb001889>, 2003.
- Roulet, N. T., Lafleur, P. M., Richard, P. J. H., Moore, T. R., Humphreys, E. R., and Bubier, J.: Contemporary carbon balance and late Holocene carbon accumulation in a northern peatland, *Global Change Biol.*, 13, 397-
695 411, <https://doi.org/10.1111/j.1365-2486.2006.01292.x>, 2007.
- Runkle, B. R. K., Wille, C., Gažovič, M., Wilmking, M., and Kutzbach, L.: The surface energy balance and its drivers in a boreal peatland fen of northwestern Russia, *J. Hydrol.*, 511, 359-373, <https://doi.org/10.1016/j.jhydrol.2014.01.056>, 2014.



- 700 Scharlemann, J. P. W., Tanner, E. V. J., Hiederer, R., and Kapos, V.: Global soil carbon: understanding and managing the largest terrestrial carbon pool, *Carbon Manag.*, 5, 81-91, <https://doi.org/10.4155/cmt.13.77>, 2014.
- Shearer, J. C.: Natural and anthropogenic influences on peat development in Waikato/Hauraki Plains restiad bogs, *J. R. Soc. N. Z.*, 27, 295-313, <https://doi.org/10.1080/03014223.1997.9517540>, 1997.
- 705 Shimoyama, K.: Seasonal and interannual variation in water vapor and heat fluxes in a West Siberian continental bog, *J. Geophys. Res.*, 108, <https://doi.org/10.1029/2003jd003485>, 2003.
- Shimoyama, K.: Controls on evapotranspiration in a west Siberian bog, *J. Geophys. Res.*,
- 710 109, <https://doi.org/10.1029/2003jd004114>, 2004.
- Smith, J.: Fluxes of Carbon Dioxide and Water Vapour at a Waikato Peat Bog, The University of Waikato, Hamilton, New Zealand, 2003.
- 715 Sonnentag, O., Van Der Kamp, G., Barr, A. G., and Chen, J. M.: On the relationship between water table depth and water vapor and carbon dioxide fluxes in a minerotrophic fen, *Global Change Biol.*, 16, 1762-1776, <https://doi.org/10.1111/j.1365-2486.2009.02032.x>, 2009.
- Speranskaya, L., Campbell, D. I., Lafleur, P. M., Humphreys, E. R.: Data sets for "Peatland evaporation across hemispheres: contrasting controls and sensitivity to climate warming driven by plant functional types", Zenodo [data set], <https://doi.org/10.5281/zenodo.8226863>, 2023.
- 720 Strilesky, S. L. and Humphreys, E. R.: A comparison of the net ecosystem exchange of carbon dioxide and evapotranspiration for treed and open portions of a temperate peatland, *Agr. Forest Meteorol.*, 153, 45-
- 725 53, <https://doi.org/10.1016/j.agrformet.2011.06.006>, 2012.
- Takagi, K., Tsuboya, T., Takahashi, H., and Inoue, T.: Effect of the invasion of vascular plants on heat and water balance in the Sarobetsu mire, northern Japan, *Wetlands*, 19, 246-254, <https://doi.org/10.1007/BF03161754>, 1999.
- 730 Thompson, M. A., Campbell, D. I., and Spronken-Smith, R. A.: Evaporation from natural and modified raised peat bogs in New Zealand, *Agr. Forest Meteorol.*, 95, 85-98, [https://doi.org/10.1016/S0168-1923\(99\)00027-1](https://doi.org/10.1016/S0168-1923(99)00027-1), 1999.



- Villegas, J. C., Breshears, D. D., Zou, C. B., and Law, D. J.: Ecohydrological controls of soil evaporation in deciduous drylands: How the hierarchical effects of litter, patch and vegetation mosaic cover interact with phenology and season, *J. Arid Environ.*, 74, 595-602, <https://doi.org/10.1016/j.jaridenv.2009.09.028>, 2010.
- Vitt, D. H.: Functional Characteristics and Indicators of Boreal Peatlands, in: *Boreal Peatland Ecosystems*, edited by: Wieder, R. K., and Vitt, D. H., Springer-Verlag, Berlin, Germany, 9-24, https://doi.org/10.1007/978-3-540-31913-9_2, 2006.
- 740 Wagstaff, S. J. and Clarkson, B. R.: Systematics and ecology of the Australasian genus *Empodisma* (Restionaceae) and description of a new species from peatlands in northern New Zealand, *PhytoKeys*, 39-79, <https://doi.org/10.3897/phytokeys.13.3259>, 2012.
- Wang, M., Wu, J., and Lafleur, P.: Comparison of energy fluxes between an undisturbed bog and an adjacent abandoned peatland pasture, *Agr. Forest Meteorol.*, 291, <https://doi.org/10.1016/j.agrformet.2020.108086>, 2020.
- 745 Woods, R. and Rowe, L.: The changing spatial variability of subsurface flow across a hillside, *J. Hydrol.*, 35, 49-84, 1996.
- Wu, J., Kutzbach, L., Jager, D., Wille, C., and Wilmking, M.: Evapotranspiration dynamics in a boreal peatland and its impact on the water and energy balance, *J. Geophys. Res.*, 115, <https://doi.org/10.1029/2009jg001075>, 2010.
- 750 Xu, J., Morris, P. J., Liu, J., and Holden, J.: PEATMAP: Refining estimates of global peatland distribution based on a meta-analysis, *Catena*, 160, 134-140, <https://doi.org/10.1016/j.catena.2017.09.010>, 2018.
- 755 Yu, Z., Loisel, J., Brosseau, D. P., Beilman, D. W., and Hunt, S. J.: Global peatland dynamics since the Last Glacial Maximum, *Geophys. Res. Lett.*, 37, n/a-n/a, <https://doi.org/10.1029/2010gl043584>, 2010.
- Zhong, Y., Jiang, M., and Middleton, B. A.: Effects of water level alteration on carbon cycling in peatlands, *Ecosystem Health and Sustainability*, 6, <https://doi.org/10.1080/20964129.2020.1806113>, 2020.
- 760