

# High Spectral Resolution Lidar – generation 2 (HSRL-2) Retrievals of Ocean Surface Wind Speed: Methodology and Evaluation

Sanja Dmitrovic<sup>1</sup>, Johnathan W. Hair<sup>2</sup>, Brian L. Collister<sup>2</sup>, Ewan Crosbie<sup>2,3</sup>, Marta A. Fenn<sup>2</sup>, Richard A. Ferrare<sup>2</sup>, David B. Harper<sup>2</sup>, Chris A. Hostetler<sup>2</sup>, Yongxiang Hu<sup>2</sup>, John A. Reagan<sup>4</sup>, Claire E. Robinson<sup>2,3,+</sup>, Shane T. Seaman<sup>2</sup>, Taylor J. Shingler<sup>2</sup>,  
5 Kenneth L. Thornhill<sup>2,3</sup>, Holger Vömel<sup>5</sup>, Xubin Zeng<sup>6</sup>, Armin Sorooshian<sup>1,6,7</sup>

<sup>1</sup>James C. Wyant College of Optical Sciences, University of Arizona, Tucson, AZ 85721, USA

<sup>2</sup>NASA Langley Research Center, Hampton, VA 23681, USA

<sup>3</sup>Analytical Mechanics Associates, Hampton, VA 23666, USA

<sup>4</sup>Department of Electrical and Computer Engineering, University of Arizona, Tucson, AZ 85721, USA

10 <sup>5</sup>National Center for Atmospheric Research, Boulder, CO 80307, USA

<sup>6</sup>Department of Hydrology and Atmospheric Sciences, University of Arizona, Tucson, AZ 85721, USA

<sup>7</sup>Department of Chemical and Environmental Engineering, University of Arizona, Tucson, AZ 85721, USA

<sup>+</sup>Deceased

*Correspondence to:* Armin Sorooshian (armin@arizona.edu)

15 **Abstract.** Ocean surface wind speed (i.e., wind speed 10 m above sea level) is a critical parameter used by atmospheric models to estimate the state of the marine atmospheric boundary layer (MABL). Accurate surface wind speed measurements in diverse locations are required to improve characterization of MABL dynamics and assess how models simulate large-scale phenomena related to climate change and global weather patterns. To provide these measurements, this study introduces and evaluates a new surface wind speed data product from NASA Langley Research Center’s High Spectral Resolution Lidar – generation 2  
20 (HSRL-2) using data collected as part of NASA’s Aerosol Cloud meTeorology Interactions oVer the western ATLantic Experiment (ACTIVATE) mission. The HSRL-2 [is a nadir-viewing lidar that](#) can directly measure vertically resolved aerosol backscatter and extinction profiles without additional constraints or assumptions, enabling the instrument to accurately derive atmospheric attenuation and directly determine surface reflectance (i.e., surface backscatter). Also, the high horizontal spatial resolution of the HSRL-2 retrievals (0.5 s or ~75 m along track) allows the instrument to probe the fine-scale spatial variability  
25 of surface wind speeds over time along the flight track and breaks in broken cloud fields. A rigorous evaluation on these retrievals is performed by comparing coincident HSRL-2 and National Center for Atmospheric Research (NCAR) AVAPS dropsonde data, owing to the joint deployment of these two instruments on ACTIVATE’s King Air aircraft. These comparisons show correlations of 0.89, slopes of 1.04 and 1.17, and y-intercepts of -0.13 m s<sup>-1</sup> and -1.05 m s<sup>-1</sup> for linear and bisector regressions, respectively and the overall accuracy is calculated to be 0.15 m s<sup>-1</sup> ± 1.80 m s<sup>-1</sup>. It is also shown that the dropsonde  
30 surface wind speed data most closely follows the HSRL-2 distribution of wave-slope variance using the distribution proposed by Hu et al. (2008) than the ones proposed by Cox and Munk (1954) and Wu (1990) for surface wind speeds below 7 m s<sup>-1</sup>, with this category comprising most of the ACTIVATE data set. The retrievals are then evaluated separately for surface wind speeds below 7 m s<sup>-1</sup> and between 7 m s<sup>-1</sup> and 13.3 m s<sup>-1</sup> and show that the HSRL-2 retrieves surface wind speeds with a bias of ~0.5 m s<sup>-1</sup> and an error of ~1.5 m s<sup>-1</sup>, a finding not apparent in the cumulative comparisons. Also, it is shown that the HSRL-

35 2 retrievals are more accurate in the summer ( $-0.18 \text{ m s}^{-1} \pm 1.52 \text{ m s}^{-1}$ ) than winter ( $0.63 \text{ m s}^{-1} \pm 2.07 \text{ m s}^{-1}$ ), but the HSRL-2 is still able to make numerous, ( $N = 236$ ) accurate retrievals in the winter. Overall, this study highlights the abilities and assesses the performance of the HSRL-2 surface wind speed retrievals and it is hoped that further evaluation of these retrievals will be performed using other airborne and satellite data sets.

## 1 Introduction

40 The layer between the ocean and free troposphere, known as the marine atmospheric boundary layer (MABL), hosts various processes such as the modulation of sensible and latent heat fluxes, the exchange of gases such as carbon dioxide, the evolution of clouds, and the transport of aerosol particles (Neukermans et al., 2018). Improved characterization of MABL dynamics is required to accurately simulate large-scale phenomena related to climate change and global weather patterns (Paiva et al., 2021). This characterization relies on a combination of global numerical weather prediction (NWP) models and real  
45 observations (Carvalho, 2019). One of the most influential parameters that drive these MABL processes is ocean surface wind speeds or wind speeds at 10 m above sea level (hereafter called surface wind speeds). Therefore, instruments such as lidar are used to provide accurate surface wind speed measurements in various geographical locations to improve estimations of the MABL state globally. For instance, satellite lidar systems that measure aerosol and cloud vertical distributions, such as the lidar on board the NASA Cloud-Aerosol Lidar and Infrared Pathfinder Observation (CALIPSO) satellite, also have the  
50 capability to provide horizontally-resolved surface wind speed data. The underlying principle of lidar surface wind speed retrievals was first derived by Cox and Munk (1954), where bidirectional reflectance measurements of sea-surface glint are used to establish a Gaussian relationship between surface wind speeds and the distribution of wind-driven wave slopes. To probe these surface wave slopes, lidar instruments emit laser pulses into the atmosphere and measure the reflectance (or backscatter) of those laser pulses from particles, molecules, and the ocean surface. The magnitude of the measured signal is  
55 then used to estimate the variance of the wave-slope distribution (i.e., wave-slope variance) and therefore surface wind speed. Note that reflectance and backscatter are used interchangeably throughout this paper.

Although many studies have expanded upon the original Cox-Munk relationship (e.g., Hu et al., 2008; Josset et al., 2008; Josset et al., 2010a; Kiliyanpilakkil and Meskhidze, 2011; Nair and Rajeev, 2014; Murphy and Hu, 2021; Sun et al., 2023),  
60 these parameterizations do not account for atmospheric attenuation by aerosols and therefore have difficulty in calibrating the measured ocean surface reflectance accurately. This presents a difficulty for elastic backscatter lidars like CALIPSO, for which the signal is typically calibrated high in the atmosphere where molecular backscatter dominates and aerosol backscatter is insignificant or can be accurately estimated. The problem lies in the transfer of this calibration to the ocean surface, which entails accounting for the attenuation of the transmitted and backscattered light by the intervening atmosphere between the calibration region and the ocean surface. If coincident aerosol optical depth (AOD) data are available (e.g., from MODIS in  
65 the case of CALIPSO detailed in Josset et al. (2008)) then they may be used to estimate the intervening attenuation and transfer

the calibration. However, such data from passive sensors including MODIS are only available during daytime, are typically not produced in the vicinity of clouds and may have unacceptably high uncertainties for accurately accounting for aerosol attenuation. Estimation of the attenuation from the lidar data alone requires an assumption of the aerosol extinction-to-backscatter ratio (or “lidar ratio”), so errors in the assumed value can lead to an incorrect estimate of attenuation, especially when AOD is high. Because of this, the surface wind speed estimates in Hu et al. (2008) were limited to scenes with no clouds and negligible aerosol loading.

This study addresses retrieving surface wind speed directly from a lidar without other assumptions or external constraints by employing the high-spectral-resolution lidar (HSRL) technique through NASA Langley Research Center’s (LaRC’s) airborne High Spectral Resolution Lidar – generation 2 (HSRL-2) instrument (Hair et al., 2008). The HSRL-2 can directly measure vertically resolved aerosol backscatter and extinction profiles without relying on an assumed lidar ratio or other external aerosol constraints, enabling accurate estimates of the attenuation of the atmosphere. Therefore, the surface reflectance can be directly determined, providing a measure of the wave-slope variance and thus surface wind speed. Note that the HSRL-2 operates at a nadir-viewing geometry, which is detailed more in Sect. 2.4. At nadir or near-nadir incidence angles, the surface contribution of the lidar surface backscatter signal is the largest and is therefore sensitive to changes in wind speed (Josset et al., 2008; Josset et al., 2010a; Josset et al., 2010b), making it possible to introduce relatively simplified models of sea surface reflectance. However, Li et al. (2010) demonstrated that at the higher incidence angle lidar systems ( $> 15^\circ$ ), the sensitivity of the lidar surface signal would rapidly decrease as these highly non-nadir incidences shift the signal towards a subsurface contribution rather than a surface one. A more recent lidar study based on the highly non-nadir ( $\sim 37^\circ$ ) Aeolus UV HSRL lidar (Labzovskii et al., 2023) indirectly confirms this phenomenon by showing low agreement between passive remote sensing reflectivity and Aeolus surface reflectivity parameters over water surfaces such as oceans. For these reasons, an opportunity to retrieve ocean surface wind speeds using lidar ocean backscattering has been shown to be effective only for nadir or near-nadir lidar systems such as the HSRL-2.

This study details the HSRL-2’s surface wind speed retrieval methodology and evaluates this surface wind speed product through comparison with measurements from National Center for Atmospheric Research (NCAR) Airborne Vertical Atmospheric Profiling System (AVAPS) dropsondes. This work leverages an extensive data set from NASA’s Aerosol Cloud meTeorology Interactions oVer the western ATlantic Experiment (ACTIVATE) mission, which had multiple scientific and technological objectives described in Sect. 2.1 (Sorooshian et al., 2019). The mission consisted of six deployments between 2020 and 2022 and featured the joint deployment of the HSRL-2 and dropsonde launcher on one of its two aircraft to enable direct comparison between the two instrument data sets. The mission, dropsonde and HSRL-2 instrumentation, HSRL-2 algorithm, and the methods and results of using/evaluating the HSRL-2 and dropsonde surface wind speed data sets are all detailed in the following discussion.

**2.1 ACTIVATE Mission Description**

The HSRL-2 ocean surface wind speed product is assessed during the ACTIVATE campaign, which is a NASA Earth Venture Suborbital-3 (EVS-3) mission. The primary aim of ACTIVATE is to improve knowledge of aerosol-cloud-meteorology interactions, which are linked to the highest uncertainty among components contributing to total anthropogenic radiative forcing (Bellouin et al., 2020). There are three major scientific objectives: (i) characterize interrelationships between aerosol particle number concentration ( $N_a$ ), CCN concentration, and cloud drop number concentration ( $N_d$ ) with the goal of decreasing uncertainty in model parameterizations of droplet activation; (ii) advance process-level knowledge and simulation of cloud microphysical and macrophysical properties, including the coupling of aerosol effects on clouds and cloud effects on aerosol particles; and (iii) assess remote sensing capabilities to retrieve geophysical variables related to aerosol-cloud interactions. This study focuses on the third objective, which has already received attention with ACTIVATE data for retrievals other than ocean surface wind speeds (Schlosser et al., 2022; Van Diedenhoven et al., 2022; Chemyakin et al., 2023; Ferrare et al., 2023). ACTIVATE built a high volume of flight data statistics over the Western North Atlantic Ocean (WNAO) by flying six deployments across three years (2020 – 2022), with a winter and summer deployment each year (Sorooshian et al., 2023). Winter deployments included the following date ranges: 14 February – 12 March (2020), 27 January – 2 April (2021), 30 November 2021 – 29 March (2022). Summer deployments were as follows: 13 August – 30 September (2020), 13 May – 30 June (2021), 3 May – 18 June (2022). Across all three years, 90 King Air flights during the winter deployment were performed with 373 dropsondes launched while 78 flights during the summer deployment took place with 412 dropsondes launched.

Two NASA Langley aircraft flew in spatial and temporal coordination for the majority of the total flights (162 of 179). A “stacked” flight strategy was developed where a low-flying ( $< 5$  km) HU-25 aircraft collected in situ data in and just above the MABL while a high-flying ( $\sim 9$  km) King Air aircraft simultaneously provided remote sensing retrievals and dropsonde measurements in the same altitude range. In doing so, the stacked aircraft would simultaneously obtain data relevant to aerosol-cloud-meteorology interactions in the same column of the atmosphere and provide a complete picture of the lower troposphere (Sorooshian et al., 2019). In situ measurements of gases, particles, meteorological variables, and cloud properties were conducted by the HU-25 Falcon. The King Air’s payload included the NASA Goddard Institute for Space Studies (GISS) Research Scanning Polarimeter (RSP) and the two instruments relevant to this work: the NASA LaRC HSRL-2 and the NCAR AVAPS dropsondes (Sorooshian et al., 2023). An advantage of the joint deployment of HSRL-2 and AVAPS dropsondes on the King Air is that the data are spatially synchronized at launch, with wind drift of the dropsondes during descent accounted for with procedures summarized in Sect. 2.3.

130

The rationale to fly over the WNAO in different seasons was to collect data across a wide range of aerosol and meteorological regimes, with the latter promoting a broad range of cloud conditions (Painemal et al., 2021). A significant meteorological

feature is the North Atlantic Oscillation, which is the oscillation between the Bermuda-Azores High (high pressure system) and the Icelandic Low (low pressure system) (Lamb and Pepler, 1987). In the summer, the Bermuda-Azores High is at its peak and introduces easterly and southwesterly trade winds (Sorooshian et al., 2020). Starting in the fall, the Icelandic Low becomes prominent and introduces westerly winds in the boundary layer. The balancing act between these pressure systems dictates the climate of the North Atlantic and the prevailing transport processes (Li et al., 2002; Creilson et al., 2003; Christoudias et al., 2012). These transport processes that vary seasonally explain why winter flights coincided with more offshore (westerly) flow containing aerosol types impacted by anthropogenic influence (e.g., Corral et al., 2022), whereas summer flights included more influence from wildfire emissions and African dust among other sources both natural and anthropogenic in nature (Mardi et al., 2021; Aldhaif et al., 2020). Winds and turbulence tend to be stronger in the winter due to higher temperature gradients between the air and the ocean (Brunke et al., 2022), resulting in a higher fraction of available aerosol particles in the MABL that activate into cloud droplets in winter coinciding with cold air outbreaks as compared to summer (Dadashazar et al., 2021; Kirschler et al., 2022; Painemal et al., 2023). Therefore, this study region allows the HSRL-2 surface wind speed retrievals to be evaluated in various meteorological and aerosol loading conditions.

## 2.2 Dropsondes

The AVAPS system deployed during the ACTIVATE mission utilized the newer, more reliable NRD41 mini sondes. Their smaller form factor along with updates to their launching hardware increased reliability for launches since these instruments could be used with more aircraft and launcher configurations (Vömel and Dunion, 2023). A variable number of dropsondes were launched per flight, usually 3 to 4 for routine flights, with more being launched for specific targeted flight opportunities. With response times much less than 1 second, AVAPS samples position, wind speed (with  $0.5 \text{ m s}^{-1}$  uncertainty) (Vömel and Dunion, 2023), and state variables such as pressure, temperature, and humidity all the way to ~6 m above the ocean surface. The data are then post-processed via NCAR's Atmospheric Sounding Processing Environment (ASPEN) software where any spurious data are removed including any data returned from the ocean surface itself (Martin and Suhr, 2021). More details on the AVAPS system and its usage on other aircraft and missions can be found in (Vömel et al., 2021) and details of its usage in ACTIVATE specifically can be found in Vömel and Dunion (2023). Not many studies exist on surface wind speed validation of aircraft instruments with dropsondes (Bedka et al., 2021), so this study also highlights the potential of using dropsondes to validate aircraft surface wind speed data.

## 2.3 HSRL-2 Instrument Description

The NASA LaRC HSRL-2 is an airborne lidar instrument designed to enable vertically resolved retrievals of aerosol properties such as aerosol backscatter and depolarization at three wavelengths (355, 532, and 1064 nm), aerosol extinction at two wavelengths (355 and 532 nm) (Hair et al., 2008; Burton et al., 2018), and aerosol classification (Burton et al., 2012). In addition to these aerosol products, other retrieval capabilities include retrievals of atmospheric mixed layer height (Scarino et al., 2014), ocean subsurface particulate backscatter and attenuation coefficients (Schulien et al., 2017), cloud optical properties

165 (in development), and 10 m surface wind speeds, the latter of which is the focus of this study. Details of the laser receiver optics and detectors are described in detail in Hair et al. (2008). This analysis utilizes the 532 nm data channels that include a total scattering channel (both molecular and particulate scattering), molecular scattering only, and the cross polarized channel, which are internally calibrated during flight. Key to determining the optical transmission and subsurface signals is a molecular channel that filters essentially all the particulate and specular scattering using the iodine notch filter as described in Hair et al. 170 (2008), determining both the laser transmission down to the surface and correction of the subsurface scattering contribution to the integrated surface backscatter signal. ~~The lidar is operated in a nadir only viewing geometry (i.e., not scanning).~~

The laser is a custom built 200 Hz repetition rate Nd:YAG laser emitting at 1064 nm, which is converted to both the second and third harmonic wavelengths of 532 nm and 355 nm, respectively. The output laser energies are nominally 34 mJ (1064 175 nm), 11 mJ (532 and 355 nm each) and each is set to a divergence ( $1/e^2$ ) of approximately 0.8 mrad, giving a beam footprint diameter on the ocean surface of  $\sim 7$  m for the nominal 9 km King Air flight altitude. The telescope is set to a full field of view of 1 mrad, giving a viewing footprint diameter of 9 m at the ocean surface at nominal flight altitude. All three wavelengths are transmitted coaxially with the telescope through a fused silica window in the bottom of the aircraft are actively boresighted to the receiver. The HSRL-2 incorporates high speed photomultiplier tubes (PMTs) and custom amplifiers to allow data collection 180 at 120 MHz sampling rates with 40 MHz bandwidths. Data are sampled at 120 MHz (1.25 m in the atmosphere and 0.94 m in the ocean) with 16-bit digitizers and single-shot profiles are summed over 100 laser shots during 0.5 s which is the fundamental acquisition interval before storing to a disk. The aircraft incorporates an Applanix Inertial Navigation System (INS) to record the aircraft altitude at 0.5 s time intervals corresponding to each 100-shot data profile. ~~For the surface wind speed calculations, data are screened to limit the pitch and roll to less than  $\pm 3^\circ$  from the median values, which are approximately  $0^\circ$  for the roll and  $3^\circ$ – $5^\circ$  for pitch on the King Air.~~ 185

## 2.4 HSRL-2 Surface Wind Speed Retrieval Method

As mentioned in the previous section, a lidar system emits laser pulses into the atmosphere and the backscattered light from particles (aerosols) and molecules is collected with a telescope and imaged onto optical detectors where the generated analog electrical signal is digitally sampled as a function of time. Backscatter is also received from the reflection of the laser pulse off 190 the ocean surface and is referred to as the “surface return” signal. To derive surface wind speeds, the surface backscattered ( $180^\circ$ ) reflected radiance ( $\beta_{surf}$ , units  $sr^{-1}$ ) is estimated from the surface return signal and related to the wave-slope variance ( $\sigma^2$ ), as detailed in Josset et al. (2010b), through

$$\beta_{surf} = \frac{C_F}{4\pi\sigma^2 \cos^5(\theta)} e^{-\frac{\tan^2(\theta)}{\sigma^2}}, \quad (2)$$

where  $C_F$  is the Fresnel coefficient and is set to 0.0205 as given in Venkata and Reagan (2016) and  $\theta$  is the angle of incidence of the laser with the ocean surface. ~~The calculations are made only when the pitch and roll of the aircraft vary less than  $3^\circ$  from the median values to limit conditions where the aircraft altitude changes rapidly within the 0.5 s data collection window. As~~ 195

noted in the Introduction, the HSRL-2 is operated in a nadir-only viewing geometry (i.e., not scanning). However, there is a small offset from this nadir incidence angle due to the pitch and roll angles of the King Air aircraft. This offset angle is measured by the Applanix INS and is then used in Eq. 2 to derive the wave-slope variance. The median pitch and roll angles depend on the flight conditions (e.g., wind and fuel loads), but ranged from 2 - 5° for pitch and < 1° for roll during ACTIVATE flights. The surface wind speed data are screened to limit the pitch and roll to less than ± 3° from the median values, resulting in HSRL-2 incidence angles of < 3° for roll and < 8° for pitch. This screening effectively selects cases where the aircraft is flying straight and level legs.

The mean wind speed at 10 m above the sea surface ( $U$ ) is then derived using a piecewise empirical relationship between surface wind speed and wave-slope variance from Hu et al. (2008), where:

$$U = \left( \frac{\langle \sigma^2 \rangle}{0.0146} \right)^2, \langle \sigma^2 \rangle < 0.0386, U < 7 \text{ m s}^{-1}, \quad (3.1)$$

$$U = \left( \frac{\langle \sigma^2 \rangle - 3.0E-3}{5.12E-3} \right), 0.0386 \leq \langle \sigma^2 \rangle < 0.0711, 7 \text{ m s}^{-1} \leq U < 13.3 \text{ m s}^{-1}, \quad (3.2)$$

$$U = 10 \left( \frac{\langle \sigma^2 \rangle + 0.084}{0.138} \right), \langle \sigma^2 \rangle \geq 0.0711, U \geq 13.3 \text{ m s}^{-1}. \quad (3.3)$$

The relationships shown in Eqs. 3.1 – 3.3 were derived by Hu et al. (2008) using the comparisons between AMSR-E surface wind speeds and CALIPSO backscatter reflectance mentioned in Sect. 1 and agree identically with the Cox-Munk relationship for surface wind speeds between 7 m s<sup>-1</sup> and 13.3 m s<sup>-1</sup> and the log-linear relationship proposed by Wu (1990) for surface wind speeds above 13.3 m s<sup>-1</sup>.

With respect to surface wind speed retrievals, the HSRL-2 instrument offers two major advantages over standard backscatter lidars such as CALIPSO: 1) it can account for atmospheric attenuation between the aircraft and the surface so retrievals can be performed without constraining the retrieval to low AOD conditions (i.e., negligible aerosol loading) or assuming the lidar ratio, and 2) it has high vertical resolution sampling (1.25 m) that enables accurate correction for ocean subsurface scattering, which makes a small but non-negligible contribution to the measured surface return. The equations for the HSRL-2 532-nm measurement channels are:

$$P_{mol}(r) = G_{mol} \frac{1}{r^2} F(r) \beta_m^{\parallel}(r) T^2(r), \quad (4.1)$$

$$P_{tot}(r) = G_{mol} G_{i2} \frac{1}{r^2} \left[ \left( \beta_p^{\parallel}(r) + \beta_m^{\parallel}(r) \right) + G_{dep} \left( \beta_p^{\perp}(r) + \beta_m^{\perp}(r) \right) \right] T^2(r) \quad (4.2)$$

where  $P_x$  is the total measured signal per sampling interval by the lidar and  $r$  denotes the range from the lidar. Here the *mol* subscript denotes the measured signal on the molecular channel, for which all particulate backscatter and the surface return is blocked by an iodine vapor filter. The *tot* subscript denotes the “total” backscatter calculated from the sum of two measurement channels, the co-polarized channel and the cross-polarized channel. These channels are essentially elastic backscatter lidar channels similar to the 532 nm channels on CALIPSO, in that they measure attenuated backscatter from both molecules and

particles. The co-polarized channel measures backscatter that is polarized parallel to the linear polarization of the transmitted laser pulses, and the cross-polarized channel measures backscatter with polarization perpendicular to the laser pulses. The volume backscatter coefficient,  $\beta$  (units  $m^{-1} sr^{-1}$ ), is separated into components arising from either molecular scattering ( $m$ ) or particulate scattering ( $p$ ) and by polarization parallel ( $\parallel$ ) and perpendicular ( $\perp$ ) to the laser. The combined collection efficiency, optical efficiency, and the overall electronic gain for the signals is denoted by  $G_x$ . The  $T^2$  factor is the two-way transmission of the atmosphere, which accounts for both molecular and particulate scattering and absorption between the lidar and range  $r$ . A full description of the instrument channels are described in Hair et al. (2008).

Eqs. 4.1 and 4.2 are generalized such that the backscatter coefficients and transmission factors can be either from the atmosphere or ocean, depending on the altitude (or depth) of the scattering volume. Also, the transmission of the molecular backscatter through the iodine vapor filter,  $F$ , is based on either the atmosphere ( $atm$ ) or the ocean ( $ocn$ ) scattering regions, as they have different backscatter spectra and thus different iodine filter transmission factors, both of which are determined by laboratory calibrations and modeled molecular scattering spectra (Hair et al., 2008). Calibration operations are conducted during each flight to provide the relative gain ratios between the molecular ( $mol$ ) and co-polarized ( $par$ ) channels,  $G_{iz}$ , and between the co-polarized and cross-polarized ( $per$ ) channels,  $G_{dep}$ , such that

$$G_{iz} = \frac{G_{par}}{G_{mol}}, \quad G_{dep} = \frac{G_{per}}{G_{par}}. \quad (5)$$

After the internal gain ratios (Eq. 5) are applied, the two signals (Eqs. 4.1 and 4.2) have the same relative gain. As will be shown below, the retrieval implements ratios of these two signals, and therefore neither the absolute gain nor any other absolute calibration factor is required to determine the surface backscatter.

To calculate the surface backscatter, the overall system response must be accounted for. The measured signal ( $P$ ) is the convolution of the normalized system response, ( $L$ ), with the ideal measured signal (i.e., infinite detection bandwidth and delta-function-like laser pulse), this signal being the gain-scaled ( $G$ ), range-scaled ( $\frac{1}{r^2}$ ), attenuated ( $T^2$ ) backscatter coefficient ( $\beta$ , units  $m^{-1}sr^{-1}$ ), which can be written as

$$P_{ideal}(r) = G \frac{1}{r^2} \beta(r) T^2(r). \quad (6a)$$

$$P(r) = G \int_{-\infty}^{\infty} L(r - \rho) P_{ideal}(\rho) d\rho. \quad (6b)$$

The system response includes the impact of the laser's temporal pulse shape, detector response, and analog electronic filter response.



To account for different scattering media and to better understand how the system response impacts the surface backscatter calculation, it is helpful to separate the total scattering channel,  $P_{tot}(r)$ , into three contributions: atmosphere [*atm*], surface [*surf*], and ocean [*ocn*] as follows:


$$P_{tot}(r) = P_{tot}^{atm}(r) + P_{tot}^{ocn}(r) + P_{tot}^{surf}(r). \quad (7)$$

250 Using Eq. 6, the last term in Eq. 7,  $P_{tot}^{surf}(r)$ , can be written as

$$P_{tot}^{surf}(r) = G_{mol}G_{i2} \int_{-\infty}^{\infty} L(r - \rho) \frac{1}{\rho^2} \beta_{surf} \delta(\rho - r_s) T^2(\rho) d\rho \quad (8a)$$

$$P_{tot}^{surf}(r) = G_{mol}G_{i2} L(r - r_s) \frac{1}{r_s^2} \beta_{surf} T^2(r_s) \quad (8b)$$

where the range to the ocean surface is  $r_s$  and the volume backscatter coefficient for the ocean surface is represented as  $\beta_{surf} \delta(\rho - r_s)$  (units  $m^{-1} sr^{-1}$ ), where  $\delta(\rho - r_s)$  is the Dirac delta function centered at  $r_s$ . Figure 1 illustrates the vertical distributions of the measured signals  $P_{tot}$  (black) and  $P_{mol}$  (blue) along with the  $P_{tot}^{surf}$  (green) component of  $P_{tot}$ . Note that zero altitude is the location of the ocean surface.



255

**Figure 1: Visualization of HSRL-2 measurement signals as described in Eqs. 6 – 8. Dashed line denotes ideal total backscatter signal from the atmosphere, surface reflection, and the ocean subsurface. Blue and black lines denote measured signals from total and molecular scattering channels, respectively. Red and green lines show the ocean corrected signal and the ocean surface backscatter, respectively. Dots indicate the altitudes of digitized samples. The sampling rate is 120 MHz, resulting in a vertical spacing of 1.25 m in the atmosphere and 0.94 m in the ocean.**

260

It is seen from Fig. 1 and Eq. 8b that the surface component  $P_{tot}^{surf}$  of the measured signal  $P_{tot}$  is not localized to the surface but is instead spread above and below the surface via convolution with the system response function. The atmosphere and ocean components of  $P_{tot}$  are also impacted by the convolution as is  $P_{mol}$ . Rearranging Eq. 8 and integrating the total surface backscatter component over the full vertical extent of the system response function (i.e., to  $\pm\Delta z$ ), the surface response function

265 can be eliminated in the representation of  $\beta_{surf}$  as shown in Eq. 9.

$$\beta_{surf} = \frac{1}{G_{mol} G_{i2}} \frac{r_s^2}{T^2(r_s)} \int_{r_s - \Delta z}^{r_s + \Delta z} P_{tot}^{surf}(r) dr \quad (9)$$

Of course, the measurement that can be accessed is  $P_{tot}$ , not the surface component  $P_{tot}^{surf}$ . If  $P_{tot}$  were substituted for  $P_{tot}^{surf}$  in Eq. 9,  $\beta_{surf}$  would be overestimated due to the contribution of ocean subsurface backscatter. The atmospheric contribution is negligible (i.e., <0.05%) and can be ignored. The magnitude of the contribution of the ocean subsurface scattering depends on the level of ocean particulate (hydrosol) and as well as molecular seawater backscatter. The magnitude of this scattering relative to the surface backscatter can impact the retrieved surface wind speed accuracy. For example, at  $U = 7 \text{ m s}^{-1}$  and assuming pure seawater (i.e., no hydrosols), the integrated total surface signal would be 5.7% higher than the integrated surface backscatter. This results in a decrease of  $0.75 \text{ m s}^{-1}$  (-11% error) in the estimated surface wind speed. At a  $20 \text{ m s}^{-1}$  surface wind speed, the error in the calculated surface wind speeds results in a decrease by  $2.7 \text{ m s}^{-1}$  (-14% error). The ocean subsurface correction becomes less as the particulate scattering (or absorption) increases due to increased attenuation in the seawater and therefore contributes less over the integration window around the ocean surface. Therefore, the ocean subsurface contribution is higher for clear water compared to turbid water. For example, in the case illustrated in Fig. 1, the seawater particulate and molecular scattering are equal, resulting in a contribution of only 3.8% to the integrated surface backscatter as compared to the no particulate scattering noted above of 5.7%. The atmospheric signal contribution is much less (~100 times smaller) than the ocean subsurface signal and therefore its contribution is considered negligible. Fortunately, the high vertical resolution of the HSRL-2 instrument enables the ocean subsurface contribution to be estimated. The separation of the molecular signal also enables estimation of the two-way transmittance,  $T^2$ , and gain factor,  $G_{mol}$ , in Eq. 9.

For the HSRL-2 instrument, the two-way transmittance is determined directly from the measured molecular channel,  $P_{mol}$ . The two-way total (particulate and molecular attenuation) transmittance to the surface can be calculated as follows,

$$T^2(r_{ns}) = \frac{1}{G_{mol}} \frac{\overline{P_{mol}(r_{ns})r_{ns}^2}}{F(r_{ns})\beta_m^{\parallel}(r_{ns})}, \quad (10)$$

where  $F$  is the iodine vapor filter function (known from lab and in-flight calibration),  $\beta_m^{\parallel}$  is the molecular backscatter coefficient for the atmosphere (computed from pressure and temperature data from a reanalysis model), and  $\overline{P_{mol}(r_{ns})r_{ns}^2}$  is the range-scaled molecular channel signal near the ocean surface (where  $r_{ns}$  is the near-surface range). In practice, this is computed by averaging data from 60 m to 180 m above the surface. This range is somewhat arbitrary but is chosen as a balance between ensuring that the signal does not include any of the surface reflectance and low enough to capture most of the attenuation down to the surface. Substituting Eq. 10 into Eq. 9, one can solve for the surface backscatter,

$$\beta_{surf} = \frac{1}{G_{iz}} \frac{\int_{r_s-\Delta z}^{r_s+\Delta z} r^2 P_{tot}^{surf}(r) dr}{\frac{\overline{P_{mol}(r_{ns})r_{ns}^2}}{F(r_{ns})\beta_m^{\parallel}(r_{ns})}}. \quad (11)$$

To account for the ocean subsurface contributions to the measured signal, Eq. 6 can be rearranged as

$$P_{tot}^{surf}(r) = P_{tot} - P_{tot}^{atm}(r) - P_{tot}^{ocn}(r). \quad (12)$$

A benefit of the HSRL-2 retrieval algorithm is that one can use the molecular channel signal to determine the ocean signal near the surface (see Fig. 1). To determine the near-surface ocean signal, an estimate of the total ocean scattering ratio (TSR) is employed, which is the ratio of molecular + hydrosol backscatter divided by molecular backscatter. An estimate of the near-surface TSR,  $(\overline{TSR}_{ocn})$  is computed using the quotient of the total and molecular channels ( $P_{tot} / P_{mol}$ ) averaged over a small range of depths just below the depth at which the surface signal response goes to zero, as follows:

$$\overline{TSR}_{ocn} \equiv \left( \frac{\beta_p + \beta_m}{\beta_m} \right) = \frac{F_{ocn}(r)}{G_{i2} \Delta r} \int_{r_s + \Delta z}^{r_s + 2\Delta z} \frac{P_{tot}(r)}{P_{mol}(r)} dr \quad (13)$$

where  $F_{ocn}$  accounts for the spectral transmission of the molecular seawater backscatter through the iodine vapor filter and is determined via in-flight and laboratory calibrations. The ocean subsurface component of the total channel backscatter is estimated as follows:

$$P_{tot}^{ocn}(r) = \overline{TSR}_{ocn} G_{i2} \frac{P_{mol}(r)}{F_{ocn}(r)}, \text{ below the surface } (r > r_s) \quad (14)$$


Here the assumption is that the TSR is vertically constant near the surface over the 0.5 s (~75 m horizontal resolution) integration of the lidar signals. Combining Eqs. 11, 12, and 14 and ignoring the atmospheric contribution  $P_{tot}^{atm}$  to the total channel signal, one can compute the absolute surface backscatter using the two measured channels as

$$\beta_{surf} = \frac{\int_{r_s - \Delta z}^{r_s + \Delta z} \left( \frac{P_{tot}(r)}{G_{i2}} - \overline{TSR}_{ocn} * P_{mol}(r) \right) r^2 dr}{\frac{P_{mol}(r_{ns}) r_{ns}^2}{F_{atm}(r_{ns}) \beta_m^{\parallel}(r_{ns})}} \quad (15)$$

The use of the molecular channel in this way cancels out absolute system gain constant ( $G_{mol}$ ), provides an estimate of the two-way transmittance of the atmosphere, and enables subtraction of ocean subsurface backscatter. It does not require precise knowledge of the system response function or any other assumptions. With Eq. 15, one can calculate the wave-slope variance through Eq. 2 and then use Eqs. 3.1 – 3.3 to derive surface wind speeds.

In addition to the specular reflection from the surface, whitecaps or sea foam can increase the lidar backscatter signal. As noted in Josset et al. (2010b), the contribution of scattering by the whitecaps on the ocean surface has been treated as Lambertian scattering. There is a wavelength dependence of the scattering at longer wavelengths due to the water absorption, based on measurements presented by Dierssen (2019) covering wavelengths from 0.4 – 2.5  $\mu\text{m}$ . Measurements presented here are at 532 nm, a region of the visible spectrum where scattering from foam is relatively constant with wavelength. The contribution of whitecaps is typically modeled with a constant average reflectance and an effective area weighted fraction that varies with surface wind speed (Whitlock et al., 1982; Koepke, 1984; Gordon and Wang, 1994; Moore et al., 2000). Following Moore et al. (2000), we have estimated the average reflectance due to the whitecaps as a function of surface wind speed and the difference becomes  $> 1 \text{ m s}^{-1}$  for surface wind speeds  $> 15 \text{ m s}^{-1}$  based on this relationship. As presented below, there are limited data (49 data points) above  $13.3 \text{ m s}^{-1}$  that can be compared to the dropsonde surface wind speeds to evaluate this

relationship. Moreover, since the correction depends on surface wind speed, an iterative calculation is required to use this  
320 relationship as the backscatter is dependent on wind speed.



**Figure 2. Estimated absolute difference in calculated surface wind speed if reflectance from whitecaps is not included. The lidar surface backscatter is higher than the specular reflectance if whitecaps are present, which results in a lower estimated surface wind speed if not accounted for in the retrieval.**

325


Alternatively, Hu et al. (2008) used a full month of CALIPSO integrated surface depolarization ratio (ratio of the integrated cross polarized channel to the integrated co-polarized channel across the surface) and applied an empirical correction to the reflectance that was determined using AMSR-E data as the ground-truth data set to increase the correlation of the data sets. The correlation was based on much more data than the ACTIVATE matchups between HSRL-2 and dropsondes, limiting the utility  
330 of a similar analysis with the HSRL-2. In addition, there are significant differences in the configurations of CALIPSO and HSRL-2 that limit implementation of the same empirical relationship. First, CALIPSO's integrated surface depolarization includes the subsurface contributions due to its 30 m vertical resolution, whereas the HSRL-2 surface depolarization is integrated over only a few meters as shown in Fig. 1. Second, the CALIPSO data is based on global data, which is dominated by oligotrophic (clear) waters, whereas a significant fraction of the HSRL-2 - dropsonde comparisons are from eutrophic and  
335 mesotrophic waters near the coast and along the shelf. Third, there is a significant difference in footprint size between HSRL-2 and CALIPSO (8 m versus 90 m), with HSRL-2's instantaneous footprint area being greater than 2 orders of magnitude

smaller and, considering HSRL-2's along-track averaging (100 laser shots) compared to CALIPSO's single shot data, greater than one order of magnitude smaller in terms of area over which surface depolarization is integrated.

## 2.5 Collocation and Statistical Procedures

340 Since surface wind speeds are the focus of this study, first the dropsonde wind speed data points closest to 10 m (altitude of  $11.56 \text{ m} \pm 3.19 \text{ m}$  for the 577 points) above sea level are recorded for each launch (multiple launches per flight) to allow meaningful comparison with the HSRL-2 surface wind speeds. Since one data point was taken per dropsonde for each flight, there are 160 recorded dropsonde measurements for 2020, 245 measurements for 2021, and 335 measurements for 2022. Then, the HSRL-2 surface wind speed retrieval closest in space and time to the corresponding dropsonde measurement is recorded.

345 Collocation between the HSRL-2 and the dropsondes is constrained to below 30 km horizontally and below 15 minutes temporally to remove outliers while trying to maximize the number of data points to be used in the study. Further constraining these distance and time conditions would eliminate more data points with negligible improvement to the statistics as shown by Figs. S1 and S2 in the supplement. Due to missing data in the HSRL-2 data set and the removal of outliers based on collocation constraints, 577 data points are available for comparison between the dropsondes and the HSRL-2 (Fig. 3).



350

**Figure 3: Map of 577 ACTIVATE dropsondes launched from the King Air between 2020 and 2022 that are used to evaluate the HSRL-2 surface wind speed retrievals introduced in this study.**

After the surface wind speed data are prepared using the procedure above, scatterplots along with the correlation coefficient (r), linear regression, and ordinary least squares bisector regression (OLS-bisector) are used to visually demonstrate how well HSRL-2 surface wind speed data match dropsonde data and show any potential variability in the data. Since OLS-bisector is less common than linear regression, a brief explanation of their differences is provided. In linear regression, X is treated as the independent variable while Y is treated as the dependent variable. In other words, one observes how Y varies with changes to fixed X values. OLS-bisector is known as an errors-in-variable regression technique, where X and Y are both dependent variables and thus both subject to error. OLS-bisector regresses Y on X (standard OLS) and then regresses X on Y (inverse OLS), then bisects the angle of these two regression lines (Ricker, 1973). Although other errors-in-variable techniques exist (e.g., Deming regression, orthogonal distance regression), OLS-bisector is chosen because it calculates the error present in both data sets using the bisector rather than assuming an error a priori like the examples mentioned (Wu and Yu, 2018). After performing these regressions, histograms of surface wind speed deltas, which are defined as HSRL-2 surface wind speed minus dropsonde surface wind speed, are created to show the distribution and spread of the data more easily. The mean and standard deviation (STD), of the surface wind speed deltas are computed and then used to define the mean error (mean  $\pm$  STD). This metric is used to evaluate how accurately the HSRL-2 retrieves surface wind speeds.


### **3 Results and Discussion**

#### **3.1 Case Studies**

Before delving into the HSRL-2 – dropsonde surface wind speed intercomparisons in full statistical detail, surface wind speed data from two ACTIVATE research flights are analyzed: Research Flight 29 on 28 August 2020 and Research Flight 14 on 1 March 2020. These flights are analyzed to demonstrate the ability of the HSRL-2 to 1) provide profiles that show the spatial variability of surface wind speed over time, which are beneficial to observe phenomena like sea-surface temperature dynamics and cloud evolution and 2) sample the surface in broken cloud scenes, showing that the retrievals are not limited to cloud- and aerosol-free conditions like in Hu et al. (2008).

##### **3.1.1 Research Flight 29 on 28 August 2020**

Research Flight 29 was a near cloud-free day where an above average number of dropsondes were launched and ACTIVATE's aircraft were coordinated with the CALIPSO satellite overpass. These conditions allow for the examination of how the high horizontal spatial resolution of the HSRL-2 (~75 m along track as mentioned in Sect. 2.4) influences its retrievals and how the data can be used to track sea surface temperature (SST) gradients common to the WNAO (Painemal et al., 2021) as seen in Fig. 4. Note that Fig. 4a uses SST data from Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2) (Gelaro et al., 2017) to contextualize the SST gradients present in the WNAO, and no comparisons with MERRA-2 surface wind speed data are performed in this study.



385 **Figure 4: a) Flight map of King Air (black line) and dropsondes (dark yellow circles) overlaid onto map of MERRA-2 mean sea-surface temperature (SST) data (Gmao, 2015) for Research Flight 29 on 28 August 2020. White dashed line corresponds to the CALIPSO overpass coincident with King Air flight path. Time stamps represent where the King Air crosses over sharp SST changes associated with the Gulf Stream. b) Time series of surface wind speed data from HSRL-2 and dropsondes for the same flight, where the black solid line signifies total HSRL-2 surface wind speed data and circles indicate collocated surface wind speed data points. Black dashed lines represent time stamps of interest as indicated in a).**


395 It is seen that changes in the HSRL-2 surface wind speeds (Fig. 4b) correspond with changes in SST (Fig. 4a), especially seen at 17:51 and 18:24 (UTC throughout paper). As the aircraft approaches and crosses the SST boundary at 17:51 (i.e., SST increasing), there is a corresponding increase in surface wind speeds. The reverse observation can be seen when the aircraft approaches and crosses the boundary at 18:24 (i.e., SST decreasing), where surface wind speeds noticeably decrease. Although further analysis is needed to rigorously examine the relationship between surface wind speed and SST, these observations show that the HSRL-2 has the high horizontal spatial resolution needed to probe the fine-scale variability of surface wind speeds and has the potential to improve atmospheric modeling of MABL processes. These profiles capture the spatial gradients in surface wind speeds that would otherwise not be available with the dropsondes alone, since these instruments can only take point measurements as they drop vertically to the surface and therefore cannot provide the horizontal spatial extent like the derived HSRL-2 surface wind speed product can.

### 3.1.2 Research Flight 14 on 1 March 2020

405 Next, Research Flight 14 is shown in Fig. 5 to demonstrate the ability of the HSRL-2 to sample in broken cloud scenes. This flight along with the associated morning flight on 1 March 2020 have been the subject of several studies owing to its coincidence with cold air outbreak conditions (see cloud streets in Fig. 5a) and a flight strategy that allowed for detailed characterization of the evolving aerosol-cloud system as a function of distance offshore (Seethala et al., 2021; Chen et al.,



2022; Li et al., 2022; Tornow et al., 2022; Sorooshian et al., 2023). The morning flight focused on a location with very detailed characterization including stacked level flight legs (i.e., termed a “wall”) with the Falcon flying below, in, and above clouds, with the King Air flying aloft to further characterize the same region. The afternoon flight consisted of both aircraft flying back to that same location, adjusting the sampling strategy to fly along the boundary layer wind direction in a quasi-Lagrangian fashion to keep studying the evolution of the air mass characterized in the morning. The afternoon flight is chosen because it shows the full range of cloud conditions from clear to completely overcast. Therefore, the HSRL-2 surface wind speed retrievals are able to be evaluated in this range of conditions.




415 **Figure 5: a) Flight map of the King Air (red line), Falcon (yellow line), and dropsondes (dark yellow circles) overlaid onto**  
**Geostationary Operational Environmental Satellite (GOES-16) cloud imagery for Research Flight 14 on 1 March 2020. Blue stars**  
**represent time stamps where the King Air crosses over from cloud-free to cloudy areas. b) Time series of surface wind speed data**  
 420 **from HSRL-2 and dropsondes for the same flight, where lines signify total HSRL-2 surface wind speed data and circles indicate**  
**collocated surface wind speed data points. Blue dashed lines represent time stamps of interest as indicated in a).**

As the aircraft approaches the cloud scene at 19:18, there is a noticeable and steady increase of HSRL-2 surface wind speeds. The reverse observation is seen when the aircraft approaches 21:15, where the HSRL-2 surface wind speeds start to decrease steadily. As highlighted in the 28 August 2020 case study, the high horizontal spatial resolution of the HSRL-2 retrievals enables these spatial gradients to be observed. Another important takeaway is the HSRL-2 is still able to sample the surface in cloud scenes, as seen by the almost complete surface wind speed profile in Fig. 5b. Although a gap in data occurs at 20:15 where cloud cover is most substantial, some retrievals are still present in that area. The reason is that the HSRL-2 can probe the surface through gaps between clouds, allowing for the surface wind speed retrievals to take place. Although the HSRL-2

retrievals would be unavailable in overcast cloud scenes, the ability of the instrument to sample the surface in broken cloud fields and not just aerosol- and cloud-free scenes is a significant benefit of the lidar and the HSRL technique.

### 430 3.2 HSRL-2 – Dropsonde Comparisons

Now, the collocated HSRL-2 retrievals and dropsonde measurements of surface wind speed are compared and the results are shown in Fig. 6.




435 **Figure 6: Scatterplots with associated histograms for HSRL-2 – dropsonde collocated surface wind speed data points using ACTIVATE’s 2020 – 2022 data set. N represents the number of data points.**

The comparison yields correlation coefficients of 0.89, slopes of 1.04 and 1.17, and y-intercepts of  $-0.13 \text{ m s}^{-1}$  and  $-1.05 \text{ m s}^{-1}$  for linear and bisector regressions, respectively. Note that the correlation coefficients are the same for linear and bisector regressions throughout this analysis, so they are listed as one value throughout Sect. 3.2. Using the mean and STD values in the same figure, the mean error or accuracy of the HSRL-2 surface wind speed retrievals is  $0.15 \text{ m s}^{-1} \pm 1.80 \text{ m s}^{-1}$ . These results show that on average, the HSRL-2 slightly overestimates surface wind speeds and the estimation can be off by about 2

440  $\text{m s}^{-1}$  in either direction.

Now that the HSRL-2 retrievals have been broadly evaluated, Fig. 7 shows how their accuracy varies per  $1 \text{ m s}^{-1}$  interval in surface wind speed. This plot also provides the opportunity to compare the Hu et al. (2008) model with the models proposed by Cox and Munk (1954) and Wu (1990) to see if some of the error in the HSRL-2 retrievals can be attributed to model


445 characteristics.



**Figure 7: HSRL-2 surface wind speed using Hu, Cox-Munk, and Wu models versus mean dropsonde surface wind speed calculated per 1 m s<sup>-1</sup> bin. A histogram of dropsonde surface wind speeds is also included to show their distribution.**

It is seen that the mean Cox-Munk and Wu surface wind speed values are higher than the mean Hu values from 0 m s<sup>-1</sup> to 7 m s<sup>-1</sup>, showing that the Cox-Munk and Wu relationships overestimate dropsonde surface wind speeds more than the Hu relationship. The variability (i.e., STD) around the mean per bin is similar between the three models, which is 1.59 m s<sup>-1</sup> for Hu, 1.43 m s<sup>-1</sup> for Cox-Munk, and 1.55 m s<sup>-1</sup> for Wu on average. Although similar, the STD of the Hu surface wind speeds found here is ~0.4 m s<sup>-1</sup> lower than the one found in Fig. 6. This could be attributed to an STD not being able to be calculated for the 17 to 18 m s<sup>-1</sup> bin since it only contained one point.

Although it is apparent Cox-Munk and Wu retrievals overestimate dropsonde observations for surface wind speeds below 7 m s<sup>-1</sup>, it is still unclear which of the models perform better overall. Therefore, the y-axis from Fig. 7 is converted to wave-slope space and the result of this modification is shown in Fig. 8. HSRL-2 wave-slope is used because it directly reports the original measurements of surface reflectance rather than estimated values of surface wind speed. Using the original data ensures that uncertainty is coming from the actual HSRL-2 – dropsonde comparisons rather than from potential errors in the conversion from wave-slope to surface wind speed.




465 **Figure 8: HSRL-2 wave-slope variance versus mean dropsonde surface wind speed calculated per  $1 \text{ m s}^{-1}$  bin. Ideal Hu, Cox-Munk, and Wu distributions are included to show how well observed dropsonde data match with each parameterization. A histogram of dropsonde surface wind speeds is also included to show their distribution.**

470 From Fig. 8, it is more easily seen how the dropsonde surface wind speed distribution compares with Hu, Cox-Munk, and Wu parameterizations. Dropsonde surface wind speeds match quite closely to Hu and Cox-Munk parameterizations as opposed to the Wu parameterization between  $7 \text{ m s}^{-1}$  and  $13.3 \text{ m s}^{-1}$ , although some divergence is seen above  $\sim 10.5 \text{ m s}^{-1}$ . However, a critical observation that is more apparent in Fig. 8 than Fig. 7 is how the dropsonde data most resemble the Hu distribution for surface wind speeds below  $7 \text{ m s}^{-1}$ . This improvement is substantial, especially since most of the surface wind speeds in ACTIVATE fall into this category. Surface wind speeds above  $13.3 \text{ m s}^{-1}$  substantially diverge from all models, especially above  $16 \text{ m s}^{-1}$ . As mentioned previously, there are few surface wind speed observations in this category, so more measurements are necessary to make meaningful comparisons between the two data sets. Overall, Figs. 7 and 8 demonstrate the benefits of using the Hu parameterization in this study and why surface wind speeds above  $13.3 \text{ m s}^{-1}$  are not the main focus of the comparisons in this section. Further analysis is warranted to rigorously compare the performance of various surface reflectance models and potentially apply corrections (i.e., whitecap correction for surface wind speeds above  $13.3 \text{ m s}^{-1}$ ), but

475

the aim of this paper is to evaluate LARC’s HSRL-2 surface wind speed retrieval algorithm using the available ground-truth dropsonde measurements.

480 Now that the Hu relationship has been deemed the more effective model through the preliminary analysis shown in Figs. 7 and 8, a more rigorous statistical analysis is performed for surface wind speeds 1) below  $7 \text{ m s}^{-1}$  and 2) between  $7 \text{ m s}^{-1}$  and  $13.3 \text{ m s}^{-1}$  to assess the overall accuracy of the HSRL-2 retrievals in these categories (Fig. 9).




485 **Figure 9: Scatterplots with associated histograms for HSRL-2 – dropsonde collocated surface wind speed data points for a) surface wind speeds  $< 7 \text{ m s}^{-1}$  and b) surface wind speeds between  $7 \text{ m s}^{-1}$  and  $13.3 \text{ m s}^{-1}$ . Note that x- and y-axis ranges vary to better showcase results in individual panels. N represents the number of data points.**

Intercomparisons for surface wind speeds below  $7 \text{ m s}^{-1}$  (Fig. 9a) show correlation coefficients of 0.66, slopes of 0.65 and 0.99, and y-intercepts of  $1.10 \text{ m s}^{-1}$  and  $-0.49 \text{ m s}^{-1}$  for linear and bisector regressions, respectively. The accuracy of the HSRL-2 retrievals is calculated to be  $-0.54 \text{ m s}^{-1} \pm 1.34 \text{ m s}^{-1}$ , showing that the HSRL-2 on average underestimates surface wind speeds and this estimation could vary by  $\pm 1.34 \text{ m s}^{-1}$ . For surface wind speeds between  $7 \text{ m s}^{-1}$  and  $13.3 \text{ m s}^{-1}$  (Fig. 9b),

490

495 correlation coefficients of 0.75, slopes of 0.64 and 0.85, and y-intercepts of  $3.80 \text{ m s}^{-1}$  and  $1.87 \text{ m s}^{-1}$  are reported for linear and bisector regressions, respectively. The mean error of  $0.56 \text{ m s}^{-1} \pm 1.49 \text{ m s}^{-1}$  shows that the HSRL-2 overpredicts surface wind speeds by about  $\sim 0.5 \text{ m s}^{-1}$  on average with a variability of  $\pm \sim 1.5 \text{ m s}^{-1}$ . Therefore, the means from both categories average out to  $\sim 0 \text{ m s}^{-1}$  since they are approximately the same but in opposite directions. Separating the data into these categories highlight an important result that could not be seen in the cumulative data (Fig. 6): one can expect bias of up to  $\sim 0.5 \text{ m s}^{-1}$  in either direction and error of up to  $\sim 1.5 \text{ m s}^{-1}$  on average for most HSRL-2 surface wind speed retrievals in ACTIVATE.

The data are then divided into winter and summer deployments (dates provided in Sect. 2.1) as shown in Fig. 10 to assess the HSRL-2's retrieval accuracy in different seasons.



500

**Figure 10: Scatterplots with associated histograms for HSRL-2 – dropsonde collocated surface wind speed data points for a) winter and b) summer deployments. Data are highlighted based on surface wind speed categories:  $7 \text{ m s}^{-1} \leq \text{Wind Speed} < 13.3 \text{ m s}^{-1}$ ,  $\text{Wind Speed} < 7 \text{ m s}^{-1}$ , and  $\text{Wind Speed} \geq 13.3 \text{ m s}^{-1}$ . -N represents the number of data points.**

As seen in Fig. 10a, the winter surface wind speed intercomparisons show correlation coefficients of 0.88, slopes of 0.95 and  
 505 1.08, and y-intercepts of  $1.03 \text{ m s}^{-1}$  and  $-0.08 \text{ m s}^{-1}$  for linear and bisector regressions, respectively. The summer surface wind  
 speed intercomparisons (Fig. 10b) have correlations of 0.87, slopes of 1.08 and 1.24, and y-intercepts of  $-0.69 \text{ m s}^{-1}$  and  $-1.68$   
 $\text{m s}^{-1}$ . Finally, the mean errors for winter and summer, respectively, are reported as  $0.63 \text{ m s}^{-1} \pm 2.07 \text{ m s}^{-1}$  and  $-0.18 \pm 1.52 \text{ m}$   
 $\text{s}^{-1}$ . It is seen that the error in the HSRL-2 estimations of surface wind speeds is larger for winter than summer, most likely due  
 to the higher fraction of surface wind speeds above  $13.3 \text{ m s}^{-1}$  and lower fraction of them below  $7 \text{ m s}^{-1}$  in the winter. This  
 510 observation makes sense because of the increased presence of clouds, precipitation, and whitecaps for the higher surface wind  
 speeds observed in the winter. These observations show that HSRL-2 retrievals of surface wind speed are more accurate in the  
 summer over the winter. However, the HSRL-2 can still make numerous accurate retrievals as shown by the Fig. 10 and the 1  
 March 2020 research flight discussions. Caution must still be exercised when using data from days featuring turbulent  
 meteorological conditions that could induce whitecaps and/or substantial cloud cover that could limit or even prevent the  
 515 HSRL-2 from sampling the surface.

Statistics evaluating the HSRL-2 surface wind speed retrievals (Figs. 6, 9, 10) are summarized in Table 1 for convenience.

**Table 1: Summary of all HSRL-2 – dropsonde surface wind speed comparison statistics shown in Figs. 6, 9, and 10. The two values  
 520 for slope and y-intercept refer to those for the linear and bisector regressions, in that order. R values are the same for both linear  
 and bisector regressions, so they are listed as one value.**

	N	r	Slope	Y-intercept [m s <sup>-1</sup> ]	Mean Error [m s <sup>-1</sup> ]
<b>Overall</b>	577	0.89	1.04/1.17	-0.13/-1.05	$0.15 \pm 1.80$
<b>Wind Speed &lt; 7 m s<sup>-1</sup></b>	292	0.66	0.65/0.99	1.10/-0.49	$-0.54 \pm 1.34$
<b>7 m s<sup>-1</sup> ≤ Wind Speed &lt; 13.3 m s<sup>-1</sup></b>	236	0.75	0.64/0.85	3.80/1.87	$0.56 \pm 1.49$
<b>Winter</b>	236	0.88	0.95/1.08	1.03/-0.08	$0.63 \pm 2.07$
<b>Summer</b>	341	0.87	1.08/1.24	-0.69/-1.68	$-0.18 \pm 1.52$

#### 4. Conclusions

This study introduces the High Spectral Resolution Lidar – generation 2 (HSRL-2) surface wind speed retrieval method,  
 demonstrates its use, and evaluates its accuracy using NCAR AVAPS dropsonde data collected during the NASA ACTIVATE  
 525 field campaign. ACTIVATE featured the joint deployment of the HSRL-2 and AVAPS dropsondes during six deployments  
 from 2020 to 2022, enabling the accuracy of the HSRL-2 surface wind speed retrievals to be assessed using the coincident  
 dropsonde measurements. Comparisons of HSRL-2 and dropsonde surface wind speeds show correlations of 0.89, slopes of  
 1.04 and 1.17, y-intercepts of  $-0.13 \text{ m s}^{-1}$  and  $-1.05 \text{ m s}^{-1}$  for linear and bisector regressions, respectively. The accuracy of the

530 HSRL-2 retrievals, as denoted by mean error, is calculated to be  $0.15 \text{ m s}^{-1} \pm 1.80 \text{ m s}^{-1}$ . It is also observed that the dropsonde surface wind speed measurements most closely match with the Hu et al. (2008) wind speed – wave-slope variance model than the Cox and Munk (1954) and Wu (1990) models for surface wind speeds below  $7 \text{ m s}^{-1}$ , which is an important finding because most ACTIVATE surface wind speeds fall into this category. After this overview of model performance, the HSRL-2 retrievals for surface wind speeds separated into below  $7 \text{ m s}^{-1}$  and between  $7 \text{ m s}^{-1}$  and  $13.3 \text{ m s}^{-1}$  categories are then evaluated in more detail. For surface wind speeds below  $7 \text{ m s}^{-1}$ , correlations of 0.66, slopes of 0.65 and 0.99, and y-intercepts of  $1.10 \text{ m s}^{-1}$  and  $-0.49 \text{ m s}^{-1}$  are found and the accuracy of the retrievals is found to be  $-0.54 \text{ m s}^{-1} \pm 1.34 \text{ m s}^{-1}$ . Surface wind speeds between  $7 \text{ m s}^{-1}$  and  $13.3 \text{ m s}^{-1}$  show correlations of 0.75, slopes of 0.64 and 0.85, and y-intercepts of  $3.80 \text{ m s}^{-1}$  and  $1.87 \text{ m s}^{-1}$  and the retrieval accuracy is shown to be  $0.56 \text{ m s}^{-1} \pm 1.49 \text{ m s}^{-1}$ . Statistics are not reported for surface wind speeds above  $13.3 \text{ m s}^{-1}$  because there are too few points in this category to make meaningful comparisons. These results showcase an important observation not seen in the cumulative results, which is that the HSRL-2 estimates surface wind speeds with a bias of  $\pm \sim 0.5$   $\text{m s}^{-1}$  and an error of  $\pm \sim 1.5 \text{ m s}^{-1}$ . Lastly, the data are divided into winter and summer deployments (dates denoted in Sect. 2.1) to assess how the HSRL-2 performs between seasons. The winter surface wind speed data comparisons show correlations of 0.88, slopes of 0.95 and 1.08, and y-intercepts of  $1.03 \text{ m s}^{-1}$  and  $-0.08 \text{ m s}^{-1}$  and the summer data show correlations of 0.87, slopes of 1.08 and 1.24, and y-intercepts of  $-0.69 \text{ m s}^{-1}$  and  $-1.68 \text{ m s}^{-1}$  (linear and bisector regressions, respectively). The accuracy of the lidar retrievals is reported as  $0.63 \text{ m s}^{-1} \pm 2.07 \text{ m s}^{-1}$  and  $-0.18 \text{ m s}^{-1} \pm 1.52 \text{ m s}^{-1}$  for winter and summer, respectively. These findings show that HSRL-2 retrievals are more accurate in the summer than in winter, but still provide substantial (N = 236) and accurate surface wind speed data in winter as well.

This retrieval method offers a new path forward in airborne field work for the acquisition of surface wind speed data at a high spatial ( $\sim 75 \text{ m}$  along track) and time (0.5 s) resolution, as demonstrated with two case study flights (Research Flight 29 on 28 August 2020 and Research Flight 14 on 1 March 2020). The high horizontal spatial resolution of the HSRL-2 allows it to probe the fine-scale variability of surface wind speeds over time. As a result, the instrument provides near-continuous profiles of surface wind speeds over time that correspond to MABL phenomena such as SST dynamics and cloud evolution. Another important conclusion about the HSRL-2 surface retrievals is that the instrument can detect the surface in broken cloud scenes and are not limited to aerosol-free conditions like in Hu et al. (2008). Overall, having such data can benefit model assimilation efforts and consequently several scientific applications related to air-sea interactions such as estimating heat fluxes, gas exchange, sea salt emissions and aerosol transport, and cloud life cycle.

Forthcoming work will continue assessments of surface wind speed measurements during ACTIVATE by comparing dropsonde data to in situ measurements taken by the Turbulent Air Motion Measurement System (TAMMS) onboard the Falcon aircraft at its various altitude flight legs (between 120 m and 5 km) (Thornhill et al., 2003). Additional work is also warranted to assess the surface wind speed retrievals performed by ACTIVATE's other remote sensor, the Research Scanning Polarimeter (RSP), to fully demonstrate ACTIVATE's remote sensing capabilities.



## Data Availability

ACTIVATE airborne data are available through <https://asdc.larc.nasa.gov/project/ACTIVATE> (ACTIVATE Science Team, 2020). MERRA-2 mean sea surface temperature data are taken from the 2d, 1-Hourly, Time-Averaged, Single-Level, Assimilation, Surface Flux Diagnostics V5.12.4 (M2T1NXFLX) product found at [https://disc.gsfc.nasa.gov/datasets/M2T1NXFLX\\_5.12.4/summary](https://disc.gsfc.nasa.gov/datasets/M2T1NXFLX_5.12.4/summary) (doi.org/10.5067/7MCPBJ41Y0K6). GOES-16 data are from [https://asdc.larc.nasa.gov/ACTIVATE/ACTIVATE-Satellite\\_1](https://asdc.larc.nasa.gov/ACTIVATE/ACTIVATE-Satellite_1) (doi: 10.5067/ASDC/SUBORBITAL/ACTIVATE-Satellite\_1).

## 570 Author Contribution

SD performed all analyses with input from all co-authors. SD, JH, RF, CH, JR, and AS prepared manuscript with all co-authors involved in review/editing. TS, DH, SS, and CR conducted flight scientist duties on the King Air and helped with preparation and deployment of dropsondes. JH, RF, MF, CH, BC, DH, SS, TS, and EC were responsible for the HSRL-2 instrumentation and KT, CR, and HV were responsible for the NCAR AVAPS dropsonde instrumentation collected and subsequent archival of surface wind speed data sets needed to conduct this analysis. JH, JR, YH, RF, CH, and BC all contributed with formulation of HSRL-2 retrieval algorithm.

## Competing Interests

The authors declare that they have no conflict of interest.

## Disclaimer

580 Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

## Acknowledgements

The work was funded by ACTIVATE, a NASA Earth Venture Suborbital-3 (EVS-3) investigation funded by NASA's Earth Science Division and managed through the Earth System Science Pathfinder Program Office. We thank pilots and aircraft maintenance personnel of NASA Langley Research Services Directorate for successfully conducting ACTIVATE flights. The MERRA-2 1-Hourly, Time-Averaged, Single-Level, Assimilation, Surface Flux Diagnostics V5.12.4 (M2T1NXFLX) data used in this effort were acquired as part of the activities of NASA's Science Mission Directorate and are archived and distributed by the Goddard Earth Sciences (GES) Data and Information Services Center (DISC).

## Financial support

590 University of Arizona investigators were funded by NASA grant no. 80NSSC19K0442.

## References

ACTIVATE Science Team: Aerosol Cloud meteorology Interactions over the western Atlantic Experiment Data [dataset], doi.org/10.5067/SUBORBITAL/ACTIVATE/DATA001, 2020.

595 Aldhaif, A. M., Lopez, D. H., Dadashazar, H., and Sorooshian, A.: Sources, frequency, and chemical nature of dust events impacting the United States East Coast, Atmospheric Environment, 231, 117456, <https://doi.org/10.1016/j.atmosenv.2020.117456>, 2020.

600 Bedka, K. M., Nehrir, A. R., Kavaya, M., Barton-Grimley, R., Beaubien, M., Carroll, B., Collins, J., Cooney, J., Emmitt, G. D., Greco, S., Kooi, S., Lee, T., Liu, Z., Rodier, S., and Skofronick-Jackson, G.: Airborne lidar observations of wind, water vapor, and aerosol profiles during the NASA Aeolus calibration and validation (Cal/Val) test flight campaign, Atmos. Meas. Tech., 14, 4305-4334, 10.5194/amt-14-4305-2021, 2021.

605 Bellouin, N., Quaas, J., Gryspeerdt, E., Kinne, S., Stier, P., Watson-Parris, D., Boucher, O., Carslaw, K. S., Christensen, M., Daniau, A.-L., Dufresne, J.-L., Feingold, G., Fiedler, S., Forster, P., Gettelman, A., Haywood, J. M., Lohmann, U., Malavelle, F., Mauritsen, T., McCoy, D. T., Myhre, G., Mülmenstädt, J., Neubauer, D., Possner, A., Rugenstein, M., Sato, Y., Schulz, M., Schwartz, S. E., Sourdeval, O., Storelvmo, T., Toll, V., Winker, D., and Stevens, B.: Bounding Global Aerosol Radiative Forcing of Climate Change, Reviews of Geophysics, 58, e2019RG000660, <https://doi.org/10.1029/2019RG000660>, 2020.

610 Brunke, M. A., Cutler, L., Urzua, R. D., Corral, A. F., Crosbie, E., Hair, J., Hostetler, C., Kirschler, S., Larson, V., Li, X.-Y., Ma, P.-L., Minke, A., Moore, R., Robinson, C. E., Scarino, A. J., Schlosser, J., Shook, M., Sorooshian, A., Lee Thornhill, K., Voigt, C., Wan, H., Wang, H., Winstead, E., Zeng, X., Zhang, S., and Ziemba, L. D.: Aircraft Observations of Turbulence in Cloudy and Cloud-Free Boundary Layers Over the Western North Atlantic Ocean From ACTIVATE and Implications for the Earth System Model Evaluation and Development, Journal of Geophysical Research: Atmospheres, 127, e2022JD036480, <https://doi.org/10.1029/2022JD036480>, 2022.

Burton, S. P., Ferrare, R. A., Hostetler, C. A., Hair, J. W., Rogers, R. R., Obland, M. D., Butler, C. F., Cook, A. L., Harper, D. B., and Froyd, K. D.: Aerosol classification using airborne High Spectral Resolution Lidar measurements – methodology and examples, Atmos. Meas. Tech., 5, 73-98, 10.5194/amt-5-73-2012, 2012.

- 615 Burton, S. P., Hostetler, C. A., Cook, A. L., Hair, J. W., Seaman, S. T., Scola, S., Harper, D. B., Smith, J. A., Fenn, M. A., Ferrare, R. A., Saide, P. E., Chemyakin, E. V., and Müller, D.: Calibration of a high spectral resolution lidar using a Michelson interferometer, with data examples from ORACLES, *Appl. Opt.*, 57, 6061-6075, 10.1364/AO.57.006061, 2018.
- Carvalho, D.: An Assessment of NASA's GMAO MERRA-2 Reanalysis Surface Winds, *Journal of Climate*, 32, 8261-8281, <https://doi.org/10.1175/JCLI-D-19-0199.1>, 2019.
- 620 Chemyakin, E., Stamnes, S., Hair, J., Burton, S. P., Bell, A., Hostetler, C., Ferrare, R., Chowdhary, J., Moore, R., Ziemba, L., Crosbie, E., Robinson, C., Shook, M., Thornhill, L., Winstead, E., Hu, Y., van Diedenhoven, B., and Cairns, B.: Efficient single-scattering look-up table for lidar and polarimeter water cloud studies, *Opt. Lett.*, 48, 13-16, 10.1364/OL.474282, 2023.
- Chen, J., Wang, H., Li, X., Painemal, D., Sorooshian, A., Thornhill, K. L., Robinson, C., and Shingler, T.: Impact of Meteorological Factors on the Mesoscale Morphology of Cloud Streets during a Cold-Air Outbreak over the Western North Atlantic, *Journal of the Atmospheric Sciences*, 79, 2863-2879, <https://doi.org/10.1175/JAS-D-22-0034.1>, 2022.
- 625 Christoudias, T., Pozzer, A., and Lelieveld, J.: Influence of the North Atlantic Oscillation on air pollution transport, *Atmos. Chem. Phys.*, 12, 869-877, 10.5194/acp-12-869-2012, 2012.
- Corral, A. F., Choi, Y., Crosbie, E., Dadashazar, H., DiGangi, J. P., Diskin, G. S., Fenn, M., Harper, D. B., Kirschler, S., Liu, H., Moore, R. H., Nowak, J. B., Scarino, A. J., Seaman, S., Shingler, T., Shook, M. A., Thornhill, K. L., Voigt, C., Zhang, B., Ziemba, L. D., and Sorooshian, A.: Cold Air Outbreaks Promote New Particle Formation Off the U.S. East Coast, *Geophysical Research Letters*, 49, e2021GL096073, <https://doi.org/10.1029/2021GL096073>, 2022.
- 630 Cox, C. and Munk, W.: Measurement of the Roughness of the Sea Surface from Photographs of the Sun's Glitter, *J. Opt. Soc. Am.*, 44, 838-850, 10.1364/JOSA.44.000838, 1954.
- Creilson, J. K., Fishman, J., and Wozniak, A. E.: Intercontinental transport of tropospheric ozone: a study of its seasonal variability across the North Atlantic utilizing tropospheric ozone residuals and its relationship to the North Atlantic Oscillation, *Atmos. Chem. Phys.*, 3, 2053-2066, 10.5194/acp-3-2053-2003, 2003.
- Dadashazar, H., Painemal, D., Alipanah, M., Brunke, M., Chellappan, S., Corral, A. F., Crosbie, E., Kirschler, S., Liu, H., Moore, R. H., Robinson, C., Scarino, A. J., Shook, M., Sinclair, K., Thornhill, K. L., Voigt, C., Wang, H., Winstead, E., Zeng, X., Ziemba, L., Zuidema, P., and Sorooshian, A.: Cloud drop number concentrations over the western North Atlantic Ocean: seasonal cycle, aerosol interrelationships, and other influential factors, *Atmos. Chem. Phys.*, 21, 10499-10526, 10.5194/acp-21-10499-2021, 2021.
- 640

- Dierssen, H. M.: Hyperspectral Measurements, Parameterizations, and Atmospheric Correction of Whitecaps and Foam From Visible to Shortwave Infrared for Ocean Color Remote Sensing, *Frontiers in Earth Science*, 7, 10.3389/feart.2019.00014, 2019.
- Ferrare, R., Hair, J., Hostetler, C., Shingler, T., Burton, S. P., Fenn, M., Clayton, M., Scarino, A. J., Harper, D., Seaman, S.,  
645 Cook, A., Crosbie, E., Winstead, E., Ziemba, L., Thornhill, L., Robinson, C., Moore, R., Vaughan, M., Sorooshian, A.,  
Schlosser, J. S., Liu, H., Zhang, B., Diskin, G., DiGangi, J., Nowak, J., Choi, Y., Zuidema, P., and Chellappan, S.: Airborne  
HSRL-2 measurements of elevated aerosol depolarization associated with non-spherical sea salt, *Frontiers in Remote Sensing*,  
4, 10.3389/frsen.2023.1143944, 2023.
- Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G.,  
650 Reichle, R., Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V., Conaty, A., da Silva, A. M., Gu, W., Kim,  
G.-K., Koster, R., Lucchesi, R., Merkova, D., Nielsen, J. E., Partyka, G., Pawson, S., Putman, W., Rienecker, M., Schubert, S.  
D., Sienkiewicz, M., and Zhao, B.: The Modern-Era Retrospective Analysis for Research and Applications, Version 2  
(MERRA-2), *Journal of Climate*, 30, 5419-5454, <https://doi.org/10.1175/JCLI-D-16-0758.1>, 2017.
- GMAO: MERRA-2 tavg1\_2d\_flux\_Nx: 2d, 1-Hourly, Time-Averaged, Single-Level, Assimilation, Surface Flux Diagnostics  
655 V5.12.4 [dataset], 2015.
- Gordon, H. R. and Wang, M.: Influence of oceanic whitecaps on atmospheric correction of ocean-color sensors, *Appl. Opt.*,  
33, 7754-7763, 10.1364/AO.33.007754, 1994.
- Hair, J. W., Hostetler, C. A., Cook, A. L., Harper, D. B., Ferrare, R. A., Mack, T. L., Welch, W., Izquierdo, L. R., and Hovis,  
F. E.: Airborne High Spectral Resolution Lidar for profiling aerosol optical properties, *Appl. Opt.*, 47, 6734-6752,  
660 10.1364/AO.47.006734, 2008.
- Hu, Y., Stamnes, K., Vaughan, M., Pelon, J., Weimer, C., Wu, D., Cisewski, M., Sun, W., Yang, P., Lin, B., Omar, A., Flittner,  
D., Hostetler, C., Treppe, C., Winker, D., Gibson, G., and Santa-Maria, M.: Sea surface wind speed estimation from space-  
based lidar measurements, *Atmos. Chem. Phys.*, 8, 3593-3601, 10.5194/acp-8-3593-2008, 2008.
- Josset, D., Pelon, J., and Hu, Y.: Multi-Instrument Calibration Method Based on a Multiwavelength Ocean Surface Model,  
665 *IEEE Geoscience and Remote Sensing Letters*, 7, 195-199, 10.1109/LGRS.2009.2030906, 2010a.
- Josset, D., Pelon, J., Protat, A., and Flamant, C.: New approach to determine aerosol optical depth from combined CALIPSO  
and CloudSat ocean surface echoes, *Geophysical Research Letters*, 35, <https://doi.org/10.1029/2008GL033442>, 2008.

- Josset, D., Zhai, P.-W., Hu, Y., Pelon, J., and Lucker, P. L.: Lidar equation for ocean surface and subsurface, *Opt. Express*, 18, 20862-20875, 10.1364/OE.18.020862, 2010b.
- 670 Kiliyanpilakkil, V. P. and Meskhidze, N.: Deriving the effect of wind speed on clean marine aerosol optical properties using the A-Train satellites, *Atmos. Chem. Phys.*, 11, 11401-11413, 10.5194/acp-11-11401-2011, 2011.
- Kirschler, S., Voigt, C., Anderson, B., Campos Braga, R., Chen, G., Corral, A. F., Crosbie, E., Dadashazar, H., Ferrare, R. A., Hahn, V., Hendricks, J., Kaufmann, S., Moore, R., Pöhlker, M. L., Robinson, C., Scarino, A. J., Schollmayer, D., Shook, M. A., Thornhill, K. L., Winstead, E., Ziemba, L. D., and Sorooshian, A.: Seasonal updraft speeds change cloud droplet number  
 675 concentrations in low-level clouds over the western North Atlantic, *Atmos. Chem. Phys.*, 22, 8299-8319, 10.5194/acp-22-8299-2022, 2022.
- Koepke, P.: Effective reflectance of oceanic whitecaps, *Appl. Opt.*, 23, 1816-1824, 10.1364/AO.23.001816, 1984.
- Labzovskii, L. D., van Zadelhoff, G. J., Tilstra, L. G., de Kloe, J., Donovan, D. P., and Stoffelen, A.: High sensitivity of Aeolus UV surface returns to surface reflectivity, *Scientific Reports*, 13, 17552, 10.1038/s41598-023-44525-5, 2023.
- 680 Lamb, P. J. and Pepler, R. A.: North Atlantic Oscillation: Concept and an Application, *Bulletin of the American Meteorological Society*, 68, 1218-1225, [https://doi.org/10.1175/1520-0477\(1987\)068<1218:NAOCAA>2.0.CO;2](https://doi.org/10.1175/1520-0477(1987)068<1218:NAOCAA>2.0.CO;2), 1987.
- Li, Q., Jacob, D. J., Bey, I., Palmer, P. I., Duncan, B. N., Field, B. D., Martin, R. V., Fiore, A. M., Yantosca, R. M., Parrish, D. D., Simmonds, P. G., and Oltmans, S. J.: Transatlantic transport of pollution and its effects on surface ozone in Europe and North America, *Journal of Geophysical Research: Atmospheres*, 107, ACH 4-1-ACH 4-21,  
 685 <https://doi.org/10.1029/2001JD001422>, 2002.
- Li, X.-Y., Wang, H., Chen, J., Endo, S., George, G., Cairns, B., Chellappan, S., Zeng, X., Kirschler, S., Voigt, C., Sorooshian, A., Crosbie, E., Chen, G., Ferrare, R. A., Gustafson, W. I., Hair, J. W., Kleb, M. M., Liu, H., Moore, R., Painemal, D., Robinson, C., Scarino, A. J., Shook, M., Shingler, T. J., Thornhill, K. L., Tornow, F., Xiao, H., Ziemba, L. D., and Zuidema, P.: Large-Eddy Simulations of Marine Boundary Layer Clouds Associated with Cold-Air Outbreaks during the ACTIVATE  
 690 Campaign. Part I: Case Setup and Sensitivities to Large-Scale Forcings, *Journal of the Atmospheric Sciences*, 79, 73-100, <https://doi.org/10.1175/JAS-D-21-0123.1>, 2022.
- Li, Z., Lemmerz, C., Paffrath, U., Reitebuch, O., and Witschas, B.: Airborne Doppler Lidar Investigation of Sea Surface Reflectance at a 355-nm Ultraviolet Wavelength, *Journal of Atmospheric and Oceanic Technology*, 27, 693-704, <https://doi.org/10.1175/2009JTECHA1302.1>, 2010.

- 695 Mardi, A. H., Dadashazar, H., Painemal, D., Shingler, T., Seaman, S. T., Fenn, M. A., Hostetler, C. A., and Sorooshian, A.: Biomass Burning Over the United States East Coast and Western North Atlantic Ocean: Implications for Clouds and Air Quality, *Journal of Geophysical Research: Atmospheres*, 126, e2021JD034916, <https://doi.org/10.1029/2021JD034916>, 2021.
- Martin, C. and Suhr, I.: NCAR/EOL Atmospheric Sounding Processing ENvironment (ASPEN) software. Version 3.4.5. [dataset], 2021.
- 700 Moore, K. D., Voss, K. J., and Gordon, H. R.: Spectral reflectance of whitecaps: Their contribution to water-leaving radiance, *Journal of Geophysical Research: Oceans*, 105, 6493-6499, <https://doi.org/10.1029/1999JC900334>, 2000.
- Murphy, A. and Hu, Y.: Retrieving Aerosol Optical Depth and High Spatial Resolution Ocean Surface Wind Speed From CALIPSO: A Neural Network Approach, *Frontiers in Remote Sensing*, 1, 10.3389/frsen.2020.614029, 2021.
- Nair, A. K. M. and Rajeev, K.: Multiyear CloudSat and CALIPSO Observations of the Dependence of Cloud Vertical Distribution on Sea Surface Temperature and Tropospheric Dynamics, *Journal of Climate*, 27, 672-683, <https://doi.org/10.1175/JCLI-D-13-00062.1>, 2014.
- 705 Neukermans, G., Harmel, T., Galí, M., Rudorff, N., Chowdhary, J., Dubovik, O., Hostetler, C., Hu, Y., Jamet, C., Knobelspiesse, K., Lehahn, Y., Litvinov, P., Sayer, A. M., Ward, B., Boss, E., Koren, I., and Miller, L. A.: Harnessing remote sensing to address critical science questions on ocean-atmosphere interactions, *Elementa: Science of the Anthropocene*, 6, 10.1525/elementa.331, 2018.
- 710 Painemal, D., Corral, A. F., Sorooshian, A., Brunke, M. A., Chellappan, S., Afzali Gorooh, V., Ham, S.-H., O'Neill, L., Smith Jr., W. L., Tselioudis, G., Wang, H., Zeng, X., and Zuidema, P.: An Overview of Atmospheric Features Over the Western North Atlantic Ocean and North American East Coast—Part 2: Circulation, Boundary Layer, and Clouds, *Journal of Geophysical Research: Atmospheres*, 126, e2020JD033423, <https://doi.org/10.1029/2020JD033423>, 2021.
- 715 Painemal, D., Chellappan, S., Smith Jr., W. L., Spangenberg, D., Park, J. M., Ackerman, A., Chen, J., Crosbie, E., Ferrare, R., Hair, J., Kirschler, S., Li, X.-Y., McComiskey, A., Moore, R. H., Sanchez, K., Sorooshian, A., Tornow, F., Voigt, C., Wang, H., Winstead, E., Zeng, X., Ziemba, L., and Zuidema, P.: Wintertime Synoptic Patterns of Midlatitude Boundary Layer Clouds Over the Western North Atlantic: Climatology and Insights From In Situ ACTIVATE Observations, *Journal of Geophysical Research: Atmospheres*, 128, e2022JD037725, <https://doi.org/10.1029/2022JD037725>, 2023.
- 720 Paiva, V., Kampel, M., and Camayo, R.: Comparison of Multiple Surface Ocean Wind Products with Buoy Data over Blue Amazon (Brazilian Continental Margin), *Advances in Meteorology*, 2021, 6680626, 10.1155/2021/6680626, 2021.

- Ricker, W. E.: Linear Regressions in Fishery Research, *Journal of the Fisheries Research Board of Canada*, 30, 409-434, 10.1139/f73-072, 1973.
- 725 Scarino, A. J., Obland, M. D., Fast, J. D., Burton, S. P., Ferrare, R. A., Hostetler, C. A., Berg, L. K., Lefer, B., Haman, C., Hair, J. W., Rogers, R. R., Butler, C., Cook, A. L., and Harper, D. B.: Comparison of mixed layer heights from airborne high spectral resolution lidar, ground-based measurements, and the WRF-Chem model during CalNex and CARES, *Atmos. Chem. Phys.*, 14, 5547-5560, 10.5194/acp-14-5547-2014, 2014.
- 730 Schlosser, J. S., Stamnes, S., Burton, S. P., Cairns, B., Crosbie, E., Van Dierenhoven, B., Diskin, G., Dmitrovic, S., Ferrare, R., Hair, J. W., Hostetler, C. A., Hu, Y., Liu, X., Moore, R. H., Shingler, T., Shook, M. A., Thornhill, K. L., Winstead, E., Ziemba, L., and Sorooshian, A.: Polarimeter + Lidar-Derived Aerosol Particle Number Concentration, *Frontiers in Remote Sensing*, 3, 10.3389/frsen.2022.885332, 2022.
- Schulien, J. A., Behrenfeld, M. J., Hair, J. W., Hostetler, C. A., and Twardowski, M. S.: Vertically- resolved phytoplankton carbon and net primary production from a high spectral resolution lidar, *Opt. Express*, 25, 13577-13587, 10.1364/OE.25.013577, 2017.
- 735 Seethala, C., Zuidema, P., Edson, J., Brunke, M., Chen, G., Li, X.-Y., Painemal, D., Robinson, C., Shingler, T., Shook, M., Sorooshian, A., Thornhill, L., Tornow, F., Wang, H., Zeng, X., and Ziemba, L.: On Assessing ERA5 and MERRA2 Representations of Cold-Air Outbreaks Across the Gulf Stream, *Geophysical Research Letters*, 48, e2021GL094364, <https://doi.org/10.1029/2021GL094364>, 2021.
- 740 Sorooshian, A., Corral, A. F., Braun, R. A., Cairns, B., Crosbie, E., Ferrare, R., Hair, J., Kleb, M. M., Hossein Mardi, A., Maring, H., McComiskey, A., Moore, R., Painemal, D., Scarino, A. J., Schlosser, J., Shingler, T., Shook, M., Wang, H., Zeng, X., Ziemba, L., and Zuidema, P.: Atmospheric Research Over the Western North Atlantic Ocean Region and North American East Coast: A Review of Past Work and Challenges Ahead, *Journal of Geophysical Research: Atmospheres*, 125, e2019JD031626, <https://doi.org/10.1029/2019JD031626>, 2020.
- 745 Sorooshian, A., Anderson, B., Bauer, S. E., Braun, R. A., Cairns, B., Crosbie, E., Dadashazar, H., Diskin, G., Ferrare, R., Flagan, R. C., Hair, J., Hostetler, C., Jonsson, H. H., Kleb, M. M., Liu, H., MacDonald, A. B., McComiskey, A., Moore, R., Painemal, D., Russell, L. M., Seinfeld, J. H., Shook, M., Smith, W. L., Thornhill, K., Tselioudis, G., Wang, H., Zeng, X., Zhang, B., Ziemba, L., and Zuidema, P.: Aerosol-Cloud-Meteorology Interaction Airborne Field Investigations: Using Lessons Learned from the U.S. West Coast in the Design of ACTIVATE off the U.S. East Coast, *Bulletin of the American Meteorological Society*, 100, 1511-1528, <https://doi.org/10.1175/BAMS-D-18-0100.1>, 2019.

- 750 Sorooshian, A., Alexandrov, M. D., Bell, A. D., Bennett, R., Betito, G., Burton, S. P., Buzanowicz, M. E., Cairns, B., Chemyakin, E. V., Chen, G., Choi, Y., Collister, B. L., Cook, A. L., Corral, A. F., Crosbie, E. C., van Diedenhoven, B., DiGangi, J. P., Diskin, G. S., Dmitrovic, S., Edwards, E. L., Fenn, M. A., Ferrare, R. A., van Gilst, D., Hair, J. W., Harper, D. B., Hilario, M. R. A., Hostetler, C. A., Jester, N., Jones, M., Kirschler, S., Kleb, M. M., Kusterer, J. M., Leavor, S., Lee, J. W., Liu, H., McCauley, K., Moore, R. H., Nied, J., Notari, A., Nowak, J. B., Painemal, D., Phillips, K. E., Robinson, C. E., Scarino, A. J., Schlosser, J. S., Seaman, S. T., Seethala, C., Shingler, T. J., Shook, M. A., Sinclair, K. A., Smith Jr, W. L., Spangenberg, D. A., Stammes, S. A., Thornhill, K. L., Voigt, C., Vömel, H., Wasilewski, A. P., Wang, H., Winstead, E. L., Zeider, K., Zeng, X., Zhang, B., Ziemba, L. D., and Zuidema, P.: Spatially-coordinated airborne data and complementary products for aerosol, gas, cloud, and meteorological studies: The NASA ACTIVATE dataset, *Earth Syst. Sci. Data Discuss.*, 2023, 1-79, 10.5194/essd-2023-109, 2023.
- 755 Sun, K., Dai, G., Wu, S., Reitebuch, O., Baars, H., Liu, J., and Zhang, S.: Correlation between marine aerosol optical properties and wind fields over remote oceans with use of spaceborne lidar observations, *EGUsphere*, 2023, 1-34, 10.5194/egusphere-2023-433, 2023.
- Thornhill, K. L., Anderson, B. E., Barrick, J. D. W., Bagwell, D. R., Friesen, R., and Lenschow, D. H.: Air motion intercomparison flights during Transport and Chemical Evolution in the Pacific (TRACE-P)/ACE-ASIA, *Journal of Geophysical Research: Atmospheres*, 108, <https://doi.org/10.1029/2002JD003108>, 2003.
- 765 Tornow, F., Ackerman, A. S., Fridlind, A. M., Cairns, B., Crosbie, E. C., Kirschler, S., Moore, R. H., Painemal, D., Robinson, C. E., Seethala, C., Shook, M. A., Voigt, C., Winstead, E. L., Ziemba, L. D., Zuidema, P., and Sorooshian, A.: Dilution of Boundary Layer Cloud Condensation Nucleus Concentrations by Free Tropospheric Entrainment During Marine Cold Air Outbreaks, *Geophysical Research Letters*, 49, e2022GL098444, <https://doi.org/10.1029/2022GL098444>, 2022.
- 770 van Diedenhoven, B., Hasekamp, O. P., Cairns, B., Schuster, G. L., Stammes, S., Shook, M. A., and Ziemba, L. D.: Remote sensing of aerosol water fraction, dry size distribution and soluble fraction using multi-angle, multi-spectral polarimetry, *EGUsphere*, 2022, 1-41, 10.5194/egusphere-2022-670, 2022.
- Venkata, S. L. and Reagan, J. A.: Aerosol Retrievals from CALIPSO Lidar Ocean Surface Returns, *Remote Sensing*, 8, 1006, 2016.
- 775 Vömel, H. and Dunion, J.: Chapter 10 - Aircraft dropsonde campaigns, in: *Field Measurements for Passive Environmental Remote Sensing*, edited by: Nalli, N. R., Elsevier, 185-194, <https://doi.org/10.1016/B978-0-12-823953-7.00021-6>, 2023.
- Vömel, H., Goodstein, M., Tudor, L., Witte, J., Fuchs-Stone, Ž., Sentić, S., Raymond, D., Martinez-Claros, J., Juračić, A., Maithel, V., and Whitaker, J. W.: High-resolution in situ observations of atmospheric thermodynamics using dropsondes



- during the Organization of Tropical East Pacific Convection (OTREC) field campaign, *Earth Syst. Sci. Data*, 13, 1107-1117, 780 10.5194/essd-13-1107-2021, 2021.
- Whitlock, C. H., Bartlett, D. S., and Gurganus, E. A.: Sea foam reflectance and influence on optimum wavelength for remote sensing of ocean aerosols, *Geophysical Research Letters*, 9, 719-722, 1982.
- Wu, C. and Yu, J. Z.: Evaluation of linear regression techniques for atmospheric applications: the importance of appropriate weighting, *Atmos. Meas. Tech.*, 11, 1233-1250, 10.5194/amt-11-1233-2018, 2018.
- 785 Wu, J.: Mean square slopes of the wind-disturbed water surface, their magnitude, directionality, and composition, *Radio Science*, 25, 37-48, <https://doi.org/10.1029/RS025i001p00037>, 1990.