

Remotely Sensed and Surface Measurement Derived Mass-Conserving Inversion of Daily High-Resolution NO_x Emissions and Inferred Combustion Technologies in Energy Rich Northern China

Xiaolu Li^{1,2}, Jason Blake Cohen^{2*}, Kai Qin², Hong Geng¹, Liling Wu⁴, Xiaohui Wu³, Chengli Yang³, Rui Zhang⁵, Liqin Zhang⁶

¹Institute of Environmental Science, Shanxi University, Taiyuan, 030006, China

²School of Environment and Spatial Informatics, China University of Mining and Technology, Xuzhou, 221116, China

³Shanxi Dadi Eco-environmental Technology Research Institute Co., Taiyuan, 030000, China

10 ⁴School of Environment, Tsinghua University, Beijing, 10084, China

⁵Shanxi Institute of Eco-environmental Planning and Technology, Taiyuan, 030002, China

⁶Shanxi Institute of Eco-environmental Monitoring and Emergency Response Center, Taiyuan, 030027, China

Correspondence to: Jason Blake Cohen (jasonbc@alum.mit.edu)

15 **Abstract.** This work presents a new model free inversion estimation framework using daily TROPOMI NO₂ columns and observed fluxes from the continuous emissions monitoring systems (CEMS) to quantify three years of daily-scale emissions of NO_x at 0.05° × 0.05° over Shanxi Province, a major world-wide energy producing and consuming region. The NO_x emissions, day-to-day variability, and uncertainty on a climatological basis are computed to be 1.86, 1.02, and 1.04 Tg per year respectively. The highest emissions are concentrated in the lower Fen River valley, which accounts for 25% of the area, 53% of the NO_x emissions, and 72% of CEMS sources. Two major forcing factors (10th to 90th percentile) are horizontal transport distance per day (63–508 km) and lifetime of NO_x (7.1–18.1 h). Both of these values are consistent with NO_x emissions to both the surface layer and the free troposphere. The third forcing factor, the ratio of NO_x/NO₂, on a pixel-by-pixel basis is demonstrated to correlate with the combustion temperature and energy efficiency of large energy consuming sources. Specifically, thermal power plants, cement, and iron and steel companies have a relatively high NO_x/NO₂ ratio, while coking, industrial boilers, and aluminium oxide show relatively low ratio. Variance maximization is applied to daily TROPOMI NO₂ columns, which facilitates identification of three orthogonal and statistically significant modes of variability, and successfully attributes them both spatially and temporally to (a) this work's computed emissions, (b) remotely sensed TROPOMI UVAI, and (c) computed transport based on TROPOMI NO₂.

30

删除了: 83

删除了: 01

删除了: 06

删除了: 52

删除了: 6

删除了: 666

删除了: 6.7

删除了: 4

删除了:

删除了: have a




删除了: significant

删除了: ion

删除了: NO₂ columns identifies


删除了:

Graphical abstract.



删除了:

带格式的: 居中



删除了:

1 Introduction

Economic growth is based in part on the emissions of air pollution, leading to serious environmental and health consequences.

55 To address and alleviate the air quality problems, the Chinese government has implemented new air pollution controls, with the aim of producing higher-quality development. Two recent examples are the Air Pollution Prevention and Control Action Plan from 2013 to 2017 and the Three-Year Action Plan for Winning the Battle in Defense of Blue Sky from 2018 to 2020 (Zhang et al., 2019; Geng et al., 2019; Jiang et al., 2021; Wang et al., 2020b; Li et al., 2022a; Wei et al., 2023), which have led to a significant reduction in annual average concentration of particulate matter (PM), sulfur dioxide (SO₂) and carbon monoxide (CO) in Shanxi Province. Shanxi is selected for this study, with geographical location and topography showed in Fig. 1, as it is a highly energy rich location that produces nearly 30% of all of China's coal, as well as having substantial industry that consumes a significant amount of coal for both local energy production, steel, cement, coke, and aluminum production, and export, among other economic activities (Li et al., 2022). Moreover, there have also been minor increases in the observed annual average concentrations of both ozone (O₃) and nitrogen dioxide (NO₂) in Shanxi between 2015 and 2020 (DEESP, 2015; DEESP, 2020). Furthermore, due to its relatively dry climate, high elevation, and mountainous geography, it has complex underlying natural factors that impact its atmospheric environment.

60 The sum of NO₂ and nitric oxide (NO) is frequently grouped as nitrogen oxides (herein termed NO_x), which is an important trace gas impacting of the earth's atmosphere because it is a strong marker of anthropogenic combustion-related pollution, a precursor to ozone (Jacob et al., 1993), secondary aerosol (Rollins et al., 2012), and acid rain (Singh and Agrawal, 2007). In order to gain a better understanding of NO_x and its impacts, precise and quantitative emissions inventories are crucial information for policy makers, air quality modelers, climate change modelers, and those who conduct pollution weather response interactions, among others (Hoesly et al., 2018; Crippa et al., 2018; Li et al., 2017a; Xing et al., 2013; McDonald et al., 2013). However, it is challenging to quantify emissions in rapidly developing and changing areas accurately as there are a variety of contributing sources, a complex underlying mixture of combustion technologies, industrial restructuring, changing population dynamics, and ongoing atmospheric environmental management, all of which may lead to substantial changes in pollution sources.

70 Presently, most emission inventories are compiled from statistics, representing emitting activities and associated typical emission factors, herein called "bottom-up" approaches (Ohara et al., 2007; Zheng et al., 2018; Li et al., 2017b). Bottom-up methods provide emissions data at finer scales. However, higher resolution emission inventories using bottom-up methods require rich, detailed, and extremely precise records of energy use, facility locations, and other socioeconomic datasets from multiple regional and temporal scales (Cai et al., 2018), which frequently have a considerable amount of uncertainty (Bond et al., 2007). With the ever-increasing tightening of environmental management, emission factors have changed significantly and will continue to do so in the future. On-site surveys for bottom-up methods are time consuming and resource demanding. When performing emission factor determination in laboratory, it is important to note that the differences between small field studies and controlled laboratory combustion experiments and real-world examples also are quite significant, with super-emitters

删除了: has always been accompanied by air pollution, with

删除了: associated with non-green economic growth are more serious...

删除了: severe

删除了: been

删除了: ing

删除了: more than 25

删除了:

删除了: steel, cement, coke, and aluminum production,

删除了: However

删除了: (2015, 2020)

删除了: also impacting

删除了: Nitrogen Monoxide

删除了: (Beirle et al., 2011)

删除了: (Rollins et al., 2012)

删除了: on

known to create large differences when using insufficiently large datasets (Zavala et al., 2006) and missing large sources leading to significant error (Wang et al., 2021). Due to the low temporal resolution and time-lag associated with many of these datasets being available, bottom-up inventories are not easy to keep up with rapid changes in industrial, economic, and pandemic, and therefore are not very good at tracking atmospheric emissions under actual existing environmental conditions, limiting their use (Mijling and Van Der A, 2012).

To overcome the disadvantages identified above, while simultaneously improving the spatial and temporal resolution of emission inventories, attempts at top-down emissions inventories using remote sensed dataset have been made by the community. Some of these attempts have focused on applications to long-lived gasses (CH₄, CFCs, and N₂O), since their

chemical decay is slow compared with their transport processes, allowing a simpler set of approximations to perform the inversion (Chen and Prinn, 2006; Tu et al., 2022b; Liu et al., 2021). Satellite observations have also been widely used to quantify short-lived species such as NO_x emissions (Martin, 2003; Beirle et al., 2019; Goldberg et al., 2019; Qu et al., 2019) by providing up-to-date and continuous time series of NO₂ columns in different regions. Gaussian plumes have been used for a long time (Green et al., 1980), with more advanced but similar approximations including the exponentially modified gaussian

model to quantify the NO_x emissions of isolated megacity sources (Beirle et al., 2011) and the probabilistic collocation method to train the emissions flux enhancement of megacities as a function of their size and shape (Cohen and Prinn, 2011). Some methods used a partial estimation of the mass balance approaches, including Beirle et al. (2019) and Kong et al. (2019). Another category based on atmospheric chemical transport or climate models and data assimilation, such as 3D-Var, 4D-Var, and Kalman filters work very well but are susceptible to underlying model and scientific uncertainty, as well as being extremely

computationally intensive (Cohen and Wang, 2014; Zhang et al., 2021). The selection of a priori inventories is crucial when using the methods above, since it has been demonstrated that missing sources can frequently not be compensated for by merely scaling or increasing other existing sources if their spatial and temporal distributions are not matched correctly (Cohen, 2014).

In recent years, continuous emissions monitoring systems (CEMS) have been introduced in China, the USA, and other countries and areas, as a means to detect in an integrated manner at the emissions effluent source, on an hourly and/or daily time scale, and contains well-known aspects of quality control and assurance. This platform provides reliable technical information to fundamentally quantify local emissions on a stack-by-stack basis, in order to improve both the temporal resolution and magnitude of emission inventories (Tang et al., 2019; Gu et al., 2022; Chen et al., 2019; Lange et al., 2022).

This study takes advantage of the respective strengths of top-down and bottom-up emissions estimation by applying a new, fast, first-order approximation of physical, chemical, and thermodynamics controlling the distribution of NO_x in-situ, and constrains these approximations using daily measurements of remotely sensed NO₂ from the Tropospheric Monitoring Instrument (TROPOMI) using a mass conserving inversion to estimate the daily NO_x emissions on a mesoscale grid at (0.05° × 0.05°) from January 2019 through December 2021. This approach is partly originated from similar box-modeling ideas in previous studies (Rigby et al., 2008; Beirle et al., 2019; Kong et al., 2019), which themselves are based on previous theory underlying the development of mass-conserving box models (Seigneur et al., 1986). In this specific work, the mass of emissions is connected with the in-situ observed column loadings through application of the following factors: the temporal rate of change

删除了: very

删除了: Kong et al. (Kong et al., 2019) and

删除了: (Beirle et al., 2019), which have assumed that some of the mass-based-terms are negligible, or some of the factors underlying the terms are very narrowly constrained

删除了: costly to run

删除了: by increasing

删除了:

删除了: if the

删除了: At the present time, some of the most accepted emissions inventories include the Multi-resolution Emission Inventory for China (MEIC) and the Emission Database for Global Atmospheric Research (EDGAR) (Crippa et al., 2018).

删除了: locations

删除了: in

in column loadings, first order chemical loss of NO_x, gradient transport of NO_x, and gradient transport of atmospheric air mass. The coefficients weighting these terms are flexibly fitted, allowing a wider range of possible driving forces and solutions to be considered, while still requiring that these parameters are consistent with observations (Rollins et al., 2012; Karl et al., 2023). The fitted relationship is formed without the use of complex models, can be run on a normal desktop computer, and the end product can be flexibly modified by the user for their own various applications. The a priori emissions used in this work come from daily CEMS observations at powerplants and other large sources, as well as the Multi-resolution Emission Inventory for China (MEIC) over Shanxi province. This unique perspective is capable of using measured emissions, fitting variable parameters that were fixed in previous approaches, and inverting emissions as a function of their month-to-month constrained driving forces, under different but realistic environmental conditions. This method has been used in different situation such as over different months, over multi-year changes in the environment, under different actinic flux and atmospheric oxidation conditions, under complex meteorological domains, and over sources which are both thermodynamically stable as well as unstable. That permits this study to explore the full range of variations. Additionally, this approach allows for results with both a robust error quantification, and emissions that compare well with both the mean and the measured spatial and temporal variation in the underlying remotely sensed NO₂ columns.




Figure 1: Location, topography and administrative division of Shanxi Province.

- 删除了: The equation
- 删除了: allow the end-user to
- 删除了: y
- 删除了: the equation
- 删除了: (Rigby et al., 2008)
- 删除了: This net combination of factors does not rely on complex models, and allows a flexible approximation which can be modified by the user for various applications.
- 删除了:
- 删除了: have used both
- 删除了: -resolution
- 删除了: and
- 删除了: MEIC
- 删除了: of
- 删除了: The fact that three years of driving data have been used over different months of the year, over multi-year changes in the environment, under high UV and low UV conditions, under complex meteorological domains, over sources which are both thermodynamically stable as well as unstable, and even through COVID-19, permits this study to fully explore the full range of variations observed.
- 删除了: ,
- 删除了: s

155

160

165

190 2 Materials and Methods

2.1 Tropospheric Vertical Column Measurements from TROPOMI

TROPOMI measures reflected solar radiation in the UV, visible, and Near IR bands following a sun-synchronous, low-earth orbit with an equator overpass time of approximately 13:30 LT, allowing daily-scale measurements across the globe (Veefkind et al., 2012; Goldberg et al., 2019; Tu et al., 2022a). Starting from August 2019, the spatial resolution of TROPOMI has been refined to 5.5 km×3.5 km (Lange et al., 2022). This study uses two distinct products measured by TROPOMI over different radiative bands, but at the same place and time: NO₂ and Ultraviolet Aerosol Index (UVAI).

Daily level-2 version 2.3.1 tropospheric NO₂ columns and version 2.2.0 UVAI over Shanxi Province has been introduced. All available days and swaths corresponding to the time period from January 2019 through December 2021 are analyzed (<https://disc.gsfc.nasa.gov/datasets>). Overlapping NO₂ and UVAI column pixels in each swath are resampled to a common latitude-longitude grid at 0.05 °×0.05 ° using weighted polygons (<http://stcorp.github.io/harp/doc/html/index.html>). Before use, it is required that all TROPOMI data is quality assured, specifically insisting that each pixel has a “qa_value” greater than 0.75, that the “cloud radiance fraction” is smaller than 0.5, and that scenes covered by snow/ice, errors and similar problematic retrievals are removed (Henk et al., 2021). Furthermore, an additional filter is applied to set all individual grid of NO₂ column which is less than 1.4×10^{15} molec cm⁻² to be NaN. This is done to avoid issues where the observed signal may be smaller than the uncertainty of the signal itself (J.H.G.M. Van Geffen, 2021; Qin et al., 2022). This combination of assumptions ensures that the data used should be of the highest possible precision based on the current available technology.

The average loading, daily variation, and number of days without TROPOMI NO₂ columns from 2019 through 2021 are shown in Fig. 2. The number of invalid days varies pixel-by-pixel from 357 to 742 d (521 d on average), with higher altitude and mountainous areas tending to be more. The higher values on the maps are consistent with known urban and industrial regions, such as Taiyuan Basin, Xinding Basin, Linfen Basin, and Yangquan City. Areas with a high variation and relatively low mean value are observed in regions where new economic development zones have been recently created or are in the process of being actively developed, including urban areas of Datong and Xinzhou. Areas with a relatively high variation and high mean value are indicative of high urbanization and developed industrial areas, corresponding with the Taiyuan Basin, and southern Yangquan. Areas with a high average value and a low variation correspond with areas which have a fewer number of temporally consistent emissions sources, as is observed in parts of the Linfen Basin, central Changzhi, Lvliang and Jincheng City, and industrial parks in Jiexiu district of Jinzhong City.

删除了: UVAI

删除了: 1

删除了: 31

删除了: (Henk Eskes, 2021)

删除了: Furthermore, in the case of NO₂, an additional filter is applied to avoid issues where the signal is possibly smaller than the uncertainty about 30% to 50% (J.H.G.M. Van Geffen, 2021; Qin et al., 2022), leading to all grids with a column loading smaller than 1.4×10^{15} molec cm⁻² being discarded.

删除了:

删除了: The TROPOMI NO₂ columns used in this study can portray the spatial and temporal distribution of sources in a high amount of detail, including being able to effectively identify spatial hotspots (Griffin et al., 2019)




Figure 2: TROPOMI daily NO₂ column loadings from 2019 through 2021: (a) mean value (unit: molec cm⁻²), (b) day by day standard deviation (unit: molec cm⁻²), and (c) the number of invalid days (unit: d).

2.2 A Prior Emissions inventories

235 2.2.1 CEMS

CEMS was introduced by the Ministry of Environmental Protection of China in 2007 to monitor and manage the emissions of certain (mainly high-emitting) plants (Schreifels et al., 2012; Karplus et al., 2018). CEMS makes actual stack flue gas measurements of the concentration of PM, SO₂ and NO_x as emitted from power plants, steel and iron plants, aluminum smelters, coke plants, coal-fired boilers and others, all in real-time (Tang et al., 2020; Zhang and Schreifels, 2011). Statistics of the emissions sites monitored in Shanxi are given in Table 1. There are two different technologies for measuring NO_x concentrations. One converts NO₂ to NO and measures the total NO concentration, the other measures NO₂ and NO separately. Both of the measured results have been converted to NO₂ mass concentration.

In this work, all available CEMS monitors of daily-scale emissions from 2019 to 2021 were obtained from the Department of Ecology and Environment of Shanxi Province (DEESP), with the government making great effort to regulate the CEMS network and to ensure the reliability of CEMS data (Tang et al., 2020). Preprocessing of these data include using google earth to correct the location of the factories, conducting quality control on measured concentrations according to the CEMS technical requirements, including eliminating negative values, outlier values, and null values. The overall percentage of abnormal values

删除了: discharged

删除了: coal

删除了: measuring

删除了: s

is found to account for 0.63%, 1.18%, and 1.55% of the raw data respectively for 2019, 2020 and 2021. The formula used to calculate NO_x emissions is given in Eq. (1):

$$E_d = \bar{C}_h \times \bar{Q}_h \times 24 \quad (1)$$

255 where \bar{C}_h is the daily average of hourly NO_x concentration, mg m⁻³; \bar{Q}_h is the daily average of hourly wet flue gas flow under actual working condition, m³ h⁻¹, and 24 is ~~used to convert units from hours to days~~. Following the “Specifications and test procedures for a continuous emission monitoring system for SO₂, NO_x, and particulate matter in flue gas emitted from stationary sources (HJ/T76-2017)”, the uncertainty of NO_x concentration (C_h) is less than or equal to 30% for the data used in this study. After quality control, the emission intensity on a grid-by-grid basis is found to be 0.64±0.08 μg m⁻² s⁻¹, 0.45±0.13
260 μg m⁻² s⁻¹, and 0.41±0.05 μg m⁻² s⁻¹ for 2019, 2020 and 2021, respectively. Probability density functions (PDFs) of daily emission intensity and a map of the multi-year mean are displayed in Fig. 3. The proportion of low values is ~~the highest in one or two months of 2020, which are closely related to the epidemic, during which many key enterprises spontaneously shut down or otherwise limited production. However, on a year-by-year basis, the values in 2021 are lower than in 2020, reflecting the fact that the long-term efforts to reduce emissions may be even more important. Similarly, the highest emission intensity is~~
265 observed in 2019.

Table 1: Summary statistics for plants included in CEMS.

Year	Number of companies	Number of sacks monitored	Number of days without public data	Percentage of days without public data (%)
2019	624	1607	102	27.9
2020	705	1819	24	6.6
2021	714	1836	0	0




Figure 3: CEMS emissions intensity from January 2019 through December 2021 (unit: μg m⁻² s⁻¹): (a) 3-year average gridded NO_x emissions, and (b) PDFs of day-by-day and grid-by-grid emissions over individual years (~~log-scale for x-axis~~).

删除了: convert

删除了: is


删除了: in 2020

删除了: ting

删除了: The

删除了: , consistent with the decade-long decrease in emissions in China...


删除了: able 1:



删除了:

2.2.2 MEIC

280 At the present time, some of the emissions inventories most widely used by the community are MEIC (Zheng et al., 2018) and
the Emission Database for Global Atmospheric Research (EDGAR) (Crippa et al., 2018). In this work, MEIC is selected since
it provides bottom-up emissions of anthropogenic air pollutants over mainland China, with a monthly time step and a
0.25°×0.25° spatial resolution. NO_x emissions are provided from January 2019 to December 2020 over five sectors: agriculture,
industry, power, residential and transportation (Zheng et al., 2021). To match with the higher resolution of TROPOMI grids,
285 all of the MEIC data in this work is mapped uniformly to a 0.05°×0.05° grid, with each TROPOMI sized grid assigned the
same flux as the underlying MEIC grid. The average emission values and PDFs of the grid-by-grid data over Shanxi from
2019 January to 2020 December are given in Fig. 4a and Fig. 4b, demonstrating little change in emissions between the two
years except for at very low values.



290 **Figure 4:** MEIC emissions intensity from January 2019 through December 2020 (unit: $\mu\text{g m}^{-2} \text{s}^{-1}$): (a) 2-year average monthly MEIC
emissions, and (b) PDFs of month-by-month and the grid-by-grid emission over individual years (log-scale for x-axis).

2.3 Wind

Wind speed and direction are from the European Centre for Medium-Range Weather Forecasts, ERA-5 reanalysis products.
This work uses the hourly 6:00 UTC u and v wind products (closest in terms of time to the TROPOMI overpass) at 850 hPa
and 0.25° × 0.25° resolution, available at <https://www.ecmwf.int/en/forecasts/dataset/ecmwf-reanalysis-v5>. The wind was
295 linearly interpolated to follow the TROPOMI 0.05° × 0.05° grid data in space and time. The reason for choosing the 850 hPa
(which is approximately 1500 m) level is two-fold. First, Shanxi has complex topography, with less than 16% of the total area
of the province under 800 m in height, 68% of the surface over 1000 m, and 17% of area over 1500 m, see Fig. 1, leading to a
significant amount of pollutant transport from near the ground to the lower free troposphere. Second, due to the relatively dry

删除了: is


删除了: . This work uses data from 2019 and 2020

删除了: uniformly

删除了: (

删除了:)

删除了: (b)



删除了:

删除了:

删除了:

删除了:

删除了:

conditions, vertical plume-based rise is thought to not be insignificant (Wang et al., 2020). Overall, we aim to use wind speed and direction that correspond to a reasonable approximation of the median of the NO_x emissions vertical profile.

2.4 Variance Maximization

To extract the spatial and temporal features of the extremes of the remotely sensed NO₂ fields in an unbiased manner, the Empirical Orthogonal Functions Principal Components Analysis (EOF) is applied. This technique decomposes the data into a set of orthogonal standing signals in space [EOF] and in time [PC], with those signals contributing the most to the overall variance of the underlying dataset being selected, representing unique phenomenon that control the overall characteristics of the NO₂ columns (Zhou et al., 2016; Lin et al., 2020). Further details including mathematical derivations are given in Björnsson and Venegas (1997), and Cohen (2014). This work retains the first three EOFs, which are found to contribute to 29.4%, 8.4%, and 4.4% of the total variation, with subsequent EOFs each contributing an insignificant amount (less than 4.0%) and therefore no longer considered in this work.

2.5 Model Free Inversion Estimation Framework

The Model Free Inversion Estimation Framework (MFIEF) is based on a mass balance assumption Eq. (2), and the detail is shown in Fig. 5. In the case where there is an observed change in the stock of NO_x in the atmosphere, herein represented as C , in a Lagrangian sense, there must be either a source or sink (Harte, 1988; Seinfeld and Pandis, 1997). When dealing with a fixed spatial grid, such as in this work, there is also a contribution of transport into or from the Lagrangian parcel. The first of these changes in the stock is due to the amount of NO_x emitted, herein represented as E , which will always increase the existing stock. The second of these is the chemical loss of NO_x, which will always lead to a decrease in the stock. The chemical sink of NO_x is dominated by the reaction between NO₂ and OH, via reactions with products formed from the actinic flux (i.e. chemistry such as RO₂), and on aerosol surfaces via heterogeneous reactions (Valin et al., 2013; Kenagy et al., 2018; Romer Present et al., 2020), which herein is described as S . The third change in the stock is the sum of pressure induced and advective transport, which may either increase or decrease the stock. The transport is herein is described as D , and is calculated by the gradient of the product of the wind vector and the NO_x column loadings, which consists of an advective portion (Wang et al., 2014) and a pressure-based portion (Mahowald et al., 2005). The mass conservation equation for NO_x is calculated as

$$dC = E - S + D \quad (2)$$

Solving Eq. (2) for emissions on a grid-by-grid basis requires knowledge of the mass change of the loading in time and space, and detailed consideration of chemical loss and transformation, and transport. An explicit formulation of these processes into a readily solvable mass balance method is derived as Eq. (3):

$$E_{NO_x} = \alpha_1 \cdot \frac{dV_{NO_2}}{dt} + 24 \cdot \frac{\alpha_1}{\alpha_2} \cdot V_{NO_2} + 0.001 \cdot \frac{\alpha_1}{\alpha_3} \cdot (\nabla(\mathbf{u} \cdot V_{NO_2}) + \nabla(\mathbf{v} \cdot V_{NO_2})) \quad (3)$$

删除了: (Björnsson and Venegas, 1997)

删除了: (Cohen, 2014)

删除了: and Eq. (3)

删除了: from

删除了: to

删除了: will

删除了: The second of these is the chemical loss of NO_x, which will always lead to a decrease in the stock. The chemical sink of NO_x is dominated by the reaction between NO₂ and OH (Valin et al., 2013; Kenagy et al., 2018; Romer Present et al., 2020),

删除了: then

删除了:

删除了:

Where E_{NO_x} represents the total atmospheric column emissions of NO_x within the troposphere on a grid-by-grid and day-by-day basis, with a unit of $\mu\text{g m}^{-2} \text{d}^{-1}$. This is the total emission into each column accounting for all sources including anthropogenic sources (industry, vehicle, and residential), biomass burning, and others. V_{NO_2} represents the tropospheric NO_2 column concentration after it has been converted into the unit of $\mu\text{g m}^{-2}$. Due to the fact that TROPOMI only measures NO_2 rather than NO_x , a transformation is required to transform NO_2 columns into NO_x , hereafter given as $NO_x = \alpha_1 \cdot NO_2$. $\alpha_1 \cdot \frac{dV_{NO_2}}{dt}$ computes the NO_x concentration change rate with a unit of $\mu\text{g m}^{-2} \text{d}^{-1}$, assuming the day-to-day temporal derivative of NO_2 exists in the TROPOMI data on the respective days. $24 \cdot \frac{\alpha_1}{\alpha_2} \cdot V_{NO_2}$ represents as the sink of NO_x concentration, where α_2 is related to the NO_x lifetime (unit h), and 24 is the unit change factor. $\nabla(\mathbf{u} \cdot V_{NO_2}) + \nabla(\mathbf{v} \cdot V_{NO_2})$ represents the gradients of daily zonal fluxes, meridional fluxes, and changes in the air column mass and density. These were derived by multiplying the gridded V_{NO_2} with the center point wind vector on a grid-by-grid basis, with unit of $\mu\text{g m}^{-2} \text{d}^{-1}$. This computation assumes that the spatial gradient of TROPOMI NO_2 and reanalysis wind both exist on the respective grids (Cohen and Prinn, 2011). In this case, α_3 is the parameter representing the transport distance (km) per day, where 0.001 is used to convert m to km. V_{NO_2} is observed as either one or two overlapping snapshots of total column information occurring at 13:30 LT (and under some conditions also 101.5 minutes either earlier or later (Tonion and Pirrotti, 2022)). In all cases, the meteorological values and CEMS values are representative of 24-hour total and/or daily average conditions respectively. Therefore, the fitted values of α_1 , α_2 , and α_3 , as presented are representative of 24-hour average or 24-hour net effect respectively, acting on the entire column of NO_x . In all cases, the values of α_1 , α_2 , and α_3 are computed month-to-month over all grids in which data is available, and by bootstrap at other grids and times.

删除了: time

删除了: time

删除了: .

删除了: .

删除了: ·day⁻¹

删除了: and not

删除了: .

删除了: represents the divergence of NO_2

删除了: with a

删除了: of

删除了: ,

删除了: ing

删除了: grid-to-grid

删除了: .



删除了: Maps of daily zonal and meridional fluxes were derived by multiplying gridded V_{NO_2} with spatially gridded wind vector

删除了: and

删除了: -by-grid based on the best fits of all individual daily data within each individual grid, or

删除了: the

删除了: method on a grid-to-grid and month-by-month basis



删除了: E

Figure 5: The framework of MFIEF methodology.

2.6 Additional Analytical Methods

This work employs multiple linear regression to fit the values of α_1 , α_2 , and α_3 on a month-by-month, grid-by-grid basis using all available daily measurements and Eq. (3). These values are further filtered based on both statistics ($p < 0.15$, and removal of outliers defined as elements more than three scaled MAD from the median, calculated by Eq. (4)) and being physically realistic ($|\alpha_1| > 1$, $\alpha_2 < 0$).

$$MAD = c \times \text{median}(\text{abs}(\alpha - \text{median}(\alpha))) \quad (4)$$

$$c = \frac{-1}{\sqrt{2} * \text{erfcinv}(\frac{3}{2})} \approx 1.483$$

α_1 , α_2 , and α_3 are brought into α in Eq. (4) respectively for outlier rejection. Bootstrapping is used as a means to create a new sample to represent the parent sample distribution through multiple repetitions of sampling (Liu and Cohen, 2022). Specifically, the distributions of α_1 , α_2 , and α_3 are sampled across the central 80% of their probability distributions, which are then used to generate a set of pseudo α_1 , α_2 , and α_3 on a grid-by-grid basis where there is no existing a priori and therefore no actual solution of these variables. These bootstrapped pseudo α_1 , α_2 , and α_3 are then used on these specific grids to approximate the emissions of NO_x using Eq. (3) on a daily basis where TROPOMI NO_2 column data and wind data is available. The mean data has been taken out as the daily results. The standard deviation of each grid was calculated as the uncertainty of this emission.

3 Results and Discussion

3.1 Computed Emissions Using CEMS and MEIC

The daily emissions of NO_x are calculated throughout the given spatial and temporal domain using bootstrapping in connection with Eq. (3) and the fitted values of α_1 , α_2 , and α_3 . Fig. 6a and Fig. 6b show the spatial distribution of daily average and variation of NO_x emissions based on CEMS (EI_{CEMS}) from January 2019 to December 2021 over Shanxi at $0.05^\circ \times 0.05^\circ$. For all subsequent emissions values displayed, the numbers correspond to the sum over the day-to-day mean \pm uncertainty. It is observed that the grids with the highest NO_x emission in Shanxi are mainly concentrated in the lower Fen River valley, which happens to also be located at the lowest altitude areas in the province as shown in Fig. 1 pink line area containing Taiyuan Basin, Xinding Basin, and LinFen Basin, which also corresponds to the area containing the highest population density. This area in total accounts for around 25% of the total area of the province and contributes 53% (0.98 ± 0.55 Tg per year) of the total NO_x emission (1.86 ± 1.04 Tg per year) in Shanxi. It is of significance to note the regions with a moderate average density of emissions, ranging from 0.3 to $0.7 \mu\text{g m}^{-2} \text{ s}^{-1}$, contribute 61% (1.14 ± 0.64 Tg per year) of the total emissions. The use of MFIEF can effectively optimize the distribution of the inventory and perform inventory correction based on satellite data, while complementing many areas where there is no existing emissions data, incomplete data, mis-characterized data, or data which may be reasonable on average but not account for daily-scale variability.

删除了: α_1 , α_2 , and α_3

删除了: of α_1 , α_2 , and α_3

删除了: α_1 , α_2 , and α_3

删除了: s

删除了: α_1 , α_2 , and α_3

删除了: α_1 , α_2 , and α_3

删除了: α_1 , α_2 , and α_3 .

删除了: Figures 6(a) and (b)

删除了: Jan

删除了:

删除了:

删除了: .

删除了: 52


删除了: 96

删除了: 83

删除了: 06

删除了: at

删除了: 11




445 **Figure 6:** Daily average emissions based on CEMS from January 2019 to December 2021 over Shanxi at $0.05^\circ \times 0.05^\circ$ (unit: $\mu\text{g m}^{-2} \text{s}^{-1}$): (a) daily average NO_x emissions. (b) day-to-day variability of NO_x emissions. (c) bootstrapping uncertainty range (10% to 90% of distribution).

MEIC was also separately used as the a priori with MFIEF to produce an additional emissions inventory from January 2019 to December 2020, herein called EI_{MEIC} . The total EI_{MEIC} over the province is 1.34 ± 0.60 Tg per year. The multi-annual mean value of EI_{CEMS} minus EI_{MEIC} and histogram of the day-to-day and grid-to-grid emissions of both EI_{CEMS} and EI_{MEIC} from 2019 to 2020 are displayed in Fig. 7. On a multi-annual average basis, EI_{CEMS} is larger than EI_{MEIC} at 98% of grids, while on a day-to-day basis it is larger at 88% of grids. Some higher value appeared in EI_{MEIC} calculated by high MEIC values in some places of Shuozhou, Yangquan, Lvliang, Changzhi, and Jincheng City may be due to mis-positioned hotspots in the existing inventories, in terms of both space and time. But on the opposite, EI_{CEMS} better captured in residential and small industrial sources than EI_{MEIC} .

This is possibly due to the fact that the a priori MEIC distribution has many low values which actually fall within the uncertainty of the bottom-up emissions processes (Cohen and Wang, 2014; Bond, 2004; Crippa et al., 2018). The low a priori emissions on a grid-by-grid basis, which in turn shift the physically filtered values of α_1 towards lower values. At the same time the values of α_2 were shifted to higher values. In tandem with these, the transport term α_3 was shifted to include larger absolute values of distance. This net combination caused the final calculation to be biased smaller overall.

460 Fig. 8 shows the differences between CEMS, EI_{CEMS} , EI_{MEIC} and MEIC, and the 30% error range for CEMS and the computed 10th to 90th percentile error ranges for EI_{CEMS} and EI_{MEIC} as computed from January 2019 to December 2020 in all 11 cities in Shanxi. It shows that while generally CEMS is larger than EI_{CEMS} , that they are always found to be within the error ranges of each other, while EI_{MEIC} only overlaps with CEMS in 6 cities. Similarly, MEIC is always found to be the lowest, while EI_{MEIC}



删除了:

删除了: Jan

删除了: Jan

删除了: 33

删除了: Jan

删除了: Dec

删除了: α_2

删除了: α_3

删除了: ure

删除了: the

删除了: 4

is found to be larger than MEIC and smaller than EI_{CEMS} . In specific EI_{CEMS} and EI_{MEIC} have error ranges which overlap in all cities, while MEIC overlaps with the EI_{MEIC} range in 8 cities.

475






Figure 7: Differences between EI_{CEMS} and EI_{MEIC} through 2019 to 2020 (unit: $\mu\text{g m}^{-2} \text{s}^{-1}$): (a) average values in space of EI_{CEMS} minus EI_{MEIC} . (b) Histogram of EI_{CEMS} and EI_{MEIC} .



480 **Figure 8:** Differences between CEMS, MEIC, EI_{CEMS} and EI_{MEIC} with their error range through 2019 to 2020 in all 11 cities in Shanxi (unit: $\mu\text{g m}^{-2} \text{s}^{-1}$).

3.2 Underlying Factors Contributing to Variance Maximized TROPOMI NO₂ Columns

485 A deeper analysis of the factors contributing to the variance in the TROPOMI NO₂ column measurements is essential to determine if the computed emissions and underlying factors are consistent with the remotely sensed fields both in terms of mean value and temporal variability. Recent practice has devised a way to ensure this consistency through the use of an EOF Analysis (Cohen et al., 2017; Cohen, 2014; Lin et al., 2020), which is applied to the daily TROPOMI NO₂ columns. The three spatial modes contributing the most variation to the observed daily TROPOMI NO₂ fields [EOF1, EOF2, and EOF3] contribute
490 29.4%, 8.4%, and 4.4% respectively, as shown in Fig. 9.

It is asserted that EOF1 is directly driven by EI_{CEMS}. The comparison of EOF1 and the emissions is shown in Fig. 10 in terms of both spatial and temporal scales. By applying four different progressively increasing cutoffs to the domain of EOF1, it is observed that as the EOF1 domain increases in magnitude, that the 3-year mean EI_{CEMS} computed over the same domains also increase in magnitude. Therefore, the more extreme the EOF1 value, the higher the emissions, demonstrating that the emissions are responsible for the first mode of the maximized variance. After detrending and removing any seasonal signal from EI_{CEMS},
495 the result correlates well with PC1 ($r=0.75$, $p<0.01$), indicating that the first mode of variation is strongly connected with the changes in EI_{CEMS} in terms of both spatial and temporal dimensions.





Figure 9: Spatial distribution map of first three modes (a) EOF1, (b) EOF2, and (c) EOF3.

500

删除了: 66

删除了: (PC1)



删除了:




Figure 10: Four different cutoffs of EOF1 to set the domains. The maps in (a-d) are plots of $EOF1/EI_{CEMS}$ where the cutoffs are given as (a) $EOF1 > 0.005$, (b) $EOF1 > 0.01$, (c) $EOF1 > 0.015$, (d) $EOF1 > 0.02$. (e) Histograms of the EI_{CEMS} over the domains given respectively in a-d. (f) Time series of weekly PC1 compare with EI_{CEMS} in whole Shanxi.


510 Second, it is asserted that EOF2 is related to TROPOMI measured UVAI, which physically makes sense, since satellite observations of UVAI are sensitive to absorption of solar radiation by aerosols. There have been numerous studies reporting that absorbing aerosols affect the downwelling surface radiative forcing in the visible (and therefore the actinic flux) (Léon, 2002) as well as OH concentrations (Hammer et al., 2016). Therefore, UVAI indirectly is one component of the chemical decay capacity of NO_x in-situ. Applying four different cutoffs to EOF2, it is observed that as the EOF2 domain increase in

515 magnitude, that the 3-year mean measured TROPOMI UVAI decreases, as demonstrated in Fig. 11. Since UVAI scales with atmospheric absorption, it therefore also scales inversely with surface UV radiation, implying that when the UVAI is lower, that there is more available UV radiation, and hence implicitly faster chemical decay of NO_x . This is consistent with the UV radiation being responsible for the second mode of the maximized variance. The negative correlation observed between absolute values of UVAI weighted by high absolute values of EOF2 ($EOF2 > 0.02$) grid-by-grid in the same EOF2 region is

520 anticorrelated with PC2 ($r = -0.35$, $p < 0.01$). While r in this case is a lot smaller, it is also consistent with theory, since in order

- 删除了: is a proxy of the absorbing aerosol loading, which itself directly constrains the UV in a cloud-free atmosphere
- 删除了: This change in UV radiation at the surface then changes the levels of OH, which then alter the chemical decay of NO_x .
- 删除了: the available
- 删除了: therefore
- 删除了: lower
- 删除了: implies higher
- 删除了: surface
- 删除了: ,
- 删除了: therefore demonstrating that the surface
- 删除了: is

to significantly affect the OH levels, the changes in UV radiation and hence UVAI must be very large, which is found to not occur frequently in time, but when it does occur, it makes a significant impact.



535

Figure 11: Four different cutoffs of EOF2 to set the domains. The maps in (a-d) are plots of EOF2/|UVAI| where the cutoffs are given as (a) EOF2 >0.005, (b) EOF2 >0.01, (c) EOF2 >0.015, (d) EOF2 >0.02. (e) Histograms of the UVAI over the domains given respectively in a-d. (f) Time series of weekly PC2 compare with |UVAI| in the (d) domain.


Finally, it is asserted that EOF3 is related to the transport of NO₂. This term has been specifically computed by taking the variance of the multiple of wind and TROPOMI NO₂ column loadings, specifically $\nabla(\mathbf{u} \cdot V_{NO_2})$. Similarly, to the above cases, it is demonstrated that as four different cutoffs are applied to EOF3, it is observed that as the EOF3 domain increase in magnitude, so does the measured transport based on TROPOMI NO₂ also increase, as observed in Fig. 12. $\nabla(\mathbf{u} \cdot V_{NO_2})$ weighted by EOF3 grid-by-grid positively correlates with PC3 ($r = 0.60$, $p < 0.01$), indicating that the third mode of variability, PC3 is strongly consistent with $\nabla(\mathbf{u} \cdot V_{NO_2})$ in the temporal dimensions. Therefore, transport is responsible for the third mode of the maximized variance.

540

545

删除了: measured

删除了: 70



550 **Figure 12:** Four different cutoffs of EOF3 to set the domains. The maps in (a-d) are plots of EOF1/NO₂-transport where the cutoffs are given as (a) EOF3 >0.005, (b) EOF3 >0.01, (c) EOF3 >0.015, (d) EOF3 >0.02. (e) Histograms of the NO₂-transport over the domains given respectively in a-d. (f) Time series of weekly PC3 compare with ∇(u·V_{NO2}) in whole Shanxi.

3.3 Uncertainty in Emission and the parameters

555 The uncertainty of the computed emissions on a grid-by-grid and day-by-day basis is due to a combination of uncertainties in the satellite data, CEMS or MEIC as a priori emissions, and the calculations involved with computing the various best-fit parameters. NO_x column concentrations and CEMS both have uncertainties about 30%. Another uncertainty comes from the parameters α₁, α₂, and α₃ generated during the regression of Eq. (3) as given in section 2.5. The fitted coefficients are computed month-by-month over the three years from January 2019 through December of 2021. Their absolute value overall mean, 10th percentile, and 90th percentile are found to be α₁=[4.0,1.3,8.2], α₂=[12.3,7.1,18.1] h, and α₃=[239,63,508] km. However, it is observed in the fits that some amount of the variability is not uniform in space and time, with the month-by-month values and standard deviations given in Fig. 13. In general, α₁ tends to be slightly higher during one or a combination of both hotter months and times when the UVAI is low (hence the UV values are high), α₂ shown the lifetime reflect all hours during the day. In general, it tends to be variable, both inter-annual and intra-annual variations seem to drive most of the change. Given

Deleted: ... [1]
Deleted: α₁
Deleted: 3..., α₂
Deleted: 7...,6.7...,1,18.4... h, and α₃
Deleted: 303...39,66...3,666...08] km. Due to the fact that the distributions are skewed to the lower end.
Deleted: the ... other months and of the year, but it also has a...imes when the UVAI is low (hence the higher variability when the ...V values are high) as well, making July and August the only months in which it statistically has fewer small values than the other times of the year... α₂ shown the lifetime reflect all hours during the day. In general, it α₂
Deleted: without any significant seasonal or month of year pattern. Instead,

600 that this is related to both the column average temperature and UV availability, the largest values are found in June or July and the lowest values are found in September. Furthermore, there are other complex forcing factors including the height of the aerosol layer, the total aerosol loadings, cloudiness, and other factors. The absolute magnitudes of α_2 and their uncertainty range are reasonable when compared with vertically integrated and 24-hour integrated chemical transport model values. In general, α_3 also seems to not have any significant seasonal or monthly pattern, with inter-annual and intra-annual terms seeming to dominate. The values tend to be slightly larger than chemical transport models account for, but are reasonable when compared with the ultra-long-range transport simulated for plumes which break the boundary layer. This range, combined with 605 the wide basins of 200 km to 400 km in length, seem to provide a reasonable bound on the output results. One possible reason for this is that the atmospheric wind patterns were slightly different in 2019 due to the El Niño pattern (Hu et al., 2020). The result of overall uncertainty of the emissions as a result of the overall bootstrapped fitting ranges from 8% to 69% on a grid-by-grid basis, as shown in Fig. 3c. In general, the larger relative uncertainty values are observed over mountain regions. A sensitivity test has been performed to test the robustness of the fits assuming that the TROPOMI NO₂ column values are varied 610 near the extreme top and bottom of their 30% uncertainty range. It is shown that the new NO_x emissions in the 70% case are always larger than 0.7 times the original emissions case (spatially annually averaged and grid-by-grid this varies from 0.89 to 1.00, while temporally domain averaged this varies from 0.77 to 1.04). Similarly, the new NO_x emissions in the 130% case over the median 90% of data is smaller than 1.3 times the original emissions case (spatially annually averaged and grid-by-grid this varies from 1.08 to 1.12 while temporally domain averaged this varies from 1.01 to 1.30).

删除了: April

删除了: α_2

删除了:

删除了: α_3

删除了: 31

删除了: 75

删除了: (

删除了:)





Figure 13: Distribution of monthly α_1 (a), α_2 (b) and α_3 (c) calculated based on CEMS.



删除了: a
删除了: a

删除了: α_1

625 **3.4 Application of α_1 to Analyze Different Combustion Technologies**

A significant finding is observed when the value of α_1 is analyzed more closely on a pixel-by-pixel level and compared with underlying CEMS combustion source type. This analysis is motivated by the fact that NO_x is produced during high temperature combustion of air, with three different major parts contributing to the overall amount of NO_x produced: thermal NO_x formation, fuel NO_x formation and chemical NO_x formation (Schwerdt, 2006; Le Bris et al., 2007). This work demonstrates clearly that the values of α_1 are significantly related to the underlying thermodynamic conditions occurring at the time of combustion. Thermal NO_x formation describes the process when N_2 reacts with O_2 in the air at high temperatures (Le Bris et al., 2007), with NO_2 forming preferentially at temperatures between 800 °C and 1200 °C and NO forming preferentially at temperatures above 1200 °C. Thermal NO_x usually dominates the overall NO_x emissions when the temperature is over 1100 °C, and reaches

删除了: allowing for many future applications of the results herein to better understand and monitor such plants around the world.
删除了: in the air

a maximum contribution when the temperature is over 1600 °C. There is additional NO_x produced due to free nitrogen in the fuel itself. Finally, chemical decay may occur when there are mixed organo-nitrides, resulting in the prompt NO_x formation. Therefore, a deeper understanding of the overall and oxygen partial pressures and temperature in the combustion chamber are all important for NO_x formation. In summary, as the temperature of combustion increases, both the amount of NO and NO₂ will increase. When the temperature exceeds 1200 °C, NO will continue to increase while NO₂ will decrease. During any time when the pressure increases, the yield of NO₂ will also decrease and NO will increase (Aho et al., 1995; Turns, 1995).

A deeper look at the various different CEMS sources reveals that the internal combustion processes are extremely important in terms of the overall value of α_1 . Production of cement is a major source of NO_x in Shanxi, with the major technology being dry process rotary kiln technology. Given that the temperature of the main burner of cement rotary kilns are higher than 1400 °C, with some peaking as high as 1800 °C to 2000 °C (Wu et al., 2020; Akgun, 2003), it is expected that there will be a large amount of thermodynamic NO_x generation. As observed at the cement CEMS sites, the computed α_1 has a value always within or above the values computed at powerplants, including both in terms of the mode, as well as at the highest emitting individual values, as show in Fig. 14. Steel and iron are produced through a set of different processes, involving combustion at a range of different temperatures. The steps involved in the blast furnaces as well as some other processes, require a high flame temperature, in the range from 1350 °C to 2000 °C. There are further processes occurring that require a relatively lower temperature, such as in the sinter bed stage, where the highest temperature is only about 1300 °C (Zhou et al., 2018). Therefore, while in general the values are relatively high compared to the other source types below, and are generally found within the range of values at power plants, there are some individual differences as well, including a larger fraction of large values (9-10) and a smaller fraction at very large values (12-15), as observed in Fig. 14. The maximum temperature of the combustion chamber of thermal power plants can reach 2000 °C. In fact, such plants are constantly finding ways to increase the combustion efficiency, so that they can be more energy efficient and produce as much energy per ton of CO₂ emitted. α_1 is relatively high at these sites, consistent with thermal production.


Boilers use a similar technology as power plants, but tend to be smaller and run at a lower temperature range and efficiency. This is because their use is to produce hot water and steam for direct residential and industrial use, not high-pressure steam to run turbines. In general, these boilers have a much smaller overall capacity, and therefore without access to CEMS, may not be otherwise be detectable. For these reasons, it is logical that there is a greater amount of NO₂ produced than the above cases, and subsequently the value of α_1 is much lower, in terms of both the mode, as well as all moderate values (5 and larger) as shown Fig. 14. Coke and aluminum oxide are both produced using a different technique from the other combustion sources, specifically focusing on creating high temperature, oven-like conditions to bake/roast their products. The average temperature of the coke oven charring chamber and aluminum oxide roasting furnace are around 1000 °C (Abyzov, 2019; Neto et al., 2021), with the material temperature continuously held in that temperature for a long period of time, e.g., one day. At the same time, the oxygen content is low. In net, there is far less thermal NO and more thermal NO₂. Aluminum is also smelted in an oven-like condition. Correspondingly, while the mode of α_1 for aluminum oxide is more similar to steel and iron, and the mode of

- 删除了: First
- 删除了: produced will increase along with
- 删除了: .
- 删除了: Furthermore, when
- 删除了: In addition, in situ processes also impact the value of α_1 since there is a rapid adjustment after emitted from a combustion stack into the atmosphere, before the parcel comes to thermodynamic equilibrium (Cohen et al., 2018; Wang et al., 2020). From Fig. 12(a) it can be seen that the values are highest in the hottest months without maximum UV (July and August) and are lowest in the coldest months with the minimum UV (November, December, and January), with both factors moderating the combustion values during the atmospheric in situ processing time. This is especially important in the case of hotter power sources, since they will contain more buoyancy, and rise to a higher height, making them more likely to be in contact with air which is more exposed to UV and also generally colder than the surface. Overall, the value of α_1 seems to rely on both the temperature under which the initial NO_x was generated, as ... [5]
- 删除了: α_1
- 删除了: α_1
- 删除了: the error range of
- 删除了: some of the
- 删除了: ly
- 删除了: highest
- 删除了: (a)
- 删除了: ↵
- 删除了: ,
- 删除了: and are usually found within the
- 删除了: s
- 删除了: values of
- 删除了: α_1
- 删除了: computed which are slightly outside the range of the # ... [6]
- 删除了: (c)
- 删除了: ↵
- 删除了: As observed in Fig. 14(b), α_1
- 删除了: Industrial b
- 删除了: (as small as one tenth of the total power output)
- 删除了: However, analyzing the values of α_1 over these sites ... [7]
- 删除了: therefore
- 删除了: as expected, the values of α_1
- 删除了: are significantly and consistently lower
- 删除了: (e)
- 删除了: ↵
- 删除了: α_1

α_1 for coke is slightly lower, they both share a generally similar distribution of α_1 with the bulk of their observations lower than 5, and a small but not insignificant tail, as detailed in Fig. 14.

740 In addition, in-situ processes also impact the value of α_1 since there is a rapid adjustment after emitted from a combustion stack into the atmosphere, before the parcel comes to thermodynamic equilibrium (Cohen et al., 2018; Wang et al., 2020). From Fig. 13a it can be seen while there is a pattern where the values are higher in certain months than others, that there is also a clear difference between the different source types. This demonstrates clearly that the atmospheric processing modulates the process to some extent, but does not dominate the signal on average. This is especially important in the case of hotter power sources, since they will contain more buoyancy, and rise to a higher height, making them more likely to be in contact with air which is more exposed to UV and also generally colder than the surface. Overall, the value of α_1 seems to rely on both the temperature under which the initial NO_x was generated, as well as any rapid processes taking place once it is emitted into the atmosphere (including photo-chemistry and vertical lofting).


745



删除了: the values of α_1 are relatively lower.

删除了: (d) and Fig. 14(f)

删除了:



755 **Figure 14:** Histogram of monthly α_1 calculated based on CEMS using MFIEF at the following source: cement factories, power plants, steel and iron factories, coke ovens, boilers, and aluminium oxide factories.

4 Conclusions

MFIEF based on daily measurements from TROPOMI and a priori daily emissions from CEMS successfully inverts daily NO_x emissions. First, the emissions computed match well with known urban, suburban, and industrial locations. Secondly, the best fit values for thermodynamics (α_1) and first order chemical decay (α_2) are both physically realistic, while the best fit term for transport (α_3) is reasonable based on the mountain and basin geography of the province. Thirdly, the variability of emissions in terms of different geographic location, source types, special events which changed the emissions levels (such as the onset of COVID-19), and general oxidative, photochemical, and transport conditions of the atmosphere on a monthly-scale, are all consistent with what is known. Fourth, the uncertainty is observed to be lower than the day-to-day variability over 30% of the region, and mostly distributed in regions with higher emission intensities, showing that the results on a day-to-day basis are significant.

MFIEF emissions computed using different a priori datasets (CEMS versus MEIC) yield significant differences from on-the-ground CEMS measurements and local geospatial knowledge of where large sources exist. EI_{MEIC} severely underestimates sources which have a lower amount of emissions, as well as newer sources, while at the same time overestimates sources in Yangquan, Xiaoyi, and enormous steel and power plants, among other similar high emissions sources. EI_{MEIC} incorporates too many low values from MEIC in the suburban and rural areas, leading to many of these grids possessing physically unreasonable or at best very low/high values of α_1/α_2 values. A site-by-site comparison with CEMS shows very large differences with EI_{MEIC} at locations which MEIC do have an a priori value, indicating that any a priori dataset must be both precise and accurate, including on a day-to-day basis in order to present a good overall model fit and emissions output product. The MFIEF method as a procedure can quantitatively ally some of these shortcomings of present-day a priori inventories.

The results of variance maximization analysis of 3-years of daily TROPOMI NO_2 columns reveal 3 patterns that drive the variability in the NO_2 columns. EI_{CEMS} is attributed as responsible for pattern 1, measured UVAI from TROPOMI (which is inversely related to photochemistry) is attributed as responsible for pattern 2, while transport (computed from the gradient of reanalysis wind and TROPOMI NO_2 columns) is attributed as responsible for pattern 3. This procedure and its results form a basis of a best-practice approach that the community can adapt to subsequently analyze the efficacy of future emissions products. It is essential to ensure that emissions not only match on average, but also match well with the observed spatial and temporal gradients of the observed remotely sensed fields.

It is observed that the calculated values of α_1 are correlated with the thermodynamic conditions of underlying large combustion sources. This offers a new and self-consistent way to quantify the underlying combustion conditions of NO_x generation using remotely sensed measurements. Here highest mode and very high values are consistently associated with α_1 at cement factories. Generally high modes of α_1 , with significant values at moderately high and high values are found respectively at steel/iron and power plants. Locations that have CEMS aluminum oxide plants, coke ovens, and boilers are generally lower, but in the case

删除了: Distribution

删除了: (mean values, 20th, 40th, 60th, and 80th percentile values)

删除了: (a)

删除了: ;

删除了: (b)

删除了: ;

删除了: (c)

删除了: ;

删除了: (d)

删除了: ;

删除了: (e)

删除了: ;

删除了: (f)

删除了: es

设置了格式: 字体: (默认) +西文正文 (Times New Roman), 非突出显示

删除了: **Thirdly, the general variability in geography, month of the year, and years before and after COVID-19 are all consistent with what is known.**

删除了: at the province's

删除了: geospatial

设置了格式: 字体: (默认) +西文正文 (Times New Roman), (中文) 中文(中国), 非突出显示

删除了: is consistently high. At locations that have CEMS

设置了格式: 字体: (默认) +西文正文 (Times New Roman), 非突出显示

设置了格式: 字体: (默认) +西文正文 (Times New Roman), 非突出显示

删除了: cement

删除了: , α_1 is very high, but also has less consistency

删除了: are

of both aluminum oxide and coke ovens, there is still a significant range of α_1 up to 5, while in the case of boilers, there is nearly no high value. There is a slight offset based on the atmospheric temperature, UV radiation, and other atmospheric mixing and chemical forcings, with both colder and lower radiation months having α_1 slightly negatively offset and hotter temperature and higher radiation conditions having α_1 slightly positively offset. While this offset is consistent across all plant types, it is slightly stronger for the hottest types (cement, iron/steel, and electricity), which are most likely to rise to a higher elevation and therefore be more impacted by the surrounding atmospheric conditions. In all cases, this offset is smaller than the difference between the different source types, indicating that attribution is possible.

The procedure introduced here offers a next step advance in terms of computing emissions from a top-down perspective. Community adaptation and use of these new results will ideally allow improvement in bottom-up inventory constraints and attribution. This work would be improved by reduction in remotely sensed measurement errors/uncertainties, increased use of and access to surface CEMS and other high quality surface flux measurements, improved a priori emissions databases, and higher frequency temporal data availability from new geostationary satellite platforms. The adaptation of day-to-day and other higher frequency quantification data sources, especially sources with well quantified errors would also improve the work herein. The ability to identify large and moderately large plants and industrial sources could be used to identify and quantify sources from many parts of the Global South where ground-based measurements may not be readily available. It is hoped that the findings herein will be improved upon, possibly in an iterative manner, allowing for more precision and predictability, so that emissions and environmental regulators can have more quantitative support to focus their efforts.

Data Availability

The satellite NO₂ datasets used in this study are available at <https://disc.gsfc.nasa.gov/datasets>. The ERA-5 reanalysis product is available at <https://doi.org/10.24381/cds.bd0915c6>. The CEMS online data is available at <https://sthjt.shanxi.gov.cn/wryjg>. The MEIC product can be accessed from <https://doi.org/10.6084/m9.figshare.c.5214920.v2>. All of the data and underlying Figures are available for download at <https://doi.org/10.6084/m9.figshare.20459889>

Author Contributions

Xiaolu Li, Jason Blake Cohen, Kai Qin and Hong Geng developed the research question and set up the whole experimental program. Xiaolu Li wrote the manuscript and performed the data analysis with input from Jason Blake Cohen, Kai Qin, Rui Zhang, Liling Wu and Liqin Zhang. Liling Wu compared this result with the existing emission inventories and environmental statistics data. Liling Wu and Xiaohui Wu rectify the deviation of CEMS location and make statistics of annual changes. Jason Blake Cohen drafted and corrected the manuscript. Hong Geng, Xiaohui Wu, Chengli Yang, Rui Zhang, and Liqin Zhang supported consultation of the local situation and CEMS data. All authors discussed the results and contributed to the final manuscript.

删除了: have a few individual high values of α_1

删除了: and

删除了: months with

删除了: T

删除了: is

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

删除了: but especially so for

删除了: This work would be improved by reduction in remotely sensed measurement errors/uncertainties, increased use of and access to surface CEMS and other high quality surface flux measurements, and improved a priori emissions databases.

Competing Interests

The authors declare that they have no conflict of interests.

Acknowledgments

The authors would like to thank the continuous emissions monitoring systems (CEMS) for the provision of their in situ data.

855 This work thanks the PIs of the TROPOMI, ERA-5, MEIC products for making their data available. The study was supported by the National Natural Science Foundation of China (42075147), the Shanxi Province Major Science and Technique Program (202101090301013), and the Shanxi Province Postgraduate Education Innovation Program (2021Y035).

References

- Abyzov, A.: Aluminum oxide and alumina ceramics (review). Part 1. Properties of Al₂O₃ and commercial production of dispersed Al₂O₃, *Refract. Ind. Ceram.*, 60, 24-32, <https://doi.org/10.1007/s11148-019-00304-2>, 2019.
- 860 Aho, M. J., Paakkinen, K. M., Pirkonen, P. M., Kilpinen, P., and Hupa, M.: The effects of pressure, oxygen partial pressure, and temperature on the formation of N₂O, NO, and NO₂ from pulverized coal, *Combust. Flame.*, 102, 387-400, [https://doi.org/10.1016/0010-2180\(95\)00019-3](https://doi.org/10.1016/0010-2180(95)00019-3), 1995.
- Akgun, F.: Investigaton of energy saving and NO_x reduction possibilities in a rotary cement kiln, *Int. J. Energ. Res.*, 27, 455-865 465, <https://doi.org/10.1002/er.888>, 2003.
- Beirle, S., Boersma, K. F., Platt, U., Lawrence, M. G., and Wagner, T.: Megacity Emissions and Lifetimes of Nitrogen Oxides Probed from Space, *Science*, 333, 1737-1739, <https://doi.org/10.1126/science.1207824>, 2011.
- Beirle, S., Borger, C., Dörner, S., Li, A., Hu, Z. K., Liu, F., Wang, Y., and Wagner, T.: Pinpointing nitrogen oxide emissions from space, *Sci. Adv.*, 5, <https://doi.org/10.1126/sciadv.aax9800>, 2019.
- 870 Björnsson, H. and Venegas, S.: A manual for EOF and SVD analyses of climatic data, Open File Rep, Department of Atmospheric and Oceanic Sciences and Centre for Climate and Global Change Research, McGill University, 1997.
- Bond, T. C.: A technology-based global inventory of black and organic carbon emissions from combustion, *J. Geophys. Res.*, <https://doi.org/10.1029/2003jd003697>, 2004.
- Bond, T. C., Bhardwaj, E., Dong, R., Jogani, R., Jung, S., Roden, C., Streets, D. G., and Trautmann, N. M.: Historical emissions 875 of black and organic carbon aerosol from energy-related combustion, 1850-2000, *Global Biogeochem. Cycles*, 21, GB2018, <https://doi.org/10.1029/2006GB002840>, 2007.
- Cai, B., Liang, S., Zhou, J., Wang, J., Cao, L., Qu, S., Xu, M., and Yang, Z.: China high resolution emission database (CHRED) with point emission sources, gridded emission data, and supplementary socioeconomic data, *Resour. Conserv. Recycl.*, 129, 232-239, <https://doi.org/10.1016/j.resconrec.2017.10.036>, 2018.

- 880 Chen, X., Liu, Q., Sheng, T., Li, F., Xu, Z., Han, D., Zhang, X., Huang, X., Fu, Q., and Cheng, J.: A high temporal-spatial emission inventory and updated emission factors for coal-fired power plants in Shanghai, China, *Sci. Total. Environ.*, 688, 94-102, <https://doi.org/10.1016/j.scitotenv.2019.06.201>, 2019.
- Chen, Y. H. and Prinn, R. G.: Estimation of atmospheric methane emissions between 1996 and 2001 using a three-dimensional global chemical transport model, *Journal of Geophysical Research: Atmospheres*, 111, <https://doi.org/10.1029/2005JD006058>, 2006.
- 885 Cohen, J. B.: Quantifying the occurrence and magnitude of the Southeast Asian fire climatology, *Environ. Res. Lett.*, 9, <https://doi.org/10.1088/1748-9326/9/11/114018>, 2014.
- Cohen, J. B. and Prinn, R. G.: Development of a fast, urban chemistry metamodel for inclusion in global models, *Atmos. Chem. Phys.*, 11, <https://doi.org/7629-7656>, 10.5194/acp-11-7629-2011, 2011.
- 890 Cohen, J. B. and Wang, C.: Estimating global black carbon emissions using a top-down Kalman Filter approach, *J. Geophys. Res. Atmos.*, 119, 307-323, <https://doi.org/10.1002/2013jd019912>, 2014.
- Cohen, J. B., Lecoœur, E., and Hui Loong Ng, D.: Decadal-scale relationship between measurements of aerosols, land-use change, and fire over Southeast Asia, *Atmos. Chem. Phys.*, 17, 721-743, <https://doi.org/10.5194/acp-17-721-2017>, 2017.
- Cohen, J. B., Ng, D. H. L., Lim, A. W. L., and Chua, X. R.: Vertical distribution of aerosols over the Maritime Continent during El Niño, *Atmos. Chem. Phys.*, 18, 7095-7108, <https://doi.org/10.5194/acp-18-7095-2018>, 2018.
- 895 Crippa, M., Guizzardi, D., Muntean, M., Schaaf, E., Dentener, F., van Aardenne, J. A., Monni, S., Doering, U., Olivier, J. G. J., Pagliari, V., and Janssens-Maenhout, G.: Gridded emissions of air pollutants for the period 1970–2012 within EDGAR v4.3.2, *Earth Syst. Sci. Data*, 10, 1987-2013, <https://doi.org/10.5194/essd-10-1987-2018>, 2018.
- DEESP: Shanxi Province Ecology and Environment Status Bulletin: <https://sthjt.shanxi.gov.cn/zwgk/hjgb/hjzkgb/index.shtml>, last access: 14 December 2022, 2015 (in Chinese).
- 900 DEESP: Shanxi Province Ecology and Environment Status Bulletin: <https://sthjt.shanxi.gov.cn/zwgk/hjgb/hjzkgb/index.shtml>, last access: 14 December 2022, 2020 (in Chinese).
- Geng, G., Xiao, Q., Zheng, Y., Tong, D., Zhang, Y., Zhang, X., Zhang, Q., He, K., and Liu, Y.: Impact of China's air pollution prevention and control action plan on PM_{2.5} chemical composition over eastern China, *Science China (Earth Sciences)*, 62, 1872-1884, <https://doi.org/10.1007/s11430-018-9353-x>, 2019.
- 905 Goldberg, D. L., Lu, Z., Streets, D. G., de Foy, B., Griffin, D., McLinden, C. A., Lamsal, L. N., Krotkov, N. A., and Eskes, H.: Enhanced Capabilities of TROPOMI NO₂: Estimating NO_x from North American Cities and Power Plants, *Environ. Sci. Technol.*, 53, 12594-12601, <https://doi.org/10.1021/acs.est.9b04488>, 2019.
- Green, A., Singhal, R., and Venkateswar, R.: Analytic extensions of the Gaussian plume model, *J. Air Pollu. Control Assoc.*, 30, 773-776, <https://doi.org/10.1080/00022470.1980.10465108>, 1980.
- 910 Gu, X., Li, B., Sun, C., Liao, H., Zhao, Y., and Yang, Y.: An improved hourly-resolved NO_x emission inventory for power plants based on continuous emission monitoring system (CEMS) database: A case in Jiangsu, China, *J. Clean. Prod.*, 369, <https://doi.org/10.1016/j.jclepro.2022.133176>, 2022.

删除了: Griffin, D., McLinden, C. A., Boersma, F., Bourassa, A., Dammers, E., Degenstein, D., Eskes, H., Fehr, L., Fioletov, V., Hayden, K., Kharol, S. K., Li, S. M., Makar, P., Martin, R. V., Mihele, C., Mittermeier, R. L., Krotkov, N., Snee, M., Lamsal, L. N., Ter Linden, M., van Geffen, J., Veefkind, P., Wolde, M., and Zhao, X.: High resolution mapping of nitrogen dioxide with TROPOMI: First results and validation over the Canadian oil sands, *Geophys. Res. Lett.*, 46, 1049-1060, <https://doi.org/10.1029/2018GL081095>, 2019.⁴⁰

- 925 [Hammer, M. S., Martin, R. V., van Donkelaar, A., Buchard, V., Torres, O., Ridley, D. A., and Spurr, R. J. D.: Interpreting the ultraviolet aerosol index observed with the OMI satellite instrument to understand absorption by organic aerosols: implications for atmospheric oxidation and direct radiative effects, *Atmos. Chem. Phys.*, 16, 2507-2523, <https://doi.org/10.5194/acp-16-2507-2016>, 2016.](#)
- Harte, J.: Consider a spherical cow: A course in environmental problem solving, University Science Books, 1988.
- 930 [Henk E., Jos v. G., Folkert B., Kai-Uwe E., Arnoud A., Mattia P., Maarten S., J. Pepijn V., and Diego L.: Sentinel-5 precursor/TROPOMI Level 2 ~~product user manual nitrogendioxide~~, Royal Netherlands Meteorological Institute Ministry of Infrastructure and Water Management, Open File Rep., 2021.](#)
- Hoesly, R. M., Smith, S. J., Feng, L., Klimont, Z., Janssens-Maenhout, G., Pitkanen, T., Seibert, J. J., Vu, L., Andres, R. J., Bolt, R. M., Bond, T. C., Dawidowski, L., Kholod, N., Kurokawa, J.-i., Li, M., Liu, L., Lu, Z., Moura, M. C. P., ~~O'Rourke, P. R.~~, and Zhang, Q.: Historical (1750–2014) anthropogenic emissions of reactive gases and aerosols from the Community Emissions Data System (CEDS), *Geosci. Model Dev.*, 11, 369-408, <https://doi.org/10.5194/gmd-11-369-2018>, 2018.
- 935 [Hu, P., Chen, W., Chen, S., Liu, Y., and Huang, R.: Extremely early summer monsoon onset in the South China Sea in 2019 following an El Niño event, *Mon. Weather Rev.*, 148, 1877-1890, <https://doi.org/10.1175/MWR-D-19-0317.1>, 2020.](#)
- Jacob, D. J., Logan, J. A., Gardner, G. M., Yevich, R. M., Spivakovsky, C. M., Wofsy, S. C., Sillman, S., and Prather, M. J.: Factors regulating ozone over the United States and its export to the global atmosphere, *J. Geophys. Res. Atmos.*, 98, 14817-14826, <https://doi.org/10.1029/98JD01224>, 1993.
- 940 [Jiang, X., Li, G., and Fu, W.: Government environmental governance, structural adjustment and air quality: A quasi-natural experiment based on the Three-year Action Plan to Win the Blue Sky Defense War, *J. Environ. Manage.*, 277, 111470, <https://doi.org/10.1016/j.jenvman.2020.111470>, 2021.](#)
- [J.H.G.M. van Geffen, H. J. Eskes, K.F. Boersma and J.P. Veefkind: TROPOMI ATBD of the total and tropospheric NO₂ data products, Open File Rep., 2021.](#)
- 945 [Karl, T., Lamprecht, C., Graus, M., Cede, A., Tiefengraber, M., Vila-Guerau de Arellano, J., Gurarie, D., and Lenschow, D.: High urban NO_x triggers a substantial chemical downward flux of ozone, *Sci. Adv.*, 9, eadd2365, <https://doi.org/doi:10.1126/sciadv.add2365>, 2023.](#)
- Karplus, V. J., Zhang, S., and Almond, D.: Quantifying coal power plant responses to tighter SO₂ emissions standards in China, *Proc. Natl. Acad. Sci. U S A*, 115, 7004-7009, <https://doi.org/10.1073/pnas.1800605115>, 2018.
- 950 [Kenagy, H. S., Sparks, T. L., Ebben, C. J., Wooldrige, P. J., Lopez-Hilfiker, F. D., Lee, B. H., Thornton, J. A., McDuffie, E. E., Fibiger, D. L., Brown, S. S., Montzka, D. D., Weinheimer, A. J., Schroder, J. C., Campuzano-Jost, P., Day, D. A., Jimenez, J. L., Dibb, J. E., Campos, T., Shah, V., Jaeglé, L., and Cohen, R. C.: NO_x lifetime and NO_x partitioning during winter, *J. Geophys. Res. Atmos.*, 123, 9813-9827, <https://doi.org/10.1029/2018jd028736>, 2018.](#)
- 955 [Kong, H., Lin, J., Zhang, R., Liu, M., Weng, H., Ni, R., Chen, L., Wang, J., Yan, Y., and Zhang, Q.: High-resolution \(0.05° × 0.05°\) NO_x emissions in the Yangtze River Delta inferred from OMI, *Atmos. Chem. Phys.*, 19, 12835-12856, <https://doi.org/10.5194/acp-19-12835-2019>, 2019.](#)

删除了: Eskes

删除了: .

删除了: Boersma

删除了: Eichmann

删除了: Apituley

删除了: Pedergnana

删除了: Sneeep

删除了: Veefkind

删除了: Loyola

删除了: Product

删除了: User

删除了: Manual

删除了: Nitrogendioxide

删除了: O'Rourke

删除了: "

- Lange, K., Richter, A., and Burrows, J. P.: Variability of nitrogen oxide emission fluxes and lifetimes estimated from Sentinel-5P TROPOMI observations, *Atmos. Chem. Phys.*, 22, 2745-2767, <https://doi.org/10.5194/acp-22-2745-2022>, 2022.
- 975 [Léon, J. F.: Aerosol direct radiative impact over the INDOEX area based on passive and active remote sensing, *J. Geophys. Res.*, 107, <https://doi.org/10.1029/2000jd000116>, 2002.](https://doi.org/10.1029/2000jd000116)
- Le Bris, T., Cadavid, F., Caillat, S., Pietrzyk, S., Blondin, J., and Baudoin, B.: Coal combustion modelling of large power plant, for NO_x abatement, *Fuel*, 86, 2213-2220, <https://doi.org/10.1016/j.fuel.2007.05.054>, 2007.
- 980 [Li, C., Hammer, M. S., Zheng, B., and Cohen, R. C.: Accelerated reduction of air pollutants in China, 2017-2020, *Science of The Total Environment*, 803, 150011, <https://doi.org/10.1016/j.scitotenv.2021.150011>, 2022a.](https://doi.org/10.1016/j.scitotenv.2021.150011)
- Li, H., Zhang, J., Wen, B., Huang, S., Gao, S., Li, H., Zhao, Z., Zhang, Y., Fu, G., and Bai, J.: Spatial-temporal distribution and variation of NO₂ and its sources and chemical sinks in Shanxi province, China, *Atmosphere*, 13, 1096, <https://doi.org/10.3390/atmos13071096>, 2022.
- Li, M., Liu, H., Geng, G., Hong, C., Liu, F., Song, Y., Tong, D., Zheng, B., Cui, H., Man, H., Zhang, Q., and He, K.: Anthropogenic emission inventories in China: a review, *Natl. Sci. Rev.*, 4, 834-866, <https://doi.org/10.1093/nsr/nwx150>, 2017a.
- 985 Li, M., Zhang, Q., Kurokawa, J. I., Woo, J. H., He, K., Lu, Z., Ohara, T., Song, Y., Streets, D. G., Carmichael, G. R., Cheng, Y., Hong, C., Huo, H., Jiang, X., Kang, S., Liu, F., Su, H., and Zheng, B.: MIX: a mosaic Asian anthropogenic emission inventory under the international collaboration framework of the MICS-Asia and HTAP, *Atmos. Chem. Phys.*, 17, 935-963, <https://doi.org/10.5194/acp-17-935-2017>, 2017b.
- 990 Lin, C., Cohen, J. B., Wang, S., and Lan, R.: Application of a combined standard deviation and mean based approach to MOPITT CO column data, and resulting improved representation of biomass burning and urban air pollution sources, *Remote. Sens. Environ.*, 241, 111720, <https://doi.org/10.1016/j.rse.2020.111720>, 2020.
- Liu, J. and Cohen, J.: Quantifying the Missing Half of Daily NO_x Emissions over South, Southeast and East Asia, nature portfolio [preprint], <https://doi.org/10.21203/rs.3.rs-1613262/v1>, 11 May, 2022.
- 995 Liu, M., van der A, R., van Weele, M., Eskes, H., Lu, X., Veeffkind, P., de Laat, J., Kong, H., Wang, J., Sun, J., Ding, J., Zhao, Y., and Weng, H.: A New Divergence Method to Quantify Methane Emissions Using Observations of Sentinel-5P TROPOMI, *Geophys. Res. Lett.*, 48, 10, <https://doi.org/1029/2021gl094151>, 2021.
- Mahowald, N. M., Baker, A. R., Bergametti, G., Brooks, N., Duce, R. A., Jickells, T. D., Kubilay, N., Prospero, J. M., and Tegen, I.: Atmospheric global dust cycle and iron inputs to the ocean, *Global Biogeochem. Cycles*, 19, <https://doi.org/10.1029/2004gb002402>, 2005.
- 1000 Martin, R. V.: Global inventory of nitrogen oxide emissions constrained by space-based observations of NO₂ columns, *J. Geophys. Res.*, 108, <https://doi.org/10.1029/2003jd003453>, 2003.
- [McDonald, B. C., Gentner, D. R., Goldstein, A. H., and Harley, R. A.: Long-term trends in motor vehicle emissions in U.S. urban areas, *Environ. Sci. Technol.*, 47, 10022-10031, <https://doi.org/10.1021/es401034z>, 2013.](https://doi.org/10.1021/es401034z)

- 1005 Mijling, B. and van der A, R. J.: Using daily satellite observations to estimate emissions of short-lived air pollutants on a mesoscopic scale, *J. Geophys. Res. Atmos.*, 117, <https://doi.org/10.1029/2012JD017817>, 2012.
- Neto, G. F., Leite, M., Marcelino, T., Carneiro, L., Brito, K., and Brito, R.: Optimizing the coke oven process by adjusting the temperature of the combustion chambers, *Energy*, 217, 119419, <https://doi.org/10.1016/j.energy.2020.119419>, 2021.
- 1010 Ohara, T., Akimoto, H., Kurokawa, J.-i., Horii, N., Yamaji, K., Yan, X., and Hayasaka, T.: An Asian emission inventory of anthropogenic emission sources for the period 1980–2020, *Atmos. Chem. Phys.*, 7, 4419–4444, <https://doi.org/10.5194/acp-7-4419-2007>, 2007.
- Qin, K., Shi, J., He, Q., Deng, W., Wang, S., Liu, J., and Cohen, J. B.: New Model-Free Daily Inversion of NO_x Emissions using TROPOMI (MCMFE-NO_x): Deducing a See-Saw of Halved Well Regulated Sources and Doubled New Sources, *ESS Open Archive* [preprint], <https://doi.org/10.1002/essoar.10512010.1>, July 26, 2022.
- 1015 Qu, Z., Henze, D. K., Theys, N., Wang, J., and Wang, W.: Hybrid Mass Balance/4D-Var Joint Inversion of NO_x and SO₂ Emissions in East Asia, *J. Geophys. Res. Atmos.*, 124, 8203–8224, <https://doi.org/10.1029/2018JD030240>, 2019.
- [Rigby, M., Prinn, R. G., Fraser, P. J., Simmonds, P. G., Langenfelds, R. L., Huang, J., Cunnold, D. M., Steele, L. P., Krummel, P. B., Weiss, R. F., O'Doherty, S., Salameh, P. K., Wang, H. J., Harth, C. M., Mühle, J., and Porter, L. W.: Renewed growth of atmospheric methane, *Geophys. Res. Lett.*, 35, <https://doi.org/10.1029/2008gl036037>, 2008.](#)
- 1020 [Rollins, A. W., Browne, E. C., Min, K. E., Pusede, S. E., Wooldridge, P. J., Gentner, D. R., Goldstein, A. H., Liu, S., Day, D. A., Russell, L. M., and Cohen, R. C.: Evidence for NO_x Control over Nighttime SOA Formation, *Science*, 337, 1210–1212, <https://doi.org/10.1126/science.1221520>, 2012.](#)
- [Romer Present, P. S., Zare, A., and Cohen, R. C.: The changing role of organic nitrates in the removal and transport of NO_x, *Atmos. Chem. Phys.*, 20, 267–279, <https://doi.org/10.5194/acp-20-267-2020>, 2020.](#)
- 1025 Schreifels, J. J., Fu, Y. L., and Wilson, E. J.: Sulfur dioxide control in China: policy evolution during the 10th and 11th Five-year Plans and lessons for the future, *Energ Policy*, 48, 779–789, <https://doi.org/10.1016/j.enpol.2012.06.015>, 2012.
- Schwerdt, C.: Modelling NO_x-formation in combustion processes, MSc Theses, 2006.
- [Seigneur, C., A. B. Hudischewskyj, J. H. Seinfeld, K. T. Whitby, E. R. Whitby, J. R. Brock, and H. M. Barnes. Simulation of aerosol dynamics: A comparative review of mathematical models, *Aerosol Sci. Technol.*, 5, 205–222, <https://doi.org/10.1080/02786828608959088>, 1986.](#)
- 1030 Seinfeld, J. and Pandis, S.: Atmospheric chemistry and physics, New York, 1997.
- Singh, A. and Agrawal, M.: Acid rain and its ecological consequences, *Journal of Environmental Biology*, 29, 15, 2007.
- S5P MPC VDAF Validation Web Article: Nitrogen Dioxide Column Data: https://mpc-vdaf.tropomi.eu/ProjectDir/reports/pdf/S5P-MPC-VDAF-WVA-L2_NO2_20180904.pdf, last access: 18 December 2022.
- 1035 [Tang, L., Xue, X., Qu, J., Mi, Z., Bo, X., Chang, X., Wang, S., Li, S., Cui, W., and Dong, G.: Air pollution emissions from Chinese power plants based on the continuous emission monitoring systems network, *Sci. Data*, 7, 325, <https://doi.org/10.1038/s41597-020-00665-1>, 2020.](#)

删除了: Shanxi Province Ecology and Environment Status Bulletin (2015, 2020): <https://sthit.shanxi.gov.cn/zwgk/hjgb/hjzkgb/index.shtml>, last access: 14 December 2022.↖

- Tang, L., Qu, J. B., Mi, Z. F., Bo, X., Chang, X. Y., Anadon, L. D., Wang, S. Y., Xue, X. D., Li, S. B., Wang, X., and Zhao, X. H.: Substantial emission reductions from Chinese power plants after the introduction of ultra-low emissions standards, *Nature Energy*, 4, 929-938, <https://doi.org/10.1038/s41560-019-0468-1>, 2019.
- 1045 [Tonion, F. and Pirotti, F.: Sentinel-5p NO₂ data: cross-validation and comparison with ground measurements, ISPRS Archives, XLIII-B3-2022, 749-756, <https://doi.org/10.5194/isprs-archives-XLIII-B3-2022-749-2022>, 2022.](#)
- Tu, Q., Schneider, M., Hase, F., Khosrawi, F., Ertl, B., Necki, J., Dubravica, D., Diekmann, C. J., Blumenstock, T., and Fang, D.: Quantifying CH₄ emissions in hard coal mines from TROPOMI and IASI observations using the wind-assigned anomaly method, *Atmos. Chem. Phys.*, 22, 9747-9765, <https://doi.org/10.5194/acp-22-9747-2022>, 2022a.
- 1050 Tu, Q., Hase, F., Schneider, M., García, O., Blumenstock, T., Borsdorff, T., Frey, M., Khosrawi, F., Lorente, A., Alberti, C., Bustos, J. J., Butz, A., Carreño, V., Cuevas, E., Curcoll, R., Diekmann, C. J., Dubravica, D., Ertl, B., Estruch, C., León-Luis, S. F., Marrero, C., Morgui, J. A., Ramos, R., Scharun, C., Schneider, C., Sepúlveda, E., Toledano, C., and Torres, C.: Quantification of CH₄ emissions from waste disposal sites near the city of Madrid using ground- and space-based observations of COCCON, TROPOMI and IASI, *Atmos. Chem. Phys.*, 22, 295-317, [https://doi.org/10.5194/acp-22-295-](https://doi.org/10.5194/acp-22-295-2022)
- 1055 [2022](#), 2022b.
- Turns, S. R.: Understanding NO_x formation in nonpremixed flames: Experiments and modeling, *Progress in Energy and Combustion Science*, 21, 361-385, [https://doi.org/10.1016/0360-1285\(94\)00006-9](https://doi.org/10.1016/0360-1285(94)00006-9), 1995.
- Valin, L. C., Russell, A. R., and Cohen, R. C.: Variations of OH radical in an urban plume inferred from NO₂ column measurements, *Geophys. Res. Lett.*, 40, 1856-1860, <https://doi.org/10.1002/grl.50267>, 2013.
- 1060 Veefkind, J. P., Aben, I., McMullan, K., Förster, H., de Vries, J., Otter, G., Claas, J., Eskes, H. J., de Haan, J. F., Kleipool, Q., van Weele, M., Hasekamp, O., Hoogeveen, R., Landgraf, J., Snel, R., Tol, P., Ingmann, P., Voors, R., Kruizinga, B., Vink, R., Visser, H., and Levelt, P. F.: TROPOMI on the ESA Sentinel-5 Precursor: A GMES mission for global observations of the atmospheric composition for climate, air quality and ozone layer applications, *Remote. Sens. Environ.*, 120, 70-83, <https://doi.org/10.1016/j.rse.2011.09.027>, 2012.
- 1065 Wang, H., Rasch, P. J., Easter, R. C., Singh, B., Zhang, R., Ma, P. L., Qian, Y., Ghan, S. J., and Beagley, N.: Using an explicit emission tagging method in global modeling of source-receptor relationships for black carbon in the Arctic: Variations, sources, and transport pathways, *Journal of Geophysical Research: Atmospheres*, 119, <https://doi.org/10.1002/2014jd022297>, 2014.
- Wang, S., Cohen, J. B., Lin, C. Y., and Deng, W. Z.: Constraining the relationships between aerosol height, aerosol optical depth and total column trace gas measurements using remote sensing and models, *Atmos. Chem. Phys.*, 20, 15401-15426, <https://doi.org/10.5194/acp-20-15401-2020>, 2020.
- 1070 Wang, S., Cohen, J. B., Deng, W., Qin, K., and Guo, J.: Using a new top-down constrained emissions inventory to attribute the previously unknown source of extreme aerosol loadings observed annually in the Monsoon Asia free troposphere, *Earth's Future*, 9, <https://doi.org/10.1029/2021ef002167>, 2021.

删除了: Geophysical Research Letters

- 1080 Wang, S., Su, H., Chen, C., Tao, W., Streets, D. G., Lu, Z., Zheng, B., Carmichael, G. R., Lelieveld, J., Pöschl, U., and Cheng, Y.: Natural gas shortages during the “coal-to-gas” transition in China have caused a large redistribution of air pollution in winter 2017, *Proc. Natl. Acad. Sci.*, 117, 31018-31025, <https://doi.org/10.1073/pnas.2007513117>, 2020b.
- Wei, J., Li, Z., Wang, J., Li, C., Gupta, P., and Cribb, M.: Ground-level gaseous pollutants (NO₂, SO₂, and CO) in China: Daily seamless mapping and spatiotemporal variations, *Atmos. Chem. Phys.*, 23, 1511-1532, <https://doi.org/10.5194/acp-23-1511-2023>, 2023.
- Wu, H., Cai, J., Ren, Q., Shi, C., Zhao, A., and Lyu, Q.: A thermal and chemical fuel pretreatment process for NO_x reduction from cement kiln, *Fuel Process. Technol.*, 210, 106556, <https://doi.org/10.1016/j.fuproc.2020.106556>, 2020.
- 1085 Xing, J., Pleim, J., Mathur, R., Pouliot, G., Hogrefe, C., Gan, C. M., and Wei, C.: Historical gaseous and primary aerosol emissions in the United States from 1990 to 2010, *Atmos. Chem. Phys.*, 13, 7531-7549, <https://doi.org/10.5194/acp-13-7531-2013>, 2013.
- Zavala, M., Herndon, S. C., Slott, R. S., Dunlea, E. J., Marr, L. C., Shorter, J. H., Zahniser, M., Knighton, W. B., Rogers, T., and Kolb, C.: Characterization of on-road vehicle emissions in the Mexico City Metropolitan Area using a mobile laboratory in chase and fleet average measurement modes during the MCMA-2003 field campaign, *Atmos. Chem. Phys.*, 6, 5129-5142, <https://doi.org/10.5194/acp-6-5129-2006>, 2006.
- 1090 Zhang, Q., Zheng, Y., Tong, D., Shao, M., Wang, S., Zhang, Y., Xu, X., Wang, J., He, H., Liu, W., Ding, Y., Lei, Y., Li, J., Wang, Z., Zhang, X., Wang, Y., Cheng, J., Liu, Y., Shi, Q., Yan, L., Geng, G., Hong, C., Li, M., Liu, F., Zheng, B., Cao, J., Ding, A., Gao, J., Fu, Q., Huo, J., Liu, B., Liu, Z., Yang, F., He, K., and Hao, J.: Drivers of improved PM_{2.5} air quality in China from 2013 to 2017, *Proc. Natl. Acad. Sci.*, 116, 24463-24469, <https://doi.org/10.1073/pnas.1907956116>, 2019.
- 1095 Zhang, X. and Schreifels, J.: Continuous emission monitoring systems at power plants in China: Improving SO₂ emission measurement, *Energ. Policy.*, 39, 7432-7438, <https://doi.org/10.1016/j.enpol.2011.09.011>, 2011.
- Zhang, Z., Zang, Z., Cheng, X., Lu, C., Huang, S., Hu, Y., Liang, Y., Jin, L., and Ye, L.: Development of three-dimensional variational data assimilation method of aerosol for the CMAQ model: an application for PM_{2.5} and PM₁₀ forecasts in the Sichuan Basin, *Earth Space Sci.*, 8, <https://doi.org/10.1029/2020EA001614>, 2021.
- 1100 Zheng, B., Zhang, Q., Geng, G., Chen, C., Shi, Q., Cui, M., Lei, Y., and He, K.: Changes in China's anthropogenic emissions and air quality during the COVID-19 pandemic in 2020, *Earth Syst. Sci. Data*, 13, 2895-2907, <https://doi.org/10.5194/essd-13-2895-2021>, 2021.
- Zheng, B., Tong, D., Li, M., Liu, F., Hong, C., Geng, G., Li, H., Li, X., Peng, L., and Qi, J.: Trends in China's anthropogenic emissions since 2010 as the consequence of clean air actions, *Atmos. Chem. Phys.*, 18, 14095-14111, <https://doi.org/10.5194/acp-18-14095-2018>, 2018.
- 1105 Zhou, H., Ma, P., Cheng, M., Zhou, M., and Li, Y.: Effects of temperature and circulating flue gas components on combustion and NO_x emissions characteristics of four types quasi-particles in iron ore sintering process, *ISIJ Int.*, 58, 1650-1658, <https://doi.org/10.2355/isijinternational.ISIJINT-2018-185>, 2018.

删除了: , e2020EA001614

- 1110 Zhou, S., Davy, P. K., Wang, X., Cohen, J. B., Liang, J., Huang, M., Fan, Q., Chen, W., Chang, M., Ancelet, T., and Trompetter, W. J.: High time-resolved elemental components in fine and coarse particles in the Pearl River Delta region of Southern China: Dynamic variations and effects of meteorology, *Sci. Total Environ.*, 572, 634-648, <https://doi.org/10.1016/j.scitotenv.2016.05.194>, 2016.