

Global estimates of 100-year return values of daily precipitation from ensemble weather prediction data

Florian Ruff and Stephan Pfahl

Freie Universität Berlin, Institute of Meteorology, Carl-Heinrich-Becker-Weg 6-10, 12165 Berlin, Germany

Correspondence: Florian Ruff (florian.ruff@met.fu-berlin.de)

Abstract. High-impact river floods are often caused by very extreme precipitation events with return periods of several decades or centuries, and the design of flood protection measures thus relies on reliable estimates of the corresponding return values. However, calculating such return values from observations is associated with large statistical uncertainties due to the limited length of observational time series, [uneven spatial distributions of rain gauges and trends associated with anthropogenic climate change](#). Here, 100-year return values of daily precipitation are estimated on a global grid based on a large data set of model-generated precipitation events from ensemble weather prediction. In this way, the statistical uncertainties of the return values can be substantially reduced compared to observational estimates due to the substantially longer time series. In spite of a general agreement of spatial patterns, the model-generated data set leads to systematically higher return values than the observations in many regions, with statistically significant differences, for instance, over the Amazon, western Africa, the Arabian Peninsula and India. This might be linked to an overestimation of tropical extreme precipitation in the model or an underestimation of very extreme precipitation events in observations, which, if true, would have important consequences for practical water management.

1 Introduction

Extreme precipitation is often associated with river floods, which are one of the most dangerous hazards for society and have caused high socio-economic losses around the globe (Merz et al., 2021; Kron, 2015; Barredo, 2007; Douben, 2006). There are many historical examples of these events such as several flash floods due to a mesoscale convective system in South America in 2022 (Alcântara et al., 2023), an extreme flood event in Central Europe in 2021 (Mohr et al., 2023) and a widespread flooding in Thailand during the monsoon season in 2011 (Gale and Saunders, 2013). River floods typically develop along one of two distinct pathways, either very quickly in form of a flash flood or as a slower increase of the runoff and water levels over several hours. On the one hand, flash floods, which often occur in smaller rivers, are associated with very high precipitation rates in a spatially limited domain, leading to a quickly developing peak discharge (Sene, 2016) that can threaten society along the river route, also downstream of the extreme precipitation event. On the other hand, larger-scale floods in larger rivers with a slower increase of water levels are typically caused by longer-lasting heavy precipitation over a larger area. A multitude of atmospheric drivers can contribute to development of such extreme precipitation events and floods in different regions around the globe, such as convective cells, mesoscale convective systems, monsoonal lows or intense upper-level troughs or cut-off

lows (Alcântara et al., 2023; Mohr et al., 2023; Ruff and Pfahl, 2023; Gaume et al., 2016; Gale and Saunders, 2013). In addition, orographic enhancement of precipitation can contribute to the development of floods in complex terrain.

The increase in global population, especially in high-risk areas, and the rise of the vulnerability of the society increases the risk of flood disasters (Kron, 2015). Moreover, climate simulations project that the frequency and intensity of extreme precipitation events is going to increase in a warmer climate (Pendergrass and Hartmann, 2014; Fischer et al., 2013; O’Gorman and Schneider, 2009), which is also associated with an increasing frequency of intense floods (see e. g. Alfieri et al., 2015). This, and a higher exposure of a growing proportion of the population to floods in a warmer climate, will drastically increase the flood risk on a global scale (Tellman et al., 2021; Alfieri et al., 2017; Jongman et al., 2012). In order to reduce or even prevent flood losses and damages, different kinds of flood protection measures have been developed. More recent extreme events have shown that flood damages have been reduced with the help of flood barrages compared to extreme precipitation that occurred several decades ago (see e. g. Merz et al., 2014; Bissolli et al., 2011), and that there is further potential to minimise flood losses in the future (Jongman et al., 2015). To this end, e. g., for the appropriate construction of dikes, it is crucial to precisely determine the precipitation amounts of potential extreme events that the protection measures should be able to withstand. In practical water management, such events are often denoted as "probable maximum precipitation" (World Meteorological Organization, 2009).

From a statistical point of view, this probable maximum precipitation can be quantified as the precipitation amount associated with extreme events with long return periods, typically on the order of 100 years, through extreme value statistics. To estimate such return values, long precipitation time series are required, which are conventionally obtained from observations. Although there are several observational data sets available with high quality and relatively high temporal and spatial resolution and coverage, this approach is affected by certain limitations (Rajulapati et al., 2020). First, the time series are often shorter than 100 years, which requires extrapolation to determine 100-year or larger return values and increases the associated statistical uncertainties. Second, in addition to the first limitation, there are different extreme value methods (type of distribution, parameter fitting) used to perform this extrapolation, which have to be selected by the users. Third, if global coverage is desired, the uneven spatial distribution of rain gauges requires combining different data sources (e. g., rain gauge and satellite data), which can lead to spatial inhomogeneity in the estimated return values and associated uncertainties. Combining different data sets with diverging representations of extreme precipitation can also lead to inconsistencies on local scales (see e. g. Rajulapati et al., 2020). ~~Third~~ And fourth, precipitation trends, e. g., due to anthropogenic climate change (Fischer et al., 2014), can compromise the extreme value statistics. Accordingly, Rajulapati et al. (2020) have shown that precipitation observations typically do not provide a consistent representation of extreme events and that 100-year return values differ significantly between observational data sets. Due to these limitations, previous studies often focused on extreme events with return periods of much less than 100 years (Rodrigues et al., 2020; Donat et al., 2013), for which observational time series are sufficient, and/or specific regions (Rodrigues et al., 2020; Maraun et al., 2011). Alternatively, model simulations, for instance from weather prediction (Ruff and Pfahl, 2023), seasonal forecasts (Kelder et al., 2020) or climate models (Mizuta and Endo, 2020) can provide long time series that allow for a statistically robust estimate also of 100-year return values. Nevertheless, these model-based estimates may of course suffer from other biases due to the imperfect representation of precipitation processes in the models.

In this study, we explore the possibility to use a model-generated data set from ensemble weather prediction for estimating 100-year return values of daily precipitation on a $1^\circ \times 1^\circ$ grid covering the entire globe, extending our previous analysis that focused on European river catchments (Ruff and Pfahl, 2023). The equivalent length of the weather prediction data set is about 1200 years, that is much longer than the 100-year return period, promoting statistical robustness. Due to the daily accumulation, relatively large spatial scale, and limited model resolution, the studied extreme precipitation events are most relevant for larger-scale river floods and not so much for local flash floods triggered by convective precipitation. The main goal of the study is to compare these model-based 100-year return values to estimates from three different observational data sets and quantify their relative biases and statistical uncertainties. This may provide a basis for using the model-generated data set also for practical estimates of probable maximum precipitation, in particular in data-sparse regions.

The following Sec. 2 describes the ensemble weather prediction data as well as three observational data sets that are applied to evaluate the differences between the model-based and observation-based estimates of 100-year return values. In Sec. 3, the statistical methods to evaluate the ensemble prediction data and determine return values and confidence intervals are explained in detail. The resulting return values, their confidence intervals and differences to observational data sets are presented in Sec. 4. Finally, conclusions and a discussion of the main findings and their limitations are provided in Sec. 5.

2 Data

100-year return values are estimated from a large global data set of daily precipitation events, which is obtained from ensemble weather prediction data. The resulting estimates are compared to observational data sets obtained from rain gauge measurements and satellites for evaluating the differences between a model-based and an observation approach. All data sets are described in the following.

2.1 Ensemble prediction data

In order to generate a large data set of realistic and daily precipitation events, ensemble weather prediction data are used. In this study, the ensemble prediction system (EPS) of the European Center for Medium-Range Weather Forecasts (ECMWF) is accessed for this purpose. The ensemble predictions from the EPS are obtained from the Integrated Forecasting System (IFS), which is a comprehensive earth system model with an atmospheric component from the ECMWF and other community models for certain other components of the earth system (ECMWF, 2023e). More details regarding the IFS and the operational EPS forecasts are described in ECMWF (2023c). An operational weather prediction model is very useful for such an approach as the model is capable of representing daily precipitation events more realistically than e. g. climate models due to a comprehensive comparison to observations (even without a surface precipitation assimilation scheme). Nevertheless, the EPS data may suffer from inter-dependence between the ensemble members, which is investigated in more detail in Sec. 3.1, and from temporal inhomogeneities due to updates of the prediction system. The latter as well as additional limitations of the application of ensemble weather prediction data for the approach in this study are discussed in more detail in Sec. 5.

Started in March 2003, ensemble simulations of the operational weather prediction model are performed twice a day, at 0 and 12 UTC, with forecasting times of at least 10 days. The ensemble contains 51 ensemble members. One member is a controlled run without any perturbations while the other 50 members represent runs with marginally changed initial conditions
95 between each other and with stochastic perturbations of the model physics. This results in 102 simulations per real day. More information about the workflow of the EPS can be found in Molteni et al. (1996).

The analyses in this study are based on daily precipitation sums, which are computed by adding up the large-scale and convective precipitation over 24 hours. From each simulation, the daily precipitation sum just of the 10th forecast day (between forecast hours 216 and 240, same procedure for every initialisation time) is selected, instead of using all forecast days. This
100 approach follows Ruff and Pfahl (2023), who used a daily precipitation data set from the EPS to investigate the atmospheric conditions during extreme precipitation events over Central Europe, and Breivik et al. (2013), who estimated return values of oceanic surface wave heights from EPS data. The basis of the approach is the assumption that, due to the advanced forecast time, the model realisation on the 10th forecast day do not significantly correlate with the conditions in the beginning of each specific simulation. In addition, an inter-dependence of consecutive days, which would also decrease the effective sample size
105 of extreme events in further analyses, is excluded by selecting just one day of this multi-day forecast. Therefore, the different realisations obtained from the ensemble members can be considered as statistically independent from each other. While the simulations of individual ensemble members are highly correlated to each other in the beginning due to very similar initial conditions, this correlation reduces with increasing forecast time. This decrease is particularly large for precipitation, compared to, e. g., geopotential height, due to its high variability in space and time and its dependence on small-scale processes. Both Ruff and Pfahl (2023) and Breivik et al. (2013) performed comprehensive statistical analyses to demonstrate the independence of the ensemble members on the 10th forecast day and to compare the statistics of daily precipitation and wave height to observational data. The data set used here is very similar to the data of Ruff and Pfahl (2023), except that they analysed spatially averaged precipitation time series over Central European river catchments, while this study uses time series on a spatial grid of $1^\circ \times 1^\circ$ spanning the entire globe. Therefore, just a short statistical evaluation of the ensemble prediction data (see Sec. 3.1), adapted to
115 time series at individual grid point, is performed in this study, while we refer to Ruff and Pfahl (2023) for other, more detailed statistical analyses.

The operational model IFS has been updated on a regular basis. Certain technical and physical schemes were changed with each implementation of a new model cycle during the years 2003-2019 in order to continuously improve the forecast skill. However, this may lead to the potential of influencing the simulated precipitation and the upcoming results of this study.
120 Although mainly minor improvements were implemented within each individual model cycle, there are some important updates that include a changed formulation of the humidity analysis (Cycle 26r1, in 2003), improved precipitation forecasts over Europe (Cycle 32r3, in 2007) and improved precipitation forecasts over coastal areas due to changes in cloud physics (Cycle 45r1, in 2018). Details of all model cycle changes are described in ECMWF (2023a) and a full documentation of each model cycle itself is available from ECMWF (2023c). Ruff and Pfahl (2023) have evaluated the influence of these model cycle updates on
125 their daily precipitation time series. They demonstrated a systematic decrease of high precipitation percentiles (99th, 99.9th and 99.99th), which corresponds to extreme precipitation events with large return periods, over Central Europe within the first

five years of the ensemble simulations (2003-2007). On the contrary, since 2008, the amplitude of these percentiles is rather constant, as discussed in more detail in Sec. 3.1. Hence, in order to avoid any temporal inconsistencies within the data set, only the ensemble simulations from the 1st of January 2008 until the 31st of December 2019 are used in this study, following Ruff and Pfahl (2023). The restricted time period of 12 years of forecasts from the EPS archive along with 102 daily simulations, provides a data set with an equivalent length of 1224 years of modelled, but realistic daily precipitation events. The data set is available on a regular lat/lon grid with increasing resolutions over time (higher than $1^\circ \times 1^\circ$ over the entire period) due to changes in the forecast model cycles. For a consistent analysis and comparison to observational data, they are downloaded on a coarser grid (here $1^\circ \times 1^\circ$). During that process, the Meteorological Interpolation and Regridding (MIR) scheme by the ECMWF "upscales" the data by linear interpolation based on a triangular mesh, which is a drawback for the current study, since it does, in general, not conserve area-mean precipitation. More details can be found in (ECMWF, 2023d, f).

2.2 Observational data sets

One observational data set based on rain gauge measurements (REGEN) and two data sets based on a combination of satellite data and rain gauges (CHIRPS and PERSIANN) are used to compare daily 100-year precipitation return values and their confidence intervals from the EPS forecasts to observations. The observational data sets mainly differ in their covered region, the type and amount of observations and the interpolation of the data to a regular lat/lon grid. The observational data sets are first regridded and then the very high return values are determined in order to provide extremes of area-averaged precipitation. This is relevant for large-scale precipitation extremes over large river catchments, on which this study focuses and provides the best comparability to the model-generated EPS data. More detailed descriptions of the observations can be found in the following. The most important information of all observational data sets as used in this study are summarised in Table 1.

2.2.1 REGEN data

The Rainfall Estimates on a Gridded Network (REGEN) is an observational data set for daily precipitation which uses quality controlled rain gauge measurements, spatially interpolated from daily precipitation data of large observational archives such as the Global Historical Climate Network Daily, provided by the National Oceanic and Atmospheric Administration, and the Global Precipitation Climatology Centre, provided by the Deutscher Wetterdienst. More information about this data set are presented in Contractor et al. (2020a). The spatial density of available rain gauges is very different between certain regions. Especially over Africa and Central Asia, the rain gauge density is considerably lower than over North America, Europe and Australia. The reliability of these precipitation observations for specific regions can be evaluated with a quality mask, which takes, e. g., the weather station density and interpolation variation measures into account (Contractor et al., 2020a). Trustworthy areas are mainly located over North America, Europe and Australia as well as over large parts of continental Asia, Brazil and South Africa. While this study only uses the daily precipitation sums (Version 1-2019), other information are additionally available on a global grid such as e. g. the number of rain gauges and the standard deviation of the precipitation sums. The precipitation data are constructed from around 135,000 rain gauges between the 1st of January 1950 and the 31st of

December 2016 while not all stations are available for the entire period and can be used on a regular lat/lon grid with a spatial
160 resolution of $1^\circ \times 1^\circ$ for all global land areas, except for Antarctica.

2.2.2 CHIRPS data

The Climate Hazards Group Infrared Precipitation with Stations (CHIRPS) data archive is a quasi-global, daily precipitation
data set, hosted by the U.S. Geological Survey Earth Resources Observation and Science Center in collaboration with the Santa
Barbara Climate Hazards Group at the University of California. The CHIRPS data result from a combination of quasi-global
165 geostationary thermal infrared satellite observations from two NOAA sources, in situ precipitation observations obtained from
a variety of national and regional meteorological services, the Tropical Rainfall Measuring Mission product from NASA,
a monthly precipitation climatology and atmospheric model rainfall fields from NOAA. A more detailed description of the
development workflow of the CHIRPS data can be found in Funk et al. (2014a). The data are available from the 1st of January
1981 until the ~~31th~~ 31st of December 2021 over land areas on a regular lat/lon grid between 50°S and 50°N . In this study, the
170 CHIRPS data (Version 2.0) with a spatial resolution of $0.25^\circ \times 0.25^\circ$ are selected and averaged over a $1^\circ \times 1^\circ$ box for further
analyses.

2.2.3 PERSIANN data

The Precipitation Estimation from Remotely Sensed Information using Artificial Neural Networks–Climate Data Record
(PERSIANN-CDR), hosted by the Center for Hydrometeorology and Remote Sensing at the University of California, are
175 satellite based, daily precipitation observations. This data sets results from gridded satellite infrared data, obtained from a com-
bination of several international geostationary satellites, in combination with an artificial neural network training using hourly
precipitation data from the National Centers for Environmental Prediction stage IV. Additionally, the monthly product of the
Global Precipitation Climatology Project is used for bias adjustments. Further details on this data set are described in Ashouri
et al. (2015a). The daily precipitation sums of this data set (Version 1) are available from the 1st of January 1983 until the ~~31th~~
180 31st of December 2021 on a regular lat/lon grid between 60°S and 60°N with a spatial resolution of $0.25^\circ \times 0.25^\circ$. In this study,
the data are averaged over a $1^\circ \times 1^\circ$ box and just data over land areas are used for further analyses. Missing values in the data
set appear in cases when satellite data are not available and on dry days when no precipitation occurred (see Supplementary
Fig. S1). As an under-representation of dry days strongly influences the evaluation of quantile distributions of the data (used

| Data sets | Coverage | Time period |
|-----------|---|-------------|
| REGEN | global, land only | 1950-2016 |
| CHIRPS | 50°N - 50°S , land only | 1981-2021 |
| PERSIANN | 60°N - 60°S , land only | 1983-2021 |

Table 1. Summary of the most important details of the observational data sets REGEN, CHIRPS and PERSIANN as used in this study.

for evaluations in Sec. 3.1), all missing values are set to 0 for further analyses, in order to improve the representation of the per-
centiles. However, this also leads to errors over areas that are regularly affected by non-availability of satellite data. Especially
at around 50°N and 70°E this is most dominant, which is why these areas should be taken into account when interpreting the
results from the PERSIANN observations.

3 Methodology

In this section, statistical analyses of the suitability of the EPS data for determining 100-year precipitation return values on a
global grid are presented. Subsequently, the method to determine the return values and their confidence intervals is described.

3.1 Statistical evaluation of the ensemble prediction data

For the determination of 100-year return values of daily precipitation on a global scale, the EPS data at the 10th forecast day
are used as a large climatological data set. This implies that the data set can be considered as a combination of realistic and
independent realisations of daily precipitation in order to suitably apply extreme value statistics to this data set. Therefore,
we evaluate if (1) the ensemble members can be considered as independent from each other, (2) each ensemble member
properly represents the statistical distribution of precipitation compared to observations and (3) no significant trend of the high
precipitation percentiles can be identified over time, following Ruff and Pfahl (2023) and Breivik et al. (2013). These criteria
are statistically evaluated and discussed in the following based on time series of daily precipitation sums on the 10th day of
each forecast and at each global grid point. This spatial coverage is the main difference between this study and Ruff and Pfahl
(2023), who analysed time series of spatially averaged precipitation in several Central European river catchments.

Beginning with the first criterion, the daily precipitation at the 10th forecast day of each individual ensemble member is
investigated with regard to its independence to each other. For this, Ruff and Pfahl (2023) analysed the statistical distribution of
Spearman correlation coefficients between all possible ensemble member combinations (5151 in total) for time series of both
all daily precipitation sums and their annual maxima (which are used to determine the 100-year return values, see Sec. 3.2).
They showed that the time series based on all daily events are weakly, but still significantly, correlated between the ensemble
members (mean correlation around 0.19, similar for all river catchments). Time series of yearly maxima of daily precipitation
sums are generally very weakly correlated as well (mean around 0), but there is almost no significant correlation identified,
supporting the assumption that daily extreme precipitation is independent between ensemble members. Here, this correlation
analysis is expanded to the global grid, leading to an analysis of multiple correlations. In order to determine the statistical
significance of the correlation coefficients in such a multi-test framework, the False Discovery Rate (FDR) test of Benjamini
and Hochberg (1995), as described in Ventura et al. (2004), is applied to all p-values of the correlation coefficients from each
combination of ensemble members at a certain grid points. Figure 1 shows the number of statistically significant correlation
coefficients from the FDR test for time series of annual maximum daily precipitation at each global land grid point, except for
Antarctica. For most of the grid points, (almost) no significant correlations are found, especially poleward of 20°N and 20°S,
supporting the hypothesis that annual maximum precipitation events are independent between ensemble members on a global




Figure 1. Number of statistical significant correlation coefficients per grid point, obtained from the FDR tests of Benjamini and Hochberg (1995), as described in Ventura et al. (2004), applied to multiple (5152) p values associated with the Pearson correlations between yearly maxima time series of individual ensemble members. Note the logarithmic colour scale.

scale. However, there are certain areas over the tropical South America and Africa as well as over the Maritime Continent, where FDR tests show high numbers of significant correlation coefficients. In these regions, extreme precipitation events in the individual ensemble members show a certain dependence, and the EPS data set cannot be considered as equivalent to a time series of 1224 years in the analysis of 100-year return values. A likely reason for this inter-dependence is the influence of internal climate modes with relatively long time scales, such as the El Niño-Southern Oscillation, on tropical precipitation events, which can lead to a synchronisation of annual maxima between ensemble members.

To evaluate the statistical distribution of daily precipitation, Ruff and Pfahl (2023) compared quantiles from the EPS data to three observational data sets and found a good agreement in the Central European river catchments. For a similar analysis on the global scale with a more pronounced focus on more intense daily precipitation, we select the 99th and 99.99th percentiles at each grid point. Note that all days of a time series, including dry days, are considered for determining these percentiles (Pfleiderer et al., 2019). Figure 2 shows the 99th percentile (ca. 0.3-year event) and the 99.99th percentile (ca. 30-year event) from the EPS data (taking all members together) in (a-b) and the differences to the observational REGEN data in (c-f). High values of the 99th percentile of daily precipitation can generally be found over the tropics and subtropics (see Fig. 2a), especially in the area of the Intertropical Convergence Zone (ITCZ), as well as near complex terrain. Over the extratropical continents, the 99th percentile is slightly lower. The differences to the REGEN data set in Fig. 2c,e reveal that EPS data show a relatively good agreement over regions where the REGEN observations are trustworthy (black dots). There is generally a slight overestimation of the percentiles in the EPS data, except for some parts of Europe, Mexico and India, where REGEN observations indicate higher 99th percentiles. Larger differences can especially be found over South America, Africa and the Maritime Continent, where the REGEN observations are not considered trustworthy, with values of up to +500% or -80%. The absolute and relative




Figure 2. Global distribution of the (a) 99th and (b) 99.99th percentile of daily precipitation from the EPS data and their (c,d) absolute and (e,f) relative difference to the REGEN observations. Stippling in (c-f) shows where the REGEN observations are trustworthy as explained in 2.2.1. Note the non-linear colour scales.

235 differences to the CHIRPS and PERSIANN observations in general show similar results (see Supplementary Fig. S2a,c,e,g). However, there are some large differences also between the observational data sets, e. g., with lower 99th percentiles over North America, China and the Maritime Continent in PERSIANN observations, lower 99th percentiles over central Asia in CHIRPS observations and large differences of the 99th percentile over large parts of Africa between all observational data sets. Similar results are obtained for the 99.99th percentile. Even higher precipitation values from the EPS data are found in

240 the tropics (see Fig. 2b), while there is still a relatively good agreement over areas with trustworthy REGEN observations (see Fig. 2d,f). However, the differences to the REGEN observations generally increase with higher percentiles. This is also evident for differences of the 99.99th percentile to the CHIRPS and PERSIANN observations (see Supplementary Fig. S2b,d,f,h), while still some large differences remain over the previously mentioned regions between the observational data sets. In summary, the percentiles calculated from EPS data show a relatively good agreement with observational estimates over North America
245 and large parts of Europe, central Asia, Australia and South America. With an increasing level of percentile, the differences to observational data sets increase, which might be due to an overestimation of the EPS data or biases in the representation of very high percentiles in the observations due to a limited length of time series. Still, a bias correction of the EPS data is not considered in this study as the observational data sets show large uncertainties over several parts of the globe. Ruff and Pfahl (2023) also compared the precipitation statistics between each individual ensemble member, but as they found no systematic
250 differences, this analysis is not repeated here.

Stationarity of the time series is important for applying the extreme value analysis described in Sec. 3.2. To evaluate the stationarity of the EPS data, Ruff and Pfahl (2023) applied the Mann-Kendall test to several high percentiles and the yearly number of 100-year precipitation events for the years 2008 to 2019 and also compared the occurrences of such events to a Poisson distribution of independent events with a constant mean rate. They did not find an indication of temporal non-
255 stationarity in this period of 12 years. Here, the same Mann-Kendall test is applied to the yearly maximum of daily precipitation at all global land grid points. In order to evaluate the statistical significance of these multiple tests, again the FDR test is applied (see Supplementary Fig. S3) as introduced earlier. No significant trend is found for the EPS data. For the observational data sets, also due to the longer time series covering 39 years or more, some areas are associated with significant trends, rather evenly distributed over the globe (for REGEN, 6.7% of grid points with significant trends), over Central Africa and Central
260 Asia (for CHIRPS, 0.7% of grid points with significant trends) or India and central Asia (for PERSIANN, 1.2% of grid points with significant trends). Nevertheless, trends are still not significant over the most part of the globe. In order to use a consistent methodology for all data sets and locations, we thus make the stationarity assumption also for the observational time series.

In summary, our analyses show that, in most regions, daily precipitation obtained from different members of the ECMWF EPS can be considered as statistically independent. Exceptions are some areas over tropical regions of South America and
265 Africa as well as over the Maritime continent (see again Fig. 1). Additionally, the model data represent different quantiles of daily precipitation quite well in comparison to three observational data (see Figs. 2 and 2S2), again with the exception of a few regions mostly in the tropics and larger differences for very high percentiles. Finally, there is no indication of non-stationarity in the data set over the time frame analysed here. Thus, we consider the EPS data to be suitable for a global analysis of 100-year return values of daily precipitation.

270 3.2 Determination of return values and confidence intervals

To determine 100-year return values of daily precipitation and their confidence intervals at each grid point, the daily precipitation sums from all ensemble members are used to build long time series. For this investigation, extreme value statistics (see

275 Coles et al., 2001) are applied in order to fit a generalised extreme value distribution (GEV) to a selected sample of block maxima, using the maximum likelihood approach. This sample of block maxima is here selected from yearly blocks of daily precipitation. Such an approach leads to 1224 block maxima, to which the GEV can be fitted at each grid point. The best fit of the GEV is accomplished by estimating the location (μ), scale (σ) and shape (ξ) parameters from the maximum likelihood approach. Following Stephenson (2002), the return value v can be computed from these estimated parameters by the following equation:

$$v = \mu + \sigma \cdot \frac{(x^\xi - 1)}{\xi} \quad (1)$$

$$x = \frac{-1}{\log(1 - \frac{1}{p})}$$

280

in which a certain yearly return period is described by p . The confidence intervals of the return values are computed by the bootstrap resampling method (see Coles et al., 2001). Taking the original set of block maxima, a new set of maxima is drawn with replacement. Then, the previously explained approach of fitting a GEV to a selected sample of block maxima is repeated with the new set of block maxima and the return value is again determined from Eq. (1). As each procedure leads to a slightly 285 changed return value, the uncertainty can be evaluated by repeating this process several times. Here, it is repeated 1000 times. From the resulting 1000 return values, the 0.025 and 0.975 quantiles are considered to be the confidence intervals of the return value from the original sample of block maxima. This process to determine return values and their confidence intervals from EPS data is also used for all the observational data sets.

At some grid points with very low precipitation amounts during the entire year (e. g., over the Sahara Desert), the best fit 290 of the GEV yields very high estimates of the shape parameter ξ (up to 3). This results in extraordinary high return values compared to neighbouring grid points with low return values. Papalexiou and Koutsoyiannis (2013) analysed estimated shape parameters from GEV fits for over 15,000 globally distributed observational records, using yearly maxima of daily precipitation as block maxima as well. Even for rather short time series of at most 10 years, which are often associated with higher shape parameters than longer time series, the shape parameters all lie between -0.6 and 0.6, independent of their location. However, 295 almost no time series of their observational data set are located in very dry regions such as the Sahara Desert. In order to prevent unrealistic high or low return values but still allow shape parameters outside the range of -0.6 and 0.6 for areas that are typically not covered by observations, grid points with estimated shape parameters above 1 and below -1 and scale parameters above 70 and below -70 are excluded from the analyses in this study.

4 Results

300 In this section, the estimated 100-year return values of daily precipitation and their confidence intervals are presented. Estimates from the EPS data are compared to the observational estimates based on REGEN, CHIRPS and PERSIANN.




Figure 3. (a) Global distribution of 100-year return values of daily precipitation, estimated from EPS data, and their difference to the observational estimates from (b) REGEN, (c) CHIRPS and (d) PERSIANN. Dark grey shading indicates grid points for which the GEV parameters are outside the allowed range and thus no return values can be estimated. Stippling in (b-d) shows where the confidence interval of the EPS data overlaps with the confidence interval from the specific observational data set. Note the non-linear colour scales.

4.1 Return values

Global estimates of 100-year return values of daily precipitation from the EPS data set are shown in Fig. 3a. Generally, higher return values can be found in tropics and parts of the subtropics with values between 400 mm/d and 600 mm/d over most parts of India and near the Himalaya as well as values around 300 mm/d for tropical regions, while return values decrease towards the poles. A 100-year event in the mid-latitudes is typically associated with values of 50-100 mm/d, while usually dry regions such as the Sahara Desert and the Taklamakan Desert are associated with very low return values of around 30 mm/d. Still, such extreme events over these very dry regions typically lead to an exceptional exceedance of the annual precipitation amount, for the two latter regions with values of about 300-700% (see Supplementary Fig. S4).

A comparison with the observational data set REGEN shows generally higher EPS estimates over large parts of the globe and very large differences in several tropical and subtropical regions (Fig. 3b). The 100-year return values from the EPS data are clearly higher, e. g., over India (by 200-400 mm/d), western Africa (by 100-300 mm/d) and over the Amazon (by 50-150 mm/d), with large differences also existing for much lower return periods at individual locations (see Supplementary

Fig. S5c,d). Additionally, the confidence intervals of the EPS data do not overlap with the REGEN data in these regions (even
315 for much lower return periods, see Supplementary Fig. S5c,d), the differences are thus statistically significant. Over other parts
of South America, the southern half of Africa and Australia, the EPS estimates are about 50 mm/d higher, but the confidence
intervals overlap in parts of these areas. The mid-latitudes do not show large differences in the estimated return values (similar
for lower return periods at individual locations, see Supplementary Fig. S5a,b). However, over parts of Chile, the Abyssinia
Plateau in East Africa and over some coastal areas in South-East Asia, the EPS data are associated with lower return values
320 than the REGEN data (mostly no overlap of confidence intervals). Overall, the confidence intervals of EPS and REGEN return
value estimates overlap at 55.6% of the grid boxes. The differences of EPS return values to the CHIRPS and PERSIANN
observational estimates (Figs. 3c,d) are very similar to each other. The confidence intervals of EPS and CHIRPS overlap at
48.6% of the grid boxes, and for EPS and PERSIANN this overlap is 51.6%. Furthermore, the CHIRPS and PERSIANN
results are mostly consistent with the differences to the REGEN estimates described above. However, the strongly negative
325 differences over Chile and the Abyssinia Plateau do not occur for the CHIRPS and PERSIANN data. Additionally, the positive
differences over India, South-East Asia and Australia are even slightly larger. It should be kept in mind here that, especially
over tropical parts of South America and the Maritime Continent, the EPS estimates are less trustworthy than in other regions
due to methodological issues associated with inter-dependence between the ensemble members (see again Sec. 3.1).

4.2 Confidence intervals

330 Confidence intervals (CIs) on a 95% level are determined for each return value estimate. To compare the confidence interval
ranges (that is, the difference between upper and lower bounds) between data sets, on the one hand, in Fig. 4a,b they are shown
relative to the corresponding return value estimate, as higher return values are typically associated with larger confidence
intervals. In this way, the relative uncertainty of the 100-year return values is quantified. On the other hand, Fig. 4c,d shows the
absolute confidence interval ranges, quantifying the absolute uncertainties.

335 The largest relative uncertainties are found in tropical and subtropical regions. Especially over the Sahara Desert and the
southeastern part of the Arabian Peninsula, the relative CIs range from 30% to 65%, while poleward of 30°N and 30°S, the
range of the relative CIs lies between 10% and 20% (see Fig. 4a). This pattern of relative CIs strongly correlates with the ratio of
a 100-year event to the annual precipitation amount in Supplementary Fig. S4. Over regions where a 100-year event represents
a multiple of the annual precipitation amount, the daily precipitation distribution is typically associated with a **long and thin**
340 **rather short and thick** "tail". An estimation of very high percentiles from such a distribution is more difficult and leads to higher
uncertainties. As mentioned before, higher return values are associated with higher CIs, as shown by the absolute values in
Fig. 4c. The tropics and several subtropical areas have the highest absolute CI ranges with typical values of 50-150 mm/d.
Absolute CIs over the mid-latitudes typically lie around 10 mm/d.

The differences of the relative CI ranges between the EPS and REGEN data sets are shown in Fig. 4b. Over almost all
345 continental areas, the relative uncertainties are reduced in the EPS data compared to REGEN, with typical differences on the
order of 50-100 percentage points (p.p.). Over specific regions such as the west coast of South America, the Sahara Desert and




Figure 4. (Left) Global distribution of confidence interval ranges on a 95% level of EPS data and (right) their differences to the observational estimates from REGEN for (a,b) the relative range, relative to the associated return values, and (c,d) the absolute range of these confidence intervals. Dark grey shading indicates grid points for which the GEV parameters are outside the allowed range and thus no return values can be estimated. Stippling in (b,d) shows where the confidence interval of the EPS data overlaps with the confidence interval from the REGEN observations. Note the non-linear colour scales.

the Arabian Peninsula, this decrease is even larger (-200 p.p. to -800 p.p.), while the CIs mainly overlap with each other in these regions. A slight increase in the relative uncertainty by around 10 p.p. is found for some grid points over the Amazon, West Africa, India and China. In terms of absolute CI ranges (Fig. 4d), the uncertainty is often reduced in the EPS data compared to REGEN by up to 300 mm/d over the tropics and subtropics and around 20 mm/d over the mid-latitudes. Note that this corresponds to a reduction by at least a factor of two, that is a substantially reduced uncertainty, in most regions. There are a few more extended (compared to the relative CI ranges) areas of larger absolute uncertainties in the EPS data over the Amazon, West Africa and India with increases of 20-100 mm/d. These are also regions where the CIs do not overlap.

Similar results are found for both relative and absolute uncertainties taking the CHIRPS data for comparison. The relative CI range is smaller in the EPS data over almost all regions (see Supplementary Fig. S6a), and the magnitude of this reduction is typically even larger (around -100 p.p.) than for REGEN, except for the Sahara Desert. Differences in the absolute uncertainty compared to CHIRPS are again similar as for REGEN, but with a more enhanced and broader increase in the southeast of the Arabian Peninsula of up to 300 mm/d (see Supplementary Fig. S6c). Finally, the uncertainty in EPS is also reduced compared




Figure 5. Boxplot distribution of 1000 return value estimates of a 100-year event obtained from subsampling the EPS block maxima with different sample sizes for a single grid point over Berlin, Germany (52°N , 13°E). The boxplot whiskers represent the data point, which is no more than 1.5 times the interquartile range from the box. The red bars indicate the 95% confidence interval. The blue line indicates the 100-year estimate from the largest sample size of 1200.

to PERSIANN, with similar patterns over the continents as for REGEN and CHIRPS (see Supplementary Fig. S6b,d). Note
 360 that a co-occurring increase of absolute uncertainty and decrease in relative uncertainty of return value estimates is linked to substantially higher return value estimates in the EPS data.

The main advantage of the model-generated EPS data compared to the observational data sets is the extraordinary length of the time series that is used for the 100-year return value estimate. However, it is not directly evident how large this effect of the length of the time series is for the return value estimate and its uncertainty. In order to quantify this effect, the determination
 365 of the 100-year return value and its confidence intervals is repeated for different sample sizes of the yearly EPS block maxima, starting with a sample size of 60 as this is roughly the length of the observational data sets. The results show that the estimated return values do not depend strongly on the sample size and increase only slightly with increasing sample sizes over all parts of the globe (not shown). Figure 5 shows the distribution of 1000 return value estimates and its confidence intervals for sample sizes of 60, 120, 240, 480 and 1200 for one grid point over Berlin, Germany (52°N , 13°E). As mentioned previously, the
 370 increase of the return value estimate with higher sample sizes is rather small. This indicates that the systematic overestimation of the EPS return values in many parts of the world are likely due to biases of the model or the observational data sets and not caused by the systematic differences in sample size. However, there is a clear reduction of the uncertainty of the 100-year

return value with a larger sample size. Thus, the sampling effect can clearly explain the reduced uncertainty in the EPS data set.

375 **5 Discussion and conclusion**

The aim of this study has been to determine 100-year return values of daily precipitation and their confidence intervals (CIs) on a global scale from a large data set of model-generated events and to evaluate the differences to three observational data sets (REGEN, CHIRPS and PERSIANN). A quantification of such extreme return values is crucial for properly setting up flood protection measures and also because such extreme events are expected to occur more frequently in a warmer climate. 380 The large set of simulated global daily precipitation fields has been obtained from operational ensemble weather prediction data by the ECMWF, following the approach of Breivik et al. (2013) and Ruff and Pfahl (2023). Our statistical analyses show that, when using the 10th forecast days from these simulations, annual precipitation maxima are independent between the different ensemble members in most regions, except for some tropical areas over South America and Africa as well as over the Maritime Continent. There, extreme precipitation events are correlated between different ensemble members, so that 385 these events can not be considered as independent. This reduces the effective sample size of events for the analyses and decreases the length of the precipitation time series, leading to less trustworthy results of 100-year return values. Biases in the climatological distribution of extreme precipitation with respect to the observational data sets, evaluated through the 99th and 99.99th percentile, are relatively small in most regions of North and South America, Europe, central Asia and Australia, but larger for other regions and generally increase for very high percentiles. Additionally, there is no significant trend in the 390 occurrence of intense precipitation over the 12 years at all grid points, and the EPS data can thus be considered as stationary in time. With the help of extreme value statistics, 100-year return values and the associated confidence intervals on a 95% level are determined and compared to estimates from the observational data sets.

Based on these EPS data, the largest 100-year return values of daily precipitation, which are computed over $1^\circ \times 1^\circ$ grid boxes, occur in the tropics and subtropics with a maximum of up to 600 mm/d (absolute CI range of 150 mm/d) over India 395 and typical values around 300 mm/d (absolute CI range around 100 mm/d) for most of the other tropical and subtropical regions. The return values decrease towards the poles with values of 50-100 mm/d (absolute CI range of 10 mm/d) over the mid-latitudes. Such an extreme event would typically amount to about 10-20% of annual mean precipitation (20-50% for Australia), but in dry regions such as the Sahara Desert the estimated 100-year return value even exceed the annual mean precipitation by a factor of up to 7. The 100-year return values from the EPS data are in general agreement with other studies of 400 multi-year precipitation extremes. For instance, Rodrigues et al. (2020) determined 10-year return values for Brazil and found slightly lower values and a local maximum of 200 mm/d further towards the east coast. Gründemann et al. (2023) studied 100-year return values over global land areas based on several statistical approaches and a data set obtained from a combination of satellite observations, reanalyses and gauge data. They found similar spatial patterns as documented here, but lower return values over, for instance, the Sahel, the east coast of the Arabian Peninsula and parts of India. Also the comparison with the 405 observational data sets in this study indicates systematically higher return values in the EPS data set over most of the globe.

In many regions, in particular in the extratropics, the confidence intervals of EPS and observational estimates still overlap. Larger, also statistically significant differences are obtained in some tropical regions where the EPS method is less robust due to inter-dependence of ensemble members and/or general biases in the precipitation climatology, but also over other areas where the data set performs well in our statistical evaluation, such as parts of northeastern South America, western tropical Africa, India and eastern China. This might be due to model deficiencies, e. g., in the representation of convective extreme precipitation in the tropics. However, it may also point to a systematic underestimation of very extreme daily precipitation events in observations. Such a potential under-estimation might have important consequences for practical applications and should be considered in estimates of "probable maximum precipitation" for designing flood protection measures.

The relative uncertainty of the 100-year return values is quantified through relative confidence interval ranges, which typically lie within 10-30% for most of the regions, but can be higher than 50% over the Sahara Desert and the Arabian Peninsula. An important result is that these relative uncertainties, and in most regions also the absolute CI ranges, are substantially reduced in the EPS data compared to observational estimates. This reduction is typically on the order of 50-100 p.p., but can locally amount to up to 500 p.p., e. g., over the Sahara Desert and the Arabian Peninsula.

The Observation-based estimates of very high percentiles of daily precipitation contain uncertainties due to the finite length of observational time series, the not unambiguous choice of the statistical method to estimate extreme values, spatial inhomogeneities from e. g. an uneven distribution of rain gauges as well as due to trends from anthropogenic climate change. The two latter limitations are reduced by using a weather prediction model for a rather short period of time (12 years). Additionally, the systematic and substantial reduction of statistical uncertainty in this study is due to the much longer time series and is the main advantage of the model-generated EPS data, compared to the observational data sets. However, we do not claim that these model data is superior to observations in any other way but rather show the advantages and disadvantages of both model-generated EPS and observational data when estimating 100-year daily precipitation events on a global scale and its uncertainties.

Our approach to estimate 100-year precipitation return values from EPS data has several limitations, as also discussed by Ruff and Pfahl (2023). First, the model-generated precipitation data are affected by model biases and the imperfect representation of specific processes in the model, especially over the tropics. Such biases are assumed to be particularly large for small-scale, convective precipitation events, which is one reason why we focus on larger-scale events on a $1^\circ \times 1^\circ$ grid. Second, the time span of the forecast data is limited to 12 years (2008-2019) which comes along with a limited sampling of large-scale boundary conditions. Therefore, not the entire range of (multi-)decadal variability of the climate system is reproduced in the data set. Third, there is a certain influence of anthropogenic forcing in the data set in specific regions, as mentioned in Sec. 3.1, which can lead to temporal inhomogeneities. However, our trend analyses shows that this is mainly restricted to very few areas. Fourth, the method does not work well in regions where the ensemble members are not independent from each other, which is mainly the tropical regions of South America and Africa as well as the Maritime Continent.

In future research, the approach used here may be applied for events with different return periods and also to other weather prediction ensemble data set, which may help in clarifying the reasons for the systematically higher return values in the EPS

440 data compared to observations. Finally, large initial condition ensemble simulations with climate models may be used to investigate the influence of climate warming on 100-year precipitation events.

Data availability. The global estimates of 100-year return values and their confidence intervals on a $1^\circ \times 1^\circ$ lat/lon grid from the EPS data and the REGEN, CHIRPS and PERSIANN observations, presented in this study, can be downloaded from <http://dx.doi.org/10.17169/refubium-39650>. The operational ensemble forecast data from the ECMWF can be downloaded from <https://apps.ecmwf.int/archive-catalogue/?type=cf&class=od&stream=enfo&expver=1> (ECMWF, 2023b) for the control run and from <https://apps.ecmwf.int/archive-catalogue/?type=pf&class=od&stream=enfo&expver=1> (ECMWF, 2023g) for the perturbed runs. The user's affiliation needs to belong to an ECMWF member state. The observational data sets are freely accessible from <https://doi.org/10.25914/5ca4c380b0d44> (Contractor et al., 2020b) for REGEN, https://data.chc.ucsb.edu/products/CHIRPS-2.0/global_daily/netcdf/p25/ (Funk et al., 2014b) for CHIRPS and <https://www.ncei.noaa.gov/data/precipitation-persiann/access/> (Ashouri et al., 2015b) for PERSIANN.

450 *Author contributions.* Florian Ruff performed the analysis, produced the figures and drafted the manuscript. Both authors designed the study, discussed results and edited the manuscript.

Competing interests. The authors have no competing interests to declare.

Acknowledgements. The German Meteorological Service DWD and ECMWF are acknowledged for providing the operational IFS model data. This work used resources of the Deutsches Klimarechenzentrum (DKRZ), which is granted by its Scientific Steering Committee (WLA) under project ID bb1152. We are grateful to Felix Fauer and Henning Rust (both Freie Universität Berlin) for their helpful comments on the statistical approaches, and to all colleagues of the ClimXtreme Module D for their technical assistance.

Financial support. This study has been funded by the German Ministry of Education and Research (Bundesministerium für Bildung und Forschung, BMBF) in its strategy Research for Sustainability (FONA) in the framework of the ClimXtreme programme, sub-project A2 (MExRain, grant number 01LP1901C).

460 References

- Alcântara, E., Marengo, J. A., Mantovani, J., Londe, L. R., San, R. L. Y., Park, E., Lin, Y. N., Wang, J., Mendes, T., Cunha, A. P., Pampuch, L., Seluchi, M., Simões, S., Cuartas, L. A., Goncalves, D., Massi, K., Alvalá, R., Moraes, O., Filho, C. S., Mendes, R., and Nobre, C.: Deadly disasters in Southeastern South America: flash floods and landslides of February 2022 in Petrópolis, Rio de Janeiro, *Natural Hazards and Earth System Sciences*, 23, 1157–1175, <https://doi.org/10.5194/nhess-23-1157-2023>, 2023.
- 465 Alfieri, L., Burek, P., Feyen, L., and Forzieri, G.: Global warming increases the frequency of river floods in Europe, *Hydrology and Earth System Sciences*, 19, 2247–2260, <https://doi.org/10.5194/hess-19-2247-2015>, 2015.
- Alfieri, L., Bisselink, B., Dottori, F., Naumann, G., de Roo, A., Salamon, P., Wyser, K., and Feyen, L.: Global projections of river flood risk in a warmer world, *Earth's Future*, 5, 171–182, <https://doi.org/10.1002/2016EF000485>, 2017.
- Ashouri, H., Hsu, K.-L., Sorooshian, S., Braithwaite, D. K., Knapp, K. R., Cecil, L. D., Nelson, B. R., and Prat, O. P.: PERSIANN-CDR: 470 Daily precipitation climate data record from multisatellite observations for hydrological and climate studies, *Bulletin of the American Meteorological Society*, 96, 69–83, <https://doi.org/10.1175/BAMS-D-13-00068.1>, 2015a.
- Ashouri, H., Hsu, K.-L., Sorooshian, S., Braithwaite, D. K., Knapp, K. R., Cecil, L. D., Nelson, B. R., and Prat, O. P.: Index of /data/precipitation-persiann/access, National Centers for Environmental Information [data set], <https://www.ncei.noaa.gov/data/precipitation-persiann/access/>, last access: 15 May 2023, 2015b.
- 475 Barredo, J. I.: Major flood disasters in Europe: 1950–2005, *Natural Hazards*, 42, 125–148, <https://doi.org/10.1007/s11069-006-9065-2>, 2007.
- Benjamini, Y. and Hochberg, Y.: Controlling the false discovery rate: a practical and powerful approach to multiple testing, *Journal of the Royal Statistical Society*, 57, 289–300, <https://doi.org/10.1111/j.2517-6161.1995.tb02031.x>, 1995.
- Bissolli, P., Friedrich, K., Rapp, J., and Ziese, M.: Flooding in eastern central Europe in May 2010—reasons, evolution and climatological assessment, *Weather*, 66, 147–153, <https://doi.org/10.1002/wea.759>, 2011.
- 480 Breivik, Ø., Aarnes, O. J., Bidlot, J.-R., Carrasco, A., and Saetra, Ø.: Wave extremes in the northeast Atlantic from ensemble forecasts, *Journal of Climate*, 26, 7525–7540, <https://doi.org/10.1175/JCLI-D-12-00738.1>, 2013.
- Coles, S., Bawa, J., Trenner, L., and Dorazio, P.: An introduction to statistical modeling of extreme values, vol. 208, Springer, London, <https://doi.org/10.1007/978-1-4471-3675-0>, 2001.
- Contractor, S., Donat, M. G., Alexander, L. V., Ziese, M., Meyer-Christoffer, A., Schneider, U., Rustemeier, E., Becker, A., Durre, I., and 485 Vose, R. S.: Rainfall Estimates on a Gridded Network (REGEN)—a global land-based gridded dataset of daily precipitation from 1950 to 2016, *Hydrology and Earth System Sciences*, 24, 919–943, <https://doi.org/10.5194/hess-24-919-2020>, 2020a.
- Contractor, S., Donat, M. G., Alexander, L. V., Ziese, M., Meyer-Christoffer, A., Schneider, U., Rustemeier, E., Becker, A., Durre, I., and Vose, R. S.: Rainfall Estimates on a Gridded Network based on all station data v1-2019, National Computational Infrastructure [data set], <https://doi.org/10.25914/5ca4c380b0d44>, last access: 15 May 2023, 2020b.
- 490 Donat, M. G., Alexander, L. V., Yang, H., Durre, I., Vose, R., Dunn, R. J. H., Willett, K. M., Aguilar, E., Brunet, M., Caesar, J., Hewitson, B., Jack, C., Klein Tank, A. M. G., Kruger, A. C., Marengo, J., Peterson, T. C., Renom, M., Oria Rojas, C., Rusticucci, M., Salinger, J., Elayah, A. S., Sekele, S. S., Srivastava, A. K., Trewin, B., Villarroel, C., Vincent, L. A., Zhai, P., Zhang, X., and Kitching, S.: Updated analyses of temperature and precipitation extreme indices since the beginning of the twentieth century: The HadEX2 dataset, *Journal of Geophysical Research*, 118, 2098–2118, <https://doi.org/10.1002/jgrd.50150>, 2013.
- 495 Douben, K.-J.: Characteristics of river floods and flooding: a global overview, 1985–2003, *Irrigation and Drainage*, 55, S9–S21, <https://doi.org/10.1002/ird.239>, 2006.

- ECMWF: Changes in ECMWF model, <https://www.ecmwf.int/en/forecasts/documentation-and-support/changes-ecmwf-model>, last accessed: 10 February 2023, 2023a.
- ECMWF: Archive Catalogue - Control forecast, ECMWF [data set], <https://apps.ecmwf.int/archive-catalogue/?type=cf&class=od&stream=enfo&expver=1>, last access: 23 February 2023, 2023b.
- ECMWF: IFS documentation, <https://www.ecmwf.int/en/publications/ifs-documentation>, last accessed: 10 February 2023, 2023c.
- ECMWF: MARS interpolation with MIR, <https://www.ecmwf.int/en/newsletter/152/computing/new-ecmwf-interpolation-package-mir>, last accessed: 08 April 2024, 2023d.
- ECMWF: Modelling and Prediction, <https://www.ecmwf.int/en/research/modelling-and-prediction>, last accessed: 10 February 2023, 2023e.
- 505 ECMWF: The new ECMWF interpolation package MIR, <https://www.ecmwf.int/en/newsletter/152/computing/new-ecmwf-interpolation-package-mir>, last accessed: 11 July 2023, 2023f.
- ECMWF: Archive Catalogue - Perturbed forecast, ECMWF [data set], <https://apps.ecmwf.int/archive-catalogue/?type=pf&class=od&stream=enfo&expver=1>, last access: 23 February 2023, 2023g.
- Fischer, E. M., Beyerle, U., and Knutti, R.: Robust spatially aggregated projections of climate extremes, *Nature Climate Change*, 3, 1033–1038, <https://doi.org/10.1038/nclimate2051>, 2013.
- Fischer, E. M., Sedláček, J., Hawkins, E., and Knutti, R.: Models agree on forced response pattern of precipitation and temperature extremes, *Geophysical Research Letters*, 41, 8554–8562, 2014.
- Funk, C. C., Peterson, P. J., Landsfeld, M. F., Pedreros, D. H., Verdin, J. P., Rowland, J. D., Romero, B. E., Husak, G. J., Michaelsen, J. C., and Verdin, A. P.: A quasi-global precipitation time series for drought monitoring, *U.S. Geological Survey Data Series*, 832, 1–12, <https://doi.org/10.3133/ds832>, 2014a.
- 515 Funk, C. C., Peterson, P. J., Landsfeld, M. F., Pedreros, D. H., Verdin, J. P., Rowland, J. D., Romero, B. E., Husak, G. J., Michaelsen, J. C., and Verdin, A. P.: Index of /products/CHIRPS-2.0/global_daily/netcdf/p25/, University of California at Santa Barbara [data set], https://data.chc.ucsb.edu/products/CHIRPS-2.0/global_daily/netcdf/p25/, last access: 15 May 2023, 2014b.
- Gale, E. L. and Saunders, M. A.: The 2011 Thailand flood: climate causes and return periods, *Weather*, 68, 233–237, <https://doi.org/10.1002/wea.2133>, 2013.
- 520 Gaume, E., Borga, M., Llassat, M. C., Maouche, S., Lang, M., and Diakakis, M.: Mediterranean extreme floods and flash floods, in: *The Mediterranean Region under Climate Change. A Scientific Update*, pp. 133–144, IRD Editions, <https://hal.science/hal-01465740>, 2016.
- Gründemann, G. J., Zorzetto, E., Beck, H. E., Schleiss, M., Van de Giesen, N., Marani, M., and van der Ent, R. J.: Extreme precipitation return levels for multiple durations on a global scale, *Journal of Hydrology*, 621, <https://doi.org/10.1016/j.jhydrol.2023.129558>, 2023.
- 525 Jongman, B., Ward, P. J., and Aerts, J. C. J. H.: Global exposure to river and coastal flooding: Long term trends and changes, *Global Environmental Change*, 22, 823–835, <https://doi.org/10.1016/j.gloenvcha.2012.07.004>, 2012.
- Jongman, B., Winsemius, H. C., Aerts, J. C. J. H., Coughlan de Perez, E., Van Aalst, M. K., Kron, W., and Ward, P. J.: Declining vulnerability to river floods and the global benefits of adaptation, *Proceedings of the National Academy of Sciences*, 112, E2271–E2280, <https://doi.org/10.1073/pnas.1414439112>, 2015.
- 530 Kelder, T., Müller, M., Slater, L. J., Marjoribanks, T. I., Wilby, R. L., Prudhomme, C., Bohlinger, P., Ferranti, L., and Nipen, T.: Using UNSEEN trends to detect decadal changes in 100-year precipitation extremes, *npj Climate and Atmospheric Science*, 3, <https://doi.org/10.1038/s41612-020-00149-4>, 2020.
- Kron, W.: Flood disasters—a global perspective, *Water Policy*, 17, 6–24, <https://doi.org/10.2166/wp.2015.001>, 2015.

- Maraun, D., Osborn, T. J., and Rust, H. W.: The influence of synoptic airflow on UK daily precipitation extremes. Part I: Observed spatio-temporal relationships, *Climate Dynamics*, 36, 261–275, <https://doi.org/10.1007/s00382-009-0710-9>, 2011.
- 535 Merz, B., Elmer, F., Kunz, M., Mühr, B., Schröter, K., and Uhlemann-Elmer, S.: The extreme flood in June 2013 in Germany, *La Houille Blanche*, 1, 5–10, <https://doi.org/10.1051/lhb/2014001>, 2014.
- Merz, B., Blöschl, G., Vorogushyn, S., Dottori, F., Aerts, J. C. J. H., Bates, P., Bertola, M., Kemter, M., Kreibich, H., Lall, U., and Macdonald, E.: Causes, impacts and patterns of disastrous river floods, *Nature Reviews Earth & Environment*, 2, 592–609, <https://doi.org/10.1038/s43017-021-00195-3>, 2021.
- 540 Mizuta, R. and Endo, H.: Projected changes in extreme precipitation in a 60-km AGCM large ensemble and their dependence on return periods, *Geophysical Research Letters*, 47, e2019GL086855, <https://doi.org/10.1029/2019GL086855>, 2020.
- Mohr, S., Ehret, U., Kunz, M., Ludwig, P., Caldas-Alvarez, A., Daniell, J. E., Ehmele, F., Feldmann, H., Franca, M. J., Gattke, C., Hundhausen, M., Knippertz, P., Küpfer, K., Mühr, B., Pinto, J. G., Quinting, J., Schäfer, A. M., Scheibel, M., Seidel, F., and Wisotzky, C.: A multi-disciplinary analysis of the exceptional flood event of July 2021 in central Europe—Part 1: Event description and analysis, *Natural Hazards and Earth System Sciences*, 23, 525–551, <https://doi.org/10.5194/nhess-23-525-2023>, 2023.
- 545 Molteni, F., Buizza, R., Palmer, T. N., and Petroliagis, T.: The ECMWF ensemble prediction system: Methodology and validation, *Quarterly Journal of the Royal Meteorological Society*, 122, 73–119, <https://doi.org/10.1002/qj.49712252905>, 1996.
- O’Gorman, P. A. and Schneider, T.: The physical basis for increases in precipitation extremes in simulations of 21st-century climate change, *Proceedings of the National Academy of Sciences*, 106, 14773–14777, <https://doi.org/10.1073/pnas.0907610106>, 2009.
- 550 Papalexiou, S. M. and Koutsoyiannis, D.: Battle of extreme value distributions: A global survey on extreme daily rainfall, *Water Resources Research*, 49, 187–201, <https://doi.org/10.1029/2012WR012557>, 2013.
- Pendergrass, A. G. and Hartmann, D. L.: Changes in the distribution of rain frequency and intensity in response to global warming, *Journal of Climate*, 27, 8372–8383, <https://doi.org/10.1175/JCLI-D-14-00183.1>, 2014.
- 555 Pfeleiderer, P., Schleussner, C.-F., Kornhuber, K., and Coumou, D.: Summer weather becomes more persistent in a 2°C world, *Nature Climate Change*, 9, 666–671, <https://doi.org/10.1038/s41558-019-0555-0>, 2019.
- Rajulapati, C. R., Papalexiou, S. M., Clark, M. P., Razavi, S., Tang, G., and Pomeroy, J. W.: Assessment of extremes in global precipitation products: How reliable are they?, *Journal of Hydrometeorology*, 21, 2855–2873, <https://doi.org/10.1175/JHM-D-20-0040.1>, 2020.
- Rodrigues, D. T., Gonçalves, W. A., Spyrides, M. H. C., Santos e Silva, C. M., and de Souza, D. O.: Spatial distribution of the level of return of extreme precipitation events in Northeast Brazil, *International Journal of Climatology*, 40, 5098–5113, <https://doi.org/10.1002/joc.6507>, 2020.
- 560 Ruff, F. and Pfahl, S.: What distinguishes 100-year precipitation extremes over Central European river catchments from more moderate extreme events?, *Weather and Climate Dynamics*, 4, 427–447, <https://doi.org/10.5194/wcd-4-427-2023>, 2023.
- Sene, K.: Flash floods, in: *Hydrometeorology*, pp. 273–312, Springer, Cham, https://doi.org/10.1007/978-3-319-23546-2_9, 2016.
- 565 Stephenson, A. G.: evd: Extreme Value Distributions, *R News*, 2, 31–32, <https://CRAN.R-project.org/doc/Rnews/>, last access: 26 April 2023, 2002.
- Tellman, B., Sullivan, J. A., Kuhn, C., Kettner, A. J., Doyle, C. S., Brakenridge, G. R., Erickson, T. A., and Slayback, D. A.: Satellite imaging reveals increased proportion of population exposed to floods, *Nature*, 596, 80–86, <https://doi.org/10.1038/s41586-021-03695-w>, 2021.
- Ventura, V., Paciorek, C. J., and Risbey, J. S.: Controlling the proportion of falsely rejected hypotheses when conducting multiple tests with climatological data, *Journal of Climate*, 17, 4343–4356, <https://doi.org/10.1175/3199.1>, 2004.
- 570

World Meteorological Organization: Manual on estimation of probable maximum precipitation (PMP), vol. 1045, World Meteorological Organization, Geneva, 2009.

Supplementary Material




Figure S1. Frequency of missing values (either dry days or days when satellite data are not available) in the original PERSIANN observations for a data period of 1983 to 2021.




Figure S2. Difference of the (left) 99th and (right) 99.99th percentile of daily precipitation between the EPS data and the (a-d) CHIRPS or (e-h) PERSIANN observations, for (a,b,e,f) absolute and (c,d,g,h) relative differences. Note the non-linear colour scales.




Figure S3. Spatial distribution of statistical significant temporal trends (red grid points), obtained from the FDR test of Benjamini and Hochberg (1995), as described in Ventura et al. (2004), applied on multiple p values associated with the Mann-Kendall trend tests applied for the 99.9th percentile of daily precipitation for (a) the EPS data and the (b) REGEN, (c) CHIRPS and (d) PERSIANN observations.




Figure S4. Ratio between the precipitation amount of a 100-year event of daily precipitation and the annual mean precipitation from the EPS data. The annual mean precipitation is averaged over all ensemble members and over the years 2008 to 2019. Dark grey shading indicates grid points for which the GEV parameters are outside the allowed range and thus no return values can be estimated. Note the non-linear colour scale.




Figure S5. Return value plots of daily precipitation at individual grid boxes in **(a)** the eastern US (79°W , 36°N), **(b)** Sweden (15°E , 58°N), **(c)** Brazil (52°W , 6°S) and **(d)** India (80°E , 21°N), for the EPS data (black) and the observational data sets REGEN (blue), CHIRPS (green) and PERSIANN (red). Confidence intervals on a 95% level are added for the EPS data and REGEN observations.




Figure S6. Difference of the **(a,b)** relative range, relative to the associated return values, and the **(c,d)** absolute range of confidence intervals on a 95% level between the EPS data and the estimates from **(a,c)** CHIRPS or **(b,d)** PERSIANN observations. Dark grey shading indicates grid points for which the GEV parameters are outside the allowed range and thus no return values can be estimated. Stippling indicates where the confidence interval of the EPS data overlaps with the confidence interval from the specific observations. Note the non-linear colour scales.