

A Possible Unaccounted Source of Nitrogen-Containing Compounds Formation in Aerosols: Amines Reacting with Secondary Ozonides

Junting Qiu^{1,2,3}, Xinlin Shen^{1,2}, Jiangyao Chen^{1,2}, Guiying Li^{1,2}, Taicheng An^{1,2}

¹Guangdong-Hong Kong-Macao Joint Laboratory for Contaminants Exposure and Health, Guangdong Key Laboratory of Environmental Catalysis and Health Risk Control, Institute of Environmental Health and Pollution control, Guangdong University of Technology, Guangzhou 510006, China.

²Guangzhou Key Laboratory of Environmental Catalysis and Pollution Control, Key Laboratory of City Cluster Environmental Safety and Green Development of the Ministry of Education, School of Environmental Science and Engineering, Guangdong University of Technology, Guangzhou 510006, China.

³Graduate School of Frontier Sciences, The University of Tokyo, 5-1-5 Kashiwanoha, Kashiwa 277-8563, Japan.

Correspondence to: Taicheng An (antc99@gdut.edu.cn)

Abstract. Nitrogen (N)-containing compounds have a significant impact on the optical and toxicological properties of aerosols. 1,2,4-Trioxolanes, known as secondary ozonides (SOZs), key products from the ozonolysis of biogenic terpenoids, are readily taken up into atmospheric aerosols and act as oxidants, potentially interacting with amines in the atmosphere. In the present work, we carefully investigated the component of the particles produced by the ozonolysis of β -caryophyllene (β -C) in the presence of ethylamine (EA), methylamine (MA), dimethylamine (DMA) or ammonia. The mass spectrometric results show that SOZ is the dominant product from the ozonolysis of β -C. It readily reacts with EA and MA but has inert reactivities toward DMA and ammonia. Similar experimental results were achieved with α -humulene (α -H), an isomer of β -C, was used in place of β -C. Additionally, D₂O and H₂¹⁸O solvents were used for the characterization of products. The results revealed an intriguing phenomenon that the products from β -C SOZ and α -H SOZ reacting with the same amine (EA or MA) possessed different functional groups, despite the fact that they are isomerized species with the identical chemical structure (1,2,4-trioxolane). That indicates the chemical conformation of SOZs has a strong influence on how they react with amines. For the first time, SOZs derived from β -C and α -H reacting with amines is reported in this study, that may represent a hitherto unrecognized source of N-containing compounds production in atmospheric aerosols.

1 Introduction

Nitrogen (N)-containing compounds are ubiquitous in atmospheric aerosols. The compounds possessing some N-containing functional groups, such as imidazole, pyrazine, and organonitrate groups, are recognized as acting as chromophores, which are closely associated with optical properties of the aerosols (Laskin et al., 2015; Moise et al., 2015). Meanwhile, some N-containing compounds are considered to be hazardous to human health. For instance, nitration of polycyclic aromatic compounds (PAHs) typically contributes more to the toxicity of ambient PM than parent PAHs (Albinet et al., 2008). Another study revealed that the compounds in some pollens will have a higher allergenic potential after being converted into

N-containing species (Cuinica et al., 2014). A main contributor to the formation of N-containing compounds is the progress that NO/NO₂ are transferred to terminal products in NO_x radical cycle, including peroxy nitrates (RO₂NO₂), alkyl nitrates (RONO₂), and nitric acid (HNO₃) (Perring et al., 2013). Additional route for N-containing species formation in the atmosphere is the oxidation of VOCs by the nitrate radical (NO₃) (Ng et al., 2017; Fry et al., 2014). On the other hand, atmospheric amine chemistry, involving new particle formation (NPF) events is another important contributor to N-containing compounds in aerosols. The substitution by one or more organic functional groups leads to stronger basicity of amines than ammonia, indicating that amines are readily to participate in NPF through acid-base reactions, which have been confirmed in numerous field researches (Yao et al., 2016; Yu et al., 2012) and laboratory works (Almeida et al., 2013; Erupe et al., 2011; Glasoe et al., 2015; Kurten et al., 2014; Tong et al., 2020).

Amines are massively emitted from various biogenic and anthropogenic sources, such as biomass burning, animal husbandry, ocean organisms, automobiles and industries (Ge et al., 2011). Aliphatic amines with low molecular weight, including methylamine (MA), dimethylamine (DMA), trimethylamine (TMA), or ethylamine (EA), are dominant species of amines in the atmosphere. Yao et al. reported a high concentration of total DMA and EA up to 130 pptv in Shanghai, China (Yao et al., 2016). In the boreal forest, the concentration of total DMA and EA were detected to be around 150 pptv (Kieloaho et al., 2013). Amines involved atmospheric chemistry regarding the formation of N-containing compounds and growth of secondary organic aerosols (SOAs) has been investigated in cohort works. For instance, amines are found to be fairly reactive toward important atmospheric aldehydes in condensed phase, such as glyoxal, methylglyoxal, glycoaldehyde, or acetaldehyde, which significantly affect the physicochemical properties of aerosols and contribute to SOAs growth (De Haan et al., 2011; Galloway et al., 2014). A recent study shows that alkylammonium carboxylates formed from the reactions of amines with organic acids have lower vapor pressures than original organic acids, implying that alkylammonium carboxylates could enhance SOAs formation (Lavi et al., 2015). Another study carried out by the same group reinforced the crucial role of alkylammonium carboxylates in determining the characteristics of aerosols, for the reason that alkylammonium carboxylates are capable of enhancing the particle hygroscopicity and the cloud condensation nuclei activity (Gomez-Hernandez et al., 2016). In addition, Duporte et al. reported a systematic research on the ozonolysis of α -pinene in the presence of DMA, and they found that DMA was able to react with aldehydes or carboxylate acids generated from the ozonolysis of α -pinene, enhancing SOAs formation (Duporte et al., 2017).

Amines can also be directly oxidized by major atmospheric oxidants, such as OH, O₃ and NO₃, in gas phase or in condensed phase (Ge et al., 2011; Qiu and Zhang, 2013; Tang et al., 2013). Recently, the interactions of Criegee intermediates with amines have been investigated in some studies, however, the actual effect of Criegee intermediates on oxidizing amines is by now unclear (Chhantyal-Pun et al., 2019b; Kumar and Francisco, 2019; Ma et al., 2020; Mull et al., 2020). 1,2,4-Trioxolanes, known as secondary ozonides (SOZs), are formed by the intramolecular reactions of the Criegee moieties with the carbonyl endgroups, as well as bimolecular reaction of Criegee intermediates with carbonyl species such as formaldehyde and acetone (Chhantyal-Pun et al., 2020; Chhantyal-Pun et al., 2019a; Cornwell et al., 2021; Wang et al., 2022). SOZs are major products from the ozonolysis of important biogenic terpenoids, such as limonene, carene, β -pinene,

β -caryophyllene and α -humulene (Vibenholt et al., 2009; Winterhalter et al., 2000; Nguyen et al., 2009; Winterhalter et al., 2009; Beck et al., 2011), and they are readily taken up into atmospheric aerosols (Yao et al., 2014). The formation of SOZs occurs not only in the gas phase but also in bulk liquid phases (Griesbaum et al., 1996) and at gas-liquid/solid interfaces (Enami et al., 2008; Karagulian et al., 2008; Coffaro and Weisel, 2022). Additionally, SOZs can also be formed via OH reactions of lipid molecules (Zeng et al., 2020; Zhang et al., 2018). Since SOZs are categorized as both organic peroxides and reactive oxygen species (Sanchez and Myers, 2000), they potentially function as oxidants and interact with amines in atmospheric condensed phase. Therefore, the aim of this study is to determine whether the interaction of SOZs with amines results in the formation of N-containing compounds.

β -Caryophyllene (β -C) and α -humulene (α -H) are representative sesquiterpenes (Arey et al., 1995; Helmig et al., 2007), and their chemical structures are listed in Figure S1. Albeit not predominant terpene species like isoprene or α -pinene, β -C and α -H are of special significance as powerful SOA makers, due to their rapid degradation in the atmosphere and the low volatility of the degradation products. SOZs are dominant products from the ozonolysis of both β -C and α -H (Nguyen et al., 2009; Winterhalter et al., 2009; Beck et al., 2011), thus, in this study, we choose the reactions of β -C and α -H with O_3 to produce SOZs, and investigate the reactivities of SOZs toward amines. Firstly, we carefully carry out the ozonolysis experiments of β -C in the absence/presence of EA in a smog chamber. The particles generated inside the smog chamber are monitored and the chemical components of the particles are detected by mass spectrometry. D_2O and $H_2^{18}O$ isotope labelling experiments were performed for the identification of the products detected. Next, chemical structures of the products formed from EA reacting with β -C SOZ and α -H SOZ are compared to know the effect of molecular conformation on the reaction mechanism of SOZs. In addition, the reactivities of SOZs toward MA, EA, DMA or ammonia are also comparably investigated.

2 Experimental

2.1 Ozonolysis experiment

All the ozonolysis experiments were carried out in a smog chamber. Considering that the details of the smog chamber has been reported in other articles (Luo et al., 2021; Luo et al., 2020), we just make a brief description herein. Two extremely same pillow-shaped Teflon reactors ($2.5\text{ m} \times 2.0\text{ m}$) mounted inside the smog chamber, and each reactor is surrounded by 3 high efficiency ionizing blower (varied from 0 to a maximum of 2000 r/min), in order to mix the air inside the reactors evenly. **In this study each experiment needs to be repeated at least 3 times. To avoid unexpected errors in the experiment processes, all the experiments were conducted in the same reactor.** Before the operation of each experiment, the reactor was filled with zero air at a volume of 1000 L, with no detectable particles, < 0.5 ppb non-methane hydrocarbon (NMHC), and < 1 ppb NO_x , O_3 and carbonyl compounds. The relative humidity (RH) inside the reactor is $\leq 5\%$, and the temperature is kept at $295 \pm 3\text{ K}$.

Schematic experimental procedure for the ozonolysis of β -C in the presence of EA is presented in Figure S2. β -C is well mixed with EA before the addition of O_3 , and the initial concentrations of chemical species inside a reactor are β -C 200 ppb, EA 80 ppb, O_3 50 ppb. Other experiments (such as 200 ppb β -C + 80 ppb MA + 50ppb O_3 , or 200 ppb α -H + 80 ppb ammonia + 50 ppb O_3), are carried out with the same method. Because the reaction rate of O_3 toward β -C ($1.4 \times 10^{-14} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$) or α -H ($1.2 \times 10^{-14} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$) (Atkinson and Arey, 2003) is much faster than amines or ammonia ($10^{18} - 10^{21} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$) (Ge et al., 2011) and the initial concentration of β -C/ α -H (200 ppb) is times of O_3 (50 ppb), O_3 will almost be consumed intermediately via the reaction with β -C/ α -H after being injected inside the reactor. The products from ozonolysis of β -C/ α -H may subsequently participate in the reactions with amines or ammonia. Ozonolysis leads to the generation of particle matters via condensation of oxidized low-volatility species, which are able to be monitored by a scanning mobility particle sizer (SMPS, TSI). All experiments were performed in dark conditions and without an OH scavenger.

2.2 Particle collection and analysis

Particles with considerable sizes were collected on 47 mm quartz filters at a timing of 3 h after the injection of ozone, and quartz filters were pretreated via 8 h baking inside a Muffle furnace at a temperature of 450 °C. All the filter samples were wrapped in aluminum foil and stored in a freezer at -18 °C until extraction. Particle phase compounds were extracted by soaking filter samples in 5 mL mixture of acetonitrile/ultrapure water (AN/W, vol/vol = 4/1) for 30 min at room temperature. D_2O and $H_2^{18}O$ were also used in extraction instead of ultrapure water for a detailed characterization of products. A high-resolution electrospray ionization mass spectrometer (ESI-MS, ThermoFisher Q Extract quadrupole-Orbitrap) was applied in the detection of chemical compounds extracted in solutions.

2.3 Materials

Gas phase chemicals, such as O_3 and MA, were directly injected into the reactor, while liquid phase chemicals, such as EA and DMA, were injected slowly through a T-junction connected to a Fluorinated Ethylene Propylene line and spread with the flow of purified dry air, by using airtight syringes (Shanghai Anting). O_3 was generated by a commercial ozone generator, and the amount of O_3 was carefully calculated according to the injection time and the power of the ozone generator. Before the operation of each experiment, O_3 was injected into the reactor filled with 1000 L zero air, and confirmed by an O_3 analyzer (Model 49i, Thermo Scientific). Ultrapure water was obtained from a Millipore Milli-Q water purification system (Xiamen Research Water Purification Technology, Unique-R20, resistivity $\geq 18.2 \text{ M}\Omega \text{ cm}$ at 298 K). Chemicals β -caryophyllene (Tokyo Chemical Industrial, > 95%), α -humulene (Tokyo Chemical Industrial, > 93%), methylamine (Wuhan Newradar, 98.6 ppm mixed in N_2 gas), ethylamine (Aladdin Industrial, 70 wt % in H_2O), dimethylamine (Aladdin Industrial, 40 wt % in H_2O), ammonia solution (Aladdin Industrial, 25 wt % in H_2O), acetonitrile (Aladdin Industrial, $\geq 99.9\%$), D_2O (J&K, > 99.8 atom % D) and $H_2^{18}O$ (Macklin, > 97 atom % ^{18}O) were used as received.

3 Results and discussion

3.1 Reaction of β -C SOZ with EA

130 The particle formation was monitored by SMPS from the ozonolysis of β -C in the absence/presence of EA, and the chemical components of the particles were analyzed, as shown in Figure 1. Because β -C is fairly reactive toward O_3 and the products generated in situ have extremely low volatility, the formation of particles can be observed intermediately after O_3 being injected inside the Teflon reactor. The number of particles decreased rapidly as presented in Figure 1A due to their coagulation to form larger particles or deposition on the wall. According to Figure 1B, the total particle volume produced by
135 the ozonolysis of β -C grew initially before beginning to decline about 10 minutes after the particle loss rate surpassed the creation rate. With the addition of EA, no discernible change in the number concentration of particles was seen in Figure 1A, however the volume of total particles slightly increased as shown in Figure 1B.

The addition of EA has limited effect on promoting particle formation, on account of the volatility of the products from the ozonolysis of β -C is sufficiently low. Nevertheless, the reaction of EA with the products from the ozonolysis of β -C
140 considerably altered the chemical components of particles as shown in Figure 1C that contrasts the positive-ion ESI mass spectra of products from the ozonolysis of β -C in the absence/presence of EA respectively. In the β -C + O_3 experiment, the prominent signal rise at m/z 275 that is assigned to Na^+ -adducted $C_{15}H_{24}O_3$ species, $275 = 252 (C_{15}H_{24}O_3) + 23 (Na^+)$. Both experimental and theoretical research have explicitly explored the mechanisms on the ozonolysis of β -C (Nguyen et al., 2009; Winterhalter et al., 2009). Major species of $C_{15}H_{24}O_3$ are SOZ, vinyl ROOH and carboxylic acid, which are isomerized
145 products of Criegee intermediate as demonstrated in Scheme 1. Furthermore, product appeared at m/z 275 was identified by the method of replacing the AN/W with AN/D₂O (vol/vol = 4/1) in the extraction process. **D₂O was used to test if a molecule contains an active H-atom. When a molecule possessing an active H-atom is dissolved in D₂O solution, the active H-atom readily exchanges with D-atom of D₂O, increasing its molecular weight by 1 unit. The results in Figure S3 show that the product appeared at m/z 275 possesses no exchangeable H-atom, implying it should be $[SOZ + Na]^+$. In contrast, the H-**
150 **atoms of vinyl ROOH and carboxylic acid are exchangeable with D-atom. Na^+ was leached into the solution from trace metals in the laboratory glassware (Greaves and Roboz, 2014). Na^+ has an affinity toward an O-atom of a species possessing R-O-R' such as ethers (Sugimura et al., 2015). In addition, a previous study reported that SOZ originated from α -terpineol was detectable as Na^+ -adducted species (Qiu et al., 2022).**

The observation that intensity of m/z 275 clearly decreased in the experiment of β -C + EA + O_3 indicates that β -C SOZ
155 readily react with EA. Intense peaks appear at m/z 280 (**P1**), which are assigned to the H^+ -adducted products from β -C SOZ reacting with EA, $280 (C_{17}H_{29}O_2N + H^+) = 252 (C_{15}H_{24}O_3) + 45 (C_2H_7N) - 18 (H_2O) + 1 (H^+)$. **Because the aim of this study is to investigate the interactions of SOZs with amines, we only present a part of the products related to this study in mass spectra. The larger products from the ozonolysis of β -C as well as their potential interactions with amines will not be discussed in this work. For the same reason, all the experiments were carried out at an extremely low humidity ($RH \leq 5\%$),**
160 **to avoid the generation of unwanted products in the presence of high concentration water vapor (Kundu et al., 2017).**

The vast majority of P1 is formed from the heterogeneous reaction of EA with SOZ in condensed phase. The evidence can be found in Figure S4, that is even when EA is added in the reactor 30 min after the start of ozonolysis (a situation that SOZ is almost in condensed phase), more than half of the P1 is still produced compared to the situation that EA is well mixed. The results in Figure S4 provides another information that P1 is not the product from EA reacting with Criegee intermediate, the precursor of SOZ, which is a highly reactive species, because Criegee intermediate could not have survived for so long.

Moreover, the particles generated from the ozonolysis of β -C in the absence of EA were sampled and dissolved in AN/W solution. EA was directly added into the solution. 30 min later the solution was analyzed by the mass spectrometer. As a result, no signal appeared at m/z 280 in mass spectra, indicating that β -C SOZ reacting with EA in liquid phase is not available. Through this experiment, the possibility that EA vapor condensed on the particles first and then react with β -C SOZ in the extraction process was ruled out, and it was determined that the reaction of β -C SOZ with EA occurred in the smog chamber.

To the best of our knowledge, the phenomenon that β -C SOZ reacting with EA leads to the production of N-containing compounds, was reported for the first time in this work. Next, chemical analysis of products appearing at m/z 280 was conducted to better comprehend the chemical structures of previously unreported N-containing compounds.

3.2 Chemical identification of P1

D_2O and $H_2^{18}O$ isotope labelling experiments were performed for the chemical identification of P1 in this work. As mentioned before, D_2O was used to confirm if a molecule contains an active H-atom, while $H_2^{18}O$ was used to test if a molecule is carrying a carbonyl group. After being dissolved in $H_2^{18}O$ solution, the compound possessing a carbonyl group forms a gem-diol via the addition of $H_2^{18}O$ which is a reversible process. Since the concentration of $H_2^{18}O$ is overwhelming, ^{16}O of the compound will be almost replaced by ^{18}O of $H_2^{18}O$, resulting in 2 units rise in molecular weight. This method was proven to be beneficial for examining the chemical structure of unidentified species in previous studies (Qiu et al., 2019; Qiu et al., 2020b; Qiu et al., 2020a). Figure 2 shows high-resolution positive-ion ESI mass spectra of products extracted in AN/W (vol/vol = 4/1), AN/ D_2O (vol/vol = 4/1) and AN/ $H_2^{18}O$ (vol/vol = 4/1) solutions, from the reaction of β -C SOZ with EA, where P1 ($C_{17}H_{29}O_3N + H^+$) appeared at m/z 280. 226 in the AN/W experiment. P1 shifted by + 3 mass units in the AN/ D_2O experiment and + 2 mass units in the AN/ $H_2^{18}O$ experiment, showing that P1 possess two exchangeable H-atoms ($H^+ \rightarrow D^+$ contributed another + 1 mass unit) and one exchangeable O-atom, respectively.

The mechanism of β -C SOZ reacting with EA can be explained as follow. The electronegativity of the neighboring oxygens induced a net positive charge on the α -carbon of β -C SOZ. EA acting as a nucleophile may add to α -carbon and cleave β -C SOZ. This theory is supported by a previously reported research by Na et al., which revealed that ammonia reacts with styrene SOZ via a nucleophilic attack at the α -carbon of styrene SOZ, producing benzaldehyde, hydrogen peroxide, and phenylmethanimine in the process (Na et al., 2006). Despite the attack of EA opened the cyclic structure of β -C SOZ, we didn't detect cleavage products as Na et al. reported. Instead, P1 detected in this work is the product from the addition reaction between β -C SOZ and EA, and a water molecule was removed in the process. Based on the molecular weight of P1

and the results of D₂O and H₂¹⁸O isotope labelling experiments, a potential structure of P1 is illustrated in Scheme 2. It has two active H-atoms in the -NH and -OH moieties, and C=¹⁶O can be transferred into C=¹⁸O via H₂¹⁸O addition reaction (see Scheme S1 in SI).

3.3 Reactions of β-C SOZ with MA, DMA and Ammonia

By replacing EA with MA, DMA or ammonia, another three smog chamber experiments were carried out. The particles were sampled and analyzed by electrospray mass spectrometry as shown in Figure 3. Obviously, in the β-C + MA + O₃ experiment, intensity of m/z 275 clearly diminished and intense peak appeared at m/z 266 (**P2**), which is assigned to the H⁺-adducted product from β-C SOZ reacting with MA, $266 (\text{C}_{16}\text{H}_{27}\text{O}_2\text{N} + \text{H}^+) = 252 (\text{C}_{15}\text{H}_{24}\text{O}_3) + 31 (\text{CH}_3\text{N}) - 18 (\text{H}_2\text{O}) + 1 (\text{H}^+)$. In sharp contrast, the intensities of m/z 275 in the other two experiments are essentially identical to that in the β-C + O₃ experiment, suggesting that DMA and ammonia have inert reactivities toward β-C SOZ. The results of D₂O and H₂¹⁸O isotope labelling experiments of P2 shown in Figure S5A. **P2 shifted by + 3 mass units in the AN/D₂O experiment and + 2 mass units in the AN/H₂¹⁸O experiment, which is identical to P1. A possible structure of P2 that is similar to P1 is presented in Scheme S2.**

Substituted by one alkyl moiety, EA or MA are considered more basic than ammonia, that potentially increased the reactivity of EA or MA toward β-C SOZ. On the other hand, the reason why DMA is less reactive than EA and MA can be explained that DMA possesses two alkyl moieties, resulting in a steric hinderance that would limit the accessibility of DMA to β-C SOZ. Na et al. also pointed out α-methylstyrene SOZ is less reactive than styrene SOZ toward ammonia, due to being sterically hindered by the methyl group attached to the α-carbon of 1,2,4-trioxolane (Na et al., 2006). Moreover, in order to obtain more information about the mechanism of SOZs reacting with amines, in the following section, we mainly report the reactions of another SOZ produced by the ozonolysis of α-humulene (α-H), an isomer of β-C, with EA, MA DMA and ammonia.

3.4 Reactions of α-H SOZ with EA, MA, DMA and Ammonia

The smog chamber experiments of α-H + amines/ammonia + O₃ were conducted by using the same procedure, and the chemical composition of particles generated in the reactor were analyzed by the mass spectrometer as shown in Figure 4. SOZ (m/z 275, C₁₅H₂₄O₃ + Na⁺) is the dominant product from the ozonolysis of α-H, which is consistent with the previous study (Beck et al., 2011). As can be seen, the behavior of α-H SOZ is comparable to that observed in the experiments of β-C, exhibiting inert reactivities toward DMA and ammonia and selectively reacting with EA and MA. The products from α-H SOZ reacting with EA and MA appeared at m/z 280 (**P3**) and 266 (**P4**). Since α-H SOZ and β-C SOZ are isomerized species, molecular formula of P3 and P4 should be same to P1 and P2 respectively, which are $280 (\text{C}_{17}\text{H}_{29}\text{O}_2\text{N} + \text{H}^+) = 252 (\text{C}_{15}\text{H}_{24}\text{O}_3) + 45 (\text{C}_2\text{H}_7\text{N}) - 18 (\text{H}_2\text{O}) + 1 (\text{H}^+)$ and $266 (\text{C}_{16}\text{H}_{27}\text{O}_2\text{N} + \text{H}^+) = 252 (\text{C}_{15}\text{H}_{24}\text{O}_3) + 31 (\text{CH}_3\text{N}) - 18 (\text{H}_2\text{O}) + 1 (\text{H}^+)$.

225 D₂O and H₂¹⁸O isotope labelling experiments were also performed for chemical identification of P3 and P4. High-resolution positive-ion ESI mass spectra of P3 extracted in AN/W (vol/vol = 4/1), AN/D₂O (vol/vol = 4/1) and AN/H₂¹⁸O (vol/vol = 4/1) solutions are demonstrated in Figure 5. The observation that P3 shifted by + 2 mass units in AN/D₂O experiment and has no mass-shift in AN/H₂¹⁸O experiment indicates that P3 generated from α -H SOZ reacting with EA possesses only one exchangeable H-atom and no exchangeable O-atom. A probable contributor to the exchangeable H-atom is the -NH moiety. Given that P3 possesses no carbonyl or hydroxyl moieties, a dioxirane structure generated by the breaking of C-O bonds appears to be plausible for P3, and a potential structure of P3 is deduced and displayed in Scheme 3. The results from isotope labelling experiments of P4 are presented in Figure S5B, and they point out a production mechanism of P4 that is similar to that of P3 as illustrated in Scheme S2. What should be mentioned is that α -H contains three endocyclic double bonds which are able to be attacked by ozone to generate different SOZs. As a result, P3 and P4 may have multiple conformations, however, to simplify the representations, just one kind is provided here as an example of each. In addition, it is worth noting that the signal intensity of P4 in the α -H + MA + O₃ experiment is rather weak, even though the majority of α -H SOZ has been consumed via its reaction with MA and contribute to the formation of P4. This phenomenon can be explained that dioxiranes are active species and their stabilities are highly dependent on their molecular structures (El-Assaad et al., 2022). In other words, P4 dissipated rapidly after it formed because it is less stable than P3.

230
235
240 An astonishing phenomenon revealed here is that the product of α -H SOZ reacting with EA (P3) bear no resemblance to that of β -C SOZ reacting with EA (P1), in spite of the fact that α -H SOZ is an isomerized species of β -C SOZ and they share a same chemical structure of 1,2,4-trioxolane, which suggests that the molecular conformations of SOZs have a substantial impact on their reaction mechanism, resulting in the formation of N-containing products processing various functional groups.

245 **4 Atmospheric implications**

The predominant source of SOZs is biogenic terpenoids, which are emitted to atmosphere at a rate of 10¹⁴ g/year. (Guenther et al., 1995) SOZs originating from terpenoids are less volatile and hence more easily taken up into aerosols. (Yao et al., 2014) Multiphase ozonolysis and OH oxidations of unsaturated organic compounds possessing C=C bond(s) also produce SOZs, which causes an accumulation of SOZs in condensed phases (Heine et al., 2017; Enami et al., 2008; Karagulian et al., 2008; Coffaro and Weisel, 2022; Zhou et al., 2022; Zhang et al., 2018). Terpenoid-derived SOZs are comparatively stable organic peroxides in condensed phase. For instance, A recent study revealed that C₁₀ and C₁₃ SOZs derived from terpineol can persist in water for weeks (Qiu et al., 2022). In addition, products from the ozonolysis of terpenoids including SOZs, are usually surface active in aerosols (Qiu et al., 2018b; Qiu et al., 2018a), which facilitated their reactions toward gas-phase amines. Thus, SOZs reacting with amines is probably a nonnegligible source of N-containing compounds formation in aerosols.

250
255 Moreover, Na et al. reported that SOZs derived from styrene and α -methylstyrene can react with ammonia (Na et al., 2006). In sharp contrast, our research suggests that both β -C SOZ and α -H SOZ exhibit inert reactivities toward ammonia but

readily react with EA and MA. Additionally, we discovered that SOZs in different conformations reacting with EA or MA produces N-containing compounds with various functional groups. The aforementioned works indicate that the interaction of SOZs with amines or ammonia is a complicated process in the real atmosphere, leading to the formation of various N-containing compounds. Due to the distinct roles that N-containing compounds with different functional groups play in the properties of aerosols (Laskin et al., 2015), a thorough investigation on the mechanism of SOZs reacting with amines is still urgently required. Nevertheless, the information obtained in the present study is just the tip of an iceberg, and detailed laboratory works combined with field researches are necessary toward a full picture of N-containing compounds originating from SOZs reacting with amines or ammonia.

Apart from SOZs, other organic peroxides like ROOH or ROOR', play more significant roles in atmospheric chemistry (Wang et al., 2023). For example, the oxidation of dissolved SO₂ by organic peroxides has been considered as a main source of sulfate formation in aerosols (Dovrou et al., 2021; Dovrou et al., 2019; Wang et al., 2021; Wang et al., 2019; Yao et al., 2019). In addition, organic peroxides can directly interact with transitional metal ions via Fenton-like reactions mechanism (Fang et al., 2020; Hu et al., 2021; Tong et al., 2016; Wei et al., 2022). However, the interaction of organic peroxides with amines has rarely been reported in previous studies. The present study recommends extensive research on organic peroxides including SOZs reacting with amines, which will deepen our understanding of the source of N-containing compounds and benefit in the works on precisely evaluating the effects of atmospheric aerosols on human health and climate (Seinfeld et al., 2016; Shiraiwa et al., 2017; Shrivastava et al., 2017).

5 Conclusion

In this study, the chamber experiments showed that the component of particles produced by the ozonolysis of both β -C and α -H was dramatically altered in addition of EA or MA, which was originated from the reactions of SOZs with EA or MA. However, both β -C SOZ and α -H SOZ were found have inert reactivities toward DMA and ammonia. Additionally, D₂O and H₂¹⁸O isotope labelling experiments revealed that the products from β -C SOZ and α -H SOZ reacting with the same amine (EA or MA) possessed different functional groups, despite β -C SOZ and α -H SOZ are isomerized species and share a same chemical structure of 1,2,4-trioxolane. The experimental results obtained in this study indicate that a variety of N-containing compounds can be generated via the interaction of SOZs with amines, which may constitute a hitherto unaccounted source of N-containing compounds formation in atmospheric aerosols.

Data availability. The data that support the results are available upon request. Please email Junting Qiu (paziqjt@gamil.com)

Supplement. The supplement related to this article is available online.

Author contributions. TA and JQ designed research. JQ and XS performed experiments. JQ analyzed the data. All the authors participated in writing the paper.

290

Competing interests. The authors declare that they have no conflict of interest.

Financial support. This work is financially supported by a National Natural Science Foundation of China (42020104001, 42107118 and 42177354), Local Innovative and Research Teams Project of Guangdong Pearl River Talents Program (2017BT01Z032), and China Postdoctoral Science Foundation (2021M700881).

295


References

- Albinet, A., Leoz-Garziandia, E., Budzinski, H., Villenave, E., and Jaffrezo, J.-L.: Nitrated and oxygenated derivatives of polycyclic aromatic hydrocarbons in the ambient air of two French alpine valleysPart 1: Concentrations, sources and gas/particle partitioning, *Atmospheric Environment*, 42, 43-54, 10.1016/j.atmosenv.2007.10.009, 2008.
- Almeida, J., Schobesberger, S., Kurten, A., Ortega, I. K., Kupiainen-Maatta, O., Praplan, A. P., Adamov, A., Amorim, A., Bianchi, F., Breitenlechner, M., David, A., Dommen, J., Donahue, N. M., Downard, A., Dunne, E., Duplissy, J., Ehrhart, S., Flagan, R. C., Franchin, A., Guida, R., Hakala, J., Hansel, A., Heinritzi, M., Henschel, H., Jokinen, T., Junninen, H., Kajos, M., Kangasluoma, J., Keskinen, H., Kupc, A., Kurten, T., Kvashin, A. N., Laaksonen, A., Lehtipalo, K., Leiminger, M., Leppa, J., Loukonen, V., Makhmutov, V., Mathot, S., McGrath, M. J., Nieminen, T., Olenius, T., Onnela, A., Petaja, T., Riccobono, F., Riipinen, I., Rissanen, M., Rondo, L., Ruuskanen, T., Santos, F. D., Sarnela, N., Schallhart, S., Schnitzhofer, R., Seinfeld, J. H., Simon, M., Sipila, M., Stozhkov, Y., Stratmann, F., Tome, A., Trostl, J., Tsagkogeorgas, G., Vaattovaara, P., Viisanen, Y., Virtanen, A., Vrtala, A., Wagner, P. E., Weingartner, E., Wex, H., Williamson, C., Wimmer, D., Ye, P., Yli-Juuti, T., Carslaw, K. S., Kulmala, M., Curtius, J., Baltensperger, U., Worsnop, D. R., Vehkamäki, H., and Kirkby, J.: Molecular understanding of sulphuric acid-amine particle nucleation in the atmosphere, *Nature*, 502, 359-363, 10.1038/nature12663, 2013.
- Arey, J., Crowley, D. E., Crowley, M., Resketo, M., and Lester, J.: Hydrocarbon emissions from natural vegetation in California's South Coast Air Basin, *Atmospheric Environment*, 29, 2977-2988, [https://doi.org/10.1016/1352-2310\(95\)00137-N](https://doi.org/10.1016/1352-2310(95)00137-N), 1995.
- Atkinson, R. and Arey, J.: Gas-phase tropospheric chemistry of biogenic volatile organic compounds: a review, *Atmospheric Environment*, 37, 197-219, 10.1016/s1352-2310(03)00391-1, 2003.
- Beck, M., Winterhalter, R., Herrmann, F., and Moortgat, G. K.: The gas-phase ozonolysis of α -humulene, *Physical Chemistry Chemical Physics*, 13, 10970, 10.1039/c0cp02379e, 2011.
- Chhantyal-Pun, R., Khan, M. A. H., Taatjes, C. A., Percival, C. J., Orr-Ewing, A. J., and Shallcross, D. E.: Criegee intermediates: production, detection and reactivity, *Int. Rev. Phys. Chem.*, 39, 383-422, 10.1080/0144235x.2020.1792104, 2020.
- Chhantyal-Pun, R., Khan, M. A. H., Martin, R., Zachhuber, N., Buras, Z. J., Percival, C. J., Shallcross, D. E., and Orr-Ewing, A. J.: Direct Kinetic and Atmospheric Modeling Studies of Criegee Intermediate Reactions with Acetone, *ACS Earth Space Chem.*, 3, 2363-2371, 10.1021/acsearthspacechem.9b00213, 2019a.
- Chhantyal-Pun, R., Shannon, R. J., Tew, D. P., Caravan, R. L., Duchi, M., Wong, C., Ingham, A., Feldman, C., McGillen, M. R., Khan, M. A. H., Antonov, I. O., Rotavera, B., Ramasesha, K., Osborn, D. L., Taatjes, C. A., Percival, C. J., Shallcross, D. E., and Orr-Ewing, A. J.: Experimental and computational studies of Criegee intermediate reactions with NH₃ and CH₃NH₂, *Phys Chem Chem Phys*, 21, 14042-14052, 10.1039/c8cp06810k, 2019b.
- Coffaro, B. and Weisel, C. P.: Reactions and Products of Squalene and Ozone: A Review, *Environ. Sci. Technol.*, 56, 7396-7411, 10.1021/acs.est.1c07611, 2022.
- Cornwell, Z. A., Harrison, A. W., and Murray, C.: Kinetics of the Reactions of CH₂O with Acetone, α -Diketones, and β -Diketones, *J. Phys. Chem. A*, 125, 8557-8571, 10.1021/acs.jpca.1c05280, 2021.
- Cuñica, L. G., Abreu, I., and Esteves da Silva, J.: Effect of air pollutant NO(2) on *Betula pendula*, *Ostrya carpinifolia* and *Carpinus betulus* pollen fertility and human allergenicity, *Environ Pollut*, 186, 50-55, 10.1016/j.envpol.2013.12.001, 2014.


- De Haan, D. O., Hawkins, L. N., Kononenko, J. A., Turley, J. J., Corrigan, A. L., Tolbert, M. A., and Jimenez, J. L.: Formation of Nitrogen-Containing Oligomers by Methylglyoxal and Amines in Simulated Evaporating Cloud Droplets, *Environmental Science & Technology*, 45, 984-991, 10.1021/es102933x, 2011.
- 335 Dovrou, E., Rivera-Rios, J. C., Bates, K. H., and Keutsch, F. N.: Sulfate Formation via Cloud Processing from Isoprene Hydroxyl Hydroperoxides (ISOPOOH), *Environmental Science & Technology*, 53, 12476-12484, 10.1021/acs.est.9b04645, 2019.
- Dovrou, E., Bates, K. H., Rivera-Rios, J. C., Cox, J. L., Shutter, J. D., and Keutsch, F. N.: Towards a chemical mechanism of the oxidation of aqueous sulfur dioxide via isoprene hydroxyl hydroperoxides (ISOPOOH), *Atmos. Chem. Phys.*, 21, 8999-9008, 10.5194/acp-21-8999-2021, 2021.
- 340 Duporte, G., Riva, M., Parshintsev, J., Heikkinen, E., Barreira, L. M. F., Myllys, N., Heikkinen, L., Hartonen, K., Kulmala, M., Ehn, M., and Riekkola, M. L.: Chemical Characterization of Gas- and Particle-Phase Products from the Ozonolysis of alpha-Pinene in the Presence of Dimethylamine, *Environ Sci Technol*, 51, 5602-5610, 10.1021/acs.est.6b06231, 2017.
- El-Assaad, T. H., Zhu, J., Sebastian, A., McGrath, D. V., Neogi, I., and Parida, K. N.: Dioxiranes: a half-century journey, *Organic Chemistry Frontiers*, 9, 5675-5725, 10.1039/D2QO01005D, 2022.
- 345 Enami, S., Hoffmann, M. R., and Colussi, A. J.: Acidity enhances the formation of a persistent ozonide at aqueous ascorbate/ozone gas interfaces, *Proc. Natl. Acad. Sci. U. S. A.*, 105, 7365-7369, 10.1073/pnas.0710791105, 2008.
- Erupe, M. E., Viggiano, A. A., and Lee, S. H.: The effect of trimethylamine on atmospheric nucleation involving H₂SO₄, *Atmospheric Chemistry and Physics*, 11, 4767-4775, 10.5194/acp-11-4767-2011, 2011.
- 350 Fang, T., Lakey, P. S. J., Rivera-Rios, J. C., Keutsch, F. N., and Shiraiwa, M.: Aqueous-Phase Decomposition of Isoprene Hydroxy Hydroperoxide and Hydroxyl Radical Formation by Fenton-like Reactions with Iron Ions, *The Journal of Physical Chemistry A*, 124, 5230-5236, 10.1021/acs.jpca.0c02094, 2020.
- Fry, J. L., Draper, D. C., Barsanti, K. C., Smith, J. N., Ortega, J., Winkle, P. M., Lawler, M. J., Brown, S. S., Edwards, P. M., Cohen, R. C., and Lee, L.: Secondary Organic Aerosol Formation and Organic Nitrate Yield from NO₃ Oxidation of Biogenic Hydrocarbons, *Environmental Science & Technology*, 48, 11944-11953, 10.1021/es502204x, 2014.
- 355 Galloway, M. M., Powelson, M. H., Sedehi, N., Wood, S. E., Millage, K. D., Kononenko, J. A., Rynaski, A. D., and De Haan, D. O.: Secondary Organic Aerosol Formation during Evaporation of Droplets Containing Atmospheric Aldehydes, Amines, and Ammonium Sulfate, *Environmental Science & Technology*, 48, 14417-14425, 10.1021/es5044479, 2014.
- Ge, X., Wexler, A. S., and Clegg, S. L.: Atmospheric amines – Part I. A review, *Atmospheric Environment*, 45, 524-546, 10.1016/j.atmosenv.2010.10.012, 2011.
- 360 Glasoe, W. A., Volz, K., Panta, B., Freshour, N., Bachman, R., Hanson, D. R., McMurry, P. H., and Jen, C.: Sulfuric acid nucleation: An experimental study of the effect of seven bases, *Journal of Geophysical Research: Atmospheres*, 120, 1933-1950, 10.1002/2014jd022730, 2015.
- Gomez-Hernandez, M., Mckeown, M., Secrest, J., Marrero-Ortiz, W., Lavi, A., Rudich, Y., Collins, D. R., and Zhang, R.: Hygroscopic Characteristics of Alkylammonium Carboxylate Aerosols, *Environmental Science & Technology*, 50, 2292-2300, 10.1021/acs.est.5b04691, 2016.
- 365 Greaves, J. and Roboz, J.: *Mass spectrometry for the novice*, CRC Press, is an imprint of the Taylor & Francis Group, Boca Raton, xxxii, 275 pages pp.2014.
- Griesbaum, K., Hilß, M., and Bosch, J.: Ozonides of mono-, bi- and tricyclic terpenes, *Tetrahedron*, 52, 14813-14826, [https://doi.org/10.1016/0040-4020\(96\)00936-2](https://doi.org/10.1016/0040-4020(96)00936-2), 1996.
- 370 Guenther, A., Hewitt, C. N., Erickson, D., Fall, R., Geron, C., Graedel, T., Harley, P., Klinger, L., Lerdau, M., McKay, W. A., Pierce, T., Scholes, B., Steinbrecher, R., Tallamraju, R., Taylor, J., and Zimmerman, P.: A global-model of natural volatile organic-compound emissions, *J. Geophys. Res. Atmos.*, 100, 8873-8892, 10.1029/94jd02950, 1995.
- Heine, N., Houle, F. A., and Wilson, K. R.: Connecting the Elementary Reaction Pathways of Criegee Intermediates to the Chemical Erosion of Squalene Interfaces during Ozonolysis, *Environ. Sci. Technol.*, 51, 13740-13748, 10.1021/acs.est.7b04197, 2017.
- 375 Helmig, D., Ortega, J., Duhl, T., Tanner, D., Guenther, A., Harley, P., Wiedinmyer, C., Milford, J., and Sakulyanontvittaya, T.: Sesquiterpene Emissions from Pine Trees – Identifications, Emission Rates and Flux Estimates for the Contiguous United States, *Environmental Science & Technology*, 41, 1545-1553, 10.1021/es0618907, 2007.
- Hu, M., Tonokura, K., Morino, Y., Sato, K., and Enami, S.: Effects of Metal Ions on Aqueous-Phase Decomposition of α -Hydroxyalkyl-Hydroperoxides Derived from Terpene Alcohols, *Environmental Science & Technology*, 55, 12893-12901, 10.1021/acs.est.1c04635, 2021.
- 380 Karagulian, F., Lea, A. S., Dilbeck, C. W., and Finlayson-Pitts, B. J.: A new mechanism for ozonolysis of unsaturated organics on solids: phosphocholines on NaCl as a model for sea salt particles, *Phys. Chem. Chem. Phys.*, 10, 528-541, 10.1039/b712715d, 2008.
- Kieloaho, A. J., Hellen, H., Hakola, H., Manninen, H. E., Nieminen, T., Kulmala, M., and Pihlatie, M.: Gas-phase alkylamines in a boreal Scots pine forest air, *Atmospheric Environment*, 80, 369-377, 10.1016/j.atmosenv.2013.08.019, 2013.
- 385 Kumar, M. and Francisco, J. S.: Elucidating the molecular mechanisms of Criegee-amine chemistry in the gas phase and aqueous surface environments, *Chemical Science*, 10, 743-751, 10.1039/c8sc03514h, 2019.
- Kundu, S., Fisseha, R., Putman, A. L., Rahn, T. A., and Mazzoleni, L. R.: Molecular formula composition of β -caryophyllene ozonolysis SOA formed in humid and dry conditions, *Atmospheric Environment*, 154, 70-81, 10.1016/j.atmosenv.2016.12.031, 2017.

- 390 Kurten, A., Jokinen, T., Simon, M., Sipila, M., Sarnela, N., Junninen, H., Adamov, A., Almeida, J., Amorim, A., Bianchi, F., Breitenlechner, M., Dommen, J., Donahue, N. M., Duplissy, J., Ehrhart, S., Flagan, R. C., Franchin, A., Hakala, J., Hansel, A., Heinritzi, M., Hutterli, M., Kangasluoma, J., Kirkby, J., Laaksonen, A., Lehtipalo, K., Leiminger, M., Makhmutov, V., Mathot, S., Onnela, A., Petaja, T., Praplan, A. P., Riccobono, F., Rissanen, M. P., Rondo, L., Schobesberger, S., Seinfeld, J. H., Steiner, G., Tome, A., Trostl, J., Winkler, P. M., Williamson, C., Wimmer, D., Ye, P. L., Baltensperger, U., Carslaw, K. S., Kulmala, M., Worsnop, D. R., and Curtius, J.: Neutral molecular cluster formation of sulfuric acid-dimethylamine observed in real time under atmospheric conditions, *P Natl Acad Sci USA*, 111, 15019-15024, 10.1073/pnas.1404853111, 2014.
- 395 Laskin, A., Laskin, J., and Nizkorodov, S. A.: Chemistry of Atmospheric Brown Carbon, *Chemical Reviews*, 115, 4335-4382, 10.1021/cr5006167, 2015.
- Lavi, A., Segre, E., Gomez-Hernandez, M., Zhang, R., and Rudich, Y.: Volatility of Atmospherically Relevant Alkylammonium Carboxylate Salts, *The Journal of Physical Chemistry A*, 119, 4336-4346, 10.1021/jp507320v, 2015.
- 400 Luo, H., Chen, J., Li, G., and An, T.: Formation kinetics and mechanisms of ozone and secondary organic aerosols from photochemical oxidation of different aromatic hydrocarbons: dependence on NO_x and organic substituents, *Atmospheric Chemistry and Physics*, 21, 7567-7578, 10.5194/acp-21-7567-2021, 2021.
- Luo, H., Li, G., Chen, J., Wang, Y., and An, T.: Reactor characterization and primary application of a state of art dual-reactor chamber in the investigation of atmospheric photochemical processes, *J Environ Sci (China)*, 98, 161-168, 10.1016/j.jes.2020.05.021, 2020.
- 405 Ma, X., Zhao, X., Wei, Y., Wang, W., Xu, F., Zhang, Q., and Wang, W.: Effect of multifunctional compound monoethanolamine on Criegee intermediates reactions and its atmospheric implications, *Sci Total Environ*, 715, 136812, 10.1016/j.scitotenv.2020.136812, 2020.
- Moise, T., Flores, J. M., and Rudich, Y.: Optical Properties of Secondary Organic Aerosols and Their Changes by Chemical Processes, *Chemical Reviews*, 115, 4400-4439, 10.1021/cr5005259, 2015.
- Mull, H. F., Aroeira, G. J. R., Turney, J. M., and Schaefer, H. F.: The atmospheric importance of methylamine additions to Criegee intermediates, *Phys Chem Chem Phys*, 22, 22555-22566, 10.1039/d0cp03781h, 2020.
- 410 Na, K., Song, C., and Cockeriii, D.: Formation of secondary organic aerosol from the reaction of styrene with ozone in the presence and absence of ammonia and water, *Atmospheric Environment*, 40, 1889-1900, 10.1016/j.atmosenv.2005.10.063, 2006.
- Ng, N. L., Brown, S. S., Archibald, A. T., Atlas, E., Cohen, R. C., Crowley, J. N., Day, D. A., Donahue, N. M., Fry, J. L., Fuchs, H., Griffin, R. J., Guzman, M. I., Herrmann, H., Hodzic, A., Iinuma, Y., Jimenez, J. L., Kiendler-Scharr, A., Lee, B. H., Luecken, D. J., Mao, J., McLaren, R., Mutzel, A., Osthoff, H. D., Ouyang, B., Picquet-Varrault, B., Platt, U., Pye, H. O. T., Rudich, Y., Schwantes, R. H., Shiraiwa, M., Stutz, J., Thornton, J. A., Tilgner, A., Williams, B. J., and Zaveri, R. A.: Nitrate radicals and biogenic volatile organic compounds: oxidation, mechanisms, and organic aerosol, *Atmos Chem Phys*, 17, 2103-2162, 10.5194/acp-17-2103-2017, 2017.
- 415 Nguyen, T. L., Winterhalter, R., Moortgat, G., Kanawati, B., Peeters, J., and Vereecken, L.: The gas-phase ozonolysis of beta-carophyllene (C(15)H(24)). Part II: A theoretical study, *Phys Chem Chem Phys*, 11, 4173-4183, 10.1039/b817913a, 2009.
- Perring, A. E., Pusede, S. E., and Cohen, R. C.: An observational perspective on the atmospheric impacts of alkyl and multifunctional nitrates on ozone and secondary organic aerosol, *Chem Rev*, 113, 5848-5870, 10.1021/cr300520x, 2013.
- 420 Qiu, C. and Zhang, R. Y.: Multiphase chemistry of atmospheric amines, *Physical Chemistry Chemical Physics*, 15, 5738-5752, 10.1039/c3cp43446j, 2013.
- Qiu, J., Tonokura, K., and Enami, S.: Proton-Catalyzed Decomposition of alpha-Hydroxyalkyl-Hydroperoxides in Water, *Environ Sci Technol*, 54, 10561-10569, 10.1021/acs.est.0c03438, 2020a.
- 425 Qiu, J., Fujita, M., Tonokura, K., and Enami, S.: Stability of Terpenoid-Derived Secondary Ozonides in Aqueous Organic Media, *The Journal of Physical Chemistry A*, 126, 5386-5397, 10.1021/acs.jpca.2c04077, 2022.
- Qiu, J., Ishizuka, S., Tonokura, K., and Enami, S.: Reactions of Criegee Intermediates with Benzoic Acid at the Gas/Liquid Interface, *J Phys Chem A*, 122, 6303-6310, 10.1021/acs.jpca.8b04995, 2018a.
- 430 Qiu, J., Ishizuka, S., Tonokura, K., Colussi, A. J., and Enami, S.: Reactivity of Monoterpene Criegee Intermediates at Gas-Liquid Interfaces, *J Phys Chem A*, 122, 7910-7917, 10.1021/acs.jpca.8b06914, 2018b.
- Qiu, J., Ishizuka, S., Tonokura, K., Colussi, A. J., and Enami, S.: Water Dramatically Accelerates the Decomposition of alpha-Hydroxyalkyl-Hydroperoxides in Aerosol Particles, *J Phys Chem Lett*, 10, 5748-5755, 10.1021/acs.jpcl.9b01953, 2019.
- 435 Qiu, J., Liang, Z., Tonokura, K., Colussi, A. J., and Enami, S.: Stability of Monoterpene-Derived alpha-Hydroxyalkyl-Hydroperoxides in Aqueous Organic Media: Relevance to the Fate of Hydroperoxides in Aerosol Particle Phases, *Environ Sci Technol*, 54, 3890-3899, 10.1021/acs.est.9b07497, 2020b.
- Sanchez, J. and Myers, T. N.: Peroxides and peroxide compounds, organic peroxides, Kirk-Othmer Encyclopedia of Chemical Technology, doi.org/10.1002/0471238961.1518070119011403.a01, 2000.
- Seinfeld, J. H., Bretherton, C., Carslaw, K. S., Coe, H., DeMott, P. J., Dunlea, E. J., Feingold, G., Ghan, S., Guenther, A. B., Kahn, R., Kraucunas, I., Kreidenweis, S. M., Molina, M. J., Nenes, A., Penner, J. E., Prather, K. A., Ramanathan, V., Ramaswamy, V., Rasch, P. J., Ravishankara, A. R., Rosenfeld, D., Stephens, G., and Wood, R.: Improving our fundamental understanding of the role of aerosol-cloud interactions in the climate system, *P Natl Acad Sci USA*, 113, 5781-5790, 10.1073/pnas.1514043113, 2016.
- 440 Shiraiwa, M., Ueda, K., Pozzer, A., Lammel, G., Kampf, C. J., Fushimi, A., Enami, S., Arangio, A. M., Fröhlich-Nowoisky, J., Fujitani, Y., Furuyama, A., Lakey, P. S. J., Lelieveld, J., Lucas, K., Morino, Y., Pöschl, U., Takahama, S., Takami, A., Tong, H., Weber, B., Yoshino,


- 445 A., and Sato, K.: Aerosol Health Effects from Molecular to Global Scales, *Environmental Science & Technology*, 51, 13545-13567, 10.1021/acs.est.7b04417, 2017.
- Shrivastava, M., Cappa, C. D., Fan, J. W., Goldstein, A. H., Guenther, A. B., Jimenez, J. L., Kuang, C., Laskin, A., Martin, S. T., Ng, N. L., Petaja, T., Pierce, J. R., Rasch, P. J., Roldin, P., Seinfeld, J. H., Shilling, J., Smith, J. N., Thornton, J. A., Volkamer, R., Wang, J., Worsnop, D. R., Zaveri, R. A., Zelenyuk, A., and Zhang, Q.: Recent advances in understanding secondary organic aerosol: Implications for global climate forcing, *Rev Geophys*, 55, 509-559, 10.1002/2016rg000540, 2017.
- 450 Sugimura, N., Furuya, A., Yatsu, T., and Shibue, T.: Prediction of adducts on positive mode electrospray ionization mass spectrometry: proton/sodium selectivity in methanol solutions, *Eur J Mass Spectrom* (Chichester), 21, 725-731, 10.1255/ejms.1389, 2015.
- Tang, X., Price, D., Praske, E., Lee, S. A., Shattuck, M. A., Purvis-Roberts, K., Silva, P. J., Asa-Awuku, A., and Cocker, D. R.: NO₃ radical, OH radical and O₃-initiated secondary aerosol formation from aliphatic amines, *Atmospheric Environment*, 72, 105-112, 10.1016/j.atmosenv.2013.02.024, 2013.
- 455 Tong, D., Chen, J., Qin, D., Ji, Y., Li, G., and An, T.: Mechanism of atmospheric organic amines reacted with ozone and implications for the formation of secondary organic aerosols, *Sci Total Environ*, 737, 139830, 10.1016/j.scitotenv.2020.139830, 2020.
- Tong, H., Arangio, A. M., Lakey, P. S. J., Berkemeier, T., Liu, F., Kampf, C. J., Brune, W. H., Pöschl, U., and Shiraiwa, M.: Hydroxyl radicals from secondary organic aerosol decomposition in water, *Atmospheric Chemistry and Physics*, 16, 1761-1771, 10.5194/acp-16-1761-2016, 2016.
- 460 Vibenholt, A., Norgaard, A. W., Clausen, P. A., and Wolkoff, P.: Formation and stability of secondary ozonides from monoterpenes studied by mass spectrometry, *Chemosphere*, 76, 572-577, 10.1016/j.chemosphere.2009.02.060, 2009.
- Wang, P.-B., Truhlar, D. G., Xia, Y., and Long, B.: Temperature-dependent kinetics of the atmospheric reaction between CH₂OO and acetone, *Phys. Chem. Chem. Phys.*, 24, 13066-13073, 10.1039/D2CP01118B, 2022.
- Wang, S., Liu, T., Jang, J., Abbatt, J. P. D., and Chan, A. W. H.: Heterogeneous interactions between SO₂ and organic peroxides in submicron aerosol, *Atmos. Chem. Phys.*, 21, 6647-6661, 10.5194/acp-21-6647-2021, 2021.
- 465 Wang, S., Zhao, Y., Chan, A. W. H., Yao, M., Chen, Z., and Abbatt, J. P. D.: Organic Peroxides in Aerosol: Key Reactive Intermediates for Multiphase Processes in the Atmosphere, *Chemical Reviews*, 123, 1635-1679, 10.1021/acs.chemrev.2c00430, 2023.
- Wang, S., Zhou, S., Tao, Y., Tsui, W. G., Ye, J., Yu, J. Z., Murphy, J. G., McNeill, V. F., Abbatt, J. P. D., and Chan, A. W. H.: Organic Peroxides and Sulfur Dioxide in Aerosol: Source of Particulate Sulfate, *Environ Sci Technol*, 53, 10695-10704, 10.1021/acs.est.9b02591, 2019.
- 470 Wei, J., Fang, T., Lakey, P. S. J., and Shiraiwa, M.: Iron-Facilitated Organic Radical Formation from Secondary Organic Aerosols in Surrogate Lung Fluid, *Environmental Science & Technology*, 56, 7234-7243, 10.1021/acs.est.1c04334, 2022.
- Winterhalter, R., Neeb, P., Grossmann, D., Kolloff, A., Horie, O., and Moortgat, G.: Products and mechanism of the gas phase reaction of ozone with beta-pinene, *J Atmos Chem*, 35, 165-197, Doi 10.1023/A:1006257800929, 2000.
- 475 Winterhalter, R., Herrmann, F., Kanawati, B., Nguyen, T. L., Peeters, J., Vereecken, L., and Moortgat, G. K.: The gas-phase ozonolysis of beta-caryophyllene (C(15)H(24)). Part I: an experimental study, *Phys Chem Chem Phys*, 11, 4152-4172, 10.1039/b817824k, 2009.
- Yao, L., Ma, Y., Wang, L., Zheng, J., Khalizov, A., Chen, M. D., Zhou, Y. Y., Qi, L., and Cui, F. P.: Role of stabilized Criegee Intermediate in secondary organic aerosol formation from the ozonolysis of alpha-cedrene, *Atmos. Environ.*, 94, 448-457, 10.1016/j.atmosenv.2014.05.063, 2014.
- 480 Yao, L., Wang, M. Y., Wang, X. K., Liu, Y. J., Chen, H. F., Zheng, J., Nie, W., Ding, A. J., Geng, F. H., Wang, D. F., Chen, J. M., Worsnop, D. R., and Wang, L.: Detection of atmospheric gaseous amines and amides by a high-resolution time-of-flight chemical ionization mass spectrometer with protonated ethanol reagent ions, *Atmospheric Chemistry and Physics*, 16, 14527-14543, 10.5194/acp-16-14527-2016, 2016.
- Yao, M., Zhao, Y., Hu, M., Huang, D., Wang, Y., Yu, J. Z., and Yan, N.: Multiphase Reactions between Secondary Organic Aerosol and Sulfur Dioxide: Kinetics and Contributions to Sulfate Formation and Aerosol Aging, *Environmental Science & Technology Letters*, 6, 768-774, 10.1021/acs.estlett.9b00657, 2019.
- 485 Yu, H., McGraw, R., and Lee, S.-H.: Effects of amines on formation of sub-3 nm particles and their subsequent growth, *Geophysical Research Letters*, 39, n/a-n/a, 10.1029/2011gl050099, 2012.
- Zeng, M., Heine, N., and Wilson, K. R.: Evidence that Criegee intermediates drive autoxidation in unsaturated lipids, *Proc. Natl. Acad. Sci. U. S. A.*, 117, 4486-4490, doi:10.1073/pnas.1920765117, 2020.
- 490 Zhang, X., Barraza, K. M., and Beauchamp, J. L.: Cholesterol provides nonsacrificial protection of membrane lipids from chemical damage at air-water interface, *Proc. Natl. Acad. Sci. U. S. A.*, 115, 3255-3260, doi:10.1073/pnas.1722323115, 2018.
- Zhou, Z., Lakey, P. S. J., von Domaros, M., Wise, N., Tobias, D. J., Shiraiwa, M., and Abbatt, J. P. D.: Multiphase Ozonolysis of Oleic Acid-Based Lipids: Quantitation of Major Products and Kinetic Multilayer Modeling, *Environ. Sci. Technol.*, 56, 7716-7728, 10.1021/acs.est.2c01163, 2022.
- 495



500 **Figure 1** Effects of EA on size and chemical composition of particles produced by the ozonolysis of β -C. (A) Number concentrations of particles, (B) Volume concentrations of total particles, (C) Positive-ion ESI mass spectra of the chemical components of particles extracted in AN/W (vol/vol = 4/1) solutions. Blue dots and red dots represent three independent experiments



505 Figure 2 High-resolution positive-ion ESI mass spectra of P1 extracted in AN/W (vol/vol = 4/1), AN/D₂O (vol/vol = 4/1) and AN/H₂¹⁸O (vol/vol = 4/1) solutions



510 **Figure 3** Positive-ion ESI mass spectra of the products extracted in AN/W (vol/vol = 4/1) from ozonolysis of β -C in the absence/presence of MA, DMA, or ammonia




Figure 4 Positive-ion ESI mass spectra of the products extracted in AN/W (vol/vol = 4/1) from ozonolysis of $\alpha\text{-H}$ in the absence/presence of MA, EA, DMA, or ammonia

515






Figure 5 High-resolution positive-ion ESI mass spectra of P3 extracted in AN/W (vol/vol = 4/1), AN/D₂O (vol/vol = 4/1) and AN/H₂¹⁸O (vol/vol = 4/1) solutions



Scheme 2 Possible structure of P1 generated from the reaction of β -C SOZ with EA



525

Scheme 3 Possible structure of P3 generated from the reaction of α -H SOZ with EA

