

# Moisture and temperature effects on the radiocarbon signature of respired carbon dioxide to assess stability of soil carbon in the Tibetan Plateau

Andrés Tangarife-Escobar<sup>1</sup>, Georg Guggenberger<sup>2</sup>, Xiaojuan Feng<sup>3</sup>, Guohua Dai<sup>3</sup>, Carolina Urbina-Malo<sup>2</sup>, Mina Azizi-Rad<sup>1</sup>, and Carlos A. Sierra<sup>1,4</sup>

<sup>1</sup>Max Planck Institute for Biogeochemistry, Jena, Germany

<sup>2</sup>Institute of Soil Science, Leibniz Universität Hannover, Hannover, Germany

<sup>3</sup>Institute of Botany, Chinese Academy of Sciences, Beijing, China

<sup>4</sup>Department of Ecology, Swedish University of Agricultural Sciences, Uppsala, Sweden

**Correspondence:** Andrés Tangarife-Escobar (atanga@bgc-jena.mpg.de)

**Abstract.** Microbial release of CO<sub>2</sub> from soils to the atmosphere reflects how environmental conditions affect the stability of soil organic matter (SOM), especially in massive organic-rich ecosystems like the peatlands and grasslands of the Qinghai-Tibetan Plateau (QTP). Radiocarbon (<sup>14</sup>C) is an important tracer of the global carbon cycle and can be used to understand SOM dynamics through the estimation of time lags between ~~C-carbon~~ fixation and respiration, often assessed with metrics such as age and transit time. In this study, we incubated peatland and grassland soils at four temperature (5, 10, 15 and 20 °C) and two water-filled pore space (WFPS) levels (60 and 95 %), and measured the <sup>14</sup>C signature of bulk soil and respired CO<sub>2</sub>. We ~~compare-compared~~ the relation between the  $\Delta^{14}\text{C}$  of the bulk soil and the  $\Delta^{14}\text{CO}_2$  of respired carbon as a function of temperature and WFPS for the two soils. To better interpret our results, we used a mathematical model to analyse how the calculated number of pools, decomposition rates of carbon ( $k$ ), transfer ( $\alpha$ ) and partitioning ( $\gamma$ ) coefficients affect the  $\Delta^{14}\text{C}$ -bulk and  $\Delta^{14}\text{CO}_2$  relation, with their respective mean age and mean transit time. From our incubations, we found that <sup>14</sup>C ~~from peatland-was~~ values in bulk and CO<sub>2</sub> from peatland were significantly more depleted (old) than from grassland soil. Our results showed that changes in temperature did not affect the  $\Delta^{14}\text{C}$  values of respired CO<sub>2</sub> in either soil. However, changes in WFPS had a small effect on the <sup>14</sup>CO<sub>2</sub> in grassland soils and a ~~strong-significant~~ influence in peatland soils, where higher WFPS levels led to more depleted  $\Delta^{14}\text{CO}_2$ . In our models, ~~we-observed-large-differences-between-slow-and-fast-cycling-systems,~~ where low values of  $k$  modified  $\Delta^{14}\text{C}$  patterns due to the incorporation of <sup>14</sup>C-bomb in the soil. Hence, the ~~the~~ correspondence between  $\Delta^{14}\text{C}$  ~~and~~ age and transit time ~~strongly-highly~~ depended on the internal dynamics of the soil ( $k$ ,  $\alpha$ ,  $\gamma$  and number of pools) as well as on model structure. We ~~conclude-observed large differences between slow and fast cycling systems, where low values of decomposition rates modified the  $\Delta^{14}\text{C}$  values in a non-linear pattern due to the incorporation of modern carbon (<sup>14</sup>C-bomb) in the soil. We concluded~~ that the stability of carbon in ~~these-systems-the peatland and grassland soils of the QTP~~ depends strongly on the direction of change in ~~temperature-and~~ moisture and how it affects the rates of SOM decomposition. ~~Finally,  $\Delta^{14}\text{C}$  modelling along with empirical data from SOM dynamics is a useful tool to improve predictions on interactions between terrestrial and atmospheric carbon-while temperature regulates the amount of fluxes. Current land cover modification~~

(desiccation) in Zoige peatlands and climate change occurring on the QTP, might largely increase CO<sub>2</sub> fluxes along with the release of old carbon to the atmosphere potentially shifting carbon sinks into sources.

## 25 1 Introduction

Studying soil organic matter (SOM) stability and persistence in a globally-changing environment is of fundamental importance to understand temporal variations of carbon cycling in the earth-climate system. Soil constitutes the largest carbon (C) stock in the terrestrial biosphere (Chen et al., 2021), making it a key component in global climate models (McGuire et al., 2001; Wieder et al., 2013; Xu et al., 2016). Physical, chemical and biological properties determine SOM decomposition, and in consequence  
30 its persistence over decades to millennia (Schmidt et al., 2011). Temperature and moisture are two of the most important abiotic variables controlling the rates of SOM decomposition (Sierra et al., 2015) and its transit time across the different ecosystem pools. Changes in these environmental conditions are, however, occurring simultaneously, highlighting the necessity to conduct multifactorial experiments to disentangle the dominant mechanisms on such cycling times. Hence, discerning the relationship between SOM persistence and C-carbon cycling times in terrestrial ecosystems is imperative to improve climate change models  
35 and to inform decisions on CO<sub>2</sub> mitigation and management strategies (Bradford et al., 2016; Mesfin et al., 2021).

The Qinghai-Tibetan Plateau (QTP) with an area of 2.5 million km<sup>2</sup> is a macroregion of global importance for the cycling of water, carbon and other biogeochemical elements (Anslan et al., 2020). Land cover is mainly dominated by alpine grasslands (44 %) (Scurlock and Hall, 1998; Gao et al., 2014), which store 23.4 % of China's total organic carbon and 2.5 % of the global  
40 soil carbon (Genxu et al., 2002). The QTP also hosts one of the largest high-mountain marshes in the world (Xiang et al., 2009; Chen et al., 2014; Ma et al., 2016), the Zoige peatlands, considered to be the most important carbon stock in peatlands for China (Liu et al., 2018). However, climate change and land cover change are currently driving-modulating the net carbon balance through CO<sub>2</sub> and CH<sub>4</sub> effluxes from the grasslands (Piao et al., 2012; Chen et al., 2014, 2017; Du and Gao, 2020)  
(Piao et al., 2012; Chen et al., 2017; Du and Gao, 2020) and the peatlands at both daily and interannual timescales (Hao et al., 2011; Kang  
45 (Hao et al., 2011; Chen et al., 2014; Kang et al., 2018; Liu et al., 2019b; Chen et al., 2021)). Anthropogenic activities in the Zoige peatlands (drainage, peat mining and overgrazing) have caused a degradation of approximately 30 % of wet and dry meadows (Chen et al., 2014; Zhou et al., 2021). Additionally, the QTP has been facing an air temperature increase of 0.2°C per decade over the past 50 years (Zhang et al., 2013; Chen et al., 2014; Yang et al., 2014; Ganjurjav et al., 2016), equivalent to two to three times faster than the world average (Yao et al., 2019; Nieberding et al., 2020), along with a moderate increase in precipitation  
50 (Dong et al., 2018). According to the IPCC (Arias et al., 2021), climate models predict an increment in heavy precipitation events and high temperature extremes for this region.

Such changes in temperature and soil moisture control the magnitude in which stabilization and de-stabilization-destabilization mechanisms enable carbon storage in the QTP (Xiang et al., 2009; Ma et al., 2016). It has been observed that warming increases  
55 soil respiration (Rustad et al., 2001; Lu et al., 2013; Pold et al., 2015) and that soil moisture modulates ecosystem and soil respi-

ration (Geng et al., 2012; Piao et al., 2012; Pan et al., 2022; Azizi-Rad et al., 2022) (Geng et al., 2012; Piao et al., 2012; Moyano et al., 2012). Although the influence of destabilization mechanisms on SOM decomposition has been already studied in the Zoige peatlands (Zhao et al., 2011; Wang et al., 2015; Liu et al., 2016, 2018; Li et al., 2018; Liu et al., 2019a), temporal scales of SOM persistence under temperature and soil moisture variations are not clearly understood yet. Therefore, a quantification of the SOM persistence is of vital importance to predict climate change feedback magnitudes and pathways.

A useful approach to quantify SOM persistence is through the theory of compartmental dynamical systems, where soil is understood as a set of interconnected pools with transformations of  $\text{C-carbon}$  to different forms (gas, dissolved or solid) (Emanuel et al., 1984; Schimel, 1995; Sierra et al., 2018a). Therefore,  $\text{C-carbon}$  can be characterized by the time it remains inside a compartment or the entire system (Eriksson, 1971; Bolin and Rodhe, 1973) through the calculation of age and transit time. Here, age is defined as the time elapsed since the  $\text{C-carbon}$  entered the system until the time of observation in the bulk soil, while transit time is defined as the time spent by the  $\text{C-carbon}$  between the entry to the system and its exit as respired  $\text{CO}_2$  (Eriksson, 1971; Bolin and Rodhe, 1973; Manzoni et al., 2009; Sierra et al., 2018b). These ~~times ( $T$ )~~ timescale metrics are however, not the same for all the  $\text{C-carbon}$  atoms in a soil, since physical, chemical and biological stabilization and destabilization processes of SOM modify the rate at which mixing and storage occurs.

Radiocarbon measurements in bulk soil and in respired  $\text{CO}_2$  are a powerful tool to approximate ages and transit times of  $\text{C-carbon}$  in soils as they trace back the trajectory of  $\text{C-carbon}$  through the different stocks on decadal to millennia timescales (Trumbore, 2000; Sierra et al., 2014; Schuur et al., 2016; Estop-Aragones et al., 2020; Chen et al., 2021). The radiocarbon content reflects the time a carbon atom has been in the soil since it was fixed by photosynthesis from the atmosphere (Trumbore, 2000). Carbon fixed since the early 1960's has higher  $^{14}\text{C}$  than carbon fixed previously due to the enrichment by thermonuclear weapons testing. By comparing the  $^{14}\text{C}$  relative abundance in the carbon pools with the atmospheric  $^{14}\text{CO}_2$  concentrations ~~produced after the nuclear weapon tests during the early 1960's~~, it is possible to model soil-atmosphere  $\text{C-carbon}$  cycling times (Trumbore, 2000).

$^{14}\text{C}$  values can be altered by SOM ~~(de)-stabilization-destabilization~~ processes and soil characteristics such as soil organic carbon (SOC) content, age and diagenetic state, thaw depth, redox state, seasonality, etc. (Gaudinski et al., 2000; Trumbore, 2000; von Lützow et al., 2008; Sierra et al., 2018b; Estop-Aragones et al., 2020). In recent decades, it has been widely observed that temperature plays a major role on the dynamics of SOM (Knorr et al., 2005; Davidson and Janssens, 2006; Feng and Simpson, 2008) and the mean age of respired  $\text{CO}_2$  (Hopkins et al., 2012; Chen et al., 2021) by increasing the decomposition rates from fast-cycling pools (Trumbore et al., 1996) and mobilizing old  $\text{C-carbon}$  in warming conditions (Dutta et al., 2006; Briones et al., 2010). In contrast, other studies have suggested that warming does not lead to release of old  $\text{C-carbon}$  (Briones et al., 2021; Dioumaeva et al., 2002) and that non-labile  $\text{C-carbon}$  decomposition is insensitive to temperature increase (Liski et al., 1999). By comparison, drying phenomena increased the release of modern  $^{14}\text{CO}_2$  from shallow soil layers but preserved the old soil carbon pools in deeper layers (Kwon et al., 2019). Observations in rich-SOC soils of the Arctic indicated the release

of old ~~C-carbon~~ from deep soil layers after thaw and drainage processes (~~Schuur et al., 2009; Estop-Aragones et al., 2020~~) (~~Schuur et al., 2009; Estop-Aragones et al., 2020; Pegoraro et al., 2021~~). Such a process might be occurring in the Zoige peatland soils due to the ~~seasonal-presence-of~~presence of seasonal frozen layers (Liu et al., 2021; Yang et al., 2022). Nonetheless, the influence of temperature and soil moisture on SOM persistence is a matter of current debate.

95

Changes in temperature and moisture contribute to the destabilization of carbon in soils from the QTP. Hence, we hypothesize that higher temperature ~~and moisture levels would contribute to increasing the~~ would increase the age of respired CO<sub>2</sub> as well as changes in soil moisture would increase or decrease (depending on the direction of moisture change) the age of respired CO<sub>2</sub> ~~for in~~ soils subjected to controlled manipulations. Greater ages of respired CO<sub>2</sub> would imply that previously stabilized ~~C~~ carbon is destabilized by changes in the manipulated environmental factors. For this purpose, we posed three specific research questions: (1) Do specific changes in temperature and moisture result in the release of old carbon in the respiration flux? (2) Are there differences in the age of respired CO<sub>2</sub> between a grassland and a peatland soil from the QTP? (3) How can radiocarbon data in bulk soil and respired CO<sub>2</sub> be interpreted to understand the effect of changes in decomposition rates on ages and transit times? To address these questions, we conducted a controlled multifactorial experiment with soils from two different ecosystems in the QTP, and measured ~~their~~the radiocarbon signature in the bulk soil and the respired CO<sub>2</sub>. In addition, we used a mathematical model to better interpret ~~our results with the concepts of the~~ interaction between decomposition rates change, expressed through internal dynamics of the soil ( $k$ ,  $\alpha$ ,  $\gamma$  and number of pools) and the  $\Delta^{14}\text{C}$  values by targeting the range found in the incubations. Thus, our observations and models strengthen each other to gain a deeper comprehension on the relationship between soil carbon stability,  $\Delta^{14}\text{C}$ , age and transit timeof carbon.

110

## 2 Materials and methods

To evaluate the influence of soil moisture and temperature on the  $^{14}\text{C}$  values from bulk soil and respired CO<sub>2</sub> of grasslands and peatlands soils, we incubated soil samples from a high elevation grassland at the Nam Co catchment collected in September 2018 and a peatland from the Zoige region sampled in July 2021, both located in the QTP.


### 115 2.1 Site description and sampling

#### 2.1.1 Nam Co grassland

The Nam Co site (Fig. 1) is located in the central Tibetan Plateau (altitude of the Nam Co lake 4,726 m a.s.l., 30°46'N, 90°59'E) and represents a frequent study location for monitoring and tracking of environmental changes over different time scales (Anslan et al., 2020). The dominant climate at Nam Co consists of cold winters and short and moist summers. Mean annual temperature (MAT) measured at the NAMORS research station was -0.6 °C (2006 – 2017) and mean annual precipitation (MAP) was 406 mm, occurring mostly during the monsoon season between May and October (Anslan et al., 2020). The sam-

120

pling area was mainly covered by *Kobresia pygmaea* and has been grazed intensively by yaks and sheep. A total amount of 40 kg of soil was collected from randomly selected spots at depths between 5 and 15 cm within an area of about 40 m<sup>2</sup>. The soil was then mixed homogeneously to produce one single sample which consisted of a sandy loam with a bulk density of 1.3 g cm<sup>-3</sup>, pH of 7.5, 3.2 % of total organic carbon, 0.05 % of inorganic carbon and 0.3 % of total nitrogen, cation exchange capacity of 89 mmol<sub>c</sub> kg<sup>-1</sup> soil and electrical conductivity of 89.6 μS cm<sup>-1</sup>.



**Figure 1.** a) Overview of the QTP; b) Location of the Nam Co lake with land cover dominated by grasslands, and c) Zoigê peatlands highlighting the location of Zoigê city. The sampling site is located in the southeastern of the peatland (Ruokeba, Hongyuan county). SRTM elevation data from Jarvis et al. (2008).

### 2.1.2 Zoigê peatlands

The Zoigê peatlands are located in the northeast of the QTP (average elevation of 3,400 m a.s.l.) and cover an area of about 4,605 km<sup>2</sup> on the headwaters of the Yellow River basin. These peatlands have been recognised as one of the largest high-mountain marshes in the world (Xiang et al., 2009; Ma et al., 2016) and store 88 % of the carbon in the QTP (Chen et al., 2014). The Zoigê region shows a MAT of 1.5 °C, where the warmest monthly temperature (11 °C) is recorded in July and

the coldest in January (-10.1 °C). The MAP is 720 mm, and according to data from the Chinese National Meteorological Information Centre (www.nmic.gov.cn), the region has faced a slight drying trend and a warming of 0.4 °C per decade over the last 40 years (Chen et al., 2014; Yang et al., 2014). The soils are unfrozen from April to October while the layer between 0 and 50 cm depth is seasonally frozen between November and March. Vegetation cover consists mainly of *Potentilla anserine*, *Blysmus sinocompressus*, *Kobresia myosuroides* and *Scirpus triqueter*. The samples (0 – 35 cm depth) were collected with a spade from an area of 50 m<sup>2</sup> at a long-term monitoring site (33°4'5''N, 102°33'52''E), then thoroughly mixed to produce a single sample. The soil bulk density was 0.3 g ~~cm<sup>3</sup>~~cm<sup>-3</sup>, contained in average 27.6 % of organic carbon, 0.06 % of inorganic carbon and 1.8 % of total nitrogen.

## 140 2.2 Incubation of peatland and grassland soils

We conducted two sets of incubation experiments, one set with the grassland soil and a second set with the peatland soil. Samples were incubated with their original roots to minimise disturbance and allow comparisons with field conditions, however stones were removed. Each of the sets ~~were was~~ placed at two different WFPS ~~values levels~~ (60 and 95 %) which were selected in order to reassemble the thaw and consequent water saturation of seasonally frozen soils in grasslands; and for peatland soils, the process of drying (through artificial desiccation) after high water saturation. Soil moisture treatments were combined with four temperature levels (5, 10, 15 and 20 °C) for a total of 69 samples, 33 for grassland and 36 for peatland (2 sites × 4 temperature levels × 2 moisture levels × ~~4-5-3-6 analytical~~ replicates, to prevent scarcity of data due to eventual failure in CO<sub>2</sub> extraction), ~~totaling 33 for grassland and 36 for peatland~~. CO<sub>2</sub> concentrations were measured at intervals of 1 to 2 weeks using a CO<sub>2</sub> analyzer LI-COR 6262. Incubations for each subset ended simultaneously until every sample had an estimated concentration of CO<sub>2</sub>-C in the headspace equivalent to ≥ 0.2-2 mg of C, enough for radiocarbon analysis. Grassland samples were incubated between 15 and 67 days, while peatland samples were incubated for 13 days (Table A1). For sampling headspace air, 50-ml vials were filled with 12 g of soil (± 1.5 g) and placed inside 0.5 L glass flasks along with 0.2 ml of water at the bottom of the flask (away from contact with the sample) to avoid possible drying (Dioumaeva et al., 2002); thereafter the flasks were sealed with rubber plugs and screwed with plastic caps. Flasks with samples were flushed with synthetic air (CO<sub>2</sub> free) to remove atmospheric CO<sub>2</sub>. This flushing marked the starting day of the incubations.

## 2.3 Radiocarbon analysis of incubated soils

~~Respired~~ Radiocarbon analysis were conducted in the bulk soil of each sample after the incubation. Soil inorganic carbon was eliminated through decalcification following the preparation protocol by Steinhof et al. (2017). Also, respired CO<sub>2</sub> accumulated in the headspace of incubation flasks was extracted and purified on a vacuum line, graphitized by Fe reduction in H<sub>2</sub> and measured for Δ<sup>14</sup>C by Accelerator Mass Spectrometry (Micadas, Ionplus, Switzerland) in the radiocarbon laboratory of the Max Planck Institute for Biogeochemistry in Jena, Germany (Steinhof et al., 2017).

165 The radiocarbon content reflects the time a C atom has been in the soil since it was fixed by photosynthesis from the atmosphere (Trumbore, 2000). Carbon fixed since the early 1960's has higher  $^{14}\text{C}$  than carbon fixed previously due to the enrichment by thermonuclear weapons testing. Samples with significant  $^{14}\text{C}$  positive values show incorporation by the bomb  $^{14}\text{C}$ , presenting  $^{14}\text{C}/^{12}\text{C}$  ratios greater than that of the 1890 wood used as  $^{14}\text{C}$  standard for pre-bomb periods. Radiocarbon data are expressed as  $\Delta^{14}\text{C}$  (the deviation in ‰ from 0.95 times the oxalic acid standard in 1950). The values were corrected to a  $\delta^{13}\text{C}$  value of -25 ‰ for differences in biological mass-dependent fractionation (Stuiver and Polach, 1977). These data are  
 170 presented as percentage of Modern Carbon (pMC) that can be converted to  $F^{14}\text{C}$  by dividing it by 100 and later to  $\Delta^{14}\text{C}$  using equation (1) (Stuiver and Polach, 1977).

$$\Delta^{14}\text{C} = \left[ F^{14}\text{C} e^{\lambda_C(1950-t)} - 1 \right] \times 1000 [\text{‰}], \quad (1)$$

where  $F^{14}\text{C}$  is the Fraction Modern, i.e. the ratio of the measured sample normalized to a  $\delta^{13}\text{C}$  value of -25 ‰, divided by  
 175 0.95 times the measured ratio of the Oxalic Acid I standard (OX-I) (Schuur et al., 2016),  $\lambda_C$  is the updated radiocarbon decay constant (equals  $1/8267 [\text{y}^{-1}]$ ), and  $t$  is the year of sampling.

The effects of soil moisture and temperature manipulation on  $\Delta^{14}\text{C}$  were evaluated for the bulk and  $\text{CO}_2$  fractions in each ecosystem separately through two-way ANOVA tests (type III). This type of ANOVA is also referred to as Partial Sum of Squares and is appropriate for unbalanced data since it does not depend on the sampling structure or the particular order in the model (Shaw and Mitchell-Olds, 1993); hence, this approach adjusts best to our data set where treatments did not have equal amount of values. As for the  $\text{CO}_2$  fluxes, rates were measured regularly for every treatment and mean  $\text{CO}_2$  respiration rates ( $\text{mg CO}_2 \text{ g soil}^{-1} \text{ day}^{-1}$ ) were calculated based on the total duration of the incubation (Fig. A1).

## 2.4 SOC decomposition models to predict $\Delta^{14}\text{C}$ as a proxy of SOM persistence

185 The representation of the SOC dynamics has been commonly described through models that can be expressed as systems of linear differential equations (Manzoni et al., 2009; Sierra et al., 2012; Sierra and Mueller, 2015; Sierra et al., 2017b) of the form:

$$\frac{d\mathbf{C}(t)}{dt} = \mathbf{I} + \mathbf{A} \cdot \mathbf{C}(t), \quad (2)$$

where the vector  $\mathbf{C}(t)$  is the rate of change of carbon over time in  $n$  different pools; the time dependent  $n$ -dimensional vector  
 190  $\mathbf{I}$  represents the total inputs of carbon to each pool; and  $\mathbf{A}$  represents the  $n \times n$  dimensional matrix with the rates of carbon processing for each pool in its main diagonal and the proportion of carbon transferred from one pool to another in the off-diagonals (Sierra et al., 2012, 2014; Metzler and Sierra, 2018; Sierra et al., 2018b). This mass balance equation has a radiocarbon counterpart:

$$\frac{d^{14}\mathbf{C}(t)}{dt} = \mathbf{I}_{14}\mathbf{C}(t) + \mathbf{A} \cdot {}^{14}\mathbf{C}(t) - \lambda {}^{14}\mathbf{C}(t) \quad (3)$$



195 where  $\lambda$  is the radiocarbon decay constant (1/8267 [y<sup>-1</sup>]).

We used a SOC decomposition model to calculate the  $\Delta^{14}\text{C}$  in bulk and  $\text{CO}_2$  as well as their equivalent mean age and mean transit time of a theoretical soil for the year 2019 calibrated with the data set for the Northern Hemisphere Zone 3 (Hua et al., 2021). The simulated  $\Delta^{14}\text{C}$  results aimed at ~~reproducing the similar~~ finding the same  $\Delta^{14}\text{C}$  ranges obtained from the incubated  
 200 soils stablishing different ranges of decomposition rates for peatland and grassland, with the objective to compare under which settings the models could describe best laboratory or field conditions and for this reason, model fitting was not necessary. For this purpose, we implemented a two-pool model (one slow and one fast pool) considering two different structures: parallel and series (Fig. 2) and modified the parameters involved in the soil E-carbon processing (Manzoni et al., 2009; Falloon and Smith, 2000) (Table 1). In the parallel structure, E-carbon enters the soil and splits among the two pools according to  $\gamma$  and  
 205 decomposition occurs independently in each pool according to their respective  $k$ . In the series structure, E-carbon enters only to the fast pool and it is either decomposed and emitted to the atmosphere or transferred to the slow pool according to the transfer coefficient  $\alpha$ .



$$\frac{d\mathbf{C}}{dt} = I \begin{pmatrix} \gamma \\ 1-\gamma \end{pmatrix} + \begin{pmatrix} -k_1 & 0 \\ 0 & -k_2 \end{pmatrix} \cdot \begin{pmatrix} C_1 \\ C_2 \end{pmatrix}$$

$$\frac{d\mathbf{C}}{dt} = I \begin{pmatrix} 1 \\ 0 \end{pmatrix} + \begin{pmatrix} -k_1 & 0 \\ \alpha_{2,1}k_1 & -k_2 \end{pmatrix} \cdot \begin{pmatrix} C_1 \\ C_2 \end{pmatrix}$$



**Figure 2.** Parallel (left) and series (right) model structure and formulas implemented in the simulations.  $\epsilon$ -Carbon is fixed into the biomass from atmospheric  $\text{CO}_2$  through photosynthesis, subsequently it is incorporated into the soil as litterfall. From this state on, it is considered as input ( $I$ ) and will split between the two pools according to the different structures ( $\gamma$  for parallel and  $\alpha$  for series). The rate at which the  $\epsilon$ -carbon is decomposed ( $k$ ) in each pool ( $C$ ) will depend on microbial activity, environmental factors (temperature and soil moisture) and physical and chemical protection of the SOM (Blanco-Canqui and Lal, 2004; von Lützow et al., 2008; Manzoni et al., 2009). Boxes represent the soil pools, which account with an initial amount of  $\epsilon$ -carbon ( $C_1=200$ ,  $C_2=5000$ ) and arrows represent directions of inputs and outputs.

Initial parameters of the model such as the initial year of simulation and initial  $\Delta^{14}\text{C}$  values of each pool were considered separately for each type of ecosystem. Moreover, we defined  $I$  and  $C$  as constant since they can be adjusted depending on the specific soil characteristics. Additionally, we assumed that decomposition rates  $k$  reflect the effect of temperature and soil moisture on  $\epsilon$ -carbon cycling timescales (Manzoni et al., 2009), and therefore on  $\Delta^{14}\text{C}$  values. In our approach, only one parameter at time was modified for each simulation.

**Table 1.** Definition of parameters used to evaluate the variation of  $\Delta^{14}\text{C}$  values, mean age and mean transit time and their ranges ~~when type of constant was variable~~ used in the simulation experiments.

Parameter	Notation	Type	Value	Definition
Litter Input	I	Constant	100	A scalar or a data.frame object specifying the amount of litter inputs by time
Decomposition rate	$k_1, k_2$	Variable	0.00001 - 0.8	A vector of length 2 with the values of the decomposition rate for pools 1 and 2
<sup>‡</sup> Partitioning coefficient	$\gamma$	Variable	0 - 1	A scalar representing the proportion of I that goes to pool 1 in a parallel structure
Transference coefficient	$\alpha$	Variable	0 - 1	A scalar with the value of the transfer rate from pool 1 to pool 2
Carbon stocks	$C_1, C_2$	Constant	200, 5000	Initial amount of C for the two pools

215 We used the R package SoilR (Sierra et al., 2014) to simulate the temporal dynamics of  $\Delta^{14}\text{C}$  in the bulk soil and the respired  $\text{CO}_2$  as well as the age and transit time distributions of carbon ~~for two different model structures (parallel and series)~~. Assuming steady-state for the carbon stocks, the probability density function (pdf) of the age (equation 4) and the transit time (equation 5) (Metzler and Sierra, 2018) as well as their means can be calculated by the following expressions (Sierra et al., 2018b):

$$220 \quad f(a) = -\mathbf{1}^T \cdot \mathbf{A} \cdot e^{a \cdot \mathbf{A}} \cdot \frac{\mathbf{C}^*}{\sum \mathbf{C}^*}, \quad a \geq 0, \quad (4)$$

where  $a$  is the random variable age,  $\mathbf{1}^T$  is the transpose of the  $n$ -dimensional vector containing ones,  $e^{a \cdot \mathbf{A}}$  is the matrix exponential for each value of  $a$ , and  $\sum \mathbf{C}^*$  is the sum of stocks of all pools at steady state.

The pdf of transit times can be obtained as

$$f(\tau) = -\mathbf{1}^T \cdot \mathbf{A} \cdot e^{\tau \cdot \mathbf{A}} \cdot \frac{\mathbf{I}}{\sum \mathbf{I}}, \quad \tau \geq 0, \quad (5)$$

225 where  $\tau$  represents the random variable transit time.

These pdfs measure the probability that a certain amount of carbon is above or below a specific age or transit time (Sierra et al., 2018b) (equations 6 and 7, respectively). The mean of the age, that is, the expected value of the pdf can be computed by the expression:

$$\mathbb{E}(a) = -\mathbf{1}^T \cdot \mathbf{A}^{-1} \cdot \frac{\mathbf{C}^*}{\sum \mathbf{C}^*}, \quad (6)$$

230 and the mean of the transit time by


$$\mathbb{E}(\tau) = -\mathbf{1}^T \cdot \mathbf{A}^{-1} \cdot \frac{\mathbf{I}}{\sum \mathbf{I}}. \quad (7)$$

Based on the age and transit time distribution for the different simulated cases, metrics such as the mean, median and quantiles can ~~indicate the~~ be used as proxies of SOM persistence. The relation between ages and transit times may present  
235 three cases:  $Ta = Tt$   $\mathbb{E}(a) = \mathbb{E}(\tau)$  (Type I - well-mixed homogeneous system),  $Ta < Tt$   $\mathbb{E}(a) < \mathbb{E}(\tau)$  (Type II - retention system) and  $Ta > Tt$   $\mathbb{E}(a) > \mathbb{E}(\tau)$  (Type III - non-retention system) (Sierra et al., 2018b). In the type I, the probability of mineralization and release as CO<sub>2</sub> is the same for every C atom. In the type II, the C-carbon is retained for a relatively long time before it is released, ~~as in soils where C is constantly recycled and reused sequentially before respiration outside the system.~~ In type III,  
240 1973). Comparisons between these distributions will provide detailed information on mixing, store, recycling, transport and transformation processes of the SOM (Sierra et al., 2018b).


### 3 Results

#### 3.1 $\Delta^{14}\text{C}$ values of incubated soils under temperature and soil moisture variation

Values of  $\Delta^{14}\text{C}$  values from the incubation experiment contrasted strongly between grassland and peatland soils (~~Tangarife-Escobar et al., 2018~~)  
245 (Fig. 3 and 4). Generally,  $\Delta^{14}\text{C}$  of bulk soil and respiration from the peatland soil was more depleted than from the grassland soil. For example, in peatland soil the  $\Delta^{14}\text{C}$  values of bulk soil (-85.9 to -60.9 ‰, mean=-80.1, n=36, including outlier of -180) were clearly more depleted than those of the respired flux (-20.9 to ~~to~~-23.9 ‰, mean=4, n=36), which indicated that the peatland behaved as a retention system (type II). In contrast, for the grassland soil, the  $\Delta^{14}\text{C}$  of bulk soil (21.1 to 73.9 ‰, mean=43.3, n=33) ~~was not too different from the~~ fell similarly around the 1:1 line compared to the  $\Delta^{14}\text{CO}_2$  (13.9 to 83.4  
250 ‰, mean=38.5, n=33, including outliers of -227.1 and -105.1) indicating that the samples behaved mostly as a well-mixed homogeneous system (type I). Such similar values can also reflect a type II system where  $\Delta^{14}\text{C}$  values in the bulk and the respired flux are equal for the year of sampling (Fig. A7).



**Figure 3.** Boxplot with the variation of the  $\Delta^{14}\text{C}$  values in  $\text{CO}_2$  (left panel) and bulk soil (right panel) for peatland and grassland incubated soils.



**Figure 4.** Relationship between  $\Delta^{14}\text{C}$  in bulk soil and  $\Delta^{14}\text{C}$  in respired  $\text{CO}_2$  for incubated grassland (green box) and peatland (brown box) soils of the QTP discriminated by temperature and WFPS. Possible types of system according to the  $\Delta^{14}\text{C}$  relations between bulk and respired carbon.

The temperature treatments did not systematically affect the radiocarbon signature of the bulk or the respired  $\text{CO}_2$  in peatland or grassland soils (Table 2). There was strong significant evidence that manipulations in WFPS resulted in changes in the  $\Delta^{14}\text{C}$  values of bulk soil and  $\text{CO}_2$  ( $p$ -values = 0.09 in grasslands and  $p$ -value = 0.01 in peatlands from an ANOVA test, Table 2, Fig. 5 A-B) except for the bulk soil in peatlands. When the interacting effects of WFPS and temperature were evaluated together, there was no evidence that their interplay affected the radiocarbon signature of both soils ( $p$ -values > 0.05 for all the analysis). However, higher temperature and WFPS caused an increase of  $\text{CO}_2$  fluxes from respiration in the treated incubated soils (Table A1). Outliers for bulk and Outliers presented extremely depleted  $\Delta^{14}\text{C}$  values and occurred both in bulk (peatland) and  $\text{CO}_2$  in both peatlands and grasslands occurred in (grassland) at the combined treatment WFPS = 60 and temperature = 10 °C. Such wide variation in  $\Delta^{14}\text{C}$  both between ecosystems and treatments could be potentially explained by intrinsic processes affecting soil carbon dynamics, which will be explored in the following sections with a SOC decomposition model.

265 Higher temperature and WFPS caused an increase of CO<sub>2</sub> fluxes from respiration in the treated incubated soils (Table A1, Fig. 5 C-D). In both ecosystems, wetter conditions showed higher respiration rates and higher slopes as the temperature increased. The absolute amounts of CO<sub>2</sub> produced from peatland soil was in average 14 times higher than from grassland soils for every independent treatment.

**Table 2.** Summary of *p*-values obtained from ANOVA tests for the  $\Delta^{14}\text{C}$  of bulk soil and the  $\Delta^{14}\text{CO}_2$  ~~from-of~~ grassland and peatland soils ~~versus-after incubation.~~ *p*-values are given for the independent effect of temperature (T) and WFPS ~~independently~~ as well as ~~versus~~ the integrated effect of temperature and WFPS (T · WFPS).

Ecosystem	Type	T	WFPS	T · WFPS
Grassland	<u><math>\Delta^{14}\text{C}</math> Bulk</u>	0.8	0.091	0.21
	<u><math>\Delta^{14}\text{CO}_2</math></u>	0.73	0.089	0.92
Peatland	<u><math>\Delta^{14}\text{C}</math> Bulk</u>	0.4	0.21	0.65
	<u><math>\Delta^{14}\text{CO}_2</math></u>	0.16	0.01	0.32

### 3.2 Effect of changes of decomposition rates (*k*) on the $\Delta^{14}\text{C}$ values


270 ~~Temperature and soil moisture manipulations showed responses~~ Changes in the vertical and horizontal direction ~~in-of~~ the  $\Delta^{14}\text{C}$ -bulk versus  $\Delta^{14}\text{C}$ -CO<sub>2</sub> space are more evident across ecosystem type, which at the same time implies specific environmental conditions for the stability of SOM. To understand possible drivers of  $\Delta^{14}\text{C}$  changes in these directions and how they can be interpreted in terms of ages and transit times, we used a SOC decomposition model. We evaluated how model structure, decomposition rates of Carbon, and the partitioning ( $\gamma$ ) and transfer ( $\alpha$ ) coefficients of a two-pool parallel and a series model affected  $\Delta^{14}\text{C}$  values, and as a consequence, the mean age and the mean transit time. ~~Simulations were made for the Our~~ simulations mimicked a fast cycling grassland ~~system-and-the-and~~ a slow cycling peatland by differentiating the ranges of decomposition rates (Table 3) and showed how the modelled conditions affected the type of system (I, II and III).

275

#### 3.2.1 SOC decomposition in fast cycling systems (grasslands)

Variation of  $k_1$  yielded  $\Delta^{14}\text{C}$  curves similar to mean age and mean transit time. In other words, in these simulations the change in the parameter  $k_1$  resulted in similar trends in  $\Delta^{14}\text{C}$ -bulk versus  $\Delta^{14}\text{C}$ -CO<sub>2</sub> space as in the mean age versus mean transit time space ((Fig. 6). In the parallel structure (Fig. 6, A and B), high values of  $k_1$  yielded more enriched  $\Delta^{14}\text{C}$  values in the bulk soil than in the respired flux. This was expressed as a fast respiration of SOC and reflected in a short transit time. Simultaneously, low values of  $k_2$  resulted in a slow respiration, which was registered in ages longer than transit times, at the initial state of the Carbon stay in the system. However, as  $k_1$  decreased and approached  $k_2$ , mean age and mean transit time became similar and converged to the 1:1 line. In the case of the series structure (Fig. 6, C and D), the total amount of inputs traveled first through

280



**Figure 5.** Comparison between  $\Delta^{14}\text{C}$  of respiration from incubated **peatland and grassland (A)** and **peatland (B)** soil at different temperature levels under WFPS = 60 and 95 %. Black points represent minimum and maximum values out of the range between quartile 1 and 3 (25 to 75 % of the data). The quartile 50 (median) represented by the line inside the box indicates the midpoint value in the frequency distribution. Box for the treatment WFPS= 60 % and T= 10 °C shows a large dispersion of the 50 % of the data, which is explained by the outliers observed in Fig. 4. Additional panels indicate the respiration rates (mg CO<sub>2</sub> g soil<sup>-1</sup> day<sup>-1</sup>) for each treatment based on the total duration of incubation (see Fig. A1). Response of mean respiration rates to temperature and WFPS treatments in grassland (C) and peatland (D) soils.

285 the fast pool, which meant that while  $k_1$  is high, most of the **E-carbon** decomposition occurs under the dynamics of the fast pool. Hence, the system showed equal mean age and mean transit time. As  $k_1$  decreased, the transfer from the fast to the slow pool becomes more relevant, then the mean transit time increased compared to the mean age.

The behavior of  $\Delta^{14}\text{C}$  values contrasted significantly with that of mean age and mean transit time when  $k_2$  tended to lower values (Fig. 7). Generally, when  $k_2$  equals  $k_1$ ,  $\Delta^{14}\text{C}$  and mean age and mean transit time showed similar values. However, as  
 290  $k_2$  decreased, the **E-carbon** stayed in the slow pool for longer time and the  $\Delta^{14}\text{C}$  enriched in the bulk soil until reaching a

**Table 3.** Parameters—Range of parameters used for simulations in a SOC decomposition model for fast and slow systems (Tangarife-Eseobar et al., 2023). These ranges explore the required conditions to describe the  $\Delta^{14}\text{C}$  variation found in the incubated grassland and peatland soils and their equivalent age and transit time relationship. The target  $\Delta^{14}\text{C}$  intervals are shown as boxes in the Fig. 6 to 9.

System	Fig.	Model structure	$k_1$	$k_2$	$\gamma$	$\alpha$	Starting year of simulation	$\Delta^{14}\text{C}$ (‰) for starting year
Fast cycling	6	Parallel	0.1 - 0.8	0.1	0.8	-	1890	-4.9 ‰
		Series	0.1 - 0.8	0.1	-	0.8	1890	-4.9 ‰
	7	Parallel	0.8	0.001 - 0.8	0.8	-	1890	-4.9 ‰
		Series	0.8	0.001 - 0.8	-	0.8	1890	-4.9 ‰
Slow cycling	8	Parallel	0.0001 - 0.8	0.0001	0.8	-	500	-22.2 ‰
		Series	0.0001 - 0.8	0.0001	-	0.8	500	-22.2 ‰
	9	Parallel	0.1	0.000001 - 0.001	0.2	-	500	-22.2 ‰
		Series	0.1	0.000001 - 0.001	-	0.8	500	-22.2 ‰


peak (112 and 89 ‰ for series and parallel structures, respectively), from where it subsequently depleted to the initial  $\Delta^{14}\text{CO}_2$  value. For the parallel structure (Fig. 7, A and B), transit time kept increasing slowly since most of the C-carbon stayed in the fast pool for a short time, so, the C-carbon remaining in the slow pool contributed to increase the mean age. As for the series structure, most of the C-carbon was transferred from the fast to the slow pool due to the high transfer rate  $\alpha$ , which contributed to increase the transit time.

### 3.2.2 SOC decomposition in slow cycling systems (peatlands)

Simulations of  $\Delta^{14}\text{C}$  values in slow cycling systems resulted in significantly more depleted values in both bulk soil and respired  $\text{CO}_2$  (Fig. 8) than in fast cycling systems. Generally,  $\Delta^{14}\text{C}$  values were more depleted in the bulk soil than in the respiration flux. Also, the results from these simulations showed very different patterns in the  $\Delta^{14}\text{C}$ -bulk versus  $\Delta^{14}\text{C}$ - $\text{CO}_2$  space than in the mean age versus mean transit time space; only with the exception of equal  $k_1$  and  $k_2$ , which resulted in similar  $\Delta^{14}\text{C}$  values as well as similar mean age and mean transit time in the parallel structure.


Additionally, we looked at the variation of  $\Delta^{14}\text{C}$  keeping the same  $k$  values but reducing  $\alpha$  and  $\gamma$  to 0.1. Simulations indicated that the behavior of  $\Delta^{14}\text{C}$  values were opposite between series and parallel structures when inputs to each pool were inversely proportional (Fig. 7 versus Fig. A3 and Fig. 8 versus Fig. A4). For example, for the case of low values of  $k_1$  and  $k_2$





**Figure 6.** Predictions of  $\Delta^{14}\text{C}$  in bulk soil vs  $\Delta^{14}\text{CO}_2$  with their equivalent simulation of mean age in bulk soil vs mean transit time in  $\text{CO}_2$  for parallel (panel A and B) and series model structure (C and D) [for a fast cycling system](#). Variation of  $k_1$  with  $\alpha = 0.8$  and  $\gamma = 0.8$ . Complementary prediction with  $\alpha = 0.1$  and  $\gamma = 0.1$  can be seen in Fig. A2. [Green box represents the range of measured  \$\Delta^{14}\text{C}\$  values obtained from incubated grassland soils for bulk \(20 to 74 ‰\) and  \$\text{CO}\_2\$  \(10 to 85 ‰\), excluding outliers.](#)

(Fig. 9) ( $\gamma = 0.2$  and  $\alpha = 0.8$ ) both model structures showed similar patterns due to relatively similar amounts of inputs going to the slow pool. When  $\gamma = 0.8$ ,  $\Delta^{14}\text{C}$  showed values out of the target range (-20 to 23 for  $\Delta^{14}\text{C}$ - $\text{CO}_2$  and -86 to -60 for  $\Delta^{14}\text{C}$ -bulk based on incubation results) since decomposition occurred as in a fast cycling system. As  $k_2$  decreased,  $\Delta^{14}\text{CO}_2$  enriched and  $\Delta^{14}\text{C}$  and mean age remained higher than mean transit time since the slow decomposition of the slow pool dominated the system response.




**Figure 7.** Predictions of  $\Delta^{14}\text{C}$  in bulk soil vs  $\Delta^{14}\text{CO}_2$  with their equivalent simulation of mean age in bulk soil vs mean transit time in  $\text{CO}_2$  for parallel (panel A and B) and series model structure (C and D) for a fast cycling system. Variation of  $k$  with  $\alpha = 0.8$  and  $\gamma = 0.8$ . Green box represents the range of measured  $\Delta^{14}\text{C}$  values obtained from incubated grassland soils for bulk (20 to 74 ‰) and  $\text{CO}_2$  (10 to 85 ‰), excluding outliers.

### 3.3 Variation in the proportion of inputs $\gamma$ and $\alpha$ modulated mean age and mean transit time

315 Our simulations showed the importance that the partitioning and transference coefficients ( $\gamma$  and  $\alpha$ ) have to define the amount of E-carbon entering to each pool, and in consequence on the mean age and mean transit time (Fig. A1). For a parallel structure, when the total amount of inputs entered only to the fast pool ( $\gamma = 1$ ), the system behaved as a one pool system and mean age and mean transit time were equal. In the extreme opposite ( $\gamma = 0$ ), all E-carbon entered only to the slow pool and mean age and mean transit time converged, although with a longer transit through the system. In contrast, for a series structure, the extreme

320 value  $\alpha = 1$  resulted in a longer mean transit time than mean age as the contribution to the respired flux comprehended the sum of the decomposition of the newly absorbed E-carbon and the initial stocks stored in the fast pool. In the case of slow cycling systems (Fig. A5 and Fig. A6),  $\Delta^{14}\text{CO}_2$  moved in a wider range including very depleted values (-100 to 70 ‰) and




**Figure 8.** Predictions of  $\Delta^{14}\text{C}$  in bulk soil vs  $\Delta^{14}\text{CO}_2$  with their equivalent simulation of mean age in bulk soil vs mean transit time in  $\text{CO}_2$  for parallel (panel A and B) and series model structure (C and D) [for a slow cycling system](#). Variation of  $k_1$  with  $\alpha = 0.8$  and  $\gamma = 0.8$ . [Brown box represents the range of measured  \$\Delta^{14}\text{C}\$  values obtained from incubated peatland soils for bulk \(-90 to -65 ‰\) and  \$\text{CO}\_2\$  \(-18 to 25 ‰\), excluding outliers.](#)

approached to a steady state of  $\Delta^{14}\text{C}$  in bulk soil from the right when  $k_1 = 0.01$  or from the left when  $k_1 = 0.1$ .

## 325 4 Discussion

### 4.1 Did changes in temperature and soil moisture result in the release of old carbon in the respiration flux?

We measured the  $\Delta^{14}\text{C}$  values of respired  $\text{CO}_2$  from soils incubated at different temperatures and WFPS levels to assess whether changes in these variables would result in destabilization of stored [carbon](#). [In general, higher temperature and WFPS levels resulted in larger  \$\text{CO}\_2\$  fluxes although with a faster increment in peatland soils, which could be explained by the high](#)



**Figure 9.** Predictions of  $\Delta^{14}\text{C}$  in bulk soil vs  $\Delta^{14}\text{CO}_2$  with their equivalent simulation of mean age in bulk soil vs mean transit time in  $\text{CO}_2$  for parallel (panel A and B) for a slow cycling system. Series model presented exactly the same behaviour. Variation of  $k_2$  with  $\alpha = 0.8$  and  $\gamma = 0.2$ . Brown box represents the range of measured  $\Delta^{14}\text{C}$  values obtained from incubated peatland soils for bulk (-90 to -65 ‰) and  $\text{CO}_2$  (-18 to 25 ‰), excluding outliers.

330 amount of total organic carbon available for decomposition. Respiration rates, however, did not correlate with the  $\Delta^{14}\text{C}$  values. Our results showed that changes in temperature did not affect systematically the radiocarbon values of respired  $\text{CO}_2$  in any of the incubated soils. Nevertheless, changes in WFPS had a moderate significant effect on the  $\Delta^{14}\text{CO}_2$  in grasslands, but a strong influence and  $\Delta^{14}\text{C}$ -bulk of grassland soils and only on the  $\Delta^{14}\text{CO}_2$  of peatlands. Our results peatland soils. Our experiments indicated that higher WFPS levels led to depleted  $\Delta^{14}\text{CO}_2$  values in peatlands at all temperature treatments, except at 10 °C.

335 In general, higher temperature and WFPS levels resulted in increased  $\text{CO}_2$  fluxes although without showing a relationship with the  $\Delta^{14}\text{C}$  values.

In contrast, higher WFPS resulted in more enriched  $\Delta^{14}\text{CO}_2$  values for grasslands. This suggests that the direction of changing WFPS depends on ecosystem characteristics. It is interesting to observe that the  $\Delta^{14}\text{C}$  in the bulk soil was always  
340 more negative than the  $\Delta^{14}\text{CO}_2$  in the peatlands, while the grasslands registered similar  $\Delta^{14}\text{C}$  values in both states. This indicates that peatlands are systems that stabilize organic matter over time and release it once the stable conditions change. In ~~contrast~~comparison, for grasslands two interpretations may arise, first that there was not organic matter stabilization under the incubated conditions since the respired flux reassembles the bulk soil  $\Delta^{14}\text{C}$  signature, or second that the incubation conditions were not strong or long enough to destabilize the existing old SOM. Our simulations indicate that similar  $\Delta^{14}\text{C}$  values in the  
345 bulk and in the respired  $\text{CO}_2$  can also occur at an specific year due to the dynamics of the atmospheric bomb curve (Fig A7) independently of the different SOM cycling times in the soil pools.

In tundra ecosystems, Kwon et al. (2019) suggested that drainage of shallow soil layers may have accelerated old carbon decomposition. In addition, Estop-Aragones et al. (2020) concluded that old ~~C-carbon~~ would increase in proportion from “cold”  
350 across “warm wet” to “warm and dry” for high arctic tundra as well as from “undisturbed” to “burnt active layer” for peatland plateau. These previous findings are in disagreement with our observations since after our experiments, lower WFPS resulted in the release of relatively enriched  $\Delta^{14}\text{CO}_2$  instead of releasing old carbon. One potential explanation for this discrepancy is that the existing old SOM could have reacted too slow to drier conditions (WFPS = ~~6560~~) compared to the fast response of “younger” SOM, causing an evident prevalence of young carbon given the short duration of the incubations for peatlands.  
355 Consistent with this, it has been found that some SOM components or compartments may be more sensitive to modified conditions than others (Feng and Simpson, 2008) and therefore, transit time would react strongly when such sensitivity is higher in the slower pool (Manzoni et al., 2009).

For this reason, our results are indicative, but not conclusive, about the influence of environmental factors on the  $\Delta^{14}\text{C}$   
360 signature of respired carbon. It is possible that our  $\Delta^{14}\text{C}$  values are limited by the short time of the incubations since the dynamics of ~~C-carbon~~ transfers cannot be properly observed in short timescales (Crow and Sierra, 2018). For example, during a 1-year incubation experiment, most of the  $\text{CO}_2$  was derived from labile SOM as the temperature increased (Leifeld and Fuhrer, 2005). Additionally, other factors such as yearly climate seasonality, daily freeze-thaw cycles and water table oscillation were not replicated in the laboratory due to equipment and time availability. An important aspect that has been observed in soil incubations, is that  $\text{CO}_2$  accumulation decreases or even stops after a certain period probably due to the  $\text{CO}_2$  saturation of the limited head-space in the incubation flasks, which depends on the SOC content. From this we could deduce that old carbon, which usually needs longer times to be decomposed, can only be recovered if soil respiration is not space limited. For example, Azizi-Rad et al. (2022) found the respiration rate to decline after 14 days holding the soil at 10 °C. In this sense, incubations with high TOC may run out of headspace soon and affect the interpretation of the  $\Delta^{14}\text{C}$  values. Finally, although temperature  
370 has been found to be the main modulator of SOC decomposition rates (Azizi-Rad et al., 2022), dominating over the effect of soil moisture and oxygen availability, it is important to consider that an increase in respiration rates, as observed in our incubations, does not necessarily involve the release of ~~old or young carbon~~  $\Delta^{14}\text{C}$  of lower or higher signatures as shown in

[our model simulations.](#)

## 375 4.2 Are there differences in the age of respired CO<sub>2</sub> between grassland and peatland?

We found that  $\Delta^{14}\text{C}$  values from peatland were strongly more depleted than those in grasslands, indicating the presence and respiration of older carbon in peatlands. Generally, low temperature, low soil microbial activity and anoxic conditions (Xiang et al., 2009; Ma et al., 2016) posed favourable conditions for the stabilization of organic matter since the early Holocene in the Zoige region (Chen et al., 2014; Sun et al., 2017). Older bulk soil than CO<sub>2</sub> demonstrated that the peatland soil behaved  
380 as a retention system (Type II) where [C-carbon](#) was stored for relatively long time before its respiration (Sierra et al., 2018b). The release of that old [C-carbon](#) may occur when the SOM destabilization dominates over the stabilization mechanisms, for example when peatlands are drained (increasing oxidation through water table reduction) and their temperature is raised (Dutta et al., 2006; Hicks Pries et al., 2013; Lupascu et al., 2014; Estop-Aragónés et al., 2018), both factors mimicked in our incubations.

385

In contrast,  $\Delta^{14}\text{CO}_2$  from grasslands were mostly similar to  $\Delta^{14}\text{C}$  in bulk soil, indicating that the soil behaved predominantly as a well-mixed homogeneous system (Type I) where most of the SOM has the same probability of being mineralized and released as CO<sub>2</sub> (Sierra et al., 2018b). However,  $\Delta^{14}\text{C}$  relation also indicated that the grasslands shift from retention to non-retention system and vice-versa. In grasslands, SOM from the topsoil is permanently under changing conditions (daily  
390 temperature and soil moisture variation, grazing and mechanical alteration, among others) that promote its fast cycling instead of its stabilization (Han et al., 2017). These factors, along with climatic, vegetation and edaphic properties caused a low SOM stability expressed in a large proportion of labile carbon (Hou et al., 2021). Nonetheless, [although-even though](#) SOM accumulation still occurs in grassland areas (Tian et al., 2009) registered in the extremely depleted outliers from the grassland soil respiration, it happens at a slower rate than in peatlands. As a result, most of the fixed atmospheric [C-carbon](#) is respired at short  
395 time scales, which is in turn registered as young (“post bomb”)  $\Delta^{14}\text{C}$  values. [Finally, although extremely depleted  \$\Delta^{14}\text{C}\$  values are rare, they are probable since the measured carbon particles represent one value, which can fall on the tail of the system-age probability distribution. We interpret that outlier results belong to carbon particles that have remained for very long time in the system \(out of the mean values\) whose specific soil characteristics were not captured in our model structure or internal soil conditions. The reasons behind the occurrence of these outliers at the specific 10 °C remain to be investigated.](#)

400

Taken together, SOM stability and ecosystem characteristics defined the relation between  $\Delta^{14}\text{C}$  in bulk soil and  $\Delta^{14}\text{CO}_2$ , which in turn explained the difference in the age of respiration between grassland and peatlands. The  $\Delta^{14}\text{CO}_2$  values obtained in our incubated peatlands indicate that such [C-carbon](#) was captured through photosynthesis during a time when the  $^{14}\text{C}$  levels of the atmosphere were depleted and have remained stored in the soil for a long enough time for radioactive decay to become  
405 relevant. However, it is imprecise to attribute an specific year to the entire bulk soil or CO<sub>2</sub> fractions, since they are formed by

SOM accumulated in multiple steps and of different qualities and their respective decomposition.

In that sense, a  $\Delta^{14}\text{C}$  value is not per se indicative of age or transit time of SOC. To disentangle the utility of radiocarbon as a tool for SOM persistence in soil and be able to shed light on the time-scales of C-carbon cycling in different ecosystems, we used SOC decomposition models. Our models contribute to understand the time component of SOM persistence through the use of mean transit time as a parameter that integrates processes of SOM dynamics (Manzoni et al., 2009).

### 4.3 How can $\Delta^{14}\text{C}$ values be interpreted to understand the effect of decomposition rates on mean age and mean transit time?

We simulated the dynamics of  $\Delta^{14}\text{C}$  values and the as decomposition rates change, and how this behavior differ from those of mean age and mean transit time of soils with high and low decomposition rates. Our simulations were able to reassemble the  $\Delta^{14}\text{C}$  values ranges obtained from incubations and showed that modelled  $\Delta^{14}\text{C}$  values differed widely between slow cycling systems and fast cycling systems. Generally, decomposition rate, transfer rates and partitioning coefficients of a given model structure modulated  $\Delta^{14}\text{C}$  values, and in consequence, mean age and mean transit time (Bruun et al., 2004; Manzoni et al., 2009). For example, fast cycling systems with high decomposition rates resulted in a fast respiration represented by short mean transit times (lower than 20 years). At the same time, low decomposition rates resulted in longer mean transit times, which in turn represented older respired  $\text{CO}_2$  (lower than 140 years). In slow cycling systems, where the decomposition rates were lower, transit times were consequently longer (lower than 20000 years). Independently of the  $k$  decomposition rates values, the series model structure increased the transit time of C-carbon compared to parallel structure due to the fact that a proportion of the C-carbon atoms had to pass through the two pools before being respired.

Other factors such as observation time and starting year of the simulation modified the response of  $\Delta^{14}\text{C}$ . This is due to the radioactive nature of the  $^{14}\text{C}$  isotope and the influence of the “bomb  $^{14}\text{C}$ ”. For instance, the  $\Delta^{14}\text{C}$  enrichment during the last 60 years was clearly observed in fast cycling systems only with low  $k_2$ . Furthermore, the inflexion points on the  $\Delta^{14}\text{C}$  curves occurred at different values for slow and fast cycling systems. Hence, we could assume that a simulation with starting year after 1962 would not show an inflexion point in the  $\Delta^{14}\text{C}$  due to the absence of the bomb peak.

Our modelling results suggest that the increase of decomposition rates contribute to the release of older carbon in the respired flux as we hypothesised, but depending on the initial state of the system. This interpretation is not straightforward since the stability of SOM depends on the specific combination of temperature and moisture for different ecosystems. For peatlands, only drier conditions and consequent increase of oxygen might increase  $k$  decomposition rates and therefore cause the destabilization of SOM. In contrast, grasslands would need an increase of temperature to facilitate SOM decomposition provided that moisture and oxygen are available (Azizi-Rad et al., 2022)(Sierra et al., 2017a; Azizi-Rad et al., 2022). SOM decomposition rates are expected to increase at higher temperatures (Leifeld and Fuhrer, 2005) and towards the extremes of the moisture

440 range (Sierra et al., 2015). However, the destabilization of SOM can occur in any direction of WFPS variation (depending on the oxygen content) and in turn affect the age and transit time of C in a nonlinear carbon in a non-linear pattern.

From our results we could observe how the relation  $\Delta^{14}\text{C}$ -bulk versus  $\Delta^{14}\text{C}$ -CO<sub>2</sub> properly represented the relation mean age versus mean transit time. We found that there is a good correspondence between both relations in the fast cycling systems as long as the decomposition rates ( $k$ ) remain high. Such correspondence did not occur as  $k$  became smaller (typical for slow cycling systems) since the appearance of the bomb peak may have introduced anomalies that modified the equivalence between the two relations. In addition, the outlier  $\Delta^{14}\text{C}$  values found in our experimental data, where  $\Delta^{14}\text{C}$  changed drastically in the vertical and the horizontal direction, may be related to this complex response of the <sup>14</sup>C tracer.

450 To quantify cycling times of carbon, radiocarbon can be used as a tool to understand SOM destabilization and persistence through the use of the concepts of age and transit time and their mutual relation. Nonetheless, it is essential to couple  $\Delta^{14}\text{C}$  measured values with a model that involves the dynamics of soil carbon in different pools and their interaction with the environment. Therefore, the acquisition of empirical data from soils (number of pools,  $I$ ,  $C$ ,  $ks$ ,  $\gamma$ ,  $\alpha$ ) along with the correct setting of model structure will improve predictions on terrestrial and atmospheric carbon interactions.

455

## 5 Conclusions

Based on the incubation results of soils from the QTP, we showed that the  $\Delta^{14}\text{C}$  values of the peatland are significantly more depleted than the ones of the grassland both in the bulk soil and the respired CO<sub>2</sub>. Our results indicated that changes in temperature did not affect systematically the radiocarbon values of respired CO<sub>2</sub> in any of the incubated soils despite increases in respired C at higher temperatures. Nevertheless, changes in WFPS had a relatively small effect on the <sup>14</sup>CO<sub>2</sub> in grasslands, but a strong significant influence on the <sup>14</sup>CO<sub>2</sub> of peatlands, where higher WFPS levels led to more depleted <sup>14</sup>CO<sub>2</sub> values except at 10 °C. In peatlands, more depleted <sup>14</sup>C-bulk values than <sup>14</sup>C-CO<sub>2</sub> indicated that SOM stabilizes over time and it is released once the stable conditions change. In grasslands, similar  $\Delta^{14}\text{C}$  values in bulk and respired CO<sub>2</sub> indicated that the soil behaved as a well-mixed homogeneous system due to either an absence of SOM stabilization or that the manipulation treatments were not long or strong enough to destabilize the existing old SOM. In this sense, the short duration of our incubations might have been an obstacle to register the influence of long term factors such as climate seasonality and water table oscillation on SOM dynamics.

From our modelling approach, we conclude that radiocarbon can be used as a tool to understand SOM persistence through the use of the concepts of mean age and mean transit time and their mutual relation. Our simulations were able to reassemble the  $\Delta^{14}\text{C}$  values obtained from incubations and showed that modelled  $\Delta^{14}\text{C}$  values differed widely between slow cycling systems and fast cycling systems. We found that low values of kdecomposition rates, more common in slow cycling systems,



modified the behaviour of  $\Delta^{14}\text{C}$  patterns due to the incorporation of  $^{14}\text{C}$ -bomb in the soil system. Hence, the correspondence between these mutual relations strongly depended on the internal dynamics of the soil and its interaction with the environment.

475 For this reason, the acquisition of empirical data from soils (number of pools,  $I$ ,  $C$ ,  $k_s$ ,  $\gamma$  and  $\alpha$ ) along with the correct setting of model structure will improve ~~predictions on terrestrial and atmospheric carbon interactions.~~ our understanding on the stability of carbon in the soils of a changing QTP. In this way, current changes in climate patterns and land cover alteration may have a larger impact on the Zoige peatlands than on the grasslands given the vulnerability of large carbon stocks to be destabilized by changes in temperature. Nevertheless, the interaction with moisture may dampen or amplify the temperature effect, adding

480 uncertainty on the future trajectories of soil carbon in the Qinghai-Tibetan Plateau.

*Code and data availability.* Lab analysis results and code are stored at <https://doi.org/10.5281/zenodo.7620008> (DOI:10.5281/zenodo.7620008) (Tangarife-Escobar et al., 2023).

*Author contributions.* The conceptualization, data curation and formal analysis were done by ATE and CS. The investigation was performed by ATE with support from GD and MAR. Methodology and validation were done by ATE and CS. ATE conducted the validation, visualization and writing of the original draft (the latter with the help of CUM). Supervision was done by GG, XF and CS. ATE wrote the manuscript with contributions from CS, GG, XF, CUM, GD and MAR.

485

*Competing interests.* The authors declare that they have no conflict of interest.

*Acknowledgements.* We thank all colleagues who contributed to this study, especially Eike Reinosch for providing the location maps used in Fig.1. and Paula Sierra for the digitization of the scheme used in Fig. 2. Special thanks to David Martini for his helpful advices to improve code writing and to Manuel Röst and Axel Steinhof for their useful training at the  $^{14}\text{C}$  laboratory. Finally, thanks to Nicole Börner for the active collaboration with the project administration and to Susan Trumbore for her feedback on the final version of this manuscript. This study was developed as part of the International Research Training Group (GRK 2309/1) Geo-ecosystems in transition on the Tibetan Plateau (TransTiP) funded by the Deutsche Forschungsgemeinschaft (DFG). The Max Planck Institute for Biogeochemistry provided permanent administrative support.

490


**Table A1.** Mean daily CO<sub>2</sub> respiration for incubated soils under temperature and WFPS variation.

Ecosystem	Temperature (°C)	WFPS (%)	Mean CO <sub>2</sub> respiration (mg CO <sub>2</sub> g soil <sup>-1</sup> day <sup>-1</sup> )	$\sigma$	<u>Incubation time (days)</u> <u>n = number of samples</u>
Grassland	20	95	<del>0.510</del> <u>0.040</u>	0.035	<u>2n = 32, 1n = 19</u>
	20	60	<del>0.371</del> <u>0.028</u>	0.016	<u>4n = 19</u>
	15	95	<del>0.330</del> <u>0.024</u>	0.008	<u>3n = 30, 1n = 15</u>
	15	60	<del>0.234</del> <u>0.016</u>	0.004	<u>4n = 30</u>
	10	95	<del>0.209</del> <u>0.011</u>	0.011	<u>3n = 66</u>
	10	60	<del>0.134</del> <u>0.008</u>	0.023	<u>6n = 66</u>
	5	95	<del>0.066</del> <u>0.005</u>	0.003	<u>5n = 67</u>
	5	60	<del>0.044</del> <u>0.004</u>	0.001	<u>4n = 67</u>
Peatland	20	95	<del>0.040</del> <u>0.510</u>	0.001	<u>5n = 13</u>
	20	60	<del>0.028</del> <u>0.371</u>	0.003	<u>4n = 13</u>
	15	95	<del>0.024</del> <u>0.330</u>	0.004	<u>5n = 13</u>
	15	60	<del>0.016</del> <u>0.234</u>	0.003	<u>3n = 13</u>
	10	95	<del>0.011</del> <u>0.209</u>	0.002	<u>5n = 13</u>
	10	60	<del>0.008</del> <u>0.134</u>	0.001	<u>5n = 13</u>
	5	95	<del>0.005</del> <u>0.066</u>	0.000	<u>4n = 13</u>
	5	60	<del>0.004</del> <u>0.044</u>	0.001	<u>5n = 13</u>


**Table A2.** ~~Parameters~~ Range of parameters used for simulations in a SOC decomposition ~~models~~ model for fast and slow systems (Tangarife-Escobar et al., 2023). These ranges explore the required conditions to describe the  $\Delta^{14}\text{C}$  variation found in the incubated grassland (fast cycling) and peatland (slow cycling) soils and their equivalent age and transit time relationship. The target  $\Delta^{14}\text{C}$  intervals are shown as boxes in the Fig. A1 to A6.

System	Fig.	Model structure	$k_1$	$k_2$	$\gamma$	$\alpha$	Starting year of simulation	$\Delta^{14}\text{C}$ (‰) for starting year
Fast cycling	A1	Parallel	0.8	0.1	0 - 1	-	1890	-4.9 ‰
		Series	0.8	0.1	-	0 - 1	1890	-4.9 ‰
	A2	Parallel	0.1 - 0.8	0.1	0.1	-	1890	-4.9 ‰
		Series	0.1 - 0.8	0.1	-	0.1	1890	-4.9 ‰
	A3	Parallel	0.8	0.0001 - 0.8	0.1	-	1890	-4.9 ‰
		Series	0.8	0.0001 - 0.8	-	0.1	1890	-4.9 ‰
Slow cycling	A4	Parallel	0.0001 - 0.8	0.0001	0.2	-	500	-22.2 ‰
		Series	0.0001 - 0.8	0.0001	-	0.1	500	-22.2 ‰
	A5	Parallel	0.01	0.0001	0 - 1	-	500	-22.2 ‰
		Series	0.01	0.0001	-	0 - 1	500	-22.2 ‰
	A6	Parallel	0.1	0.0001	0 - 1	-	500	-22.2 ‰
		Series	0.1	0.0001	-	0 - 1	500	-22.2 ‰


-  $\Delta^{14}\text{C}$  variation in fast cycling systems



**Figure A1.** Predictions of  $\Delta^{14}\text{C}$  in bulk soil vs  $\Delta^{14}\text{CO}_2$  with their equivalent simulation of mean age in bulk soil vs mean transit time in  $\text{CO}_2$  for parallel (panel A and B) and series model structure (C and D) [for a fast cycling system](#).  $k_1 = 0.8$ ,  $k_2 = 0.1$ , with  $\alpha = 0 - 1$  and  $\gamma = 0 - 1$ . Green box represents the range of measured  $\Delta^{14}\text{C}$  values obtained from incubated grassland soils for bulk (20 to 74 ‰) and  $\text{CO}_2$  (10 to 85 ‰), excluding outliers.




**Figure A2.** Predictions of  $\Delta^{14}\text{C}$  in bulk soil vs  $\Delta^{14}\text{CO}_2$  with their equivalent simulation of mean age in bulk soil vs mean transit time in  $\text{CO}_2$  for parallel (panel A and B) and series model structure (C and D) [for a fast cycling system](#). Variation of  $k_1$  with  $\alpha = 0.1$  and  $\gamma = 0.1$ . [Green box represents the range of measured  \$\Delta^{14}\text{C}\$  values obtained from incubated grassland soils for bulk \(20 to 74 ‰\) and  \$\text{CO}\_2\$  \(10 to 85 ‰\), excluding outliers.](#)




**Figure A3.** Predictions of  $\Delta^{14}\text{C}$  in bulk soil vs  $\Delta^{14}\text{CO}_2$  with their equivalent simulation of mean age in bulk soil vs mean transit time in  $\text{CO}_2$  for parallel (panel A and B) and series model structure (C and D) [for a fast cycling system](#). Variation of  $k_2$  with  $\alpha = 0.1$  and  $\gamma = 0.1$ . [Green box represents the range of measured  \$\Delta^{14}\text{C}\$  values obtained from incubated grassland soils for bulk \(20 to 74 ‰\) and  \$\text{CO}\_2\$  \(10 to 85 ‰\), excluding outliers.](#)

-  $\Delta^{14}\text{C}$  variation in slow cycling systems




**Figure A4.** Predictions of  $\Delta^{14}\text{C}$  in bulk soil vs  $\Delta^{14}\text{CO}_2$  with their equivalent simulation of mean age in bulk soil vs mean transit time in  $\text{CO}_2$  for parallel (panel A and B) and series model structure (C and D) for a slow cycling system. Variation of  $k_1$  with  $\alpha = 0.1$  and  $\gamma = 0.2$ . Brown box represents the range of measured  $\Delta^{14}\text{C}$  values obtained from incubated peatland soils for bulk (-90 to -65 ‰) and  $\text{CO}_2$  (-18 to 25 ‰), excluding outliers.




**Figure A5.** Predictions of  $\Delta^{14}\text{C}$  in bulk soil vs  $\Delta^{14}\text{CO}_2$  with their equivalent simulation of mean age in bulk soil vs mean transit time in  $\text{CO}_2$  for parallel (panel A and B) and series model structure (C and D) for a slow cycling system.  $k_1 = 0.01$ ,  $k_2 = 0.0001$ , with  $\alpha = 0 - 1$  and  $\gamma = 0 - 1$ . Brown box represents the range of measured  $\Delta^{14}\text{C}$  values obtained from incubated peatland soils for bulk (-90 to -65 ‰) and  $\text{CO}_2$  (-18 to 25 ‰), excluding outliers.





**Figure A6.** Predictions of  $\Delta^{14}\text{C}$  in bulk soil vs  $\Delta^{14}\text{CO}_2$  with their equivalent simulation of mean age in bulk soil vs mean transit time in  $\text{CO}_2$  for parallel (panel A and B) and series model structure (C and D) [for a slow cycling system](#).  $k_1 = 0.1$ ,  $k_2 = 0.0001$ , with  $\alpha = 0 - 1$  and  $\gamma = 0 - 1$ . [Brown box represents the range of measured  \$\Delta^{14}\text{C}\$  values obtained from incubated peatland soils for bulk \(-90 to -65 ‰\) and  \$\text{CO}\_2\$  \(-18 to 25 ‰\), excluding outliers.](#)

-  $\Delta^{14}\text{C}$  curves for bulk soil, respired  $\text{CO}_2$  and atmosphere for a two-pool system in parallel and series structure



**Figure A7.** Variation of  $\Delta^{14}\text{C}$  values in atmosphere, bulk soil and respired  $\text{CO}_2$  for a two-pool soil during the period 1950 - 2019. The similarity between  $\Delta^{14}\text{C}$  values may indicate both a well-mixed homogeneous system (type I) or a retention system (type II) where the bulk soil and the respired flux record the same value. Nonetheless, the mean transit time for the bulk and the soil respiration might be different due to the contrasting decomposition rates of the fast ( $k_1 = 0.8$ ) and the slow pool ( $k_2 = 0.1$ ). Simulation conducted with  $\alpha = 0.8$  and  $\gamma = 0.8$ .

## References

- 500 Anslan, S., Rad, M. A., Buckel, J., Galindo, P. E., Kai, J., Kang, W., Keys, L., Maurischat, P., Nieberding, F., Reinosch, E., Tang, H., Tran, T. V., Wang, Y., and Schwalb, A.: Reviews and syntheses: How do abiotic and biotic processes respond to climatic variations in the Nam Co catchment (Tibetan Plateau)?, *Biogeosciences*, 17, 1261–1279, <https://doi.org/10.5194/bg-17-1261-2020>, 2020.
- Arias, P., Bellouin, N., Coppola, E., Jones, R., Krinner, G., Marotzke, J., Naik, V., Palmer, M., Plattner, G.-K., Rogelj, J., et al.: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change; Technical Summary*, 2021.
- 505 Azizi-Rad, M., Guggenberger, G., Ma, Y., and Sierra, C. A.: Sensitivity of soil respiration rate with respect to temperature, moisture and oxygen under freezing and thawing, *Soil Biology and Biochemistry*, 165, 108488, <https://doi.org/https://doi.org/10.1016/j.soilbio.2021.108488>, 2022.
- Blanco-Canqui, H. and Lal, R.: Mechanisms of carbon sequestration in soil aggregates, *Critical reviews in plant sciences*, 23, 481–504, 2004.
- 510 Bolin, B. and Rodhe, H.: A note on the concepts of age distribution and transit time in natural reservoirs, *Tellus*, 25, 58–62, <https://doi.org/https://doi.org/10.1111/j.2153-3490.1973.tb01594.x>, 1973.
- Bradford, M. A., Wieder, W. R., Bonan, G. B., Fierer, N., Raymond, P. A., and Crowther, T. W.: Managing uncertainty in soil carbon feedbacks to climate change, *Nature Climate Change*, 6, 751–758, 2016.
- Briones, M. J., Garnett, M. H., and Ineson, P.: No evidence for increased loss of old carbon in a temperate organic soil after 13 years of  
515 simulated climatic warming despite increased CO<sub>2</sub> emissions, *Global Change Biology*, 2021.
- Briones, M. J. I., Garnett, M. H., and Ineson, P.: Soil biology and warming play a key role in the release of ‘old C’ from organic soils, *Soil Biology and Biochemistry*, 42, 960–967, <https://doi.org/10.1016/j.soilbio.2010.02.013>, 2010.
- Bruun, S., Six, J., and Jensen, L. S.: Estimating vital statistics and age distributions of measurable soil organic carbon fractions based on their pathway of formation and radiocarbon content, *Journal of theoretical biology*, 230, 241–250, 2004.
- 520 Chen, H., Yang, G., Peng, C., Zhang, Y., Zhu, D., Zhu, Q., Hu, J., Wang, M., Zhan, W., Zhu, E., Bai, Z., Li, W., Wu, N., Wang, Y., Gao, Y., Tian, J., Kang, X., Zhao, X., and Wu, J.: The carbon stock of alpine peatlands on the Qinghai-Tibetan Plateau during the Holocene and their future fate, *Quaternary Science Reviews*, 95, 151–158, <https://doi.org/10.1016/j.quascirev.2014.05.003>, 2014.
- Chen, L., Jing, X., Flynn, D. F., Shi, Y., Kühn, P., Scholten, T., and He, J.-S.: Changes of carbon stocks in alpine grassland soils from 2002 to 2011 on the Tibetan Plateau and their climatic causes, *Geoderma*, 288, 166–174, 2017.
- 525 Chen, L., Fang, K., Wei, B., Qin, S., Feng, X., Hu, T., Ji, C., and Yang, Y.: Soil carbon persistence governed by plant input and mineral protection at regional and global scales, *Ecology Letters*, 24, 1018–1028, 2021.
- Crow, S. E. and Sierra, C. A.: Dynamic, intermediate soil carbon pools may drive future responsiveness to environmental change, *Journal of environmental quality*, 47, 607–616, 2018.
- Davidson, E. and Janssens, I.: Temperature sensitivity of soil carbon decomposition and feedbacks to climate change, *Nature*, 440, 165–173, <https://doi.org/10.1038/nature04514>, 2006.
- 530 Dioumaeva, I., Trumbore, S., Schuur, E., Goulden, M., Litvak, M., and Hirsch, A.: Decomposition of peat from upland boreal forest: Temperature dependence and sources of respired carbon, *Journal of Geophysical Research: Atmospheres*, 108, <https://doi.org/10.1029/2001JD000848>, 2002.
- Dong, S., Peng, F., You, Q., Guo, J., and Xue, X.: Lake dynamics and its relationship to climate change on the Tibetan Plateau over the last  
535 four decades, *Regional environmental change*, 18, 477–487, 2018.

- Du, C. and Gao, Y.: Opposite patterns of soil organic and inorganic carbon along a climate gradient in the alpine steppe of northern Tibetan Plateau, *Catena*, 186, 104366, 2020.
- Dutta, K., Schuur, E. A. G., Neff, J. C., and Zimov, S. A.: Potential carbon release from permafrost soils of Northeastern Siberia, *Global Change Biology*, 12, 2336–2351, <https://doi.org/10.1111/j.1365-2486.2006.01259.x>, 2006.
- 540 Emanuel, W., Killough, G., Post, W., and Shugart, H.: Modeling terrestrial ecosystems in the global carbon-cycle with shifts in carbon storage capacity by land-use change, *Ecology*, 65, 970–983, <https://doi.org/10.2307/1938069>, 1984.
- Eriksson, E.: Compartment Models and Reservoir Theory, *Annual Review of Ecology and Systematics*, 2, 67–84, <https://doi.org/10.1146/annurev.es.02.110171.000435>, 1971.
- Estop-Aragones, C., Czimczik, C. I., Heffernan, L., Gibson, C., Walker, J. C., Xu, X., and Olefeldt, D.: Respiration of aged soil carbon during  
545 fall in permafrost peatlands enhanced by active layer deepening following wildfire but limited following thermokarst, *Environmental Research Letters*, 13, 085002, 2018.
- Estop-Aragones, C., Olefeldt, D., Abbott, B. W., Chanton, J. P., Czimczik, C. I., Dean, J. F., Egan, J. E., Gandois, L., Garnett, M. H., Hartley, I. P., Hoyt, A., Lupascu, M., Natali, S. M., O'Donnell, J. A., Raymond, P. A., Tanentzap, A. J., Tank, S. E., Schuur, E. A. G., Turetsky, M., and Anthony, K. W.: Assessing the Potential for Mobilization of Old Soil Carbon After Permafrost Thaw: A Synthesis of  
550 C-14 Measurements From the Northern Permafrost Region, *Global Biogeochemical Cycles*, 34, <https://doi.org/10.1029/2020GB006672>, 2020.
- Falloon, P. and Smith, P.: Modelling refractory soil organic matter, *Biology and fertility of soils*, 30, 388–398, 2000.
- Feng, X. and Simpson, M. J.: Temperature responses of individual soil organic matter components, *Journal of Geophysical Research: Biogeosciences*, 113, 2008.
- 555 Ganjurjav, H., Gao, Q., Gornish, E. S., Schwartz, M. W., Liang, Y., Cao, X., Zhang, W., Zhang, Y., Li, W., Wan, Y., Li, Y., Danjiu, L., Guo, H., and Lin, E.: Differential response of alpine steppe and alpine meadow to climate warming in the central Qinghai-Tibetan Plateau, *Agricultural and Forest Meteorology*, 223, 233–240, <https://doi.org/10.1016/j.agrformet.2016.03.017>, 2016.
- Gao, Q.-z., Li, Y., Xu, H.-m., Wan, Y.-f., and Jiangcun, W.-z.: Adaptation strategies of climate variability impacts on alpine grassland ecosystems in Tibetan Plateau, *Mitigation and Adaptation Strategies for Global Change*, 19, 199–209, 2014.
- 560 Gaudinski, J., Trumbore, S., Davidson, E., and Zheng, S.: Soil carbon cycling in a temperate forest: radiocarbon-based estimates of residence times, sequestration rates and partitioning of fluxes, *Biogeochemistry*, 51, 33–69, <https://doi.org/10.1023/A:1006301010014>, 2000.
- Geng, Y., Wang, Y., Yang, K., Wang, S., Zeng, H., Baumann, F., Kuehn, P., Scholten, T., and He, J.-S.: Soil respiration in Tibetan alpine grasslands: belowground biomass and soil moisture, but not soil temperature, best explain the large-scale patterns, *PLoS one*, 7, e34968, 2012.
- 565 Genxu, W., Ju, Q., Guodong, C., and Yuanmin, L.: Soil organic carbon pool of grassland soils on the Qinghai-Tibetan Plateau and its global implication, *Science of the Total Environment*, 291, 207–217, 2002.
- Han, C., Wang, Z., Si, G., Lei, T., Yuan, Y., and Zhang, G.: Increased precipitation accelerates soil organic matter turnover associated with microbial community composition in topsoil of alpine grassland on the eastern Tibetan Plateau, *Canadian journal of microbiology*, 63, 811–821, 2017.
- 570 Hao, Y. B., Cui, X. Y., Wang, Y. F., Mei, X. R., Kang, X. M., Wu, N., Luo, P., and Zhu, D.: Predominance of Precipitation and Temperature Controls on Ecosystem CO<sub>2</sub> Exchange in Zoige Alpine Wetlands of Southwest China, *Wetlands*, 31, 413–422, <https://doi.org/10.1007/s13157-011-0151-1>, 2011.

- Hicks Pries, C. E., Schuur, E. A., and Crummer, K. G.: Thawing permafrost increases old soil and autotrophic respiration in tundra: Partitioning ecosystem respiration using D13C an D14C, *Global Change Biology*, 19, 649–661, 2013.
- 575 Hopkins, F. M., Torn, M. S., and Trumbore, S. E.: Warming accelerates decomposition of decades-old carbon in forest soils, *Proceedings of the National Academy of Sciences of the United States of America*, 109, E1753–E1761, <https://doi.org/10.1073/pnas.1120603109>, 2012.
- Hou, Y., He, K., Chen, Y., Zhao, J., Hu, H., and Zhu, B.: Changes of soil organic matter stability along altitudinal gradients in Tibetan alpine grassland, *Plant and Soil*, 458, 21–40, 2021.
- Hua, Q., Turnbull, J. C., Santos, G. M., Rakowski, A. Z., Ancapichún, S., De Pol-Holz, R., Hammer, S., Lehman, S. J., Levin, I., Miller, 580 J. B., et al.: Atmospheric radiocarbon for the period 1950–2019, *Radiocarbon*, pp. 1–23, 2021.
- Jarvis, A., Reuter, H. I., Nelson, A., and Guevara, E.: Hole-filled SRTM for the globe Version 4. available from the CGIAR-CSI SRTM 90m Database, 15, (25–54), <http://srtm.csi.cgiar.org>, 2008.
- Kang, X., Yan, L., Cui, L., Zhang, X., Hao, Y., Wu, H., Zhang, Y., Li, W., Zhang, K., Yan, Z., Li, Y., and Wang, J.: Reduced Carbon Dioxide Sink and Methane Source under Extreme Drought Condition in an Alpine Peatland, *Sustainability*, 10, 585 <https://doi.org/10.3390/su10114285>, 2018.
- Knorr, W., Prentice, I., House, J., and Holland, E.: Long-term sensitivity of soil carbon turnover to warming, *Nature*, 433, 298–301, <https://doi.org/10.1038/nature03226>, 2005.
- Kwon, M. J., Natali, S. M., Pries, C. E. H., Schuur, E. A. G., Steinhof, A., Crummer, K. G., Zimov, N., Zimov, S. A., Heimann, M., Kolle, O., and Goeckede, M.: Drainage enhances modern soil carbon contribution but reduces old soil carbon contribution to ecosystem respiration 590 in tundra ecosystems, *Global Change Biology*, 25, 1315–1325, <https://doi.org/10.1111/gcb.14578>, 2019.
- Leifeld, J. and Fuhrer, J.: The temperature response of CO<sub>2</sub> production from bulk soils and soil fractions is related to soil organic matter quality, *Biogeochemistry*, 75, 433–453, 2005.
- Li, J., Yan, D., Pendall, E., Pei, J., Noh, N. J., He, J.-S., Li, B., Nie, M., and Fang, C.: Depth dependence of soil carbon temperature sensitivity across Tibetan permafrost regions, *Soil Biology and Biochemistry*, 126, 82–90, 2018.
- 595 Liski, J., Ilvesniemi, H., Makela, A., and Westman, C.: CO<sub>2</sub> emissions from soil in response to climatic warming are overestimated - The decomposition of old soil organic matter is tolerant of temperature, *Ambio*, 28, 171–174, 1999.
- Liu, L., Chen, H., Zhu, Q., Yang, G., Zhu, E., Hu, J., Peng, C., Jiang, L., Zhan, W., Ma, T., He, Y., and Zhu, D.: Responses of peat carbon at different depths to simulated warming and oxidizing, *Science of the Total Environment*, 548, 429–440, <https://doi.org/10.1016/j.scitotenv.2015.11.149>, 2016.
- 600 Liu, L., Chen, H., Liu, X., Yang, Z., Zhu, D., He, Y., and Liu, J.: Contemporary, modern and ancient carbon fluxes in the Zoige peatlands on the Qinghai-Tibetan Plateau, *Geoderma*, 352, 138–149, <https://doi.org/10.1016/j.geoderma.2019.06.008>, 2019a.
- Liu, X., Zhu, D., Zhan, W., Chen, H., Zhu, Q., Zhang, J., Wu, N., and He, Y.: Dominant influence of non-thawing periods on annual CO<sub>2</sub> emissions from Zoige peatlands: Five-year eddy covariance analysis, *Ecological Indicators*, 129, 107 913, 2021.
- Liu, X., Chen, H., Zhu, Q., Wu, J., Froelking, S., Zhu, D., Wang, M., Wu, N., Peng, C., and He, Y.: Holocene peatland development and carbon stock of Zoige peatlands, Tibetan Plateau: a modeling approach, *Journal of Soils and Sediments*, 18, 2032–2043, 605 <https://doi.org/10.1007/s11368-018-1960-0>, 2018.
- Liu, X., Zhu, D., Zhan, W., Chen, H., Zhu, Q., Hao, Y., Liu, W., and He, Y.: Five-Year Measurements of Net Ecosystem CO<sub>2</sub> Exchange at a Fen in the Zoige Peatlands on the Qinghai-Tibetan Plateau, *Journal of Geophysical Research: Atmospheres*, 124, 11 803–11 818, <https://doi.org/10.1029/2019JD031429>, 2019b.

- 610 Lu, M., Zhou, X., Yang, Q., Li, H., Luo, Y., Fang, C., Chen, J., Yang, X., and Li, B.: Responses of ecosystem carbon cycle to experimental warming: a meta-analysis, *Ecology*, 94, 726–738, 2013.
- Lupascu, M., Welker, J., Xu, X., and Czimczik, C.: Rates and radiocarbon content of summer ecosystem respiration in response to long-term deeper snow in the High Arctic of NW Greenland, *Journal of Geophysical Research: Biogeosciences*, 119, 1180–1194, 2014.
- Ma, K., Zhang, Y., Tang, S., and Liu, J.: Spatial distribution of soil organic carbon in the Zoige alpine wetland, northeastern Qinghai-Tibet Plateau, *Catena*, 144, 102–108, <https://doi.org/10.1016/j.catena.2016.05.014>, 2016.
- 615 Manzoni, S., Katul, G. G., and Porporato, A.: Analysis of soil carbon transit times and age distributions using network theories, *Journal of Geophysical Research: Biogeosciences*, 114, 2009.
- McGuire, A., Sitch, S., Clein, J. S., Dargaville, R., Esser, G., Foley, J., Heimann, M., Joos, F., Kaplan, J., Kicklighter, D., et al.: Carbon balance of the terrestrial biosphere in the twentieth century: Analyses of CO<sub>2</sub>, climate and land use effects with four process-based ecosystem models, *Global biogeochemical cycles*, 15, 183–206, 2001.
- 620 Mesfin, S., Gebresamuel, G., Haile, M., and Zenebe, A.: Modelling spatial and temporal soil organic carbon dynamics under climate and land management change scenarios, northern Ethiopia, *European Journal of Soil Science*, 72, 1298–1311, 2021.
- Metzler, H. and Sierra, C. A.: Linear Autonomous Compartmental Models as Continuous-Time Markov Chains: Transit-Time and Age Distributions, *Mathematical Geosciences*, 50, 1–34, <https://doi.org/10.1007/s11004-017-9690-1>, 2018.
- 625 Moyano, F. E., Manzoni, S., and Chenu, C.: Responses of soil heterotrophic respiration to moisture availability: An exploration of processes and models, *Soil Biology and Biochemistry*, 59, 72–85, 2013.
- Nieberding, F., Wille, C., Fratini, G., Asmussen, M. O., Wang, Y., Ma, Y., and Sachs, T.: A long-term (2005–2019) eddy covariance data set of CO<sub>2</sub> and H<sub>2</sub>O fluxes from the Tibetan alpine steppe, *Earth System Science Data*, 12, 2505–2524, <https://doi.org/10.5194/essd-12-2705-2020>, 2020.
- 630 Pan, Y., Li, X., Li, S., and Li, Z.: Different responses of soil respiration to climate change in permafrost and non-permafrost regions of the Tibetan plateau from 1979 to 2018, *International Journal of Climatology*, 2022.
- Pegoraro, E. F., Mauritz, M. E., Ogle, K., Ebert, C. H., and Schuur, E. A.: Lower soil moisture and deep soil temperatures in thermokarst features increase old soil carbon loss after 10 years of experimental permafrost warming, *Global change biology*, 27, 1293–1308, 2021.
- Piao, S., Tan, K., Nan, H., Ciais, P., Fang, J., Wang, T., Vuichard, N., and Zhu, B.: Impacts of climate and CO<sub>2</sub> changes on the vegetation growth and carbon balance of Qinghai–Tibetan grasslands over the past five decades, *Global and Planetary Change*, 98, 73–80, 2012.
- 635 Pold, G., Melillo, J. M., and DeAngelis, K. M.: Two decades of warming increases diversity of a potentially lignolytic bacterial community, *Frontiers in microbiology*, 6, 480, 2015.
- Rustad, L., Campbell, J., Marion, G., Norby, R., Mitchell, M., Hartley, A., Cornelissen, J., and Gurevitch, J.: A meta-analysis of the response of soil respiration, net nitrogen mineralization, and aboveground plant growth to experimental ecosystem warming, *Oecologia*, 126, 543–562, 2001.
- 640 Schimel, D.: Terrestrial ecosystems and the carbon-cycle, *Global Change Biology*, 1, 77–91, <https://doi.org/10.1111/j.1365-2486.1995.tb00008.x>, 1995.
- Schmidt, M. W., Torn, M. S., Abiven, S., Dittmar, T., Guggenberger, G., Janssens, I. A., Kleber, M., Kögel-Knabner, I., Lehmann, J., Manning, D. A., et al.: Persistence of soil organic matter as an ecosystem property, *Nature*, 478, 49–56, 2011.
- 645 Schuur, E. A. G., Vogel, J. G., Crummer, K. G., Lee, H., Sickman, J. O., and Osterkamp, T. E.: The effect of permafrost thaw on old carbon release and net carbon exchange from tundra, *Nature*, 459, 556–559, <https://doi.org/10.1038/nature08031>, 2009.

- Schuur, E. A. G., Druffel, E., and Trumbore, S. E.: Radiocarbon and Climate Change: Mechanisms, Applications and Laboratory Techniques, pp. 1–315, Springer International, <https://doi.org/10.1007/978-3-319-25643-6>, 2016.
- Scurlock, J. and Hall, D.: The global carbon sink: a grassland perspective, *Global Change Biology*, 4, 229–233, 1998.
- 650 Shaw, R. G. and Mitchell-Olds, T.: ANOVA for unbalanced data: an overview, *Ecology*, 74, 1638–1645, 1993.
- Sierra, C. A. and Mueller, M.: A general mathematical framework for representing soil organic matter dynamics, *Ecological Monographs*, 85, 505–524, <https://doi.org/10.1890/15-0361.1>, 2015.
- Sierra, C. A., Trumbore, S. E., Davidson, E. A., Vicca, S., and Janssens, I.: Sensitivity of decomposition rates of soil organic matter with respect to simultaneous changes in temperature and moisture, *Journal of Advances in Modeling Earth Systems*, 7, 335–356, 2015.
- 655 Sierra, C. A., Malghani, S., and Loescher, H. W.: Interactions among temperature, moisture, and oxygen concentrations in controlling decomposition rates in a boreal forest soil, *Biogeosciences*, 14, 703–710, 2017a.
- Sierra, C. A., Müller, M., Metzler, H., Manzoni, S., and Trumbore, S. E.: The muddle of ages, turnover, transit, and residence times in the carbon cycle, *Global Change Biology*, 23, 1763–1773, 2017b.
- Sierra, C. A., Mueller, M., and Trumbore, S. E.: Models of soil organic matter decomposition: the SoilR package, version 1.0, *Geoscientific Model Development*, 5, 1045–1060, <https://doi.org/10.5194/gmd-5-1045-2012>, 2012.
- 660 Sierra, C. A., Mueller, M., and Trumbore, S. E.: Modeling radiocarbon dynamics in soils: SOILR version 1.1, *Geoscientific Model Development*, 7, 1919–1931, <https://doi.org/10.5194/gmd-7-1919-2014>, 2014.
- Sierra, C. A., Ceballos-Nunez, V., Metzler, H., and Mueller, M.: Representing and Understanding the Carbon Cycle Using the Theory of Compartmental Dynamical Systems, *Journal of Advances in Modeling Earth Systems*, 10, 1729–1734, <https://doi.org/10.1029/2018MS001360>,
- 665 2018a.
- Sierra, C. A., Hoyt, A. M., He, Y., and Trumbore, S. E.: Soil Organic Matter Persistence as a Stochastic Process: Age and Transit Time Distributions of Carbon in Soils, *Global Biogeochemical Cycles*, 32, 1574–1588, <https://doi.org/10.1029/2018GB005950>, 2018b.
- Steinhof, A., Altenburg, M., and Machts, H.: Sample preparation at the Jena C-14 laboratory, *Radiocarbon*, 59, 815–830, <https://doi.org/10.1017/RDC.2017.50>, 2017.
- 670 Stuiver, M. and Polach, H.: Reporting of C-14 data - Discussion, *Radiocarbon*, 19, 355–363, <https://doi.org/10.1017/S0033822200003672>, 1977.
- Sun, X., Zhao, Y., and Li, Q.: Holocene peatland development and vegetation changes in the Zoige Basin, eastern Tibetan Plateau, *Science China Earth Sciences*, 60, 1826–1837, 2017.
- Tangarife-Escobar, A., Guggenberger, G., Feng, X., Dai, G., Urbina-Malo, C., Azizi-Rad, M., and Sierra, C.: Moisture and temperature effects on the radiocarbon signature of respired carbon dioxide to assess stability of soil carbon in the Tibetan Plateau, <https://doi.org/10.5281/zenodo.7620008>, 2023.
- 675 Tian, Y.-Q., Xu, X.-L., Song, M.-H., Zhou, C.-P., Gao, Q., and Ouyang, H.: Carbon sequestration in two alpine soils on the Tibetan Plateau, *Journal of integrative plant biology*, 51, 900–905, 2009.
- Trumbore, S.: Age of soil organic matter and soil respiration: Radiocarbon constraints on belowground C dynamics, *Ecological Applications*, 10, 399–411, [https://doi.org/10.1890/1051-0761\(2000\)010\[0399:AOSOMA\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2000)010[0399:AOSOMA]2.0.CO;2), 2000.
- 680 Trumbore, S., Chadwick, O., and Amundson, R.: Rapid exchange between soil carbon and atmospheric carbon dioxide driven by temperature change, *Science*, 272, 393–396, <https://doi.org/10.1126/science.272.5260.393>, 1996.

- von Lützow, M., Kögel-Knabner, I., Ludwig, B., Matzner, E., Flessa, H., Ekschmitt, K., Guggenberger, G., Marschner, B., and Kalbitz, K.: Stabilization mechanisms of organic matter in four temperate soils: Development and application of a conceptual model, *Journal of Plant Nutrition and Soil Science*, 171, 111–124, 2008.
- 685 Wang, M., Yang, G., Gao, Y., Chen, H., Wu, N., Peng, C., Zhu, Q., Zhu, D., Wu, J., He, Y., Tian, J., Zhao, X., and Zhang, Y.: Higher recent peat C accumulation than that during the Holocene on the Zoige Plateau, *Quaternary Science Reviews*, 114, 116–125, <https://doi.org/10.1016/j.quascirev.2015.01.025>, 2015.
- Wieder, W. R., Bonan, G. B., and Allison, S. D.: Global soil carbon projections are improved by modelling microbial processes, *Nature* 690 *climate change*, 3, 909–912, 2013.
- Xiang, S., Guo, R., Wu, N., and Sun, S.: Current status and future prospects of Zoige Marsh in Eastern Qinghai-Tibet Plateau, *Ecological Engineering*, 35, 553–562, <https://doi.org/10.1016/j.ecoleng.2008.02.016>, 2009.
- Xu, X., Shi, Z., Chen, X., Lin, Y., Niu, S., Jiang, L., Luo, R., and Luo, Y.: Unchanged carbon balance driven by equivalent responses of production and respiration to climate change in a mixed-grass prairie, *Global Change Biology*, 22, 1857–1866, 2016.
- 695 Yang, G., Chen, H., Wu, N., Tian, J., Peng, C., Zhu, Q., Zhu, D., He, Y., Zheng, Q., and Zhang, C.: Effects of soil warming, rainfall reduction and water table level on CH<sub>4</sub> emissions from the Zoige peatland in China, *Soil Biology and Biochemistry*, 78, 83–89, <https://doi.org/10.1016/j.soilbio.2014.07.013>, 2014.
- Yang, Z., Zhu, D., Liu, L., Liu, X., and Chen, H.: The effects of freeze-thaw cycles on methane emissions from peat soils of a high altitude peatland, *Frontiers in Earth Science*, p. 452, 2022.
- 700 Yao, T., Xue, Y., Chen, D., Chen, F., Thompson, L., Cui, P., Koike, T., Lau, W. K.-M., Lettenmaier, D., Mosbrugger, V., Zhang, R., Xu, B., Dozier, J., Gillespie, T., Gu, Y., Kang, S., Piao, S., Sugimoto, S., Ueno, K., Wang, L., Wang, W., Zhang, F., Sheng, Y., Guo, W., Ailikun, Yang, X., Ma, Y., Shen, S. S. P., Su, Z., Chen, F., Liang, S., Liu, Y., Singh, V. P., Yang, K., Yang, D., Zhao, X., Qian, Y., Zhang, Y., and Li, Q.: Recent Third Pole’s Rapid Warming Accompanies Cryospheric Melt and Water Cycle Intensification and Interactions between Monsoon and Environment: Multidisciplinary Approach with Observations, Modeling, and Analysis, *Bulletin of the American Meteorological Society*, 100, 423–444, <https://doi.org/10.1175/BAMS-D-17-0057.1>, 2019.
- 705 Zhang, G., Zhang, Y., Dong, J., and Xiao, X.: Green-up dates in the Tibetan Plateau have continuously advanced from 1982 to 2011, *Proceedings of the National Academy of Sciences of the United States of America*, 110, 4309–4314, <https://doi.org/10.1073/pnas.1210423110>, 2013.
- Zhao, Y., Yu, Z., and Zhao, W.: Holocene vegetation and climate histories in the eastern Tibetan Plateau: controls 710 by insolation-driven temperature or monsoon-derived precipitation changes?, *Quaternary Science Reviews*, 30, 1173–1184, <https://doi.org/10.1016/j.quascirev.2011.02.006>, 2011.
- Zhou, W., Cui, L., Wang, Y., Li, W., and Kang, X.: Carbon emission flux and storage in the degraded peatlands of the Zoige alpine area in the Qinghai–Tibetan Plateau, *Soil Use and Management*, 37, 72–82, 2021.