

# Moisture and temperature effects on the radiocarbon signature of respired carbon dioxide to assess stability of soil carbon in the Tibetan Plateau

Andrés Tangarife-Escobar<sup>1</sup>, Georg Guggenberger<sup>2</sup>, Xiaojuan Feng<sup>3</sup>, Guohua Dai<sup>3</sup>, Carolina Urbina-Malo<sup>2</sup>, Mina Azizi-Rad<sup>1</sup>, and Carlos A. Sierra<sup>1,4</sup>

<sup>1</sup>Max Planck Institute for Biogeochemistry, Jena, Germany

<sup>2</sup>Institute of Soil Science, Leibniz Universität Hannover, Hannover, Germany

<sup>3</sup>Institute of Botany, Chinese Academy of Sciences, Beijing, China

<sup>4</sup>Department of Ecology, Swedish University of Agricultural Sciences, Uppsala, Sweden

**Correspondence:** Andrés Tangarife-Escobar (atanga@bgc-jena.mpg.de)

**Abstract.** Microbial release of CO<sub>2</sub> from soils to the atmosphere reflects how environmental conditions affect the stability of soil organic matter (SOM), especially in massive organic-rich ecosystems like the peatlands and grasslands of the Qinghai-Tibetan Plateau (QTP). Radiocarbon (<sup>14</sup>C) is an important tracer of the global carbon cycle and can be used to understand SOM dynamics through the estimation of time lags between carbon fixation and respiration, often assessed with metrics such as age and transit time. In this study, we incubated peatland and grassland soils at four temperature (5, 10, 15 and 20 °C) and two water-filled pore space (WFPS) levels (60 and 95 %), and measured the <sup>14</sup>C signature of bulk soil and [heterotrophic](#) respired CO<sub>2</sub>. We compared the relation between the  $\Delta^{14}\text{C}$  of the bulk soil and the  $\Delta^{14}\text{CO}_2$  of respired carbon as a function of temperature and WFPS for the two soils. To better interpret our results, we used a mathematical model to analyse how the calculated number of pools, decomposition rates of carbon ( $k$ ), transfer ( $\alpha$ ) and partitioning ( $\gamma$ ) coefficients affect the  $\Delta^{14}\text{C}$ -bulk and  $\Delta^{14}\text{CO}_2$  relation, with their respective mean age and mean transit time. From our incubations, we found that <sup>14</sup>C values in bulk and CO<sub>2</sub> from peatland were significantly more depleted (old) than from grassland soil. Our results showed that changes in temperature did not affect the  $\Delta^{14}\text{C}$  values of [heterotrophic](#) respired CO<sub>2</sub> in either soil. However, changes in WFPS had a small effect on the <sup>14</sup>CO<sub>2</sub> in grassland soils and a significant influence in peatland soils, where higher WFPS levels led to more depleted  $\Delta^{14}\text{CO}_2$ . In our models, the correspondence between  $\Delta^{14}\text{C}$ , age and transit time highly depended on the internal dynamics of the soil ( $k$ ,  $\alpha$ ,  $\gamma$  and number of pools) as well as on model structure. We observed large differences between slow and fast cycling systems, where low values of decomposition rates modified the  $\Delta^{14}\text{C}$  values in a non-linear pattern due to the incorporation of modern carbon (<sup>14</sup>C-bomb) in the soil. We concluded that the stability of carbon in the peatland and grassland soils of the QTP depends strongly on the direction of change in moisture and how it affects the rates of SOM decomposition while temperature regulates the amount of fluxes. Current land cover modification (desiccation) in Zoige peatlands and climate change occurring on the QTP, might largely increase CO<sub>2</sub> fluxes along with the release of old carbon to the atmosphere potentially shifting carbon sinks into sources.

## 1 Introduction

Studying soil organic matter (SOM) stability and persistence in a globally-changing environment is of fundamental importance to understand temporal variations of carbon cycling in the earth-climate system. Soil constitutes the largest carbon (C) stock in the terrestrial biosphere (Chen et al., 2021), making it a key component in global climate models (McGuire et al., 2001; Wieder et al., 2013; Xu et al., 2016). Physical, chemical and biological properties determine SOM decomposition, and in consequence its persistence over decades to millennia (Schmidt et al., 2011). Temperature and moisture are two of the most important abiotic variables controlling the rates of SOM decomposition (Sierra et al., 2015) and its transit time across the different ecosystem pools. Changes in these environmental conditions are, however, occurring simultaneously, highlighting the necessity to conduct multifactorial experiments to disentangle the dominant mechanisms on such cycling times. Hence, discerning the relationship between SOM persistence and carbon cycling times in terrestrial ecosystems is imperative to improve climate change models and to inform decisions on CO<sub>2</sub> mitigation and management strategies (Bradford et al., 2016; Mesfin et al., 2021).

The Qinghai-Tibetan Plateau (QTP) with an area of 2.5 million km<sup>2</sup> is a macroregion of global importance for the cycling of water, carbon and other biogeochemical elements (Anslan et al., 2020). Land cover is mainly dominated by alpine grasslands (44 %) (Scurlock and Hall, 1998; Gao et al., 2014), which store 23.4 % of China's total organic carbon and 2.5 % of the global soil carbon (Genxu et al., 2002). The QTP also hosts one of the largest high-mountain marshes in the world (Xiang et al., 2009; Chen et al., 2014; Ma et al., 2016), the Zoige peatlands, considered to be the most important carbon stock in peatlands for China (Liu et al., 2018). However, climate change and land cover change are currently modulating the net carbon balance through CO<sub>2</sub> and CH<sub>4</sub> effluxes from the grasslands (Piao et al., 2012; Chen et al., 2017; Du and Gao, 2020) and the peatlands at both daily and interannual timescales (Hao et al., 2011; Chen et al., 2014; Kang et al., 2018; Liu et al., 2019b; Chen et al., 2021). Anthropogenic activities in the Zoige peatlands (drainage, peat mining and overgrazing) have caused a degradation of approximately 30 % of wet and dry meadows (Chen et al., 2014; Zhou et al., 2021). Additionally, the QTP has been facing an air temperature increase of 0.2°C per decade over the past 50 years (Zhang et al., 2013; Chen et al., 2014; Yang et al., 2014; Ganjurjav et al., 2016), equivalent to two to three times faster than the world average (Yao et al., 2019; Nieberding et al., 2020), along with a moderate increase in precipitation (Dong et al., 2018). According to the IPCC (Arias et al., 2021), climate models predict an increment in heavy precipitation events and high temperature extremes for this region.

Such changes in temperature and soil moisture control the magnitude in which stabilization and destabilization mechanisms enable carbon storage in the QTP (Xiang et al., 2009; Ma et al., 2016). It has been observed that warming increases soil respiration (Rustad et al., 2001; Lu et al., 2013; Pold et al., 2015) and that soil moisture modulates ecosystem and soil respiration (Geng et al., 2012; Piao et al., 2012; Moyano et al., 2013; Pan et al., 2022; Azizi-Rad et al., 2022). Although the influence of destabilization mechanisms on SOM decomposition has been already studied in the Zoige peatlands (Zhao et al., 2011; Wang et al., 2015; Liu et al., 2016, 2018; Li et al., 2018; Liu et al., 2019a), temporal scales of SOM persistence under temperature and soil moisture variations are not clearly understood yet. Therefore, a quantification of the SOM persistence is of vital importance

to predict climate change feedback magnitudes and pathways.

A useful approach to quantify SOM persistence is through the theory of compartmental dynamical systems, where soil is understood as a set of interconnected pools with transformations of carbon to different forms (gas, dissolved or solid) (Emanuel et al., 1984; Schimel, 1995; Sierra et al., 2018a). Therefore, carbon can be characterized by the time it remains inside a compartment or the entire system (Eriksson, 1971; Bolin and Rodhe, 1973) through the calculation of age and transit time. Here, age is defined as the time elapsed since the carbon entered the system until the time of observation in the bulk soil, while transit time is defined as the time spent by the carbon between the entry to the system and its exit as respired CO<sub>2</sub> (Eriksson, 1971; Bolin and Rodhe, 1973; Manzoni et al., 2009; Sierra et al., 2018b). These timescale metrics are however, not the same for all the carbon atoms in a soil, since physical, chemical and biological stabilization and destabilization processes of SOM modify the rate at which mixing and storage occurs.

Radiocarbon measurements in bulk soil and in respired CO<sub>2</sub> are a powerful tool to approximate ages and transit times of carbon in soils as they trace back the trajectory of carbon through the different stocks on decadal to millennia timescales (Trumbore, 2000; Sierra et al., 2014; Schuur et al., 2016; Estop-Aragones et al., 2020; Chen et al., 2021). The radiocarbon content reflects the time a carbon atom has been in the soil since it was fixed by photosynthesis from the atmosphere (Trumbore, 2000). Carbon fixed since the early 1960's has higher <sup>14</sup>C than carbon fixed previously due to the enrichment by thermonuclear weapons testing. By comparing the <sup>14</sup>C relative abundance in the carbon pools with the atmospheric <sup>14</sup>CO<sub>2</sub> concentrations, it is possible to model soil-atmosphere carbon cycling times (Trumbore, 2000).

<sup>14</sup>C values can be altered by SOM destabilization processes and soil characteristics such as soil organic carbon (SOC) content, age and diagenetic state, thaw depth, redox state, seasonality, etc. (Gaudinski et al., 2000; Trumbore, 2000; von Lützow et al., 2008; Sierra et al., 2018b; Estop-Aragones et al., 2020). In recent decades, it has been widely observed that temperature plays a major role on the dynamics of SOM (Knorr et al., 2005; Davidson and Janssens, 2006; Feng and Simpson, 2008) and the mean age of respired CO<sub>2</sub> (Hopkins et al., 2012; Chen et al., 2021) by increasing the decomposition rates from fast-cycling pools (Trumbore et al., 1996) and mobilizing old carbon in warming conditions (Dutta et al., 2006; Briones et al., 2010). In contrast, other studies have suggested that warming does not lead to release of old carbon (Briones et al., 2021; Dioumaeva et al., 2002) and that non-labile carbon decomposition is insensitive to temperature increase (Liski et al., 1999). By comparison, drying phenomena increased the release of modern <sup>14</sup>CO<sub>2</sub> from shallow soil layers but preserved the old soil carbon pools in deeper layers (Kwon et al., 2019). Observations in rich-SOC soils of the Arctic indicated the release of old carbon from deep soil layers after thaw and drainage processes (Schuur et al., 2009; Estop-Aragones et al., 2020; Pegoraro et al., 2021). Such a process might be occurring in the Zoige peatland soils due to the presence of seasonal frozen layers (Liu et al., 2021; Yang et al., 2022). Nonetheless, the influence of temperature and soil moisture on SOM persistence is a matter of current debate.

90 Changes in temperature and moisture contribute to the destabilization of carbon in soils from the QTP. Hence, we hypothesize that higher temperature would increase the age of respired CO<sub>2</sub> as well as changes in soil moisture would increase or decrease (depending on the direction of moisture change) the age of respired CO<sub>2</sub> in soils subjected to controlled manipulations. Greater ages of respired CO<sub>2</sub> would imply that previously stabilized carbon is destabilized by changes in the manipulated environmental factors. For this purpose, we posed three specific research questions: (1) Do specific changes in temperature and  
95 moisture result in the release of old carbon in the respiration flux? (2) Are there differences in the age of [heterotrophic](#) respired CO<sub>2</sub> between a grassland and a peatland soil from the QTP? (3) How can radiocarbon data in bulk soil and [heterotrophic](#) respired CO<sub>2</sub> be interpreted to understand the effect of changes in decomposition rates on ages and transit times? To address these questions, we conducted a controlled multifactorial experiment with soils from two different ecosystems in the QTP, and measured the radiocarbon signature in the bulk soil and the [heterotrophic](#) respired CO<sub>2</sub>. In addition, we used a mathematical  
100 model to better interpret the interaction between decomposition rates change, expressed through internal dynamics of the soil ( $k$ ,  $\alpha$ ,  $\gamma$  and number of pools) and the  $\Delta^{14}\text{C}$  values by targeting the range found in the incubations. Thus, our observations and models strengthen each other to gain a deeper comprehension on the relationship between soil carbon stability,  $\Delta^{14}\text{C}$ , age and transit time.


## 105 2 Materials and methods

To evaluate the influence of soil moisture and temperature on the  $^{14}\text{C}$  values from bulk soil and [heterotrophic](#) respired CO<sub>2</sub> of grasslands and peatlands soils, we incubated soil samples from a high elevation grassland at the Nam Co catchment collected in September 2018 and a peatland from the Zoige region sampled in July 2021, both located in the QTP.

### 2.1 Site description and sampling

#### 110 2.1.1 Nam Co grassland

The Nam Co site (Fig. 1) is located in the central Tibetan Plateau (altitude of the Nam Co lake 4,726 m a.s.l., 30°46'N, 90°59'E) and represents a frequent study location for monitoring and tracking of environmental changes over different time scales (Anslan et al., 2020). The dominant climate at Nam Co consists of cold winters and short and moist summers. Mean annual temperature (MAT) measured at the NAMORS research station was -0.6 °C (2006 – 2017) and mean annual precipitation  
115 (MAP) was 406 mm, occurring mostly during the monsoon season between May and October (Anslan et al., 2020). The sampling area was mainly covered by *Kobresia pygmaea* and has been grazed intensively by yaks and sheep. A total amount of 40 kg of soil was collected from randomly selected spots at depths between 5 and 15 cm within an area of about 40 m<sup>2</sup>. The soil was then mixed homogeneously to produce one single sample which consisted of a sandy loam with a bulk density of 1.3 g cm<sup>-3</sup>, pH of 7.5, 3.2 % of total organic carbon, 0.05 % of inorganic carbon and 0.3 % of total nitrogen, cation exchange  
120 capacity of 89 mmol<sub>c</sub> kg<sup>-1</sup> soil and electrical conductivity of 89.6 μS cm<sup>-1</sup>.



**Figure 1.** a) Overview of the QTP; b) Location of the Nam Co lake with land cover dominated by grasslands, and c) Zoigê peatlands highlighting the location of Zoigê city. The sampling site is located in the southeastern of the peatland (Ruokeba, Hongyuan county). SRTM elevation data from Jarvis et al. (2008).

### 2.1.2 Zoigê peatlands

The Zoigê peatlands are located in the northeast of the QTP (average elevation of 3,400 m a.s.l.) and cover an area of about 4,605 km<sup>2</sup> on the headwaters of the Yellow River basin. These peatlands have been recognised as one of the largest high-mountain marshes in the world (Xiang et al., 2009; Ma et al., 2016) and store 88 % of the carbon in the QTP (Chen et al., 125 2014). The Zoigê region shows a MAT of 1.5 °C, where the warmest monthly temperature (11 °C) is recorded in July and the coldest in January (-10.1 °C). The MAP is 720 mm, and according to data from the Chinese National Meteorological Information Centre (www.nmic.gov.cn), the region has faced a slight drying trend and a warming of 0.4 °C per decade over the last 40 years (Chen et al., 2014; Yang et al., 2014). The soils are unfrozen from April to October while the layer between 0 and 50 cm depth is seasonally frozen between November and March. Vegetation cover consists mainly of *Potentilla anserine*, 130 *Blysmus sinocompressus*, *Kobresia myosuroides* and *Scirpus triqueter*. The samples (0 – 35 cm depth) were collected with a

spade from an area of 50 m<sup>2</sup> at a long-term monitoring site (33°4'5''N, 102°33'52''E), then thoroughly mixed to produce a single sample. The soil bulk density was 0.3 g cm<sup>-3</sup>, contained in average 27.6 % of organic carbon, 0.06 % of inorganic carbon and 1.8 % of total nitrogen.

## 2.2 Incubation of peatland and grassland soils

135 We conducted two sets of incubation experiments without pre-incubation period, one set with the grassland soil and a second set with the peatland soil. Samples were incubated with their original roots to minimise disturbance and allow comparisons with field conditions, however stones were removed. Briefly, 50-ml vials were filled with 12 g of soil ( $\pm$  1.5 g) and placed inside glass flasks along with 0.2 ml of water at the bottom of the flask (away from contact with the sample) to avoid possible drying (Dioumaeva et al., 2002). Thereafter the flasks were sealed with rubber plugs and screwed with plastic caps. Each of  
140 the sets was placed at two different WFPS levels (60 and 95 %) which were selected in order to reassemble the thaw and consequent water saturation of seasonally frozen soils in grasslands; and for peatland soils, the process of drying (through artificial desiccation) after high water saturation. Despite the high WFPS, the soil samples had still contact with air inside the vials, which guaranteed microbial decomposition of the organic matter and accumulation of heterotrophic CO<sub>2</sub> respiration. Soil moisture treatments were combined with four temperature levels (5, 10, 15 and 20 °C) for a total of 69 samples, 33 for  
145 grassland and 36 for peatland (2 sites  $\times$  4 temperature levels  $\times$  2 moisture levels  $\times$  3-6 analytical replicates, to prevent scarcity of data due to eventual failure in CO<sub>2</sub> extraction). Flasks with samples were flushed with synthetic air (CO<sub>2</sub> concentrations were measured at intervals of 1 to 2 weeks using a free) to remove atmospheric CO<sub>2</sub> analyzer LI-COR 6262-. This flushing marked the starting day of the incubations.

150 Incubations for each subset ~~ended simultaneously until every sample had an estimated concentration of~~ concluded concurrently once all the samples reached a C concentration (from CO<sub>2</sub>-C) in the headspace equivalent to  $\geq$  estimated to be equal to or exceeding 2 mg of C, enough for, sufficient for subsequent radiocarbon analysis. ~~Grassland This approach was not possible in two of the 16 subsets due to lab material limitations and therefore, grassland~~ This approach was not possible in two of the 16 subsets due to lab material limitations and therefore, grassland samples were incubated between 15 and 67 days, while peatland samples were incubated for 13 days (Table A1). ~~For sampling headspace air, 50-ml vials were filled with 12~~  
155 ~~g of soil ( $\pm$  1.5 g) and placed inside 0.5 L glass flasks along with 0.2 ml of water at the bottom of~~ The headspace volume of the incubation flasks was measured as 587 ml, which was corrected after adding the soil. In average, the final headspace was 575.6 and 533 ml for the incubated grassland and peatland soils, respectively. These values were used to calculate the fluxes of heterotrophic CO<sub>2</sub> respiration.

160 The rate of accumulation of CO<sub>2</sub> in the headspace of our incubations represents the diffusion rate of heterotrophic CO<sub>2</sub> respiration released from the flask (away from contact with the sample) to avoid possible drying (Dioumaeva et al., 2002) ; thereafter the flasks were sealed with rubber plugs and screwed with plastic caps. Flasks with samples were flushed with synthetic air (incubated soils. We refer throughout the manuscript to heterotrophic respiration, but we acknowledge that our measurements better capture how this heterotrophic CO<sub>2</sub> respiration flux diffuses out of the soil. Rates were measured at

165 intervals of 1 to 2 weeks using a CO<sub>2</sub> analyzer LI-COR 6262 for every treatment and mean heterotrophic CO<sub>2</sub> respiration rates (mg CO<sub>2</sub> free) to remove atmospheric g soil<sup>-1</sup> day<sup>-1</sup>) were calculated through the division of CO<sub>2</sub> concentration in the headspace by the product of the accumulation duration (days) (Table A1) and the mass of the introduced soil (g).

### 170 2.3 Radiocarbon analysis of incubated soils

Radiocarbon analysis were conducted in the bulk soil of each sample after the incubation. Soil inorganic carbon was eliminated through decalcification following the preparation protocol by Steinhof et al. (2017). Also, respired heterotrophic CO<sub>2</sub> respiration accumulated in the headspace of incubation flasks was extracted and purified on a vacuum line, graphitized by Fe reduction in H<sub>2</sub> and measured for Δ<sup>14</sup>C by Accelerator Mass Spectrometry (Micadas, Ionplus, Switzerland) in the radiocarbon laboratory of the Max Planck Institute for Biogeochemistry in Jena, Germany (Steinhof et al., 2017).

Radiocarbon data are expressed as Δ<sup>14</sup>C (the deviation in ‰ from 0.95 times the oxalic acid standard in 1950). The values were corrected to a δ<sup>13</sup>C value of -25 ‰ for differences in biological mass-dependent fractionation (Stuiver and Polach, 1977). These data are presented as percentage of Modern Carbon (pMC) that can be converted to F<sup>14</sup>C by dividing it by 100 and later to Δ<sup>14</sup>C using equation (1) (Stuiver and Polach, 1977).

$$\Delta^{14}C = \left[ F^{14}C e^{\lambda_C(1950-t)} - 1 \right] \times 1000 [\text{‰}], \quad (1)$$

where F<sup>14</sup>C is the Fraction Modern, i.e. the ratio of the measured sample normalized to a δ<sup>13</sup>C value of -25 ‰, divided by 0.95 times the measured ratio of the Oxalic Acid I standard (OX-I) (Schoor et al., 2016), λ<sub>C</sub> is the updated radiocarbon decay constant (equals 1/8267 [y<sup>-1</sup>]), and t is the year of sampling.

185

The effects of soil moisture and temperature manipulation on Δ<sup>14</sup>C were evaluated for the bulk and CO<sub>2</sub> fractions in each ecosystem separately through two-way ANOVA tests (type III). This type of ANOVA is also referred to as Partial Sum of Squares and is appropriate for unbalanced data since it does not depend on the sampling structure or the particular order in the model (Shaw and Mitchell-Olds, 1993); hence, this approach adjusts best to our data set where treatments did not have equal amount of values. As for the CO<sub>2</sub> fluxes, rates were measured regularly for every treatment and mean CO<sub>2</sub> respiration rates (mg CO<sub>2</sub> g soil<sup>-1</sup> day<sup>-1</sup>) were calculated based on the total duration of the incubation (Fig. A1).

### 190 2.4 SOC decomposition models to predict Δ<sup>14</sup>C as a proxy of SOM persistence

The representation of the SOC dynamics has been commonly described through models that can be expressed as systems of linear differential equations (Manzoni et al., 2009; Sierra et al., 2012; Sierra and Mueller, 2015; Sierra et al., 2017b) of the form:

195

$$\frac{d\mathbf{C}(t)}{dt} = \mathbf{I} + \mathbf{A} \cdot \mathbf{C}(t), \quad (2)$$

where the vector  $\mathbf{C}(t)$  is the rate of change of carbon over time in  $n$  different pools; the time dependent  $n$ -dimensional vector  $\mathbf{I}$  represents the total inputs of carbon to each pool; and  $\mathbf{A}$  represents the  $n \times n$  dimensional matrix with the rates of carbon processing for each pool in its main diagonal and the proportion of carbon transferred from one pool to another in the off-diagonals (Sierra et al., 2012, 2014; Metzler and Sierra, 2018; Sierra et al., 2018b). This mass balance equation has a radiocarbon counterpart:

$$\frac{d^{14}\mathbf{C}(t)}{dt} = \mathbf{I}_{14}\mathbf{C}(t) + \mathbf{A} \cdot {}^{14}\mathbf{C}(t) - \lambda {}^{14}\mathbf{C}(t) \quad (3)$$

where  $\lambda$  is the radiocarbon decay constant ( $1/8267 \text{ [y}^{-1}\text{]}$ ).



We used a SOC decomposition model to calculate the  $\Delta^{14}\text{C}$  in bulk and  $\text{CO}_2$  as well as their equivalent mean age and mean transit time of a theoretical soil for the year 2019 calibrated with the data set for the Northern Hemisphere Zone 3 (Hua et al., 2021). The simulated  $\Delta^{14}\text{C}$  results aimed at finding the same  $\Delta^{14}\text{C}$  ranges obtained from the incubated soils stabilising different ranges of decomposition rates for peatland and grassland, with the objective to compare under which settings the models could describe best laboratory or field conditions and for this reason, model fitting was not necessary. For this purpose, we implemented a two-pool model (one slow and one fast pool) considering two different structures: parallel and series (Fig. 2) and modified the parameters involved in the soil carbon processing (Manzoni et al., 2009; Falloon and Smith, 2000) (Table 1). In the parallel structure, carbon enters the soil and splits among the two pools according to  $\gamma$  and decomposition occurs independently in each pool according to their respective  $k$ . In the series structure, carbon enters only to the fast pool and it is either decomposed and emitted to the atmosphere or transferred to the slow pool according to the transfer coefficient  $\alpha$ .

215



$$\frac{d\mathbf{C}}{dt} = I \begin{pmatrix} \gamma \\ 1-\gamma \end{pmatrix} + \begin{pmatrix} -k_1 & 0 \\ 0 & -k_2 \end{pmatrix} \cdot \begin{pmatrix} C_1 \\ C_2 \end{pmatrix}$$

$$\frac{d\mathbf{C}}{dt} = I \begin{pmatrix} 1 \\ 0 \end{pmatrix} + \begin{pmatrix} -k_1 & 0 \\ \alpha_{2,1}k_1 & -k_2 \end{pmatrix} \cdot \begin{pmatrix} C_1 \\ C_2 \end{pmatrix}$$



**Figure 2.** Parallel (left) and series (right) model structure and formulas implemented in the simulations. Carbon is fixed into the biomass from atmospheric CO<sub>2</sub> through photosynthesis, subsequently it is incorporated into the soil as litterfall. From this state on, it is considered as input ( $I$ ) and will split between the two pools according to the different structures ( $\gamma$  for parallel and  $\alpha$  for series). The rate at which the carbon is decomposed ( $k$ ) in each pool ( $C$ ) will depend on microbial activity, environmental factors (temperature and soil moisture) and physical and chemical protection of the SOM (Blanco-Canqui and Lal, 2004; von Lützow et al., 2008; Manzoni et al., 2009). Boxes represent the soil pools, which account with an initial amount of carbon ( $C_1 = 200$ ,  $C_2 = 5000$ ) and arrows represent directions of inputs and outputs.

Initial parameters of the model such as the initial year of simulation and initial  $\Delta^{14}\text{C}$  values of each pool were considered separately for each type of ecosystem. Moreover, we defined  $I$  and  $C$  as constant since they can be adjusted depending on the specific soil characteristics. Additionally, we assumed that decomposition rates  $k$  reflect the effect of temperature and soil moisture on carbon cycling timescales (Manzoni et al., 2009), and therefore on  $\Delta^{14}\text{C}$  values. In our approach, only one parameter at time was modified for each simulation.

**Table 1.** Definition of parameters used to evaluate the variation of  $\Delta^{14}\text{C}$  values, mean age and mean transit time and their ranges used in the simulation experiments.

Parameter	Notation	Type	Value	Definition
Litter Input	I	Constant	100	A scalar or a data.frame object specifying the amount of litter inputs by time
Decomposition rate	$k_1, k_2$	Variable	0.00001 - 0.8	A vector of length 2 with the values of the decomposition rate for pools 1 and 2
Partitioning coefficient	$\gamma$	Variable	0 - 1	A scalar representing the proportion of I that goes to pool 1 in a parallel structure
Transference coefficient	$\alpha$	Variable	0 - 1	A scalar with the value of the transfer rate from pool 1 to pool 2
Carbon stocks	$C_1, C_2$	Constant	200, 5000	Initial amount of C for the two pools

We used the R package SoilR (Sierra et al., 2014) to simulate the temporal dynamics of  $\Delta^{14}\text{C}$  in the bulk soil and the respired  $\text{CO}_2$  as well as the age and transit time distributions of carbon. Assuming steady-state for the carbon stocks, the probability density function (pdf) of the age (equation 4) and the transit time (equation 5) (Metzler and Sierra, 2018) as well as their means  
225 can be calculated by the following expressions (Sierra et al., 2018b):

$$f(a) = -\mathbf{1}^T \cdot \mathbf{A} \cdot e^{a \cdot \mathbf{A}} \cdot \frac{\mathbf{C}^*}{\sum \mathbf{C}^*}, \quad a \geq 0, \quad (4)$$

where  $a$  is the random variable age,  $\mathbf{1}^T$  is the transpose of the  $n$ -dimensional vector containing ones,  $e^{a \cdot \mathbf{A}}$  is the matrix exponential for each value of  $a$ , and  $\sum \mathbf{C}^*$  is the sum of stocks of all pools at steady state.

The pdf of transit times can be obtained as

$$230 \quad f(\tau) = -\mathbf{1}^T \cdot \mathbf{A} \cdot e^{\tau \cdot \mathbf{A}} \cdot \frac{\mathbf{I}}{\sum \mathbf{I}}, \quad \tau \geq 0, \quad (5)$$

where  $\tau$  represents the random variable transit time.

These pdfs measure the probability that a certain amount of carbon is above or below a specific age or transit time (Sierra et al., 2018b) (equations 6 and 7, respectively). The mean of the age, that is, the expected value of the pdf can be computed by the expression:

$$235 \quad \mathbb{E}(a) = -\mathbf{1}^T \cdot \mathbf{A}^{-1} \cdot \frac{\mathbf{C}^*}{\sum \mathbf{C}^*}, \quad (6)$$

and the mean of the transit time by


$$\mathbb{E}(\tau) = -\mathbf{1}^T \cdot \mathbf{A}^{-1} \cdot \frac{\mathbf{I}}{\sum \mathbf{I}}. \quad (7)$$

Based on the age and transit time distribution for the different simulated cases, metrics such as the mean, median and quantiles can be used as proxies of SOM persistence. The relation between ages and transit times may present three cases:  $\mathbb{E}(a) = \mathbb{E}(\tau)$  (Type I - well-mixed homogeneous system),  $\mathbb{E}(a) < \mathbb{E}(\tau)$  (Type II - retention system) and  $\mathbb{E}(a) > \mathbb{E}(\tau)$  (Type III - non-retention system) (Sierra et al., 2018b). In the type I, the probability of mineralization and release as  $\text{CO}_2$  is the same for every C atom. In the type II, the carbon is retained for a relatively long time before it is released. In type III, most of the C atoms stay in the system for a short period of time, but some atoms remain for a long time (Bolin and Rodhe, 1973). Comparisons between these distributions will provide detailed information on mixing, store, recycling, transport and transformation processes of the SOM (Sierra et al., 2018b).


### 3 Results

#### 3.1 $\Delta^{14}\text{C}$ values of incubated soils under temperature and soil moisture variation

Values of  $\Delta^{14}\text{C}$  from the incubation experiment contrasted strongly between grassland and peatland soils (Fig. 3 and 4). Generally,  $\Delta^{14}\text{C}$  of bulk soil and [heterotrophic  \$\text{CO}\_2\$](#)  respiration from the peatland soil was more depleted than from the grassland soil. For example, in peatland soil the  $\Delta^{14}\text{C}$  values of bulk soil (-85.9 to -60.9 ‰, mean=-80.1, n=36, including outlier of -180) were clearly more depleted than those of the [heterotrophic](#) respired flux (-20.9 to 23.9 ‰, mean=4, n=36), which indicated that the peatland behaved as a retention system (type II). In contrast, for the grassland soil, the  $\Delta^{14}\text{C}$  of bulk soil (21.1 to 73.9 ‰, mean=43.3, n=33) fell similarly around the 1:1 line compared to the  $\Delta^{14}\text{CO}_2$  (13.9 to 83.4 ‰, mean=38.5, n=33, including outliers of -227.1 and -105.1) indicating that the samples behaved mostly as a well-mixed homogeneous system (type I). Such similar values can also reflect a type II system where  $\Delta^{14}\text{C}$  values in the bulk and the respired flux are equal for the year of sampling (Fig. A7).



**Figure 3.** Boxplot with the variation of the  $\Delta^{14}\text{C}$  values in  $\text{CO}_2$  (left panel) and bulk soil (right panel) for peatland and grassland incubated soils.



**Figure 4.** Relationship between  $\Delta^{14}\text{C}$  in bulk soil and  $\Delta^{14}\text{C}$  in [heterotrophic](#) respired CO<sub>2</sub> for incubated grassland (green box) and peatland (brown box) soils of the QTP discriminated by temperature and WFPS. Possible types of system according to the  $\Delta^{14}\text{C}$  relations between bulk and respired carbon.

The temperature treatments did not systematically affect the radiocarbon signature of the bulk or the [heterotrophic](#) respired CO<sub>2</sub> in peatland or grassland soils (Table 2). There was significant evidence that manipulations in WFPS resulted in changes in the  $\Delta^{14}\text{C}$  values of bulk soil and CO<sub>2</sub> ( $p$ -values = 0.09 in grasslands and  $p$ -value = 0.01 in peatlands from an ANOVA test, Table 2, Fig. 5 A-B) except for the bulk soil in peatlands. When the interacting effects of WFPS and temperature were evaluated together, there was no evidence that their interplay affected the radiocarbon signature of both soils ( $p$ -values > 0.05 for all the analysis). Outliers presented extremely depleted  $\Delta^{14}\text{C}$  values and occurred both in bulk (peatland) and CO<sub>2</sub> (grassland) at the combined treatment WFPS = 60 and temperature = 10 °C. Such wide variation in  $\Delta^{14}\text{C}$  both between ecosystems and treatments could be potentially explained by intrinsic processes affecting soil carbon dynamics, which will be explored in the following sections with a SOC decomposition model.

Higher temperature and WFPS caused an increase of CO<sub>2</sub> fluxes from heterotrophic respiration in the treated incubated soils (Table A1, Fig. 5 C-D). In both ecosystems, wetter conditions showed higher respiration rates and higher slopes as the temperature increased. The absolute amounts of CO<sub>2</sub> produced from peatland soil was in average 14 times higher than from grassland soils for every independent treatment.

**Table 2.** Summary of *p*-values obtained from ANOVA tests for the  $\Delta^{14}\text{C}$  of bulk soil and the  $\Delta^{14}\text{CO}_2$  of grassland and peatland soils after incubation. *p*-values are given for the independent effect of temperature (T) and WFPS as well as the integrated effect of temperature and WFPS (T · WFPS).


Ecosystem	Type	T	WFPS	T · WFPS
Grassland	$\Delta^{14}\text{C}$ Bulk	0.8	0.091	0.21
	$\Delta^{14}\text{CO}_2$	0.73	0.089	0.92
Peatland	$\Delta^{14}\text{C}$ Bulk	0.4	0.21	0.65
	$\Delta^{14}\text{CO}_2$	0.16	0.01	0.32

### 3.2 Effect of changes of decomposition rates (*k*) on the $\Delta^{14}\text{C}$ values

Changes in the vertical and horizontal direction of the  $\Delta^{14}\text{C}$ -bulk versus  $\Delta^{14}\text{C}$ -CO<sub>2</sub> space are more evident across ecosystem type, which at the same time implies specific environmental conditions for the stability of SOM. To understand possible drivers of  $\Delta^{14}\text{C}$  changes in these directions and how they can be interpreted in terms of ages and transit times, we used a SOC decomposition model. We evaluated how model structure, decomposition rates of carbon, and the partitioning ( $\gamma$ ) and transfer ( $\alpha$ ) coefficients of a two-pool parallel and a series model affected  $\Delta^{14}\text{C}$  values, and as a consequence, the mean age and the mean transit time. Our simulations mimicked a fast cycling grassland and a slow cycling peatland by differentiating the ranges of decomposition rates (Table 3) and showed how the modelled conditions affected the type of system (I, II and III).

#### 3.2.1 SOC decomposition in fast cycling systems (grasslands)

Variation of  $k_1$  yielded  $\Delta^{14}\text{C}$  curves similar to mean age and mean transit time. In other words, in these simulations the change in the parameter  $k_1$  resulted in similar trends in  $\Delta^{14}\text{C}$ -bulk versus  $\Delta^{14}\text{C}$ -CO<sub>2</sub> space as in the mean age versus mean transit time space ((Fig. 6). In the parallel structure (Fig. 6, A and B), high values of  $k_1$  yielded more enriched  $\Delta^{14}\text{C}$  values in the bulk soil than in the heterotrophic respired flux. This was expressed as a fast respiration of SOC and reflected in a short transit time. Simultaneously, low values of  $k_2$  resulted in a slow respiration, which was registered in ages longer than transit times, at the initial state of the carbon stay in the system. However, as  $k_1$  decreased and approached  $k_2$ , mean age and mean transit time became similar and converged to the 1:1 line. In the case of the series structure (Fig. 6, C and D), the total amount of inputs traveled first through the fast pool, which meant that while  $k_1$  is high, most of the carbon decomposition occurs under



**Figure 5.** Comparison between  $\Delta^{14}\text{C}$  of [heterotrophic](#) respiration from incubated grassland (A) and peatland (B) soil at different temperature levels under WFPS = 60 and 95 %. Black points represent minimum and maximum values out of the range between quartile 1 and 3 (25 to 75 % of the data). The quartile 50 (median) represented by the line inside the box indicates the midpoint value in the frequency distribution. Box for the treatment WFPS= 60 % and T= 10 °C shows a large dispersion of the 50 % of the data, which is explained by the outliers observed in Fig. 4. Additional panels indicate the [heterotrophic](#) respiration rates ( $\text{mg CO}_2 \text{ g soil}^{-1} \text{ day}^{-1}$ ) for each treatment based on the [total duration of incubation the flux accumulation](#) (see [Fig.-A+Table A1](#)). Response of mean [heterotrophic](#) respiration rates to temperature and WFPS treatments in grassland (C) and peatland (D) soils.

the dynamics of the fast pool. Hence, the system showed equal mean age and mean transit time. As  $k_1$  decreased, the transfer from the fast to the slow pool becomes more relevant, then the mean transit time increased compared to the mean age.

The behavior of  $\Delta^{14}\text{C}$  values contrasted significantly with that of mean age and mean transit time when  $k_2$  tended to lower values (Fig. 7). Generally, when  $k_2$  equals  $k_1$ ,  $\Delta^{14}\text{C}$  and mean age and mean transit time showed similar values. However, as  $k_2$  decreased, the carbon stayed in the slow pool for longer time and the  $\Delta^{14}\text{C}$  enriched in the bulk soil until reaching a peak (112

**Table 3.** Range of parameters used for simulations in a SOC decomposition model for fast and slow systems. These ranges explore the required conditions to describe the  $\Delta^{14}\text{C}$  variation found in the incubated grassland and peatland soils and their equivalent age and transit time relationship. The target  $\Delta^{14}\text{C}$  intervals are shown as boxes in the Fig. 6 to 9.

System	Fig.	Model structure	$k_1$	$k_2$	$\gamma$	$\alpha$	Starting year of simulation	$\Delta^{14}\text{C}$ (‰) for starting year
Fast cycling	6	Parallel	0.1 - 0.8	0.1	0.8	-	1890	-4.9 ‰
		Series	0.1 - 0.8	0.1	-	0.8	1890	-4.9 ‰
	7	Parallel	0.8	0.001 - 0.8	0.8	-	1890	-4.9 ‰
		Series	0.8	0.001 - 0.8	-	0.8	1890	-4.9 ‰
Slow cycling	8	Parallel	0.0001 - 0.8	0.0001	0.8	-	500	-22.2 ‰
		Series	0.0001 - 0.8	0.0001	-	0.8	500	-22.2 ‰
	9	Parallel	0.1	0.000001 - 0.001	0.2	-	500	-22.2 ‰
		Series	0.1	0.000001 - 0.001	-	0.8	500	-22.2 ‰

295 and 89 ‰ for series and parallel structures, respectively), from where it subsequently depleted to the initial  $\Delta^{14}\text{CO}_2$  value. For the parallel structure (Fig. 7, A and B), transit time kept increasing slowly since most of the carbon stayed in the fast pool for a short time, so, the carbon remaining in the slow pool contributed to increase the mean age. As for the series structure, most of the carbon was transferred from the fast to the slow pool due to the high transfer rate  $\alpha$ , which contributed to increase the transit time.


300

### 3.2.2 SOC decomposition in slow cycling systems (peatlands)

Simulations of  $\Delta^{14}\text{C}$  values in slow cycling systems resulted in significantly more depleted values in both bulk soil and respired  $\text{CO}_2$  (Fig. 8) than in fast cycling systems. Generally,  $\Delta^{14}\text{C}$  values were more depleted in the bulk soil than in the respiration flux. Also, the results from these simulations showed very different patterns in the  $\Delta^{14}\text{C}$ -bulk versus  $\Delta^{14}\text{C}$ - $\text{CO}_2$  space than in the mean age versus mean transit time space; only with the exception of equal  $k_1$  and  $k_2$ , which resulted in similar  $\Delta^{14}\text{C}$  values as well as similar mean age and mean transit time in the parallel structure.

310 Additionally, we looked at the variation of  $\Delta^{14}\text{C}$  keeping the same  $k$  values but reducing  $\alpha$  and  $\gamma$  to 0.1. Simulations indicated that the behavior of  $\Delta^{14}\text{C}$  values were opposite between series and parallel structures when inputs to each pool were inversely proportional (Fig. 7 versus Fig. A3 and Fig. 8 versus Fig. A4). For example, for the case of low values of  $k_1$  and  $k_2$  (Fig. 9) ( $\gamma = 0.2$  and  $\alpha = 0.8$ ) both model structures showed similar patterns due to relatively similar amounts of inputs going






**Figure 6.** Predictions of  $\Delta^{14}\text{C}$  in bulk soil vs  $\Delta^{14}\text{CO}_2$  with their equivalent simulation of mean age in bulk soil vs mean transit time in  $\text{CO}_2$  for parallel (panel A and B) and series model structure (C and D) for a fast cycling system. Variation of  $k_1$  with  $\alpha = 0.8$  and  $\gamma = 0.8$ . Complementary prediction with  $\alpha = 0.1$  and  $\gamma = 0.1$  can be seen in Fig. A2. Green box represents the range of measured  $\Delta^{14}\text{C}$  values obtained from incubated grassland soils for bulk (20 to 74 ‰) and  $\text{CO}_2$  (10 to 85 ‰), excluding outliers.

to the slow pool. When  $\gamma = 0.8$ ,  $\Delta^{14}\text{C}$  showed values out of the target range (-20 to 23 for  $\Delta^{14}\text{C-CO}_2$  and -86 to -60 for  $\Delta^{14}\text{C-bulk}$  based on incubation results) since decomposition occurred as in a fast cycling system. As  $k_2$  decreased,  $\Delta^{14}\text{CO}_2$  enriched and  $\Delta^{14}\text{C}$  and mean age remained higher than mean transit time since the slow decomposition of the slow pool dominated the system response.

315

### 3.3 Variation in the proportion of inputs $\gamma$ and $\alpha$ modulated mean age and mean transit time


Our simulations showed the importance that the partitioning and transference coefficients ( $\gamma$  and  $\alpha$ ) have to define the amount of carbon entering to each pool, and in consequence on the mean age and mean transit time (Fig. A1). For a parallel structure,



**Figure 7.** Predictions of  $\Delta^{14}\text{C}$  in bulk soil vs  $\Delta^{14}\text{CO}_2$  with their equivalent simulation of mean age in bulk soil vs mean transit time in  $\text{CO}_2$  for parallel (panel A and B) and series model structure (C and D) for a fast cycling system. Variation of  $k$  with  $\alpha = 0.8$  and  $\gamma = 0.8$ . Green box represents the range of measured  $\Delta^{14}\text{C}$  values obtained from incubated grassland soils for bulk (20 to 74 ‰) and  $\text{CO}_2$  (10 to 85 ‰), excluding outliers.

320 when the total amount of inputs entered only to the fast pool ( $\gamma = 1$ ), the system behaved as a one pool system and mean age and mean transit time were equal. In the extreme opposite ( $\gamma = 0$ ), all carbon entered only to the slow pool and mean age and mean transit time converged, although with a longer transit through the system. In contrast, for a series structure, the extreme value  $\alpha = 1$  resulted in a longer mean transit time than mean age as the contribution to the respired flux comprehended the sum of the decomposition of the newly absorbed carbon and the initial stocks stored in the fast pool. In the case of slow cycling systems

325 (Fig. A5 and Fig. A6),  $\Delta^{14}\text{CO}_2$  moved in a wider range including very depleted values (-100 to 70 ‰) and approached to a steady state of  $\Delta^{14}\text{C}$  in bulk soil from the right when  $k_1 = 0.01$  or from the left when  $k_1 = 0.1$ .



**Figure 8.** Predictions of  $\Delta^{14}\text{C}$  in bulk soil vs  $\Delta^{14}\text{CO}_2$  with their equivalent simulation of mean age in bulk soil vs mean transit time in  $\text{CO}_2$  for parallel (panel A and B) and series model structure (C and D) for a slow cycling system. Variation of  $k_1$  with  $\alpha = 0.8$  and  $\gamma = 0.8$ . Brown box represents the range of measured  $\Delta^{14}\text{C}$  values obtained from incubated peatland soils for bulk (-90 to -65 ‰) and  $\text{CO}_2$  (-18 to 25 ‰), excluding outliers.


## 4 Discussion

### 4.1 Did changes in temperature and soil moisture result in the release of old carbon in the heterotrophic respiration flux?

330

We measured the  $\Delta^{14}\text{C}$  values of heterotrophic respired  $\text{CO}_2$  from soils incubated at different temperatures and WFPS levels to assess whether changes in these variables would result in destabilization of stored carbon. In general, higher temperature and WFPS levels resulted in larger  $\text{CO}_2$  fluxes although with a faster increment in peatland soils, which could be explained by the high amount of total organic carbon available for decomposition. Respiration Heterotrophic respiration rates, however, did not correlate with the  $\Delta^{14}\text{C}$  values. Our results showed that changes in temperature did not affect systematically the radiocarbon

335



**Figure 9.** Predictions of  $\Delta^{14}\text{C}$  in bulk soil vs  $\Delta^{14}\text{CO}_2$  with their equivalent simulation of mean age in bulk soil vs mean transit time in  $\text{CO}_2$  for parallel (panel A and B) for a slow cycling system. Series model presented exactly the same behaviour. Variation of  $k_2$  with  $\alpha = 0.8$  and  $\gamma = 0.2$ . Brown box represents the range of measured  $\Delta^{14}\text{C}$  values obtained from incubated peatland soils for bulk (-90 to -65 ‰) and  $\text{CO}_2$  (-18 to 25 ‰), excluding outliers.

values of [heterotrophic](#) respired  $\text{CO}_2$  in any of the incubated soils. Nevertheless, changes in WFPS had a significant effect on the  $\Delta^{14}\text{CO}_2$  and  $\Delta^{14}\text{C}$ -bulk of grassland soils and only on the  $\Delta^{14}\text{CO}_2$  of peatland soils. Our experiments indicated that higher WFPS levels led to depleted  $\Delta^{14}\text{CO}_2$  values in peatlands at all temperature treatments, except at 10 °C.

340 In contrast, higher WFPS resulted in more enriched  $\Delta^{14}\text{CO}_2$  values for grasslands. This suggests that the direction of changing WFPS depends on ecosystem characteristics. It is interesting to observe that the  $\Delta^{14}\text{C}$  in the bulk soil was always more negative than the  $\Delta^{14}\text{CO}_2$  in the peatlands, while the grasslands registered similar  $\Delta^{14}\text{C}$  values in both states. This indicates that peatlands are systems that stabilize organic matter over time and release it once the stable conditions change. In comparison, for grasslands two interpretations may arise, first that there was not organic matter stabilization under the incubated

345 conditions since the respired flux reassembles the bulk soil  $\Delta^{14}\text{C}$  signature, or second that the incubation conditions were not strong or long enough to destabilize the existing old SOM. Our simulations indicate that similar  $\Delta^{14}\text{C}$  values in the bulk and in the respired  $\text{CO}_2$  can also occur at a specific year due to the dynamics of the atmospheric bomb curve (Fig A7) independently of the different SOM cycling times in the soil pools.

350 In tundra ecosystems, Kwon et al. (2019) suggested that drainage of shallow soil layers may have accelerated old carbon decomposition. In addition, Estop-Aragones et al. (2020) concluded that old carbon would increase in proportion from “cold” across “warm wet” to “warm and dry” for high arctic tundra as well as from “undisturbed” to “burnt active layer” for peatland plateau. These previous findings are in disagreement with our observations since after our experiments, lower WFPS resulted in the release of relatively enriched  $\Delta^{14}\text{CO}_2$  instead of releasing old carbon. One potential explanation for this discrepancy is that  
355 the existing old SOM could have reacted too slow to drier conditions (WFPS = 60) compared to the fast response of “younger” SOM, causing an evident prevalence of young carbon given the short duration of the incubations for peatlands. Consistent with this, it has been found that some SOM components or compartments may be more sensitive to modified conditions than others (Feng and Simpson, 2008) and therefore, transit time would react strongly when such sensitivity is higher in the slower pool (Manzoni et al., 2009).

360

For this reason, our results are indicative, but not conclusive, about the influence of environmental factors on the  $\Delta^{14}\text{C}$  signature of [heterotrophic](#) respired carbon. It is possible that our  $\Delta^{14}\text{C}$  values are limited by the short time of the incubations since the dynamics of carbon transfers cannot be properly observed in short timescales (Crow and Sierra, 2018). For example, during a 1-year incubation experiment, most of the  $\text{CO}_2$  was derived from labile SOM as the temperature increased (Leifeld  
365 and Fuhrer, 2005). Additionally, other factors such as yearly climate seasonality, daily freeze-thaw cycles and water table oscillation were not replicated in the laboratory due to equipment and time availability. An important aspect that has been observed in soil incubations, is that  $\text{CO}_2$  accumulation decreases or even stops after a certain period probably due to the  $\text{CO}_2$  saturation of the limited head-space in the incubation flasks, which depends on the SOC content. From this we could deduce that old carbon, which usually needs longer times to be decomposed, can only be recovered if soil respiration is not space limited. For  
370 example, Azizi-Rad et al. (2022) found the respiration rate to decline after 14 days holding the soil at 10 °C. In this sense, incubations with high TOC may run out of headspace soon and affect the interpretation of the  $\Delta^{14}\text{C}$  values. Finally, although temperature has been found to be the main modulator of SOC decomposition rates (Azizi-Rad et al., 2022), dominating over the effect of soil moisture and oxygen availability, it is important to consider that an increase in respiration rates, as observed in our incubations, does not necessarily involve the release of  $\Delta^{14}\text{C}$  of lower or higher signatures as shown in our model  
375 simulations.

## 4.2 Are there differences in the age of heterotrophic respired CO<sub>2</sub> between grassland and peatland?

We found that  $\Delta^{14}\text{C}$  values from peatland were strongly more depleted than those in grasslands, indicating the presence and respiration of older carbon in peatlands. Generally, low temperature, low soil microbial activity and anoxic conditions (Xiang et al., 2009; Ma et al., 2016) posed favourable conditions for the stabilization of organic matter since the early Holocene in the Zoige region (Chen et al., 2014; Sun et al., 2017). Older bulk soil than CO<sub>2</sub> demonstrated that the peatland soil behaved as a retention system (Type II) where carbon was stored for relatively long time before its respiration (Sierra et al., 2018b). The release of that old carbon may occur when the SOM destabilization dominates over the stabilization mechanisms, for example when peatlands are drained (increasing oxidation through water table reduction) and their temperature is raised (Dutta et al., 2006; Hicks Pries et al., 2013; Lupascu et al., 2014; Estop-Aragónés et al., 2018), both factors mimicked in our incubations.

In contrast,  $\Delta^{14}\text{CO}_2$  from grasslands were mostly similar to  $\Delta^{14}\text{C}$  in bulk soil, indicating that the soil behaved predominantly as a well-mixed homogeneous system (Type I) where most of the SOM has the same probability of being mineralized and released as CO<sub>2</sub> (Sierra et al., 2018b). However,  $\Delta^{14}\text{C}$  relation also indicated that the grasslands shift from retention to non-retention system and vice-versa. In grasslands, SOM from the topsoil is permanently under changing conditions (daily temperature and soil moisture variation, grazing and mechanical alteration, among others) that promote its fast cycling instead of its stabilization (Han et al., 2017). These factors, along with climatic, vegetation and edaphic properties caused a low SOM stability expressed in a large proportion of labile carbon (Hou et al., 2021). Nonetheless, even though SOM accumulation still occurs in grassland areas (Tian et al., 2009) registered in the extremely depleted outliers from the grassland soil respiration, it happens at a slower rate than in peatlands. As a result, most of the fixed atmospheric carbon is respired at short time scales, which is in turn registered as young (“post bomb”)  $\Delta^{14}\text{C}$  values. Finally, although extremely depleted  $\Delta^{14}\text{C}$  values are rare, they are probable since the measured carbon particles represent one value, which can fall on the tail of the system-age probability distribution. We interpret that outlier results belong to carbon particles that have remained for very long time in the system (out of the mean values) whose specific soil characteristics were not captured in our model structure or internal soil conditions. The reasons behind the occurrence of these outliers at the specific 10 °C remain to be investigated.

Taken together, SOM stability and ecosystem characteristics defined the relation between  $\Delta^{14}\text{C}$  in bulk soil and  $\Delta^{14}\text{CO}_2$ , which in turn explained the difference in the age of heterotrophic respiration between grassland and peatlands. The  $\Delta^{14}\text{CO}_2$  values obtained in our incubated peatlands indicate that such carbon was captured through photosynthesis during a time when the  $^{14}\text{C}$  levels of the atmosphere were depleted and have remained stored in the soil for a long enough time for radioactive decay to become relevant. However, it is imprecise to attribute an specific year to the entire bulk soil or CO<sub>2</sub> fractions, since they are formed by SOM accumulated in multiple steps and of different qualities and their respective decomposition.

In that sense, a  $\Delta^{14}\text{C}$  value is not per se indicative of age or transit time of SOC. To disentangle the utility of radiocarbon as a tool for SOM persistence in soil and be able to shed light on the time-scales of carbon cycling in different ecosystems, we

used SOC decomposition models. Our models contribute to understand the time component of SOM persistence through the use of mean transit time as a parameter that integrates processes of SOM dynamics (Manzoni et al., 2009).

### 4.3 How can $\Delta^{14}\text{C}$ values be interpreted to understand the effect of decomposition rates on mean age and mean transit time?

415

We simulated the dynamics of  $\Delta^{14}\text{C}$  as decomposition rates change, and how this behavior differ from those of mean age and mean transit time. Our simulations were able to reassemble the  $\Delta^{14}\text{C}$  ranges obtained from incubations and showed that modelled  $\Delta^{14}\text{C}$  values differed widely between slow cycling systems and fast cycling systems. Generally, decomposition rate, transfer rates and partitioning coefficients of a given model structure modulated  $\Delta^{14}\text{C}$  values, and in consequence, mean age and mean transit time (Bruun et al., 2004; Manzoni et al., 2009). For example, fast cycling systems with high decomposition rates resulted in a fast respiration represented by short mean transit times (lower than 20 years). At the same time, low decomposition rates resulted in longer mean transit times, which in turn represented older respired  $\text{CO}_2$  (lower than 140 years). In slow cycling systems, where the decomposition rates were lower, transit times were consequently longer (lower than 20000 years). Independently of the decomposition rates values, the series model structure increased the transit time of carbon compared to parallel structure due to the fact that a proportion of the carbon atoms had to pass through the two pools before being respired.

420

425

Other factors such as observation time and starting year of the simulation modified the response of  $\Delta^{14}\text{C}$ . This is due to the radioactive nature of the  $^{14}\text{C}$  isotope and the influence of the “bomb  $^{14}\text{C}$ ”. For instance, the  $\Delta^{14}\text{C}$  enrichment during the last 60 years was clearly observed in fast cycling systems only with low  $k_2$ . Furthermore, the inflexion points on the  $\Delta^{14}\text{C}$  curves occurred at different values for slow and fast cycling systems. Hence, we could assume that a simulation with starting year after 1962 would not show an inflexion point in the  $\Delta^{14}\text{C}$  due to the absence of the bomb peak.

430

Our modelling results suggest that the increase of decomposition rates contribute to the release of older carbon in the respired flux as we hypothesised, but depending on the initial state of the system. This interpretation is not straightforward since the stability of SOM depends on the specific combination of temperature and moisture for different ecosystems. For peatlands, only drier conditions and consequent increase of oxygen might increase decomposition rates and therefore cause the destabilization of SOM. In contrast, grasslands would need an increase of temperature to facilitate SOM decomposition provided that moisture and oxygen are available (Sierra et al., 2017a; Azizi-Rad et al., 2022). SOM decomposition rates are expected to increase at higher temperatures (Leifeld and Fuhrer, 2005) and towards the extremes of the moisture range (Sierra et al., 2015). However, the destabilization of SOM can occur in any direction of WFPS variation (depending on the oxygen content) and in turn affect the age and transit time of carbon in a non-linear pattern.

440

From our results we could observe how the relation  $\Delta^{14}\text{C}$ -bulk versus  $\Delta^{14}\text{C}$ - $\text{CO}_2$  properly represented the relation mean  
445 age versus mean transit time. We found that there is a good correspondence between both relations in the fast cycling systems  
as long as the decomposition rates ( $k$ ) remain high. Such correspondence did not occur as  $k$  became smaller (typical for slow  
cycling systems) since the appearance of the bomb peak may have introduced anomalies that modified the equivalence between  
the two relations. In addition, the outlier  $\Delta^{14}\text{C}$  values found in our experimental data, where  $\Delta^{14}\text{C}$  changed drastically in the  
vertical and the horizontal direction, may be related to this complex response of the  $^{14}\text{C}$  tracer.

450

To quantify cycling times of carbon, radiocarbon can be used as a tool to understand SOM destabilization and persistence  
through the use of the concepts of age and transit time and their mutual relation. Nonetheless, it is essential to couple  $\Delta^{14}\text{C}$   
measured values with a model that involves the dynamics of soil carbon in different pools and their interaction with the envi-  
ronment. Therefore, the acquisition of empirical data from soils (number of pools,  $I$ ,  $C$ ,  $k$ ,  $\gamma$ ,  $\alpha$ ) along with the correct setting  
455 of model structure will improve predictions on terrestrial and atmospheric carbon interactions.

## 5 Conclusions

Based on the incubation results of soils from the QTP, we showed that the  $\Delta^{14}\text{C}$  values of the peatland are significantly more  
depleted than the ones of the grassland both in the bulk soil and the [heterotrophic](#) respired  $\text{CO}_2$ . Our results indicated that  
460 changes in temperature did not affect systematically the radiocarbon values of [heterotrophic](#) respired  $\text{CO}_2$  in any of the incu-  
bated soils despite increases in [heterotrophic](#) respired C at higher temperatures. Nevertheless, changes in WFPS had a relatively  
small effect on the  $^{14}\text{CO}_2$  in grasslands, but a significant influence on the  $^{14}\text{CO}_2$  of peatlands, where higher WFPS levels led  
to more depleted  $^{14}\text{CO}_2$  values except at 10 °C. In peatlands, more depleted  $^{14}\text{C}$ -bulk values than  $^{14}\text{C}$ - $\text{CO}_2$  indicated that  
SOM stabilizes over time and it is released once the stable conditions change. In grasslands, similar  $\Delta^{14}\text{C}$  values in bulk and  
465 [heterotrophic](#) respired  $\text{CO}_2$  indicated that the soil behaved as a well-mixed homogeneous system due to either an absence of  
SOM stabilization or that the manipulation treatments were not long or strong enough to destabilize the existing old SOM. In  
this sense, the short duration of our incubations might have been an obstacle to register the influence of long term factors such  
as climate seasonality and water table oscillation on SOM dynamics.

470 From our modelling approach, we conclude that radiocarbon can be used as a tool to understand SOM persistence through  
the use of the concepts of mean age and mean transit time and their mutual relation. Our simulations were able to reassemble  
the  $\Delta^{14}\text{C}$  values obtained from incubations and showed that modelled  $\Delta^{14}\text{C}$  values differed widely between slow cycling  
systems and fast cycling systems. We found that low values of decomposition rates, more common in slow cycling systems,  
modified the behaviour of  $\Delta^{14}\text{C}$  patterns due to the incorporation of  $^{14}\text{C}$ -bomb in the soil system. Hence, the correspondence  
475 between these mutual relations strongly depended on the internal dynamics of the soil and its interaction with the environment.  
For this reason, the acquisition of empirical data from soils (number of pools,  $I$ ,  $C$ ,  $k$ ,  $\gamma$  and  $\alpha$ ) along with the correct setting of



model structure will improve our understanding on the stability of carbon in the soils of a changing QTP. In this way, current changes in climate patterns and land cover alteration may have a larger impact on the Zoige peatlands than on the grasslands given the vulnerability of large carbon stocks to be destabilized by changes in temperature. Nevertheless, the interaction with moisture may dampen or amplify the temperature effect, adding uncertainty on the future trajectories of soil carbon in the Qinghai-Tibetan Plateau.

*Code and data availability.* Lab analysis results and code are stored at <https://doi.org/10.5281/zenodo.10537332> (DOI:10.5281/zenodo.10537332) (Tangarife-Escobar et al., 2024).

*Author contributions.* The conceptualization, data curation and formal analysis were done by ATE and CS. The investigation was performed by ATE with support from GD and MAR. Methodology and validation were done by ATE and CS. ATE conducted the validation, visualization and writing of the original draft (the latter with the help of CUM). Supervision was done by GG, XF and CS. ATE wrote the manuscript with contributions from CS, GG, XF, CUM, GD and MAR.

*Competing interests.* The authors declare that they have no conflict of interest.

*Acknowledgements.* We thank all colleagues who contributed to this study, especially Eike Reinosch for providing the location maps used in Fig.1. and Paula Sierra for the digitization of the scheme used in Fig. 2. Special thanks to David Martini for his helpful advices to improve code writing and to Manuel Röst and Axel Steinhof for their useful training at the <sup>14</sup>C laboratory. Finally, thanks to Nicole Börner for the active collaboration with the project administration and to Susan Trumbore for her feedback on the final version of this manuscript. This study was developed as part of the International Research Training Group (GRK 2309/1) Geo-ecosystems in transition on the Tibetan Plateau (TransTiP) funded by the Deutsche Forschungsgemeinschaft (DFG). The Max Planck Institute for Biogeochemistry provided permanent administrative support.

## Appendix A: Appendix A


**Table A1.** Mean daily CO<sub>2</sub> heterotrophic respiration for incubated soils under temperature and WFPS variation.

Ecosystem	<del>Temperature</del> Temperature (°C)	<del>WFPS</del> WFPS (%)	Mean CO <sub>2</sub> respiration (mg CO <sub>2</sub> g soil <sup>-1</sup> day <sup>-1</sup> )	$\sigma$	<u>Flux duration</u> (day)	Incubation time n = number of
Grassland	20	95	<del>0.040</del> <u>0.044</u>	<del>0.035</del> <u>0.009</u>	<u>19</u>	2n = <del>32</del> <u>19</u> , 1
	20	60	0.028	<del>0.016</del> <u>0.003</u>	<u>19</u>	4n =
	15	95	<del>0.024</del> <u>0.023</u>	<del>0.008</del> <u>0.004</u>	<u>15</u>	3n = 30, 1
	15	60	0.016	<del>0.004</del> <u>0.003</u>	<u>15</u>	4n =
	10	95	0.011	<del>0.011</del> <u>0.001</u>	<u>60</u>	3n =
	10	60	<del>0.008</del> <u>0.007</u>	<del>0.023</del> <u>0.001</u>	<u>60</u>	6n =
	5	95	0.005	<del>0.003</del> <u>0.000</u>	<u>60</u>	5n =
	5	60	0.004	0.001	<u>60</u>	4n =
Peatland	20	95	<del>0.510</del> <u>0.462</u>	<del>0.001</del> <u>0.032</u>	<u>7</u>	5n =
	20	60	<del>0.371</del> <u>0.337</u>	<del>0.003</del> <u>0.014</u>	<u>7</u>	4n =
	15	95	<del>0.330</del> <u>0.299</u>	<del>0.004</del> <u>0.007</u>	<u>7</u>	5n =
	15	60	<del>0.234</del> <u>0.212</u>	<del>0.003</del> <u>0.004</u>	<u>7</u>	3n =
	10	95	<del>0.209</del> <u>0.189</u>	<del>0.002</del> <u>0.010</u>	<u>9</u>	5n =
	10	60	<del>0.134</del> <u>0.121</u>	<del>0.001</del> <u>0.020</u>	<u>9</u>	5n =
	5	95	<del>0.066</del> <u>0.061</u>	<del>0.000</del> <u>0.004</u>	<u>7</u>	4n =
	5	60	<del>0.044</del> <u>0.040</u>	0.001	<u>7</u>	5n =


**Table A2.** Range of parameters used for simulations in a SOC decomposition model for fast and slow systems (↻) (Tangarife-Escobar et al., 2024). These ranges explore the required conditions to describe the  $\Delta^{14}\text{C}$  variation found in the incubated grassland and peatland soils and their equivalent age and transit time relationship. The target  $\Delta^{14}\text{C}$  intervals are shown as boxes in the Fig. A1 to A6.

System	Fig.	Model structure	$k_1$	$k_2$	$\gamma$	$\alpha$	Starting year of simulation	$\Delta^{14}\text{C}$ (‰) for starting year
Fast cycling	A1	Parallel	0.8	0.1	0 - 1	-	1890	-4.9 ‰
		Series	0.8	0.1	-	0 - 1	1890	-4.9 ‰
	A2	Parallel	0.1 - 0.8	0.1	0.1	-	1890	-4.9 ‰
		Series	0.1 - 0.8	0.1	-	0.1	1890	-4.9 ‰
	A3	Parallel	0.8	0.0001 - 0.8	0.1	-	1890	-4.9 ‰
		Series	0.8	0.0001 - 0.8	-	0.1	1890	-4.9 ‰
Slow cycling	A4	Parallel	0.0001 - 0.8	0.0001	0.2	-	500	-22.2 ‰
		Series	0.0001 - 0.8	0.0001	-	0.1	500	-22.2 ‰
	A5	Parallel	0.01	0.0001	0 - 1	-	500	-22.2 ‰
		Series	0.01	0.0001	-	0 - 1	500	-22.2 ‰
	A6	Parallel	0.1	0.0001	0 - 1	-	500	-22.2 ‰
		Series	0.1	0.0001	-	0 - 1	500	-22.2 ‰


-  $\Delta^{14}\text{C}$  variation in fast cycling systems



**Figure A1.** Predictions of  $\Delta^{14}\text{C}$  in bulk soil vs  $\Delta^{14}\text{CO}_2$  with their equivalent simulation of mean age in bulk soil vs mean transit time in  $\text{CO}_2$  for parallel (panel A and B) and series model structure (C and D) for a fast cycling system.  $k_1 = 0.8$ ,  $k_2 = 0.1$ , with  $\alpha = 0 - 1$  and  $\gamma = 0 - 1$ . Green box represents the range of measured  $\Delta^{14}\text{C}$  values obtained from incubated grassland soils for bulk (20 to 74 ‰) and  $\text{CO}_2$  (10 to 85 ‰), excluding outliers.




**Figure A2.** Predictions of  $\Delta^{14}\text{C}$  in bulk soil vs  $\Delta^{14}\text{CO}_2$  with their equivalent simulation of mean age in bulk soil vs mean transit time in  $\text{CO}_2$  for parallel (panel A and B) and series model structure (C and D) for a fast cycling system. Variation of  $k_1$  with  $\alpha = 0.1$  and  $\gamma = 0.1$ . Green box represents the range of measured  $\Delta^{14}\text{C}$  values obtained from incubated grassland soils for bulk (20 to 74 ‰) and  $\text{CO}_2$  (10 to 85 ‰), excluding outliers.




**Figure A3.** Predictions of  $\Delta^{14}\text{C}$  in bulk soil vs  $\Delta^{14}\text{CO}_2$  with their equivalent simulation of mean age in bulk soil vs mean transit time in  $\text{CO}_2$  for parallel (panel A and B) and series model structure (C and D) for a fast cycling system. Variation of  $k_2$  with  $\alpha = 0.1$  and  $\gamma = 0.1$ . Green box represents the range of measured  $\Delta^{14}\text{C}$  values obtained from incubated grassland soils for bulk (20 to 74 ‰) and  $\text{CO}_2$  (10 to 85 ‰), excluding outliers.

-  $\Delta^{14}\text{C}$  variation in slow cycling systems




**Figure A4.** Predictions of  $\Delta^{14}\text{C}$  in bulk soil vs  $\Delta^{14}\text{CO}_2$  with their equivalent simulation of mean age in bulk soil vs mean transit time in  $\text{CO}_2$  for parallel (panel A and B) and series model structure (C and D) for a slow cycling system. Variation of  $k_1$  with  $\alpha = 0.1$  and  $\gamma = 0.2$ . Brown box represents the range of measured  $\Delta^{14}\text{C}$  values obtained from incubated peatland soils for bulk (-90 to -65 ‰) and  $\text{CO}_2$  (-18 to 25 ‰), excluding outliers.




**Figure A5.** Predictions of  $\Delta^{14}\text{C}$  in bulk soil vs  $\Delta^{14}\text{CO}_2$  with their equivalent simulation of mean age in bulk soil vs mean transit time in  $\text{CO}_2$  for parallel (panel A and B) and series model structure (C and D) for a slow cycling system.  $k_1 = 0.01$ ,  $k_2 = 0.0001$ , with  $\alpha = 0 - 1$  and  $\gamma = 0 - 1$ . Brown box represents the range of measured  $\Delta^{14}\text{C}$  values obtained from incubated peatland soils for bulk (-90 to -65 ‰) and  $\text{CO}_2$  (-18 to 25 ‰), excluding outliers.





**Figure A6.** Predictions of  $\Delta^{14}\text{C}$  in bulk soil vs  $\Delta^{14}\text{CO}_2$  with their equivalent simulation of mean age in bulk soil vs mean transit time in  $\text{CO}_2$  for parallel (panel A and B) and series model structure (C and D) for a slow cycling system.  $k_1 = 0.1$ ,  $k_2 = 0.0001$ , with  $\alpha = 0 - 1$  and  $\gamma = 0 - 1$ . Brown box represents the range of measured  $\Delta^{14}\text{C}$  values obtained from incubated peatland soils for bulk (-90 to -65 ‰) and  $\text{CO}_2$  (-18 to 25 ‰), excluding outliers.

-  $\Delta^{14}\text{C}$  curves for bulk soil, respired  $\text{CO}_2$  and atmosphere for a two-pool system in parallel and series structure



**Figure A7.** Variation of  $\Delta^{14}\text{C}$  values in atmosphere, bulk soil and respired  $\text{CO}_2$  for a two-pool soil during the period 1950 - 2019. The similarity between  $\Delta^{14}\text{C}$  values may indicate both a well-mixed homogeneous system (type I) or a retention system (type II) where the bulk soil and the respired flux record the same value. Nonetheless, the mean transit time for the bulk and the soil respiration might be different due to the contrasting decomposition rates of the fast ( $k_1 = 0.8$ ) and the slow pool ( $k_2 = 0.1$ ). Simulation conducted with  $\alpha = 0.8$  and  $\gamma = 0.8$ .

## 500 **References**

- Anslan, S., Rad, M. A., Buckel, J., Galindo, P. E., Kai, J., Kang, W., Keys, L., Maurischat, P., Nieberding, F., Reinosch, E., Tang, H., Tran, T. V., Wang, Y., and Schwalb, A.: Reviews and syntheses: How do abiotic and biotic processes respond to climatic variations in the Nam Co catchment (Tibetan Plateau)?, *Biogeosciences*, 17, 1261–1279, <https://doi.org/10.5194/bg-17-1261-2020>, 2020.
- Arias, P., Bellouin, N., Coppola, E., Jones, R., Krinner, G., Marotzke, J., Naik, V., Palmer, M., Plattner, G.-K., Rogelj, J., et al.: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change; Technical Summary, 2021.
- Azizi-Rad, M., Guggenberger, G., Ma, Y., and Sierra, C. A.: Sensitivity of soil respiration rate with respect to temperature, moisture and oxygen under freezing and thawing, *Soil Biology and Biochemistry*, 165, 108488, <https://doi.org/https://doi.org/10.1016/j.soilbio.2021.108488>, 2022.
- Blanco-Canqui, H. and Lal, R.: Mechanisms of carbon sequestration in soil aggregates, *Critical reviews in plant sciences*, 23, 481–504, 2004.
- Bolin, B. and Rodhe, H.: A note on the concepts of age distribution and transit time in natural reservoirs, *Tellus*, 25, 58–62, <https://doi.org/https://doi.org/10.1111/j.2153-3490.1973.tb01594.x>, 1973.
- Bradford, M. A., Wieder, W. R., Bonan, G. B., Fierer, N., Raymond, P. A., and Crowther, T. W.: Managing uncertainty in soil carbon feedbacks to climate change, *Nature Climate Change*, 6, 751–758, 2016.
- Briones, M. J., Garnett, M. H., and Ineson, P.: No evidence for increased loss of old carbon in a temperate organic soil after 13 years of simulated climatic warming despite increased CO<sub>2</sub> emissions, *Global Change Biology*, 2021.
- Briones, M. J. I., Garnett, M. H., and Ineson, P.: Soil biology and warming play a key role in the release of ‘old C’ from organic soils, *Soil Biology and Biochemistry*, 42, 960–967, <https://doi.org/10.1016/j.soilbio.2010.02.013>, 2010.
- Bruun, S., Six, J., and Jensen, L. S.: Estimating vital statistics and age distributions of measurable soil organic carbon fractions based on their pathway of formation and radiocarbon content, *Journal of theoretical biology*, 230, 241–250, 2004.
- Chen, H., Yang, G., Peng, C., Zhang, Y., Zhu, D., Zhu, Q., Hu, J., Wang, M., Zhan, W., Zhu, E., Bai, Z., Li, W., Wu, N., Wang, Y., Gao, Y., Tian, J., Kang, X., Zhao, X., and Wu, J.: The carbon stock of alpine peatlands on the Qinghai-Tibetan Plateau during the Holocene and their future fate, *Quaternary Science Reviews*, 95, 151–158, <https://doi.org/10.1016/j.quascirev.2014.05.003>, 2014.
- Chen, L., Jing, X., Flynn, D. F., Shi, Y., Kühn, P., Scholten, T., and He, J.-S.: Changes of carbon stocks in alpine grassland soils from 2002 to 2011 on the Tibetan Plateau and their climatic causes, *Geoderma*, 288, 166–174, 2017.
- Chen, L., Fang, K., Wei, B., Qin, S., Feng, X., Hu, T., Ji, C., and Yang, Y.: Soil carbon persistence governed by plant input and mineral protection at regional and global scales, *Ecology Letters*, 24, 1018–1028, 2021.
- Crow, S. E. and Sierra, C. A.: Dynamic, intermediate soil carbon pools may drive future responsiveness to environmental change, *Journal of environmental quality*, 47, 607–616, 2018.
- Davidson, E. and Janssens, I.: Temperature sensitivity of soil carbon decomposition and feedbacks to climate change, *Nature*, 440, 165–173, <https://doi.org/10.1038/nature04514>, 2006.
- Dioumaeva, I., Trumbore, S., Schuur, E., Goulden, M., Litvak, M., and Hirsch, A.: Decomposition of peat from upland boreal forest: Temperature dependence and sources of respired carbon, *Journal of Geophysical Research: Atmospheres*, 108, <https://doi.org/10.1029/2001JD000848>, 2002.
- Dong, S., Peng, F., You, Q., Guo, J., and Xue, X.: Lake dynamics and its relationship to climate change on the Tibetan Plateau over the last four decades, *Regional environmental change*, 18, 477–487, 2018.

- Du, C. and Gao, Y.: Opposite patterns of soil organic and inorganic carbon along a climate gradient in the alpine steppe of northern Tibetan Plateau, *Catena*, 186, 104366, 2020.
- Dutta, K., Schuur, E. A. G., Neff, J. C., and Zimov, S. A.: Potential carbon release from permafrost soils of Northeastern Siberia, *Global Change Biology*, 12, 2336–2351, <https://doi.org/10.1111/j.1365-2486.2006.01259.x>, 2006.
- 540 Emanuel, W., Killough, G., Post, W., and Shugart, H.: Modeling terrestrial ecosystems in the global carbon-cycle with shifts in carbon storage capacity by land-use change, *Ecology*, 65, 970–983, <https://doi.org/10.2307/1938069>, 1984.
- Eriksson, E.: Compartment Models and Reservoir Theory, *Annual Review of Ecology and Systematics*, 2, 67–84, <https://doi.org/10.1146/annurev.es.02.110171.000435>, 1971.
- 545 Estop-Aragones, C., Czimczik, C. I., Heffernan, L., Gibson, C., Walker, J. C., Xu, X., and Olefeldt, D.: Respiration of aged soil carbon during fall in permafrost peatlands enhanced by active layer deepening following wildfire but limited following thermokarst, *Environmental Research Letters*, 13, 085002, 2018.
- Estop-Aragones, C., Olefeldt, D., Abbott, B. W., Chanton, J. P., Czimczik, C. I., Dean, J. F., Egan, J. E., Gandois, L., Garnett, M. H., Hartley, I. P., Hoyt, A., Lupascu, M., Natali, S. M., O'Donnell, J. A., Raymond, P. A., Tanentzap, A. J., Tank, S. E., Schuur, E. A. G., Turetsky, M., and Anthony, K. W.: Assessing the Potential for Mobilization of Old Soil Carbon After Permafrost Thaw: A Synthesis of C-14 Measurements From the Northern Permafrost Region, *Global Biogeochemical Cycles*, 34, <https://doi.org/10.1029/2020GB006672>, 2020.
- 550 Falloon, P. and Smith, P.: Modelling refractory soil organic matter, *Biology and fertility of soils*, 30, 388–398, 2000.
- Feng, X. and Simpson, M. J.: Temperature responses of individual soil organic matter components, *Journal of Geophysical Research: Biogeosciences*, 113, 2008.
- 555 Ganjurjav, H., Gao, Q., Gornish, E. S., Schwartz, M. W., Liang, Y., Cao, X., Zhang, W., Zhang, Y., Li, W., Wan, Y., Li, Y., Danjiu, L., Guo, H., and Lin, E.: Differential response of alpine steppe and alpine meadow to climate warming in the central Qinghai-Tibetan Plateau, *Agricultural and Forest Meteorology*, 223, 233–240, <https://doi.org/10.1016/j.agrformet.2016.03.017>, 2016.
- Gao, Q.-z., Li, Y., Xu, H.-m., Wan, Y.-f., and Jiangcun, W.-z.: Adaptation strategies of climate variability impacts on alpine grassland ecosystems in Tibetan Plateau, *Mitigation and Adaptation Strategies for Global Change*, 19, 199–209, 2014.
- 560 Gaudinski, J., Trumbore, S., Davidson, E., and Zheng, S.: Soil carbon cycling in a temperate forest: radiocarbon-based estimates of residence times, sequestration rates and partitioning of fluxes, *Biogeochemistry*, 51, 33–69, <https://doi.org/10.1023/A:1006301010014>, 2000.
- Geng, Y., Wang, Y., Yang, K., Wang, S., Zeng, H., Baumann, F., Kuehn, P., Scholten, T., and He, J.-S.: Soil respiration in Tibetan alpine grasslands: belowground biomass and soil moisture, but not soil temperature, best explain the large-scale patterns, *PLoS one*, 7, e34968, 2012.
- 565 Genxu, W., Ju, Q., Guodong, C., and Yuanmin, L.: Soil organic carbon pool of grassland soils on the Qinghai-Tibetan Plateau and its global implication, *Science of the Total Environment*, 291, 207–217, 2002.
- Han, C., Wang, Z., Si, G., Lei, T., Yuan, Y., and Zhang, G.: Increased precipitation accelerates soil organic matter turnover associated with microbial community composition in topsoil of alpine grassland on the eastern Tibetan Plateau, *Canadian journal of microbiology*, 63, 811–821, 2017.
- 570 Hao, Y. B., Cui, X. Y., Wang, Y. F., Mei, X. R., Kang, X. M., Wu, N., Luo, P., and Zhu, D.: Predominance of Precipitation and Temperature Controls on Ecosystem CO<sub>2</sub> Exchange in Zoige Alpine Wetlands of Southwest China, *Wetlands*, 31, 413–422, <https://doi.org/10.1007/s13157-011-0151-1>, 2011.

- Hicks Pries, C. E., Schuur, E. A., and Crummer, K. G.: Thawing permafrost increases old soil and autotrophic respiration in tundra: Partitioning ecosystem respiration using D13C an D14C, *Global Change Biology*, 19, 649–661, 2013.
- Hopkins, F. M., Torn, M. S., and Trumbore, S. E.: Warming accelerates decomposition of decades-old carbon in forest soils, *Proceedings of the National Academy of Sciences of the United States of America*, 109, E1753–E1761, <https://doi.org/10.1073/pnas.1120603109>, 2012.
- Hou, Y., He, K., Chen, Y., Zhao, J., Hu, H., and Zhu, B.: Changes of soil organic matter stability along altitudinal gradients in Tibetan alpine grassland, *Plant and Soil*, 458, 21–40, 2021.
- 580 Hua, Q., Turnbull, J. C., Santos, G. M., Rakowski, A. Z., Ancapichún, S., De Pol-Holz, R., Hammer, S., Lehman, S. J., Levin, I., Miller, J. B., et al.: Atmospheric radiocarbon for the period 1950–2019, *Radiocarbon*, pp. 1–23, 2021.
- Jarvis, A., Reuter, H. I., Nelson, A., and Guevara, E.: Hole-filled SRTM for the globe Version 4. available from the CGIAR-CSI SRTM 90m Database, 15, (25–54), <http://srtm.csi.cgiar.org>, 2008.
- Kang, X., Yan, L., Cui, L., Zhang, X., Hao, Y., Wu, H., Zhang, Y., Li, W., Zhang, K., Yan, Z., Li, Y., and Wang, J.: Reduced Carbon Dioxide Sink and Methane Source under Extreme Drought Condition in an Alpine Peatland, *Sustainability*, 10, <https://doi.org/10.3390/su10114285>, 2018.
- 585 Knorr, W., Prentice, I., House, J., and Holland, E.: Long-term sensitivity of soil carbon turnover to warming, *Nature*, 433, 298–301, <https://doi.org/10.1038/nature03226>, 2005.
- Kwon, M. J., Natali, S. M., Pries, C. E. H., Schuur, E. A. G., Steinhof, A., Crummer, K. G., Zimov, N., Zimov, S. A., Heimann, M., Kolle, O., and Goeckede, M.: Drainage enhances modern soil carbon contribution but reduces old soil carbon contribution to ecosystem respiration in tundra ecosystems, *Global Change Biology*, 25, 1315–1325, <https://doi.org/10.1111/gcb.14578>, 2019.
- Leifeld, J. and Fuhrer, J.: The temperature response of CO<sub>2</sub> production from bulk soils and soil fractions is related to soil organic matter quality, *Biogeochemistry*, 75, 433–453, 2005.
- Li, J., Yan, D., Pendall, E., Pei, J., Noh, N. J., He, J.-S., Li, B., Nie, M., and Fang, C.: Depth dependence of soil carbon temperature sensitivity across Tibetan permafrost regions, *Soil Biology and Biochemistry*, 126, 82–90, 2018.
- 595 Liski, J., Ilvesniemi, H., Makela, A., and Westman, C.: CO<sub>2</sub> emissions from soil in response to climatic warming are overestimated - The decomposition of old soil organic matter is tolerant of temperature, *Ambio*, 28, 171–174, 1999.
- Liu, L., Chen, H., Zhu, Q., Yang, G., Zhu, E., Hu, J., Peng, C., Jiang, L., Zhan, W., Ma, T., He, Y., and Zhu, D.: Responses of peat carbon at different depths to simulated warming and oxidizing, *Science of the Total Environment*, 548, 429–440, <https://doi.org/10.1016/j.scitotenv.2015.11.149>, 2016.
- 600 Liu, L., Chen, H., Liu, X., Yang, Z., Zhu, D., He, Y., and Liu, J.: Contemporary, modern and ancient carbon fluxes in the Zoige peatlands on the Qinghai-Tibetan Plateau, *Geoderma*, 352, 138–149, <https://doi.org/10.1016/j.geoderma.2019.06.008>, 2019a.
- Liu, X., Zhu, D., Zhan, W., Chen, H., Zhu, Q., Zhang, J., Wu, N., and He, Y.: Dominant influence of non-thawing periods on annual CO<sub>2</sub> emissions from Zoige peatlands: Five-year eddy covariance analysis, *Ecological Indicators*, 129, 107 913, 2021.
- 605 Liu, X., Chen, H., Zhu, Q., Wu, J., Frolking, S., Zhu, D., Wang, M., Wu, N., Peng, C., and He, Y.: Holocene peatland development and carbon stock of Zoige peatlands, Tibetan Plateau: a modeling approach, *Journal of Soils and Sediments*, 18, 2032–2043, <https://doi.org/10.1007/s11368-018-1960-0>, 2018.
- Liu, X., Zhu, D., Zhan, W., Chen, H., Zhu, Q., Hao, Y., Liu, W., and He, Y.: Five-Year Measurements of Net Ecosystem CO<sub>2</sub> Exchange at a Fen in the Zoige Peatlands on the Qinghai-Tibetan Plateau, *Journal of Geophysical Research: Atmospheres*, 124, 11 803–11 818, <https://doi.org/10.1029/2019JD031429>, 2019b.
- 610

- Lu, M., Zhou, X., Yang, Q., Li, H., Luo, Y., Fang, C., Chen, J., Yang, X., and Li, B.: Responses of ecosystem carbon cycle to experimental warming: a meta-analysis, *Ecology*, 94, 726–738, 2013.
- Lupascu, M., Welker, J., Xu, X., and Czimczik, C.: Rates and radiocarbon content of summer ecosystem respiration in response to long-term deeper snow in the High Arctic of NW Greenland, *Journal of Geophysical Research: Biogeosciences*, 119, 1180–1194, 2014.
- 615 Ma, K., Zhang, Y., Tang, S., and Liu, J.: Spatial distribution of soil organic carbon in the Zoige alpine wetland, northeastern Qinghai-Tibet Plateau, *Catena*, 144, 102–108, <https://doi.org/10.1016/j.catena.2016.05.014>, 2016.
- Manzoni, S., Katul, G. G., and Porporato, A.: Analysis of soil carbon transit times and age distributions using network theories, *Journal of Geophysical Research: Biogeosciences*, 114, 2009.
- McGuire, A., Sitch, S., Clein, J. S., Dargaville, R., Esser, G., Foley, J., Heimann, M., Joos, F., Kaplan, J., Kicklighter, D., et al.: Carbon  
620 balance of the terrestrial biosphere in the twentieth century: Analyses of CO<sub>2</sub>, climate and land use effects with four process-based ecosystem models, *Global biogeochemical cycles*, 15, 183–206, 2001.
- Mesfin, S., Gebresamuel, G., Haile, M., and Zenebe, A.: Modelling spatial and temporal soil organic carbon dynamics under climate and land management change scenarios, northern Ethiopia, *European Journal of Soil Science*, 72, 1298–1311, 2021.
- Metzler, H. and Sierra, C. A.: Linear Autonomous Compartmental Models as Continuous-Time Markov Chains: Transit-Time and Age  
625 Distributions, *Mathematical Geosciences*, 50, 1–34, <https://doi.org/10.1007/s11004-017-9690-1>, 2018.
- Moyano, F. E., Manzoni, S., and Chenu, C.: Responses of soil heterotrophic respiration to moisture availability: An exploration of processes and models, *Soil Biology and Biochemistry*, 59, 72–85, 2013.
- Nieberding, F., Wille, C., Fratini, G., Asmussen, M. O., Wang, Y., Ma, Y., and Sachs, T.: A long-term (2005–2019) eddy covariance data set of CO<sub>2</sub> and H<sub>2</sub>O fluxes from the Tibetan alpine steppe, *Earth System Science Data*, 12, 2505–2524, <https://doi.org/10.5194/essd-12-2705-2020>, 2020.  
630
- Pan, Y., Li, X., Li, S., and Li, Z.: Different responses of soil respiration to climate change in permafrost and non-permafrost regions of the Tibetan plateau from 1979 to 2018, *International Journal of Climatology*, 2022.
- Pegoraro, E. F., Mauritz, M. E., Ogle, K., Ebert, C. H., and Schuur, E. A.: Lower soil moisture and deep soil temperatures in thermokarst features increase old soil carbon loss after 10 years of experimental permafrost warming, *Global change biology*, 27, 1293–1308, 2021.
- 635 Piao, S., Tan, K., Nan, H., Ciais, P., Fang, J., Wang, T., Vuichard, N., and Zhu, B.: Impacts of climate and CO<sub>2</sub> changes on the vegetation growth and carbon balance of Qinghai–Tibetan grasslands over the past five decades, *Global and Planetary Change*, 98, 73–80, 2012.
- Pold, G., Melillo, J. M., and DeAngelis, K. M.: Two decades of warming increases diversity of a potentially lignolytic bacterial community, *Frontiers in microbiology*, 6, 480, 2015.
- Rustad, L., Campbell, J., Marion, G., Norby, R., Mitchell, M., Hartley, A., Cornelissen, J., and Gurevitch, J.: A meta-analysis of the response  
640 of soil respiration, net nitrogen mineralization, and aboveground plant growth to experimental ecosystem warming, *Oecologia*, 126, 543–562, 2001.
- Schimel, D.: Terrestrial ecosystems and the carbon-cycle, *Global Change Biology*, 1, 77–91, <https://doi.org/10.1111/j.1365-2486.1995.tb00008.x>, 1995.
- Schmidt, M. W., Torn, M. S., Abiven, S., Dittmar, T., Guggenberger, G., Janssens, I. A., Kleber, M., Kögel-Knabner, I., Lehmann, J.,  
645 Manning, D. A., et al.: Persistence of soil organic matter as an ecosystem property, *Nature*, 478, 49–56, 2011.
- Schuur, E. A. G., Vogel, J. G., Crummer, K. G., Lee, H., Sickman, J. O., and Osterkamp, T. E.: The effect of permafrost thaw on old carbon release and net carbon exchange from tundra, *Nature*, 459, 556–559, <https://doi.org/10.1038/nature08031>, 2009.

- Schuur, E. A. G., Druffel, E., and Trumbore, S. E.: Radiocarbon and Climate Change: Mechanisms, Applications and Laboratory Techniques, pp. 1–315, Springer International, <https://doi.org/10.1007/978-3-319-25643-6>, 2016.
- 650 Scurluck, J. and Hall, D.: The global carbon sink: a grassland perspective, *Global Change Biology*, 4, 229–233, 1998.
- Shaw, R. G. and Mitchell-Olds, T.: ANOVA for unbalanced data: an overview, *Ecology*, 74, 1638–1645, 1993.
- Sierra, C. A. and Mueller, M.: A general mathematical framework for representing soil organic matter dynamics, *Ecological Monographs*, 85, 505–524, <https://doi.org/10.1890/15-0361.1>, 2015.
- Sierra, C. A., Trumbore, S. E., Davidson, E. A., Vicca, S., and Janssens, I.: Sensitivity of decomposition rates of soil organic matter with  
655 respect to simultaneous changes in temperature and moisture, *Journal of Advances in Modeling Earth Systems*, 7, 335–356, 2015.
- Sierra, C. A., Malghani, S., and Loescher, H. W.: Interactions among temperature, moisture, and oxygen concentrations in controlling decomposition rates in a boreal forest soil, *Biogeosciences*, 14, 703–710, 2017a.
- Sierra, C. A., Müller, M., Metzler, H., Manzoni, S., and Trumbore, S. E.: The muddle of ages, turnover, transit, and residence times in the carbon cycle, *Global Change Biology*, 23, 1763–1773, 2017b.
- 660 Sierra, C. A., Mueller, M., and Trumbore, S. E.: Models of soil organic matter decomposition: the SoilR package, version 1.0, *Geoscientific Model Development*, 5, 1045–1060, <https://doi.org/10.5194/gmd-5-1045-2012>, 2012.
- Sierra, C. A., Mueller, M., and Trumbore, S. E.: Modeling radiocarbon dynamics in soils: SOILR version 1.1, *Geoscientific Model Development*, 7, 1919–1931, <https://doi.org/10.5194/gmd-7-1919-2014>, 2014.
- Sierra, C. A., Ceballos-Nunez, V., Metzler, H., and Mueller, M.: Representing and Understanding the Carbon Cycle Using the Theory of Com-  
665 partmental Dynamical Systems, *Journal of Advances in Modeling Earth Systems*, 10, 1729–1734, <https://doi.org/10.1029/2018MS001360>, 2018a.
- Sierra, C. A., Hoyt, A. M., He, Y., and Trumbore, S. E.: Soil Organic Matter Persistence as a Stochastic Process: Age and Transit Time Distributions of Carbon in Soils, *Global Biogeochemical Cycles*, 32, 1574–1588, <https://doi.org/10.1029/2018GB005950>, 2018b.
- Steinhof, A., Altenburg, M., and Mächts, H.: Sample preparation at the Jena C-14 laboratory, *Radiocarbon*, 59, 815–830,  
670 <https://doi.org/10.1017/RDC.2017.50>, 2017.
- Stuiver, M. and Polach, H.: Reporting of C-14 data - Discussion, *Radiocarbon*, 19, 355–363, <https://doi.org/10.1017/S0033822200003672>, 1977.
- Sun, X., Zhao, Y., and Li, Q.: Holocene peatland development and vegetation changes in the Zoige Basin, eastern Tibetan Plateau, *Science China Earth Sciences*, 60, 1826–1837, 2017.
- 675 Tangarife-Escobar, A., Guggenberger, G., Feng, X., Dai, G., Urbina-Malo, C., Azizi-Rad, M., and Sierra, C.: Moisture and temperature effects on the radiocarbon signature of respired carbon dioxide to assess stability of soil carbon in the Tibetan Plateau, <https://doi.org/10.5281/zenodo.10537332>, 2024.
- Tian, Y.-Q., Xu, X.-L., Song, M.-H., Zhou, C.-P., Gao, Q., and Ouyang, H.: Carbon sequestration in two alpine soils on the Tibetan Plateau, *Journal of integrative plant biology*, 51, 900–905, 2009.
- 680 Trumbore, S.: Age of soil organic matter and soil respiration: Radiocarbon constraints on belowground C dynamics, *Ecological Applications*, 10, 399–411, [https://doi.org/10.1890/1051-0761\(2000\)010\[0399:AOSOMA\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2000)010[0399:AOSOMA]2.0.CO;2), 2000.
- Trumbore, S., Chadwick, O., and Amundson, R.: Rapid exchange between soil carbon and atmospheric carbon dioxide driven by temperature change, *Science*, 272, 393–396, <https://doi.org/10.1126/science.272.5260.393>, 1996.

- von Lützow, M., Kögel-Knabner, I., Ludwig, B., Matzner, E., Flessa, H., Ekschmitt, K., Guggenberger, G., Marschner, B., and Kalbitz, K.:  
685 Stabilization mechanisms of organic matter in four temperate soils: Development and application of a conceptual model, *Journal of Plant  
Nutrition and Soil Science*, 171, 111–124, 2008.
- Wang, M., Yang, G., Gao, Y., Chen, H., Wu, N., Peng, C., Zhu, Q., Zhu, D., Wu, J., He, Y., Tian, J., Zhao, X., and Zhang, Y.:  
Higher recent peat C accumulation than that during the Holocene on the Zoige Plateau, *Quaternary Science Reviews*, 114, 116–125,  
<https://doi.org/10.1016/j.quascirev.2015.01.025>, 2015.
- 690 Wieder, W. R., Bonan, G. B., and Allison, S. D.: Global soil carbon projections are improved by modelling microbial processes, *Nature  
climate change*, 3, 909–912, 2013.
- Xiang, S., Guo, R., Wu, N., and Sun, S.: Current status and future prospects of Zoige Marsh in Eastern Qinghai-Tibet Plateau, *Ecological  
Engineering*, 35, 553–562, <https://doi.org/10.1016/j.ecoleng.2008.02.016>, 2009.
- Xu, X., Shi, Z., Chen, X., Lin, Y., Niu, S., Jiang, L., Luo, R., and Luo, Y.: Unchanged carbon balance driven by equivalent responses of  
695 production and respiration to climate change in a mixed-grass prairie, *Global Change Biology*, 22, 1857–1866, 2016.
- Yang, G., Chen, H., Wu, N., Tian, J., Peng, C., Zhu, Q., Zhu, D., He, Y., Zheng, Q., and Zhang, C.: Effects of soil warming, rain-  
fall reduction and water table level on CH<sub>4</sub> emissions from the Zoige peatland in China, *Soil Biology and Biochemistry*, 78, 83–89,  
<https://doi.org/10.1016/j.soilbio.2014.07.013>, 2014.
- Yang, Z., Zhu, D., Liu, L., Liu, X., and Chen, H.: The effects of freeze-thaw cycles on methane emissions from peat soils of a high altitude  
700 peatland, *Frontiers in Earth Science*, p. 452, 2022.
- Yao, T., Xue, Y., Chen, D., Chen, F., Thompson, L., Cui, P., Koike, T., Lau, W. K.-M., Lettenmaier, D., Mosbrugger, V., Zhang, R., Xu,  
B., Dozier, J., Gillespie, T., Gu, Y., Kang, S., Piao, S., Sugimoto, S., Ueno, K., Wang, L., Wang, W., Zhang, F., Sheng, Y., Guo, W.,  
Ailikun, Yang, X., Ma, Y., Shen, S. S. P., Su, Z., Chen, F., Liang, S., Liu, Y., Singh, V. P., Yang, K., Yang, D., Zhao, X., Qian, Y., Zhang,  
Y., and Li, Q.: Recent Third Pole’s Rapid Warming Accompanies Cryospheric Melt and Water Cycle Intensification and Interactions  
705 between Monsoon and Environment: Multidisciplinary Approach with Observations, Modeling, and Analysis, *Bulletin of the American  
Meteorological Society*, 100, 423–444, <https://doi.org/10.1175/BAMS-D-17-0057.1>, 2019.
- Zhang, G., Zhang, Y., Dong, J., and Xiao, X.: Green-up dates in the Tibetan Plateau have continuously advanced from 1982 to 2011, *Proceed-  
ings of the National Academy of Sciences of the United States of America*, 110, 4309–4314, <https://doi.org/10.1073/pnas.1210423110>,  
2013.
- 710 Zhao, Y., Yu, Z., and Zhao, W.: Holocene vegetation and climate histories in the eastern Tibetan Plateau: controls  
by insolation-driven temperature or monsoon-derived precipitation changes?, *Quaternary Science Reviews*, 30, 1173–1184,  
<https://doi.org/10.1016/j.quascirev.2011.02.006>, 2011.
- Zhou, W., Cui, L., Wang, Y., Li, W., and Kang, X.: Carbon emission flux and storage in the degraded peatlands of the Zoige alpine area in  
the Qinghai–Tibetan Plateau, *Soil Use and Management*, 37, 72–82, 2021.