



Moisture and temperature effects on the radiocarbon signature of respired carbon dioxide to assess stability of soil carbon in the Tibetan Plateau

Andrés Tangarife-Escobar¹, Georg Guggenberger², Xiaojuan Feng³, Guohua Dai³, Carolina Urbina-Malo², Mina Azizi-Rad¹, and Carlos A. Sierra^{1,4}

¹Max Planck Institute for Biogeochemistry, Jena, Germany

²Institute of Soil Science, Leibniz Universität Hannover, Hannover, Germany

³Institute of Botany, Chinese Academy of Sciences, Beijing, China

⁴Department of Ecology, Swedish University of Agricultural Sciences, Uppsala, Sweden

Correspondence: Andrés Tangarife-Escobar (atanga@bgc-jena.mpg.de)

Abstract. Microbial release of CO₂ from soils to the atmosphere reflects how environmental conditions affect the stability of soil organic matter (SOM), especially in massive organic-rich ecosystems like the peatlands and grasslands of the Qinghai-Tibetan Plateau (QTP). Radiocarbon (¹⁴C) is an important tracer of the global carbon cycle and can be used to understand SOM dynamics through the estimation of time lags between C fixation and respiration, often assessed with metrics such as age and transit time. In this study, we incubated peatland and grassland soils at four temperature (5, 10, 15 and 20 °C) and two water-filled pore space (WFPS) levels (60 and 95 %), and measured the ¹⁴C signature of bulk soil and respired CO₂. We compare the relation between the $\Delta^{14}\text{C}$ of the bulk soil and the $\Delta^{14}\text{CO}_2$ of respired carbon as a function of temperature and WFPS for the two soils. To better interpret our results, we used a mathematical model to analyse how the calculated number of pools, decomposition rates of carbon (k), transfer (α) and partitioning (γ) coefficients affect the $\Delta^{14}\text{C}$ -bulk and $\Delta^{14}\text{CO}_2$ relation, with their respective mean age and mean transit time. From our incubations, we found that ¹⁴C from peatland was significantly more depleted (old) than from grassland soil. Our results showed that changes in temperature did not affect the $\Delta^{14}\text{C}$ values of respired CO₂ in either soil. However, changes in WFPS had a small effect on the ¹⁴CO₂ in grassland soils and a strong influence in peatland soils, where higher WFPS levels led to more depleted $\Delta^{14}\text{CO}_2$. In our models, we observed large differences between slow and fast cycling systems, where low values of k modified $\Delta^{14}\text{C}$ patterns due to the incorporation of ¹⁴C-bomb in the soil. Hence, the correspondence between $\Delta^{14}\text{C}$ and age and transit time strongly depended on the internal dynamics of the soil (k , α , γ and number of pools) as well as on model structure. We conclude that the stability of carbon in these systems depends strongly on the direction of change in temperature and moisture and how it affects the rates of SOM decomposition. Finally, $\Delta^{14}\text{C}$ modelling along with empirical data from SOM dynamics is a useful tool to improve predictions on interactions between terrestrial and atmospheric carbon.



20 1 Introduction

Studying soil organic matter (SOM) stability and persistence in a globally-changing environment is of fundamental importance to understand temporal variations of carbon cycling in the earth-climate system. Soil constitutes the largest carbon (C) stock in the terrestrial biosphere (Chen et al., 2021), making it a key component in global climate models (McGuire et al., 2001; Wieder et al., 2013; Xu et al., 2016). Physical, chemical and biological properties determine SOM decomposition and in consequence its persistence over decades to millennia (Schmidt et al., 2011). Temperature and moisture are two of the most important abiotic variables controlling the rates of SOM decomposition (Sierra et al., 2015) and its transit time across the different ecosystem pools. Changes in these environmental conditions are, however, occurring simultaneously, highlighting the necessity to conduct multifactorial experiments to disentangle the dominant mechanisms on such cycling times. Hence, discerning the relationship between SOM persistence and C cycling times in terrestrial ecosystems is imperative to improve climate change models and to inform decisions on CO₂ mitigation and management strategies (Bradford et al., 2016; Mesfin et al., 2021).

The Qinghai-Tibetan Plateau (QTP) with an area of 2.5 million km² is a macroregion of global importance for the cycling of water, carbon and other biogeochemical elements (Anslan et al., 2020). Land cover is mainly dominated by alpine grasslands (44 %) (Scurlock and Hall, 1998; Gao et al., 2014) which store 23.4 % of China's total organic carbon and 2.5 % of the global soil carbon (Genxu et al., 2002). The QTP also hosts one of the largest high-mountain marshes in the world (Xiang et al., 2009; Chen et al., 2014; Ma et al., 2016), the Zoige peatlands, considered to be the most important carbon stock in peatlands for China (Liu et al., 2018). However, climate change and land cover change are currently driving the net carbon balance through CO₂ and CH₄ effluxes from the grasslands (Piao et al., 2012; Chen et al., 2014, 2017; Du and Gao, 2020) and the peatlands at both daily and interannual timescales (Hao et al., 2011; Kang et al., 2018; Liu et al., 2019b; Chen et al., 2021). Anthropogenic activities in the Zoige peatlands (drainage, peat mining and overgrazing) have caused a degradation of approximately 30 % of wet and dry meadows (Chen et al., 2014; Zhou et al., 2021). Additionally, the QTP has been facing an air temperature increase of 0.2°C per decade over the past 50 years (Zhang et al., 2013; Chen et al., 2014; Yang et al., 2014; Ganjurjav et al., 2016), equivalent to two to three times faster than the world average (Yao et al., 2019; Nieberding et al., 2020), along with a moderate increase in precipitation (Dong et al., 2018). According to the IPCC (Arias et al., 2021), climate models predict an increment in heavy precipitation events and high temperature extremes for this region.

Such changes in temperature and soil moisture control the magnitude in which stabilization and de-stabilization mechanisms enable carbon storage in the QTP (Xiang et al., 2009; Ma et al., 2016). It has been observed that warming increases soil respiration (Rustad et al., 2001; Lu et al., 2013; Pold et al., 2015) and that soil moisture modulates ecosystem and soil respiration (Geng et al., 2012; Piao et al., 2012; Pan et al., 2022; Azizi-Rad et al., 2022). Although the influence of destabilization mechanisms on SOM decomposition has been already studied in the Zoige peatlands (Zhao et al., 2011; Wang et al., 2015; Liu et al., 2016, 2018; Li et al., 2018; Liu et al., 2019a), temporal scales of SOM persistence under temperature and soil moisture variations are not clearly understood yet. Therefore, a quantification of the SOM persistence is of vital importance to predict



climate change feedback magnitudes and pathways.

55

A useful approach to quantify SOM persistence is through the theory of compartmental dynamical systems, where soil is understood as a set of interconnected pools with transformations of C to different forms (gas, dissolved or solid) (Emanuel et al., 1984; Schimel, 1995; Sierra et al., 2018a). Therefore, C can be characterized by the time it remains inside a compartment or the entire system (Eriksson, 1971; Bolin and Rodhe, 1973) through the calculation of age and transit time. Here, age is defined as the time elapsed since the C entered the system until the time of observation in the bulk soil, while transit time is defined as the time spent by the C between the entry to the system and its exit as respired CO₂ (Eriksson, 1971; Bolin and Rodhe, 1973; Manzoni et al., 2009; Sierra et al., 2018b). These times (*T*) are however, not the same for all the C atoms in a soil, since physical, chemical and biological stabilization and destabilization processes of SOM modify the rate at which mixing and storage occurs.

60

65

Radiocarbon measurements in bulk soil and in respired CO₂ are a powerful tool to approximate ages and transit times of C in soils as they trace back the trajectory of C through the different stocks on decadal to millennia timescales (Trumbore, 2000; Sierra et al., 2014; Schuur et al., 2016; Estop-Aragones et al., 2020; Chen et al., 2021). By comparing the ¹⁴C relative abundance in the carbon pools with the atmospheric ¹⁴CO₂ concentrations produced after the nuclear weapon tests during the early 1960's, it is possible to model soil-atmosphere C cycling times (Trumbore, 2000).

70

¹⁴C values can be altered by SOM (de)-stabilization processes and soil characteristics such as soil organic carbon (SOC) content, age and diagenetic state, thaw depth, redox state, seasonality, etc. (Gaudinski et al., 2000; Trumbore, 2000; von Lützow et al., 2008; Sierra et al., 2018b; Estop-Aragones et al., 2020). In recent decades, it has been widely observed that temperature plays a major role on the dynamics of SOM (Knorr et al., 2005; Davidson and Janssens, 2006; Feng and Simpson, 2008) and the mean age of respired CO₂ (Hopkins et al., 2012; Chen et al., 2021) by increasing the decomposition rates from fast-cycling pools (Trumbore et al., 1996) and mobilizing old C in warming conditions (Dutta et al., 2006; Briones et al., 2010). In contrast, other studies have suggested that warming does not lead to release of old C (Briones et al., 2021; Dioumaeva et al., 2002) and that non-labile C decomposition is insensitive to temperature increase (Liski et al., 1999). By comparison, drying phenomena increased the release of modern ¹⁴CO₂ from shallow soil layers but preserved the old soil carbon pools in deeper layers (Kwon et al., 2019). Observations in rich-SOC soils of the Arctic indicated the release of old C from deep soil layers after thaw and drainage processes (Schuur et al., 2009; Estop-Aragones et al., 2020). Such a process might be occurring in the Zoige peatland soils due to the seasonal presence of frozen layers (Liu et al., 2021; Yang et al., 2022). Nonetheless, the influence of temperature and soil moisture on SOM persistence is a matter of current debate.

80

85

Changes in temperature and moisture contribute to the destabilization of carbon in soils from the QTP. Hence, we hypothesize that higher temperature and moisture levels would contribute to increasing the age of respired CO₂ for soils subjected to controlled manipulations. Greater ages of respired CO₂ would imply that previously stabilized C is destabilized by changes in the manipulated environmental factors. For this purpose, we posed three specific research questions: (1) Do specific changes



in temperature and moisture result in the release of old carbon in the respiration flux? (2) Are there differences in the age of
90 respired CO₂ between a grassland and a peatland soil from the QTP? (3) How can radiocarbon data in bulk soil and respired
CO₂ be interpreted to understand the effect of changes in decomposition rates on ages and transit times? To address these ques-
tions, we conducted a controlled multifactorial experiment with soils from two different ecosystems in the QTP, and measured
their radiocarbon signature in the bulk soil and the respired CO₂. In addition, we used a mathematical model to better interpret
our results with the concepts of age and transit time of carbon.

95

2 Materials and methods

To evaluate the influence of soil moisture and temperature on the ¹⁴C values from bulk soil and respired CO₂ of grasslands and
peatlands soils, we incubated soil samples from a high elevation grassland at the Nam Co catchment collected in September
2018 and a peatland from the Zoige region sampled in July 2021, both located in the QTP.

100 2.1 Site description and sampling

2.1.1 Nam Co grassland

The Nam Co site (Fig. 1) is located in the central Tibetan Plateau (altitude of the Nam Co lake 4,726 m a.s.l.) and represents
a frequent study location for monitoring and tracking of environmental changes over different time scales (Anslan et al.,
2020). The dominant climate at Nam Co consists of cold winters and short and moist summers. Mean annual temperature
105 (MAT) measured at the NAMORS research station was -0.6 °C (2006 – 2017) and mean annual precipitation (MAP) was 406
mm, occurring mostly during the monsoon season between May and October (Anslan et al., 2020). The sampling area was
mainly covered by *Kobresia pygmaea* and has been grazed intensively by yaks and sheep. A total amount of 40 kg of soil was
collected from randomly selected spots at depths between 5 and 15 cm within an area of about 40 m². The soil was then mixed
homogeneously to produce one single sample which consisted of a sandy loam with a bulk density of 1.3 g/cm³, pH of 7.5, 3.2
110 % of total organic carbon, 0.05 % of inorganic carbon and 0.3 % of total nitrogen, cation exchange capacity of 89 mmol_c/kg
soil and electrical conductivity of 89.6 μS/cm.




Figure 1. a) Overview of the QTP; b) Location of the Nam Co lake with land cover dominated by grasslands, and c) Zoigê peatlands highlighting the location of Zoigê city. The sampling site is located in the southeastern of the peatland (Ruokeba, Hongyuan county). SRTM elevation data from Jarvis et al. (2008).

2.1.2 Zoigê peatlands

The Zoigê peatlands are located in the northeast of the QTP (average elevation of 3,400 m a.s.l.) and cover an area of about 4,605 km² on the headwaters of the Yellow River basin. These peatlands have been recognised as one of the largest high-
115 mountain marshes in the world (Xiang et al., 2009; Ma et al., 2016) and store 88 % of the carbon in the QTP (Chen et al., 2014). The Zoigê region shows a MAT of 1.5 °C, where the warmest monthly temperature (11 °C) is recorded in July and the coldest in January (-10.1 °C). The MAP is 720 mm, and according to data from the Chinese National Meteorological Information Centre (www.nmic.gov.cn), the region has faced a slight drying trend and a warming of 0.4 °C per decade over
120 and 50 cm depth is seasonally frozen between November and March. Vegetation cover consists mainly of *Potentilla anserine*, *Blysmus sinocompressus*, *Kobresia myosuroides* and *Scirpus triqueter*. The samples (0 – 35 cm depth) were collected from an



area of 50 m² at a long-term monitoring site (33°4'5''N, 102°33'52''E), then thoroughly mixed to produce a single sample. The soil bulk density was 0.3 g/cm³, contained in average 27.6 % of organic carbon, 0.06 % of inorganic carbon and 1.8 % of total nitrogen.

125 2.2 Incubation of peatland and grassland soils

We conducted two sets of incubation experiments, one set with the grassland soil and a second set with the peatland soil. Samples were incubated with their original roots to minimise disturbance and allow comparisons with field conditions, however stones were removed. Each of the sets were placed at two different WFPS values (60 and 95 %) combined with four temperature levels (5, 10, 15 and 20 °C) for a total of 69 samples (2 sites × 4 temperature levels × 2 moisture levels × 4-5 replicates, 130 to prevent scarcity of data due to eventual failure in CO₂ extraction), totaling 33 for grassland and 36 for peatland. CO₂ concentrations were measured at intervals of 1 to 2 weeks using a CO₂ analyzer LI-COR 6262. Incubations for each subset ended simultaneously until every sample had an estimated concentration of CO₂-C in the headspace equivalent to ≥ 0.2 mg of C, enough for radiocarbon analysis. Grassland samples were incubated between 15 and 67 days, while peatland samples were 135 incubated for 13 days. For sampling headspace air, 50-ml vials were filled with 12 g of soil (± 1.5 g) and placed inside 0.5 L glass flasks along with 0.2 ml of water at the bottom of the flask (away from contact with the sample) to avoid possible drying (Dioumaeva et al., 2002); thereafter the flasks were sealed with rubber plugs and screwed with plastic caps. Flasks with samples were flushed with synthetic air (CO₂ free) to remove atmospheric CO₂. This flushing marked the starting day of the incubations.

2.3 Radiocarbon analysis of incubated soils

140 Respired CO₂ accumulated in the headspace of incubation flasks was extracted and purified on a vacuum line, graphitized by Fe reduction in H₂ and measured for Δ¹⁴C by Accelerator Mass Spectrometry (Micadas, Ionplus, Switzerland) in the radiocarbon laboratory of the Max Planck Institute for Biogeochemistry in Jena, Germany (Steinhof et al., 2017).

The radiocarbon content reflects the time a C atom has been in the soil since it was fixed by photosynthesis from the 145 atmosphere (Trumbore, 2000). Carbon fixed since the early 1960's has higher ¹⁴C than carbon fixed previously due to the enrichment by thermonuclear weapons testing. Samples with significant ¹⁴C positive values show incorporation by the bomb ¹⁴C, presenting ¹⁴C/¹²C ratios greater than that of the 1890 wood used as ¹⁴C standard for pre-bomb periods. Radiocarbon data are expressed as Δ¹⁴C (the deviation in ‰ from 0.95 times the oxalic acid standard in 1950). The values were corrected to a δ¹³C value of -25 ‰ for differences in biological mass-dependent fractionation (Stuiver and Polach, 1977). These data are 150 presented as percentage of Modern Carbon (pMC) that can be converted to F¹⁴C by dividing it by 100 and later to Δ¹⁴C using equation (1) (Stuiver and Polach, 1977).



$$\Delta^{14}C = \left[F^{14}C e^{\lambda_C(1950-t)} - 1 \right] \times 1000 [\text{‰}], \quad (1)$$

where $F^{14}C$ is the Fraction Modern, i.e. the ratio of the measured sample normalized to a $\delta^{13}C$ value of -25‰ , divided by
155 0.95 times the measured ratio of the Oxalic Acid I standard (OX-I) (Schuur et al., 2016), λ_C is the updated radiocarbon decay
constant (equals $1/8267 \text{ [y}^{-1}\text{]}$), and t is the year of sampling.

2.4 SOC decomposition models to predict $\Delta^{14}C$ as a proxy of SOM persistence

The representation of the SOC dynamics has been commonly described through models that can be expressed as systems of
160 linear differential equations (Manzoni et al., 2009; Sierra et al., 2012; Sierra and Mueller, 2015; Sierra et al., 2017) of the form:

$$\frac{d\mathbf{C}(t)}{dt} = \mathbf{I} + \mathbf{A} \cdot \mathbf{C}(t), \quad (2)$$

where the vector $\mathbf{C}(t)$ is the rate of change of carbon over time in n different pools; the time dependent n -dimensional vector
 \mathbf{I} represents the total inputs of carbon to each pool; and \mathbf{A} represents the $n \times n$ dimensional matrix with the rates of car-
bon processing for each pool in its main diagonal and the proportion of carbon transferred from one pool to another in the
165 off-diagonals (Sierra et al., 2012, 2014; Metzler and Sierra, 2018; Sierra et al., 2018b). This mass balance equation has a
radiocarbon counterpart:

$$\frac{d^{14}\mathbf{C}(t)}{dt} = \mathbf{I}_{14}\mathbf{C}(t) + \mathbf{A} \cdot {}^{14}\mathbf{C}(t) - \lambda {}^{14}\mathbf{C}(t) \quad (3)$$

where λ is the radiocarbon decay constant ($1/8267 \text{ [y}^{-1}\text{]}$).

170 We used a SOC decomposition model to calculate the $\Delta^{14}C$ in bulk and CO_2 as well as their equivalent mean age and mean
transit time of a theoretical soil for the year 2019 calibrated with the data set for the Northern Hemisphere Zone 3 (Hua et al.,
2021). The simulated $\Delta^{14}C$ results aimed at reproducing the similar $\Delta^{14}C$ ranges obtained from the incubated soils with the
objective to compare under which settings the models could describe best laboratory or field conditions. For this purpose, we
implemented a two-pool model (one slow and one fast pool) considering two different structures: parallel and series (Fig. 2)
175 and modified the parameters involved in the soil C processing (Manzoni et al., 2009; Falloon and Smith, 2000) (Table 1). In
the parallel structure, C enters the soil and splits among the two pools according to γ and decomposition occurs independently
in each pool according to their respective k . In the series structure, C enters only to the fast pool and it is either decomposed
and emitted to the atmosphere or transferred to the slow pool according to the transfer coefficient α .



$$\frac{d\mathbf{C}}{dt} = I \begin{pmatrix} \gamma \\ 1-\gamma \end{pmatrix} + \begin{pmatrix} -k_1 & 0 \\ 0 & -k_2 \end{pmatrix} \cdot \begin{pmatrix} C_1 \\ C_2 \end{pmatrix}$$

$$\frac{d\mathbf{C}}{dt} = I \begin{pmatrix} 1 \\ 0 \end{pmatrix} + \begin{pmatrix} -k_1 & 0 \\ \alpha_2 k_1 & -k_2 \end{pmatrix} \cdot \begin{pmatrix} C_1 \\ C_2 \end{pmatrix}$$





Figure 2. Parallel (left) and series (right) model structure and formulas implemented in the simulations. C is fixed into the biomass from atmospheric CO_2 through photosynthesis, subsequently it is incorporated into the soil as litterfall. From this state on, it is considered as input (I) and will split between the two pools according to the different structures (γ for parallel and α for series). The rate at which the C is decomposed (k) in each pool (C) will depend on microbial activity, environmental factors (temperature and soil moisture) and physical and chemical protection of the SOM (Blanco-Canqui and Lal, 2004; von Lütow et al., 2008; Manzoni et al., 2009). Boxes represent the soil pools, which account with an initial amount of C ($C_1 = 200$, $C_2 = 5000$) and arrows represent directions of inputs and outputs.

180 Initial parameters of the model such as the initial year of simulation and initial $\Delta^{14}\text{C}$ values of each pool were considered separately for each type of ecosystem. Moreover, we defined I and C as constant since they can be adjusted depending on the specific soil characteristics. Additionally, we assumed that decomposition rates k reflect the effect of temperature and soil moisture on C cycling timescales (Manzoni et al., 2009), and therefore on $\Delta^{14}\text{C}$ values. In our approach, only one parameter at time was modified for each simulation.

185



Table 1. Definition of parameters used to evaluate the variation of $\Delta^{14}\text{C}$ values, mean age and mean transit time and their ranges when type of constant was variable.

Parameter	Notation	Type	Value	Definition
Litter Input	I	Constant	100	A scalar or a data.frame object specifying the amount of litter inputs by time
Decomposition rate	k_1, k_2	Variable	0.00001 - 0.8	A vector of length 2 with the values of the decomposition rate for pools 1 and 2
Partitioning coefficient	γ	Variable	0 - 1	A scalar representing the proportion of I that goes to pool 1 in a parallel structure
Transference coefficient	α	Variable	0 - 1	A scalar with the value of the transfer rate from pool 1 to pool 2
Carbon stocks	C_1, C_2	Constant	200, 5000	Initial amount of C for the two pools

We used the R package SoilR (Sierra et al., 2014) to simulate the temporal dynamics of $\Delta^{14}\text{C}$ in the bulk soil and the respired CO_2 as well as the age and transit time distributions of carbon for two different model structures (parallel and series). Assuming steady-state for the carbon stocks, the probability density function (pdf) of the age (equation 4) and the transit time (equation 5) (Metzler and Sierra, 2018) as well as their means can be calculated by the following expressions (Sierra et al., 2018b):

$$f(a) = -\mathbf{1}^T \cdot \mathbf{A} \cdot e^{a \cdot \mathbf{A}} \cdot \frac{\mathbf{C}^*}{\sum \mathbf{C}^*}, \quad a \geq 0, \quad (4)$$

where a is the random variable age, $\mathbf{1}^T$ is the transpose of the n -dimensional vector containing ones, $e^{a \cdot \mathbf{A}}$ is the matrix exponential for each value of a , and $\sum \mathbf{C}^*$ is the sum of stocks of all pools at steady state.

The pdf of transit times can be obtained as

$$f(\tau) = -\mathbf{1}^T \cdot \mathbf{A} \cdot e^{\tau \cdot \mathbf{A}} \cdot \frac{\mathbf{I}}{\sum \mathbf{I}}, \quad \tau \geq 0, \quad (5)$$

where τ represents the random variable transit time.

These pdfs measure the probability that a certain amount of carbon is above or below a specific age or transit time (Sierra et al., 2018b) (equations 6 and 7, respectively). The mean of the age, that is, the expected value of the pdf can be computed by the expression:

$$\mathbb{E}(a) = -\mathbf{1}^T \cdot \mathbf{A}^{-1} \cdot \frac{\mathbf{C}^*}{\sum \mathbf{C}^*}, \quad (6)$$



and the mean of the transit time by

$$\mathbb{E}(\tau) = -\mathbf{1}^T \cdot \mathbf{A}^{-1} \cdot \frac{\mathbf{I}}{\sum \mathbf{I}}. \quad (7)$$

Based on the age and transit time distribution for the different simulated cases, metrics such as the mean, median and
205 quantiles can indicate the SOM persistence. The relation between ages and transit times may present three cases: $T_a = T_t$ (Type
I - well-mixed homogeneous system), $T_a < T_t$ (Type II - retention system) and $T_a > T_t$ (Type III - non-retention system)
(Sierra et al., 2018b). In the type I, the probability of mineralization and release as CO_2 is the same for every C atom. In the
type II, the C is retained for a relatively long time before it is released, as in soils where C is constantly recycled and reused
sequentially before respiration outside the system. In type III, most of the C atoms stay in the system for a short period of
210 time, but some atoms remain for a long time (Bolin and Rodhe, 1973). Comparisons between these distributions will provide
detailed information on mixing, store, recycling, transport and transformation processes of the SOM (Sierra et al., 2018b).

3 Results

3.1 $\Delta^{14}\text{C}$ values of incubated soils under temperature and soil moisture variation

$\Delta^{14}\text{C}$ values contrasted strongly between grassland and peatland soils (Tangarife-Escobar et al., 2023) (Fig. 3 and 4). Generally,
215 $\Delta^{14}\text{C}$ of bulk soil and respiration from the peatland soil was more depleted than from the grassland soil. For example, in
peatland soil the $\Delta^{14}\text{C}$ values of bulk soil (-85.9 to -60.9 ‰, mean=-80.1, n=36, including outlier of -180) were clearly more
depleted than those of the respired flux (-20.9 to 23.9 ‰, mean=4, n=36) which indicated that the peatland behaved as a
retention system (type II). In contrast, for the grassland soil, the $\Delta^{14}\text{C}$ of bulk soil (21.1 to 73.9 ‰, mean=43.3, n=33) was not
too different from the $\Delta^{14}\text{CO}_2$ (13.9 to 83.4 ‰, mean=38.5, n=33, including outliers of -227.1 and -105.1) indicating that the
220 samples behaved mostly as a well-mixed homogeneous system (type I). Such similar values can also reflect a type II system
where $\Delta^{14}\text{C}$ values in the bulk and the respired flux are equal for the year of sampling (Fig. A7).




Figure 3. Boxplot with the variation of the $\Delta^{14}\text{C}$ values in CO_2 (left panel) and bulk soil (right panel) for peatland and grassland incubated soils.




Figure 4. Relationship between $\Delta^{14}\text{C}$ in bulk soil and $\Delta^{14}\text{C}$ in respired CO_2 for incubated grassland (green box) and peatland (brown box) soils of the QTP discriminated by temperature and WFPS. Possible types of system according to the $\Delta^{14}\text{C}$ relations between bulk and respired carbon.

The temperature treatments did not systematically affect the radiocarbon signature of the bulk or the respired CO_2 in peatland or grassland soils (Table 2). There was strong evidence that manipulations in WFPS resulted in changes in the $\Delta^{14}\text{C}$ values of bulk soil and CO_2 (p -values = 0.09 in grasslands and p -value = 0.01 in peatlands from an ANOVA test, Table 2, Fig. 5) except for the bulk soil in peatlands. When the interacting effects of WFPS and temperature were evaluated together, there was no evidence that their interplay affected the radiocarbon signature of both soils (p -values > 0.05 for all the analysis). However, higher temperature and WFPS caused an increase of CO_2 fluxes from respiration in the treated incubated soils (Table A1). Outliers for bulk and $\Delta^{14}\text{CO}_2$ in both peatlands and grasslands occurred in the combined treatment WFPS = 60 and temperature = 10 °C. Such wide variation in $\Delta^{14}\text{C}$ both between ecosystems and treatments could be potentially explained by intrinsic processes affecting soil carbon dynamics, which will be explored in the following sections with a SOC decomposition model.



Table 2. Summary of p -values obtained from ANOVA tests for the $\Delta^{14}\text{C}$ of bulk soil and the $\Delta^{14}\text{CO}_2$ from grassland and peatland soils versus temperature (T) and WFPS independently as well as versus the integrated effect of temperature and WFPS (T · WFPS).

Ecosystem	Type	T	WFPS	T · WFPS
Grassland	^{14}C Bulk	0.8	0.091	0.21
	$^{14}\text{CO}_2$	0.73	0.089	0.92
Peatland	^{14}C Bulk	0.4	0.21	0.65
	$^{14}\text{CO}_2$	0.16	0.01	0.32

Table 3. Parameters used for simulations in a SOC decomposition model for fast and slow systems (Tangarife-Escobar et al., 2023).

System	Fig.	Model structure	k_1	k_2	γ	α	Starting year of simulation	$\Delta^{14}\text{C}$ (‰) for starting year
Fast cycling	6	Parallel	0.1 - 0.8	0.1	0.8	-	1890	-4.9 ‰
		Series	0.1 - 0.8	0.1	-	0.8	1890	-4.9 ‰
	7	Parallel	0.8	0.001 - 0.8	0.8	-	1890	-4.9 ‰
		Series	0.8	0.001 - 0.8	-	0.8	1890	-4.9 ‰
Slow cycling	8	Parallel	0.0001 - 0.8	0.0001	0.8	-	500	-22.2 ‰
		Series	0.0001 - 0.8	0.0001	-	0.8	500	-22.2 ‰
	9	Parallel	0.1	0.000001 - 0.001	0.2	-	500	-22.2 ‰
		Series	0.1	0.000001 - 0.001	-	0.8	500	-22.2 ‰

3.2 Effect of changes of decomposition rates (k) on the $\Delta^{14}\text{C}$ values

Temperature and soil moisture manipulations showed responses in the vertical and horizontal direction in the $\Delta^{14}\text{C}$ -bulk versus $\Delta^{14}\text{C}$ - CO_2 space. To understand possible drivers of changes in these directions and how they can be interpreted in terms of ages and transit times, we used a SOC decomposition model. We evaluated how model structure, decomposition rates of C, and the partitioning (γ) and transfer (α) coefficients of a two-pool parallel and a series model affected $\Delta^{14}\text{C}$ values, and as a consequence, the mean age and the mean transit time. Simulations were made for the fast cycling grassland system and the slow cycling peatland (Table 3).




Figure 5. Comparison between $\Delta^{14}\text{C}$ of respiration from incubated peatland and grassland soil at different temperature levels under WFPS = 60 and 95 %. Black points represent minimum and maximum values out of the range between quartile 1 and 3 (25 to 75 % of the data). The quartile 50 (median) represented by the line inside the box indicates the midpoint value in the frequency distribution. Box for the treatment WFPS= 60 % and T= 10 °C shows a large dispersion of the 50 % of the data, which is explained by the outliers observed in Fig. 4.

3.2.1 SOC decomposition in fast cycling systems (grasslands)

240 Variation of k_1 yielded $\Delta^{14}\text{C}$ curves similar to mean age and mean transit time. In other words, in these simulations the change
in the parameter k_1 resulted in similar trends in $\Delta^{14}\text{C}$ -bulk versus $\Delta^{14}\text{C}$ -CO₂ space as in the mean age versus mean transit time
space ((Fig. 6). In the parallel structure (Fig. 6, A and B), high values of k_1 yielded more enriched $\Delta^{14}\text{C}$ values in the bulk soil
than in the respired flux. This was expressed as a fast respiration of SOC and reflected in a short transit time. Simultaneously,
low values of k_2 resulted in a slow respiration, which was registered in ages longer than transit times, at the initial state of the C
245 stay in the system. However, as k_1 decreased and approached k_2 , mean age and mean transit time became similar and converged
to the 1:1 line. In the case of the series structure (Fig. 6, C and D), the total amount of inputs traveled first through the fast
pool, which meant that while k_1 is high, most of the C decomposition occurs under the dynamics of the fast pool. Hence, the




Figure 6. Predictions of $\Delta^{14}\text{C}$ in bulk soil vs $\Delta^{14}\text{CO}_2$ with their equivalent simulation of mean age in bulk soil vs mean transit time in CO_2 for parallel (panel A and B) and series model structure (C and D). Variation of k_1 with $\alpha = 0.8$ and $\gamma = 0.8$. Complementary prediction with $\alpha = 0.1$ and $\gamma = 0.1$ can be seen in Fig. A2.

system showed equal mean age and mean transit time. As k_1 decreased, the transfer from the fast to the slow pool becomes more relevant, then the mean transit time increased compared to the mean age.

250

The behavior of $\Delta^{14}\text{C}$ values contrasted significantly with that of mean age and mean transit time when k_2 tended to lower values (Fig. 7). Generally, when k_2 equals k_1 , $\Delta^{14}\text{C}$ and mean age and mean transit time showed similar values. However, as k_2 decreased, the C stayed in the slow pool for longer time and the $\Delta^{14}\text{C}$ enriched in the bulk soil until reaching a peak (112 and 89 ‰ for series and parallel structures, respectively), from where it subsequently depleted to the initial $\Delta^{14}\text{CO}_2$ value.

255 For the parallel structure (Fig. 7, A and B), transit time kept increasing slowly since most of the C stayed in the fast pool for a short time, so, the C remaining in the slow pool contributed to increase the mean age. As for the series structure, most of




Figure 7. Predictions of $\Delta^{14}\text{C}$ in bulk soil vs $\Delta^{14}\text{C}$ CO_2 with their equivalent simulation of mean age in bulk soil vs mean transit time in CO_2 for parallel (panel A and B) and series model structure (C and D). Variation of k with $\alpha = 0.8$ and $\gamma = 0.8$.

the C was transferred from the fast to the slow pool due to the high transfer rate α , which contributed to increase the transit time.

3.2.2 SOC decomposition in slow cycling systems (peatlands)

260 Simulations of $\Delta^{14}\text{C}$ values in slow cycling systems resulted in significantly more depleted values in both bulk soil and respired CO_2 (Fig. 8) than in fast cycling systems. Generally, $\Delta^{14}\text{C}$ values were more depleted in the bulk soil than in the respiration flux. Also, the results from these simulations showed very different patterns in the $\Delta^{14}\text{C}$ -bulk versus $\Delta^{14}\text{C}$ - CO_2 space than in the mean age versus mean transit time space; only with the exception of equal k_1 and k_2 , which resulted in similar $\Delta^{14}\text{C}$ values as well as similar mean age and mean transit time in the parallel structure.

265




Figure 8. Predictions of $\Delta^{14}\text{C}$ in bulk soil vs $\Delta^{14}\text{C}\text{CO}_2$ with their equivalent simulation of mean age in bulk soil vs mean transit time in CO_2 for parallel (panel A and B) and series model structure (C and D). Variation of k_1 with $\alpha = 0.8$ and $\gamma = 0.8$.

270 Additionally, we looked at the variation of $\Delta^{14}\text{C}$ keeping the same k values but reducing α and γ to 0.1. Simulations indicated that the behavior of $\Delta^{14}\text{C}$ values were opposite between series and parallel structures when inputs to each pool were inversely proportional (Fig. 7 versus Fig. A3 and Fig. 8 versus Fig. A4). For example, for the case of low values of k_1 and k_2 (Fig. 9) ($\gamma = 0.2$ and $\alpha = 0.8$) both model structures showed similar patterns due to relatively similar amounts of inputs going to the slow pool. When $\gamma = 0.8$, $\Delta^{14}\text{C}$ showed values out of the target range (-20 to 23 for $\Delta^{14}\text{C}\text{-CO}_2$ and -86 to -60 for $\Delta^{14}\text{C}$ -bulk based on incubation results) since decomposition occurred as in a fast cycling system. As k_2 decreased, $\Delta^{14}\text{CO}_2$ enriched and $\Delta^{14}\text{C}$ and mean age remained higher than mean transit time since the slow decomposition of the slow pool dominated the system response.




Figure 9. Predictions of $\Delta^{14}\text{C}$ in bulk soil vs $\Delta^{14}\text{C}\text{CO}_2$ with their equivalent simulation of mean age in bulk soil vs mean transit time in CO_2 for parallel (panel A and B). Series model presented exactly the same behaviour. Variation of k_2 with $\alpha = 0.8$ and $\gamma = 0.2$.

275 3.3 Variation in the proportion of inputs γ and α modulated mean age and mean transit time

Our simulations showed the importance that the partitioning and transference coefficients (γ and α) have to define the amount of C entering to each pool and in consequence on the mean age and mean transit time (Fig. A1). For a parallel structure, when the total amount of inputs entered only to the fast pool ($\gamma = 1$), the system behaved as a one pool system and mean age and mean transit time were equal. In the extreme opposite ($\gamma = 0$), all C entered only to the slow pool and mean age and mean transit time converged, although with a longer transit through the system. In contrast, for a series structure, the extreme value $\alpha = 1$ resulted in a longer mean transit time than mean age as the contribution to the respired flux comprehended the sum of the decomposition of the newly absorbed C and the initial stocks stored in the fast pool. In the case of slow cycling systems (Fig. 280 A5 and Fig. A6), $\Delta^{14}\text{CO}_2$ moved in a wider range including very depleted values (-100 to 70 ‰) and approached to a steady



state of $\Delta^{14}\text{C}$ in bulk soil from the right when $k_1 = 0.01$ or from the left when $k_1 = 0.1$.

285

4 Discussion

4.1 Did changes in temperature and soil moisture result in the release of old carbon in the respiration flux?

We measured the $\Delta^{14}\text{C}$ values of respired CO_2 from soils incubated at different temperatures and WFPS levels to assess whether changes in these variables would result in destabilization of stored C. Our results showed that changes in temperature
290 did not affect systematically the radiocarbon values of respired CO_2 in any of the incubated soils. Nevertheless, changes in WFPS had a moderate effect on the $\Delta^{14}\text{CO}_2$ in grasslands, but a strong influence on the $\Delta^{14}\text{CO}_2$ of peatlands. Our results indicated that higher WFPS levels led to depleted $\Delta^{14}\text{CO}_2$ values in peatlands at all temperature treatments, except at 10 °C. In general, higher temperature and WFPS levels resulted in increased CO_2 fluxes although without showing a relationship with the $\Delta^{14}\text{C}$ values.

295

In contrast, higher WFPS resulted in more enriched $\Delta^{14}\text{CO}_2$ values for grasslands. This suggests that the direction of changing WFPS depends on ecosystem characteristics. It is interesting to observe that the $\Delta^{14}\text{C}$ in the bulk soil was always more negative than the $\Delta^{14}\text{CO}_2$ in the peatlands, while the grasslands registered similar $\Delta^{14}\text{C}$ values in both states. This indicates that peatlands are systems that stabilize organic matter over time and release it once the stable conditions change. In contrast,
300 for grasslands two interpretations may arise, first that there was not organic matter stabilization under the incubated conditions since the respired flux reassembles the bulk soil $\Delta^{14}\text{C}$ signature, or second that the incubation conditions were not strong enough to destabilize the existing old SOM. Our simulations indicate that similar $\Delta^{14}\text{C}$ values in the bulk and in the respired CO_2 can occur at an specific year (Fig A7) independently of the different SOM cycling times in the soil pools.

305 In tundra ecosystems, Kwon et al. (2019) suggested that drainage of shallow soil layers may have accelerated old carbon decomposition. In addition, Estop-Aragones et al. (2020) concluded that old C would increase in proportion from “cold” across “warm wet” to “warm and dry” for high arctic tundra as well as from “undisturbed” to “burnt active layer” for peatland plateau. These previous findings are in disagreement with our observations since after our experiments, lower WFPS resulted in the release of relatively enriched $\Delta^{14}\text{CO}_2$ instead of releasing old carbon. One potential explanation for this discrepancy is that
310 the existing old SOM could have reacted too slow to drier conditions (WFPS = 65) compared to the fast response of “younger” SOM, causing an evident prevalence of young carbon given the short duration of the incubations. Consistent with this, it has been found that some SOM components or compartments may be more sensitive to modified conditions than others (Feng and Simpson, 2008) and therefore, transit time would react strongly when such sensitivity is higher in the slower pool (Manzoni et al., 2009).

315



For this reason, our results are indicative, but not conclusive, about the influence of environmental factors on the $\Delta^{14}\text{C}$ signature of respired carbon. It is possible that our $\Delta^{14}\text{C}$ values are limited by the short time of the incubations since the dynamics of C transfers cannot be properly observed in short timescales (Crow and Sierra, 2018). For example, during a 1-year incubation experiment, most of the CO_2 was derived from labile SOM as the temperature increased (Leifeld and Fuhrer, 2005).
320 Additionally, other factors such as yearly climate seasonality, daily freeze-thaw cycles and water table oscillation were not replicated in the laboratory due to equipment and time availability. Finally, although temperature has been found to be the main modulator of SOC decomposition rates (Azizi-Rad et al., 2022), dominating over the effect of soil moisture and oxygen availability, it is important to consider that an increase in respiration rates, as observed in our incubations, does not necessarily involve the release of old or young carbon.

325

4.2 Are there differences in the age of respired CO_2 between grassland and peatland?

We found that $\Delta^{14}\text{C}$ values from peatland were strongly more depleted than those in grasslands, indicating the presence and respiration of older carbon in peatlands. Generally, low temperature, low soil microbial activity and anoxic conditions (Xiang et al., 2009; Ma et al., 2016) posed favourable conditions for the stabilization of organic matter since the early Holocene in
330 the Zoige region (Chen et al., 2014; Sun et al., 2017). Older bulk soil than CO_2 demonstrated that the peatland soil behaved as a retention system (Type II) where C was stored for relatively long time before its respiration (Sierra et al., 2018b). The release of that old C may occur when the SOM destabilization dominates over the stabilization mechanisms, for example when peatlands are drained (increasing oxidation through water table reduction) and their temperature is raised (Dutta et al., 2006; Hicks Pries et al., 2013; Lupascu et al., 2014; Estop-Aragonés et al., 2018), both factors mimicked in our incubations.

335

In contrast, $\Delta^{14}\text{CO}_2$ from grasslands were mostly similar to $\Delta^{14}\text{C}$ in bulk soil, indicating that the soil behaved predominantly as a well-mixed homogeneous system (Type I) where most of the SOM has the same probability of being mineralized and released as CO_2 (Sierra et al., 2018b). However, $\Delta^{14}\text{C}$ relation also indicated that the grasslands shift from retention to non-retention system and vice-versa. In grasslands, SOM from the topsoil is permanently under changing conditions (daily
340 temperature and soil moisture variation, grazing and mechanical alteration, among others) that promote its fast cycling instead of its stabilization (Han et al., 2017). These factors, along with climatic, vegetation and edaphic properties caused a low SOM stability expressed in a large proportion of labile carbon (Hou et al., 2021). Nonetheless, although SOM accumulation still occurs in grassland areas (Tian et al., 2009) registered in the extremely depleted outliers from the grassland soil respiration, it happens at a slower rate than in peatlands. As a result, most of the fixed atmospheric C is respired at short time scales, which
345 is in turn registered as young (“post bomb”) $\Delta^{14}\text{C}$ values.

Taken together, SOM stability and ecosystem characteristics defined the relation between $\Delta^{14}\text{C}$ in bulk soil and $\Delta^{14}\text{CO}_2$, which in turn explained the difference in the age of respiration between grassland and peatlands. The $\Delta^{14}\text{CO}_2$ values obtained in our incubated peatlands indicate that such C was captured through photosynthesis during a time when the ^{14}C levels of



350 the atmosphere were depleted and have remained stored in the soil for a long enough time for radioactive decay to become relevant. However, it is imprecise to attribute an specific year to the entire bulk soil or CO₂ fractions, since they are formed by SOM accumulated in multiple steps and of different qualities and their respective decomposition.

In that sense, a $\Delta^{14}\text{C}$ value is not per se indicative of age or transit time of SOC. To disentangle the utility of radiocarbon as a tool for SOM persistence in soil and be able to shed light on the time-scales of C cycling in different ecosystems, we used SOC decomposition models. Our models contribute to understand the time component of SOM persistence through the use of mean transit time as a parameter that integrates processes of SOM dynamics (Manzoni et al., 2009).

360 **4.3 How can $\Delta^{14}\text{C}$ values be interpreted to understand the effect of decomposition rates on mean age and mean transit time?**

We simulated the $\Delta^{14}\text{C}$ values and the mean age and mean transit time of soils with high and low decomposition rates. Our simulations were able to reassemble the $\Delta^{14}\text{C}$ values obtained from incubations and showed that modelled $\Delta^{14}\text{C}$ values differed widely between slow cycling systems and fast cycling systems. Generally, decomposition rate, transfer rates and partitioning coefficients of a given model structure modulated $\Delta^{14}\text{C}$ values, and in consequence, mean age and mean transit time (Bruun et al., 2004; Manzoni et al., 2009). For example, fast cycling systems with high decomposition rates resulted in a fast respiration represented by short mean transit times (lower than 20 years). At the same time, low decomposition rates resulted in longer mean transit times, which in turn represented older respired CO₂ (lower than 140 years). In slow cycling systems, where the decomposition rates were lower, transit times were consequently longer (lower than 20000 years). Independently of the k values, the series model structure increased the transit time of C compared to parallel structure due to the fact that a proportion of the C atoms had to pass through the two pools before being respired.

Other factors such as observation time and starting year of the simulation modified the response of $\Delta^{14}\text{C}$. This is due to the radioactive nature of the ¹⁴C isotope and the influence of the “bomb ¹⁴C”. For instance, the $\Delta^{14}\text{C}$ enrichment during the last 60 years was clearly observed in fast cycling systems only with low k_2 . Furthermore, the inflexion points on the $\Delta^{14}\text{C}$ curves occurred at different values for slow and fast cycling systems. Hence, we could assume that a simulation with starting year after 1962 would not show an inflexion point in the $\Delta^{14}\text{C}$ due to the absence of the bomb peak.

Our modelling results suggest that the increase of decomposition rates contribute to the release of older carbon in the respired flux as we hypothesised, but depending on the initial state of the system. This interpretation is not straightforward since the stability of SOM depends on the specific combination of temperature and moisture for different ecosystems. For peatlands, only drier conditions and consequent increase of oxygen might increase k and therefore cause the destabilization of SOM. In contrast, grasslands would need an increase of temperature to facilitate SOM decomposition provided that moisture and oxygen are available (Azizi-Rad et al., 2022). SOM decomposition rates are expected to increase at higher temperatures (Leifeld and



Fuhrer, 2005) and towards the extremes of the moisture range (Sierra et al., 2015). However, the destabilization of SOM can
385 occur in any direction of WFPS variation (depending on the oxygen content) and in turn affect the age and transit time of C in
a nonlinear pattern.

From our results we could observe how the relation $\Delta^{14}\text{C}$ -bulk versus $\Delta^{14}\text{C}$ - CO_2 properly represented the relation mean
age versus mean transit time. We found that there is a good correspondence between both relations in the fast cycling systems
390 as long as the decomposition rates k remain high. Such correspondence did not occur as k became smaller (typical for slow
cycling systems) since the appearance of the bomb peak may have introduced anomalies that modified the equivalence between
the two relations. In addition, the outlier $\Delta^{14}\text{C}$ values found in our experimental data, where $\Delta^{14}\text{C}$ changed drastically in the
vertical and the horizontal direction, may be related to this complex response of the ^{14}C tracer.

395 To quantify cycling times of carbon, radiocarbon can be used as a tool to understand SOM destabilization and persistence
through the use of the concepts of age and transit time and their mutual relation. Nonetheless, it is essential to couple $\Delta^{14}\text{C}$
measured values with a model that involves the dynamics of soil carbon in different pools and their interaction with the envi-
ronment. Therefore, the acquisition of empirical data from soils (number of pools, I , C , ks , γ , α) along with the correct setting
of model structure will improve predictions on terrestrial and atmospheric carbon interactions.

400

5 Conclusions

Based on the incubation results of soils from the QTP, we showed that the $\Delta^{14}\text{C}$ values of the peatland are significantly more
depleted than the ones of the grassland both in the bulk soil and the respired CO_2 . Our results indicated that changes in temper-
ature did not affect systematically the radiocarbon values of respired CO_2 in any of the incubated soils. Nevertheless, changes
405 in WFPS had a relatively small effect on the $^{14}\text{CO}_2$ in grasslands, but a strong influence on the $^{14}\text{CO}_2$ of peatlands, where
higher WFPS levels led to more depleted $^{14}\text{CO}_2$ values except at 10 °C. In peatlands, more depleted ^{14}C -bulk values than
 ^{14}C - CO_2 indicated that SOM stabilizes over time and it is released once the stable conditions change. In grasslands, similar
 $\Delta^{14}\text{C}$ values in bulk and respired CO_2 indicated that the soil behaved as a well-mixed homogeneous system due to either an
absence of SOM stabilization or that the manipulation treatments were not long or strong enough to destabilize the existing
410 old SOM. In this sense, the short duration of our incubations might have been an obstacle to register the influence of long term
factors such as climate seasonality and water table oscillation on SOM dynamics.

From our modelling approach, we conclude that radiocarbon can be used as a tool to understand SOM persistence through
the use of the concepts of mean age and mean transit time and their mutual relation. Our simulations were able to reassemble the
415 $\Delta^{14}\text{C}$ values obtained from incubations and showed that modelled $\Delta^{14}\text{C}$ values differed widely between slow cycling systems
and fast cycling systems. We found that low values of k , more common in slow cycling systems, modified the behaviour of



$\Delta^{14}\text{C}$ patterns due to the incorporation of ^{14}C -bomb in the soil system. Hence, the correspondence between these mutual relations strongly depended on the internal dynamics of the soil and its interaction with the environment. For this reason, the acquisition of empirical data from soils (number of pools, I , C , k_s , γ and α) along with the correct setting of model structure will improve predictions on terrestrial and atmospheric carbon interactions.

Code and data availability. Lab analysis results and code are stored at <https://doi.org/10.5281/zenodo.7620008> (DOI:10.5281/zenodo.7620008) (Tangarife-Escobar et al., 2023).

Author contributions. The conceptualization, data curation and formal analysis were done by ATE and CS. The investigation was performed by ATE with support from GD and MAR. Methodology and validation were done by ATE and CS. ATE conducted the validation, visualization and writing of the original draft (the latter with the help of CUM). Supervision was done by GG, XF and CS. ATE wrote the manuscript with contributions from CS, GG, XF, CUM, GD and MAR.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. We thank all colleagues who contributed to this study, especially Eike Reinosch for providing the location maps used in Fig.1. and Paula Sierra for the digitization of the scheme used in Fig. 2. Special thanks to David Martini for his helpful advices to improve code writing and to Manuel Röst and Axel Steinhof for their useful training at the ^{14}C laboratory. Finally, thanks to Nicole Börner for the active collaboration with the project administration and to Susan Trumbore for her feedback on the final version of this manuscript. This study was developed as part of the International Research Training Group (GRK 2309/1) Geo-ecosystems in transition on the Tibetan Plateau (TransTiP) funded by the Deutsche Forschungsgemeinschaft (DFG). The Max Planck Institute for Biogeochemistry provided permanent administrative support.



435 **Appendix A: Appendix A**

Table A1. Mean daily CO₂ respiration for incubated soils under temperature and WFPS variation.

Ecosystem	Temperature (°C)	WFPS (%)	Mean CO ₂ respiration (mg CO ₂ g soil ⁻¹ day ⁻¹)	σ
Grassland	20	95	0.510	0.035
	20	60	0.371	0.016
	15	95	0.330	0.008
	15	60	0.234	0.004
	10	95	0.209	0.011
	10	60	0.134	0.023
	5	95	0.066	0.003
	5	60	0.044	0.001
Peatland	20	95	0.040	0.001
	20	60	0.028	0.003
	15	95	0.024	0.004
	15	60	0.016	0.003
	10	95	0.011	0.002
	10	60	0.008	0.001
	5	95	0.005	0.000
	5	60	0.004	0.001



Table A2. Parameters used for SOC decomposition models in grassland (fast cycling) and peatland (slow cycling).

System	Fig.	Model structure	k_1	k_2	γ	α	Starting year of simulation	$\Delta^{14}\text{C}$ (‰) for starting year
Fast cycling	A1	Parallel	0.8	0.1	0 - 1	-	1890	-4.9 ‰
		Series	0.8	0.1	-	0 - 1	1890	-4.9 ‰
	A2	Parallel	0.1 - 0.8	0.1	0.1	-	1890	-4.9 ‰
		Series	0.1 - 0.8	0.1	-	0.1	1890	-4.9 ‰
	A3	Parallel	0.8	0.0001 - 0.8	0.1	-	1890	-4.9 ‰
		Series	0.8	0.0001 - 0.8	-	0.1	1890	-4.9 ‰
Slow cycling	A4	Parallel	0.0001 - 0.8	0.0001	0.2	-	500	-22.2 ‰
		Series	0.0001 - 0.8	0.0001	-	0.1	500	-22.2 ‰
	A5	Parallel	0.01	0.0001	0 - 1	-	500	-22.2 ‰
		Series	0.01	0.0001	-	0 - 1	500	-22.2 ‰
	A6	Parallel	0.1	0.0001	0 - 1	-	500	-22.2 ‰
		Series	0.1	0.0001	-	0 - 1	500	-22.2 ‰



- $\Delta^{14}\text{C}$ variation in fast cycling systems




Figure A1. Predictions of $\Delta^{14}\text{C}$ in bulk soil vs $\Delta^{14}\text{CO}_2$ with their equivalent simulation of mean age in bulk soil vs mean transit time in CO_2 for parallel (panel A and B) and series model structure (C and D). $k_1 = 0.8$, $k_2 = 0.1$, with $\alpha = 0 - 1$ and $\gamma = 0 - 1$.




Figure A2. Predictions of $\Delta^{14}\text{C}$ in bulk soil vs $\Delta^{14}\text{CO}_2$ with their equivalent simulation of mean age in bulk soil vs mean transit time in CO_2 for parallel (panel A and B) and series model structure (C and D). Variation of k_1 with $\alpha = 0.1$ and $\gamma = 0.1$.




Figure A3. Predictions of $\Delta^{14}\text{C}$ in bulk soil vs $\Delta^{14}\text{CO}_2$ with their equivalent simulation of mean age in bulk soil vs mean transit time in CO_2 for parallel (panel A and B) and series model structure (C and D). Variation of k_2 with $\alpha = 0.1$ and $\gamma = 0.1$.



- $\Delta^{14}\text{C}$ variation in slow cycling systems




Figure A4. Predictions of $\Delta^{14}\text{C}$ in bulk soil vs $\Delta^{14}\text{CO}_2$ with their equivalent simulation of mean age in bulk soil vs mean transit time in CO_2 for parallel (panel A and B) and series model structure (C and D). Variation of k_1 with $\alpha = 0.1$ and $\gamma = 0.2$.




Figure A5. Predictions of $\Delta^{14}\text{C}$ in bulk soil vs $\Delta^{14}\text{CO}_2$ with their equivalent simulation of mean age in bulk soil vs mean transit time in CO_2 for parallel (panel A and B) and series model structure (C and D). $k_1 = 0.01$, $k_2 = 0.0001$, with $\alpha = 0 - 1$ and $\gamma = 0 - 1$.




Figure A6. Predictions of $\Delta^{14}\text{C}$ in bulk soil vs $\Delta^{14}\text{CO}_2$ with their equivalent simulation of mean age in bulk soil vs mean transit time in CO_2 for parallel (panel A and B) and series model structure (C and D). $k_1 = 0.1$, $k_2 = 0.0001$, with $\alpha = 0 - 1$ and $\gamma = 0 - 1$.



- $\Delta^{14}\text{C}$ curves for bulk soil, respired CO_2 and atmosphere for a two-pool system in parallel and series structure




Figure A7. Variation of $\Delta^{14}\text{C}$ values in atmosphere, bulk soil and respired CO_2 for a two-pool soil during the period 1950 - 2019. The similarity between $\Delta^{14}\text{C}$ values may indicate both a well-mixed homogeneous system (type I) or a retention system (type II) where the bulk soil and the respired flux record the same value. Nonetheless, the mean transit time for the bulk and the soil respiration might be different due to the contrasting decomposition rates of the fast ($k_1 = 0.8$) and the slow pool ($k_2 = 0.1$). Simulation conducted with $\alpha = 0.8$ and $\gamma = 0.8$.



References

- 440 Anslan, S., Rad, M. A., Buckel, J., Galindo, P. E., Kai, J., Kang, W., Keys, L., Maurischat, P., Nieberding, F., Reinosch, E., Tang, H., Tran, T. V., Wang, Y., and Schwalb, A.: Reviews and syntheses: How do abiotic and biotic processes respond to climatic variations in the Nam Co catchment (Tibetan Plateau)?, *Biogeosciences*, 17, 1261–1279, <https://doi.org/10.5194/bg-17-1261-2020>, 2020.
- Arias, P., Bellouin, N., Coppola, E., Jones, R., Krinner, G., Marotzke, J., Naik, V., Palmer, M., Plattner, G.-K., Rogelj, J., et al.: Climate Change 2021: The Physical Science Basis. Contribution of Working Group 14 I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change; Technical Summary, 2021.
- 445 Azizi-Rad, M., Guggenberger, G., Ma, Y., and Sierra, C. A.: Sensitivity of soil respiration rate with respect to temperature, moisture and oxygen under freezing and thawing, *Soil Biology and Biochemistry*, 165, 108488, <https://doi.org/https://doi.org/10.1016/j.soilbio.2021.108488>, 2022.
- Blanco-Canqui, H. and Lal, R.: Mechanisms of carbon sequestration in soil aggregates, *Critical reviews in plant sciences*, 23, 481–504, 2004.
- 450 Bolin, B. and Rodhe, H.: A note on the concepts of age distribution and transit time in natural reservoirs, *Tellus*, 25, 58–62, <https://doi.org/https://doi.org/10.1111/j.2153-3490.1973.tb01594.x>, 1973.
- Bradford, M. A., Wieder, W. R., Bonan, G. B., Fierer, N., Raymond, P. A., and Crowther, T. W.: Managing uncertainty in soil carbon feedbacks to climate change, *Nature Climate Change*, 6, 751–758, 2016.
- Briones, M. J., Garnett, M. H., and Ineson, P.: No evidence for increased loss of old carbon in a temperate organic soil after 13 years of simulated climatic warming despite increased CO₂ emissions, *Global Change Biology*, 2021.
- 455 Briones, M. J. I., Garnett, M. H., and Ineson, P.: Soil biology and warming play a key role in the release of ‘old C’ from organic soils, *Soil Biology and Biochemistry*, 42, 960–967, <https://doi.org/10.1016/j.soilbio.2010.02.013>, 2010.
- Bruun, S., Six, J., and Jensen, L. S.: Estimating vital statistics and age distributions of measurable soil organic carbon fractions based on their pathway of formation and radiocarbon content, *Journal of theoretical biology*, 230, 241–250, 2004.
- 460 Chen, H., Yang, G., Peng, C., Zhang, Y., Zhu, D., Zhu, Q., Hu, J., Wang, M., Zhan, W., Zhu, E., Bai, Z., Li, W., Wu, N., Wang, Y., Gao, Y., Tian, J., Kang, X., Zhao, X., and Wu, J.: The carbon stock of alpine peatlands on the Qinghai-Tibetan Plateau during the Holocene and their future fate, *Quaternary Science Reviews*, 95, 151–158, <https://doi.org/10.1016/j.quascirev.2014.05.003>, 2014.
- Chen, L., Jing, X., Flynn, D. F., Shi, Y., Kühn, P., Scholten, T., and He, J.-S.: Changes of carbon stocks in alpine grassland soils from 2002 to 2011 on the Tibetan Plateau and their climatic causes, *Geoderma*, 288, 166–174, 2017.
- 465 Chen, L., Fang, K., Wei, B., Qin, S., Feng, X., Hu, T., Ji, C., and Yang, Y.: Soil carbon persistence governed by plant input and mineral protection at regional and global scales, *Ecology Letters*, 24, 1018–1028, 2021.
- Crow, S. E. and Sierra, C. A.: Dynamic, intermediate soil carbon pools may drive future responsiveness to environmental change, *Journal of environmental quality*, 47, 607–616, 2018.
- Davidson, E. and Janssens, I.: Temperature sensitivity of soil carbon decomposition and feedbacks to climate change, *Nature*, 440, 165–173, <https://doi.org/10.1038/nature04514>, 2006.
- 470 Dioumaeva, I., Trumbore, S., Schuur, E., Goulden, M., Litvak, M., and Hirsch, A.: Decomposition of peat from upland boreal forest: Temperature dependence and sources of respired carbon, *Journal of Geophysical Research: Atmospheres*, 108, <https://doi.org/10.1029/2001JD000848>, 2002.
- Dong, S., Peng, F., You, Q., Guo, J., and Xue, X.: Lake dynamics and its relationship to climate change on the Tibetan Plateau over the last four decades, *Regional environmental change*, 18, 477–487, 2018.
- 475



- Du, C. and Gao, Y.: Opposite patterns of soil organic and inorganic carbon along a climate gradient in the alpine steppe of northern Tibetan Plateau, *Catena*, 186, 104–366, 2020.
- Dutta, K., Schuur, E. A. G., Neff, J. C., and Zimov, S. A.: Potential carbon release from permafrost soils of Northeastern Siberia, *Global Change Biology*, 12, 2336–2351, <https://doi.org/10.1111/j.1365-2486.2006.01259.x>, 2006.
- 480 Emanuel, W., Killough, G., Post, W., and Shugart, H.: Modeling terrestrial ecosystems in the global carbon-cycle with shifts in carbon storage capacity by land-use change, *Ecology*, 65, 970–983, <https://doi.org/10.2307/1938069>, 1984.
- Eriksson, E.: Compartment Models and Reservoir Theory, *Annual Review of Ecology and Systematics*, 2, 67–84, <https://doi.org/10.1146/annurev.es.02.110171.000435>, 1971.
- Estop-Aragonés, C., Czimczik, C. I., Heffernan, L., Gibson, C., Walker, J. C., Xu, X., and Olefeldt, D.: Respiration of aged soil carbon during
485 fall in permafrost peatlands enhanced by active layer deepening following wildfire but limited following thermokarst, *Environmental Research Letters*, 13, 085002, 2018.
- Estop-Aragones, C., Olefeldt, D., Abbott, B. W., Chanton, J. P., Czimczik, C. I., Dean, J. F., Egan, J. E., Gandois, L., Garnett, M. H., Hartley, I. P., Hoyt, A., Lupascu, M., Natali, S. M., O'Donnell, J. A., Raymond, P. A., Tanentzap, A. J., Tank, S. E., Schuur, E. A. G., Turetsky, M., and Anthony, K. W.: Assessing the Potential for Mobilization of Old Soil Carbon After Permafrost Thaw: A Synthesis of
490 C-14 Measurements From the Northern Permafrost Region, *Global Biogeochemical Cycles*, 34, <https://doi.org/10.1029/2020GB006672>, 2020.
- Falloon, P. and Smith, P.: Modelling refractory soil organic matter, *Biology and fertility of soils*, 30, 388–398, 2000.
- Feng, X. and Simpson, M. J.: Temperature responses of individual soil organic matter components, *Journal of Geophysical Research: Biogeosciences*, 113, 2008.
- 495 Ganjurjav, H., Gao, Q., Gornish, E. S., Schwartz, M. W., Liang, Y., Cao, X., Zhang, W., Zhang, Y., Li, W., Wan, Y., Li, Y., Danjiu, L., Guo, H., and Lin, E.: Differential response of alpine steppe and alpine meadow to climate warming in the central Qinghai-Tibetan Plateau, *Agricultural and Forest Meteorology*, 223, 233–240, <https://doi.org/10.1016/j.agrformet.2016.03.017>, 2016.
- Gao, Q.-z., Li, Y., Xu, H.-m., Wan, Y.-f., and Jiangcun, W.-z.: Adaptation strategies of climate variability impacts on alpine grassland ecosystems in Tibetan Plateau, *Mitigation and Adaptation Strategies for Global Change*, 19, 199–209, 2014.
- 500 Gaudinski, J., Trumbore, S., Davidson, E., and Zheng, S.: Soil carbon cycling in a temperate forest: radiocarbon-based estimates of residence times, sequestration rates and partitioning of fluxes, *Biogeochemistry*, 51, 33–69, <https://doi.org/10.1023/A:1006301010014>, 2000.
- Geng, Y., Wang, Y., Yang, K., Wang, S., Zeng, H., Baumann, F., Kuehn, P., Scholten, T., and He, J.-S.: Soil respiration in Tibetan alpine grasslands: belowground biomass and soil moisture, but not soil temperature, best explain the large-scale patterns, *PLoS one*, 7, e34968, 2012.
- 505 Genxu, W., Ju, Q., Guodong, C., and Yuanmin, L.: Soil organic carbon pool of grassland soils on the Qinghai-Tibetan Plateau and its global implication, *Science of the Total Environment*, 291, 207–217, 2002.
- Han, C., Wang, Z., Si, G., Lei, T., Yuan, Y., and Zhang, G.: Increased precipitation accelerates soil organic matter turnover associated with microbial community composition in topsoil of alpine grassland on the eastern Tibetan Plateau, *Canadian journal of microbiology*, 63, 811–821, 2017.
- 510 Hao, Y. B., Cui, X. Y., Wang, Y. F., Mei, X. R., Kang, X. M., Wu, N., Luo, P., and Zhu, D.: Predominance of Precipitation and Temperature Controls on Ecosystem CO₂ Exchange in Zoige Alpine Wetlands of Southwest China, *Wetlands*, 31, 413–422, <https://doi.org/10.1007/s13157-011-0151-1>, 2011.



- Hicks Pries, C. E., Schuur, E. A., and Crummer, K. G.: Thawing permafrost increases old soil and autotrophic respiration in tundra: Partitioning ecosystem respiration using D13C and D14C, *Global Change Biology*, 19, 649–661, 2013.
- 515 Hopkins, F. M., Torn, M. S., and Trumbore, S. E.: Warming accelerates decomposition of decades-old carbon in forest soils, *Proceedings of the National Academy of Sciences of the United States of America*, 109, E1753–E1761, <https://doi.org/10.1073/pnas.1120603109>, 2012.
- Hou, Y., He, K., Chen, Y., Zhao, J., Hu, H., and Zhu, B.: Changes of soil organic matter stability along altitudinal gradients in Tibetan alpine grassland, *Plant and Soil*, 458, 21–40, 2021.
- Hua, Q., Turnbull, J. C., Santos, G. M., Rakowski, A. Z., Ancapichún, S., De Pol-Holz, R., Hammer, S., Lehman, S. J., Levin, I., Miller, J. B., et al.: Atmospheric radiocarbon for the period 1950–2019, *Radiocarbon*, pp. 1–23, 2021.
- 520 Jarvis, A., Reuter, H. I., Nelson, A., and Guevara, E.: Hole-filled SRTM for the globe Version 4. available from the CGIAR-CSI SRTM 90m Database, 15, (25–54), <http://srtm.csi.cgiar.org>, 2008.
- Kang, X., Yan, L., Cui, L., Zhang, X., Hao, Y., Wu, H., Zhang, Y., Li, W., Zhang, K., Yan, Z., Li, Y., and Wang, J.: Reduced Carbon Dioxide Sink and Methane Source under Extreme Drought Condition in an Alpine Peatland, *Sustainability*, 10, <https://doi.org/10.3390/su10114285>, 2018.
- 525 Knorr, W., Prentice, I., House, J., and Holland, E.: Long-term sensitivity of soil carbon turnover to warming, *Nature*, 433, 298–301, <https://doi.org/10.1038/nature03226>, 2005.
- Kwon, M. J., Natali, S. M., Pries, C. E. H., Schuur, E. A. G., Steinhof, A., Crummer, K. G., Zimov, N., Zimov, S. A., Heimann, M., Kolle, O., and Goeckede, M.: Drainage enhances modern soil carbon contribution but reduces old soil carbon contribution to ecosystem respiration in tundra ecosystems, *Global Change Biology*, 25, 1315–1325, <https://doi.org/10.1111/gcb.14578>, 2019.
- 530 Leifeld, J. and Fuhrer, J.: The temperature response of CO₂ production from bulk soils and soil fractions is related to soil organic matter quality, *Biogeochemistry*, 75, 433–453, 2005.
- Li, J., Yan, D., Pendall, E., Pei, J., Noh, N. J., He, J.-S., Li, B., Nie, M., and Fang, C.: Depth dependence of soil carbon temperature sensitivity across Tibetan permafrost regions, *Soil Biology and Biochemistry*, 126, 82–90, 2018.
- 535 Liski, J., Ilvesniemi, H., Makela, A., and Westman, C.: CO₂ emissions from soil in response to climatic warming are overestimated - The decomposition of old soil organic matter is tolerant of temperature, *Ambio*, 28, 171–174, 1999.
- Liu, L., Chen, H., Zhu, Q., Yang, G., Zhu, E., Hu, J., Peng, C., Jiang, L., Zhan, W., Ma, T., He, Y., and Zhu, D.: Responses of peat carbon at different depths to simulated warming and oxidizing, *Science of the Total Environment*, 548, 429–440, <https://doi.org/10.1016/j.scitotenv.2015.11.149>, 2016.
- 540 Liu, L., Chen, H., Liu, X., Yang, Z., Zhu, D., He, Y., and Liu, J.: Contemporary, modern and ancient carbon fluxes in the Zoige peatlands on the Qinghai-Tibetan Plateau, *Geoderma*, 352, 138–149, <https://doi.org/10.1016/j.geoderma.2019.06.008>, 2019a.
- Liu, X., Zhu, D., Zhan, W., Chen, H., Zhu, Q., Zhang, J., Wu, N., and He, Y.: Dominant influence of non-thawing periods on annual CO₂ emissions from Zoige peatlands: Five-year eddy covariance analysis, *Ecological Indicators*, 129, 107 913, 2021.
- Liu, X., Chen, H., Zhu, Q., Wu, J., Frohking, S., Zhu, D., Wang, M., Wu, N., Peng, C., and He, Y.: Holocene peatland development and carbon stock of Zoige peatlands, Tibetan Plateau: a modeling approach, *Journal of Soils and Sediments*, 18, 2032–2043, <https://doi.org/10.1007/s11368-018-1960-0>, 2018.
- 545 Liu, X., Zhu, D., Zhan, W., Chen, H., Zhu, Q., Hao, Y., Liu, W., and He, Y.: Five-Year Measurements of Net Ecosystem CO₂ Exchange at a Fen in the Zoige Peatlands on the Qinghai-Tibetan Plateau, *Journal of Geophysical Research: Atmospheres*, 124, 11 803–11 818, <https://doi.org/10.1029/2019JD031429>, 2019b.



- 550 Lu, M., Zhou, X., Yang, Q., Li, H., Luo, Y., Fang, C., Chen, J., Yang, X., and Li, B.: Responses of ecosystem carbon cycle to experimental warming: a meta-analysis, *Ecology*, 94, 726–738, 2013.
- Lupascu, M., Welker, J., Xu, X., and Czimczik, C.: Rates and radiocarbon content of summer ecosystem respiration in response to long-term deeper snow in the High Arctic of NW Greenland, *Journal of Geophysical Research: Biogeosciences*, 119, 1180–1194, 2014.
- Ma, K., Zhang, Y., Tang, S., and Liu, J.: Spatial distribution of soil organic carbon in the Zoige alpine wetland, northeastern Qinghai-Tibet Plateau, *Catena*, 144, 102–108, <https://doi.org/10.1016/j.catena.2016.05.014>, 2016.
- 555 Manzoni, S., Katul, G. G., and Porporato, A.: Analysis of soil carbon transit times and age distributions using network theories, *Journal of Geophysical Research: Biogeosciences*, 114, 2009.
- McGuire, A., Sitch, S., Clein, J. S., Dargaville, R., Esser, G., Foley, J., Heimann, M., Joos, F., Kaplan, J., Kicklighter, D., et al.: Carbon balance of the terrestrial biosphere in the twentieth century: Analyses of CO₂, climate and land use effects with four process-based ecosystem models, *Global biogeochemical cycles*, 15, 183–206, 2001.
- 560 Mesfin, S., Gebresamuel, G., Haile, M., and Zenebe, A.: Modelling spatial and temporal soil organic carbon dynamics under climate and land management change scenarios, northern Ethiopia, *European Journal of Soil Science*, 72, 1298–1311, 2021.
- Metzler, H. and Sierra, C. A.: Linear Autonomous Compartmental Models as Continuous-Time Markov Chains: Transit-Time and Age Distributions, *Mathematical Geosciences*, 50, 1–34, <https://doi.org/10.1007/s11004-017-9690-1>, 2018.
- 565 Nieberding, F., Wille, C., Fratini, G., Asmussen, M. O., Wang, Y., Ma, Y., and Sachs, T.: A long-term (2005–2019) eddy covariance data set of CO₂ and H₂O fluxes from the Tibetan alpine steppe, *Earth System Science Data*, 12, 2505–2524, <https://doi.org/10.5194/essd-12-2705-2020>, 2020.
- Pan, Y., Li, X., Li, S., and Li, Z.: Different responses of soil respiration to climate change in permafrost and non-permafrost regions of the Tibetan plateau from 1979 to 2018, *International Journal of Climatology*, 2022.
- 570 Piao, S., Tan, K., Nan, H., Ciais, P., Fang, J., Wang, T., Vuichard, N., and Zhu, B.: Impacts of climate and CO₂ changes on the vegetation growth and carbon balance of Qinghai–Tibetan grasslands over the past five decades, *Global and Planetary Change*, 98, 73–80, 2012.
- Pold, G., Melillo, J. M., and DeAngelis, K. M.: Two decades of warming increases diversity of a potentially lignolytic bacterial community, *Frontiers in microbiology*, 6, 480, 2015.
- Rustad, L., Campbell, J., Marion, G., Norby, R., Mitchell, M., Hartley, A., Cornelissen, J., and Gurevitch, J.: A meta-analysis of the response of soil respiration, net nitrogen mineralization, and aboveground plant growth to experimental ecosystem warming, *Oecologia*, 126, 543–562, 2001.
- 575 Schimel, D.: Terrestrial ecosystems and the carbon-cycle, *Global Change Biology*, 1, 77–91, <https://doi.org/10.1111/j.1365-2486.1995.tb00008.x>, 1995.
- Schmidt, M. W., Torn, M. S., Abiven, S., Dittmar, T., Guggenberger, G., Janssens, I. A., Kleber, M., Kögel-Knabner, I., Lehmann, J., Manning, D. A., et al.: Persistence of soil organic matter as an ecosystem property, *Nature*, 478, 49–56, 2011.
- 580 Schuur, E. A. G., Vogel, J. G., Crummer, K. G., Lee, H., Sickman, J. O., and Osterkamp, T. E.: The effect of permafrost thaw on old carbon release and net carbon exchange from tundra, *Nature*, 459, 556–559, <https://doi.org/10.1038/nature08031>, 2009.
- Schuur, E. A. G., Druffel, E., and Trumbore, S. E.: Radiocarbon and Climate Change: Mechanisms, Applications and Laboratory Techniques, pp. 1–315, Springer International, <https://doi.org/10.1007/978-3-319-25643-6>, 2016.
- 585 Scurlock, J. and Hall, D.: The global carbon sink: a grassland perspective, *Global Change Biology*, 4, 229–233, 1998.
- Sierra, C. A. and Mueller, M.: A general mathematical framework for representing soil organic matter dynamics, *Ecological Monographs*, 85, 505–524, <https://doi.org/10.1890/15-0361.1>, 2015.



- Sierra, C. A., Trumbore, S. E., Davidson, E. A., Vicca, S., and Janssens, I.: Sensitivity of decomposition rates of soil organic matter with respect to simultaneous changes in temperature and moisture, *Journal of Advances in Modeling Earth Systems*, 7, 335–356, 2015.
- 590 Sierra, C. A., Müller, M., Metzler, H., Manzoni, S., and Trumbore, S. E.: The muddle of ages, turnover, transit, and residence times in the carbon cycle, *Global Change Biology*, 23, 1763–1773, 2017.
- Sierra, C. A., Mueller, M., and Trumbore, S. E.: Models of soil organic matter decomposition: the SoilR package, version 1.0, *Geoscientific Model Development*, 5, 1045–1060, <https://doi.org/10.5194/gmd-5-1045-2012>, 2012.
- Sierra, C. A., Mueller, M., and Trumbore, S. E.: Modeling radiocarbon dynamics in soils: SOILR version 1.1, *Geoscientific Model Development*, 7, 1919–1931, <https://doi.org/10.5194/gmd-7-1919-2014>, 2014.
- 595 Sierra, C. A., Ceballos-Nunez, V., Metzler, H., and Mueller, M.: Representing and Understanding the Carbon Cycle Using the Theory of Compartmental Dynamical Systems, *Journal of Advances in Modeling Earth Systems*, 10, 1729–1734, <https://doi.org/10.1029/2018MS001360>, 2018a.
- Sierra, C. A., Hoyt, A. M., He, Y., and Trumbore, S. E.: Soil Organic Matter Persistence as a Stochastic Process: Age and Transit Time Distributions of Carbon in Soils, *Global Biogeochemical Cycles*, 32, 1574–1588, <https://doi.org/10.1029/2018GB005950>, 2018b.
- Steinhof, A., Altenburg, M., and Machts, H.: Sample preparation at the Jena C-14 laboratory, *Radiocarbon*, 59, 815–830, <https://doi.org/10.1017/RDC.2017.50>, 2017.
- Stuiver, M. and Polach, H.: Reporting of C-14 data - Discussion, *Radiocarbon*, 19, 355–363, <https://doi.org/10.1017/S0033822200003672>, 1977.
- 600 Sun, X., Zhao, Y., and Li, Q.: Holocene peatland development and vegetation changes in the Zoige Basin, eastern Tibetan Plateau, *Science China Earth Sciences*, 60, 1826–1837, 2017.
- Tangarife-Escobar, A., Guggenberger, G., Feng, X., Dai, G., Urbina-Malo, C., Azizi-Rad, M., and Sierra, C.: Moisture and temperature effects on the radiocarbon signature of respired carbon dioxide to assess stability of soil carbon in the Tibetan Plateau, <https://doi.org/10.5281/zenodo.7620008>, 2023.
- 610 Tian, Y.-Q., Xu, X.-L., Song, M.-H., Zhou, C.-P., Gao, Q., and Ouyang, H.: Carbon sequestration in two alpine soils on the Tibetan Plateau, *Journal of integrative plant biology*, 51, 900–905, 2009.
- Trumbore, S.: Age of soil organic matter and soil respiration: Radiocarbon constraints on belowground C dynamics, *Ecological Applications*, 10, 399–411, [https://doi.org/10.1890/1051-0761\(2000\)010\[0399:AOSOMA\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2000)010[0399:AOSOMA]2.0.CO;2), 2000.
- Trumbore, S., Chadwick, O., and Amundson, R.: Rapid exchange between soil carbon and atmospheric carbon dioxide driven by temperature change, *Science*, 272, 393–396, <https://doi.org/10.1126/science.272.5260.393>, 1996.
- 615 von Lütow, M., Kögel-Knabner, I., Ludwig, B., Matzner, E., Flessa, H., Ekschmitt, K., Guggenberger, G., Marschner, B., and Kalbitz, K.: Stabilization mechanisms of organic matter in four temperate soils: Development and application of a conceptual model, *Journal of Plant Nutrition and Soil Science*, 171, 111–124, 2008.
- Wang, M., Yang, G., Gao, Y., Chen, H., Wu, N., Peng, C., Zhu, Q., Zhu, D., Wu, J., He, Y., Tian, J., Zhao, X., and Zhang, Y.: Higher recent peat C accumulation than that during the Holocene on the Zoige Plateau, *Quaternary Science Reviews*, 114, 116–125, <https://doi.org/10.1016/j.quascirev.2015.01.025>, 2015.
- 620 Wieder, W. R., Bonan, G. B., and Allison, S. D.: Global soil carbon projections are improved by modelling microbial processes, *Nature climate change*, 3, 909–912, 2013.
- Xiang, S., Guo, R., Wu, N., and Sun, S.: Current status and future prospects of Zoige Marsh in Eastern Qinghai-Tibet Plateau, *Ecological Engineering*, 35, 553–562, <https://doi.org/10.1016/j.ecoleng.2008.02.016>, 2009.
- 625



- Xu, X., Shi, Z., Chen, X., Lin, Y., Niu, S., Jiang, L., Luo, R., and Luo, Y.: Unchanged carbon balance driven by equivalent responses of production and respiration to climate change in a mixed-grass prairie, *Global Change Biology*, 22, 1857–1866, 2016.
- Yang, G., Chen, H., Wu, N., Tian, J., Peng, C., Zhu, Q., Zhu, D., He, Y., Zheng, Q., and Zhang, C.: Effects of soil warming, rainfall reduction and water table level on CH₄ emissions from the Zoige peatland in China, *Soil Biology and Biochemistry*, 78, 83–89, <https://doi.org/10.1016/j.soilbio.2014.07.013>, 2014.
- 630 Yang, Z., Zhu, D., Liu, L., Liu, X., and Chen, H.: The effects of freeze-thaw cycles on methane emissions from peat soils of a high altitude peatland, *Frontiers in Earth Science*, p. 452, 2022.
- Yao, T., Xue, Y., Chen, D., Chen, F., Thompson, L., Cui, P., Koike, T., Lau, W. K.-M., Lettenmaier, D., Mosbrugger, V., Zhang, R., Xu, B., Dozier, J., Gillespie, T., Gu, Y., Kang, S., Piao, S., Sugimoto, S., Ueno, K., Wang, L., Wang, W., Zhang, F., Sheng, Y., Guo, W., Ailikun, Yang, X., Ma, Y., Shen, S. S. P., Su, Z., Chen, F., Liang, S., Liu, Y., Singh, V. P., Yang, K., Yang, D., Zhao, X., Qian, Y., Zhang, Y., and Li, Q.: Recent Third Pole's Rapid Warming Accompanies Cryospheric Melt and Water Cycle Intensification and Interactions between Monsoon and Environment: Multidisciplinary Approach with Observations, Modeling, and Analysis, *Bulletin of the American Meteorological Society*, 100, 423–444, <https://doi.org/10.1175/BAMS-D-17-0057.1>, 2019.
- 635 Zhang, G., Zhang, Y., Dong, J., and Xiao, X.: Green-up dates in the Tibetan Plateau have continuously advanced from 1982 to 2011, *Proceedings of the National Academy of Sciences of the United States of America*, 110, 4309–4314, <https://doi.org/10.1073/pnas.1210423110>, 2013.
- Zhao, Y., Yu, Z., and Zhao, W.: Holocene vegetation and climate histories in the eastern Tibetan Plateau: controls by insolation-driven temperature or monsoon-derived precipitation changes?, *Quaternary Science Reviews*, 30, 1173–1184, <https://doi.org/10.1016/j.quascirev.2011.02.006>, 2011.
- 645 Zhou, W., Cui, L., Wang, Y., Li, W., and Kang, X.: Carbon emission flux and storage in the degraded peatlands of the Zoige alpine area in the Qinghai–Tibetan Plateau, *Soil Use and Management*, 37, 72–82, 2021.