

Microphysical modelling of aerosol scavenging by different types of clouds.

Description and validation of the approach

Pascal Lemaître¹, Arnaud Quérel², Alexis Dépée¹, Alice Guerra Devigne¹, Marie Monier^{3,4}, Thibault Hiron³, Chloé Soto Minguez¹, Daniel Hardy⁵, Andrea Flossmann^{3,4}

[1] { Institut de Radioprotection et de Sûreté Nucléaire (IRSN), PSN-RES/SCA/LPMA, F-91400, Saclay, France}

[2] { Institut de Radioprotection et de Sûreté Nucléaire (IRSN), PSE-SANTE/SESUC/BMCA, F-92260, Fontenay-aux-Roses, France}

[3] {Clermont University, Blaise Pascal University, Physical Meteorology Laboratory, Clermont-Ferrand, France}

[4] {CNRS, INSU, UMR 6016, LaMP, Aubière, France}

[⁵] {School of Chemistry, University of Bristol, Bristol, BS8 1TS, United Kingdom.}

Correspondence to: P. Lemaître (pascal.lemaître@irsn.fr & arnaud.querel@irsn.fr)

Short abstract

A new in-cloud scavenging scheme is proposed. It is based on a microphysical model of cloud formation and may be applied to long-distance atmospheric transport models (>100 km) and climatic models. This model is applied to for the two most extreme precipitating cloud types, in terms of both relative humidity and vertical extension: cumulonimbus and stratus.

Abstract

With dry deposition and below-cloud scavenging, in-cloud scavenging is one of the three components of aerosol transfer from the atmosphere to the ground. There is no experimental validation of in-cloud particle scavenging models for all cloud types that is not impacted by uncertainties concerning below-cloud scavenging. In this article, the choice was made to start with a recognised and validated microphysical cloud formation model (DESCAM) to extract a scheme of aerosol scavenging by clouds, valid for different cloud types. The resulting model works for the two most extreme precipitation clouds: from cumulonimbus to stratus. It is based on data accessible a priori from Numerical Weather Prediction (NWP) outputs, i.e., the intensity of the rain and the relative humidity in the cloud. The description of the altitude of the cloud base proves to be a key parameter, and accuracy in this regard is vital. This new in-cloud scavenging scheme is intended for use in long-distance (> 100 km) Atmospheric Transport Models (ATMs) or Global Climate Models (GCMs).

Introduction

Clouds are an essential component of the troposphere. They play a central role in meteorological forecasting and in the water cycle on the planet (Zhang *et al.*, 2020). Similarly, by interacting with solar radiation, they make a significant contribution to the terrestrial radiation balance (Twomey, 1974; Wang and Su, 2013). Moreover, they are often cited as one of the main sources of uncertainty in climate prediction models (Bony and Dufresne, 2005; Palmer, 2014). They can seriously disrupt air traffic, and even produce aircraft crashes (e. g. Air France flight 447 Rio-Paris air disaster).

By scavenging aerosols, they contribute to improve air quality (Leaith *et al.*, 1987; Sievering *et al.*, 1984), but also to impact soil pollution, through the deposition of atmospheric pollutants via precipitation (Clark and Smith, 1988; Flossmann, 1998). In case of severe nuclear accidents, radioactive aerosol particles might be released into the troposphere (De Cort, 1998 ; Baklanov and Sørensen, 2001 ; Adachi *et al.*, 2013). When radionuclides are emitted into the environment, it is essential, in order to protect populations, to jointly assess the concentrations of radioactive aerosols in the atmospheric boundary layer, as well as their transfer to the ground. Thus, during and after the occurrence of an accident, it is necessary to accurately assess the exposures of populations, both by inhalation and by ingestion (Mathieu *et al.*, 2004; Quélo *et al.*, 2007; Quérel *et al.*, 2012).

In nature, deposition of aerosols (and therefore a fortiori of particulate radionuclides) on the ground consists of the contribution of dry deposition and of wet deposition (Slinn 1977). Dry deposition is approximately 1000 times less effective than wet deposition but is the only mechanism operating when there is no precipitation. To date, there are still many uncertainties about the modelling of these two deposition pathways (Petroff *et al.*, 2008, Croft *et al.*, 2010 and Ervens, 2015).

44 Flossmann (1998) used the DESCAM model (Flossmann *et al.*, 1985, 1987, 1988) to assess that, for a droplet from a convective cloud, about 70% of the mass
 45 of particles the droplet contains when deposited on the ground was incorporated into the droplet in the cloud. This result is consistent with the
 46 measurements in the environment of Laguionie *et al.*, (2014), which estimate the cloud to be 60% responsible for the total downwash of particles.

47 Our objective in this article is to establish theoretically a scavenging coefficient applicable to clouds. **Scavenging by clouds is more challenging to model than**
 48 **scavenging by rain, under the cloud, as it is much more sensitive to certain input parameters.** Rain scavenging is only controlled by a single microphysical
 49 mechanism: collection by raindrops (Kerker and Hampl, 1974; Beard, 1974; Grover *et al.*, 1977; Wang and Pruppacher, 1977; Lai *et al.*, 1978; Pranesha and
 50 Kamra, 1996; Vohl *et al.*, 1999; Quérel *et al.*, 2014; Lemaitre *et al.*, 2017), whereas cloud scavenging encompasses a set of mechanisms which will firstly
 51 make it possible to incorporate aerosols into the cloud droplets (activation, collection, ice nucleation, collection by crystals) then, secondly, to convert a
 52 fraction of the cloud hydrometeors into raindrops (condensation, coalescence, Bergeron effects). Only after raindrops have been deposited on the ground
 53 is the atmosphere washed out and, by **the same process**, the soil contaminated. Furthermore, **most** atmospheric transport models use significantly different
 54 schemes to model scavenging by cloud and by rain. Quérel *et al.*, 2021 summarised **a few of them** in Table 3 of their article.

55 **Therefore, in theoretically assessing cloud scavenging, the use of a cloud formation model such as DESCAM (DEtailed SCAvenging Model), forms a good**
 56 **foundation.** This model, developed by Andrea Flossmann and her group since the mid-80s (Flossmann *et al.*, 1985; Flossmann, 1998; Monier *et al.*, 2006;
 57 Leroy *et al.*, 2007; Planche *et al.*, 2007; Flossmann and Wobrock, 2010; Hiron and Flossmann, 2015; Dépée, 2019), makes it possible, through a detailed
 58 microphysical description, to model clouds from their formation through to precipitation, and to monitor the aerosols and what becomes of them once
 59 incorporated into the droplets.

60 In this article, we will show how, using a model like DESCAM, it is possible to theoretically calculate a scavenging coefficient in the cloud, on the scale of the
 61 cloud system. We will apply this approach to two extreme types of cloud: a cumulonimbus (CCOPE, Dye *et al.*, 1986) and a stratus (Zhang *et al.*, 2004). This
 62 approach will then be compared to the models derived from the deposits observed following the Fukushima nuclear accident (Leadbetter *et al.*, 2015; Quérel
 63 *et al.*, 2021). Finally, in the last part **we will present**, a theoretical scheme of the scavenging coefficient, applicable to any type of cloud, will be proposed.
 64 **We** begin by considering some **key** elements of **the** theoretical context and some definitions.

65 **1. Definitions and theoretical context**

66 **1.1. Definition of cloud scavenging**

67 In large range transport models, the description of scavenging shall remain simple, the operational scientific community models it through a parametrisation
 68 involving the cloud scavenging coefficient (Λ_{cloud}). It is defined as the fraction of pollutants that is transferred from the atmosphere to precipitation (then
 69 to the ground) per unit of time. In this article, we will focus on pollutants carried by aerosols (and not gaseous pollutants). The scavenging coefficient is
 70 therefore defined spectrally thus:

$$71 \quad \left. \frac{d\mathcal{N}(d_{ap})}{\mathcal{N}(d_{ap})} \right|_{cloud} = \left. \frac{d\mathcal{M}(d_{ap})}{\mathcal{M}(d_{ap})} \right|_{cloud} = -\Lambda_{cloud}(d_{ap}) \cdot dt \quad \text{Equation 1}$$

72 In this equation $\mathcal{N}(d_{ap})$ and $\mathcal{M}(d_{ap})$ are respectively the concentrations in number and in mass of aerosols of diameter d_{ap} per unit of air volume;
 73 likewise, $d\mathcal{N}(d_{ap})$ and $d\mathcal{M}(d_{ap})$ are respectively the variations in concentration in number and in mass of aerosols of diameter d_{ap} , in relation to their
 74 transfer into precipitation, per unit of time. **The two definitions are considered equal when expressed spectrally, assuming a uniform density of all aerosol**
 75 **particles. In this study, we specifically considered aerosol particles composed of ammonium sulphate, thus confirming this equality. However, in real**
 76 **atmospheric conditions, aerosol particle density is typically not uniform. In such cases, it becomes crucial to specify whether we are referring to a mass**
 77 **scavenging coefficient or a numerical one. To prevent any confusion, in the following article, the exponent m is introduced to the scavenging coefficient**
 78 **Λ_{cloud}^m .**

79 **The approach we use** is to apply this definition to an elementary volume of cloud (volume outlined in red in Figure 1). This volume is bounded at its base by
 80 an arbitrary section (dS) aligned with the base of the cloud, with this volume extending vertically to the cloud summit.




Figure 1. Definition of the scavenging coefficient at the scale of a cloud

81

82

83

84 In this elementary cloud volume, it is elementary to calculate the variation in the average mass concentration of aerosols of diameter d_{ap} , in relation to
85 their transfer into precipitation:

86

87

$$d\mathcal{M}(d_{ap}) = - \frac{\phi_{ap,precip}(d_{ap}) \cdot dS \cdot dt}{dV_{cloud}} \quad \text{Equation 2}$$

88

89 In this equation, $\phi_{ap,precip}(d_{ap})$ is the mass flow of dry particles of diameter (d_{ap}) leaving the cloud via precipitation (solids and liquids), and dV_{cloud} is the
90 elementary volume of cloud considered ($dV_{cloud} = H_{cloud} \cdot dS$).

$$\Lambda_{cloud}^m(d_{ap}) = - \frac{1}{dt} \frac{d\mathcal{M}(d_{ap})}{\langle \mathcal{M}(d_{ap}) \rangle_{cloud}} \Big|_{cloud} = \frac{\phi_{ap,precip}(d_{ap})|_{dS}}{\langle \mathcal{M}(d_{ap}) \rangle_{cloud} \cdot H_{cloud}} \quad \text{Equation 3}$$

91 In this equation, $\langle \mathcal{M}(d_{ap}) \rangle$ is the average mass concentration (over the thickness of the cloud) of dry particles of diameter d_{ap} . **It is noteworthy that**
92 $\langle \mathcal{M}(d_{ap}) \rangle$ is not the average concentration of interstitial aerosols in the cloud but the average concentration of particles, which includes, in addition to the
93 interstitial aerosols, all the particles included in the droplets and potentially in the ice phase. Thus, if we jointly determine $\langle \mathcal{M}(d_{ap}) \rangle$, $\phi_{ap,precip}$ as well
94 as the thickness of the cloud H_{cloud} , it is possible to deduce a scavenging coefficient.

95 The average particle concentration is calculated, using Equation 4, by spatially averaging, over the entire thickness of the cloud, the concentrations of
96 interstitial aerosols (of diameters d_{ap}) $\mathcal{M}_{int}(z, d_{ap})$, the concentrations of particles in the drops ($\mathbb{M}(z, d_{ap})$), and the concentrations in the ice phase
97 ($\mathfrak{M}(z, d_{ap})$).

$$\langle \mathcal{M}(d_{ap}) \rangle = \frac{1}{H_{cloud}} \int_{cloud\ base}^{cloud\ summit} (\mathcal{M}_{int}(z, d_{ap}) + \mathbb{M}(z, d_{ap}) + \mathfrak{M}(z, d_{ap})) dz \quad \text{Equation 4}$$

98

99 Finally, in order to evaluate the mass flow of particles exiting the cloud at its base ($\phi_{ap,precip}(d_{ap})$), it is necessary to evaluate the cloud volume
100 ($\mathcal{V}(\mathcal{D}_{drop})$) which contains all the droplets whose drop velocity $w_{\infty}(\mathcal{D}_{drop})$ is sufficient for them to pass through the section dS , during the time dt . Using
101 the velocity composition law, we can deduce Equation 5.

$$\mathcal{V}(\mathcal{D}_{drop}) = \max(0, w_{\infty}(\mathcal{D}_{drop}) - w_{air}(Z_{cloud\ base})) \cdot dS \quad \text{Equation 5}$$

102

103 In this equation, $w_{air}(Z_{cloud\ base})$ is the velocity of the air parcel at the base of the cloud. By convention, w_{air} is positive for an updraft and negative for
104 a downdraft.

105 It is then immediately possible to deduce the flow of particles passing dS through liquid precipitation:

$$\phi_{ap,rain}(d_{ap}) = \int_{d_{droplet}=0}^{\infty} \max(0, w_{\infty}(\mathcal{D}_{drop}) - w_{air}(Z_{cloud\ base})) \cdot \mathbb{M}(z, d_{ap}, \mathcal{D}_{drop}) \cdot d\mathcal{D}_{drop} \quad \text{Equation 6}$$

106 In this equation, $\mathbb{M}(z, d_{ap}, \mathcal{D}_{drop})$ is the concentration of particles contained in the droplets and of dry diameter d_{ap} .

107 The same applies to solid precipitation $\phi_{ap, ice}(d_{ap})$:

$$\phi_{ap, ice}(d_{ap}) = \int_{d_{ice}=0}^{\infty} \max(0, w_{\infty}(d_{ice}) - w_{air}(Z_{cloud\ base})) \cdot \mathfrak{M}(z, d_{ap}, d_{ice}) \cdot d_{ice} \quad \text{Equation 7}$$

108 By adding these two flows together, it is possible to deduce the total flow of particles (of diameter d_{ap}) exiting the cloud through all precipitation:

$$\phi_{ap, precip}(d_{ap}) = \phi_{ap, rain}(d_{ap}) + \phi_{ap, ice}(d_{ap}) \quad \text{Equation 8}$$

110

111 Thus, to theoretically evaluate the cloud scavenging coefficient, it is first and foremost essential to be able to evaluate its contours, but also to be able to
112 determine the mass concentrations of particles in the droplets $\mathbb{M}(z, d_{ap}, \mathcal{D}_{drop})$, in the ice phase $\mathfrak{M}(z, d_{ice}, d_{ap})$, and in the interstitial aerosol $\mathcal{M}_{int}(z, d_{ap})$.

113 To evaluate the contours of the cloud, it seems necessary in the first instance to consider once again its definition.

114 **1.2. What is a cloud? (how to define its boundaries?)**

115 The World Meteorological Organization defines clouds as: “an aggregation of minute particles of liquid water or ice, or of both, suspended in the atmosphere
116 and usually not touching the ground” (WMO, 2014). This definition would appear to be very inadequate for enabling the contours of a cloud to be
117 determined. Clouds, although very commonly talked about in everyday life and subject to numerous scientific studies, have contours that remain very
118 blurred. It is therefore always difficult to define them rigorously, and above all non-recursively. Spankuch *et al.*, (2022) further emphasised that, depending
119 on the scope of the authors’ expertise (meteorology, climate, satellite observations, airborne, from ground radars, or using microphysical models), these
120 authors use significantly differing definitions and thresholds.

121 For example, Wood and Field, 2011 proposed criteria with respect to liquid water content (LWC), ice water content (IWC), or total concentrations in numbers
122 of hydrometeors (droplets and crystals).

123 Hiron (2018) proposed separating cloud water and precipitation water based on the criterion of the size of hydrometeors (hydrometeors with a diameter of
124 less than 64 μm are considered part of the cloud; larger than that are considered part of the rain); then, if the total cloud water content is greater than 0.1
125 g/cm^3 , the air parcel is considered part of the cloud.

126 Other authors proposed contours based on relative humidity (Del Genio *et al.*, 1996) or total water content (TWC), whereas meteorologists and
127 climatologists tend to prefer optical thickness (Sassen and Cho, 1992), each with arbitrarily established thresholds.

128 Although these definitions can be linked mathematically to each other, these relationships are most often highly non-linear. **Therefore, in this article, a
129 variety of cloud definitions will be considered**, and we will examine the criteria that are most relevant for studying in-cloud scavenging and distinguishing it
130 from below-cloud scavenging. This relevance **of the definition what constitutes a cloud** will be analysed from two perspectives. Firstly, we consider a purely
131 physical **perspective** and, secondly, a more pragmatic **perspective** linked rather to applicability in an atmospheric dispersion model dedicated to crisis
132 management.

133 **1.3. The DESCAM model**

134 To simulate clouds of different types and theoretically evaluate their scavenging coefficient, it is necessary to have a model that makes it possible to simulate
135 all the water phase changes, considering the catalyst role of aerosols in most of these state changes (activation, ice nucleation, etc.). It is also necessary to
136 calculate the sink terms of interstitial aerosols (related to droplet collection or activation) and associate them with the source terms of particles in droplets
137 and ice, in order to calculate the mass of particles in droplets ($\mathbb{M}(d_{drop})$) and in ice ($\mathfrak{M}(d_{ice})$) throughout the simulation (Equation 6, Equation 7).

138 The DESCAM model meets these specifications. This detailed microphysical model classifies droplets ($\mathcal{D}_{drop} \in [1 \mu\text{m}, 6.5 \text{mm}]$), ice ($d_{ice} \in$
139 $[1 \mu\text{m}, 6.5 \text{mm}]$), and aerosols ($d_{ap} \in [2 \text{nm}, 12.7 \mu\text{m}]$), each into 39 logarithmically distributed size classes. This makes it possible to explicitly monitor
140 their respective particle size distributions $\mathbb{N}(\mathcal{D}_{drop})$, $\mathfrak{N}(d_{ice})$ and $\mathcal{N}(d_{ap})$, spatially and temporally. This model can be coupled with various dynamic models
141 that allow consideration of atmospheric flows. In this article, we will only consider a dynamic called 1D1/2 (Asai and Kasahara, 1967), implemented in the
142 DESCAM model by Monier, 2003. More realistic 3D dynamics (Clark and Hall, 1991) are implemented in DESCAM (Leroy 2007) but will not be considered in
143 this article.

144 **Description of the microphysical models modelled in DESCAM**

145 All the microphysical processes considered in the DESCAM model are presented in Figure 2.




Figure 2. Modelling of microphysical processes in the DESCAM model.

In this figure, we can see the central role of aerosols in most water phase changes. The explicit resolution of all these microprocesses enables calculation of the particle size distributions of aerosols in each grid cell and at each time step (in number: $\mathcal{N}(d_{ap})$ (Equation 9) and in mass: $\mathcal{M}(d_{ap})$), as well as the particle size distributions of the droplets (in number $\mathcal{N}(\mathcal{D}_{drop})$, Equation 10) and of the ice (in number $\mathcal{N}(\mathcal{d}_{ice})$, Equation 11). In addition, in order to preserve the total mass of particles, the model also calculates two other quantities that are used to determine the masses of particles in droplets of diameter d_{drop} ($\mathcal{M}(\mathcal{D}_{drop})$), and in ice crystals of diameter d_{ice} ($\mathcal{M}(\mathcal{d}_{ice})$).

$$\frac{d\mathcal{N}(d_{ap})}{dt} = \frac{d\mathcal{N}(d_{ap})}{dt}\Big|_{dyn} + \frac{d\mathcal{N}(d_{ap})}{dt}\Big|_{coll} + \frac{d\mathcal{N}(d_{ap})}{dt}\Big|_{hygro} + \frac{d\mathcal{N}(d_{ap})}{dt}\Big|_{act, deact} \quad \text{Equation 9}$$

$$\begin{aligned} \frac{d\mathcal{N}(\mathcal{D}_{drop})}{dt} = & \frac{d\mathcal{N}(\mathcal{D}_{drop})}{dt}\Big|_{dyn} + \frac{d\mathcal{N}(\mathcal{D}_{drop})}{dt}\Big|_{act, deact} + \frac{d\mathcal{N}(\mathcal{D}_{drop})}{dt}\Big|_{coal} \\ & + \frac{d\mathcal{N}(\mathcal{D}_{drop})}{dt}\Big|_{cond, vap} + \frac{d\mathcal{N}(\mathcal{D}_{drop})}{dt}\Big|_{riming} + \frac{d\mathcal{N}(\mathcal{D}_{drop})}{dt}\Big|_{frz, melt} \end{aligned} \quad \text{Equation 10}$$

$$\begin{aligned} \frac{d\mathcal{N}(\mathcal{d}_{ice})}{dt} = & \frac{d\mathcal{N}(\mathcal{d}_{ice})}{dt}\Big|_{dyn} + \frac{d\mathcal{N}(\mathcal{d}_{ice})}{dt}\Big|_{nucleation} + \frac{d\mathcal{N}(\mathcal{d}_{ice})}{dt}\Big|_{frz, sub} + \frac{d\mathcal{N}(\mathcal{d}_{ice})}{dt}\Big|_{agg} \\ & + \frac{d\mathcal{N}(\mathcal{d}_{ice})}{dt}\Big|_{riming} + \frac{d\mathcal{N}(\mathcal{d}_{ice})}{dt}\Big|_{frz, melt} \end{aligned} \quad \text{Equation 11}$$

$$\begin{aligned} \frac{d\mathcal{M}(\mathcal{D}_{drop})}{dt} = & \frac{d\mathcal{M}(\mathcal{D}_{drop})}{dt}\Big|_{dyn} + \frac{d\mathcal{M}(\mathcal{D}_{drop})}{dt}\Big|_{act, deact} + \frac{d\mathcal{M}(\mathcal{D}_{drop})}{dt}\Big|_{coal} \\ & + \frac{d\mathcal{M}(\mathcal{D}_{drop})}{dt}\Big|_{cond, vap} + \frac{d\mathcal{M}(\mathcal{D}_{drop})}{dt}\Big|_{riming} + \frac{d\mathcal{M}(\mathcal{D}_{drop})}{dt}\Big|_{frz, melt} \end{aligned} \quad \text{Equation 12}$$

$$\begin{aligned} \frac{d\mathcal{M}(\mathcal{d}_{ice})}{dt} = & \frac{d\mathcal{M}(\mathcal{d}_{ice})}{dt}\Big|_{dyn} + \frac{d\mathcal{M}(\mathcal{d}_{ice})}{dt}\Big|_{nucleation} + \frac{d\mathcal{M}(\mathcal{d}_{ice})}{dt}\Big|_{frz, sub} + \frac{d\mathcal{M}(\mathcal{d}_{ice})}{dt}\Big|_{agg} \\ & + \frac{d\mathcal{M}(\mathcal{d}_{ice})}{dt}\Big|_{riming} + \frac{d\mathcal{M}(\mathcal{d}_{ice})}{dt}\Big|_{frz, melt} \end{aligned} \quad \text{Equation 13}$$

In these equations, except for the index ($|_{dyn}$) denoting variations due to atmospheric transport, each term corresponds to one of the microphysical processes outlined in Figure 2. For instance, ($|_{act, deact}$) denotes to activation and deactivation processes. The subscripts *coll*, *hygro*, *frz*, *sub*, *coal*, *cond*, *vap* and *agg* respectively refers to the processes of aerosol collection by droplets, hygroscopicity of aerosol particles, freezing, sublimation,

163 coalescence, condensation, vaporization, and aggregation. The hygroscopic growth of aerosol particles is calculated assuming they are in thermodynamic
 164 equilibrium with the air supersaturation S (Equation 14). This equilibrium is modelled after the κ -Köhler theory.

$$\left. \frac{d\mathcal{N}(d_{ap})}{dt} \right|_{hygro} = \left. \frac{\partial \mathcal{N}(d_{ap})}{\partial S} \right|_{\kappa\text{-Köhler}} \left. \frac{\partial S}{\partial t} \right|_{dyn} \quad \text{Equation 14}$$

165

166 Concerning the cold microphysics (Figure 2), this simulation integrates homogeneous freezing mechanisms (i.e., which do not require aerosol contribution)
 167 and heterogeneous freezing mechanisms (for which aerosols act as a catalyst for phase change). For homogeneous freezing, we consider the parametrisation
 168 of Koop *et al.*, (2000) adapted to DESCAM by Monier *et al.*, (2006). To model heterogeneous ice nucleation, we consider all the mechanisms described by
 169 Vali *et al.*, (2015). The Biggs formula (1953) is used to describe immersion freezing and the model of Meyers *et al.*, (1992) for condensation and contact
 170 freezing, as well as deposition nucleation. All these mechanisms have recently been incorporated into the DESCAM model by Hiron and Flossmann, (2015).


171 The main mechanism responsible for the flow of particles exiting the cloud via precipitation ($\phi_{ap, precip}(d_{ap})$, Equation 3) is activation. The collection of
 172 aerosols by droplets is only second order (Flossmann and Wobrock, 2010; Dépée, 2019). In the DESCAM model, activation is modelled by the κ -Köhler theory
 173 (Petters and Kreidenweis, 2007). This model makes it possible to determine equilibrium vapour pressure in the vicinity of a droplet of diameter \mathcal{D}_{drop} , as a
 174 function of the mass and type of solute (modelled by the κ value) it contains, and therefore the supersaturation for this droplet. For a given mass and
 175 chemical nature of the pristine dry particle, one can compute the corresponding supersaturations for given size of solution droplets. This curve has a unique
 176 maximum, called critical supersaturation (Figure 3). The diameter associated with this critical supersaturation is called the activation diameter. Aerosols
 177 with a diameter smaller than the activation diameter are brought into thermodynamic equilibrium with their environment by hygroscopicity, with aerosols
 178 of a diameter greater than the activation diameter being converted into droplets and growing by means of vapour diffusion (by condensation).

179 In the DESCAM code, the microphysical process of collection (i.e. the process by which, during falling, droplets encounter impact and capture interstitial
 180 aerosol particles), is modelled in Equation 12 by the term $\left. \frac{d\mathbb{M}(\mathcal{D}_{drop})}{dt} \right|_{coll}$ which is calculated by solving Equation 15. In this equation, the central term is the
 181 collection efficiencies ($E(d_{ap}, \mathcal{D}_{drop}, \mathcal{RH})$). This is calculated by the model developed and validated by Dépée (Dépée *et al.*, 2019; Dépée *et al.*, 2021, Part I;
 182 Dépée *et al.*, 2020, Part II).

$$\left. \frac{d\mathbb{M}(\mathcal{D}_{drop})}{dt} \right|_{coll} = \int_{d_{ap}=0}^{\infty} \mathcal{N}(d_{ap}) \frac{\pi \mathcal{D}_{drop}^2}{4} \mathcal{U}_{\infty, droplet}(\mathcal{D}_{drop}) E(d_{ap}, \mathcal{D}_{drop}, \mathcal{RH}) \frac{\pi d_{ap}^3}{6} \rho_{ap} dd_{ap} \quad \text{Equation 15}$$

183 In this equation $\mathcal{U}_{\infty, droplet}(\mathcal{D}_{drop})$ corresponds to the terminal velocity of a droplet of diameter \mathcal{D}_{drop} (calculated after the Beard (1973) model), ρ_{ap} is
 184 the density of the aerosol particle, and \mathcal{RH} refers to the relative humidity of air in the parcel.

185



186

187 Figure 3. Thermodynamic equilibrium of an aerosol calculated as a function of particle diameter and nature of the initial dry particle. This
 188 calculation is made using the κ -Köhler theory (for a temperature of 293 K and a surface tension between solution and air of $72 \times 10^{-3} \text{ N.m}^{-1}$).

189 (—) initial dry radius of the particles set at 4 nm; (—) initial dry radius set at 20 nm; (—) initial dry radius set at 100 nm;
 190 (···) $\kappa = 0$, insoluble aerosol; (--) $\kappa = 0.61$, moderately hygroscopic, as $(\text{NH}_4)_2\text{SO}_4$; (—) $\kappa = 1.28$, highly hygroscopic, as NaCl;
 191 ★ critical supersaturations needed to activate the aerosol and convert it into a cloud droplet.

192

193 Note that Equation 6 and Equation 7 are not directly calculable by the DESCAM model. This is because the model makes an inventory of the mass of particles
194 in the droplets and crystals according to the size of the hydrometeors ($M(\mathcal{D}_{drop})$, Equation 12 and $\mathfrak{M}(\mathbb{d}_{ice})$, Equation 13), but without memorising the size
195 of the aerosols before their incorporation. The scavenging coefficients calculated in this article are therefore averaged, in mass, over the particle size
196 distribution of the aerosols. **In this article we focus on validating the approach described by applying it to different types of cloud. Current work and future
197 publications will focus on demonstrating that the scavenging coefficient may be simply spectrally calculated, without modifying the model.**

198

199 **Modelling of atmospheric dynamics**

200 As stated previously, in this article we have limited our study to the 1.5D Dynamic framework developed by Asai and Kasahara, (1967). This has been regularly
201 used (Monier *et al.*, 2005; Leroy *et al.*, 2006; Quérel *et al.*, 2014; Hiron & Flossmann, 2015) to study the microphysical processes involved in the life cycle of
202 cumulus clouds. This model considers two concentric cylinders. The inner cylinder has a radius 10 times smaller than the outer cylinder. In the inner cylinder,
203 the vertical velocity of the flows is determined by solving a simplified form of the Navier-Stokes equations, coupled with the energy conservation equation.
204 The outer cylinder serves primarily for guaranteeing the condition of zero velocity divergence (continuity equation for incompressible flow). To this end, a
205 radial velocity component is introduced at the interface between these two cylinders (hence the expression of a 1.5D model), **calculated** from the
206 convergence or divergence layers and allowing entrainment from the environment. In this environment the only variable updated in this outer cylinder is
207 the vertical velocity to evaluate the radial gradient in vertical velocity and the subsequent turbulent flux; all the other variables are assumed to be unaffected
208 by the cloud processes within the inner cylinder and kept constant throughout the simulation.

209 All the microphysical processes detailed in the previous section and summarised in Figure 2 are calculated only in the central cylinder. Thus, this is also in
210 the inner cylinder, for each grid layer, that phase changes in the water will be computed with the subsequent absorption or release of latent heat that will
211 alter the buoyancy of the air and which ultimately generate the updraft and downdraft motions.

212 **2. Application to example cloud types**

213 To establish a theoretical scavenging coefficient scheme, the methodology **described above** is applied to two very different idealised case studies
214 representative of two different types of clouds. First, we will model a vigorous cumulonimbus, then a shallow stratus. These two cloud types were selected
215 as they present, respectively, the higher and lower values, in terms of vertical extension, relative humidity, and rainfall intensity. Furthermore, while the
216 stratus that we simulate is **thin and low altitude** enough to be a warm cloud, cold microphysical processes are essential to capture the development of the
217 cumulonimbus situation.

218

219 **2.1. Application to a cumulonimbus**

220 **Description of the cumulonimbus considered**

221 The cloud selected to model the cumulonimbus is the episode of 19 July 1981 of the CCOPE campaign (Cooperative Convective Precipitation Experiment;
222 Knight 1982 and Dye *et al.*, 1986), which took place near Miles City in Montana (USA). This episode was selected because it is very finely documented and
223 this episode is a test case for many codes simulating the formation of convective clouds, and in particular the DESCAM model (Flossmann & Wobrock, 2010).
224 A radio sounding taken at Miles City at 16h00 local time (just before the storm) is used to initialise the thermodynamic conditions of the atmospheric column:
225 temperature and humidity. For the vertical pressure profiles, standard conditions are assumed. In addition, we used the observations of both two Doppler
226 radars measured high-resolution reflectivity as well as the movements of the cloud and five aircrafts that were able to make numerous passes through the
227 cloud throughout its maturation and through to the precipitation episodes. The spatial-temporal evolution of the thermodynamic conditions, associated
228 with the microphysical properties of the cloud system, and the atmospheric flows are therefore recorded in fine detail for the entire life of this cumulonimbus
229 and can be used to evaluate the model performance to capture the clouds physics. For this article, we therefore used the same modelling hypotheses as
230 those detailed by Leroy *et al.*, (2006). Convection was triggered by +2.3°C heating of the ground during the first 10 minutes of the simulation. During this
231 campaign, no physicochemical measurements were made of the aerosols, hence we consider that they consisted of ammonium **sulfate** ($\kappa = 0.61$ and $\rho_{dry} =$
232 $1.77 \cdot 10^3 \text{ kg} \cdot \text{m}^{-3}$ Petters and Kreidenweis, 2007) with an initial particle size distribution of the Jaenicke continental type (1988). We considered a homogeneous
233 distribution in the atmospheric boundary layer (i.e., over the first 3 kilometres), then above the concentration is assumed to decrease exponentially with a
234 scale height of 3000 m (Figure 4).





Figure 4. Initial particle size distribution of aerosols considered for this simulation.

235

236

237 The simulation lasted 3600 s, on a 10 km high column. The spatial and temporal resolutions were set to 100 m and three seconds respectively.

238 Figure 5 and Figure 6 show respectively the spatial-temporal evolutions of the vertical flows in the central cylinder and the liquid water and ice content.




239

240 Figure 5. Spatial-temporal distribution of the vertical components of atmospheric flows. The thick line in red separates the updraft ($w_{air} > 0$) flows from
241 the downdraft flows ($w_{air} < 0$).

242

243 In these figures, we can see that the initial superheating of the air layer at ground level induces an updraft flow due to buoyancy forces. Approximately 500 s
244 after the start of the simulation, the air reaches critical supersaturation at an altitude of 3000 m and the aerosols are gradually converted into droplets. The
245 spatial-temporal distribution shown on the left of Figure 6 highlights the appearance of a cloud at the spatial-temporal coordinate (500 s, 3000 m). Vapour
246 condensation induces a latent heat release, which in turn increases buoyancy of the air parcel, accelerating the updraft flows (approx. 15 m.s^{-1} at 4000 m).
247 This flow transports the vapour at altitude and by cooling this induces the progressive activation of the aerosols, and a vertical extension of the cloud. Near
248 7000 m, the first ice crystals are formed. The coexistence of ice crystals and supercooled droplets will allow rapid crystal growth with a corresponding
249 reduction of liquid water content (best known as the **WBF process** for: Wegener, 1911; Bergeron, 1928; Findeisen, 1938). Then, the crystals begin to
250 precipitate at around 1700 s since they reach sizes large enough for their gravitational settling velocity to supersede updraft speed. In precipitating, the
251 larger crystals collect the suspended droplets. Hence, under the coupled influence of the **WBF process** and, above all, the collection of droplets by the ice
252 particles, after 2200 seconds of simulation the cloud only contains ice. Finally, below an altitude of 3000 m, the solid hydrometeors melt and liquid
253 precipitation forms. Figure 6 shows on the right the rainfall intensities, as well as the cumulative precipitation calculated by the model at ground level. This
254 timeline presents two local maxima: the first at 2750 s after the start of the simulation, corresponding to an intensity of 18 mm.h^{-1} and the second 300 s
255 later, with an intensity of about 46 mm.h^{-1} .



256


257 Figure 6. On the left: Spatial-temporal distributions of liquid water content (LWC, greyscale) and ice content (IWC, iso-contours).
 258 On the right: Temporal evolution of rainfall intensity and cumulative precipitation at ground level

259 **Calculation of the scavenging coefficient**

260 Based on this modelling results, we applied all the methodology described in section 1.1. The first step consisted in establishing the contours of the cloud,
 261 in order then, using Equation 3 integrated over the entire aerosol distribution, to be able to calculate the equivalent cloud scavenging coefficient.

$$A_{cloud}^m = - \left. \frac{1}{dt} \frac{d\mathcal{M}}{\langle \mathcal{M} \rangle} \right|_{cloud} = \frac{\Phi_{ap, precip}}{\langle \mathcal{M} \rangle \cdot H_{cloud}} \quad \text{Equation 16}$$

262 As already indicated in section 1.2, there is no strict definition of the boundaries of the cloud, particularly at the interface with the precipitation, and it can
 263 also be observed in Figure 6 that the total water content does not show any demarcation between the cloud and the precipitation. Moreover, as Spänkuch
 264 *et al.*, (2022) point out, the physical phenomenon studied will determine which contours are the most relevant. Therefore, for this study, we examined three
 265 of the physical parameters to establish this contour. These three criteria are the relative humidity of the air parcel (calculated in relation to liquid water),
 266 the mass concentration of hydrometeors with a diameter of less than 64 μm and, lastly, the concentration of hydrometeors. The contours of this
 267 cumulonimbus are presented in Figure 7 for each of these criteria, each with two thresholds considered.



268

269 Figure 7. Test of different criteria to establish the contours of the simulated cumulonimbus.


270 a) threshold based on relative humidity: (---) $RH > 85\%$; (—) $RH > 80\%$.

271 b) threshold based on the total water content of hydrometeors with a diameter less than 64 μm : (---) mass concentration of cloud
 272 hydrometeors $> 0.1 \text{ g.m}^{-3}$; (—) mass concentration of cloud hydrometeors $> 0.001 \text{ g.m}^{-3}$.

273 c) threshold based on number concentration of total hydrometeors: (---) $\int dN + d\mathfrak{N} > 0.03 \text{ cm}^{-3}$
 274 (—) $\int dN + d\mathfrak{N} > 0.003 \text{ cm}^{-3}$


275 We can see in this figure that, apart from the criterion based on the concentration of hydrometeors (Equation 9), with a threshold of 0.003 hydrometeors
 276 per cubic centimetre, all the other criteria yield very similar contours. Thus, the cloud forms close to an altitude of 3000 m and its base remains constant for
 277 up to 2500 s of simulation. During these 2500 s, the cloud thickens vertically until it reaches the tropopause (considered to be at 10,000 m in this calculation).
 278 As shown in Figure 6 (on the right), 2500 s corresponds to the start of precipitation. This moment corresponds to an elevation of the base of the cloud up to
 279 about 7000 m, except for the last criterion ($\mathcal{N}_{hydrometeor} > 0.003 \text{ cm}^{-3}$), for which the height of the base of the cloud remains constant close to the
 280 altitude of 3000 m, even during rain.

281 Initially, our objective was to find a bijective relationship between a set of meteorological parameters available in DESCAM and the scavenging coefficient
 282 calculated by this methodology. Most often in the literature, cloud scavenging is described as a power function of precipitation intensity (Hertel *et al.*, 1995;
 283 MRI, 2015; Leadbetter *et al.*, 2015; Groell *et al.*, 2014; Quérel *et al.*, 2021). Figure 8, Figure 9 and Figure 10 respectively present the contours of the cloud
 284 established on the basis of the three criteria previously introduced (Figure 7). Within these contours, we calculated the total mass concentration of
 285 ammonium sulfate ($\mathcal{M}(z)$), adding together the respective concentrations of the aerosol phases ($\mathcal{M}_{int}(z)$), in the droplets ($\mathcal{M}_l(z)$) and in the crystals
 286 ($\mathcal{M}_c(z)$). Knowing the flux of ammonium sulfate that is within the precipitative hydrometeors through the base of the cloud (Equation 6, Equation 7 and
 287 Equation 8), we could deduce the scavenging coefficient, which we plotted according to the precipitation intensity calculated at the base of the cloud. Like
 288 Costa *et al.*, (2010), Stephan *et al.*, (2008) and Quérel *et al.* (2021), a threshold of 0.1 mm.h^{-1} was considered to limit noise. In Figure 8, 9, 10, the
 289 correspondence of the dots can be deduced with the colour codes of the points. On the left-hand side, the identification of the spatial-temporal coordinates
 290 where precipitation and scavenging coefficient are calculated is plotted. On the right-hand side, the corresponding relationship between scavenging
 291 coefficient and precipitation intensity can be read. These results are of great importance because they show that the relationship between the scavenging
 292 coefficient and the rainfall intensity is the same at the beginning and the end of the rainfall episode. In addition, an adjustment by a power law is determined
 293 for each contour. The coefficients for these adjustments are shown in Table 1.




294

295 Figure 8. On the left: spatial-temporal distribution of the ammonium sulfate concentration in the cloud contour (— cloud contour for a
 296 relative humidity greater than 80%). On the right: correlation between the scavenging coefficient and the precipitation intensity determined
 297 at the base of the cloud, (—) adjusted by a power law. Coloured points indicated where and when the scavenging coefficient is calculated,
 298 colours are time increasing from the start (blue) to the end (red) of the rainfall.



299

300 Figure 9. On the left: spatial-temporal distribution of the ammonium sulfate concentration in the cloud contour (— cloud contour for a mass
 301 concentration of cloud hydrometeors greater than $0.001 \text{ g}\cdot\text{m}^{-3}$). On the right: correlation between the scavenging coefficient and the
 302 precipitation intensity determined at the base of the cloud, (—) adjusted by a power law. Coloured points indicated where and when the
 303 scavenging coefficient is calculated, colours are time increasing from the start (blue) to the end (red) of the rainfall.



304

305 Figure 10. On the left: spatial-temporal distribution of the ammonium sulfate concentration in the cloud contour (— cloud contour for a
 306 concentration in number of hydrometeors greater than $0.003 \text{ particles}\cdot\text{cm}^{-3}$). On the right: correlation between the scavenging coefficient
 307 and the precipitation intensity determined at the base of the cloud, (—) adjusted by a power law. Coloured points indicated where and when
 308 the scavenging coefficient is calculated, colours are time increasing from the start (blue) to the end (red) of the rainfall.

309

310 In these three figures, we observe that the relationship linking the intensity of precipitation to the scavenging coefficient by the cloud is fairly insensitive to
 311 the definition selected to describe its contour. Moreover, the power law adjustments plotted in Figure 8, Figure 9 and Figure 10 are very similar (Table 1).
 312 Nevertheless, only the last contour, based on the hydrometeor concentration (and with a threshold of 0.003 cm^{-3}), gives a perfectly bijective relationship
 313 between the precipitation intensity at the base of the defined contour and the scavenging coefficient. This result is surprising because, as previously
 314 mentioned in section 1.3, the driving mechanism for in-cloud scavenging is dominated by the activation (which is driven by the supersaturation level and
 315 physical-chemical properties of the aerosols, Flossmann and Wobrock, 2010). It would therefore seem logical that a criterion based on the relative humidity
 316 in the grid cell would be the most relevant. However, it is the criterion based on the concentration of hydrometeors that is the more reliable. This is because
 317 there are zones in the cloud where the humidity is too low to activate the aerosols (e.g., at 4000 m at 2500 s where $\text{RH} < 85\%$, or indeed Figure 7a), but
 318 where there is a significant number of droplets and crystals ($> 0.03 \text{ cm}^{-3}$). These droplets and crystals have been activated elsewhere and previously, but

319 they nevertheless continue to collect aerosols around them – for example by Brownian capture, contributing to scavenging. It therefore **justifiable** to define
 320 a cloud contour based on a diagnostic of the numeric concentration of hydrometeors.

321

322 Table 1: Power law adjustment associated with each of the cloud contours studied.

Contour type	Power law adjustments
Based on relative humidity (Figure 8)	$\Lambda_{cloud}^m = 7.6 \times 10^{-5} I^{0.92}$
Based on mass concentration of cloud hydrometeors (Figure 9)	$\Lambda_{cloud}^m = 7.2 \times 10^{-5} I^{0.9}$
Based on numeric concentration of hydrometeors (Figure 10)	$\Lambda_{cloud}^m = 8.6 \times 10^{-5} I^{0.6}$

323

324 However, this numeric concentration criterion, although more precise for theoretically assessing the scavenging coefficient, is not easily accessible in a crisis
 325 code. Nevertheless, detailed analysis of the results of these simulations seems to show that it would be wise to define the cloud base as being constant and
 326 equal to the altitude at which critical supersaturation was first reached, i.e., the altitude at which the cloud began its formation.

327


328 2.2. Application to a stratus

329 Description of the stratus considered

330 The same approach as above was considered for modelling scavenging by a shallow stratus cloud. The main differences with the previous modelling (i.e., of
 331 the cumulonimbus) beyond the initialisation of thermodynamical profile is the treatment of the vertical advection within the cloud. Whereas, for the
 332 previous modelling, differences in air buoyancy (related to the initial thermal gradients and latent heat released by water phase changes) were the cause of
 333 vertical velocities and could be described and captured by the dynamics of the model. **When it comes to modelling stratus clouds, the dynamics are governed**
 334 **by large scales features that are not included in the 1.5D model. Therefore, the idea is to completely prescribe the time evolving profile for vertical velocity**
 335 **to model this forcing. Since convection is forced rather than triggered by buoyancy, it's reasonable to prescribe it and not calculate** the microphysical
 336 feedback on dynamics. For the scenario, we considered the vertical advection model proposed by Zhang *et al.*, (2004) and recapitulated in Equation 17. We
 337 therefore imposed a sinusoidal profile vertical velocity, the maximum oscillating from positive to negative values with a period of 1800 s. The maximum of
 338 the velocities was located at the altitude (z_c) of 1000 m and vertical motions allowed between 700 and 1300 m ($h_c = 600$ m, Figure 11). Like Zhang *et al.*,
 339 (2014), in the advection model, we imposed an average updraft velocity (w_0) of 0.2 m.s^{-1} and an oscillation amplitude (w_1) of 0.8 m.s^{-1} at an altitude of
 340 1000 m. Figure 5 shows the spatial-temporal distribution of vertical flows prescribed in the central cylinder. The temperature profile follows a dry adiabatic
 341 lapse rate with a temperature of 15°C on the ground so that there are no negative temperatures in the cloud. Above 1300 m, like Zhang *et al.* (2014), we
 342 imposed an inversion of the thermal profile. At altitudes between 700 and 1300 m, the relative humidity was initialised at 98.5%, and 95% outside of this
 343 range. For the aerosols, the initial conditions were identical to those for cumulonimbus (Equation 5).

$$\left\{ \begin{array}{l} w(z, t) = \cos\left(\pi \frac{z - z_c}{h_c}\right) \left[w_0 + w_1 \sin\left(\frac{2\pi t}{t_c}\right) \right] \quad \text{if } |z - z_c| \leq \frac{h_c}{2} \\ w(z, t) = 0 \quad \text{if } |z - z_c| > \frac{h_c}{2} \end{array} \right. \quad \text{Equation 17}$$


$z_c = 1000 \text{ m}; h_c = 600 \text{ m}; t_c = 1800 \text{ s}; w_0 = 0.2 \text{ m.s}^{-1}; w_1 = 0.8 \text{ m.s}^{-1}$



344

345 Figure 11. Spatial-temporal distribution of the vertical components of atmospheric flows (Zhang *et al.*, 2004)

346 Figure 12 shows on the left the spatial-temporal distribution of the water content calculated by DESCAM. The critical supersaturation was reached close to
 347 the altitude of 700 m from the first updraft phase (0–1000 s). The LWC then increased with altitude throughout the phase where the atmospheric flows
 348 were ascending. At the cloud summit, the liquid water content reached approximately 1.6 g.m^{-3} . Conversely, during the downdraft phases, the supply of dry
 349 air to lower altitudes induced, due to the temperature profile considered, a drop in the relative humidity **occurs**, which in turn induced evaporation of the
 350 droplets, resulting in a significant reduction in the LWC. These downdraft phases also had the effect of advecting droplets below the cloud band. During the
 351 period of velocity oscillations ($t < t_c$), precipitation completely evaporated before reaching the ground. However, from the second period onwards, rain
 352 was diagnosed at ground level (Figure 6. b).



353


354 Figure 12. On the left: Spatial-temporal distribution of the liquid water content calculated by DESCAM (LWC, in greyscale).
 355 On the right: Temporal evolution of rainfall intensity and cumulative precipitation diagnosed by DESCAM at ground level.

356 The DESCAM model predicts intermittent precipitation at ground level with flurries of precipitation in the order of a millimetre per hour. Over a precipitation
 357 period of approximately four hours, the cumulative precipitation was only approximately 3 mm.

358

359 Calculation of the stratus scavenging coefficient

360 As before in the case of cumulonimbus, it is necessary to define the contours of the cloud. We therefore used the three criteria previously introduced and
 361 look for the one with the clearest demarcation line between the cloud zone and the precipitation zone, to apply a dedicated scavenging coefficient (Figure
 362 13). As before, we observe from Figure 12.a that water content is not a good indicator to outline the cloud boundaries. Indeed, no discontinuity is observed
 363 for this parameter enabling demarcation between the cloud and the precipitation.



364

365

Figure 13. Test of different criteria to establish the contours of the simulated stratus.

366

a) threshold based on relative humidity: (---) $RH > 99\%$; (—) $RH > 100\%$.

367

b) threshold based on the mass concentration of hydrometeors with a diameter less than $64 \mu\text{m}$: (---) mass concentration of cloud hydrometeors $> 0.01 \text{ g} \cdot \text{m}^{-3}$; (—) mass concentration of cloud hydrometeors $> 0.001 \text{ g} \cdot \text{m}^{-3}$;

368

c) threshold based on the concentration of hydrometeors: (---); $\int dN > 0.1 \text{ cm}^{-3}$; (—) $\int dN + dN > 0.03 \text{ cm}^{-3}$.

369

370

371 Based on these results, it is more difficult to delineate the contours of this stratus than for the cumulonimbus. This is because, for these three criteria, only

372 the droplets concentration shows a clear demarcation between precipitation and cloud zone. Moreover, only this criterion gives stable cloud contours,

373 regardless of the threshold value selected. This difference with respect to the cumulonimbus is mainly due to the size of the precipitating hydrometeors,

374 which are much larger in the case of cumulonimbus. Figure 14 shows that the particle size distribution mode in number of raindrops, for the cumulonimbus,

375 is close to a diameter of 1 mm , whereas it is $100 \mu\text{m}$ for the stratus. It is therefore easier with a cumulonimbus than with a stratus to define a size threshold

376 distinguishing droplet (belonging to the cloud) from raindrops (belonging to precipitation). The criterion based on the mass concentration of hydrometeors


377 exceeding $64 \mu\text{m}$ is therefore less effective under a stratus than under a cumulonimbus. To explain the poor performance of the criterion based on relative

378 humidity, again it is the particle size that counts. As the droplets under the stratus are smaller than under the cumulonimbus, their velocities are lower, and

379 they reside longer in the atmosphere – about 10 times longer. This longer residence time promotes the increase in relative humidity under the cloud, and

380 humidity saturation under the cloud. This makes it difficult to use this criterion to determine the boundary between rain and cloud for a stratus.

381



382

Figure 14. Particle size distributions of raindrops determined by the DESCAM model at ground level

383

Left: For cumulonimbus at time $t=3000 \text{ s}$

384


Right: For cumulonimbus at time $t=8200 \text{ s}$

385

386


387 As previously, for the cumulonimbus, we search for a criterion to delimit cloud from rain. The same parameters as in section 2.1 are investigated and

388 presented in Figure 15, 16 and 17.




389

390 Figure 15. On the left: spatial-temporal distribution of the ammonium sulfate concentration in the cloud contour (— cloud contour for a
 391 relative humidity above 99%). On the right: correlation between the scavenging coefficient and the precipitation intensity determined at the
 392 base of the cloud, (—) adjusted by a power law.



393

394 Figure 16. On the left: spatial-temporal distribution of the ammonium sulfate concentration in the cloud contour (—) criterion based on the
 395 mass concentration of hydrometeors with a diameter greater than $64 \mu m$ with a threshold set at $0.01 g \cdot m^{-3}$). On the right: correlation
 396 between the scavenging coefficient and the precipitation intensity determined at the base of the cloud, (—) adjusted by a power law.



397

398 Figure 17. On the left: spatial-temporal distribution of the ammonium sulfate concentration in the contour of the cloud (criterion based on
 399 the concentration of hydrometeors with a threshold set at $0.01 particle \cdot cm^{-3}$). On the right: correlation between the scavenging coefficient
 400 and the precipitation intensity determined at the base of the cloud, (—) adjusted by a power law.

401

402 In these three figures, we observe that the contour introduced by Hiron (2017) for cumulonimbus (based on a separation between cloud water and
 403 precipitation water, on the basis of a criterion on the size of hydrometeors, cf. section 1.2) is no longer applicable for the stratus, and gives highly dispersed
 404 scavenging coefficient results, particularly for low rain intensity ($I < 2 mm \cdot h^{-1}$). This is because, for this stratus, it is difficult to establish a strict boundary

405 between a raindrop and a cloud droplet, based on their size. However, the other two criteria yield bijective and similar relationships, both in terms of the
 406 cloud contours (Figure 15.a and Figure 17.a) and the adjusted power laws (Table 2). Unlike cumulonimbus, stratus contours appear to be reliable using a
 407 criterion based on relative humidity. This difference is related to the intensities of vertical flows in the cumulonimbus. Indeed, we observe in Figure 5 that,
 408 in the simulated cumulonimbus, the downdraft flows can be very intense (up to 5 m.s^{-1}), transporting to the base of the cloud air masses with a lower mixing
 409 ratio and hence lower relative humidity.


410 Table 2. Adjustment of scavenging coefficients by power laws for the three types of contours studied.

Contour type	Power law adjustments
Based on relative humidity (Figure 15)	$\Lambda_{cloud}^m = 7.03 \times 10^{-4} I^{0.94}$
Based on mass concentration of cloud hydrometeors (Figure 16)	$\Lambda_{cloud}^m = 2.10 \times 10^{-3} I^{1.16}$
Based on hydrometeor concentration (Figure 17)	$\Lambda_{cloud}^m = 6.24 \times 10^{-4} I^{0.86}$

411
 412 These calculations show that, regardless of the type of simulated cloud, cumulonimbus or stratus, the criterion based on the hydrometeor concentration
 413 makes it possible to yield cloud contours that are both stable (with little variation when the threshold value is varied), and for which the relationship between
 414 the scavenging coefficient and the rainfall intensity is the most biunivocal (Figure 8, Figure 9 and Figure 10 for cumulonimbus and Figure 15, Figure 16 and
 415 Figure 17 for stratus). This criterion is not directly accessible in meteorological models; however, examination of Figure 10 and Figure 17 suggests that the
 416 cloud base remains stable over time. It would therefore be possible to assess the altitude at which critical supersaturation is reached, and to consider this
 417 altitude constant over a period that depends on the ratio between the size of the grid cell and the velocity of the horizontal flows.

418 419 **2.3.Comparison with the literature and unification of the scavenging coefficient scheme for a cumulonimbus** 420 **and a stratus**

421 **There is insufficient data to compare our theoretical findings with field study data. Few experimental data have established *in situ* scavenging coefficients**
 422 **for different types of clouds. Based on caesium-137 deposition measured following the Fukushima Daiichi accident, Leadbetter *et al.*, (2015) used the Met**
 423 **Office dispersion model: NAME (Numerical Atmospheric-dispersion Modelling Environment) for the dispersion of the radioactive plume emitted during the**
 424 **accident, considering the meteorological data from the ECMWF model. The authors managed to determine the cloud scavenging coefficient which best suits**
 425 **the ground measurements of deposition (Kinoshita *et al.*, 2011). In the same general approach, but using the IRSN LdX dispersion model (Quérel *et al.*, 2007**
 426 **and Groëll *et al.*, 2014) and meteorological data from MRI (Sekiyama *et al.*, 2017), Quérel *et al.* (2021) established a very similar scavenging coefficient.**
 427 **These two schemes are compared in Figure 18. The comparison is made using the κ value of ammonium sulfate. This decision is based on the findings of the**
 428 **Kaneyasu *et al.* (2012) study, which demonstrated the long-distance transport of cesium-137 by these particles - a distance particularly relevant for in-cloud**
 429 **scavenging.**



430
 431 Figure 18. Comparison of the parameterisations established respectively for a cumulonimbus (left) and a stratus (right) with the
 432 parameterisations established by Leadbetter *et al.*, 2015 and Quérel *et al.*, 2021 following the Fukushima accident.
 433

434 In this figure, we observe that the application of our scheme to a stratus (Figure 18, on the right), concurs excellently with the parametrisation of scavenging
 435 by clouds established following the Fukushima accident; in particular the parametrisation of [Quérel *et al.*, \(2021\)](#). However, the application of our approach
 436 to cumulonimbus presents much greater differences. Indeed, over the entire rainfall intensity range, our results are on average six times lower than the
 437 correlations of [Leadbetter *et al.*, \(2015\)](#) and [Quérel *et al.*, \(2021\)](#). Two questions therefore arise:


438 **First**, was there scavenging by cumulonimbus during the Fukushima accident? This would explain why it is difficult to compare our parametrisation of
 439 scavenging by cumulonimbus with those deduced during the Fukushima accident.

440 Next, why, for the same rainfall intensity, do our calculations show that cumulonimbus scavenges less than stratus?

441 We will therefore address these two questions.

442 **Was there scavenging by cumulonimbus during the Fukushima accident?**

443 To answer this question, let us consider the distribution of rainfall intensities diagnosed from radar measurements by [Saito *et al.*, \(2015\)](#) during March 2011
 444 in the Fukushima region (Figure 19). These results show that 80% of rain episodes diagnosed corresponded to rainfall intensities of less than 1.5 mm.h^{-1} ,
 445 and 97% to intensities of less than 3.5 mm.h^{-1} (range of rainfall intensity produced at the base of the simulated stratus, Figure 15) and less than 0.01% had
 446 intensities of more than 10 mm.h^{-1} . In view of these results, it is not possible to completely exclude the presence of rain issuing from cumulonimbus over
 447 the period of the accident; however, if there was any, its contribution to the construction of the parameterisation of [Leadbetter *et al.*, \(2015\)](#) and [Quérel *et al.*, \(2021\)](#)
 448 *et al.* (2021) is negligible.



449

450 Figure 19. Distribution of rainfall intensity measured in the Fukushima region during March 2011 ([Sekiyama *et al.*, 2017](#)).

451

452 **How to unify the scavenging coefficient scheme for a cumulonimbus and a stratus?**

453 Our calculations show that cumulonimbus scavenges less than stratus under the same rainfall intensity. How can we understand this and unify the two
 454 equations? This phenomenon can be attributed to the significantly higher level of supersaturation observed in cumulonimbus clouds (Figure 7.a) compared
 455 to that in stratus cloud (Figure 13.a). Hence, if the supersaturation is higher, as is the case for cumulonimbus, for the same activated aerosol mass, these
 456 particles are diluted in a larger mass of water, as the condensation is also much greater (in reality, the activated aerosol mass increases significantly since,
 457 as we have indicated previously, the activation diameter of the aerosols decreases as supersaturation increases). Let us therefore examine the impact of
 458 this effect of vapour condensation on the deduced parameterisation. In the DESCAM model, condensation is modelled by Equation 18. This equation is taken
 459 from [Pruppacher *et al.*, \(1998, chapter 13, section 2\)](#). It results from the vapour diffusion equation on a droplet of diameter \mathcal{D}_{drop} in the air with
 460 supersaturation \mathcal{S} and temperature T_∞ , considering the thermodynamic equilibrium of the suspended drop within air using the κ -Köhler theory.

$$\left\{ \begin{array}{l} \frac{d\mathcal{D}_{drop}}{dt} = \frac{4}{\mathcal{D}_{drop}} \frac{\rho_w \mathcal{R} T_\infty}{p_{sat,w} \mathcal{D}_v M_w} \frac{\mathcal{S} - y}{1 + \frac{\mathcal{O} n \rho_w}{k T_\infty} (\frac{\mathcal{O} n \rho_w}{\mathcal{R} T_\infty} - 1)} \\ y = \frac{4 \sigma_w \mathcal{D}_{drop} M_w}{\rho_w \mathcal{R} T_\infty \mathcal{D}_{drop}} - \kappa \frac{a_{ap}^{dry^3}}{\mathcal{D}_{drop}^3} \end{array} \right. \quad \text{Equation 18}$$

461

462 In this equation \mathcal{R} is the **ideal** gas constant, $p_{sat,w}$ the saturating vapour pressure, d_{ap}^{dry} the dry diameter of the aerosol, \mathcal{O}_n the latent heat of vaporisation
 463 of the water, $\sigma_{w,D_{drop}}$ the surface tension, k the thermal conductivity of the air, M_w and ρ_w the molar mass and density of the water vapour, and finally \mathcal{D}_v
 464 the diffusion coefficient of the vapour in water in the air. As the Kelvin effect (linked to the curvature of the interface) and the solute effect become very
 465 quickly negligible after activation of the aerosol, this equation can be greatly simplified and reduced to: $\mathcal{D}_{drop} d\mathcal{D}_{drop} = \mathbb{C} \cdot \mathcal{S} dt$, where \mathbb{C} is a constant,
 466 enabling it to be integrated analytically, **Δt the lifetime of the drop, to give: $\mathcal{D}_{drop} = \sqrt{\mathcal{D}_{drop,t_0}^2 + 2\mathbb{C} \cdot \mathcal{S} \cdot \Delta t}$** . Thus, to assess the effect of dilution of the
 467 aerosol in the droplet due to condensation, we can write:

$$\frac{\mathcal{D}_{drop}^{stratus^2}}{\mathcal{D}_{drop}^{cumulonimbus^2}} = \frac{\mathcal{D}_{drop,t_0}^2 + 2\mathbb{C} \cdot \langle \mathcal{S} \rangle_{stratus} \cdot \Delta t_{stratus}}{\mathcal{D}_{drop,t_0}^2 + 2\mathbb{C} \cdot \langle \mathcal{S} \rangle_{cumulonimbus} \cdot \Delta t_{cumulonimbus}} \quad \text{Equation 19}$$

468

469 **$\Delta t_{stratus}$ and $\Delta t_{cumulonimbus}$ are respectively the lifetime of the drop into a stratus and a cumulonimbus. Attention,** in this equation, $\mathcal{D}_{drop}^{stratus}$ and
 470 $\mathcal{D}_{drop}^{cumulonimbus}$ are not the diameters of the droplets in the stratus and in the cumulonimbus, but the diameters they would have had, if only the condensation
 471 mechanism had caused them to grow. We are in fact seeking to assess how large will be the dilution of aerosol material in the droplets related to vapour
 472 condensation. There are other mechanisms modelled in DESCAM (such as coalescence or riming, **Figure 2**) that lead to the growth of hydrometeors, without
 473 necessarily diluting the aerosols in the droplets. If there had only been the condensation mechanism, we could have used Figure 14 directly to assess this
 474 dilution. **When precipitation begins,** further simplification can still be made because $\mathcal{D}_{drop,t_0}^2 \ll \mathcal{D}_{drop}^{stratus^2} < \mathcal{D}_{drop}^{cumulonimbus^2}$. Finally, we can write:

$$\frac{\mathcal{D}_{drop}^{stratus^2}}{\mathcal{D}_{drop}^{cumulonimbus^2}} = \frac{\langle \mathcal{S} \rangle_{stratus} \cdot \Delta t_{stratus}}{\langle \mathcal{S} \rangle_{cumulonimbus} \cdot \Delta t_{cumulonimbus}} \quad \text{Equation 20}$$

475 In this equation, the **lifetimes $\Delta t_{stratus}$ and $\Delta t_{cumulonimbus}$** are therefore the **durations** necessary for the formation of precipitation under the cloud. **These**
 476 **durations are assimilated to the duration of precipitation formation.** For each of the types of cloud, we observe in Figure 6.a and Figure 12.a that these
 477 **durations** are very similar (≈ 2200 s), which allows us to write:


$$\frac{\mathcal{D}_{drop}^{stratus}}{\mathcal{D}_{drop}^{cumulonimbus}} = \sqrt{\frac{\langle \mathcal{S} \rangle_{stratus}}{\langle \mathcal{S} \rangle_{cumulonimbus}}} \quad \text{Equation 21}$$

478 The numerical application of this equation highlights a condensation growth ratio of a factor 2.3 between cumulonimbus and stratus. In mass, this coefficient
 479 corresponds to a dilution factor of 12. However, Figure 18 shows that, with this new approach, we can calculate that cumulonimbus scavenges 6 times less
 480 than stratus. This explanation is therefore satisfactory in view of all the hypotheses that have been made, especially since we have considered that the
 481 activated aerosol mass remained constant when supersaturation increased. We therefore propose a new generic parametrisation to any type of cloud,
 482 which this time **considers** this condensation-related dilution effect, Equation 22. This scavenging scheme is therefore corrected by a coefficient $1/\langle \mathcal{S} \rangle_{cloud}^{3/2}$
 483 which characterises the dilution related to the growth of droplets by condensation:

$$\Lambda_{cloud}^m = \frac{5 \times 10^{-8}}{\langle \mathcal{S} \rangle_{cloud}^{3/2}} I^{0.75} \quad \text{Equation 22}$$

484

485 The application of this new correlation, presented in Figure 20 shows an excellent match both for the cumulonimbus and for the simulated stratus. It
 486 remains to be considered whether supersaturation is accessible in the NWP's and, if so, if the horizontal resolutions of 1 to 10 km of such models are
 487 sufficiently representative of a real cloud.



488

489 Figure 20. Comparison of the newly developed correlation Equation 22 with the scavenging modelled by DESCAM for a cumulonimbus and a
490 stratus.

491

492

493 **3. Conclusions**

494

495 The in-cloud scavenging scheme established in this article shows a dependence on rain intensity and average supersaturation in the cloud. Supersaturation
496 allows the scheme to be applicable to both cumulonimbus and stratus clouds. If supersaturation in the cloud is not accessible, it is still possible to apply a
497 different scheme for convective clouds and stratiform clouds. But, since this boundary between the two types of cloud may be ambiguous, it will be
498 preferable to apply the scheme with supersaturation if available.

499 This scavenging scheme is based on the DESCAM microphysical cloud model. This model allows fine-scale description of the life cycle of a cloud up to
500 precipitation development. It tracks particles, crystals, and droplets particle size distributions and models all the water phase changes and, above all, how
501 aerosol particles impact them. The in-cloud scavenging scheme is established by calculating the mass fluxes of particle material exiting the cloud being
502 included in precipitation hydrometeors (both liquid and solid) and based on the mass of particles initially present in the cloud volume.

503 This calculation of cloud volume has proved to be a complex issue, in particular for establishing the altitude of the cloud base, especially when rain occurs.
504 The most relevant method to identify cloud base in this study has been proven to be the one using the number of hydrometeors, rather than the relative
505 humidity or the mass of the hydrometeors. The problem with this method is that this information on the number is not available for most of the NWP. The
506 use of the in-cloud scavenging scheme must be based on a diagnostic independent of the altitude of the base – and the summit – of the cloud.

507 In the case of stratus cloud, the parametrization obtained with DESCAM is close to those currently used in the NAME and LdX atmospheric dispersion models,
508 which were established on the basis of the Fukushima accident. As the precipitation that caused deposition of radioactive particles following the accident
509 was largely generated by stratiform clouds, this study confirms *a posteriori* the choice of the in-cloud deposition scheme used to study radioactive deposition
510 following the Fukushima accident and can be extended to all types of cloud.

511 In future works, this deposition scheme will be used with confidence to study deposition. As an example, it can be used for the deposition of radon progeny
512 (Quérel *et al.*, 2022), in order to statistically measure the impact of this scheme in relation to the existing corpus.

513 Beyond the applications and validations of the scheme described in this article, the scheme itself is currently being refined. First of all, we are working on
514 establishing an in-cloud scavenging rate that will depend on particle size. This important issue was discussed in section 1.3, and requires some modifications
515 to the model to establish a model spectrally. This will make it possible to apply a finer-scale scheme to the atmospheric models with a spectral representation
516 of the particles.

517 The influence of the coefficient κ of the κ -Köhler theory can be also examined. This will make it possible to measure the importance of the physical-chemical
518 properties of the particles: what error is made by applying the same scavenging rate for a hygroscopic aerosol (salt or sulfate) and a non-hygroscopic aerosol
519 (soot, desert dust).

520 The initial particle size distribution of aerosols could also have a significant influence on the final scavenging rate. A distribution centred on 100 nm will not
521 create the same cloud as the same total mass centred around 5 μm particles. This aspect must be assessed.

522 The question of evaporation of droplets between the cloud base and the ground has not yet been addressed. The scheme developed is based on the
523 precipitation intensity at the cloud base, but in the models the precipitation intensity is diagnosed on the ground. This is important for the applicability of
524 the scheme and this difference can lead to errors, especially in the event of high droplet evaporation.

525 Finally, it is not yet established that this scheme is as effective when applied to a model whose spatial resolution is lower than that of DESCAM, as is the
526 case for all Climate (GCM) and Transport (ATM) models.

527 The work still to be carried out will make it possible to best define the scope of validity of this new scheme for in-cloud aerosol scavenging, as well as the
528 uncertainties associated with this model. This will enable the scheme to be used in full knowledge of the facts and according to the highest scientific
529 standards.

530

531

532 **Competing interests**

533 The contact author has declared that none of the authors has any competing interests.

534

535 **References**

- 536 Adachi K, Kajino M, Zaizen Y, Igarashi Y. Emission of spherical cesium-bearing particles from an early stage of the Fukushima nuclear accident. *Sci Rep.*
537 2013;3:2554. doi: 10.1038/srep02554. PMID: 23989894; PMCID: PMC3757362.
- 538 Asai, T., & Kasahara, A. (1967). A theoretical study of the compensating downward motions associated with cumulus clouds. *Journal of Atmospheric*
539 *Sciences*, 24(5), 487-496.
- 540 American Meteorological Society, 2020b: Glossary of meteorology: cloud. Website, accessed on October 21, 2020, URL [http://](http://glossary.ametsoc.org/wiki/cloud)
541 glossary.ametsoc.org/wiki/cloud.
- 542 Baklanov, A., & Sørensen, J. H. (2001). Parameterisation of radionuclide deposition in atmospheric long-range transport modelling. *Physics and Chemistry*
543 *of the Earth, Part B: Hydrology, Oceans and Atmosphere*, 26(10), 787-799.
- 544 Bigg, E. K. (1953). The formation of atmospheric ice crystals by the freezing of droplets. *Quarterly Journal of the Royal Meteorological Society*, 79(342), 510-
545 519.
- 546 Beard, K. V. (1974). Experimental and numerical collision efficiencies for submicron particles scavenged by small raindrops. *Journal of Atmospheric*
547 *Sciences*, 31(6), 1595-1603.
- 548 Bergeron, T. (1928). *Über die dreidimensional verknüpfende Wetteranalyse. 1. Prinzipielle Einführung in das Problem der Luftmassen-und Frontenbildung.*
549 na. Bony, S., & Dufresne, J. L. (2005). Marine boundary layer clouds at the heart of tropical cloud feedback uncertainties in climate models. *Geophysical*
550 *Research Letters*, 32(20).
- 551 Clark, M. J., & Smith, F. B. (1988). Wet and dry deposition of Chernobyl releases. *Nature*, 332(6161), 245-249.
- 552 Costa, M. J., Salgado, R., Santos, D., Levizzani, V., Bortoli, D., Silva, A. M., & Pinto, P. (2010). Modelling of orographic precipitation over Iberia: a springtime
553 case study. *Advances in Geosciences*, 25, 103-110.
- 554 Croft, B., Lohmann, U., Martin, R. V., Stier, P., Wurzler, S., Feichter, J., ... & Ferrachat, S. (2010). Influences of in-cloud aerosol scavenging parameterizations
555 on aerosol concentrations and wet deposition in ECHAM5-HAM. *Atmospheric Chemistry and Physics*, 10(4), 1511-1543.
- 556 De Cort, M. (1998). Atlas of caesium deposition on Europe after the Chernobyl accident.
- 557 Dépée, A., Lemaitre, P., Gelain, T., Mathieu, A., Monier, M., & Flossmann, A. (2019). Theoretical study of aerosol particle electroscavenging by clouds. *Journal*
558 *of Aerosol Science*, 135, 1-20.
- 559 Dépée, A. (2019). *Etude expérimentale et théorique des mécanismes microphysiques mis en jeu dans la capture des aérosols radioactifs par les*
560 *nuages* (Doctoral dissertation, Université Clermont Auvergne [2017-2020]).
- 561 Dépée, A., Lemaitre, P., Gelain, T., Monier, M., & Flossmann, A. (2021). Laboratory study of the collection efficiency of submicron aerosol particles by cloud
562 droplets—Part I: Influence of relative humidity. *Atmospheric Chemistry and Physics*, 21(9), 6945-6962.
- 563 Dépée, A., Lemaitre, P., Gelain, T., Monier, M., & Flossmann, A. (2021). Laboratory study of the collection efficiency of submicron aerosol particles by cloud
564 droplets—Part II: Influence of electric charges. *Atmospheric Chemistry and Physics*, 21(9), 6963-6984.
- 565 Dye, J. E., Jones, J. J., Winn, W. P., Cerni, T. A., Gardiner, B., Lamb, D., ... & Saunders, C. P. R. (1986). Early electrification and precipitation development in a
566 small, isolated Montana cumulonimbus. *Journal of Geophysical Research: Atmospheres*, 91(D1), 1231-1247.
- 567 Ervens, B. (2015). Modeling the processing of aerosol and trace gases in clouds and fogs. *Chemical reviews*, 115(10), 4157-4198.
- 568 Findeisen, Z. (1938). Kolloid meteorologische vorgänge bei neiderschlags-bildung. *Meteorologische Zeitschrift*, 55, 121.
- 569 Flossmann, A. I., Hall, W. D., & Pruppacher, H. R. (1985). A theoretical study of the wet removal of atmospheric pollutants. Part I: The redistribution of aerosol
570 particles captured through nucleation and impaction scavenging by growing cloud drops. *Journal of Atmospheric Sciences*, 42(6), 583-606.
- 571 Flossmann, A. I., Pruppacher, H. R., & Topalian, J. H. (1987). A theoretical study of the wet removal of atmospheric pollutants. Part II: The uptake and
572 redistribution of (NH₄)₂SO₄ particles and SO₂ gas simultaneously scavenged by growing cloud drops. *Journal of Atmospheric Sciences*, 44(20), 2912-2923.
- 573 Flossmann, A. I., & Pruppacher, H. R. (1988). A theoretical study of the wet removal of atmospheric pollutants. Part III: The uptake, redistribution, and
574 deposition of (NH₄)₂SO₄ particles by a convective cloud using a two-dimensional cloud dynamics model. *Journal of Atmospheric Sciences*, 45(13), 1857-
575 1871.
- 576 Flossmann, A. I. (1998). Clouds and pollution. *Pure and applied chemistry*, 70(7), 1345-1352.

577 Flossmann, A. I., & Wobrock, W. (2010). A review of our understanding of the aerosol–cloud interaction from the perspective of a bin resolved cloud scale
578 modelling. *Atmospheric Research*, 97(4), 478-497.

579 Groëll, J., Quélo, D., & Mathieu, A. (2014). Sensitivity analysis of the modelled deposition of 137 Cs on the Japanese land following the Fukushima
580 accident. *International Journal of Environment and Pollution*, 55(1-4), 67-75.

581 Hertel, O., Christensen, J., Runge, E. H., Asman, W. A., Berkowicz, R., Hovmand, M. F., & Hov, Ø. (1995). Development and testing of a new variable scale air
582 pollution model—ACDEP. *Atmospheric Environment*, 29(11), 1267-1290.

583 Hirsch, E., Koren, I., Levin, Z., Altaratz, O., & Agassi, E. (2014). On transition-zone water clouds. *Atmospheric Chemistry and Physics*, 14(17), 9001-9012.

584 Hiron, T., & Flossmann, A. I. (2015). A study of the role of the parameterization of heterogeneous ice nucleation for the modeling of microphysics and
585 precipitation of a convective cloud. *Journal of the Atmospheric Sciences*, 72(9), 3322-3339.

586 Hiron, T. (2017). *Experimental and modeling study of heterogeneous ice nucleation on mineral aerosol particles and its impact on a convective cloud* (Doctoral
587 dissertation, Université Clermont Auvergne).

588 Kaneyasu, N., Ohashi, H., Suzuki, F., Okuda, T., & Ikemori, F. (2012). Sulfate aerosol as a potential transport medium of radiocesium from the Fukushima
589 nuclear accident. *Environmental science & technology*, 46(11), 5720-5726.

590 Kerker, M., & Hampl, V. (1974). Scavenging of Aerosol Particles by a Falling Water Drop and Calculation of Washout Coefficients. *Journal of Atmospheric
591 Sciences*, 31(5), 1368-1376.

592 Kinoshita, N., Sueki, K., Sasa, K., Kitagawa, J. I., Ikarashi, S., Nishimura, T., ... & Yamagata, T. (2011). Assessment of individual radionuclide distributions from
593 the Fukushima nuclear accident covering central-east Japan. *Proceedings of the National Academy of Sciences*, 108(49), 19526-19529.

594 Knight, C. A. (1982). The cooperative convective precipitation experiment (CCOPE), 18 May–7 August 1981. *Bulletin of the American Meteorological
595 Society*, 63(4), 386-398.

596 Koop, T., Luo, B., Tsias, A., & Peter, T. (2000). Water activity as the determinant for homogeneous ice nucleation in aqueous solutions. *Nature*, 406(6796),
597 611-614.

598 Laguionie, P., Roupsard, P., Maro, D., Solier, L., Rozet, M., Hébert, D., & Connan, O. (2014). Simultaneous quantification of the contributions of dry, washout
599 and rainout deposition to the total deposition of particle-bound 7Be and 210Pb on an urban catchment area on a monthly scale. *Journal of aerosol
600 science*, 77, 67-84.

601 Lai, K. Y., Dayan, N., & Kerker, M. (1978). Scavenging of aerosol particles by a falling water drop. *Journal of Atmospheric Sciences*, 35(4), 674-682.

602 Leadbetter, S. J., Hort, M. C., Jones, A. R., Webster, H. N., & Draxler, R. R. (2015). Sensitivity of the modelled deposition of Caesium-137 from the Fukushima
603 Dai-ichi nuclear power plant to the wet deposition parameterisation in NAME. *Journal of environmental radioactivity*, 139, 200-211.

604 Leaitch, W., Strapp, J. W., Isaac, G. A., & Hudson, J. G. (1986). Cloud droplet nucleation and cloud scavenging of aerosol sulphate in polluted
605 atmospheres. *Tellus B: Chemical and Physical Meteorology*, 38(5), 328-344.

606 Lemaitre, P., Quérel, A., Monier, M., Menard, T., Porcheron, E., & Flossmann, A. I. (2017). Experimental evidence of the rear capture of aerosol particles by
607 raindrops. *Atmospheric Chemistry and Physics*, 17(6), 4159-4176.

608 Leroy, D., Monier, M., Wobrock, W., & Flossmann, A. I. (2006). A numerical study of the effects of the aerosol particle spectrum on the development of the
609 ice phase and precipitation formation. *Atmospheric Research*, 80(1), 15-45.

610 Leroy, D., Wobrock, W., & Flossmann, A. I. (2007). On the influence of the treatment of aerosol particles in different bin microphysical models: a comparison
611 between two different schemes. *Atmospheric research*, 85(3-4), 269-287.

612 Meyers, M. P., DeMott, P. J., & Cotton, W. R. (1992). New primary ice-nucleation parameterizations in an explicit cloud model. *Journal of Applied Meteorology
613 and Climatology*, 31(7), 708-721.

614 Monier, M., Wobrock, W., Gayet, J.-F., Flossmann, A., (2006). Development of a detailed microphysics cirrus model for interpretation of the recent INCA
615 campaign. *J. Atmos. Sci.* 163.

616 Mathieu, A., Korsakissok, I., Quélo, D., Groëll, J., Tombette, M., Didier, D., ... & Isnard, O. (2012). Atmospheric dispersion and deposition of radionuclides
617 from the Fukushima Daiichi nuclear power plant accident. *Elements*, 8(3), 195-200.

618 MRI, 2015. JMA-RATM Technical Report, vol. 76.

619 Palmer, T. (2014). Climate forecasting: Build high-resolution global climate models. *Nature*, 515(7527), 338-339.

620 Petters, M. D., & Kreidenweis, S. M. (2007). A single parameter representation of hygroscopic growth and cloud condensation nucleus activity. *Atmospheric*
621 *Chemistry and Physics*, 7(8), 1961-1971.

622 Petroff, A., Mailliat, A., Amielh, M., & Anselmet, F. (2008). Aerosol dry deposition on vegetative canopies. Part I: review of present knowledge. *Atmospheric*
623 *Environment*, 42(16), 3625-3653.

624 Pranesha, T. S., & Kamra, A. K. (1996). Scavenging of aerosol particles by large water drops: 1. Neutral case. *Journal of Geophysical Research:*
625 *Atmospheres*, 101(D18), 23373-23380.

626 Pruppacher, H. R., Klett, J. D., & Wang, P. K. (1998). Microphysics of clouds and precipitation. 2nd ed., Atmospheric and Oceanographic Sciences Library,
627 Kluwer Academic Publishers, doi:10.1080/02786829808965531.

628 Quélo, D., Krysta, M., Bocquet, M., Isnard, O., Minier, Y., & Sportisse, B. (2007). Validation of the Polyphemus platform on the ETEX, Chernobyl and Algeciras
629 cases. *Atmospheric environment*, 41(26), 5300-5315.

630 Quérel, A., Monier, M., Flossmann, A. I., Lemaitre, P., & Porcheron, E. (2014). The importance of new collection efficiency values including the effect of rear
631 capture for the below-cloud scavenging of aerosol particles. *Atmospheric research*, 142, 57-66.

632 Quérel, A., Quelo, D., Roustan, Y., & Mathieu, A. (2021). Sensitivity study to select the wet deposition scheme in an operational atmospheric transport
633 model. *Journal of Environmental Radioactivity*, 237, 106712.

634 Quérel, A., Meddouni, K., Quélo, D., Doursout, T., Chuzel, S., 2022. Statistical approach to assess radon-222 long-range atmospheric transport modelling and
635 its associated gamma dose rate peaks, in: Advances in Geosciences. Presented at the Geoscience applications of environmental radioactivity - EGU General
636 Assembly 2021, Online, 19–30 April 2021, Copernicus GmbH, pp. 109–124.

637 Saito, K., Shimbori, T., & Draxler, R. (2015). JMA's regional atmospheric transport model calculations for the WMO technical task team on meteorological
638 analyses for Fukushima Daiichi Nuclear Power Plant accident. *Journal of environmental radioactivity*, 139, 185-199.

639 Sassen, K., and B. S. Cho, 1992: Subvisual-thin cirrus lidar dataset for satellite verification and climatological research. *J. Appl. Meteorol.*, 31, 1275–1285,
640 doi:10.1175/1520-0450(1992) 031<1275:STCLDF>2.0.CO;2, [https://doi.org/10.1175/1520-0450\(1992\)031<1275:STCLDF>](https://doi.org/10.1175/1520-0450(1992)031<1275:STCLDF>2.0.CO;2)
641 2.0.CO;2.

642 Saito, K., Shimbori, T., & Draxler, R. (2015). JMA's regional atmospheric transport model calculations for the WMO technical task team on meteorological
643 analyses for Fukushima Daiichi Nuclear Power Plant accident. *Journal of environmental radioactivity*, 139, 185-199.

644 Sievering, H., Van Valin, C. C., Barrett, E. W., & Pueschel, R. F. (1984). Cloud scavenging of aerosol sulfur: Two case studies. *Atmospheric Environment*
645 (1967), 18(12), 2685-2690.

646 Slinn, W. G. N.: Some approximations for the wet and dry removal of particles and gases from the atmosphere, *Water Air Soil Poll.*, 7, 513–543, 1977.

647 Spänkuch, D., Hellmuth, O., & Görtsdorf, U. (2022). What is a cloud? Toward a more precise definition?. *Bulletin of the American Meteorological Society*.

648 Stephan, K., Klink, S., & Schraff, C. (2008). Assimilation of radar-derived rain rates into the convective-scale model COSMO-DE at DWD. *Quarterly Journal of*
649 *the Royal Meteorological Society: A journal of the atmospheric sciences, applied meteorology and physical oceanography*, 134(634), 1315-1326.

650 **Textor, C., Schulz, M., Guibert, S., Kinne, S., Balkanski, Y., Bauer, S., ... & Tie, X. (2006). Analysis and quantification of the diversities of aerosol life cycles**
651 **within AeroCom. *Atmospheric Chemistry and Physics*, 6(7), 1777-1813.**

652 Wegener, A. (1911). *Thermodynamik der atmosphäre*. JA Barth.

653 Vali, G., DeMott, P. J., Möhler, O., & Whale, T. F. (2015). A proposal for ice nucleation terminology. *Atmospheric Chemistry and Physics*, 15(18), 10263-10270.

654 Vohl, O., Mitra, S. K., Wurzler, S. C., & Pruppacher, H. R. (1999). A wind tunnel study of the effects of turbulence on the growth of cloud drops by collision
655 and coalescence. *Journal of the atmospheric sciences*, 56(24), 4088-4099.

656 Wang, H., & Su, W. (2013). Evaluating and understanding top of the atmosphere cloud radiative effects in Intergovernmental Panel on Climate Change (IPCC)
657 Fifth Assessment Report (AR5) Coupled Model Intercomparison Project Phase 5 (CMIP5) models using satellite observations. *Journal of Geophysical*
658 *Research: Atmospheres*, 118(2), 683-699.

659 WMO. (2014). Guide to Meteorological Instruments and Methods of Observation (CIMO Guide, WMO-No. 8. Wood, R., & Field, P. R. (2011). The distribution
660 of cloud horizontal sizes. *Journal of Climate*, 24 (18), 4800-4816.

661 Zhang, L., Michelangeli, D. V., & Taylor, P. A. (2004). Numerical studies of aerosol scavenging by low-level, warm stratiform clouds and
662 precipitation. *Atmospheric Environment*, 38(28), 4653-4665.

663 Zhang, S., Xiang, M., Xu, Z., Wang, L., & Zhang, C. (2020). Evaluation of water cycle health status based on a cloud model. *Journal of cleaner production*, 245,
664 118850.