

# Influence of Ocean Alkalinity Enhancement with Olivine or Steel Slag on a Coastal Plankton Community in Tasmania

Jiaying A. Guo<sup>1,2</sup>, Robert F. Strzepek<sup>2</sup>, Kerrie M. Swadling<sup>1,2</sup>, Ashley T. Townsend<sup>3</sup>, Lennart T. Bach<sup>1</sup>

<sup>1</sup>Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, Tasmania, 7000 Australia

<sup>2</sup>Australian Antarctic Program Partnership (AAPP), Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, Tasmania, 7000 Australia

<sup>3</sup>Central Science Laboratory, University of Tasmania, Sandy Bay, Tasmania, 7005 Australia


Correspondence to: Jiaying A. Guo (Jiaying.guo@utas.edu.au)

**Abstract.** Ocean alkalinity enhancement (OAE) aims to increase atmospheric CO<sub>2</sub> sequestration in the oceans through the acceleration of chemical rock weathering. This could be achieved by grinding rocks containing alkaline minerals and adding the rock powder to the surface ocean where it dissolves and chemically locks CO<sub>2</sub> in seawater as bicarbonate. However, CO<sub>2</sub> sequestration during dissolution coincides with the release of potentially bio-active chemicals and may induce side effects. Here, we used 53 L microcosms to test how coastal plankton communities from Tasmania respond to OAE with olivine (mainly Mg<sub>2</sub>SiO<sub>4</sub>) or steel slag (mainly CaO and Ca(OH)<sub>2</sub>) as alkalinity sources. Three microcosms were left unperturbed and served as a control, three were enriched with olivine powder (1.9 g L<sup>-1</sup>), and three with steel slag powder (0.038 g L<sup>-1</sup>). Olivine and steel slag powders were of similar grain size. Olivine was added in a higher amount than the steel slag ~~since previous tests evidenced that it would have released less with the aim to compensate for the lower efficiency of olivine to deliver~~ alkalinity over the 3-week experiment. Phytoplankton and zooplankton community responses as well as some biogeochemical parameters were monitored. Olivine and steel slag additions increased total alkalinity by 29 μmol kg<sup>-1</sup> and 361 μmol kg<sup>-1</sup> respectively, which corresponds to a theoretical increase of 0.9 % and 14.8 % of the seawater storage capacity for atmospheric CO<sub>2</sub>. Olivine and steel slag released silicate nutrients into the seawater, but steel slag released considerably more and also significant amounts of phosphate. After 21 days, no significant difference was found in dissolved iron concentrations (>100 nmol L<sup>-1</sup>) in the treatments and the control. The slag addition increased dissolved manganese concentrations (771 nmol L<sup>-1</sup>), while olivine increased dissolved nickel concentrations (37 nmol L<sup>-1</sup>). There was no significant difference in total chlorophyll *a* concentrations between the treatments and the control, likely due to nitrogen limitation of the phytoplankton community. However, flow cytometry results indicated an increase in the cellular abundance of several smaller (<20 μm) phytoplankton groups in the olivine treatment. The abundance of larger phytoplankton (>20 μm) decreased much more in the control than in the treatments after day 10. Furthermore, the maximum quantum yields of photosystem II (F<sub>v</sub>/F<sub>m</sub>) were higher in slag and olivine treatments, suggesting that mineral additions increased photosynthetic performance. The zooplankton community composition was also affected with the most notable changes being observed in the dinoflagellate *Noctiluca scintillans* and the appendicularian *Oikopleura* sp. in the olivine treatment. Overall, steel slag is much more efficient for CO<sub>2</sub> removal with OAE than olivine and appears to induce less change in the plankton community when relating the CO<sub>2</sub> removal potential to the level of environmental impact that was observed here.

## 37 1 Introduction


38 Keeping global warming below 2 °C requires immediate emissions reduction. Additionally, between 450-1100 Gigatonnes  
39 of carbon dioxide (CO<sub>2</sub>) need to be removed from the atmosphere by 2100 (Smith et al., 2023). This could be achieved  
40 with a portfolio of terrestrial and marine Carbon Dioxide Removal (CDR) methods. Ocean alkalinity enhancement (OAE)  
41 is a marine CDR method that could theoretically contribute significantly to the global CDR portfolio (Ilyina et al., 2013;  
42 Feng et al., 2017; Lenton et al., 2018).

43  
44 Alkalinity is generated naturally when rock weathers and it has control on the ocean's chemical capacity to store CO<sub>2</sub>  
45 (Schuiling and Krijgsman, 2006). Natural rock weathering is currently responsible for about 0.5 Gt of atmospheric CO<sub>2</sub>  
46 sequestration every year (Renforth and Henderson, 2017). The idea behind OAE is to accelerate natural rock weathering  
47 by extracting calcium- or magnesium-rich rocks, such as olivine, pulverizing them, and spreading them onto the sea surface  
48 to increase chemical weathering rates (Hartmann et al., 2013). The weathering (i.e., dissolution) of these alkaline minerals  
49 will consume protons (H<sup>+</sup>), which shifts the carbonate chemistry equilibrium in seawater from CO<sub>2</sub> towards increasing  
50 bicarbonate (HCO<sub>3</sub><sup>-</sup>) and carbonate ion (CO<sub>3</sub><sup>2-</sup>) concentrations:




52  
53  
54 thereby making new space for atmospheric CO<sub>2</sub> to be dissolved in seawater and permanently stored. Previous model studies  
55 have shown that OAE can mitigate climate change significantly by increasing the oceanic uptake of CO<sub>2</sub> from the  
56 atmosphere (Kohler et al., 2010; Paquay and Zeebe, 2013; Keller et al., 2014; Lenton et al., 2018). For example, the study  
57 by Burt et al. (2021) suggested that the total global mean dissolved inorganic carbon (DIC) inventories would increase by  
58 156 GtC after total alkalinity is enhanced at a rate of 0.25 Pmol year<sup>-1</sup> in 75-year simulations.

59  
60 There are a variety of alkaline minerals that could be used for OAE. A widely considered naturally occurring mineral is  
61 forsterite, a (Mg<sub>2</sub>SiO<sub>4</sub>)-rich olivine. This type of olivine is abundant in ultramafic rock such as dunite, constituting at least  
62 88 % of the rock composition (Ackerman et al., 2009; Su et al., 2016). Olivine occurs in the Earth's crust but is more  
63 abundant in the upper mantle. There are at least several billion tons of olivine resources on Earth (Caserini et al., 2022).  
64 However, the extraction of olivine in 2017 was only around 8.4 Mt year<sup>-1</sup> (Reichl et al., 2018), which is about two orders  
65 of magnitude below the mass needed for climate-relevant OAE with olivine (Caserini et al., 2022). The net reaction for  
66 CO<sub>2</sub> sequestration with Mg<sub>2</sub>SiO<sub>4</sub> is:



68  
69  
70 Another potential OAE source material is steel slag (Renforth, 2019), a by-product of steel manufacturing. During steel  
71 manufacturing, high-purity calcium oxide (CaO) is used to improve the quality of the steel through accumulation of  
72 unwanted materials such as sulphur and phosphorus. Steel slag mainly contains CaO, SiO<sub>2</sub>, Al<sub>2</sub>O<sub>3</sub>, Fe<sub>2</sub>O<sub>3</sub>, MgO, and MnO  
73 (Kourounis et al., 2007), and the chemical composition can vary depending on the manufacturing process (Wang et al.,  
74 2011). Due to the presence of CaO and potentially other alkaline components, steel slag can increase alkalinity when  
75 dissolved in seawater. The chemical reaction for CO<sub>2</sub> sequestration with CaO is:

76



78

79 Some of the steel slag that is produced during steel manufacturing is further used (e.g., for road construction and civil  
80 engineering) but in some countries like China, 70.5 % of steel slag is left unused and stored in dumps (Guo et al., 2018).  
81 In 2016, more than 300 million tons of steel slag was not used effectively, thereby occupying the land and raising  
82 environmental concerns (Guo et al., 2018). The effective alkaline composition, availability, and relatively low cost of the  
83 raw materials make olivine and steel slag potential source materials for OAE.

84

85 To assess whether OAE is viable, it needs to be understood how its application may affect marine biota such as plankton  
86 and the biogeochemical fluxes they drive. Some data on the effects of OAE with sodium hydroxide (NaOH) on plankton  
87 communities have recently been published (Ferderer et al., 2022; Subhas et al., 2022), but to the best of our knowledge, no  
88 such data are available for olivine- and/or slag-based OAE. Chemical perturbations via olivine and slag should be like  
89 those by NaOH in that they increase seawater pH and shift the carbonate chemistry equilibrium (see Eq. 1). However, there  
90 would be additional chemical perturbations because minerals contain a variety of potentially bioactive elements that are  
91 released into the environment when they dissolve in seawater (Bach et al., 2019). One particular concern is that natural and  
92 anthropogenic minerals such as olivine and steel slag are rich in bioactive metals that are usually scarce in the ocean, such  
93 as iron (Fe), copper (Cu), nickel (Ni), manganese (Mn), zinc (Zn), cadmium (Cd), and chromium (Cr). Many of these trace  
94 metals are essential micronutrients for phytoplankton growth (Sunda, 2000; Sunda, 2012), such as being co-factors for  
95 various metalloenzymes (summarized by Twining and Baines, 2013). It is possible that the addition of alkaline minerals  
96 may benefit phytoplankton by providing trace metals currently limiting phytoplankton growth (Falkowski, 1994; Basu and  
97 Mackey, 2018). For instance, the addition of Fe is well known to stimulate phytoplankton blooms in those vast ocean  
98 regions where Fe levels limit growth (Boyd et al., 2007; Moore et al., 2013). However, some trace metals can also inhibit  
99 phytoplankton growth, and different phytoplankton species have different requirements and tolerances for trace metals  
100 (Sunda, 2012) so the addition of trace metals via OAE may change phytoplankton community composition.


101

102 Here, we describe a microcosm experiment with coastal Tasmanian plankton communities that was used to investigate: (1)  
103 how effectively OAE via the application of finely ground olivine and steel slag could sequester atmospheric CO<sub>2</sub>, and (2)  
104 if/how olivine and steel slag additions affect various components of the plankton community.

105

106 **2 Methodology**

107 **2.1 Microcosm setup**



108

109 **Fig. 1.** Experimental design and alkalinity sources. (a) Raw materials used as alkalinity sources: olivine (left) and steel slag (right).  
110 Olivine and steel slag were originally larger than 20 mm. (b) Ground minerals observed with a scanning electron microscope (SEM). (c)  
111 Microcosm setup: each microcosm enclosed ~ 53 L of surface seawater with natural plankton communities. Olivine and steel slag  
112 treatments and the control were kept in a temperature-controlled room and two heat belts were attached to the bottom of each microcosm  
113 to create convective circulation.

114

115 We used nine 53 L transparent Kegland® Fermzilla conical unitank fermenters (polyethylene terephthalate) (Fig. 1) as  
116 microcosms to incubate natural plankton communities. All microcosms were prewashed with hydrochloric acid (10 % v/v)  
117 and rinsed five times with 18.2 MΩ Milli-Q water. Seawater with coastal plankton communities was collected at Battery  
118 Point, Tasmania (42.892°S, 147.337°E) within 2 hours by lowering the microcosms into the ocean with a crane and filling  
119 them in a manner similar to a Niskin bottle, as described in detail in Ferderer et al. (2022). A sieve with a mesh size of 2  
120 mm was attached to the top and bottom of the microcosms during filling to avoid the entrapment of large and patchily  
121 distributed organisms in the microcosms. The enclosed seawater weight was initially between 52.35-54.70 kg. After  
122 seawater collection, filled microcosms were immediately transported back to the Institute for Marine and Antarctic Studies  
123 (University of Tasmania) on a truck and transferred within 75 min into a temperature-controlled room set to 7.5-8 °C. Two  
124 heat belts were attached to the bottom of each microcosm to induce a convective mixing current (Ferderer et al., 2022).  
125 Seawater temperature inside the microcosms was about 13.5 °C due to the heating effects of the heat belts and was the  
126 same as the sampled region. LED light strips were used to provide an average light intensity of 236  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$   
127 (ranging from 208 to 267  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ) with a daily light-dark cycle of 10:14 hours. The light intensity was the  
128 average light intensity in each microcosm measured with a LICOR light meter at 0.15 m depth within the microcosm.  
129 Microcosms positioned in the temperature-controlled room were shuffled anti-clockwise every day to ensure similar light

130 intensity for each microcosm throughout the experiment. Treatments were established 24 hours after collecting the seawater.  
131 The total alkalinity released per amount of mineral powder added was much higher for the slag powder than the olivine  
132 powder in our preliminary test trials. So, three microcosms were enriched with 100 g of olivine powder, three microcosms  
133 with 2 g of steel slag powder, while the remaining three microcosms were left unperturbed and served as controls.

134

135

## 136 **2.2 Preparation of olivine and steel slag powder**

137 The olivine rocks were provided by Moyne Shire Council who sourced the mineral from a quarry in Mortlake, Victoria,  
138 Australia. The Basic Oxygen Slag (hereafter referred to as “slag”) was provided by Bradley Mansell who sourced the  
139 material from Liberty Primary Steel Whyalla Steelworks in Whyalla, South Australia, Australia. Upon delivery, the olivine  
140 rocks were 40-80 mm in diameter, and slag aggregates were 20-50 mm in diameter. These were crushed to smaller than 10  
141 mm pieces using a hydraulic crusher. The crushed material was further ground with a ring mill with a chrome milling pot.  
142 Afterwards, finely-ground samples were sieved to get samples with 150 ~ 250  $\mu\text{m}$  grain size. The sieved olivine and slag  
143 grains were inspected for their appearance and elemental composition using a Hitachi SU-70 analytical field emission  
144 scanning electron microscope (SEM), and energy dispersive spectrometers (Central Science Laboratory (CSL), University  
145 of Tasmania). Grain size spectra were determined with a Sympatec QICPIC particle size analyser LIXCELL (CSL,  
146 University of Tasmania).

147

## 148 **2.3 Seawater sampling**

149 Seawater was transferred with a peristaltic pump from the microcosms at a depth of about 0.15 m into 1 L acid-washed  
150 sampling bottles (LDPE) using an acid-washed silicon tube. Seawater in these bottles was then subsampled for dissolved  
151 trace metal samples, filtrations, Fast Repetition Rate fluorometry (FRRf), and flow cytometry analysis. Samples for  
152 nutrients and total alkalinity (TA) were transferred using the same pump but through a silicone tube into 80 mL HDPE  
153 bottles. Total alkalinity and macronutrient samples were filtered during this process through a 0.2  $\mu\text{m}$  nylon filter attached  
154 to the silicone tube to remove all particles and organisms  $> 0.2 \mu\text{m}$ .

155

## 156 **2.4 Salinity, nutrients, carbonate chemistry, and trace metal analysis**

157 Salinity was measured before and at the end of the experiment using a HACH HQ40d portable meter. The  $\text{pH}_T$  (total scale)  
158 and temperatures were measured daily (2-3 hours after the onset of the light period) using a pH meter (914  
159 pH/Conductometer Metrohm). We recorded voltages and temperature from the pH meter and calibrated the  $\text{pH}_T$  at original  
160 temperature at sampled time using the certified reference material (CRM) Tris buffer following the method described in  
161 SOP6a by Dickson et al. (2007). Briefly, the standard buffer’s pH and voltage at different temperature gradients were  
162 recorded, and temperature vs. voltage polynomial regression data were generated for calculating calibrated pH values ( $\text{pH}_T$ )  
163 (refer to Eq. 3 in SOP6a of Dickson et al. (2007)). The regression could then be used to obtain a CRM pH value for each  
164 temperature and to calibrate the pH measured in the microcosms to the total pH scale.

165

166 Total alkalinity was sampled every four days. It was measured in duplicate using a Metrohm 862 Compact Titrosampler  
167 coupled with an Aquatrode Plus with PT1000 temperature sensor following the SOP3b open-cell titration protocol  
168 described in Dickson et al. (2007). Filtered TA samples were stored at 8 °C for a maximum of 23 days before measurement.  
169 Titration curves were evaluated using the “calculate” script within PyCO2sys by Humphreys et al. (2022). The carbon  
170 chemistry equilibrium was calculated with the R package “seacarb” Gattuso et al. (2023) from  $pH_T$ , TA, phosphate, silicate,  
171 temperature, and salinities using stoichiometric equilibrium constants from Lueker et al. (2000). Dissolved macronutrients  
172 were measured every second day using standard spectrophotometric methods developed by Hansen and Koroleff (1999)  
173 on the day the samples were taken from the microcosms.

174

175 Dissolved trace metal concentrations were measured four times during the experiment: a few hours before olivine and slag  
176 were added, a few hours after these minerals were added on day 2, near the middle of the experiment on day 13, and at the  
177 end of the experiment on day 22. Sixty mL of seawater was collected using an acid-washed 60 mL syringe, and the seawater  
178 was filtered through 25 mm diameter 0.2  $\mu\text{m}$  pore size polycarbonate filters. Unfortunately, we did not notice that 0.2  $\mu\text{m}$   
179 pore size nylon filters (acid washed) were used during sampling on days 1 and 2 so we refiltered these seawater samples  
180 again using 0.2  $\mu\text{m}$  pore size polycarbonate filters after one month. All seawater samples were diluted approximately 20-  
181 fold by weight using Milli-Q water (18.2  $\text{M}\Omega\cdot\text{cm}$  grade) and acidified using 1 % ultrapure HCl. These samples were  
182 analysed using Sector Field Inductively Coupled Plasma Mass Spectrometry (SF-ICP-MS) employing multiple resolution  
183 settings to overcome major spectral interferences. Due to the presence of abundant major metal ions in our samples, such  
184 as Na and Mg, natural open-ocean seawater from the Southern Ocean with very low trace metal concentrations was diluted  
185 20 times with Milli-Q water and used as a representative blank. The same Southern Ocean seawater was enriched with  
186 different gradients of trace metal standards to calculate the samples’ trace metal concentrations. Five of the total 36 samples  
187 had abnormal trace metal concentrations, and 2 of them were from day 1. We considered values as outliers using the  
188 interquartile range (IQR) criterion on pre-addition data, and if values are more than 10 times higher than replicates, they  
189 are also considered as outliers. These samples containing outliers were excluded from the data analysis (Table S1.). The  
190 major likely source of these metal contaminations is sampling in the temperature control room, where precautions were  
191 insufficiently implemented.

192

## 193 **2.5 Particulate matter and plankton community analysis**

194 Chlorophyll *a* was sampled every second day by filtering the seawater through glass fibre filters (GF/F, pore size = 0.7  $\mu\text{m}$ ,  
195 diameter = 25 mm), and filters were stored in 15 mL polypropylene tubes wrapped with aluminium foil and stored at -80 °C  
196 for 50-70 days before measurement. Each filter was immersed in 10 mL 100 % methanol for 18-20 h to extract chlorophyll  
197 from phytoplankton and these samples were analysed on a Turner fluorometer (Model 10-AU) following the method  
198 described by Evans et al. (1987).

199

200 Phytoplankton flow cytometry samples were fixed with 40  $\mu\text{L}$  of a mixture of formaldehyde-hexamine (18 %:10 % v/w)  
201 added to 1400  $\mu\text{L}$  of seawater sample. All bacteria samples (700  $\mu\text{L}$ ) were fixed with 14  $\mu\text{L}$  glutaraldehyde (Electron-  
202 microscope grade, 25 %). After mixing samples with fixatives, samples were stored for 25 minutes at 10 °C, then flash-  
203 frozen in liquid nitrogen, and stored at -80 °C until measurement 83-86 days later. Directly before the measurement,  
204 samples were thawed at 37 °C. Bacteria samples were stained with SYBR green I (diluted in dimethylsulfoxide) at a final

205 ratio of 1:10000 (SYBR Green I: sample).

206

207 A Cytex Aurora flow cytometer (Cytex Biosciences) was used to quantify the abundance of fluorescing particles such as  
208 phytoplankton or stained bacteria. Phytoplankton groups were distinguished based on their fluorescence signal intensity of  
209 different laser excitation/emission wavelength combinations and forward scatter (FSC). The yellow-green laser (centre  
210 wavelength: 577 nm), in combination with FSC signal strength, was used to separate cyanobacteria and cryptophytes from  
211 other phytoplankton. The violet laser (centre wavelength: 664 nm) in combination with FSC was used to distinguish  
212 picoeukaryotes, nanoeukaryotes, and microphytoplankton. The blue laser (centre wavelength: 508 nm) in combination with  
213 FSC was used to distinguish bacteria from other living (i.e., DNA-containing) particles (Fig. S. 1).

214

215 The biovolume of each classified flow cytometry phytoplankton type was calculated using the equation:

216

$$217 \text{ Biovolume} = \text{Cell number count} \times \left(\frac{\text{FSC}}{10248}\right)^{2.14} \quad (4)$$

218

219 where biovolume is the biovolume of the phytoplankton ( $\mu\text{m}^3$ ), cell number is the cell count per mL of sample, and the  
220 FSC is the forward scatter signal value from the flow cytometry. This equation is calculated based on the relationship  
221 between biovolume and FSC for different phytoplankton species (Selfe, 2022). The biovolume of each phytoplankton type  
222 was then divided by the total biovolume of all phytoplankton type to calculate the biovolume proportion of each  
223 phytoplankton type (Biovolume prop.). This derived value was used to estimate the phytoplankton composition in each  
224 microcosm.

225

226 Phytoplankton photosynthetic performance was estimated from the rapid light curves measured with an FRRf (FastOcean  
227 Sensor FRRf3, Chelsea Instruments Group) every second day following the protocol adapted from Schallenberg et al.  
228 (2020). Samples were kept in the dark for 20 minutes before the measurement and then added to the FRR fluorometry  
229 cuvette, which was temperature-controlled at 13.5 °C. Filtered natural seawater was used for blank correction. A channel  
230 with three light wavelengths (450, 530, and 624 nm) was used in each acquisition sequence. At least 10 acquisitions were  
231 measured for each sample. The maximum electron transport rate ( $\text{ETR}_{\text{max}}$ ), initial slope of the rapid light curve ( $\alpha$ ), and the  
232 light-saturation parameter ( $E_k$ ) were calculated using the equation described by Platt et al. (1980) without photoinhibition:

233

$$234 \text{ ETR} = \text{ETR}_{\text{max}} \left[1 - e^{-\frac{\alpha E}{\text{ETR}_{\text{max}}}}\right] \quad (5)$$

235

236 These parameters together with the maximum quantum yield of PSII ( $F_v/F_m$ ) were used to compare the photosynthetic  
237 performance of the phytoplankton communities in different microcosms.

238

239 Seawater was sampled before the treatment and at the end of the experiment for particulate trace metal concentrations.  
240 Samples of 100 mL were filtered through an acid-cleaned polycarbonate filter (25 mm diameter, 0.8  $\mu\text{m}$  pore size) and  
241 placed in an acid-cleaned polypropylene filter holder in a trace metal-clean laminar flow bench. The filters were washed  
242 with the EDTA-oxalate reagent (1.4 mL) twice (8 min total) and rinsed with chelexed NaCl solution (0.6 mol L<sup>-1</sup> with 2.38  
243 mmol L<sup>-1</sup> of HCO<sub>3</sub><sup>-</sup>, pH=8.2) 10 times (1.5 mL aliquots) (Tovar-Sanchez et al., 2003; Tang and Morel, 2006). Filters were  
244 stored in acid-washed well plates at -20 °C before analysis. The digestion process followed the method reported by Bowie

245 et al. (2010). Briefly, all samples and triplicate certified reference materials plankton standards (50 mg/vial) were digested  
246 in a mixture of strong ultrapure acids (750  $\mu\text{L}$  12 mol  $\text{L}^{-1}$  HCl, 250  $\mu\text{L}$  40 % HF, 250  $\mu\text{L}$  14 mol  $\text{L}^{-1}$   $\text{HNO}_3$ ) in 15 mL Teflon  
247 perfluoroalkoxy (PFA) vials on a 95 °C hot plate for 12 h in a fume hood. They were then dry evaporated for 4 h and re-  
248 suspended in 10 % v-v ultrapure  $\text{HNO}_3$ . All prepared solutions had indium as internal standard added to a final  
249 concentration of 10  $\mu\text{g L}^{-1}$ . Three pre-mixed multi-element standard solutions (MISA) were prepared as external calibration  
250 standards.

251

252 Particulate organic carbon (POC) was sampled by filtering 100 mL of seawater from each microcosm. Glass fibre filters  
253 (Whatman GF/F, pore size = 0.7  $\mu\text{m}$ , diameter = 13 mm) were pre-combusted at 400 °C for 6 h. Filters were stored at -20 °C  
254 before measurement. Samples were treated via fuming with 2N HCl to remove carbonates overnight and dried in the oven  
255 for 4h. Finally, filters were folded into silver cups and stored in a desiccator until analysis. Samples were analysed for  
256 carbon with a Thermo Finnigan EA 1112 Series Flash Elemental Analyser (CSL, University of Tasmania).

257

258 Biogenic silica (BSi) concentrations were analysed every 4 days by filtering 100 mL of seawater from each microcosm.  
259 Mixed Cellulose Ester (MCE) membrane filters (diameter = 25 mm, pore size = 0.8  $\mu\text{m}$ ) were used for BSi samples. BSi  
260 filters were placed in a plastic petri dish and stored at -20 °C before measurement. Filters were processed using the hot  
261 NaOH digestion method of Nelson et al. (1989). The final solution was measured using the same process as the dissolved  
262 silicate (see section 2.4).

263

264 A self-made plastic zooplankton net (20 mm height and 15 mm width) with a 210  $\mu\text{m}$  mesh size was acid-washed first and  
265 then used to collect zooplankton from microcosms before mineral addition on day 2, near the middle (day 13), and at the  
266 end of the experiment (day 23). Samples were stored in 10 % formalin seawater solutions and kept at room temperature  
267 until measurements. Zooplankton were quantified and identified under a Leica M165C microscope fitted with a Canon 5D  
268 camera. The number of zooplankton from one mini-trawl in each collection was converted to the unit of individual  $\text{L}^{-1}$  and  
269 used for data analysis. The diversity of zooplankton communities was estimated with the Shannon Diversity Index (H)  
270 calculated as:

271

$$272 \quad H = -\sum(pi \times \ln(pi)) \quad (6)$$

273

274 where pi is the proportion of the entire zooplankton community made up of individual species abundance, and ln is the  
275 natural logarithm.

276

277

## 278 **2.6 Statistic analysis**

279 R studio was used for data analyses. Generalized additive models (GAMs) from the package “mgcv” were fitted to the data  
280 to predict the changes over time. The GAMs all shared the same equations:

281

$$282 \quad Y = s(\text{Day}), \quad (7)$$

283



284 in which Y presents the dependent variable and s(Day) is the smooth term of the day of the experiment. Another GAM was  
285 used to detect significant differences between treatments and the control:

$$287 \quad Y = Treatment + s(Day) + s(Day, by = oTreatment) \quad (8)$$

288  
289 In this equation, the variable “Treatment” includes three conditions: “Control”, “Slag” and “Olivine”; while “oTreatment”  
290 is the ordered factor of the variable “Treatment” which allowed us to compare the GAMs smooth terms from different  
291 treatments and the control (Simpson, 2017).

292  
293 When comparing GAMs, P-means represent the p-value obtained from comparing two GAMs, such as the control and the  
294 olivine treatment. If P-means is below 0.05, it indicates that the mean values of the two GAMs exhibit significant  
295 differences over the course of the experiment. Conversely, if P-means is equal to or greater than 0.05, it suggests that the  
296 two GAMs have similar mean values. In contrast, P-smooths represents the p-value derived from comparing the smooth  
297 terms of two GAMs. If P-smooths is below 0.05, it indicates that the two GAMs demonstrate significantly different trends  
298 in their change over time.

299  
300 For the analysis of trace metal concentrations and zooplankton abundance, Generalized Linear Models (GLMs) from the  
301 'stats' package were fitted to the data to determine significant differences between treatments and the control. The selection  
302 of specific GLMs was based on the distribution of the raw data. One GLM equation is

$$304 \quad Y = Treatment + \frac{Day}{22} + \left(\frac{Day}{22}\right)^2 \quad (9)$$

305  
306 with family = Gamma, where Y represents the measured parameter (abundance of a zooplankton species and dissolved  
307 trace metal concentrations); treatment is the conditions (“Control”, “Slag” and “Olivine”); and Day represents the day of  
308 the experiment. The other GLM equation,

$$310 \quad Y = Treatment + Day \quad (10)$$

311  
312 with family = Gaussian, was employed for particulate trace metal data and the Shannon Diversity Index. To compare the  
313 contribution of the three treatments on the measured parameters, Tukey's significant difference test was conducted on the  
314 GLMs using the 'glht' function.

315

### 316 **3. Results**

#### 317 **3.1 Elemental composition and grain size of the finely-ground minerals**

318 SEM analysis revealed the approximate elemental composition of olivine and slag powder (Table 1). Based on this analysis  
319 the olivine composition resembles the Mg-rich olivine mineral “forsterite” (Mg<sub>2</sub>SiO<sub>4</sub>). The particle size spectrum of olivine  
320 powder is shown in detail in Fig. S2. Roughly 69 % of the olivine particles, when measured by volume, fell within the

321 diameter range of 35 - 300  $\mu\text{m}$ . Additionally, SEM analysis revealed high levels of Ca and O in the slag, indicative of the  
 322 considerable  $\text{Ca}(\text{OH})_2$  and CaO content of the powder (Table 1; please note that H cannot be measured with the applied  
 323 method). The particle size measurement (Fig. S2) showed that 78 % of the ground slag particles were between 35 - 300  
 324  $\mu\text{m}$ .

325

326 **Table 1.** The weight percentage of elements from two minerals. Unit: wt %.

Element	O	Ca	Mn	Si	Mg	Fe	Al	Ti	Cr	Ni
Olivine	39.9	0.4		19.9	26.4	13.0	1.0			0.8
Steel slag	41.9	36.0	7.0	6.5	4.3	3.7	3.4	1.7	1.6	


327

328

### 329 3.2 Physical and chemical conditions over the course of the experiment.

330 On day 2 of the experiment, when olivine particles were introduced into the microcosms, the smallest fraction of the powder  
 331 remained suspended, causing the seawater to become highly turbid for several days. The resulting milky appearance of the  
 332 seawater eventually faded over a period of approximately five days, and by day 5, the turbidity had visually become like  
 333 the slag treatment and the control. This effect was not anticipated, and as a result, we decided to investigate its impact on  
 334 light intensity. To do so, a test was conducted after the main experiment in which olivine powder was added to a microcosm  
 335 identical to those used in the experiment, and light intensity was measured daily at a depth of 0.15 m. The results showed  
 336 that the addition of olivine caused an initial reduction in light intensity of 18.5 % at 15 mins after addition, which declined  
 337 to 7.4 %, 3.7 %, 3.7 % and 0 % after 1, 2, 3, and 4 days, respectively. These findings indicate that olivine additions can  
 338 significantly affect the light environment in the microcosms, whereas no such effect was observed in the slag treatment.

339



340

341 **Fig. 2.** Carbonate chemistry conditions. The temporal development of (a)  $\text{pH}_T$ , (b) temperature, (c) total alkalinity (TA), (d)  $\text{CO}_2$  fugacity

342 ( $f\text{CO}_2$ ) computed at *in situ* temperature and atmospheric pressure, (e) dissolved inorganic carbon (DIC), and (f) aragonite saturation state  
343 ( $\Omega_{\text{aragonite}}$ ). The dots represent the raw data ( $n=3$  for each treatment per sampling time), and the fitted curve is the generalized additive  
344 model (GAM). The shading represents the 95 % confidence interval of the fitted GAM.

345

346 The  $\text{pH}_T$  of all microcosms increased from day 1 to day 5 (Fig. 2a). This was due to photosynthetic  $\text{CO}_2$  drawdown in the  
347 control or photosynthetic  $\text{CO}_2$  drawdown in combination with alkalinity release from minerals in the treatments. During  
348 the peak of the bloom,  $\text{pH}_T$  was  $8.037 \pm 0.010$  in the control (average values  $\pm$  standard error),  $8.054 \pm 0.014$  in the olivine  
349 treatment and  $8.411 \pm 0.015$  in the slag treatment. The  $\text{pH}_T$  was significantly higher in the slag than the olivine treatment  
350 and the control throughout the experiment (control and olivine  $\text{pH}_T$  were not significantly different). The  $\text{pH}_T$  on day 23 of  
351 the control, olivine, and slag treatments were  $7.893 \pm 0.012$ ,  $7.978 \pm 0.015$ , and  $8.309 \pm 0.019$ , respectively. The temperature  
352 inside of the microcosms varied between replicates, which may have added noise in the biological response data. However,  
353 on average there was no statistically significant difference between control/treatments during the experiment.

354


355 In our data analysis, all the fitted GAMs from the treatments and the control exhibited significant differences in  $\text{pH}_T$  from  
356 each other, as evidenced by the p-values of both P-means and P-smooths being smaller than 0.001. For detailed results of  
357 the GAM p-values, please refer to Table S2.

358

359 Total alkalinity increased marginally from  $2255 \pm 2$  to  $2262 \pm 13 \mu\text{mol kg}^{-1}$  within the first 6 days after olivine addition  
360 while it increased more substantially from  $2259 \pm 1$  to  $2522 \pm 11 \mu\text{mol kg}^{-1}$  in the same time span in the slag treatment (Fig.  
361 2c). The TA in the control decreased from  $2261 \pm 2 \mu\text{mol kg}^{-1}$  to  $2240 \pm 7 \mu\text{mol kg}^{-1}$  from day 1 to day 6 but remained  
362 stable thereafter. The TA reached  $2279 \pm 6 \mu\text{mol kg}^{-1}$  in the olivine treatment and  $2611 \pm 9 \mu\text{mol kg}^{-1}$  in the slag treatment  
363 on day 22. The slag treatment reached a significantly higher TA than the olivine treatment and the control (P-smooths <  
364 0.001). The mean TA from GAM in olivine treatment was higher than the control (P-means < 0.001).

365

366 The  $\text{CO}_2$  fugacity ( $f\text{CO}_2$ ) computed at *in situ* temperature and atmospheric pressure decreased continuously in the first 6  
367 days in all microcosms (Fig. 2d). Then it increased again in the control and olivine treatments while staying lower in the  
368 slag treatment (P-means and P-smooths  $\leq 0.001$  between either treatment or the control). Dissolved inorganic carbon (Fig.  
369 2e) and the aragonite saturation state ( $\Omega_{\text{aragonite}}$ ; Fig. 2f) revealed a similar trend over the course of the experiment in the  
370 control and the olivine treatment. In contrast, the slag treatment had higher DIC and  $\Omega_{\text{aragonite}}$  values throughout the  
371 experiment (P-means < 0.001).




372

373 **Fig. 3.** Macronutrients concentrations over the course of the study. (a) Nitrate and nitrite concentrations. (b) Phosphate concentrations.  
 374 (c) Silicic acid concentrations. The dots represent the raw data ( $n=3$  for each treatment per collection), and the fitted curve is the  
 375 generalized additive model.

376

377 Initial nitrate and nitrite ( $\text{NO}_x^-$ ), phosphate ( $\text{PO}_4^{3-}$ ), and silicic acid ( $\text{Si}(\text{OH})_4$ ) concentrations were  $1.58 \pm 0.12$ ,  $0.69 \pm 0.59$ ,  
 378 and  $8.04 \pm 0.10 \mu\text{mol L}^{-1}$ , respectively (Fig. 3).  $\text{NO}_x^-$  declined rapidly in all microcosms once the experiment had  
 379 commenced to values below  $0.5 \mu\text{mol L}^{-1}$  and no significant difference was detected between treatments and control ( $P$ -  
 380 smooths  $>0.05$ ; Fig. 3a). In both the olivine treatment and the control, the  $\text{PO}_4^{3-}$  concentration decreased in the first six  
 381 days (Fig. 3b). In the slag treatment,  $\text{PO}_4^{3-}$  increased to a maximum of  $2.65 \pm 0.01 \mu\text{mol L}^{-1}$ , which was significantly higher  
 382 than in the olivine treatment and the control ( $P$ -means  $<0.001$ ). The  $\text{Si}(\text{OH})_4$  concentration increased to a maximum of  
 383  $15.99 \pm 0.87 \mu\text{mol L}^{-1}$  in the olivine treatment, increased to a maximum of  $41.92 \pm 1.75 \mu\text{mol L}^{-1}$  in the slag treatment, but  
 384 decreased below the detection limit in the control (Fig. 3c). Significant differences were observed in the development of  
 385  $\text{Si}(\text{OH})_4$  between all treatments and the control (Table S2).

386



387

388

389 **Fig. 4.** Dissolved and particulate trace metal concentrations in microcosm seawater. (a)-(d) are dissolved trace metal concentrations, and  
 390 (e)-(h) are total particulate trace metal concentrations. The error bars represent the standard error from measured samples. The pre-  
 391 addition data shown in (a)-(d) represent the average of 7 microcosms before addition of slag or olivine. The data for the control on day  
 392 22 in (a)-(d) and for the pre-addition on day 1 in (e)-(h) were based on two of three microcosm replicates. The remaining data were based  
 393 on all three microcosm replicates.

394

395 The dissolved trace metal concentrations measured from microcosms are presented in Fig. S3. While the mass of olivine  
 396 added to the microcosms was 50-fold greater than in steel slag (100 g vs 2 g), it's noteworthy that the variation in dissolved  
 397 trace metal concentrations between the two treatments were much smaller than 50 folds. After 21 days of experiment, the  
 398 treatments showed an increase in dissolved Al concentrations from  $920 \pm 286$  to  $970 \pm 228 \text{ nmol L}^{-1}$  in olivine treatment,  
 399 and from  $920 \pm 286$  to  $1093 \pm 77 \text{ nmol L}^{-1}$  in slag treatment, while in the control dissolved Al decreased to  $230 \pm 10 \text{ nmol}$

400 L<sup>-1</sup> (Fig. S3). The fitted GLMs were compared, and the p-value revealed how much influence a treatment had on the  
401 dissolved metal concentrations (Table S3). The results indicate that the slag and olivine additions led to significantly higher  
402 Al concentrations than in the control (p-values < 0.05), but no significant difference was found between the two treatments  
403 (p-value = 0.189). The Cu concentration in the olivine on day 22 was significantly higher than the slag treatment and the  
404 control (p-value < 0.05) (Fig. S3). The addition of olivine and slag released some dissolved Fe, but overall, the concentration  
405 of Fe did not differ significantly between treatments (Fig. 4a, Table S3). The slag released a substantial amount of dissolved  
406 Mn (maximum 810 ± 10 nmol L<sup>-1</sup> on day 22) (Fig. 4b), leading to significantly higher concentrations than in the olivine  
407 treatment and the control (p-values < 0.001). A significant amount of dissolved Ni (maximum 77 ± 3 nmol L<sup>-1</sup> on day 22)  
408 was released from the olivine powder (p-values < 0.001) (Fig. 4c). The initial concentration of dissolved Zn in seawater  
409 was much higher than on day 22 in all microcosms, and no significant difference in Zn concentrations was found between  
410 the treatments and the control.

411

412 Particulate concentrations of some trace metals also differed between treatments. The total particulate Fe decreased in all  
413 microcosms on day 22 comparing with the pre-addition level, but both mineral addition treatments had higher particulate  
414 Fe concentrations than the control (Fig. 4e). The addition of slag elevated particulate Mn concentrations to a level higher  
415 than the pre-addition and the control on day 22 (Fig. 4f), while the addition of olivine increased the particulate Ni  
416 concentrations to a level higher than the slag, the control, and the pre-addition (Fig. 4g). The particulate Zn concentrations  
417 in general decreased by the end of the experiment (Fig. 4h), and no significant differences were found between the  
418 treatments and the control.


419

420 The POC on day 1 and day 22 from all microcosms were very similar, 10.99 ± 0.58 and 11.03 ± 0.41 μmol L<sup>-1</sup> respectively  
421 (Fig. S4) so the metal:POC results were consistent with the particulate trace metal results (Fig. 4 e-h). In general, the non-  
422 surface metal:POC are positively correlated with the total metal:POC ratios (Fig. S5). The ratio of non-surface to total  
423 particulate trace metal concentrations is summarized in Table S5. Both non-surface and total Fe concentrations decreased  
424 in microcosms on day 22 compared with the pre-addition level. Iron:POC ratios were significantly higher in the treatments  
425 than in the control on day 22 (p-values < 0.05. Table S3), and there was no significant difference between mineral addition  
426 treatments. The non-surface to total Fe:POC ratios were > 0.94 in all microcosms on both day 1 and day 22. The total and  
427 non-surface Mn:POC ratio was the highest in the slag treatment. These ratios were higher than the pre-addition level and  
428 the control at the end of the experiment. The total particulate Ni concentrations in the olivine treatment were significantly  
429 higher than before olivine addition. The olivine treatment led to a >22-fold higher Ni:POC ratio compared to the other two  
430 treatments (p-value < 0.001).

431

432

433



435  
 436 **Fig. 5.** Temporal development of chlorophyll a concentration (chl-a), BSi, and different eukaryotic and bacterial plankton groups as  
 437 determined with flow cytometry. (a) chlorophyll a; (b) BSi; cell concentrations of (c) heterotrophic bacteria, (d) microphytoplankton, (e)  
 438 nanoeukaryotes2, (f) nanoeukaryotes1 (g) picoeukaryotes, (h) cyanobacteria, and (i) cryptophytes; biovolume proportion of (j)  
 439 microphytoplankton, (k) nanoeukaryotes2, (l) nanoeukaryotes1 (m) picoeukaryotes, (n) cyanobacteria, and (o) cryptophytes. The figure  
 440 data points represent the raw data, and the fitted curve is the generalized additive model. The shaded area represents the 95 % confidence  
 441 interval.

442

443 The chl-a concentration in all microcosms increased from day 1 to day 4 from  $1 \mu\text{g L}^{-1}$  to  $3\text{-}4 \mu\text{g L}^{-1}$  (Fig. 5a). The chl-a  
444 concentration then decreased rapidly from day 4 to day 8, then continued to decrease, though more slowly, to  $<0.3 \mu\text{g L}^{-1}$   
445 until the end of the experiment. The GAMs of chl-a did not show any difference between treatments and the control (both  
446 P-means and P-smooths  $>0.05$ , see Table S2).

447

448 The BSi concentration increased from day 1 to day 6 in all microcosms (Fig. 5b). In the olivine treatments, BSi  
449 concentrations decreased slightly after the peak until day 12 but then increased again. In the slag treatment, BSi  
450 concentrations remained relatively stable after the initial phytoplankton bloom. In contrast, BSi concentration decreased  
451 continuously in the control after the initial peak. Olivine particles suspended in seawater after the mineral addition (see  
452 section 3.2) partially ended up on BSi filters during filtration. This led to extremely high BSi measurements on days 2 and  
453 4 that were removed from Fig. 5b. Without these outliers, the mean of fitted BSi GAM in the olivine treatment was lower  
454 than the control and the slag treatment (Table S2), and the slag treatment had the highest average BSi over the course of  
455 the experiment. Overall, the BSi trends in the two treatments were similar (P-smooths = 0.269), and both were significantly  
456 different from the control (P-smooths  $<0.05$ ).

457


458 The development of the phytoplankton community composition showed significant differences between the treatments and  
459 the control. In general, most phytoplankton groups exhibited similar patterns to chl-a, with peak cell numbers occurring on  
460 day 4 (Fig. 5f-i) apart from microphytoplankton and nanoeukaryotes2 which had the peak delayed for 1-2 days (Fig. 5d-  
461 e). Please be aware that flow cytometers may not capture some large and chain-forming phytoplankton. After reaching  
462 peak values during the bloom, phytoplankton abundance generally decreased steadily. Microphytoplankton displayed  
463 similar trends to the results for BSi. Before day 10, all microcosms had similar microphytoplankton abundances (Fig. 5d).  
464 However, in the control, microphytoplankton abundance declined continuously and at a faster rate compared to the two  
465 treatments (P-smooths values  $<0.03$ ). From day 2 to day 6, the abundance of nanoeukaryotes1, nanoeukaryotes2,  
466 picoeukaryotes, and cryophytes was higher in the olivine treatment compared to the slag treatment and the control. After  
467 day 8, their abundance in the olivine treatment decreased to a similar level as the slag treatment and the control. Notably,  
468 there were few significant differences observed between the slag treatment and the control in terms of the abundances of  
469 nanoeukaryotes1, nanoeukaryotes2, picoeukaryotes, cyanobacteria, and cryptophytes throughout the experiment. In the  
470 olivine treatment, cyanobacteria experienced a second bloom after day 10, which was significantly different from the other  
471 two groups (P-smooths  $<0.01$ ). Heterotrophic bacteria exhibited an increase and decline pattern following the  
472 phytoplankton bloom until day 8 (Fig. 5c). Subsequently, bacteria abundance increased again, reaching a second peak  
473 during days 12-14, followed by a decline until the end of the experiment. The decline in bacteria abundance was slower in  
474 the olivine treatment, although no significant differences were detected between treatments (Table S2).

475

476 Among all the microcosms, microphytoplankton consistently accounted for the largest proportion of biovolume. From the  
477 perspective of biovolume proportion, the mineral addition mainly influenced the microphytoplankton and nanoeukaryotes.  
478 The control had similar phytoplankton biovolume distribution as the treatments from day 1 to day 15, but after that the  
479 proportion of microphytoplankton biovolume decreased to a level significantly lower than the treatments. In the control  
480 treatment, the proportion of nanoeukaryotes' biovolume increased as the proportion of microphytoplankton decreased. The  
481 biovolume of picoeukaryotes, cyanobacteria and cryptophytes increased during the phytoplankton bloom and then  
482 decreased drastically after the bloom. There were no significant differences in biovolume proportion observed for

483 picoeukaryotes, cyanobacteria and cryptophytes between the treatments and the control.

484



485

486 **Fig. 6.** The photosynthetic performance of the phytoplankton community. (a)  $F_v/F_m$ , the maximum quantum yield of photosynthesis II.  
487 (b)  $\alpha$ , the initial slope of the rapid light curves. (c)  $ETR_{max}$  is the maximum electron transport rate, the maximum potential photosynthetic  
488 rate. (d)  $E_k$  is light-saturation parameter, Unit:  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ .


489

490 The temporal development of  $F_v/F_m$ ,  $\alpha$ ,  $ETR_{max}$ , and  $E_k$  is illustrated in Fig. 6. The  $F_v/F_m$  values of the phytoplankton  
491 community were approximately  $0.42 \pm 0.01$  and increased to levels  $> 0.5$  during the peak of the phytoplankton bloom on  
492 day 4 (Fig. 6a). Following the bloom,  $F_v/F_m$  values dropped below 0.3 in the control. However, the decline in  $F_v/F_m$  after  
493 the bloom was less pronounced in the two mineral addition treatments with the olivine treatment maintaining higher  $F_v/F_m$   
494 values than the slag treatment (P-smooths  $< 0.05$ ). At the end of the experiment,  $F_v/F_m$  was  $0.22 \pm 0.04$  in the control,  $0.35$   
495  $\pm 0.01$  in the slag treatment, and  $0.42 \pm 0.02$  in the olivine treatment. The temporal development of  $\alpha$  aligned with the  
496 patterns observed for  $F_v/F_m$  (compare Fig. 6a and 6b). The maximum values of  $ETR_{max}$  were observed on day 4 in the  
497 control and the slag treatment, while in the olivine treatment, it occurred on day 5 (Fig. 6c). Subsequently,  $ETR_{max}$   
498 continuously decreased until day 10 and then stabilized until the end of the experiment. However,  $ETR_{max}$  exhibited a  
499 subsequent increase in the mineral treatments around day 12. The  $ETR_{max}$  values were higher in the mineral treatments  
500 compared to the control group (P-means  $< 0.001$ , Table S2). The parameter  $E_k$  decreased from  $246 \pm 17 \mu\text{mol photons m}^{-2}$   
501  $\text{s}^{-1}$  on day 1 to  $121 \pm 7 \mu\text{mol photons m}^{-2} \text{s}^{-1}$  on day 10, and then it increased again to approximately  $200 \mu\text{mol photons m}^{-2}$   
502  $\text{s}^{-1}$  by the end of the experiment (Fig. 6d). The change in  $E_k$  did not exhibit significant differences between the treatments  
503 and the control (both P-means and P-smooths  $> 0.05$ ).

504

505





506

507 **Fig. 7.** The dominant zooplankton abundance and community diversity from different treatments. Abundance of dominant zooplankton  
 508 in microcosms: (a) calanoid copepod; (b) *Noctiluca scintillans*; (c) *Oikopleura* sp.; (d) *Oithona* sp.; (e) *Penilia* sp.; and (f) the Shannon  
 509 diversity index (H) of different treatments and the control. Error bars represent the standard error calculated from three microcosm  
 510 replicates. Photographs of each zooplankton group are shown on the corresponding graphs.

511

512 Thirteen zooplankton taxonomic groups were identified in the microcosms. The dominant taxa were the appendicularian  
 513 *Oikopleura* sp., the cyclopoid copepod *Oithona* sp., the cladoceran *Penilia* sp., the heterotrophic dinoflagellate *Noctiluca*  
 514 *scintillans* and several calanoid copepods including *Acartia* sp., *Paracalanus* sp. and *Gladioferens* sp. The larvae and eggs  
 515 of *Oikopleura*, *Penilia* and copepod were also observed under the microscope. In general, higher zooplankton numbers  
 516 were observed after the bloom on day 13 (Fig. 7). The abundance of calanoid copepods and *Oithona* sp. increased after  
 517 day 2 (Fig. 7a, d), and there was no significant difference between treatments and the control (p-values >0.05, Table S4).  
 518 The abundance of *N. scintillans* increased significantly more in the olivine treatment than in the control and the slag  
 519 treatment, with highest abundance of  $13 \pm 9$  individual  $L^{-1}$  observed in the olivine treatment on the last day (Fig. 7b). The  
 520 abundance of *Oikopleura* in the control and the slag treatment was higher than the olivine treatment on day 13 but was  
 521 higher in the olivine treatment on day 22 (Fig. 7c). A higher abundance of *Penilia* sp. was found in the slag treatment on  
 522 day 13 and in the control on day 23 (Fig. 7e). Due to the patchy distribution of zooplankton, these data have large standard  
 523 errors and only the differences in the numbers of *N. scintillans* in the olivine treatment were statistically significantly  
 524 different from the slag treatment and the control (p-value <0.05, Table S4).

525

526 Considering the control and slag treatment, the Shannon Diversity Index (H) increased from day 2 to day 13 and declined  
 527 on day 23, while in the olivine treatment, H was lower on day 13 than on day 2 and day 23 (Fig. 7f). The GLMs revealed  
 528 that the olivine treatment had significantly lower H on day 13 than the control and the slag treatment (p-values <0.001).  
 529 There were no significant differences in H between the control and the slag treatment (Table S4). The addition of olivine  
 530 decreased the zooplankton community's diversity. This is mainly driven by distinct trends observed in the abundance of  
 531 *Oikopleura* sp., *Penilia* sp., and *N. scintillans* (Fig. 7).

532

## 533 4. Discussion

### 534 4.1 CO<sub>2</sub> removal potential of slag and olivine

535 The slag powder created significantly higher CO<sub>2</sub> removal potential than the olivine powder over the course of the study.  
536 Ca(OH)<sub>2</sub> and CaO in slag and Mg<sub>2</sub>SiO<sub>4</sub> in olivine are likely to be the main functional minerals driving the measured  
537 alkalinity enhancement. Total alkalinity increased by 361 μmol kg<sup>-1</sup> in the slag treatment while it increased by only 29  
538 μmol kg<sup>-1</sup> in the olivine treatment, equivalent to a potential increase in marine inorganic carbon by 14.7 and 0.9% within  
539 3 weeks of their application. When normalizing these alkalinity increases to the same material weight, 1 g of slag would  
540 release 9626 μmol TA while 1 g of olivine would release 16 μmol TA. Thus, over 3 weeks of experimental incubation, slag  
541 is ~600-fold more efficient in releasing alkalinity for particles of this size class (please note that particle size spectra of  
542 olivine and slag were similar but not identical; Fig. S1). We can also use these values to make a rough estimate of how  
543 much CO<sub>2</sub> these two minerals could potentially sequester. One mole of alkalinity from olivine and slag can sequester  
544 approximately 0.85 mole of CO<sub>2</sub>. Thus, one tonne of slag and olivine powder as used here could sequester 360 and 0.6 kg,  
545 respectively, within 3 weeks. It is likely that optimization of particle size and application method may lead to higher  
546 efficiencies. Nevertheless, the slag showed potential as an OAE source mineral, even when applied as relatively coarse  
547 powder in this experiment.

548

### 549 4.2 Environmental implications of slag and olivine additions

550 The amount of olivine and slag powder added to the treatments differed significantly (100 g of olivine powder were added  
551 while only 2 g of slag powder were added to the 53 L microcosms). Our rationale for these different mass additions was to  
552 yield somewhat similar amounts of detectable alkalinity enhancement in the dissolved phase, since we already knew from  
553 tests before the experiment that slag elevates alkalinity faster than olivine. However, olivine was less efficient in releasing  
554 alkalinity than we had anticipated so that even a 50-fold higher addition of olivine (in mass) did not compensate for this  
555 difference. As such, our experiments are associated with an “apples and oranges issue” in that our perturbation with  
556 minerals and associated OAE differs. ~~We note that an adjusted addition of minerals depending on the alkalinity~~  
557 ~~enhancement rate would be consistent with what OAE practitioners may do under real-world conditions. Presumably, OAE~~  
558 ~~deployments may have to adjust the amounts of minerals to detect alkalinity enhancement in the dissolved phase for~~  
559 ~~verification purposes. Nevertheless, to account for this “apples and oranges issue”, the following discussion mainly~~  
560 relates the observed environmental effects with the alkalinity enhancement achieved over the course of the study.

#### 561 4.2.1. OAE effects on phytoplankton physiology and community

562 Previous research has hypothesised that OAE-induced changes in seawater carbonate chemistry could delay phytoplankton  
563 bloom formation due to reductions in seawater pCO<sub>2</sub> in the aftermath of an OAE deployment (Bach et al., 2019). The build-  
564 up of chlorophyll *a* concentration as observed here was indistinguishable between treatments and the control, suggesting  
565 no effect of slag- or olivine-based OAE on phytoplankton bloom dynamics under these experimental settings. A lack of  
566 bloom delay due to carbonate chemistry is unsurprising for the olivine treatment where the release of alkalinity was small

567 (29  $\mu\text{mol kg}^{-1}$  alkalinity release), but somewhat more surprising in the slag treatment where alkalinity was quite rapidly  
568 increased by 361  $\mu\text{mol kg}^{-1}$ . However, the release was still lower than in a very similar study by Ferderer et al., (2022)  
569 where alkalinity was increased by 500  $\mu\text{mol kg}^{-1}$  using sodium hydroxide and even there they did not observe a bloom  
570 delay. Based on this very limited evidence, it seems that bloom delays do not occur consistently under OAE within the  
571 alkalinity ranges tested in this study.

572

573 The nutrient data show that the phytoplankton community was most likely N-limited after day 4 so that the release of  
574  $\text{Si(OH)}_4$  from olivine and  $\text{Si(OH)}_4$  and  $\text{PO}_4^{3-}$  from slag did not stimulate a further increase in chlorophyll-*a* concentration  
575 in the treatments. The development of BSi concentrations is indicative of the prevalence of diatoms in the microcosms but  
576 differences between treatments and the control were small. The release of  $\text{Si(OH)}_4$  through olivine and slag will most likely  
577 benefit diatoms but this fertilization effect did not manifest in this specific experiment because N was limiting diatom  
578 growth. However, when new N is supplied then diatoms will likely take a bigger share of the limiting N pool when olivine  
579 or slag are used for OAE, as has been shown in  $\text{Si(OH)}_4$  manipulation experiments in and outside the context of OAE  
580 research (Egge and Jacobsen, 1997; Ferderer et al., 2023). In the case of slag, the release of  $\text{PO}_4^{3-}$  will likely be another  
581 driver that affects plankton productivity and community composition. As for  $\text{Si(OH)}_4$ , however, the effect of additional  
582  $\text{PO}_4^{3-}$  did likely not materialise in this experiment because  $\text{PO}_4^{3-}$  was not limiting over the course of the study. However, in  
583 ecosystems where  $\text{PO}_4^{3-}$  is a limiting resource, the application of slag could enhance productivity with associated benefits  
584 for higher trophic levels. In contrast, excessive applications of slag and concomitant  $\text{PO}_4^{3-}$  release could also pose a risk of  
585 eutrophication. Future studies may need to investigate what the most sustainable dose of OAE via olivine and/or slag  
586 applications could be and the suitable regions for application.

587

588 The flow cytometry results further revealed the change in phytoplankton community composition. Both the olivine and  
589 slag treatments sustained higher microphytoplankton abundances after the peak of the phytoplankton bloom. This trend is  
590 consistent with higher  $F_v/F_m$  values in the treatments than in the control so that it is tempting to assume that  
591 photophysiological fitness gain measured with the FRRf led to higher competitiveness of microphytoplankton in the  
592 community. Indeed, calculations of the contribution of different phytoplankton groups to total biovolume based on flow  
593 cytometry indicate that microphytoplankton were predominantly contributing to the phytoplankton community biovolume  
594 so that the responses measured by the FRRf were probably to a large extent driven by this group.

595

596 Apart from the increased microphytoplankton abundance, for the slag treatment, other phytoplankton groups distinguished  
597 with flow cytometry did not deviate considerably from the control. The olivine addition, however, triggered more  
598 pronounced shifts in the phytoplankton community. In particular, the nanoeukaryotes (roughly between 2-20  $\mu\text{m}$ ),  
599 picoeukaryotes and the cryptophytes showed relatively higher abundance during the peak of the phytoplankton bloom, and  
600 the abundance of cyanobacteria was higher after the bloom. We speculate that this shift following olivine treatment may  
601 be attributable to a top-down effect from the decrease in zooplankton grazing effects in microcosms, which will be  
602 discussed in section 4.2.2.

603

604 The measurement of photophysiological parameters revealed that the phytoplankton had generally better photosynthetic  
605 performance in the slag and olivine treatments than in the control, especially after the phytoplankton bloom. During the  
606 first 5 days, the changes in phytoplankton photosynthetic performance were indistinguishable between the control and the

607 slag treatment, while the values of  $\alpha$ ,  $ETR_{max}$  and  $F_v/F_m$  were lower in olivine treatment. At this time all microcosms had  
608 similar health because of the relatively high  $NO_x^-$  concentrations and Fe supply (around  $500 \text{ nmol L}^{-1}$ ), but the suspended  
609 particles in the olivine treatment may have led to artifacts in the measuring of photophysiology by FRRf. Scattering and/or  
610 absorption of light by suspended olivine particles is the most parsimonious explanation for the simultaneous depression in  
611  $\alpha$ ,  $ETR_{max}$  and  $F_v/F_m$ . After day 5, the  $F_v/F_m$ ,  $\alpha$  and  $ETR_{max}$  values decreased significantly faster in the control than in the  
612 treatments, and to values lower than the initial condition. A decrease of  $F_v/F_m$  is commonly associated with physiological  
613 stress, such as nutrient limitation, and high light stress (Bhagooli, et al., 2021), with Fe limitation causing a more  
614 pronounced decline in  $F_v/F_m$  than nitrogen limitation (Gorbunov, et al., 2021). The  $ETR_{max}$ , which represents the maximum  
615 electron transport rate, has also been shown to be negatively affected when phytoplankton experience nitrogen or Fe  
616 limitation (Kolber et al., 1994; Gorbunov & Falkowski 2021). Furthermore, the change in photosynthesis performance  
617 after day 10 was suspected to be driven by the microphytoplankton because the decrease of  $F_v/F_m$ ,  $\alpha$ , and  $ETR_{max}$  in the  
618 control was coupled with the decrease in microphytoplankton abundance while the other phytoplankton groups were in  
619 low abundance as in the mineral addition treatments, and the microphytoplankton contributed significantly (75 %) to  
620 community biovolume. All microcosms were similarly  $NO_x^-$  limited from day 5 onward (Fig. 3) so that N-limitation is  
621 unlikely to explain different trends in photophysiological parameters between the control and OAE treatments. Trace metals,  
622 especially Fe, released through slag and olivine additions could potentially explain these differences.

623

624 Several of the trace metals released from slag and olivine are required for photosynthesis. For example, Fe is required for  
625 many proteins functioning in photosynthesis, such as cytochromes, ferredoxin, and superoxide dismutase (SOD) (Twining  
626 and Baines, 2013), and the addition of Fe can stimulate the growth of phytoplankton (Sunda and Huntsman, 1997) and  
627 increase  $F_v/F_m$  (Behrenfeld et al., 2006). The dissolved and particulate Fe concentrations were higher in mineral addition  
628 treatments than in the control indicating potentially more Fe available to sustain phytoplankton photosynthesis. While this  
629 explanation is intriguing for the observed trends in photophysiology, it remains unclear why such strong differences  
630 occurred between mineral addition and control treatments despite dissolved Fe concentrations of  $\sim 500 \text{ nmol L}^{-1}$  at the end  
631 of the experiment in the control. In Fe-limited ocean regions, dissolved Fe is at least two orders of magnitude lower, and  
632 the enhancement of Fe to  $\sim 1.5 \text{ nmol L}^{-1}$  can induce major phytoplankton blooms and relieve photophysiological stress (De  
633 Baar et al., 2005). It is possible that these coastal phytoplankton species have higher Fe requirements than those from the  
634 open ocean where Fe is limiting (Strzepek and Harrison, 2004). Our findings suggest that Fe perturbations may not only  
635 be relevant for low Fe open ocean regions but could also be relevant for coastal ocean locations.

636

637 Alternatively, the addition of Mn, Ni and other trace metals from mineral addition may have benefited photosynthesis.  
638 Manganese is required for the water-splitting reaction of photosystem II (Armstrong, 2008), and both Mn and Ni are  
639 common bioactive trace metals for SODs in marine phytoplankton. The noxious superoxide anion radical ( $O_2^-$ ) generated  
640 from aerobic respiration and oxygenic photosynthesis could be harmful to phytoplankton physiology, and SOD removes  
641  $O_2^-$ , thus improving photosynthesis (Wafar et al., 1995; Wolfe-Simon et al., 2005). This is consistent with our  
642 photosynthetic measurements. Interestingly, although the amounts and types of trace metals released from the slag and  
643 olivine powders were different, they led to relatively similar  $F_v/F_m$  values with only slightly higher  $F_v/F_m$  in the olivine  
644 than the slag treatment from days 10-21. Over this time, these trace metal additions could have fertilized different  
645 phytoplankton species (Pausch et al., 2019; Balaguer et al., 2022; Guo et al., 2022) possibly because different  
646 phytoplankton could have different trace metal requirements, such as for SOD. For example, cyanobacteria have NiSOD,

647 diatoms have MnSOD, dinoflagellates have both FeSOD and MnSOD (Wolfe-Simon et al., 2005). Another explanation is  
648 that phytoplankton in the control were limited by bicarbonate while the treatments had sufficient bicarbonate from added  
649 minerals. However, we were unable to determine the species-level changes in the phytoplankton community, and hence  
650 whether these trace metals, individually or combined, could account for the observed phytoplankton community  
651 photosynthetic performance.

652

#### 653 **4.2.2. OAE impacts on the zooplankton community**

654 Slag-based OAE did not significantly influence the zooplankton community composition while olivine-based OAE induced  
655 some statistically significant effects, including a lower Shannon diversity. The increase in *N. scintillans* abundance and the  
656 decrease in *Penilia* sp. and *Oikopleura* sp. in the olivine treatment indicate that the zooplankton response to OAE can vary  
657 among different zooplankton types.

658

659 The observed lower abundance of *Oikopleura* sp. on day 13 in the olivine treatment may indicate a temporary suppression  
660 or a slower growth rate of this zooplankton species in response to the olivine addition. This could be attributed to the  
661 potential effects of olivine on the availability of essential nutrients or changes in the physicochemical environment of the  
662 water. However, the subsequent increase in *Oikopleura* sp. abundance by day 22 suggests that the growth of this species  
663 may have recovered or accelerated in the olivine treatment, leading to a higher abundance compared to the slag treatment  
664 and the control on day 22. As discussed in section 4.2.1, reduced *Oikopleura* sp. abundance was unlikely due to reduced  
665 food availability since phytoplankton within the preferred edible size spectrum, such as cyanobacteria and nanoeukaryotes,  
666 were even more abundant in the olivine treatment. Instead, we hypothesize it to be an effect of the suspended olivine  
667 particles that occurred for approximately the first 5 days of the study that were so plentiful that they turned the enclosed  
668 seawater milky and may have clogged the mucous feeding mesh of *Oikopleura* sp. (Lombard et al., 2011).

669

670 The abundance of *Penilia* sp. and *Oikopleura* sp. was lower in the olivine treatment than the other two groups throughout  
671 the experiment while the abundance of *N. scintillans* was consistently higher. The second bloom of cyanobacteria in olivine  
672 is potentially the results of decreased predators, like *Penilia* sp. and *Oikopleura* sp.. We cannot provide a particularly  
673 convincing hypothesis about what specifically drove these in these zooplankton species, although it is tempting to speculate  
674 that suspended particles present in the olivine treatment at the beginning may have played a role also for those organisms  
675 since this was the only apparent systematic difference to the control and slag treatment. The proliferation of *N. scintillans*  
676 can be problematic since heterotrophic dinoflagellate blooms can regulate phytoplankton communities, cause toxicity to  
677 aquatic fish, and create a hypoxic sub-surface zone (Baliarsingh et al., 2016; Zhang et al., 2020; Al-Azri et al., 2007),  
678 although a bloom of *N. scintillans* in southeast Australia only induced ichthyotoxicity when the cell concentration reached  
679 2,000,000 cells L<sup>-1</sup> (Hallegraeff et al., 2019). For comparison, we observed a maximum of 32 cells L<sup>-1</sup> in one microcosm  
680 replicate of the olivine treatment.

681

682 In comparison to olivine, steel slag seemed to have less potential to affect zooplankton community composition. The  
683 abundance of all groups of phytoplankton, apart from microphytoplankton after day 10, was similar in the slag treatment  
684 and the control through the experiment. This is probably because the amount of slag powder added in the treatment was  
685 much less than the olivine powder resulting in fewer physical particle perturbations to zooplankton. In addition, the

686 chemistry perturbations such as enhanced alkalinity concentration and various dissolved trace metals, especially Mn, from  
687 the slag powder did not seem to have a notable direct influence on zooplankton abundance over the three-week period.  
688 Even though we did not observe drastic changes in zooplankton abundance during the experiment, considering there was  
689 higher microphytoplankton abundance in the slag treatment after day 10, slag powder may benefit some zooplankton  
690 especially those who feed on large phytoplankton on a longer time scale.  
691

#### 692 **4.2.3. Dissolved trace metal accumulation in seawater and its environmental implications**

693 The addition of olivine and slag as OAE source minerals released trace metals into the seawater, predominantly Al, Fe, Ni,  
694 and Cu (olivine) as well as Al, Fe, and Mn (slag). The maximum measured concentrations for dissolved Al, Fe, Ni, Cu, and  
695 Mn were 1093, 253, 77, 27, and 810 nmol L<sup>-1</sup>, respectively. The threshold values for drinking water with health or aesthetic  
696 considerations by the Australian Drinking Water Guidelines for Al, Fe, Ni, Cu, and Mn are 7400, 5360, 340, 15600, and  
697 1800 nmol L<sup>-1</sup>, respectively (NRMMC, 2022). All dissolved trace metal concentrations measured herein are well below  
698 these health and aesthetic threshold values. In natural freshwater sources, the concentrations of Al, Fe, Ni, Cu and Mn are  
699 generally less than 44000, 71400, 510, 156, and 25400 nmol L<sup>-1</sup> (NRMMC, 2022). Although these natural water data were  
700 primarily derived from rivers and streams, they serve as valuable references for evaluating trace metal release in our  
701 experiment. Thus, mineral additions to the microcosms as simulated here did not increase thresholds for any of the  
702 measured trace metals beyond those that are considered safe for drinking water quality, and they were within the trace  
703 metal concentration range in natural water. However, while these guidelines on drinking water provide a good starting point  
704 on how to quantify what OAE perturbation could be considered “safe” and “unsafe” with regards to trace metals, it must  
705 be recognized that seawater is not drinking water and that critical thresholds may be different in the latter.  
706

707 The release of trace metals from OAE materials is considered to have relatively strong effects on biology, particularly in  
708 the open ocean where trace metals usually occur in lower concentrations. For example, oceanic Al, Fe, Ni, and Mn  
709 concentrations are about 2, 0.5, 8, and 0.3 nmol L<sup>-1</sup> (Bruland and Lohan, 2003; Sohrin and Bruland, 2011). Previous  
710 research on OAE-associated trace metal impacts on individual phytoplankton species grown in laboratory environments  
711 has shown that concentration thresholds beyond which trace metal induces negative effects on fitness likely differ between  
712 species (Guo et al., 2022; Hutchins et al., 2023; Xin et al., 2023). Indeed, our experiment with plankton communities  
713 provides further support that several components of the planktonic food web are affected by OAE. However, our experiment  
714 does not allow determining whether observed effects were primarily invoked by carbonate chemistry, macronutrient (P and  
715 Si), or trace metal perturbations. Thus, dedicated experiments isolating the impact of these factors on plankton will be  
716 required in the future.

#### 717 **4.2.4. Particulate trace metal accumulation in seawater and its environmental implications**

718 The Derwent Estuary (where we collected our plankton communities) was highly metal polluted due to industrial practice  
719 (Macleod and Coughanowr, 2019). Both our dissolved and particulate trace metal data indicated high background metal  
720 concentrations, especially for Fe and Zn. Furthermore, the metal:POC ratios found here are higher than reported for open  
721 ocean studies or lab cultures. For example, the Fe:POC can vary from 2-136  $\mu\text{mol mol}^{-1}$  depending on the cultured  
722 phytoplankton species and the environmental dissolved Fe concentration (Kulkarni et al., 2006; Sunda and Huntsman,  
723 1995; King et al., 2012; Boyd et al., 2015). In our results the Fe:POC values ranged from 1200 to 39 000  $\mu\text{mol mol}^{-1}$ , which

724 may be due to the particulate trace metal richness of the Derwent Estuary (control) and/or the addition of lithogenic particles  
725 (slag and olivine treatment). The presence of abiotic particulate metal sources creates challenges to quantify metal quotas  
726 and then to evaluate metal accumulation effects on biological organisms.

727

728 Our study reveals that the added minerals enriched the particulate trace metal pools to various degrees. Consistent with the  
729 dissolved trace metal data, the slag treatment was enriched with particulate Fe and Mn while the olivine treatment was  
730 enriched with particulate Fe and Ni. The enhanced particulate Ni and Mn concentrations were higher than before mineral  
731 additions and the control levels. This is in line with previous research which indicates a positive correlation between  
732 particulate and dissolved trace metal concentrations (Gaulier et al., 2019).

733

734 Based on the amounts released through OAE as simulated herein, it appears that Ni and Mn have the highest potential to  
735 cause toxicity in certain marine organisms (Jakimska et al., 2011). These trace metals have the potential to accumulate in  
736 marine organisms over time (bioaccumulation effects), and their increased concentrations in the food chain can lead to  
737 adverse effects on the health and well-being of organisms at higher trophic levels (biomagnification effects). One crucial  
738 next step will be to investigate whether the enhanced dissolved/particulate trace metal will affect higher trophic levels to  
739 estimate the environmental risks of OAE on other marine organisms.

740

## 741 **5 Conclusions**

742 Our study aimed to assess the environmental impacts of two ground OAE minerals, olivine and steel slag, on coastal  
743 plankton communities. Both minerals released alkalinity, leading to an elevation in  $\text{pH}_T$ . However, the addition of steel  
744 slag exhibited significantly higher efficiency in elevating alkalinity compared to olivine.

745

746 Approximately  $1.9 \text{ g L}^{-1}$  of olivine powder were added in the olivine treatments, leading to a  $29 \text{ } \mu\text{mol kg}^{-1}$  increase in  
747 alkalinity and increased concentrations of  $\text{Si(OH)}_4$  and trace metals (Fe and Ni). Compared to this relatively modest  
748 increase of alkalinity and associated  $\text{CO}_2$  removal potential, the impacts on the plankton community appeared to be  
749 relatively pronounced. Thus, although our experiment ran for only 3 weeks, and olivine powder may slowly release more  
750 alkalinity, the short-term response monitored here suggests that the immediate climatic benefit is relatively small compared  
751 to a relatively pronounced environmental effect.

752

753 Only  $0.038 \text{ g L}^{-1}$  of slag were added to the treatment but this led to an alkalinity enhancement of  $361 \text{ } \mu\text{mol kg}^{-1}$  and the  
754 increased concentrations of macronutrients (P and Si) and trace metals (Mn and Fe) additions as well as changes in  
755 carbonate chemistry. Although limited environmental impacts were observed from the slag treatment in our experiment,  
756 some aspects require further study. For example, the pronounced release of P could cause eutrophication and the relatively  
757 rapid increase in pH may be a detrimental aspect if organisms cannot acclimate fast enough. Furthermore, it is essential to  
758 consider that the composition of steel slag can vary depending on the source factory (Wang et al., 2011; Proctor et al.,  
759 2000), which may affect the efficiency of carbon removal and change the trace metal perturbation. Nevertheless, just based  
760 on our experiment, the comparison between the immediate climatic benefit and environmental effect appears to be more  
761 favourable for slag than olivine.

762

763 Based on our findings, it can be concluded that steel slag powder exhibited fewer environmental impacts on plankton  
764 communities compared to olivine powder relative to its capacity for alkalinity enhancement. The results highlight the  
765 importance of carefully assessing the environmental consequences of using specific OAE minerals, particularly when  
766 considering their potential effects on plankton communities.

767

768 **Data availability.** Data are available in the Institute for Marine and Antarctic Studies (IMAS) data catalogue, University  
769 of Tasmania (UTAS) (<https://doi.org/10.25959/X6FH-9K15>, Guo, J., & Bach, L. (2023).).

770

771 **Author contributions.** LTB, RFS, KMS and JAG designed the experiments and JAG carried them out. LTB, RFS and  
772 KMS supervised the study. ATT analysed the dissolved/particulate trace metal samples. JAG conducted statistical analyses.  
773 JAG prepared the manuscript with contributions from all authors.

774

775 **Competing interests.** The contact author has declared that none of the authors has any competing interests.

776

777 **Disclaimer.** Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims in published  
778 maps and institutional affiliations.

779

780 **Acknowledgements.** We would like to thank Steve Van Orsouw from Moyne Shire Council, Victoria, Australia for  
781 providing olivine rocks. We also thank Bradley Mansell who provided the Basic Oxygen Slag from Liberty Primary Steel  
782 Whyalla Steelworks in Whyalla, South Australia, Australia. We thank Sandrin Feig and Thomas Rodemann for their support  
783 on scanning electron microscopy and particulate organic matter. We appreciate the assistance of Pam Quayle and Axel  
784 Durand (IMAS) in the lab, particularly with particulate metal digestions.

785

786 **Financial support.** This research has been supported by the Australian Research Council through a Future Fellowship  
787 project (FT200100846 to LTB), the Carbon to Sea Initiative (LTB), and by the Australian Antarctic Program Partnership  
788 (ASCI000002 to RFS, KMS and JAG). Access to SF-ICP-MS instrumentation was facilitated through ARC LIEF funding  
789 (LE0989539) awarded to ATT. JAG thanks the Australian Research Training Program (RTP) for her scholarship.

## 790 **References**

- 791 Ackerman, L., Jelínek, E., Medaris, G., Ježek, J., Siebel, W., and Strnad, L.: Geochemistry of Fe-rich peridotites and  
792 associated pyroxenites from Horní Bory, Bohemian Massif: Insights into subduction-related melt–rock reactions,  
793 *Chem. Geol.*, 259, 152-167, <https://doi.org/10.1016/j.chemgeo.2008.10.042>, 2009.
- 794 Al-Azri, A., Al-Hashmi, K., Goes, J., Gomes, H., Rushdi, A. I., Al-Habsi, H., Al-Khusaibi, S., Al-Kindi, R., and Al-Azri, N.:  
795 Seasonality of the bloom-forming heterotrophic dinoflagellate *Noctiluca scintillans* in the Gulf of Oman in relation  
796 to environmental conditions, *Int. J. Oceans Oceanogr.*, 2, 51-60, 2007.
- 797 Armstrong, F. A.: Why did nature choose manganese to make oxygen?, *Philos Trans R Soc Lond B Biol Sci*, 363, 1263-1270,  
798 <https://doi.org/10.1098/rstb.2007.2223>, 2008.
- 799 Bach, L. T., Gill, S. J., Rickaby, R. E. M., Gore, S., and Renforth, P.: CO2 removal with enhanced weathering and ocean  
800 alkalinity enhancement: potential risks and co-benefits for marine pelagic ecosystems, *Front Clim*, 1, 1-21,  
801 <http://doi.org/10.3389/fclim.2019.00007>, 2019.
- 802 Balaguer, J., Koch, F., Hassler, C. et al.: Iron and manganese co-limit the growth of two phytoplankton groups dominant at  
803 two locations of the Drake Passage. *Commun Biol* 5, 207, <https://doi.org/10.1038/s42003-022-03148-8>, 2022.



804 Baliarsingh, S. K., Lotliker, A. A., Trainer, V. L., Wells, M. L., Parida, C., Sahu, B. K., Srichandan, S., Sahoo, S., Sahu, K. C., and  
805 Kumar, T. S.: Environmental dynamics of red *Noctiluca scintillans* bloom in tropical coastal waters, *Mar. Pollut. Bull.*,  
806 111, 277-286, <https://doi.org/10.1016/j.marpolbul.2016.06.103>, 2016.

807 Basu, S. and Mackey, K. R. M.: Phytoplankton as key mediators of the biological carbon pump: their responses to a  
808 changing climate, *Sustainability*, 10, 869, <https://doi.org/10.3390/su10030869>, 2018.

809 Behrenfeld, M. J., Worthington, K., Sherrell, R. M., Chavez, F. P., Strutton, P., McPhaden, M., and Shea, D. M.: Controls on  
810 tropical Pacific Ocean productivity revealed through nutrient stress diagnostics, *Nature*, 442, 1025-1028,  
811 <https://doi.org/10.1038/nature05083>, 2006.

812 Bowie, A. R., Townsend, A. T., Lannuzel, D., Remenyi, T. A., and van der Merwe, P.: Modern sampling and analytical  
813 methods for the determination of trace elements in marine particulate material using magnetic sector inductively  
814 coupled plasma-mass spectrometry, *Anal Chim Acta*, 676, 15-27, <https://doi.org/10.1016/j.aca.2010.07.037>, 2010.

815 Boyd, P. W., Strzepek, R. F., Ellwood, M. J., Hutchins, D. A., Nodder, S. D., Twining, B. S., and Wilhelm, S. W.: Why are biotic  
816 iron pools uniform across high- and low-iron pelagic ecosystems?, *Global Biogeochem. Cycles*, 29, 1028-1043,  
817 <http://doi.org/10.1002/2014gb005014>, 2015.

818 Boyd, P. W., Jickells, T., Law, C., Blain, S., Boyle, E., Buesseler, K., Coale, K., Cullen, J., De Baar, H. J., and Follows, M.:  
819 Mesoscale iron enrichment experiments 1993-2005: synthesis and future directions, *Science*, 315, 612-617,  
820 <http://doi.org/10.1126/science.1131669>, 2007.

821 Bruland, K. W. and Lohan, M. C.: 6.02 Controls of Trace Metals in Seawater, in: *Treatise on Geochemistry*, edited by:  
822 Elderfield, H., Holland, H. D., and Turekian, K. K., Elsevier Pergamon, 23-47, [http://doi.org/10.1016/b0-08-043751-](http://doi.org/10.1016/b0-08-043751-6/06105-3)  
823 6/06105-3, 2003.

824 Burt, D. J., Fröb, F., and Ilyina, T.: The sensitivity of the marine carbonate system to regional ocean alkalinity enhancement,  
825 *Front Clim*, 3, <http://doi.org/10.3389/fclim.2021.624075>, 2021.

826 Caserini, S., Storni, N., and Grosso, M.: The availability of limestone and other raw materials for ocean alkalinity  
827 enhancement, *Global Biogeochem. Cycles*, 36, <http://doi.org/10.1029/2021gb007246>, 2022.

828 De Baar, H. J., Boyd, P. W., Coale, K. H., Landry, M. R., Tsuda, A., Assmy, P., Bakker, D. C., Bozec, Y., Barber, R. T., and Brzezinski,  
829 M. A.: Synthesis of iron fertilization experiments: from the iron age in the age of enlightenment, *J Geophys Res*  
830 *Oceans*, 110, <https://doi.org/10.1029/2004JC002601>, 2005.

831 Dickson, A. G., Sabine, C. L., and Christian, J. R.: *Guide to best practices for ocean CO<sub>2</sub> measurements*, North Pacific Marine  
832 Science Organization, Canada2007.

833 Egge, J. and Jacobsen, A.: Influence of silicate on particulate carbon production in phytoplankton, *Mar. Ecol. Prog. Ser.*,  
834 147, 219-230, <http://doi.org/10.3354/meps147219>, 1997.

835 Evans, C., O'Reilly, J. E., and Thomas, J.: *A handbook for the measurement of chlorophyll and primary production*, 1987.

836 Falkowski, P. G.: The role of phytoplankton photosynthesis in global biogeochemical cycles, *Photosynthesis Research*, 39,  
837 235-258, <https://doi.org/10.1007/BF00014586>, 1994.

838 Feng, E. Y., Koeve, W., Keller, D. P., and Oschlies, A.: Model-based assessment of the CO<sub>2</sub> sequestration potential of coastal  
839 ocean alkalization, *Earth's Future*, 5, 1252-1266, <http://doi.org/10.1002/2017ef000659>, 2017.

840 Ferderer, A., Chase, Z., Kennedy, F., Schulz, K. G., and Bach, L. T.: Assessing the influence of ocean alkalinity enhancement  
841 on a coastal phytoplankton community, *Biogeosciences*, 19, 5375-5399, <http://doi.org/10.5194/bg-19-5375-2022>,  
842 2022.

843 Ferderer, A., Schulz, K. G., Riebesell, U., Baker, K. G., Chase, Z., and Bach, L. T.: Investigating the effect of silicate and calcium  
844 based ocean alkalinity enhancement on diatom silicification, *Biogeosciences Discuss.* [preprint],  
845 <https://doi.org/10.5194/bg-2023-144>, in review, 2023.

846 Package 'seacarb'-Seawater Carbonate Chemistry: <https://cran.r-project.org/web/packages/seacarb/index.html>, last  
847 access: 2023/6/1.

848 Gaulier, C., Zhou, C., Guo, W., Bratkic, A., Superville, P. J., Billon, G., Baeyens, W., and Gao, Y.: Trace metal speciation in  
849 North Sea coastal waters, *Sci. Total Environ.*, 692, 701-712, <http://doi.org/10.1016/j.scitotenv.2019.07.314>, 2019.

850 Guo, J., Bao, Y., and Wang, M.: Steel slag in China: Treatment, recycling, and management, *Waste Management*, 78, 318-  
851 330, <https://doi.org/10.1016/j.wasman.2018.04.045>, 2018.

852 Guo, J. A., Strzepek, R., Willis, A., Ferderer, A., and Bach, L. T.: Investigating the effect of nickel concentration on  
853 phytoplankton growth to assess potential side-effects of ocean alkalinity enhancement, *Biogeosciences*, 19, 3683-  
854 3697, <https://doi.org/10.5194/bg-19-3683-2022>, 2022.

855 Hallegraeff, G. M., Albinsson, M. E., Dowdney, J., Holmes, A. K., Mansour, M. P., and Seger, A.: Prey preference,  
856 environmental tolerances and ichthyotoxicity by the red-tide dinoflagellate *Noctiluca scintillans* cultured from  
857 Tasmanian waters, *J. Plankton Res.*, 41, 407-418, <https://doi.org/10.1093/plankt/fbz037>, 2019.

858 Hansen, H. P. and Koroleff, F.: Determination of nutrients, in: *Methods of seawater analysis*, edited by: Grasshoff, K.,  
859 Kremling, K., and Ehrhardt, M., 159-228, <https://doi.org/10.1002/9783527613984.ch10>, 1999.

860 Hartmann, J., West, A. J., Renforth, P., Köhler, P., De La Rocha, C. L., Wolf-Gladrow, D. A., Dürr, H. H., and Scheffran, J.:

861 Enhanced chemical weathering as a geoengineering strategy to reduce atmospheric carbon dioxide, supply  
862 nutrients, and mitigate ocean acidification, *Rev. Geophys.*, 51, 113-149, <http://doi.org/10.1002/rog.20004>, 2013.

863 Humphreys, M. P., Lewis, E. R., Sharp, J. D., and Pierrot, D.: PyCO2SYS v1. 8: marine carbonate system calculations in  
864 Python, *Geosci Model Dev*, 15, 15-43, <https://doi.org/10.5194/gmd-15-15-2022>, 2022.

865 Hutchins, D. A., Fu, F.-X., Yang, S.-C., John, S. G., Romaniello, S. J., Andrews, M. G., and Walworth, N. G.: Responses of  
866 globally important phytoplankton species to olivine dissolution products and implications for carbon dioxide  
867 removal via ocean alkalinity enhancement, *Biogeosciences*, 20, 4669–4682, [https://doi.org/10.5194/bg-20-4669-](https://doi.org/10.5194/bg-20-4669-2023)  
868 2023, 2023.

869 Ilyina, T., Wolf-Gladrow, D., Munhoven, G., and Heinze, C.: Assessing the potential of calcium-based artificial ocean  
870 alkalization to mitigate rising atmospheric CO<sub>2</sub> and ocean acidification, *Geophys. Res. Lett.*, 40, 5909-5914,  
871 <https://doi.org/10.1002/2013GL057981>, 2013.

872 Jakimska, A., Konieczka, P., Skóra, K., and Namieśnik, J.: Bioaccumulation of metals in tissues of marine animals, Part II:  
873 metal concentrations in animal tissues, *Pol J Environ Stud*, 20, 2011.

874 Keller, D. P., Feng, E. Y., and Oschlies, A.: Potential climate engineering effectiveness and side effects during a high carbon  
875 dioxide-emission scenario, *Nat. Commun.*, 5, 3304, <https://doi.org/10.1038/ncomms4304>, 2014.

876 King, A. L., Sañudo-Wilhelmy, S. A., Boyd, P. W., Twining, B. S., Wilhelm, S. W., Breene, C., Ellwood, M. J., and Hutchins, D.  
877 A.: A comparison of biogenic iron quotas during a diatom spring bloom using multiple approaches, *Biogeosciences*,  
878 9, 667-687, <http://doi.org/10.5194/bg-9-667-2012>, 2012.

879 Kohler, P., Hartmann, J., and Wolf-Gladrow, D. A.: Geoengineering potential of artificially enhanced silicate weathering of  
880 olivine, *Proc. Natl. Acad. Sci. USA*, 107, 20228-20233, <https://doi.org/10.1073/pnas.1000545107>, 2010.

881 Kourounis, S., Tsvilis, S., Tsakiridis, P. E., Papadimitriou, G. D., and Tsi bouki, Z.: Properties and hydration of blended  
882 cements with steelmaking slag, *Cem. Concr. Res.*, 37, 815-822, <https://doi.org/10.1016/j.cemconres.2007.03.008>,  
883 2007.

884 Kulkarni, P. P., She, Y. M., Smith, S. D., Roberts, E. A., and Sarkar, B.: Proteomics of metal transport and metal-associated  
885 diseases, *Chemistry*, 12, 2410-2422, <http://doi.org/10.1002/chem.200500664>, 2006.

886 Lenton, A., Matear, R. J., Keller, D. P., Scott, V., and Vaughan, N. E.: Assessing carbon dioxide removal through global and  
887 regional ocean alkalization under high and low emission pathways, *Earth. Syst. Dyn.*, 9, 339-357,  
888 <https://doi.org/10.5194/esd-9-339-2018>, 2018.

889 Lombard, F., Selander, E., and Kjørboe, T.: Active prey rejection in the filter-feeding appendicularian *Oikopleura dioica*,  
890 *Limnol. Oceanogr.*, 56, 1504-1512, <http://doi.org/10.4319/lo.2011.56.4.1504>, 2011.

891 Lueker, T. J., Dickson, A. G., and Keeling, C. D.: Ocean pCO<sub>2</sub> calculated from dissolved inorganic carbon, alkalinity, and  
892 equations for K<sub>1</sub> and K<sub>2</sub>: validation based on laboratory measurements of CO<sub>2</sub> in gas and seawater at equilibrium,  
893 *Mar. Chem.*, 70, 105-119, [https://doi.org/10.1016/S0304-4203\(00\)00022-0](https://doi.org/10.1016/S0304-4203(00)00022-0), 2000.

894 Macleod, C. and Coughanowr, C.: Heavy metal pollution in the Derwent estuary: History, science and management, *Reg.*  
895 *Stud. Mar. Sci.*, 32, <http://doi.org/10.1016/j.rsma.2019.100866>, 2019.

896 Moore, C. M., Mills, M. M., Arrigo, K. R., Berman-Frank, I., Bopp, L., Boyd, P. W., Galbraith, E. D., Geider, R. J., Guieu, C.,  
897 Jaccard, S. L., Jickells, T. D., La Roche, J., Lenton, T. M., Mahowald, N. M., Marañón, E., Marinov, I., Moore, J. K.,  
898 Nakatsuka, T., Oschlies, A., Saito, M. A., Thingstad, T. F., Tsuda, A., and Ulloa, O.: Processes and patterns of oceanic  
899 nutrient limitation, *Nat. Geosci.*, 6, 701-710, <http://doi.org/10.1038/ngeo1765>, 2013.

900 Nelson, D. M., Smith Jr, W. O., Muench, R. D., Gordon, L. I., Sullivan, C. W., and Husby, D. M.: Particulate matter and nutrient  
901 distributions in the ice-edge zone of the Weddell Sea: relationship to hydrography during late summer, *Deep. Sea.*  
902 *Res. A*, 36, 191-209, [https://doi.org/10.1016/0198-0149\(89\)90133-7](https://doi.org/10.1016/0198-0149(89)90133-7), 1989.

903 NRMCC, N. a.: The Australian Drinking Water Guidelines (2011) - Version 3.8 Updated 2022, 2022.

904 Pausch, F., Bischof, K., Trimborn, S.: Iron and manganese co-limit growth of the Southern Ocean diatom *Chaetoceros debilis*.  
905 *PLOS ONE* 14, e0221959. <https://doi.org/10.1371/journal.pone.0221959>, 2019.

906 Paquay, F. S. and Zeebe, R. E.: Assessing possible consequences of ocean liming on ocean pH, atmospheric CO<sub>2</sub>  
907 concentration and associated costs, *Int. J. Greenh. Gas Control.*, 17, 183-188,  
908 <https://doi.org/10.1016/j.ijggc.2013.05.005>, 2013.

909 Platt, T., Gallegos, C. L., and Harrison, W. G.: Photoinhibition of photosynthesis in natural assemblages of marine  
910 phytoplankton, *J. Mar. Res.*, 38, 687-701, 1980.

911 Proctor, D. M., Fehling, K. A., Shay, E. C., Wittenborn, J. L., Green, J. J., Avent, C., Bigham, R. D., Connolly, M., Lee, B.,  
912 Shepker, T. O., and Zak, M. A.: Physical and chemical characteristics of blast furnace, basic oxygen furnace, and  
913 electric arc furnace steel industry slags, *Environ. Sci. Technol.*, 34, 1576-1582, <http://doi.org/10.1021/es9906002>,  
914 2000.

915 Reichl, C., Schatz, M., and Zsak, G.: World mining data, 1-261, 2018.

916 Renforth, P.: The negative emission potential of alkaline materials, *Nat. Commun.*, 10, [http://doi.org/10.1038/s41467-](http://doi.org/10.1038/s41467-019-09475-5)  
917 019-09475-5, 2019.

918 Renforth, P. and Henderson, G.: Assessing ocean alkalinity for carbon sequestration, *Rev. Geophys.*, 55, 636-674,  
919 <http://doi.org/10.1002/2016rg000533>, 2017.

920 Schallenberg, C., Strzepek, R. F., Schuback, N., Clementson, L. A., Boyd, P. W., and Trull, T. W.: Diel quenching of Southern  
921 Ocean phytoplankton fluorescence is related to iron limitation, *Biogeosciences*, 17, 793-812,  
922 <https://doi.org/10.5194/bg-17-793-2020>, 2020.

923 Schuiling, R. D. and Krijgsman, P.: Enhanced weathering: an effective and cheap tool to sequester CO<sub>2</sub>, *Clim. Change*, 74,  
924 349-354, <https://doi.org/10.1007/s10584-005-3485-y>, 2006.

925 Selfe, C.: Developing Transfer Function to Measuring Phytoplankton Cellular Properties with Flow Cytometry, Master's  
926 thesis, Institute of Marine and Antarctic Studies, University of Tasmania, Australia, 2022.

927 Comparing smooths in factor-smooth interactions II ordered factors:  
928 <https://fromthebottomoftheheap.net/2017/12/14/difference-splines-ii/>, last access: March 2023.

929 Smith, S. M., Geden, O., Nemet, G. F., Gidden, M. J., Lamb, W. F., Powis, C., Bellamy, R., Callaghan, M. W., Cowie, A., Cox,  
930 E., Fuss, S., Gasser, T., Grassi, G., Greene, J., Lück, S., Mohan, A., Müller-Hansen, F., Peters, G. P., Pratama, Y., Repke,  
931 T., Riahi, K., Schenuit, F., Steinhauser, J., Strefler, J., Valenzuela, J. M., and Minx, J. C.: The State of Carbon Dioxide  
932 Removal - 1st Edition, <http://doi.org/10.17605/OSF.IO/W3B4Z>, 2023.

933 Sohrin, Y. and Bruland, K. W.: Global status of trace elements in the ocean, *TrAC, Trends Anal. Chem.*, 30, 1291-1307,  
934 <https://doi.org/10.1016/j.trac.2011.03.006>, 2011.

935 Strzepek, R. F. and Harrison, P. J.: Photosynthetic architecture differs in coastal and oceanic diatoms, *Nature*, 431, 689-  
936 692, <http://doi.org/10.1038/nature02954>, 2004.

937 Su, B., Chen, Y., Guo, S., and Liu, J.: Origins of orogenic dunites: petrology, geochemistry, and implications, *Gondwana Res.*,  
938 29, 41-59, <https://doi.org/10.1016/j.gr.2015.08.001>, 2016.

939 Subhas, A. V., Marx, L., Reynolds, S., Flohr, A., Mawji, E. W., Brown, P. J., and Cael, B.: Microbial ecosystem responses to  
940 alkalinity enhancement in the North Atlantic Subtropical Gyre, *Front Clim*, 4,  
941 <https://doi.org/10.3389/fclim.2022.784997>, 2022.

942 Sunda, W. G.: Trace metal-phytoplankton interactions in aquatic systems, in: *Environmental Microbe-Metal Interactions*,  
943 edited by: Lovley, D. R., 79-107, <https://doi.org/10.1128/9781555818098.ch4>, 2000.

944 Sunda, W. G.: Feedback interactions between trace metal nutrients and phytoplankton in the ocean, *Frontiers in*  
945 *Microbiology*, 3, 1-22, <http://doi.org/10.3389/fmicb.2012.00204>, 2012.

946 Sunda, W. G. and Huntsman, S. A.: Iron Uptake and Growth Limitation in Oceanic and Coastal Phytoplankton, *Marine*  
947 *Chemistry*, 50, 189-206, [Doi 10.1016/0304-4203\(95\)00035-P](https://doi.org/10.1016/0304-4203(95)00035-P), 1995.

948 Sunda, W. G. and Huntsman, S. A.: Interrelated influence of iron, light and cell size on marine phytoplankton growth,  
949 *Nature*, 390, 389-392, <http://doi.org/10.1038/37093>, 1997.

950 Tang, D. G. and Morel, F. M. M.: Distinguishing between cellular and Fe-oxide-associated trace elements in phytoplankton,  
951 *Mar. Chem.*, 98, 18-30, <http://doi.org/10.1016/j.marchem.2005.06.003>, 2006.

952 Tovar-Sanchez, A., Sanudo-Wilhelmy, S. A., Garcia-Vargas M., Weaver R. S., Popels L., C., and Hutchins D. A.: A trace metal  
953 clean reagent to remove surface-bound iron from marine phytoplankton. *Mar. Chem.*, 82, 1-2, 91-99,  
954 [https://doi.org/10.1016/S0304-4203\(03\)00054-9](https://doi.org/10.1016/S0304-4203(03)00054-9), 2003.

955 Twining, B. S. and Baines, S. B.: The trace metal composition of marine phytoplankton, *Ann. Rev. Mar. Sci.*, 5, 191-215,  
956 <http://doi.org/10.1146/annurev-marine-121211-172322>, 2013.

957 Wafar, M., Le Corre, P., and L'Helguen, S.: f-Ratios calculated with and without urea uptake in nitrogen uptake by  
958 phytoplankton, *Deep Sea Res. I Oceanogr. Res. Pap.*, 42, 1669-1674, [https://doi.org/10.1016/0967-0637\(95\)00066-F](https://doi.org/10.1016/0967-0637(95)00066-F), 1995.

960 Wang, Q., Yan, P., and Feng, J.: A discussion on improving hydration activity of steel slag by altering its mineral  
961 compositions, *J. Hazard. Mater.*, 186, 1070-1075, <https://doi.org/10.1016/j.jhazmat.2010.11.109>, 2011.

962 Wang, W.: Interactions of trace metals and different marine food chains, *Mar. Ecol. Prog. Ser.*, 243, 295-309,  
963 <http://doi.org/10.3354/meps243295>, 2002.

964 Wolfe-Simon, F., Grzebyk, D., Schofield, O., and Falkowski, P. G.: The role and evolution of superoxide dismutases in algae,  
965 *J. Phycol.*, 41, 453-465, <https://doi.org/10.1111/j.1529-8817.2005.00086.x>, 2005.

966 Xin, X., Faucher, G., and Riebesell, U.: Phytoplankton response to Increased nickel in the context of ocean alkalinity  
967 enhancement, *Biogeosciences* [preprint], <https://doi.org/10.5194/bg-2023-130>, 2023.

968 Zhang, W., Dong, Z., Zhang, C., Sun, X., Hou, C., Liu, Y., Wang, L., Ma, Y., and Zhao, J.: Effects of physical-biochemical  
969 coupling processes on the *Noctiluca scintillans* and *Mesodinium* red tides in October 2019 in the Yantai nearshore,  
970 China, *Mar. Pollut. Bull.*, 160, 111609, <https://doi.org/10.1016/j.marpolbul.2020.111609>, 2020.

971