

Response to Reviewer #2 (EGUSPHERE-2023-2122)

First of all, we would like to thank the editor and reviewers for their valuable comments. We have taken all the suggested changes into consideration and revised the manuscript accordingly. The reviewers' comments are copied here as texts in BLACK, our responses are followed in BLUE, and the major corrections are marked in RED in the manuscript.

The manuscript employs a hybrid method to estimate brown carbon (BrC) radiative effects using a combination of aethalometer measurements of aerosol absorption, an optical separation method, simulated BrC optical properties, and a radiative transfer model. As BrC is still poorly characterized in the field and largely ignored by many chemical transport models and climate models, the presented results contribute to our understanding of its radiation and climate significance. The manuscript fits the scope of ACP very well. I have some comments below for the authors to address.

Response: Thanks so much for your constructive comments. We have implemented all suggestions for improvement in the revised manuscript. Please find our point-by-point responses listed below.

Line 143-145: the authors set the $AAE_{BC}=1$ to calculate the absorption contributions of BrC at different wavelengths. However, existing research suggests that the AAE of BC in the atmosphere varies within a certain range due to factors such as mixing state and morphology (Lack and Cappa, 2010; Zhang et al., 2020b). To comprehensively assess BrC's absorption contributions, different AAE_{BC} values could be set, and some sensitivity analyses are needed to evaluate the impact of this parameter on the study results.

Lack, D. A. and Cappa, C. D.: Impact of brown and clear carbon on light absorption enhancement, single scatter albedo and absorption wavelength dependence of black carbon, *Atmos. Chem. Phys.*, 10, 4207–4220, <https://doi.org/10.5194/acp-10-4207-2010>, 2010.

Zhang, X., Mao, M., Yin, Y., and Tang, S.: The absorption Ångstrom exponent of black carbon with brown coatings: effects of aerosol microphysics and parameterization, *Atmos. Chem. Phys.*, 20, 9701–9711, <https://doi.org/10.5194/acp-20-9701-2020>, 2020.

Response: Per your suggestion, we added a new section (Section 5) to discuss the influences of AAE_{BC} uncertainties on our radiative effect estimation. We followed the AAE_{BC} values suggested by Liu et al. (2018), and found that a change of BC AAE from 1.0 to 0.8 may introduce BC radiative effects of approximately 10% and BrC effects of over 40% (due to an obvious increase in BrC amount estimation).

Line 355 – Line 359 in the main text,

“The BC AAE assumption showed influences on BC radiative effects with relative differences of approximately 10%, but much stronger influences on BrC radiative effects, i.e., ~40%. This result was ascribed to smaller BC AAE causing weaker BC absorption in shorter wavelengths, leading to smaller radiative effects. However, larger BrC absorption coefficients and amounts during the segregation could cause larger BrC radiative effects.”

Line 178-179: In the optical closure calculation, a three-component aerosol model (BrC, BC, and pure-scattering components) was chosen with the assumption of an external mixing state. However, internal mixing between chemical components has been confirmed as a crucial factor influencing aerosol optical properties. It is recommended that the authors consider the internal mixing state to comprehensively evaluate the aerosol optical properties. Alternatively, the authors could explicitly state the reasons for choosing the external mixing model. This discussion will enhance the reliability of its results.

Response: This is an excellent question and suggestion. The mixing structure plays an important role in aerosol optical and radiative property estimation, which has been extensively studied as well. Considering the high complexity of aerosol mixing states, the general observations considered in this study can hardly provide any meaningful information on our model, so we considered this external mixing assumption for the sake of developing a more practical and general method with existing observations. However, we highly agree with the suggestion, and will further investigate the influences of internal mixing on this work in future work. The discussions above have also been added in the revision.

Line 209-210: In this study, it is crucial to address whether the Mie numerical simulation is based on the spherical assumption, as this may introduce biases in optical calculations for non-spherical particles. Particularly, a spherical Mie model tends to significantly overestimate the light absorption of fractal BC particles. The numerical simulation of aerosol optical properties is intricately linked to the accuracy of mass distribution among different components. It is recommended that the potential biases introduced by the spherical assumption should be discussed.

Response: As mentioned above, we added a systematic sensitive study on the influences of our assumptions on BC/BrC radiative effect estimations, which includes the influences due to BC geometries. To achieve this, we used the BC optical property database with fractal aggregate structures we developed (Liu et al., 2019), and discussed the differences in the radiative effects between spherical and fractal-aggregate-based

BC particles. The corresponding results and discussions are presented in Section 5 of the revision.

Line 359 – Line 364 in the main text,

“The assumption regarding BC spherical particles showed minor influences on BrC radiative effects, with the relative differences being less than 5%, but much stronger influences on BC radiative effects, particularly on the RF(TOA) with an average deviation of almost 50%, caused mainly by weaker scattering owing to BC non-sphericity (Li et al., 2016). In other words, the assumption regarding spherical BC led to overestimation of BC scattering, increasing the upward radiation reaching the TOA. However, the influence of BC non-sphericity on BrC/BC segregation was lower and the BrC radiative effects change was less than 5%.”

As shown in Figure 4, the uncertainty in the Refractive Index of BrC is notable. The authors should provide additional details explaining the reason for choosing the RI reported by Shamjad et al. (2016). It would be beneficial to illustrate the factors such as the similarity in organic aerosol composition and sources between the two cities. This additional information will enhance the understanding of the selection criteria for the RI values.

Response: Yes, there are significant uncertainties on the refractive indices of BrC. As suggested by both reviewers, we added a sensitive study on the influences of BrC refractive indices (imaginary part) on our BC and BrC radiative effect estimation. We found that BrC refractive index variations may introduce uncertainties up to over 50% on the BrC TOA RF, and more details were added in Section 5.

Line 364 – Line 367 in the main text,

“The right panel shows the influences deriving from uncertainties of the absorptivity of BrC. Except for the RF (TOA) with a relatively larger difference of 62%, the rest of the BrC radiative effects (average absolute values) were all below 30%, and, as expected, all the differences were close to zero.”

Line 205-207: The data utilized in the study are relatively dated, and it is recommended to consider relevant data from recent studies. The inclusion of more recent data would enhance the timeliness and relevance of the findings.

Response: Thanks for the suggestion. As you may notice, even this “general” model requires a large number of observations for aerosol constraints. Currently, the relatively dated results are used due to their well-maintained quality and longer observational period. Meanwhile, this study mainly focuses on introducing the method for BrC

radiative effects, and the date of the observations will not influence the application of the methods. Of course, we will further improve and validate our method for more recent data in future studies.

Line 207: The density of BC is commonly reported as $1.8 \text{ g}\cdot\text{cm}^{-3}$ in the literature. It is advised to verify the accuracy of the statement indicating a density of $1.0 \text{ g}\cdot\text{cm}^{-3}$ for BC in this context.

Response: Thanks for suggestion. After carefully review, we agree that $1.8 \text{ g}/\text{cm}^3$ may be a more appropriate value, and have updated our results by assuming a BC density of $1.8 \text{ g}/\text{cm}^3$. The BC density mostly influences the estimated BC mass concentration/amount, while having less effect on our optical closure study.

Line 216 in the main text,

“The densities of BC and scattering aerosol were assumed at 1.8 and $1.7 \text{ g}\cdot\text{cm}^{-3}$, respectively.”

The calculation of radiative forcing (RF) in this study is unclear regarding whether it considers only direct radiative forcing or also includes indirect radiative forcing. Some clarification is needed.

Response: Thanks for the suggestion. We have clarified the RF considers only direct radiative forcing in this study.

Line 78 in the main text,

“Radiative forcing (RF, only direct radiative forcing) and influences on the photosynthetically active radiation (PAR) and actinic flux (AF) were considered and discussed.”

The English language needs to be further polished. Some necessary edits in the abstract:

- L15, convenience -> efficient, concise -> available
- L23, BrC induces a warming effect with an average instantaneous radiative forcing (RF) of $6.4 \pm 3.4 \text{ W m}^{-2}$, corresponding to 29.2% of the BC RF.
that of black carbon (BC).
- L26, you may want to say “PAR attenuation”.

Response: Sorry for the language problems, we have corrected the mistakes, and all authors carefully proofread the revision. Meanwhile, the revision was further edited for proper English language, grammar, punctuation, spelling, and overall style by a qualified native English-speaking editor.

Line 15 in the main text,

“This study proposes an efficient method for estimating BrC radiative effects based on the available observational data.”

Line 23 in the main text,

“In the atmosphere, BrC plays a warming role, with its average instantaneous radiative forcing (RF) and standard deviation of $4.0 \pm 2.3 \text{ W m}^{-2}$ corresponding to $15 \pm 4.2\%$ of the black carbon (BC) RF.”

Line 26 in the main text,

“The photosynthetically active radiation (PAR) attenuated by BrC was approximately $33.5 \pm 9.4\%$ of that attenuated by BC.”