

1 **Running title:** Global biocrusts distribution

2 **Advancing studies on global biocrusts distribution**

3 Siqing Wang^{1,2}, Li Ma^{1,2}, Liping Yang^{1,2}, Yali Ma^{1,2}, Yafeng Zhang³, Changming Zhao^{1,2}, Ning
4 Chen^{1,2,4*}

5 ¹ State Key Laboratory of Herbage Improvement and Grassland Agro-ecosystems, College of
6 Ecology, Lanzhou University, Lanzhou, Gansu 730000, China

7 ² Yuzhong Mountain Ecosystems Observation and Research Station, Lanzhou University,
8 Lanzhou, Gansu 730000, China

9 ³ Shapotou Desert Research and Environment Station, Northwest Institute of Eco-Environment
10 and Resources, Chinese Academy of Sciences, No.320, Donggang West Road,
11 Lanzhou, Gansu 730000, China

12 ⁴ Instituto Multidisciplinar para el Estudio del Medio “Ramon Margalef”, Universidad de
13 Alicante, Carretera de San Vicente del Raspeig s/n, San Vicente del Raspeig Alicante 03690,
14 Spain

15 ***Correspondence to Ning Chen, Email: cn@lzu.edu.cn. Tel.: +86-931-8912551. ORCID: 0000-**
16 **0002-1779-915X.**

17

18 **Abstract:** Biological soil crusts (biocrusts hereafter) cover a substantial proportion of dryland
19 ecosystem and play crucial roles in ecological processes such as biogeochemical cycles, water
20 distribution and soil erosion. Consequently, studying spatial distribution of biocrusts holds great
21 significance for drylands, which is still lacking, especially in a global scale. This study aimed
22 to stimulate global-scale investigations of biocrusts distribution by introducing three major
23 approaches: spectral characterization indices, dynamic vegetation models, and geospatial
24 models, while discussing their applicability. Then, we summarized present understandings of
25 biocrusts distribution. Finally, to further advance this field, we proposed several potential
26 research topics and aspects, including building standardized database of biocrusts, enhancing
27 non-vascular vegetation dynamic models, integrating multi-sensor monitoring, making full use
28 of machine learning, and focusing on regional research co-development. This work is supposed
29 to significantly contribute to mapping biocrusts distribution, and thereby to advance our
30 understandings of dryland ecosystem management and restoration.

31 **Key words:** biological soil crusts; distribution; drylands; global scales; regional scales

32

33 **1. Introduction**

34 Biological soil crusts (biocrusts hereafter) are continuous [biotic](#) complexes that live in the

35 topsoil, which are formed by different proportions of photosynthetic autotrophic (e.g.
36 cyanobacteria, algae, lichens, mosses) and heterotrophic (e.g. bacteria, fungi, archaea)
37 organisms colloidal with soil particles, usually with a thickness of a few millimeters to a few
38 centimeters (Weber et al., 2022). They [are able to occupy a wide ecological niche in mid](#)
39 [latitudes, polar and alpine regions, covering approximately 11% of the global land area](#) (Porada
40 et al., 2019). [In particular, biocrusts can be adapted to](#) water-limited, nutrient-poor and hostile
41 environments, [such as](#) arid and semi-arid areas characterized by low ratios of precipitation to
42 potential evaporation (0.05-0.5 mm_{mm⁻¹}) (Pravalie, 2016; Read et al., 2014; Weber et al., 2016).

43 As vital components of dryland ecosystems, biocrusts fulfill [many](#) essential ecological
44 functions. They contribute to stabilizing the soil surface and improving soil permeability and
45 water-holding capacity within the upper [few](#) centimeters (Sun et al., 2023; Shi et al., 2023; Gao
46 et al., 2017). By participating in a suite of biogeochemical cycles, biocrusts [were estimated to](#)
47 contribute [to](#) 15% of terrestrial net primary productivity and 40-85% of biological nitrogen
48 fixation (Elbert et al., 2012; Rodriguez-Caballero et al., 2018). They also impact
49 ecohydrological processes by altering soil microclimate and redistributing soil water (Kidron
50 et al., 2022; Tucker et al., 2017). Moreover, biocrusts influence seed capture and soil seed banks
51 (Kropfl et al., 2022), thereby mediating plant growth and community assembly (Havrilla and
52 Barger, 2018; Song et al., 2022). The extent and magnitude of these ecological functions and
53 services depend on the spatial distribution of biocrusts. Therefore, it is crucial to understand
54 their distribution.

55 Despite the significance of biocrusts, previous studies have primarily focused on [their](#)
56 [contributions to C and N cycling in varying habitats and climates](#) (Hu et al., 2019; Morillas and
57 Gallardo, 2015), interspecific interactions and biocrusts biodiversity (Machado De Lima et al.,
58 2021; Munoz-Martin et al., 2019), rather than their spatial distribution, particularly at the global
59 scale. Consequently, a systematic and accurate assessment of biocrusts' ecological roles
60 remains challenging. In this study, we firstly sorted out the main research methods for studying
61 biocrusts distribution (section 1), then reviewed the existing knowledge (section 2), and finally
62 proposed strategies to advance the study of large-scale biocrusts distribution (section 3). This
63 work is expected to deepen our understandings of dryland ecosystem processes, and to provide

64 a scientific basis for dryland ecosystem conservation and their responses to global change.

65 **2. Research Methods**

66 In the study of biocrusts distribution, three methods are commonly used: spectral
67 characterization, vegetation dynamic modeling, and geospatial modeling. This section provides
68 an overview of these methods, including their basic principles, adaptability, and limitations.

69 **2.1 Spectral characterization index**

70 [With advances in remote sensing and geo-information technology, spectroscopy provides](#)
71 [a feasible method of characterizing distribution features from a physical point of view.](#)

72 Differences in absorption or reflection of specific wavelengths of different ground covers can
73 effectively identify soil surface objects (Rodriguez-Caballero et al., 2015). By identifying
74 biocrust-specific bands from the reflectance spectral images (Karnieli et al., 1999), it is possible
75 to construct a presence-absence map of biocrusts distribution (Fig. 1a).

76 The crust index (CI) and biological soil crust index (BSCI) are two of spectral
77 characterization indexes and have been successfully applied in Negev Desert (cyanobacteria-
78 dominated) (Karnieli, 1997; Noy et al., 2021) and Gurbantunggut Desert (lichen-dominated)
79 (Chen et al., 2005). For such indicators, it is critical to determine the threshold of spectral bands
80 that represent biocrusts. For instance, at an aerosol optical depth of 0.2, the BSCI ranges from
81 4.13 to 6.23, and narrows to 4.58-5.69 with increasingly poor atmospheric conditions. Overly
82 strict or loose threshold ranges can easily lead to biocrusts omission or misidentification. In
83 order to improve the accuracy of biocrusts identification, some researchers took advantage of
84 the hyperspectral sensor's continuous waveband, and created Continuum Removal Crust
85 Identification Algorithm (CRCIA) (Chamizo et al., 2012b; Weber et al., 2008). Baxter et al.
86 (2021) innovatively applied the random forest algorithm to spectral feature classification, and
87 achieved an accuracy of 78.5% in biocrusts recognition. Another two indexes, i.e., the sandy
88 land ratio crust index (SRCI) and the desert ratio crust index (DRCI), were also introduced by
89 taking into account the differences between sandy land (vegetation cover FVC <20%) and
90 desert environments, which could improve the accuracy of the mapping by ~6% (Wang et al.,
91 2022b).

92 The spectral characterization method is easy to use, and thus, facilitates the access to

93 continuous long-term dynamics of biocrusts distribution. However, mosses and vascular plants
94 are generally mixed up in this case because their reflectance characteristics are close to each
95 other in all wavelengths especially when mosses are wet, which makes them indistinguishable
96 (Fang et al., 2015). Therefore, spectral characterization method is mainly applicable to
97 situations where biocrusts cover is >30% and plants cover is <10% (Beaugendre et al., 2017).
98 It should be noted that, the existing indexes mostly correspond to biocrusts cover consisting of
99 specific dominant groups in specific environments, which cannot be directly extrapolated to
100 areas with high heterogeneous environments (Table 1). Wetting or disturbance may also lead to
101 large fluctuations in reflectance of different land types and interfere with biocrusts distribution
102 monitoring (Rodriguez-Caballero et al., 2015; Weber and Hill, 2016).


103 **2.2 Dynamic global vegetation models (DGVMs)**

104 Dynamic global vegetation model is another major method to obtain vegetation cover
105 (Deng et al., 2022). Dynamic global vegetation model mainly **focuses** on simulating the
106 biogeochemical processes (e.g. carbon, water cycles), metabolic and hydrological processes of
107 organisms (Lenton et al., 2016; Porada et al., 2017). The method possesses significant
108 advantages to map biocrusts distribution because its assumptions have clear biological
109 implications (Cuddington et al., 2013), yet may lead to poor predictions of global-scale
110 distributions due to subjective regional experience and insufficient amounts of biocrust data
111 (Table 1) (Quillet et al., 2010). To use DGVMs, the following procedures need to be taken. The
112 first step is building model framework, simulating important and interesting processes such as
113 biocrusts growth and death, nutrient cycle, and water cycle (Jia et al., 2019). So far, there is
114 only one dynamic global vegetation model targeting at biocrusts - the Lichen and Bryophyte
115 Model (LiBry) (Fig. **1b**) (Porada et al., 2019; Porada et al., 2013). The second step is
116 parameterization. Generally, literature and open databases are used to assign physiological
117 strategies for different types of biocrust, such as photosynthesis, **respiration**, and nitrogen
118 emission under the influence of temperature, precipitation, radiation, biological water
119 saturation, etc. The third step is importing environmental data into the model to obtain the
120 biocrusts cover at grid points over a specific study region. At last, the results are tuned and
121 validated against observation data of biocrusts distribution (obtained by literature comparison

122 and local field observations), and thereby biocrusts distribution map is produced.

123 **2.3 Geospatial models**

124 Directly relating vegetation presence or cover to environmental data instead of indirectly
125 via biological processes in DGVMs is another important way to obtain biocrust's distribution
126 (Beaugendre et al., 2017; Fischer and Subbotina, 2014; Skidmore et al., 2011). Classic
127 statistical models can serve for this purpose. However, they still require comprehensive expert
128 knowledge of how environmental factors affect biocrusts (Pearce et al., 2001), which is hard to
129 get and prone to be biased. Geospatial models which integrating machine learning tools with
130 field survey data and remote sensing data is supposed to hold the most promise (Crego et al.,
131 2022). Geospatial model is also known as species distribution model or ecological niche model
132 (Brown and Anderson, 2014; Jiménez-Valverde et al., 2008; Soberon and Nakamura, 2009).
133 The procedures of how to use geospatial modelling are illustrated in Figure 1c(Rodriguez-
134 Caballero et al., 2018): 1) extracting environmental data for the sites where biocrusts
135 observation data are reported, 2) importing the extracted environmental data into the machine
136 learning framework and obtaining the relationship between biocrusts distribution and
137 environmental variables through a specific algorithm (e.g., decision tree algorithms, Bayesian
138 algorithms, artificial neural networks, etc.), 3) simulating biocrusts distribution by
139 extrapolating to the whole study region using the constructed relationships. Nevertheless,
140 geospatial models are black-boxes and largely non-interpretable, and thus, less capable of
141 capturing key mechanisms behind phenomenon, which may limit its applications. One should
142 note that, to avoid confounding model predictions, inclusion of environmental factors should
143 be based on relevance of environmental factors to biocrusts, and still need expert knowledge to
144 a certain degree(Mäkinen et al., 2022). Besides, not only natural conditions such as climate,
145 topography, soil, etc. that affect biocrust distribution, but also data about human activities such
146 as afforestation, trampling, population density, etc. also need to be considered as environmental
147 indicators in the model. Supplied with sufficient computing power and observation data of
148 biocrust distribution, and suitable environmental, geospatial models are supposed to be able to
149 predict biocrust distribution accurately. Therefore, geospatial modelling is considered to be one
150 of the most appropriate methods available (Table 1).



151

152 **Fig. 1** Summary of three major approaches of studying biocrust distribution. Illuminations of
 153 applying spectral characterization method **(a)**, vegetation dynamics model **(b)** and geospatial
 154 model **(c)** in biocrusts distribution study. See main text for more detailed introduction to these
 155 methods.

156 **Table 1** Comparison among the three main types of methods to predict biocrusts distribution


	Spectral characteristic index	Vegetation dynamics model	Geospatial model
Principle	Differences in wavelength reflectance of surface features	Differences in the physiological processes of different biocrusts types	Remote sensing information-driven and survey data-based machine learning framework
Advantages	Convenience and ease of use	Clear ecological significance	Machine training simulation, without subjective interference
Disadvantages	Reflectivity is affected by climate change, disturbances; Mosses and vascular plants have similar reflectance characteristics;	Experience-based promotion with significant human intervention; Experiments need to be supported by big data	Large amount of computing power; Adequate number of sample points to support accuracy

	The results only show the presence or absence of biocrusts, without coverage		
Applicable scales	Regional scale (Desert and sandy land with <20% vegetation cover)	Regional scale Global scale	Regional scale Global scale

157

158 **3. Current State of Knowledge**

159 Since 1990, studies on the distribution of biocrusts [have](#) been continuously increasing. A
160 group of ecologists, represented by Fernando Maestre (Maestre et al., 2021), David
161 Eldridge (Eldridge and Delgado-Baquerizo, 2019; Eldridge et al., 2023), Matthew Bowker (Qiu
162 et al., 2023), Emilio Rodríguez-Caballero (Rodríguez-Caballero et al., 2018) and others, [have](#)
163 [actively promoted progress in the field \(Fig. 2\(a\)\). The topic has gradually received attention](#)
164 [from all over the world, particularly the countries with extensive dryland areas such as China,](#)
165 [United States, Spain, United Kingdom, Germany, **Australia**, and Israel \(Fig. 2\(b\)\). However,](#)
166 [some other dryland countries and regions, such as central and southern Africa, where biocrust](#)
167 [distribution has been reported still there is a paucity of studies and the amount of data on](#)
168 [biocrusts is far from adequate \(Fig. 2\(c\)\). These areas may be potential areas of widespread](#)
169 [distribution of biocrusts in the future.](#)



170

171 Fig. 2 Literature review of biocrust distribution studies. (a) Representative authors associated
 172 frameworks for biocrusts distribution studies (1990 to 2022). The time series is the average
 173 time of the year of publication, e.g., if the number of articles is 2 in 2004 and 8 in 2019, the
 174 node in this figure shows the year as $(2004 \times 2 + 2019 \times 8) / 10 = 2016$. (b) Map of hotspot
 175 countries for biocrust distribution research, with the top 12 countries in terms of number of
 176 publications shown; The database is Web of Science, TS = ("biogenic crust*" OR "biological

177 crust*" OR "biological soil crust*" OR "biocrust*" OR "microphytic crust*" OR "microbiotic
178 crust*" OR "cyanobacterial*" OR "algal*" OR "lichen*" OR "moss*" OR "biotic crust*") AND
179 ("mapping*" OR "distribution*" OR "spatial pattern*") AND (“dryland” OR “hyper*arid*” OR
180 “arid*” OR “semi*arid*” OR “dry subhumid*”), with research interests in Environmental
181 Sciences/Ecology and a total of 700 papers. [\(c\) Global biocrust data distribution, based on field](#)
182 [surveys and literature compilation. Data have been collected and expanded from the published](#)
183 [database](#) (Chen et al., 2020; Rodriguez-Caballero et al., 2018) [to 3848 items.](#)

184 **3.1 Local-regional scales**

185 At local-regional scales, numerous studies have provided valuable insights into the
186 distribution patterns of biocrusts in different regions around the world [\(Fig. 3\)](#). In the Mojave
187 Desert, biocrusts distribution was closely related to geological age, surface stability, topography,
188 and dust transport (Miller et al., 2004). [Lichen](#), moss, and dark algal crusts patchily distributed
189 on [the desert](#), averaging 8% cover, though in some bar and shrub zones, the cover could be as
190 high as [26%](#) (Pietrasiak et al., 2014). In the Colorado Plateau, highly heterogeneous soil matrix
191 determined the fragmented biocrusts distribution of different types and the wide disparity in
192 relative abundance and cover (Reynolds et al., 2006; Steven et al., 2013). Collier et al. (2022)
193 trained drone imagery in the Hawaiian region using timely data collected by cameras and then
194 successfully mapped watershed-scale biocrusts distribution, predicting cover of ~15-23%. In
195 the Gurbantunggut Desert, biocrusts cover 28.7% of the area, with a high and uniform biocrusts
196 cover in the southern part of the desert and a scattered distribution of biocrusts in other areas
197 (Chen et al., 2005; Zhang et al., 2007). [In the Loess Plateau](#), RGB image-based biocrusts
198 monitoring showed that variability in biocrusts cover decreased logarithmically with increasing
199 plot size until a critical size of 1m² after which biocrusts cover remained approximately constant
200 (Wang et al., 2022a). In Qatar, 26% of the country is covered by biocrusts, with cyanobacterial
201 biocrusts cover showing a decreasing trend from north-east to south-west (Richer et al., 2012).
202 In the northern Negev, the distribution of biocrusts is well developed, by simulating the spectral
203 characteristics of the different components of biocrusts after seasonal precipitation, especially
204 the chlorophyll absorption characteristics, it was found that the greening of the biocrusts was
205 obvious, which could provide a basis for the subsequent tracking of their distribution (Panigada

206 et al., 2019). At the Sinai Peninsula (Egypt) – Negev desert (Israel) border, the distribution
207 dynamics of cyanobacterial biocrusts over a 31-year period has now been obtained from the
208 crust index (Noy et al., 2021). Filamentous cyanobacteria grow in the African Sahara (Issa et
209 al., 1999), where statistical models of combined environmental indicators showed biocrusts
210 cover of 1 ~ 48% and 0 ~ 65% in Banizoumbou and Tamou respectively (Beaugendre et al.,
211 2017).

212 **3.2 Global scale**

213 To date, there are only two global-scale studies of biocrusts distribution. Porada et al. (2013)
214 focused on CO₂ diffusion rates and photosynthetic processes under dynamic water content
215 saturation in dryland biocrusts. By parameterizing long-term climate and disturbance intervals
216 and averaging simulation results for the past 20 years for each grid point, they estimated that
217 biocrusts cover 11% of global terrestrial land surface (Porada et al., 2019). Their results also
218 showed that the light and dark cyanobacteria were widely distributed in deserts, savannas,
219 grasslands, and Mediterranean woodlands at low latitudes, and increase to some extent with
220 increasing dryness, while mosses were mostly distributed in middle and high latitudes and polar
221 regions (Fig. **3a**). Rodriguez-Caballero et al. (2018) fitted biocrusts presence-only data to the
222 bioclimate, soil properties, land use data, and then extrapolated to a continuous global
223 distribution of biocrusts using the Maxent model. This work assessed the total area covered by
224 biocrusts to be 1.79×10^7 km², 12.2% of global drylands, in other words, biocrusts cover was
225 1.2% larger compared to the area predicted by Porada et al. (2019) (Fig. **3b**). After comparing
226 two maps, biocrust distribution is generally consistent in the large deserts of Asia, western
227 America, Europe and Oceania, while some semi-arid regions, such as the northern and southern
228 margins of African Sahara Desert, South Asia and central North America, have significantly
229 higher biocrusts cover in the latter projection of Rodriguez-Caballero et al. (2018). We estimate
230 that the reason may be that geospatial modeling focuses more on the influence of climate, as
231 the Mediterranean climate and tropical desert climate in the Sahara Desert, as well as the
232 tropical desert climate of northwestern South Asia, which is suitable for biocrust surviving.
233 Additionally, the large number and high cover of biocrust training sets in the central North
234 America could have contributed to the generally high predicted cover in machine learning.




Fig. 3 Maps of global biocrusts distribution. (a) **Prediction** based on vegetation dynamic model (Porada et al., 2019). (b) **Prediction** based on geospatial model (Rodriguez-Caballero et al., 2018).

4. Influencing **Factors** of **Biocrust** Distribution

Numerous experimental observations and modelling (Kidron and Xiao, 2023; Li et al., 2023; Rodriguez-Caballero et al., 2018) have proved that, on the global scale, biocrust distribution is mainly influenced by water conditions, temperature, soil properties, fire and [disturbance](#) (Bowker et al., 2016).

[Water conditions.](#) In general, [total precipitation \(Fig. 4\(b\)\)](#) is considered to be critical in determining the distribution of biocrusts (Eldridge and Tozer, 1997). Increased precipitation could lead to higher levels of lichen and moss, while algal cover may first increase and then decrease (Budel et al., 2009; Marsh et al., 2006; Zhao et al., 2014). It should be noted that, precipitation can also promote the growth of vascular plants, and the continuous and high cover of vascular plants and litterfall will limit the space available to biocrusts (Bowker et al., 2005). In addition to the total amount of precipitation, the seasonality and frequency of precipitation cannot be ignored (Budel et al., 2009). [Winter precipitation and/or smaller rain events benefit biocrusts, especially when mean annual precipitation is <500 mm, and high frequency of precipitation can lead to the dominance of biocrusts over vascular plants](#) (Chamizo et al., 2016; Jia et al., 2019). [It was experimentally proven that precipitation events of 5 mm were able to maintain normal physiological and ecological functions of the biocrust on the Colorado Plateau, USA, while ever lower precipitation event of 1.2 mm could rapidly kill the moss biocrust](#) (Reed et al., 2012). Non-precipitation water input is another important water resource type. The Namib Desert receives little rainfall, but lichens and moss biocrusts can reach a relatively high cover (~70%) (Budel et al., 2009). This is because local water vapour tends to condense into fog or

260 dew, which facilitates the survival of three-dimensional species (such as leafy lichens) by
261 trapping air moisture (Eldridge et al., 2020; Kidron, 2019; Li et al., 2021). Similarly, lichen
262 biocrusts are widely distributed in the western U.S. along with the Mexican coast, as the result
263 of the high air humidity (dew formation for almost 1/3 of the year) (McCune et al., 2022;
264 Miranda - González and McCune, 2020).

265 Temperature. Relatively high soil temperature can create an environment of high
266 evaporation that impedes biocrusts colonization (García-Pichel et al., 2013). Regarding air
267 temperature, warming by 4°C could alter biocrust community structure, resulting in a sharp
268 decrease in moss biocrust cover and an increase in cyanobacterial biocrust cover, which became
269 even more significant when warming was interacting with time and precipitation treatments
270 (Ferrenberg et al., 2015). Recent studies have shown that historical and future temperature
271 changes also affect biocrust distribution. For example, the climate legacy over the last 20,000
272 years could indirectly affect the distribution and relative species richness of biocrusts through
273 changing vegetation cover and soil pH (Eldridge and Delgado-Baquerizo, 2019). Additionally,
274 under future scenarios of increased temperature and aridity, biocrusts cover is predicted to
275 decrease by approximately 25% by the end of the century, with communities shifting towards
276 early cyanobacterial biocrusts (Rodríguez-Caballero et al., 2022).

277 Soil properties. For a long time, it was commonly believed that finer soils benefit biocrusts
278 growth (Belnap et al., 2014; Williams et al., 2013). However, this has been challenged by some
279 scientists (Fig. 4(c)). For example, Kidron (2018) argued that soils with high dust or fine grains
280 were not a necessary condition for biocrusts distribution. Qiu et al. (2023) suggested that soils
281 with small amounts of gravel (0.04-22.34% content, 0.58% best) are more favorable for
282 biocrusts. Another study had shown that the soil parent material determines the degree of
283 surface weathering and the water-holding capacity of the soil, thus indirectly changing the
284 distribution of biocrusts (Bowker and Belnap, 2008). Gypsum or calcareous soils tend to
285 develop mosses and lichens (Elbert et al., 2012), while sandy soils tend to develop
286 cyanobacteria (Root and McCune, 2012).

287 Fire. Grassland is one major lifeform in dryland ecosystems, making it of great
288 significance to explore the effects of fire events on biocrusts distribution (Palmer et al., 2022).


289 Fire-induced soil warming can alter the resource allocation and dynamic growth mechanisms
290 between biocrusts and vascular plants (Mccann et al., 2021), potentially leading to a reduction
291 in species richness and cover of biocrusts, especially cyanobacteria and algae (Abella et al.,
292 2020; Palmer et al., 2020). (Condon and Pyke, 2018) showed that moss cover increases with
293 time after fire, with no significant change in lichen cover.

294 Disturbance. Activities such as grazing, agricultural practices and land development can
295 significantly impact biocrust distribution. Studies have demonstrated that grazing intensity can
296 lead to substantial changes in biocrust cover. For instance, in Patagonian rangelands, biocrust
297 cover decreased by 85%, 89%, and 98% under light, medium, and heavy grazing, respectively
298 (Velasco Ayuso et al., 2019). In the Loess Plateau, total biocrust cover remained almost
299 unchanged under light grazing (< 30.00 goat dung / m²), but there were variations in community
300 structure, with an increase in cyanobacteria biocrusts (23.1%) and a decrease in moss biocrusts
301 (42.2%) due to the decrease in vascular plant cover (Ma et al., 2023). Tillage practices can
302 distrupt the soil surface, leading to a reduction in biocrust cover (6% on average) and diversity,
303 and lichens tend to struggle to survive in tilled fields compared to mosses (Durham et al., 2018).
304 Additionally, late successional biocrusts exhibit higher tolerance compared to pre-successional
305 biocrusts. Moss biocrusts, for instance, can maintain soil microbial biomass and nematode
306 abundance better under trampling disturbance compared to cyanobacteria and lichen biocrusts
307 (Yang et al., 2018). However, contrary to this view, it has been observed that cyanobacterial
308 biocrusts increased in cover from 81% to 99% after trampling, while lichen and moss biocrusts
309 decreased from 1.5% and 18% to less than 0.5%. Furthermore, mining activities can
310 significantly reduce the photosynthetic potential of biocrusts, particularly affecting the recovery
311 of cyanobacterial biocrusts (Gabay et al., 2022).

312 Other factors. At the global scale, biocrust distribution is also closely linked to
313 biogeographic isolation. Strong spatial heterogeneity accompanied with spatial distance can set
314 up barriers to the dispersal of propagules (spores, fungal bodies), which indirectly impedes
315 colonization of the biocrusts (Garcia-Pichel et al., 2013). In addition, factors such as vascular
316 plant cover, topography, and solar radiation also influence biocrust distribution, albert to a lesser
317 extent than the factors mentioned in the above paragraphs. For further insights, readers are

318 [encouraged to consult Chapter 10 of *Biological Soil Crusts: An Organizing Principle in*](#)
319 [Drylands, which overview of the control and distribution patterns of biocrust from micro to](#)
320 [global scales](#) (Bowker et al., 2016).

321 [To sum, climate is the most important factor of influencing global biocrust distribution,](#)
322 [especially in drylands where water is precious to the organisms. But exploration of the roles of](#)
323 [climatic factors such as rainfall seasonality and atmospheric drought still needs much more](#)
324 [further efforts](#) (Wright and Collins, 2024), [especially context of global climate change.](#)
325 [Although more attention has been paid to physical properties of soils, the roles of its chemical](#)
326 [properties such as the N, P content need to be taken more seriously. Fire and disturbance are](#)
327 [usually ignored. Whereas due to the trend towards warmer and drier environments, as well as](#)
328 [increasing population and the need to sustain livelihoods, their influences on biocrust](#)
329 [distribution may become more important. As one of the basic processes on global scale,](#)
330 [biogeographic isolation or changes in land use should be paid more attentions. As amounting](#)
331 [data points of biocrust, we can expect this aspect will see a surge in research.](#)



332 [Fig. 4 Biocrust distribution and its critical influencing factors. \(a\) Biocrust cover map and its](#)
333 [influencing factors. \(a\) Global biocrust distribution, by random forest modelling. Based on a](#)
334 [global biocrust database constructed by Chen et al., we expanded the biocrust data to 3848](#)
335 [entries through literature compilation and field surveys and fitted them with four types of](#)

336 [remotely sensed environmental data, including climate, land use, soil properties, and elevation,](#)
337 [to finally predict the suitable areas for the biocrust distribution and quantify the biocrust cover.](#)
338 [\(b\) Global average annual precipitation \(1970-2020\), data from the WorldClim database](#)
339 [\(version 2.1\). \(c\) Global soil texture distribution, data from HWSD \(Harmonized World Soil](#)
340 [Database, version 1.2\). Precipitation and soil texture were taken as examples of environmental](#)
341 [factors.](#)

342 **5. Challenges and Perspectives**

343 Biocrusts are very important for dryland ecosystems, and thus, [it is of outstanding](#)
344 [significance to understand the current status and dynamics of biocrusts distribution. For the](#)
345 [influencing factors \(Chapter 4\), traditional observation and control experiments provide us with](#)
346 [multiple perspectives of basic knowledge. For assessing biocrust distribution patterns \(Chapter](#)
347 [3\), the methods](#) are shifting from traditional [approaches](#) to [spectral index, vegetation dynamics](#)
348 [and geospatial model, that span multiple subjects like](#) ecology, biology, geology and computer
349 science. However, high-precision biocrusts distribution data across geographic units are still
350 lacking, and research methods are still limited. To further advance studies of biocrusts
351 distribution, we raise the following aspects.

352 **5.1 Building standardized biocrusts database**

353 [Currently, biocrust data are fragmented, low in volume and accessed from narrow sources,](#)
354 [largely limiting spatial prediction from points to areas. Thus, we suggest that a global effort](#)
355 [should build a](#) standardized and specialized biocrusts database, consisting of the same data
356 items (main types and cover of biocrusts, latitude, longitude and cover, etc.) and using the same
357 inclusion criteria. [It](#) is an important infrastructure for mapping global biocrusts distribution,
358 serving as the benchmark to train and validate spectral characteristics, DGVM, and geospatial
359 models (Engel et al., 2023). Due to the difficulty of conducting field surveys worldwide,
360 compiling biocrusts data from the published literatures or other [sources would be](#) the major
361 approach (Fig. 4(a)). [To date, several published studies have assembled 900 ~ 1,000 data on](#)
362 [biocrust presence or absence from the literature \(including 584 data on biocrust cover\)](#) (Chen
363 et al., 2020; Eldridge et al., 2020; Havrilla et al., 2019; Rodriguez-Caballero et al., 2018).
364 However, compiling from literatures largely comes to its limitation and is still far from building

365 a standardized and specialized biocrusts database. While open databases are not specialized to
366 biocrusts, some of them may be important additions [\(Fig. 5\)](#). For example, the biodiversity and
367 specimen datasets such as GBIF and the Atlas of Living Australia (Belbin and Williams, 2015;
368 García-Roselló et al., 2015), which contain a large amount of information on species, of course
369 including mosses and lichens (Table 2), which can contain hundreds or even thousands of
370 entries of biocrusts occurrence or cover. Similarly, global, national and regional plant flora may
371 significantly contribute to building the standardized and specialized biocrusts database. For
372 example, sPlot includes ~2 millions of vegetation plot data (Sabatini et al., 2021), and the
373 European Vegetation Archive (*EVA*) also possesses 1.6 million entries over globe or Europe
374 (Chytrý et al., 2016). Regional datasets like Environmental Monitoring of arid and Semiarid
375 Regions (*MARAS*) surveyed 426 sites (up to September 2020), and provided regular access to
376 624.50 km² of rangeland vegetation spatial patterns, species diversity data, soil functional
377 indices, climatic data and landscape photographs in Patagonia region straddling Argentina and
378 Chile (Oliva et al., 2020). [Concerns about land use products are also necessary. Global land use
379 maps, based on the PROBA-V sensor, which contain spatial information for Moss & Lichen
380 layer, have an annual update frequency and a resolution of 100 m.](#) In addition to above channels,
381 an increasing number of amateurs are getting involved in science through species identification
382 apps with clean, easy-to-use apps, contributing significantly to global species information
383 entries. The citizen science project *iNaturalist* is a very good example (Wolf et al., 2022).
384 Furthermore, when collecting and collating data from the non-academic sources, the
385 combination of web crawlers and text analysis can help in obtaining biocrusts data and solving
386 key ecological issues._



387

388 **Fig. 5** Potential approaches to build a standardized biocrusts database. (a) Distribution of
 389 lichens in the GBIF database with an example photo, (b) environmental monitors distribution
 390 map of MARAS database, (c) distribution of "mosses and lichens" in the PROBAV_LC100
 391 database (light yellow area) in northern Asia, for instance.

392

Table 2 References for biocrusts database expansion channels

5.2 Improving non-vascular vegetation dynamic models

Data type	Data source	Extend	Biocrust type	Georeferenced records	Presence	Coverage	Link
Biodiversity data	the Global Biodiversity Information Facility(GBIF)	Worldwide	Cyanobacteria	~780000	✓	--	https://www.gbif.org/
			Lichen	~19000			
			Moss	~90000			
	Atlas of Living Australia(ALA)	Australia	Cyanobacteria	~53000	✓	--	https://www.ala.org.au/
			Lichen	~12000			
			Moss	~20000			
	Chinese Virtual Herbarium	China	Moss and lichen	--	✓	--	https://www.cvh.ac.cn/
	Global Plants on JSTOR	Worldwide	Lichen	~2000	✓	--	https://plants.jstor.org/
			Moss	~480			
Citizen Science	iNaturalist	Worldwide	All	--	✓	--	https://www.inaturalist.org/
	MARAS	Argentina and Chile	All	426	✓	✓	https://springernature.figshare.com/collections/The_MARAS_dataset_vegetation_and_soil_characteristics_of_dryland_rangelands_across_Patagonia/4789113
Survey data	sPlot	Worldwide	Lichen	6801	✓	✓	https://www.idiv.de/en/splot.html
			Moss	11001	✓	✓	
	GrassPlot	Worldwide	Non-vascular plants	6623	✓	✓	https://edgg.org/databases/GrassPlot/
			Moss and lichen	~15000	✓	✓	http://vegbank.org/
	BLM_AIM	Canada and the United States	Moss and lichen	5200	✓	✓	https://gbp-blm-egis.hub.arcgis.com/pages/aim
Landcover data	TERN AEKOS	Australia	All	~300			http://www.aekos.org.au/
	PROBAV_LC100	Worldwide	Moss and lichen	--			https://land.copernicus.eu/global/products/lc

395 There are only two DGVMs applicable to non-vascular organisms – LiBry and ECHAM6-
396 HAM2-BIOCRUST (Rodriguez-Caballero et al., 2022). Their performances still need to be
397 improved. Considering spatial self-organization of non-vascular organisms (Gassmann et al.,
398 2000), the effects of fire (Thonicke et al., 2001), vegetation-environment feedback processes
399 (Quillet et al., 2010), functional traits (Boulangeat et al., 2012), intraspecific-interspecific
400 interactions (Boulangeat et al., 2014) and seasonal dynamics in current and/or new DGVMs
401 can be future directions. In addition, how the physical properties, photosynthetic capacity,
402 carbon and nitrogen allocation of biocrusts change along environmental gradient are complex
403 and context-dependent, which should be incorporated into DGVMs (Faticchi et al., 2019).
404 Spatial-explicit DGVMs may be one key to effectively improving the accuracy of simulations
405 in future studies, but are data-consuming. Also, biocrusts are significantly influenced by
406 hydrological processes and vice versa (Chen et al., 2018; Whitney et al., 2017), while
407 ecohydrological models based on hydrological processes are rarely connected to global
408 biocrusts distribution predictions. (Jia et al., 2019) tried to incorporate biocrusts cover as a
409 system state variable in an ecohydrological model and investigated biocrusts cover under
410 rainfall gradient. If fed with global data of environmental variables (mainly hydrological
411 relevant ones), ecohydrological models may be a new approach to predicting biocrusts
412 distribution at global scales.

413 **5.3 Integrated application of high-quality sensors**

414 Spectral characterization method lies on the differences in spectral reflectance of biocrusts
415 and other land types at different wavelengths, and thus, the accuracy of the results depends on
416 the quality of the sensors. However, previous studies have often used a single sensor with
417 constant band intervals for distinguishing biocrusts, which may result in missed spectral feature
418 identification of land types (Chamizo et al., 2012a). If the biocrusts index can be constructed
419 by combining and comparing the full-band spectral data from multiple terrestrial sensors and
420 infrared cameras and other devices, the errors will be reduced to a certain extent, thus improving
421 the classification accuracy (Wang et al., 2022b). In addition, the unique advantages of
422 hyperspectral data with large data volume and narrow band allow it to be used to combine with
423 observation data to co-develop new biocrusts discrimination standard. If further estimation of

424 biocrusts cover can be achieved on this basis, it will be an important contribution to the study
425 of large scale biocrusts distribution (Rodríguez-Caballero et al., 2017). [To date](#), high resolution
426 sensors have been found to be successful in monitoring lichens and mosses (Blanco-Sacristan
427 et al., 2021) and the release of such products is something important to look out for in the future.

428 **[5.4 Making full use of machine learning](#)**

429 Machine learning can be combined with remote sensing products to find complex features
430 from big data to predict global biocrust distribution (Collier et al., 2022). This data-driven
431 approach has powerful predictive capabilities, especially for mapping species distribution, and
432 can largely circumvent the mistakes of missing or misidentifying biocrusts caused by traditional
433 methods (relying on field measurements to determine threshold ranges) (Wang et al., 2022b).
434 In the remote sensing image classification problem, mature machine learning algorithms
435 include support vector machines, single decision trees, random forests, artificial neural
436 networks, etc. (Yu et al., 2020). Ensemble models combining multiple algorithms have been
437 widely used in the field of species distribution, but relatively few applications exist in field of
438 biocrusts prediction. In the future, using machine learning to find parameters for dynamic
439 models of biocrusts, which may be one of the most promising method to predict biocrusts
440 distribution (Perry et al., 2022).

441 **[5.5 Regional research synergy development](#)**

442 [Research of biocrust distribution have shown significant spatial and climate imbalances.](#)
443 Spatially, the study areas that have been conducted are relatively concentrated in countries such
444 as China, USA, Spain, [Australia](#), and Israel. Although there are large areas of dryland
445 distributed in Africa (other than South Africa), central Asia, central South America and northern
446 North America, research on biocrusts in these regions are scarce. Unbalanced regional research
447 efforts are one of the constraints to advancing studies of global biocrusts distribution. Therefore,
448 how to coordinate and promote the common progress of regional research is an urgent issue at
449 present. [Climatically](#), in addition to the drylands, the cold zones may be another important area
450 to explore biocrusts distribution (Pushkareva et al., 2016). On the Tibetan Plateau, studies have
451 investigated the spatial variation of different types of biocrusts communities across climatic
452 gradients and their effects on soil temperature features and freezing duration (Ming et al., 2022;

453 Wei et al., 2022). This urges the studies of biocrusts distribution in the alpine areas.

454 **6. Conclusion**

455 [This work aims to advance global knowledge of biocrust distribution for better ecosystem](#)
456 [management and sustainable development in drylands. We firstly compared the advantages,](#)
457 [disadvantages, and applicability among three methods, spectral characterization index,](#)
458 [dynamic global vegetation models and geospatial models, in order to provide the most](#)
459 [appropriate methodological suggestions for biocrust distribution studies at different scales and](#)
460 [needs. Then, we systematically sorted out the regional-global biocrust distribution cases, and](#)
461 [drew a map of global biocrust distribution hotspots and a map of spatial distribution of data](#)
462 [points. Further, we tried to clarify the causes of biocrust distribution from several aspects, such](#)
463 [as precipitation, temperature, soil, fire, and other anthropogenic factors. Finally, from a personal](#)
464 [point of view, we would like to focus more on the following points in the future: database](#)
465 [construction, model performance enhancement, big data processing, and synergistic progress](#)
466 [of potential distribution area studies.](#)

467 **Acknowledgements**

468 [We are grateful to Zhiguang Zhao, Lixun Zhang, Chao Guan for helpful discussions when](#)
469 [preparing the first draft.](#) The study was supported by the National Natural Science Foundation
470 of China (grant number [32271620, 31971452](#)), [National Key Research and Development](#)
471 [Program of China \(2023YFF0805603\)](#), and the Central University Basic Research Funds
472 [\(lzujbky-2021-ey16, lzujbky-2023-eyt01\)](#).

473

474 **Reference**

475 Abella, S. R., Gentilcore, D. M., and Chiquoine, L. P.: Resilience and alternative stable states after desert
476 wildfires, *Ecological Monographs*, 91, 1-19, 10.1002/ecm.1432, 2020.
477 Baxter, C., Mallen-Cooper, M., Lyons, M. B., and Cornwell, W. K.: Measuring reflectance of tiny
478 organisms: The promise of species level biocrust remote sensing, *Methods in Ecology and Evolution*, 12,
479 2174-2183, 10.1111/2041-210x.13690, 2021.
480 Beaugendre, N., Malam Issa, O., Choné, A., Cerdan, O., Desprats, J.-F., Rajot, J. L., Sannier, C., and
481 Valentin, C.: Developing a predictive environment-based model for mapping biological soil crust patterns
482 at the local scale in the Sahel, *Catena*, 158, 250-265, 10.1016/j.catena.2017.06.010, 2017.
483 Belbin, L. and Williams, K. J.: Towards a national bio-environmental data facility: experiences from the
484 Atlas of Living Australia, *International Journal of Geographical Information Science*, 30, 108-125,
485 10.1080/13658816.2015.1077962, 2015.
486 Belnap, J., Miller, D. M., Bedford, D. R., and Phillips, S. L.: Pedological and geological relationships

487 with soil lichen and moss distribution in the eastern Mojave Desert, CA, USA, *Journal of Arid*
488 *Environments*, 106, 45-57, 10.1016/j.jaridenv.2014.02.007, 2014.

489 Blanco-Sacristan, J., Panigada, C., Gentili, R., Tagliabue, G., Garzonio, R., Martin, M. P., Ladron de
490 Guevara, M., Colombo, R., Dowling, T. P. F., and Rossini, M.: UAV RGB, thermal infrared and
491 multispectral imagery used to investigate the control of terrain on the spatial distribution of dryland
492 biocrust, *Earth Surf Process Landf*, 46, 2466-2484, 10.1002/esp.5189, 2021.

493 Boulangeat, I., Georges, D., and Thuiller, W.: FATE-HD: a spatially and temporally explicit integrated
494 model for predicting vegetation structure and diversity at regional scale, *Global Change Biology*, 20,
495 2368-2378, 10.1111/gcb.12466, 2014.

496 Boulangeat, I., Philippe, P., Abdulhak, S., Douzet, R., Garraud, L., Lavergne, S., Lavorel, S., van Es, J.,
497 Vittoz, P., and Thuiller, W.: Improving plant functional groups for dynamic models of biodiversity: at the
498 crossroads between functional and community ecology, *Global Change Biology*, 18, 3464-3475,
499 10.1111/j.1365-2486.2012.02783.x, 2012.

500 Bowker, M. A. and Belnap, J.: A simple classification of soil types as habitats of biological soil crusts on
501 the Colorado Plateau, USA, *Journal of Vegetation Science*, 19, 831-840, 10.3170/2008-8-18454, 2008.

502 Bowker, M. A., Belnap, J., Davidson, D. W., and Phillips, S. L.: Evidence for Micronutrient Limitation
503 of Biological Soil Crusts: Importance to Arid-Lands Restoration, *Ecological Applications*, 15, 1941-1951,
504 10.1890/04-1959, 2005.

505 Bowker, M. A., Belnap, J., Büdel, B., Sannier, C., Pietrasiak, N., Eldridge, D. J., and Rivera-Aguilar, V.:
506 Controls on Distribution Patterns of Biological Soil Crusts at Micro- to Global Scales, in: *Biological Soil*
507 *Crusts: An Organizing Principle in Drylands*, edited by: Weber, B., Büdel, B., and Belnap, J., Springer
508 International Publishing, Cham, 173-197, 10.1007/978-3-319-30214-0_10, 2016.

509 Brown, J. L. and Anderson, B.: SDMtoolbox: a python-based GIS toolkit for landscape genetic,
510 biogeographic and species distribution model analyses, *Methods in Ecology and Evolution*, 5, 694-700,
511 10.1111/2041-210x.12200, 2014.

512 Budel, B., Darienko, T., Deuschewitz, K., Dojani, S., Friedl, T., Mohr, K. I., Salisch, M., Reisser, W.,
513 and Weber, B.: Southern African biological soil crusts are ubiquitous and highly diverse in drylands,
514 being restricted by rainfall frequency, *Microb Ecol*, 57, 229-247, 10.1007/s00248-008-9449-9, 2009.

515 Chamizo, S., Cantón, Y., Miralles, I., and Domingo, F.: Biological soil crust development affects
516 physicochemical characteristics of soil surface in semiarid ecosystems, *Soil Biology and Biochemistry*,
517 49, 96-105, 10.1016/j.soilbio.2012.02.017, 2012a.

518 Chamizo, S., Belnap, J., Eldridge, D. J., Cantón, Y., and Malam Issa, O.: The Role of Biocrusts in Arid
519 Land Hydrology, in: *Biological Soil Crusts: An Organizing Principle in Drylands*, edited by: Weber, B.,
520 Büdel, B., and Belnap, J., Springer International Publishing, Cham, 321-346, 10.1007/978-3-319-30214-
521 0_17, 2016.

522 Chamizo, S., Stevens, A., Canton, Y., Miralles, I., Domingo, F., and Van Wesemael, B.: Discriminating
523 soil crust type, development stage and degree of disturbance in semiarid environments from their spectral
524 characteristics, *European Journal of Soil Science*, 63, 42-53, 10.1111/j.1365-2389.2011.01406.x, 2012b.

525 Chen, J., Zhang, M. Y., Wang, L., Shimazaki, H., and Tamura, M.: A new index for mapping lichen-
526 dominated biological soil crusts in desert areas, *Remote Sensing of Environment*, 96, 165-175,
527 10.1016/j.rse.2005.02.011, 2005.

528 Chen, N., Jayaprakash, C., Yu, K., and Guttal, V.: Rising Variability, Not Slowing Down, as a Leading
529 Indicator of a Stochastically Driven Abrupt Transition in a Dryland Ecosystem, *Am Nat*, 191, E1-E14,
530 10.1086/694821, 2018.

531 Chen, N., Yu, K. L., Jia, R. L., Teng, J. L., and Zhao, C. M.: Biocrust as one of multiple stable states in
532 global drylands, *Science Advances*, 6, 10.1126/sciadv.aay3763, 2020.

533 Chytrý, M., Hennekens, S. M., Jiménez-Alfaro, B., Knollová, I., Dengler, J., Jansen, F., Landucci, F.,
534 Schaminée, J. H. J., Acíć, S., Agrillo, E., Ambarlı, D., Angelini, P., Apostolova, I., Attorre, F., Berg, C.,
535 Bergmeier, E., Biurrun, I., Botta-Dukát, Z., Brisse, H., Campos, J. A., Carlón, L., Čarni, A., Casella, L.,
536 Csiky, J., Čušterevska, R., Dajić Stevanović, Z., Danihelka, J., De Bie, E., de Ruffray, P., De Sanctis, M.,
537 Dickoré, W. B., Dimopoulos, P., Dubyna, D., Dziuba, T., Ejrnaes, R., Ermakov, N., Ewald, J., Fanelli, G.,
538 Fernández-González, F., FitzPatrick, Ú., Font, X., García-Mijangos, I., Gavilán, R. G., Golub, V.,
539 Guarino, R., Haveman, R., Indreica, A., Işık Gürsoy, D., Jandt, U., Janssen, J. A. M., Jiroušek, M., Kački,
540 Z., Kavgacı, A., Kleikamp, M., Kolomiychuk, V., Krstivojević Ćuk, M., Krstonošić, D., Kuzemko, A.,
541 Lenoir, J., Lysenko, T., Marcenò, C., Martynenko, V., Michalcová, D., Moeslund, J. E., Onyshchenko,
542 V., Pedashenko, H., Pérez-Haase, A., Peterka, T., Prokhorov, V., Rašomavičius, V., Rodríguez-Rojo, M.
543 P., Rodwell, J. S., Rogova, T., Ruprecht, E., Rūsiņa, S., Seidler, G., Šibík, J., Šilc, U., Škvorc, Ž.,
544 Sopotlieva, D., Stančić, Z., Svenning, J.-C., Swacha, G., Tsiripidis, I., Turtureanu, P. D., Uğurlu, E.,
545 Uogintas, D., Valachovič, M., Vashenyak, Y., Vassilev, K., Venanzoni, R., Virtanen, R., Weekes, L.,
546 Willner, W., Wohlgemuth, T., Yamalov, S., and Pärtel, M.: European Vegetation Archive (EVA): an
547 integrated database of European vegetation plots, *Applied Vegetation Science*, 19, 173-180,
548 10.1111/avsc.12191, 2016.

549 Collier, E. A., Perroy, R. L., Reed, S. C., and Price, J. P.: Mapping biological soil crusts in a Hawaiian
550 dryland, *International Journal of Remote Sensing*, 43, 484-509, 10.1080/01431161.2021.2003904, 2022.

551 Condon, L. A. and Pyke, D. A.: Fire and Grazing Influence Site Resistance to *Bromus tectorum* Through
552 Their Effects on Shrub, Bunchgrass and Biocrust Communities in the Great Basin (USA), *Ecosystems*,
553 21, 1416-1431, 10.1007/s10021-018-0230-8, 2018.

554 Crego, R. D., Stabach, J. A., Connette, G., and Leroy, B.: Implementation of species distribution models
555 in Google Earth Engine, *Diversity and Distributions*, 28, 904-916, 10.1111/ddi.13491, 2022.

556 Cuddington, K., Fortin, M., Gerber, L., Hastings, A., Liebhold, A., O'Connor, M., and Ray, C.: Process-
557 based models are required to manage ecological systems in a changing world, *Ecosphere*, 4, 1-12, 2013.

558 Deng, M., Meng, X., Lu, Y., Li, Z., Zhao, L., Niu, H., Chen, H., Shang, L., Wang, S., and Sheng, D.: The
559 Response of Vegetation to Regional Climate Change on the Tibetan Plateau Based on Remote Sensing
560 Products and the Dynamic Global Vegetation Model, *Remote Sensing*, 14, 3337, 10.3390/rs14143337,
561 2022.

562 Durham, R. A., Doherty, K. D., Antoninka, A. J., Ramsey, P. W., and Bowker, M. A.: Insolation and
563 disturbance history drive biocrust biodiversity in Western Montana rangelands, *Plant and Soil*, 430, 151-
564 169, 10.1007/s11104-018-3725-3, 2018.

565 Elbert, W., Weber, B., Burrows, S., Steinkamp, J., Buedel, B., Andreae, M. O., and Poeschl, U.:
566 Contribution of cryptogamic covers to the global cycles of carbon and nitrogen, *Nature Geoscience*, 5,
567 459-462, 10.1038/ngeo1486, 2012.

568 Eldridge, D. J. and Delgado-Baquerizo, M.: The influence of climatic legacies on the distribution of
569 dryland biocrust communities, *Global Change Biology*, 25, 327-336, 10.1111/gcb.14506, 2019.

570 Eldridge, D. J. and Tozer, M. E.: Environmental factors relating to the distribution of terricolous
571 bryophytes and lichens in semi-arid eastern Australia, *Bryologist*, 100, 28-39, 1997.

572 Eldridge, D. J., Reed, S., Travers, S. K., Bowker, M. A., Maestre, F. T., Ding, J., Havrilla, C., Rodríguez-
573 Caballero, E., Barger, N., Weber, B., Antoninka, A., Belnap, J., Chaudhary, B., Faist, A., Ferrenberg, S.,
574 Huber-Sannwald, E., Malam Issa, O., and Zhao, Y.: The pervasive and multifaceted influence of biocrusts

575 on water in the world's drylands, *Global Change Biology*, 26, 6003-6014, 10.1111/gcb.15232, 2020.

576 Eldridge, D. J., Guirado, E., Reich, P. B., Ochoa-Hueso, R., Berdugo, M., Sáez-Sandino, T., Blanco-
577 Pastor, J. L., Tedersoo, L., Plaza, C., Ding, J., Sun, W., Mamet, S., Cui, H., He, J.-Z., Hu, H.-W., Sokoya,
578 B., Abades, S., Alfaro, F., Bamigboye, A. R., Bastida, F., de los Ríos, A., Durán, J., Gaitan, J. J., Guerra,
579 C. A., Grebenc, T., Illán, J. G., Liu, Y.-R., Makhalanyane, T. P., Mallen-Cooper, M., Molina-Montenegro,
580 M. A., Moreno, J. L., Nahberger, T. U., Peñaloza-Bojacá, G. F., Picó, S., Rey, A., Rodríguez, A., Siebe,
581 C., Teixido, A. L., Torres-Díaz, C., Trivedi, P., Wang, J., Wang, L., Wang, J., Yang, T., Zaady, E., Zhou,
582 X., Zhou, X.-Q., Zhou, G., Liu, S., and Delgado-Baquerizo, M.: The global contribution of soil mosses
583 to ecosystem services, *Nature Geoscience*, 10.1038/s41561-023-01170-x, 2023.

584 Engel, T., Bruelheide, H., Hoss, D., Sabatini, F. M., Altman, J., Arfin-Khan, M. A. S., Bergmeier, E.,
585 Černý, T., Chytrý, M., Dainese, M., Dengler, J., Dolezal, J., Field, R., Fischer, F. M., Huygens, D., Jandt,
586 U., Jansen, F., Jentsch, A., Karger, D. N., Kattge, J., Lenoir, J., Lens, F., Loos, J., Niinemets, Ü., Overbeck,
587 G. E., Ozinga, W. A., Penuelas, J., Peyre, G., Phillips, O., Reich, P. B., Römermann, C., Sandel, B.,
588 Schmidt, M., Schrod, F., Velez-Martin, E., Violle, C., and Pillar, V.: Traits of dominant plant species
589 drive normalized difference vegetation index in grasslands globally, *Global Ecology and Biogeography*,
590 32, 695-706, 10.1111/geb.13644, 2023.

591 Fang, S., Yu, W., and Qi, Y.: Spectra and vegetation index variations in moss soil crust in different seasons,
592 and in wet and dry conditions, *International Journal of Applied Earth Observation and Geoinformation*,
593 38, 261-266, 10.1016/j.jag.2015.01.018, 2015.

594 Fatichi, S., Pappas, C., Zscheischler, J., and Leuzinger, S.: Modelling carbon sources and sinks in
595 terrestrial vegetation, *New Phytologist*, 221, 652-668, 10.1111/nph.15451, 2019.

596 Ferrenberg, S., Reed, S. C., and Belnap, J.: Climate change and physical disturbance cause similar
597 community shifts in biological soil crusts, *Proceedings of the National Academy of Sciences of the*
598 *United States of America*, 112, 12116-12121, 10.1073/pnas.1509150112, 2015.

599 Fischer, T. and Subbotina, M.: Climatic and soil texture threshold values for cryptogamic cover
600 development: a meta analysis, *Biologia*, 69, 1520-1530, 10.2478/s11756-014-0464-7, 2014.

601 Gabay, T., Rotem, G., Gillor, O., and Ziv, Y.: Understanding changes in biocrust communities following
602 phosphate mining in the Negev Desert, *Environmental Research*, 207, 10.1016/j.envres.2021.112200,
603 2022.

604 Gao, L., Bowker, M. A., Xu, M., Sun, H., Tuo, D., and Zhao, Y.: Biological soil crusts decrease erodibility
605 by modifying inherent soil properties on the Loess Plateau, China, *Soil Biology & Biochemistry*, 105,
606 49-58, 10.1016/j.soilbio.2016.11.009, 2017.

607 Garcia-Pichel, F., Loza, V., Marusenko, Y., Mateo, P., and Potrafka, R. M.: Temperature Drives the
608 Continental-Scale Distribution of Key Microbes in Topsoil Communities, *Science*, 340, 1574-1577,
609 10.1126/science.1236404, 2013.

610 García-Roselló, E., Guisande, C., Manjarrés-Hernández, A., González-Dacosta, J., Heine, J., Pelayo-
611 Villamil, P., González-Vilas, L., Vari, R. P., Vaamonde, A., Granado-Lorencio, C., and Lobo, J. M.: Can
612 we derive macroecological patterns from primary Global Biodiversity Information Facility data?, *Global*
613 *Ecology and Biogeography*, 24, 335-347, 10.1111/geb.12260, 2015.

614 Gassmann, F., Klotzli, F., and Walther, G. R.: Simulation of observed types of dynamics of plants and
615 plant communities, *Journal of Vegetation Science*, 11, 397-408, 10.2307/3236632, 2000.

616 Havrilla, C. A. and Barger, N. N.: Biocrusts and their disturbance mediate the recruitment of native and
617 exotic grasses from a hot desert ecosystem, *Ecosphere*, 9, e02361, 10.1002/ecs2.2361, 2018.

618 Havrilla, C. A., Chaudhary, V. B., Ferrenberg, S., Antoninka, A. J., Belnap, J., Bowker, M. A., Eldridge,

619 D. J., Faist, A. M., Huber-Sannwald, E., Leslie, A. D., Rodriguez-Caballero, E., Zhang, Y., Barger, N.
620 N., and Vries, F.: Towards a predictive framework for biocrust mediation of plant performance: A meta-
621 analysis, *Journal of Ecology*, 107, 2789-2807, 10.1111/1365-2745.13269, 2019.

622 Hu, P., Zhang, W., Xiao, L., Yang, R., Xiao, D., Zhao, J., Wang, W., Chen, H., and Wang, K.: Moss-
623 dominated biological soil crusts modulate soil nitrogen following vegetation restoration in a subtropical
624 karst region, *Geoderma*, 352, 70-79, 10.1016/j.geoderma.2019.05.047, 2019.

625 Issa, O. M., Trichet, J., Defarge, C., Coute, A., and Valentin, C.: Morphology and microstructure of
626 microbiotic soil crusts on a tiger bush sequence (Niger, Sahel), *Catena*, 37, 175-196, 10.1016/S0341-
627 8162(99)00052-1, 1999.

628 Jia, R., Chen, N., Yu, K., and Zhao, C.: High rainfall frequency promotes the dominance of biocrust under
629 low annual rainfall, *Plant and Soil*, 435, 257-275, 10.1007/s11104-018-3880-6, 2019.

630 Jiménez-Valverde, A., Lobo, J. M., and Hortal, J.: Not as good as they seem: the importance of concepts
631 in species distribution modelling, *Diversity and Distributions*, 14, 885-890, 10.1111/j.1472-
632 4642.2008.00496.x, 2008.

633 Karnieli, A.: Development and implementation of spectral crust index over dune sands, *International*
634 *Journal of Remote Sensing*, 18, 1207-1220, 10.1080/014311697218368, 1997.

635 Karnieli, A., Kidron, G. J., Glaesser, C., and Ben-Dor, E.: Spectral characteristics of cyanobacteria soil
636 crust in semiarid environments, *Remote Sensing of Environment*, 69, 67-75, 10.1016/s0034-
637 4257(98)00110-2, 1999.

638 Kidron, G. J.: Biocrust research: A critical view on eight common hydrological-related paradigms and
639 dubious theses, *Ecohydrology*, 12, e2061, 10.1002/eco.2061, 2018.

640 Kidron, G. J.: The enigmatic absence of cyanobacterial biocrusts from the Namib fog belt: Do dew and
641 fog hold the key?, *Flora*, 257, 151416, 10.1016/j.flora.2019.06.002, 2019.

642 Kidron, G. J. and Xiao, B.: A false paradigm? Do biocrust types necessarily reflect 'successional stages'?,
643 *Ecohydrology*, 17, 10.1002/eco.2610, 2023.

644 Kidron, G. J., Lichner, L., Fischer, T., Starinsky, A., and Or, D.: Mechanisms for biocrust-modulated
645 runoff generation – A review, *Earth-Science Reviews*, 231, 104100, 10.1016/j.earscirev.2022.104100,
646 2022.

647 Kropfl, A. I., Distel, R. A., Cecchi, G. A., and Villasuso, N. M.: Functional Role of Moss Biocrust in
648 Disturbed Semiarid Shrublands of North-Eastern Patagonia, Argentina, *Applied Ecology and*
649 *Environmental Research*, 20, 905-917, 10.15666/aeer/2001_905917, 2022.

650 Lenton, T. M., Dahl, T. W., Daines, S. J., Mills, B. J., Ozaki, K., Saltzman, M. R., and Porada, P.: Earliest
651 land plants created modern levels of atmospheric oxygen, *Proceedings of the National Academy of*
652 *Sciences of the United States of America*, 113, 9704-9709, 10.1073/pnas.1604787113, 2016.

653 Li, S., Bowker, M. A., and Xiao, B.: Biocrusts enhance non-rainfall water deposition and alter its
654 distribution in dryland soils, *Journal of Hydrology*, 595, 126050, 10.1016/j.jhydrol.2021.126050, 2021.

655 Li, X., Sun, J., Zhang, H., Tan, H., Hui, R., Qi, J., Zhang, P., and Ward, N. D.: Warming decreases desert
656 ecosystem functioning by altering biocrusts in drylands, *Journal of Applied Ecology*, 10.1111/1365-
657 2664.14528, 2023.

658 Ma, X., Zhao, Y., Yang, K., Ming, J., Qiao, Y., Xu, M., and Pan, X.: Long-term light grazing does not
659 change soil organic carbon stability and stock in biocrust layer in the hilly regions of drylands, *Journal*
660 *of Arid Land*, 15, 940-959, 10.1007/s40333-023-0064-x, 2023.

661 Machado de Lima, N. M., Muñoz-Rojas, M., Vázquez-Campos, X., and Branco, L. H. Z.: Biocrust
662 cyanobacterial composition, diversity, and environmental drivers in two contrasting climatic regions in

663 Brazil, *Geoderma*, 386, 114914, 10.1016/j.geoderma.2020.114914, 2021.

664 Maestre, F. T., Benito, B. M., Berdugo, M., Concostrina-Zubiri, L., Delgado-Baquerizo, M., Eldridge, D.
665 J., Guirado, E., Gross, N., Kefi, S., Le Bagousse-Pinguet, Y., Ochoa-Hueso, R., and Soliveres, S.:
666 Biogeography of global drylands, *New Phytologist*, 231, 540-558, 10.1111/nph.17395, 2021.

667 Mäkinen, J., Numminen, E., Niittynen, P., Luoto, M., and Vanhatalo, J.: Spatial confounding in Bayesian
668 species distribution modeling, *Ecography*, 2022, e06183, 10.1111/ecog.06183, 2022.

669 Marsh, J., Nouvet, S., Sanborn, P., and Coxson, D.: Composition and function of biological soil crust
670 communities along topographic gradients in grasslands of central interior British Columbia (Chilcotin)
671 and southwestern Yukon (Kluane), *Canadian Journal of Botany*, 84, 717-736, 10.1139/b06-026, 2006.

672 McCann, E., Reed, S. C., Saud, P., Reibold, R. H., Howell, A., and Faist, A. M.: Plant growth and
673 biocrust-fire interactions across five North American deserts, *Geoderma*, 401, 115325,
674 10.1016/j.geoderma.2021.115325, 2021.

675 McCune, B., Yang, S., Jovan, S., and Root, H. T.: Climate and epiphytic macrolichen communities in the
676 Four Corners region of the USA, *Bryologist*, 125, 70-90, 10.1639/0007-2745-125.1.070, 2022.

677 Miller, D. M., Bedford, D. R., Hughson, D. L., McDonald, E. V., Robinson, S. E., and Schmidt, K. M.:
678 Mapping Mojave Desert Ecosystem Properties with Surficial Geology, 3rd Mojave Desert Science
679 Symposium, Univ Redlands, Redlands, CA, Nov 16-18, WOS:000266552000012, 225-251, 2009.

680 Ming, J., Zhao, Y., Wu, Q., He, H., and Gao, L.: Soil temperature dynamics and freezing processes for
681 biocrustal soils in frozen soil regions on the Qinghai-Tibet Plateau, *Geoderma*, 409, 115655,
682 10.1016/j.geoderma.2021.115655, 2022.

683 Miranda-González, R. and McCune, B.: The weight of the crust: Biomass of crustose lichens in tropical
684 dry forest represents more than half of foliar biomass, *Biotropica*, 52, 1298-1308, 10.1111/btp.12837,
685 2020.

686 Morillas, L. and Gallardo, A.: Biological soil crusts and wetting events: Effects on soil N and C cycles,
687 *Applied Soil Ecology*, 94, 1-6, 10.1016/j.apsoil.2015.04.015, 2015.

688 Munoz-Martin, M. A., Becerra-Absalon, I., Perona, E., Fernandez-Valbuena, L., Garcia-Pichel, F., and
689 Mateo, P.: Cyanobacterial biocrust diversity in Mediterranean ecosystems along a latitudinal and climatic
690 gradient, *New Phytologist*, 221, 123-141, 10.1111/nph.15355, 2019.

691 Noy, K., Ohana-Levi, N., Panov, N., Silver, M., and Karnieli, A.: A long-term spatiotemporal analysis of
692 biocrusts across a diverse arid environment: The case of the Israeli-Egyptian sandfield, *Science of The
693 Total Environment*, 774, 145154, 10.1016/j.scitotenv.2021.145154, 2021.

694 Oliva, G., Dos Santos, E., Sofia, O., Umana, F., Massara, V., Garcia Martinez, G., Caruso, C., Cariac, G.,
695 Echevarria, D., Fantozzi, A., Butti, L., Bran, D., Gaitan, J., Ferrante, D., Paredes, P., Dominguez, E., and
696 Maestre, F. T.: The MARAS dataset, vegetation and soil characteristics of dryland rangelands across
697 Patagonia, *Scientific Data*, 7, 327, 10.1038/s41597-020-00658-0, 2020.

698 Palmer, B., Hernandez, R., and Lipson, D. A.: The fate of biological soil crusts after fire: A meta-analysis,
699 *Global Ecology and Conservation*, 24, e01380, 10.1016/j.gecco.2020.e01380, 2020.

700 Palmer, B., Lawson, D., and Lipson, D. A.: Years After a Fire, Biocrust Microbial Communities are
701 Similar to Unburned Communities in a Coastal Grassland, *Microbial Ecology*, 10.1007/s00248-022-
702 02137-y, 2022.

703 Panigada, C., Tagliabue, G., Zaady, E., Rozenstein, O., Garzonio, R., Di Mauro, B., De Amicis, M.,
704 Colombo, R., Cogliati, S., Miglietta, F., and Rossini, M.: A new approach for biocrust and vegetation
705 monitoring in drylands using multi-temporal Sentinel-2 images, *Progress in Physical Geography: Earth
706 and Environment*, 43, 496-520, 10.1177/0309133319841903, 2019.

707 Pearce, J. L., Cherry, K., M, D., S, F., and Wish, G.: Incorporating expert opinion and fine-scale
708 vegetation mapping into statistical models of faunal distribution, *Journal of Applied Ecology*, 38, 412-
709 424, 10.1046/j.1365-2664.2001.00608.x, 2001.

710 Perry, G. L. W., Seidl, R., Bellvé, A. M., and Rammer, W.: An Outlook for Deep Learning in Ecosystem
711 Science, *Ecosystems*, 25, 1700-1718, 10.1007/s10021-022-00789-y, 2022.

712 Pietrasiak, N., Drenovsky, R. E., Santiago, L. S., and Graham, R. C.: Biogeomorphology of a Mojave
713 Desert landscape — Configurations and feedbacks of abiotic and biotic land surfaces during landform
714 evolution, *Geomorphology*, 206, 23-36, 10.1016/j.geomorph.2013.09.015, 2014.

715 Porada, P., Pöschl, U., Kleidon, A., Beer, C., and Weber, B.: Estimating global nitrous oxide emissions
716 by lichens and bryophytes with a process-based productivity model, *Biogeosciences*, 14, 1593-1602,
717 10.5194/bg-14-1593-2017, 2017.

718 Porada, P., Weber, B., Elbert, W., Pöschl, U., and Kleidon, A.: Estimating global carbon uptake by lichens
719 and bryophytes with a process-based model, *Biogeosciences*, 10, 6989-7033, 10.5194/bg-10-6989-2013,
720 2013.

721 Porada, P., Tamm, A., Raggio, J., Cheng, Y., Kleidon, A., Pöschl, U., and Weber, B.: Global NO and
722 HONO emissions of biological soil crusts estimated by a process-based non-vascular vegetation model,
723 *Biogeosciences*, 16, 2003-2031, 10.5194/bg-16-2003-2019, 2019.

724 Pravalie, R.: Drylands extent and environmental issues. A global approach, *Earth-Science Reviews*, 161,
725 259-278, 10.1016/j.earscirev.2016.08.003, 2016.

726 Pushkareva, E., Johansen, J. R., and Elster, J.: A review of the ecology, ecophysiology and biodiversity
727 of microalgae in Arctic soil crusts, *Polar Biology*, 39, 2227-2240, 10.1007/s00300-016-1902-5, 2016.

728 Qiu, D., Bowker, M. A., Xiao, B., Zhao, Y., Zhou, X., and Li, X.: Mapping biocrust distribution in China's
729 drylands under changing climate, *Science of The Total Environment*, 905,
730 10.1016/j.scitotenv.2023.167211, 2023.

731 Quillet, A., Peng, C., and Garneau, M.: Toward dynamic global vegetation models for simulating
732 vegetation-climate interactions and feedbacks: recent developments, limitations, and future challenges,
733 *Environmental Reviews*, 18, 333-353, 10.1139/a10-016, 2010.

734 Read, C. F., Duncan, D. H., Vesk, P. A., Elith, J., and Wan, S.: Biocrust morphogroups provide an
735 effective and rapid assessment tool for drylands, *Journal of Applied Ecology*, 51, 1740-1749,
736 10.1111/1365-2664.12336, 2014.

737 Reed, S. C., Coe, K. K., Sparks, J. P., Housman, D. C., Zelikova, T. J., and Belnap, J.: Changes to dryland
738 rainfall result in rapid moss mortality and altered soil fertility, *Nature Climate Change*, 2, 752-755,
739 10.1038/nclimate1596, 2012.

740 Reynolds, R., Neff, J., Reheis, M., and Lamothe, P.: Atmospheric dust in modern soil on aeolian
741 sandstone, Colorado Plateau (USA): Variation with landscape position and contribution to potential plant
742 nutrients, *Geoderma*, 130, 108-123, 10.1016/j.geoderma.2005.01.012, 2006.

743 Richer, R., Anchassi, D., El-Assaad, I., El-Matbouly, M., Ali, F., Makki, I., and Metcalf, J. S.: Variation
744 in the coverage of biological soil crusts in the State of Qatar, *Journal of Arid Environments*, 78, 187-190,
745 10.1016/j.jaridenv.2011.10.009, 2012.

746 Rodriguez-Caballero, E., Knerr, T., and Weber, B.: Importance of biocrusts in dryland monitoring using
747 spectral indices, *Remote Sensing of Environment*, 170, 32-39, 10.1016/j.rse.2015.08.034, 2015.

748 Rodriguez-Caballero, E., Belnap, J., Budel, B., Crutzen, P. J., Andreae, M. O., Pöschl, U., and Weber, B.:
749 Dryland photoautotrophic soil surface communities endangered by global change, *Nature Geoscience*,
750 11, 185-189, 10.1038/s41561-018-0072-1, 2018.

751 Rodríguez-Caballero, E., Stanelle, T., Egerer, S., Cheng, Y., Su, H., Canton, Y., Belnap, J., Andreae, M.
752 O., Tegen, I., and Reick, C. H.: Global cycling and climate effects of aeolian dust controlled by biological
753 soil crusts, *Nature Geoscience*, 15, 458-463, 10.1038/s41561-022-00942-1, 2022.

754 Rodríguez-Caballero, E., Escribano, P., Olehowski, C., Chamizo, S., Hill, J., Cantón, Y., and Weber, B.:
755 Transferability of multi- and hyperspectral optical biocrust indices, *ISPRS Journal of Photogrammetry
756 and Remote Sensing*, 126, 94-107, 10.1016/j.isprsjprs.2017.02.007, 2017.

757 Rodríguez-Caballero, E., Reyes, A., Kratz, A., Caesar, J., Guirado, E., Schmiedel, U., Escribano, P.,
758 Fiedler, S., and Weber, B.: Effects of climate change and land use intensification on regional biological
759 soil crust cover and composition in southern Africa, *Geoderma*, 406, 115508,
760 10.1016/j.geoderma.2021.115508, 2022.

761 Root, H. T. and McCune, B.: Regional patterns of biological soil crust lichen species composition related
762 to vegetation, soils, and climate in Oregon, USA, *Journal of Arid Environments*, 79, 93-100,
763 10.1016/j.jaridenv.2011.11.017, 2012.

764 Sabatini, F. M., Lenoir, J., Hattab, T., Arnst, E. A., Chytrý, M., Dengler, J., De Ruffray, P., Hennekens, S.
765 M., Jandt, U., Jansen, F., Jiménez-Alfaro, B., Kattge, J., Levesley, A., Pillar, V. D., Purschke, O., Sandel,
766 B., Sultana, F., Aavik, T., Ačić, S., Acosta, A. T. R., Agrillo, E., Alvarez, M., Apostolova, I., Arfin Khan,
767 M. A. S., Arroyo, L., Attorre, F., Aubin, I., Banerjee, A., Bauters, M., Bergeron, Y., Bergmeier, E., Biurrun,
768 I., Bjorkman, A. D., Bonari, G., Bondareva, V., Brunet, J., Čarni, A., Casella, L., Cayuela, L., Černý, T.,
769 Chepinoga, V., Csiky, J., Čušterevska, R., De Bie, E., Gasper, A. L., De Sanctis, M., Dimopoulos, P.,
770 Dolezal, J., Dziuba, T., El-Sheikh, M. A. E. R. M., Enquist, B., Ewald, J., Fazayeli, F., Field, R., Finckh,
771 M., Gachet, S., Galán-de-Mera, A., Garbolino, E., Gholizadeh, H., Giorgis, M., Golub, V., Alsos, I. G.,
772 Grytnes, J. A., Guerin, G. R., Gutiérrez, A. G., Haider, S., Hatim, M. Z., Hérault, B., Hinojos Mendoza,
773 G., Hölzel, N., Homeier, J., Hubau, W., Indreica, A., Janssen, J. A. M., Jedrzejek, B., Jentsch, A., Jürgens,
774 N., Kaçki, Z., Kapfer, J., Karger, D. N., Kavgacı, A., Kearsley, E., Kessler, M., Khanina, L., Killeen, T.,
775 Korolyuk, A., Kreft, H., Kühl, H. S., Kuzemko, A., Landucci, F., Lengyel, A., Lens, F., Lingner, D. V.,
776 Liu, H., Lysenko, T., Mahecha, M. D., Marcenò, C., Martynenko, V., Moeslund, J. E., Monteagudo
777 Mendoza, A., Mucina, L., Müller, J. V., Munzinger, J., Naqinezhad, A., Noroozi, J., Nowak, A.,
778 Onyshchenko, V., Overbeck, G. E., Pärtel, M., Pauchard, A., Peet, R. K., Peñuelas, J., Pérez-Haase, A.,
779 Peterka, T., Petřík, P., Peyre, G., Phillips, O. L., Prokhorov, V., Rašomavičius, V., Revermann, R., Rivas-
780 Torres, G., Rodwell, J. S., Ruprecht, E., Rüşiņa, S., Samimi, C., Schmidt, M., Schrodte, F., Shan, H.,
781 Shirokikh, P., Šibík, J., Šilc, U., Sklenář, P., Škvorc, Ž., Sparrow, B., Sperandii, M. G., Stančić, Z.,
782 Svenning, J. C., Tang, Z., Tang, C. Q., Tsiropidis, I., Vanselow, K. A., Vásquez Martínez, R., Vassilev, K.,
783 Véllez-Martin, E., Venanzoni, R., Vibrans, A. C., Violle, C., Virtanen, R., Wehrden, H., Wagner, V.,
784 Walker, D. A., Waller, D. M., Wang, H. F., Wesche, K., Whitfeld, T. J. S., Willner, W., Wiser, S. K.,
785 Wohlgemuth, T., Yamalov, S., Zobel, M., Bruelheide, H., and Bates, A.: sPlotOpen – An
786 environmentally balanced, open - access, global dataset of vegetation plots, *Global Ecology and
787 Biogeography*, 30, 1740-1764, 10.1111/geb.13346, 2021.

788 Shi, W., Pan, Y.-x., Zhang, Y.-f., Hu, R., and Wang, X.-p.: The effect of different biocrusts on soil
789 hydraulic properties in the Tengger Desert, China, *Geoderma*, 430, 116304,
790 10.1016/j.geoderma.2022.116304, 2023.

791 Skidmore, A. K., Franklin, J., Dawson, T. P., and Pilesjö, P.: Geospatial tools address emerging issues in
792 spatial ecology: a review and commentary on the Special Issue, *International Journal of Geographical
793 Information Science*, 25, 337-365, 10.1080/13658816.2011.554296, 2011.

794 Soberon, J. and Nakamura, M.: Niches and distributional areas: Concepts, methods, and assumptions,

795 Proceedings of the National Academy of Sciences of the United States of America, 106, 19644-19650,
796 2009.

797 Song, G., Hui, R., Yang, H., Wang, B., and Li, X.: Biocrusts mediate the plant community composition
798 of dryland restoration ecosystems, *Science of the Total Environment*, 844, 157135,
799 10.1016/j.scitotenv.2022.157135, 2022.

800 Steven, B., Gallegos-Graves, L. V., Belnap, J., and Kuske, C. R.: Dryland soil microbial communities
801 display spatial biogeographic patterns associated with soil depth and soil parent material, *FEMS*
802 *Microbiology Ecology*, 86, 101-113, 10.1111/1574-6941.12143, 2013.

803 Sun, F., Xiao, B., Kidron, G. J., and Tuller, M.: Towards the effects of moss-dominated biocrusts on
804 surface soil aeration in drylands: Air permeability analysis and modeling, *Catena*, 223, 106942,
805 10.1016/j.catena.2023.106942, 2023.

806 Thonicke, K., Venevsky, S., Sitch, S., and Cramer, W.: The role of fire disturbance for global vegetation
807 dynamics: coupling fire into a Dynamic Global Vegetation Model, *Global Ecology and Biogeography*,
808 10, 661-677, 10.1046/j.1466-822x.2001.00175.x, 2001.

809 Tucker, C. L., McHugh, T. A., Howell, A., Gill, R., Weber, B., Belnap, J., Grote, E., and Reed, S. C.: The
810 concurrent use of novel soil surface microclimate measurements to evaluate CO₂ pulses in biocrusted
811 interspaces in a cool desert ecosystem, *Biogeochemistry*, 135, 239-249, 10.1007/s10533-017-0372-3,
812 2017.

813 Velasco Ayuso, S., Oñatibia, G. R., Maestre, F. T., and Yahdjian, L.: Grazing pressure interacts with
814 aridity to determine the development and diversity of biological soil crusts in Patagonian rangelands,
815 *Land Degradation & Development*, 31, 488-499, 10.1002/ldr.3465, 2019.

816 Wang, S., Liu, B., Zhao, Y., Gao, L., Yin, B., Yang, K., and Ji, J.: Determination of the representative
817 elementary area (REA) of biocrusts: A case study from the Hilly Loess Plateau region, China, *Geoderma*,
818 406, 115502, 10.1016/j.geoderma.2021.115502, 2022a.

819 Wang, Z., Wu, B., Zhang, M., Zeng, H., Yang, L., Tian, F., Ma, Z., and Wu, H.: Indices enhance biological
820 soil crust mapping in sandy and desert lands, *Remote Sensing of Environment*, 278, 113078,
821 10.1016/j.rse.2022.113078, 2022b.

822 Weber, B. and Hill, J.: Remote Sensing of Biological Soil Crusts at Different Scales, in: *Biological Soil*
823 *Crusts: An Organizing Principle in Drylands*, edited by: Weber, B., Büdel, B., and Belnap, J., Springer
824 International Publishing, Cham, 215-234, 10.1007/978-3-319-30214-0_12, 2016.

825 Weber, B., Budel, B., and Belnap, J., Weber, B., Budel, B., and Belnap, J. (Eds.): *Biological Soil Crusts:*
826 *An Organizing Principle in Drylands*, Springer Nature, 10.1007/978-3-319-30214-0, 2016.

827 Weber, B., Olehowski, C., Knerr, T., Hill, J., Deutschewitz, K., Wessels, D. C. J., Eitel, B., and Buedel,
828 B.: A new approach for mapping of Biological Soil Crusts in semidesert areas with hyperspectral imagery,
829 *Remote Sensing of Environment*, 112, 2187-2201, 10.1016/j.rse.2007.09.014, 2008.

830 Weber, B., Belnap, J., Budel, B., Antoninka, A. J., Barger, N. N., Chaudhary, V. B., Darrouzet-Nardi, A.,
831 Eldridge, D. J., Faist, A. M., Ferrenberg, S., Havrilla, C. A., Huber-Sannwald, E., Malam Issa, O.,
832 Maestre, F. T., Reed, S. C., Rodriguez-Caballero, E., Tucker, C., Young, K. E., Zhang, Y., Zhao, Y., Zhou,
833 X., and Bowker, M. A.: What is a biocrust? A refined, contemporary definition for a broadening research
834 community, *Biological Reviews*, 97, 1768-1785, 10.1111/brv.12862, 2022.

835 Wei, X., Qin, F., Han, B., Zhou, H., Liu, M., and Shao, X.: Spatial variations of bacterial communities
836 associated with biological soil crusts along a climatic gradient in alpine grassland ecosystems, *Plant and*
837 *Soil*, 480, 493-506, 10.1007/s11104-022-05595-y, 2022.

838 Whitney, K. M., Vivoni, E. R., Duniway, M. C., Bradford, J. B., Reed, S. C., and Belnap, J.:

839 Ecohydrological role of biological soil crusts across a gradient in levels of development, *Ecohydrology*,
840 10, e1875, 10.1002/eco.1875, 2017.

841 Williams, A. J., Buck, B. J., Soukup, D. A., and Merkler, D. J.: Geomorphic controls on biological soil
842 crust distribution: A conceptual model from the Mojave Desert (USA), *Geomorphology*, 195, 99-109,
843 10.1016/j.geomorph.2013.04.031, 2013.

844 Wolf, S., Mahecha, M. D., Sabatini, F. M., Wirth, C., Bruelheide, H., Kattge, J., Moreno Martinez, A.,
845 Mora, K., and Kattenborn, T.: Citizen science plant observations encode global trait patterns, *Nature*
846 *Ecology & Evolution*, 6, 1850-1859, 10.1038/s41559-022-01904-x, 2022.

847 Wright, A. J. and Collins, S. L.: Drought experiments need to incorporate atmospheric drying to better
848 simulate climate change, *BioScience*, 74, 65-71, 10.1093/biosci/biad105, 2024.

849 Yang, H., Liu, C., Liu, Y., and Xing, Z.: Impact of human trampling on biological soil crusts determined
850 by soil microbial biomass, enzyme activities and nematode communities in a desert ecosystem, *European*
851 *Journal of Soil Biology*, 87, 61-71, 10.1016/j.ejsobi.2018.05.005, 2018.

852 Yu, H., Cooper, A. R., and Infante, D. M.: Improving species distribution model predictive accuracy
853 using species abundance: Application with boosted regression trees, *Ecological Modelling*, 432, 109202,
854 10.1016/j.ecolmodel.2020.109202, 2020.

855 Zhang, Y. M., Chen, J., Wang, L., Wang, X. Q., and Gu, Z. H.: The spatial distribution patterns of
856 biological soil crusts in the Gurbantunggut Desert, Northern Xinjiang, China, *Journal of Arid*
857 *Environments*, 68, 599-610, 10.1016/j.jaridenv.2006.06.012, 2007.

858 Zhao, Y., Qin, N., Weber, B., and Xu, M.: Response of biological soil crusts to raindrop erosivity and
859 underlying influences in the hilly Loess Plateau region, China, *Biodiversity and Conservation*, 23, 1669-
860 1686, 10.1007/s10531-014-0680-z, 2014.

861