

1 **Running title:** Global biocrusts distribution

2 **Advancing studies on global biocrusts distribution**

3 Siqing Wang<sup>1,2</sup>, Li Ma<sup>1,2</sup>, Liping Yang<sup>1,2</sup>, Yali Ma<sup>1,2</sup>, Yafeng Zhang<sup>3</sup>, Changming Zhao<sup>1,2</sup>, Ning  
4 Chen<sup>1,2,4\*</sup>

5 <sup>1</sup> State Key Laboratory of Herbage Improvement and Grassland Agro-ecosystems, College of  
6 Ecology, Lanzhou University, Lanzhou, Gansu 730000, China

7 <sup>2</sup> Yuzhong Mountain Ecosystems Observation and Research Station, Lanzhou University,  
8 Lanzhou, Gansu 730000, China

9 <sup>3</sup> Shapotou Desert Research and Environment Station, Northwest Institute of Eco-Environment  
10 and Resources, Chinese Academy of Sciences, No.320, Donggang West Road,  
11 Lanzhou, Gansu 730000, China

12 <sup>4</sup> Instituto Multidisciplinar para el Estudio del Medio “Ramon Margalef”, Universidad de  
13 Alicante, Carretera de San Vicente del Raspeig s/n, San Vicente del Raspeig Alicante 03690,  
14 Spain

15 \*Correspondence to Ning Chen, Email: cn@lzu.edu.cn. Tel.: +86-931-8912551. ORCID: 0000-  
16 0002-1779-915X.

17

18 **Abstract:** Biological soil crusts (biocrusts hereafter) cover a substantial proportion of the  
19 dryland ecosystem and play crucial roles in ecological processes such as biogeochemical cycles,  
20 water distribution, and soil erosion. Consequently, studying the spatial distribution of biocrusts

21 holds great significance for drylands, especially on a global scale, but it still needs to be  
22 improved. This study aimed to stimulate global-scale investigations of biocrust distribution by

23 introducing three major approaches: spectral characterization indices, dynamic vegetation  
24 models, and geospatial models, while discussing their applicability. We then summarized the

25 present understanding of the factors influencing biocrust distribution. Finally, to further  
26 advance this field, we proposed several potential research topics and directions, including the

27 development of a standardized biocrust database, enhancement of non-vascular vegetation  
28 dynamic models, integration of multi-sensor monitoring, extensive use of machine learning,

29 and a focus on regional research co-development. This work is supposed to significantly  
30 contribute to mapping the biocrust distribution and thereby advance our understanding of

31 dryland ecosystem management and restoration.

32 **Key words:** biological soil crusts; distribution; drylands; global scales; regional scales

33

34 **1. Introduction**

删除了: , which is still lacking

删除了: remains limited

删除了: ,

删除了: Then, w

删除了: biocrusts distribution

删除了: aspect

删除了: including building a standardized database of biocrusts, enhancing non-vascular vegetation dynamic models, integrating multi-sensor monitoring, making full use of machine learning, and focusing on regional research co-development....

删除了: biocrusts

47 Biological soil crusts (biocrusts hereafter) are continuous biotic complexes that live in the  
48 topsoil, which are formed by different proportions of photosynthetic autotrophic (e.g.  
49 cyanobacteria, algae, lichens, mosses) and heterotrophic (e.g. bacteria, fungi, archaea)  
50 organisms colloidal with soil particles, usually with a thickness of a few millimeters to a few  
51 centimeters (Weber et al., 2022). Biocrusts occupy a wide range of ecological niches in mid  
52 latitudes, polar and alpine regions, covering approximately 11% of the global land area (Porada  
53 et al., 2019). In particular, biocrusts are well-adapted to water-limited, nutrient-poor, and hostile  
54 environments, such as arid and semi-arid areas characterized by low ratios of precipitation to  
55 potential evaporation (0.05-0.5 mm mm<sup>-1</sup>) (Pravalie, 2016; Read et al., 2014; Weber et al., 2016).

56 As vital components of dryland ecosystems, biocrusts fulfill many essential ecological  
57 functions. They contribute to stabilizing the soil surface, improving soil permeability, and  
58 enhancing water-holding capacity within the upper few centimeters of soil (Sun et al., 2023;  
59 Shi et al., 2023; Gao et al., 2017). By participating in various biogeochemical cycles, biocrusts  
60 were estimated to contribute to 15% of terrestrial net primary productivity and 40-85% of  
61 biological nitrogen fixation (Elbert et al., 2012; Rodriguez-Caballero et al., 2018). They also  
62 impact ecohydrological processes by altering soil microclimate and redistributing soil water  
63 (Kidron et al., 2022; Tucker et al., 2017). Moreover, biocrusts influence seed capture and soil  
64 seed banks (Kropfl et al., 2022), thereby mediating plant growth and community assembly  
65 (Havrilla and Barger, 2018; Song et al., 2022). The extent and magnitude of these ecological  
66 functions and services depend on the spatial distribution of biocrusts. Therefore, it is crucial to  
67 understand their distribution.

68 Despite the significance of biocrusts, previous studies have primarily focused on their  
69 contributions to carbon and nitrogen cycling across various habitats and climates (Hu et al.,  
70 2019; Morillas and Gallardo, 2015), as well as interspecific interactions and biocrust  
71 biodiversity (Machado De Lima et al., 2021; Munoz-Martin et al., 2019), rather than their  
72 spatial distribution. A group of ecologists, including Fernando Maestre (Maestre et al., 2021),  
73 David Eldridge (Eldridge and Delgado-Baquerizo, 2019; Eldridge et al., 2023), Matthew  
74 Bowker (Qiu et al., 2023), Emilio Rodríguez-Caballero (Rodríguez-Caballero et al., 2018) and  
75 others, have actively promoted progress in the field (Fig. 1a). Countries with extensive dryland

删除了: They are able to

删除了: can be

删除了:

删除了: and

删除了: a suite of

删除了: C

删除了: N

删除了: in

删除了: varying

删除了: s

移动了(插入) [1]

删除了: , particularly at the global scale

删除了:

删除了: Since 1990, studies on the distribution of biocrusts have been continuously increasing. A

删除了: represented by

域代码已更改

域代码已更改

域代码已更改

删除了: 2(


删除了:)

删除了: . The topic has gradually received attention from all over the world, particularly the

删除了: c

96 areas, such as China, the United States, Spain, the United Kingdom, Germany, Australia, and  
 97 Israel, have attempted to make breakthroughs on this issue. (Fig. 1b). However, other dryland  
 98 countries and regions, such as central and southern Africa, where the biocrust distribution has  
 99 been reported, still suffer from a paucity of studies and data on biocrusts (Fig. 1c). This  
 100 geographical imbalance in biocrust distribution studies has resulted in most knowledge  
 101 remaining at local to regional scales, with very limited discoveries on a global scale.

- 删除了: 2(
- 删除了: )
- 删除了: some
- 删除了: keep
- 删除了: the amount of
- 删除了: (
- 删除了: )
- 删除了: is far from adequate (Fig. 2(c))
- 删除了: global-scale
- 删除了: These areas may be potential areas of widespread distribution of biocrusts in the future.



102

114 Fig. 1 Literature review of biocrust distribution studies. (a) Representative authors associated  
 115 frameworks for biocrusts distribution studies (1990 to 2022). The time series is the average  
 116 time of the year of publication, e.g., if the number of articles is 2 in 2004 and 8 in 2019, the  
 117 node in this figure shows the year as  $(2004 \times 2 + 2019 \times 8)/10 = 2016$ . (b) Map of hotspot  
 118 countries for biocrust distribution research, with the top 12 countries in terms of number of  
 119 publications shown: The database is Web of Science, TS = ("biogenic crust\*" OR "biological  
 120 crust\*" OR "biological soil crust\*" OR "biocrust\*" OR "microphytic crust\*" OR "microbiotic  
 121 crust\*" OR "cyanobacterial\*" OR "algal\*" OR "lichen\*" OR "moss\*" OR "biotic crust\*") AND  
 122 ("mapping\*" OR "distribution\*" OR "spatial pattern\*") AND ("dryland" OR "hyper\*arid\*" OR  
 123 "arid\*" OR "semi\*arid\*" OR "dry subhumid\*"), with research interests in Environmental  
 124 Sciences/Ecology and a total of 700 papers. (c) Global biocrust data distribution, based on field  
 125 surveys and literature compilation. Data have been collected and expanded from the published  
 126 database (Chen et al., 2020; Rodriguez-Caballero et al., 2018) to 3848 items.

127 In this study, we aimed to sort out and advance the understanding of biocrust distribution  
 128 from three perspectives: the applicability and comparison of research methods (section 2),  
 129 clarification of factors influencing biocrust distribution (section 3), and challenges and  
 130 strategies for future studies on biocrust distribution (section 4). This work is expected to deepen  
 131 our understanding of dryland ecosystem processes and provide a scientific basis for conserving  
 132 dryland ecosystems, and their responses to global change.

## 133 2. Research Methods

134 Three methods are commonly used to study biocrust distribution: spectral characterization,  
 135 vegetation dynamic modeling, and geospatial modeling. This section provides an overview of  
 136 these methods, including their basic principles, case studies, adaptability, and limitations.

### 137 2.1 Spectral characterization index

138 With advances in remote sensing and geo-information technology, spectroscopy offers a  
 139 feasible method of characterizing distribution features from a physical point of view.  
 140 Differences in absorption or reflection of specific wavelengths by different ground covers can  
 141 effectively identify soil surface objects (Rodriguez-Caballero et al., 2015). By identifying  
 142 biocrust-specific bands from reflectance spectral images (Karnieli et al., 1999), it is possible to

删除了: 2

域代码已更改

#### 删除了: 3.1 Local-regional scales

At local-regional scales, numerous studies have provided valuable insights into the distribution patterns of biocrusts in different regions around the world (Fig. 3). In the Mojave Desert, biocrusts distribution was closely related to geological age, surface stability, topography, and dust transport (Miller et al., 2004). Lichen, moss, and dark algal crusts patchily distributed on the desert, averaging 8% cover, though in some bar and shrub zones, the cover could be as high as 26% (Pietrasiak et al., 2014). In the Colorado Plateau, highly heterogeneous soil matrix determined the fragmented biocrusts distribution of different types and the wide disparity in relative abundance and cover (Reynolds et al., 2006; Steven et al., 2013). Collier et al. (2022) trained drone imagery in the Hawaiian region using timely data collected by cameras and then successfully mapped watershed-scale biocrusts distribution, predicting cover of ~15-23%. In the Gurbantunggut Desert, biocrusts cover 28.7% of the area, with a high and uniform biocrusts cover in the southern part of the desert and a scattered distribution of biocrusts in other areas (Chen et al., 2005; Zhang et al., 2007). In the Loess Plateau, RGB image-based biocrusts monitoring showed that variability in biocrusts cover decreased logarithmically with increasing plot size until a critical size of 1m<sup>2</sup> after which biocrusts cover remained approximately constant (Wang et al., 2022a). In Qatar, 26% of the country is covered by biocrusts, with cyanobacterial biocrusts cover showing a decreasing trend from north-east to south-west (Richer et al., 2022).

删除了: In this study, we firstly sorted out the main research methods for studying biocrusts distribution (section 1), then reviewed the existing knowledge (section 2), and finally

删除了: to

删除了: the conservation of

删除了: conservation

带格式的: 缩进: 首行缩进: 2 字符

删除了:

删除了: provide

删除了: of

删除了: the

241 construct a presence-absence map of biocrust distribution (Fig. 2a),

242 Currently, spectral characterization indices have been widely applied in many areas of

243 drylands. For example, cyanobacterial biocrusts are widely distributed in the Sahara region of

244 Africa (Beaugendre et al., 2017) and the Negev Desert of Israel(Panigada et al., 2019), where

245 the study inverted the Biocrust Index (CI) based on remotely sensed imagery to access the

246 characteristics of localized changes in biocrust distribution over 31 years (Karnieli, 1997; Noy

247 et al., 2021). Sun et al. (2024) developed the fraction biocrust cover index (FBCI) based on

248 radiative transfer and mapped biocrust distribution over a desert area at 10 m resolution,

249 showing well-matched results between the model and field observations (RMSE of 0.0774,

250 systematic deviation of -4.05%). In the Gurbantunggut Desert, a study constructed the

251 Biological Soil Crust Index (BSCI) with lichen biocrust as the dominant group and mapped the

252 distribution of biocrusts with high accuracy (accuracy of 94.7%, kappa coefficient of 0.82)

253 (Chen et al., 2005), spatially, biocrusts cover 28.7% of the area, with a high and uniform cover

254 in the southern part of the desert and a scattered distribution in other regions (Zhang et al.,

255 2007). In the Loess Plateau, RGB image-based biocrust monitoring showed that variability in

256 biocrusts cover decreased logarithmically with increasing plot size until a critical size of 1m<sup>2</sup>,

257 after which biocrusts cover remained approximately constant (Wang et al., 2022a).

258 For the spectral characterization method, it is critical to determine the threshold of spectral

259 bands that represent biocrusts. For instance, at an aerosol optical depth of 0.2, the BSCI ranges

260 from 4.13 to 6.23, and narrows to 4.58-5.69 with increasingly poor atmospheric conditions.

261 Overly strict or loose threshold ranges can easily lead to biocrust omission or misidentification.

262 To improve the accuracy of biocrust identification, some researchers have utilized the

263 hyperspectral sensor's continuous waveband capabilities and created the Continuum Removal

264 Crust Identification Algorithm (CRCIA) (Chamizo et al., 2012b; Weber et al., 2008). Baxter et

265 al. (2021) innovatively applied the random forest algorithm to spectral feature classification,

266 achieving an accuracy of 78.5% in biocrusts recognition. Additionally, two other indices, the

267 Sandy Land Ratio Crust Index (SRCI) and the Desert Ratio Crust Index (DRCI), were

268 introduced to account for differences between sandy land (vegetation cover FVC <20%) and

269 desert environments, improving mapping accuracy by approximately 6% (Wang et al., 2022b).

- 删除了: biocrusts
- 删除了: (Fig. 1a)
- 删除了: a
- 删除了: areas
- 删除了: The crust index (CI) and biological soil crust index (BSCI) are two of spectral characterization indexes and hav...
- 删除了: such
- 删除了: indicators
- 删除了: ,
- 删除了: s
- 删除了: In order t
- 删除了: took advantage of
- 删除了: a
- 删除了: and
- 删除了: ed
- 删除了: nother two indexes
- 删除了: i.e.
- 删除了: s
- 删除了: l
- 删除了: r
- 删除了: c
- 删除了: i
- 删除了: d
- 删除了: r
- 删除了: c
- 删除了: i
- 删除了: also
- 删除了: by taking in
- 删除了: the
- 删除了: which could
- 删除了: e the
- 删除了: of the mapping
- 删除了: ~

308 The spectral characterization method is easy to use and, thus, facilitates access to  
 309 continuous long-term dynamics of biocrusts distribution. However, mosses and vascular plants  
 310 are generally mixed up in this method, because their reflectance characteristics are similar across  
 311 all wavelengths, especially when mosses are wet, which makes them indistinguishable (Fang  
 312 et al., 2015). Therefore, the spectral characterization method mainly applies to situations where  
 313 biocrust cover is greater than 30% and plant cover is less than 10% (Beaugendre et al., 2017).  
 314 It should be noted that the existing indexes mostly correspond to biocrusts cover consisting of  
 315 specific dominant groups in specific environments, which cannot be directly extrapolated to  
 316 areas with highly heterogeneous environments (Table 1). Wetting or disturbance may also lead  
 317 to large fluctuations in the reflectance of different land types, interfering with biocrust  
 318 distribution monitoring (Rodriguez-Caballero et al., 2015; Weber and Hill, 2016).

## 319 2.2 Dynamic global vegetation models (DGVMs)

320 Dynamic global vegetation models are another major method for estimating vegetation  
 321 cover (Deng et al., 2022). These models mainly focus on simulating the biogeochemical  
 322 processes (e.g. carbon and water cycles) and the metabolic and hydrological processes of  
 323 organisms (Fig. 2b) (Lenton et al., 2016; Porada et al., 2017). DGVMs have significant  
 324 advantages in mapping biocrust distribution because their assumptions have clear biological  
 325 implications (Cuddington et al., 2013). Porada et al. (2013) focused on CO<sub>2</sub> diffusion rates and  
 326 photosynthetic processes under dynamic water content saturation in dryland biocrusts. By  
 327 parameterizing long-term climate data and disturbance intervals, and averaging simulation  
 328 results for the past 20 years for each grid point, they estimated that biocrusts cover 11% of the  
 329 global terrestrial land surface (Fig. 3a) (Porada et al., 2019). Specifically, the light and dark  
 330 cyanobacteria were widely distributed in deserts, savannas, grasslands, and Mediterranean  
 331 woodlands at low latitudes, with their presence increasing to some extent with increasing  
 332 dryness. In contrast, mosses were mainly distributed in middle and high latitudes and polar  
 333 regions.

334 Dynamic vegetation models can be combined with cross-scale remotely sensed data to  
 335 quantify the geographic distribution and biogeochemical effects of plants, replacing traditional  
 336 measurements. However, the uneven distribution density of biocrust data points along the

删除了: case

删除了: close to each other in

删除了: is mainly applicable

删除了: >

删除了: <

删除了: and

删除了: e

删除了: is

删除了: to

删除了: e

删除了: ,

删除了: and

删除了: es

删除了: carbon, water cycles),

删除了: (

删除了: )

删除了: The method possesses

删除了: its

删除了: ,

删除了: ,

删除了: ,

删除了: were mostly

设置了格式: 字体颜色: 自动设置

359 aridity gradient or a small amount of data may lead to poor prediction of global-scale  
360 distributions (Quillet et al., 2010). So far, non-vascular vegetation has not received enough  
361 attention, and only the Lichen and Bryophyte Model (LiBry) used in the above case is uniquely  
362 suited to emulating biocrust distribution (Porada et al., 2019; Porada et al., 2013). The LiBry  
363 model includes variations in biocrust cover strategy under disturbance and its growth, but it  
364 relies heavily on subjective experience and model parameterization, which is still immature  
365 compared to dynamic models of vascular vegetation.

### 366 2.3 Geospatial models

367 Directly relating vegetation presence or cover to environmental data, instead of indirectly  
368 via biological processes, is another important way to obtain biocrust distribution (Beaugendre  
369 et al., 2017; Fischer and Subbotina, 2014; Skidmore et al., 2011). Classic statistical models can  
370 serve this purpose. However, they still require comprehensive expert knowledge of how  
371 environmental factors affect biocrusts (Pearce et al., 2001), which is hard to obtain, and prone  
372 to bias. Geospatial models, which integrate machine learning tools with field survey data and  
373 remote sensing data, hold the most promise (Fig. 2c) (Crego et al., 2022). They are also known  
374 as species distribution models or ecological niche models (Brown and Anderson, 2014;  
375 Jiménez-Valverde et al., 2008; Soberon and Nakamura, 2009). At the global scale, there has  
376 been only one study that predicted biocrust distribution patterns using geospatial modeling  
377 (Rodríguez-Caballero et al., 2018), which found that biocrust covers 12.2% of the global land  
378 surface area, which is about  $1.79 \times 10^7 \text{ km}^2$  (Fig. 3b).

删除了: yet may lead to poor predictions of global-scale distributions due to subjective regional experience and insufficient amounts of biocrust data (Table 1)

设置了格式: 字体颜色: 自动设置

删除了: ,

删除了: To use DGVMs, the following procedures need to be taken. The first step is building model framework, simulating important and interesting processes such as biocrusts growth and death, nutrient cycle, and water cycle (Jia et al., 2019). So far, there is only one dynamic global vegetation model targeting at biocrusts - the Lichen and Bryophyte Model (LiBry) (Fig. 1b)

设置了格式: 字体颜色: 自动设置

域代码已更改

删除了: . The second step is parameterization. Generally, literature and open databases are used to assign physiological strategies for different types of biocrust, such as photosynthesis, respiration, and nitrogen emission under the influence of temperature, precipitation, radiation, biological water saturation, etc. The third step is importing environmental data into the model to obtain the biocrusts cover at grid points over a specific study region. At last, the results are tuned and validated against observation data of biocrusts distribution (obtained by literature comparison and local field observations), and thereby biocrusts distribution map is produced.

删除了: in DGVMs

删除了: get

删除了: be

删除了: ed


删除了: are supposed to hold

删除了: and

删除了: The procedures of how to use geospatial modelling are illustrated in Figure 1c

设置了格式: 上标

设置了格式: 上标




带格式的: 缩进: 首行缩进: 0 字符

移动了(插入) [2]

删除了::

删除了:



设置了格式: 上标


删除了: that

410

411 **Fig. 2** Summary of three major approaches to studying biocrust distribution. Illuminations of  
 412 applying spectral characterization method (a), dynamic vegetation model (b), and geospatial  
 413 model (c) in biocrusts distribution study. See the main text for a more detailed introduction to  
 414 these methods.

415 Compared with the result of the dynamic vegetation model, the simulation accuracy  
 416 ( $R^2 \sim 0.8$ ) and mapping resolution ( $0.5^\circ \times 0.5^\circ$ ) of the geospatial model were improved.  
 417 Biocrust distribution is generally consistent in the large deserts of Asia, western America,  
 418 Europe, and Oceania, while some semi-arid regions, such as the northern and southern margins  
 419 of the African Sahara Desert, South Asia, and central North America, have significantly higher  
 420 biocrust cover in the projection by Rodriguez-Caballero et al. (2018). We estimate that this may  
 421 be because geospatial modeling focuses more on the influence of climate, as the Mediterranean  
 422 climate and tropical desert climate in the Sahara Desert, as well as the tropical desert climate  
 423 of northwestern South Asia, are suitable for biocrust survival. Additionally, the large number  
 424 and high cover of biocrust training sets in central North America could have contributed to the  
 425 generally high predicted cover in machine learning.





**Fig. 3** Maps of global biocrusts distribution. (a) Prediction based on vegetation dynamic model (Porada et al., 2019). (b) Prediction based on geospatial model (Rodriguez-Caballero et al., 2018).

As black-boxes, geospatial models are largely non-interpretable and, thus, less capable of capturing the key mechanisms behind phenomena, which may limit their applications. Under this methodological framework, only the direct effects of various environmental indicators are considered. For example, it focuses on the direct effect of precipitation on biocrust distribution while ignoring the indirect effects, such as interactions among shrubs, grasses, and biocrusts (Wang et al., 2024). In addition, to avoid confounding model predictions, the inclusion of environmental factors should be based on their relevance to biocrusts, and expert knowledge should still be needed to a certain degree (Mäkinen et al., 2022). Not only natural conditions such as climate, topography, and soil, but also data on human activities such as afforestation, trampling, and population density need to be considered as environmental indicators in the model. It should be noted that the superimposition of environmental layers of different resolutions may cause deviations in results to some extent, which is unavoidable (Zhao et al., 2024). Despite the above limitations of geospatial modeling, with sufficient computing power, observation data of biocrust distribution, and suitable environmental information, geospatial models are supposed to be relatively optimal solutions for predicting biocrust distribution (Table 1).

**Table 1** Comparison among the three main types of methods to predict biocrust distribution

	Spectral characteristic index	Vegetation dynamics model	Geospatial model
Principle	Differences in wavelength reflectance of surface features	Differences in the physiological processes of different biocrust	Remote sensing information-driven and survey data-based

带格式的: 缩进: 首行缩进: 0 厘米

删除了: 1) extracting environmental data for the sites where biocrusts observation data are reported, 2) importing the extracted environmental data into the machine learning framework and obtaining the relationship between biocrusts distribution and environmental variables through a specific

删除了: black-boxes and

带格式的: 两端对齐, 缩进: 首行缩进: 0.74 厘米

删除了: One should note that

删除了: of environmental factors

删除了: Besides, n

删除了: . that affect biocrust distribution

删除了: as well as

删除了: about

删除了: . also


删除了: supplied with

删除了: and

删除了: able

删除了: to predict

删除了: accurately. Therefore, geospatial modelling is considered to be one of the most appropriate methods



上移了 [2]:

删除了:

删除了: Fig. 1 Summary of three major approaches of studying biocrust distribution. Illustrations of applying

		types	machine learning framework
Advantages	Convenience and ease of use	Clear ecological significance	Machine training simulation, without subjective interference
Disadvantages	Reflectivity is affected by climate change, disturbances; Mosses and vascular plants have similar reflectance characteristics; The results only show the presence or absence of biocrusts without coverage	Experience-based promotion with significant human intervention; Experiments need to be supported by big data	<a href="#">A large</a> amount of computing power; Adequate number of sample points to support accuracy
Applicable scales	Regional scale (Desert and sandy land with <20% vegetation cover)	Regional scale Global scale	Regional scale Global scale

492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505

### 3. Influencing Factors of Biocrust Distribution

It is of great importance to clarify the environmental variables associated with biocrust distribution. On the one hand, it helps to frame the range of data selection before modeling and on the other hand, it aids in identifying patterns of biocrust distribution in the context of dynamic changes and various types of environmental information, thereby facilitating the prediction of distributed evolution on longer time scales. Numerous modelling studies (Kidron and Xiao, 2023; Li et al., 2023; Rodriguez-Caballero et al., 2018) have demonstrated that, on the global scale, biocrust distribution is mainly influenced by water conditions, temperature, soil properties, fire and disturbance (Bowker et al., 2016).

*Water conditions.* In general, total precipitation (Fig. 4b) is considered to be critical in determining the distribution of biocrusts (Eldridge and Tozer, 1997). Increased precipitation can lead to higher levels of lichen and moss cover, while algal cover may initially increase and then decrease (Budel et al., 2009; Marsh et al., 2006; Zhao et al., 2014). It should be noted that

上移了 [1]; <#>Since 1990, studies on the distribution of biocrusts have been continuously increasing. A group of ecologists, represented by Fernando Maestre (Maestre et al., 2021), David Eldridge (Eldridge and Delgado-Baquerizo, 2019; Eldridge et al., 2023), Matthew Bowker (Qiu et al., 2023), Emilio Rodriguez-Caballero (Rodriguez-Caballero et al., 2018) and others, have actively promoted progress in the field (Fig. 2(a)). The topic has gradually received attention from all over the world, particularly the countries with extensive dryland areas such as China, United States, Spain, United Kingdom, Germany, Australia, and Israel (Fig. 2(b)). However, some other dryland countries and regions, such as central and southern Africa, where biocrust distribution has been reported still there is a paucity of studies and the amount of data on biocrusts is far from adequate (Fig. 2(c)). These areas may be potential areas of widespread distribution of biocrusts in the future.

- 删除了: <#>Current State of Knowledge
- 删除了: it helps
- 删除了: experimental observations and
- 删除了: proved
- 删除了: (
- 删除了: )
- 删除了: could
- 删除了: first

572 precipitation can also promote the growth of vascular plants, and continuous high cover of  
 573 vascular plants and litterfall will limit the space available to biocrusts (Bowker et al., 2005). In  
 574 addition to the total amount of precipitation, the seasonality and frequency of precipitation  
 575 cannot be ignored (Budel et al., 2009). Winter precipitation and/or smaller rain events benefit  
 576 biocrusts, especially when mean annual precipitation is less than 500 mm. Meanwhile, a high  
 577 frequency of precipitation can lead to the dominance of biocrusts over vascular plants (Chamizo  
 578 et al., 2016; Jia et al., 2019). Experimental evidence shows that precipitation events of 5 mm  
 579 are able to maintain normal physiological and ecological functions of the biocrust on the  
 580 Colorado Plateau, USA, while ever lower precipitation events of 1.2 mm can rapidly kill moss  
 581 biocrust (Reed et al., 2012). Non-precipitation water input is another important water resource  
 582 type. The Namib Desert receives little rainfall, but lichens and moss biocrusts can reach a  
 583 relatively high cover (~70%) (Budel et al., 2009). This is because local water vapor tends to  
 584 condense into fog or dew, which facilitates the survival of three-dimensional species (such as  
 585 leafy lichens) by trapping air moisture (Eldridge et al., 2020; Kidron, 2019; Li et al., 2021).  
 586 Similarly, lichen biocrusts are widely distributed in the western U.S. along the Mexican coast  
 587 due to the high air humidity (dew formation for almost 1/3 of the year) (McCune et al., 2022;  
 588 Miranda - González and McCune, 2020).

589 *Temperature.* Relatively high soil temperature can create an environment of high  
 590 evaporation that impedes biocrusts colonization (Garcia-Pichel et al., 2013). Regarding air  
 591 temperature, warming by 4°C could alter biocrust community structure, resulting in a sharp  
 592 decrease in moss biocrust cover and an increase in cyanobacterial biocrust cover. This effect  
 593 becomes even more significant when warming interacts with time and precipitation treatments  
 594 (Ferrenberg et al., 2015). Recent studies have shown that historical and future temperature  
 595 changes also affect biocrust distribution. For example, the climate legacy over the last 20,000  
 596 years could indirectly affect the distribution and relative species richness of biocrusts by  
 597 altering vegetation cover and soil pH (Eldridge and Delgado-Baquerizo, 2019). Additionally,  
 598 under future scenarios of increased temperature and aridity, biocrust cover is predicted to  
 599 decrease by approximately 25% by the end of the century, with communities shifting towards  
 600 early cyanobacterial biocrusts (Rodríguez-Caballero et al., 2022).

删除了: the

删除了: and

删除了: <

删除了: and

删除了: It was e

删除了: ly

删除了: proven

删除了: were

删除了: could

删除了: the

删除了: u

删除了: as the result of

删除了: ,

删除了: which became even more significant when warming was interacting with time and precipitation treatments

删除了: through changing

删除了: s

618 *Soil properties.* For a long time, it was commonly believed that finer soils benefit [biocrust](#)  
619 growth (Belnap et al., 2014; Williams et al., 2013). However, [some scientists have challenged](#)  
620 [this notion](#) (Fig. 4c). For example, Kidron (2018) argued that soils with high dust or fine grains  
621 [are](#), not a necessary condition for [biocrust](#) distribution. Qiu et al. (2023) suggested that soils  
622 with small amounts of gravel (0.04-22.34% content, 0.58% [being optimal](#)) are more favorable  
623 for biocrusts. Another study [has](#) shown that the soil parent material determines the degree of  
624 surface weathering and the water-holding capacity of the soil, thus indirectly [influencing](#) the  
625 distribution of biocrusts (Bowker and Belnap, 2008). Gypsum or calcareous soils tend to  
626 develop mosses and lichens (Elbert et al., 2012), while sandy soils tend to develop  
627 cyanobacteria (Root and Mccune, 2012).

628 *Fire.* Grassland is [a major life form](#) in dryland ecosystems, making it [crucial](#) to explore  
629 the effects of fire events on [biocrust](#) distribution (Palmer et al., 2022). Fire-induced soil  
630 warming can alter the resource allocation and dynamic growth mechanisms between biocrusts  
631 and vascular plants (McCann et al., 2021), potentially leading to a reduction in species richness  
632 and cover of biocrusts, especially cyanobacteria, and algae (Abella et al., 2020; Palmer et al.,  
633 2020). (Condon and Pyke, 2018) showed that moss cover increases with time after [the](#) fire, with  
634 no significant change in lichen cover.

635 *Disturbance.* Activities such as grazing, agricultural practices, and land development can  
636 significantly impact biocrust distribution. Studies have demonstrated that grazing intensity can  
637 lead to substantial changes in biocrust cover. For instance, in Patagonian rangelands, biocrust  
638 cover decreased by 85%, 89%, and 98% under light, medium, and heavy grazing, respectively  
639 (Velasco Ayuso et al., 2019). In the Loess Plateau, total biocrust cover remained almost  
640 unchanged under light grazing (< 30.00 goat dung / m<sup>2</sup>), but there were variations in community  
641 structure, with an increase in cyanobacteria biocrusts (23.1%) and a decrease in moss biocrusts  
642 (42.2%) due to [reduction](#) in vascular plant cover (Ma et al., 2023). Tillage practices can disrupt  
643 the soil surface, leading to a reduction in biocrust cover ( 6% on average) and diversity, [with](#),  
644 lichens [struggling](#) to survive in tilled fields compared to mosses (Durham et al., 2018).  
645 Additionally, [late-successional](#) biocrusts exhibit higher tolerance compared to pre-successional  
646 biocrusts. Moss biocrusts, for instance, can maintain soil microbial biomass and nematode

删除了: y

删除了: (

删除了: )

删除了: were

删除了: best

删除了: changing

删除了: one

删除了: of great significance

删除了: the decrease

删除了: and

删除了: tend to

删除了: e

659 abundance better under trampling disturbance compared to cyanobacteria and lichen biocrusts  
660 (Yang et al., 2018). However, contrary to this view, it has been observed that cyanobacterial  
661 biocrusts increased in cover from 81% to 99% after trampling, while lichen and moss biocrusts  
662 decreased from 1.5% and 18% to less than 0.5%. Furthermore, mining activities can  
663 significantly reduce the photosynthetic potential of biocrusts, particularly affecting the recovery  
664 of cyanobacterial biocrusts (Gabay et al., 2022).

665 *Other factors.* On a global scale, biocrust distribution is also closely linked to  
666 biogeographic isolation. Strong spatial heterogeneity, accompanied by spatial distance, can  
667 create barriers to the dispersal of propagules (spores, fungal bodies), which indirectly impedes  
668 colonization of the biocrusts (Garcia-Pichel et al., 2013). In addition, factors such as vascular  
669 plant cover, topography, and solar radiation also influence biocrust distribution, albeit to a lesser  
670 extent than the factors mentioned above. For further insights, readers are encouraged to consult  
671 Chapter 10 of *Biological Soil Crusts: An Organizing Principle in Drylands*, which provides an  
672 overview of the control and distribution patterns of biocrusts from micro to global scales  
673 (Bowker et al., 2016).

674 To sum up, climate is the most important factor influencing global biocrust distribution,  
675 especially in drylands where water is precious to the organisms. However, exploration of the  
676 roles of climatic factors such as rainfall seasonality and atmospheric drought still needs much  
677 further effort (Wright and Collins, 2024), especially in the context of global climate change.  
678 Although more attention has been paid to the physical properties of soils, the roles of their  
679 chemical properties, such as the nitrogen (N) and phosphorus (P) content, need to be taken  
680 more seriously. Fire and disturbance are usually ignored. However, due to the trend towards  
681 warmer and drier environments, as well as increasing population and the need to sustain  
682 livelihoods, their influences on biocrust distribution may become more important. As one of  
683 the basic processes on a global scale, biogeographic isolation or changes in land use should be  
684 paid more attention to. With the increasing number of biocrust data points, we can expect this  
685 aspect will see a surge in research.

删除了: set up

删除了: ,

删除了: A

删除了: ,

删除了: in the


删除了: paragraphs

删除了: But

删除了: more

删除了: s

删除了: As amounting data points of biocrust



696 Fig. 4 Biocrust distribution and its critical influencing factors. (a) Biocrust cover map and its  
 697 influencing factors. (a) Global biocrust distribution, by random forest modelling. Based on a  
 698 global biocrust database constructed by Chen et al., we expanded the biocrust data to 3848  
 699 entries through literature compilation and field surveys and fitted them with four types of  
 700 remotely sensed environmental data, including climate, land use, soil properties, and elevation,  
 701 to finally predict the suitable areas for the biocrust distribution and quantify the biocrust cover.  
 702 (b) Global average annual precipitation (1970-2020), data from the WorldClim database  
 703 (version 2.1). (c) Global soil texture distribution, data from HWSD (Harmonized World Soil  
 704 Database, version 1.2). Precipitation and soil texture were taken as examples of environmental  
 705 factors.

706 **4. Challenges and Perspectives**

707 Biocrusts play a crucial role in dryland ecosystems, making it essential to understand their  
 708 current status and distribution dynamics. For influencing factors (Chapter 3), traditional  
 709 observational studies and controlled experiments offer multiple perspectives of foundational  
 710 knowledge. For assessing biocrust distribution patterns (Chapter 2), the methods shift from  
 711 traditional approaches to spectral index, vegetation dynamics, and geospatial models that span  
 712 multiple subjects like ecology, biology, geology, and computer science. However, high-

- 删除了: B
- 删除了: iocrusts are very important for dryland ecosystems, and thus, it is of outstanding significance to understand the current status and dynamics of biocrust distribution.
- 删除了: the
- 删除了: 4
- 删除了: provide us with
- 删除了: basic
- 删除了: 3

722 precision biocrust distribution data across geographic units remain scarce, and current research  
723 methods are still limited. To further advance studies of biocrust distribution, we propose the  
724 following aspects for consideration.

删除了: s

删除了: are still lacking

删除了: raise

### 725 5.1 Building standardized biocrusts database

726 Currently, biocrust data are fragmented, low in volume, and derived from narrow sources,  
727 largely limiting spatial prediction from points to broader areas. Thus, we suggest that a global  
728 effort to build a standardized and specialized biocrusts database. This database should include  
729 consistent data items (such as main types and cover of biocrusts, latitude, longitude, and cover)  
730 and adhere to uniform inclusion criteria. Such a database is an important infrastructure for  
731 mapping global biocrust distribution, serving as the benchmark for training and validating  
732 spectral characteristics, DGVM, and geospatial models (Engel et al., 2023). Given the difficulty  
733 of conducting field surveys worldwide, compiling biocrust data from the published literature  
734 or other sources would be a primary approach (Fig. 4(a)). To date, several published studies  
735 have assembled 900 ~ 1,000 data on biocrust presence or absence from the literature (including  
736 584 data on biocrust cover) (Chen et al., 2020; Eldridge et al., 2020; Havrilla et al., 2019;  
737 Rodriguez-Caballero et al., 2018). However, compiling from literature largely comes to its  
738 limitations and is still far from building a standardized and specialized biocrusts database.

删除了: accessed

删除了: should

删除了: consisting of the same data items (main types and cover of biocrusts, latitude, longitude, and cover, etc.) and using the same inclusion criteria.

删除了: It

删除了: to

删除了: e

删除了: Due to

删除了: s

删除了: the major

739 While open databases are not specialized to biocrusts, some of them may provide valuable  
740 additions (Fig. 5). For instance, the biodiversity and specimen datasets such as GBIF and the  
741 Atlas of Living Australia (Belbin and Williams, 2015; García-Roselló et al., 2015), contain a  
742 large amount of information on species, including mosses and lichens (Table 2), potentially  
743 offering hundreds or even thousands of entries of biocrusts occurrence or cover. Similarly,  
744 global, national, and regional plant flora can significantly contribute to building the  
745 standardized and specialized biocrusts database. For example, sPlot includes ~2 million  
746 vegetation plot data (Sabatini et al., 2021), and the European Vegetation Archive (EVA) also  
747 holds 1.6 million entries over the globe or Europe (Chytrý et al., 2016). Regional datasets like  
748 the Environmental Monitoring of Arid and Semiarid Regions (MARAS) have surveyed 426 sites  
749 (up to September 2020) and provided regular access to 624.50 km<sup>2</sup> of rangeland vegetation  
750 spatial patterns, species diversity, soil functional indices, climatic data, and landscape

删除了: be

删除了: important

删除了: example

删除了: , which

删除了: of course

删除了: which can contain

删除了: may

删除了: s of

删除了: possesses

删除了: data

775 photographs in [the Patagonia region of Argentina and Chile](#) (Oliva et al., 2020). Concerns about  
 776 land use products are also necessary. Global land use maps, based on the PROBA-V sensor,  
 777 which contain spatial information for [the Moss & Lichen layer](#), have an annual update  
 778 frequency and a resolution of 100 m. [Additionally, an increasing number of amateurs contribute](#)  
 779 [significantly to global species information entries through species identification apps, which](#)  
 780 [are user-friendly and widely accessible.](#) The citizen science project *iNaturalist* is a very good  
 781 example (Wolf et al., 2022). Furthermore, when collecting and collating data from non-  
 782 academic sources, the combination of web crawlers and text analysis can help in obtaining  
 783 biocrusts data and [addressing](#) key ecological issues.

删除了: straddling

删除了: In addition to the above channels, an increasing number of amateurs are getting involved in science through species identification apps with clean, easy-to-use apps, contributing significantly to global species information entries....

删除了: solving



784  
 785 **Fig. 5** Potential approaches to [building](#) a standardized biocrusts database. (a) Distribution of  
 786 lichens in the GBIF database with an example photo, (b) environmental monitors distribution  
 787 map of MARAS database, (c) distribution of "mosses and lichens" in the PROBAV\_LC100  
 788 database (light yellow area) in northern Asia, for instance.

**Table 2** References for biocrusts database expansion channels



797

798

## 5.2 Improving non-vascular vegetation dynamic models

17

Data type	Data source	Extend	Biocrust type	Georeferenced records	Presence	Coverage	Link	
Biodiversity data	the Global Biodiversity Information Facility (GBIF)	Worldwide	Cyanobacteria	~780000	✓	--	<a href="https://www.gbif.org/">https://www.gbif.org/</a>	
			Lichen	~19000				
			Moss	~90000				
	Atlas of Living Australia (ALA)	Australia	Cyanobacteria	~53000	✓	--	<a href="https://www.ala.org.au/">https://www.ala.org.au/</a>	
			Lichen	~12000				
			Moss	~20000				
	Chinese Virtual Herbarium	China	Moss and lichen	--	✓	--	<a href="https://www.cvh.ac.cn/">https://www.cvh.ac.cn/</a>	
			Lichen	~2000	✓	--	<a href="https://plants.jstor.org/">https://plants.jstor.org/</a>	
	Citizen Science Survey data	Global Plants on JSTOR	Worldwide	Moss	~480	✓	--	<a href="https://www.inaturalist.org/">https://www.inaturalist.org/</a>
				All	--	✓	--	<a href="https://springernature.figshare.com/collections/The_MARAS_dataset_vegetation_and_soil_characteristics_of_dryland_rangelands_across_Patagonia/4789113">https://springernature.figshare.com/collections/The_MARAS_dataset_vegetation_and_soil_characteristics_of_dryland_rangelands_across_Patagonia/4789113</a>
iNaturalist		Worldwide	All	426	✓	✓	<a href="https://www.idiv.de/en/splot.html">https://www.idiv.de/en/splot.html</a>	
			All	--	✓	✓	<a href="https://edg.org/databases/GrassPlot/">https://edg.org/databases/GrassPlot/</a>	
MARAS		Argentina and Chile	Lichen	6801	✓	✓	<a href="http://vegbank.org/">http://vegbank.org/</a>	
			Moss	11001	✓	✓	<a href="https://gpp-blm-egis.hub.arsgis.com/pages/aim">https://gpp-blm-egis.hub.arsgis.com/pages/aim</a>	
sPlot		Worldwide	Non-vascular plants	6623	✓	✓	<a href="http://vegbank.org/">http://vegbank.org/</a>	
			Moss and lichen	~15000	✓	✓		
GrassPlot		Worldwide	Moss and lichen	5200	✓	✓		
			Moss and lichen	--	✓	✓		
Vegbank	Canada and the United States	All	~300			<a href="http://www.aekos.org.au/">http://www.aekos.org.au/</a>		
		Moss and lichen	--			<a href="https://land.copernicus.eu/global/products/lc">https://land.copernicus.eu/global/products/lc</a>		
BLM_AIM	The United States	All	--					
		Moss and lichen	--					
Landcover data	TERN AEKOS PROBA_V_LC100	Australia	~300					
		Worldwide	--					

799 There are only two DGVMs applicable to non-vascular organisms – LiBry and ECHAM6-  
 800 HAM2-BIOCRUST (Rodriguez-Caballero et al., 2022). ~~Despite their utility, these models still~~  
 801 ~~require performance improvements. Future directions for enhancing these models could include~~  
 802 ~~incorporating spatial self-organization of non-vascular organisms~~ (Gassmann et al., 2000), the  
 803 effects of fire (Thonicke et al., 2001), vegetation-environment feedback processes (Quillet et  
 804 al., 2010), functional traits (Boulangeat et al., 2012), intraspecific-interspecific interactions  
 805 (Boulangeat et al., 2014) and seasonal dynamics. ~~Moreover, the physical properties,~~  
 806 ~~photosynthetic capacity, and carbon and nitrogen allocation of biocrusts change along~~  
 807 ~~environmental gradients in complex and context-dependent ways. These factors~~ should be  
 808 incorporated into DGVMs (Fatichi et al., 2019). Spatial-explicit DGVMs may be one key to  
 809 effectively improving the accuracy of simulations in future studies, ~~although they~~ are data-  
 810 ~~intensive~~. Also, biocrusts are significantly influenced by hydrological processes ~~and, in turn,~~  
 811 ~~affect these processes~~ (Chen et al., 2018; Whitney et al., 2017). ~~However,~~ ecohydrological  
 812 models, ~~which focus~~ on hydrological processes are rarely connected to global biocrust  
 813 distribution predictions. (Jia et al., 2019) ~~attempted~~ to incorporate biocrusts cover as a system  
 814 state variable in an ecohydrological model, ~~investigating~~ biocrusts cover under ~~varying~~ rainfall  
 815 gradients. ~~By feeding ecohydrological models with global environmental data, particularly~~  
 816 ~~hydrological variables, these models could offer a new approach to predicting biocrust~~  
 817 ~~distribution on a global scale.~~

### 818 5.3 Integrated application of high-quality sensors

819 ~~The spectral~~ characterization method lies ~~in~~ the differences in spectral reflectance of  
 820 biocrusts and other land types at ~~various~~ wavelengths. ~~Consequently,~~ the accuracy of the results  
 821 ~~is contingent~~ on the quality of the sensors ~~used~~. ~~Previous studies~~ often ~~employed~~ a single sensor  
 822 with ~~fixed~~ band intervals for distinguishing biocrusts, ~~potentially missing critical~~ spectral  
 823 features ~~of different~~ land types (Chamizo et al., 2012a). If the biocrusts index can be constructed  
 824 by combining and comparing the full-band spectral data from multiple terrestrial sensors and  
 825 infrared cameras, and other devices, the errors will be reduced to a certain extent, thus  
 826 improving the classification accuracy (Wang et al., 2022b). In addition, the unique advantages  
 827 of hyperspectral data, ~~which include large data volumes and narrow bands,~~ allow for the

- 删除了: Their performances still need to be improved.
- 删除了: Considering spatial self-organization of non-vascular organisms
- 删除了: in current and/or new DGVMs can be future directions
- 删除了: In addition
- 删除了: , how the physical properties, photosynthetic capacity, and carbon and nitrogen allocation of biocrusts change along environmental gradient are complex and context-dependent,
- 删除了: which
- 删除了: but
- 删除了: consuming
- 删除了: and vice versa
- 删除了: , while
- 删除了: based
- 删除了: s
- 删除了: tried
- 区域代码已更改
- 删除了: and
- 删除了: ed
- 删除了: If fed with global data on environmental variables (mainly hydrological relevant ones), ecohydrological models may be a new approach to predicting biocrust distribution at global scales
- 删除了: different
- 删除了: , and thus
- 删除了: depends
- 删除了: However, p
- 删除了: have
- 删除了: used
- 删除了: constant
- 删除了: which may result in missed
- 删除了: identification
- 删除了: with large data volume and narrow band

862 ~~development of new biocrust discrimination standards when combined with observational data,~~  
863 If further estimation of biocrust cover can be achieved on this basis, it will be a ~~significant~~  
864 contribution to the study of ~~large-scale biocrust~~ distribution (Rodríguez-Caballero et al., 2017).  
865 To date, high-resolution sensors have ~~proven~~ successful in monitoring lichens and mosses  
866 (Blanco-Sacristan et al., 2021), and the release of such products is something important to look  
867 out for in the future.

删除了: allow it to be combined with observation data to co-develop new biocrusts discrimination standards

删除了: n

删除了: important

删除了: been found to be

#### 868 5.4 Making full use of machine learning

869 Machine learning can be combined with remote sensing products to ~~uncover~~ complex  
870 features from big data, ~~enabling the prediction of global biocrust distribution,~~ (Collier et al.,  
871 2022). This data-driven approach has powerful predictive capabilities, especially for mapping  
872 species distribution, and can largely ~~avoid the errors~~ of missing or misidentifying biocrusts  
873 caused by traditional methods (relying on field measurements to determine threshold ranges)  
874 (Wang et al., 2022b). In the remote sensing image classification, ~~mature machine learning~~  
875 algorithms include support vector machines, single decision trees, random forests, artificial  
876 neural networks, etc. (Yu et al., 2020). Ensemble models combining multiple algorithms have  
877 been widely used in the field of species distribution, ~~but have seen relatively few applications~~  
878 ~~in biocrust prediction,~~ In the future, using machine learning to ~~identify~~ parameters for dynamic  
879 models of biocrusts may be one of the most promising ~~methods~~ to predict biocrust distribution  
880 (Perry et al., 2022).

删除了: find

删除了: to predict global biocrust distribution

删除了: circumvent

删除了: mistakes

删除了: problem

删除了: but relatively few applications exist in the field of biocrusts prediction.

删除了: find

#### 881 5.5 Regional research synergy development

882 Research ~~on~~ biocrust distribution ~~has~~ shown significant spatial and climatic imbalances.  
883 ~~The~~ study areas that have been conducted are relatively concentrated in countries such as China,  
884 ~~the United States,~~ Spain, Australia, and Israel. Although there are large areas of dryland  
885 distributed in Africa (other than South Africa), central Asia, central South America, and  
886 northern North America, research on biocrusts in these regions ~~is~~ scarce. ~~These unbalanced~~  
887 regional research efforts ~~constraint~~ ~~the advancement of studies on~~ global ~~biocrust~~ distribution.  
888 Therefore, how to coordinate and promote the common progress of regional research is an  
889 urgent issue at present. Climatically, in addition to the drylands, the cold zones may be another  
890 important area to explore biocrust distribution (Pushkareva et al., 2016). On the Tibetan Plateau,

删除了: e

删除了: USA,

删除了: U

删除了: are one of the

删除了: s

删除了: to

删除了: ing

删除了: of

912 studies have investigated the spatial variation of different types of biocrust communities across  
913 climatic gradients and their effects on soil temperature features and freezing duration (Ming et  
914 al., 2022; Wei et al., 2022). These findings highlight the need for more studies on biocrust  
915 distribution in the alpine areas.

## 916 **5. Conclusion**

917 Biocrusts are of great significance to the ecohydrological processes, soil material cycling,  
918 landscape shaping, and biodiversity conservation in drylands. To date, numerous studies have  
919 tried to fill the knowledge gap in biocrust distribution at the regional scale. However, global-  
920 scale research remains scarce, and mapping accuracy is still insufficient, directly leading to  
921 ambiguities in ecological function assessment and prediction. Therefore, advancing global-  
922 scale biocrust distribution research requires a more comprehensive consideration of the  
923 applicability of previous methods and a broader knowledge base to help select environmental  
924 indicators. For future work in this field, we advocate for closer cooperation among scientists to  
925 build a global standardized database incorporating multiple sources of biocrust data. This effort  
926 should primarily focus on expanding biocrust data items in understudied regions where  
927 biocrusts have been reported, thereby creating a larger, multi-habitat training set. Meanwhile,  
928 modern learning tools, such as deep learning, should be broadly applied to high-quality sensor  
929 image segmentation, data classification, and model parameter tuning. Finally, long-term  
930 monitoring and simulation are necessary to better understand the dynamics of ecological  
931 restoration in drylands and the response of biocrusts to environmental changes.

## 932 **Author contribution**

933 Siqing Wang co-conceived the idea, collected data on the biological soil crust, wrote the first  
934 draft, prepared the figures, and revised the manuscript. Li Ma, Liping Yang, Yali Ma, and  
935 Yafeng Zhang collected data on biological soil crust and revised the manuscript. Changming  
936 Zhao co-conceived the idea. Ning Chen co-conceived the idea, collected data on the biological  
937 soil crust, revised the manuscript, and provided support for funding.

## 938 **Acknowledgements**

删除了: is urges the

删除了: of

删除了: s

删除了: T

删除了: his work aims to advance global knowledge of biocrust distribution for better ecosystem management and sustainable development in drylands.

删除了: ed

删除了: of

删除了: the distribution of dominant biocrust communities

删除了: scales

删除了: thethe of XX

删除了: area

删除了: a

删除了: especially

删除了: We firstly compared the advantages, disadvantages, and applicability among three methods, spectral characterization index, dynamic global vegetation models and geospatial models, in order to provide the most appropriate methodological suggestions for biocrust distribution studies at different scales and needs. Then, we systematically sorted out the regional-global biocrust distribution cases, and drew a map of global biocrust distribution hotspots and a map of spatial distribution of data points. Further, we tried to clarify the causes of biocrust distribution from several aspects, such as precipitation, temperature, soil, fire, and other anthropogenic factors. Finally, from a personal point of view, we would like to focus more on the following points in the future: database construction, model performance enhancement, big data processing, and synergistic progress of potential distribution area studies.

设置了格式: 字体: 非加粗

972 We are grateful to Zhiguang Zhao, Lixun Zhang, [and](#) Chao Guan for [their helpful discussions](#)  
973 [while](#) preparing the first draft. The study was supported by the National Natural Science  
974 Foundation of China (grant numbers [32271620](#), [31971452](#)), [the](#) National Key Research and  
975 Development Program of China (2023YFF0805603), and the Central University Basic  
976 Research Funds (lzujbky-2021-ey16, lzujbky-2023-eyt01).

977

### 978 **Competing interests**

979 All authors declare no conflict of interest.

980

### 981 **Reference**

- 982 Abella, S. R., Gentilcore, D. M., and Chiquoine, L. P.: Resilience and alternative stable states after desert  
983 wildfires, *Ecological Monographs*, 91, 1-19, 10.1002/ecm.1432, 2020.
- 984 Baxter, C., Mallen-Cooper, M., Lyons, M. B., and Cornwell, W. K.: Measuring reflectance of tiny  
985 organisms: The promise of species level biocrust remote sensing, *Methods in Ecology and Evolution*, 12,  
986 2174-2183, 10.1111/2041-210x.13690, 2021.
- 987 Beaugendre, N., Malam Issa, O., Choné, A., Cerdan, O., Desprats, J.-F., Rajot, J. L., Sannier, C., and  
988 Valentin, C.: Developing a predictive environment-based model for mapping biological soil crust patterns  
989 at the local scale in the Sahel, *Catena*, 158, 250-265, 10.1016/j.catena.2017.06.010, 2017.
- 990 Belbin, L. and Williams, K. J.: Towards a national bio-environmental data facility: experiences from the  
991 Atlas of Living Australia, *International Journal of Geographical Information Science*, 30, 108-125,  
992 10.1080/13658816.2015.1077962, 2015.
- 993 Belnap, J., Miller, D. M., Bedford, D. R., and Phillips, S. L.: Pedological and geological relationships  
994 with soil lichen and moss distribution in the eastern Mojave Desert, CA, USA, *Journal of Arid  
995 Environments*, 106, 45-57, 10.1016/j.jaridenv.2014.02.007, 2014.
- 996 Blanco-Sacristan, J., Panigada, C., Gentili, R., Tagliabue, G., Garzonio, R., Martin, M. P., Ladron de  
997 Guevara, M., Colombo, R., Dowling, T. P. F., and Rossini, M.: UAV RGB, thermal infrared and  
998 multispectral imagery used to investigate the control of terrain on the spatial distribution of dryland  
999 biocrust, *Earth Surface Processes and Landforms*, 46, 2466-2484, 10.1002/esp.5189, 2021.
- 1000 Boulangeat, I., Georges, D., and Thuiller, W.: FATE-HD: a spatially and temporally explicit integrated  
1001 model for predicting vegetation structure and diversity at regional scale, *Global Change Biology*, 20,  
1002 2368-2378, 10.1111/gcb.12466, 2014.
- 1003 Boulangeat, I., Philippe, P., Abdulhak, S., Douzet, R., Garraud, L., Lavergne, S., Lavorel, S., van Es, J.,  
1004 Vittoz, P., and Thuiller, W.: Improving plant functional groups for dynamic models of biodiversity: at the  
1005 crossroads between functional and community ecology, *Global Change Biology*, 18, 3464-3475,  
1006 10.1111/j.1365-2486.2012.02783.x, 2012.
- 1007 Bowker, M. A. and Belnap, J.: A simple classification of soil types as habitats of biological soil crusts on  
1008 the Colorado Plateau, USA, *Journal of Vegetation Science*, 19, 831-840, 10.3170/2008-8-18454, 2008.
- 1009 Bowker, M. A., Belnap, J., Davidson, D. W., and Phillips, S. L.: Evidence for Micronutrient Limitation  
1010 of Biological Soil Crusts: Importance to Arid-Lands Restoration, *Ecological Applications*, 15, 1941-1951,  
1011 10.1890/04-1959, 2005.

删除了: helpful discussions when

带格式的: 缩进: 首行缩进: 0 字符

设置了格式: 字体: 加粗

1013 Bowker, M. A., Belnap, J., Büdel, B., Sannier, C., Pietrasiak, N., Eldridge, D. J., and Rivera-Aguilar, V.:  
1014 Controls on Distribution Patterns of Biological Soil Crusts at Micro- to Global Scales, in: *Biological Soil*  
1015 *Crusts: An Organizing Principle in Drylands*, edited by: Weber, B., Büdel, B., and Belnap, J., Springer  
1016 International Publishing, Cham, 173-197, 10.1007/978-3-319-30214-0\_10, 2016.

1017 Brown, J. L. and Anderson, B.: SDMtoolbox: a python-based GIS toolkit for landscape genetic,  
1018 biogeographic and species distribution model analyses, *Methods in Ecology and Evolution*, 5, 694-700,  
1019 10.1111/2041-210x.12200, 2014.

1020 Budel, B., Darienko, T., Deuschewitz, K., Dojani, S., Friedl, T., Mohr, K. I., Salisch, M., Reisser, W.,  
1021 and Weber, B.: Southern African biological soil crusts are ubiquitous and highly diverse in drylands,  
1022 being restricted by rainfall frequency, *Microbial Ecology*, 57, 229-247, 10.1007/s00248-008-9449-9,  
1023 2009.

1024 Chamizo, S., Cantón, Y., Miralles, I., and Domingo, F.: Biological soil crust development affects  
1025 physicochemical characteristics of soil surface in semiarid ecosystems, *Soil Biology and Biochemistry*,  
1026 49, 96-105, 10.1016/j.soilbio.2012.02.017, 2012a.

1027 Chamizo, S., Belnap, J., Eldridge, D. J., Cantón, Y., and Malam Issa, O.: The Role of Biocrusts in Arid  
1028 Land Hydrology, in: *Biological Soil Crusts: An Organizing Principle in Drylands*, edited by: Weber, B.,  
1029 Büdel, B., and Belnap, J., Springer International Publishing, Cham, 321-346, 10.1007/978-3-319-30214-  
1030 0\_17, 2016.

1031 Chamizo, S., Stevens, A., Canton, Y., Miralles, I., Domingo, F., and Van Wesemael, B.: Discriminating  
1032 soil crust type, development stage and degree of disturbance in semiarid environments from their spectral  
1033 characteristics, *European Journal of Soil Science*, 63, 42-53, 10.1111/j.1365-2389.2011.01406.x, 2012b.

1034 Chen, J., Zhang, M. Y., Wang, L., Shimazaki, H., and Tamura, M.: A new index for mapping lichen-  
1035 dominated biological soil crusts in desert areas, *Remote Sensing of Environment*, 96, 165-175,  
1036 10.1016/j.rse.2005.02.011, 2005.

1037 Chen, N., Jayaprakash, C., Yu, K., and Guttal, V.: Rising Variability, Not Slowing Down, as a Leading  
1038 Indicator of a Stochastically Driven Abrupt Transition in a Dryland Ecosystem, *Am Nat*, 191, E1-E14,  
1039 10.1086/694821, 2018.

1040 Chen, N., Yu, K. L., Jia, R. L., Teng, J. L., and Zhao, C. M.: Biocrust as one of multiple stable states in  
1041 global drylands, *Science Advances*, 6, 10.1126/sciadv.aay3763, 2020.

1042 Chytrý, M., Hennekens, S. M., Jiménez-Alfaro, B., Knollová, I., Dengler, J., Jansen, F., Landucci, F.,  
1043 Schaminée, J. H. J., Acic, S., Agrillo, E., Ambarlı, D., Angelini, P., Apostolova, I., Attorre, F., Berg, C.,  
1044 Bergmeier, E., Biurrun, I., Botta-Dukát, Z., Brisse, H., Campos, J. A., Carlón, L., Čarni, A., Casella, L.,  
1045 Csiky, J., Čušterevska, R., Dajić Stevanović, Z., Danihelka, J., De Bie, E., de Ruffray, P., De Sanctis, M.,  
1046 Dickoré, W. B., Dimopoulos, P., Dubyna, D., Dziuba, T., Ejmaes, R., Ermakov, N., Ewald, J., Fanelli, G.,  
1047 Fernández-González, F., FitzPatrick, Ú., Font, X., García-Mijangos, I., Gavilán, R. G., Golub, V.,  
1048 Guarino, R., Haveman, R., Indreica, A., Işık Gürsoy, D., Jandt, U., Janssen, J. A. M., Jiroušek, M., Kącki,  
1049 Z., Kavğacı, A., Kleikamp, M., Kolomiychuk, V., Krstivojević Čuk, M., Krstonošić, D., Kuzemko, A.,  
1050 Lenoir, J., Lysenko, T., Marcenò, C., Martynenko, V., Michalčová, D., Moeslund, J. E., Onyshchenko,  
1051 V., Pedashenko, H., Pérez-Haase, A., Peterka, T., Prokhorov, V., Rašomavičius, V., Rodríguez-Rojo, M.  
1052 P., Rodwell, J. S., Rogova, T., Ruprecht, E., Růsiņa, S., Seidler, G., Šibík, J., Šilc, U., Škvorc, Ž.,  
1053 Sopotlieva, D., Stančić, Z., Svenning, J.-C., Swacha, G., Tsiripidis, I., Turtureanu, P. D., Uğurlu, E.,  
1054 Uogintas, D., Valachovič, M., Vashenyak, Y., Vassilev, K., Venanzoni, R., Virtanen, R., Weekes, L.,  
1055 Willner, W., Wohlgemuth, T., Yamalov, S., and Pärtel, M.: European Vegetation Archive (EVA): an  
1056 integrated database of European vegetation plots, *Applied Vegetation Science*, 19, 173-180,

1057 10.1111/avsc.12191, 2016.

1058 Collier, E. A., Perroy, R. L., Reed, S. C., and Price, J. P.: Mapping biological soil crusts in a Hawaiian  
1059 dryland, *International Journal of Remote Sensing*, 43, 484-509, 10.1080/01431161.2021.2003904, 2022.

1060 Condon, L. A. and Pyke, D. A.: Fire and Grazing Influence Site Resistance to *Bromus tectorum* Through  
1061 Their Effects on Shrub, Bunchgrass and Biocrust Communities in the Great Basin (USA), *Ecosystems*,  
1062 21, 1416-1431, 10.1007/s10021-018-0230-8, 2018.

1063 Crego, R. D., Stabach, J. A., Connette, G., and Leroy, B.: Implementation of species distribution models  
1064 in Google Earth Engine, *Diversity and Distributions*, 28, 904-916, 10.1111/ddi.13491, 2022.

1065 Cuddington, K., Fortin, M., Gerber, L., Hastings, A., Liebhold, A., O'Connor, M., and Ray, C.: Process-  
1066 based models are required to manage ecological systems in a changing world, *Ecosphere*, 4, 1-12, 2013.

1067 Deng, M., Meng, X., Lu, Y., Li, Z., Zhao, L., Niu, H., Chen, H., Shang, L., Wang, S., and Sheng, D.: The  
1068 Response of Vegetation to Regional Climate Change on the Tibetan Plateau Based on Remote Sensing  
1069 Products and the Dynamic Global Vegetation Model, *Remote Sensing*, 14, 3337, 10.3390/rs14143337,  
1070 2022.

1071 Durham, R. A., Doherty, K. D., Antoninka, A. J., Ramsey, P. W., and Bowker, M. A.: Insolation and  
1072 disturbance history drive biocrust biodiversity in Western Montana rangelands, *Plant and Soil*, 430, 151-  
1073 169, 10.1007/s11104-018-3725-3, 2018.

1074 Elbert, W., Weber, B., Burrows, S., Steinkamp, J., Buedel, B., Andreae, M. O., and Poeschl, U.:  
1075 Contribution of cryptogamic covers to the global cycles of carbon and nitrogen, *Nat. Geosci.*, 5, 459-462,  
1076 10.1038/ngeo1486, 2012.

1077 Eldridge, D. J. and Delgado-Baquerizo, M.: The influence of climatic legacies on the distribution of  
1078 dryland biocrust communities, *Global Change Biology*, 25, 327-336, 10.1111/gcb.14506, 2019.

1079 Eldridge, D. J. and Tozer, M. E.: Environmental factors relating to the distribution of terricolous  
1080 bryophytes and lichens in semi-arid eastern Australia, *Bryologist*, 100, 28-39, 1997.

1081 Eldridge, D. J., Reed, S., Travers, S. K., Bowker, M. A., Maestre, F. T., Ding, J., Havrilla, C., Rodriguez-  
1082 Caballero, E., Barger, N., Weber, B., Antoninka, A., Belnap, J., Chaudhary, B., Faist, A., Ferrenberg, S.,  
1083 Huber-Sannwald, E., Malam Issa, O., and Zhao, Y.: The pervasive and multifaceted influence of biocrusts  
1084 on water in the world's drylands, *Global Change Biology*, 26, 6003-6014, 10.1111/gcb.15232, 2020.

1085 Eldridge, D. J., Guirado, E., Reich, P. B., Ochoa-Hueso, R., Berdugo, M., Sáez-Sandino, T., Blanco-  
1086 Pastor, J. L., Tedersoo, L., Plaza, C., Ding, J., Sun, W., Mamet, S., Cui, H., He, J.-Z., Hu, H.-W., Sokoya,  
1087 B., Abades, S., Alfaro, F., Bamigboye, A. R., Bastida, F., de los Ríos, A., Durán, J., Gaitan, J. J., Guerra,  
1088 C. A., Grebenc, T., Illán, J. G., Liu, Y.-R., Makhalyane, T. P., Mallen-Cooper, M., Molina-Montenegro,  
1089 M. A., Moreno, J. L., Nahberger, T. U., Peñaloza-Bojacá, G. F., Picó, S., Rey, A., Rodríguez, A., Siebe,  
1090 C., Teixido, A. L., Torres-Díaz, C., Trivedi, P., Wang, J., Wang, L., Wang, J., Yang, T., Zaady, E., Zhou,  
1091 X., Zhou, X.-Q., Zhou, G., Liu, S., and Delgado-Baquerizo, M.: The global contribution of soil mosses  
1092 to ecosystem services, *Nat. Geosci.*, 10.1038/s41561-023-01170-x, 2023.

1093 Engel, T., Bruehlheide, H., Hoss, D., Sabatini, F. M., Altman, J., Arfin-Khan, M. A. S., Bergmeier, E.,  
1094 Černý, T., Chytrý, M., Dainese, M., Dengler, J., Dolezal, J., Field, R., Fischer, F. M., Huygens, D., Jandt,  
1095 U., Jansen, F., Jentsch, A., Karger, D. N., Kattge, J., Lenoir, J., Lens, F., Loos, J., Niinemets, Ü., Overbeck,  
1096 G. E., Ozinga, W. A., Penuelas, J., Peyre, G., Phillips, O., Reich, P. B., Römermann, C., Sandel, B.,  
1097 Schmidt, M., Schrod, F., Velez-Martin, E., Violle, C., and Pillar, V.: Traits of dominant plant species  
1098 drive normalized difference vegetation index in grasslands globally, *Global Ecology and Biogeography*,  
1099 32, 695-706, 10.1111/gcb.13644, 2023.

1100 Fang, S., Yu, W., and Qi, Y.: Spectra and vegetation index variations in moss soil crust in different seasons,

1101 and in wet and dry conditions, *International Journal of Applied Earth Observation and Geoinformation*,  
1102 38, 261-266, 10.1016/j.jag.2015.01.018, 2015.

1103 Faticchi, S., Pappas, C., Zscheischler, J., and Leuzinger, S.: Modelling carbon sources and sinks in  
1104 terrestrial vegetation, *New Phytologist*, 221, 652-668, 10.1111/nph.15451, 2019.

1105 Ferrenberg, S., Reed, S. C., and Belnap, J.: Climate change and physical disturbance cause similar  
1106 community shifts in biological soil crusts, *Proceedings of the National Academy of Sciences of the*  
1107 *United States of America*, 112, 12116-12121, 10.1073/pnas.1509150112, 2015.

1108 Fischer, T. and Subbotina, M.: Climatic and soil texture threshold values for cryptogamic cover  
1109 development: a meta analysis, *Biologia*, 69, 1520-1530, 10.2478/s11756-014-0464-7, 2014.

1110 Gabay, T., Rotem, G., Gillor, O., and Ziv, Y.: Understanding changes in biocrust communities following  
1111 phosphate mining in the Negev Desert, *Environmental Research*, 207, 10.1016/j.envres.2021.112200,  
1112 2022.

1113 Gao, L., Bowker, M. A., Xu, M., Sun, H., Tuo, D., and Zhao, Y.: Biological soil crusts decrease erodibility  
1114 by modifying inherent soil properties on the Loess Plateau, China, *Soil Biology & Biochemistry*, 105,  
1115 49-58, 10.1016/j.soilbio.2016.11.009, 2017.

1116 García-Pichel, F., Loza, V., Marusenko, Y., Mateo, P., and Potrafka, R. M.: Temperature Drives the  
1117 Continental-Scale Distribution of Key Microbes in Topsoil Communities, *Science*, 340, 1574-1577,  
1118 10.1126/science.1236404, 2013.

1119 García-Roselló, E., Guisande, C., Manjarrés-Hernández, A., González-Dacosta, J., Heine, J., Pelayo-  
1120 Villamil, P., González-Vilas, L., Vari, R. P., Vaamonde, A., Granado-Lorencio, C., and Lobo, J. M.: Can  
1121 we derive macroecological patterns from primary Global Biodiversity Information Facility data?, *Global*  
1122 *Ecology and Biogeography*, 24, 335-347, 10.1111/geb.12260, 2015.

1123 Gassmann, F., Klotzli, F., and Walther, G. R.: Simulation of observed types of dynamics of plants and  
1124 plant communities, *Journal of Vegetation Science*, 11, 397-408, 10.2307/3236632, 2000.

1125 Havrilla, C. A. and Barger, N. N.: Biocrusts and their disturbance mediate the recruitment of native and  
1126 exotic grasses from a hot desert ecosystem, *Ecosphere*, 9, e02361, 10.1002/ecs2.2361, 2018.

1127 Havrilla, C. A., Chaudhary, V. B., Ferrenberg, S., Antoninka, A. J., Belnap, J., Bowker, M. A., Eldridge,  
1128 D. J., Faist, A. M., Huber-Sannwald, E., Leslie, A. D., Rodriguez-Caballero, E., Zhang, Y., Barger, N.  
1129 N., and Vries, F.: Towards a predictive framework for biocrust mediation of plant performance: A meta-  
1130 analysis, *Journal of Ecology*, 107, 2789-2807, 10.1111/1365-2745.13269, 2019.

1131 Hu, P., Zhang, W., Xiao, L., Yang, R., Xiao, D., Zhao, J., Wang, W., Chen, H., and Wang, K.: Moss-  
1132 dominated biological soil crusts modulate soil nitrogen following vegetation restoration in a subtropical  
1133 karst region, *Geoderma*, 352, 70-79, 10.1016/j.geoderma.2019.05.047, 2019.

1134 Jia, R., Chen, N., Yu, K., and Zhao, C.: High rainfall frequency promotes the dominance of biocrust under  
1135 low annual rainfall, *Plant and Soil*, 435, 257-275, 10.1007/s11104-018-3880-6, 2019.

1136 Jiménez-Valverde, A., Lobo, J. M., and Hortal, J.: Not as good as they seem: the importance of concepts  
1137 in species distribution modelling, *Diversity and Distributions*, 14, 885-890, 10.1111/j.1472-  
1138 4642.2008.00496.x, 2008.

1139 Karnieli, A.: Development and implementation of spectral crust index over dune sands, *International*  
1140 *Journal of Remote Sensing*, 18, 1207-1220, 10.1080/014311697218368, 1997.

1141 Karnieli, A., Kidron, G. J., Glaesser, C., and Ben-Dor, E.: Spectral characteristics of cyanobacteria soil  
1142 crust in semiarid environments, *Remote Sensing of Environment*, 69, 67-75, 10.1016/s0034-  
1143 4257(98)00110-2, 1999.

1144 Kidron, G. J.: Biocrust research: A critical view on eight common hydrological-related paradigms and



1145 dubious theses, *Ecohydrology*, 12, e2061, 10.1002/eco.2061, 2018.

1146 Kidron, G. J.: The enigmatic absence of cyanobacterial biocrusts from the Namib fog belt: Do dew and  
1147 fog hold the key?, *Flora*, 257, 151416, 10.1016/j.flora.2019.06.002, 2019.

1148 Kidron, G. J. and Xiao, B.: A false paradigm? Do biocrust types necessarily reflect ‘successional stages’?,  
1149 *Ecohydrology*, 17, 10.1002/eco.2610, 2023.

1150 Kidron, G. J., Lichner, L., Fischer, T., Starinsky, A., and Or, D.: Mechanisms for biocrust-modulated  
1151 runoff generation – A review, *Earth-Science Reviews*, 231, 104100, 10.1016/j.earscirev.2022.104100,  
1152 2022.

1153 Kropfl, A. I., Distel, R. A., Cecchi, G. A., and Villasuso, N. M.: Functional Role of Moss Biocrust in  
1154 Disturbed Semi-arid Shrublands of North-Eastern Patagonia, Argentina, *Applied Ecology and  
1155 Environmental Research*, 20, 905-917, 10.15666/aeer/2001\_905917, 2022.

1156 Lenton, T. M., Dahl, T. W., Daines, S. J., Mills, B. J., Ozaki, K., Saltzman, M. R., and Porada, P.: Earliest  
1157 land plants created modern levels of atmospheric oxygen, *Proceedings of the National Academy of  
1158 Sciences of the United States of America*, 113, 9704-9709, 10.1073/pnas.1604787113, 2016.

1159 Li, S., Bowker, M. A., and Xiao, B.: Biocrusts enhance non-rainfall water deposition and alter its  
1160 distribution in dryland soils, *Journal of Hydrology*, 595, 126050, 10.1016/j.jhydrol.2021.126050, 2021.

1161 Li, X., Sun, J., Zhang, H., Tan, H., Hui, R., Qi, J., Zhang, P., and Ward, N. D.: Warming decreases desert  
1162 ecosystem functioning by altering biocrusts in drylands, *Journal of Applied Ecology*, 10.1111/1365-  
1163 2664.14528, 2023.

1164 Ma, X., Zhao, Y., Yang, K., Ming, J., Qiao, Y., Xu, M., and Pan, X.: Long-term light grazing does not  
1165 change soil organic carbon stability and stock in biocrust layer in the hilly regions of drylands, *Journal  
1166 of Arid Land*, 15, 940-959, 10.1007/s40333-023-0064-x, 2023.

1167 Machado de Lima, N. M., Muñoz-Rojas, M., Vázquez-Campos, X., and Branco, L. H. Z.: Biocrust  
1168 cyanobacterial composition, diversity, and environmental drivers in two contrasting climatic regions in  
1169 Brazil, *Geoderma*, 386, 114914, 10.1016/j.geoderma.2020.114914, 2021.

1170 Maestre, F. T., Benito, B. M., Berdugo, M., Concostrina-Zubiri, L., Delgado-Baquerizo, M., Eldridge, D.  
1171 J., Guirado, E., Gross, N., Kefi, S., Le Bagousse-Pinguet, Y., Ochoa-Hueso, R., and Soliveres, S.:  
1172 Biogeography of global drylands, *New Phytologist*, 231, 540-558, 10.1111/nph.17395, 2021.

1173 Mäkinen, J., Numminen, E., Niittynen, P., Luoto, M., and Vanhatalo, J.: Spatial confounding in Bayesian  
1174 species distribution modeling, *Ecography*, 2022, e06183, 10.1111/ecog.06183, 2022.

1175 Marsh, J., Nouvet, S., Sanborn, P., and Coxson, D.: Composition and function of biological soil crust  
1176 communities along topographic gradients in grasslands of central interior British Columbia (Chilcotin)  
1177 and southwestern Yukon (Kluane), *Canadian Journal of Botany*, 84, 717-736, 10.1139/b06-026, 2006.

1178 McCann, E., Reed, S. C., Saud, P., Reibold, R. H., Howell, A., and Faist, A. M.: Plant growth and  
1179 biocrust-fire interactions across five North American deserts, *Geoderma*, 401, 115325,  
1180 10.1016/j.geoderma.2021.115325, 2021.

1181 McCune, B., Yang, S., Jovan, S., and Root, H. T.: Climate and epiphytic macrolichen communities in the  
1182 Four Corners region of the USA, *Bryologist*, 125, 70-90, 10.1639/0007-2745-125.1.070, 2022.

1183 Ming, J., Zhao, Y., Wu, Q., He, H., and Gao, L.: Soil temperature dynamics and freezing processes for  
1184 biocrustal soils in frozen soil regions on the Qinghai–Tibet Plateau, *Geoderma*, 409, 115655,  
1185 10.1016/j.geoderma.2021.115655, 2022.

1186 Miranda-González, R. and McCune, B.: The weight of the crust: Biomass of crustose lichens in tropical  
1187 dry forest represents more than half of foliar biomass, *Biotropica*, 52, 1298-1308, 10.1111/btp.12837,  
1188 2020.

1189 Morillas, L. and Gallardo, A.: Biological soil crusts and wetting events: Effects on soil N and C cycles,  
1190 *Applied Soil Ecology*, 94, 1-6, 10.1016/j.apsoil.2015.04.015, 2015.

1191 Munoz-Martin, M. A., Becerra-Absalon, I., Perona, E., Fernandez-Valbuena, L., Garcia-Pichel, F., and  
1192 Mateo, P.: Cyanobacterial biocrust diversity in Mediterranean ecosystems along a latitudinal and climatic  
1193 gradient, *New Phytologist*, 221, 123-141, 10.1111/nph.15355, 2019.

1194 Noy, K., Ohana-Levi, N., Panov, N., Silver, M., and Karnieli, A.: A long-term spatiotemporal analysis of  
1195 biocrusts across a diverse arid environment: The case of the Israeli-Egyptian sandfield, *Science of The  
1196 Total Environment*, 774, 145154, 10.1016/j.scitotenv.2021.145154, 2021.

1197 Oliva, G., Dos Santos, E., Sofia, O., Umana, F., Massara, V., Garcia Martinez, G., Caruso, C., Cariac, G.,  
1198 Echevarria, D., Fantozzi, A., Butti, L., Bran, D., Gaitan, J., Ferrante, D., Paredes, P., Dominguez, E., and  
1199 Maestre, F. T.: The MARAS dataset, vegetation and soil characteristics of dryland rangelands across  
1200 Patagonia, *Scientific Data*, 7, 327, 10.1038/s41597-020-00658-0, 2020.

1201 Palmer, B., Hernandez, R., and Lipson, D. A.: The fate of biological soil crusts after fire: A meta-analysis,  
1202 *Global Ecology and Conservation*, 24, e01380, 10.1016/j.gecco.2020.e01380, 2020.

1203 Palmer, B., Lawson, D., and Lipson, D. A.: Years After a Fire, Biocrust Microbial Communities are  
1204 Similar to Unburned Communities in a Coastal Grassland, *Microbial Ecology*, 10.1007/s00248-022-  
1205 02137-y, 2022.

1206 Panigada, C., Tagliabue, G., Zaady, E., Rozenstein, O., Garzonio, R., Di Mauro, B., De Amicis, M.,  
1207 Colombo, R., Cogliati, S., Miglietta, F., and Rossini, M.: A new approach for biocrust and vegetation  
1208 monitoring in drylands using multi-temporal Sentinel-2 images, *Progress in Physical Geography: Earth  
1209 and Environment*, 43, 496-520, 10.1177/0309133319841903, 2019.

1210 Pearce, J. L., Cherry, K., M, D., S, F., and Whish, G.: Incorporating expert opinion and fine-scale  
1211 vegetation mapping into statistical models of faunal distribution, *Journal of Applied Ecology*, 38, 412-  
1212 424, 10.1046/j.1365-2664.2001.00608.x, 2001.

1213 Perry, G. L. W., Seidl, R., Bellvé, A. M., and Rammer, W.: An Outlook for Deep Learning in Ecosystem  
1214 Science, *Ecosystems*, 25, 1700-1718, 10.1007/s10021-022-00789-y, 2022.

1215 Porada, P., Pöschl, U., Kleidon, A., Beer, C., and Weber, B.: Estimating global nitrous oxide emissions  
1216 by lichens and bryophytes with a process-based productivity model, *Biogeosciences*, 14, 1593-1602,  
1217 10.5194/bg-14-1593-2017, 2017.

1218 Porada, P., Weber, B., Elbert, W., Pöschl, U., and Kleidon, A.: Estimating global carbon uptake by lichens  
1219 and bryophytes with a process-based model, *Biogeosciences*, 10, 6989-7033, 10.5194/bg-10-6989-2013,  
1220 2013.

1221 Porada, P., Tamm, A., Raggio, J., Cheng, Y., Kleidon, A., Pöschl, U., and Weber, B.: Global NO and  
1222 HONO emissions of biological soil crusts estimated by a process-based non-vascular vegetation model,  
1223 *Biogeosciences*, 16, 2003-2031, 10.5194/bg-16-2003-2019, 2019.

1224 Pravalie, R.: Drylands extent and environmental issues. A global approach, *Earth-Science Reviews*, 161,  
1225 259-278, 10.1016/j.earscirev.2016.08.003, 2016.

1226 Pushkareva, E., Johansen, J. R., and Elster, J.: A review of the ecology, ecophysiology and biodiversity  
1227 of microalgae in Arctic soil crusts, *Polar Biology*, 39, 2227-2240, 10.1007/s00300-016-1902-5, 2016.

1228 Qiu, D., Bowker, M. A., Xiao, B., Zhao, Y., Zhou, X., and Li, X.: Mapping biocrust distribution in China's  
1229 drylands under changing climate, *Science of The Total Environment*, 905,  
1230 10.1016/j.scitotenv.2023.167211, 2023.

1231 Quillet, A., Peng, C., and Garneau, M.: Toward dynamic global vegetation models for simulating  
1232 vegetation-climate interactions and feedbacks: recent developments, limitations, and future challenges,

1233 Environmental Reviews, 18, 333-353, 10.1139/a10-016, 2010.

1234 Read, C. F., Duncan, D. H., Vesk, P. A., Elith, J., and Wan, S.: Biocrust morphogroups provide an  
1235 effective and rapid assessment tool for drylands, *Journal of Applied Ecology*, 51, 1740-1749,  
1236 10.1111/1365-2664.12336, 2014.

1237 Reed, S. C., Coc, K. K., Sparks, J. P., Housman, D. C., Zelikova, T. J., and Belnap, J.: Changes to dryland  
1238 rainfall result in rapid moss mortality and altered soil fertility, *Nature Climate Change*, 2, 752-755,  
1239 10.1038/nclimate1596, 2012.

1240 Rodríguez-Caballero, E., Knerr, T., and Weber, B.: Importance of biocrusts in dryland monitoring using  
1241 spectral indices, *Remote Sensing of Environment*, 170, 32-39, 10.1016/j.rse.2015.08.034, 2015.

1242 Rodríguez-Caballero, E., Belnap, J., Budel, B., Crutzen, P. J., Andreae, M. O., Poschl, U., and Weber, B.:  
1243 Dryland photoautotrophic soil surface communities endangered by global change, *Nat. Geosci.*, 11, 185-  
1244 189, 10.1038/s41561-018-0072-1, 2018.

1245 Rodríguez-Caballero, E., Stanelle, T., Egerer, S., Cheng, Y., Su, H., Canton, Y., Belnap, J., Andreae, M.  
1246 O., Tegen, I., and Reick, C. H.: Global cycling and climate effects of aeolian dust controlled by biological  
1247 soil crusts, *Nat. Geosci.*, 15, 458-463, 10.1038/s41561-022-00942-1, 2022.

1248 Rodríguez-Caballero, E., Escribano, P., Olehowski, C., Chamizo, S., Hill, J., Cantón, Y., and Weber, B.:  
1249 Transferability of multi- and hyperspectral optical biocrust indices, *ISPRS Journal of Photogrammetry  
1250 and Remote Sensing*, 126, 94-107, 10.1016/j.isprsjprs.2017.02.007, 2017.

1251 Rodríguez-Caballero, E., Reyes, A., Kratz, A., Caesar, J., Guirado, E., Schmiedel, U., Escribano, P.,  
1252 Fiedler, S., and Weber, B.: Effects of climate change and land use intensification on regional biological  
1253 soil crust cover and composition in southern Africa, *Geoderma*, 406, 115508,  
1254 10.1016/j.geoderma.2021.115508, 2022.

1255 Root, H. T. and McCune, B.: Regional patterns of biological soil crust lichen species composition related  
1256 to vegetation, soils, and climate in Oregon, USA, *Journal of Arid Environments*, 79, 93-100,  
1257 10.1016/j.jaridenv.2011.11.017, 2012.

1258 Sabatini, F. M., Lenoir, J., Hattab, T., Arnst, E. A., Chytrý, M., Dengler, J., De Ruffray, P., Hennekens, S.  
1259 M., Jandt, U., Jansen, F., Jiménez-Alfaro, B., Kattge, J., Levesley, A., Pillar, V. D., Purschke, O., Sandel,  
1260 B., Sultana, F., Aavik, T., Acic, S., Acosta, A. T. R., Agrillo, E., Alvarez, M., Apostolova, I., Arfin Khan,  
1261 M. A. S., Arroyo, L., Attorre, F., Aubin, I., Banerjee, A., Bauters, M., Bergeron, Y., Bergmeier, E., Biurrun,  
1262 I., Bjorkman, A. D., Bonari, G., Bondareva, V., Brunet, J., Čarni, A., Casella, L., Cayuela, L., Černý, T.,  
1263 Chepinoga, V., Csiky, J., Čušterevska, R., De Bie, E., Gasper, A. L., De Sanctis, M., Dimopoulos, P.,  
1264 Dolezal, J., Dziuba, T., El-Sheikh, M. A. E. R. M., Enquist, B., Ewald, J., Fazayeli, F., Field, R., Finckh,  
1265 M., Gachet, S., Galán-de-Mera, A., Garbolino, E., Gholizadeh, H., Giorgis, M., Golub, V., Alsos, I. G.,  
1266 Grytnes, J. A., Guerin, G. R., Gutiérrez, A. G., Haider, S., Hatim, M. Z., Hérault, B., Hinojos Mendoza,  
1267 G., Hölzel, N., Homeier, J., Hubau, W., Indreica, A., Janssen, J. A. M., Jedrzejek, B., Jentsch, A., Jürgens,  
1268 N., Kaçki, Z., Kapfer, J., Karger, D. N., Kavgaci, A., Kearsley, E., Kessler, M., Khanina, L., Killeen, T.,  
1269 Korolyuk, A., Kreft, H., Kühn, H. S., Kuzemko, A., Landucci, F., Lengyel, A., Lens, F., Lingner, D. V.,  
1270 Liu, H., Lysenko, T., Mahecha, M. D., Marcenò, C., Martynenko, V., Moeslund, J. E., Monteagudo  
1271 Mendoza, A., Mucina, L., Müller, J. V., Munzinger, J., Naqinezhad, A., Noroozi, J., Nowak, A.,  
1272 Onyshchenko, V., Overbeck, G. E., Pärtel, M., Pauchard, A., Peet, R. K., Peñuelas, J., Pérez-Haase, A.,  
1273 Peterka, T., Petřík, P., Peyre, G., Phillips, O. L., Prokhorov, V., Rašomavičius, V., Revermann, R., Rivas-  
1274 Torres, G., Rodwell, J. S., Ruprecht, E., Rüşiņa, S., Samimi, C., Schmidt, M., Schrod, F., Shan, H.,  
1275 Shirokikh, P., Šibík, J., Šilc, U., Sklenář, P., Škvorec, Ž., Sparrow, B., Sperandii, M. G., Stančić, Z.,  
1276 Svenning, J. C., Tang, Z., Tang, C. Q., Tsiripidis, I., Vanselow, K. A., Vázquez Martínez, R., Vassilev, K.,

1277 Vélez-Martin, E., Venanzoni, R., Vibrans, A. C., Violle, C., Virtanen, R., Wehrden, H., Wagner, V.,  
1278 Walker, D. A., Waller, D. M., Wang, H. F., Wesche, K., Whitfeld, T. J. S., Willner, W., Wiser, S. K.,  
1279 Wohlgenuth, T., Yamalov, S., Zobel, M., Bruelheide, H., and Bates, A.: sPlotOpen – An  
1280 environmentally balanced, open - access, global dataset of vegetation plots, *Global Ecology and*  
1281 *Biogeography*, 30, 1740-1764, 10.1111/geb.13346, 2021.

1282 Shi, W., Pan, Y.-x., Zhang, Y.-f., Hu, R., and Wang, X.-p.: The effect of different biocrusts on soil  
1283 hydraulic properties in the Tengger Desert, China, *Geoderma*, 430, 116304,  
1284 10.1016/j.geoderma.2022.116304, 2023.

1285 Skidmore, A. K., Franklin, J., Dawson, T. P., and Pilesjö, P.: Geospatial tools address emerging issues in  
1286 spatial ecology: a review and commentary on the Special Issue, *International Journal of Geographical*  
1287 *Information Science*, 25, 337-365, 10.1080/13658816.2011.554296, 2011.

1288 Soberon, J. and Nakamura, M.: Niches and distributional areas: Concepts, methods, and assumptions,  
1289 *Proceedings of the National Academy of Sciences of the United States of America*, 106, 19644-19650,  
1290 2009.

1291 Song, G., Hui, R., Yang, H., Wang, B., and Li, X.: Biocrusts mediate the plant community composition  
1292 of dryland restoration ecosystems, *Science of the Total Environment*, 844, 157135,  
1293 10.1016/j.scitotenv.2022.157135, 2022.

1294 Sun, F., Xiao, B., Kidron, G. J., and Tuller, M.: Towards the effects of moss-dominated biocrusts on  
1295 surface soil aeration in drylands: Air permeability analysis and modeling, *Catena*, 223, 106942,  
1296 10.1016/j.catena.2023.106942, 2023.

1297 Sun, H., Ma, X., Liu, Y., Zhou, G. Y., Ding, J. L., Lu, L., Wang, T. J., Yang, Q. L., Shu, Q. T., and Zhang,  
1298 F.: A New Multiangle Method for Estimating Fractional Biocrust Coverage From Sentinel-2 Data in Arid  
1299 Areas, *Ieee Transactions on Geoscience and Remote Sensing*, 62, 10.1109/tgrs.2024.3361249, 2024.

1300 Thonicke, K., Venevsky, S., Sitch, S., and Cramer, W.: The role of fire disturbance for global vegetation  
1301 dynamics: coupling fire into a Dynamic Global Vegetation Model, *Global Ecology and Biogeography*,  
1302 10, 661-677, 10.1046/j.1466-822x.2001.00175.x, 2001.

1303 Tucker, C. L., McHugh, T. A., Howell, A., Gill, R., Weber, B., Belnap, J., Grote, E., and Reed, S. C.: The  
1304 concurrent use of novel soil surface microclimate measurements to evaluate CO<sub>2</sub> pulses in biocrusted  
1305 interspaces in a cool desert ecosystem, *Biogeochemistry*, 135, 239-249, 10.1007/s10533-017-0372-3,  
1306 2017.

1307 Velasco Ayuso, S., Oñatibia, G. R., Maestre, F. T., and Yahdjian, L.: Grazing pressure interacts with  
1308 aridity to determine the development and diversity of biological soil crusts in Patagonian rangelands,  
1309 *Land Degradation & Development*, 31, 488-499, 10.1002/ldr.3465, 2019.

1310 Wang, S., Liu, B., Zhao, Y., Gao, L., Yin, B., Yang, K., and Ji, J.: Determination of the representative  
1311 elementary area (REA) of biocrusts: A case study from the Hilly Loess Plateau region, China, *Geoderma*,  
1312 406, 115502, 10.1016/j.geoderma.2021.115502, 2022a.

1313 Wang, Y., Xiao, B., Wang, W., and Kidron, G. J.: Interactions between biocrusts and herbaceous  
1314 communities are divergent in dry and wet semiarid ecosystems, *Science of The Total Environment*, 941,  
1315 173759, 10.1016/j.scitotenv.2024.173759, 2024.

1316 Wang, Z., Wu, B., Zhang, M., Zeng, H., Yang, L., Tian, F., Ma, Z., and Wu, H.: Indices enhance biological  
1317 soil crust mapping in sandy and desert lands, *Remote Sensing of Environment*, 278, 113078,  
1318 10.1016/j.rse.2022.113078, 2022b.

1319 Weber, B. and Hill, J.: Remote Sensing of Biological Soil Crusts at Different Scales, in: *Biological Soil*  
1320 *Crusts: An Organizing Principle in Drylands*, edited by: Weber, B., Büdel, B., and Belnap, J., Springer

1321 International Publishing, Cham, 215-234, 10.1007/978-3-319-30214-0\_12, 2016.

1322 Weber, B., Budel, B., and Belnap, J., Weber, B., Budel, B., and Belnap, J. (Eds.): *Biological Soil Crusts:*

1323 *An Organizing Principle in Drylands*, Springer Nature, 10.1007/978-3-319-30214-0, 2016.

1324 Weber, B., Olehowski, C., Knerr, T., Hill, J., Deuschewitz, K., Wessels, D. C. J., Eitel, B., and Buedel,

1325 B.: A new approach for mapping of Biological Soil Crusts in semidesert areas with hyperspectral imagery,

1326 *Remote Sensing of Environment*, 112, 2187-2201, 10.1016/j.rse.2007.09.014, 2008.

1327 Weber, B., Belnap, J., Budel, B., Antoninka, A. J., Barger, N. N., Chaudhary, V. B., Darrouzet-Nardi, A.,

1328 Eldridge, D. J., Faist, A. M., Ferrenberg, S., Havrilla, C. A., Huber-Sannwald, E., Malam Issa, O.,

1329 Maestre, F. T., Reed, S. C., Rodriguez-Caballero, E., Tucker, C., Young, K. E., Zhang, Y., Zhao, Y., Zhou,

1330 X., and Bowker, M. A.: What is a biocrust? A refined, contemporary definition for a broadening research

1331 community, *Biological Reviews*, 97, 1768-1785, 10.1111/brv.12862, 2022.

1332 Wei, X., Qin, F., Han, B., Zhou, H., Liu, M., and Shao, X.: Spatial variations of bacterial communities

1333 associated with biological soil crusts along a climatic gradient in alpine grassland ecosystems, *Plant and*

1334 *Soil*, 480, 493-506, 10.1007/s11104-022-05595-y, 2022.

1335 Whitney, K. M., Vivoni, E. R., Duniway, M. C., Bradford, J. B., Reed, S. C., and Belnap, J.:

1336 Ecohydrological role of biological soil crusts across a gradient in levels of development, *Ecohydrology*,

1337 10, e1875, 10.1002/eco.1875, 2017.

1338 Williams, A. J., Buck, B. J., Soukup, D. A., and Merkler, D. J.: Geomorphic controls on biological soil

1339 crust distribution: A conceptual model from the Mojave Desert (USA), *Geomorphology*, 195, 99-109,

1340 10.1016/j.geomorph.2013.04.031, 2013.

1341 Wolf, S., Mahecha, M. D., Sabatini, F. M., Wirth, C., Bruelheide, H., Kattge, J., Moreno Martinez, A.,

1342 Mora, K., and Kattenborn, T.: Citizen science plant observations encode global trait patterns, *Nature*

1343 *Ecology & Evolution*, 6, 1850-1859, 10.1038/s41559-022-01904-x, 2022.

1344 Wright, A. J. and Collins, S. L.: Drought experiments need to incorporate atmospheric drying to better

1345 simulate climate change, *BioScience*, 74, 65-71, 10.1093/biosci/biad105, 2024.

1346 Yang, H., Liu, C., Liu, Y., and Xing, Z.: Impact of human trampling on biological soil crusts determined

1347 by soil microbial biomass, enzyme activities and nematode communities in a desert ecosystem, *European*

1348 *Journal of Soil Biology*, 87, 61-71, 10.1016/j.ejsobi.2018.05.005, 2018.

1349 Yu, H., Cooper, A. R., and Infante, D. M.: Improving species distribution model predictive accuracy

1350 using species abundance: Application with boosted regression trees, *Ecological Modelling*, 432, 109202,

1351 10.1016/j.ecolmodel.2020.109202, 2020.

1352 Zhang, Y. M., Chen, J., Wang, L., Wang, X. Q., and Gu, Z. H.: The spatial distribution patterns of

1353 biological soil crusts in the Gurbantunggut Desert, Northern Xinjiang, China, *Journal of Arid*

1354 *Environments*, 68, 599-610, 10.1016/j.jaridenv.2006.06.012, 2007.

1355 Zhao, Y., Qin, N., Weber, B., and Xu, M.: Response of biological soil crusts to raindrop erosivity and

1356 underlying influences in the hilly Loess Plateau region, China, *Biodiversity and Conservation*, 23, 1669-

1357 1686, 10.1007/s10531-014-0680-z, 2014.

1358 Zhao, Y., Ji, J., Zhang, W., Ming, J., Huang, W., and Gao, L.: Characteristics of spatial and temporal

1359 variability in the distribution of biological soil crusts on the Loess Plateau, China, *Chinese Journal of*

1360 *Applied Ecology*, 35, 739-748, 2024.

1361