

The organisation of subglacial drainage during the demise of the Finnish Lake District Ice-Lobe

Adam J. Hepburn^{1,2*}, Christine F. Dow¹, Antti Ojala³, Joni Mäkinen³, Elina Ahokangas³, Jussi Hovikoski⁴, Jukka-Pekka Palmu⁴, and Kari Kajuutti³

¹Department of Geography and Environmental Management, University of Waterloo, Waterloo, ON, Canada

²European Space Astronomy Centre, European Space Agency, Madrid, Spain

³Department of Geography and Geology, University of Turku, Turku, Finland

⁴Geological Survey of Finland, Espoo, Finland

Correspondence: Adam J. Hepburn, now at Aberystwyth University, Aberystwyth, Wales, UK (adam.hepburn@aber.ac.uk)

Abstract. Unknown basal characteristics limit our ability to simulate the subglacial hydrology of rapidly melting contemporary ice sheets. Sediment-based landforms generated beneath Late Pleistocene ice sheets, together with detailed digital elevation models, offer a valuable means of testing basal hydrology models, which describe the flow and dynamics of water in the subglacial system. However, to date no work has evaluated how well process-based subglacial hydrology models represent the hypothesised conditions associated with glaciofluvial landform formation in the palaeo setting. Previous work comparing model output to geomorphological evidence has typically done so using models which do not resolve subglacial processes and instead express likely subglacial water pathways. Here, we explore the ability of the Glacier Drainage System model (GlaDS), a process based subglacial hydrology model, to represent the genesis conditions associated with a specific glaciofluvial landform termed ‘murtoos’. Distinctive triangular landforms found throughout Finland and Sweden, murtoos are hypothesised to form 40–60 km from the former Fennoscandian Ice Sheet margin within a ‘semi-distributed’ system at the onset of channelised drainage, in small cavities where water pressure is equal to or exceeds ice overburden pressure. Concentrating within a specific ice lobe of the former Fennoscandian Ice Sheet and using digital elevation models with a simulated former ice surface geometry, we forced GlaDS with transient surface melt and explored the sensitivity of our model outcomes to parameter decisions such as the system conductivity and bed topography. Our model outputs closely match the general spacing, direction and complexity of eskers and mapped assemblages of features related to subglacial drainage in ‘meltwater routes’. Many of the predictions for murtoo formation are produced by the model, including the location of water pressure equal to ice overburden, the onset of channelised drainage, the transition in drainage modes, and importantly, the seasonal sequence of drainage conditions inferred from murtoo sedimentology. These conclusions are largely robust to a range of parameter decisions and we explore seasonal and inter-annual drainage behaviour associated with murtoo zones and meltwater pathways. Our results demonstrate that examining palaeo basal topography alongside subglacial hydrology model outputs holds promise for the mutually beneficial analyses of palaeo and contemporary ice sheets to assess the controls of hydrology on ice dynamics and subglacial landform evolution.

1 Introduction

The changing configuration of the basal hydrological system beneath ice sheets throughout the melt season is primarily responsible for modulating the response of ice flow to meltwater input (Schoof, 2010). Subglacial water is typically conceptualised as being routed through either *distributed*, inefficient, and high water pressure systems (Weertman, 1972; Kamb, 1987; Boulton and Jones, 1979), or *channelised*, efficient, and lower water pressure systems (Nye, 1972; Röthlisberger, 1972; Hooke, 1989), transitioning between the two as a function of discharge (Schoof, 2010). The seasonal delivery of meltwater to the bed of ice sheets, and transition of subglacial hydrological systems in response, exerts a first order control on ice flow by modifying the frictional resistance to ice flow (Schoof, 2010). Understanding where and when transitions between distributed and channelised drainage occur beneath ice sheets is critical if such processes are going to be faithfully represented in ice sheet models used to predict the rate and timing of ice sheet mass loss (Andrews et al., 2014; Nienow et al., 2017), particularly in response to more widespread and prolonged atmospheric warming (e.g., van den Broeke et al., 2023).

To date, most analyses of seasonal subglacial hydrological development have been applied to contemporary ice sheets and glaciers. However, these sites lack key information about basal characteristics, such as basal topography, underlying sedimentology, and the hydraulic properties of the subsurface material (Chu, 2014). Subglacial hydrology models are often used to analyse these systems at the catchment scale but, given the absence of more detailed information, basal characteristics are often reduced to parameterisations, or simplifications, of what is likely a more complex reality (e.g., Schoof, 2010; Werder et al., 2013; Flowers, 2018; Kazmierczak et al., 2022).

Glaciated Late Pleistocene terrains may provide a valuable insight into the subglacial hydrological processes operating beneath ice sheets (Stokes et al., 2015; Greenwood et al., 2016). Numerical ice sheet models are already evaluated against the rich geomorphological record that Late Pleistocene ice sheets left behind, particularly landforms relating to ice flow direction or ice marginal position (e.g., Boulton and Hagdorn, 2006; Kleman et al., 2006; Tarasov et al., 2012; Gandy et al., 2019, 2021; Archer et al., 2023; García-Ruiz et al., 2023). Glaciofluvial landforms are especially common in the landform record (e.g., Clark and Walder, 1994; Cofaigh, 1996; Rampton, 2000; Utting et al., 2009; Coughlan et al., 2020; Dewald et al., 2021, 2022) and may represent ideal targets against which to evaluate subglacial hydrology models, potentially including processes variable at sub-annual scales and across the distributed–channelised transition (Kleman et al., 2006; Bingham et al., 2010; Stokes et al., 2015). However, landform genesis uncertainty arises from both fundamentally different concepts of how a landform is formed and the variable spatial and temporal scales of formation (Cofaigh, 1996; Stokes et al., 2015; Greenwood et al., 2016).

Previous work has largely used relatively simple models to subglacial hydrology to explore the spatial expression of channelised glaciofluvial landforms. These models often make assumptions about the configuration of the drainage system (Stokes et al., 2015) and do not explicitly resolve channel formation or exchanges between drainage systems. Typically, such models operate by prescribing a fixed water pressure at or near overburden everywhere, resulting in an expression of likely flow routing but not physical channel development or evolution (e.g., Livingstone et al., 2013a, b, 2015; Karlsson and Dahl-Jensen, 2015; Shackleton et al., 2018; Kirkham et al., 2022). Alternatively, in models where water pressure is allowed to vary, channels are assumed to form where water pressure is equal to ice overburden pressure but the process of channel formation is not explicitly

resolved (e.g., Boulton et al., 2007a, b, 2009). These models are unable to capture dynamic drainage changes, and are instead intended to represent long-term, interannual, ‘steady-state’ conditions (Banwell et al., 2013). More complex models have been used to address esker formation over shorter timescales in 1D (e.g., Beaud et al., 2018; Hewitt and Creyts, 2019) while 2D models which include exchanges between a distributed system and a single channel have been used to interrogate esker length and spacing scaling relationships (e.g., Hewitt, 2011) and erosion rates associated with subglacial drainage efficiency (e.g., Beaud et al., 2014). However, these more complex models use idealised parabolic ice surfaces, often with a flat bed. In contrast, modern subglacial hydrology models (i.e., those capable of resolving transitions between distributed and channelised drainage in both space and time) are widely applied to contemporary ice sheets (e.g., Flowers, 2018; Indrigo et al., 2021; Dow et al., 2022; Sommers et al., 2022; Ehrenfeucht et al., 2023). However, despite the critical need to evaluate and improve modern subglacial hydrology models using all available sources of data (Dow, 2023), we are not aware of previous work which has evaluated the ability of such models to reproduce the subglacial conditions (e.g., water pressure, channel location) associated with glaciofluvial landform formation.

In this paper we apply the Glacier Drainage System model (GlaDS Werder et al., 2013)—a modern subglacial hydrology model capable of resolving the transition between channelised and distributed drainage—to a palaeo ice sheet terrain. We evaluate the ability of GlaDS to represent the conditions giving rise to specific glaciofluvial landforms by making comparisons between model output and the spatial expression and predicted generation of ‘murtoos’ (singular: murtoo, Mäkinen et al., 2017; Ojala et al., 2019) recently identified in Fennoscandia. More widely described channelised features such as eskers (e.g., Storrar and Livingstone, 2017) and tunnel valleys (e.g., Kirkham et al., 2022, 2024) often exceed 10 km in length and likely represent time-transgressive formation over decades–millennia of ice-sheet margin retreat (Mäkinen, 2003). In contrast, murtoos are small (<100 m) glaciofluvial landforms thought to represent the spatiotemporal transition from distributed to channelised subglacial drainage over as little as one melt season (Hovikoski et al., 2023; Mäkinen et al., 2023). The size, formation rate, spatial distribution, and sedimentological architecture of murtoos provide a unique set of predictions against which a subglacial hydrology model can be tested (Hovikoski et al., 2023), including the location of a persistent area of high water pressure, the evolution of discharge through the year, and the spatial onset of channelised drainage. The aim of this paper is to explore the ability of GlaDS, a process based subglacial hydrology model, to explain murtoo formation in both space and time.

2 The glaciofluvial significance of murtoos

Clearly distinguishable from other glaciofluvial landforms, murtoos are small (30–100 m in width/length), low relief (~5 m high) features orientated parallel to ice flow with a distinctive, broadly triangular morphology (See Figure 1, & Ojala et al., 2019, 2021, 2022; Ahokangas et al., 2021; Peterson Becher and Johnson, 2021; Vérité et al., 2022; Van Boeckel et al., 2022). Mapping across Finland and Sweden (Mäkinen et al., 2017; Peterson et al., 2017; Ahokangas et al., 2021) reveals a preferential clustering of murtoos in swarms or fields (e.g., Figure 1B) along subglacial meltwater routes—integrated assemblages of multiple landforms associated with subglacial meltwater (Lewington et al., 2020; Ahokangas et al., 2021; Dewald et al., 2022). Subglacial meltwater routes containing murtoos, or ‘*murtoo routes*’ (Ahokangas et al., 2021) are concentrated in faster flowing,




Figure 1. The study area. **A)** The extent of the GlaDS model domain (solid line) in the Finnish Lake District Ice Lobe (FLDIL) and the Salpausselkä (dashed line). Salpausselkä II (Ss II) marks the 12 cal. ka ice extent. Murtoo fields identified by Ahokangas et al. (2021) within the FLDIL are shown as inverted triangles. **B)** A murtoo field. Adjacent to the murtoos, a large esker is visible in the lake. **C)** Murtoo fields amongst ribbed moraines in the north of the FLDIL. **D)** An abrupt downstream transition from murtoo fields to a large esker. Additional murtoo fields are found directly adjacent to the esker. **E)** Three murtoo fields amongst streamlined terrain within at the centre of the FLDIL. All panels show a multi-directional oblique weighted hillshade based on 2 m LiDAR data (see Ahokangas et al., 2021, for details). In panels B–E, black arrows in the upper left corner of each panel indicate the approximate ice flow direction and the inverted triangles are aligned with the orientation of murtoo fields.

warm-based sectors of the Fennoscandian ice sheet (FIS) and are often adjacent to, or downstream of, drumlin fields or ribbed moraines (e.g., V^{er}it^e et al., 2022, and Figure 1C). Murtoo routes are also often located upstream of, and appear to transition into, eskers (e.g., Ahokangas et al., 2021, and Figure 1D). Crucially, murtoo routes are rarely found closer than 40–60 km to the former FIS ice margins (Mäkinen et al., 2023), aligning well with the maximum observed length of channels (~50 km) in contemporary Greenland (e.g., Chandler et al., 2013, 2021; Dow et al., 2015).

The sedimentological sequence of a murtoo (as described in Hovikoski et al., 2023, and summarised in Table 1) is characterised by abrupt changes in sedimentary structure and grain size and charts the spatiotemporal transition from distributed to increasingly channelised flow within a single melt season during deglaciation (Mäkinen et al., 2023). Murtoos typically comprise a core unit containing sorted sediments, which develops at the end of meltwater pulses within a rapidly enlarging broad subglacial conduit. This core represents the first phase of murtoo formation and evidences at least partial ice contact and periodic deformation (Hovikoski et al., 2023). Following the onset of spring melt, pulses of water deposit a main body unit (murtoo developmental phase 2 in Table 1 and referred to as Unit 2 by Mäkinen et al., 2023) that i) distally is comprised of alternating facies of heterogeneous diamicton, with strong fabrics interbedded with sorted gravelly and sandy sediment and ii) proximally is comprised of alternating sequences of glaciofluvial deposits, with current ripples (formed in low discharge, lower flow regimes) giving way to transitional cross-bedding (transitional flow regimes), and antidunal sinusoidal lamination (formed in higher discharge, upper flow regimes; Hovikoski et al., 2023). The proximal transition from lower to upper flow regimes represents a rapid increase in water flow velocity and depth through a melt season and transition from inefficient distributed flow to the development of an enlarged, water-filled cavity. Boulder size-distributions suggests that this cavity reached a maximum flow space of 1 m (developmental phase 3, Hovikoski et al., 2023). The development of this enlarged cavity/pond and subsequent water pressure drop encourages localised creep closure at the broadest part of the murtoo (developmental phase 4), evidenced by a disappearance of sorted sediment (interpreted as non-deposition rather than erosion), and in some murtoos this is succeeded by compacted interbedded diamicton—indicating ice-bed recoupling (Hovikoski et al., 2023). Meanwhile, closer to the margins of the murtoo body, meltwater flow continues and is routed obliquely towards the tip, forming boulder-rich proto-channels. These deposits indicate that the ice-bed recoupling at the broadest part of the murtoo coincided with intense and increasingly erosional channelised flow at the murtoo margins. The final stage of murtoo development (developmental phase 5) is commonly represented by the generation of boulder-rich marginal channels that finalise the triangular shape of the murtoos (Peterson Becher and Johnson, 2021). Murtoo deposition is abruptly terminated and marginal channels are abandoned. The sedimentation within these marginal channels is characterised by suspension settling and laminated muds, indicating that the depositional space (0.6–0.8 m tall) remained open and water filled but no longer hydraulically connected to the active meltwater system (Ojala et al., 2022; Hovikoski et al., 2023).

Importantly, murtoo morphometry (Mäkinen et al., 2017; Ojala et al., 2021), their sedimentological architecture (Peterson Becher and Johnson, 2021; Hovikoski et al., 2023; Mäkinen et al., 2023), and close spatial association with eskers, ribbed tracts, and putative subglacial lakes (Ojala et al., 2021; Ahokangas et al., 2021; V^{er}it^e et al., 2022; Mäkinen et al., 2023) suggests that murtoos represent a transition between distributed and channelised drainage. Their formation occurs within broad and low conduits, subject to increasing water discharge throughout a melt season, at water pressures close to or exceeding ice

overburden pressure, and with short sediment transport distances such as might be found at the spatial onset of channelisation
125 in a 'semi-distributed' transitional drainage system (Hovikoski et al., 2023).

3 Study area

Our study area comprises the Finnish Lake District Ice-Lobe (FLDIL, Putkinen et al., 2017; Palmu et al., 2021), central
Finland (Figure 1A). The FLDIL is one ice lobe amongst several which comprised the eastern margin of the FIS, and contains
the highest density of murtoo fields in the region (Figure 1A, Ahokangas et al., 2021). Murtoo distribution within the FLDIL is
130 representative of their distribution across the wider FIS. In the upstream FLDIL trunk, murtoo fields occur amongst ribbed and
hummocky moraines (Figure 1C) in two longitudinal bands, each bounded by a dense assemblage of streamlined forms. In the
northeastern longitudinal bands, eskers are particularly clearly associated with murtoo routes (Figure 1D). Downstream, where
the FLDIL broadens into a lobe (Figure 1A), murtoo distribution is more fragmented with less clustering evident. Murtoos
are sparse in the centre of the ice lobe where thin sediment cover (Figure A1) may have limited the material necessary for
135 murtoo formation, and the high density of water bodies may act to mask existing murtoo fields (Ahokangas et al., 2021). The
FLDIL geology is predominantly crystalline bedrock, dominated by Precambrian schists, gneisses, and granitoids (Lehtinen
et al., 2005) with a thin Quaternary overburden (Lunkka et al., 2021).

In total, the FLDIL encompasses an area of $\sim 57,600$ km² and its distal margin is clearly defined by the first and second
Salpausselkäs (Figure 1A). These large ice-marginal complexes mark the Younger Dryas extent of the FIS (12.7 to 11.7 cal. ka,
140 Donner, 2010; Lunkka et al., 2021). Upstream of the Salpausselkäs there is no clearly defined ice marginal complexes suggest-
ing the FLDIL collapsed continuously and rapidly following the Younger Dryas (Kleman et al., 1997), retreating northwest
towards the Scandes mountains (Sweden and Norway) and ice-free by 9–10 cal. ka (Hughes et al., 2016; Stroeven et al., 2016;
Regnéll et al., 2019). Accordingly, to avoid arbitrarily demarcating an ice margin, we bound our model domain at the sec-
ond Salpausselkä, which marks the FLDIL extent at ~ 12 cal. ka (Putkinen et al., 2017). With a fixed domain bound at the
145 second Salpausselkä, we are effectively representing a single time slice ~ 12 cal. ka. Shoreline data indicates that the second
Salpausselkä terminated in a shallow water body ranging in depth from <5 m to ~ 50 m (Lunkka and Erikkilä, 2012). The
speed of the FLDIL retreat, together with the complex and dense assemblage of glaciofluvial landforms (see Palmu et al.,
2021; Dewald et al., 2021), suggest that during deglaciation, our study area was characterised by high and spatially extensive
atmospheric-driven surface melting delivered to the bed, proximal to significant calving of the FIS into the Baltic Sea Basin
150 (Greenwood et al., 2017; Patton et al., 2017; Boswell et al., 2019). Conditions within the FLDIL, and the FIS more broadly,
were likely comparable to conditions prevalent in land- or shallow-water terminating portions of the Greenland Ice Sheet today
(Greenwood et al., 2016; Ojala et al., 2019).

4 Methods

To explore how well a process-based model of basal hydrology can explain murtoo genesis, we applied the Glacier Drainage System Model, GlaDS (Werder et al., 2013), to our study area in the FLDIL. With a representative ice sheet surface and palaeo basal topography, together with a baseline set of GlaDS parameters derived from previous work beneath contemporary ice sheets (e.g., Dow et al., 2018a, 2020, 2022), we compared model output from GlaDS to the subglacial hydrological conditions proposed for murtoo genesis. Model output includes water pressure expressed as a percentage of ice overburden, $overburden\%$, sheet discharge, q_s , channel discharge, Q_c , and water velocity, V_W . We examined catchment-scale hydrology outputs and compare these to murtoo formation predictions as well as the distribution of channelised landforms (eskers) in the FLDIL. At individual nodes, we compared the evolution of nodes across our domain against the developmental phases recorded within murtoo sediment excavations (see Table 1, Hovikoski et al., 2023). We then explored seasonal and inter-annual model outputs in the area 40–60 km from the ice margin, at the upglacier limit of channelisation, where murtoos are hypothesised to form (Ahokangas et al., 2021; Ojala et al., 2021). Finally, we go on to investigate differences in model outputs between observed murtoo and meltwater routes, and terrain in which there are no glaciofluvial landforms. We sensitivity tested the robustness of all these findings to a range of input parameters. Readers interested in specific details regarding our modelling approaches are referred to Sections 4.1–4.1.2 in which we describe GlaDS; our climate forcing and ice sheet/bed topography; model parameters; and sensitivity testing in detail.

Table 1. Murtoo developmental phases (see Figure 10 in Hovikoski et al., 2023), their sedimentological signature, and anticipated model outcomes, where q_s is sheet discharge ($\text{m}^2 \text{s}^{-1}$), Q_c is channel discharge ($\text{m}^3 \text{s}^{-1}$), $overburden\%$ is water pressure expressed as a percentage of ice overburden pressure, and V_W is water velocity (m s^{-1}). See Section 4.1 and Appendix A for the model description.

Murtoo developmental phase	Sedimentological evidence and interpretation	Expected model outcomes
1	Sorted sediment core within rapidly enlarging cavity, partial ice contact	Sharp increases in q_s , increase in $overburden\%$
2	Onset of spring melt, cavity continuing to enlarge, with deposition of sinusoidal stratification and cobbles	Peak in $overburden\%$, continued increase in Q_c
3	Increasing grain size indicates high water velocity and boulder deposition indicates maximum cavity size of 1 m	Peak in V_W , drop in $overburden\%$, peak in Q_c approaching $1 \text{ m}^3 \text{ s}^{-1}$
4	Enlarged cavity leads to water pressure drop and ice-bed recoupling	Continued drop in $overburden\%$ and Q_c
5	Abrupt termination of discharge with appearance of laminated mud	$overburden\%$ approaching winter values

4.1 Model description

170 We used the Ice-sheet and Sea-Level System Model (ISSM, Larour et al., 2012, Revision 27448) and the implementation of
the GlaDS model (Werder et al., 2013) contained therein. The GlaDS model (described further in Appendix A and in full in
Werder et al., 2013) is a 2D finite element model, which has been widely applied to contemporary ice sheets in Greenland
(e.g., Dow et al., 2018a; Cook et al., 2020, 2022; Ehrenfeucht et al., 2023) and Antarctica (e.g., Dow et al., 2018b, 2020;
175 Indrigo et al., 2021; Dow et al., 2022; McArthur et al., 2023; Hayden and Dow, 2023) as well as glaciers in Svalbard (e.g.,
Scholzen et al., 2021). The GlaDS model operates on an unstructured mesh and includes a model of distributed flow through
linked cavities (Hewitt, 2011) represented by a continuous ‘sheet’ of water with variable thickness at mesh elements, and
channelised flow—describing semi-circular R othlisberger channels (R-channels) that are allowed to change diameter—along
element edges (Schoof, 2010). Sheet elements exchange water with channels, and the cross-sectional area of these channels
evolves through time due to the dissipation of potential energy, sensible heat exchange, and cavity closure rates due to viscous
180 ice creep. Here, flow in both the sheet and channel is assumed to be fully turbulent (cf., Hill et al., 2023). Unlike in other models
previously applied to the palaeo setting, GlaDS does not require a predetermined drainage system, the growth and restriction of
channels is entirely due to drainage dynamics (Dow et al., 2020). Following Werder et al. (2013), we set a threshold discharge
of $Q_c = 1 \text{ m}^3 \text{ s}^{-1}$ above which an element edge is classified as a ‘meaningful’ channel for our subsequent analysis.

4.1.1 Model setup: boundary conditions and forcings

185 As model inputs, GlaDS requires bed elevation, z_b , ice thickness, H , basal velocity, U_b , boundary conditions, and meltwater
input. We anticipate that the modern topography is not representative of bed elevation ~ 12 cal. ka. Therefore, as the baseline
 z_b , we account for these anticipated changes, particularly in terrain associated with the second Salpausselk a ice-marginal
formation, by subtracting Quaternary sediment thickness estimates (GTK, Finland, 2010) from the 25 m/pixel EU-Digital
Elevation Model V1.1 (available at: <https://www.eea.europa.eu/data-and-maps/data/copernicus-land-monitoring-service-e>
190 u-dem). Lake bathymetry was only partially available in the FLDIL, so we did not subtract this from our input DEM in the
baseline model. We also did not adjust our model to account for differences in elevation due to glacial isostatic adjustment
(GIA) since ~ 12 cal. ka. To ensure the numerical stability of GlaDS the input DEM was smoothed using a low-pass filter.
Finally, within steep terrain, an anisotropic mesh ($n_{nodes} \approx 19,000$) was refined based on z_b such that element edges were
shortest (to a minimum edge length of 400 m) in rougher terrain and longer where terrain was flatter (to a maximum edge
195 length of 2 km).

We generated H using the 2D Shallow-Shelf Approximation (SSA, MacAyeal, 1989) within ISSM (Larour et al., 2012). An
initial estimate of H was given using a parabolic profile as a function of distance from the terminus, and initialisation values
for basal velocity were calculated using a stress balance solution for this ice surface. Dirichlet conditions were imposed at the
mesh edges along the boundary with zero inflow. Basal motion was modelled using a viscous sliding law (Budd et al., 1979)
200 and following  akesson et al. (2018) we used a spatially variable basal drag coefficient, a , proportional to z_b , given by:

$$a = 120 \frac{\min(\max(0, z_b + 800), 2000)}{2000}. \quad (1)$$

Ice was assumed to be isothermal with a viscosity, B , equivalent to an ice temperature of -5°C (from Cuffey and Paterson, 2010, p.73; rate factor, A , listed in Table A1). In reality, ice temperature is both spatially and temporally variable. However, without using a more detailed thermomechanical ice model, we follow the previous ad-hoc assumptions of Nick et al. (2013) for the Greenland Ice Sheet and Åkesson et al. (2018) for the FIS, by setting our ice temperature to -5°C . The 12 cal. ka climate for the ice sheet model was estimated using a modern (1981-2010) reanalysis dataset (see Abatzoglou et al., 2018). Precipitation was kept at the contemporary monthly value, but we depressed monthly temperature by 15°C , approximately the temperature differential indicated by NGRIP $\delta^{18}\text{O}$ records (Johnsen et al., 1997). To calculate surface mass balance efficiently in our long term ice sheet model we used a simple positive degree day (PDD) model (as in Cuzzone et al., 2019) allowed to vary about a fixed Gaussian distribution with standard deviation, $\sigma_{PDD} = 5.5^\circ\text{C}$ around the monthly mean and a lapse rate of $7.5^\circ\text{C km}^{-1}$. To reach volumetric steady state, defined for our ice sheet model as differences in ice volume between successive iterations of less than 10^{-6} km^3 , we ran the ice sheet model for 20,000 years with an adaptive timestep, allowed to vary between 1 day and 1 year. The final H was stored and used as the ice sheet input to GlaDS.

As boundary conditions for GlaDS, we imposed a zero flux condition on the domain edge everywhere except at the ice terminus, where given spatial variability in water depth (Lunkka and Erikkilä, 2012), an outlet Dirichlet condition equivalent to atmospheric pressure was prescribed in the baseline model. By enforcing zero input flux we neglect to include basal water input from beyond the model domain and we also do not account for any exchange of water between adjacent ice lobe provinces. To promote model stability, we used an adaptive timestep that was allowed to vary between one hour and ~ 90 seconds and all of our transient models were run for 10,000 days, or ~ 27 years. To approximate winter conditions and avoid suddenly overwhelming our initial system with sudden surface meltwater inputs, we first ran GlaDS to steady state with basal meltwater input but no surface melt. To guarantee the majority of elements were pressurised at the end of each steady state run, we used a low, fixed basal velocity of 30 m yr^{-1} to limit the rate of cavity expansion (see Equation A3). We judged the system to be in steady state once the median difference in sheet thickness between two successive steps was less than 10^{-6} m . This occurred within 20,000 days in all runs, and the majority of nodes reached 90% of overburden pressure¹ with no channel formation.

For the subsequent transient model runs, melt input to GlaDS was estimated using the same depressed monthly average temperature and precipitation record as with the ice sheet model. In simply depressing the climate by 15°C we do not represent the complex seasonality (short, warm summers with extreme winters) that characterised the Younger Dryas cold reversal in Fennoscandia (Schenk et al., 2018; Amon et al., 2022). However in fixing our domain to the second Salpausselkä our hydrology model is representative of the end of the Younger Dryas (~ 12 cal. ka) at which time this complex seasonality rapidly gave way to a markedly warmer climate with similar seasonality to the present day (Mangerud et al., 2023). Compared to the ice sheet model, we did use a modified PDD scheme for GlaDS to more faithfully reflect daily temperature variability over the shorter

¹Borehole measurements of overwinter water pressure in the distributed drainage system have been measured at 80-90% of overburden pressure (e.g., Harper et al., 2021)

maximum timestep. It is commonly assumed that the total monthly positive degree days can be represented by a fixed Gaussian distribution with $\sigma_{PDD} \approx 5.5^\circ\text{C}$ (e.g., Braithwaite and Olesen, 1989). However, field measurements suggest that this does not hold for the Greenland Ice Sheet (Wake and Marshall, 2015), particularly at temperatures $\geq -5^\circ\text{C}$. Instead, Wake and
235 Marshall (2015) suggest monthly variability in temperature, σ_M , is more accurately described by a quadratic function:

$$\sigma_M = -0.0042T_M^2 - 0.3T_M + 2.64, \quad (2)$$

where T_M is the mean monthly temperature. This function accounts for the reduction in variability with increasing temperature (Gardner et al., 2009; Marshall and Sharp, 2009; Fausto et al., 2011) due to heat buffering, which promotes a more stable boundary layer (Wake and Marshall, 2015). We used σ_M from Wake and Marshall (2015) but did not take into account vari-
240 ations in kurtosis and skewness with temperature, as these become significant where $T_M < -20^\circ\text{C}$ (see Wake and Marshall, 2015), temperatures below those derived from our depressed MAT. Instead we used the calculated σ_M to add Gaussian noise to a daily temperature record estimated by linearly interpolating our depressed MAT record. The number of positive degree days per month, PDD_M was taken as $PDD_M \geq -5^\circ\text{C}$. Following van den Broeke et al. (2010) we used -5°C as our threshold (rather than the more commonly used 0°C threshold) to account for melt which may occur even for days with an average
245 temperature of 0°C . Finally, we used melt rate factors $\gamma_{ice} = 17.22$ mm per PDD and $\gamma_{snow} = 2.65$ mm per PDD following Cuzzone et al. (2019) keeping these consistent between our ice sheet model and GlaDS model. We did not prescribe any inter-annual variability in average monthly temperature. Melt varied in absolute terms between individual simulations but the mean melt and standard distribution remained identical throughout.

The total monthly melt was routed to the bed via a series of moulins. Following Werder et al. (2013) we divided our domain
250 using Voronoi tessellation on a randomly distributed series of points. Within each Voronoi cell, acting as a ‘catchment zone’, the lowest elevation node was identified and used as the location for a moulin towards which all melt from all other catchment nodes flow. Surface melt rate was integrated over each catchment and converted to instantaneous moulin discharge, Q_m^k . Without a detailed record of daily melt variability we neglect to include daily and diurnal changes in melt, which are known to drive rapid changes in hydraulic head on the Greenland Ice Sheet (Andrews et al., 2014). Smoothing melt variability reduced
255 model size and improved the stability of GlaDS over the ~ 27 year model runs, and we note that the inclusion of an englacial storage term in GlaDS acts to restrict the influence of diurnal variability to within 2 km of moulins with a limited influence on the overall pattern of channelised drainage (see Werder et al., 2013).

4.1.2 Model parameters in the baseline model and sensitivity testing

The GlaDS model has been extensively sensitivity tested for contemporary ice sheets where model results can be compared with
260 geophysical evidence to determine the most plausible model output (e.g., Werder et al., 2013; Dow et al., 2018b, 2020, 2022; Indrigo et al., 2021; Scholzen et al., 2021), and as such we do not conduct a detailed review here. We set the parameters in our baseline model (default values listed in Table A1) following the default values in these studies, which provide a reasonable approximation of contemporary ice sheet subglacial conditions. However, because several parameters in GlaDS have uncertain

physical values, we did test the robustness of our findings from the baseline scenario throughout the ranges indicated in
265 Table A1. We sensitivity tested for basal melt rate; moulin density and distribution; sheet and channel conductivity terms; basal
bump height; the englacial void ratio; basal ice velocity; terminus boundary conditions; bed topography; and mesh geometry.
We can assign higher confidence to our baseline model when similar model outputs (e.g., similar channel lengths or patterns
of water pressure) are evident across multiple sensitivity tests (e.g., Dow, 2023).

Given uncertainty regarding the spatial variability of basal melt rates beneath current and former ice sheets, which vary
270 as a function of geothermal heat and frictional heating, we used a spatially and temporally constant basal water input (as
in Dow et al., 2018a, c, 2020; Poinar et al., 2019). Basal melt rates beneath the Greenland Ice Sheet typically range between
 $1\text{--}7 \times 10^{-3} \text{ m yr}^{-1}$ (see Karlsson et al., 2021) and we used $5 \times 10^{-3} \text{ m yr}^{-1}$ for our baseline model configuration and the major-
ity of the subsequent transient runs. We tested the influence of basal melt rate by running a low basal melt rate ($1 \times 10^{-3} \text{ m yr}^{-1}$)
and high basal melt rate scenario ($7 \times 10^{-3} \text{ m yr}^{-1}$).

275 As default, surface meltwater was routed through ~ 2500 moulins, a density of 0.04 moulins per km^2 . Measured moulin
density varies between 0.02 to 0.09 moulins per km^2 in Greenland (Yang and Smith, 2016). To test the sensitivity of our
system to moulin density we also ran models with ~ 1000 (0.02 per km^2), ~ 4000 (0.06 per km^2), and two further randomly
generated configurations of the default ~ 2500 (0.04 per km^2). We also tested an additional configuration in which melt at every
node was routed directly to the bed.

280 Further sensitivity testing (parameters listed in bold in Table A1) was carried out for several poorly constrained parameters
in GlaDS, as well as for the basal geometry and moulin density. The conductivity of both the sheet, k_s and channels, k_c are key
controls on the pressure of the system, and alter the relative efficiency of each system, and in turn alter the spacing, length, and
upstream pressure influence of channels. The conductivity terms in GlaDS are poorly constrained, and following previous work
we tested at magnitude limits up to the point at which the model failed to converge (see Dow, 2023). The baseline value for
285 k_s was $10^{-4} \text{ m}^{7/4} \text{ kg}^{-1/2}$, and we tested additional setups where $k_s = 10^{-2}, 10^{-3}$, and $10^{-5} \text{ m}^{7/4} \text{ kg}^{-1/2}$. For k_c the baseline
value was 10^{-1} and we tested $k_c = 5 \times 10^{-1}, 5 \times 10^{-2}$, and $10^{-3} \text{ m}^{3/2} \text{ kg}^{-1/2}$.

The basal bump height, h_r alters how readily cavities open in the distributed system. Our default value for h_r was 0.085 m
and we additionally tested $h_r = 0.05$ and 0.1 m. We tested values of englacial storage, $E_{vr} = 10^{-3}$ and $E_{vr} = 10^{-5}$. For basal
velocity, U_b , we tested prescribed values of between $100\text{--}200 \text{ m yr}^{-1}$ chosen to be comparable to surface velocity across land-
290 terminating sectors of the Greenland Ice Sheet (e.g., Tedstone et al., 2015). We tested both a temporally fixed and temporally
variable U_b , with the transient U_b varying between 85% and 140% of the mean to approximate speed-ups at the onset of
the melt season and winter slowdowns commonly observed in Greenland (e.g., Sole et al., 2013). Without a more detailed
understanding of past ice dynamics, U_b was kept spatially uniform throughout.

Although the default configuration describes a terrestrial margin, we also tested the influence of a shallow body of water at
295 the ice margin by prescribing Dirichlet conditions at the ice margin where water pressure is equivalent to that of a uniform 30 m
water depth (a simplification of the variable 5–50 m water depth from Lunkka and Erikkilä, 2012). To explore the influence
of our modified topography boundary condition, we ran tests with a uniformly flat bed, one representing contemporary terrain
(without Quaternary sediment thickness removed), and one with the available partial lake bathymetry removed. Finally, we also

explored the dependency of our results on mesh geometry, including using a coarser mesh (maximum edge length of 5 km), a
300 mesh not refined by elevation in any way, and a mesh in which a coarse mesh (edge length between 5–8 km) was prescribed
>80 km from the ice margin and a much finer mesh (edge length ≈ 300 m) was prescribed <80 km from the ice margin.

5 Results and discussion

In total, 30 simulations were carried out and for each model run, we examined the subglacial water pressure, expressed as a
percentage of the overburden pressure, $overburden\%$, sheet discharge, q_s on element faces; channel discharge Q_c on element
305 edges, and water velocity, V_W . In order to examine how well GlaDS is able to explain murtoo genesis we first describe the
catchment scale hydrology across our model runs and examine the evolution of the model through time (Sections 5.1–5.3). In
Section 5.4 we describe the limitations of our approach and suggest possible future research.

5.1 Catchment scale hydrology

In the baseline model (see Figures 2, & 3), model outcomes at the catchment scale suggest that GlaDS matches several of
310 the spatial predictions for murtoo genesis. After an initial adjustment from steady state to transient forcing over five years,
the baseline model reached a quasi-steady state configuration in which the system responded seasonally to summer meltwater
input (Figure A3). Following this adjustment, a clear and sharply demarcated transition in drainage modes develops 40–60 km
from the ice margin during summer. Murtoos are predicted to have formed at this distance from the ice margin of the FIS,
as widespread distributed drainage gave way to channelised drainage within a semi-distributed system (see Mäkinen et al.,
315 2017, 2023; Peterson Becher and Johnson, 2021; Ojala et al., 2019, 2021, 2022; Ahokangas et al., 2021; Vérité et al., 2022;
Hovikoski et al., 2023). Additionally, weak to moderate deformation of murtoo sediments suggests that water pressure remained
at or close to overburden pressure for sustained periods during the melt season (Peterson Becher and Johnson, 2021; Vérité
et al., 2022; Mäkinen et al., 2023; Hovikoski et al., 2023). In our results, a persistent area of $overburden\% > 100\%$ develops
across the full width of the domain 40–60 km from the ice margin during summer (Figures 2,3, & Movie A1). Outside this
320 area (<40 km of the ice margin, and >70 km from the ice margin) $overburden\%$ is 10–30% lower during summer. By winter,
 $overburden\%$ drops by up to 30% across the domain as melt ceases (Figure 2A).

Approximately 35 modelled channels (edges where $Q_c > 1 \text{ m}^3 \text{ s}^{-1}$) extend up to 50 km perpendicular from the ice margin,
into but not beyond the hypothesised zone of murtoo formation (Ojala et al., 2021) and terminate where $overburden\%$ exceeds
100% during summer. Murtoo fields (Ahokangas et al., 2021) are evident at the head of many modelled channels, particularly
325 in the western and eastern areas of the ice-lobe (Figure 4A). Within 40–60 km of the ice margin, the median cross-sectional
area of edges is 2.8 m^2 (equivalent to a semi-circle with radius of 1.3 m). Closer to the ice margin, the median summer Q_c
of channels reaches up to $100 \text{ m}^3 \text{ s}^{-1}$ (Figure 2D) and they are comparable in both sinuosity and spacing to esker deposits in
the FLDIL (Figure 4, Palmu et al., 2021). Channel spacing and length is also comparable to the theoretical spacing of eskers
derived from the modelling results of Boulton et al. (2009) and Hewitt (2011). The development of channels during summer
330 also strongly influences q_s and V_W , and each are highest adjacent to active channels close to the ice margin ($q_s \approx 10^{-3} \text{ m}^2 \text{ s}^{-1}$,

$V_W \approx 1.4 \times 10^{-3} \text{ m s}^{-1}$), and remain high until 40–60 km from the ice margin ($q_s \approx 10^{-4} \text{ m}^2 \text{ s}^{-1}$, $V_W \approx 8 \times 10^{-4} \text{ m s}^{-1}$) where values drop to a low upglacier ($q_s \approx 10^{-5} \text{ m}^2 \text{ s}^{-1}$, $V_W \approx 1 \times 10^{-4} \text{ m s}^{-1}$).

Without independent constraint against which to compare our results, we ran additional sensitivity tests to explore the parameter dependency of our findings, assigning higher confidence to model outcomes present across the majority of tests (see Dow, 2023, and section 4.1.2). The catchment scale hydrology described in our baseline model remains consistent across most of the additional sensitivity tests. Furthermore, sensitivity test results remain consistent with predictions for moraine genesis. This includes for: all moulin density tests (Figures A10, & A12–A14), except the highest moulin density; basal melt rate (Figures A15, & A16); mesh geometry and bed elevation (Figures A23–A22); the addition of lake bathymetry (Figure A24); a shallow proglacial water body (Figure A25); the englacial void ratio (Figures A26–A27); and basal velocity (Figures A28–A32).

While catchment-scale trends are largely robust, the exact location of channels, and their exact length and local *overburden*_%, does vary between sensitivity tests. Because GlaDS operates on a fixed mesh (cf. Felden et al., 2023), the resolution of which is a balance of suitable fidelity against the increased computational cost of resolving finer details, the exact location of where channels may form does vary between sensitivity tests. These minor differences in channel location change the spatial expression of summer *overburden*_%, with the area of *overburden*_% $\approx 100\%$ changing by up 10 km and differences of between 5–10% for any given location. Channel location is particularly sensitive to mesh geometry, but differences also arise because of moulin density and location, bed topography, basal velocity, and basal bump height. However, while the absolute position of channels does vary, channel spacing remains consistent (~ 15 km), and changes in channel length are limited to a maximum of ~ 10 km.

Although consistent across the majority of tests, channel length and *overburden*_% does vary considerably at the tested limits of k_s and k_c parameters, describing the sheet and channel conductivity respectively (Werder et al., 2013). For both the maximum tested sheet conductivity ($k_s = 10^{-2} \text{ m}^{7/4} \text{ kg}^{-1/2}$, Figure A4) and the minimum tested channel conductivity ($k_c = 10^{-3} \text{ m}^{3/2} \text{ kg}^{-1/2}$, Figure A9), there are major changes to the catchment hydrology. Channels are restricted to within ~ 1 km of the ice boundary and *overburden*_% within and near to these channels is $\approx 50\%$ when $k_s = 10^{-2} \text{ m}^{7/4} \text{ kg}^{-1/2}$ and approaches 120% when $k_c = 10^{-3} \text{ m}^{3/2} \text{ kg}^{-1/2}$.

For both the minimum sheet conductivity ($k_s = 10^{-5} \text{ m}^{7/4} \text{ kg}^{-1/2}$, Figure A6) and the maximum channel conductivity ($k_c = 5 \times 10^{-1} \text{ m}^{3/2} \text{ kg}^{-1/2}$, Figure A7) channel length increases to between 50–60 km, only a 10 km increase in channel length compared to the baseline scenario, but there are additional major changes in *overburden*_%. At the minimum k_s , an area of *overburden*_% $\approx 100\%$ extends up to 150 km from the ice margin during summer, and at the maximum k_c there is no area of *overburden*_% $> 90\%$ during summer. When $k_s = 10^{-3} \text{ m}^{7/4} \text{ kg}^{-1/2}$ (Figure A5) and when moulin density is highest (Figure A11), channels are restricted to within 20 km and there is an area of high *overburden*_% $> 100\%$ between 10–70 km of the ice margin.

Changing the channel and sheet conductivity by orders of magnitude strongly modifies the efficiency with which either the distributed or channelised system can transmit water, limiting the influence of the other system (Werder et al., 2013). At the highest tested moulin density, the reduced discharge associated with any one moulin resulted in a higher density of short channels as fewer reach the threshold discharge at which wall-melt exceeds creep-driven channel closure. Excessively




Figure 2. Median summer and winter system states in the baseline model run. **A)** Water pressure as a percentage of overburden pressure, $overburden\%$. In summer and winter, **B)** Sheet discharge, q_s . **C)** Water velocity, V_w . **D)** Channel discharge, Q_c . For each output, we took the median from model years 5–27 disregarding the initial period of adjustment to transient forcing. *Summer* extends from May to September, all other points fall into *winter*. Note that the scales for panels B and D are logarithmic. Dashed lines in all panels indicate contours of 40 and 60 km from the ice margin. Murtoo fields (Ahokangas et al., 2021) are shown as inverted triangles in all plots. Channels are shown as black solid lines in panels A–C.

long (>50 km) or short (<10 km) channels are considered to be invalid on the basis of modern Greenland observations (e.g., Chandler et al., 2013; Dow et al., 2015) and an anomalous $overburden\%$ is considered invalid on the basis of the conceptual model for murtoo distribution and genesis (e.g., Ahokangas et al., 2021; Hovikoski et al., 2023). Therefore, our baseline conductivity terms are the considered the most plausible parameters.

370 Finally, given the parameters in our baseline model are set following existing work on contemporary ice sheets (Section 4.1.2), it is no surprise that at the catchment scale, the modelling outputs appear similar to the subglacial hydrological system evident in land-terminating sectors of the Greenland Ice Sheet. Tracer transit times (e.g., Chandler et al., 2013) and basal hydrology modelling indicate efficient channelisation extends up to 40–50 km from the ice margin in Greenland, transitioning between channelised and distributed drainage modes where ice is ~ 900 – 1200 m thick (De Fleurian et al., 2016; Dow
375 et al., 2015) as it does here. However, the pressure conditions within large channels close to the ice margin are notably different




Figure 3. The evolution of water pressure as a percentage of overburden pressure, ($overburden\%$), sheet discharge, (q_s), water velocity, (V_w), and maximum channel discharge, (Q_{cmax}) at four nodes over model years 15–25 in the baseline model run. **A)** Node No. 6,277 located ~ 120 km from the ice margin. **B)** Node No. 18,517 located ~ 17 km from the ice margin. **C)** Overburden at the end of the melt season in model year 19 (arbitrarily selected). Channels are represented as black lines, murtoo fields as inverted black triangles, and the location of panels A, B, D, and E as coloured points. Inset shows overburden at every node as a function of distance, D from the ice margin with a smoothing function shown in red and vertical lines at 40 and 60 km from the ice margin. **D)** Node No. 16,402 located 0.7 km from a murtoo field and ~ 45 km from the ice margin. **E)** Node No. 3,842 located ~ 54 km from the ice margin at the head of a channel system without an adjacent murtoo field. The time slice shown in panel C is represented as a vertical dashed line in panels A, B, D, and E. Note the logarithmic scale for q_s and Q_{cmax} .




Figure 4. Modelled channel location compared to esker deposits (mapped by Palmu et al., 2021) and murtoo fields (mapped by Ahokangas et al., 2021) at the ice margin of the FLDIL. **A)** Modelled channels during summer in the baseline run (black lines) across the full width of the domain compared to esker deposits (blue polygons) and murtoo fields (inverted black triangles). Each channel is defined where the median channel discharge, Q_c , exceeds $1 \text{ m}^3 \text{ s}^{-1}$ during summer. **B & C)** Detailed comparison of two large esker systems against model channels.

in our model results compared to the Greenland settings (e.g., Van de Wal et al., 2015). In Greenland, channels exist at lower water pressures than the surrounding distributed system (Davison et al., 2019), and the resultant hydraulic potential gradient forces large volumes of water from the surrounding distributed system towards channels, in turn lowering water pressure in the distributed system and increasing basal traction (Schoof, 2010). Even as meltwater delivery to the bed subsequently increases through the melt season, these channels can act to reduce ice velocity (Nienow et al., 2017) and reduce ice mass loss. In contrast, the channels modelled here remain at relatively high *overburden*_% throughout the year ($> 60\%$), with a lower hydraulic potential gradient between channelised and distributed systems. The FLDIL is low-relief compared to the steep margins of the Greenland Ice Sheet (e.g., Wright et al., 2016), and the shallow topography may act to reduce the hydraulic gradient between distributed and channelised drainage. As a result, the influence of channelisation on basal velocity would be relatively limited in the FLDIL. Lower rates of water exchange between distributed and channelised drainage would permit more of the bed to remain closer to *overburden*_% $\approx 100\%$, sustaining higher velocities for extended periods of time (Dow et al., 2022), highlighting the sensitivity of low-relief areas of the FIS to extensive atmospheric melting.

5.2 Comparison to the murtoo developmental phases

The widespread and time-integrated distribution of murtoos throughout our model domain complicates model validation as murtoo formation conditions remain uncertain. The ability of GlaDS to reproduce the hypothesised spatial pattern of murtoo formation (i.e., summer *overburden*_% $\approx 100\%$ 40-60 km from the ice margin) alone cannot definitively confirm or refute the

hypothesized formation process because murtoos are distributed across our model domain. In this context, comparing seasonal model evolution to murtoo sedimentology (e.g., Hovikoski et al., 2023; Mäkinen et al., 2023) becomes particularly important as there are multiple predictions in sequence that the model must achieve (Table 1).

395 Examined at individual nodes through time, the baseline model (and most sensitivity tests) agrees well, though not perfectly, with the hypothesised conditions and location of murtoo genesis. Internal murtoo sediments chart an overall increase in meltwater discharge throughout the melt season followed by an abrupt termination (Table 1), possibly within the same year (Hovikoski et al., 2023). Against this backdrop, the alternating sequences of glaciofluvial deposits in the main body of murtoos suggests that the system was also subject to repeated pulses of meltwater and rapid changes in flow regime, marking
400 the rerouting and periodic isolation of cavities within a developing, semi-distributed drainage system (Hovikoski et al., 2023; Mäkinen et al., 2023). Figure 3 shows the evolution of the system through time at four representative nodes across the study area. At node 6,277, 120 km from the ice margin and upglacier of any significant surface melt inputs, the system is effectively inert, with $overburden_{\%}$ remaining $\approx 80\%$ and only small periodic perturbations in q_s , Q_c , and V_W . However, closer to the ice margin, the system responds annually to meltwater inputs and is more variable through time.

405 Figure 3D & E demonstrates the seasonal evolution of two nodes between 40–60 km from the ice margin, each nearby to channel systems. Both nodes fall within the hypothesised zone of murtoo formation and both nodes display drainage behaviour which could accommodate murtoo formation, with a seasonal increase in $overburden_{\%}$ up to a maximum of approximately 120% and a more gradual decrease thereafter. However, the two nodes show different interannual behaviour, and only one is located close to a murtoo field. At node 3,842, ~ 54 km from the ice margin and chosen to be representative of surrounding
410 nodes at the onset of a channel (Figure 3E), the pattern of drainage repeats annually—every year the increase and decrease in $overburden_{\%}$ is accompanied by peaks in q_s , Q_c , and V_W and the nearby development of channels throughout the meltwater season. At the onset of channelisation the maximum Q_c approaches but never exceeds $1 \text{ m}^3 \text{ s}^{-1}$. However, although this evolution through time does appear consistent with each of the murtoo developmental phases (Table 1), node 3,842 is not located near to a murtoo field.

415 At node 16,402, located 0.7 km from a murtoo field and chosen to be representative of murtoo fields, a more complex signal is evident. At the maximum upglacier extent of two adjacent channels, ~ 45 km from the ice margin a biannual signal is evident (Figure 3D). Every year, there is a sharp increase in $overburden_{\%}$ at the start of the melt season to $overburden_{\%} \geq 100\%$. However, the subsequent drop in $overburden_{\%}$ varies every other year. Either $overburden_{\%}$ spikes and then drops rapidly over 1-2 months to the winter value ($\sim 80\%$) until the following melt season, or the drop in $overburden_{\%}$ is initially shallower
420 before quickly dropping to an $overburden_{\%}$ ($\sim 90\%$) that is elevated relative to the previous winter. Years with an elevated winter $overburden_{\%}$ are also associated with lower Q_c and flatter peaks in q_s . In contrast, years that have a rapid drop in $overburden_{\%}$ after the start of the melt season are associated with higher values of Q_c , approaching $1 \text{ m}^3 \text{ s}^{-1}$, and sharper peaks in q_s . We consider that the latter case is more consistent with the murtoo developmental phases (Table 1) because the higher values of Q_c indicate discharge approached that necessary to form channels.

425 A similarly biannual pattern is evident at node 18,517, 17 km from the ice margin and collocated with a channel (e.g., Figure 3B). Here, close to the ice margin the maximum Q_c associated with edges connected to the node exceeds the threshold

for a meaningful channel every summer, reaching a peak of $200 \text{ m}^3 \text{ s}^{-1}$ \sim June each year with a maximum cross-sectional channel area of 42 m^2 (equivalent to a half-circle with radius, $r \approx 5 \text{ m}$). However, every other year, the maximum Q_c of edges connected to node 18,517 remains $\geq 1 \text{ m}^3 \text{ s}^{-1}$ i.e., the channel remains active over winter. Preexisting channels dampen the influence of melt in the following summer by providing an already established efficient drainage pathway with only a small increase in *overburden*_% and little change in q_s (Figure 3B). Overwinter channels form across the width of the domain in the baseline model, but an alternating spatial pattern of overwinter channel persistence is evident. In any given year, channels will persist through winter in either the central third of the lobe or in the remaining two thirds of the lobe (Figure A2 & Movie A1).

Finally, although individual nodes do track the overall increase in meltwater discharge throughout some melt seasons, as well as the evolution of *overburden*_% consistent with limited cavity expansion, our modelling fails to reproduce the sharp drop in discharge at the end of the melt season or the rapidly changing flow regimes within a single melt season that have been invoked to explain murtoo sedimentology (see Mäkinen et al., 2023; Hovikoski et al., 2023, and Section 2). We did not include diurnal variability in our modelling on the grounds of model stability and the limited influence diurnal forcing has on catchment scale drainage in GlaDS (Section 4.1.1 & Werder et al., 2013). Diurnal forcing would be critical in order to represent rapid changes in the flow regime within murtoo-forming cavities. However, as with other subglacial hydrology models, GlaDS is a model in which the subglacial system is assumed to be pervasively hydraulically connected (see Rada Giacaman and Schoof, 2023), and there is no mechanism which can lead to the hydraulic isolation of specific areas of the bed (e.g., Rada and Schoof, 2018; Hoffman et al., 2016). As a result, even if diurnal forcing were to be included, we do not expect to be able to reproduce the rapid changes in meltwater discharge necessary to form upper and lower flow regime deposits (see Section 2, & Hovikoski et al., 2023) or laminated muds in marginal murtoo channels (e.g., Ojala et al., 2022). Including spatially variable system conductivity is likely to be important in future work which seeks to evaluate the ability of process-based subglacial hydrology models to represent landform formation.

5.3 Comparing murtoo and meltwater route hydrology

Extensive geomorphological mapping has identified meltwater pathways across the FIS and within the FLDIL in particular (see Dewald et al., 2021; Ahokangas et al., 2021). We explore drainage behaviour in the area of anticipated murtoo formation by isolating and taking a spatial median of nodes in the baseline model 40–60 km from the ice margin. We grouped nodes by their relation to i) murtoo routes ($n = 244$), ii) meltwater routes ($n = 951$), and iii) neither ($n = 1205$), using 500 m buffers to approximate the lateral extent of murtoo/meltwater routes along 2D polylines from Ahokangas et al. (2021). Thresholding by distance is necessary to exclude nodes which either i) do not respond seasonally to meltwater input (e.g., those $> 60 \text{ km}$ from the ice margin, Figure 3A), or ii) nodes closer to the ice margin (within 0–40 km) where channelised drainage dominates in our modelling. Nodes 40–60 km were selected because this is both the hypothesised area of murtoo formation (Ahokangas et al., 2021) and the location in our modelling identified as one in which conditions for murtoo formation are met (Section 5.1). We note that we also include eskers within 40–60 km of the ice margin (mapped as ‘channelised routes’ by Ahokangas et al., 2021) which likely postdate $\sim 12 \text{ cal. ka}$. We do not have age-control on any individual landform and many channelised routes intersect or overprint murtoo/meltwater routes (Ahokangas et al., 2021), so we therefore classified them accordingly.

Plotting the median evolution of these groups through time (Figure 5) and as probability density functions (Figure 6) reveals clear differences between nodes in murtoo or meltwater routes and nodes which do not intersect any mapped glaciofluvial geomorphology. Compared to individual nodes (Section 5.2) the average evolution of nodes in murtoo and meltwater routes follows a regular pattern year-on-year (Figure 5). At the onset of the melt season, following winter minima, $overburden\%$, q_s , Q_c and V_W all begin to increase (Figure 5). The increase in $overburden$, q_s , and V_W is sharp, with a more gradual increase in Q_c through time. As cavity expansion promotes lower water pressure and more efficient discharge, $overburden\%$ peaks earliest \sim June each year, but remains $>100\%$ until \sim August, at which point V_W and q_s peak. As $overburden\%$, q_s , and V_W all decrease towards winter minima following peak melt, Q_c peaks in September, decreasing steadily towards a minimum before the next melt season. By contrast, the pattern across all other nodes (i.e., those which do not intersect mapped glaciofluvial geomorphology) remains relatively stable through time, and $overburden\%$, q_s , Q_c and V_W are lower throughout the year, suggesting there is limited evolution of the hydrological system in these nodes. In clearly distinguishing murtoo and meltwater routes from nodes which do not intersect mapped glaciofluvial geomorphology, GlaDS appears to be faithfully representing the drainage pathways active beneath the FLDIL at the end of the Younger Dryas ~ 12 cal. ka.

The probability density functions of murtoo routes and meltwater routes is also clearly distinct from terrain without glaciofluvial landforms (Figure 6). However, the probability density functions of murtoo routes and meltwater routes are also different from one another, particularly at the lower tail of the distributions (Figure 6). Murtoo routes have a $overburden\%$ distribution with a more tightly constrained lower tail than meltwater routes, with fewer nodes dropping below $overburden\% = 80\%$. There is a bimodal distribution of both q_s and V_W within murtoo routes that is not evident in meltwater routes at the lower tail of the distribution. Both meltwater routes and murtoo routes have a bimodal Q_c distribution, but the lower murtoo route peak is offset towards higher channel discharges. One-way ANOVA testing indicates that the difference in distribution between murtoo and meltwater routes is statistically significant ($p < 0.05$ at the 95% confidence interval). Additional Tukey-Kramer testing indicates that the significant difference between murtoo and meltwater routes varies throughout the year (Tables A2–A5). In June and July, $overburden\%$ is significantly lower in murtoo routes than in meltwater routes by 1–3%, throughout the rest of the year, $overburden\%$ is significantly higher in murtoo routes than in meltwater routes by the same amount. Between January and May there are no significant differences in q_s between murtoo routes and meltwater routes. However, between June and December, q_s is significantly lower in murtoo routes than in meltwater routes by $\sim 1 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$. There are no statistically significant differences in Q_c between murtoo routes and meltwater routes throughout the year. Statistically significant differences between murtoo/meltwater routes and all other nodes is limited to between June and October when Q_c is significantly higher in murtoo and meltwater routes than beyond by $\sim 1 \times 10^{-5} \text{ m}^3 \text{ s}^{-1}$. Finally, throughout the year, V_W is higher in murtoo and meltwater routes than beyond them by $\sim 1 \times 10^{-3} \text{ m s}^{-1}$. In murtoo routes V_W is significantly lower for each month than in murtoo free meltwater routes by $\sim 1 \times 10^{-4} \text{ m s}^{-1}$.

The differences between meltwater routes and murtoo routes are subtle, and the annual evolution of model outputs in both closely match the murtoo developmental phases (Table 1 & Hovikoski et al., 2023). We anticipate that this statistical difference has a strong spatial component linked to the overwinter channels reported in Section 5.2—murtoo routes are notably absent 40–60 km from the ice margin in the centre of the FLDIL, whereas meltwater routes are more evenly distributed across the full




Figure 5. Median output during model years 15–20 at nodes between 40–60 km from the ice margin. In all panels, nodes that fall within murtoo routes are shown in blue, those which fall within meltwater routes (without murtoos) are shown in orange, and all other nodes are shown in purple. The black line in each panel represents the residual difference between the median of murtoo routes and meltwater routes. A positive residual indicates higher median values in murtoo routes, and vice versa for negative residual values. **A)** Water pressure expressed as a percentage of overburden, $overburden_{\%}$. **B)** Sheet discharge, q_s . **C)** Water velocity, V_W . **D)** Channel discharge, Q_c . Note panels B and D have logarithmic scales.

width of the domain (see Figure A1A). Murtoo distribution closely overlaps with the distribution of overwinter channels and when channels in the central third of the FLDIL persist over one winter, those in the outer two thirds do not—and vice versa in the following winter (Figure A2). As a result, when overwinter channels persist nearby to murtoo routes, the following summers are characterised by sharper peaks in $overburden_{\%}$ and lower values of Q_c , V_W and q_s as melt is quickly accommodated by an established efficient drainage pathway (Section 5.2). In contrast, meltwater routes—distributed more evenly—are less sensitive to the spatial pattern of winter channels. We therefore anticipate that the statistically significant difference between murtoo routes and meltwater routes arises because of the repetitive expression of overwinter channels, and does not necessarily indicate that GlADS is resolving differences between the landforms themselves.

On the Greenland Ice Sheet, winter slowdowns following high-melt summers have been linked to the sustained persistence of larger and more extensive channels into winter months (Sole et al., 2013) and their existence alone in our baseline model is not




Figure 6. Probability density estimates from kernel smoothing of model output during all model years at nodes between 40–60 km from the ice margin. As in Figure 5, nodes that fall within murtoo routes are shown in blue, those which fall within meltwater routes (without murtos) are shown in orange, and all other nodes are shown in purple. **A)** Water pressure expressed as a percentage of overburden, *overburden%*. **B)** Sheet discharge, q_s . **C)** Water velocity, V_W . **D)** Channel discharge, Q_c . Note panels B and D have logarithmic scales.

necessarily surprising. Our fixed repetitive model forcing, fixed ice sheet geometry, and constant basal velocity (see Section 4) likely explains their repetitive interannual expression, however the the spatial pattern of overwinter channels is unexpected. The lobate geometry of the ice lobe is one possible explanation, and together with spatial variability in the climate input may result in a non-uniform concentration of meltwater within the lobe which is allowed to persist because of fixed model forcing. In a more sophisticated model setup—one including sediment dynamics; coupled ice flow and subglacial hydrology; and/or more realistic meltwater variability—the repetitive pattern of overwinter channels is unlikely to persist and overwinter channels may be absent altogether. Murtoo route distribution meanwhile appears to be a complex interplay of upstream sediment availability; meltwater input location and timing; local geology; ice velocity; and drainage characteristics (see Ahokangas et al., 2021). By predicting the apparent conditions for murtoo formation where no murtoo routes are present (e.g., in the centre of the FLDIL, 40–60 km from the ice margin, Figure 3B) it is clear that although GlaDS is capturing the broader patterns of meltwater drainage, it is failing to reproduce the exact spatial pattern of murtos themselves, likely due to an incomplete description of subglacial processes in our modelling (see Section 5.4). Nonetheless, our results clearly demonstrate the potential for GlaDS to represent subglacial hydrology beneath former ice sheets, but further research is necessary to more accurately resolve the specific spatial distribution of murtos.

520 5.4 Limitations and future work

This work represents, to the best of our knowledge, the first comparison of a process based subglacial hydrology model to specific glaciofluvial landforms and we view it as necessarily exploratory. To ensure models could run to completion with wall-time of 1–2 days and remain numerically stable across the tested range of parameters we make a number of simplifying assumptions. These include smoothing of the bed topography below the maximum resolution available, and using a relatively

525 large mesh. However, sensitivity testing indicates our conclusions are largely insensitive to topography, including a total absence of relief, and that the ice surface gradient instead imposes the dominant control on basal hydrology. Similarly, changing the mesh resolution also appears to have limited impact on our conclusions. We did not account for changes in elevation due to glacial isostatic adjustment (GIA) since 12 cal. ka. Assuming this area has been uplifted by a maximum of ~ 100 m (Ojala et al., 2013; Rosentau et al., 2021), the volume of melt delivered to the bed would have been higher during the Younger Dryas due
530 to higher temperatures ($\leq 0.75^\circ\text{C}$) at lower altitude. Accounting for this would result in higher discharge channels that persist further upglacier of those high-uplift areas. Additional uncertainty arises from our estimated (and constant) meltwater and basal melt inputs, lack of diurnal forcing, fixed and spatially uniform basal velocity, fixed conductivity parameters (in both space and time), fixed semi-circular channel geometry, assumed water turbulence, pervasive hydraulic connectivity, lack of water flux from abutting ice, and randomly seeded moulin inputs. Initial sensitivity testing of basal velocity forced to change seasonally
535 does indicate that changes in basal velocity throughout the year is important for repressurising the system each winter to more closely match borehole records (e.g., Doyle et al., 2018, 2022). Changes in geometry are also known to be important in synthetic experiments of GlaDS (see Hayden and Dow, 2023) whereas we kept ice geometry fixed here. Finally, we note that in its uncoupled configuration, GlaDS does not account for a reduction in the frictional resistance to ice flow where *overburden%* exceeds 100% or the increase in cavity closure rates that would accompany the increase in basal velocity associated with such
540 a change in friction. In reality, sustained summer *overburden%* $\geq 100\%$ would result in the decoupling of the ice from the underlying bed as is suggested to be the reason for the limited observations of deformational structures within murtoo sediment exposures (e.g., Peterson Becher and Johnson, 2021; Mäkinen et al., 2023; Hovikoski et al., 2023). Future work should seek to address some of these limitations by including, for example, a more variable climate or coupled ice dynamics whereby the frictional resistance to ice flow is allowed to vary in response to changes *overburden%* (as in Ehrenfeucht et al., 2023).

545 **6 Summary and conclusions**

In this paper we present the first application of a modern, process based subglacial hydrology model to the palaeo setting. We compared model outputs from the Glacier Drainage System model (GlaDS) against the predicted conditions associated with murtoo genesis. Murtoos are a unique glaciofluvial landform, identified throughout Finland and Sweden in terrain formerly occupied by the Fennoscandian Ice Sheet (FIS). The alternating sedimentological sequence of upper and lower flow regimes
550 preserved within murtoos suggest that they formed amongst a network of small channels and cavities subject to rapid changes in water discharge and where water pressure met or exceeded ice overburden pressure. Further, their spatial distribution, rarely found closer than 40 km from the ice margin and often found downstream of ribbed moraines and upstream of eskers, suggests that murtoos represent the glaciofluvial imprint of a spatial and/or temporal transition between distributed and channelised drainage. We modelled this system using a setup representative of the Finnish Lake District Ice Lobe (FLDIL) at the end of the
555 Younger Dryas, ~ 12 cal. ka. Our model was forced with a positive degree model representative of the palaeo climate, as well as a modified digital elevation model and reconstructed ice surface elevation representative of the same time period.

Our model outputs reproduce many of the conditions predicted for murtoo genesis including:

- i. An extensive area of water pressure at or equal to ice overburden pressure 40–60 km from the ice margin, largely robust to the range of parameters tested here.
- 560 ii. The annual evolution of a semi-distributed drainage system, which matches many of the anticipated conditions for murtoo genesis.
- iii. Modelled channels which extend 40–50 km from the ice margin upglacier into the hypothesised transitional drainage zone associated with murtoo formation. These channels also have a similar spacing and geometry to mapped eskers in the region.
- 565 iv. A statistically meaningful difference between areas of the bed without any indication of meltwater flow and areas of the bed with meltwater routes or murtoo routes.

Additionally, we find a statistically meaningful difference in water pressure, water velocity, and sheet/channel discharge, between meltwater routes and murtoo routes. We interpret this as a combination of patchy murtoo distribution and the presence of overwinter channels in our model outputs. Murtoo fields are not universally present where the conditions for their formation are predicted in our model, particularly within the centre of the FLDIL lobe, this may arise from necessary simplifications in our modelling which include the lack of sediment dynamics and/or coupling to ice flow. Nonetheless, many of our model outcomes from the baseline model, in particular the area of high water pressure 40–60 km from the ice margin, are robust across the majority of 29 sensitivity tests carried out here, in which various values for model parameters and boundary conditions were tested within a range of numerical stability. At extremely high and low values of conductivity, parameters controlling how readily water flows through the distributed or channelised system, water was evacuated from the system too easily or slowly to form meaningful channels. Channels are also restricted when the highest moulin density was tested. However, across all other tests, including random mesh geometries, alternate bed topographies, changing basal velocity, and changing moulin density, similar patterns of modelled channels and water pressures emerge. Although our system is necessarily an idealised representation of the study area—not including adjacent and abutting ice lobes, an upstream catchment area, or a coupled representation of ice dynamics and basal hydrology—this work nonetheless demonstrates the potential application of modern process based hydrology models to the palaeo setting, where model outputs can be directly compared to geomorphology and specific models of landform genesis.

570
575
580

Code and data availability. All geophysical data used to parameterise the modelling (e.g., Quaternary sediment thickness, geothermal heat flux, lake bathymetry) is available from Finnish Geological Survey's 'Hakku' service (<https://hakku.gtk.fi/?locale=en>, last accessed on 06-09-2023). The Copernicus DEM used as basal elevation is available from: <https://spacedata.copernicus.eu/collections/copernicus-digital-elevation-model> (last accessed on 06-09-2023). For our modelling we used the Ice-sheet and Sea-level System Model (Larour et al., 2012) revision 27448 available from: <https://issm.jpl.nasa.gov/> (last accessed on 06-09-2023). Murtoo field locations from Ahokangas et al. (2021), glacial landforms shapefile data from Palmu et al. (2021), model results, and example input scripts used to produce and plot those results are available at the repository linked to this manuscript (<https://doi.org/10.5281/zenodo.8344208>, Hepburn et al., 2023)

585

590 *Video supplement.* Movie A1 is available at the online repository linked to this article (<https://doi.org/10.5281/zenodo.8344208>, Hepburn
et al., 2023).

Author contributions. A.O, J.M, and C.F.D conceived the study, A.J.H designed and carried out the study and wrote the manuscript, all authors commented on the writing and helped with the analysis and interpretation.

Competing interests. The authors declare that no competing interests are present

595 *Acknowledgements.* This work forms part of the RewarD project (MUST consortium, University of Turku), funded by the Academy of
Finland (grant numbers 322243/J.M and 322252/A.O). A.J.H is funded by a 150th Anniversary Vice Chancellor's Fellowship at Aberystwyth
University, C.F.D is funded by the Canada Research Chair program (950-231237). All simulations were run on the Digital Research Alliance
of Canada compute cluster, and we thank the European Union and the Finnish Geological Survey for enabling access to the data used to
parameterise our model. We thank M. Werder for making the GlaDS model available, and we also thank M. Morlighem, J. Quinn, and J.
600 Cuzzone for their help with ISSM. Finally, we thank L. Tarasov, S. Greenwood, and an anonymous reviewer for their suggestions to improve
this paper.

References

- Abatzoglou, J. T., Dobrowski, S. Z., Parks, S. A., and Hegewisch, K. C.: TerraClimate, a high-resolution global dataset of monthly climate and climatic water balance from 1958–2015, *Scientific data*, 5, 1–12, 2018.
- 605 Ahokangas, E., Ojala, A. E., Tuunainen, A., Valkama, M., Palmu, J.-P., Kajuutti, K., and Mäkinen, J.: The distribution of glacial meltwater routes and associated murtoo fields in Finland, *Geomorphology*, 389, 107 854, 2021.
- Åkesson, H., Morlighem, M., Nisancioglu, K. H., Svendsen, J. I., and Mangerud, J.: Atmosphere-driven ice sheet mass loss paced by topography: Insights from modelling the south-western Scandinavian Ice Sheet, *Quaternary Science Reviews*, 195, 32–47, 2018.
- Amon, L., Wagner-Cremer, F., Vassiljev, J., and Veski, S.: Spring onset and seasonality patterns during the Late Glacial period in the eastern
610 Baltic region, *Climate of the Past*, 18, 2143–2153, 2022.
- Andrews, L. C., Catania, G. A., Hoffman, M. J., Gullely, J. D., Lüthi, M. P., Ryser, C., Hawley, R. L., and Neumann, T. A.: Direct observations of evolving subglacial drainage beneath the Greenland Ice Sheet, *Nature*, 514, 80–83, 2014.
- Archer, R., Ely, J. C., Heaton, T., Butcher, F. E., Hughes, A. L., and Clark, C. D.: Assessing ice sheet models against the landform record: The Likelihood of Accordant Lineations Analysis (LALA) tool, *Earth Surface Processes and Landforms*, 48, 2754–2771, 2023.
- 615 Banwell, A. F., Willis, I. C., and Arnold, N. S.: Modeling subglacial water routing at Paakitsoq, W Greenland, *Journal of Geophysical Research: Earth Surface*, 118, 1282–1295, 2013.
- Beaud, F., Flowers, G. E., and Pimentel, S.: Seasonal-scale abrasion and quarrying patterns from a two-dimensional ice-flow model coupled to distributed and channelized subglacial drainage, *Geomorphology*, 219, 176–191, 2014.
- Beaud, F., Flowers, G. E., and Venditti, J. G.: Modeling sediment transport in ice-walled subglacial channels and its implications for esker
620 formation and proglacial sediment yields, *Journal of Geophysical Research: Earth Surface*, 123, 3206–3227, 2018.
- Bingham, R. G., King, E. C., Smith, A. M., and Pritchard, H. D.: Glacial geomorphology: towards a convergence of glaciology and geomorphology, *Progress in Physical Geography*, 34, 327–355, 2010.
- Boswell, S. M., Toucanne, S., Pitel-Roudaut, M., Creyts, T. T., Eynaud, F., and Bayon, G.: Enhanced surface melting of the Fennoscandian Ice Sheet during periods of North Atlantic cooling, *Geology*, 47, 664–668, 2019.
- 625 Boulton, G. and Hagdorn, M.: Glaciology of the British Isles Ice Sheet during the last glacial cycle: form, flow, streams and lobes, *Quaternary Science Reviews*, 25, 3359–3390, 2006.
- Boulton, G. and Jones, A.: Stability of temperate ice caps and ice sheets resting on beds of deformable sediment, *Journal of Glaciology*, 24, 29–43, 1979.
- Boulton, G., Lunn, R., Vidstrand, P., and Zatsepin, S.: Subglacial drainage by groundwater-channel coupling, and the origin of esker systems: part Iglaciological observations, *Quaternary Science Reviews*, 26, 1067–1090, 2007a.
- 630 Boulton, G., Lunn, R., Vidstrand, P., and Zatsepin, S.: Subglacial drainage by groundwater–channel coupling, and the origin of esker systems: part IItheory and simulation of a modern system, *Quaternary Science Reviews*, 26, 1091–1105, 2007b.
- Boulton, G., Hagdorn, M., Maillot, P., and Zatsepin, S.: Drainage beneath ice sheets: groundwater–channel coupling, and the origin of esker systems from former ice sheets, *Quaternary Science Reviews*, 28, 621–638, 2009.
- 635 Braithwaite, R. J. and Olesen, O. B.: Calculation of glacier ablation from air temperature, West Greenland, in: *Glacier Fluctuations and Climatic Change: Proceedings of the Symposium on Glacier Fluctuations and Climatic Change, held in Amsterdam, 1–5 June 1987*, pp. 219–233, Springer, 1989.
- Budd, W., Keage, P., and Blundy, N.: Empirical studies of ice sliding, *Journal of glaciology*, 23, 157–170, 1979.

- Chandler, D., Wadham, J., Lis, G., Cowton, T., Sole, A., Bartholomew, I., Telling, J., Nienow, P., Bagshaw, E., Mair, D., et al.: Evolution of
640 the subglacial drainage system beneath the Greenland Ice Sheet revealed by tracers, *Nature Geoscience*, 6, 195–198, 2013.
- Chandler, D. M., Wadham, J. L., Nienow, P. W., Doyle, S. H., Tedstone, A. J., Telling, J., Hawkings, J., Alcock, J. D., Linhoff, B., and
Hubbard, A.: Rapid development and persistence of efficient subglacial drainage under 900 m-thick ice in Greenland, *Earth and Planetary
Science Letters*, 566, 116982, 2021.
- Chu, V. W.: Greenland ice sheet hydrology: A review, *Progress in Physical Geography*, 38, 19–54, 2014.
- 645 Clark, P. U. and Walder, J. S.: Subglacial drainage, eskers, and deforming beds beneath the Laurentide and Eurasian ice sheets, *Geological
Society of America Bulletin*, 106, 304–314, 1994.
- Cofaigh, C. Ó.: Tunnel valley genesis, *Progress in physical geography*, 20, 1–19, 1996.
- Cook, S. J., Christoffersen, P., Todd, J., Slater, D., and Chauché, N.: Coupled modelling of subglacial hydrology and calving-front melting
at Store Glacier, West Greenland, *The Cryosphere*, 14, 905–924, 2020.
- 650 Cook, S. J., Christoffersen, P., and Todd, J.: A fully-coupled 3D model of a large Greenlandic outlet glacier with evolving subglacial hydrology,
frontal plume melting and calving, *Journal of Glaciology*, 68, 486–502, 2022.
- Coughlan, M., Tóth, Z., Van Landeghem, K. J., Mccarron, S., and Wheeler, A. J.: Formational history of the Wicklow Trough: a marine-
transgressed tunnel valley revealing ice flow velocity and retreat rates for the largest ice stream draining the late-Devensian British–Irish
Ice Sheet, *Journal of Quaternary Science*, 35, 907–919, 2020.
- 655 Cuffey, K. M. and Paterson, W. S. B.: *The physics of glaciers*, Academic Press, 2010.
- Cuzzone, J. K., Schlegel, N.-J., Morlighem, M., Larour, E., Briner, J. P., Seroussi, H., and Caron, L.: The impact of model resolution on
the simulated Holocene retreat of the southwestern Greenland ice sheet using the Ice Sheet System Model (ISSM), *The Cryosphere*, 13,
879–893, 2019.
- Davison, B. J., Sole, A. J., Livingstone, S. J., Cowton, T. R., and Nienow, P. W.: The influence of hydrology on the dynamics of land-
660 terminating sectors of the Greenland ice sheet, *Frontiers in Earth Science*, 7, 10, 2019.
- De Fleurian, B., Morlighem, M., Seroussi, H., Rignot, E., van den Broeke, M. R., Kuipers Munneke, P., Mougintot, J., Smeets, P. C., and
Tedstone, A. J.: A modeling study of the effect of runoff variability on the effective pressure beneath Russell Glacier, West Greenland,
Journal of Geophysical Research: Earth Surface, 121, 1834–1848, 2016.
- Dewald, N., Lewington, E. L., Livingstone, S. J., Clark, C. D., and Storrar, R. D.: Distribution, characteristics and formation of esker
665 enlargements, *Geomorphology*, 392, 107919, 2021.
- Dewald, N., Livingstone, S. J., and Clark, C. D.: Subglacial meltwater routes of the Fennoscandian Ice Sheet, *Journal of Maps*, 18, 382–396,
2022.
- Donner, J.: The Younger Dryas age of the Salpausselkä moraines in Finland, *Bulletin of the Geological Society of Finland*, 82, 69–80, 2010.
- Dow, C., Werder, M., Nowicki, S., and Walker, R.: Modeling Antarctic subglacial lake filling and drainage cycles., *Cryosphere Discussions*,
670 9, 2015.
- Dow, C., McCormack, F., Young, D., Greenbaum, J., Roberts, J., and Blankenship, D.: Totten Glacier subglacial hydrology determined from
geophysics and modeling, *Earth and Planetary Science Letters*, 531, 115961, 2020.
- Dow, C., Ross, N., Jeofry, H., Siu, K., and Siegert, M.: Antarctic basal environment shaped by high-pressure flow through a subglacial river
system, *Nature Geoscience*, 15, 892–898, 2022.
- 675 Dow, C. F.: The role of subglacial hydrology in Antarctic ice sheet dynamics and stability: a modelling perspective, *Annals of Glaciology*,
pp. 1–6, 2023.

- Dow, C. F., Karlsson, N. B., and Werder, M. A.: Limited impact of subglacial supercooling freeze-on for Greenland ice sheet stratigraphy, *Geophysical Research Letters*, 45, 1481–1489, 2018a.
- Dow, C. F., Lee, W. S., Greenbaum, J. S., Greene, C. A., Blankenship, D. D., Poinar, K., Forrest, A. L., Young, D. A., and Zappa, C. J.: Basal
680 channels drive active surface hydrology and transverse ice shelf fracture, *Science Advances*, 4, eaao7212, 2018b.
- Dow, C. F., Werder, M., Babonis, G., Nowicki, S., Walker, R. T., Csathó, B., and Morlighem, M.: Dynamics of active subglacial lakes in
Recovery Ice Stream, *Journal of Geophysical Research: Earth Surface*, 123, 837–850, 2018c.
- Doyle, S. H., Hubbard, B., Christoffersen, P., Young, T. J., Hofstede, C., Bougamont, M., Box, J., and Hubbard, A.: Physical conditions of
fast glacier flow: 1. Measurements from boreholes drilled to the bed of Store Glacier, West Greenland, *Journal of Geophysical Research:*
685 *Earth Surface*, 123, 324–348, 2018.
- Doyle, S. H., Hubbard, B., Christoffersen, P., Law, R., Hewitt, D. R., Neufeld, J. A., Schoonman, C. M., Chudley, T. R., and Bougamont,
M.: Water flow through sediments and at the ice-sediment interface beneath Sermeq Kujalleq (Store Glacier), Greenland, *Journal of*
Glaciology, 68, 665–684, 2022.
- Ehrenfeucht, S., Morlighem, M., Rignot, E., Dow, C. F., and Mouginot, J.: Seasonal acceleration of Petermann Glacier, Greenland, from
690 changes in subglacial hydrology, *Geophysical Research Letters*, 50, e2022GL098 009, 2023.
- Fausto, R. S., Ahlstrøm, A. P., Van As, D., and Steffen, K.: Present-day temperature standard deviation parameterization for Greenland,
Journal of Glaciology, 57, 1181–1183, 2011.
- Felden, A. M., Martin, D. F., and Ng, E. G.: SUHMO: an adaptive mesh refinement SUBglacial Hydrology MOdel v1. 0, *Geoscientific Model*
Development, 16, 407–425, 2023.
- 695 Flowers, G. E.: Hydrology and the future of the Greenland Ice Sheet, *Nature communications*, 9, 2729, 2018.
- Gandy, N., Gregoire, L. J., Ely, J. C., Cornford, S. L., Clark, C. D., and Hodgson, D. M.: Exploring the ingredients required to successfully
model the placement, generation, and evolution of ice streams in the British-Irish Ice Sheet, *Quaternary Science Reviews*, 223, 105 915,
2019.
- Gandy, N., Gregoire, L. J., Ely, J. C., Cornford, S. L., Clark, C. D., and Hodgson, D. M.: Collapse of the last Eurasian Ice Sheet in the North
700 Sea modulated by combined processes of ice flow, surface melt, and marine ice sheet instabilities, *Journal of Geophysical Research: Earth*
Surface, 126, e2020JF005 755, 2021.
- García-Ruiz, J. M., Hughes, P. D., Palacios, D., and Andrés, N.: The European glacial landscapes from the main deglaciation, in: *European*
Glacial Landscapes, pp. 243–259, Elsevier, 2023.
- Gardner, A. S., Sharp, M. J., Koerner, R. M., Labine, C., Boon, S., Marshall, S. J., Burgess, D. O., and Lewis, D.: Near-surface temperature
705 lapse rates over Arctic glaciers and their implications for temperature downscaling, *Journal of Climate*, 22, 4281–4298, 2009.
- Greenwood, S. L., Clason, C. C., Helanow, C., and Margold, M.: Theoretical, contemporary observational and palaeo-perspectives on ice
sheet hydrology: processes and products, *Earth-Science Reviews*, 155, 1–27, 2016.
- Greenwood, S. L., Clason, C. C., Nyberg, J., Jakobsson, M., and Holmlund, P.: The Bothnian Sea ice stream: early Holocene retreat dynamics
of the south-central Fennoscandian Ice Sheet, *Boreas*, 46, 346–362, 2017.
- 710 GTK, Finland: Superficial deposits of Finland 1:200 000 (sediment polygons), [https://tupa.gtk.fi/paikkatieto/meta/maapera_200k.html#tun
nistamistiedo](https://tupa.gtk.fi/paikkatieto/meta/maapera_200k.html#tun
nistamistiedo), 2010.
- Harper, J., Meierbachtol, T., Humphrey, N., Saito, J., and Stansberry, A.: Generation and fate of basal meltwater during winter, western
Greenland Ice Sheet, *The Cryosphere*, 15, 5409–5421, 2021.

- Hayden, A.-M. and Dow, C. F.: Examining the effect of ice dynamic changes on subglacial hydrology through modelling of a synthetic Antarctic glacier, *Journal of Glaciology*, pp. 1–14, 2023.
- Hepburn, A., Dow, C., Ojala, A., Mäkinen, J., Ahokangas, E., Hovikoski, J., Jukka-Pekka, P., and Kajuutti, K.: Supplementary material for 'Reorganisation of subglacial drainage processes during rapid melting of the Fennoscandian Ice Sheet, <https://doi.org/10.5281/zenodo.8344208>, 2023.
- Hewitt, I. J.: Modelling distributed and channelized subglacial drainage: the spacing of channels, *Journal of Glaciology*, 57, 302–314, 2011.
- Hewitt, I. J. and Creyts, T. T.: A model for the formation of eskers, *Geophysical Research Letters*, 46, 6673–6680, 2019.
- Hill, T., Flowers, G. E., Hoffman, M. J., Bingham, D., and Werder, M. A.: Improved representation of laminar and turbulent sheet flow in subglacial drainage models, *Journal of Glaciology*, pp. 1–14, 2023.
- Hoffman, M. J., Andrews, L. C., Price, S. F., Catania, G. A., Neumann, T. A., Lüthi, M. P., Gulley, J., Ryser, C., Hawley, R. L., and Morriss, B.: Greenland subglacial drainage evolution regulated by weakly connected regions of the bed, *Nature communications*, 7, 13 903, 2016.
- Hooke, R. L.: Englacial and subglacial hydrology: a qualitative review, *Arctic and Alpine Research*, 21, 221–233, 1989.
- Hovikoski, J., Mäkinen, J., Winsemann, J., Soini, S., Kajuutti, K., Hepburn, A., and Ojala, A.: Upper-flow regime bedforms in a subglacial triangular-shaped landform (murtoo), late Pleistocene, SW Finland: Implications for flow dynamics and sediment transport in (semi-) distributed subglacial meltwater drainage systems, *Sedimentary Geology*, p. 106448, 2023.
- Hughes, A. L., Gyllencreutz, R., Lohne, Ø. S., Mangerud, J., and Svendsen, J. I.: The last Eurasian ice sheets—a chronological database and time-slice reconstruction, *DATED-1, Boreas*, 45, 1–45, 2016.
- Indrigo, C., Dow, C. F., Greenbaum, J. S., and Morlighem, M.: Drygalski Ice Tongue stability influenced by rift formation and ice morphology, *Journal of Glaciology*, 67, 243–252, 2021.
- Johnsen, S. J., Clausen, H. B., Dansgaard, W., Gundestrup, N. S., Hammer, C. U., Andersen, U., Andersen, K. K., Hvidberg, C. S., Dahl-Jensen, D., Steffensen, J. P., et al.: The $\delta^{18}\text{O}$ record along the Greenland Ice Core Project deep ice core and the problem of possible Eemian climatic instability, *Journal of Geophysical Research: Oceans*, 102, 26 397–26 410, 1997.
- Kamb, B.: Glacier surge mechanism based on linked cavity configuration of the basal water conduit system, *Journal of Geophysical Research: Solid Earth*, 92, 9083–9100, 1987.
- Karlsson, N. B. and Dahl-Jensen, D.: Response of the large-scale subglacial drainage system of Northeast Greenland to surface elevation changes, *The Cryosphere*, 9, 1465–1479, 2015.
- Karlsson, N. B., Solgaard, A. M., Mankoff, K. D., Gillet-Chaulet, F., MacGregor, J. A., Box, J. E., Citterio, M., Colgan, W. T., Larsen, S. H., Kjeldsen, K. K., et al.: A first constraint on basal melt-water production of the Greenland ice sheet, *Nature Communications*, 12, 3461, 2021.
- Kazmierczak, E., Sun, S., Coulon, V., and Pattyn, F.: Subglacial hydrology modulates basal sliding response of the Antarctic ice sheet to climate forcing, *The Cryosphere*, 16, 4537–4552, 2022.
- Kirkham, J. D., Hogan, K. A., Larter, R. D., Arnold, N. S., Ely, J. C., Clark, C. D., Self, E., Games, K., Huuse, M., Stewart, M. A., et al.: Tunnel valley formation beneath deglaciating mid-latitude ice sheets: Observations and modelling, *Quaternary Science Reviews*, p. 107680, 2022.
- Kirkham, J. D., Hogan, K. A., Larter, R. D., Self, E., Games, K., Huuse, M., Stewart, M. A., Ottesen, D., Le Heron, D. P., Lawrence, A., et al.: The infill of tunnel valleys in the central North Sea: Implications for sedimentary processes, geohazards, and ice-sheet dynamics, *Marine Geology*, 467, 107 185, 2024.

- Kleman, J., Hättestrand, C., Borgström, I., and Stroeven, A.: Fennoscandian palaeoglaciology reconstructed using a glacial geological inversion model, *Journal of glaciology*, 43, 283–299, 1997.
- Kleman, J., Hättestrand, C., Stroeven, A. P., Jansson, K. N., De Angelis, H., and Borgström, I.: Reconstruction of Palaeo-Ice Sheets-Inversion of their Glacial Geomorphological Record, *Glacier science and environmental change*, pp. 192–198, 2006.
- 755 Larour, E., Seroussi, H., Morlighem, M., and Rignot, E.: Continental scale, high order, high spatial resolution, ice sheet modeling using the Ice Sheet System Model (ISSM), *Journal of Geophysical Research: Earth Surface*, 117, 2012.
- Lehtinen, M., Nurmi, P. A., and Ramo, O.: *Precambrian Geology of Finland*, Elsevier, 2005.
- Lewington, E. L., Livingstone, S. J., Clark, C. D., Sole, A. J., and Storrar, R. D.: A model for interaction between conduits and surrounding hydraulically connected distributed drainage based on geomorphological evidence from Keewatin, Canada, *The Cryosphere*, 14, 2949–2976,
- 760 2020.
- Livingstone, S., Clark, C., Woodward, J., and Kingslake, J.: Potential subglacial lakes and meltwater drainage pathways beneath the Antarctic and Greenland ice sheets, *The Cryosphere*, 7, 1721–1740, 2013a.
- Livingstone, S. J., Clark, C. D., and Tarasov, L.: Modelling North American palaeo-subglacial lakes and their meltwater drainage pathways, *Earth and Planetary Science Letters*, 375, 13–33, 2013b.
- 765 Livingstone, S. J., Storrar, R. D., Hillier, J. K., Stokes, C. R., Clark, C. D., and Tarasov, L.: An ice-sheet scale comparison of eskers with modelled subglacial drainage routes, *Geomorphology*, 246, 104–112, 2015.
- Lunkka, J. P. and Erikkilä, A.: Behaviour of the lake district ice lobe of the Scandinavian ice sheet during the younger dryas chronozone (ca. 12 800–11 500 years ago), Tech. rep., Posiva Oy, 2012.
- Lunkka, J. P., Palmu, J.-P., and Seppänen, A.: Deglaciation dynamics of the Scandinavian Ice Sheet in the Salpausselkä zone, southern
- 770 Finland, *Boreas*, 50, 404–418, 2021.
- MacAyeal, D. R.: Large-scale ice flow over a viscous basal sediment: Theory and application to ice stream B, Antarctica, *Journal of Geophysical Research: Solid Earth*, 94, 4071–4087, 1989.
- Mäkinen, J.: Time-transgressive deposits of repeated depositional sequences within interlobate glaciofluvial (esker) sediments in Köyliö, SW Finland, *Sedimentology*, 50, 327–360, 2003.
- 775 Mäkinen, J., Kajuutti, K., Palmu, J.-P., Ojala, A., and Ahokangas, E.: Triangular-shaped landforms reveal subglacial drainage routes in SW Finland, *Quaternary Science Reviews*, 164, 37–53, 2017.
- Mäkinen, J., Kajuutti, K., Ojala, A. E., Ahokangas, E., Tuunainen, A., Valkama, M., and Palmu, J.-P.: Genesis of subglacial triangular-shaped landforms (murtoos) formed by the Fennoscandian Ice Sheet, *Earth Surface Processes and Landforms*, 2023.
- Mangerud, J., Hughes, A. L., Johnson, M. D., and Lunkka, J. P.: The Fennoscandian Ice Sheet during the Younger Dryas Stadial, in: *European*
- 780 *Glacial landscapes*, pp. 437–452, Elsevier, 2023.
- Marshall, S. J. and Sharp, M. J.: Temperature and melt modeling on the Prince of Wales ice field, Canadian High Arctic, *Journal of Climate*, 22, 1454–1468, 2009.
- McArthur, K., McCormack, F. S., and Dow, C. F.: Basal conditions of Denman Glacier from glacier hydrology and ice dynamics modeling, *The Cryosphere Discussions*, pp. 1–29, 2023.
- 785 Nick, F. M., Vieli, A., Andersen, M. L., Joughin, I., Payne, A., Edwards, T. L., Pattyn, F., and van de Wal, R. S.: Future sea-level rise from Greenland's main outlet glaciers in a warming climate, *Nature*, 497, 235–238, 2013.
- Nienow, P., Sole, A., Slater, D. A., and Cowton, T.: Recent advances in our understanding of the role of meltwater in the Greenland Ice Sheet system, *Current Climate Change Reports*, 3, 330–344, 2017.

- Nye, J.: Water at the bed of a glacier, in: International Glaciological Society, pp. 189–194, 1972.
- 790 Ojala, A. E., Palmu, J.-P., Åberg, A., Åberg, S., and Virkki, H.: Development of an ancient shoreline database to reconstruct the Litorina Sea maximum extension and the highest shoreline of the Baltic Sea basin in Finland, *Bulletin of the Geological Society of Finland*, 2013.
- Ojala, A. E., Peterson Becher, G., Mäkinen, J., Johnson, M. D., Kajuutti, K., Palmu, J.-P., Ahokangas, E., and Öhrling, C.: Ice-sheet scale distribution and morphometry of triangular-shaped hummocks (murtoos): a subglacial landform produced during rapid retreat of the Scandinavian Ice Sheet, *Annals of Glaciology*, 60, 115–126, 2019.
- 795 Ojala, A. E., Mäkinen, J., Ahokangas, E., Kajuutti, K., Valkama, M., Tuunainen, A., and Palmu, J.-P.: Diversity of murtoos and murtoo-related subglacial landforms in the Finnish area of the Fennoscandian Ice Sheet, *Boreas*, 50, 1095–1115, 2021.
- Ojala, A. E., Mäkinen, J., Kajuutti, K., Ahokangas, E., and Palmu, J.-P.: Subglacial evolution from distributed to channelized drainage: evidence from the Lake Murtoo area in SW Finland, *Earth Surface Processes and Landforms*, 47, 2877–2896, 2022.
- Palmu, J.-P., Ojala, A. E., Virtasalo, J., Putkinen, N., Kohonen, J., and Sarala, P.: Classification system of superficial (quaternary) geological units in Finland, *Developments in Map Data Management and Geological Unit Nomenclature in Finland*, 412, 115–169, 2021.
- 800 Patton, H., Hubbard, A., Andreassen, K., Auriac, A., Whitehouse, P. L., Stroeven, A. P., Shackleton, C., Winsborrow, M., Heyman, J., and Hall, A. M.: Deglaciation of the Eurasian ice sheet complex, *Quaternary Science Reviews*, 169, 148–172, 2017.
- Peterson, G., Johnson, M. D., and Smith, C. A.: Glacial geomorphology of the south Swedish uplands—focus on the spatial distribution of hummock tracts, *Journal of Maps*, 13, 534–544, 2017.
- 805 Peterson Becher, G. and Johnson, M. D.: Sedimentology and internal structure of murtoos-V-shaped landforms indicative of a dynamic subglacial hydrological system, *Geomorphology*, 380, 107 644, 2021.
- Poinar, K., Dow, C. F., and Andrews, L. C.: Long-term support of an active subglacial hydrologic system in Southeast Greenland by firm aquifers, *Geophysical Research Letters*, 46, 4772–4781, 2019.
- Putkinen, N., Eyles, N., Putkinen, S., Ojala, A. E., Palmu, J.-P., Sarala, P., Väänänen, T., Räisänen, J., Saarelainen, J., Ahtonen, N., et al.: High-resolution LiDAR mapping of glacial landforms and ice stream lobes in Finland., *Bulletin of the Geological Society of Finland*, 89, 2017.
- 810 Rada, C. and Schoof, C.: Channelized, distributed, and disconnected: subglacial drainage under a valley glacier in the Yukon, *The Cryosphere*, 12, 2609–2636, 2018.
- Rada Giacaman, C. A. and Schoof, C.: Channelized, distributed, and disconnected: spatial structure and temporal evolution of the subglacial drainage under a valley glacier in the Yukon, *The Cryosphere*, 17, 761–787, 2023.
- 815 Rampton, V.: Large-scale effects of subglacial meltwater flow in the southern Slave Province, Northwest Territories, Canada, *Canadian Journal of Earth Sciences*, 37, 81–93, 2000.
- Regnéll, C., Mangerud, J., and Svendsen, J. I.: Tracing the last remnants of the Scandinavian Ice Sheet: Ice-dammed lakes and a catastrophic outburst flood in northern Sweden, *Quaternary Science Reviews*, 221, 105 862, 2019.
- 820 Rosentau, A., Klemann, V., Bennike, O., Steffen, H., Wehr, J., Latinović, M., Bagge, M., Ojala, A., Berglund, M., Peterson Becher, G., et al.: A Holocene relative sea-level database for the Baltic Sea, *Quaternary Science Reviews*, 266, 107 071, 2021.
- Röthlisberger, H.: Water pressure in intra-and subglacial channels, *Journal of Glaciology*, 11, 177–203, 1972.
- Schenk, F., Välranta, M., Muschitiello, F., Tarasov, L., Heikkilä, M., Björck, S., Brandefelt, J., Johansson, A. V., Näslund, J.-O., and Wohlfarth, B.: Warm summers during the Younger Dryas cold reversal, *Nature Communications*, 9, 1634, 2018.
- 825 Scholzen, C., Schuler, T. V., and Gilbert, A.: Sensitivity of subglacial drainage to water supply distribution at the Kongsfjord basin, Svalbard, *The Cryosphere*, 15, 2719–2738, 2021.

- Schoof, C.: Ice-sheet acceleration driven by melt supply variability, *Nature*, 468, 803–806, 2010.
- Shackleton, C., Patton, H., Hubbard, A., Winsborrow, M., Kingslake, J., Esteves, M., Andreassen, K., and Greenwood, S. L.: Subglacial water storage and drainage beneath the Fennoscandian and Barents Sea ice sheets, *Quaternary Science Reviews*, 201, 13–28, 2018.
- 830 Sole, A., Nienow, P., Bartholomew, I., Mair, D., Cowton, T., Tedstone, A., and King, M. A.: Winter motion mediates dynamic response of the Greenland Ice Sheet to warmer summers, *Geophysical Research Letters*, 40, 3940–3944, 2013.
- Sommers, A., Meyer, C., Morlighem, M., Rajaram, H., Poinar, K., Chu, W., and Mejjia, J.: Subglacial hydrology modeling predicts high winter water pressure and spatially variable transmissivity at Helheim Glacier, Greenland, *Journal of Glaciology*, pp. 1–13, 2022.
- Stokes, C. R., Tarasov, L., Blomdin, R., Cronin, T. M., Fisher, T. G., Gyllencreutz, R., Hättestrand, C., Heyman, J., Hindmarsh, R. C., Hughes, 835 A. L., et al.: On the reconstruction of palaeo-ice sheets: Recent advances and future challenges, *Quaternary Science Reviews*, 125, 15–49, 2015.
- Storrar, R. D. and Livingstone, S. J.: Glacial geomorphology of the northern Kivalliq region, Nunavut, Canada, with an emphasis on meltwater drainage systems, *Journal of Maps*, 13, 153–164, 2017.
- Stroeven, A. P., Hättestrand, C., Kleman, J., Heyman, J., Fabel, D., Fredin, O., Goodfellow, B. W., Harbor, J. M., Jansen, J. D., Olsen, L., 840 et al.: Deglaciation of fennoscandia, *Quaternary Science Reviews*, 147, 91–121, 2016.
- Tarasov, L., Dyke, A. S., Neal, R. M., and Peltier, W. R.: A data-calibrated distribution of deglacial chronologies for the North American ice complex from glaciological modeling, *Earth and Planetary Science Letters*, 315, 30–40, 2012.
- Tedstone, A. J., Nienow, P. W., Gourmelen, N., Dehecq, A., Goldberg, D., and Hanna, E.: Decadal slowdown of a land-terminating sector of the Greenland Ice Sheet despite warming, *Nature*, 526, 692–695, 2015.
- 845 Utting, D. J., Ward, B. C., and Little, E. C.: Genesis of hummocks in glaciofluvial corridors near the Keewatin Ice Divide, Canada, *Boreas*, 38, 471–481, 2009.
- Van Boeckel, M., Van Boeckel, T., and Hall, A. M.: Late erosion pulse triggered by rapid melt in the cold-based interior of the last Fennoscandian Ice Sheet, an example from Rogan, *Earth Surface Processes and Landforms*, 47, 3376–3394, 2022.
- Van de Wal, R., Smeets, C., Boot, W., Stoffelen, M., Van Kampen, R., Doyle, S. H., Wilhelms, F., van den Broeke, M. R., Reijmer, C., 850 Oerlemans, J., et al.: Self-regulation of ice flow varies across the ablation area in south-west Greenland, *The Cryosphere*, 9, 603–611, 2015.
- van den Broeke, M., Bus, C., Ettema, J., and Smeets, P.: Temperature thresholds for degree-day modelling of Greenland ice sheet melt rates, *Geophysical Research Letters*, 37, 2010.
- van den Broeke, M. R., Kuipers Munneke, P., Noël, B., Reijmer, C., Smeets, P., van de Berg, W. J., and van Wessem, J. M.: Contrasting 855 current and future surface melt rates on the ice sheets of Greenland and Antarctica: Lessons from in situ observations and climate models, *PLOS Climate*, 2, e0000203, 2023.
- Vérité, J., Ravier, É., Bourgeois, O., Bessin, P., Livingstone, S. J., Clark, C. D., Pochat, S., and Mourgues, R.: Formation of murtoos by repeated flooding of ribbed bedforms along subglacial meltwater corridors, *Geomorphology*, 408, 108248, 2022.
- Wake, L. and Marshall, S.: Assessment of current methods of positive degree-day calculation using in situ observations from glaciated 860 regions, *Journal of Glaciology*, 61, 329–344, 2015.
- Walder, J. S.: Hydraulics of subglacial cavities, *Journal of Glaciology*, 32, 439–445, 1986.
- Weertman, J.: General theory of water flow at the base of a glacier or ice sheet, *Reviews of Geophysics*, 10, 287–333, 1972.
- Werder, M. A., Hewitt, I. J., Schoof, C. G., and Flowers, G. E.: Modeling channelized and distributed subglacial drainage in two dimensions, *Journal of Geophysical Research: Earth Surface*, 118, 2140–2158, 2013.

- 865 Wright, P. J., Harper, J. T., Humphrey, N. F., and Meierbachtol, T. W.: Measured basal water pressure variability of the western Greenland Ice Sheet: Implications for hydraulic potential, *Journal of Geophysical Research: Earth Surface*, 121, 1134–1147, 2016.
- Yang, K. and Smith, L. C.: Internally drained catchments dominate supraglacial hydrology of the southwest Greenland Ice Sheet, *Journal of Geophysical Research: Earth Surface*, 121, 1891–1910, 2016.

Appendix: Contents

870 This file contains supplementary information for ‘*The organisation of subglacial drainage during the demise of the Finnish Lake District Ice-Lobe*’

Appendix A. Additional model description for the Glacier Drainage System model (GlaDS Werder et al., 2013).

Table A1. List of input values for GlaDS, values highlighted in bold indicate those used for sensitivity testing and a range of values is provided. Note, in all instances *sheet* refers to the subglacial drainage system.

875 **Movie A1.** Evolution of the system with respect to *overburden*_% through time in the baseline model run. Model years 15–17 were arbitrarily chosen to illustrate the transient state of the system through several melt season cycles. Channels are shown as black lines where Q_c exceeds $1 \text{ m}^3 \text{ s}^{-1}$.

Figure A1. The distribution of meltwater routes, murtoo routes, and sediment in the Finnish Lake District Ice Lobe. **A)** Meltwater routes and murtoo routes as mapped by Ahokangas et al. (2021). There is a general absence of murtoos in the
880 centre of the lobe 40–60 km from the ice margin. **B)** Sediment cover (GTK, Finland, 2010) showing thin sediment thickness in the terrain from which murtoos appear absent.

Figure A2 Overwinter channels and their influence on subsequent summer drainage. **A)** Modelled channels in January, model year 15 in the baseline run (black lines). Channels are present overwinter in the outer two thirds of the FLDIL. **B)** Modelled channels in June, model year 15, near to the start of the melt season. Note that channels in the outer two thirds are already
885 well established compared to those in the central third of the FLDIL. **C)** Modelled channels in January, model year 2016. Note that winter channels are now visible in the central third of the FLDIL. In all panels, background panels shows water pressure as a percentage of ice overburden pressure. Murtoo fields are shown as inverted black triangles. Each channel is defined where the median channel discharge, Q_c , exceeds $1 \text{ m}^3 \text{ s}^{-1}$.

Figure A3. Median *overburden*_%, channel discharge, Q_c , and sheet discharge, q_s per timestep over the full length of the
890 baseline model run.

Figure A4–A32. Comparison of the median summer system for the range of sensitivity parameters against the baseline model run. **A)** Water pressure expressed as a percentage of overburden pressure, *overburden*_%. Channels are shown as black lines where median discharge exceeds $1 \text{ m}^3 \text{ s}^{-1}$. **B)** Baseline median summer *overburden*_% minus the tested median summer *overburden*_%. The same figure caption applies for Figures A4–A32.

895 **Figure A33.** Boxplots of model parameters grouped by month for overburden (*overburden*_%, **A**), sheet discharge (q_s , **B**), water velocity (V_W , **C**), and channel discharge (Q_c , **D**) during all model years at nodes between 40–60 km from the ice margin. As in Figure 5, nodes that fall within meltwater routes which do host murtoos (murtoo routes) are shown in blue, nodes which fall within mapped meltwater routes that do not contain murtoo fields (meltwater routes) are shown in orange, and all other nodes are shown in purple. Medians for each group are shown as black circles, and ‘outliers’—defined as points more than 150% of
900 the interquartile range away from the upper and lower quartile—are shown as crosses.

Tables A2–A5. Tukey-Kramer HSD test results for *overburden%* (Table A2), q_s (Table A3), Q_c (Table A4), and V_w (Table A5) in meltwater routes, murtoo routes, and non-meltwater routes between 40–60 km from the ice margin. The upper and lower limits describe the 95% confidence intervals for the true mean difference, A-B is the difference between group means.

Appendix A: Additional model description

905 In GlaDS (Werder et al., 2013) water flux, q_s , through the distributed system is driven by the hydraulic potential gradient, $\nabla\phi$, along with the sheet conductivity, k_s

$$q_s = -k_s h^\alpha |\nabla\phi|^\beta \nabla\phi, \quad (\text{A1})$$

where the flow exponents, $\alpha = 5/4$ and $\beta = 3/2$ describe fully turbulent flow in the Darcy-Weishbach law, and h is the sheet thickness. The sheet thickness evolves through time given by

$$910 \quad \frac{\delta h}{\delta t} = w - v, \quad (\text{A2})$$

for functions w and v which describe the cavity opening and closing rate respectively (Walder, 1986; Kamb, 1987). Basal sliding opens cavities at a rate given by the basal sliding speed, U_b acting over basal bumps with a height, h_r through

$$w(h) = \begin{cases} U_b (h_r - h) / l_r & \text{if } h < h_r \\ 0 & \text{otherwise} \end{cases} \quad (\text{A3})$$

where l_r is the typical horizontal cavity spacing. In turn, viscous ice deformation leads to cavity closure, which is related to the
915 effective pressure, N by

$$v(h, N) = Ah|N|^{n-1}N, \quad (\text{A4})$$

where A is the rate factor, or the rheological constant of ice, multiplied by a first order geometrical factor, and n is the Glen's flow law exponent. Sheet elements exchange water with channels and the cross sectional area of these channels S , evolves through time due to the dissipation of potential energy, Π , sensible heat exchange, Ξ , and cavity closure rates due to viscous
920 ice creep v_c

$$\frac{\partial S}{\partial t} = \frac{\Xi - \Pi}{\rho_i L} - v_c, \quad (\text{A5})$$

where ρ_i is the ice density and L is the latent heat of fusion. The default parameters used here, as well as those sensitivity tested, are listed in Table A1.

Table A1. List of input values for GlADS, values highlighted in bold indicate those used for sensitivity testing and a range of values is provided. Note, in all instances *sheet* refers to the subglacial drainage system.

Symbol	Description	Default value	Tested range	Units
ρ_i	ice density	918		kg m ³
ρ_w	water density	1000		kg m ³
g	gravitational acceleration	9.81		m s ⁻²
n	Glen's flow law exponent	3		
a	basal friction coefficient	0–120		(Pa a ⁻¹) ^{1/2}
A	rate factor	1.7×10^{-24}		s ⁻¹ Pa ⁻³
L	latent heat	3.34×10^5		J kg ⁻¹
c_t	pressure melt coefficient	7.5×10^{-8}		KPa ⁻¹
c_w	heat capacity of water	4.22×10^3		Jkg ⁻¹ K ⁻¹
α	first sheet flow exponent	5/4		
β	second sheet flow exponent	3/2		
α_c	first channel flow exponent	5/4		
β_c	second channel flow exponent	3/2		
k_s	sheet conductivity	10^{-4}	10^{-2} – 10^{-5}	m ^{7/4} kg ^{-1/2}
k_c	channel conductivity	10^{-1}	5×10^{-1} – 10^{-3}	m ^{3/2} kg ^{-1/2}
E_{vr}	englacial void ratio	10^{-4}	10^{-3} – 10^{-5}	
l_c	sheet width below channel	2		m
A_m	moulin cross-sectional area	10		m ²
l_r	cavity spacing	2		m
h_r	basal bump height	0.085	0.05–0.1	m
b_{melt}	basal melt rate	5×10^{-3}	1 – 7×10^{-3}	m yr ⁻¹
U_b	mean annual basal velocity [†]	150	100–200	m yr ⁻¹
$N_{moulins}$	number of moulins [*]	2500	1000–4000	

[†] We tested both a transient and temporally constant basal velocity within these given ranges for mean annual basal velocity

^{*} We also ran an experiment in which melt was routed directly to the bed at each node (SHEET)




Figure A1. The distribution of meltwater routes, murtoo routes, and sediment in the Finnish Lake District Ice Lobe. **A)** Meltwater routes and murtoo routes as mapped by Ahokangas et al. (2021). There is a general absence of murtoos in the centre of the lobe 40–60 km from the ice margin. **B)** Sediment cover (GTK, Finland, 2010) showing thin sediment thickness in the terrain from which murtoos appear absent.




Figure A2. Overwinter channels and their influence on subsequent summer drainage. **A)** Modelled channels in January, model year 15 in the baseline run (black lines). Channels are present overwinter in the outer two thirds of the FLDIL. **B)** Modelled channels in June, model year 15, near to the start of the melt season. Note that channels in the outer two thirds are already well established compared to those in the central third of the FLDIL. **C)** Modelled channels in January, model year 2016. Note that winter channels are now visible in the central third of the FLDIL. In all panels, background panels shows water pressure as a percentage of ice overburden pressure. Murtoo fields are shown as inverted black triangles. Each channel is defined where the median channel discharge, $Q_{c,m}$, exceeds $1 \text{ m}^3 \text{ s}^{-1}$.




Figure A3. Median *overburden*%, channel discharge, Q_c , and sheet discharge, q_s over the full length of the baseline model run.




Figure A4. Comparison of the median summer system for sheet conductivity, $k_s = 10^{-2} \text{ m}^{7/4} \text{ kg}^{-1/2}$ against the baseline model run ($k_s = 10^{-4} \text{ m}^{7/4} \text{ kg}^{-1/2}$). **A)** Water pressure expressed as a percentage of overburden pressure, $overburden_{\%}$. Channels are shown as black lines where median discharge exceeds $1 \text{ m}^3 \text{ s}^{-1}$. **B)** Baseline median summer $overburden_{\%}$ minus the $k_s = 10^{-2} \text{ m}^{7/4} \text{ kg}^{-1/2}$ median summer $overburden_{\%}$. The same figure caption applies for Figures A4–A32




Figure A5. Comparison of the median summer system for sheet conductivity, $k_s = 10^{-3} \text{ m}^{7/4} \text{ kg}^{-1/2}$ against the baseline model run ($k_s = 10^{-4} \text{ m}^{7/4} \text{ kg}^{-1/2}$). The same figure caption as Figure A4 applies.




Figure A6. Comparison of the median summer system for sheet conductivity, $k_s = 10^{-5} \text{ m}^{7/4} \text{ kg}^{-1/2}$ against the baseline model run ($k_s = 10^{-4} \text{ m}^{7/4} \text{ kg}^{-1/2}$). The same figure caption as Figure A4 applies.




Figure A7. Comparison of the median summer system for channel conductivity, $k_c = 5 \times 10^{-1} \text{ m}^{3/2} \text{ kg}^{-1/2}$ against the baseline model run ($k_c = 10^{-1} \text{ m}^{3/2} \text{ kg}^{-1/2}$). The same figure caption as Figure A4 applies.




Figure A8. Comparison of the median summer system for channel conductivity, $k_c = 5 \times 10^{-2} \text{ m}^{3/2} \text{ kg}^{-1/2}$ against the baseline model run ($k_c = 10^{-1} \text{ m}^{3/2} \text{ kg}^{-1/2}$). The same figure caption as Figure A4 applies.




Figure A9. Comparison of the median summer system for channel conductivity, $k_c = 10^{-3} \text{ m}^{3/2} \text{ kg}^{-1/2}$ against the baseline model run ($k_c = 10^{-1} \text{ m}^{3/2} \text{ kg}^{-1/2}$). The same figure caption as Figure A4 applies.




Figure A10. Comparison of the median summer system for moulin frequency, $N_{moulins} = 1000$ against the baseline model run ($N_{moulins} = 2500$). The same figure caption as Figure A4 applies.




Figure A11. Comparison of the median summer system for moulin frequency, $N_{moulins} = 4000$ against the baseline model run ($N_{moulins} = 2500$). The same figure caption as Figure A4 applies.




Figure A12. Comparison of the median summer system for where water was directly input at every nodes against the baseline model run ($N_{moulins} = 2500$). The same figure caption as Figure A4 applies.




Figure A13. Comparison of the median summer system for a second random distribution of moulin frequency, $N_{moulins} = 2500$ against the baseline model run ($N_{moulins} = 2500$). The same figure caption as Figure A4 applies.




Figure A14. Comparison of the median summer system for a third random distribution of moulin frequency, $N_{moulins} = 2500$ against the baseline model run ($N_{moulins} = 2500$). The same figure caption as Figure A4 applies.




Figure A15. Comparison of the median summer system for basal melt rate, $b_{melt} = 1 \times 10^{-3} \text{ m yr}^{-1}$ against the baseline model run ($b_{melt} = 5 \times 10^{-3} \text{ m yr}^{-1}$). The same figure caption as Figure A4 applies.




Figure A16. Comparison of the median summer system for basal melt rate, $b_{melt} = 1 \times 10^{-3} \text{ m yr}^{-1}$ against the baseline model run ($b_{melt} = 5 \times 10^{-3} \text{ m yr}^{-1}$). The same figure caption as Figure A4 applies.




Figure A17. Comparison of the median summer system for basal bump height, $h_r = 0.1 \text{ m}$ against the baseline model run ($h_r = 0.085 \text{ m}$). The same figure caption as Figure A4 applies.




Figure A18. Comparison of the median summer system for basal bump height, $h_r = 0.05$ m against the baseline model run ($h_r = 0.085$ m). The same figure caption as Figure A4 applies.




Figure A19. Comparison of a mesh that is not refined with respect to elevation against the baseline model run. The same figure caption as Figure A4 applies.




Figure A20. Comparison of a coarser mesh (edge length ~ 5 km) against the baseline model run. The same figure caption as Figure A4 applies.




Figure A21. Comparison of a refined mesh (minimum edge length ≈ 300 m) < 80 km from the ice margin against the baseline model run. The same figure caption as Figure A4 applies.




Figure A22. Comparison of a flat bed against the baseline model run. The same figure caption as Figure A4 applies.




Figure A23. Comparison of a modern mesh (without subtracting Quaternary sediment thickness) against the baseline model run. The same figure caption as Figure A4 applies.




Figure A24. Comparison including lake bathymetry against the baseline model run. The same figure caption as Figure A4 applies.




Figure A25. Comparison of a 30 m deep water body at the ice margin boundary against the baseline model run (land-terminating). The same figure caption as Figure A4 applies.




Figure A26. Comparison of the median summer system for an englacial void ratio, $E_{vr} = 10^{-3}$ against the baseline model run ($E_{vr} = 10^{-4}$). The same figure caption as Figure A4 applies.




Figure A27. Comparison of the median summer system for an englacial void ratio, $E_{vr} = 10^{-5}$ against the baseline model run ($E_{vr} = 10^{-4}$). The same figure caption as Figure A4 applies.




Figure A28. Comparison of the median summer system for a fixed basal velocity, $U_b = 100 \text{ m yr}^{-1}$ against the baseline model run ($U_b = 150 \text{ m yr}^{-1}$). The same figure caption as Figure A4 applies.




Figure A29. Comparison of the median summer system for a fixed basal velocity, $U_b = 200 \text{ m yr}^{-1}$ against the baseline model run ($U_b = 150 \text{ m yr}^{-1}$). The same figure caption as Figure A4 applies.




Figure A30. Comparison of the median summer system for a transient basal velocity, U_b with a median $U_b = 150 \text{ m yr}^{-1}$ against the fixed baseline model run ($U_b = 150 \text{ m yr}^{-1}$). The same figure caption as Figure A4 applies.




Figure A31. Comparison of the median summer system for a transient basal velocity, U_b with a median $U_b = 100 \text{ m yr}^{-1}$ against the fixed baseline model run ($U_b = 150 \text{ m yr}^{-1}$). The same figure caption as Figure A4 applies.




Figure A32. Comparison of the median summer system for a transient basal velocity, U_b with a median $U_b = 200 \text{ m yr}^{-1}$ against the fixed baseline model run ($U_b = 150 \text{ m yr}^{-1}$). The same figure caption as Figure A4 applies.




Figure A33. Boxplots of model parameters grouped by month for overburden ($overburden\%$, **A**), sheet discharge (q_s , **B**), water velocity (V_w , **C**), and channel discharge (Q_c , **D**) during all model years at nodes between 40–60 km from the ice margin. As in Figure 5, nodes that fall within meltwater routes which do host murtoos (Murtoo free MRs) are shown in blue, nodes which fall within mapped meltwater routes that do not contain murtoo fields (Murtoo hosting MRs) are shown in orange, and all other nodes are shown in purple. Medians for each group are shown as black circles, and ‘outliers’—defined as points more than 150% of the interquartile range away from the upper and lower quartile—are shown as crosses.

Table A2. Tukey-Kramer HSD test of *overburden%* in meltwater routes, murtoo routes, and non-meltwater routes between 40–60 km from the ice margin. The upper and lower limits describe the 95% confidence intervals for the true mean difference, A-B is the difference between group means.

Month	Group A	Group B	Lower Limit	A-B	Upper limit	P-Value
January	meltwater route	murtoo route	-1.71	-1.30	-0.89	0.00
	all other nodes	murtoo route	-10.49	-10.12	-9.75	0.00
	meltwater route	all other nodes	8.58	8.82	9.06	0.00
February	meltwater route	murtoo route	-1.62	-1.21	-0.80	0.00
	all other nodes	murtoo route	-9.87	-9.50	-9.12	0.00
	meltwater route	all other nodes	8.05	8.29	8.52	0.00
March	meltwater route	murtoo route	-1.58	-1.17	-0.76	0.00
	all other nodes	murtoo route	-9.30	-8.93	-8.55	0.00
	meltwater route	all other nodes	7.52	7.75	7.99	0.00
April	meltwater route	murtoo route	-1.55	-1.14	-0.73	0.00
	all other nodes	murtoo route	-8.33	-7.96	-7.58	0.00
	meltwater route	all other nodes	6.58	6.82	7.06	0.00
May	meltwater route	murtoo route	-1.10	-0.69	-0.28	0.00
	all other nodes	murtoo route	-5.26	-4.89	-4.51	0.00
	meltwater route	all other nodes	3.96	4.20	4.43	0.00
June	meltwater route	murtoo route	0.70	1.12	1.54	0.00
	all other nodes	murtoo route	-8.56	-8.17	-7.79	0.00
	meltwater route	all other nodes	9.05	9.30	9.54	0.00
July	meltwater route	murtoo route	0.10	0.52	0.93	0.00
	all other nodes	murtoo route	-18.72	-18.34	-17.96	0.00
	meltwater route	all other nodes	18.61	18.85	19.10	0.00
August	meltwater route	murtoo route	-2.01	-1.59	-1.18	0.00
	all other nodes	murtoo route	-18.81	-18.43	-18.04	0.00
	meltwater route	all other nodes	16.59	16.83	17.08	0.00
September	meltwater route	murtoo route	-2.30	-1.88	-1.47	0.00
	all other nodes	murtoo route	-14.47	-14.09	-13.71	0.00
	meltwater route	all other nodes	11.97	12.21	12.45	0.00
October	meltwater route	murtoo route	-2.14	-1.72	-1.31	0.00
	all other nodes	murtoo route	-12.49	-12.11	-11.73	0.00
	meltwater route	all other nodes	10.15	10.39	10.63	0.00
November	meltwater route	murtoo route	-1.99	-1.57	-1.15	0.00
	all other nodes	murtoo route	-12.06	-11.68	-11.30	0.00
	meltwater route	all other nodes	9.87	10.11	10.36	0.00
December	meltwater route	murtoo route	-1.88	-1.46	-1.04	0.00
	all other nodes	murtoo route	-11.56	-11.17	-10.79	0.00
	meltwater route	all other nodes	9.47	9.72	9.96	0.00

Table A3. Tukey-Kramer HSD test of q_s in meltwater routes, murtoo routes, and non-meltwater routes between 40–60 km from the ice margin. The upper and lower limits describe the 95% confidence intervals for the true mean difference, A-B is the difference between group means.

Month	Group A	Group B	Lower Limit	A-B	Upper limit	P-Value
January	meltwater route	murtoo route	-5.07×10^{-7}	2.89×10^{-6}	6.28×10^{-6}	0.27
	all other nodes	murtoo route	-1.57×10^{-5}	-1.26×10^{-5}	-9.54×10^{-7}	0.00
	meltwater route	all other nodes	1.36×10^{-5}	1.55×10^{-5}	1.75×10^{-5}	0.00
February	meltwater route	murtoo route	-1.23×10^{-6}	2.19×10^{-6}	5.6×10^{-6}	0.88
	all other nodes	murtoo route	-1.39×10^{-5}	-1.08×10^{-5}	-7.69×10^{-6}	0.00
	meltwater route	all other nodes	1.1×10^{-5}	1.3×10^{-5}	1.5×10^{-5}	0.00
March	meltwater route	murtoo route	-2.05×10^{-6}	1.36×10^{-6}	4.77×10^{-6}	0.99
	all other nodes	murtoo route	-1.26×10^{-5}	-9.49×10^{-6}	-6.37×10^{-6}	0.00
	meltwater route	all other nodes	8.86×10^{-6}	1.08×10^{-5}	1.28×10^{-5}	0.00
April	meltwater route	murtoo route	-2.66×10^{-6}	7.6×10^{-7}	4.18×10^{-6}	0.99
	all other nodes	murtoo route	-1.14×10^{-5}	-8.28×10^{-6}	-5.15×10^{-6}	0.00
	meltwater route	all other nodes	7.05×10^{-6}	9.04×10^{-6}	1.10×10^{-5}	0.00
May	meltwater route	murtoo route	-2.92×10^{-6}	4.99×10^{-7}	3.92×10^{-6}	0.99
	all other nodes	murtoo route	-7.11×10^{-6}	-3.98×10^{-6}	-8.53×10^{-5}	0.00
	meltwater route	all other nodes	2.49×10^{-6}	4.48×10^{-6}	6.47×10^{-6}	0.00
June	meltwater route	murtoo route	9.57×10^{-7}	4.46×10^{-6}	7.97×10^{-6}	0.00
	all other nodes	murtoo route	6.04×10^{-6}	9.25×10^{-6}	1.25×10^{-5}	0.00
	meltwater route	all other nodes	-6.83×10^{-6}	-4.79×10^{-6}	-2.75×10^{-6}	0.00
July	meltwater route	murtoo route	1.36×10^{-5}	1.7×10^{-5}	2.05×10^{-5}	0.00
	all other nodes	murtoo route	9.95×10^{-6}	1.31×10^{-5}	1.63×10^{-5}	0.00
	meltwater route	all other nodes	1.89×10^{-6}	3.9×10^{-6}	5.91×10^{-6}	0.00
August	meltwater route	murtoo route	1.82×10^{-5}	2.17×10^{-5}	2.52×10^{-5}	0.00
	all other nodes	murtoo route	-9.99×10^{-6}	-6.79×10^{-6}	-3.59×10^{-6}	0.00
	meltwater route	all other nodes	2.65×10^{-5}	2.85×10^{-5}	3.05×10^{-5}	0.00
September	meltwater route	murtoo route	1.06×10^{-5}	1.41×10^{-5}	1.75×10^{-5}	0.00
	all other nodes	murtoo route	-2.33×10^{-5}	-2.02×10^{-5}	1.7×10^{-5}	0.00
	meltwater route	all other nodes	3.22×10^{-5}	3.42×10^{-5}	3.62×10^{-5}	0.00
October	meltwater route	murtoo route	5.67×10^{-6}	9.15×10^{-6}	1.26×10^{-5}	0.00
	all other nodes	murtoo route	-2.44×10^{-5}	-2.12×10^{-5}	-1.8×10^{-5}	0.00
	meltwater route	all other nodes	2.84×10^{-5}	3.04×10^{-5}	3.24×10^{-5}	0.00
November	meltwater route	murtoo route	2.52×10^{-6}	6×10^{-6}	9.48×10^{-6}	0.00
	all other nodes	murtoo route	-2.14×10^{-5}	-1.82×10^{-5}	-1.50×10^{-5}	0.00
	meltwater route	all other nodes	2.22×10^{-5}	2.42×10^{-5}	2.62×10^{-5}	0.00
December	meltwater route	murtoo route	7.23×10^{-7}	4.22×10^{-6}	7.71×10^{-6}	0.00
	all other nodes	murtoo route	-1.87×10^{-5}	-1.55×10^{-5}	1.23×10^{-5}	0.00
	meltwater route	all other nodes	1.77×10^{-5}	1.97×10^{-5}	2.18×10^{-5}	0.00

Table A4. Tukey-Kramer HSD test of Q_c in meltwater routes, murtoo routes, and non-meltwater routes between 40–60 km from the ice margin. The upper and lower limits describe the 95% confidence intervals for the true mean difference, A-B is the difference between group means.

Month	Group A	Group B	Lower Limit	A-B	Upper limit	P-Value
January	meltwater route	murtoo route	-7.57×10^{-3}	6.51×10^{-4}	8.87×10^{-3}	0.99
	all other nodes	murtoo route	-6.19×10^{-3}	1.34×10^{-3}	8.86×10^{-3}	0.99
	meltwater route	all other nodes	-5.47×10^{-3}	-6.86×10^{-4}	4.1×10^{-3}	0.99
February	meltwater route	murtoo route	-7.78×10^{-3}	5.08×10^{-4}	8.79×10^{-3}	0.99
	all other nodes	murtoo route	-6.72×10^{-3}	5.08×10^{-4}	8.44×10^{-3}	0.99
	meltwater route	all other nodes	-5.18×10^{-3}	-3.53×10^{-4}	4.47×10^{-3}	0.99
March	meltwater route	murtoo route	-7.86×10^{-3}	4.06×10^{-4}	8.677×10^{-3}	0.99
	all other nodes	murtoo route	-6.93×10^{-3}	6.3×10^{-4}	8.19×10^{-3}	0.99
	meltwater route	all other nodes	-5.04×10^{-3}	-2.24×10^{-4}	4.59×10^{-5}	0.99
April	meltwater route	murtoo route	-7.98×10^{-3}	3.07×10^{-4}	8.59×10^{-3}	0.99
	all other nodes	murtoo route	-7.10×10^{-3}	4.83×10^{-4}	8.06×10^{-3}	0.99
	meltwater route	all other nodes	-5×10^{-3}	-1.76×10^{-4}	4.65×10^{-3}	0.99
May	meltwater route	murtoo route	-8.02×10^{-3}	2.60×10^{-4}	8.54×10^{-3}	0.99
	all other nodes	murtoo route	-5.58×10^{-3}	2.01×10^{-3}	9.59×10^{-3}	0.99
	meltwater route	all other nodes	-6.57×10^{-3}	-1.75×10^{-3}	3.08×10^{-3}	0.99
June	meltwater route	murtoo route	-7.93×10^{-3}	5.65×10^{-4}	9.06×10^{-3}	0.99
	all other nodes	murtoo route	7.28×10^{-3}	1.51×10^{-2}	2.28×10^{-2}	0.00
	meltwater route	all other nodes	-1.94×10^{-2}	-1.45×10^{-2}	-9.54×10^{-3}	0.00
July	meltwater route	murtoo route	-6.22×10^{-3}	2.14×10^{-3}	1.05×10^{-2}	0.99
	all other nodes	murtoo route	3.20×10^{-2}	3.97×10^{-2}	4.73×10^{-2}	0.00
	meltwater route	all other nodes	-4.24×10^{-2}	-3.75×10^{-2}	-3.27×10^{-2}	0.00
August	meltwater route	murtoo route	-4.45×10^{-3}	4.02×10^{-3}	1.25×10^{-2}	0.99
	all other nodes	murtoo route	3.97×10^{-2}	4.74×10^{-2}	5.52×10^{-2}	0.00
	meltwater route	all other nodes	-4.84×10^{-2}	-4.34×10^{-2}	-3.85×10^{-2}	0.00
September	meltwater route	murtoo route	-4.64×10^{-3}	3.75×10^{-3}	1.21×10^{-2}	0.99
	all other nodes	murtoo route	2.24×10^{-2}	3.01×10^{-2}	3.78×10^{-2}	0.00
	meltwater route	all other nodes	-3.12×10^{-2}	-2.63×10^{-2}	-2.15×10^{-2}	0.00
October	meltwater route	murtoo route	-6.24×10^{-3}	2.19×10^{-3}	1.06×10^{-2}	0.99
	all other nodes	murtoo route	2.22×10^{-3}	9.94×10^{-3}	1.77×10^{-2}	0.00
	meltwater route	all other nodes	-1.27×10^{-2}	-7.75×10^{-3}	-2.84×10^{-3}	0.00
November	meltwater route	murtoo route	-7.16×10^{-3}	1.27×10^{-3}	9.70×10^{-3}	0.99
	all other nodes	murtoo route	-4.08×10^{-3}	3.63×10^{-3}	1.13×10^{-2}	0.99
	meltwater route	all other nodes	-7.27×10^{-3}	-2.36×10^{-3}	2.55×10^{-3}	0.99
December	meltwater route	murtoo route	-7.56×10^{-3}	9.10×10^{-4}	9.38×10^{-3}	0.99
	all other nodes	murtoo route	-5.57×10^{-3}	2.18×10^{-3}	9.94×10^{-3}	0.99
	meltwater route	all other nodes	-6.21×10^{-3}	-1.27×10^{-3}	3.66×10^{-3}	0.99

Table A5. Tukey-Kramer HSD test of V_W in meltwater routes, murtoo routes, and non-meltwater routes between 40–60 km from the ice margin. The upper and lower limits describe the 95% confidence intervals for the true mean difference, A-B is the difference between group means.

Month	Group A	Group B	Lower Limit	A-B	Upper limit	P-Value
January	meltwater route	murtoo route	1.41×10^{-6}	7.42×10^{-6}	1.34×10^{-5}	0.00
	all other nodes	murtoo route	-5.63×10^{-5}	-5.08×10^{-5}	-4.53×10^{-5}	0.00
	meltwater route	all other nodes	5.47×10^{-5}	5.82×10^{-5}	6.17×10^{-5}	0.00
February	meltwater route	murtoo route	1.11×10^{-6}	7.17×10^{-6}	1.32×10^{-5}	0.00
	all other nodes	murtoo route	-4.63×10^{-5}	-4.08×10^{-5}	-3.53×10^{-5}	0.00
	meltwater route	all other nodes	4.44×10^{-5}	4.80×10^{-5}	5.15×10^{-5}	0.00
March	meltwater route	murtoo route	7.90×10^{-7}	6.83×10^{-6}	1.29×10^{-5}	0.01
	all other nodes	murtoo route	-3.88×10^{-5}	-3.32×10^{-5}	-2.77×10^{-5}	0.00
	meltwater route	all other nodes	3.65×10^{-5}	4.01×10^{-5}	4.36×10^{-5}	0.00
April	meltwater route	murtoo route	1.06×10^{-6}	7.11×10^{-6}	1.32×10^{-5}	0.00
	all other nodes	murtoo route	-3.26×10^{-5}	-2.71×10^{-5}	-2.16×10^{-5}	0.00
	meltwater route	all other nodes	3.07×10^{-5}	3.42×10^{-5}	3.77×10^{-5}	0.00
May	meltwater route	murtoo route	1.73×10^{-6}	7.78×10^{-6}	1.38×10^{-5}	0.00
	all other nodes	murtoo route	-6.47×10^{-6}	-9.33×10^{-7}	4.61×10^{-6}	1.00
	meltwater route	all other nodes	5.19×10^{-6}	8.72×10^{-6}	1.22×10^{-5}	0.00
June	meltwater route	murtoo route	1.22×10^{-5}	1.84×10^{-5}	2.46×10^{-5}	0.00
	all other nodes	murtoo route	4.12×10^{-5}	4.69×10^{-5}	5.26×10^{-5}	0.00
	meltwater route	all other nodes	-3.21×10^{-5}	-2.85×10^{-5}	-2.48×10^{-5}	0.00
July	meltwater route	murtoo route	2.47×10^{-5}	3.09×10^{-5}	3.70×10^{-5}	0.00
	all other nodes	murtoo route	7.34×10^{-7}	6.33×10^{-6}	1.19×10^{-5}	0.01
	meltwater route	all other nodes	2.10×10^{-5}	2.45×10^{-5}	2.81×10^{-5}	0.00
August	meltwater route	murtoo route	1.46×10^{-5}	2.08×10^{-5}	2.70×10^{-5}	0.00
	all other nodes	murtoo route	-1.05×10^{-4}	-9.90×10^{-5}	-9.34×10^{-5}	0.00
	meltwater route	all other nodes	1.16×10^{-4}	1.20×10^{-4}	1.23×10^{-4}	0.00
September	meltwater route	murtoo route	2.52×10^{-6}	8.65×10^{-6}	1.48×10^{-5}	0.00
	all other nodes	murtoo route	-1.19×10^{-4}	-1.14×10^{-4}	-1.08×10^{-4}	0.00
	meltwater route	all other nodes	1.19×10^{-4}	1.22×10^{-4}	1.26×10^{-4}	0.00
October	meltwater route	murtoo route	8.15×10^{-6}	1.43×10^{-5}	2.05×10^{-5}	0.00
	all other nodes	murtoo route	-9.90×10^{-5}	-9.34×10^{-5}	-8.77×10^{-5}	0.00
	meltwater route	all other nodes	1.04×10^{-4}	1.08×10^{-4}	1.11×10^{-4}	0.00
November	meltwater route	murtoo route	6.68×10^{-6}	1.28×10^{-5}	1.90×10^{-5}	0.00
	all other nodes	murtoo route	-8.03×10^{-5}	-7.46×10^{-5}	-6.90×10^{-5}	0.00
	meltwater route	all other nodes	8.39×10^{-5}	8.75×10^{-5}	9.10×10^{-5}	0.00
December	meltwater route	murtoo route	3.95×10^{-6}	1.01×10^{-5}	1.63×10^{-5}	0.00
	all other nodes	murtoo route	-6.79×10^{-5}	-6.22×10^{-5}	-5.65×10^{-5}	0.00
	meltwater route	all other nodes	6.87×10^{-5}	7.23×10^{-5}	7.59×10^{-5}	0.00