

# Variability and Predictability of a reduced-order land atmosphere coupled model

Anupama K Xavier<sup>1,2</sup>, Jonathan Demaeyer<sup>1</sup>, and Stéphane Vannitsem<sup>1</sup>

<sup>1</sup>Royal Meteorological Institute of Belgium, Avenue Circulaire 3, 1180, Brussels, Belgium

<sup>2</sup>Université libre de Bruxelles, de la Plaine 155, 1050, Brussels, Belgium

**Correspondence:** Anupama K Xavier (xavier.anupamak@meteo.be)

**Abstract.** This study delves into the predictability of atmospheric blocking, zonal, and transition patterns utilizing a simplified coupled model. This model, implemented in Python, emulates midlatitude atmospheric dynamics with a two-layer quasi-geostrophic channel atmosphere on a beta-plane, encompassing simplified land effects. Initially, we comprehensively scrutinize the model's responses to environmental parameters like solar radiation, surface friction, and atmosphere-ground heat exchange.

5 Our findings confirm that the model faithfully replicates real-world Earth-like flow regimes, establishing a robust foundation for further analysis. Subsequently, employing Gaussian mixture clustering, we successfully delineate distinct blocking, zonal, and transition flow regimes, unveiling their dependencies on surface friction. To gauge predictability and persistence, we compute the averaged local Lyapunov exponents for each regime. Our investigation uncovers the presence of zonal, blocking, and transition regimes, particularly under conditions of reduced surface friction. As surface friction increases further, the system

10 transitions to a state characterized by two blocking regimes and a transition regime. Intriguingly, periodic behavior emerges under specific surface friction values, returning to patterns observed under low friction coefficients. Model resolution increase impacts the system in a way that only two regimes are then obtained with the clustering: the transition phase disappears and the predictability drops to roughly 2 days for both of the remaining regimes. In accordance with previous research findings, our study underscores that when all three regimes coexist, zonal patterns exhibit a more extended predictability horizon compared

15 to blocking patterns. Remarkably, transition patterns exhibit reduced predictability when coexisting with the other regimes. In addition, within a specified range of surface friction values where two blocking regimes are found, it is observed that blocked atmospheric situations in the west of the applied topography are marked by instabilities and reduced predictability in contrast to the blockings appearing on the eastern side of the topography.

## 1 Introduction and motivation

20 Low-Frequency Variability (LFV) encompasses a wide range of atmospheric and climate processes, including atmospheric blockings, heat waves, cold spells, and long-term oscillations like the Madden-Julian Oscillation (MJO), the North Atlantic Oscillation (NAO), and the El-Niño - Southern Oscillation (ENSO). Despite extensive research, a comprehensive understanding of the nature of these LFVs remains elusive. In practical terms, exploiting these LFVs to achieving accurate extended-range forecasts beyond two weeks at midlatitudes remains a formidable challenge. On the climate front, comprehending how climate change affects the low-frequency variability of the atmosphere also remains an area of incomplete knowledge. Previous works, as reported in Ghil and Robertson (2002), Lucarini and Gritsun (2020), have highlighted this gap in understanding.

Blocking systems - a notable form of LFV observed in the atmosphere - can be described as long-lasting, quasi-stationary flow patterns in the troposphere (Liu, 1994). These patterns are characterized by a significant meridional flow component, leading to a disruption or deceleration of the zonal westerly flow at midlatitudes (Nakamura and Huang, 2018). While the blocking systems persist, strong zonal flows may simultaneously exist to the north and south of them. The evolution of blocking systems involves transitions between more zonal and more meridional flow patterns during their onset and decay phases, posing challenges for forecast models (Frederiksen et al., 2004). Moreover, the dynamics of blocking systems are complex, involving interactions across different spatial and temporal scales, both internally within the system and with the surrounding flow environment (Shutts, 1983; Lupo and Smith, 1995). Researchers have highlighted the intricate nature of these dynamics and the connections between various scales, contributing to the challenges in understanding and predicting the behavior of blocking systems. As blocking systems have the potential to induce weather extreme like heatwaves, there is a notable interest in understanding how the characteristics of these blocking events might evolve in the future and how such changes could subsequently impact the occurrence and features of surface extreme weather events. The investigation of these potential changes is of significant importance to assess the risks associated with extreme weather events and to enhance our understanding of the complex interactions between blocking patterns and surface weather conditions in a changing climate context (Kautz et al., 2022).

Eventhough the concerns above matters, identifying and evaluating LFVs in GCMs is computationally expensive, so in this study an idealised reduced order coupled model is used. It is a climate model 'stripped to the bone', which links theoretical understanding to the complexity of more realistic models, made by key ingredients and approximations; which hence helps us to study a particular phenomenon by tweaking the parameters affecting them with less computational cost. ~~These types-~~The pursuit of simplified models age back from Lorenz (1963a) where he demonstrated the presence of the property of sensitivity to initial conditions in a simple system, that subsequently led to the development of chaos theory for atmospheric phenomena has a long history, dating back to Lorenz's seminal work in the early 1960s (Lorenz, 1960, 1962, 1963a, b). This approach recognizes the value of sacrificing some detail in exchange for a deeper grasp of fundamental physical processes.

50 ~~Later on the~~Lorenz demonstrated the power of this strategy by leveraging Fourier series to distill the barotropic vorticity equation into three ordinary differential equations (Lorenz, 1960). These equations, while omitting smaller scales of motion, yielded valuable insights into atmospheric scenarios such as flow interactions and current stability. Subsequently, he developed

55 a simplified geostrophic model using truncated Fourier-Bessel series (Lorenz, 1962). This eight-equation model captured baroclinic instability, a critical process in atmospheric dynamics, while maintaining key energy relationships. Notably, the model successfully reproduced observed flow regimes and transitions in rotating fluids, suggesting its effectiveness in studying large-scale atmospheric behavior. Lorenz's 1963 research yielded significant advancements in our understanding of atmospheric dynamics through two key publications (Lorenz, 1963a). The first introduced a now-iconic system of three differential equations, derived from a further simplified model for fluid flow. This groundbreaking work unveiled the concept of sensitive dependence on initial conditions, a cornerstone of chaos theory.

60 In the same year, Lorenz explored a separate avenue by investigating a simplified model for symmetrically heated rotating viscous fluids (Lorenz, 1963b). This work resulted in a system of fourteen ordinary differential equations governed by two external parameters: the thermal Rossby number and the Taylor number. Analytical solutions revealed the existence of purely zonal flow and superimposed steady waves, while numerical integration unveiled a richer tapestry of flow behaviors. Oscillatory solutions with periodic shape changes and irregular non-periodic flow emerged. Interestingly, increasing the Taylor number generally led to greater flow complexity, except at very high values where the model's truncations became unrealistic.

65 Perhaps most intriguing was the coexistence of unstable purely zonal, steady-wave, and oscillatory solutions. This suggests intricate flow dynamics, with transitions between symmetric and unsymmetric vacillation occurring independently of instability. These findings highlight the ability of simplified models to unveil complex and nuanced behaviors in atmospheric dynamics (Shen et al., 2023). Lorenz's pioneering work in the early 1960s demonstrated the power of simplified models for understanding atmospheric dynamics. By strategically neglecting certain complexities, he was able to capture key phenomena like baroclinic instability and chaos. However, for large-scale atmospheric simulations, computational efficiency becomes paramount. This is where Qg (quasi-geostrophic) models come in. Qg models prioritize large-scale features by making specific approximations, allowing for rapid simulations and analyses of broad atmospheric circulation patterns. While they may not capture the intricate details explored by Lorenz's models, Qg models remain a workhorse for studying large-scale atmospheric phenomena. Hence  
70 the trend continued with Charney and DeVore (1979) in which a quasi-geostrophic model, projected onto Fourier modes for a more efficient and concise representation, has been developed. It also incorporates an idealized parameterization closure to account for subgrid-scale processes. By imposing in addition a meridional temperature gradient over a topography, the model becomes a forced-dissipative system, which exhibits multiple stable equilibria, representing distinct atmospheric flow patterns. Charney and Devore hypothesized that the transitions between these solutions were primarily influenced by small-scale  
80 perturbations or the presence of baroclinic instability within the system.

Charney and Straus (1980) partially confirmed this hypothesis and found out that these transitions were indeed the result of baroclinic instability. Their study sheds light on the complex interactions between atmospheric flow, orography and propagating planetary waves in baroclinic systems. They discovered that the interactions between atmospheric flow and orography induces form-drag instability, generating eddies and perturbations, and leading to multiple stable equilibria with distinct flow patterns  
85 under consistent forcing conditions.

Charney's seminal study sparked significant interest in the low-order spectral model and the theory of multiple flow equilibria. Zhengxin and Baozhen (1982) and Zhu (1985) employed a two-layer low-order spectral model, discovering stable

equilibrium states resembling actual blocking, with zonally asymmetric thermal and topographic forcings and flow nonlinearity playing critical roles in blocking dynamics. The summarized version of the evolution of numerical weather prediction and predictability tools are included in Yoden (1983a, b, 2007).

Reinhold and Pierrehumbert (1982, 1985) extended Charney's model, incorporating additional synoptic-scale waves, revealing two distinct weather regime states influenced by wave-wave interactions causing transitions between equilibrium states.

Cehelsky and Tung (1987) demonstrated that the behavior of a reduced-order model exhibits notable disparities at higher resolutions, primarily attributed to the inadequate representation of energy upscaling and vorticity downscaling pathways. They coined this phenomenon as '*spurious chaos*', denoting the emergence of irregular dynamics that are not genuinely representative of the underlying physical processes. Although this is a valid point, high resolution models are usually hard to analyze in detail. There is therefore a need for investigating first reduced coupled model in order to get qualitative conclusions on a problem at hand. We started this journey by using a reduced order land atmospheric model to investigate the impact of coupling between the land and the atmosphere on LFV.

Legras and Ghil (1985) employed a higher-order barotropic spectral spherical model to investigate blocking and zonal flow regimes dynamics, suggesting that their model displayed properties akin to an index cycle, and later stochastic forcing was introduced to Charney's deterministic model, leading to transitions between high- and low-index states (Benzi et al., 1984; Egger, 1981; Sura, 2002). The impact of stochastic forcing on the stability of atmospheric regimes was also recently considered in a highly-truncated barotropic model by Dorrington and Palmer (2023), where they provide a mechanism to explain the increased persistence of blocking due to the noise in such simple models.

[Schubert and Lucarini \(2016\) recent Legras and Ghil \(1985\) also discussed the realistic existence of blocked and zonal flow regimes which are obtained as unstable stationary solutions due to the barotropic influence of the LFVs in the atmosphere. More persistent zonal flows are also identified in several occasions which seems to be a deviation from the earlier studies. Later the stability studies by Weeks et al. \(1997\) recreating zonal and blocked regimes in an experimental annulus setup further substantiated the findings of Legras and Ghil \(1985\).](#)

[Schubert and Lucarini \(2016\)'s](#) numerical investigation employing a QG model revealed a counter-intuitive finding that during blocking events, the global growth rates of the fastest growing covariant Lyapunov vectors (CLVs) are significantly higher, indicating stronger instability compared to typical zonal conditions. The difficulty in predicting the specific timing of blocking onset and decay further contributes to the observed instability behavior, aligning with Kwasniok (2019) findings associating anomalously high values of finite time largest Lyapunov exponents with blocked atmospheric flows.

[Lucarini and Gritsun \(2020\) demonstrated that blocking phenomena exhibit higher instability compared to typical atmospheric conditions, irrespective of whether they occur in the Atlantic, Pacific, or globally. This analysis utilized the simplified atmospheric model proposed by Marshall and Molteni \(1993\) and assessed stability based on unstable periodic orbits \(UPOs\). Importantly, this research dispelled the misconception that the increased stability of zonal flows solely resulted from the barotropic nature of the model in the study of Legras and Ghil \(1985\) and Weeks et al. \(1997\) apparatus.](#) Consistent results were obtained by Faranda et al. (2016, 2017), utilizing extreme value theory for dynamical systems, which identified blocking regimes with unstable fixed points in a heavily reduced phase space. Their findings indicated that blockings exhibit higher instability in the cir-



125 culation, linked to an increased effective dimensionality of the system. This agreement with [Schubert and Lucarini \(2016\)](#),  
[Lucarini and Gritsun \(2020\)](#) study further supports the notion that blocking events display stronger turbulence and instability,  
challenging conventional expectations.

We here aim at extensively investigating the predictability of blocking, zonal, and transition regimes utilizing backward Lyapunov exponents (BLVs) in the context of a recently developed reduced-order land-atmosphere model, providing a more comprehensive understanding of the system's behavior and regime predictability.

130 The classic Charney's model lacks feedback from atmospheric flow to the artificially specified "thermal forcing," leading  
to potential unrealistic effects on large-scale atmospheric motions. To address this limitation, a new land atmospheric coupled  
model is proposed in Li et al. (2018), which incorporate an energy balance scheme to allow atmospheric motions to influ-  
ence the land temperature distribution and vice-versa. By considering horizontally inhomogeneous radiative input fields as the  
driving force for land-atmosphere dynamics, this coupled model offers a more realistic representation of the interactions be-  
tween the land and the atmosphere. The model bears resemblance to the low-order coupled ocean-atmosphere model proposed  
135 by Vannitsem et al. (2015), but with a heat bath featuring the land and an idealized topography.

Prior to conducting the investigation on the predictability of blocking, zonal, and transition regimes using backward Lyapunov exponents (BLVs), we performed a characterization of the sensitivity of the quasi-geostrophic land-atmosphere coupled model embedded in the `qgs` framework (Demaeyer et al., 2020) with respect to various environmental parameters that are essential for the functioning of the atmosphere.

140 The structure of the paper is as follows. Section 2 introduces the model, outlining its structure, main properties, and the  
parameters employed in the study. Additionally, this section includes a discussion on the stability properties of the system and  
temporal evolution of the modes (barotropic stream function, baroclinic streamfunction and ground temperature). In Section 3,  
the methodology used for the investigation is explained. Section 4 presents the stability and the Lyapunov properties of the  
model corresponding to various environmental factors, and in Section 5, predictability properties of different weather regimes  
145 are discussed. Effects of the model resolution are presented in section 6 and the conclusions drawn from the research are  
provided in section 7, along with future perspectives for further studies.

## 2 Land atmosphere coupled model

### 2.1 Model Characteristics

150 `qgs` is a Python framework in which several reduced-order climate models are implemented for midlatitudes (Demaeyer et al.,  
2020). It models the dynamics of a 2-layer quasi-geostrophic (QG) channel atmosphere on a beta-plane, coupled to a simple  
surface component that could be a land or an ocean. In the current study, we are using the quasi-geostrophic land-atmosphere  
coupled model version (Li et al., 2018).

The atmospheric part of the model is represented as a 2-layered, quasi-geostrophic flow defined on a  $\beta$  plane within the zonal  
walls  $y = 0$  and  $\pi L$  (Reinhold and Pierrehumbert, 1982). The thermodynamic equations of the baroclinic atmosphere includes  
155 the energy exchanges between land, atmosphere and space similar to the radiative and heat flux scheme provided in Barsugli

and Battisti (1998). The coupling of the atmospheric components with the ground is constituted by the surface friction and the radiative and heat exchanges between the atmosphere and the ground. As usual in such types of models, channel atmosphere is considered with no-flux boundary conditions on the north and south borders and periodic boundary conditions on the east and west border.

160 The equations governing the time evolution of barotropic and baroclinic streamfunction of the atmospheric part are as follows:

$$\begin{aligned} \frac{\partial}{\partial t} (\nabla^2 \psi_a) + J(\psi_a, \nabla^2 \psi_a) + J(\theta_a, \nabla^2 \theta_a) + \frac{1}{2} J(\psi_a - \theta_a, f_0 h / H_a) + \beta \frac{\partial \psi_a}{\partial x} \\ = -\frac{k_d}{2} \nabla^2 (\psi_a - \theta_a) \end{aligned} \quad (1)$$

$$\begin{aligned} 165 \frac{\partial}{\partial t} (\nabla^2 \theta_a) + J(\psi_a, \nabla^2 \theta_a) + J(\theta_a, \nabla^2 \psi_a) - \frac{1}{2} J(\psi_a - \theta_a, f_0 h / H_a) + \beta \frac{\partial \theta_a}{\partial x} \\ = -2k'_d \nabla^2 \theta_a + \frac{k_d}{2} \nabla^2 (\psi_a - \theta_a) + \frac{f_0}{\Delta p} \omega \end{aligned} \quad (2)$$

where  $\omega$  is the vertical velocity of the system.  $\psi_a$  is the barotropic streamfunction and  $\theta_a$  is the baroclinic streamfunction of the atmosphere. The constants  $k_d$  and  $k'_d$  multiply the surface friction term and the internal friction between layers, respectively. The temperature equation of the baroclinic atmosphere and ground are :

$$170 \gamma_a \left( \frac{\partial T_a}{\partial t} + J(\psi_a, T_a) - \sigma \omega \frac{p}{R} \right) = -\lambda(T_a - T_g) + \epsilon_a \sigma_B T_g^4 - 2\epsilon_a \sigma_B T_a^4 + R_a \quad (3)$$

$$\gamma_g \frac{\partial T_g}{\partial t} = -\lambda(T_g - T_a) - \sigma_B T_g^4 + \epsilon_a \sigma_B T_a^4 + R_g \quad (4)$$

where  $T_g$  and  $T_a$  are the ground and atmospheric temperature respectively.  $\sigma$  is the static stability with  $p$  as the pressure.  $R$  is the gas constant for dry air.  $\gamma_a$  is the heat capacity of the atmosphere for a 1000-hPa deep column where as  $\gamma_g$  is the heat capacity of the active layer of the land for a mean thickness of 10 m (Monin, 1986).  $\lambda$  is the heat transfer coefficient between the land and atmosphere.  $\sigma_B$  is the Stefan-Boltzmann constant and  $\epsilon_a$  is the longwave emissivity of the atmosphere.  $R_a$  is the shortwave solar radiation directly absorbed by the atmosphere whereas  $R_g$  is the shortwave solar radiation absorbed by the land.

180 The hydrostatic relation in pressure coordinates  $\frac{\partial \Phi}{\partial p} = \frac{-1}{\rho_a}$  where the geopotential height  $\Phi^i = f_0 \psi_a^i$  and the ideal gas relation  $p = \rho_a R T_a$  allow one to write the spatially dependent atmospheric temperature anomaly  $\delta T_a = \frac{2f_0 \theta_a}{R}$ , with  $\theta_a$  as the baroclinic streamfunction. This can be used to eliminate the vertical velocity  $\omega$ . This changes the independent dynamical field

185 to the streamfunction field  $\psi_a$  and the spatially dependent temperatures  $\delta T_a$  and  $\delta T_g$ . The dimensional meridional differential shortwave solar radiation absorbed by the land and the atmosphere are given by  $\delta R_g = \sqrt{2}C_g \cos \frac{y}{L}$  and  $\delta R_a = \sqrt{2}C_a \cos \frac{y}{L}$  respectively. Hence we decide to provide  $C_a = 0.4C_g$ . The variable  $C_g$  is a dimensional parameter, which is an indicator of the meridional difference in solar heating absorbed by the land between the walls, and it is the crucial parameter in our land atmosphere coupled model. As in Vannitsem et al. (2015) and De Cruz et al. (2016), quartic terms of the temperature equations are linearised. Upon nondimensionalization, the `qgs` framework represents the above equations as ordinary differential equations by projecting them on to a set of basis functions, a procedure which is also known as Galerkin expansion. We investigate the (2, 2) resolution configuration of the model for the current study, which means that basis functions up to wavenumber 2 in each coordinate of the model are used. Consequently, the following list of 10 basis functions was used for the study:

$$\begin{aligned}
F_1(x, y) &= \sqrt{2} \cos(y), \\
F_2(x, y) &= 2 \cos(nx) \sin(y), \\
F_3(x, y) &= 2 \sin(nx) \sin(y), \\
F_4(x, y) &= \sqrt{2} \cos(2y), \\
F_5(x, y) &= 2 \cos(nx) \sin(2y), \\
F_6(x, y) &= 2 \sin(nx) \sin(2y), \\
F_7(x, y) &= 2 \cos(2nx) \sin(y), \\
F_8(x, y) &= 2 \sin(2nx) \sin(y), \\
F_9(x, y) &= 2 \cos(2nx) \sin(2y), \\
F_{10}(x, y) &= 2 \sin(2nx) \sin(2y),
\end{aligned} \tag{5}$$

This configuration yields therefore a set of 30 variables, including 10 barotropic variables, 10 baroclinic variables, and 10 ground temperature variables. Note that as explained in the `qgs` documentation, the basis functions of the model can be easily altered.


## 195 2.2 Model Parameters

Even though the model that we are using is a highly truncated spectral model, its comparability towards the original atmosphere and ground coupling is of great importance. Simulations produced by the model will be more meaningful if it characterizes *earthlike* properties. This can be obtained by tweaking and tuning the model parameters. In the present scenario, the parameters used are derived from Reinhold and Pierrehumbert (1982), where they specifically estimated realistic parameter ranges that result in midlatitude terrestrial flow characteristics and regimes. The typical dimensional parameter values used for this study are displayed in Table 1.

**Table 1.** Typical values of the model used for the study

Parameter	value	Parameter	value
$a$	6371 km	$\sigma_B$	$5.67 \times 10^{-8} \text{ Wm}^{-2}\text{K}^{-4}$
$\pi L$	5000 km	$\gamma_a$	$1.0 \times 10^7 \text{ Jm}^{-2}\text{K}^{-1}$
$H$	8.5 km	$\gamma_g$	$1.6 \times 10^7 \text{ Jm}^{-2}\text{K}^{-1}$
$\phi_o$	$50^\circ \text{ N}$	$\lambda$	$10 \text{ Wm}^{-2}\text{K}^{-1}$
$n$	1.3	$f_o$	$0.0001032 \text{ s}^{-1}$
$\epsilon$	0.76	$R$	$287.058 \text{ Jkg}^{-1}\text{K}^{-1}$
$k_d$	$1.2384 \times 10^{-5} \text{ s}^{-1}$	$k'_d$	$2.068 \times 10^{-6} \text{ s}^{-1}$
$\sigma$	$2.158 \times 10^{-6} \text{ m}^2\text{s}^{-2}\text{Pa}^{-2}$		

Values of  $\lambda, n, k_d$  are varied beyond the displayed value to study the model sensitivity.




**Figure 1.** Temporal evolution of barotropic ( $\psi_{a,1}$ ) and baroclinic ( $\theta_{a,1}$ ) atmospheric streamfunction and the ground temperature ( $T_{g,1}$ ) for  $C_g = 300 \text{ Wm}^{-2}$  and  $k_d = 0.085$ .

### 2.3 Model trajectories and mean fields

Figure 1 displays the time evolution of the first barotropic ( $\psi_{a,1}$ ) and baroclinic ( $\theta_{a,1}$ ) streamfunction modes of the atmosphere and the ground temperature ( $T_{g,1}$ ) for about 10 years starting after 10000 days of transient integration. Fluctuations are more erratic in the atmospheric part, which denotes its key role in the dynamics of the system. The variable representing the land component of the system is comparatively slower and less erratic. This difference suggests that the land component has a longer typical time scale than the atmosphere in this system.

Figure 2(a) emphasizes the observation above, where the autocorrelation of the first barotropic atmospheric mode and the first ground temperature mode for  $C_g = 300 \text{ Wm}^{-2}$  and  $k_d = 0.085$  are displayed. These, evaluated on a time series of 10000 days, helps us estimate the memory loss of the variables.

Indeed, the typical timescale of the processes at hand can be evaluated by the  $e$ -folding time which is the time beyond which the correlation has decreased by  $1/e$ . As expected the  $e$ -folding time of the atmospheric part is approximately 1.9 days which is comparatively lower than that of the land part ( $\approx 7.6$  days) indicating that the system is a multiscale model with a typical timescale ratio of 10 for the specific parameter values considered.



**Figure 2.** (a) Autocorrelation of barotropic ( $\psi_{a,1}$ ) atmospheric streamfunction and the ground temperature ( $T_{g,1}$ ) for  $C_g = 300 \text{ W m}^{-2}$  and  $k_d = 0.085$ . (b) Powerspectrum of the same variables.

215 Figure 2(b) is depicting the power spectra of modes  $\psi_{a,1}$  and  $T_{g,1}$  calculated by the Fourier transformation of the auto-  
correlation function using again timeseries of 10000 days. Atmospheric mode has a flat spectrum for lower frequencies and  
decays rapidly for higher frequencies. The spectrum for the ground mode is initially following the path of the atmosphere  
(lower frequencies up to 0.001) but starts to decay earlier which indicates more structured variabilities at lower frequencies  
than the atmosphere. The existence of a substantial continuous part in the spectrum is an indication of the complexity of the  
220 deterministic dynamics in time and suggests the presence of a chaotic dynamics (Arbabi and Mezić, 2017; Mezić, 2020).

### 3 Methodology

The objective of this research - besides introducing the basic equations, energy balance scheme and its sensitivity - is to explore  
the predictability of blocking and zonal weather regimes. By analyzing the peculiarities and the patterns of the land atmospheric  
coupled model, the study concludes that, when utilizing the parameters described in section 2.2, the system shows a qualita-  
225 tively similar behaviour as the large scale actual atmosphere at midlatitudes (examples are illustrated in Appendix B). More-  
over, varying the surface friction term,  $k_d$ , within the range of 0.06 to 0.12 yields numerous instances of realistic flow regimes,  
including blocking, zonal, and transition phases between these two states. To isolate between these flow regimes, a machine  
learning algorithm called Gaussian Mixture Clustering (GMC) is employed, which will be described in [appendix-A](#)  
230 [Appendix A](#). After classifying the data, the average geopotential height at 500 hPa of each cluster is calculated to identify different flow  
regimes. ~~The number of clusters is fixed to three using a trial and error method.~~ Through experimentation encompassing 2 to  
6 clusters, discernible flow regimes emerged for 2 and 3 clusters, showcasing significant distinctions. However, as the cluster  
count reached 4, we noted convergence between two clusters, leading to identical structures and flow patterns. This trend  
persisted with additional cluster increments. Hence, we inferred that the attractor attained optimal clustering with evident and  
nearly uniform data point distribution when employing 3 clusters. Each flow regime is equally important due to its presence  
235 in the actual atmosphere. The cluster containing the lowest fraction of points in percentage being designated as the transition

regime. The predictability horizon of each regime is evaluated by computing the inverse of the average largest local Lyapunov exponents of the clusters, which are calculated at each point of the attractor before clustering. Although the attractor exhibits multiple positive Lyapunov exponents, our investigation focused solely on the first exponent, as it governs the dynamics of the error and is therefore considered the most influential.

## 240 4 Regime stability and Lyapunov properties of the low order land atmospheric coupled model

Characterizing the instability properties of different flow regimes and their dependence with respect to important parameters are investigated as follows: The first part is exploring the sensitivity of the model about essential parameters which play a key role in structuring the output of the system. In the second part, the predictability properties of zonal, blocking and transition flow regimes using the Lyapunov exponents is investigated.

### 245 4.1 Stability properties of the model

Stability properties of the land-atmosphere coupled model at their equilibrium states are well depicted in Li et al. (2018). They also defined high-index equilibria and low-index equilibria based on the value of the streamfunction in the upper and lower atmospheric layer when the model solutions are equilibrium states. Even though stability of the model's equilibrium states is interesting and insightful, the actual atmosphere displays time-dependent solutions. Moreover, realistic atmospheric models are chaotic and acutely sensitive to the initial conditions. Hence in this section, we investigated the stability properties of the land atmosphere model when its behavior is similar to *earthlike* situations with erratic dynamics.

Chaotic dynamical systems which exhibit sensitivity to the initial conditions can be qualitatively analysed by computing Lyapunov exponents and vectors.

### 4.2 Theory

255 Sensitivity to initial conditions is usually estimated using Lyapunov exponents. Let us consider an initial state,  $\mathbf{x}(t_o) = \mathbf{x}_o$ , a small perturbation  $\delta\mathbf{x}_o$  is added to it which produces eventually a completely different trajectory. The dynamics of the error growth of the system can be linearized provided the perturbation is infinitesimally small as

$$\frac{d\delta\mathbf{x}}{dt} = \left. \frac{\partial \mathbf{f}}{\partial \mathbf{x}} \right|_{\mathbf{x}(t)} \delta\mathbf{x} \quad (6)$$

and its solution is

$$260 \quad \delta\mathbf{x}(t) = \mathbf{M}(t, \mathbf{x}(t_o))\delta\mathbf{x}(t_o) \quad (7)$$

where  $\mathbf{M}$  is known as the resolvent or propagator matrix. The Euclidean norm of the error can be computed as

$$E_t = |\delta \mathbf{x}(t)|^2 = \delta \mathbf{x}(t)^T \delta \mathbf{x}(t)$$

$$E_t = \delta \mathbf{x}(t_o)^T \mathbf{M}(t, \mathbf{x}(t_o)) \mathbf{M}(t, \mathbf{x}(t_o))^T \delta \mathbf{x}(t_o) \quad (8)$$

hence indicating that the error growth is provided by the eigenvalues of  $\mathbf{M}^T \mathbf{M}$ , where  $\mathbf{M}^T$  denotes the transpose of the  
 265 resolvent matrix  $\mathbf{M}$ . By the multiplicative ergodic theorem of Oseledec (Eckmann and Ruelle, 1985; Kuptsov and Parlitz, 2012), a double limit is considered with perturbation amplitude going to 0 and time going to infinity. The logarithm of the eigenvalues of matrix  $(\mathbf{M}^T \mathbf{M})^{2(t-t_o)}$  within these limits, which are known as Lyapunov exponents, quantifies the divergence of the initially close trajectories. The complete set of Lyapunov exponents that are usually represented in decreasing order constitutes the Lyapunov spectrum. Different types of Lyapunov vectors existed based on the method of calculation like forward  
 270 Lyapunov vectors, backward Lyapunov vectors and covariant Lyapunov vectors whose properties are described in details in (Kuptsov and Parlitz, 2012; Legras and Vautard, 1996).


In the current study, we are using backward Lyapunov vectors (BLVs) which are obtained by considering the eigenvalues of the matrix  $(\mathbf{M}^T \mathbf{M})^{2(t-t_o)}$  by taking initial state  $t_o \rightarrow -\infty$ . Several numerical techniques exist for the calculation of the Lyapunov exponents. The most common method is using Gram-Schmidt orthonormalization (Shimada and Nagashima, 1979;  
 275 Parker and Chua, 2012): A set of orthonormal random vectors are propagated in the tangent space of the trajectory, according to equation (6), frequently re-orthonormalizing this basis to avoid the collapse of all the vectors towards the most unstable direction which is associated with the largest Lyapunov exponent. After a transient, these vectors provide the BLVs and concurrently the sought Lyapunoc exponents. The complete set of these vectors give the full picture of the instability of the trajectory in phase space.

### 280 4.3 Lyapunov spectra and averaged variance of the model


Figure 3 displays the Lyapunov spectrum of the land-atmosphere coupled model when  $C_g = 300 \text{ W m}^{-2}$  and  $k_d = 0.085$ . All other parameters are provided in Table 1. The system has 3 positive, 1 zero and 26 negative Lyapunov exponents. Similarly to the ocean-atmospheric coupled model (Vannitsem et al., 2015; Vannitsem and Lucarini, 2016; Vannitsem, 2017), the spectrum contains a set of Lyapunov exponents forming a plateau close to 0, but the amplitude of the Lyapunov exponents around  
 285 this plateau is however quite substantial as compared to the coupled ocean-atmosphere model. This plateau is expected to be associated with the presence of the land whose typical time scales of variability are slower than the atmosphere.

The coupling between land and atmosphere plays a key role in the behaviour of the model. Hence quantifying the extent of this coupling and its instabilities is essential for interpreting the properties of the model. Therefore, the averaged variance of the Lyapunov vectors is displayed in figure 4 to elucidate this information along each variable.

290 The first ten variables represent the barotropic streamfunction of the atmosphere, the next ten represent the baroclinic streamfunction of the atmosphere, and the last ten represent the ground temperature. The variance of the BLVs is primarily residing in the atmospheric part, indicating that this part primarily contributes to the system's dynamics. It should also be noted that



**Figure 3.** Lyapunov spectra of the land atmospheric coupled model for  $C_g = 300 \text{ Wm}^{-2}$  and  $k_d = 0.085$ . The values of the Lyapunov exponents are given in  $\text{days}^{-1}$ .




**Figure 4.** Values of the time-averaged and normalized variance of the BLVs as a function of the variables of the model ( $\log_{10}$  scale). The 20 first modes correspond to the variables of the atmosphere and the next 10 ones correspond to the temperature of the ground. Parameters' value:  $C_g = 300 \text{ Wm}^{-2}$  and  $k_d = 0.085$ . The Euclidean norm is used for all BLVs, and their squared norm is normalized to 1


the atmospheric part includes the most unstable BLVs (corresponding to the first two Lyapunov exponents) as well as the most stable BLVs (21 to 30 corresponds to large negative LEs).

295 Conversely, variance is predominantly projected on the temperature variables of the ground part for the BLVs 3 to 20. These BLVs are associated with the plateau formed by the near-zero LEs visible in Fig. 3. The observation of comparable variance in both the atmospheric and ground parts describes (horizontally) the coupling in the model, which is thus represented for BLVs





**Figure 5.** Lyapunov spectra of the land atmospheric coupled model for different values of  $C_g$ . Each color represents corresponding  $C_g$  values used for calculating the spectrum. The values of the Lyapunov exponents are given in days<sup>-1</sup>



**Figure 6.** First and second Lyapunov exponents of the land atmospheric coupled model for different values of  $C_g$  for a fixed value of  $k_d = 0.08$ . blue line represents first and orange line represents second Lyapunov exponents respectively. The values of the Lyapunov exponents are given in days<sup>-1</sup>


3-7. For BLVs 21 to 30, variance projection is almost non-existent in the ground part, indicating that the ground part makes almost no contribution to the stabilization of the system.

300 Figure 5 displays Lyapunov spectra calculated for various energy input levels ( $C_g$ ) ranging from 300 to 400  $Wm^{-2}$ .

When  $C_g$  is lower, the model exhibits chaotic behavior, indicated by two positive exponents, one zero exponent, and 26 negative exponents. Surprisingly, the amount of incoming shortwave radiation doesn't significantly affect the spectrum between  $C_g$  values of 300 and 360  $Wm^{-2}$ .

305 However, as  $C_g$  increases, the system shifts to periodic behavior, characterized by one zero exponent and 29 Lyapunov exponents, before switching back to chaos.

This is illustrated in Figure 6 where the first and second Lyapunov exponents are positive for the system for the lower values of  $C_g$  and then system enters in to a periodic window for the  $C_g$  values and revert back again to chaotic behaviour later.



**Figure 7.** Lyapunov spectra of the land atmosphere coupled model for different values of  $k_d$ . Each color represents corresponding  $k_d$  values used for calculating the spectrum. The values of the Lyapunov exponents are given in days<sup>-1</sup>.

The surface friction  $k_d$  is also affecting stability of the system to a great extent. In order to investigate the sensitivity of the model towards  $k_d$ , Lyapunov spectra were drawn with the parameter values exhibited in table 1 with different  $k_d$  values as displayed in Fig. 7.

The non-dimensional  $k_d$  values depicted in the figure were obtained from Reinhold and Pierrehumbert (1982), where they asserted that flow regimes generated using these values exhibit realistic midlatitude terrestrial properties. Among the system configurations with  $k_d$  values of 0.06, 0.075, 0.09, and 0.12, there are 3 positive, 1 zero, and 26 negative Lyapunov exponents. Up to the point of plateau formation, all these spectra display similar behavior. Note that the spectrum corresponding to  $k_d = 0.12$  exhibits notably strong negative Lyapunov exponents, while the spectrum for  $k_d = 0.06$  demonstrates comparatively weaker negative values, as expected from the increased associated dissipation.


From Figure 4, we identified that BLVs 20 - 30 are actually depicting the variance of temperature of the atmosphere. Given the significant variation in BLVs 20 - 30 within the current context, it can be inferred that the atmospheric temperature gradient, and hence baroclinic instability is becoming weaker and the system is stabilizing due to the changes in the surface friction.

As thoroughly explained in section 4.2, the largest Lyapunov exponent serves as an indicator of the system's highest degree of instability, while the second positive Lyapunov exponent represents the second most unstable characteristic, and so forth. In Fig. 8, the results demonstrate that at lower values of  $k_d$ , the system exhibits a highly chaotic nature. However, as we increase  $k_d$  towards higher values, the system stabilizes, revealing a periodic window. This can also be seen with the Lyapunov spectrum for  $k_d = 0.105$  in Fig. 7.


Subsequently, with further increment in  $k_d$ , the system returns to a more chaotic behavior, illustrating the complex role of dissipative features.

The primary interaction between land and atmosphere in the model is facilitated through heat exchange denoted by  $\lambda$ . Therefore, it is crucial to understand and analyze how alterations in the heat exchange mechanism influence the behavior of the model. This is illustrated in figure 9.

When  $\lambda$  is set to zero, the system displays a periodic behavior characterized by one zero and 29 negative Lyapunov exponents. However, for all other non-zero  $\lambda$  values, the system exhibits chaos with 3 positive exponents, 1 zero, and 26 negative Lyapunov exponents. Smaller values of  $\lambda$  (e.g.,  $\lambda = 10, 25$ , and 50) yield a smooth spectrum with a plateau which is similar to the typical




**Figure 8.** First and second Lyapunov exponents of the land-atmosphere coupled model for different non-dimensional values of  $k_d$ . Blue line represents the first and orange line represents the second Lyapunov exponents, respectively.  $C_g = 300 \text{ Wm}^{-2}$ . The values of the Lyapunov exponents are given in  $\text{days}^{-1}$



**Figure 9.** Lyapunov spectra of the land atmospheric coupled model for different values of  $\lambda$ . Each color represents corresponding  $\lambda$  values used for calculating the spectrum. The values of the Lyapunov exponents are given in  $\text{days}^{-1}$ .

Lyapunov spectrum encountered before. Note that with higher  $\lambda$  values, an anomalous bend is observed in the spectrum, specifically from the 20th exponent onward, indicating unrealistic stability. It is also interesting to note that the spectrum associated with the intersection of the land part (between 20th and 21st exponent) is becoming steeper with the increase of  $\lambda$ .  
 335 The increased heat exchange leads to a reduced temperature difference between the atmosphere and the ground, thereby giving rise to this particular situation. This can be further explained by the averaged variance of Lyapunov exponents for different values of heat exchange  $\lambda$  in Figure 10.

A clear separation between land and atmosphere exists at  $\lambda = 0$ , when there is no exchange of heat between them, resulted in  
 340 to respective spectrum. For  $\lambda = 10, 25$ , and  $50$ , the distribution remains similar, with variances concentrated predominantly in the atmospheric component for the first three BLVs and the last ten BLVs. BLVs 3-20 represent a plateau-like pattern, indicating a coupling between the atmosphere and the ground. Despite variance distribution, it's notable that there is a uniform spread



**Figure 10.** Values of the time-averaged and normalized variance of the BLVs as a function of the variables of the model ( $\log_{10}$  scale) for different  $\lambda$  values in  $\text{Wm}^{-2}\text{K}^{-1}$ . The 20 first modes corresponds to the variables of the atmosphere and the next 10 ones corresponds to the temperature of the ground. Parameters' value:  $C_g = 300 \text{ Wm}^{-2}$  and  $k_d = 0.085$ . The Euclidean norm is used for all BLVs, and their squared norm is normalized to 1.

rather than strong concentration or absence of variance. For  $\lambda = 75$  and 100, the variance distribution becomes compartmentalized. Variance is now concentrated within the first 20 BLVs for the barotropic streamfunction and ground temperature, while  
 345 the last 10 BLVs primarily represent atmospheric temperature or baroclinic streamfunction. In contrast to earlier cases, there is a clear absence of variance within the middle portion, specifically confined to BLVs 20-30. This compartmentalized energy distribution leads to a characteristic jump in the corresponding spectrum.

## 5 Predictability properties of zonal, blocking and transition flow regimes

Upon concluding our investigation of the land-atmosphere coupled model's utility in studying low-frequency variability in  
 350 the atmosphere, we have identified zonal, blocking, and transition regimes concerning different  $k_d$  values, as explained in section 3. For the range of  $k_d$  values between 0.06 and 0.12, the lower values (0.06 to 0.075) encompass all three regimes: zonal, blocking, and transition. As we progress from 0.08 onward, we observe two blocking regimes and a transitional regime between them until  $k_d = 0.10$ . Subsequently, the system exhibits periodic behavior. At  $k_d = 0.115$ , two blocking regimes are observed, and further, at  $k_d = 0.12$ , three flow regimes are identified. These findings demonstrate the model's capability to  
 355 capture various flow regimes and their transitions based on the selected range of  $k_d$  values.

The predictability horizon of the zonal regimes is found to be longer compared to the blocking regimes in the range of  $k_d$  values where all three regimes coexist ( $k_d = 0.06$  to 0.075). This implies that the blocking regimes exhibit higher instability in agreement with previous findings (Schubert and Lucarini, 2016; Faranda et al., 2016, 2017; Lucarini et al., 2016). The

transition regimes, on the other hand, show notably lower predictability in comparison to the other regimes. Within the interval  
360 of  $k_d$  values encompassing only two blocking regimes and a transition regime ( $k_d = 0.08$  to  $0.10$ ), both blocking regimes  
display significantly different predictability horizons. At the outset ( $k_d = 0.08$ ), they demonstrate a predictability difference  
of approximately 1 day. Subsequently, this difference increases and reaches about 10 days at  $k_d = 0.095$ . However, it then  
decreases again to a difference of 1 day when  $k_d = 0.10$ . Moreover, it is observed that the stability of the transition regime  
surpasses that of one of the blocking regimes. This indicates the occurrence of a qualitative change in the predictability of the  
365 blocking regimes within this particular range of  $k_d$  values.

In these instances, it is noteworthy that situations characterized by lower predictability or significant instability tend to occur  
when atmospheric blocking takes place on the western side of topographical features. Conversely, when blocking occurs on the  
eastern side of such topography, it exhibits greater stability and a longer predictability horizon. This observation draws parallels  
with real-world scenarios, such as the persistence of North Pacific blocking patterns (Breedon et al., 2020; Kim and Kim, 2019).  
370 The morphology of the identified blocking events bears resemblance to North Pacific blocks, where a high-pressure system is  
situated either to the west or east of the underlying topography. These locations correspond to the windward and leeward sides  
of the mountain ranges in the model.


The system is entering a periodic window after  $k_d = 0.10$ . Then it again become chaotic with 2 blocking regimes and a  
transition regime and later on with 3 distinct regimes. Figure 11 depicts all the findings obtained from this study.

## 375 **6 Impact of model resolution**


As per Cehelsky and Tung (1987), resolution of the model can affect the output in various ways. For instance, the nonlinear  
interaction between the modes (as number of modes increases, resolution of the system increases) is quite altered if different  
number of modes are considered, resulting in entirely different dynamics for the same set of parameters. Hence, analysis of high  
resolution runs and its Lyapunov properties is inevitable. In this section, the system is ran with a higher resolution configuration  
380 (5,5) with 55 modes being used for both the atmospheric and land part, giving a system with a total of 165 variables. Parameter  
values are the same as that of the earlier analysis listed in Table 1. Figure 12 is depicting the Lyapunov spectrum and Figure 13  
is representing the time-averaged variance of the BLVs of the high resolution run.

The Lyapunov spectrum exhibits structural similarity when compared to the low-resolution spectra, except that more positive  
exponents are present. Specifically, it comprises 12 positive, one zero, and 152 negative exponents. Notably, there is a distinctive  
385 plateau observable within the range of 20 to 75, which is a characteristic feature attributed to the interaction between the land  
and the atmosphere in the model. This plateau phenomenon arises due to the disparity in timescales resulting from the intricate  
interplay between land and atmosphere dynamics.

In Fig. 13, higher variances in the atmosphere are observed on first 40 BLVs and also at the BLVs greater than 70 indicating  
that the most chaotic and stable dynamics are resulting from the atmospheric component. Furthermore, it is noteworthy that a  
390 greater concentration of variance is observed to be shifted towards the variables that specifically represent the ground temper-  
ature, particularly within the range of BLVs 20 to 75. This shift is a key factor contributing to the formation of a plateau within




**Figure 11.** Predictability horizon of zonal, blocking and transition flow regimes for different non-dimensional  $k_d$ . The numbers represents the percentage of the total number of points included in the respective regime. Predictability horizon is depicted in days



**Figure 12.** Lyapunov spectrum of the land-atmosphere coupled model for  $C_g = 300 \text{ Wm}^{-2}$  and  $k_d = 0.085$  in (5,5) model configuration. The values of the Lyapunov exponents are given in days<sup>-1</sup>.

the spectrum in question. It becomes evident that there is a distinct compartmentalization between each set of variables, includ-



**Figure 13.** Values of the time-averaged and normalized variance of the BLVs as a function of the variables of the model ( $\log_{10}$  scale) in (5,5) model configuration. The 110 first modes corresponds to the variables of the atmosphere and the next 55 ones corresponds to the temperature of the ground. Parameters' value:  $C_g = 300 \text{ Wm}^{-2}$  and  $k_d = 0.085$ . The Euclidean norm is used for all BLVs, and their squared norm is normalized to 1.

ing the barotropic, baroclinic, and ground temperature variables, which is more pronounced when compared to the variance illustration at lower resolutions.

395 The outcome of the clustering analysis has delineated two clearly defined patterns: one marked by zonal flow and the other by instances of blocking. The intermediate pattern that once existed between these two regimes is now absent. Notably, both the zonal and blocking events exhibit an identical predictability horizon, specifically spanning a period of two days. This feature is contrasting with what is found at the lower resolution and also in the current literature on this topic. This aspect is worth investigating further in the future by exploring other sets of parameters and other resolutions.

## 400 7 Discussions

This study focused on characterizing the variability and instability properties of different flow regimes and their dependence on important parameters in an idealized coupled model, namely the quasi-geostrophic land atmosphere coupled model. The investigation also aimed at exploring the predictability of zonal, blocking, and transition flow regimes using Lyapunov exponents.

405 The analysis revealed that the model is less sensitive to variations in meridional differences in solar heating absorbed by  
the land, represented by  $C_g$ . Based on observations, a fixed value of  $300 \text{ Wm}^{-2}$  for  $C_g$  was chosen for further analysis. The  
study found that the model's stability is significantly affected by surface friction  $k_d$ . Different values of  $k_d$  were explored,  
and it was observed that at lower  $k_d$  values, the system exhibits chaotic behavior, while at higher  $k_d$  values, periodic windows  
alternate with chaotic behaviors. Within this range, the systems solutions meander between various flow regimes, including  
410 zonal, blocking, and transition.

The heat exchange mechanism, represented by  $\lambda$ , was also analyzed, and it was found that when  $\lambda$  is set to zero, the  
system displays periodic behavior, while for non-zero  $\lambda$  values, the system exhibits considerable chaos. Overall, the model  
demonstrated the capability to capture various flow regimes and their transitions based on the selected range of  $k_d$  values,  
providing insights into the potential behavior of the atmosphere.

415 The predictability properties of three distinct flow regimes were investigated: zonal, blocking, and transition, which were  
the fundamental components of low-frequency variability (LFV) in the model. The predictability horizon of the zonal regimes  
was found to be longer compared to the blocking regimes which is consistent with earlier results (Schubert and Lucarini, 2016;  
Faranda et al., 2016, 2017; Lucarini et al., 2016), when all three regimes coexist. The transition regimes showed notably lower  
predictability compared to the other regimes. Within a specific range of  $k_d$  values, the blocking regimes displayed different  
420 predictability horizons, indicating the occurrence of a qualitative change in the predictability of the blocking regimes.

Weather patterns that involve atmospheric blocking to the west of a given topographical feature tend to have reduced pre-  
dictability and show instability when contrasted with blocking occurrences situated to the east of such topographical elements.  
This finding aligns with actual meteorological occurrences, such as the persistence of North Pacific blocking patterns (Breedon  
et al., 2020; Kim and Kim, 2019). The shape and characteristics of the identified blocking events closely resemble North Pacific  
425 blocks, where a high-pressure system exists either on the western or eastern side of the underlying topography. In the physical  
world, these positions correspond to the windward and leeward sides of mountain ranges composed of rocky terrain. Despite  
qgs models being regarded as less comprehensive, their utilization in this study allows for a more relevant impact, akin to  
real-world situations as described above.

Upon increasing the model resolution, Lyapunov properties exhibited a remarkable resemblance to those observed at lower  
430 resolutions. However, the distribution of points on the attractor gave rise to two distinct clusters, delineating the blocking and  
zonal regimes, thereby extinguishing the potential for the transition regime. Notably, both the blocking and zonal regimes  
displayed a predictability horizon limited to 2 days.

In the backdrop of rising global temperatures and the escalation of climate extremes, comprehending the intricate dynam-  
ics governing atmospheric blocking occurrences and their predictability becomes paramount, given that blocking events are  
435 consistently linked to extreme weather phenomena. The impact of climate change can also be explored in the current study by  
modifying the emissivity of the atmosphere. This will be explored in the future.

The knowledge acquired through this study holds potential significance for climate and weather prediction models, con-  
tributing to the advancement of our understanding of the crucial atmospheric dynamics shaping the Earth's climate system. In  
future investigations, we will assess the influence of the same parameters within more complex models, enabling us to conduct



440 comparisons that will aid in identifying alterations in land-atmosphere interactions as atmospheric complexity intensifies. This undertaking will further our comprehension of how the interaction between land and the atmosphere evolves with increasing intricacies in atmospheric systems and how it affect the predictability of the weather regimes.

*Code availability.* The code used to obtain the results is a new version (v0.2.7) of `qgs` (Demayer et al., 2020) that was recently released on GitHub.

## 445 **Appendix A: Gaussian Mixture Clustering (GMC)**

Gaussian Mixture Clustering (GMC) is a prevalent unsupervised machine learning technique utilized for partitioning data points into clusters by modeling their underlying distribution. ~~The method assumes that the data arises from a combination of multiple Gaussian distributions~~ While the distribution of data points on the attractor may not align with a Gaussian distribution, the algorithm proceeds by approximating the distribution as Gaussian and subsequently identifies clusters based on this approximation.

450 approximation. Each cluster is characterized by a Gaussian component, and the primary objective is to accurately estimate the parameters of these components to optimally describe the data. It comprises several steps.

- **Initialization:** GMC starts by randomly selecting  $K$  initial centers (means)  $\mu_i$  for each cluster. Additionally, it initializes the covariance matrices  $\Sigma_i$  and the mixing coefficients  $\pi_i$ , which represent the probabilities of data points belonging to each cluster, where  $i = 1, 2, \dots, K$ . The mixing coefficients must sum up to 1, and each must be between 0 and 1.

455 – **Expectation - Maximization (EM) Algorithm:** GMC employs the EM algorithm (Dempster et al., 1977) to iteratively estimate the parameters of the Gaussian components. The algorithm comprises two steps, the Expectation step (E-step) and the Maximization step (M-step).

- **Expectation Step (E-Step):** During this step, the algorithm computes the responsibility ( $\gamma_{i,j}$ ) of each data point  $x_j$  for each cluster  $i$ . The responsibility represents the probability that data point  $x_j$  belongs to cluster  $i$ , given the current parameters. This is calculated using Bayes' theorem:

460

$$\gamma_{i,j} = \frac{\pi_i \cdot \mathcal{N}(x_j; \mu_i, \Sigma_i)}{\sum_{k=1}^K \pi_k \cdot \mathcal{N}(x_j; \mu_k, \Sigma_k)} \quad (\text{A1})$$

where  $\gamma_{i,j}$  is the responsibility of cluster  $i$  for data point  $x_j$ ,  $\pi_i$  is the mixing coefficient for cluster  $i$ .  $\mu_i$  and  $\Sigma_i$  are the mean and covariance matrix for cluster  $i$ , respectively.  $\mathcal{N}(x_j; \mu_i, \Sigma_i)$  is the Gaussian probability density function for data point  $x_j$  with mean  $\mu_i$  and covariance  $\Sigma_i$ .

465

- **Maximization Step (M-Step):** In this step, the algorithm updates the parameters of the Gaussian distributions (mean, covariance, and mixing coefficients) based on the responsibilities calculated in the E-step:

$$\text{New mean } , \mu_i = \frac{\sum_{j=1}^N \gamma_{i,j} \cdot x_j}{\sum_{j=1}^N \gamma_{i,j}} \quad (\text{A2})$$

$$470 \quad \text{New covariance matrix } , \Sigma_i = \frac{\sum_{j=1}^N \gamma_{i,j} \cdot (x_j - \mu_i) \cdot (x_j - \mu_i)^T}{\sum_{j=1}^N \gamma_{i,j}} \quad (\text{A3})$$

$$\text{New mixing coefficient } , \pi_i = \frac{1}{N} \sum_{j=1}^N \gamma_{i,j} \quad (\text{A4})$$


475 where  $N$  is the number of data points.  $\mu_i$ ,  $\Sigma_i$ , and  $\pi_i$  are the updated mean, covariance matrix, and mixing coefficient for cluster  $i$ , respectively.  $\gamma_{i,j}$  is the responsibility of cluster  $i$  for data point  $x_j$  (computed in the E-step).  $x_j$  is the  $j$ -th data point.

- **Convergence:** The E-step and M-step are repeated iteratively until the algorithm converges. Convergence happens when the change in the likelihood of the data between iterations becomes very small or when a predefined number of iterations is reached.
- **Cluster Assignment:** Once the algorithm converges, each data point is assigned to the cluster with the highest probability (highest responsibility).  
480
- **Number of Clusters ( $K$ ):** The number of clusters,  $K$ , is typically determined either by the user based on prior knowledge or by using techniques like the Bayesian Information Criterion (BIC) or cross-validation to find the optimal number of clusters.


485 In our study, we used cross-validation method and decided  $K$  as 3 for obtaining realistic results. Further explanation regarding the clustering algorithm can be obtained from recent literature on that subject (Hastie et al., 2009; Bishop and Nasrabadi, 2006; Dempster et al., 1977).

## Appendix B: Examples of blocking and zonal patterns evolved from the study

As previously indicated, employing the parameter configuration outlined in Table 1, we have successfully generated *earthlike* flow patterns. These flow regimes are visually represented in figures B2 and B1, corresponding to different values of the  
490 parameter  $k_d$ . Specifically, for lower values of  $k_d$ , we observe the coexistence of zonal, blocking, and transitional flow regimes,




**Figure B1.** Phase space dynamics of the model projected on the  $(\psi_{a,3}, \psi_{a,2})$ -plane is shown in panel (a) for  $k_d = 0.065$ . Gaussian mixture clusters covariance are represented with orange, green and red ellipsis. Orange cluster is associated with a zonal regime which can be identified by the geopotential height at 500 hPa in panel (b). Red and green clusters are respectively transition and blocking regimes attributed in panel (c) and (d), also at 500hPa geopotential height. The orographic profile of the domain is depicted in panels (b), (c) and (d).




**Figure B2.** Same as Fig. B1 but for  $k_d = 0.08$ . Here the red (panel (c)) and green clusters (panel (d)) are both blocking regimes and the orange one (panel (b)) is a transition regime.

as depicted in figure B1. Conversely, when  $k_d$  assumes higher values, we observe the emergence of two blocking flow regimes along with a transition regime, as illustrated in figure B2.



**Figure B3.** Same as Fig. B1 but for  $k_d = 0.105$  where the system exhibits periodic behavior results in to non identifiable flow regimes and indistinguishable clusters



**Figure B4.** Phase space dynamics of the model projected on the  $(\psi_{a,3}, \psi_{a,2})$ -plane is shown in panel (a) for  $k_d = 0.08$ . The figure illustrates the Gaussian mixture clustering results, where each data point is colored according to its corresponding cluster or flow regime. Panels (b), (c), and (d) depict the temporal evolution of barotropic ( $\psi_{a,1}$ ) and baroclinic ( $\theta_{a,1}$ ) atmospheric streamfunction, along with ground temperature ( $T_{g,1}$ ) for  $C_g = 300 \text{ W m}^{-2}$  and  $k_d = 0.08$ , with colors corresponding to their respective clusters.

*Author contributions.* AKX contributed to conceptualisation, method development, method implementation and data analysis and writing and visualisation. JD contributed to conceptualisation, model and method development, and text improvements. JD is also the developer of  
495 qgs land-atmosphere coupled model used for the study. SV contributed to supervision, conceptualisation and writing.

*Competing interests.* The contact author has declared that none of the authors has any competing interests.

*Acknowledgements.* We would like to thank Thomas Gilbert, Francesco Ragone and Oisín Hamilton for the helpful discussions. This project received funding from the European Union's Horizon 2020 research and innovation programme under Marie Skłodowska-Curie grant No. 956396.

## 500 **References**

- Arbabi, H. and Mezić, I.: Study of dynamics in post-transient flows using Koopman mode decomposition, *Physical Review Fluids*, 2, 124 402, 2017.
- Barsugli, J. J. and Battisti, D. S.: The basic effects of atmosphere–ocean thermal coupling on midlatitude variability, *Journal of the Atmospheric Sciences*, 55, 477–493, 1998.
- 505 Benzi, R., Hansen, A. R., and Sutera, A.: On stochastic perturbation of simple blocking models, *Quarterly Journal of the Royal Meteorological Society*, 110, 393–409, 1984.
- Bishop, C. M. and Nasrabadi, N. M.: *Pattern recognition and machine learning*, Springer, 2006.
- Breeden, M. L., Hoover, B. T., Newman, M., and Vimont, D. J.: Optimal North Pacific blocking precursors and their deterministic subseasonal evolution during boreal winter, *Monthly Weather Review*, 148, 739–761, 2020.
- 510 Cehelsky, P. and Tung, K. K.: Theories of multiple equilibria and weather regimes—A critical reexamination. Part II: Baroclinic two-layer models, *J. Atmos. Sci*, 44, 3282–3303, 1987.
- Charney, J. G. and DeVore, J. G.: Multiple flow equilibria in the atmosphere and blocking, *Journal of Atmospheric Sciences*, 36, 1205–1216, 1979.
- Charney, J. G. and Straus, D. M.: Form-drag instability, multiple equilibria and propagating planetary waves in baroclinic, orographically  
515 forced, planetary wave systems, *Journal of Atmospheric Sciences*, 37, 1157–1176, 1980.
- De Cruz, L., Demaeyer, J., and Vannitsem, S.: The modular arbitrary-order ocean-atmosphere model: MAOOAM v1. 0, *Geoscientific Model Development*, 9, 2793–2808, 2016.
- Demaeyer, J., De Cruz, L., and Vannitsem, S.: qgs: A flexible Python framework of reduced-order multiscale climate models, *Journal of Open Source Software*, 5, 2020.
- 520 Dempster, A. P., Laird, N. M., and Rubin, D. B.: Maximum likelihood from incomplete data via the EM algorithm, *Journal of the royal statistical society: series B (methodological)*, 39, 1–22, 1977.
- Dorrington, J. and Palmer, T.: On the interaction of stochastic forcing and regime dynamics, *Nonlinear Processes in Geophysics*, 30, 49–62, 2023.
- Eckmann, J.-P. and Ruelle, D.: Ergodic theory of chaos and strange attractors, *Reviews of modern physics*, 57, 617, 1985.
- 525 Egger, J.: Stochastically driven large-scale circulations with multiple equilibria, *Journal of Atmospheric Sciences*, 38, 2606–2618, 1981.
- Faranda, D., Masato, G., Moloney, N., Sato, Y., Daviaud, F., Dubrulle, B., and Yiou, P.: The switching between zonal and blocked mid-latitude atmospheric circulation: a dynamical system perspective, *Climate Dynamics*, 47, 1587–1599, 2016.
- Faranda, D., Messori, G., and Yiou, P.: Dynamical proxies of North Atlantic predictability and extremes, *Scientific reports*, 7, 41 278, 2017.
- Frederiksen, J. S., Collier, M. A., and Watkins, A. B.: Ensemble prediction of blocking regime transitions, *Tellus A: Dynamic Meteorology  
530 and Oceanography*, 56, 485–500, 2004.
- Ghil, M. and Robertson, A. W.: “Waves” vs. “particles” in the atmosphere’s phase space: A pathway to long-range forecasting?, *Proceedings of the National Academy of Sciences*, 99, 2493–2500, 2002.
- Hastie, T., Tibshirani, R., Friedman, J. H., and Friedman, J. H.: *The elements of statistical learning: data mining, inference, and prediction*, vol. 2, Springer, 2009.
- 535 Kautz, L.-A., Martius, O., Pfahl, S., Pinto, J. G., Ramos, A. M., Sousa, P. M., and Woollings, T.: Atmospheric blocking and weather extremes over the Euro-Atlantic sector—a review, *Weather and climate dynamics*, 3, 305–336, 2022.

- Kim, S.-H. and Kim, B.-M.: In search of winter blocking in the western North Pacific Ocean, *Geophysical Research Letters*, 46, 9271–9280, 2019.
- Kuptsov, P. V. and Parlitz, U.: Theory and computation of covariant Lyapunov vectors, *Journal of nonlinear science*, 22, 727–762, 2012.
- 540 Kwasniok, F.: Fluctuations of finite-time Lyapunov exponents in an intermediate-complexity atmospheric model: a multivariate and large-deviation perspective, *Nonlinear Processes in Geophysics*, 26, 195–209, 2019.
- Legras, B. and Ghil, M.: Persistent anomalies, blocking and variations in atmospheric predictability, *Journal of Atmospheric Sciences*, 42, 433–471, 1985.
- Legras, B. and Vautard, R.: A guide to Liapunov vectors, in: *Proceedings 1995 ECMWF seminar on predictability*, vol. 1, pp. 143–156, 1996.
- 545 Li, D., He, Y., Huang, J., Bi, L., and Ding, L.: Multiple equilibria in a land–atmosphere coupled system, *Journal of Meteorological Research*, 32, 950–973, 2018.
- Liu, J. S.: The collapsed Gibbs sampler in Bayesian computations with applications to a gene regulation problem, *Journal of the American Statistical Association*, 89, 958–966, 1994.
- Lorenz, E. N.: Maximum simplification of the dynamic equations, *Tellus*, 12, 243–254, 1960.
- 550 Lorenz, E. N.: Simplified dynamic equations applied to the rotating-basin experiments, *Journal of the Atmospheric Sciences*, 19, 39–51, 1962.
- Lorenz, E. N.: Deterministic nonperiodic flow, *Journal of atmospheric sciences*, 20, 130–141, 1963a.
- Lorenz, E. N.: The mechanics of vacillation, *Journal of the atmospheric sciences*, 20, 448–465, 1963b.
- Lucarini, V. and Gritsun, A.: A new mathematical framework for atmospheric blocking events, *Climate Dynamics*, 54, 575–598, 2020.
- 555 Lucarini, V., Faranda, D., de Freitas, J. M. M., Holland, M., Kuna, T., Nicol, M., Todd, M., Vaienti, S., et al.: *Extremes and recurrence in dynamical systems*, John Wiley & Sons, 2016.
- Lupo, A. R. and Smith, P. J.: Planetary and synoptic-scale interactions during the life cycle of a mid-latitude blocking anticyclone over the North Atlantic, *Tellus A*, 47, 575–596, 1995.
- Marshall, J. and Molteni, F.: Toward a dynamical understanding of planetary-scale flow regimes, *Journal of the atmospheric sciences*, 50, 1792–1818, 1993.
- 560 Mezić, I.: Spectrum of the Koopman operator, spectral expansions in functional spaces, and state-space geometry, *Journal of Nonlinear Science*, 30, 2091–2145, 2020.
- Monin, A. S.: *An introduction to the theory of climate*, Hingham, MA USA: Sold and distributed in the USA and Canada by Kluwer Academic Publishers, c1986, 261, 1986.
- 565 Nakamura, N. and Huang, C. S.: Atmospheric blocking as a traffic jam in the jet stream, *Science*, 361, 42–47, 2018.
- Parker, T. S. and Chua, L.: *Practical numerical algorithms for chaotic systems*, Springer Science & Business Media, 2012.
- Reinhold, B. and Pierrehumbert, R.: Corrections to “Dynamics of weather regimes: Quasi-stationary waves and blocking.”, *Mon. Wea. Rev.*, 113, 2055–2056, 1985.
- Reinhold, B. B. and Pierrehumbert, R. T.: Dynamics of weather regimes: Quasi-stationary waves and blocking, *Monthly Weather Review*, 570 110, 1105–1145, 1982.
- Schubert, S. and Lucarini, V.: Dynamical analysis of blocking events: spatial and temporal fluctuations of covariant Lyapunov vectors, *Quarterly Journal of the Royal Meteorological Society*, 142, 2143–2158, 2016.
- Shen, B.-W., Pielke Sr, R. A., and Zeng, X.: The 50th anniversary of the metaphorical butterfly effect since Lorenz (1972): Multistability, multiscale predictability, and sensitivity in numerical models, 2023.



- 575 Shimada, I. and Nagashima, T.: A numerical approach to ergodic problem of dissipative dynamical systems, *Progress of theoretical physics*, 61, 1605–1616, 1979.
- Shutts, G.: The propagation of eddies in diffluent jetstreams: Eddy vorticity forcing of ‘blocking’ flow fields, *Quarterly Journal of the Royal Meteorological Society*, 109, 737–761, 1983.
- Sura, P.: Noise-induced transitions in a barotropic  $\beta$ -plane channel, *Journal of the atmospheric sciences*, 59, 97–110, 2002.
- 580 Vannitsem, S.: Predictability of large-scale atmospheric motions: Lyapunov exponents and error dynamics, *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 27, 2017.
- Vannitsem, S. and Lucarini, V.: Statistical and dynamical properties of covariant Lyapunov vectors in a coupled atmosphere-ocean model—Multiscale effects, geometric degeneracy, and error dynamics, *Journal of Physics A: Mathematical and Theoretical*, 49, 224 001, 2016.
- 585 Vannitsem, S., Demaeyer, J., De Cruz, L., and Ghil, M.: Low-frequency variability and heat transport in a low-order nonlinear coupled ocean–atmosphere model, *Physica D: Nonlinear Phenomena*, 309, 71–85, 2015.
- Weeks, E. R., Tian, Y., Urbach, J., Ide, K., Swinney, H. L., and Ghil, M.: Transitions between blocked and zonal flows in a rotating annulus with topography, *Science*, 278, 1598–1601, 1997.
- Yoden, S.: Nonlinear Interactions in a Two-layer, Quasi-geostrophic, Low-order Model with Topography Part I: Zonal Flow-Forced Wave
- 590 Interactions, *Journal of the Meteorological Society of Japan. Ser. II*, 61, 1–18, 1983a.
- Yoden, S.: Nonlinear Interactions in a Two-layer, Quasi-geostrophic, Low-order Model with Topography Part II: Interactions between Zonal Flow, Forced Waves and Free Waves, *Journal of the Meteorological Society of Japan. Ser. II*, 61, 19–35, 1983b.
- Yoden, S.: Atmospheric predictability, *Journal of the Meteorological Society of Japan. Ser. II*, 85, 77–102, 2007.
- Zhengxin, Z. and Baozhen, Z.: Equilibrium states of ultra-long waves driven by non-adiabatic heating and blocking situation, *Science in*
- 595 *China (Series B)*, 4, 361–371, 1982.
- Zhu, Z.: Equilibrium states of planetary waves forced by topography and perturbation heating and blocking situation, *Advances in Atmospheric Sciences*, 2, 359–367, 1985.