

1 ~~18-2128~~ Dec 2023

2 **Fractionation of stable carbon isotopes during formate consumption in**
3 **anoxic rice paddy soils and lake sediments**

4

5 Ralf Conrad¹, Peter Claus¹

6

7 ¹Max Planck Institute for Terrestrial Microbiology, Karl-von-Frisch-Str. 10, 35043 Marburg,
8 Germany

9

10

11 Correspondence to: Ralf Conrad (Conrad@mpi-marburg.mpg.de)

12

13 **Running head:** Isotope fractionation by anaerobic formate consumption

14

15

16 **Abstract.**


17 Formate is energetically equivalent to hydrogen and thus, is an important intermediate during
18 the breakdown of organic matter in anoxic rice paddy soils and lake sediments. Formate is a
19 common substrate for methanogenesis, homoacetogenesis and sulfate reduction. However,
20 how much these processes contribute to formate degradation and fractionate carbon stable
21 isotopes is largely unknown. Therefore, we measured the conversion of formate to acetate,
22 CH₄ and CO₂ and the δ¹³C of these compounds in samples of paddy soils from Vercelli
23 (Italy) and the International Rice Research Institute (IRRI, the Philippines) and of sediments
24 from the NE and SW basins of Lake Fuchskuhle (Germany). The samples were suspended in
25 phosphate buffer (pH 7.0) both in the absence and presence of sulfate (gypsum) and of
26 methyl fluoride (CH₃F), an inhibitor of acetoclastic methanogenesis. In the paddy soils,
27 formate was mainly converted to acetate both under methanogenic and sulfidogenic
28 conditions. Methane was only a minor product and was mainly formed from the acetate. In
29 the lake sediments, the product spectrum was similar, but only under methanogenic
30 conditions. In the presence of sulfate, however, acetate and CH₄ were only minor products.
31 The isotopic enrichment factors (ε_{form}) of formate consumption, determined by Mariotti plots,
32 were in the low range of -8‰ to -2.5‰ when sulfate was absent and formate was mainly
33 converted to acetate and CH₄. However, no enrichment factor was detectable when formate
34 was degraded with sulfate to mainly CO₂. The δ¹³C of acetate was by about 25-50‰ more
35 negative than that of formate indicating acetate production by chemolithotrophic
36 homoacetogenesis. Hence, formate seems to be an excellent substrate for homoacetogenesis
37 in anoxic soils and sediments, so that this process is competing well with methanogenesis and
38 sulfate reduction.


39

40 **1 Introduction**


41 Formate is energetically almost equivalent to H₂ (Schink et al. 2017) and thus, is an
42 important intermediate in the anaerobic degradation of organic matter. Formate is a product
43 of microbial fermentation, where it is for example produced in pyruvate cleavage by pyruvate
44 formate lyase (Thauer et al., 1977) or by reduction of CO₂ (Schuchmann and Müller, 2013).
45 Formate can also be produced in secondary fermentation, such as oxidation of butyrate or
46 propionate (Dong et al., 1994; Sieber et al., 2014). In fact, formate and H₂ may equivalently
47 be used as electron shuttles between secondary fermenting bacteria and methanogens
48 (Montag and Schink, 2018; Schink et al., 2017)

49 Formate can serve alternatively to H₂ as a substrate for methanogenesis (Zinder, 1993),
50 (homo)acetogenesis (Drake, 1994) or sulfate reduction (Widdel, 1988), i.e.:





54 Formate may also be a substrate for syntrophic bacteria, which live from the little Gibbs free
55 energy ($\Delta G^{0'} = -3.4 \text{ kJ mol}^{-1}$) that is generated by the conversion of formate to H_2 plus CO_2
56 (Dolfing et al., 2008; Kim et al., 2010; Martins et al., 2015), i.e.



58 Formate can also be enzymatically equilibrated with H_2 and CO_2 without energy generation.
59 This reaction happens in any organism possessing the suitable enzymes, such as formate
60 hydrogen lyase or hydrogen-dependent carbon dioxide reductase, and in anoxic sediments
61 (DeGraaf and Cappenberg, 1996; Peters et al., 1999; Schuchmann et al., 2018):



63 Formate has been identified as an important substrate for methanogenesis,
64 homoacetogenesis or sulfate reduction in lake sediments (DeGraaf and Cappenberg, 1996;
65 Lovley and Klug, 1982; Phelps and Zeikus, 1985), soils (Kotsyurbenko et al., 1996; Küsel
66 and Drake, 1999; Rothfuss and Conrad, 1993), mires (Hausmann et al., 2016; Hunger et al.,
67 2011; Liebner et al., 2012; Wüst et al., 2009) and marine sediments (Glombitza et al., 2015).
68 However, it is not very clear to which extent formate-dependent methanogenesis,
69 homoacetogenesis and sulfate reduction are actually operative and to which extent formate
70 affects stable carbon isotope fractionation. The $\delta^{13}\text{C}$ values of compounds involved in the
71 degradation process of organic matter provide valuable information on the metabolic
72 pathways involved (Conrad, 2005; Elsner et al., 2005; Hayes, 1993). However, for correct
73 interpretation the knowledge of the enrichment factors (ϵ) of the major metabolic processes is
74 also important. The ϵ values of methanogenesis or homoacetogenesis from H_2 plus CO_2 are
75 large (Blaser and Conrad, 2016). However, our knowledge of carbon isotope fractionation
76 with formate as substrate is scarce. In cultures of homoacetogenic bacteria the carbon in the
77 acetate produced from formate was strongly depleted in ^{13}C ($\epsilon = -56.5\text{‰}$) almost similarly as
78 with CO_2 as carbon source (Freude and Blaser, 2016). However, it is not known which
79 enrichment factors operate in methanogenic or sulfidogenic environmental samples.

80 Therefore, we measured isotope fractionation in methanogenic and sulfidogenic rice paddy
81 soils and lake sediments amended with formate. We recorded the consumption of formate
82 along with the production of acetate, CH_4 and CO_2 and measured the $\delta^{13}\text{C}$ of these
83 compounds. We also used the treatment with methyl fluoride (CH_3F) to inhibit the
84 consumption of acetate by methanogenic archaea (Janssen and Frenzel, 1997). We used the
85 same environmental samples as for the study of carbon isotope fractionation during
86 consumption of acetate (Conrad et al., 2021) and propionate (Conrad and Claus, 2023), i.e.,
87 rice paddy soils from Vercelli, Italy and the International Rice Research Institute (IRRI, the
88 Philippines) and sediments from the NE and SW basins of Lake Fuchskuhle (Germany). The
89 molecular data characterizing the microbial community compositions in these samples are
90 found in Conrad et al. (2021).

91

92 **2 Materials and Methods**

93 *2.1 Environmental samples and incubation conditions*

94 The soil samples were from the research stations in Vercelli, Italy and the International Rice
95 research Institute (IRRI) in the Philippines. Sampling and soil characteristics were described
96 before (Liu et al., 2018). The lake sediments (top 10 cm layer) were from the NE and SW
97 basins of Lake Fuchskuhle, [an acidic-bog \(pH 4.2 – 4.6\) lake in Northeastern Germany](#)
98 (Casper et al., 2003). [The lake was artificially divided into four compartments in 1987 and](#)
99 [finally in 1991, resulting in four nearly equal sized compartments, each with a different](#)
100 [catchment area. The NE basin is characterized by higher biomass and activity throughout all](#)
101 [trophic levels in the water column than the SW basin.](#) ~~They~~ [The lake sediments](#) were sampled
102 in July 2016 using a gravity core sampler as described before (Kanaparthi et al., 2013). [The](#)
103 [experiments with rice field soil were carried out in 2016, those with sediments of Lake](#)
104 [Fuchskuhle in 2017.](#)

105 The experimental setup was exactly the same as during previous studies of acetate
106 consumption (Conrad et al., 2021) and propionate consumption (Conrad and Claus, 2023).
107 For methanogenic conditions, paddy soil was mixed with autoclaved anoxic H₂O (prepared
108 under N₂) at a ratio of 1:1 and incubated under N₂ at 25°C for 4 weeks. In a second
109 incubation, for sulfidogenic conditions, paddy soil was mixed with autoclaved anoxic H₂O at
110 a ratio of 1:1, was amended with 0.07 g CaSO₄·2H₂O, and then incubated under N₂ at 25°C
111 for 4 weeks. These two preincubated soil slurries were sampled and stored at -20°C for later
112 molecular analysis (see data in Conrad et al. (2021)). The preincubated soil slurries were also
113 used (in 3 replicates) for the following incubation experiments. Two different sets of
114 incubations were prepared. In the first set (resulting in methanogenic conditions), 5 mL soil
115 slurry preincubated without sulfate was incubated at 25°C with 40 mL of 20 mM potassium
116 phosphate buffer (pH 7.0) in a 150-mL bottle under an atmosphere of N₂. The bottles were
117 ~~the~~-amended with (i) 5 mL H₂O; (ii) 5 mL H₂O + 4.5 mL CH₃F; (iii) 5 mL 200 mM sodium
118 formate; (iv) 5 mL 200 mM sodium formate + 4.5 mL CH₃F. In the second set (resulting in
119 sulfidogenic conditions), 5 mL soil slurry preincubated with sulfate was incubated at 25°C
120 with 40 mL of 20 mM potassium phosphate buffer (pH 7.0) in a 150-mL bottle under an
121 atmosphere of N₂. The amendments were the same as above, but with the addition of 200 µl
122 of a CaSO₄ suspension corresponding to a concentration of 2.5 M (giving a final
123 concentration of 10 mM sulfate).

124 For lake sediments under methanogenic conditions, 5 ml sediment was incubated in 3
125 replicates at 10°C (which is close to the in-situ temperature) with 40 ml of 20 mM potassium
126 phosphate buffer (pH 7.0) in a 150-ml bottle under an atmosphere of N₂. The bottles were ~~the~~
127 amended with (i) 5 ml H₂O; (ii) 5 ml H₂O + 4.5 ml CH₃F; (iii) 5 ml 200 mM sodium formate;
128 (iv) 5 ml 200 mM sodium formate + 4.5 ml CH₃F. For sulfidogenic conditions, lake

129 sediments were preincubated with sulfate by adding 0.1 g CaSO₄·2H₂O (gypsum) to 50 ml
130 sediment and incubating at 10°C for 4 weeks. For sulfidogenic conditions, 5 ml of the
131 preincubated sediment was incubated in 3 replicates at 10°C with 40 ml of 20 mM potassium
132 phosphate buffer (pH 7.0) in a 150-ml bottle under an atmosphere of N₂. The bottles were
133 amended as above, but in addition also with 200 µl of a CaSO₄ suspension giving a final
134 concentration of 10 mM sulfate. Samples for later molecular analysis were taken from the
135 original lake sediment and from the lake sediment preincubated with sulfate. The samples
136 were stored at -20°C (see data in Conrad et al. (2021)).

137

138 2.2 Chemical and isotopic analyses

139 Gas samples for analysis of partial pressures of CH₄ and CO₂ were taken from the
140 headspace of the incubation bottles after vigorous manual shaking for about 30 s using a gas-
141 tight pressure-lock syringe, which had been flushed with N₂ before each sampling. Soil
142 slurries were sampled, centrifuged and filtered through a 0.2 µm cellulose membrane filter
143 and stored frozen at -20°C for later fatty acid analysis. Chemical and isotopic analyses were
144 performed as described in detail previously (Govert and Conrad, 2009). Methane was
145 analyzed by gas chromatography (GC) with flame ionization detector. Carbon dioxide was
146 analyzed after conversion to CH₄ with a Ni catalyst. Stable isotope analyses of ¹³C/¹²C in gas
147 samples were performed using GC-combustion isotope ratio mass spectrometry (GC-C-
148 IRMS). Formate and acetate were measured using high-performance liquid chromatography
149 (HPLC) linked via a Finnigan LC IsoLink to an IRMS. The isotopic values are reported in the
150 delta notation (δ¹³C) relative to the Vienna Peedee Belemnite standard having a ¹³C/¹²C ratio
151 (R_{standard}) of 0.01118: δ¹³C = 10³ (R_{sample}/R_{standard} - 1). The precision of the GC-C-IRMS was
152 ± 0.2‰, that of the HPLC-IRMS was ± 0.3‰.

153

154 2.3 Calculations

155 Millimolar concentrations of CH₄ were calculated from the mixing ratios (1 ppmv = 10⁻⁶
156 bar) measured in the gas phase of the incubation bottles: 1000 ppmv CH₄ correspond to 0.09
157 µmol per mL of liquid. Note, that this is the total amount of CH₄ in the gas phase relative to
158 the liquid phase.

159 Fractionation factors for reaction A → B are defined after Hayes (Hayes, 1993) as:

$$160 \alpha_{A/B} = (\delta_A + 1000) / (\delta_B + 1000) \quad (7)$$

161 also expressed as $\epsilon \equiv 1000 (1 - \alpha)$ in permil. The carbon isotope enrichment factor ϵ_{form}
162 associated with formate consumption was calculated from the temporal change of δ¹³C of
163 formate as described by Mariotti et al. (Mariotti et al., 1981) from the residual reactant

$$164 \delta_r = \delta_{ri} + \epsilon [\ln(1-f)] \quad (8)$$

165 where δ_{ri} is the isotopic composition of the reactant (formate) at the beginning, and δ_r is the
166 isotopic composition of the residual formate, both at the instant when f is determined. f_{form} is

167 the fractional yield of the products based on the consumption of formate ($0 < f_{\text{form}} < 1$).
168 Linear regression of $\delta^{13}\text{C}$ of formate against $\ln(1 - f)$ yields ϵ_{form} as the slope of best fit lines.
169 The regressions of $\delta^{13}\text{C}$ of formate were done for data in the range of $f_{\text{form}} < 0.7$. The linear
170 regressions were done individually for each experimental replicate ($n = 3$) and were only
171 accepted if $r^2 > 0.7$. The ϵ values resulting from the replicate experiments were then averaged
172 (\pm SE).

174 **3 Results**

175 *3.1 Conversion of formate under methanogenic and sulfidogenic conditions*

176 The rice paddy soils were submerged and preincubated to create methanogenic or
177 sulfidogenic conditions. Samples of these soils were suspended in buffer at pH 7 and
178 amended with formate. In the Vercelli soil, formate was consumed after a lag phase of 4 days
179 under methanogenic and 5 days under sulfidogenic conditions (Fig. 1a). During this time the
180 pH increased from pH 7 up to pH 8 despite buffering. Formate consumption was not inhibited
181 by CH_3F (Fig. 1a). Similar results were obtained with IRRI soil (Fig. S1). Acetate was
182 produced concomitantly with formate consumption, again without effect by CH_3F (Fig. 1b).
183 The production of acetate under sulfidogenic conditions was smaller than under
184 methanogenic conditions. Methane was also produced under both methanogenic and
185 sulfidogenic conditions concomitantly with formate consumption (Fig. 1c; S1c). It is
186 noteworthy that CH_3F inhibited the production of CH_4 (Fig. 1c; S1c). Finally, CO_2 was
187 produced under all conditions without lag phase and without effect by CH_3F (Fig. 1de). In
188 Vercelli soil, CO_2 production was about ~~twice-two times larger~~ under sulfidogenic than under
189 methanogenic conditions (Fig. 1de). In IRRI soil, it was only slightly larger (Fig. S1de). The
190 accumulation of acetate plus CH_4 was equimolar to the consumption of formate in terms of
191 electron equivalents, while the accumulation of CH_4 alone accounted only for $<30\%$, in the
192 presence of CH_3F even less (Fig. 2a; S2a). Hence, acetate was the more important product of
193 formate consumption. Under sulfidogenic conditions, accumulation of acetate plus CH_4 was
194 less than equimolar, especially in Vercelli soil (Fig. 2b), probably since formate was instead
195 converted to CO_2 . However, acetate formation was still substantial accounting for 60-80% of
196 formate consumption (Fig. 2b; S2b).

197 The sediments from Lake Fuchskuhle were methanogenic in-situ so that preincubation of
198 the samples was not required. However, sulfidogenic conditions were created analogously to
199 the paddy soils by preincubation with sulfate (gypsum). Substantial formate depletion did not
200 start before about 20 days of incubation both in sediments from the NE basin (Fig. 3) and the
201 SW basin (Fig. S3). Again, CH_3F only inhibited the production of CH_4 but not that of acetate
202 or CO_2 (Fig. 3; S3). The main difference to the paddy soils was that CH_4 was not produced
203 concomitantly with formate consumption, but started right from the beginning. However, the
204 amounts of CH_4 produced were only small and were apparently due to the little formate that

205 was consumed in the beginning of incubation (i.e., before day 20), as seen by the fact that
206 CH₄ production in the water control (not amended with formate) was negligible (Fig. 3c;
207 S3c). Production of CO₂ started without lag phase but accelerated together with formate
208 consumption (Fig. 3d; S3d). In the lake sediments, CH₄ accounted only for <10% of formate
209 consumption, while acetate was the main product when sulfate was absent (Fig. 4a, S4a). In
210 contrast to the paddy soils, formate consumption in both lake sediments was much slower
211 under sulfidogenic than under methanogenic conditions (Fig. 3a; S3a). In the sediment from
212 SW basin, formate consumption was very slow so that less than half of the formate was
213 consumed during 80 days of incubation and consumption was not completed until the end of
214 the experiment (Fig. S3a). Very little acetate was produced and no CH₄ was formed from
215 formate in both lake sediments, when sulfate was present (Fig. 4b, S4b).

216

217 3.2 Isotope fractionation during formate consumption

218 In the rice paddy soils, ~~values of~~ $\delta^{13}\text{C}$ -~~values of formate~~ increased when formate was
219 ~~being~~ consumed indicating discrimination against the heavy carbon isotope. This process was
220 not affected by CH₃F and was similar without and with sulfate (Fig. 1e; S1e). The same was
221 the case with the sediment from the NE lake basin, but only in the absence of sulfate (Fig.
222 3e). With sulfate, the $\delta^{13}\text{C}$ of formate slowly decreased with time (Fig. 3e). In the sediment
223 from the SW basin, $\delta^{13}\text{C}$ of formate slowly decreased (without sulfate) or stayed constant
224 with time (with sulfate) (Fig. S3e). Note that formate was not completely consumed in the
225 SW sediment when sulfate was present (Fig. S3a).

226 Mariotti plots of $\delta^{13}\text{C}$ of formate as function of f_{form} resulted in negative slopes (Fig. 4;
227 S5). Hence, the enrichment factors (ϵ_{form}) for the paddy soils, both without and with sulfate,
228 and for the sediments from the NE basin of Lake Fuchskuhle without sulfate showed that the
229 light isotope of formate carbon was preferred. Values of ϵ_{form} were in the range of -8.5 to -
230 2.5‰ (Fig. 6). Under sulfidogenic conditions, however, the Mariotti plots of the sediments
231 from the NE basin (Fig. 5) did not show a negative slope and ϵ_{form} could not be determined.
232 The same was the case for the sediments from the SW basin (Fig. 6).

233 The negative ϵ_{form} indicates that products of formate should be depleted in ^{13}C . Indeed the
234 $\delta^{13}\text{C}$ of acetate and CH₄ were generally more negative than the $\delta^{13}\text{C}$ of formate. This was the
235 case in the paddy soils from Vercelli (Fig. 1f) and the IRRI (Fig. S1f) as well as in the
236 sediments from the NE basin (Fig. 3f) and the SW basin (Fig. S3f) of Lake Fuchskuhle. In the
237 sediment of the NE basin, the $\delta^{13}\text{C}$ of acetate increased from very low -95‰ to finally about -
238 57‰ in parallel with formate consumption (Fig. 3f). CO₂ was also produced during formate
239 degradation to various extent (equ. 1, 2 and 3). Since the pH was in a range of pH 7 to pH 8,
240 CO₂ was also converted to bicarbonate. The $\delta^{13}\text{C}$ of bicarbonate is generally by about 10‰
241 more positive than the $\delta^{13}\text{C}$ of CO₂ (Stumm and Morgan, 1996). The $\delta^{13}\text{C}$ of the gaseous CO₂
242 was always close to the $\delta^{13}\text{C}$ of formate or was more positive. In the paddy soils and the NE

243 basin of Lake Fuchskuhle, the $\delta^{13}\text{C}$ of CO_2 increased in parallel with the increasing $\delta^{13}\text{C}$ of
244 formate (Fig. 1h, 3h; S1h). The $\delta^{13}\text{C}$ of the gaseous CO_2 produced from the formate-amended
245 samples was initially more negative than that from the unamended samples, but eventually
246 the $\delta^{13}\text{C}$ increased above these values when formate was completely consumed (Fig. 1h, 3h;
247 S3h).

248 The $\delta^{13}\text{C}$ values of the initial formate were about -24‰ (Fig. 5). When formate was
249 completely consumed, the $\delta^{13}\text{C}$ values of the products acetate and CH_4 were always more
250 negative. The average $\delta^{13}\text{C}$ values of the products after complete consumption of formate are
251 shown in Fig. 7. In the absence of sulfate, $\delta^{13}\text{C}$ of acetate was in a range of -51‰ to -49‰
252 and -70‰ to -63‰, in the paddy soils and lake sediments, respectively (Fig. 7). In the
253 presence of sulfate, $\delta^{13}\text{C}$ of acetate was in a range of -57‰ to -52‰ and -78‰ to -72‰, in
254 the paddy soils and lake sediments (only NE basin), respectively (Fig. 7). The $\delta^{13}\text{C}$ of CH_4
255 was in a range of -70‰ to -54‰ and -60‰ to -54‰, in the absence and presence of sulfate,
256 respectively (Fig. 7). The $\delta^{13}\text{C}$ of gaseous CO_2 (for bicarbonate plus 10‰) was in a range of -
257 23‰ to -11‰ and -24‰ to -19‰, in the absence and presence of sulfate, respectively (Fig.
258 7).

259

260 **4 Discussion**

261 *4.1 Formate degradation under acetogenic/methanogenic conditions*

262 In rice paddy soils formate was consumed within <10 days. The absence of sulfate did not
263 allow sulfidogenic (equ.3) degradation, but allowed the operation of methanogenic (equ.1),
264 homoacetogenic (equ.2) or syntrophic (equ.4) degradation. Syntrophic degradation is still
265 disputed, since many microorganisms are able to enzymatically equilibrate H_2 and formate
266 and thus prohibit generation-exploitation of the difference in the energy content (Montag and
267 Schink, 2018; Schink et al., 2017). Syntrophic formate degradation generates-is exergonic by
268 only a few kilojoules of Gibbs free energy per mole and requires the coupling with
269 methanogenesis or other efficient hydrogen (electron) scavengers. Although formate-driven
270 CH_4 production was observed in our study, the production was sensitive to inhibition by
271 CH_3F indicating that CH_4 was predominantly produced from acetate rather than from H_2 .
272 Therefore, syntrophic formate oxidation coupled to CH_4 production was probably not a major
273 pathway.

274 Acetate was the most important product of formate degradation in the paddy soils as well
275 as in the lake sediments. Methane also was a product, but was much less important than
276 acetate. Furthermore, it was predominantly produced from acetate as shown by the inhibition
277 by CH_3F and the concomitant decrease of $\delta^{13}\text{C}$ of CH_4 , which is characteristic for
278 hydrogenotrophic methanogenesis that is not inhibited by CH_3F (Conrad et al., 2010). Hence,
279 formate was apparently primarily degraded by homoacetogenesis (equ.1). Only part of the
280 produced acetate was immediately used by aceticlastic methanogenesis generating CH_4 as

281 secondary product. Although formate is a perfect substrate for homoacetogenic bacteria
282 operating the Wood-Ljungdahl pathway (WLP) (Drake, 1994), the yield of Gibbs free energy
283 per mole formate is less for homoacetogenic than for methanogenic degradation (Dolfing et
284 al., 2008). Thus, it is surprising that formate-driven homoacetogenesis prevailed over
285 methanogenesis. Nevertheless, simultaneous operation of homoacetogenesis and
286 methanogenesis from formate has been observed before in a fen soil (Hunger et al., 2011).
287 Homoacetogenesis prevailing over methanogenesis has also frequently been observed with
288 H₂/CO₂ as substrate (Conrad et al., 1989; Nozhevnikova et al., 1994), indicating that
289 homoacetogens can take particular advantage from low temperatures (Conrad, 2023) or the
290 availability of secondary substrates (Peters et al., 1998). It is noteworthy that homoacetogens
291 have to invest ATP for fixation of formate, while methanogens are able to bypass this step
292 (Lemaire et al., 2020). Perhaps it is such energy investment which makes the homoacetogens
293 to competitive formate utilizers.

294 Formate consumption was recorded upon addition of formate to initial concentrations of
295 about 15 mM, which was much higher than the in-situ concentration being typically on the
296 order of a few micromolar (Montag and Schink, 2018). However, the increased concentration
297 allowed stable isotope fractionation, which would not occur under formate limitation. The
298 $\delta^{13}\text{C}$ of the produced acetate was by about 24-33‰ lower than that of formate. This isotopic
299 discrimination between formate and acetate is similar to that measured in a culture of the
300 homoacetogen *Thermoanaerobacter kivui* (Freude and Blaser, 2016). However, this
301 discrimination is much larger than the isotopic enrichment factors (ϵ_{form} of -8‰ to -2.5‰)
302 determined from the change of $\delta^{13}\text{C}$ during formate consumption. There are two conceivable
303 explanations for this observation. (1) Formate is disproportionated to CO₂ and acetate. In the
304 WLP three formate are oxidized to CO₂, one formate is reduced to the methyl group of
305 acetate and one of the produced CO₂ is reduced to the carboxyl group of acetate. The
306 disproportionation of formate to acetate and 2 CO₂ is possibly a branch point (Fry, 2003;
307 Hayes, 2001), at which the carbon flow is split into the production of ¹³C-enriched CO₂ and
308 ¹³C-depleted acetate, which together result in the ϵ_{form} observed. (2) Formate first is
309 completely converted to CO₂ plus H₂ (equ.5) or other electron equivalents. This reaction
310 displays the ϵ_{form} determined by the Mariotti plots. Acetate is then produced via the WLP by
311 the chemolithotrophic reduction of 2 CO₂ to acetate, of which the isotopic enrichment factor
312 is typically on the order of about -55‰ (Blaser and Conrad, 2016). In any case, it is plausible
313 to assume that acetate was formed via the WLP. In the WLP, oxidation of formate is
314 catalyzed by a formate dehydrogenase, which provides CO₂ to the carboxyl branch of the
315 WLP. The methyl branch of the WLP normally starts with formate being converted to
316 formyl-THF. However, it can also start with the reduction of CO₂ to formate with a
317 hydrogen-dependent carbon dioxide reductase (HDCD). Homoacetogens (e.g., *Acetobacter*
318 *woodii*, *T. kivui*) contain such a HDCD, which allows the interconversion of formate and H₂

319 plus CO₂ (Jain et al., 2020; Schuchmann et al., 2018). The isotope discrimination in our
320 experiments indicates that the CO₂ produced from formate has been enriched in ¹³C rather
321 than depleted, thus supporting the first explanation. The δ¹³C of CO₂ produced from formate
322 was initially lower than that of the unamended soil or sediment being on the order of -20‰ to
323 -10‰ (Fig. 1h, 3h, S1h, S3h). Eventually, however, δ¹³C of CO₂ reached values of -25‰ to -
324 10‰ (Fig. 7). The δ¹³C of bicarbonate is 10‰ more positive than that of CO₂. This mixed
325 inorganic carbon would be the CO₂ substrate for WLP, which together with formate generates
326 the acetate having a δ¹³C of about -70‰ to -50‰ (Fig. 7).

327 Methane was a minor product of formate degradation in all soils and sediments. Since CH₄
328 formation was strongly inhibited by CH₃F, it was most likely produced from acetate by
329 aceticlastic methanogens. Since CH₄ production from the soils or sediments was much lower
330 without formate amendment, the CH₄ must have primarily been produced from the acetate
331 that was generated from formate. The δ¹³C of CH₄ in the soil incubations was more negative
332 than that of acetate (Fig. 7). The difference between the δ¹³C of CH₄ and the δ¹³C of acetate
333 indicated an isotopic enrichment factor of ε_{ac-CH₄} = -10‰ to -8‰, which is close to the
334 enrichment factor of aceticlastic *Methanosaeta* (*Methanotherix*) *concilii* (Penning et al., 2006).
335 In the lake sediments, the δ¹³C of CH₄ and acetate were not much different indicating that
336 acetate was instantaneously consumed by methanogens as it was produced by homoacetogens
337 so that carbon isotopes were not discriminated. Both, paddy soils and lake sediments
338 contained *mcrA* genes (coding for a subunit of methyl CoM reductase) of *Methanosaetaceae*
339 (*Methanotrichaceae*) (Conrad et al., 2021).

340

341 4.2 Formate degradation under sulfidogenic conditions

342 In the rice paddy soils, formate was consumed within ten days when sulfate was present,
343 not quite as fast as without sulfate. In the lake sediments, however, sulfidogenic formate
344 consumption was much slower. Formate degradation by sulfate reduction normally results in
345 complete oxidation to CO₂ (equ.3). In the lake sediments, CO₂ was indeed the main
346 degradation product. However, in the paddy soils substantial amounts of acetate and even
347 CH₄ were also produced. The homoacetogenic bacteria in these soils apparently competed
348 well with the sulfate reducing bacteria, although the soils had been adapted by preincubation
349 in the presence of sulfate. The production of acetate and CH₄ was dependent on formate
350 degradation, since no production was observed in the unamended control. Production of CH₄
351 was inhibited by CH₃F indicating that aceticlastic methanogenesis was the main process of
352 CH₄ production. The carbon isotope fractionation of formate was similar as under non-
353 sulfidogenic conditions, exhibiting a small ε_{form} of -8‰ to -3.5‰ (Fig. 5) and displaying a
354 strong isotope effect with the formation of acetate (δ¹³C = -57--52‰) and CH₄ (δ¹³C = -60--
355 58‰). The mechanism of fractionation is probably the same (see above).

356 In the lake sediments, however, sulfidogenic degradation of formate was much slower
357 than methanogenic/acetogenic degradation. In the sediment of the SW basin, formate was not
358 even completely degraded within 80 days. In the sediments of both lake basins, neither
359 acetate nor CH₄ was a major product of sulfidogenic formate degradation. Hence, formate
360 was apparently degraded according to equ.3 forming CO₂ as main carbon product. This
361 formation process displayed no depletion of the heavy carbon isotope, as the Mariotti plots of
362 δ¹³C of formate did not exhibit a negative slope. The δ¹³C of the CO₂ slowly decreased with
363 increasing fraction of formate consumed (Fig. 3h; 5c), probably involving isotope exchange
364 between formate and CO₂ (DeGraaf and Capenberg, 1996). The little acetate, which was
365 formed, displayed a δ¹³C of -77‰ (Fig. 7b) indicating that it was produced by a similar
366 mechanism as in the absence of sulfate, presumably via the WLP.

367 The strong differences between rice paddy soils and lake sediments were possibly caused
368 by their different microbial communities (Conrad et al., 2021). The differences were seen in
369 the composition of the *mcrA* and *dsrB* genes coding for methyl CoM reductase and
370 dissimilatory sulfate reductase, respectively, as well as the gene coding for the bacterial 16S
371 rRNA (data are shown in Conrad et al. (2021)). The microbial community structures based
372 on composition of these genes was/were similar whether the soils and sediments were
373 amended with sulfate or not. However, they were strongly different between soils and
374 sediments (Conrad et al., 2021). Unfortunately, these data do not allow to discriminate for
375 particular taxa of homoacetogenic bacteria. Nevertheless, it is possible that formate-
376 consuming homoacetogens were more prevalent in the soils than in the sediments and
377 accordingly competed more or less with the formate-consuming sulfate reducers.

378

379 4.3 Conclusions

380 Formate was found to be an excellent substrate for acetate formation in the paddy soils as
381 well as in the lake sediments, confirming and extending similar observations in a fen soil
382 (Hunger et al., 2011). In the anoxic soils, acetate was the major product even in the presence
383 of sulfate, which would have allowed sulfate reduction. The acetate was strongly depleted in
384 ¹³C relative to formate, but the consumption of formate itself displayed only a small isotopic
385 enrichment factor. Therefore, it is likely that formate was disproportionated to ¹³C-depleted
386 acetate and ¹³C-enriched CO₂. The δ¹³C of CO₂ was indeed slightly higher than that of
387 formate. Acetate was most likely produced by homoacetogenesis via the WLP. The produced
388 acetate was then used by aceticlastic methanogens (probably by *Methanotherix*), but only to
389 minor extent, resulting in further depletion of ¹³C. The homoacetogenic bacteria in the paddy
390 soils apparently competed well with both methanogenic and sulfate-reducing
391 microorganisms, when formate was the substrate. The preference of homoacetogenesis as
392 degradation pathway is unexpected, since other substrates, such as acetate and propionate, are
393 degraded in these paddy soils ~~degraded~~ by methanogenesis or sulfate reduction (Conrad et al.,

394 2021) (Conrad and Claus, 2023). Only in the lake sediments, formate oxidation by sulfate
395 reduction was more prevalent than homoacetogenesis.

396

397 **Supplement link**

398

399 **Author contribution:** RC designed the experiments, evaluated the data and wrote the
400 manuscript. PC conducted the experiments.

401

402 **Conflicting interests:** The contact authors has declared that neither of the authors has any
403 competing interests.

404

405 **Financial Support**

406 This research has been supported by the Fonds der Chemischen Industrie (grant no. 163468).

407

408

409

References

- 410
411
412 Blaser, M. and Conrad, R.: Stable carbon isotope fractionation as tracer of carbon
413 cycling in anoxic soil ecosystems, *Curr. Opin. Biotechnol.*, 41, 122-129,
414 2016.
- 415 Casper, P., Chan, O. C., Furtado, A. L. S., and Adams, D. D.: Methane in an
416 acidic bog lake: The influence of peat in the catchment on the biogeochemistry
417 of methane, *Aquat. Sci.*, 65, 36-46, 2003.
- 418 Conrad, R.: Quantification of methanogenic pathways using stable carbon
419 isotopic signatures: a review and a proposal, *Org. Geochem.*, 36, 739-752,
420 2005.
- 421 Conrad, R.: Complexity of temperature dependence in methanogenic microbial
422 environments, *Front. Microbiol.*, 14, 1232946-
423 doi:10.3389/fmicb.2023.1232946, 2023.
- 424 Conrad, R., Bak, F., Seitz, H. J., Thebrath, B., Mayer, H. P., and Schütz, H.:
425 Hydrogen turnover by psychrotrophic homoacetogenic and mesophilic
426 methanogenic bacteria in anoxic paddy soil and lake sediment, *FEMS*
427 *Microbiol. Ecol.*, 62, 285-294, 1989.
- 428 Conrad, R. and Claus, P.: Fractionation of stable carbon isotopes during microbial
429 propionate consumption in anoxic rice paddy soils, *Biogeosciences*, 20, 3625-
430 3635, 2023.
- 431 Conrad, R., Klose, M., Claus, P., and Enrich-Prast, A.: Methanogenic pathway,
432 ¹³C isotope fractionation, and archaeal community composition in the sediment
433 of two clearwater lakes of Amazonia, *Limnol. Oceanogr.*, 55, 689-702, 2010.
- 434 Conrad, R., Liu, P., and Claus, P.: Fractionation of stable carbon isotopes during
435 acetate consumption by methanogenic and sulfidogenic microbial communities
436 in rice paddy soils and lake sediments, *Biogeosciences*, 18, 6533-6546, 2021.
- 437 DeGraaf, W. and Cappenberg, T. E.: Evidence for isotopic exchange during
438 metabolism of stable- isotope-labeled formate in a methanogenic sediment,
439 *Appl. Environ. Microbiol.*, 62, 3535-3537, 1996.
- 440 Dolfing, J., Jiang, B., Henstra, A. M., Stams, A. J. M., and Plugge, C. M.:
441 Syntrophic growth on formate: a new microbial niche in anoxic environments,
442 *Appl. Environ. Microbiol.*, 74, 6126-6131, 2008.
- 443 Dong, X. Z., Plugge, C. M., and Stams, A. J. M.: Anaerobic degradation of
444 propionate by a mesophilic acetogenic bacterium in coculture and triculture
445 with different methanogens, *Appl. Environ. Microbiol.*, 60, 2834-2838, 1994.
- 446 Drake, H. L.: Acetogenesis, acetogenic bacteria, and the acetyl-CoA
447 "Wood/Ljungdahl" pathway: past and current perspectives, in: *Acetogenesis*,
448 edited by: Drake, H. L., Chapman & Hall, New York, 3-60, 1994.
- 449 Elsner, M., Zwank, L., Hunkeler, D., and Schwarzenbach, R. P.: A new concept
450 linking observable stable isotope fractionation to transformation pathways of
451 organic pollutants [review], *Environ. Sci. Technol.*, 39, 6896-6916, 2005.
- 452 Freude, C. and Blaser, M.: Carbon isotope fractionation during catabolism and
453 anabolism in acetogenic bacteria growing on different substrates, *Appl.*
454 *Environ. Microbiol.*, 82, 2728-2737, 2016.
- 455 Glombitza, C., Jaussi, M., Roy, H., Seidenkrantz, M. S., Lomstein, B. A., and
456 Joergensen, B. B.: Formate, acetate, and propionate as substrates for sulfate
457 reduction in sub-arctic sediments of Southwest Greenland, *Frontiers*
458 *Microbiol.*, 6, 846-doi: 10.3389/fmicb.2015.00846 , 2015.

459 Goevert, D. and Conrad, R.: Effect of substrate concentration on carbon isotope
460 fractionation during acetoclastic methanogenesis by *Methanosarcina barkeri*
461 and *M. acetivorans* and in rice field soil, *Appl. Environ. Microbiol.*, *75*, 2605-
462 2612, 2009.

463 Hausmann, B., Knorr, K. H., Schreck, K., Tringe, S. G., DelRio, T. G., Loy, A.,
464 and Pester, M.: Consortia of low-abundance bacteria drive sulfate reduction-
465 dependent degradation of fermentation products in peat soil microcosms,
466 *ISME J.*, *10*, 2365-2375, 2016.

467 Hayes, J. M.: Factors controlling ¹³C contents of sedimentary organic
468 compounds: principles and evidence, *Mar. Geol.*, *113*, 111-125, 1993.

469 Hunger, S., Schmidt, O., Hilgarth, M., Horn, M. A., Kolb, S., Conrad, R., and
470 Drake, H. L.: Competing formate- and carbon dioxide-utilizing prokaryotes in
471 an anoxic methane-emitting fen soil, *Appl. Environ. Microbiol.*, *77*, 3773-
472 3785, 2011.

473 Jain, S., Dietrich, H. M., Müller, V., and Basen, M.: Formate is required for
474 growth of the thermophilic acetogenic bacterium *Thermoanaerobacter kivui*
475 lacking hydrogen-dependent carbon dioxide reductase (HDCR), *Frontiers*
476 *Microbiol.*, *11*, 59-doi: 10.3389/fmicb.2020.00059, 2020.

477 Janssen, P. H. and Frenzel, P.: Inhibition of methanogenesis by methyl fluoride -
478 studies of pure and defined mixed cultures of anaerobic bacteria and archaea,
479 *Appl. Environ. Microbiol.*, *63*, 4552-4557, 1997.

480 Kanaparathi, D., Pommerenke, B., Casper, P., and Dumont, M. G.:
481 Chemolithotrophic nitrate-dependent Fe(II)-oxidizing nature of actinobacterial
482 subdivision lineage TM3, *ISME J.*, *7*, 1582-1594, 2013.

483 Kim, Y. J., Lee, H. S., Kim, E. S., Bae, S. S., Lim, J. K., Matsumi, R.,
484 Lebedinsky, A. V., Sokolova, T. G., Kozhevnikova, D. A., Cha, S. S., Kim, S.
485 J., Kwon, K. K., Imanaka, T., Atomi, H., Bonch-Osmolovskaya, E. A., Lee, J.
486 H., and Kang, S. G.: Formate-driven growth coupled with H₂ production,
487 *Nature*, *467*, 352-U137, 2010.

488 Kotsyurbenko, O. R., Nozhevnikova, A. N., Soloviova, T. I., and Zavarzin, G. A.:
489 Methanogenesis at low temperatures by microflora of tundra wetland soil, *Ant.*
490 *Leeuwenhoek*, *69*, 75-86, 1996.

491 Küsel, K. and Drake, H. L.: Microbial turnover of low molecular weight organic
492 acids during leaf litter decomposition, *Soil Biol. Biochem.*, *31*, 107-118, 1999.

493 Lemaire, O. N., Jespersen, M., and Wagner, T.: CO₂-fixation strategies in energy
494 extremophiles: What can we learn from acetogens?, *Frontiers Microbiol.*, *11*,
495 2020.

496 Liebner, S., Schwarzenbach, S. P., and Zeyer, J.: Methane emissions from an
497 alpine fen in central Switzerland, *Biogeochem.*, *109*, 287-299, 2012.

498 Liu, P. F., Klose, M., and Conrad, R.: Temperature effects on structure and
499 function of the methanogenic microbial communities in two paddy soils and
500 one desert soil, *Soil Biol. Biochem.*, *124*, 236-244, 2018.

501 Lovley, D. R. and Klug, M. J.: Intermediary metabolism of organic matter in the
502 sediments of a eutrophic lake, *Appl. Environ. Microbiol.*, *43*, 552-560, 1982.

503 Mariotti, A., Germon, J. C., Hubert, P., Kaiser, P., Letolle, R., Tardieux, A., and
504 Tardieux, P.: Experimental determination of nitrogen kinetic isotope
505 fractionation: some principles; illustration for the denitrification and
506 nitrification processes, *Plant and Soil*, *62*, 413-430, 1981.

507 Martins, M., Mourato, C., and Pereira, I. A.: *Desulfovibrio vulgaris* growth
508 coupled to formate-driven H₂ production, *Environ. Sci. Technol.*, 49, 14655-
509 14662, 2015.

510 Montag, D. and Schink, B.: Formate and hydrogen as electron shuttles in terminal
511 fermentations in an oligotrophic freshwater lake sediment, *Appl. Environ.*
512 *Microbiol.*, 84, e01572-18-[https://doi.org/10.1128/AEM .01572-18](https://doi.org/10.1128/AEM.01572-18), 2018.

513 Nozhevnikova, A. N., Kotsyurbenko, O. R. and Simankova, M. V.: Acetogenesis
514 at low temperature, in: *Acetogenesis*, edited by: Drake, H. L., Chapman &
515 Hall, New York, 416-431, 1994.

516 Penning, H., Claus, P., Casper, P., and Conrad, R.: Carbon isotope fractionation
517 during acetoclastic methanogenesis by *Methanosaeta concilii* in culture and a
518 lake sediment, *Appl. Environ. Microbiol.*, 72, 5648-5652, 2006.

519 Peters, V., Janssen, P. H., and Conrad, R.: Efficiency of hydrogen utilization
520 during unithrophic and mixotrophic growth of *Acetobacterium woodii* on
521 hydrogen and lactate in the chemostat, *FEMS Microbiol. Ecol.*, 26, 317-324,
522 1998.

523 Peters, V., Janssen, P. H., and Conrad, R.: Transient production of formate during
524 chemolithotrophic growth of anaerobic microorganisms on hydrogen, *Curr.*
525 *Microbiol.*, 38, 285-289, 1999.

526 Phelps, T. J. and Zeikus, J. G.: Effect of fall turnover on terminal carbon
527 metabolism in Lake Mendota sediments, *Appl. Environ. Microbiol.*, 50, 1285-
528 1291, 1985.

529 Rothfuss, F. and Conrad, R.: Vertical profiles of CH₄ concentrations, dissolved
530 substrates and processes involved in CH₄ production in a flooded Italian rice
531 field, *Biogeochem.*, 18, 137-152, 1993.

532 Schink, B., Montag, D., Keller, A., and Müller, N.: Hydrogen or formate:
533 Alternative key players in methanogenic degradation [review], *Environ.*
534 *Microbiol. Reports*, 9, 189-202, 2017.

535 Schuchmann, K., Chowdhury, N. P., and Müller, V.: Complex multimeric [FeFe]
536 hydrogenases: biochemistry, physiology and new opportunities for the
537 hydrogen economy [review], *Frontiers Microbiol.*, 9, 2911-[doi:](https://doi.org/10.3389/fmicb.2018.02911)
538 10.3389/fmicb.2018.02911, 2018.

539 Schuchmann, K. and Müller, V.: Direct and reversible hydrogenation of CO₂ to
540 formate by a bacterial carbon dioxide reductase, *Science*, 342, 1382-1385,
541 2013.

542 Sieber, J. R., Le, H. M., and McInerney, M. J.: The importance of hydrogen and
543 formate transfer for syntrophic fatty, aromatic and alicyclic metabolism,
544 *Environ. Microbiol.*, 16, 177-188, 2014.

545 Stumm, W. and Morgan, J. J.: *Aquatic Chemistry*, 3. ed. Wiley, New York, 1996.

546 Thauer, R. K., Jungermann, K., and Decker, K.: Energy conservation in
547 chemotrophic anaerobic bacteria, *Bacteriol. Rev.*, 41, 100-180, 1977.

548 Widdel, F.: Microbiology and ecology of sulfate- and sulfur-reducing bacteria, in:
549 *Biology of Anaerobic Microorganisms*, edited by: Zehnder, A. J. B., Wiley,
550 New York, 469-585, 1988.

551 Wüst, P. K., Horn, M. A., and Drake, H. L.: Trophic links between fermenters
552 and methanogens in a moderately acidic fen soil, *Environ. Microbiol.*, 11,
553 1395-1409, 2009.

554 Zinder, S. H.: Physiological ecology of methanogens, in: *Methanogenesis.*
555 *Ecology, Physiology, Biochemistry and Genetics*, edited by: Ferry, J. G.,
556 Chapman & Hall, New York, 128-206, 1993.

557

558

559 **Figure legends**

560

561 **Figure 1.** Formate conversion to acetate, CH₄ and CO₂ in suspensions of paddy soil from
562 Vercelli (Italy) after addition of formate without sulfate (blue squares) or formate plus sulfate
563 (gypsum) (red triangles) without CH₃F (open symbols) or with CH₃F (closed symbols).
564 Controls with addition of only water (blue or red X crosses) are only shown occasionally. The
565 panels show the temporal change of (a) concentrations of formate, (b) concentrations of
566 acetate, (c) mixing ratios of CH₄ (1 ppmv = 10⁻⁶ bar), (d) mixing ratios of CO₂, (e) δ¹³C of
567 formate, (f) δ¹³C of acetate, (g) δ¹³C of CH₄, and (h) δ¹³C of CO₂. Means ± SE.

568 **Figure 2.** Balance of produced acetate plus CH₄ (blue symbols) and of only CH₄ (red
569 symbols) against the consumed formate in (a) the absence and (b) the presence of sulfate in
570 paddy soil from Vercelli (Italy). Acetate and CH₄ are each equivalent to 4 H₂, formate to 1
571 H₂. The open and closed symbols denote conditions in the absence and the presence of CH₃F,
572 respectively. The different symbols indicate three different replicates. The line indicate
573 equimolarity (in terms of reducing equivalents between substrate and product).

574 **Figure 3.** Formate conversion to acetate, CH₄ and CO₂ in suspensions of sediment from the
575 NE basin of Lake Fuchskuhle after addition of formate without sulfate (blue squares) or
576 formate plus sulfate (gypsum) (red triangles) without CH₃F (open symbols) or with CH₃F
577 (closed symbols). Controls with addition of only water (blue or red X crosses) are only shown
578 occasionally. The panels show the temporal change of (a) concentrations of formate, (b)
579 concentrations of acetate, (c) mixing ratios of CH₄ (1 ppmv = 10⁻⁶ bar), (d) mixing ratios of
580 CO₂, (e) δ¹³C of formate, (f) δ¹³C of acetate, (g) δ¹³C of CH₄, and (h) δ¹³C of CO₂. Means ±
581 SE.

582 **Figure 4.** Balance of produced acetate plus CH₄ (blue symbols) and of only CH₄ (red
583 symbols) against the consumed formate in (a) the absence and (b) the presence of sulfate in
584 sediment from the NE basin of Lake Fuchskuhle. Acetate and CH₄ are each equivalent to 4
585 H₂, formate to 1 H₂. The open and closed symbols denote conditions in the absence and the
586 presence of CH₃F, respectively. The different symbols indicate three different replicates. The
587 line indicate equimolarity (in terms of reducing equivalents between substrate and product).

588 **Figure 5.** Mariotti plots of formate consumption in (a, b) paddy soil from Vercelli and (b, c,
589 d) sediment from the NE basin of Lake Fuchskuhle under methanogenic (a, c, blue symbols)
590 and sulfidogenic (b, d, red symbols) conditions both in the absence (open symbols) and in the
591 presence (closed symbols) of CH₃F. ~~The different symbols indicate three different~~
592 replicates.

593 **Figure 6.** Isotopic enrichment factors (ϵ_{form} , given as negative values) in paddy soils from
594 Vercelli and the IRRI (the Philippines) and in lake sediments from the NE and SW basins of

595 Lake Fuchskuhle without (left panel) and with (right panel) addition of sulfate (gypsum) and
596 CH₃F. Means ± SE.

597 **Figure 7.** Average δ¹³C of formate (at the beginning of incubation) and of CO₂, acetate and
598 CH₄ (after the depletion of formate) in paddy soils ~~or sediments~~ from Vercelli (blue); and the
599 IRRI (green), and in sediments from the NE basin (red) and the SW basin (yellow) of Lake
600 Fuchskuhle in the absence (filled bars) and the presence (dotted bars) of CH₃F. Means ± SE.
601