

# Hydrogen solubility of stishovite provides insights into water transportation to deep Earth

Mengdan Chen<sup>1</sup>, Changxin Yin<sup>1</sup>, Danling Chen<sup>1</sup>, Long Tian<sup>1</sup>, Liang Liu<sup>1</sup>, and Lei Kang<sup>1</sup>

<sup>1</sup>Department of Geology, Northwest University/State Key Laboratory of Continental Dynamics, Xi'an, 710069, China

5 Correspondence to: Lei Kang ([kanglei@nwu.edu.cn](mailto:kanglei@nwu.edu.cn)); Liang Liu ([liuliang@nwu.edu.cn](mailto:liuliang@nwu.edu.cn))

**Abstract.** Water dissolved in nominally anhydrous minerals (NAMs) can be transported to deep regions of the Earth through subducting slabs, thereby significantly influencing the physicochemical properties of deep Earth materials and impacting dynamic processes in the deep Earth. Stishovite, a prominent mineral ~~found-present~~ in subducting slabs, remains stable at mantle ~~conditions ranging from~~ pressures of 9-50 GPa and can incorporate ~~varying-various~~ amounts of water (H<sup>+</sup>, OH<sup>-</sup>, and H<sub>2</sub>O) ~~within~~ its crystal structure. Consequently, stishovite ~~can playplays~~ a crucial role in transporting water into deep Earth through subducting slabs. This paper provides a comprehensive review of the research progress concerning water (hydrogen) solubility in stishovite. The key factors that govern water solubility in stishovite are summarized as temperature, pressure, water fugacity and aluminum content. Combined with published results on the dependence of water solubility on the aforementioned parameters, this paper proposes a new equation ~~for water solubility to describe the solubility of water~~ in Al-bearing stishovite ~~can be described by the relationship~~. Calculation results based on this equation suggest that stishovite may effectively ~~absorb accommodate~~ water released from processes such as hydrous mineral breakdown, ~~dehydration reactions, and volume reduction, which could~~ ultimately contributeing to the presence of a water-rich transition zone.

## 1 Introduction

Water plays a crucial role not only in the origin and evolution of life but also in various Earth processes, including slab subduction, crust-mantle reaction and recycling. During slab subduction, water is transported from the Earth's surface to its interior through subduction zones, and a significant amount of water returns to the surface mainly through magmatism, thus forming a large-scale water cycle (Fig. 1). In subducting slabs, ~~water-hydrogen~~ exists ~~both explicitly and implicitly. Explicit water components include H<sub>2</sub>O, OH<sup>-</sup> and H<sup>+</sup>~~ in melts, fluids, fluid inclusions, and hydrous minerals (e.g., such as amphibole and mica) ~~and nominally anhydrous minerals (NAMs)~~ (Bell et al., 1995; Rossman, 1996; Xia et al., 1998; Johnson, 2006; Libowitzky, 2006; Litasov and Ohtani, 2007; Ni et al., 2017), ~~in forms of both molecular water (H<sub>2</sub>O) and structurally bound water (H<sub>3</sub>O<sup>+</sup>, H<sup>+</sup> and OH<sup>-</sup>).~~ ~~Implicit water primarily occurs as hydroxyl (OH) in nominally anhydrous minerals (NAMs) at defects in crystal structures.~~ Most hydrous minerals in subducting slabs decompose and release water before reaching a depth of 300 km (Poli and Schmidt, 2002). Part of this water returns to the surface through magmatism, while the remainder is

incorporated into NAMs in ultrahigh-pressure metamorphic rocks (Magni et al., 2014; Walter, 2021) and subsequently transported to deep mantle by subducting slabs (Ishii et al., 2022).




Figure 1: Water transport, distribution, and cycling in the Earth (after Litasov and Ohtani, 2007)

Numerous studies indicate that major constituent minerals in deep mantle, such as olivine, pyroxene, and garnet, as well as their high-pressure polymorphs, are NAMs (Litasov and Ohtani, 2007; Liu et al., 2016). Although the phase of  $\text{SiO}_2$  does not typically appear in the mantle (Kaminsky, 2012), it is a significant constituent mineral in silica-riched slabs and can stably exist at upper to lower mantle depths through high-pressure polymorphic phase transitions (Fig. 2). Both experimental and computational studies have demonstrated that stishovite's crystal structure can incorporate a certain amount of water. Given its prevalence as a major mineral in subducting slabs at mantle depths (>300 km), stishovite likely plays a significant role in water transportation to deep earth (Spektor et al., 2011; Lin et al., 2022; Ishii et al., 2022). In this paper, we provide a systematic review of previous understandings and research progress regarding water solubility, water incorporation mechanisms, and factors influencing water solubility in stishovite. Considered stishovite in subducting slabs always contains amounts of Al in its crystal structure (Ono, 1999; Lakshtanov et al., 2007a). Combined with published results, this paper establishes an empirical model for water solubility in Al-stishovite based on published results, and discuss the significance of stishovite in transporting water from subducting slabs to the deep mantle and its implications for deep Earth dynamics. Finally, we highlight key unresolved scientific questions in the research on water solubility in stishovite.

## 2 Crystal structure and stability of stishovite

Phase of  $\text{SiO}_2$  undergoes multiple high-pressure polymorphic transitions as pressure increases (Petersen et al., 2021) (Fig. 3). Under a subduction slab average geothermal gradient, coesite is the dominant phase of  $\text{SiO}_2$  at pressures ranging from 2.7 to 9 GPa. At approximately 10 GPa, coesite transforms into the denser stishovite (Ono et al., 2017). The transition pressure from coesite to stishovite is dependent on temperature, as described by the equation  $P \text{ (GPa)} = 4.7 + 0.0032 \times T \text{ (K)}$  (Ono et al.,

2017). This phase transition is believed to contribute to the seismic discontinuity observed at around 300 km depth (Ono et al., 2017). As pressure increases further to 23–50–24 GPa, stishovite undergoes a transformation into the CaCl<sub>2</sub>-type SiO<sub>2</sub> structure (commonly referred to as post-stishovite) due to the incorporation of Al and H in stishovite (Ishii et al., 2022), commonly referred to as post-stishovite. However, the transition pressure is still debated. Fischer et al. (2018) pointed out that the minimum transition pressure could be as high as 70 GPa in Earth; the phase transition between stishovite and CaCl<sub>2</sub>-type silica should occur at pressures of 68–78 GPa in the Earth, depending on the temperature in subducting slabs.

Formatted: Font: 10 pt

Field Code Changed





Figure 2: Mantle Mineral assemblages in peridotite, subducted oceanic crust, and subducted continental crust. The red line indicates the mantle geotherm (Kaminsky, 2012). Modified after Smith et al., 2018 and Wu et al., 2009. (Fp—Ferropericlas; Opx—Orthopyroxene; Cpx—Clinopyroxene; Ca-pv—Ca-perovskite; NAL—Na-Al phase; Ep—Epidote; Law—lawsonite; K-Holl—K-Hollandite)

Previous studies show that stishovite constitutes 10 vol. % and 20 vol. % of subducting mid-ocean ridge basalt (MORB) in the upper and lower mantle, respectively (Irifune et al., 1986; Ono et al., 2001). In subducted continental crust at upper mantle depths (<660 km), stishovite can reach approximately 20–25 vol. % (Irifune et al., 1994; Poli and Schmidt, 2002; Ishii et al., 2012). However, due to the exhumation of subducted slabs from ~300 km depth (termed the referred as "depth of no return" in literatures, e.g., Irifune et al., 1994; Liu et al., 2007; Wu et al., 2009) to the surface is extremely difficult, and/or the preservation conditions of stishovite are very harsh, stishovite is unstable and easily transforms to lower pressure SiO<sub>2</sub> polymorphs, naturally formed stishovite that can be observed is extremely rare. Previously, Yang et al. (2007) identified polycrystalline coesite as a potential pseudomorphic replacement of stishovite in Tibetan chromitites. The micro-exsolution microstructure (Liu et al., 2007) and pseudomorphs after stishovite (Liu et al., 2018) were only found in the South Altyn ultra-high pressure metamorphic belt in western China. Recently, Thomas et al. (2022) identified coesite and stishovite inclusions in the Waldheim granulite, and Gu et al. (2022) showed that coesite (former stishovite) was present within a natural super-

deep diamond formed at the boundary between the transition zone and the lower mantle. In addition, other naturally occurring stishovite is found sporadically in meteorite impact craters (e.g., Chao et al., 1962) or meteorites (e.g., Holtstam et al., 2003). Therefore, the current research on our current understanding of stishovite mainly relies on high-temperature and high-pressure experiments and theoretical calculations (e.g., Litasov et al., 2007; Lin et al., 2022; Ishii et al., 2022).

80 Stishovite was first synthesized by Stishov and Popova (1961) at 20 GPa (equivalent to approximately 600 km depth in Earth's interior) and 1100 °C from  $\alpha$ -quartz. It possesses a rutile-type structure with tetragonal symmetry ( $P4_2/mmm$ ). The Si atoms are coordinated by six O atoms in octahedral arrangements (Pawley et al., 1993; Spektor et al., 2011; Lin et al., 2022). These  $\text{SiO}_6$  octahedra align and form linear chains along the c-axis. This arrangement results in a highly dense packing of O atoms, with slightly elongated Si-O bonds compared to  $\text{SiO}_4$  tetrahedra. Previous studies have demonstrated that the density of stishovite is 46% higher than coesite and 60% higher than  $\alpha$ -quartz, respectively (Keskar et al., 1991). Therefore, the formation of stishovite during slab subduction at depths exceeding 300 km can dramatically increase the density of subducting slabs (Lin et al., 2022; Ishii et al., 2022), and enhancing enhance their negative buoyancy for further subduction into the mantle transition zone (410-660 km) or even deeper regions.

Formatted: Font: Italic






Figure 3: SiO<sub>2</sub> phase diagram as recommended in the reference literature (density values are indicated in brackets) (Gutzow et al., 2014). The gray dashed line represents the subduction slab geothermal gradient (Zheng et al., 2016). The diagram also illustrates the SiO<sub>4</sub> tetrahedron and SiO<sub>6</sub> octahedron, along with their corresponding distances.

### 3 Water solubility and incorporation mechanisms in stishovite

#### 3.1 Water solubility

Research on water solubility in stishovite has primarily been conducted through high-pressure experiments (Chung and Kagi, 2002; Bromiley et al., 2006; Litasov et al., 2007; Spektor et al., 2011; Nisr et al., 2017; Lin et al., 2022; Ishii et al., 2022). Fourier transform infrared spectroscopy (FTIR) is the primary technique used to determine water content in experimental products, although results from different studies often show significant discrepancies, as detailed in Yan and Liu (2021). In this paper, we compile the experimental conditions (temperature, pressure, initial water content) and results (water and Al<sub>2</sub>O<sub>3</sub> contents in stishovite products) from previous studies (Table 1).

Table 1 Water contents in stishovite from previous studies

Temperature (°C)	Pressure (GPa)	Initial Water content (wt. %)	Synthesis method	Al <sub>2</sub> O <sub>3</sub> (wt. %)	Water content (wt. ppm)	Measurement method	From References
1200	10	saturated	LVP	0-1.51	7-82	FTIR <sup>a</sup>	Pawley et al., 1993
1200-1500	15-21	saturated	LVP	0	2.5-72	FTIR <sup>b</sup>	Bolfan-

1200-1400	10-15	saturated	LVP	0.612-1.341	210-759	FTIR <sup>b</sup>	Casanova et al., 2000
1740-3700	30-62.9	0.2	DAC	unspecified	76-487	FTIR <sup>a</sup>	Chung and Kagi, 2002
1227	25	saturated	unspecified	unspecified	3000*	DFT	Panero et al. 2003
1500	15	saturated	LVP	0-2.95	3-456	FTIR <sup>a</sup>	Panero and Stixrude, 2004
1800	20-25	saturated	LVP	0-7.62	25-3010	FTIR <sup>b</sup>	Bromiley et al., 2006
800-1240	8-12.3	1-4	LVP	trace	53-187	FTIR <sup>c</sup>	Litasov et al., 2007
450-550	10	saturated	LVP	<60	0.9-1.75 wt. % <sup>‡</sup>	TGA/SIMS	Thomas et al., 2009
627-1654	12	0-1.73	LVP	0-2.24	108-1354	FTIR <sup>b</sup>	Spektor et al., 2011
350-550	10	unspecified=	LVP	0=	0.5-3 wt. % <sup>‡</sup>	TGA	Yoshino et al., 2014
450	9	unspecified=	LVP	unspecified 0	3.2 wt. % <sup>‡</sup>	LP	Spektor et al., 2016
950	8.4-9.1	1.5-6=	LVP	unspecified	246-376	FTIR <sup>c</sup>	Nisir et al., 2017
967-1562	32.5-52	unspecified=	DAC	0	4.6-9.9 wt. % <sup>‡</sup>	From Nisir et al., 2017	Frigo et al., 2019
1700	20	saturated	LVP	3.43-5.37	2500-2700	FTIR <sup>b</sup>	Lin et al., 2020
1700	28	saturated <sup>5</sup>	LVP	4.36	2700	FTIR <sup>b</sup>	Zhang et al., 2022
973-1592	53-72.5	0.5-15.2	DAC	0	0.62-3.61 wt. % <sup>‡</sup>	DFT	Ishii et al., 2022
450	9	8	LVP	0	1.57-2.25 wt. % <sup>‡</sup>	Mass balance/LOI average	Lin et al., 2022
1627	23	saturated	LVP	0	0.43-0.62 wt. %	nano-SIMS/ NMR	Kueter et al., 2023
							Li et al., 2023

Note: DAC, Diamond anvil cell; LVP, large volume press; TGA, Thermogravimetric analysis; SIMS, Secondary ion mass spectrometry. LP, water content determined from lattice parameters, DFT, density functional theory calculation, nano-SIMS/NMR, nanoscale secondary ion mass spectroscopy and solid-state nuclear magnetic resonance.

The calculation methods of water contents based on FTIR are from <sup>a</sup>: Pawley et al. (1993); <sup>b</sup>: Paterson (1982); and <sup>c</sup>: Libowitzky and Rossman (1997).

Most studies used Fourier transform infrared spectroscopy (FTIR) to determine water contents in stishovite with the calculation method from a. Pawley et al. (1993); b. Paterson (1982); and c. Libowitzky and Rossman (1997). \* represents other methods.

As shown in Table 1, water solubility in stishovite is significantly influenced by temperature, pressure, and Al content, with large variations more than one order of magnitude from a few wt. ppm to a few wt. percent. However, even under specific pressure conditions, water contents obtained by different studies researchers can differ by more than one order of magnitude. For example, at 20-25 GPa, some studies obtained 25-2700 wt. ppm water in stishovite (Litasov et al., 2007; Zhang et al., 2022b), while others reported 0.5-3 wt. % water (Spektor et al., 2011; 2016). These discrepancies may partially be attributed

to different water content measurement methods. In Table 1, Most FTIR studies obtained water contents at the wt. ppm level, while other methods yielded water contents at the wt. % level. For example, Nisr et al. (2017, 2020) suggested a calibration based on unit cell volume dependency on water in stishovite from DFT calculations. Lin et al. (2020) applied Nisr et al. (2017)'s formulation to calculate H<sub>2</sub>O contents in their high-pressure samples resulting in exceptionally high H<sub>2</sub>O concentrations exceeding 10 wt. % in many instances. However, Lin et al. (2022) noted that estimating H<sub>2</sub>O contents using unit cell volumes from samples synthesized in either the LVP or DAC is highly uncertain.

Additionally, the discrepancies may relate to variations in experimental conditions of temperature, and Al content in stishovite, as detailed in Sect. 4.

### 3.2 Hydrogen incorporation mechanisms

Previous studies indicate two primary mechanisms for water incorporation in stishovite: (1) "hydrogarnet" substitution by  $4\text{H}^+ \rightarrow \text{Si}^{4+}$ ; and (2) coupled substitution of  $\text{H}^+$  and  $\text{Al}^{3+}$  for  $\text{Si}^{4+}$  ( $\text{Al}^{3+} + \text{H}^+ \rightarrow \text{Si}^{4+}$ ). Hydrogarnet substitution is a common mechanism in pure SiO<sub>2</sub>-stishovite. Studies by Litasov et al. (2007) have shown that stishovite can dissolve up to 5 wt. % Al<sub>2</sub>O<sub>3</sub>, which can further increase to 9 wt. % with the presence of water (Ono, 1999). Numerous high-pressure experiments demonstrate that Al can directly couple with H by substituting Si to increase water solubility in Al-bearing stishovite, with the coupled substitution mechanism of  $\text{Al}^{3+} + \text{H}^+ \rightarrow \text{Si}^{4+}$  (Pawley et al., 1993; Chung and Kagi, 2002; Bromiley et al., 2006; Lakshatnov et al., 2007a). In addition to Al<sup>3+</sup>, stishovite contains minor amounts of other trivalent cations such as B<sup>3+</sup>, Fe<sup>3+</sup>, V<sup>3+</sup>, and Cr<sup>3+</sup>, which can also facilitate H incorporation into the stishovite structure by similar mechanisms as Al<sup>3+</sup> (Irifune and Ringwood, 1993; Pawley et al., 1993). However, Litasov et al. (2007) found that H<sup>+</sup> (OH) can only co-replace Si<sup>4+</sup> with up to 40% Al<sup>3+</sup>, which makes the Al/H ratio in stishovite far greater than 1/1, that is, Al in stishovite is much higher than H. Bromiley et al. (2006) also reported excess Al in stishovite, not charge balanced by hydrogen. Thus, the incorporation of aluminum cannot be explained solely by the charge-coupled substitution with protons in stishovite and other substitution mechanisms have also been proposed. For instance, At this point, most Al will occupy the oxygen vacancy (O<sub>v</sub>) and balance the charge by  $2\text{Al}^{3+} + \text{O}_v^{2-} \rightarrow 2\text{Si}^{4+}$  (Pawley et al., 1993; Chung and Kagi, 2002; Litasov et al., 2007; Zhang et al., 2022b).

Recently, molecular water (H<sub>2</sub>O) has been identified in stishovite (Lin et al., 2020, 2022; Kueter et al., 2023). The incorporation mechanism is summarized as interstitial H<sub>2</sub>O substitution (Lin et al., 2022). Li et al. (2023) presented a new mechanism (one-dimensional (1D) water channels) for molecular water incorporation into stishovite using high-dimensional neural network (HDNN) potential. Both mechanisms for incorporation of molecular water could explain the weight percent level water observed in Al-free stishovite. Notably, the initial material in these studies is pure SiO<sub>2</sub> which means H is impossible coupled with Al to substitute Si. Therefore, even though multiple substitution mechanisms exist for H incorporation in stishovite, the dominant mechanisms in the natural environment remain highly uncertain (Lin et al., 2022). Due to stishovite in subducting slabs contains Al as mentioned above (e.g., Litasov et al., 2007), we emphasize that  $\text{Al}^{3+} + \text{H}^+ \rightarrow \text{Si}^{4+}$  is still very important (e.g., Pawley et al., 1993; Chung and Kagi, 2002; Lakshatnov et al., 2007a; Zhang et al., 2022b; Ishii et al., 2022).

In addition to  $\text{Al}^{3+}$ , stishovite contains minor amounts of other trivalent cations such as  $\text{B}^{3+}$ ,  $\text{Fe}^{3+}$ ,  $\text{V}^{3+}$ , and  $\text{Cr}^{3+}$ , which can also facilitate H incorporation into the stishovite structure by similar mechanisms as  $\text{Al}^{3+}$  (Trifunac and Ringwood, 1993; Pawley et al., 1993). In summary, multiple mechanisms exist for H incorporation in stishovite with  $\text{Al}^{3+} + \text{H}^+ \rightarrow \text{Si}^{4+}$  being the most important.

#### 4 Factors Dominating Water Solubility in Stishovite

Numerous studies demonstrate that water solubility in stishovite is primarily controlled by temperature, pressure, and Al content.

##### 4.1 Temperature and Pressure


Numerous experiments have constrained the water solubility in Al-bearing stishovite as a function of temperature and pressure. Panero et al. (2003) showed that at 30-38 GPa, water contents in stishovite significantly increase with increasing temperature from ~100 wt. ppm at 1500 K to 400 wt. ppm at 3500 K. Litasov et al. (2007) measured the hydrogen contents of stishovite samples synthesized at 20-25 GPa and 1200-1800 °C from several starting materials containing up to 10 wt. %  $\text{Al}_2\text{O}_3$  and found that at 20-25 GPa and 1000-1200 °C,  $\text{H}_2\text{O}$  contents are positively correlated with temperature between 1000 and 1200 °C, but decrease with increasing temperature from at 1200-1800 °C, likely due to increased melt proportions. This can be simply explained by the fact that water-hydrogen, as an incompatible element, preferably dissolves into the melt (Smyth, 2006; Litasov et al., 2007). Therefore, the occurrence of melt at high-temperature temperature can significantly reduce  $\text{H}_2\text{O}$  contents in stishovite. Panero et al. (2003) showed that at 10-60 GPa, water solubility in stishovite increases with increasing pressure. Panero and Stixrude (2004) suggested that the water solubility in stishovite is ~0.3 wt. %  $\text{H}_2\text{O}$  at 25 GPa and 1500 K, and increases to 1.15 wt. %  $\text{H}_2\text{O}$  at 60 GPa, indicating increasing solubility with pressure.

By comparing previous studies, it is evident that water solubility in Al-bearing stishovite shows an overall weak positive correlation with temperature but a significant decrease when the melt is present at >1600 °C. Whereas, the pressure dependence is more complex, with a positive correlation below 20 GPa but a negative correlation above 20 GPa, suggesting an optimal water solubility in stishovite at ~20 GPa (Fig. 4). In addition, Lakshtanov et al. (2007b) suggested that stishovite undergoes a transition to  $\text{CaCl}_2$ -structured phase (post-stishovite) at ~30 GPa. Weight percent levels of water (0.85 to 1.1 wt. %) in  $\text{CaCl}_2$ -structured Al-rich post-stishovite at 24 to 28 GPa and 1000 to 2000 °C have been reported by Ishii et al. (2022). This implies stishovite can retain its water content through phase transition with increasing pressure. However, Fischer et al. (2018) suggested the transition to post-stishovite occurs at pressures of 68–78 GPa in the Earth. Therefore, if the post-stishovite can accommodate water at ~20 GPa where the Al-bearing stishovite has a peak water solubility remains unclear.



## 4.2 Al Content


Given that natural systems always contain some Al, its effect on water solubility in stishovite has been a focus of research. Pawley et al. (1993) found that for stishovite with >1 wt. %  $\text{Al}_2\text{O}_3$ , the  $\text{H}_2\text{O}$  content (82 wt. ppm) is ten times higher than Al-free stishovite (7 wt. ppm). Chung and Kagi (2002) showed that at 15 GPa and 1400 °C, with increasing trivalent cations (mostly Al) from 0.612 wt. % to 1.341 wt. %,  $\text{H}_2\text{O}$  contents in stishovite increase from within the range of 128 wt. ppm to 536 wt. ppm. Litasov et al. (2007) obtained the maximum 3010 wt. ppm  $\text{H}_2\text{O}$  in Al-bearing stishovite (4.4 wt. %  $\text{Al}_2\text{O}_3$ ) versus 16-30 wt. ppm  $\text{H}_2\text{O}$  in Al-free stishovite at 20 GPa-GPa and 1400 °C. Recent work by Ishii et al. (2022) also demonstrated the positive correlation between water solubility and  $\text{Al}_2\text{O}_3$  content in stishovite. Therefore, increasing Al significantly enhances water solubility in stishovite. Statistically compiling previous data demonstrates athe positive correlation between Al and water contents in stishovite (Fig. 5).



**Figure 4: Previous studies on water solubility in Al-bearing stishovite. The dimension of circles represents the Al content. (a) Water content versus temperature; (b) Water content versus pressure (Only FTIR data are presented (Data from Pawley et al., 1993; Chung and Kagi, 2002; Bromiley et al., 2006; Litasov et al., 2007; Ishii et al., 2022; Zhang et al., 2022b). The relation of water solubility and Al content is shown in Fig. 5.**


In contrast, Lin et al. (2022) showed decreasing water solubility in pure stishovite with increasing temperature and pressure, markedly different from Al-bearing systems. Also, Lin et al. (2022)'s experimental data indicates a  $\text{H}_2\text{O}$  storage capacity in Al-free stishovite of ~3.5 wt. % at ~50 GPa and 1800 K. Kueter et al. (2023) reported the Al-free stishovite contains on average 1.69 wt. % water at 9 GPa and 450 °C. However, previous LVP studies synthesized Al-free (or Al-poor at the wt. ppm level) stishovite under water-saturated conditions is nearly anhydrous and commonly contains less than 100 wt. ppm  $\text{H}_2\text{O}$ , which is significantly different from wt. %  $\text{H}_2\text{O}$  in recent studies (e.g., Spektor et al., 2011, 2016; Lin et al., 2022; Kueter et al., 2023). It is challenging to understand how different studies can produce such widely disparate results, with differences of

200 several orders of magnitude for the H<sub>2</sub>O capacity of stishovite (Lin et al., 2022). The reason for the difference is unclear and there are various possible explanations. One possible cause is water loss from the capsule (e.g., Litasov et al., 2007). Moreover, the large discrepancy of water solubility between Al-stishovite (wt. ppm level) and Al-free stishovite (wt. % level from recent studies) has partly been explained by a hydrogarnet substitution mechanism ( $\text{Si}^{4+} \leftrightarrow 4\text{H}^+$ ) and/or the incorporation of interstitial molecular water (Kueter et al., 2023). However, the discrepancy caused by different measurement methods should be taken  
205 into consideration as well. Statistically compiling previous data demonstrates the positive correlation between Al and water contents in stishovite (Fig. 5).



210 **Figure 4: Previous experimental studies on water solubility in Al-bearing stishovite. The dimension of circles represents the Al content. (a) Water content versus temperature; (b) Water content versus pressure (Only FTIR data are presented included) (Data from Pawley et al., 1993; Bolfan-Casanova, 2000; Chung and Kagi, 2002; Panero et al., 2003; Bromiley et al., 2006; Litasov et al., 2007; Frigo et al., 2019; Ishii et al., 2022; Zhang et al., 2022b). The relation of water solubility and Al content is shown in Fig. 5.**

215 However, It should be noted that the behavior of Al itself also depends on pressure and temperature. Al can remarkably increase the hydrogen solubility of stishovite (Fig. 5), hydrogen can also increase Al solubility of stishovite in return. However, previous studies mainly focus on the water solubility of Al-free or Al-saturated stishovite. Studies on the P-T dependence of Al solubility in stishovite are very limited (e.g., Liu et al., 2006; Litasov et al., 2007). Liu et al. (2006) showed investigated that Al solubility in dry stishovite decreases with pressure but increases with temperature at <2000 °C in anhydrous experiments at 15-25 GPa and 1350 °C-2150 °C. Liu et al. (2006) found that Al<sub>2</sub>O<sub>3</sub> solubility in dry stishovite is slightly but consistently reduced by pressure increase, however, its response to temperature increase, is more complicated: increases at low temperatures, maximizes at around 2000 °C, and perhaps decreases at higher temperatures. Therefore, quantitatively  
220 constraining water solubility in natural stishovite requires considering the effects of variable Al solubility under different P-T conditions. Additionally, Al expands the unit-cell volume of stishovite (Litasov et al., 2007), enhancing the diffusion rate and facilitating water loss. In summary, the role of Al and other impurities in the water content and stability of hydrous stishovite remains poorly understood (Kueter et al., 2023).



225 Figure 5: The correlation between H<sub>2</sub>O and Al<sub>2</sub>O<sub>3</sub> in stishovite. Data are from Chung and Kagi (2002), Bromiley et al. (2006), Litasov et al. (2007), Ishii et al. (2022), and Zhang et al. (2022b), respectively. In Al-free stishovite, water contents range from 0.0003-0.0108 wt. % (3-108 wt. ppm). In Al-bearing stishovite, water contents increase to 0.0082-0.301 wt. % (82-3010 wt. ppm) with increasing Al.

### 230 4.3 Other Factors

Apart from aluminum (Al), other impurity ions likely play a role in affecting water solubility in stishovite. Several studies have suggested (Ono et al., 2002; Chung and Kagi, 2002; Panero et al., 2003; Panero and Stixrude, 2004) that stishovite formed from subducted MORB (~~Mid-Ocean Ridge Basalt~~) might incorporate a variety of trivalent cations, such as Al<sup>3+</sup>, Cr<sup>3+</sup>, Fe<sup>3+</sup>, ~~and V<sup>3+</sup> and Ti<sup>3+</sup>~~. These cations could potentially increase water content through coupled substitution. However, the work of

235 Litasov et al. (2007) ~~demonstrated-reported a that a small discrepancy in~~ the H<sub>2</sub>O content ~~is similar~~ between Fe-bearing stishovite (likely containing Fe<sup>3+</sup>) and Al-free and Fe-free stishovite. Consequently, the implications of Fe<sup>3+</sup>, Ti<sup>3+</sup>, and other elements on water solubility in stishovite need further investigation.




Figure 5: The correlation between  $H_2O$  and  $Al_2O_3$  in stishovite. Data are from Chung and Kagi (2002), Bromiley et al. (2006), Litasov et al. (2007), Ishii et al. (2022), and Zhang et al. (2022), respectively. In Al-free stishovite, water contents range from 0.0003–0.0108 wt. % (3–108 wt. ppm). In Al-bearing stishovite, water contents increase to 0.0082–0.301 wt. % (82–3010 wt. ppm) with increasing Al.

In addition, oxygen fugacity ( $f_{O_2}$ ) stands as a crucial thermodynamic parameter that describes the oxidation state within-in the deep Earth. It has a strong impact on the water solubility of those NAMs containing valency elements (e.g., Fe and Ti). Its impact on water solubility within NAMs is considerable. For instance, the water solubility in olivine and garnet displays positive and negative correlations with  $f_{O_2}$ , respectively (Yang, 2016; Zhang et al., 2022a). Since a small amount of Fe and Ti determined in a pseudomorph of former stishovite from a continental subduction slab (Liu et al., 2018),  $f_{O_2}$  could theoretically affect the water solubility of stishovite. However, the specific influence of  $f_{O_2}$  dependence on stishovite's water solubility of stishovite remains unknown-to-be-accurately-determined and requires further investigation to provide clear insights.

### 5 Water solubility in stishovite as a function of pressure, temperature, water fugacity, and Al content

Numerous studies have revealed that the water content in NAMs basically follows a thermodynamic relationship with temperature, pressure and water fugacity as shown in Eq. (1) (e.g., Kohlstedt et al., 1996; Zhao et al., 2004; Karato, 2010):

$$C_{OH} = A \cdot f_{H_2O}^n \cdot \exp\left(-\frac{\Delta E + \Delta V \cdot P}{R \cdot T}\right), \quad (1)$$

where  $C_{OH}$  is the water content,  $A$  is a constant,  $f_{H_2O}^n$  is the water fugacity, the value of  $n$  depends on the hydrogen incorporation mechanism, and  $\Delta E$  and  $\Delta V$  denote the energy and volume changes associated with hydrogen dissolution.

Lin et al. (2022) suggested a relationship for water solubility in Al-free stishovite as a function of pressure and temperature. Considering the stishovite in a subducted slab contains Al up to 5% (Litasov et al., 2007), and the notable-remarkable influence of aluminum (Al) on hydrogen solubility in-of stishovite ( $\text{Al}^{3+} + \text{H}^+ \rightarrow \text{Si}^{4+}$ ), therefore, we here propose a new model for water solubility in Al-bearing stishovite based-on-as Eq. (2):

$$C_{\text{OH}}^{\text{St}} = A \cdot f_{\text{H}_2\text{O}}^n \cdot \exp\left(-\frac{\Delta E + \Delta V \cdot P}{R \cdot T}\right) \cdot \exp\left(\frac{B \cdot X_{\text{Al}}}{R \cdot T}\right) \quad (2)$$

In Eq. (2),  $C_{\text{OH}}^{\text{St}}$  is water solubility in Al-bearing stishovite,  $B$  is a constant, and  $X_{\text{Al}}$  is the molar fraction of Al in stishovite. Given that the dominant hydrogen incorporation mechanism corresponds to  $\text{Al}^{3+} + \text{H}^+ \rightarrow \text{Si}^{4+}$  (Pawley et al., 1993), yielding

$C_{\text{OH}}^{\text{St}} \propto f_{\text{H}_2\text{O}}^{0.5}$ , thus  $n = 0.5$  (Kohlstedt et al., 1996; Karato, 2010).

To eliminate large disparities arising from diverse analytical methods as discussed above, we excluded the published water solubility in Al-bearing stishovite obtained by methods such as TGA/SIMS and only included collected published FTIR data on  $\text{H}_2\text{O}$  solubility contents in Al-bearing stishovite from FTIR measurements (Chung and Kagi, 2002; Litasov et al., 2007; Ishii et al., 2022; Zhang et al., 2022b, as shown in Table 1). We used a non-linear least square fitting method to fit Eq. (2) for fitting Eq. (2). The and obtained parameters are as follows:  $n = 0.5$ ,  $A = 0.24 \pm 0.13 \text{ ppm/GPa}^{0.5}$ ,  $\Delta E = -3.06 \pm 0.88 \text{ kJ/mol}$ ,  $\Delta V = 4.29 \pm 0.27 \text{ cm}^3/\text{mol}$ ,  $B = 7.69 \pm 1.12 \text{ kJ/mol}$ . The uncertainty is one standard deviation.  $n = 0.5$ ,  $A = 0.239 \text{ ppm/bar}^{0.5}$ ,  $\Delta E = -3.065 \text{ kJ/mol}$ ,  $\Delta V = 4.29 \text{ cm}^3/\text{mol}$ ,  $B = 7.69 \text{ kJ/mol}$ . As shown in Fig 6, a very good correlation can be found between the experimental data and the calculated results from Eq. (2) (Fig. A1). Due to the data we used to fit Eq. (2) are from experiments carried out at wide conditions of 10-28 GPa and 1200-1800 °C (close to the condition of MTZ), consequently, therefore, Eq. (2) can now reasonably be used to estimate the water solubility in Al-bearing stishovite across various-temperature, pressure, and  $f_{\text{H}_2\text{O}}$  conditions of MTZ.

We here-conducted calculations to determine the water solubility in stishovite containing 3 mol. % aluminum at temperatures ranging from 800 °C to 1400-1600 °C and pressures from 10 GPa to 40 GPa (Fig. 6). The results indicate that up to 30 GPa, water content experiences a marginal decline as temperature rises, while beyond 30 GPa, a positive correlation emerges between water content and temperature. This could be attributed to a lowered solidus under reduced pressures, leading to diminished water solubility as high-temperature, water-rich melts form. Conversely, at pressures surpassing 30 GPa, the heightened solidus counteracts the melt effect. Moreover, the solubility of water experiences a marked increase at pressures below 22 GPa to 32 GPa, followed by a decrease at pressures beyond this range, signifying an optimal solubility window. This aligns well with the overarching experimental data trend (Fig. 6b).

## 6 Implications for water transport to the deep Earth

Subduction zones serve as the exclusive pathway for surface water to infiltrate the Earth's deep interior. The transport and quantities of water within subducting slabs have profound impacts on surface environments, deep Earth properties and dynamics (Shillington, 2018). Existing experimental and geophysical data suggest the mantle transition zone distributes

290 potentially water-rich regions with >1 wt. % H<sub>2</sub>O (Pearson et al., 2014). However, the origins of this water remain a subject  
of debate. One perspective posits that the deep mantle has retained intrinsic wetness since Earth's formation. Alternatively,  
NAMs in subducting slabs may carry ~~a significant an~~ amount of water to ~~the~~ deep mantle (Peslier et al., 2017). However, it is  
essential to note that most hydrous minerals within subducting slabs will break down at various depths, typically having  
stability limits below 9 GPa, which limits the transport of water to greater mantle depths (>300 km) (Zheng et al., 2016). For  
instance, minerals like antigorite/phlogopite and lawsonite, which can survive to maximum depths in cold slabs, tend to  
295 dehydrate at around 300 km (Poli and Schmidt, 2002; [Walter, 2021](#)). Therefore, a fundamental question arises: can the released  
water from these hydrous minerals be further carried to even greater depths by NAMs?




Figure 6: Water solubility in stishovite versus (a) temperature (~~1000-1400°C~~) and (b) pressure (~~10-40 GPa~~)

300 Existing research demonstrates that water storage capacities within upper mantle NAMs generally increase with depth (Yang and Li, 2016). Nevertheless, it is imperative to investigate how phase transitions and breakdown reactions of NAMs influence

305 the transport of water in subducting slabs at greater depths. For instance, Gavrilenko (2008) showed that water solubility in clinopyroxene (Cpx) decreases along a mantle geotherm (Litasov and Ohtani, 2007)(Kaminsky, 2012) from 585-517 wt. ppm at 10 GPa to 135-176 wt. ppm at 16 GPa. Given the concurrent decomposition of Cpx from approximately 65 vol. % at 9 GPa to 0 vol. % at 16 GPa, an estimated 322-310 wt. ppm of water would be released from Cpx within the subducted oceanic slab crust.

We further conducted calculations to assess the evolution of water storage capacities in major minerals within subducted Mid-Ocean Ridge Basalt (MORB) and continental sediments along a representative cold subduction slab geotherm suggested by Litasov and Ohtani (2007). The results indicate that garnet and stishovite serve as primary carriers of water, with higher water storage capacities observed in subducted oceanic crust compared to sediments (Fig. 7). In subducted MORB and continental crusts-sediments down to mantle depths, stishovite comprises 10-20 vol. % (Irifune et al., 1986; Ono et al., 2001) and >20 vol. % (Ishii et al., 2019), respectively. Our model demonstrates that stishovite water solubility increases from 642 998 wt. ppm at 14 GPa to 906-1317 wt. ppm at 22 GPa along the cold subduction slab geotherm (Fig. 7). Consequently, 315 stishovite may theoretically accommodate water released from the breakdown of hydrous minerals (Lin et al., 2020; Nisr et al., 2020; Walter, 2021) dehydration during mineral decomposition, and volume reduction. Due to a slight negative temperature dependence on water solubility of stishovite at 20 GPa (Fig. 6a), the water in stishovite will decrease with increasing temperature. That indicates a water outflow from stishovite along with the subducted slab being heated in MTZ. Comparison with a hot subduction geotherm (Table S2 and Fig. B1), slabs will reduce more water in a cold subduction geotherm. 320 Additionally, stishovite water solubility peaks at 22-32 GPa and 800-1600 °C (corresponding conditions of MTZ or topmost of the lower mantle) (Fig. 6) and declines at higher pressure-temperature (P-T) conditions. Previous studies suggest hydration of a large region of the transition zone and that dehydration melting may act to trap H<sub>2</sub>O in the transition zone (e.g., Schmandt et al., 2014). Taking into account the occurrence of melts, the optimal solubility likely falls within the water-rich mantle transition zone at pressures below 22 GPa. Consequently, stishovite emerges as a key transporter supplying water to the mantle transition zone (Walter, 2021). 325

Formatted: Indent: First line: 1 ch




Figure 7: Water solubility in a cold subducted slab. Dashed lines represent water solubility of minerals and solid lines show the actual water contents considering mineral modal fractions (from Litasov and Ohtani, 2007) in subduction slabs. A cold subduction geotherm gradient from Litasov and Ohtani (2007) is used for the calculation (Table S2). The calculated result along with a hot subduction geotherm is presented in Fig. B1 in the Appendix. Water solubility models: Grt from Lu and Keppler (1997), Cpx from Gavrilenko (2008), St from this study.

330



## 7 Conclusions and outlook

335 To sum up, the comprehensive review and exploration in this paper have encompassed water solubility in stishovite, hydrogen incorporation mechanisms and governing factors. We've constructed a new model for water solubility in Al-bearing stishovite and delved into stishovite's role in transporting water through a subducting ~~erust~~ slab and implications for water distribution in deep Earth. The key conclusions can be summarized as follows:

340 1. Water solubility in stishovite exhibits a positive correlation with Al content, which increases water concentrations via coupled substitution  $\text{Al}^{3+} + \text{H}^+ \rightarrow \text{Si}^{4+}$ , as well as enhances the incorporation of interstitial  $\text{H}_2\text{O}$  solubility by Al substitution-induced expansion of the stishovite structure.

2. The relationship between water solubility of Al-bearing stishovite and water fugacity, temperature, pressure, and Al content can be described as  $C_{\text{OH}}^{\text{St}} = A \cdot f_{\text{H}_2\text{O}}^n \cdot \exp\left(-\frac{\Delta E + \Delta V \cdot P}{R \cdot T}\right) \cdot \exp\left(\frac{B \cdot X_{\text{Al}}}{R \cdot T}\right)$  with  $n = 0.5$ ,  $A = 0.239\text{-}24 \pm 13 \text{ ppm/GPa}^{0.5}$ ,  $\Delta E = -3.065 \pm 0.88 \text{ kJ/mol}$ ,  $\Delta V = 4.29 \pm 0.27 \text{ cm}^3/\text{mol}$ ,  $B = 7.69 \pm 1.12 \text{ kJ/mol}$ .

345 3. The calculation results from Eq. (2) demonstrate that stishovite's water solubility strongly depends on pressure, exhibiting a positive correlation below 22-32 GPa and a negative correlation above, indicating a peak solubility range at 22-32 GPa. While temperature dependence is weakly negative up to 30 GPa and weakly positive beyond.

350 4. Following the cold subduction slab mantle geotherm, stishovite's water solubility increases from 998642 wt. ppm at 14 GPa to 9061317 wt. ppm at 22 GPa. This suggests that stishovite can potentially absorb/accommodate water released during the breakdown of hydrous mineral breakdown, dehydration reactions, and volume reduction, contributing to the water-rich transition zone.

355 However, several crucial unanswered questions persist in research on water solubility of stishovite water solubility research: 1. The behavior/solubility of Al and its impact on water solubility across varying pressure-temperature conditions requires clarification; 2. The roles of minor elements like Fe and Ti, as well as oxygen fugacity, remain unclear; 3. Partitioning coefficients governing water distribution between stishovite, and other major minerals-phases (e.g., Cpx, Grt, melts) and melts during slab subduction are not yet understood; 4. Equation (2) proposed in this paper needs refinement through systematic experiments conducted under controlled conditions. These issues are essential for understanding how water transports into deep Earth during subduction and require further detailed experimental and simulation-based investigations to address.

360 **Appendix A: Comparison between experimental and calculated results**





Figure A1: Comparison between experimental and calculated results (Table S1).

**Appendix B: Water solubility of minerals in hot subducted slabs at 10-25 GPa**



365 **Figure B1: Water solubility in hot subducted slabs. Dashed lines represent water solubility of minerals and solid lines show the actual water contents considering mineral modal fractions (from Litasov and Ohtani, 2007). Water solubility models: Grt from Lu and Keppler (1997), Cpx from Gavrilenko (2008), St from this study.**

*Data availability.* [Data have been made available in the Supplement.](#)

370

*Author contribution.* Mengdan Chen: Data curation, formal analysis and writing-original draft. Lei Kang: Writing - review & editing, foundation support. Danling Chen and Liang Liu: Writing - review & editing. Changxin Yin and Long Tian: Review and language modification.

375 *Competing interest.* The authors declare that they have no conflict of interest.

*Acknowledgments.* We are grateful to [Professor Shun Karato and Dr. Xin Li](#) for the discussions on Al behavior in stishovite. [Thanks to three anonymous reviewers for their constructive comments that significantly improved this article.](#)

380 *Financial support.* This study is supported by the Natural Science Foundation of China (Grant No. 42030307, 41972058, 42372064) and the MOST Special Fund from the State Key Laboratory of Continental Dynamics, Northwest University (Grant No. 201210233).

## References

- ell, D. R., Ihinger, P. D., and Rossman, G. R.: Quantitative analysis of trace OH in garnet and pyroxenes, *American Mineralogist*, 80, 465–474, <https://doi.org/10.2138/am-1995-5-607>, 1995.
- 385 Bolfan-Casanova, N., Keppler, H., and Rubie, D. C.: Water partitioning between nominally anhydrous minerals in the MgO–SiO<sub>2</sub>–H<sub>2</sub>O system up to 24 GPa: implications for the distribution of water in the Earth’s mantle, *Earth and Planetary Science Letters*, 182, 209–221, [https://doi.org/10.1016/S0012-821X\(00\)00244-2](https://doi.org/10.1016/S0012-821X(00)00244-2), 2000.
- 390 Bromiley, G. D., Bromiley, F. A., and Bromiley, D. W.: On the mechanisms for H and Al incorporation in stishovite, *Phys Chem Minerals*, 33, 613–621, <https://doi.org/10.1007/s00269-006-0107-9>, 2006.
- Chao, E. C. T., Fahey, J. J., Littler, J., and Milton, D. J.: Stishovite, SiO<sub>2</sub>, a very high pressure new mineral from Meteor Crater, Arizona, *J. Geophys. Res.*, 67, 419–421, <https://doi.org/10.1029/JZ067i001p00419>, 1962.
- Chung, J. I. and Kagi, H.: High concentration of water in stishovite in the MORB system, *Geophys. Res. Lett.*, 29, 2020, <https://doi.org/10.1029/2002GL015579>, 2002.
- 395 Fischer, R. A., Campbell, A. J., Chidester, B. A., Reaman, D. M., Thompson, E. C., Pigott, J. S., Prakapenka, V. B., and Smith, J. S.: Equations of state and phase boundary for stishovite and CaCl<sub>2</sub>-type SiO<sub>2</sub>, *American Mineralogist*, 103, 792–802, <https://doi.org/10.2138/am-2018-6267>, 2018.
- Frigo, C., Stalder, R., and Ludwig, T.: OH defects in coesite and stishovite during ultrahigh-pressure metamorphism of continental crust, *Phys Chem Minerals*, 46, 77–89, <https://doi.org/10.1007/s00269-018-0987-5>, 2019.
- 400 Gavrilenko, P.: Water solubility in diopside, Ph.D. thesis, Universität Bayreuth, Germany, 2008.
- Gu, T., Pamato, M. G., Novella, D., Alvaro, M., Fournelle, J., Brenker, F. E., Wang, W., and Nestola, F.: Hydrous peridotitic fragments of Earth’s mantle 660 km discontinuity sampled by a diamond, *Nat. Geosci.*, 15, 950–954, <https://doi.org/10.1038/s41561-022-01024-y>, 2022.
- 405 Gutzow, I., Pascova, R., Jordanov, N., Gutzow, S., Penkov, I., Markovska, I., Schmelzer, J. W. P., and Ludwig, F. P.: 3. Crystalline and Amorphous Modifications of Silica: Structure, Thermodynamic Properties, Solubility, and Synthesis, in: *Glass*, edited by: Schmelzer, J. W. P., De Gruyter, 137–196, <https://doi.org/10.1515/9783110298581.137>, 2014.
- Holtstam, D., Broman, C., Söderhielm, J., and Zetterqvist, A.: First discovery of stishovite in an iron meteorite, *Meteoritics & Planetary Science*, 38, 1579–1583, <https://doi.org/10.1111/j.1945-5100.2003.tb00002.x>, 2003.

- 410 Irifune, T. and Ringwood, A. E.: Phase transformations in subducted oceanic crust and buoyancy relationships at depths of  
600–800 km in the mantle, *Earth and Planetary Science Letters*, 117, 101–110, [https://doi.org/10.1016/0012-821X\(93\)90120-X](https://doi.org/10.1016/0012-821X(93)90120-X), 1993.
- Irifune, T., Sekine, T., Ringwood, A. E., and Hibberson, W. O.: The eclogite-garnetite transformation at high pressure and  
some geophysical implications, *Earth and Planetary Science Letters*, 77, 245–256, [https://doi.org/10.1016/0012-821X\(86\)90165-2](https://doi.org/10.1016/0012-821X(86)90165-2), 1986.
- 415 Irifune, T., Ringwood, A. E., and Hibberson, W. O.: Subduction of continental crust and terrigenous and pelagic sediments:  
an experimental study, *Earth and Planetary Science Letters*, 126, 351–368, [https://doi.org/10.1016/0012-821X\(94\)90117-1](https://doi.org/10.1016/0012-821X(94)90117-1), 1994.
- Ishii, T., Kojitani, H., and Akaogi, M.: High-pressure phase transitions and subduction behavior of continental crust at  
pressure–temperature conditions up to the upper part of the lower mantle, *Earth and Planetary Science Letters*, 357–358,  
420 31–41, <https://doi.org/10.1016/j.epsl.2012.09.019>, 2012.
- Ishii, T., Kojitani, H., and Akaogi, M.: Phase Relations of Harzburgite and MORB up to the Uppermost Lower Mantle  
Conditions: Precise Comparison With Pyrolite by Multisample Cell High-Pressure Experiments With Implication to  
Dynamics of Subducted Slabs, *J. Geophys. Res. Solid Earth*, 124, 3491–3507, <https://doi.org/10.1029/2018JB016749>,  
2019.
- 425 Ishii, T., Criniti, G., Ohtani, E., Purevjav, N., Fei, H., Katsura, T., and Mao, H.: Superhydrous aluminous silica phases as major  
water hosts in high-temperature lower mantle, *Proc. Natl. Acad. Sci. U.S.A.*, 119, e2211243119,  
<https://doi.org/10.1073/pnas.2211243119>, 2022.
- Johnson, E. A.: Water in Nominally Anhydrous Crustal Minerals: Speciation, Concentration, and Geologic Significance,  
Reviews in Mineralogy and Geochemistry, 62, 117–154, <https://doi.org/10.2138/rmg.2006.62.6>, 2006.
- 430 Kaminsky, F.: Mineralogy of the lower mantle: A review of ‘super-deep’ mineral inclusions in diamond, *Earth-Science  
Reviews*, 110, 127–147, <https://doi.org/10.1016/j.earscirev.2011.10.005>, 2012.
- Karato, S.: Rheology of the deep upper mantle and its implications for the preservation of the continental roots: A review,  
*Tectonophysics*, 481, 82–98, <https://doi.org/10.1016/j.tecto.2009.04.011>, 2010.
- 435 Kesar, N. R., Troullier, N., Martins, J. L., and Chelikowsky, J. R.: Structural properties of SiO<sub>2</sub> in the stishovite structure,  
*Phys. Rev. B*, 44, 4081–4088, <https://doi.org/10.1103/PhysRevB.44.4081>, 1991.
- Kohlstedt, D. L., Keppler, H., and Rubie, D. C.: Solubility of water in the  $\alpha$ ,  $\beta$  and  $\gamma$  phases of (Mg,Fe)<sub>2</sub>SiO<sub>4</sub>, *Contributions to  
Mineralogy and Petrology*, 123, 345–357, <https://doi.org/10.1007/s004100050161>, 1996.
- Kueter, N., Brugman, K., Miozzi, F., Cody, G. D., Yang, J., Strobel, T. A., and Walter, M. J.: Water speciation and hydrogen  
isotopes in hydrous stishovite: implications for the deep Earth water cycle, *Contrib Mineral Petrol*, 178, 48,  
440 <https://doi.org/10.1007/s00410-023-02028-6>, 2023.
- Lakshtanov, D. L., Litasov, K. D., Sinogeikin, S. V., Hellwig, H., Li, J., Ohtani, E., and Bass, J. D.: Effect of Al<sup>3+</sup> and H<sup>+</sup> on  
the elastic properties of stishovite, *American Mineralogist*, 92, 1026–1030, <https://doi.org/10.2138/am.2007.2294>, 2007a.
- Lakshtanov, D. L., Sinogeikin, S. V., Litasov, K. D., Prakupenka, V. B., Hellwig, H., Wang, J., Sanches-Valle, C., Perrillat, J.  
P., Chen, B., Somayazulu, M., Li, J., Ohtani, E., and Bass, J. D.: The post-stishovite phase transition in hydrous alumina-  
445 bearing SiO<sub>2</sub> in the lower mantle of the earth, *Proc. Natl. Acad. Sci. U.S.A.*, 104, 13588–13590,  
<https://doi.org/10.1073/pnas.0706113104>, 2007b.
- Li, J., Lin, Y., Meier, T., Liu, Z., Yang, W., Mao, H., Zhu, S., and Hu, Q.: Silica-water superstructure and one-dimensional  
superionic conduit in Earth’s mantle, *Sci. Adv.*, 9, eadh3784, <https://doi.org/10.1126/sciadv.adh3784>, 2023.
- Libowitzky, E.: The Structure of Hydrous Species in Nominally Anhydrous Minerals: Information from Polarized IR  
Spectroscopy, *Reviews in Mineralogy and Geochemistry*, 62, 29–52, <https://doi.org/10.2138/rmg.2006.62.2>, 2006.
- 450 Libowitzky, E. and Rossman, G. R.: An IR absorption calibration for water in minerals, *American Mineralogist*, 82, 1111–  
1115, <https://doi.org/10.2138/am-1997-11-1208>, 1997.
- Lin, Y., Hu, Q., Meng, Y., Walter, M., and Mao, H. K.: Evidence for the stability of ultrahydrous stishovite in Earth’s lower  
mantle, *Proc. Natl. Acad. Sci. U.S.A.*, 117, 184–189, <https://doi.org/10.1073/pnas.1914295117>, 2020.
- 455 Lin, Y., Hu, Q., Walter, M. J., Yang, J., Meng, Y., Feng, X., Zhuang, Y., Cohen, R. E., and Mao, H. K.: Hydrous SiO<sub>2</sub> in  
subducted oceanic crust and H<sub>2</sub>O transport to the core-mantle boundary, *Earth and Planetary Science Letters*, 594, 117708,  
<https://doi.org/10.1016/j.epsl.2022.117708>, 2022.

- Litasov, K. D. and Ohtani, E.: Effect of water on the phase relations in Earth's mantle and deep water cycle, in: *Advances in High-Pressure Mineralogy*, Geological Society of America, [https://doi.org/10.1130/2007.2421\(08\)](https://doi.org/10.1130/2007.2421(08)), 2007.
- 460 Litasov, K. D., Kagi, H., Shatskiy, A., Ohtani, E., Lakshatnov, D. L., Bass, J. D., and Ito, E.: High hydrogen solubility in Al-rich stishovite and water transport in the lower mantle, *Earth and Planetary Science Letters*, 262, 620–634, <https://doi.org/10.1016/j.epsl.2007.08.015>, 2007.
- 465 Liu, L., Zhang, J., Green, H. W., Jin, Z., and Bozhilov, K. N.: Evidence of former stishovite in metamorphosed sediments, implying subduction to > 350 km, *Earth and Planetary Science Letters*, 263, 180–191, <https://doi.org/10.1016/j.epsl.2007.08.010>, 2007.
- Liu, L., Zhang, J. F., Cao, Y. T., Green, H. W., Yang, W. Q., Xu, H. J., Liao, X. Y., and Kang, L.: Evidence of former stishovite in UHP eclogite from the South Altyn Tagh, western China, *Earth and Planetary Science Letters*, 484, 353–362, <https://doi.org/10.1016/j.epsl.2017.12.023>, 2018.
- 470 Liu, X., Nishiyama, N., Sanehira, T., Inoue, T., Higo, Y., and Sakamoto, S.: Decomposition of kyanite and solubility of Al<sub>2</sub>O<sub>3</sub> in stishovite at high pressure and high temperature conditions, *Phys Chem Minerals*, 33, 711–721, <https://doi.org/10.1007/s00269-006-0122-x>, 2006.
- Liu, X. W., Xie, Z. J., Wang, L., Xu, W., and Jin, Z. M.: Water incorporation in garnets from ultrahigh pressure eclogites at Shuanghe, Dabieshan, *Mineral. mag.*, 80, 959–975, <https://doi.org/10.1180/minmag.2016.080.034>, 2016.
- 475 Lu, R. and Keppler, H.: Water solubility in pyrope to 100 kbar, *Contributions to Mineralogy and Petrology*, 129, 35–42, <https://doi.org/10.1007/s004100050321>, 1997.
- Magni, V., Bouilhol, P., and Van Hunen, J.: Deep water recycling through time, *Geochem. Geophys. Geosyst.*, 15, 4203–4216, <https://doi.org/10.1002/2014GC005525>, 2014.
- Ni, H., Zheng, Y. F., Mao, Z., Wang, Q., Chen, R.-X., and Zhang, L.: Distribution, cycling and impact of water in the Earth's interior, *National Science Review*, 4, 879–891, <https://doi.org/10.1093/nsr/nwx130>, 2017.
- 480 Nisr, C., Shim, S. H., Leinenweber, K., and Chizmeshya, A.: Raman spectroscopy of water-rich stishovite and dense high-pressure silica up to 55 GPa, *American Mineralogist*, 102, 2180–2189, <https://doi.org/10.2138/am-2017-5944>, 2017.
- Nisr, C., Chen, H., Leinenweber, K., Chizmeshya, A., Prakapenka, V. B., Prescher, C., Tkachev, S. N., Meng, Y., Liu, Z., and Shim, S. H.: Large H<sub>2</sub>O solubility in dense silica and its implications for the interiors of water-rich planets, *Proc. Natl. Acad. Sci. U.S.A.*, 117, 9747–9754, <https://doi.org/10.1073/pnas.1917448117>, 2020.
- 485 Ono, S.: High temperature stability limit of phase egg, AlSiO<sub>3</sub>(OH), *Contributions to Mineralogy and Petrology*, 137, 83–89, <https://doi.org/10.1007/s004100050583>, 1999.
- Ono, S., Ito, E., and Katsura, T.: Mineralogy of subducted basaltic crust (MORB) from 25 to 37 GPa, and chemical heterogeneity of the lower mantle, *Earth and Planetary Science Letters*, 190, 57–63, [https://doi.org/10.1016/S0012-821X\(01\)00375-2](https://doi.org/10.1016/S0012-821X(01)00375-2), 2001.
- 490 Ono, S., Suto, T., Hirose, K., Kuwayama, Y., Komabayashi, T., and Kikegawa, T.: Equation of state of Al-bearing stishovite to 40 GPa at 300 K, *American Mineralogist*, 87, 1486–1489, <https://doi.org/10.2138/am-2002-1026>, 2002.
- Ono, S., Kikegawa, T., Higo, Y., and Tange, Y.: Precise determination of the phase boundary between coesite and stishovite in SiO<sub>2</sub>, *Physics of the Earth and Planetary Interiors*, 264, 1–6, <https://doi.org/10.1016/j.pepi.2017.01.003>, 2017.
- 495 Panero, W. R. and Stixrude, L. P.: Hydrogen incorporation in stishovite at high pressure and symmetric hydrogen bonding in δ-AlOOH, *Earth and Planetary Science Letters*, 221, 421–431, [https://doi.org/10.1016/S0012-821X\(04\)00100-1](https://doi.org/10.1016/S0012-821X(04)00100-1), 2004.
- Panero, W. R., Benedetti, L. R., and Jeanloz, R.: Transport of water into the lower mantle: Role of stishovite: TRANSPORT OF WATER INTO THE LOWER MANTLE, *J. Geophys. Res.*, 108, <https://doi.org/10.1029/2002JB002053>, 2003.
- Paterson, M. S.: The determination of hydroxyl by infrared absorption in quartz, silicate glasses and similar materials, *bulmi*, 105, 20–29, <https://doi.org/10.3406/bulmi.1982.7582>, 1982.
- 500 Pawley, A. R., McMillan, P. F., and Holloway, J. R.: Hydrogen in Stishovite, with Implications for Mantle Water Content, *Science*, 261, 1024–1026, <https://doi.org/10.1126/science.261.5124.1024>, 1993.
- Pearson, D. G., Brenker, F. E., Nestola, F., McNeill, J., Nasdala, L., Hutchison, M. T., Matveev, S., Mather, K., Silversmit, G., Schmitz, S., Vekemans, B., and Vincze, L.: Hydrous mantle transition zone indicated by ringwoodite included within diamond, *Nature*, 507, 221–224, <https://doi.org/10.1038/nature13080>, 2014.
- 505 Peslier, A. H., Schönbacher, M., Busemann, H., and Karato, S. I.: Correction to: Water in the Earth's Interior: Distribution and Origin, *Space Sci Rev*, 212, 811–811, <https://doi.org/10.1007/s11214-017-0420-2>, 2017.

- Petersen, S. E., Hoisch, T. D., and Porter, R. C.: Assessing the Role of Water in Alaskan Flat-Slab Subduction, *Geochem Geophys Geosyst*, 22, e2021GC009734, <https://doi.org/10.1029/2021GC009734>, 2021.
- 510 Poli, S. and Schmidt, M. W.: Petrology of Subducted Slabs, *Annu. Rev. Earth Planet. Sci.*, 30, 207–235, <https://doi.org/10.1146/annurev.earth.30.091201.140550>, 2002.
- Rossmann, G. R.: Studies of OH in nominally anhydrous minerals, *Phys Chem Minerals*, 23, <https://doi.org/10.1007/BF00207777>, 1996.
- Schmandt, B., Jacobsen, S. D., Becker, T. W., Liu, Z., and Dueker, K. G.: Dehydration melting at the top of the lower mantle, *Science*, 344, 1265–1268, <https://doi.org/10.1126/science.1253358>, 2014.
- 515 Shillington, D. J.: Water takes a deep dive into an oceanic tectonic plate, *Nature*, 563, 335–336, <https://doi.org/10.1038/d41586-018-07335-8>, 2018.
- Smith, E. M., Shirey, S. B., Richardson, S. H., Nestola, F., Bullock, E. S., Wang, J., and Wang, W.: Blue boron-bearing diamonds from Earth's lower mantle, *Nature*, 560, 84–87, <https://doi.org/10.1038/s41586-018-0334-5>, 2018.
- Smyth, J. R.: Hydrogen in High Pressure Silicate and Oxide Mineral Structures, *Reviews in Mineralogy and Geochemistry*, 520 62, 85–115, <https://doi.org/10.2138/rmg.2006.62.5>, 2006.
- Spektor, K., Nylene, J., Stoyanov, E., Navrotsky, A., Hervig, R. L., Leinenweber, K., Holland, G. P., and Häussermann, U.: Ultrahydrous stishovite from high-pressure hydrothermal treatment of SiO<sub>2</sub>, *Proc. Natl. Acad. Sci. U.S.A.*, 108, 20918–20922, <https://doi.org/10.1073/pnas.1117152108>, 2011.
- 525 Spektor, K., Nylene, J., Mathew, R., Edén, M., Stoyanov, E., Navrotsky, A., Leinenweber, K., and Häussermann, U.: Formation of hydrous stishovite from coesite in high-pressure hydrothermal environments, *American Mineralogist*, 101, 2514–2524, <https://doi.org/10.2138/am-2016-5609>, 2016.
- Stishov, S. M. and Popova, S. V.: A new dense modification of silica, *Geokhimiya*, 10, 923–926, 1961.
- Thomas, R., Davidson, P., Rericha, A., and Recknagel, U.: Discovery of Stishovite in the Prismatic-Bearing Granulite from Waldheim, Germany: A Possible Role of Supercritical Fluids of Ultrahigh-Pressure Origin, *Geosciences*, 12, 196, <https://doi.org/10.3390/geosciences12050196>, 2022.
- 530 Thomas, S. M., Koch-Müller, M., Reichart, P., Rhede, D., Thomas, R., Wirth, R., and Matsyuk, S.: IR calibrations for water determination in olivine, r-GeO<sub>2</sub>, and SiO<sub>2</sub> polymorphs, *Phys Chem Minerals*, 36, 489–509, <https://doi.org/10.1007/s00269-009-0295-1>, 2009.
- Walter, M. J.: Water transport to the core–mantle boundary, *National Science Review*, 8, nwab007, <https://doi.org/10.1093/nsr/nwab007>, 2021.
- 535 Wu, Y., Fei, Y., Jin, Z., and Liu, X.: The fate of subducted Upper Continental Crust: An experimental study, *Earth and Planetary Science Letters*, 282, 275–284, <https://doi.org/10.1016/j.epsl.2009.03.028>, 2009.
- Xia, Q. K., Chen, D. G., Delouie, E., and Zhi, X. C.: Hydrogen isotopic compositions of mantle-derived megacrysts from cenozoic basalts, Eastern China, *Chin.Sci.Bull.*, 43, 146–146, <https://doi.org/10.1007/BF02891632>, 1998.
- 540 Yan, B., Liu, S., Chastain, M. L., Yang, S., and Chen, J.: A new FTIR method for estimating the firing temperature of ceramic bronze-casting moulds from early China, *Sci Rep*, 11, 3316, <https://doi.org/10.1038/s41598-021-82806-z>, 2021.
- Yang, J. S., Dobrzhinetskaya, L., Bai, W. J., Fang, Q.-S., Robinson, P. T., Zhang, J., and Green, H. W.: Diamond- and coesite-bearing chromitites from the Luobusa ophiolite, Tibet, *Geol*, 35, 875, <https://doi.org/10.1130/G23766A.1>, 2007.
- 545 Yang, X.: Effect of oxygen fugacity on OH dissolution in olivine under peridotite-saturated conditions: An experimental study at 1.5–7 GPa and 1100–1300 °C, *Geochimica et Cosmochimica Acta*, 173, 319–336, <https://doi.org/10.1016/j.gca.2015.11.007>, 2016.
- Yang, X. and Li, Y.: High-P/T experimental studies and water in the silicate mantle, *Sci. China Earth Sci.*, 59, 683–695, <https://doi.org/10.1007/s11430-015-5241-0>, 2016.
- 550 Yoshino, T., Shimojuku, A., and Li, D.: Electrical conductivity of stishovite as a function of water content, *Physics of the Earth and Planetary Interiors*, 227, 48–54, <https://doi.org/10.1016/j.pepi.2013.12.003>, 2014.
- Zhang, K., Liu, H., Ionov, D. A., and Yang, X.: Effects of Oxygen Fugacity on Hydroxyl Incorporation in Garnet at 1–3 GPa and 800–1000 °C and Implications for Water Storage in the Mantle, *JGR Solid Earth*, 127, e2022JB023948, <https://doi.org/10.1029/2022JB023948>, 2022a.
- 555 Zhang, Y., Fu, S., Karato, S., Okuchi, T., Chariton, S., Prakapenka, V. B., and Lin, J.: Elasticity of Hydrated Al-Bearing Stishovite and Post-Stishovite: Implications for Understanding Regional Seismic  $V_S$  Anomalies Along Subducting Slabs in the Lower Mantle, *JGR Solid Earth*, 127, <https://doi.org/10.1029/2021JB023170>, 2022b.

Zhao, Y. H., Ginsberg, S. B., and Kohlstedt, D. L.: Solubility of hydrogen in olivine: dependence on temperature and iron content, *Contributions to Mineralogy and Petrology*, 147, 155–161, <https://doi.org/10.1007/s00410-003-0524-4>, 2004.

Zheng, Y., Chen, R., Xu, Z., and Zhang, S.: The transport of water in subduction zones, *Sci. China Earth Sci.*, 59, 651–682, <https://doi.org/10.1007/s11430-015-5258-4>, 2016.

560