

<https://doi.org/10.5194/egusphere-2023-2343>

Preprint. Discussion started: 3 November 2023

© Author(s) 2023. CC BY 4.0 License.



Derivation of Atmospheric Reaction Mechanisms for Volatile Organic Compounds by the SAPRC Mechanism Generation System (MechGen)

William P. L. Carter¹, Jia Jiang², John J. Orlando³, Kelley C. Barsanti^{1,3,4}

¹College of Engineering Center for Environmental Research and Technology (CE-CERT)
University of California, Riverside, CA 92521

²Department of Civil and Environmental Engineering, University of California, Davis, CA 95616

³Atmospheric Chemistry Observations & Modeling Lab, National Center for Atmospheric Research, P.O. Box 3000,
Boulder, CO 80307

⁴Department of Chemical and Environmental Engineering University of California, Riverside, CA 92521

Correspondence to: William Carter (william.carter@ucr.edu)



1

ABSTRACT

2 This paper describes the methods that are used in the SAPRC mechanism generation system, MechGen,
3 to estimate rate constants and derive mechanisms for gas-phase reactions of volatile organic compounds
4 (VOCs) in the lower atmosphere. Earlier versions of this system have been used for over 20 years in the
5 development of the SAPRC mechanisms for air quality models and ozone (O₃) reactivity assessments.
6 This is the first complete documentation of MechGen as it currently exists. MechGen can be used to
7 derive explicit mechanisms for most compounds with C, H, O, or N atoms. Included are reactions of
8 stable compounds with hydroxy (OH) and nitrate (NO₃) radicals, O₃, O³P, or by photolysis or
9 unimolecular reactions, and the reactions of the radicals they form in the presence of O₂ and oxides of
10 nitrogen (NO_x) at atmospheric temperatures and pressures. Measured or theoretically calculated rate
11 constants and branching ratios are used when data are available, but in most cases rate constants and
12 branching ratios are estimated using various structure-reactivity or other estimation methods. Types of
13 reactions that are estimated include initial reactions of organics with atmospheric oxidants or by
14 photolysis, and unimolecular and bimolecular reactions of carbon-centered, alkoxy, and peroxy radicals
15 and Criegee and other intermediates that are formed. This manuscript summarizes the capabilities of the
16 MechGen system for deriving chemical mechanisms and the methods, assignments, and estimates used to
17 derive them. Examples of MechGen predictions also are provided. Many of the estimation methods
18 discussed here have not been published previously, and others have not been used previously in
19 developing comprehensive mechanisms. This paper concludes with a discussion of the areas of
20 uncertainty where additional measurements or theoretical estimates are needed and recommendations for
21 future mechanism development and application.



22 1. Introduction

23 1.1. Background

24 When most volatile organic compounds (VOCs) are introduced into the atmosphere in the
25 presence of sunlight, they can react to form various radicals that then further react to form oxidized
26 organic products, including gas-phase toxics and secondary organic aerosol (SOA). In the presence of
27 oxides of nitrogen (NO_x) from combustion sources, these radicals also promote the formation of ozone
28 (O_3) and oxidized nitrogen compounds that further affect air quality. Many hundreds of types of organic
29 compounds are emitted, from both anthropogenic and biogenic sources. The atmospheric reaction
30 mechanisms for these compounds are complex, and for larger molecules can involve an extremely large
31 number of reactive intermediates and result in a large number of stable oxidized organic products. Most
32 of these organic products also react in the atmosphere, forming even more intermediates and products. In
33 most cases these mechanisms involve reactions whose rate constants, and in some cases mechanisms, are
34 unknown and thus have to be estimated. Because of the complexity, it is necessary either to greatly
35 simplify the mechanisms for most VOCs, use extensive lumping or condensations in representations of
36 VOCs, or use an automated chemical mechanism generation system to derive the mechanisms.

37 In the case of tropospheric reaction mechanisms of VOCs, existing automated chemical
38 mechanism generation systems include the Generator of Explicit Chemistry and Kinetics of Organics in
39 the Atmosphere (GECKO-A) developed by Aumont and co-workers (Aumont et al., 2005) and the
40 SAPRC mechanism generation system, MechGen, that is the subject of this paper. GECKO-A has been
41 used in a number of studies of varying chemical complexity (e.g., Aumont et al., 2005; Camredon et al.,
42 2007; Lee-Taylor et al., 2011; Aumont et al., 2012; Lannuque et al., 2018; Afreh et al., 2021) and is
43 designed primarily to derive and carry out model simulations using multi-generation mechanisms of
44 selected compounds and all of their oxidation products. The SAPRC MechGen system was developed
45 primarily to support the development of versions of the SAPRC atmospheric gas-phase chemical
46 mechanisms (Carter, 2000, 2010a, b, 2016, 2020), and has been focused on single-generation reactions of
47 individual compounds, with mechanisms for representative organic oxidation products being derived
48 separately. Therefore, the objectives and operations of the two systems are somewhat different. In
49 addition, although the two systems employ similar structure-activity relationships (SARs) and can give
50 very similar predictions, there are some differences in the mechanisms they derive, reflecting both
51 different treatments of uncertainties and also different areas where updates are needed to incorporate
52 recent results. For example, GECKO-A does not yet predict autoxidation reactions of peroxy radicals that
53 are believed to occur (and are predicted by MechGen), while GECKO-A employs more detailed and
54 updated estimates for bimolecular reactions of peroxy radicals and photolysis reactions of larger
55 molecules.

56 Perhaps the most important difference between GECKO-A and MechGen is that GECKO-A is
57 described in the peer-reviewed literature (e.g., Aumont et al., 2005), while the description of MechGen is
58 largely incomplete in the literature, with only a preliminary description in online documentation (Carter,
59 2019) and abbreviated description in applied studies (Jiang et al., 2020; Li et al., 2022). Different versions
60 of MechGen, incorporating our changing and evolving understanding and ability to estimate the
61 underlying chemistry, have been used in the development of SAPRC-99 (Carter, 2000), SAPRC-07
62 (Carter, 2010a, b), SAPRC-11 (Carter and Heo, 2013), SAPRC-16 (Carter, 2016; Venecek et al., 2018)
63 and SAPRC-18 (Carter, 2020). Additional updates to the system have been made since the release of
64 SAPRC-18. The lack of a stable and peer-reviewed version has inhibited the use of MechGen for
65 atmospheric chemistry research beyond the development of SAPRC mechanisms, despite its potential
66 utility as an alternative to or for comparison with GECKO-A or the semi-explicit Master Chemical
67 Mechanism (Jenkin et al., 1997, 2003; Bloss et al., 2005; MCM, 2023). The purpose of this paper is to



68 document the chemical basis of the estimates and assignments as it currently exists, so it can be
69 appropriately cited and more widely used. The version of MechGen described here has been used in the
70 preparation of the recently-completed SAPRC-22 mechanisms (Carter, 2023a). This paper can also be
71 used to as a starting point for future updates and to illustrate areas of uncertainty where experimental or
72 theoretical studies are needed.

73 1.2. Scope

74 MechGen is capable of generating fully explicit mechanisms for the atmospheric reactions of
75 most types of organic compounds emitted into the atmosphere and the intermediate radicals they form. It
76 is designed to generate mechanisms for lower tropospheric modeling only, so its mechanisms are
77 applicable primarily for temperatures at or near 300 K and pressures at or near 1 atmosphere. Although
78 temperature-dependent rate constants are assigned or estimated for many reactions, for others the rate
79 constants or branching ratios are only applicable for 300 K and 1 atmosphere. In particular, MechGen is
80 not currently designed for estimating mechanisms for combustion modeling or for low temperature or low
81 pressure systems.

82 Table 1 lists the types of stable compounds and the categories of initial atmospheric reactions that
83 can be generated for those compounds. Table 2 lists the types of reactions within each category that can
84 be generated, including reactions of intermediate radicals and reactions of stable compounds. These
85 include H-atom abstractions from stable compounds by OH, NO₃, and Cl radicals; additions to double
86 bonds by these radicals and by O₃ and O³P; and photolyses at various groups. The types of radicals that
87 can be generated include carbon-centered radicals that in most cases react primarily with O₂; peroxy
88 radicals that in most cases react with NO, NO₂, NO₃, HO₂, or other peroxy radicals and in some cases also
89 have unimolecular reactions; alkoxy radicals that can react with O₂ or by various types of unimolecular
90 reactions; and excited and stabilized Criegee intermediates (CI). More information about their reactions
91 and how they are generated are described below, with additional detail given in the Supplementary
92 Information (SI) as referenced below.

93 Although the MechGen system can represent chemical structures containing halogen atoms and
94 estimate some types of reactions of compounds with Cl atoms, not all types of reactions of halogen-
95 containing compounds are currently supported. Therefore, a discussion of reactions of halogen-containing
96 compounds is not included here.

97 The operations and capabilities of MechGen as a software system are described in a separate
98 manuscript and user's manual (Carter, In Prep.), while this paper focuses on the chemical basis for the
99 derived mechanisms.

100 1.3. Designation of Structures

101 Information on how to input and designate structures for MechGen is given in the software
102 description and users' manual (Carter, In Prep.). Briefly, the structure of an organic reactant or radical is
103 specified by giving the "groups" in the molecule or radical and indicating the groups each are bonded to,
104 the type of bond, and in some cases (e.g., *cis* or *trans* isomers) the orientation of groups around the bonds.
105 Groups are parts of molecules that are treated as units in the system and are used to determine which
106 types of reactions can occur and what products are formed and used by the SARs or other methods to
107 estimate rate constants. These groups contain no more than one carbon or nitrogen atom, and also



108 Table 1. Types of stable compounds whose reactions are supported by MechGen.

Type of Compound	Reactions	Exceptions not Supported
Alkanes	OH, NO ₃	Some estimates for polycyclic compounds are not reliable
Alkenes (including multiple double bonds)	OH, O ₃ , NO ₃ , O ³ P	See above; not all reactions of Cl-addition radicals can be generated
Alkynes	OH, NO ₃	See above
Aromatic hydrocarbons	OH	PAHs other than naphthalenes
Aldehydes, ketones, hydroperoxides, organic nitrates	OH, NO ₃ , hv	Photolysis reactions of larger compounds are over-simplified
Alcohols, ethers, esters, other oxygenates	OH, NO ₃	
Unsaturated aldehydes, ketones, hydroperoxides, and nitrates	OH, O ₃ , NO ₃ , O ³ P, hv	Ketenes
Phenols	OH, NO ₃	
Furans	OH, O ₃ , NO ₃ , O ³ P	
Other unsaturated oxygenates	OH, O ₃ , NO ₃ , O ³ P	
Amines	OH, NO ₃	Amines with no α or β hydrogens; imines
Peroxy nitrates	Unimolecular, OH, NO ₃ , hv	Formation of alkyl peroxy nitrates from RO ₂ + NO ₂ is not generated because of assumed rapid decomposition back to reactants
Bi- and polyfunctional compounds	OH, O ₃ , NO ₃ , O ³ P, hv, as applicable	Estimates are generally much more uncertain and less reliable

109

110 contain zero to three hydrogen atoms and zero to three oxygen atoms. Structure designations are
111 summarized in Section A1.1 of the SI.

112 The generated reactions and estimated rate constants depend not only on the group(s) where the
113 reaction occurs but also on the immediate neighbor group and in some cases groups some distance away
114 in the molecule. In many cases, different groups or combination of groups are estimated to react similarly
115 or have similar effects on reactions at neighboring groups, so this documentation uses designations that
116 refer to such combinations. Examples include -CX₂H- to refer to carbon-centered groups with at least one
117 hydrogen (where "X" designates either H or any neighboring group bonded to the carbon with a single
118 bond), -CH_x-OH to any carbon-centered group bonded to an OH group. G refers to any group and -G-G-
119 to chains of groups, etc. Some group designations used in the text will be somewhat different than those
120 used by MechGen in order to be more familiar to chemists. Generally the group designations are noted in
121 cases where they may not be obvious.



122 Table 2. Summary of types of reactions supported by MechGen. "VOC" indicates any stable
123 compound supported by the system (see Table 1).

Reactant(s)	Type of Reactions
VOC + OH	H-atom abstraction Addition to double and triple bonds Addition to aromatic rings (OH only)
VOC + O ₃	Addition to double bonds followed by Criegee intermediate formation Excited adduct addition to amines, followed by decomposition forming OH
VOC + NO ₃	H-atom abstraction Addition to double bonds
VOC + O ³ P	Addition to double bonds
VOC + hv	Breaking the weakest bond in aldehydes, ketones, hydroperoxides, α -dicarbonyls, PAN compounds, and organic nitrates More rapid photolysis of dinitrates and carbonyl nitrates and carbonyl hydroperoxides
VOC Uni.	Unimolecular decompositions of peroxy nitrates
Carbon-centered Radicals	Unimolecular decompositions of radicals with α -nitro, α -nitrate, or α -peroxy groups Reaction with O ₂ with H-abstraction from α -OH groups Addition of O ₂ to radicals with allylic resonance Reactions of O ₂ with aromatic - OH adducts Addition of O ₂ to other alkyl radicals Unimolecular reactions of excited OH + alkene and aromatic adducts
Peroxy or Acyl Peroxy Radicals	Reactions with NO forming the corresponding alkoxy radical or organic nitrate Cyclization of aromatic OH-O ₂ adducts Unimolecular H-shift reactions forming hydroperoxides Reactions with NO ₂ forming the corresponding peroxy nitrate or PAN Reaction with HO ₂ forming the corresponding hydroperoxide or other products Reaction with NO ₃ forming NO ₂ and the corresponding alkoxy radical Reaction with the total of other peroxy or acyl peroxy radicals forming the corresponding alkoxy radical, carbonyl compound, or alcohol
Alkoxy Radicals	α -H abstraction by O ₂ forming the corresponding carbonyl compound β -scission decompositions H-shift isomerizations Ester rearrangement
Excited Criegee Intermediates	Decompositions, stabilization, or rearrangements of saturated biradicals Internal addition to the double bond of unsaturated biradicals, followed by O-O scission and epoxide formation



124 2. Initial Reactions of Organic Compounds

125 2.1. Bimolecular reactions with Atmospheric Oxidants


126 The methods MechGen uses to estimate rate constants for reactions of organic compounds with
127 OH, NO₃, O₃, O³P, and Cl have been documented by Carter (2021), with updates as discussed below and
128 in the SI. In most cases the rate constant estimates are made separately for each reaction route, with the
129 total rate constant being the sum of the estimated rate constants for each route. The estimates are made for
130 the various types of reactions using SARs based on the groups in the molecule, the neighboring groups
131 bonded to them (in some cases the 2nd neighbor as well), and the structure of the molecule around the
132 group(s) where the reaction occurs (e.g., groups in a ring). The equations and parameters used for
133 estimating the rate constants are summarized in Table 3 and the parameter values are given in various
134 tables in Section A1.2 of the SI. Most adjustable parameters needed for mechanism generation were
135 derived by optimizations to fit the measured OH, NO₃, O₃, and Cl rate constants tabulated by McGillen et
136 al. (2020), with the rate constants for most O³P reactions taken from Calvert et al. (2000). There were
137 insufficient data to derive all parameters using this approach, so some parameters were estimated based
138 on assumed relationships with the adjustable parameters.

139 The derivation or estimation of most of the parameters and the ability of the estimates to predict
140 the experimental data were given by Carter (2021). However, parameters for predicting rate constants for
141 reactions of phenols with OH and NO₃ were recently updated to improve the ability to model
142 environmental chamber experiments with generated mechanisms, as discussed in the SI. Rate constants
143 for most hydrocarbons and monofunctional compounds can be estimated to within ±30%, though
144 predictions are not as good for multifunctional compounds and predictions for ~15% of the rate constants
145 are off by more than a factor of 2. Estimates are more uncertain in the case of NO₃ and O₃ reactions
146 compared to OH, Cl, and O³P. This reflects the current state of the science in estimating these rate
147 constants with empirical structure-reactivity methods. Note that our estimates perform comparably to
148 those developed for the GECKO-A system by Jenkin et al. (2018a, b) for OH, Kerdouci et al. (2014) for
149 O₃, which were developed independently using similar methods (see Carter (2021)).

150 The following sections describe the mechanisms generated for the various types of reactions and
151 how branching ratios, if applicable, are derived. Note that some compounds may have more than one
152 reaction for a given oxidant, so branching ratios are estimated using ratios of estimated rate constants,
153 unless the branching ratios have been assigned explicitly for the compound and oxidant, as indicated in
154 the tables in Section A1.2 of the SI. If only the total rate constant is assigned, the branching ratios are
155 unaffected because they are derived from ratios of estimated rate constants. Note that additional
156 branching ratio estimates are needed for additions of O₃ or O³P to double or triple bond systems, as
157 discussed below.

158 2.1.1. H-Atom Abstractions by OH or NO₃

159 OH and NO₃ radicals can react with groups containing H atoms as follows, with rate constants
160 estimated as indicated in Table 3a and the parameters as indicated in Table A-3.





161 Table 3. Structure-reactivity equations and parameters used to estimate rate constants for reactions
 162 of VOCs with OH, NO₃, Cl, or O³P. Parameter values are given in the SI. Rate constants
 163 are per group unless otherwise indicated.


<p>a) <u>H-Atom abstractions from a group by OH or NO₃</u></p> $kH(\text{group, site, nbrs}) = kH^{\text{base}}(\text{group}) \times FH^{\text{ring}}(\text{site}) \times FH^{\text{nbrs}}(\text{nbrs, group})$ $FH^{\text{nbrs}}(\text{nbrs, group}) =$ <p style="margin-left: 20px;">If no more than one non-alkyl substituent $FH^{\text{nbr}}(\text{nbr, group})$</p> <p style="margin-left: 20px;">If $\prod_{\text{nbrs}} FH^{\text{nbr}}(\text{nbr, group}) \leq 1$ $\prod_{\text{HCnbrs}} FH^{\text{nbr}}(\text{nbr, group}) \times \prod_{\text{nHCnbrs}} FH^{\text{nbr}}(\text{nbr, group})$</p> <p style="margin-left: 20px;">If $\prod_{\text{nbrs}} FH^{\text{nbr}}(\text{nbr, group}) > 1$ $\prod_{\text{HCnbrs}} FH^{\text{nbr}}(\text{nbr, group}) \times \text{Avg}_{\text{nHCnbrs}} [FH^{\text{nbr}}(\text{nbr, group})]$</p> <p>$kH^{\text{base}}(\text{group})$: Base rate constant for abstraction from group $FH^{\text{ring}}(\text{site})$: Ring strain correction based on the smallest ring containing the group $FH^{\text{nbrs}}(\text{nbrs, group})$: Correction factors for all substituents on group (defined above)</p>
<p>b) <u>Addition to a group on isolated or conjugated double or triple bonds by OH or NO₃</u> (Addition to group G₁ on a bond defined by G₁=G₂ or G₁≡G₂; nC = number of carbons)</p> $kA(G_1, G_2, \text{site, nbrs}) = kA^{\text{base}}(G_1, G_2) \times FA^{\text{struct}}(\text{site}) \times \prod_{\text{nbrs}1} FA_1^{\text{nbr}}(\text{nbr}) \times \prod_{\text{nbrs}2} FA_2^{\text{nbr}}(\text{nbr})$ $FA^{\text{struct}}(\text{nC}) = \{1 - \exp(-\min(\text{nC}, 12) \times FS^{\text{radical}}) / \{1 - \exp(-4 \times FS^{\text{radical}})\}; \text{ where } FS^{\text{NO}_3} = 0.101$ <p style="margin-left: 40px;">and $FS^{\text{OH}} = FS^{\text{Cl}} = 0$ (FA^{struct} correction for NO₃ only)</p> <p>$kA^{\text{base}}(G_1, G_2)$: Base rate constant for addition to this group on the double or triple bond $FA^{\text{struct}}(\text{site})$: Structural correction factor for addition to this site (NO₃ only, see below) $FA_1^{\text{nbr}}(\text{nbr})$: Correction factor for each substituent on G₁ (other than G₂) $FA_2^{\text{nbr}}(\text{nbr})$: Correction factor for each substituent on G₂ (other than G₁)</p>
<p>c) <u>Additions to groups on aromatic rings by OH or NO₃</u></p> $kAro(\text{aGrp, aSubs}) = kAro^{\text{base}}(\text{aGrp}) \times FA^{\text{ipso}}(\text{aSub}_1) \times FA^{\text{ortho}}(\text{aSub}_2) \times FA^{\text{meta}}(\text{aSub}_3) \times$ $FA^{\text{para}}(\text{aSub}_4) \times FA^{\text{meta}}(\text{aSub}_5) \times FA^{\text{ortho}}(\text{aSub}_6)$ <p>$kAro^{\text{base}}(\text{aGrp})$: Base rate constant for addition to this aromatic group $FA^{\text{ipso}}(\text{aSub}_1)$: Correction for substituents on the same group where the addition occurs $FA^{\text{ortho}}(\text{aSub}_2)$, etc: Correction factors for ortho, meta, or para substituents on the aromatic ring</p>
<p>d) <u>Additions to non-aromatic double or triple bonds by O₃ or O³P (rate constant per bond)</u></p> $kA'(\text{bond, site, nbrs}) = kA'^{\text{base}}(\text{bond}) \times FA'^{\text{ring}}(\text{site}) \times FA'^{\text{struct}}(\text{site}) \times \prod_{\text{nbrs}} FA'^{\text{nbr}}(\text{nbr})$ $FA'^{\text{struct}}(\text{n}\beta) = 1 - (F\beta^{\text{O}_3} \times \{(\min[4, \max(1, \text{n}\beta)] - 1)\}, \text{ where } F\beta^{\text{O}_3} = 0.268, F\beta^{\text{O}_3\text{P}} = 0, \text{ and } \text{n}\beta \text{ is}$ <p style="margin-left: 40px;">the number of β substituents other than H</p> <p>$kA'^{\text{base}}(\text{bond})$: Base rate constant for addition to this type of bond (O₃ only) $FA'^{\text{ring}}(\text{site})$: Ring strain correction factor for addition to this site (O₃ only) $FA'^{\text{struct}}(\text{site})$: Correction factor used for branched structures (O₃ only, see below) $FA'^{\text{nbr}}(\text{nbr})$: Correction factor for each substituent on groups at ends of the bond</p>
<p>e) <u>Additions to the center group with cumulated double bonds by OH or NO₃, and Additions to the N in amino groups by OH, NO₃, O₃, or O³P</u></p> $kA''(\text{group, nbrs}) = kA''^{\text{base}}(\text{group, nbrs}) \text{ (substituent effects incorporated in } kA''^{\text{base}})$



164 Here "G" is any group where H is removed, "X" is OH or NO₃, and "aG" refers to groups in a resonance
 165 structure (bonded to alternating double and single bonds). Separate estimates are made for each H-
 166 containing group in the molecule, so the branching ratio can be derived from the ratio of estimated rate
 167 constants to the sum of all the estimated rate constants for the VOC, including addition reactions (see
 168 below) where applicable. The subsequent reactions of the carbon-centered radicals are discussed in
 169 Section 3.

170 2.1.2. Additions of OH or NO₃ to Groups with Unsaturated Bonds


171 The OH and NO₃ radicals can also add to groups with double, triple, or aromatic bonds as
 172 follows, with rate constants estimated as indicated in Table 3b for alkenes and alkynes and Table 3c for
 173 aromatics. The parameters are given in the SI as indicated in Table A-3.



178 Here "G" refers to any group with an unsaturated bond, "X" refers to the reacting radical, "{excited}"
 179 indicates vibrational excitation, and "*" indicates ring closure. Separate estimates are made for addition to
 180 each group so these can also be used to make the branching ratio estimates. Most of the excited carbon-
 181 centered radicals are stabilized and react with O₂ as discussed in Section 3.1, but some have fast
 182 unimolecular reactions, as discussed in Section 3.3 for radical + alkene and alkyne adducts and Section
 183 3.4 for radical and aromatic adducts.

184 2.1.3. Additions of OH or NO₃ to Amines

185 Both OH and NO₃ radicals are known to react rapidly with amines (McGillen et al., 2020; Carter,
 186 2021), but the fact that the reaction with tertiary amines is about as rapid as reaction with secondary or
 187 primary amines indicates that the reaction is due to initial addition of the radical to the amino group,
 188 rather than direct H-abstraction from the amino group [see discussion in Section S1.9 of Carter (2021)].
 189 Based on this, MechGen assumes that the rate-determining step is addition of the radical to the amino
 190 group forming an addition complex that rapidly decomposes. An example is shown below.




191 In the case of tertiary amines, where reactions analogous to (1) are not possible, the adduct is assumed to
 192 decompose by abstraction of an α hydrogen if one is present, forming the same products that would be
 193 formed if the radical abstracted from the α hydrogen directly. Both decomposition modes are possible for
 194 primary or secondary amines with α hydrogens, but the formation of the α -amino carbon-centered radical
 195 (analogous to 2) is estimated to be more exothermic and is assumed to dominate. If there is no α
 196 hydrogen, then only the reaction forming the amino radical (analogous to 1) is generated.

197 Neither of these decomposition modes are possible for tertiary amines with no α hydrogens (e.g.,
 198 tri-*t*-butyl amine), so MechGen assumes that the adduct decomposes back to reactants and there is no net
 199 reaction at the amino group. We are aware of no measurements or calculations of rate constants for these
 200 reactions, but they would be interesting to study.



201 2.1.4. Additions of O₃ to Unsaturated Bonds

202 O₃ is assumed to react only by additions to double or triple bond systems as follows, with rate
 203 constants estimated as indicated in Table 3d and with parameters as indicated in Table A-3. Reactions of
 204 O₃ with aromatics or cumulated double bonds are assumed to be negligible based on low measured rate
 205 constants (McGillen et al., 2020). Note that only total rate constants for additions to unsaturated bond
 206 systems are estimated, so it is necessary to assume branching ratios for reactions of the initially formed
 207 adducts or, in the case of additions to conjugated bond systems, for the initial reaction. The branching
 208 ratios used are shown in brackets where applicable.





209 Here α and $(1-\alpha)$ are branching ratios assigned for the additions to separated double bonds, $G(=O)$ refers
 210 to a carbonyl ($-CO-$ or $-CHO$) group, and $G[OO] \{\text{excited}\}$ refers to an excited Criegee intermediate (CI)
 211 whose subsequent reactions are discussed in Section 7, and "*" indicates rings. The additions to double
 212 bonds are assumed to proceed via initial formation of excited primary ozonides, which decompose to the
 213 products as shown above; while additions to alkynes are assumed to form the excited CIs directly. In most
 214 cases it is assumed that formation of all the possible Criegee and carbonyl combinations are equally
 215 likely, as indicated above. However, following the recommendations of Jenkin et al. (2020), we estimate
 216 that $\alpha = 0.1$ if G_3 is $-HCO-$, $-CO-$, $-O-$, or $-OH$ and no such group is bonded to G_1 , or 0.5 otherwise (or if
 217 G_3 is absent). In other words, formation of the carbonyl bonded to the oxygenated group is favored.

218 In the case of alkynes, the primary ozone adduct is assumed to directly form excited CIs with α -
 219 carbonyl substituents, with the two possible CIs formed in equal yields. However, it is assumed that the
 220 intermediates can rapidly interconvert by O shifting from the Criegee to the carbonyl group, so reactions
 221 of both of the isomers in this equilibrium need to be considered. The Criegee reactions are discussed in
 222 Section 7.

223 2.1.5. Additions of O³P to Unsaturated Bonds

224 O³P is assumed to react only by additions to double bonds as follows, with rate constants
 225 estimated as indicated in Table 3d for isolated or conjugated double bonds and in Table 3e for additions to
 226 cumulated double bonds. As with O₃ additions, it is necessary to assign branching ratios for some of the
 227 reactions, and these are shown to the right in brackets.





228 Here G^H refers to a group with the H removed, with the reaction only being possible if group G has an H.


229 Addition reactions involving O^3P are expected to initially form a highly excited oxirane
 230 compound or biradical, which can stabilize, rearrange, or decompose to radicals. Based on model
 231 simulation results of high NO_x environmental chamber experiments performed when developing the
 232 SAPRC mechanisms for alkenes (e.g., Carter, 2000, 2010b), it is assumed that radical formation from
 233 reactions of O^3P with C_3+ organics are negligible, so only stabilization or rearrangements forming stable
 234 compounds are assumed to occur (as shown above). This is assumed for conjugated alkenes and alkynes
 235 as well, though this assumption is more uncertain.

236 The only type of rearrangement considered for the initially formed adducts are H-shifts from one
 237 carbon to another, forming a carbonyl compound. In the case of additions to double bonds, it is assumed
 238 that stabilization occurs 50% of the time if at least one rearrangement is possible (i.e., $\alpha_1=\alpha_4=0.5$,
 239 $\alpha_2+\alpha_3=\alpha_5=0.5$) and all of the time if it is not (i.e., $\alpha_1=\alpha_4=1$ and $\alpha_2=\alpha_3=\alpha_5=0$). If more than one
 240 rearrangement is possible, they are assumed to be equally likely ($\alpha_2=\alpha_3=0.25$). In the case of additions to
 241 triple bonds, it is assumed that stabilization (not shown) is unlikely, and that there is no net reaction if the
 242 rearrangement shown is not possible ($\alpha_6=0$) and no back decomposition of the adduct otherwise ($\alpha_6=1$).

243 These assumptions are uncertain, but they have relatively little effect on model simulations of
 244 most atmospheric systems because reactions of O^3P are generally not important unless NO_2 is very high,
 245 such as in high NO_x environmental chamber simulations or in combustion/biomass burning plumes.

246 2.1.6. Reactions of O_3 and O^3P with Amines

247 The reactions of amines with O_3 or O^3P are assumed to form excited N-oxides, with rate constants
 248 estimated as shown in Table 3e, and parameters for other addition reactions as indicated in Table A-3.



249 In the case of primary or secondary amines, these excited N-oxides are assumed to rapidly rearrange and
 250 then stabilize to form -N-OH compounds ($\alpha_7=1$). In the case of tertiary amines, it is assumed that the N-
 251 oxide is stabilized ($\alpha_7=0$). The current system does not handle reactions of stabilized N-oxides so it treats
 252 them as unreactive. Their reactions and mechanisms are unknown.



253 2.2. Photolysis Reactions

254 Compounds with the groups -CHO, -CO-, -ONO₂, or the group pairs -O-OH, -O-O- are expected
255 to undergo photolysis at non-negligible rates under lower tropospheric conditions. These groups or group
256 pairs are referred to as "photoreactive groups" hereafter. Compounds not containing these photoreactive
257 groups may photolyze with UV wavelengths less than the atmospheric cutoff of ~290 nm, but photolyses
258 at this low wavelength regime are not currently considered in MechGen. Photolysis of radicals and CIs
259 are also ignored because it is assumed that loss of these intermediates by other reactions will dominate
260 over photolysis under atmospheric conditions.

261 Rate constants for photolysis reactions depend on the compound, reaction, and lighting
262 environment, and can be calculated by:

$$263 \quad J(\text{reaction, environment}) = J(\text{photolysis set, environment}) \times \phi_{\text{overall}}(\text{reaction})$$
$$264 \quad J(\text{photolysis set, environment}) = \sum_{\lambda} \sigma_{\lambda}(\text{photolysis set}) \times \phi_{\lambda}(\text{photolysis set}) \times F_{\lambda}(\text{environment})$$

265 where J is the first order rate constant for a photolysis reaction in a particular lighting environment,
266 "photolysis set" is a set of wavelength-dependent absorption cross-sections and quantum yields assigned
267 to various types of reactions, ϕ_{overall} is an overall wavelength-dependent quantum yield used for the
268 reaction of interest; if needed, λ is the wavelength, σ_{λ} and ϕ_{λ} are the wavelength-dependent absorption
269 cross-sections and quantum yields at wavelength λ in the photolysis set, and F_{λ} is the intensity of the light
270 source at wavelength λ , which is a function only of the environment, not the reaction.

271 MechGen makes no assumptions about the light source (other than having no significant intensity
272 at wavelengths below ~290 nm) and therefore does not output estimated or assigned photolysis rate
273 constants when generating photolysis reactions. Instead, it outputs the name of the photolysis set and the
274 value of overall quantum yield (ϕ_{overall}), if applicable, assigned to the reaction. The photolysis sets
275 assigned by MechGen are a subset of those incorporated in the SAPRC-16, -18 and -22 mechanisms
276 (Carter, 2016, 2020, 2023a), and are summarized in the SI on Table A-5. That table also indicates how
277 they were derived and gives half lives (with $\phi_{\text{overall}}=1$) at direct overhead sun using the actinic fluxes in the
278 reactivity scenarios of Carter (1994), and orders them by increasing half life. The files containing the data
279 for these photolysis sets are available online at the SAPRC website (Carter, 2023a, b). Photolysis rate
280 constants can be calculated using these data and are required for model application of mechanisms
281 generated using MechGen. Photolysis rate constants should be calculated for the light conditions of the
282 experimental or atmospheric conditions of interest. The mechanisms, photolysis sets, and (where
283 applicable) overall quantum yields for the various types of photolysis reactions are summarized below.
284 The photolyses of compounds with only a single photolysis group are described first, followed by a
285 discussion of treatments of photolyses of compounds with more than one such group. Note that if there is
286 more than one reaction, the branching ratios shown are implemented as overall wavelength-independent
287 quantum yields and are assumed not to depend on the spectrum of the light source.

288 In most cases the estimated mechanisms, absorption cross-sections, and quantum yields for
289 compounds with photoreactive groups are based on those derived for the smallest C₃₊ or C₄₊ compound
290 with the photoreactive group(s), with all the other groups being alkyl (-CH_x-) groups. This is clearly an
291 oversimplification, especially since the structure and size of the molecule, and the presence of non-alkyl
292 substituents, can affect the absorption cross-sections or quantum yields for at least some of the different
293 types of photoreactive compounds (Calvert et al., 2011). Although MechGen currently does not take these
294 differences into account, this needs to be considered in future versions.




295 **2.2.1. Compounds with a Single Aldehyde Group**

296 Aldehydes with a single -CHO group that are not directly bonded to any unsaturated or
 297 photoreactive group, other than acetaldehyde and glycolaldehyde for which photolysis assignments are
 298 given explicitly (see SI), are assumed to photolyze as follows:



300 where "R" is any carbon-centered group other than methyl that does not contain a double bond on the
 301 group next to the -CHO. The absorption cross sections used are those recommended by IUPAC (2023) for
 302 propionaldehyde, and are given in photolysis set "C2CHOabs". IUPAC makes no specific
 303 recommendations concerning quantum yields or branching ratios, but it appears likely that the quantum
 304 yield $\cdot\text{CHO}$ formation for $\lambda > \sim 295$ nm is near unity, based on the data of Chen and Zhu (2001). MechGen
 305 assumes that this is applicable for higher aldehydes of this type as well.


306 Aldehydes where the -CHO group is bonded to a C=C double bond group are assumed to
 307 photolyze with a mechanism based on that derived for acrolein, which is taken as representative for all α -
 308 unsaturated aldehydes, including those with conjugated or cumulated double bonds, and with non-alkyl
 309 (but non-photoreactive) substituents. The photolysis set giving the absorption cross sections and quantum
 310 yields used for acrolein is ACROL-16, while MACR-15 (based on methacrolein) is used for the others,
 311 though the acrolein mechanism is used in all cases to predict the subsequent reactions that occur. This
 312 gives half lives of ~ 50 and ~ 60 hours, respectively, for loss by photolysis for direct overhead sunlight,
 313 which means that photolysis is relatively slow but not necessarily negligible. The assumed mechanism is
 314 as follows:



315 where $x=0-2$, $x'=0-1$, and $[\cdot]$ indicates a carbene radical group, whose subsequent reactions are discussed
 316 in Section 6.3. These branching ratios are based on the data of Gardner et al. (1987) for acrolein, as
 317 shown on Figure IX-C-7 of Calvert et al. (2011) for air at 1 atm. These are assumed to apply to photolyses
 318 of other unsaturated aldehydes as well.

319 Aldehydes where the -CHO is bonded to aromatic groups are assumed to photolyze with the same
 320 rate constant and a similar mechanism as estimated for benzaldehyde, regardless of substituents. The
 321 photolysis set used is "BALD-11", where absorption cross-sections are those recommended for
 322 benzaldehyde by Calvert et al. (2002). In addition, $\phi_{\text{overall}} = 0.06$ is used to approximately fit consumption
 323 rates of benzaldehyde in chamber experiments (Carter, 2000).

324 Benzaldehyde acts as an efficient radical terminator when added to environmental chamber
 325 experiments (Carter et al., 1982; Carter, 2000), suggesting that photolyses of aromatic aldehydes do not
 326 form radicals. For mechanism generation purposes, these are represented as follows:



328 where $\text{-aC}_x\text{-}$ refers to either -aC- or -aCH- and "*" indicates ring closure. (The reaction with OH radicals
 329 is also predicted to be radical terminating, but that would not be sufficient to compensate for radical
 330 initiation due to photolysis if the quantum yield for radical production were non-negligible.) The actual
 331 photolysis reaction is highly uncertain, and it is likely that other products are actually formed, since we
 332 know of no evidence that aromatic hydrocarbons are formed in high yields from the photolyses of
 333 aromatic aldehydes.



334 Aldehydes where the -CHO is bonded to an alkyne group, e.g., -C≡C-CHO, are assumed not to
 335 photolyze. We could not find data concerning absorption cross-sections or photolyses of these
 336 compounds, and assume that they either do not photolyze, or do so relatively slowly. Data are needed to
 337 test this assumption.

338 2.2.2. Compounds with a Single Ketone Group


339 C₄₊ ketones with a single -CO- group that is not directly bonded to any unsaturated or
 340 photoreactive group are assigned photolysis sets giving absorption cross-sections and overall quantum
 341 yields as indicated in Table 4. In the case of acetone, the wavelength-dependent quantum yields
 342 recommended by IUPAC (2023) for atmospheric temperature and pressure are used. In the case of the
 343 higher ketones, the photolysis set only has absorption cross-sections, and overall quantum yield factors
 344 are derived so that the SAPRC mechanism simulates environmental chamber data for ketones (Carter,
 345 2010b). These data suggest that the photolysis rate constants, and therefore overall quantum yield factors,
 346 decrease with carbon number for C₄₊ ketones. The structure of the ketone and the presence of non-alkyl
 347 but non-photoreactive substituents, other than -O- groups next to the carbonyl (i.e., esters), are assumed
 348 not to affect the absorption cross-sections or quantum yields.

349
 350 Table 4. Assignments used to estimate photolysis rate constants for compounds with single ketone
 351 groups based on the numbers of carbons in the ketone.

C's	Photolysis Set	ϕ_{overall}	$t_{1/2}$ [a]	Based on Data for Compound
3	ACET-06 (σ_{λ} and ϕ_{λ} for acetone)	1	~180 hours	Acetone
4	MEK-06 (σ_{λ} 's for MEK)	0.175	~70 hours	methyl ethyl ketone (MEK)
5	Same as above	0.1	~120 hours	2-pentanone
6	Same as above	0.1	~120 hours	methyl isobutyl ketone
7+	Same as above	0.02	~600 hours	2-heptanone

352 [a] Half life relative to the photolysis rate constant for ketone photolysis calculated for direct overhead
 353 sunlight ($z=0$) using the actinic fluxes used in the reactivity scenarios of Carter (1994).
 354
 355

356 The photolysis of these ketones are assumed to proceed only by breaking one of the two C-CO
 357 bonds, forming an acyl and an alkyl radical.



358 Here G is any non-photoreactive group, and α_1 is the branching ratio for the formation of radical $G_1\cdot$. The
 359 relative branching ratios for breaking CO-alkyl bonds are assumed to depend on whether the radical
 360 formed is a methyl, secondary, primary, or tertiary, with equal branching ratios if both radicals are of the
 361 same type. Based on IUPAC (2023) data for methyl ethyl ketone, we assume methyl vs. ethyl branching
 362 ratios of 15% and 85%, respectively, for that compound. Extrapolating this to other higher compounds,
 363 we use the following branching ratios, where $\alpha=0.5$ is used if both are of the same type:

\underline{G}_1	\underline{G}_2	α_1	\underline{G}_1	\underline{G}_2	α_1	\underline{G}_1	\underline{G}_2	α_1
-CH ₂ -	-CH ₃	0.85	-CH()-	-CH ₃	1	-C()-	-CH ₃	1
-CH()-	-CH ₂ -	0.85	-C()-	-CH ₂ -	1	-C()-	-CH()-	0.85

364 Note that this is somewhat arbitrary and uncertain (except for methyl ethyl ketone), and is assumed not to
 365 be affected by the presence of non-alkyl substituents, which is even more uncertain.



366 If the ketone group is in a ring, the reaction would form excited biradicals that should re-form the
367 starting ketone if stabilized. In this case, it is assumed that the excited biradical primarily decomposes by
368 CO elimination, i.e.:




370 where "*" indicates closure for any size ring.

371 Esters, carbonates, and acids, i.e., compounds with -CO-O- or -CO-OH group combinations, do
372 not appear to have significant absorption in the atmospheric wavelength region, and are assumed not to
373 photolyze.


374 Estimations of photolyses of ketones with an adjacent double bond group are based on data for
375 methyl vinyl ketone (MVK). These reactions are assigned the photolysis set "MVK-16", which contains
376 IUPAC (2023) recommendations for absorption cross-sections and quantum yields of methyl vinyl ketone
377 at 1 atm. This gives an estimated half life of ~36 hours for photodecompositions of these unsaturated
378 ketones with direct overhead sunlight, and gives reasonably good fits to NO oxidation rates in MVK -
379 NO_x chamber experiments.

380 The photolysis reactions for ketones with only a single adjacent double bond are assumed to be as
381 follows:



382 Here, $x=0-2$, $x'=0-1$, and G is any non-photoreactive group with only saturated bonds. This is based on
383 IUPAC (2023) recommendations for MVK but is assumed to apply to all compounds of this type.

384 Ketones with double bond groups on both sides of the carbonyl group are estimated to photolyze
385 with the same absorption cross-sections and quantum yields as those bonded to only one double bond
386 group, but it is assumed that formation of vinylic radicals is not as favorable based on thermochemical
387 estimates. These reactions are assumed to occur only via CO formation, as shown in the 2nd reaction
388 above. This is also assumed for unsaturated ketones where the -CO- group is in a ring, where the first
389 reaction is not expected to occur for the reasons discussed above.



392 Estimations of photolyses of ketones with an adjacent aromatic or triple bond group are not
393 supported by MechGen, so no photolysis reactions are generated for these compounds. The need to
394 represent photolyses of these compounds has not been investigated, but it is expected that
395 photodecompositions of aromatic ketones will be relatively slow, and ketones with triple bonds are not
396 expected to be important in atmospheric systems.


397 **2.2.3. Organic Nitrates, Carbonyl Nitrates and Carbonyl Peroxynitrates**

398 Compounds with a single organic nitrate group and no other photoreactive groups are assumed to
399 photolyze entirely by scission of the weak O-NO₂ bond, forming NO₂ and an alkoxy or acyloxy radical
400 whose reactions are discussed in Section 5.



402 These reactions are assigned the "IC3ONO2" photolysis set with quantum yields of 1 at all wavelengths,
 403 which gives a half life of ~50 hours for loss due to photolysis with direct overhead sunlight. This is based
 404 on the IUPAC (2023) recommendation for isopropyl nitrate, but is assumed to apply to all organic nitrates
 405 without other photoreactive groups, regardless of structure, the presence of non-alkyl substituents, and the
 406 presence of unsaturated bonds. This is an approximation because isopropyl nitrate has stronger
 407 absorptions than for n-propyl and smaller nitrates, but is taken as representative. Although the accuracy of
 408 this approximation is uncertain in some cases, loss by photolysis is relatively unimportant compared to
 409 other reactions.

410 In the case of acyl peroxy nitrates (e.g., PANs), the photolyses are assigned "PAN-11" or
 411 "PPN-11" photolysis sets, with the former used for PAN itself and the latter used for all other PAN
 412 analogues based on peroxypropionyl nitrate (PPN). These contain only absorption cross-sections, since
 413 unit quantum yields are assumed. They give approximate photodecomposition half lives of ~200 and
 414 ~100 hours, respectively, for direct overhead sunlight. The assumed mechanism is as follows:



417 This is based on Sander et al. (2006) recommendations for PAN, but is applied to all PAN analogues
 418 regardless of substituents or the presence of multiple bonds.


419 In the case of non-acyl (e.g., alkyl) peroxy nitrates, photolysis is assumed to proceed primarily by
 420 NO_2 formation,



422 with the same rate constant assignments as used for simple organic nitrates. However photolysis is not
 423 important for these compounds because they are assumed to undergo rapid thermal unimolecular
 424 decomposition, forming the same products, as discussed in Section 2.3.

425 **2.2.4. Compounds with a Single set of Peroxide Groups**

426 Compounds with peroxy or hydroperoxy groups and no other photoreactive group are assumed to
 427 photolyze entirely by scission of the weak O-O bond, forming alkoxy radicals and (for hydroperoxides)
 428 OH.



431 The reactions are assumed to have unit quantum yields at all wavelengths, and are assigned the photolysis
 432 set "COOH", which contains the absorption cross-section of methyl hydroperoxide (IUPAC, 2023), and
 433 gives a half life of ~30 hours for loss by photolysis with direct overhead sun. Note that it is assumed that
 434 organic peroxides (which tend to be relatively less important than hydroperoxides in atmospheric
 435 systems) photolyze with approximately the same rate constant as hydroperoxides, and that the size,
 436 structure, and presence of non-photoreactive substituents or unsaturated bonds do not have a significant
 437 effect. The most questionable assumption in this case is the assumption that the presence of adjacent
 438 unsaturated bonds has no effect on hydroperoxide or peroxide photolysis.



439 2.2.5. **Compounds with More than One Photoreactive Group**

440 The photolysis sets, quantum yields, and photolysis reactions used for compounds with various
441 combinations of photoreactive groups are summarized in Table 5. If the ϕ_{overall} column is blank, then no
442 overall quantum yield correction is applied and the photolysis rate constant is the same as that calculated
443 for the photolysis set. The half lives relative to photodecomposition for direct overhead sun, which
444 incorporate the overall quantum yield corrections if applicable, are also shown in the table. No corrections
445 are made for non-alkyl substituents or unsaturated bonds on the molecule except as indicated in the table.
446 The considerations used when making these assignments are discussed below.

447 The assignments for some (but not all) types of reactants with adjacent photoreactive groups are
448 made based on available data for the simplest or representative molecule of the types, i.e., methyl glyoxal,
449 biacetyl, peroxy acetic acid, PANs, carbonyl nitrates, dinitrates, and various unsaturated 1,4-dicarbonyls.
450 The applicability to higher molecular-weight compounds, compounds with non-alkyl substituents, or
451 compounds with multiple bonds is uncertain, but not as uncertain as the assignments for the reactants in
452 the table where data for representative compounds are either not available or were not considered.

453 It is assumed that -CO-O- groups are not photoreactive, so the presence of this group combination
454 is not considered when making photolysis assignments. The exception is compounds with -CO-O-ONO₂
455 group combinations (PAN analogues), where assignments are made based on data for PAN and
456 peroxypropynyl nitrate (PPN) as discussed above.

457 The unsaturated 1,4-dicarbonyls with at least one aldehyde group are highly photoreactive
458 aromatic ring opening products (Calvert et al., 2011; see also Section 3.4, below) that account for the
459 relatively high reactivity of aromatic hydrocarbons observed in environmental chamber experiments. The
460 AFGS photolysis file is used to give absorption cross sections for all these compounds, with an overall
461 wavelength-independent quantum yield adjusted to give satisfactory fit to results of aromatic - NO_x
462 chamber experiments, which are highly sensitive to this photolysis (e.g., Carter, 2010b; Carter and Heo,
463 2013). The SAPRC-22 mechanism (Carter, 2023a) that was derived using this version of MechGen best
464 fit the data using effective quantum yields close to 25%. However, the quantum yields in the AFGS file
465 do not take into account absorption cross sections reported for 4-oxo-2-pentenal reported by Xiang et al.
466 (2007), which are the basis of the current IUPAC (2023) recommendations for this compound. This is
467 because these quantum yields, which are much lower than the previous data for these compounds, could
468 not be used as a basis for a predictive mechanism for aromatics because fitting the chamber data required
469 using an effective quantum yield of greater than 3. Therefore, either there are additional sources of
470 radicals in aromatic photooxidation systems that are not taken into account in the current estimated
471 mechanisms for aromatics, the absorption cross sections of these 1,4-dicarbonyls are highly dependent on
472 structure, or the data of Xiang et al. (2007) have systemic issues. Until this situation is resolved and
473 MechGen can be updated to derive a more predictive mechanism, we will continue to use the present
474 cross sections in AFGS to derive aromatics mechanisms for airshed models.



475 Table 5. Summary of photolysis sets, overall quantum yields, approximate photolysis half lives,
476 and products formed for compounds with more than one photoreactive group.

Reactant [a]	Photolysis Set [b]	ϕ_{overall}	$t_{1/2}$ [c]	Products Formed [d]
<u>Adjacent photoreactive groups</u>				
HCO-CO-G-	MGLY-13 (methyl glyoxal)		40 min.	HCO[·] + G-CO[·]
-G ₁ -CO-CO-G ₂ -	BACL-11 (biacetyl)		25 min.	G ₁ -CO[·] + G ₂ -CO[·]
-G-CO-ONO ₂	IC3ONO2 (isopropyl nitrate σ 's)		50 hrs.	G-CO[O.] + NO ₂
-G-CO-O-OH	PAA (peroxy acetic acid σ 's)		200 hrs.	G-CO[O.] + OH
-G ₁ -CO-O-O-G ₂ -	COOH (methyl hydroperoxide σ 's)		30 hrs.	G ₁ -CO[O.] + -G ₂ O[·]
-G-CO-O-ONO ₂	PPN-11 (PPN σ 's) [e]		200 hrs.	See Section 2.2.3
<u>Separated photoreactive groups (x>0, no upper limit)</u>				
HCO-G ₁ =G ₂ -CHO	AFGS (various 1,4-unsaturated dicarbonyl σ 's, see text)	0.45	4 min.	Gs ₁ -CO[·] + H., & Gs ₂ -CO[·] + H. (equal)
HCO-(G) _x -CHO [f]	C2CHO (propionaldehyde)		8 hrs.	Same as -Gs-CHO
HCO-(G) _x -CO-	C2CHO (propionaldehyde)		8 hrs.	Same as -Gs-CHO
HCO-G ₁ =G ₂ -CO-G ₃ .	AFGS (see above)	0.45	4 min.	Gs ₂ -CO[·] + G ₃ [·]
HCO-(G) _x -ONO ₂	C2CHOabs (propionaldehyde σ 's)		3 hrs.	Same as -Gs-ONO ₂
xCO-G ₁ =G ₂ -(G) _x -O-OX	HPALDS (acroleins σ 's, see text)	0.1	3 hrs	Same at -Gs-O-OX
HCO-(G) _x -O-OX	C2CHOa (propionaldehyde σ 's)		3 hrs.	Same at -Gs-O-OX
HCO-(G) _x -O-ONO ₂	C2CHOa (propionaldehyde σ 's)		3 hrs.	Same as -Gs-O-ONO ₂
G ₁ -CO-G ₂ =G ₃ -CO-G ₄ -	Does not photolyze (see text)			
-G ₁ -CO-(G) _x -CO-G ₂ - [f]	MEK-06 (methyl ethyl ketone [MEK] σ 's)	6 C's 0.1 7+ C's 0.02	70 hrs. 120 hrs.	Same as G ₁ -CO-Gs-, & G ₂ -CO-Gs- (equal)
-G ₁ -CO-(G) _x -ONO ₂	CRBNIT (carbonyl nitrate σ 's)		4 hrs.	Same as -Gs-ONO ₂
-G-CO-(G) _x -O-OX	MEK-06 (MEK σ 's)		12 hrs.	Same at -Gs-O-OX
-G-CO-(G) _x -CO-ONO ₂	CRBNIT (carbonyl nitrate σ 's)		4 hrs.	Same as Gs-CO-ONO ₂
-G-CO-(G) _x -CO-O-ONO ₂	CRBNIT (carbonyl nitrate σ 's)		4 hrs.	Same as Gs-CO-O-ONO ₂
X ₁ O-O-(G) _x -O-OX ₂	COOH (methyl hydroperoxide σ 's)		30 hrs.	Same as Gs-O-OX ₁ + Gs-O-OX ₂ , each equal

477 [a] Group combinations that give unique compounds for which specific assignments are used (e.g.,
478 glyoxal and HNO₃) are not shown. "-OX" is -O- or -OH. "xCO-" is HCO- or -CO-.

479 [b] The derivations of the photolysis sets are given in Table A-5 in the SI. " σ 's" means that the photolysis
480 set contains only absorption cross-sections. "(equal)" means that both possible reactions occur with
481 equal probability.

482 [c] Approximate half lives relative to loss of the compound by photolysis in direct sunlight, including the
483 contribution by the assigned wavelength-independent quantum yield, if applicable.

484 [d] "Gs" refers to a combination of other groups on the molecule (other than G, G₁ or G₂), including the
485 other photoreactive group, which is assumed not to be transformed by the reaction.

486 [e] The photolysis set "PAN-11", giving an approximate photolysis half live of 100 hours for direct
487 overhead sun, is used for PAN itself (G = CH₃).

488 [f] Not applicable if -(G)_x- is -CH_x=CH_x', as shown for previous reactant.



489 It is assumed that unsaturated 1,4-diketones do not photolyze sufficiently rapidly in ambient
490 sunlight that it is necessary to include their photolysis in atmospheric models. This is consistent with the
491 discussion of photolyses studies of 3-hexene, 2,5-dione given by Calvert et al. (2011), which indicate that
492 radical formation is relatively minor compared to isomerization. This is also consistent with the fact that
493 para-substituted dialkyl benzenes, whose reactions can form these compounds, tend to be less reactive
494 than other isomers, all else being equal (Carter and Heo, 2013).

495 If a reactant has more than one non-adjacent photoreactive group of the same type it is assumed
496 that the photolysis reaction at each group occurs with equal probability, with the mechanism the same as
497 if there were no other photoreactive group. However, the total photolysis rate constant for reactions at
498 both groups is estimated to be the same as if the compound had only one group since the same photolysis
499 set is used. This is uncertain.


500 If a reactant has more than one non-adjacent photoreactive group of different types it is assumed
501 that the absorption cross-sections can be approximated by those of compounds with the single group that
502 has the highest absorption cross-section at atmospheric wavelengths. On the other hand, the mechanisms
503 (e.g., photolysis products) are assumed to be those resulting from reaction at the group with the weakest
504 bond. This is based on the assumption that once the photon is absorbed the energy is rapidly distributed
505 around the molecule. This is consistent with the data of Wolfe et al. (2012), which suggest that α -
506 unsaturated carbonyls with hydroperoxide groups photolyze at rates consistent with those calculated using
507 absorption cross-sections of α -unsaturated carbonyls, but with unit quantum yields and with the reaction
508 breaking the peroxy bond forming OH. This gives a photolysis rate that is about 100 times faster than
509 simple hydroperoxides. However isoprene and 1,3-butadiene NO_x experiments are not well simulated
510 with this high of a photolysis rate for α -unsaturated hydroperoxy carbonyls (unpublished results), so we
511 arbitrarily cut the rate by a factor of ~ 10 by using an effective quantum yield of 0.1 for these compounds.
512 The current system assigns unit quantum yields for saturated hydroperoxy or nitrate-substituted
513 carbonyls, though this is highly uncertain and the estimates are probably upper limits.

514 Despite the differences in assigned overall quantum yields, the photolysis rate constants
515 calculated for saturated carbonyls with other photoreactive groups are about the same as for those that are
516 unsaturated (see Table 5). This is due to the higher absorption cross-sections assigned to unsaturated
517 aldehydes.

518 2.3. Unimolecular Reactions

519 2.3.1. Peroxynitrates

520 Peroxynitrates are formed from the reactions of peroxy or acyl peroxy radicals with NO_2 , but the
521 O-ONO_x bond is weak enough that these can thermally decompose back to reactants at significant rates at
522 atmospheric temperatures.




523 Here, "G" is any saturated or unsaturated group and the rate constants are in $\text{cm}^3 \text{ molec}^{-1} \text{ s}^{-1}$, T is the
524 temperature in K, and $R = 1.99 \times 10^{-3} \text{ deg K mole kcal}^{-1}$. The presence of non-alkyl substituents,
525 unsaturated bonds or other structural factors are assumed not to affect these rate constant estimates or
526 decomposition mechanisms. The OO-NO₂ bonds are currently believed to be the only bonds in
527 compounds formed in atmospheric systems to be weak enough to undergo simple scission at non-
528 negligible rates at atmospheric temperatures. (The weak O-O bonds in peroxy compounds are at least 10
529 kcal/mole stronger.)



530 The decompositions of the non-acyl peroxy nitrates are estimated to have a half life of $\sim 5 \text{ sec}^{-1}$ at
531 298K, which is sufficiently fast that the formation of these compounds is rapidly reversed. Therefore, the
532 default is for MechGen to ignore the formation and therefore the decomposition of these compounds
533 when multi-step mechanisms are generated for atmospheric conditions. However, they do show up when
534 single step mechanisms are generated (Carter, In Prep.). On the other hand, the acyl peroxy nitrates
535 (PANs) decompose much slower and are observed as photooxidation products in atmospheric systems, so
536 their formation cannot be ignored. It cannot be ruled out that some α substituents other than -CO- may
537 slow down the decomposition sufficiently that peroxy nitrates for some non-acyl peroxy nitrates
538 may be non-negligible. However, this is not currently considered in MechGen.

539 2.3.2. Aromatic formation from Cyclohexadienones

540 The other type of non-radical unimolecular reaction that is currently considered involves
541 formation of aromatic rings from 2,4-cyclohexadienone compounds containing a -CHX-CO- group pair
542 in the ring, where an H-shift from the -CHX to the carbonyl group is expected to be highly favorable
543 energetically.



545 MechGen predicts that cyclohexadienones are formed in high yields in the reactions of phenolic
546 compounds if this reaction does not occur (see Section 3.4), but they are not observed as products. On the
547 other hand, catechols (the products of this type of isomerization reaction) are observed in high yields in
548 the reactions of OH with phenols (e.g., Olariu et al., 2002; Berndt and Böge, 2003), which would not be
549 expected if this reaction did not occur. The nature of the "X" group should not have an obvious effect on
550 how fast this reaction occurs, so if this is fast then it is reasonable to expect compounds with other "X"
551 substituents, including X=alkyl or H, are also fast. MechGen assumes that this is the case. The rate
552 constant is unknown but it is assumed to be high in order to account for the observed formations of
553 catechols from phenols. This assumption is implemented by treating such compounds like rapidly
554 reacting radicals when the reactions are generated.

555 It is possible that this reaction may require the presence of surfaces and/or water to occur at
556 sufficient rates. The H-shift involves a strained 4-member ring transition state, which may be catalyzed by
557 surfaces, while the involvement of water would result in the same H-shift but with a 6-member ring
558 transition state. Given surface- or water-free environments rarely occur in the ambient atmosphere it is
559 probably better that these reactions be included for modeling applications than being neglected. For
560 example, the predicted formation of catechols from phenols due to this reaction has implications in
561 predictions of SOA from aromatics.

562 2.3.3. Other Compounds

563 Decompositions of peroxy nitrates and rearrangements of cyclohexadienones are currently the
564 only types of non-radical unimolecular reaction considered by MechGen. In particular, MechGen does not
565 consider other types of water- or surface-catalyzed conversion of non-radical products that could affect
566 the environmental fate of multi-functional products. For example, geminal diols, compounds with HO-
567 CX₂-OH groups, are predicted to be formed in some cases and would be expected to be in equilibrium
568 with H₂O and XC(=O)X in the environment. It is uncertain whether this or similar reactions would occur
569 at significant rates in the gas phase, and MechGen currently assumes that this is not the case. However,
570 ignoring these potentially heterogeneous or H₂O-catalyzed reactions may also have implications in SOA
571 predictions.



572 3. Reactions of Carbon-Centered Radicals



573 Carbon-centered radicals are formed in the initial reactions of most VOCs with atmospheric
 574 oxidants, from H-shift isomerization reactions of peroxy and alkoxy radicals, and from most alkoxy
 575 radical decompositions. Under lower atmospheric conditions, their major fate is expected to be reaction
 576 with O₂, which is estimated to occur with a pseudo-unimolecular rate constant of ~3.8 x 10⁷ sec⁻¹ (see
 577 Table A-6 for measured radical + O₂ rate constants and associated text in the SI). However, rapid
 578 unimolecular reactions are estimated to dominate for some types of carbon-centered radicals formed in
 579 generated mechanisms, as discussed below. Note that there are currently no cases where MechGen
 580 assumes that both O₂ and unimolecular reactions of carbon-centered radicals occur at competitive rates --
 581 one or the other is assumed to dominate. This means that estimates of carbon-centered radical + O₂ rate
 582 constants are not currently needed or used in the generated mechanism, though estimates of these rate
 583 constants are used when considering when unimolecular reactions may dominate.

584 3.1. Reaction with O₂

585 Most carbon-centered radicals are assumed to primarily add O₂ and form the corresponding
 586 peroxy radical, whose reactions are discussed in Section 4.




588 However, the peroxy radicals formed from α-OH substituted radicals are assumed to rapidly decompose
 589 to form HO₂ and the corresponding carbonyl.




593 MechGen treats these as concerted processes when reactions of α-OH substituted carbon-centered
 594 radicals are generated, since stabilization and other reactions are expected to be unimportant.

595 If the carbon-centered radical is adjacent to one or more double bond groups, it is assumed to be a
 596 resonance structure where O₂ can add to any of the radical positions, e.g.:



599 where α(G_a,G_b) is the fraction of adding to the G_a position when G_b is the other position on the radical.
 600 Additions to radicals with more than one double bond are assumed to form only those adducts with the
 601 most conjugated structures, e.g.:



602 Because of lack of data indicating otherwise, we assume that the branching ratios, α, depend only on the
 603 nature of the radical groups where the addition occurs, regardless of the number of double bond groups in
 604 the radical. Although this has not been examined comprehensively, we believe that all allylic radical
 605 structures that may arise in generated mechanisms have only two radical centers where O₂ addition forms
 606 the most conjugated structures. This is supported by the fact that we have not found examples of where
 607 this is not the case when generating many mechanisms.



608 Available information and estimates concerning branching for allylic additions is discussed in
609 Section A1.4.2. There are very limited data, but what is known does not support the assumption that
610 additions at each position are equally likely. Product data for the reactions of 1,3-butadiene and isoprene
611 with OH indicate that in radicals with both primary (-aCH₂[·]) and secondary (-aCH[·-]) radical positions,
612 the addition occurs ~65% of the time at the secondary position. This is consistent with the fact that
613 thermochemical analysis indicates that addition to the secondary radical is ~2 kcal/mole more favorable
614 (see SI). In addition, theoretical calculations of Xu and Wang (2013) on phenol mechanisms and of Yuan
615 et al. (2017) on furan mechanisms both indicate that O₂ preferentially adds to radical centers with -OH or
616 -OR substituents, instead of those with unsubstituted secondary radical centers. However, if this is
617 assumed then the mechanisms significantly underpredict reactivities observed in environmental chamber
618 experiments with phenolic compounds, and better fits to the data are obtained if it is assumed that
619 addition to non-OH-substituted secondary radical centers occur approximately 15% of the time
620 (unpublished results). This is highly uncertain, but it provides an additional data point for deriving general
621 estimation method incorporating -OH substituted radicals.

622 As discussed in Section A1.4.2, we assume that the branching ratios can be estimated based on
623 the differences in heats of addition of O₂ to the different radical positions, with a correction included if
624 one of the radical centers is -OH or -OR substituted. The estimate used is as follows:

$$625 \alpha(G_1, G_2) = \text{WtF}(G_1) / [\text{WtF}(G_1) + \text{WtF}(G_2)]$$

$$626 \text{ where } \text{WtF}(G) = \text{Weighting factor for } G = e^{-\beta \times [\Delta\Delta\text{Hr}(G) - O_{\text{corr}}(G)]}$$


627 and $\Delta\Delta\text{Hr}(G)$ is the difference of heats of reaction for O₂ addition to G compared to a primary radical
628 center in a radical that has both, β is a parameter assigned a value of 0.33 to be consistent with the
629 product data for isoprene and 1,3-butadiene, and O_{corr} is set at 5.78 kcal/mole if G is -OH or -OR
630 substituted to be consistent with modeling results for phenols, or zero if there are no such substituents.
631 The weighting factors so derived are as follows (see Section A1.4.2 for derivations):

<u>G·</u>	<u>WtF(G)</u>	<u>G·</u>	<u>WtF(G)</u>	<u>G·</u>	<u>WtF(G)</u>
-aCH ₂ [·]	1	-aCH[·]-CO-X	0.35	-aCH[·]-OX	21.9 *
-aCH[·]-R	1.9	-aC[·](R)-CO-X	0.28	-aC[·](R)-OX	13.6 *
-aC[·](R)-R	1.13	(O _{corr} not used)		* O _{corr} used	

632 Note that use of this methodology to estimate branching ratios in general is highly uncertain because only
633 two data points are available to derive the two parameters used (β and O_{corr}), with no data available to test
634 this estimation method. However, this is the best that can be done at the present time.

635 The above discussion is not applicable to radicals that are estimated to have rapid unimolecular
636 reactions as discussed in Section 3.2. If such a rapid reaction is possible for one of the radical centers in
637 an allylic radical, it is assumed that all the reaction goes by this route unless a rapid reaction is possible at
638 the other radical center, in which case it is assumed that they occur with equally probability. The above
639 discussion is also only applicable to stabilized radicals, and not necessarily to excited adducts formed
640 when OH or NO₃ adds to double bonds. In most cases excited adducts formed following additions to
641 double bonds are assumed to be primarily stabilized and then react as discussed above, but that is not
642 always the case, as discussed in Section 3.3.

643 A different mechanism is assumed for the addition of O₂ to vinylic radicals where the radical
644 center is on an unsaturated carbon. In this case, the reaction with O₂ is expected to involve the O₂ adding
645 to each side of the bond, with the bonds rearranging to form a carbonyl product and a carbonyl radical,
646 i.e.,



647 This is based on data of Slagle et al. (1984) for vinyl and Slagle et al. (1988) for methylvinyl and $i\text{-C}_4\text{H}_9\cdot$.
 648 More recently, Matsugi and Miyoshi (2014) directly measured the yield of HCO radicals from vinyl, and
 649 observed that the yields were only about 0.2, independent of pressure, with the remainder of the reaction
 650 being prompt dissociation to $H + CO$. The issue of whether the HCO promptly decomposes is not relevant
 651 to atmospheric modeling, since the same products (HO_2 and CO) are formed under atmospheric
 652 conditions in either case. Therefore, MechGen represents the reactions of $CX_2=CH\cdot$ as forming $H\cdot + CO$
 653 alone, since this is the major overall process.


654 However, the issue of prompt dissociation is relevant in the case of $RC(=O)\cdot$, since the stabilized
 655 radical is expected to primarily add O_2 to form an acyl peroxy radical, which can give different products
 656 than if $RC(=O)$ decomposes to $R\cdot + CO$. Decomposition is expected to be more favorable because the
 657 lower estimated heat of reaction, though the rate of stabilization of the $RC(=O)\cdot$ should also be greater
 658 than for $HC(=O)\cdot$. MechGen assumes that the energetics is the larger factor in the case of additions of O_2
 659 to vinylic radicals, assumes that the prompt dissociation dominates, and also represents the overall
 660 reaction as forming the carbonyl, CO , and $R\cdot$ as a single process.

661 3.2. Unimolecular Reactions of Stabilized Radicals


662 Several types of carbon-centered radicals are expected to undergo unimolecular reactions
 663 sufficiently rapidly that unimolecular reaction is assumed to dominate over reaction with O_2 , even for
 664 thermalized radicals. Note that if these reactions are assumed to be fast for thermalized radicals, they are
 665 necessarily assumed to be fast when the radicals are excited.

666 3.2.1. Rapid α -Scission Decompositions

667 Carbon-centered radicals with adjacent hydroperoxy, peroxide, nitrate, or dicarbonyl groups are
 668 expected to rapidly undergo decompositions by simple scission of an adjacent bond, forming products
 669 that are more energetically favorable.



674 We expect that there should not be large entropy or energy barriers to these reactions, so MechGen
 675 assumes they all dominate over O_2 addition whenever radicals of these types are formed. Note that if the
 676 peroxy, nitrate, or carbonyl substituent is next to an allylic radical center, the bond scission reaction is still
 677 expected to dominate over O_2 addition,




679 so there is no O_2 addition to the other radical centers. Reactions of allylic radicals with peroxy, nitrate, or
 680 dicarbonyl groups are treated similarly.



681 **3.2.2. Cyclopropyl Ring Opening Decompositions**

682 Carbon-centered radicals with the radical center in a cyclopropyl ring can decompose by breaking
683 the bond opposite the radical center, relieving the ring strain and forming an energetically favorable
684 allylic stabilized radical.



686 This reaction is not expected to have large or any barriers, so is assumed to dominate over O₂ addition.
687 On the other hand, the analogous reactions when the radical center is in a cyclobutane ring is estimated to
688 be much less energetically favorable and is only expected to occur in chemically activated radicals, as
689 discussed in Section 3.3. The reactions for larger rings are estimated to be endothermic and are assumed
690 to be unimportant relative to stabilization and/or O₂ addition even for excited radicals.


691 **3.2.3. Cyclizations forming 5-Member Ring Cyclic Esters**

692 As discussed in Section A1.4.3, Curran et al. (1998) provided estimates for the Arrhenius
693 parameters for decompositions of various QOOH radicals of the type,




695 as a function of cyclic ether ring size. These can be used as a basis for estimating the approximate
696 importance of these and similar reactions in atmospheric systems. In general, both the estimated A factors
697 and activation energies decreased with ring size, with the net result of these opposing factors being that
698 formation of 5-member cyclic ethers is generally estimated to be most favorable for atmospheric
699 conditions. Nevertheless, these cyclization reactions are estimated to be too slow for such radicals with
700 only alkyl or hydroperoxide groups to compete with reaction with O₂ at atmospheric temperatures.

701 However, as discussed in Section A1.4.4 in the SI, extrapolating the relation between the rate
702 constant and heats of reaction derived from the results of Curran et al. (1998) results in estimates that
703 cyclizations of carbonyl-containing radicals forming 5-member ring esters are ~10 times faster than O₂
704 addition for atmospheric conditions.



705 In addition, analogous cyclizations forming cyclic esters can also occur with carbon-centered radicals
706 with peroxy, nitrate, or peroxyxynitrate substitution in the δ position, are also estimated to be faster than O₂
707 addition, i.e.,



708 where "X" = G or NO₂, and "XO." is G-O· or NO₃.

709 However, as also discussed in A1.4.3, Miyoshi (2011) reports results of theoretical calculations of
710 hydroperoxyalkyl rate constants that extrapolate to much lower rate constants at atmospheric
711 temperatures, suggesting that the cyclizations forming 5-member ring esters may not as fast as estimated
712 using the results of Curran et al. (1998). These reactions may still be important for excited radicals. This
713 will need to be re-examined in a subsequent version of MechGen, but presently this reaction is still
714 assumed to be fast even for thermalized radicals.



715 These reactions are assumed to be fast for mechanism generation purposes if (1) there is a
716 carbonyl group either at the radical center or next to the peroxy oxygen that becomes bonded to the
717 radical center, (2) the cyclic ester, carbonate, or anhydride has a 5-member ring, and (3) the radical is not
718 stabilized by resonance. O₂ addition is assumed to dominate if not all of these conditions are met. For
719 simplicity and to avoid the necessity of making highly uncertain quantitative rate constant estimates, we
720 assume that either decomposition or O₂ addition dominates, with the other being negligible. This
721 assumption can be revisited in the future if the ability to estimate these decomposition rate constants for
722 atmospheric conditions improves.

723 Although it is not likely in reactions of most VOCs and their oxidation products, in some cases
724 carbon-centered radicals may have more than one peroxy group in positions where decompositions are
725 estimated to be rapid. In these cases, it is assumed that the reactions occur at equal rates.

726 3.3. Unimolecular Reactions of Chemically Activated Radicals

727 Carbon-centered radicals formed by the addition of OH to double bonds are initially formed with
728 internal excitation, and in some cases might undergo unimolecular reactions that may not be important for
729 stabilized radicals. Reactions of the chemically activated radicals formed following additions to aromatic
730 rings are discussed separately in Section 3.4. Reactions that are assumed to be fast even for thermalized
731 radicals were discussed in the previous section, and they are also assumed to dominate for excited radicals
732 as well. Reactions that are assumed to dominate for radicals excited by additions of OH radicals to double
733 bonds, but not be important for stabilized radicals, are listed in Table 6 and are discussed in more detail in
734 Sections A1.4.3 and A1.4.4 in the SI.

735 Note that additions of NO₃ to double bonds also result in the formation of excited radicals, but the
736 excitation energy is estimated to be ~10 kcal/mole less than OH addition. Although this is uncertain, we
737 currently assume that stabilization is correspondingly more important for NO₃ adducts, and the
738 unimolecular reactions shown in Table 6 are assumed not to occur when mechanisms for reactions of NO₃
739 with alkenes are generated.

740 3.4. Reactions of Radicals Formed by Additions to Aromatic Rings

741 OH radicals can react by adding to various positions around the aromatic ring, with the branching
742 ratios derived using SARs for OH reactions with aromatics, as discussed in Section 2.1.2 and by Carter
743 (2021). These excited OH-aromatic adducts can either decompose back to the aromatic + OH or be
744 stabilized and then react with O₂. The effect of the back decomposition is taken into account in the SARs
745 for OH addition, which are based on total rate constants, so only reaction with O₂ is considered.
746 Theoretical calculations suggest that O₂ abstraction forming HO₂ and a phenol (or catechol) is negligible
747 compared to O₂ addition, so only addition forming an OH-aromatic-O₂ adduct is considered.

748 The OH-aromatic adduct is an allylic carbon-centered radical with three possible radical centers
749 where O₂ can add. Although these adducts have some excitation, we assume that the branching ratios for
750 O₂ addition to the different positions are the same as estimated for stabilized allylic radicals, as discussed
751 in Section 3.1 and Section A1.4.2. This means that all the addition is assumed to occur at the *ortho*
752 positions since this forms the adducts with the most conjugation, that addition to an alkyl-substituted
753 *ortho* position is ~60% less likely than adding to the other *ortho* position, and addition to an -OH



754 Table 6. Summary of types of unimolecular reactions that are assumed to be non-negligible for
 755 chemically activated carbon-centered radicals formed following additions of OH to
 756 double bonds.

Type [a]	Reactant [b]	Reaction	Ratio [c]
<i>Exo</i> ring opening (A1.4.3)	$-G_1[\cdot]-*G_2-G_3-G_4^*-\{\text{excited}\}$	$\rightarrow -G_1=G_2-G_3-G_4[\cdot]-$	
	$-G_1[\cdot]-*G_2-G_3-G_4-G_5^*-\{\text{excited}\}$	$\rightarrow -G_1=G_2-G_3-G_4-G_5[\cdot]-$ $+ M \rightarrow -G_1[\cdot]-*G_2-G_3-G_4-G_5^*-\ + M$	70% 30%
<i>Endo</i> ring opening (A1.4.3)	$*G_1-G_2[\cdot]-G_3^*-\{\text{excited}\}$	$\rightarrow -G_1=G_2-G_3[\cdot]-$	
	$*G_1-G_2[\cdot]-G_3-G_4^*-\{\text{excited}\}$	$\rightarrow -G_1=G_2-G_3-G_4[\cdot]-$ $+ M \rightarrow *G_1-G_2[\cdot]-G_3-G_4^*-\ + M$	70% 30%
Peroxy Cyclizations (A1.4.4)	$-G_1[\cdot]-G_2-O-OX\{\text{OHadd}\}$	$*G_1-G_2-O^* + [\cdot]OX$	
	$-G_1[\cdot]-G_2-ONO_2\{\text{OHadd}\}$	$*G_1-G_2-O^* + NO_2$	
	$-G_1(O'H)-G_2[\cdot]-CO-O-OH\{\text{OHadd}\}$	$\rightarrow G_1(O'H)-*G_2-CO-O^* + OH$ $\rightarrow *O'-G_1-G_2^*-CO-OH + OH$ $+ M \rightarrow G_1(O'H)-G_2[\cdot]-CO-O-OH + M$	14% 5% 82%
	$-G_1(O'H)-G_2[\cdot]-CO-O-ONO_2\{\text{OHadd}\}$	$\rightarrow G_1(O'H)-*G_2-CO-O^* + NO_3$ $\rightarrow *O'-G_1-G_2^*-CO-OH + NO_3$ $+ M \rightarrow G_1(O'H)-G_2[\cdot]-CO-O-ONO_2 + M$	60% 20% 20%
	$-G_1(O'H)-G_2[\cdot]-CO-O-O-G_3-\{\text{OHadd}\}$	$\rightarrow G_1(O'H)-*G_2-CO-O^* + -G_3[O\cdot]$ $\rightarrow *O'-G_1-G_2^*-CO-OH + -G_3[O\cdot]$ $+ M \rightarrow G_1(O'H)-G_2[\cdot]-CO-O-O-G_3-\ + M$	45% 15% 40%

757 [a] The section in the SI where the reaction and the estimated branching ratios are discussed is shown in
 758 parentheses.

759 [b] "*" indicates ring closure, "{excited}" indicates excitation, {OHadd} indicates excitation by OH
 760 addition only, "+ M" indicates stabilization, and "X" is H or any group and ·OX is OH, or G-O·.


761 [c] Branching ratio assumed for 1 atmosphere pressure (the default) if stabilization is assumed to be non-
 762 negligible. Note that MechGen allows for changes in atmospheric pressure, in which case this ratio
 763 would change assuming that the stabilization rate constant is proportional to pressure.

764
 765

766 substituted *ortho* position (as occurs in reactions with phenols) is ~7 times more likely, assuming that the
 767 other *ortho* position is unsubstituted.

768 The aromatic-OH-O₂ adducts are expected to be formed with some excitation, and are assumed to
 769 undergo various types of unimolecular reactions as discussed below. Although peroxy radical reactions
 770 are discussed separately in Section 4, MechGen incorporates the rapid aromatic-OH-O₂ isomerization
 771 reactions into the overall reactions generated for excited OH-aromatic adducts, so these reactions are
 772 discussed in this section.

773 The subsequent reactions of the excited adducts are assumed to be as shown in Figure 1, which
 774 uses benzene as an example. It shows the initial formation of the OH-aromatic adducts, their subsequent
 775 reactions with O₂, and the fast unimolecular reactions assumed for the excited OH-aromatic-O₂ adducts.
 776 The codes for the branching ratios, which depend on the substituents about the aromatic ring are also



777

778

Figure 1. Initial reactions following the addition of OH to aromatic rings, using benzene as an example, showing the formation of the OH-aromatic adduct, followed by unimolecular reactions assumed for the OH-aromatic-O₂ adduct. Major products following the reactions of the radicals formed are also shown.

779

780

781

782

783

shown. Three types of overall reactions are assumed. These include formation of a phenol + HO₂ via a 6-member ring transition state where the H transfers from the -OH to the peroxy group, followed by elimination of HO₂, with branching ratio designated *fPhen*; cyclization to form a bicyclic allylic-stabilized carbon-centered radical with branching ratio *fBcyc*; and the third involving a different type of cyclization where a peroxy O transfers to the opposite double bond, forming an epoxide and an alkoxy radical, with branching ratio *fEpoxy*. The alkoxy radical can form two different products depending on where the O₂ added and the substituents about the aromatic ring, which are assumed to be equally likely. Therefore, up to four overall reactions can be generated following addition of OH to aromatic rings.

784

785

786

787

788

789

790

791

The mechanisms for the subsequent reactions of the carbon-centered and alkoxy radicals formed are estimated using the same procedures discussed elsewhere in this paper. The major stable products predicted to be formed are also indicated in Figure 1. Note that as discussed in Section 2.2.5, the unsaturated 1,4-dicarbonyl aldehydes or aldehyde-ketones are predicted to be highly photoreactive, so the yields of these products significantly affect predictions of reactivity in aromatic hydrocarbon - NO_x environmental chamber experiments (e.g., Carter and Heo, 2013).

792

793

794

795

796

797

The branching ratios were derived based on reported yields of phenolic products and α-dicarbonyls from the reactions of OH with benzene and various alkylbenzenes, as well as simulations of environmental chamber experiments. Values of *fPhen* were derived based on phenolic yields, *fBcyc* values are based on 1,2 dicarbonyl yields and modeling environmental chamber experiments, and *fEpoxy* are obtained as 1 - *fPhen* - *fBcyc*. It was sufficient to assume that the branching ratios depend only on the presence of substituents in the *ipso* and *ortho* positions, with substituents in the *meta* or *para* positions assumed not to have an effect on these branching ratios. The branching ratios used for the various

798

799

800

801

802

803



804 substituent cases are summarized in Table 7. Note that f_{Phen} is necessarily zero if there is an *ipso*
 805 substituent, and the f_{Bcyc} / f_{Epox} ratios for such compounds are estimated to be the same as those for
 806 compounds without *ipso* substituents, but with the magnitudes increased so they sum up to 1. The ratios
 807 in Table 7 are used regardless of substituents, though estimates for compounds with substituents other
 808 than alkyl or -OH, and therefore generated mechanisms for such compounds, should be considered to be
 809 unreliable and probably should not be used for mechanism development.

810

811 Table 7. Branching ratios derived for the reactions of aromatic + OH and aromatic + OH + O₂
 812 adducts, based on the presence or absence of substituents in the *ipso* and *ortho* positions.

Branching Ratio	No <i>ipso</i> substituents			1 <i>ipso</i> substituent [a]		
	No <i>ortho</i>	1 <i>ortho</i>	2 <i>ortho</i>	No <i>ortho</i>	1 <i>ortho</i>	2 <i>ortho</i>
f_{Phen}	57%	16%	5%	-	-	-
f_{Bcyc}	31%	57%	83%	73%	68%	88%
f_{Epox}	12%	27%	12%	27%	32%	12%

813

[a] Derived from f_{Bcyc} and f_{Epox} assigned for adducts with no *ipso* substituents.

814

815 The above discussion is applicable only to reactions following additions of OH radicals to
 816 aromatic rings, not to additions by NO₃ radicals. As discussed by Carter (2021), the rate constants for
 817 these additions are estimated to be slow, based on the relatively low total rate constants reactions for
 818 aromatic + NO₃ reactions. MechGen currently assumes that all aromatic-NO₃ adducts rapidly decompose
 819 back to reactants, resulting in no net reaction. However, this cannot strictly speaking be true, since the
 820 experimental rate constants give SARs that predict nonzero, though low, rate constants for these reactions
 821 (Carter, 2021; see also Section 2.1.2). However, the predicted rate constants are too low to be important in
 822 atmospheric systems, so this oversimplification has no practical consequences for atmospheric
 823 mechanisms.

824

The reactions of radicals formed by the additions of OH radicals to the aromatic rings in phenols
 825 are expected to be analogous to those discussed above, but the presence of OH substitution affects
 826 branching ratios for the addition to the allylic OH + phenol adduct as discussed in Section 3.1, and some
 827 of the OH + phenolic + O₂ adducts form cyclohexadienones, which undergo are assumed to undergo
 828 rapid unimolecular reactions to form catechols as discussed in Section 2.3.2. The mechanisms assumed
 829 for OH + phenolic adducts are discussed in Section A1.8 of the SI.

830 4. Reactions of Peroxy Radicals

831

Peroxy radicals are formed primarily by the reactions of alkyl radicals with O₂ and are critical
 832 intermediates in atmospheric oxidation mechanisms. Peroxy radicals can be classified alkyl peroxy,
 833 -G[OO.] or RO₂, or acyl peroxy, -G-CO[OO.] or RCO₃ (where "G" is any group that does not have an α
 834 carbonyl substituent). In the following discussion, we will use the term "acyl" peroxy radical to refer to
 835 any peroxy radical bonded to a carbonyl group and "alkyl" peroxy radicals otherwise, including peroxy
 836 radicals with non-alkyl groups or substituents.

837

Peroxy radicals can react with NO, NO₂, NO₃, HO₂, and other peroxy radicals, and some can also
 838 undergo unimolecular reactions at significant rates under atmospheric conditions. The rate constants
 839 assigned when generating bimolecular rate constants are summarized in Table 8, and their derivations are
 840 summarized in the following subsections where the mechanisms for the different types of reactions are



841 discussed. More details concerning the derivations of the bimolecular rate are given in Section A1.5.1 of
842 the SI. The unimolecular reactions are discussed separately below.

843

844 Table 8. Summary of bimolecular rate constants used when generating reactions of peroxy
845 radicals under atmospheric conditions.

Reactant	Rate constant around 300K (cm ³ molec ⁻¹ s ⁻¹) [a]						RCO ₃
	Non OH-Substituted RO ₂			OH-Substituted RO ₂			
	Primary	Secondary	Tertiary	Primary	Secondary	Tertiary	
NO			9.13x10 ⁻¹²				2.10x10 ⁻¹¹
NO ₂			8.8x10 ⁻¹²				7.7x10 ⁻¹²
NO ₃			2.3x10 ⁻¹²				4.0x10 ⁻¹²
HO ₂	10 ⁻¹² x min [20.3, 2.48 + (2.48 x nC)]			10 ⁻¹² x min [21.1, 9.37 + (1.68 x nC)]			2.1x10 ⁻¹¹
Generic RO ₂	2.9x10 ⁻¹⁴	1.6x10 ⁻¹⁴	3.7x10 ⁻¹⁶	9.1x10 ⁻¹⁴	1.7x10 ⁻¹³	1.3x10 ⁻¹⁴	1.6x10 ⁻¹¹
Generic RCO ₃			1.6x10 ⁻¹¹				1.4x10 ⁻¹¹

846 [a] See Table A-45 in the SI for the sources of the rate constants used. Note that the same rate constant is
847 used for reactions of alkyl peroxy radicals with NO_x species and with acyl peroxy radicals, while the
848 rate constants depend on the carbon number for reactions with HO₂, and on the structure of the radical
849 for reactions with other alkyl peroxy radicals (See Sections 4.4 and 4.5, respectively).


850

851

852 4.1. Reaction with NO

853 The major atmospheric loss process of most peroxy radicals when formed in the presence of NO_x
854 is reaction with NO. Available data on the rate constants for peroxy + NO reactions are given in Table A-
855 45 in the SI, and the rate constants measured for ethyl peroxy and propionyl peroxy are estimated to apply
856 to all peroxy radicals for mechanism generation. There is some variability in the measured rate constants
857 for the alkyl peroxy radicals, but the data are insufficient to derive estimates for dependences of the rate
858 constant on the structure or size of the radicals. As indicated in Table 8, although different rate constants
859 are used for reactions of acyl peroxy vs peroxy radicals with NO, because of insufficient information we
860 assume that other structural effects have no effects on the rate constant.

861 The reactions of peroxy radicals with NO can occur via two possible routes, either forming NO₂
862 and the corresponding alkoxy radical, or forming a stable organic nitrate in the presence of a bath gas.




865 Both reactions are known to be important in the case of alkyl peroxy radicals, and the nitrate yield,
866 $y_{\text{N}} = k_2 / (k_1 + k_2)$, is an important parameter affecting model predictions in atmospheric systems since the
867 conversion of NO to NO₂ in Reaction (1) contributes to O₃ formation and the formation of alkoxy radicals
868 results in radical propagation, while nitrate formation in Reaction (2) causes radical termination and is
869 also a NO_x sink, at least for the intermediate term.

870 Available data and estimates concerning nitrate yields from various peroxy radicals are discussed
871 in Section A1.5.2 of the SI. There are extensive measurements of nitrate yields from various unsubstituted
872 peroxy radicals formed from the reactions of alkanes, including data at various temperatures and



873 pressures, allowing a parameterization to be derived that fits these data as a function of temperature,
 874 pressure, and carbon number. The parameterization used is similar to that employed previously (e.g.,
 875 Atkinson et al., 1983), updated based on the mechanism considerations discussed by Zhang et al. (2004)
 876 and re-optimized to fit available data as discussed in the SI. The predicted nitrate yields at around 298K
 877 and 1 atmosphere pressure are given in Table 9 and are plotted against carbon number in Figure 2, where
 878 they are compared to available data as discussed below.



879
 880

881 Figure 2. Measured and estimated nitrate yields from reactions of various types of peroxy radicals
 882 with NO against numbers of carbon atoms in the radicals.

883

884 Table 9. Estimated nitrate yields as a function of carbon number calculated for ~298K and 1
 885 atmosphere pressure.

Carbon Number	2	3	4	5	6	7	8	9	10	11	12	13	14+
Default nitrate yield	2%	4%	7%	11%	15%	19%	22%	25%	27%	28%	28%	28%	29%
Nitrate yield if α - or β -O substituted	2%	3%	5%	8%	11%	14%	16%	18%	19%	20%	20%	21%	21%

886

887 Figure 2 shows that the parameterization derived to fit measured nitrate yields for unsubstituted
 888 peroxy radicals from alkanes (black line) fit those data reasonably well, but its applicability for
 889 substituted radicals from other VOCs is much more uncertain. The limited available measurements of
 890 nitrate yields from β OH-substituted radicals are inconsistent. Data obtained from Shepson's (Muthuramu
 891 et al., 1993; Shepson et al., 1985; O'Brien et al., 1998) and Ziemann's (Matsunaga and Ziemann, 2009,
 892 2010) groups gave yields that are about a factor of two lower than from unsubstituted radicals, while more
 893 recent data from Teng et al. (2015) gave yields that agree with those from the unsubstituted radicals, at
 894 least at carbon numbers <10. The estimated nitrate yields from the OH-substituted radicals significantly
 895 affect results of model simulations of alkene - NO_x chamber experiments (Carter, 2000, 2010b), and we
 896 found that some data are better fit using the higher yields estimated for radicals from alkanes, while some
 897 are better fit using the yields that are a factor of ~2 lower (see Section A1.5.2 in the SI).




898 Figure 2 also shows scatter in the available data concerning nitrate yields from peroxy radicals
899 formed from ethers, esters, and ketones, though they tend to be lower than observations for unsubstituted
900 peroxy radicals (Jenkin et al., 2019 and references therein). As discussed in the SI, these yields are subject
901 to greater uncertainties because they are derived either from unspiciated total nitrate measurements in
902 complex chemical systems, or by fits to environmental chamber data, where uncertainties are even
903 greater.

904 Until more information is available to resolve these discrepancies and provide a better basis for
905 making nitrate yield estimates for non-alkyl-substituted peroxy radicals, MechGen uses a
906 parameterization (red dotted line in Figure 2) that predicts yields halfway between those derived for
907 radicals from alkanes and the lower yields for the β OH-substituted radicals, when estimating nitrate
908 yields for peroxy radicals bonded to oxygen groups in the α or β positions, e.g., -OH, ether, ester,
909 peroxides, or nitrates. This is similar to the approach used in the nitrate yield estimation method
910 developed by Jenkin et al. (2019) for alcohols, ethers, and esters. Based on the limited data, Jenkin et al.
911 (2019) also assumed carbonyl groups in the α or β positions caused even lower nitrate yields, but the
912 current version of MechGen does not yet incorporate corrections for these cases.

913 The nitrate yields predicted using this parameterization, derived as discussed in Section A1.5.2 of
914 the SI, are included in Table 9. Note that we assume that substituents farther away than the β position are
915 too far away from the reaction site to affect the nitrate yield. However, the estimates for nitrate yields
916 from substituted peroxy radicals are uncertain by at least a factor of two, and are more likely to be biased
917 high rather than low.

918 There is no evidence for nitrate formation in the case of acyl peroxy radicals, so the formation of
919 NO_2 and the corresponding acyl oxy radical is assumed to be the only pathway of significance (e.g., see
920 IUPAC, 2023).





922 As discussed above, the acyl oxy radicals are expected to rapidly decompose to form CO_2 and the
923 corresponding alkyl radical, making the overall process for acyl + NO reactions as indicated above.

924 **4.2. Reaction with NO_2**

925 Alkyl and acyl peroxy radicals also react with NO_2 under atmospheric conditions, and the rate
926 constants assigned to them are included in Table 8. The assigned rate constants are based on measured
927 high pressure rate constants for ethyl peroxy and ethyl acyl peroxy radicals, since we could not find data
928 for larger peroxy radicals (see Table A-45 in the SI). Although the reactions of ethyl and ethyl acyl
929 peroxy radicals are not in the high pressure limit under atmospheric conditions, most of the radicals are
930 larger and their rate constants should be closer or at the high pressure limit, so the high pressure rate
931 constant is used for general estimates.

932 The reactions for both alkyl and acyl peroxy radicals with NO_2 are assumed to result entirely in
933 the formation of the corresponding alkyl or acyl peroxyxynitrate.



936 Reactions forming NO_3 + alkoxy or acyloxy radicals are estimated to be endothermic, and are assumed
937 not to be important. The peroxyxynitrate compounds formed are thermally unstable at atmospheric
938 temperatures, decomposing back to NO_2 and the peroxy radical, as discussed in Section 2.3. It is assumed



939 that the presence of non-alkyl substituents on the "R" does not affect either the formation or
940 decomposition rate constants, but this is uncertain.

941 **4.3. Reaction with NO₃**

942 Reactions with NO₃ can be a potentially important sink for peroxy radicals at nighttime and
943 potentially other conditions when NO concentrations are low. The rate constants used for mechanism
944 generation (Table 8) are those measured for ethyl peroxy and acyl peroxy radicals because we could not
945 find data for larger radicals (see Table A-45).

946 Peroxy radicals react with NO₃ to form the corresponding alkoxy radical, NO₂ and O₂, with no
947 evidence of other reaction routes or products being reported (e.g., IUPAC, 2023).



950 Note that the same organic products are formed in this reaction as in the reaction of peroxy radicals with
951 NO, other than the lack of the nitrate formation route in the NO₃ reaction. Although nitrate formation in
952 this reaction is energetically possible, there is no evidence that it occurs under atmospheric conditions.
953 The presence of non-alkyl substituents on "R" is assumed not to affect the rate constants or mechanisms.

954 **4.4. Reaction with HO₂**


955 Reactions with HO₂ can be the major sink for peroxy radicals under conditions when NO_x is low,
956 being generally much more important than reactions with other peroxy radicals. Information concerning
957 measurements and estimates for mechanisms and rate constants for the reactions of peroxy radicals with
958 HO₂ are discussed in Section A1.5.3 of the SI. In the case of alkyl peroxy radicals, the data suggest that
959 the rate constants depend on the size of the radical and the presence of OH substitution on the radical,
960 with the following equations being derived for estimation purposes:


$$961 k(\text{RO}_2 + \text{HO}_2) = 10^{-12} \times \min [20.3, 2.48 + (2.48 \times nC)] \text{ cm}^3 \text{ molec}^{-1} \text{ s}^{-1}$$
$$962 k(\text{HO-RO}_2 + \text{HO}_2) = 10^{-12} \times \min [21.1, 9.37 + (1.68 \times nC)] \text{ cm}^3 \text{ molec}^{-1} \text{ s}^{-1}$$

963 where "HO-RO₂" refers to alkyl peroxy with OH substituents anywhere on the radical, and "RO₂" refers
964 to those without. Note that there is some inconsistency in the measured rate constants for OH substituted
965 radicals (see SI), so there is some uncertainty in these estimates.

966 The only data we found for reactions of acyl peroxy radicals with HO₂ concerned acetyl peroxy
967 radicals, so the rate constant measured for that reaction (on Table A-45) is used for estimation purposes.
968 This is uncertain, especially in view of the variability of the measured rate constants for the alkyl peroxy
969 radicals.

970 Available information concerning the products of the reactions of various types of peroxy radicals
971 with HO₂ have been evaluated by IUPAC (2023), and their recommendations of branching ratios for
972 methyl and acetyl peroxy radicals, and for methyl peroxy substituted with -CH₃, -OH, -OCH₃, and
973 -C(O)CH₃ are given in A-12 of the SI. Their recommendations include the following four routes:





974 with the "ozone" route only considered for acyl peroxy radicals. The "carbonyl" route occurs via an H
975 shift from an α hydrogen to the HO₂ via a 6-member ring transition state, and is not possible for tertiary
976 alkyl or acyl peroxy radicals. Although there are data only for the C₁ or C₂ peroxy radicals, they are
977 assumed to be applicable to all alkyl peroxy radicals with the same types of α substituents for mechanism
978 generation purposes, and are the basis for the branching ratios that are summarized in Table 10. However,
979 the applications of these recommendations to the many types of larger peroxy radicals predicted in
980 generated mechanisms is highly uncertain.

981

982 Table 10. Branching ratios assigned for the four possible routes for the reactions of HO₂ with
983 various types of peroxy radicals.

Route	CH ₃ O ₂	RO ₂ [a]	HO-RO ₂ [b,c]		RO-RO ₂ [b,c]		RCO-RO ₂ [b]	RCO ₃
			P, S	T	P, S	T		
Hydroperoxide	0.9	1.0	0.5	0.8	0.6	1.0	0.85	0.37
Carbonyl	0.1	-	0.3	-	0.4	-	-	-
Hydroxyl	-	-	0.2	0.2	-	-	0.15	0.5
Ozone	-	-	-	-	-	-	-	0.13

984 [a] "RO₂" refers to peroxy radicals with no -OH, -OR, or carbonyl substituents in the α position. Note
985 that they can have these and other non-alkane substituents or groups elsewhere on the molecule.

986 [b] "The "HO-", "RO-" or "RCO-" refer to the substituent in the α position only. "RCO-" refers to either
987 -CHO or -CO-, though data are only available for -CO- substitution.

988 [c] "P", "S", and "T" refer to primary, secondary, and tertiary radicals, respectively. Since the carbonyl
989 route is not possible for tertiary radicals, the branching ratio estimate for this route for primary or
990 secondary radicals is added to the estimate for the hydroperoxy route for such radicals.

991

992 4.5. Reactions with Other Peroxy Radicals


993 Organic peroxy radicals can also be consumed under atmospheric conditions by reactions with
994 other organic peroxy radicals. Although this is generally a less important loss process under atmospheric
995 conditions than reactions with NO_x or HO₂, such reactions can be non-negligible under relatively high
996 VOC and low NO_x atmospheric conditions, and also in some laboratory experiments. Available
997 information concerning measured rate constants for these reactions is given in Table A-45 of the SI. The
998 rate constants vary by orders of magnitude, with the most important factor being whether the alkyl peroxy
999 radicals are primary, secondary, or tertiary. However, the presence of α -OH substitution also appears to
1000 affect the rate constant, and this is also taken into account. Table A-45 in the SI includes a summary of
1001 the averages or estimated rate constants for various types of radicals that can be derived from the
1002 experimental data, and the rate constants used by MechGen when generating mechanisms for peroxy
1003 radicals are included in Table 8.

1004 Because it is not practical to represent all of the peroxy + peroxy combinations in large
1005 mechanisms, MechGen does not assign rate constants or generate mechanisms for all combinations of the
1006 different types of radicals, but instead represents the radicals as reacting with a generic alkyl (RO₂) and a



1007 generic acyl (RCO₃) peroxy radical. The rate constants and also mechanism assignments for the generic
 1008 RO₂ are based on those estimated for secondary peroxy radicals because they are intermediate in
 1009 reactivity and are also expected to be most frequently formed. The generic acyl peroxy radical is based on
 1010 acetyl or propionyl peroxy radicals, the only type for which data are available. Although this
 1011 representation can be inaccurate because it lumps reactions whose rate constants vary by orders of
 1012 magnitude, it is considered to be relatively unbiased, and is better than the alternative of ignoring these
 1013 reactions altogether. Fortunately this is a relatively unimportant loss process for peroxy radicals under
 1014 most conditions, so this is not considered to be a significant problem compared to other uncertainties.

1015 Available information concerning the mechanisms of peroxy + peroxy reactions is discussed in
 1016 Section A1.5.4 of the SI, and recommended and assigned branching are given in Table A-14. Three types
 1017 of generic reactions are considered.



1018 The products formed from the generic radicals are not shown because they are not generated because they
 1019 are represented separately by the reactions of the individual radicals. (This is an approximation because it
 1020 undercounts the contributions of self-reactions in the kinetic differential equations.) Note that the
 1021 "carbonyl" and "alcohol" routes are two instances of the disproportionation reaction, where an α H
 1022 migrates from the carbon next to the peroxide group on one of the radicals to a peroxy oxygen on the
 1023 other radical. Formation of carbonyl products is not possible for acyl radicals and tertiary alkyl radicals,
 1024 and the alcohol route is not possible for generic acyl radicals because they lack the necessary α hydrogen.

1025 The branching ratios used when generating reactions of the various types of peroxy radicals are
 1026 summarized in Table 11. These are based roughly on the branching ratios for individual reactions given in
 1027 Table A-14 in the SI, and extrapolating them to other types of radicals where necessary as discussed in
 1028 Section A1.5.4. These assignments are uncertain when applied to larger radicals and radicals with non-
 1029 alkyl substituents or groups, but the effects of this uncertainty are probably less than the uncertainty and
 1030 necessary inaccuracies when estimating the rate constants.

1031

1032 Table 11. Branching ratios used when generating mechanisms of peroxy + peroxy reactions.

Peroxy radical type	Reaction with Generic RO ₂			Reaction with Generic RCO ₃	
	Alkoxy	Carbonyl	Alcohol	Alkoxy	Alcohol
Primary or secondary RO ₂	0.5	0.25	0.25	0.8	0.2
Tertiary RO ₂	0.5	-	0.5	1.0	-
Any RCO ₃	0.8	-	0.2	1.0	-


1033

1034



1035 **4.6. H-shift Isomerizations**

1036 Organic peroxy radicals can undergo H-shift isomerizations where the radical center abstracts a
1037 hydrogen from elsewhere in the molecule, via a cyclic transition state, forming a hydroperoxide and a
1038 carbon-centered radical.



1040 This reaction has been known to occur at higher temperatures in combustion systems, where the α -
1041 hydroperoxy carbon-centered radicals formed are designated "QOOH", though they had not been
1042 expected to be important at atmospheric temperatures until recently. Recent experimental data indicate
1043 that these reactions occur at measurable rates and may be non-negligible under atmospheric conditions for
1044 certain radicals (e.g., Crouse et al., 2012; Teng et al., 2015; Praske et al., 2018, 2019; Nozière and
1045 Vereecken, 2019). This has been supported by quantum theoretical calculations of rate constants for a
1046 wide variety of such reactions (e.g., Davis and Francisco, 2010; Møller et al., 2019; Vereecken and
1047 Nozière, 2020 and references therein), whose results are reasonably consistent with the available
1048 experimental data and indicate that in some cases these isomerizations would be the dominant fate of such
1049 radicals.

1050 Vereecken and Nozière (2020) reviewed the available theoretical and experimental rate constant
1051 data for peroxy H-shift isomerizations for a wide variety of peroxy radicals, and provided recommended
1052 SARs for predicting their rate constants for automated mechanism generation applications. These consist
1053 of lookup tables giving the rate constant for various pairs of peroxy radical types and H-groups (the
1054 groups with the H that is transferred) involved in the H-shift, giving the rate constants as a function of
1055 transition state ring size, with correction factors for a few types of substituents around the H-group. The
1056 effects of -OH, -OOH, -OR, carbonyl groups, and double bonds at or near the H-group were incorporated
1057 into the lookup tables; separate rate constant recommendations were given for reactions of $-CH_2OO\cdot$, $-$
1058 $>CH(OO\cdot)$ - and $-C(OO\cdot)<$ radicals, and rate constants were given for ring sizes up to 8 or 9. The
1059 recommended correction factors covered the effects of α -ONO₂, β -OH, β -endo oxo and β -exo oxo
1060 substituents on the H-group. These recommendations were adopted as the starting point for use in this
1061 work. However, the recommended lookup tables did not cover all the types of peroxy radicals that might
1062 be formed in atmospheric photooxidation systems, including reactions of acyl peroxy radicals, of radicals
1063 with both β double bonds and α -OH groups, abstractions from an HCO group with a β double bond, H-
1064 shifts with transition state rings larger than 8 or 9, and several other cases. Therefore, it was necessary in
1065 this work to extrapolate or extend the work of Vereecken and Nozière (2020) to allow predictions for
1066 these other types of radicals.

1067 The adaptation of the peroxy H-shift SARs for use in this work is described in detail in Section
1068 A1.5.5 of the SI. In order to provide a basis for estimating rate constants, the various factors affecting the
1069 rate constants are split up into components that are somewhat more straightforward to estimate when
1070 expanding the lookup tables to cover the other cases. The expression used for estimation purposes was:

1071
$$k(T) \approx nH \times A(n,ts) \times e^{-\{Ea^0(H,R) + Ea^{\text{strain}}(H,R,n) + Ea^{\text{corr}}(\text{subs},n)\} / RT} \quad (I)$$

1072 where nH is the number of equivalent abstractable hydrogens; A(n,ts) is an Arrhenius A factor assumed to
1073 be dependent only on the size of the transition state ring, n, and the type of transition state, ts; $Ea^0(H,R)$ is
1074 the activation energy in cases where there is no strain or substituent corrections, and is assumed to be
1075 dependent only on the type of H-group, H, and the type of radical, R; $Ea^{\text{strain}}(H,R,n)$ is the ring strain that
1076 depends on the ring size as well as the types of groups involved; and $Ea^{\text{corr}}(\text{subs},n)$ are corrections for
1077 substituents on the H-group, some of which depend on ring size. The temperature-dependence
1078 recommendations given by Vereecken and Nozière (2020) were used to derive approximate A factors
1079 which were then used, in conjunction with the recommended rate constants in the lookup tables, to create




1080 lookup tables for E_a^0 and for E_a^{strain} for the various ring sizes. These were then used to estimate E_a^0 and
1081 E_a^{strain} for cases not covered by the original SARs. The recommended substituent correction factors were
1082 used to derive E_a^{corr} to use in Equation (I), where applicable. No new types of substituent correction
1083 factors were added in this work.

1084 The range of magnitudes of estimated rate constants for the peroxy H-shift isomerization for the
1085 various types of radicals is presented in Figure 3, which shows the rate constants for the transition state
1086 ring size (generally 6-8) that gives the highest rate constant. These maximum rate constants for the
1087 different types of reaction range in magnitude from $\sim 3 \times 10^{-6}$ to $\sim 7 \times 10^6 \text{ sec}^{-1}$, and the lookup tables cover
1088 25 types of H-groups and transition states, as indicated on the x-axis of the figure. The figure also has
1089 lines giving the upper and lower limits of these rate constants used by MechGen when generating peroxy
1090 radical reactions -- reactions faster the upper limit are assumed to be the only fate of the peroxy radical,
1091 while those that are below the lower limit are treated as negligible and not generated. For those with rate
1092 constants between these limits there is a competition between unimolecular and bimolecular reactions of
1093 these radicals, so the estimated rate constant would have an impact in predicted mechanisms.

1094 The H-shift reaction from unsubstituted $-\text{CH}_3$ and $-\text{CH}_2-$ groups by in alkyl peroxy radicals was
1095 generally negligible, while abstractions from hydroperoxy groups and aldehyde groups and abstractions
1096 of allylic hydrogens by acyl peroxy groups were generally fast. Most of the other types were in the
1097 intermediate range where there is a competition between unimolecular and bimolecular reactions of these
1098 peroxy radicals. This indicates that this is an important process for many radicals that must be considered
1099 when implementing mechanisms into models.

1100 The rate constant estimates that were derived directly from the SARs of Vereecken and Nozière
1101 (2020) are based on results of quantum theory calculations and may be good to within a factor of ~ 10 , and
1102 can predict the experimental 298 K rate constants to within a factor of 3, as shown in Figure A-12 in the
1103 SI. Those that are based on our extrapolations or estimates are much more uncertain, though many of
1104 these are estimated to be high enough that the actual fate of the peroxy radical may not be sensitive to this
1105 estimate. However, experimental and theoretical data are needed to verify or improve our estimated
1106 extensions of the Vereecken and Nozière (2020) SARs as discussed in Section 1.5.5 of the SI, as well as
1107 more experimental data to validate the extensive set of theoretical calculations that form the primary basis
1108 for the SARs that are currently employed.

1109 No attempt was made to make separate estimates of rate constants for H-shift isomerizations of
1110 cyclic peroxy radicals by either Vereecken and Nozière (2020) or in this work. For the purpose of making
1111 general estimates, we assume that the isomerization reaction does not occur if three or more of the atoms
1112 in the transition state ring are in another ring, since they would usually (but not necessarily always) result
1113 in a highly strained structure. Although these are not unreasonable assumptions, some isomerizations will
1114 be overestimated and others overlooked. The practical effects of these uncertainties have not been
1115 assessed, but such an assessment may give an indication for priorities for future theoretical or
1116 experimental studies of unimolecular reactions of peroxy radicals formed from terpenes and other
1117 cycloalkenes.



Nomenclature: H-group with substituents (if applicable), where -Ox = α -OH or -OR substituents; -CH_x=/En: β double bond by the H-group, with at least one double bond group in the transition state ring; -CH_x/Ex: β double bond not in the transition state; -OxDB = both α -Ox and a double bond substituent; /EnEx = two β double bonds by the H-group.


1118

1119 Figure 3. Estimated peroxy H-shift isomerization rate constants for the most favorable ring size for
 1120 various H- and R-group pairs without β -substitution corrections. Also shown are upper
 1121 and lower rate constant limits used during mechanism generation.

1122

1123 **4.7. Ring Closure Reactions of Unsaturated Peroxy Radicals.**

1124 The current estimated mechanisms for aromatics, discussed above in Section 3.4, assume two
 1125 additional types of unimolecular reactions of OH-aromatic-O₂ adducts to account for observed aromatic
 1126 product formation and reactivities, both involving ring formation. These are shown as reactions labeled
 1127 "[b]" and "[c]" in Figure 1. These reactions are accounted for in the process of generating reactions of
 1128 OH-aromatic adducts as discussed in Section 3.4, and thus their rate constants do not need to be estimated
 1129 to generate mechanisms for aromatics. However, these reactions may also be non-negligible in non-
 1130 aromatic systems, since neither the reactants, products, nor the transition states have aromatic rings. The
 1131 reactions shown in Figure 1 can be generalized as follows:




1136 where X is H or any group, and * indicates ring closure. The reactions labeled [b] involve the peroxy
 1137 group adding to a double bond forming a cyclic ether and an allylic carbon-centered radical, while those
 1138 labeled "[c]" have a similar cyclic transition state but the O-O bond breaks, with one O adding to the
 1139 double bond forming an epoxide and the other becoming an alkoxy radical center. Note that the reactions
 1140 labeled [b5] and [c5] involve 5-member ring products or transition states, while those labeled [b7] and
 1141 [c7] involve a 7-member ring. Since these reactions apparently dominate over bimolecular reactions when



1142 formation or intermediacy of both 5 and 7 member rings are involved, one would reasonably expect them
1143 to be even faster in acyclic systems where only one new ring is involved.

1144 Since 6- or 7-member rings tend to have less ring strain than 5 member rings, (see discussion of
1145 peroxy H-shift reactions in Section 1.5.5 of the SI) one would expect the analogue involving formation or
1146 intermediacy of 6 member rings also to be fast, e.g.,



1149 Therefore, to be consistent with the aromatics mechanisms, the above six types of reactions are generated
1150 for peroxy radicals with conjugated double bonds in the α -, β -, or γ - positions, relative to the carbon
1151 bonded to the peroxy group, and are assumed to dominate over bimolecular reactions and H-shift
1152 isomerizations. Because of this, it is not necessary to estimate their total rate constants, only the branching
1153 ratios for the [b] and [c] reaction routes, or the "fEpoxy/fBcyc" ratio using the terminology of Section 3.4
1154 in the context of the aromatic system. The fractions reacting via the [b] or Bcyc route used when
1155 generating aromatics mechanisms ranged from 68% to 88%, depending on the number of *ortho*
1156 substituents in the OH-aromatic adduct. It is highly uncertain whether this branching ratio is an
1157 appropriate basis for estimations for acyclic peroxy radicals with conjugated double bonds, but no
1158 information is available to justify other assumptions. Therefore, when generating mechanisms for peroxy
1159 radicals with conjugated double bonds in the α -, β -, or γ - positions, we assume that:

$$1160 \quad \text{Cyclic peroxide Fraction} = k_{b5}/(k_{b5} + k_{c5}) = k_{b8}/(k_{b6} + k_{c6}) = k_{b7}/(k_{b7} + k_{c7}) \approx 0.75$$

$$1161 \quad \text{Epoxy alkoxy Fraction} = 1 - \text{Cyclic peroxide Fraction} \approx 0.25$$

1162 where the fractions are approximately the averages of those given in Table 7. This is highly uncertain, but
1163 at least it is reasonably consistent with our treatment of aromatic ring addition reactions. However,
1164 formation of these types of peroxy radicals is probably relatively rare in atmospheric systems, so the
1165 practical effect of this uncertainty on atmospheric mechanisms may be relatively small.

1166 5. Reactions of Alkoxy Radicals

1167 Alkoxy radicals are formed in atmospheric oxidation mechanisms primarily from the reactions of
1168 peroxy radicals with NO and NO₃, and to a lesser extent by their reactions with other peroxy radicals, as
1169 discussed above. These can undergo a variety of reactions in atmospheric systems, and this leads to much
1170 of the complexity in atmospheric reactions of organic compounds. The types of alkoxy radical reactions
1171 that MechGen considers, and how their rate constants are estimated, are summarized in Table 12. These
1172 reactions are discussed further below.

1173 Note that alkoxy radicals formed in the reactions of peroxy radicals with NO could be formed
1174 with up to ~12 kcal/mole of excitation energy, which would affect estimates for unimolecular reactions
1175 that are not highly thermochemically favorable (Orlando et al., 2003). This is not taken into account in
1176 current estimates of unimolecular reactions of these radicals, but will need to be considered for future
1177 updates of the system.

1178



1179 Table 12. Summary and examples of reactions of alkoxy radicals as derived using MechGen.

Types	Reactions [a]	Rate Constant Estimation
Reaction with O ₂	-CH ₂ O· + O ₂ → HO ₂ + -CHO -CH(O·) + O ₂ → HO ₂ + -C(O)-	See Sections 5.1 and A1.7.1
	HC(O)O· + O ₂ → HO ₂ + CO ₂	Assumed to be fast [b]
	<u>X</u> ₂ C=C <u>X</u> O· + O ₂ → <u>X</u> ₂ C(OO·)C(O) <u>X</u>	Assumed to be fast [b, c]
β-Scission decompositions	XCH ₂ O· → X· + HCHO XCHX'O· → X· + X'CHO XCX' ₂ O· → X· + X'C(O)X'	See Sections 5.2 and A1.7.2.
	XC(O)O· → X· + CO ₂	Assumed to be fast [b]
H-Elimination decompositions	XCH ₂ O· → XCHO + H· XCH(O·)X' → XC(O)X' + H·	See Sections 5.3 and A1.7.3.
H-Shift isomerizations	HC <u>X</u> ₂ -(Y) _n -C <u>X</u> ₂ O· → ·C <u>X</u> ₂ -(Y) _n -C <u>X</u> ₂ OH	See Sections 5.4 and A1.6.
α Ester and α Nitrate Rearrangement	<u>X</u> CH(O·)OC(O) <u>X</u> ' → <u>X</u> C(O)· + <u>X</u> 'C(O)OH <u>X</u> CH(O·)ONO ₂ → <u>X</u> C(O)· + HONO ₂	See Sections 5.5 and A1.7.4.

1180 [a] The following codes used for unspecified groups: X = Any non-radical group except -H; X = -H or
1181 any non-radical group; Y = Any stable compound group that is bonded to at least two other groups,
1182 except =C=, ≡C-, and -aC_x; Y = As above, but also excluding =CH- and =C<

1183 [b] This is assumed to dominate over competing reactions of this radical, so the rate constant does not
1184 need to be estimated.

1185 [c] Vinyloxy radicals are expected to exist in resonance with a carbon-centered radical form, e.g.,
1186 X₂C=CXO· ↔ X₂C[·]C(O)X, which rapidly adds O₂ under atmospheric conditions.

1187

1188

1189 5.1. Bimolecular Reactions

1190 Primary and secondary alkoxy radicals can react with O₂ via abstraction of an α hydrogen to form
1191 HO₂ and the corresponding carbonyl compound, as shown for the first three reactions in Table 12.
1192 Available data concerning rate constants for this reaction are discussed in Section A1.7.1 of the SI. These
1193 data suggest that the rate constant may only depend on whether the radical is primary or secondary, and
1194 the following rate constants are assigned for estimation purposes:

1195 $k(-\text{CH}_2\text{O}\cdot + \text{O}_2) = 2.38 \times 10^{-14} \exp(-0.38/RT) = 1.25 \times 10^{-14} \text{ cm}^3 \text{ molec}^{-1} \text{ s}^{-1}$ ($k^{\text{uni}} = 6.4 \times 10^4 \text{ sec}^{-1}$)

1196 $k(-\text{CH}[\text{O}\cdot] + \text{O}_2) = 1.95 \times 10^{-14} \exp(-0.59/RT) = 7.27 \times 10^{-14} \text{ cm}^3 \text{ molec}^{-1} \text{ s}^{-1}$ ($k^{\text{uni}} = 3.8 \times 10^4 \text{ sec}^{-1}$)





1197 where the rate constant is given for 298 K and k^{uni} is the pseudo-first order rate constant for 1 atmosphere
1198 of air. Note that separate assignments are given for rate constants for reactions of O₂ with methoxy or
1199 ethoxy radicals (see Table A-56 in the SI), but there is no indication of effects of radical size for larger
1200 radicals, though this cannot be ruled out for C₆₊ radicals. There are also no data on effects of non-alkyl
1201 substituents or groups on this rate constant, so these estimates are highly uncertain for non-alkyl radicals.



1202 Although alkoxy radicals can also react with NO_x and radical species with relatively high rate
 1203 constants, the atmospheric levels of NO_x are many orders of magnitude less than those of O₂, and the rate
 1204 constants are not sufficiently high that these reactions are competitive. Therefore, reaction with O₂ is the
 1205 only type of bimolecular reaction generated for alkoxy radicals. The only exceptions for the types of
 1206 radicals handled by MechGen are tertiary alkoxy radicals, where β-scission decompositions are estimated
 1207 to be favorable, or phenoxy radicals, which are discussed separately in Section 6.1.

1208 5.2. β-Scission Decompositions

1209 Most alkoxy radicals can undergo β-scission reactions, where a bond on the carbon bonded to the
 1210 oxy radical center breaks, forming a carbonyl group and a fragment radical. These types of reactions can
 1211 be represented as:



1216 where the R's are any groups next to the alkoxy radical group except -H. Decompositions of acyl oxy
 1217 radicals (Reaction 4) are expected to be fast and thus are the only fate of these radicals considered in
 1218 MechGen. However, this is not the case for alkoxy radicals in general (Reactions 1-3) where, depending
 1219 on the structure of the radical, these reactions can be very fast and dominate over other reactions, or they
 1220 can be negligible compared to reaction with O₂ or other types of unimolecular reactions such as H-shifts.
 1221 Note that the relative importance of the decomposition reactions affects the extent to which the reactions
 1222 of a VOC leads to fragmentation to lower molecular-weight products, as opposed to becoming more
 1223 oxidized and forming lower volatility products that may contribute to SOA formation.

1224 Available information concerning rate constants for β-scission reactions, and methods for their
 1225 estimation, are discussed in Section A1.7.2 of the SI. These include experimentally measured rate
 1226 constants given by Orlando et al. (2003), theoretically calculated rate constant parameters from
 1227 Vereecken and Peeters (2009), and rate constants derived from results of experimental product studies.
 1228 Based on the data and estimates in these resources, MechGen uses the following equations to estimate
 1229 rate constants for β-scission reactions of alkoxy radicals:

$$1230 \quad k(T) \text{ (sec}^{-1}\text{)} = n \times 1.0 \times 10^{14} \times \exp(-E_a/RT)$$

$$1231 \quad E_a(\text{kcal/mole}) = E_aR(\text{Rad}) + E_aP(\text{ProdType}) + E_aRc(\text{R.Subst}) + E_aPc(\text{P.Subst}) + R.\text{Corr}(\text{Ring})$$

1232 Here k is the unimolecular rate constant at temperature T in sec⁻¹; n is the number of equivalent R₁ groups
 1233 on the radical; 1.0 × 10¹⁴ sec⁻¹ is the estimated A factor for all β-scission reactions; E_a is the calculated
 1234 activation energy that depends on the reaction; and E_aR, E_aP, E_aRc, E_aPc, and R.Corr are estimated or
 1235 adjustable parameters that depend on the reaction. In addition, "Rad" refers to the radical R₁· formed in
 1236 the reaction; "ProdType" refers to whether the carbonyl product formed is formaldehyde, a higher
 1237 aldehyde, or a ketone (reactions 1-3, respectively); "R.Subst" refers to substituents on R₁; "P.Subst" refers
 1238 to substituents R₂ or R₃ if applicable; and "Ring" refers to the presence of rings on the alkoxy radical at or
 1239 adjacent to the radical center. The derived or estimated values of these parameters are given in various
 1240 tables in Section A1.7.2 of the SI. Values of representative parameters are given in Table 13.

1241



1242 Table 13. Representative values of parameters used to estimate activation energy of β -scission
1243 reactions.

Radical Formed [a] Type	EaR	Product Formed		Substituent Corrections			Ring Corrections		
		Type	EaP	Substituent	EaRc	EaRp	Type	Size	R.Corr
Methyl	18.5	HCHO	0	-Alkyl	0	0	Ring is	3	-7.1
Primary	15.5	C ₂₊ Aldehyde	-2.2	-Allyl	-5.0	-2.0	broken	6	0.0
Secondary	12.8	Ketone	-3.9	-C(=O)X	-1.3	0.8			
Tertiary	10.7			-OX	-3 [b]	-4 [b]	Ring on	3	2.0
HC(O).	11.4			-ONO2	0 [b]	-1.1	radical	6	1.8
RC(O).	9.3			-CH _x -O-	0	1.1			
Alkoxy	18 [b]			-CH _x -OH	0	-0 [b]	Ring on	3	7.2
							product	6	-0.3

1244 [a] Parameters are all in units of kcal/mole. "X" refers to -H or any other substituent.




1245 [b] The value of this parameter varies somewhat depending on the specific groups involved, and the
1246 average value is given. See the tables in Section A1.7.2 of the SI for the specific values.
1247
1248

1249 From Table 13 it can be seen that the estimated activation energies range from near zero to ~20
1250 kcal/mole or higher. Note that an activation energy of ~12 kcal/mole would give a rate constant that is
1251 comparable to the pseudo-unimolecular rate constant for the reactions of primary or secondary alkoxy
1252 radicals with O₂, so reactions with higher activation energies would generally be negligible, while those
1253 with lower Ea values could potentially be the dominant process unless another type of unimolecular
1254 reaction (e.g., a β -scission involving a different group or an H-shift) is also estimated to be fast.
1255 Decompositions are usually important for tertiary alkoxy radicals not only because reactions with O₂ are
1256 not possible, but because their β -scission decompositions are generally estimated to be relatively fast.
1257 Decompositions can also be important for primary or secondary radicals.

1258 As discussed in the SI, there are sufficient data on rate constants for decompositions of alkoxy
1259 radicals formed from alkane systems that estimates in these cases are not considered to be highly
1260 uncertain. However, if non-alkyl groups are present, data to derive estimates are highly limited or lacking
1261 and thus needed. One uncertainty is the lack of measured or theoretically calculated rate constants for
1262 decompositions forming alkoxy radicals, which might occur in atmospheric photooxidation mechanisms
1263 of ethers and esters and compete with other reactions discussed below.

1264 5.3. H-Elimination Decompositions

1265 Primary and secondary alkoxy radicals can also undergo β -scission reactions by H-elimination,
1266 forming a carbonyl product and a hydrogen atom. Available information and estimates for the rate
1267 constants for these reactions are discussed in Section A1.7.3 of the SI. The rate constants for these
1268 reactions appear to be enhanced by the presence of α -oxygen substituents, so three types of these
1269 reactions are considered.



1273 Here X is -H or any group, and R is any group except -OX. The data are limited, but sufficient to derive
1274 the following expression to estimate the rate constant:



$$\begin{aligned} 1275 \quad & k(T) \text{ (sec}^{-1}\text{)} = n \times 3.3 \times 10^{13} \times \exp(-E_a/RT) \\ 1276 \quad & E_{a1} = E_{a2} \text{ (kcal/mole)} = 13.5 + 0.52 \Delta H_r \\ 1277 \quad & E_{a3} \text{ (kcal/mole)} = 11.5 + 0.52 \Delta H_r \end{aligned}$$

1278 where E_{a1} , E_{a2} , E_{a3} refer to the activation energies for reactions of types 1, 2, and 3, respectively. The
1279 estimate for E_{a3} is uncertain and is strictly speaking an upper limit -- i.e., rate constants for H-elimination
1280 reactions of alkoxy radicals with two -OX may be higher than estimated.

1281 Note that under atmospheric conditions, the H-elimination reactions yield exactly the same
1282 products as the reactions the alkoxy radicals with O_2 , because the eliminated H \cdot rapidly forms HO_2 . These
1283 estimates indicate that the H-elimination reaction is negligible compared to O_2 reaction if there are no α -
1284 O substituents, non-negligible but slower than O_2 reaction if there is only one, and faster than the O_2
1285 reaction if there are two such substituents. Therefore, this reaction is only important in the third case, so
1286 only estimates for E_{a3} have a practical effect on generated mechanisms.

1287 **5.4. H-Shift isomerizations**

1288 Alkoxy radicals with sufficiently long chains ($\approx C_4$ or larger) can also undergo H-shift
1289 isomerization reactions, where the alkoxy group abstracts an H from elsewhere in the molecule via a
1290 cyclic transition state, forming a carbon-centered radical with an -OH substituent. These are similar to the
1291 H-shift isomerizations that were previously discussed for peroxy radicals (Section 4.6), but in this case
1292 the reactions are much more exothermic, and their importance in atmospheric systems has been
1293 recognized for some time. Methods to estimate their rate constant have been developed for previous
1294 versions of the SAPRC mechanism, with the version used for developing SAPRC-99 being the most
1295 completely documented (Carter, 2000). The method used for SAPRC-99 has been subsequently updated
1296 for SAPRC-18 to incorporate evaluated experimental data (Atkinson, 2007; IUPAC, 2023), results of
1297 theoretical quantum chemistry calculations of Vereecken and Peeters (2010), and upper limit rate
1298 constants derived from various product studies, as given in Table A-50 of the SI. This updated method is
1299 documented in the SI in Section A1.6.3.

1300 The rate constants for alkoxy H-shift reactions are estimated in the same way as for peroxy
1301 radicals, using Equation (I) in Section 4.6, which is based on the types of groups from which the H is
1302 abstracted, their substituents, and also substituents elsewhere in the transition state ring. Because of more
1303 limited amount of theoretically calculated data available at the time this method was last updated in the
1304 mid-2010's, only a single set of ring-size-dependent A factors and E_a^{strain} values were derived, and E_a^0
1305 values were derived using only four types of H-groups (-CH₃, -CH₂-, -CH<, and -OOH), with no
1306 distinction being made between primary, secondary, or tertiary alkoxy radicals. To compensate for this, a
1307 larger number of substituent correction factors were employed. The A factors and strain energies used are
1308 given in Table A-25, the E_a^0 values are given in Table A-26, and the E_a^{corr} values are given in Table A-16
1309 and Tables A-27 and A-28 in the SI. Footnotes to these tables indicate how these were derived or
1310 estimated.

1311 Estimated 298 K rate constants for H-shift isomerizations of various unsubstituted alkoxy radicals
1312 are listed in Table 14, which also shows the ratios of these rate constants to the pseudo-unimolecular rate
1313 constant for the reaction of secondary alkoxy radicals with O_2 in the atmosphere. It can be seen that 1,4-H
1314 shift isomerizations, with 6-member ring transition states, are ~ 10 -350 times faster than O_2 reaction, while
1315 those involving 1,5-H shifts are ~ 4 -130 times faster, while 1,2, 1,3, and 1,8+ H shifts are negligible to
1316 minor.



1317 Note that the current estimation methods do not consider effects of functional groups that are
 1318 between the radical center and the H atom being transferred. This may be considered in the future as more
 1319 experimental or theoretical data become available.

1320 Although there are uncertainties in these estimates, particularly if non-alkyl groups or substituents
 1321 are present, their effects on generated mechanisms would be minor if these reactions are estimated to be
 1322 either very slow or very fast compared to competing processes. Uncertainties will affect generated
 1323 mechanisms primarily when there are competing β -scission decompositions with comparable estimated
 1324 rate constants.

1325

1326

1327 Table 14. Representative 298 K rate constants for H-shift isomerizations of unsubstituted alkoxy
 1328 radicals.


Ring size	Shift from -CH ₃		Shift from -CH ₂ -		Shift from -CH<		Shift from -CHO	
	k(sec-1)	k/k[O ₂]	k(sec-1)	k/k[O ₂]	k(sec-1)	k/k[O ₂]	k(sec-1)	k/k[O ₂]
5	8.9e-3	2e-7	8.5e-2	2e-6	2.8e-1	8e-6	3.3e-1	9e-6
6	3.5e+5	9.30	3.3e+6	89	1.1e+7	295	1.3e+7	346
7	1.3e+5	3.48	1.2e+6	33	4.1e+6	110	4.9e+6	130
8	4.3e+2	0.01	4.1e+3	0.1	1.4e+4	0.4	1.6e+4	0.4
9	7.2e+1	2e-3	6.8e+2	0.02	2.3e+3	0.06	2.7e+3	0.07

1329

1330



1331 5.5. Ester and Nitrate Rearrangements

1332 Alkoxy radicals with an α hydrogen and either an ester or nitrate group in the α position can
 1333 undergo a second type of H-shift isomerization, where the α hydrogen transfers to the carbonyl or nitrate
 1334 oxygen and the bonds rearrange to form a carbonyl radical and a carboxylic or nitric acid, via a 5-member
 1335 ring transition state, e.g.,:



1336

1337 Evidence for these reactions and their measured or estimated rate constants are discussed in Section
 1338 A1.7.4 of the SI. Based on limited data, the following rate constants are estimated:



1342 This predicts that the ester rearrangement of primary alkoxy radicals with ester substituents occurs at a
 1343 similar rate as reaction with O₂ under atmospheric conditions, while it is the dominant fate for most
 1344 secondary alkoxy radicals with ester substituents, being predicted to be faster than possible competing H-
 1345 shift or β -scission decompositions, as well as the reaction with O₂. On the other hand, the nitrate
 1346 rearrangement is predicted to be slower than the O₂ reaction, so it is not expected to be important in
 1347 atmospheric systems.



1348 Formation of α -nitrate alkoxy radicals that might undergo the nitrate rearrangement is not
1349 expected to be as important in atmospheric oxidation systems, because α -nitrate carbon-centered radicals
1350 formed in reactions of organic nitrates that could be precursors to these radicals rapidly decompose by
1351 eliminating NO_2 , as discussed above in Section 3.2.1, rather than reacting to form the alkoxy radical.
1352 Therefore, any uncertainties in estimating rate constants for this reaction are not expected to be important.

1353 Radicals that can undergo the ester rearrangement are predicted to be formed in the atmospheric
1354 oxidation of esters, so this reaction is a factor in predicting their atmospheric mechanisms. The
1355 uncertainties in the estimated rate constant have a significant effect on predictions for esters forming
1356 primary alkoxy radicals because it is competitive with the O_2 reaction, but is not as important for those
1357 forming secondary radicals because the ester rearrangement is predicted to dominate.


1358 6. Reactions of Other Types of Radicals

1359 The previous sections focused on the reactions of unsubstituted and substituted alkyl radicals, and
1360 the peroxy and alkoxy radicals formed from them in the presence of O_2 and NO_x , which are the major
1361 types of radicals formed in atmospheric oxidation systems. Other types of radicals are also formed in
1362 some circumstances and the treatments of these are discussed in the subsections below. Note that CIs are
1363 discussed separately in Section 7.

1364 6.1. Phenoxy Radicals

1365 Phenoxy radicals are formed in the reactions of phenols, aromatic aldehydes, and potentially
1366 other aromatic compounds. They cannot react with O_2 or undergo any of the unimolecular alkoxy radical
1367 reactions discussed in Section 5. Therefore, their only atmospheric fate is expected to be reactions with
1368 NO_x species or other radicals. MechGen assumes that their major sinks in atmospheric systems are either
1369 reaction with O_3 , NO_2 or HO_2 . Reactions with NO are ignored because they are expected to form nitroso
1370 species or nitrites that would rapidly photolyze to re-form the reactants, and because the concentrations of
1371 NO are usually lower than NO_2 in the atmosphere.

1372 MechGen assumes that the mechanisms for the reactions of phenoxy with O_3 and HO_2 are as
1373 expected for general alkoxy systems and involve simple O or H-transfers, as follows:




1376 The rate constants are in $\text{cm}^3 \text{ molec}^{-1} \text{ s}^{-1}$ and that for O_3 is from Tao et al. (1999) and that for HO_2 is
1377 estimated to be the same as for methoxy + HO_2 (Assif et al., 2018). The O_3 reaction is not a net sink for
1378 phenoxy radicals because they can be regenerated by reactions of phenyl peroxy radicals with NO . This
1379 reaction will also convert NO to NO_2 , which can also regenerate O_3 under atmospheric conditions. On the
1380 other hand, the reaction with HO_2 is a net sink for both HO_2 and phenoxy, though it will regenerate the
1381 phenol whose reactions are the source of the phenoxy radicals.

1382 The reactions of phenoxy radicals with NO_2 are expected to be the main sink for these radicals in
1383 the presence of NO_x . Alkoxy radicals are expected to react with NO_2 forming the corresponding nitrate,
1384 but for most alkoxy radicals this is minor compared to reaction with O_2 or unimolecular reactions, so
1385 these reactions can be ignored. In the case of phenoxy, the analogous formation of benzyl nitrate is not
1386 expected to be an overall sink for the phenoxy radical because the BzO-NO_2 bond is so weak (Batiha et
1387 al., 2012) that the phenyl nitrate would be expected to rapidly decompose to reactants resulting in no net



1388 reaction, analogous to the formation of alkyl peroxy nitrates discussed in Section 2.3. Instead, the reaction
1389 is assumed to occur as follows:



1390 where the rate constant is from Platz et al. (1998). If the initially formed product has an H atom on the
1391 same carbon where the NO₂ added it will undergo a fast unimolecular rearrangement to a nitrophenol as
1392 shown above and discussed in Section 2.3.2. This is consistent with theoretical studies (e.g., Batiha et al.,
1393 2012) and products formed in reactions of phenols with NO₃ (e.g., Calvert et al., 2002, 2011). However,
1394 this fast reaction is not possible if NO₂ adds to a carbon with another substituent, in which case a 2-nitro-
1395 3,5-cyclohexadienone with a substituent in the 2-position is predicted to be the final product.


1396 The presence of substituents on the aromatic rings are assumed not to affect the rate constants or
1397 mechanisms of the bimolecular reactions of phenoxy radicals. Note that the *ortho* NO₂ addition can occur
1398 in two different positions for unsymmetrical molecules, and currently MechGen assumes that the reaction
1399 at each position is equally likely, regardless of substituents.

1400 6.2. Nitrogen-Centered Radicals

1401 Nitrogen centered radicals can be formed in the atmospheric reactions of some amines, as
1402 discussed in Section 2.1.3 above. Those with α hydrogens are assumed to react primarily with O₂,
1403 forming HO₂ and the corresponding amide,



1405 while those lacking α hydrogens are assumed to undergo only bimolecular reactions, as follows:




1409 where the rate constants are in units of cm³ molec⁻¹ s⁻¹ and are at the high pressure limit where applicable.
1410 The high pressure rate constants for the reactions with NO and NO₂ are estimated to be the same as the
1411 IUPAC (2023) recommendation for their reactions with ethoxy radicals, while the rate constant for the
1412 reaction with HO₂ is estimated to be approximately the same as given in the NIST kinetics database for
1413 methoxy + HO₂ (Sander et al., 2009). MechGen does not generate reactions of amino radicals without α
1414 hydrogens, but they would only be formed from reactions of amines with more than one tertiary
1415 substituent, which do not appear to be important in current atmospheric emissions.

1416 Note that this mechanism predicts that formation of N-centered radicals in the reactions of
1417 amines without α hydrogens is a radical terminating process, which means that such amines tend to be
1418 radical inhibitors when they react in the atmosphere. This is consistent with results of environmental
1419 chamber experiments with t-butyl amine and 2-amino-2-methyl-1-propanol (Carter, 2008). This is not
1420 expected to be the case with other amines, where results of chamber experiments do not indicate
1421 significant radical inhibition in their reactions.



1422 **6.3. Carbenes**


1423 Carbenes are predicted to form in some photolysis reactions, so their reactions also need to be
 1424 considered. It is assumed that their major fate is the addition of O₂ to form highly excited Criegee
 1425 intermediates (CIs), e.g.,



1427 where "{excited}" indicates excitation. Reactions of excited and stabilized CIs are discussed in the
 1428 following section. Note that CIs formed from reactions of O₂ with carbenes are expected to have much
 1429 higher levels of excitation than those formed from the reactions of O₃ with alkenes, which affects
 1430 estimates of stabilization in some cases.

1431 **7. Reactions of Criegee Intermediates**

1432 CIs are highly reactive intermediates with general structure X₂COO that are formed primarily in
 1433 the reactions of O₃ with alkenes. These can have either zero, one, or two substituents and those with one
 1434 or two different substituents can exist in either the *syn* or *anti* form depending on the configuration of the
 1435 substituent(s) relative to the two oxygens:



1437 They are also formed to a lesser extent from the reactions of O₃ with alkynes, and in the reactions of O₂
 1438 with carbenes formed in some photolysis reactions. In all cases they are expected to be formed in a
 1439 vibrationally excited state, with the amount of excitation depending on the formation reaction, but
 1440 sufficient to allow rapid interconversion between the *syn* and *anti* forms where applicable. The
 1441 interconversion between *syn* and *anti* is assumed to be slow once the intermediate is stabilized. The
 1442 excited intermediates can undergo unimolecular decompositions or be collisionally stabilized, and the
 1443 stabilized intermediates can also undergo unimolecular reactions or react with water, though reactions of
 1444 stabilized intermediates involving interactions between the substituent and the Criegee oxygens requires
 1445 that the substituent be in the *syn* configuration.

1446 Information and estimates about the reactions of excited and stabilized CIs are discussed in
 1447 Section A1.9 of the SI. Briefly, the mechanisms adopted for MechGen are based largely on the IUPAC
 1448 (2023) reviews of the reactions of O₃ with ethene and alkenes with methyl substituents, and on the
 1449 theoretical calculations of Vereecken et al. (2017) on the possible unimolecular reactions of the
 1450 intermediates. Table 15 lists the types of intermediates and the reactions and branching ratios considered
 1451 for them. The types of reactions considered are summarized below.

1452 Excited intermediates with an alkyl group with α hydrogens, and also stabilized intermediates
 1453 with such a group in the *syn* position, are assumed to undergo the vinyl hydroperoxide (VHP)
 1454 rearrangement, where the α hydrogen transfers to the OO group forming an excited VHP, which then
 1455 decomposes to OH and an α-carbonyl radical (e.g., reactions 8, 24, 25-27, 36, and 38 in Table 15).
 1456 Vereecken et al. (2017) calculated these were relatively fast even for stabilized intermediates.



1457 Table 15. Summary of types of Criegee intermediates and the reactions generated for them

Criegee Type [a]	Reactions [b]	Branching Ratios, notes [c]		
<u>Excited CH₂[OO]</u>	1 Stabilization	0.42	1	
	2 → HCO· + OH	0.17		
	3 → CO ₂ + H ₂	0.18		
	4 → CO + H ₂ O	0.18		
	5 → CO ₂ + 2 H	0.05		
<u>Excited RCH[OO]</u> , where R = X ₂ CH-	6 Stab. to X ₂ CHCH[OO <i>syn</i>]	0.33 x SF ₁	2, 3, 4	
	7 Stab. to X ₂ CHCH[OO <i>anti</i>]	0.67 x SF ₁		
	8 → ·CX ₂ CHO + OH	0.49 x (1 - SF ₁)	2	
	9 → R· + ·C(=O)OH	0.10 x (1 - SF ₁)		
	10 → RH + CO ₂	0.26 x (1 - SF ₁)		
	11 → ROH + CO	0.14 x (1 - SF ₁)		
	R = OH	12 → HC(=O)O· + OH	Reaction fast	5
	R = CHX ₂ -CX'=CX"-	13 → ·CX ₂ CX'=CX"CHO + OH	Reaction fast	5
	R = CX ₂ =CX-	14 Stab. to CX ₂ =CXCH[OO <i>syn</i>]	0.33 x SF ₁	3, 4
		15 Stab. to CX ₂ =CXCH[OO <i>anti</i>]	0.67 x SF ₁	
	R = none of the above	16 → ·CX ₂ C(=O)X + ·C(=O)H	1 - SF ₁	6
17 Stab. to RCH[OO <i>syn</i>]		0.33 x SF ₁	3, 4	
18 Stab. to RCH[OO <i>anti</i>]		0.67 x SF ₁		
19 → R· + ·C(=O)OH		0.10 x (1 - SF ₁)	7	
20 → RH + CO ₂		0.26 x (1 - SF ₁)		
21 → ROH + CO		0.14 x (1 - SF ₁)		
SF ₁ (nC) = min (1.0, 0.21 + 0.065 x nC); SF ₁ (2) = 0.35; SF ₁ (12) = 1.0			4	
<u>Excited R₁C[OO]R₂</u> , where R ₁ = X ₂ CH- and R ₂ = X ₂ CH-	22 Stab. to R ₁ -syn-C[OO]R ₂	0.5 x SF ₂	4	
	23 Stab. to R ₁ C[OO]-syn-R ₂	0.5 x SF ₂		
	24 → ·CX ₂ C(=O)R ₂ + OH	0.5 x (1 - SF ₂)	8	
	25 → R ₁ C(=O)CX' ₂ · + OH	0.5 x (1 - SF ₂)		
	R ₁ = OH, any R ₂	26 → R ₂ C(=O)O· + OH	Reaction fast	8
R ₁ = CHX ₂ -CX'=CX"-, R ₂ not OH	27 → ·CX ₂ CX'=CX"C(=O)R ₂ + OH	Reaction fast	8	
R ₁ = CX ₂ =CX-; R ₂ = not OH or CHX ₂ -CX=CX-	28 Stab. to CX ₂ =CX-syn-C[OO]-R ₂	0.5 x SF ₂	4	
	29 Stab. to CX ₂ =CX-C[OO]-syn-R ₂	0.5 x SF ₂		
	30 → ·CX ₂ C(=O)X + ·C(=O)R ₂	1 - SF ₂	6	
R ₁ = X ₂ CH- and R ₂ none of above	31 Stab. to R ₁ -syn-C[OO]R ₂	0.5 x SF ₂	4	
	32 Stab. to R ₁ C[OO]-syn-R ₂	0.5 x SF ₂	8	
	33 → ·CX ₂ C(=O)R ₂ + OH	1 - SF ₂		
R ₁ , R ₂ = none of the above	34 Stab. to R ₁ -syn-C[OO]R ₂	0.5	9	
	35 Stab. to R ₁ C[OO]-syn-R ₂	0.5	9	
SF ₂ (nC) = min (1.0, 0.13 + 0.072 x nC); SF ₂ (3) = 0.35; SF ₂ (12) = 1.0				
<u>Stabilized X₂CHCH[OO <i>syn</i>]</u>	36 → ·CX ₂ CHO + OH	Reaction fast	5	
<u>Stabilized X₂C=CXCH[OO <i>syn</i>]</u>	37 → ·CX ₂ C(=O)X + HC(=O)·	Reaction fast	6	
<u>Stabilized X₂CH-syn-C[OO]R₂</u>	38 → ·CX ₂ C(=O)R ₂ + OH	Reaction fast	6	



Table 15 (continued)

Criegee Type [a]	Reactions [b]	Branching Ratios, notes [c]
Stabilized $X_2C=CX\text{-syn-C[OO]R}$	39 $\rightarrow \cdot CX_2C(=O)X + \cdot C(=O)R$	Reaction fast 6
Other stabilized $XC[OO]X'$	40 $+ H_2O \rightarrow X_2C(OH)OOH$	If H_2O option 10
	41 $\rightarrow *X_2COO*$ (dioxirane)	If not H_2O option 10

1458 [a] X = H or any group; R = any group other than -H or as indicated below; "*syn*" and "*anti*" refer to the
 1459 orientation of the group with respect to the C-OO bond; "*" refers to ring closure; SF is stabilization
 1460 fraction; and nC is the number of carbons in the intermediate.

1461 [b] The first column gives the reaction numbers, which are referenced in the discussion in the text.

1462 [c] Notes for the derivations of the branching ratios are as follows.

1463 1 The reactions and branching ratios of excited $CH_2[OO]$ are based on the recommendations of
 1464 IUPAC (2023) in their review of ethene + O_3 reactions.

1465 2 The reactions and branching ratios of excited $CH_3CH[OO]$ are based primarily on the
 1466 recommendations of IUPAC (2023) in their review of the reactions of O_3 with propene and the 2-
 1467 butenes, with some adjustments as discussed in the SI.

1468 3 The assumption that the *anti* isomer is favored is necessary to improve simulations of OH yields
 1469 from some alkenes. The assumption that the *syn* isomer is formed $\sim 1/3$ of the time gives
 1470 predictions that are reasonably consistent with the data.

1471 4 The stabilization fractions are assumed to increase linearly with the size of the intermediate from
 1472 35% for methyl-substituted intermediates to 100% for intermediates with 16 carbons. This
 1473 improves model predictions but is somewhat arbitrary and other methods for estimating size
 1474 dependence may perform as well or better. It is also assumed that the stabilization fraction is not
 1475 affected by how the CI are formed, except that SF is assumed to be zero for the highly excited
 1476 intermediates formed from $O_2 +$ carbene reactions.

1477 5 These reactions are assumed to be fast based on the calculations of Vereecken et al. (2017) and
 1478 are assumed to dominate over stabilization or competing reactions of the excited intermediates
 1479 when they are possible.

1480 6 These reactions are assumed to be fast for stabilized as well as excited intermediates based on the
 1481 calculations of Vereecken et al. (2017), though not so fast that they dominate over stabilization of
 1482 excited radicals where these are possible. Vereecken et al. (2017) also calculated that they are
 1483 more rapid than the VHP rearrangement, so they dominate in excited disubstituted radicals where
 1484 both are possible.

1485 7 The branching ratios for the "hot acid" decompositions are derived from those used for methyl-
 1486 substituted intermediates, with the yield of the vinyl hydroperoxide (VHP) reaction set to zero
 1487 and the yields of the others adjusted upward to take this into account.

1488 8 The VHP reaction is assumed to be the only fate of this excited intermediate other than
 1489 stabilization. If two different reactions are possible they are assumed to have equal probability;
 1490 otherwise the only possible VHP reaction dominates.

1491 9 Stabilization is assumed to dominate if the VHP, "hot acid" or the fast decomposition reactions
 1492 are not possible.

1493 10 Rate constants for these reactions are not estimated for this version, so the " H_2O " option
 1494 determines the major fate assumed for unreactive stabilized intermediates. Note that this ignores
 1495 loss by photolysis, which may be non-negligible.




1496 Vereecken et al. (2017) calculated that intermediates with -OH substituents, rare in the
1497 atmosphere, will very rapidly rearrange by the O from the -OH group transferring to the [OO] group and
1498 then decomposing to OH and an acyl oxy radical (which then decomposes to form CO₂ and a radical)
1499 (reactions 12 and 26 in Table 15). These reactions are assumed to be so rapid that they dominate over
1500 stabilization.

1501 Excited intermediates with a double bond group in the α position, and also stabilized
1502 intermediates with such a group in the *syn* position, can react in one of two ways depending on whether
1503 there is an abstractable hydrogen in the γ position. If there is such a hydrogen, it is assumed that the
1504 Criegee group abstracts the γ hydrogen to form a highly excited allylic hydroperoxide, which then rapidly
1505 decomposes to form OH and an allylic carbonyl; shown as a single process in Table 15, reactions 13 and
1506 27. Vereecken et al. (2017) calculated this to be extremely fast, and we assume that this is sufficiently fast
1507 that stabilization does not have a chance to occur. If there are no γ hydrogens, it is assumed that the
1508 Criegee group adds to the double bond forming a highly excited unsaturated peroxy ether, which then
1509 rearranges and decomposes to ultimately form the ring opening products shown as reactions 16, 30, 37
1510 and 39 in Table 15. Vereecken et al. (2017) calculates that internal cyclization reactions are more rapid
1511 than the VHP rearrangement, so it is assumed that cyclization is the major process in excited disubstituted
1512 radicals where both are possible. Note, however, that Vereecken et al. (2017) only calculated rate
1513 constants for the formation of the excited unsaturated peroxy ether, but we assume that the subsequent
1514 rearrangement and decomposition is also fast (see Section A1.9.2 in the SI).

1515 Excited intermediates with zero or one substituent can undergo "hot acid" rearrangements where
1516 an H bonded to the Criegee group is involved in a rearrangement forming a highly excited acid, which
1517 then decomposes in several ways, forming either radicals or stable compounds (e.g., reactions 1-5, 9-11,
1518 and 19-21 on Table 15). These reactions are assumed only to occur when the intermediate is excited.

1519 Excited CIs with α -carbonyl groups are assumed to be in rapid equilibrium with the primary
1520 ozonide and the isomer where the O is transferred to the other group, forming a different carbonyl-
1521 substituted intermediate if the groups bonded to them are different, i.e.:



1522

1523 It is assumed that the most reactive isomer determines the subsequent reactions, or that reaction of either
1524 form is equally likely if they undergo the same types of reactions. Although Vereecken et al. (2017)
1525 discussed a possible reaction forming primary ozonides, they apparently did not consider this O-atom
1526 rearrangement reaction.

1527 How MechGen processes stabilized intermediates that cannot undergo the above reactions
1528 depends on whether the "H₂O" option is specified in MechGen. If it is, they are assumed to react with
1529 H₂O, forming an α -hydroxy hydroperoxide (reaction 40 in Table 15). If not, they are assumed to
1530 isomerize to the corresponding dioxirane (reaction 41). MechGen treats both of these as non-radical
1531 products when generating mechanisms. MechGen does not consider effects of substituents in these cases.

1532 MechGen currently does not consider photolysis of stabilized CIs, but loss by photolysis may
1533 well be non-negligible for the less reactive intermediates. IUPAC (2023) evaluated data concerning
1534 photolyses of the simpler Criegee intermediates and gave recommended cross sections for unsubstituted
1535 and methyl substituted intermediates. They also recommended assuming unit quantum yields for the
1536 photolysis forming O³P and the corresponding carbonyl. The recommended absorption cross sections
1537 correspond to rate constants for loss by photolysis of $\sim 0.15 \text{ sec}^{-1}$ with direct overhead sun, which is faster



1538 than the estimated rate constants calculated by Vereecken et al. (2017) for the dioxirane-forming reaction,
1539 but may not be faster than loss by reaction with H₂O, depending on the intermediate and the environment.
1540 Therefore, photolysis of the less reactive stabilized CIs may need to be considered in future versions of
1541 MechGen.

1542 The stabilization factors and branching ratios are given in Table 15 for the various types of CIs,
1543 and footnotes indicate how they were derived. More details are given in Section A3.7 of the SI. The
1544 branching ratios used for unsubstituted and methyl-substituted intermediates are based primarily on
1545 IUPAC (2023) recommendations and include VHP and "hot acid" reactions of the excited intermediate,
1546 with the stabilization fraction being 42% for H₂C[OO] and 35% for the two methyl-substituted
1547 intermediates. It is assumed that the VHP reactions also occur with stabilized intermediates if the
1548 substituent with the group with the α hydrogen is in the *syn* orientation.

1549 Estimates for larger alkyl-substituted intermediates are more uncertain, but the reactions are
1550 assumed to be analogous to those of the methyl-substituted intermediates as long as the VHP reaction is
1551 possible. If the VHP reaction is not possible, then only the "hot acid" reactions are assumed for
1552 monosubstituted intermediates, and only stabilization is assumed to occur for disubstituted intermediates,
1553 where the hot acid reactions are not possible.

1554 The largest uncertainties concern stabilization fractions for the larger CIs, or intermediates
1555 formed in reactions where there is not a separate carbonyl co-product, such as reactions of O₃ with double
1556 bonds in a ring or with triple bonds. As discussed in the SI, best fits to the observed OH yields in O₃ +
1557 alkene reactions (Calvert et al., 2000) and in simulations of NO oxidation and O₃ formation rates in some
1558 chamber experiments with alkenes (e.g., see Carter, 2000, 2010b) are obtained if it is assumed that (1)
1559 stabilization increases with the size of the molecule, (2) stabilization is the same even if all of the energy
1560 is in the intermediate formed, and (3) formation of the *anti* isomer is favored when monosubstituted
1561 Criegees are stabilized, with the enhancement (somewhat arbitrarily) being assumed to be a factor of two.
1562 The assumption that stabilization energies do not affect stabilization of intermediates formed with higher
1563 excitation energies is questionable but is necessary to avoid significantly overpredicting OH yields from
1564 cycloalkenes.

1565 On the other hand, we assume that CIs formed from reactions of carbenes with O₂ are so highly
1566 excited that they react before they are stabilized. In the case of unsubstituted carbenes (reactions 1-5 in
1567 Table 15), we use a branching ratio of 0 for reaction 1 and increase the others accordingly. In the case of
1568 substituted intermediates, the branching ratios are as shown in A-33, calculated using SF₁ or SF₂=0. This
1569 is uncertain, but carbene formation is generally not predicted except in some photolysis reactions (see
1570 Section 2.2.1).

1571 MechGen does not consider all the many possible reactions of CIs or complications considered by
1572 Vereecken et al. (2017) or other more recent theoretical or experimental results, somewhat arbitrary
1573 assumptions had to be made in some cases, and treatments of unreactive stabilized intermediates is
1574 probably over-simplistic. MechGen also does not consider bimolecular reactions with species other than
1575 H₂O, though such reactions may be non-negligible in some circumstances, such as some laboratory
1576 experiments. Therefore, treatments of reactions of CIs must be considered to be uncertain, and are an
1577 appropriate area to update in future versions of MechGen.

1578 **8. Examples of Results**

1579 Although application and analysis of mechanisms for individual compounds are beyond the scope
1580 of the present paper, examples of generated mechanisms, including numbers of reactions and numbers




1581 and types of products, are given for illustrative purposes. We generated mechanisms and product yields at
1582 representative atmospheric NO_x and radical levels (see Section A3 of the SI) for 38 representative
1583 compounds. These include propane, the C₄, C₈, C₁₂ and C₁₆ n-alkanes, 2-methyl alkanes, 1-alkenes, 2-
1584 alkanes, other representative C₈ branched alkanes, alkenes, aromatics, and oxygenates, and representative
1585 terpenes. Note that the generated mechanisms considered only the reactions of the starting compounds
1586 and the radicals they form, not the subsequent reaction of the non-radical products formed. Therefore,
1587 these are only single generation mechanisms; multi-generation mechanisms, such as those derived using
1588 GECKO-A (Aumont et al., 2005) will predict many more products. The specific compounds and selected
1589 results for these compounds are given in Table A-36.

1590 The actual number of first-generation products whose impacts need to be considered, and whose
1591 mechanisms should be included when deriving multi-generation mechanisms, will depend on the
1592 maximum total yield of "negligible" products considered acceptable to ignore. The mechanisms derived
1593 for these examples were generated using a minimum yield cutoff parameter that determines the minimum
1594 branching ratios at a single step, but because of multi-step mechanisms this could still result in predictions
1595 of large numbers of very low yield products whose total yields may actually be negligible. An alternative
1596 approach that deals with this is to determine which products are needed to account for a set fraction of the
1597 carbon of the reacting molecule.

1598 Regardless of the cutoff criteria employed, the determination of which pathways can be neglected
1599 when generating mechanisms also depends on the environment where the compound reacts. That is
1600 because the mechanisms have many competitions between different bimolecular reactions and
1601 bimolecular and unimolecular reactions that depend on concentrations of atmospheric species such as OH,
1602 O₃, NO_x, HO₂ and other peroxy radicals. The conditions used in these examples are based roughly on
1603 results of representing moderately low NO_x levels where peroxy + NO reactions are important, but peroxy
1604 + peroxy reactions are non-negligible, representing conditions where changing VOCs and NO_x inputs
1605 have approximately the same relative effects on ozone. The details of these conditions and the mechanism
1606 generation procedures are provided in Section A3.1 of the SI.

1607 The method used to derive product yields for the selected set of environmental conditions could
1608 not handle cases where reactive radical intermediates re-formed themselves in a sequence of fast radical
1609 reactions. This situation occurs in mechanisms where phenoxy and phenyl peroxy radicals interconvert
1610 due to their reactions with O₃ and NO, (Section 6.1) and when HO₂-substituted peroxy radicals undergo
1611 rapid peroxy-hydroperoxy scrambling reactions (Section 4.6). This problem was avoided when generating
1612 these example mechanisms by disabling predictions of the reactions of phenoxy with O₃ and by disabling
1613 the peroxy/hydroperoxy scrambling reactions. The former affected predictions of nitrophenol / phenolc
1614 product yields from phenols and catechols, while the latter could affect predictions of HO₂-containing
1615 product compounds formed from HO₂-substituted peroxy radicals to some (usually minor) extent. The
1616 overall importance of ignoring the peroxy/hydroperoxy scrambling reactions has not been assessed, but it
1617 is expected to be relatively minor.

1618 Selected results of the example mechanism generations are given in Table A-36 in the SI and in
1619 figures in this section. Figure 4 shows numbers of products generated using various methods against
1620 numbers of carbons in the compounds; Figure 4a shows the total number of products generated using



1621
1622 Figure 4. Numbers of products generated (or ratios thereof) as a function of carbon number for two
1623 different minimum yield parameters, 0.5% (a) and 0.5%/5.0% b); plots of number of
1624 products giving 99% of the total carbon (c); and ratio of numbers of products giving
1625 99.99% of the carbon to numbers giving 99% as a function of carbon number.

1626

1627 the cutoff criterion of 0.5%, Figure 4b shows the ratio of the numbers of products generated using the
1628 0.5% criterion against the much looser criterion of 5%, Figure 4c shows the numbers of products that sum
1629 up to 95% of the reacting carbon (derived from the results with the 0.5% cutoff), and Figure 4d shows the
1630 ratio of this number of products to the total number generated using the 0.5% criterion. As expected, the
1631 numbers of products increase almost exponentially with the size of the molecules, ranging from less than
1632 10 for propane to many hundreds for C₁₄ compounds, with the increase in product numbers with carbon
1633 number being somewhat greater if the 95% carbon criterion is used. Changing the cutoff criterion from
1634 0.5% to 5% affects the numbers of predicted products by about a factor of 2 for the larger alkanes, but by
1635 factors of 5 or more for the larger alkenes, apparently increasing with the size of the molecule. In any
1636 case, this indicates that the numbers of very low yield products are usually much greater than the numbers
1637 of products that make significant contributions to the reacting carbon.


1638 It should be emphasized that these results are not strictly comparable to published descriptions of
1639 mechanisms derived by GECKO-A (e.g., Aumont et al., 2005), which is used to derive multi-generation
1640 mechanisms where all organic products are reacted until they degrade entirely to CO or CO₂ or form non-
1641 volatile products. For example, Figure 4 of Aumont et al. (2005) indicates that a multi-generation
1642 mechanism for n-octane alone contains over 10⁶ species, even after a 5% single step cutoff is employed.
1643 Many of these 10⁶ species are radical intermediates that are not counted as products in this work, but even



1644 if these are eliminated (estimated to reduce the number by about a factor of 2-3), there are many orders of
 1645 magnitude more than the ~100 stable products predicted in the single generation mechanism for n-octane
 1646 in this work (Table A-36) using a single step cutoff parameter that is an order of magnitude lower (0.5%).
 1647 MechGen is also capable of deriving multi-generation mechanisms that can be more directly compared
 1648 with those derived using GECKO-A, but further discussion of this topic is beyond the scope of the
 1649 present paper.

1650 Figure 5 illustrates the variability in the types of products formed from representative C₈
 1651 compounds and terpenes, showing total yields of compounds containing the types of groups formed in the
 1652 highest yields. Many or most of the products contain more than one such group on the molecule. The
 1653 category labeled "multiple groups" gives the total yields of compounds with more than one non-alkyl
 1654 group bonded to a carbon, which are types of compounds whose mechanisms and estimated rate constants
 1655 are particularly uncertain. The yields of such multifunctional compounds are the highest for some of the
 1656 terpenes, but are non-negligible for most types of compounds except perhaps phenols. Different symbols
 1657 are used for products derived using the cutoff parameters of 0.5% and 5%; it can be seen that the total
 1658 yields of various types of products are not strongly affected by this parameter.

1659
 1660



1661
 1662
 1663
 1664
 1665
 1666
 1667
 1668
 1669
 1670


Notes: [a] Minimum yield parameter used when generating the mechanisms; [b] Sum of carbons in products with the indicated groups; [c] Sum of carbons in multifunctional products with two or more different non-alkyl groups on the same carbon; [d] Sum of carbons in products x the fraction of the product in equilibrium with the condensed phase for atmospheric organic aerosol levels of 50 μg/m³.

Figure 5. Comparisons of total mole carbon yields of various types of products for generated mechanisms for reactions of representative C₈ compounds and terpenes.

1671 MechGen has the capability to estimate vapor pressures of generated compounds, using either the
 1672 SIMPOL (Pankow and Asher, 2008) or EVAPORATION (Compernelle et al., 2011) group contribution



1673 methods. The SIMPOL method was used here for illustrative purposes because it provides estimates for a
1674 wider range of compounds. Estimated vapor pressures were used to estimate the equilibrium fractions of
1675 the first-generation products in the condensed phase at an assumed total organic aerosol level of 50
1676 $\mu\text{g}/\text{m}^3$. The estimated total fractions of the products in the condensed phase (sum of product yields \times the
1677 fraction of each in the condensed phase) are given in Table A-36 in the SI and included in Figure 5.
1678 Figure 6 shows the fraction of products in the condensed phase as a function of carbon number for all 38
1679 compounds. As expected, the yields of condensed-phase products increase with carbon number, with the
1680 yields becoming significant for most C_{12+} compounds, though highly variable with type of compound.
1681 These volatility estimates are based on uncertain vapor pressure estimates in many cases (Bilde et al.,
1682 2015), and, most importantly, ignore the contributions from products formed in subsequent generation
1683 reactions, which tend to form lower volatility products than first generation reactions (e.g., see Aumont et
1684 al., 2012, 2013). A comprehensive investigation of the use of MechGen to predict low volatility products
1685 is beyond the scope of this paper, but this is a potentially useful area of future study and comparison with
1686 GECKO-A predictions.



1687
1688 Figure 6. Estimated fraction of products in the condensed phase at 298 K and with 50 $\mu\text{g}/\text{m}^3$ of
1689 background organic aerosol as a function of carbon number for representative
1690 compounds.

1691

1692 9. Discussion and Conclusions

1693 The MechGen system provides a means to derive nearly fully explicit mechanisms for the
1694 reactions of most organic compounds under lower atmospheric conditions. These can be used as a basis
1695 for deriving more condensed mechanisms for use in urban, regional, and global models, and serve as
1696 standards for developing and evaluating mechanism reduction approaches. The level of chemical detail
1697 that generated mechanisms potentially provide may not be necessary for modeling urban O_3 formation,
1698 but may well be necessary for developing or evaluating chemically-based models for formation of SOA in
1699 the atmosphere. MechGen, like GECKO-A (Aumont et al., 2005) also provides a potentially useful
1700 alternative to the widely-used MCM (MCM, 2023) for evaluating chemical systems for laboratory and
1701 other studies. Like MCM, it is available to the general user online; but unlike MCM, it can be used to
1702 derive mechanisms for compounds whose reactions have not been previously estimated.

1703 Explicit chemical detail in itself does not guarantee chemical accuracy or predictive capability.
1704 Detailed mechanisms that incorporate incorrect rate constants and branching ratios, incorporate reactions
1705 that do not occur, omit important reactions, or incorporate inappropriate assumptions, cannot be expected



1706 to give reliable predictions and may provide an illusion of accuracy and understanding that does not exist.
 1707 On the other hand, if the mechanism generation process appropriately incorporates our current knowledge
 1708 and estimates based on best available expert judgment, then the generated mechanism provides our best
 1709 estimate for the reactions of the compound(s) of interest, and, if our knowledge is sufficient, has a
 1710 reasonable chance of predictive capability. However, our current knowledge of atmospheric reactions of
 1711 organics is not complete, so our best judgment as to the most likely detailed mechanism will evolve over
 1712 time, and any mechanism generation system will need to evolve as well so its predictions continue to
 1713 represent the state of the science.

1714 9.1. Summary of Areas of Mechanism Uncertainty

1715 Most of the types of chemical reactions discussed in this paper have at least some areas of significant
 1716 uncertainty or areas where simplifying assumptions have to be made to avoid excessive complexity that is
 1717 either impractical to currently implement, or that is not justified by available knowledge. The impacts of
 1718 these uncertainties are expected to vary widely depending on the compounds whose mechanisms are
 1719 being generated, the environment where the mechanism will be applied, and the objectives of the model
 1720 application. Table 16 lists what are considered to be the major areas of uncertainty in the current
 1721 mechanism generation system and gives very brief indications of the associated problems and impacts.
 1722 Sections of this paper and SI where the issues are discussed in more detail are also given. This can serve
 1723 both as a guideline for areas of research that could be most beneficial for improving detailed mechanisms
 1724 and also areas of MechGen that need to be the focus of future updates. These are certainly not the only
 1725 areas of uncertainty in developing comprehensive mechanisms but appear based on this work to be
 1726 priority areas of research. However, the greatest uncertainty may be reactions or reaction mechanisms that
 1727 are not currently considered and are therefore omitted entirely in atmospheric mechanisms or models.

1728 Table 16. Summary of major areas of uncertainty in the current mechanism generation system
 1729

Uncertainty	Discussion (Section discussed)
Organic + oxidant rate constants	Affects VOC lifetimes and product formation. Not a large uncertainty for the major VOCs or mechanisms for mixtures, but potentially significant for multifunctional products. (Secs. 2.1, A1.2)
Organic photolysis reactions	Affects radical initiation rates as well as VOC lifetimes and fates. Major oversimplifications for larger and multifunctional compounds. Priority area for system update. Photolytic radical from aromatic ring fragmentation products are uncertain. (Secs. 2.2, A1.3)
Unimolecular reactions of non-radical compounds	Affects product formation. MechGen includes one type of reaction that may actually be heterogeneous or H ₂ O catalyzed but ignores others that may be equally important in real environments. (Sec. 2.3)
O ₂ additions to carbon-centered allylic radicals	Affects product formation and predictions of effects of phenolic compounds on radical levels. Insufficient data are available to reliably estimate branching ratios for additions to the different allylic radical centers. (Sec. 3.1)
Cyclization of carbon-centered radicals forming cyclic esters	Affects product formation. Experimental data needed. Existing theoretical studies are inconsistent. Reaction may not be as fast as assumed. (Secs. 3.2.3, A1.4.3)
Other unimolecular reactions of carbon-centered radicals	Affects product formation. Current assumptions are considered reasonable but some need experimental or theoretical verification. (Secs. 3.2, 3.3, A1.4)



Table 16 (continued)

<u>Uncertainty</u>	<u>Discussion</u> (Section discussed)
Aromatic ring opening reactions	Affects product formation and radical initiation. Current estimates likely oversimplifications and not consistent with all product studies. Arbitrary assumptions had to be used in some cases. (Sec. 3.4)
Nitrate yields from NO + substituted peroxy radicals	Affects radical and NO _x sink and recycling for almost all compounds. Experimental data for OH-substituted radicals are inconsistent and there are no direct data for effects of other substituents. (Secs. 4.1, A1.5.2)
H-shift isomerization reactions of peroxy radicals	Affects product formation, particularly hydroperoxides and low volatility compounds. Estimates are uncertain for multifunctional radicals and are especially uncertain for radicals formed from reactions of cyclic compounds such as terpenes. (Secs. 4.6, A1.5.5)
Ring closure reactions of unsaturated peroxy radicals	Affects product formation. Current estimates are based on analysis of uncertain aromatic ring-opening reactions. Estimates need experimental or theoretical verification. (Sec. 4.7)
β-scission decompositions of alkoxy radicals	Affects amounts of fragmentation to lower molecular weight products. Data insufficient to derive reliable estimates for decompositions forming alkoxy radicals or effects of some substituents. Current estimates do not take into account chemical activation effects. (Secs. 5.2, A1.7.2)
Other alkoxy radical reactions	Affects product formation. Uncertainty is variable, but even small estimation errors can affect product predictions. Chemical activation effects uncertain. (Secs. 5, A1.7.3, A1.7.4)
Reactions of Criegee intermediates	Affects radical initiation and product formation from alkenes. Stabilization and reactions of larger and substituted intermediates are particularly uncertain. Simplifications and arbitrary assumptions had to be used. (Secs. 7, A1.9)
Thermochemical estimates	Affects assessments of what reactions need to be considered and some estimates of radical reactions. Certain thermochemical group values used are extremely uncertain and possibly inappropriate. A complete review and update of the thermochemical estimates for mechanism development is required. (Sec. A2).
Appropriate cutoff rules for mechanism generation efficiency	Affects the size of the mechanisms that are generated and the number of products that have to be reacted in multi-generation mechanisms. Too high a cutoff will affect mechanism accuracy, and too low will affect mechanism efficiency and simulation times. (Secs 8; Carter, 2023b)

1730

1731

9.2. Recommendations

1732

1733

1734

1735

1736

1737

1738

1739

1740

1741

Mechanism generation systems such as MechGen provide an important link between basic kinetic and mechanistic data and theories, and applications such as air quality modeling for research and regulatory applications and analysis of laboratory and field data. With regard to regulatory modeling, Kaduwela et al. (2015) proposed an approach for developing atmospheric chemical mechanisms for the future, in which the development and applications of detailed chemical mechanism generation systems played an essential role. Although generated mechanisms are much too large for full three-dimensional atmospheric modeling, they can serve as a basis for deriving more condensed mechanisms for modeling application that still have a direct link to the underlying basic data and theories, and where the effects of the simplifications and condensations can be systematically characterized. Without these links, research and regulatory modeling may not benefit from the emerging science in atmospheric chemistry.



1742 Collection and evaluation of relevant laboratory and theoretical data are essential to detailed
1743 mechanism development, both for direct incorporation into the mechanisms and to support the
1744 development of the many types of SARs needed for comprehensive mechanism development. The
1745 ongoing IUPAC (2023) and NASA (Sander et al., 2006, 2009) evaluations provide an essential role in
1746 this effort, as have the books by Calvert et al. (2000, 2002, 2008, 2011, 2015) on atmospheric chemistry,
1747 and efforts like these need to continue. Recently McGillen et al. (2020) made available a comprehensive
1748 collection of rate constants for reactions of organics with OH, O₃, NO₃, and Cl atoms, and this data
1749 collection effort is continuing, with the latest update being available at [https://data.eurochamp.org/data-](https://data.eurochamp.org/data-access/kin/)
1750 [access/kin/](https://data.eurochamp.org/data-access/kin/). Similar collections are needed for other types of reactions, particularly data to support SAR
1751 development for the many types of radical reactions. The NIST kinetics database (NIST Chemical
1752 Kinetics Database, 2023) is a useful resource, but recommendations are not provided and it is limited to
1753 reactions of simpler molecules and radicals. Quantum theory has become an increasingly important tool
1754 in developing SARs for radical reactions, since experimental data are insufficient for this purpose.
1755 Compilations and evaluations of available theoretical data are needed to take full advantage of this
1756 ongoing work.

1757 Vereecken et al. (2018) reviewed the current status and research needs for development of SARs
1758 for detailed chemical mechanisms. In addition to discussing data needs and the importance of data
1759 collection, they also discussed the need to evaluate and improve existing SARs, particularly with regard
1760 to multifunctional compounds. Many of the SARs used in this work may not incorporate all the relevant
1761 data and need to be updated or possibly replaced by newer versions. Work on SAR development and
1762 improvement needs to be ongoing for chemical mechanisms to continue to represent the state of the
1763 science and retain their link to laboratory data and theories.

1764 The predictive capabilities of atmospheric chemical mechanisms, whether detailed or condensed,
1765 need to be evaluated for comparing their predictions against laboratory data. In some cases, MechGen had
1766 to be modified so that developed mechanisms gave predictions of NO oxidation and O₃ formation
1767 consistent with results of environmental chamber experiments. Consistency with O₃ formation in well-
1768 characterized environmental chamber data needs to be a priority for mechanisms used for regulatory
1769 modeling, though this is not a sufficient evaluation for detailed mechanisms. Comparison of experimental
1770 vs. predicted product yields is necessary for a comprehensive evaluation, but this is a major effort with
1771 many components and experimental difficulties. This would include comparisons with ambient data as
1772 well as results of laboratory and environmental chamber studies. Doing this comprehensively is a multi-
1773 year effort but it needs to be carried out.

1774 Many if not most of the areas of uncertainty in generated mechanisms may not be reduced
1775 significantly in the near term. These uncertainties have variable effects on model predictions of interest,
1776 with some highly uncertain or questionable estimates having almost no effects on predictions, and in
1777 some cases only moderate uncertainties in rate constants having large effects. A systematic study of the
1778 effects of the various types of uncertainties on predictions of interest is needed not only to prioritize areas
1779 where basic research is needed, but also to prioritize which SARs and estimates used in systems like
1780 MechGen should be given the highest priorities for future work.

1781 The research needs do not end once a detailed mechanism is developed and its predictive
1782 capabilities have been evaluated. Mechanisms output by systems such as MechGen or GECKO-A are far
1783 too large and complex for most modeling applications and need to be reduced to be useful. The reduction
1784 method will depend on the intended application, with, for example, detailed product predictions being a
1785 relatively low priority in O₃ models but a high priority in models for SOA and toxics. Further discussion
1786 of reduction issues is beyond the scope of this work, though it should be noted that MechGen as
1787 documented in this paper has been used to develop updated versions of the SAPRC mechanisms for use in



1788 airshed models for predicting O₃ and toxics, designated SAPRC-22 (Carter, 2023a). It can also be used as
1789 a basis for developing future versions of the mechanism for predicting SOA.

1790 **Author Contributions**

1791 This manuscript largely represents the body of work on chemical mechanism development led by
1792 W. P. L. Carter, and he is the main contributor to this publication. J. J. Orlando provided consultation on
1793 the methods, including rate constant and branching ratio estimates, and the resulting mechanism
1794 predictions. J. J. Orlando and K. C. Barsanti contributed to writing and editing, and J. Jiang contributed to
1795 editing and accuracy checking of the IUPAC references.

1796 **Competing Interests**

1797 Authors J. J. Orlando and K. C. Barsanti are members of the editorial board of Atmospheric
1798 Chemistry and Physics.

1800 **Acknowledgements**

1801 This work was supported in part by the California Air Resources Board primarily through
1802 contract no 11-761 and in part by the University of California Retirement system. W. P. L. Carter wishes
1803 to thank Dr. Ajith Kaduwela, the CARB project officer, for his support and helpful discussions.

1804 This work has also been supported by grant from the U.S. Environmental Protection Agency's
1805 Science to Achieve Results (STAR) program. This publication was developed in part under Assistance
1806 Agreement No. 84000701 awarded by the U.S. EPA. It has not been formally reviewed by EPA. EPA
1807 does not endorse any products or commercial services mentioned in this publication.

1808 The opinions and conclusions in this report are entirely those of the authors.

1809



- 1810 References
1811 Afreh, I. K., Aumont, B., Camredon, M., and Barsanti, K. C.: Using GECKO-A to derive mechanistic
1812 understanding of secondary organic aerosol formation from the ubiquitous but understudied camphene,
1813 Atmospheric Chemistry and Physics, 21, 11467–11487, <https://doi.org/10.5194/acp-21-11467-2021>,
1814 2021.
- 1815 Atkinson, R.: Rate constants for the atmospheric reactions of alkoxy radicals: An updated estimation
1816 method, Atmospheric Environment, 41, 8468–8485, <https://doi.org/10.1016/j.atmosenv.2007.07.002>,
1817 2007.
- 1818 Atkinson, R., Carter, W. P. L., and Winer, A. M.: Effects of temperature and pressure on alkyl nitrate
1819 yields in the nitrogen oxide (NO_x) photooxidations of n-pentane and n-heptane, J. Phys. Chem., 87,
1820 2012–2018, <https://doi.org/10.1021/j100234a034>, 1983.
- 1821 Aumont, B., Szopa, S., and Madronich, S.: Modelling the evolution of organic carbon during its gas-
1822 phase tropospheric oxidation: development of an explicit model based on a self generating approach,
1823 Atmospheric Chemistry and Physics, 5, 2497–2517, <https://doi.org/10.5194/acp-5-2497-2005>, 2005.
- 1824 Aumont, B., Valorso, R., Mouchel-Vallon, C., Camredon, M., Lee-Taylor, J., and Madronich, S.:
1825 Modeling SOA formation from the oxidation of intermediate volatility *n*-alkanes, Atmospheric Chemistry
1826 and Physics, 12, 7577–7589, <https://doi.org/10.5194/acp-12-7577-2012>, 2012.
- 1827 Aumont, B., Camredon, M., Mouchel-Vallon, C., La, S., Ouzebidour, F., Valorso, R., Lee-Taylor, J., and
1828 Madronich, S.: Modeling the influence of alkane molecular structure on secondary organic aerosol
1829 formation, Faraday Discuss., 165, 105–122, <https://doi.org/10.1039/C3FD00029J>, 2013.
- 1830 Batiha, M., Al-Muhtaseb, A. H., and Altarawneh, M.: Theoretical study on the reaction of the phenoxy
1831 radical with O₂, OH, and NO₂, International Journal of Quantum Chemistry, 112, 848–857,
1832 <https://doi.org/10.1002/qua.23074>, 2012.
- 1833 Berndt, T. and Böge, O.: Gas-phase reaction of OH radicals with phenol, Phys. Chem. Chem. Phys., 5,
1834 342–350, <https://doi.org/10.1039/B208187C>, 2003.
- 1835 Bilde, M., Barsanti, K., Booth, M., Cappa, C. D., Donahue, N. M., Emanuelsson, E. U., McFiggans, G.,
1836 Krieger, U. K., Marcolli, C., Topping, D., Ziemann, P., Barley, M., Clegg, S., Dennis-Smith, B.,
1837 Hallquist, M., Hallquist, A. M., Khlystov, A., Kulmala, M., Mogensen, D., Percival, C. J., Pope, F., Reid,
1838 J. P., da Silva, M. A. V. R., Rosenoern, T., Salo, K., Soonsin, V. P., Yli-Juuti, T., Prisle, N. L., Pagels, J.,
1839 Rarey, J., Zardini, A. A., and Riipinen, I.: Saturation Vapor Pressures and Transition Enthalpies of Low-
1840 Volatility Organic Molecules of Atmospheric Relevance: From Dicarboxylic Acids to Complex Mixtures,
1841 Chemical Reviews, 115, 4115–4156, <https://doi.org/10.1021/cr5005502>, 2015.
- 1842 Bloss, C., Wagner, V., Jenkin, M. E., Volkamer, R., Bloss, W. J., Lee, J. D., Heard, D. E., Wirtz, K.,
1843 Martin-Reviejo, M., Rea, G., Wenger, J. C., and Pilling, M. J.: Development of a detailed chemical
1844 mechanism (MCMv3.1) for the atmospheric oxidation of aromatic hydrocarbons, Atmospheric Chemistry
1845 and Physics, 5, 641–664, <https://doi.org/10.5194/acp-5-641-2005>, 2005.
- 1846 Calvert, J., Mellouki, A., Orlando, J., Pilling, M., and Wallington, and T.: Mechanisms of Atmospheric
1847 Oxidation of the Oxygenates, Oxford University Press, Oxford, New York, 1634 pp., 2011.



- 1848 Calvert, J. G., Atkinson, R., Kerr, J. A., Madronich, S., Moortgat, G. K., Wallington, T. J., and Yarwood,
1849 and G.: The Mechanisms of Atmospheric Oxidation of the Alkenes, Oxford University Press, Oxford,
1850 New York, 560 pp., 2000.
- 1851 Calvert, J. G., Atkinson, R., Becker, K. H., Kamens, R. M., Seinfeld, J. H., Wallington, T. H., and
1852 Yarwood, and G.: The Mechanisms of Atmospheric Oxidation of the Aromatic Hydrocarbons, Oxford
1853 University Press, Oxford, New York, 566 pp., 2002.
- 1854 Calvert, J. G., Derwent, R. G., Orlando, J. J., Tyndall, G. S., and Wallington, and T. J.: Mechanisms of
1855 Atmospheric Oxidation of the Alkanes, Oxford University Press, Oxford, New York, 1008 pp., 2008.
- 1856 Calvert, J. G., Orlando, J. J., Stockwell, W. R., and Wallington, and T. J.: The Mechanisms of Reactions
1857 Influencing Atmospheric Ozone, Oxford University Press, Oxford, New York, 608 pp., 2015.
- 1858 Camredon, M., Aumont, B., Lee-Taylor, J., and Madronich, S.: The SOA/VOC/NO_x system: an explicit
1859 model of secondary organic aerosol formation, *Atmospheric Chemistry and Physics*, 7, 5599–5610,
1860 <https://doi.org/10.5194/acp-7-5599-2007>, 2007.
- 1861 Carter, W. P. L.: Development of Ozone Reactivity Scales for Volatile Organic Compounds, *Journal of*
1862 *Air and Waste Management*, 44, 881–899, <https://doi.org/10.1080/1073161X.1994.10467290>, 1994.
- 1863 Carter, W. P. L.: Documentation of the SAPRC-99 Chemical Mechanism for VOC Reactivity
1864 Assessment, 2000. Available at <http://www.cert.ucr.edu/~carter/absts.htm#saprc99>.
- 1865 Carter, W. P. L.: Development of the SAPRC-07 Chemical Mechanism, *Atmospheric Environment*, 44,
1866 5324–5335, <https://doi.org/10.1016/j.atmosenv.2010.01.026>, 2010a.
- 1867 Carter, W. P. L.: Development of the SAPRC-07 Chemical Mechanism and Updated Ozone Reactivity
1868 Scales, 2010b. Available at <http://www.cert.ucr.edu/~carter/absts.htm#aminrep>.
- 1869 Carter, W. P. L.: Preliminary Documentation of the SAPRC-16 Mechanism, 2016. Available at:
1870 <https://intra.cert.ucr.edu/~carter/SAPRC/16/>.
- 1871 Carter, W. P. L.: Documentation of the SAPRC-16 Mechanism Generation System, 2019. Available at
1872 <https://intra.engr.ucr.edu/~carter/SAPRC/18/MechGen.pdf>
- 1873 Carter, W. P. L.: Documentation of the SAPRC-18 Mechanism, 2020. Available at
1874 <https://intra.cert.ucr.edu/~carter/SAPRC/18/S18doc.pdf>.
- 1875 Carter, W. P. L.: Estimation of Rate Constants for Reactions of Organic Compounds under Atmospheric
1876 Conditions, *Atmosphere*, 12, 1250, <https://doi.org/10.3390/atmos12101250>, 2021.
- 1877 Carter, W. P. L.: Documentation of the SAPRC-22 Mechanisms, 2023a. Available at
1878 <https://intra.engr.ucr.edu/~carter/SAPRC/22/>
- 1879 Carter, W. P. L.: SAPRC Chemical Mechanisms, Test Simulations, and Environmental Chamber
1880 Simulation Files, 2023b. <https://intra.engr.ucr.edu/~carter/SAPRC/SAPRCfiles.htm>. Updated September
1881 9.
- 1882 Carter, W. P. L.: The SAPRC Mechanism Generation System, in preparation.



- 1883 Carter, W. P. L. and Heo, G.: Development of revised SAPRC aromatics mechanisms, *Atmospheric*
1884 *Environment*, 77, 404–414, <https://doi.org/10.1016/j.atmosenv.2013.05.021>, 2013.
- 1885 Carter, W. P. L., Atkinson, R., Winer, A. M., and Pitts Jr., J. N.: Experimental investigation of chamber-
1886 dependent radical sources, *International Journal of Chemical Kinetics*, 14, 1071–1103,
1887 <https://doi.org/10.1002/kin.550141003>, 1982.
- 1888 Chen, Y. and Zhu, L.: The Wavelength Dependence of the Photodissociation of Propionaldehyde in the
1889 280–330 nm Region, *J. Phys. Chem. A*, 105, 9689–9696, <https://doi.org/10.1021/jp011445s>, 2001.
- 1890 Compernelle, S., Ceulemans, K., and Müller, J.-F.: EVAPORATION: a new vapour pressure estimation
1891 method for organic molecules including non-additivity and intramolecular interactions, *Atmospheric*
1892 *Chemistry and Physics*, 11, 9431–9450, <https://doi.org/10.5194/acp-11-9431-2011>, 2011.
- 1893 Crouse, J. D., Knap, H. C., Ørnsø, K. B., Jørgensen, S., Paulot, F., Kjaergaard, H. G., and Wennberg, P.
1894 O.: Atmospheric Fate of Methacrolein. 1. Peroxy Radical Isomerization Following Addition of OH and
1895 O₂, *J. Phys. Chem. A*, 116, 5756–5762, <https://doi.org/10.1021/jp211560u>, 2012.
- 1896 Curran, H. J., Gaffuri, P., Pitz, W. J., and Westbrook, C. K.: A Comprehensive Modeling Study of n-
1897 Heptane Oxidation, *Combustion and Flame*, 114, 149–177, <https://doi.org/10.1016/S0010->
1898 [2180\(97\)00282-4](https://doi.org/10.1016/S0010-2180(97)00282-4), 1998.
- 1899 Davis, A. C. and Francisco, J. S.: Ab Initio Study of Hydrogen Migration in 1-Alkylperoxy Radicals, *J.*
1900 *Phys. Chem. A*, 114, 11492–11505, <https://doi.org/10.1021/jp1042393>, 2010.
- 1901 Gardner, E. P., Sperry, P. D., and Calvert, J. G.: Photodecomposition of acrolein in oxygen-nitrogen
1902 mixtures, *J. Phys. Chem.*, 91, 1922–1930, <https://doi.org/10.1021/j100291a048>, 1987.
- 1903 International Union of Pure and Applied Chemistry (IUPAC): <https://iupac.aeris-data.fr/en/home-english/>,
1904 last access: 26 September 2023.
- 1905 Jenkin, M. E., Saunders, S. M., and Pilling, M. J.: The tropospheric degradation of volatile organic
1906 compounds: a protocol for mechanism development, *Atmospheric Environment*, 31, 81–104,
1907 [https://doi.org/10.1016/S1352-2310\(96\)00105-7](https://doi.org/10.1016/S1352-2310(96)00105-7), 1997.
- 1908 Jenkin, M. E., Saunders, S. M., Wagner, V., and Pilling, M. J.: Protocol for the development of the
1909 Master Chemical Mechanism, MCM v3 (Part B): tropospheric degradation of aromatic volatile organic
1910 compounds, *Atmospheric Chemistry and Physics*, 3, 181–193, <https://doi.org/10.5194/acp-3-181-2003>,
1911 2003.
- 1912 Jenkin, M. E., Valorso, R., Aumont, B., Rickard, A. R., and Wallington, T. J.: Estimation of rate
1913 coefficients and branching ratios for gas-phase reactions of OH with aliphatic organic compounds for use
1914 in automated mechanism construction, *Atmospheric Chemistry and Physics*, 18, 9297–9328,
1915 <https://doi.org/10.5194/acp-18-9297-2018>, 2018a.
- 1916 Jenkin, M. E., Valorso, R., Aumont, B., Rickard, A. R., and Wallington, T. J.: Estimation of rate
1917 coefficients and branching ratios for gas-phase reactions of OH with aromatic organic compounds for use
1918 in automated mechanism construction, *Atmospheric Chemistry and Physics*, 18, 9329–9349,
1919 <https://doi.org/10.5194/acp-18-9329-2018>, 2018b.



- 1920 Jenkin, M. E., Valorso, R., Aumont, B., and Rickard, A. R.: Estimation of rate coefficients and branching
1921 ratios for reactions of organic peroxy radicals for use in automated mechanism construction, *Atmospheric*
1922 *Chemistry and Physics*, 19, 7691–7717, <https://doi.org/10.5194/acp-19-7691-2019>, 2019.
- 1923 Jenkin, M. E., Valorso, R., Aumont, B., Newland, M. J., and Rickard, A. R.: Estimation of rate
1924 coefficients for the reactions of O₃ with unsaturated organic compounds for use in automated mechanism
1925 construction, *Atmospheric Chemistry and Physics*, 20, 12921–12937, [https://doi.org/10.5194/acp-20-](https://doi.org/10.5194/acp-20-12921-2020)
1926 [12921-2020](https://doi.org/10.5194/acp-20-12921-2020), 2020.
- 1927 Jiang, J., Carter, W. P. L., Cocker, D. R., III, and Barsanti, K. C.: Development and Evaluation of a
1928 Detailed Mechanism for Gas-Phase Atmospheric Reactions of Furans, *ACS Earth and Space Chemistry*,
1929 4, 1254–1268, <https://doi.org/10.1021/acsearthspacechem.0c00058>, 2020.
- 1930 Kaduwela, A., Luecken, D., Carter, W., and Derwent, R.: New directions: Atmospheric chemical
1931 mechanisms for the future, *Atmospheric Environment*, 122, 609–610,
1932 <https://doi.org/10.1016/j.atmosenv.2015.10.031>, 2015.
- 1933 Kerdouci, J., Picquet-Varrault, B., and Doussin, J.-F.: Structure–activity relationship for the gas-phase
1934 reactions of NO₃ radical with organic compounds: Update and extension to aldehydes, *Atmospheric*
1935 *Environment*, 84, 363–372, <https://doi.org/10.1016/j.atmosenv.2013.11.024>, 2014.
- 1936 Lannuque, V., Camredon, M., Couvidat, F., Hodzic, A., Valorso, R., Madronich, S., Bessagnet, B., and
1937 Aumont, B.: Exploration of the influence of environmental conditions on secondary organic aerosol
1938 formation and organic species properties using explicit simulations: development of the VBS-GECKO
1939 parameterization, *Atmospheric Chemistry and Physics*, 18, 13411–13428, [https://doi.org/10.5194/acp-18-](https://doi.org/10.5194/acp-18-13411-2018)
1940 [13411-2018](https://doi.org/10.5194/acp-18-13411-2018), 2018.
- 1941 Lee-Taylor, J., Madronich, S., Aumont, B., Baker, A., Camredon, M., Hodzic, A., Tyndall, G. S., Apel,
1942 E., and Zaveri, R. A.: Explicit modeling of organic chemistry and secondary organic aerosol partitioning
1943 for Mexico City and its outflow plume, *Atmospheric Chemistry and Physics*, 11, 13219–13241,
1944 <https://doi.org/10.5194/acp-11-13219-2011>, 2011.
- 1945 Li, Q., Jiang, J., Afreh, I. K., Barsanti, K. C., and Cocker, D. R., III: Secondary organic aerosol formation
1946 from camphene oxidation: measurements and modeling, *Atmospheric Chemistry and Physics*, 22, 3131–
1947 3147, <https://doi.org/10.5194/acp-22-3131-2022>, 2022.
- 1948 Master Chemical Mechanism (MCM): <http://chmlin9.leeds.ac.uk/MCM/roots.htm>, last access: 26
1949 September 2023.
- 1950 Matsugi, A. and Miyoshi, A.: Yield of Formyl Radical from the Vinyl + O₂ Reaction, *International*
1951 *Journal of Chemical Kinetics*, 46, 260–274, <https://doi.org/10.1002/kin.20823>, 2014.
- 1952 Matsunaga, A. and Ziemann, P. J.: Yields of β -Hydroxynitrates and Dihydroxynitrates in Aerosol Formed
1953 from OH Radical-Initiated Reactions of Linear Alkenes in the Presence of NO_x, *J. Phys. Chem. A*, 113,
1954 599–606, <https://doi.org/10.1021/jp807764d>, 2009.
- 1955 Matsunaga, A. and Ziemann, P. J.: Yields of β -hydroxynitrates, dihydroxynitrates, and trihydroxynitrates
1956 formed from OH radical-initiated reactions of 2-methyl-1-alkenes, *Proceedings of the National Academy*
1957 *of Sciences*, 107, 6664–6669, <https://doi.org/10.1073/pnas.0910585107>, 2010.



- 1958 McGillen, M. R., Carter, W. P. L., Mellouki, A., Orlando, J. J., Picquet-Varrault, B., and Wallington, T.
1959 J.: Database for the kinetics of the gas-phase atmospheric reactions of organic compounds, *Earth System
1960 Science Data*, 12, 1203–1216, <https://doi.org/10.5194/essd-12-1203-2020>, 2020.
- 1961 Miyoshi, A.: Systematic Computational Study on the Unimolecular Reactions of Alkylperoxy (RO₂),
1962 Hydroperoxyalkyl (QOOH), and Hydroperoxyalkylperoxy (O₂QOOH) Radicals, *J. Phys. Chem. A*, 115,
1963 3301–3325, <https://doi.org/10.1021/jp112152n>, 2011.
- 1964 Møller, K. H., Bates, K. H., and Kjaergaard, H. G.: The Importance of Peroxy Radical Hydrogen-Shift
1965 Reactions in Atmospheric Isoprene Oxidation, *J. Phys. Chem. A*, 123, 920–932,
1966 <https://doi.org/10.1021/acs.jpca.8b10432>, 2019.
- 1967 Muthuramu, K., Shepson, P. B., and O'Brien, J. M.: Preparation, analysis, and atmospheric production of
1968 multifunctional organic nitrates, *Environ. Sci. Technol.*, 27, 1117–1124,
1969 <https://doi.org/10.1021/es00043a010>, 1993.
- 1970 National Institute of Standards and Technology (NIST) Chemical Kinetics Database:
1971 <https://kinetics.nist.gov/kinetics/>, last access: 26 September 2023.
- 1972 Nozière, B. and Vereecken, L.: Direct Observation of Aliphatic Peroxy Radical Autoxidation and Water
1973 Effects: An Experimental and Theoretical Study, *Angewandte Chemie International Edition*, 58, 13976–
1974 13982, <https://doi.org/10.1002/anie.201907981>, 2019.
- 1975 O'Brien, J. M., Czuba, E., Hastie, D. R., Francisco, Joseph. S., and Shepson, P. B.: Determination of the
1976 Hydroxy Nitrate Yields from the Reaction of C₂–C₆ Alkenes with OH in the Presence of NO, *J. Phys.
1977 Chem. A*, 102, 8903–8908, <https://doi.org/10.1021/jp982320z>, 1998.
- 1978 Olariu, R. I., Klotz, B., Barnes, I., Becker, K. H., and Mocanu, R.: FT–IR study of the ring-retaining
1979 products from the reaction of OH radicals with phenol, o-, m-, and p-cresol, *Atmospheric Environment*,
1980 36, 3685–3697, [https://doi.org/10.1016/S1352-2310\(02\)00202-9](https://doi.org/10.1016/S1352-2310(02)00202-9), 2002.
- 1981 Orlando, J. J., Tyndall, G. S., and Wallington, T. J.: The Atmospheric Chemistry of Alkoxy Radicals,
1982 *Chem. Rev.*, 103, 4657–4690, <https://doi.org/10.1021/cr020527p>, 2003.
- 1983 Pankow, J. F. and Asher, W. E.: SIMPOL.1: a simple group contribution method for predicting vapor
1984 pressures and enthalpies of vaporization of multifunctional organic compounds, *Atmospheric Chemistry
1985 and Physics*, 8, 2773–2796, <https://doi.org/10.5194/acp-8-2773-2008>, 2008.
- 1986 Platz, J., Nielsen, O. J., Wallington, T. J., Ball, J. C., Hurley, M. D., Straccia, A. M., Schneider, W. F.,
1987 and Sehested, J.: Atmospheric Chemistry of the Phenoxy Radical, C₆H₅O(•): UV Spectrum and Kinetics
1988 of Its Reaction with NO, NO₂, and O₂, *J. Phys. Chem. A*, 102, 7964–7974,
1989 <https://doi.org/10.1021/jp982221i>, 1998.
- 1990 Praske, E., Otkjær, R. V., Crouse, J. D., Hethcox, J. C., Stoltz, B. M., Kjaergaard, H. G., and Wennberg,
1991 P. O.: Atmospheric autoxidation is increasingly important in urban and suburban North America,
1992 *Proceedings of the National Academy of Sciences*, 115, 64–69, <https://doi.org/10.1073/pnas.1715540115>,
1993 2018.
- 1994 Praske, E., Otkjær, R. V., Crouse, J. D., Hethcox, J. C., Stoltz, B. M., Kjaergaard, H. G., and Wennberg,
1995 P. O.: Intramolecular Hydrogen Shift Chemistry of Hydroperoxy-Substituted Peroxy Radicals, *J. Phys.
1996 Chem. A*, 123, 590–600, <https://doi.org/10.1021/acs.jpca.8b09745>, 2019.



- 1997 Sander, S. P., Friedl, R. R., and Ravishankara, A. R.: Chemical Kinetics and Photochemical Data for Use
1998 in Atmospheric Studies Evaluation Number 15, 2006.
- 1999 Sander, S. P., Friedl, R. R., and Barker, J. R.: Supplement to Evaluation 15: Update of Key Reactions,
2000 2009.
- 2001 Shepson, P. B., Edney, E. O., Kleindienst, T. E., Pittman, J. H., and Namie, G. R.: Production of organic
2002 nitrates from hydroxide and nitrate reaction with propylene, *Environ. Sci. Technol.*, 19, 849–854,
2003 <https://doi.org/10.1021/es00139a014>, 1985.
- 2004 Teng, A. P., Crouse, J. D., Lee, L., St. Clair, J. M., Cohen, R. C., and Wennberg, P. O.: Hydroxy nitrate
2005 production in the OH-initiated oxidation of alkenes, *Atmospheric Chemistry and Physics*, 15, 4297–4316,
2006 <https://doi.org/10.5194/acp-15-4297-2015>, 2015.
- 2007 Venecek, M. A., Cai, C., Kaduwela, A., Avise, J., Carter, W. P. L., and Kleeman, M. J.: Analysis of
2008 SAPRC16 chemical mechanism for ambient simulations, *Atmospheric Environment*, 192, 136–150,
2009 <https://doi.org/10.1016/j.atmosenv.2018.08.039>, 2018.
- 2010 Vereecken, L. and Nozière, B.: H migration in peroxy radicals under atmospheric conditions,
2011 *Atmospheric Chemistry and Physics*, 20, 7429–7458, <https://doi.org/10.5194/acp-20-7429-2020>, 2020.
- 2012 Vereecken, L. and Peeters, J.: Decomposition of substituted alkoxy radicals—part I: a generalized
2013 structure–activity relationship for reaction barrier heights, *Phys. Chem. Chem. Phys.*, 11, 9062–9074,
2014 <https://doi.org/10.1039/B909712K>, 2009.
- 2015 Vereecken, L. and Peeters, J.: A structure–activity relationship for the rate coefficient of H-migration in
2016 substituted alkoxy radicals, *Phys. Chem. Chem. Phys.*, 12, 12608–12620,
2017 <https://doi.org/10.1039/C0CP00387E>, 2010.
- 2018 Vereecken, L., Novelli, A., and Taraborrelli, D.: Unimolecular decay strongly limits the atmospheric
2019 impact of Criegee intermediates, *Phys. Chem. Chem. Phys.*, 19, 31599–31612,
2020 <https://doi.org/10.1039/C7CP05541B>, 2017.
- 2021 Vereecken, L., Aumont, B., Barnes, I., Bozzelli, J. w., Goldman, M. j., Green, W. h., Madronich, S.,
2022 McGillen, M. r., Mellouki, A., Orlando, J. j., Picquet-Varrault, B., Rickard, A. r., Stockwell, W. r.,
2023 Wallington, T. j., and Carter, W. p. l.: Perspective on Mechanism Development and Structure-Activity
2024 Relationships for Gas-Phase Atmospheric Chemistry, *International Journal of Chemical Kinetics*, 50,
2025 435–469, <https://doi.org/10.1002/kin.21172>, 2018.
- 2026 Wolfe, G. M., Crouse, J. D., Parrish, J. D., Clair, J. M. S., Beaver, M. R., Paulot, F., Yoon, T. P.,
2027 Wennberg, P. O., and Keutsch, F. N.: Photolysis, OH reactivity and ozone reactivity of a proxy for
2028 isoprene-derived hydroperoxyenals (HPALDs), *Phys. Chem. Chem. Phys.*, 14, 7276–7286,
2029 <https://doi.org/10.1039/C2CP40388A>, 2012.
- 2030 Xiang, B., Zhu, L., and Tang, Y.: Photolysis of 4-Oxo-2-pentenal in the 190–460 nm Region, *J. Phys.*
2031 *Chem. A*, 111, 9025–9033, <https://doi.org/10.1021/jp0739972>, 2007.
- 2032 Xu, C. and Wang, L.: Atmospheric Oxidation Mechanism of Phenol Initiated by OH Radical, *J. Phys.*
2033 *Chem. A*, 117, 2358–2364, <https://doi.org/10.1021/jp308856b>, 2013.



- 2034 Yuan, Y., Zhao, X., Wang, S., and Wang, L.: Atmospheric Oxidation of Furan and Methyl-Substituted
2035 Furans Initiated by Hydroxyl Radicals, *J. Phys. Chem. A*, 121, 9306–9319,
2036 <https://doi.org/10.1021/acs.jpca.7b09741>, 2017.
- 2037 Zhang, J., Dransfield, T., and Donahue, N. M.: On the Mechanism for Nitrate Formation via the Peroxy
2038 Radical + NO Reaction, *J. Phys. Chem. A*, 108, 9082–9095, <https://doi.org/10.1021/jp048096x>, 2004.