



Merged Observatory Data Files (MODFs): An Integrated Observational Data Product Supporting Process-Oriented Investigations and Diagnostics

5 Taneil Uttal^{1,a,*}, Leslie M. Hartten^{2,1,a,*}, Siri Jodha Khalsa^{2,3}, Barbara Casati⁴, Gunilla Svensson⁵,
Jonathan Day⁶, Jareth Holt⁵, Elena Akish^{1,2,b}, Sara Morris^{1,7}, Ewan O'Connor⁸, Roberta Pirazzini⁸,
Laura X. Huang⁹, Robert Crawford⁹, Zen Mariani⁹, Øystein Godøy¹⁰, Johanna A.K. Tjernström^{11,10}, Giri
Prakash¹², Nicki Hickmon¹³, Marion Maturilli¹⁴ and Christopher J. Cox¹

¹Physical Science Laboratory, National Oceanic and Atmospheric Administration, Boulder, CO 80305-3328, USA

²Cooperative Institute for Research in the Environmental Sciences, Boulder, CO 80309, USA

10 ³National Snow Ice and Data Center, Boulder, CO 80309-0449, USA

⁴Meteorological Research Division, Environment and Climate Change Canada, Dorval, QC H9P-1J3, Canada

⁵Department of Meteorology and Bolin Centre for Climate Change, Stockholm University, 106 91 Stockholm, Sweden

⁶European Center for Medium-Range Weather Forecasts, Reading, RG2 9AX, United Kingdom

⁷Global Monitoring Laboratory, National Oceanic and Atmospheric Administration, Boulder, CO 80305-3337, USA

15 ⁸Finnish Meteorological Institute, Helsinki, FI-00101, Finland

⁹Meteorological Research Division, Environment and Climate Change Canada, Toronto, ON M3H-5T4, Canada

¹⁰Norwegian Meteorological Institute, Oslo, N-0313, Norway

¹¹Swedish Meteorological and Hydrological Institute, Norrköping, SE-60176, Sweden

¹²Department of Energy, Oak Ridge National Laboratory, Oak Ridge, TN 37830, USA

20 ¹³Department of Energy, Argonne National Laboratory, Lemont, IL 60439, USA

¹⁴Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Potsdam, D-14473, Germany

^a retired

^b currently at: Broomfield, CO 80020, USA


Correspondence to: Taneil Uttal (taneil.uttal@gmail.com), Leslie M. Hartten (Leslie.Hartten@colorado.edu)




Abstract. A large and ever-growing body of geophysical information is measured on campaigns and at specialized observatories as a part of scientific expeditions/experiments. These collections of observed data include many essential climate variables (as defined by the World Meteorological Organization), but are often distinguished by a wide range of additional non-routine measurements that are designed to not only document the state of the environment, but also the drivers that contribute to that state. These field data are not only used to further understand the environmental processes through observation-based studies, but also to provide baseline data to test model performance and to codify understanding to improve predictive capabilities. To address the considerable barriers and difficulty in utilizing these diverse and complex data for observation-model research, the Merged Observatory Data File (MODF) concept has been developed. The MODF combines measurements from multiple instruments into a single file that complies with well-established data format and metadata practices and has been designed to parallel development of corresponding Merged Model Data Files (MMDFs). Using MODF and MMDF protocols will facilitate the evolution of Model Intercomparison Projects into Model Intercomparison and Improvement Projects by putting observation and model data ‘on the same page’ in a timely manner. The MODF concept was developed especially for weather forecast model studies in the Arctic. The surprisingly complex process of implementing MODFs in that context refined the concept itself. Thus this article explains the concept of MODFs by providing details on the issues that were revealed and resolved during this first specific implementation. Detailed instructions are provided on how to make MODFs, and this article can thus be considered a MODF creation manual.

MDFs (MODFs + MMDFs) = MIPs → MIIPs

Merged Observatory
Data Files
(MODFs)




Merged Data Files (MDFs)



Model Intercomparison
Projects (MIPs)


Model Intercomparison and
Improvement Projects (MIIPs)

Model 1




Merged Model
Data Files
(MMDFs)

Model ...



Model n





1 Introduction

45 The Merged Observatory Data File (MODF) concept is based on the simple principle of combining measurements made by multiple, co-located instruments from research observatories and campaigns into a single file that complies with already established data stewardship standards. ‘Observatory’ here refers to a facility that measures an extensive inventory of collocated geophysical variables that have been chosen with the intention of investigating specific, usually interrelated, physical processes in order to answer hypothesis-driven science questions. In this context, an observatory could be a land site
50 or a research vessel. While it is standard scientific operating procedure to co-locate research grade instruments both continuously at observatories and episodically for field campaigns (often side-by-side with routine operational station instruments with long operational histories) there are generally no standard procedures for coordinated data management such as those that have been developed for operational data. Thus, the data from separate instruments can be scattered between separate files with different authors, formats, metadata, physical archive locations, and use restrictions.

55

The specific MODF realization presented here is for Arctic observatories and field campaigns that resulted from initiatives established during the Year of Polar Prediction (YOPP; Goessling et al., 2016; Jung et al., 2016; Wilson et al., 2023). One key YOPP activity was the YOPP supersite Model Intercomparison Project (YOPPsiteMIP; Day et al., submitted), which was designed to facilitate process-based validation of numerical weather prediction (NWP) models at polar locations during
60 Special Observing Periods (SOPs). The concept of MODFs and their forecasting analogs, Merged Model Data Files (MMDFs), was motivated by the YOPPsiteMIP community’s desire to have the same variables from observations and models in easy-to-use files of the same structure in order to explore small-scale parameterized processes that are not represented well in the forecast models. These MODFs thus provide an integrated observation database to support model process representation through parameterization improvements for weather forecasts in the polar regions. At the same time,
65 they also facilitate comparative observational studies across Arctic sites.

The MODF concept addresses the problem that research-grade, process-level observations are currently underutilized for model evaluation of parameterization deficiencies. As weather forecasting models increase in complexity and include detailed representation of land, ocean, ice, and snow in addition to the atmosphere, it is increasingly important to evaluate
70 processes using observations recorded through the whole earth system column, including the fluxes at their interfaces, to inform model development. This requires multi-variate process-oriented diagnostic methods utilizing data that spans these multiple components to understand model error. An essential component of the MODF concept is that analogous Merged Model Data Files (MMDFs) can be developed with extracted model data near and around the observatory sites. Together, these are defined as Merged Data Files (MDFs). The MDFs, which bring together observations from different earth system
75 components as well as model output in a standard file format, provide the basis for this and will support Model Intercomparison and Improvement Projects (MIIPs) as an evolution from Model Intercomparison Projects (MIPs). The idea



of MIIPs (a new acronym that we define here) does not imply that previous MIPs did not have productive results. However, through the development of matched MODF and MMDF data sets, it is implied that what have often been intercomparison studies can be smoothly extended to model improvements.

80 **2 Background**

Using observations and model outputs symbiotically is an area of ongoing effort and research with recognized challenges (Holtzlag, et al., 2013). Many levels of MIPs have been in progress for decades (Stephens et al., 2023). In addition, existing efforts and methodologies facilitate the usage of increasingly heterogenous data sets in model–observation fusion efforts through assimilation (Gettleman et al., 2022) and for inputs into multivariate artificial intelligence analyses (Boukabara et al., 2021). There are well-organized systems for managing the data from operational surface networks, upper air networks and satellites that are uploaded into the Global Telecommunications System (GTS) that is overseen by the World Meteorological Organization (WMO) (see Global Telecommunications System (GTS), 2023; WMO, 2020). However, GTS data are only readily available directly to national forecasting centers (which presumably have developed institutionally specific reading and ingesting routines) and via products developed by WMO institutional repositories¹ (Bojinski et al., 85 2014; Lavergne et al., 2022). The latter have necessarily gone through various quality control, formatting, averaging and sometimes interpolation to create globally uniform products. As a result of the standardized processing, it is likely that information on high-resolution, rapid and extreme events (Sardeshmukh et al., 2015) may have sometimes been lost. 90

The MODF schema has been specifically developed for managing observatory and campaign research observations as opposed to operational observations. Research observations target local-scale and often rapid or extreme processes that are intended to lead to discovery of the physics within the atmosphere as well as the physics that govern the coupling processes between the atmosphere and the underlying surface. The surface can be land, ocean, ice or any of the three, often with obfuscating layers of plant and/or snow cover that are themselves components of the system and separate objects of study. Research-grade data pose many additional challenges regarding data latency, accessibility and uptake issues, and institutional ownership, compared to data that have been managed more systematically specifically for operational purposes. 95 100

The data science community is aware of these issues and in response has developed FAIR (Findable, Accessible, Interoperable, Reusable) data principles (Wilkinson et al., 2016) that can be applied to individual data sets. However, Wilkinson et al. explicitly stated that “These high-level FAIR Guiding Principles precede implementation choices, and do not suggest any specific technology, standard, or implementation–solution; moreover, the principles are not, themselves, a standard or a specification.” Whereas many FAIR solutions are implemented by web services (e.g., Buck et al., 2019), the MODF concept described here can be considered an alternative: an integrated data product that is based on the same 105

¹ <https://climatedata-catalogue.wmo.int/homepage>



110 metadata conventions that have been developed for web service solutions. The considerations and steps described here for creating MODFs can therefore be considered a particular FAIR implementation—solution specifically for observatory and campaign data.

3 The MODF Concept

115 Figure 1 is a conceptual schematic of the end-to-end process involved with data collection, data quality control (QC) and processing, metadata information collection and data amalgamation into netCDF files (Unidata, 2023) that follow the NetCDF Climate and Forecast (CF) Metadata Conventions (Eaton et al., 2022, hereafter the CF Metadata Conventions). The process for turning model forecast output into MMDFs is qualitatively similar², including the use of a particular set of global and variable attributes, since the specifications were developed by modellers as well as observationalists.

² Differences generally arise from the fact that forecast models can produce tendencies on timestep scales that are typically not available from field instruments, and that model output is typically regularly distributed in time and space rather than coming from discrete instruments which may move irregularly in space, which operate at different optimized cadences, and which are subject to physical or power disruptions.




Figure 1: Instruments, measurements, data processing and metadata generation for (A1) Surface Atmospheric State, (A2) Broadband Surface Radiation, (A3) Surface Turbulent Fluxes, (A4) Upper Atmospheric Profiles, (A5) Terrestrial Sub-Surface and (A6) Ocean and Sea Ice/Snow collected into netCDF files with specified (B) Global Attributes and (C) Variable Attributes that are compliant with FAIR principles.

120

The different types of geophysical data (A1 thru A6) that typically can compose observatory and campaign research data are described below.



125 3.1 Surface Atmospheric Measurements

Surface atmospheric measurements at observatories or during campaigns are made with research-grade thermometers, hygrometers and anemometers, which often measure at higher frequency and with more carefully calibrated sensors than operational weather stations. Such observations are sometimes side-by-side with operational weather service stations, providing context to multi-decadal operational records. Research meteorological measurements are frequently redundant,
130 deployed at multiple locations across a site (on a scale smaller than an NWP model grid cell) or at different levels on towers to get detailed profiles in the near-surface boundary layer. Resulting variables measured are temperature, pressure, relative humidity, and the eastward and northward components of the windspeed. Both operational and research measurements of these atmospheric state variables can be considered to have well-quantified uncertainties determined by instrument calibration and tolerances.

135

Broadband surface radiation is measured by radiometers that measure incoming and outgoing, shortwave and longwave radiation and associated variables such as surface temperature, sky brightness temperature and derived variables such as albedo. There are multiple commercial and experimental instrument options for measuring broadband radiation, and in the Arctic a particularly wide range of methods must be applied to keep glass domes clear of obstructions (e.g., ice, snow, dust
140 or sea-salt) such as heating, ventilation and manual cleaning (Cox et al., 2021). The Baseline Surface Radiation Network (Ohmura et al., 1998) is a global repository for standardized radiation products that are traceable to the WMO World Radiation Radiometric Reference. However, BSRN is not designed to accommodate short-term campaign data sets and does not account for different methods for data QC and processing (Matsui et al., 2012; Long and Shi, 2006; Long and Shi, 2008).

145 Surface Turbulent Flux variables are calculated and inferred by a number of different methods. Eddy-correlation techniques (Kaimal and Finnigan, 1994) are based on measurements by fast-response sonic anemometers and hygrometers with built-in fast-response temperature sensors. It is often necessary to make considerable site-specific adjustments to processing methods accounting for local surface roughness, sensor height, and obstructed wind-direction sectors. In the polar regions, sensors operate near the thresholds of instrument ratings for detection and environmental conditions and have to be quality-
150 checked for periods of riming. There are specific challenges for the cold snow- and ice-covered surface conditions over land (Grachev et al., 2018) and over the icepack of the central Arctic Ocean (Andreas et al., 2010; Cox et al., 2023). Eddy-correlation methods are only valid under stringent environmental conditions, which results in frequent data gaps. Alternatively, turbulence variables can be calculated by bulk-aerodynamic methods (Monin and Obukhov, 1954; Mahrt and Sun, 1995), but results are not as meaningful for comparing to models as it becomes a model-to-model rather than a model-
155 to-observation comparison. Commercial packages have been developed and compared for calculating latent, sensible and gas fluxes (Mauder et al., 2008; Fratini and Mauder, 2014), but caution must be exercised in using results and it is important to understand the basis on which variables are derived by proprietary commercial or custom software. The resulting flux



160 variables are latent and sensible heat fluxes, friction velocity, surface stress (momentum flux), drag coefficient, kinematic temperature scale, Monin-Obhukov stability parameter and dissipation rate of the turbulent kinetic energy. Given the variety of physical conditions and complex methodologies for calculating turbulent fluxes, an assessment of the interoperability and consistency between flux products from different data collections is still necessary despite the existence of operational flux data networks such as FLUXNET (Baldocchi et al. 2001) and AmeriFlux (Baldocchi et al. 1996; Boden et al. 2013).

165 Solid and liquid precipitation are both measured at the surface but are notoriously difficult to characterize accurately. Precipitation accumulation with hourly to 6-hour frequency is traditionally measured by gauges within a surface precipitation network, usually operated by operational weather centers or hydrological agencies. Measuring solid precipitation presents a number of unique challenges; snow pillows and the Double-Fence Automated Reference (DFAR) configuration around gauges provide reliable estimations of precipitating snow. However, snow precipitation measurements from WMO standard installations with Single-Alter-shielded and unshielded gauges are affected by the undercatch of solid precipitation in windy conditions (Nitu et al., 2018; Kochendorfer et al., 2022). The WMO Solid Precipitation Intercomparison Experiment (SPICE) analyzed this undercatch and developed adjustment functions for correcting it (Kochendorfer et al., 2018; Wolff et al., 2015) which are now being used in verification practices (Køltzow et al., 2020; Buisan et al. 2020, Casati et al., 2023).

175 3.2 Upper Atmospheric Profiles

Radiosonde data are typically treated as if they are instantaneous vertical profiles of the troposphere by collapsing the balloon-borne trajectory of the instrument package into a profile that is time indexed with the launch time. The measurements of temperature, dewpoint temperature, pressure, humidity, and winds are high cadence (on the order of seconds) during the balloon ascent with individual launches being low cadence (every 12 hours standard launch intervals at 180 00 and 12 UTC). During intensive campaign periods, the launch frequency is often increased to 4–6 sondes per day. A full sounding can take 2 hours to ascend and can travel up to 200 km horizontally, and important fine-scale information is available if full, original trajectory information (time–height/pressure level–latitude–longitude) coordinates are maintained to assess the spatial displacement.

185 Many observatories and campaigns support the operation of radars, lidars, sodars, profilers and microwave radiometers which separately and in combination can remotely infer properties through the depth of the planetary boundary layer (PBL) and free atmosphere. The systems use active (transmission and interpretation of reflected signal) and passive (detection of natural atmospheric signal) sensing techniques. Through a significant body of research on retrieval methods, the systems can determine properties such as cloud base, cloud liquid water path, cloud ice water path, cloud liquid water content, cloud ice 190 water content, hydrometeor sizes and shapes, snowfall rates (Matrosov et al. 2022), degree of riming, aerosol extinction



coefficients, winds, temperature and humidity. These advanced products may be obtained from diverse instrument hardware configurations and technologies as well as site-specific scanning and collection schedules. Robust site-independent retrieval methodologies that are consistent across networks such as the products produced by Cloudnet (Illingworth et al., 2007) and the ARM Active Remote Sensing of Clouds product (Clothiaux et al., 2001) have been developed, however, they may not be consistently implemented.

3.3 Terrestrial Surface and Subsurface

The terrestrial surface and subsurface is composed of complex layers of different soil, rock, ice (permafrost) and vegetation layers that can be covered with surface snow and ice with evolving density, heat capacity, conductivity and chemistry. Thermistor strings can be co-located with tensiometers; the variables measured are gradients of temperature and moisture respectively with thermal conductivity and heat capacity determined as fixed intrinsic properties from soil samples. Thermistor strings can also be co-located with moisture probes, providing vertical profiles of soil temperature and moisture respectively down to depths ranging from 0.1 to 2 m depending on the soil type. Snow depth, like precipitation, is a difficult quantity to measure representatively; techniques vary from using simple measurement stakes to downward-looking mast- or tower-mounted acoustic devices.

3.4 Ocean, Sea Ice, and Snow Surface and Subsurface

Ocean thermistor measurements are accompanied with conductivity and pressure measurements with resulting variables determined for salinity, temperature and depth. Additional current meters allow determination of turbulent fluxes using eddy correlation techniques similar in principle to those used for atmospheric fluxes. Measurement of sea ice and snow macro- and microphysical properties below the atmosphere/ice or atmosphere/snow interface (i.e. subsurface) is increasingly sophisticated with upward and downward looking acoustic devices on ice buoys determining ice thickness and snow depth (Zuo et al., 2018). Measurements are made of snow and ice density, crystalline structure and salinity via manual sampling.

4 MODFs for the Year of Polar Prediction

The World Meteorological Organization (WMO) Polar Prediction Project (PPP) organized the Year of Polar Prediction (YOPP; PPP Steering Group & Co-authors, 2019; Jung et al., 2016) which concluded in 2022 (Wilson et al., 2023). The YOPPsiteMIP (Svensson et al., 2020; Day et al., submitted) working group envisioned matched sets of observation and model data to support model process diagnostics. The efforts of this working group resulted in a metadata schema for MODFs and MMDFs (collectively known as MDFs) as well as an iterative production workflow. The YOPPsiteMIP effort focused on Polar terrestrial stations, but because it was anticipated that a similar strategy would be applied to the YOPP endorsed Multidisciplinary drifting Observatory for the Study of Arctic Climate (MOSAIC) expedition (Shupe et al., 2022,



220 Nicolaus et al., 2022), the schema was developed to also accommodate expected additional ocean and sea ice observations from ships and on-ice platforms in the central Arctic Ocean.

The first steps in MODF production are to assemble the available data files that will feed into the MODF, extract the desired data, and acquire the corresponding metadata while noting that each individual instrument and instrument group requires
225 both unique quality control and processing together with attribution tracking to produce and document the provenance of the geophysical variables. Given the heterogeneity and often research-grade nature of campaign data, the amount and quality of metadata for different variables is likely to be inconsistent. Although required metadata are ideally harvested from the internally documented data files, it is likely that many data sets will need additional metadata that will require interviewing original data collectors. (For an example of this other than our own, see Papoutsoglou et al., 2023). Once individual data and
230 any available metadata from individual instruments and instruments suites are created and or assembled, the merging process can begin.

As we developed the schema and workflow, we identified the following challenges and solutions.

235 Semantics: Data semantics address the issue of the same variable being given different names drawn from multiple or ad hoc naming conventions. A significant part of the MODF solution has been the development of an extensive schema based on already existing vocabulary standards.

Units: Variables' units are frequently absent entirely from data sets or expressed with nonstandard abbreviations (Hanisch et al., 2022). The MODF solution is to associate each variable with recommended units, typically as identified in the CF Standard Name Table (2023). These are meant to be compatible with and generally recognized by the Unidata (2020) UDUNITS package (Eaton et al., 2022).

Attribution: Perhaps the greatest MODF challenge is not technical but rather cultural. When data for a single MODF
245 product come from instruments operated from multiple institutions and researchers, there are complex issues with acknowledging the original sources of the data (Pierce et al., 2019; Nature Editorial, 2022). Data from campaigns or programs usually have multiple institutions and individual researchers involved, all of whom have different performance metrics for original research. This often leads to official or implicit data embargos so that the researchers who collected the data will have the first opportunity to publish research results. The design of MODFs is responsive to these issues by
250 providing a high level of attribution metadata including links to original data and data producers, supporting copious data citations (Vannen et al., 2020).


Data Heterogeneity: Measurement heterogeneity is generated by how data are collected and processed. The MODF solution is to document variable derivation as much as possible from original data metadata: type of instrument, calibration
255 information, method of deployment, quality control and processing histories, and original licenses.

Inconsistent Cadences: Different variables are collected on a range of native cadences varying from Hz for the fast response sensors needed to record phenomena that can change on short time scales (typically in the atmosphere) to sensors for which it is sufficient to sample on hourly or even daily time scales (typically in the terrestrial sub-surface) with a wide range in-
260 between (typically for the ocean and ice sub-surfaces). The MODF solution is for each variable to retain its native recording cadence and to minimize any temporal averaging beyond that necessary for sensible data processing. In addition, original data sets are not interpolated to fill data gaps.

Redundancy: It is common during campaigns or at intensive observatory sites for there to be multiple measurements of the
265 same variable (e.g., temperature). There is important information contained in redundant measurements when evaluating how representative a variable is of site characteristics or when comparing the in-situ observations to model grid cells and satellite footprints to evaluate local variability. The MODF solution is to include as many redundant measurements as possible, maintaining high-accuracy location information including for example soil and vegetation characteristics that help define the observations' microclimates.

Processing Levels: Research-grade data sets typically require unique quality control and often extensive post processing which rarely can be fully automated. Frequently the state variables (e.g. temperature) are readily available in near real-time, whereas more complex variables such as turbulent fluxes can have a wide range in availability depending on whether they are output by commercial software (Fratini and Mauder, 2014), calculated from bulk variables, or based on customized
275 calculations for complex environments. The MODF solution is to include full processing, data quality and usability metadata.

Versions: Since it is common for observed data to have versions, the MODF concept accommodates multiple product releases. Examples of data versions are, raw (sometimes just voltages); minimally quality checked by automated elimination
280 of extreme physically unrealistic outliers; subjectively curated for unusual and if possible correctable operational or environmental conditions; and different levels of processing to produce retrieved variables (those based on more direct measurements). Frequently, highly processed and certified data are available months or even years after the original collection period. In some cases, all processing levels of data are available and archived. Two obvious ways of handling the situation are to either replace original data with more processed data as it becomes available (Figure 2A) or to keep all
285 versions of data (Figure 2B). MODFs' rich metadata includes careful version tracking for individual variables which encourages the latter.



290 **Figure 2: There are 2 approaches for MODF augmentation (as new variables become available) and modification (as variables proceed through processing levels). In both approaches, variables that are immediately available (Var 1, Var 2) can be immediately ingested and variables can be added as they become available (Var 3, Var 4). The 2 approaches accommodate two strategies for variables that undergo post processing resulting in products with quality-control (QC) levels. In (a), original data (Var 3 v1.0) can be replaced with QC'ed data (Var3 v2.0). In (b), original data (Var3 v1.0) can be retained in the MODF and QC'ed data (Var3, 2.0) can be added.**

295 5 The H-K Variable Schema Table

The H-K Variable Schema Table, developed for the YOPPSiteMIP (Hartten and Khalsa, 2022, hereafter the H-K Schema), provides guidelines for creating both MODF and MMDFs as netCDF files (Unidata, 2022) with consistent variable names and metadata. Hereafter the discussion in this paper centers on those entries which are relevant to MODFs. The H-K Schema follows the Attribute Convention for Data Discovery (ACDD; Earth Science Information Partners (ESIP), 2022), the 300 NetCDF Climate and Forecast (CF) Metadata Conventions (Eaton et al., 2022), and the ISO/TC 211 19100 series of standards for digital geographic information. Short variable names are compliant with the World Climate Research Program (WCRP) Coupled Model Intercomparison Project, Phase 6 (CMIP6; Taylor et al., 2022) whenever possible as these are in wide use in the geophysical research community. For global attributes, the H-K Schema specifies identifying the file feature type, the file maker, the license, the location, information on when data was collected, and a version number associated with the MODF and its permanent identifier allowing for MODF augmentation. For variable attributes, the H-K Schema specifies 305 identifying vocabularies, units, individual variable attribution (who originally collected the data) and variable provenance, and also presents a method for differentiation of multiple measurements of the same geophysical variable (redundancy). The H-K Schema is available in both json (machine readable) and pdf (human readable) formats. The A-M Variable and



310 Attribute Template Table (Morris and Akish, 2022) was developed iteratively in conjunction with the H-K Schema for collating metadata and was used as a direct input to the MODF creation process.

5.1 Global Attributes

315 Global attributes are the descriptive metadata that are relevant for the entire MODF file. Table 1 lists recommended global attributes for MODFs. These were chosen because they are highly recommended by ACDD; because they are recommended by ACDD and made sense for our purpose, and seemed not overly burdensome for those most likely to be making MODFs; and/or because we felt they would help MODFs be FAIR. Some global attributes whose inclusion and consistent use is particularly important are described below. We use italics for the names of attributes and single quotes around variable names.

320 *id*: The global unique Persistent Identifier (PID). This global attribute does not supersede references to the PIDs associated with individual variables (Prakesh, et al., 2016), which can be provided within variable attribute *references*. The MODF *id* remains constant as modifications or additions are made to the MODF file.

325 *license*: Specifies terms of distribution. As with *id*, individual variables and groups of variables in the MODF may have unique license requirements that can be recorded in the variable attribute *comment*.

330 *creator_name*: The names of individuals who should be in the citation for the MODF (c.f. Jones et al. 2018, §2.2.7.3.3). The DataCite Metadata Working Group (2021) defines creators in this context as the main producers of the MODF file and also links them to authorship. (Note that the main producers of component data sets are credited within the variable attributes coming from their individual data sets.) Enabling those who create MODFs to receive appropriate credit for their work is a key element of making MODFs FAIR; data reuse depends in part on data accessibility, which in turn is more likely when data are “considered legitimate, citable products of research” (Data Citation Synthesis Group, 2014).

335 *featureType*: Although the original vision was to have a single MODF for a site during a campaign that would include all relevant variables, when considering the practicalities of archive submission, it became apparent that file *featureType* needed to be specified to facilitate data services such as visualization tools. Most observatory data fits into *featureTypes* timeSeries, timeSeriesProfile and timeSeriesTrajectory. These *featureTypes* are defined by the temporal-spatial dimensions that are associated with individual variables: time only for timeSeries, time and height (or depth) for timeSeriesProfile, and time–height (or depth)–latitude–longitude for timeSeriesTrajectory.

340 *missing_value*: Establishes one consistent *missing_value* for all data in the MODF.



Table 1: H-K Schema Recommended Global Attributes (adapted from Hartten and Khalsa, 2022)

Attribute	Description of Required Information
<i>title</i>	Short human-readable phrase or sentence describing the MODF
<i>date_created</i>	Date on which current version of the MODF was created or modified
<i>Conventions</i>	List of the conventions that are followed by the MODF
<i>standard_name_vocabulary</i>	Name and version of the controlled vocabulary from which variable Standard Names come
<i>creator_name</i>	Name of the main individual(s) involved in producing the MODF, or the authors of the publication, in priority order
<i>creator_email</i>	Email of the main individual(s) involved in producing the MODF, or the authors of the publication (if there is one) describing the MODF
<i>institution</i>	Institution where the original MODF was produced
<i>id</i>	Persistent Identifier (PID) for the MODF
<i>naming_authority</i>	Naming authority for PID (preferably using reverse-DNS naming, e.g. 'edu.ucar.unidata', although URIs may be used)
<i>license</i>	Terms of distribution and use
<i>time_coverage_start</i>	Time of the first data point in the MODF
<i>time_coverage_end</i>	Time of the last data point in the MODF
<i>featureType</i>	One of the following: 'point', 'timeSeries', 'trajectory', 'profile', 'timeSeriesProfile', or 'trajectoryProfile'
<i>contributor_name</i>	Name of any individual(s) responsible for collecting, managing, distributing, or otherwise contributing to the development of the MODF; may indicate those who helped with the development but who were not so "key" as to be listed as an author
<i>contributor_email</i>	Email of any individual(s) responsible for collecting, managing, distributing, or otherwise contributing to the development of the MODF
<i>project</i>	Name of the project(s) principally responsible for originating this MODF
<i>summary</i>	A paragraph describing the MODF
<i>source</i>	The method of production of the MODF
<i>metadata_link</i>	URL to detailed documentation (DOIs expressed as http://doi.org/...)
<i>history</i>	Provides an audit trail for modifications to the MODF
<i>references</i>	Published or web-based references that describe the MODF or methods used to produce it
<i>keywords</i>	Comma-separated list of key words and/or phrases, preferably drawn from a controlled vocabulary, e.g. the GCMD at https://earthdata.nasa.gov/earth-observation-data/find-data/gcmd/gcmd-keywords



5.2 Time, Space and Site Variables

In the H-K Schema, variables have been divided into subgroups. There are three subcategories with time, space and site information: temporal dimensions and variables; spatial dimensions and variables; single level fixed variables. Per the NetCDF Users Guide (Unidata, 2023), dimensions “may be used to represent a real physical dimension, for example, time, latitude, longitude, or height... [or] to index other quantities, for example station or model-run-number.” Examples of these indexing variables are listed in Table 2 and discussed further below. Per the H-K Schema, all dimensional and geophysical variables are given a short CMIP or CMIP-like variable name and are characterized by variable attributes *long_name*, *standard_name*, *units*, additional recommended attributes, and any other attributes the MODF creators feel necessary to fully document the data.

Table 2: Examples of dimensions and variables related to time, space, and site information, together with the associated *long_name*, *standard_name*, *units* and recommended attributes (extracted from Hartten and Khalsa, 2022).

Variable Name (CMIP or CMIP-like)	<i>long_name</i> Attribute	<i>standard_name</i> Attribute	<i>units</i> Attribute	Minimum Recommended Additional Attributes
TEMPORAL DIMENSIONS AND VARIABLES				
time	Valid Time	time	hours since ...	
time15	Valid Time for Observations with 15-Minute Cadence	time	hours since ...	<i>delta_t</i> ; <i>calendar</i>
time_sonde	Radiosonde Valid Time	time	hours since ...	
SPATIAL DIMENSIONS AND VARIABLES				
height_tower	Tower Observation Height	height	m	_sonde variables only:
lat_sonde	Radiosonde Latitude	latitude	degrees_north	<i>missing_value</i> ;
lon_sonde	Radiosonde Longitude	longitude	degrees_east	<i>actual_range</i> ; <i>instrument</i> ;
alt_sonde	Radiosonde Altitude	altitude	m	<i>source</i>
SINGLE-LEVEL FIXED VARIABLES				
orog	Surface Altitude	surface_altitude	m	<i>missing_value</i> ; <i>source</i> ; <i>references</i> ; <i>comment</i>



355 ‘time’, ‘time15’, ‘time_sonde’: MODFs are intended to retain as much high-resolution information from the original data
collection as possible. Averaging is limited to that necessary for processing (e.g. for eddy-correlation flux calculations) and
data are not interpolated. Therefore, temporal dimensions support maintaining original data collection cadences for
individual variables. For instance, in MODFs ‘time’ is a generic temporal dimension whereas ‘time15’ is a time dimension
associated with variables collected at 15 min intervals and ‘time_sonde’ is the time dimension associated with the data
360 collected during a radiosonde ascent. In the YOPPsiteMIP implementation of MODFs, we have chosen to append ‘N’ or
‘_platform’ to a generic CMIP name such as ‘time’ in order to indicate a time array with a particular cadence in minutes or a
time array tied to a particular instrument or platform. In keeping with CF Metadata guidelines, all the differently named time
variables can and should have the same long_name attribute.

365 ‘height_tower’: A spatial coordinate can be a single scalar value or a set of values (a dimension) that describes the location
where geophysical measurements are collected. This can be an array of fixed values in the case when measurements are
made at set levels, for instance on an instrumented tower. A tower may also consist of multiple measurements of the same
geophysical variable collected at different heights above the ground. For instance, observations of air temperature collected
by three sensors located at the surface (2 m AGL), on a pole (10 m AGL), and on top of a building (20 m AGL) could be
370 combined into a single array designated as ‘_tower’ so long as the sensors’ horizontal positions were co-located.”

‘lat_sonde’: Spatial coordinates can also be a dimension variable or a scalar value. For instance, if information is retained on
the latitude, longitude and/or altitude of a sonde during its ascent, this information should be put into coordinate variables
that provide the location in space for the geophysical variables collected by the sonde. If the sonde output includes only
375 varying altitudes and a fixed launch time, the geophysical data from each sonde flight should be put into a file with the
featureType “profile”. If the sonde output includes varying altitudes and associated times, the geophysical data should be put
into a file with the *featureType* “timeSeriesProfile”. In both cases a scalar altitude variable should be used as the vertical
dimension for each flight. However, if varying latitude and longitude are also provided, the geophysical data from each
sonde should be put into a file with the *featureType* “timeSeriesTrajectory” with the reported altitude, latitude and longitude
380 being provided as coordinate variables for each flight.

‘orog’: Certain fixed variables are descriptive of the site or the platform from which measurements are taken. An example of
the former is the variable with the surface altitude (CMIP name ‘orog’), which describes the site’s altitude above the surface
defined as the lower boundary of the atmosphere. Other common single-level fixed variables are the ‘lat’ (latitude) and the
385 ‘lon’ (longitude) chosen to generally identify a site location. If the site has distributed measurements, individual variables
may have more refined ‘lat_platform’ and ‘lon_platform’ variables or dimensions.



5.3 Geophysical Variable Attributes and Examples

There are six categories of geophysical variables: Single-level atmosphere variables; surface and Top-of-Atmosphere (TOA³) variables; atmospheric variables on model or instrument levels; sub-surface terrestrial variables; oceanic variables on model or instrument levels; sea ice variables. Examples of *long_name*, *standard_name*, and *units* attributes for observed geophysical variables that have been catalogued in the H-K Schema are shown in Tables 3 and 4. A full listing of all the single-level atmospheric variables, the surface and TOA variables, and the atmospheric variables on instrument levels in the H-K Schema version 1.2 is presented in Table A1, while a listing of the current H-K Schema oceanic variables on model or instrument levels, sub-surface terrestrial variables, oceanic single-level variables, and sea ice variables is found in Table A2. Some discussion of the examples in Tables 3 and 4 follows.

CMIP name: The Coupled Model Intercomparison Project (CMIP) names are taken from the CMIP6 Participation Guidance for Modelers, a program supported by the NASA Program for Climate Model Diagnosis & Intercomparison. CMIP names prioritize terse, presumably code-efficient abbreviations such as ‘ta’ (air temperature). To clearly identify the variables that were observed or derived redundantly from different methods or platforms, a suffix can be appended (e.g., *_tower*, *_radar*, *_8m*). Some standard measurements, such as 2m temperature and 10m winds, have unique CMIP names such as ‘tas’, ‘uas’ and ‘vas’ (near-surface air temperature, eastward wind, and northward wind, respectively). In the case where a CMIP variable name has not been defined in the CMIP6 vocabulary, a CMIP-like name has been composed.

long_name: Fully describes the physical quantity and can be thought of as useful attribute for labeling plot axis; in other words, it is the name that best communicates with humans about what the variable is (note that the *long_name* does not necessarily need to be in English). The H-K Schema provides ad hoc *long_name* definitions for all variables.

standard_name: Taken from the CF Standard Name Table (2023), which is periodically updated based on community requests and discussion. Standard names are constructed in conformance with the CF Metadata Conventions (Hassell et al., 2017). Different variables can have the same standard name. For instance, ‘albs’ and ‘albsn’ both have the *standard_name* of *surface_albedo*, but ‘albsn’ is restricted to use over snow-covered areas. Redundant variables (such as multiple measurements of temperature, or fluxes computed by bulk versus eddy-correlation methods) will have the same standard name and will require differentiation by a suffix added to the CMIP name, additional descriptors in the *long_name*, and possibly accompanying spatial-temporal indices.

³ TOA variables are only relevant to MMDFs, but are grouped with surface variables in the H-K Schema and therefore included for completeness.



Table 3: Examples of atmospheric variables, together with the associated *long_name*, *standard_name*, and *units* attributes (extracted from Hartten and Khalsa, 2022).

Variable Name (CMIP or CMIP-like)	<i>long_name</i> Attribute	<i>standard_name</i> Attribute	<i>units</i> Attribute
SINGLE-LEVEL ATMOSPHERIC VARIABLES			
tas	Near-Surface (2m) Air Temperature	air_temperature	K
tas_site1	Near-Surface (2m) Air Temperature at Site1	air_temperature	K
uas	Near-Surface (10m) Eastward Wind	eastward_wind	m s-1
SURFACE AND TOA VARIABLES			
snd	Surface Snow Thickness	surface_snow_thickness	K
ts	Surface (Skin) Temperature where Land or Sea Ice	surface_temperature	K
rsds	Downward Shortwave Radiation at the Surface	surface_downwelling_ shortwave_flux_in_air	W m-2
ATMOSPHERIC VARIABLES ON MODEL OR INSTRUMENT LEVELS			
ta	Temperature	air_temperature	K
rsd	Downward Shortwave Radiation	downwelling_ shortwave_flux_in_air	W m-2
ua	Eastward Wind Component	eastward_wind	m s-1



420 **Table 4: Examples of non-atmospheric variables, together with the associated *long_name*, *standard_name*, and *units* attributes (extracted from Hartten and Khalsa, 2022).**

Variable Name (CMIP or CMIP-like)	<i>long_name</i> Attribute	<i>standard_name</i> Attribute	<i>units</i> Attribute
SUB-SURFACE TERRESTRIAL VARIABLES			
gtsl	Bulk Soil Temperature	soil_temperature	K
mrsl	Layer-Average Soil Moisture	moisture_content_of_soil_layer	kg m ⁻²
OCEANIC SINGLE-LEVEL VARIABLES			
tos	Sea Surface Temperature	sea_surface_temperature	K
mlost	Ocean Mixed-Layer Depth	ocean_mixed_layer_thickness	m
rsntds	Net Downward	net_downward_	W m ⁻²
	Shortwave Radiation at Sea Water Surface	shortwave_flux_ at_sea_water_surface	
OCEANIC VARIABLES ON MODEL OR INSTRUMENT LEVEL			
to	Ocean Temperature	sea_water_temperature	K
so	Sea Water Salinity	sea_water_salinity	1e-3
SEA ICE VARIABLES			
sithick	Sea Ice Thickness	sea_ice_thickness	m
sisali	Sea Ice Salinity	sea_ice_salinity	1e-3

Table 5 lists the other attributes that should be included in MODFs for geophysical variables. Some of the variable attributes in Table 5 are listed in the CF Conventions as being for use as either global or data attributes, but the *history* attribute is listed as for global use only. We have encouraged its use with variables (data) in MODFs because we believe that maintains the spirit if not the letter of the CF Conventions; like *institution*, *references*, *source*, and *title*, *history* helps document the provenance and nature of the data included in these multi-institution, multi-sourced files. We also encourage MODF (and MMDF) makers to make use of additional data attributes to share information about the variable contents with users.

430 *original_name*: Refers to the name of the variable in the original file from which it was extracted. This provides an important cross reference in the case where a user may need to refer back to the original source data set.



435 *instrument*: Tracking instrumentation characteristics and in many cases calibration coefficients is critical to provide information for users who may be in the process of developing refined data processing, developing higher-order products, or doing instrument intercomparison studies.

references: Published material that describes either the data or the methods used to produce it should be listed here. Best practice is to include a URI (a DOI for a paper, or a URL for a website).


440 *source*: Both the CF Conventions and ACDD describe this as “the method of production of the original data”, by which they mean either the model that produced it or the instrument that gathered it. In either case, the idea is to give the user information that will help them understand what they’re working with, what assumptions or methods are inherent to the data. Therefore, the name and version of a numerical model, or the type and perhaps make and model of an instrument, would be appropriate here.

445 *history*: Provides an audit trail for the data, documenting its provenance. When used as a variable attribute, this should trace what has been done to the data from its raw state until it was put into this file. Ideally, each step should be documented by a line containing a timestamp, the person or entity who did the action, a brief description of the action, and any program(s) which accomplished the action (including relevant settings or command arguments). When used as a global attribute, this is
450 the attribute in which changes to the file should be recorded.

Table 5: Minimum recommended additional variable attributes for geophysical variables (adapted from Hartten and Khalsa, 2022)

Additional Variable Attributes	
<i>missing_value</i>	<i>contributor_name</i>
<i>actual_range</i>	<i>contributor_email</i>
<i>instrument</i>	<i>creator_name</i>
<i>source</i>	<i>creator_email</i>
<i>references</i>	<i>institution</i>
<i>history</i>	<i>comment</i>
<i>original_name</i>	

455 Figure 3 describes the work flow and process that developed through the efforts of the YOPPsiteMIP team of observers, modellers, data scientists, and data managers. This is a highly iterative process; note the two-way arrows between the H-K Schema and both the A-M Table and the MODF Checker.



460 **Figure 3: The MODF workflow process. A: Gather input files, B: Interview contributing data principals for necessary metadata**
not digitally encoded in data files, C: Utilize the A-M MODF template to construct the specific MODF framework based inputs
from A and B compliant with the H-K Schema (D). MODF Makers create the MODF (E) with inputs from (C) and the data values
from individual files collected (A). F: MODF file. G: MODF checking. H: Upon ‘pass’, send MODF to archive. I: Upon ‘fail’ assess
MODF. J: Major fails, return MODF Makers. K: Minor fails, archivist corrects and sends MODF to archive after rechecking (G).
 465 **L: Iterative developments between H-K schema (D) and MODF checking (L). People icons adapted from one designed by**
vectorstock (Image #45970079 at VectorStock.com).

6 Discussion

We have presented the H-K Schema (Hartten and Khalsa, 2022) and a production framework for organizing complex campaign and observatory data from multiple instruments into Merged Observatory Data Files (MODFs). The H-K Schema
 470 can also be used to format forecast model output into corresponding Merged Model Data Files (MMDFs). MODFs and
 MMDFs are compliant with existing metadata and data standards that support FAIR principles. The Schema and the
 framework were developed by a YOPPSiteMIP working group of observers, modelers and data managers. MODFs address
 the mundane but complex issues that arose from the YOPPSiteMIP vision to confront polar weather forecasting models with
 observations from richly instrumented sites during special observing periods. The issues addressed include semantics,
 475 attribution for original data, data provenance, different cadences, multiple measurements of the same variable from the same
 site (including local, subgrid-scale spatial distribution), versioning strategies to account for different levels of data

processing, unit conventions, and missing data indicators. Because of the high cadence of many measurements (seconds to minutes), MODFs can be used to evaluate accumulating biases at sub-model timestep increments.

480 Although providing site, instrument, processing, and attribution metadata documentation is good practice, it is often neglected. Since this essential MODF information is difficult to assemble after the fact, we recommend that observers use datagrams during development, deployment, and operation of sensors as a standard practice. Datagrams “are designed to document the life story of a data value from start to finish and provide a guide for humans to design, deploy, troubleshoot, repair, record, transmit, process, and archive data collected with measuring devices.” (Morris and Uttal, 2022)

485
Currently the H-K Schema includes geophysical variables across the atmosphere (state, clouds, turbulence, radiation), snow, ice, terrestrial, and ocean systems. Appendix A lists the MODF (and MMDF) values that are currently in the H-K Schema (version 1.2). These are not exhaustive, and we expect that MODF users and makers will augment their MODF files to accommodate individual campaigns by including, for example, atmospheric aerosols, constituent gases, additional ocean and
490 terrestrial variables, and ecosystem and biogeochemical data. Additional variables should be incorporated with the metadata standards described here.

A first set of MODF and MMDF files implementing the H-K Schema from concept to production for YOPP Special Observing Periods have been archived by the Norwegian Meteorological Service where a collection⁴ of matched MODF and
495 MMDF files are available for several Arctic ground stations. An initial analysis (Day et al., submitted) using the YOPPsiteMIP MODF–MMDF collection demonstrates through a number of examples and case studies how process-describing variables such as surface fluxes (longwave, sensible and latent) and ground fluxes can be used to gain deeper understanding into the nature of forecast errors. Although we strongly encouraged YOPPsiteMIP MODF and MMDF Makers to check their files for compliancy before submitting, we have found it expedient to have the recipient data managers
500 at the host archive take the final responsibility for inspecting, certifying, and in some cases making editorial corrections to bring the MODFs and MMDFs into compliance with the H-K Schema. Unfortunately when archives have rigorous standards for submittal, data providers often do not submit at all because they do not have the resources to support that level of data management.

505 Generating MODFs will expand usage of data from field campaigns by increasing data uptake and decreasing data latency. This will promote usage of non-operational data that are currently underutilized and difficult to access comprehensively, specifically for environmental services requiring near real-time environmental intelligence. The U.S. Interagency Arctic Research Policy Committee (IARPC) has defined environmental intelligence as “a system through which information about

⁴ https://thredds.met.no/thredds/catalog/alertness/YOPP_supersite/catalog.html



a particular region or process is collected for the benefit of decision makers through the use of more than one inter-related source”. Furthermore, IARPC notes that “Traditionally, researchers collect data, develop models, and communicate results through well-established channels that are often slow and inefficient. While the vetting of scientific results ensures that the conclusions are of highest quality, the process is not well-aligned with the need for rapid information delivery in the face of environmental transitions that are putting stress on ecosystems and human populations.” MODFs will not only accelerate timely and relevant data access and scientific results for the primary researchers that collected the data, but will also support the iterative observations, modeling and data systems that connect researchers, stakeholders and decision-makers to allow informed responses to environmental events. To this end, MODFs are designed to be living files that can be created in a timely manner with near real-time variables and then augmented when additional variables become available. This is particularly important for situations in which observational data becomes obsolete before it can be utilized by environmental awareness and short-term prediction services for extreme events. MODFs can then be augmented with variables that are only available after extensive human-assisted processing (e.g., surface energy balance fluxes) for research purposes, as well as with new versions of variables that require detailed quality control to produce higher level and higher reliability products. This addresses the concern that many data providers express about only releasing the most highly curated values (often resulting in data embargos and release time lags) and recognizes that the level of necessary post processing depends on the application.

525

We expect that MODFs and MMDFs, in other words Merged Data File (MDF) collections, will also be used to facilitate the concept of system science. The NSF workshop report Opportunities and Challenges of Arctic System Science (Vorosmarty et al., 2018) introduced the concept of “... multiple ‘currencies’ that link the Arctic climate and environment—geophysical entities such as water, energy, carbon, and nutrients with quantifiable properties—and how they interact to produce and illuminate systems-level behaviors”. MODFs, by quantifying the currencies throughout a system with internally consistent standards, can serve to break down barriers between individual studies of separate components of the system that are typically divided along disciplinary lines, thereby advancing multidisciplinary process studies.

530

The concept of organizing multivariate observational data sets from a single site or platform into a unified product is not unique. Wei et al. (2021) describes how ICARTT data format conventions (Aknan et al., 2013) can be used to create merged in-situ data products from research aircraft carrying multiple sensors. The U.S. DOE ARM program (Stokes and Schwartz, 1994) creates Climate Modeling Best Estimate (CMBE) data files (Xie et al., 2010) for their Climate Reference Sites. Hogan and O’Connor (2004) describe how the Cloudnet project preprocesses cloud data from multiple remote sensors using appropriate ancillary observations and model forecasts, then makes the output available in netCDF format with rich metadata about the preprocessing. What is unique about MODFs is the strategy of creating corresponding matched MMDF data files to support modeling verification and process evaluations. This addresses a long-standing issue with communication silos between observing and modeling sciences (Holloway et al., 2014, Sprintall et al., 2021, Neang et al., 2021).

540



Appendix A: Current Measurable Geophysical MODF and MMDF Variables and Their “Essential Climate Variable” Status

545 The H-K Schema is a set of tables that is expected to be expanded and adjusted as the need arises for additional MODFs and MMDFs and for additions to existing ones. Therefore, researchers interested in using the H-K Schema to guide file creation should always refer to the current online version; the DOI listed in the References of this article will always land on the most recent version of the Schema, and when a new version of the H-K Schema is published the earlier versions will be prefaced with a note including a link to the latest version.

550

In this Appendix we list the *long_name* attribute for all atmospheric (Table A1) and non-atmospheric (Table A2) geophysical variables, current as of the time of this manuscript’s submission. We do this for two reasons: to show the variety of variables incorporated in the H-K Schema so far, and also to highlight the variables in the Schema which the WMO has identified as Essential Climate Variables (ECVs; Essential Climate Variables, 2023). Note that some of the ECVs are defined by the WMO in a manner other than the point measurements typically made at field sites. Those details are identified by footnotes in the tables. Lavergne et al. (2022) have proposed expanding the list of ECVs related to sea ice; proposed sea ice ECVs that are in the H-K Schema are separately highlighted in Table A2.



Table A1: Atmospheric variables (extracted from Hartten and Khalsa, 2022) that are measured at Arctic YOPP supersites or included in Arctic YOPP MMDFs (prefaced by “•”). Underlined variables are ECVs.

Variables' long_name Attribute	
SINGLE-LEVEL ATMOSPHERIC VARIABLES	
<u>Surface pressure</u>	Height of atmospheric boundary layer
Mean sea level pressure	<u>Total precipitation of water in all phases per unit area</u>
<u>Near-surface eastward wind</u>	<u>Total cloud cover</u>
<u>Near-surface northward wind</u>	<u>Cloud optical thickness</u>
<u>Near-surface wind speed</u>	<u>Total column water vapour</u>
<u>Direction near-surface wind from</u>	<u>Total column cloud water in liquid phase</u>
Near-surface wind gust	<u>Total column icewater</u>
<u>Near-surface air temperature</u>	Surface horizontal visibility
Near-surface dew point temperature	<u>Ozone concentration in air^a</u>
Near-surface specific humidity	Photosynthetic photon flux density
Near-surface relative humidity	Reflected photosynthetic photon flux density
Surface roughness for momentum	
Surface roughness for heat	
SURFACE AND TOP-OF-ATMOSPHERE VARIABLES	
<u>Surface snow thickness</u>	• <u>Top-of-atmosphere incoming shortwave radiation</u>
• <u>Surface snow area fraction</u>	• <u>Top-of-atmosphere outgoing shortwave radiation</u>
<u>Snow water equivalent</u>	• <u>Top-of-atmosphere outgoing long wave radiation</u>
Snow density	<u>Upward surface shortwave radiation</u>
Surface (skin) temperature where land or sea ice	<u>Downward shortwave radiation at the surface</u>
Snow surface skin temperature	<u>Net shortwave radiation at the surface</u>
Snow temperature	<u>Upward surface longwave radiation</u>
Ground skin temperature	<u>Downward surface longwave radiation</u>
<u>Surface albedo^b</u>	<u>Net longwave radiation at the surface</u>
<u>Snow albedo^b</u>	• <u>Surface turbulent latent heat flux</u>
Surface downward heat flux in snow	<u>Surface turbulent latent heat flux^c (bulk method)</u>
Downward heat flux at snow bottom	<u>Surface turbulent latent heat flux (eddy covariance method)</u>
• <u>Canopy area fraction</u>	• <u>Surface turbulent sensible heat flux</u>
Time-average eastward turbulent surface stress	<u>Surface turbulent sensible heat flux (bulk method)</u>
Time-average northward turbulent surface stress	<u>Surface turbulent sensible heat flux</u>
	(eddy covariance method)
	Ground heat flux

^a Total column ozone is the listed ECV, while observations from a site are typically point values.

^b The ECV listing is for maps, not point measurements, of albedo; snow albedo is not explicitly listed.

^c The ECV listing specifies “Land-Biosphere Evaporation from Land”.



Variables' long_name Attribute

ATMOSPHERIC VARIABLES ON MODEL OR INSTRUMENT LEVELS

•Geopotential height	<u>Upward longwave radiation</u>
Geopotential height on half levels	<u>Downward longwave radiation</u>
Atmospheric pressure	<u>Upward shortwave radiation</u>
Pressure on full levels	<u>Downward shortwave radiation</u>
Pressure on half levels	•Vertical eddy diffusivity coefficient for momentum due to parameterized turbulence
<u>Eastward wind component</u>	•Vertical eddy diffusion coefficient for temperature due to parameterized turbulence
<u>Northward wind component</u>	Turbulent sensible heat flux based on virtual potential temperature
<u>Wind speed</u>	Turbulent moisture flux based on vapor content
<u>Direction wind from</u>	Eastward turbulent momentum flux
Vertical velocity	Northward turbulent momentum flux
•Vertical large-scale wind in pressure coordinates	Turbulent kinetic energy
<u>Air temperature</u>	•Percentage cloud cover, including both large-scale and convective cloud
Dew-point temperature	Mass fraction of cloud liquid water
Specific humidity	Mass fraction of cloud ice
Relative humidity	Snowfall flux per unit area
Wet-bulb potential temperature	<u>Cloud base height</u>
•Tendency of air temperature	
•Tendency of air temperature due to advection	
•Tendency of specific humidity	
•Tendency of specific humidity due to advection	
•Tendency of eastward wind	
•Tendency of northward wind	



565 **Table A2: Non-atmospheric variables (extracted from Hartten and Khalsa, 2022) that are measured at Arctic YOPP supersites or included in Arctic YOPP MDMFs (prefaced by “•”). Underlined variables are ECVs, with a dashed underline used for newly proposed sea ice ECVs (Lavergne et al., 2022).**

Variables' long_name Attribute	
SUB-SURFACE TERRESTRIAL VARIABLES	
Temperature of soil	<u>Average layer soil moisture</u>
OCEAN SINGLE LEVEL VARIABLES	
<u>Sea surface temperature</u>	<u>Atmosphere-ocean sensible heat flux</u>
Ocean mixed-layer depth	<u>Atmosphere-ocean latent heat flux</u>
<u>Ocean surface x-stress</u>	Net downward shortwave radiation at sea water surface
<u>Ocean surface y-stress</u>	Net downward longwave radiation at sea water surface
<u>Significant wave height^d</u>	Fresh water flux into sea water •Water flux into sea water due to sea ice thermodynamics
OCEAN VARIABLES ON MODEL OR INSTRUMENT LEVELS	
Ocean temperature	Ocean u-velocity
Sea water salinity	Ocean v-velocity Ocean w-velocity
SEA ICE VARIABLES, REPORTED ON ATMOSPHERIC GRID	
<u>Sea ice concentration (area fraction)</u>	Rainfall rate over sea ice
•Sea ice concentration (area fraction) in categories	<u>Sea ice surface temperature</u> (at the interface of sea ice or the snow on it and the overlying air)
<u>Sea ice thickness</u>	Temperature at snow-ice interface
•Sea ice thickness in thickness categories	Temperature at ice-ocean interface
<u>Snow thickness on sea ice</u>	<u>Sea ice/snow albedo</u>
<u>Sea ice age</u>	Ocean-ice net sensible heat flux
Sea ice u-velocity	Net upward sensible heat flux over sea ice
Sea ice v-velocity	Net upward latent heat flux over sea ice
Sea ice salinity	Downwelling shortwave flux over sea ice
•Sea ice normal stress (pressure)	Upwelling shortwave flux over sea ice
•Compressive sea ice strength	Downwelling longwave flux over sea ice
Fast ice concentration (area fraction)	Upwelling longwave flux over sea ice
Fast ice thickness	Net conductive heat flux in ice at the surface

^d ECV listing is for just wave height.



570 **Author Contributions**

TU conceptualized the original MODF vision; CJC and MM participated in conceptualization by providing valuable perspectives on the practicality, utility and application of the MODF concept. BC, GS and JD provided methodology requirements from a numerical modeling perspective. LMH and SJK, in developing the H-K Schema, provided methodology requirements from an observational perspective and also enhanced data curation. JT and OG guided development of the H-K
575 Schema and MODF formats by providing methodology requirements from a data repository standpoint.

Software contributions were provided by JH and JT, who wrote python code functions to facilitate MODF creation, checking and modification. Data curation, investigation, and software contributions were provided by EA, SM, NH, LXH, RC, ZM, EO, RP, JH and MM, who served as MODF Makers and through that practical application of the MODF concept developed
580 the workflow indicated in Figure 3. They also iteratively worked with LMH and SJK to further develop the H-K Schema. JT contributed to data curation, software, and validation by serving as the MODF Curator and by iteratively working with LMH and SJK to further develop the H-K Schema. GP and NH enhanced data curation by developing strategies to have MODFs include comprehensive formal attribution for individuals and institutions.

585 TU drafted the original manuscript, and TU, LMH, and SJK revised the draft, with commentary and revisions from BC, GS, JD, SM, EO, RP, ZM, JT, MM, and CJC. Visualizations, in the form of figures and tables, were drafted by TU and revised by TU and LMH.

Competing interests

The authors declare that they have no conflict of interest.

590 **Code and data availability**

The H-K Variable Schema Table developed for the YOPPsitesMIP is archived with Zenodo:

<https://zenodo.org/records/6463464>

The Table is licensed under the Creative Commons Attribution 4.0 International License

595 A preliminary set of MODF and MMDF files developed for the YOPPsitesMIP is available at:

https://thredds.met.no/thredds/catalog/alertness/YOPP_supersite/catalog.html



The catalog link provides information on licensing and crediting which is not repeated here as the example files are not the subject of this manuscript. Acknowledgements

Acknowledgements

600 This is a contribution to the Year of Polar Prediction (YOPP), a flagship activity of the Polar Prediction Project (PPP), initiated by the World Weather Research Programme (WWRP) of the World Meteorological Organization (WMO).

This work was supported in part by NOAA's Global Ocean Monitoring and Observing Program (FundRef <https://doi.org/10.13039/100018302>), the NOAA Physical Sciences Laboratory (TU, LMH, EA, SM, CJC), and
605 the NOAA Global Monitoring Laboratory (SM). LMH, EA, and SM were supported in part by NOAA cooperative agreements NA17OAR4320101 and NA22OAR4320151; LMH was also supported by NOAA's Climate Program Office, Climate Observations and Monitoring Program (FundRef 100007298). This work was also supported in part by the U.S. Department of Energy's Atmospheric System Research, an Office of Science Biological and Environmental Research program. JD was supported by European Union's Horizon 2020 Research and Innovation program through Grant
610 Agreement 871120 (INTERACTIII). RP was partly supported by the European Union's Horizon 2020 Research and Innovation program projects INTAROS (grant 727890) and PolarRES (grant 101003590).

Michael Gallagher (Univ. of Colorado/CIRES and NOAA Physical Sciences Laboratory) established the MODF Makers GitLab and Slack channels and wrote python functions to facilitate MODF creation. We are grateful to Dave Allured (Univ.
615 of Colorado/CIRES and NOAA Physical Sciences Laboratory) for extensive discussions about the CF Conventions, which greatly improved the MODF and MMDF projects. We also thank Scott Landolt (NCAR/Research Applications Lab) for discussions about measuring snow.

References

Aknan, A., Chen, G., Crawford, J., and Williams, E.: ICARTT File Format Standards V1.1, National Aeronautics and Space
620 Administration (NASA), ESDS-RFC-019v1.1, 21 pp., https://espoarchive.nasa.gov/sites/default/files/archive/ESDS-RFC-019-v1.1_0.pdf, 2013.

Andreas, E. L., Persson, P. O. G., Grachev, A. A., Jordan, R. E., Horst, T. W., Guest, P. S., and Fairall, C. W.:
Parameterizing Turbulent Exchange over Sea Ice in Winter, *Journal of Hydrometeorology*, 11, 87-104,
625 <https://doi.org/10.1175/2009JHM1102.1>, 2010.



630 Baldocchi, D., Falge, E., Gu, L., Olson, R., Hollinger, D., Running, S., Anthoni, P., Bernhofer, C., Davis, K., Evans, R.,
Fuentes, J., Goldstein, A., Katul, G., Law, B., Lee, X., Malhi, Y., Meyers, T., Munger, W., Oechel, W., Paw U, K. T.,
Pilegaard, K., Schmid, H. P., Valentini, R., Verma, S., Vesala, T., Wilson, K., and Wofsy, S.: FLUXNET: A New Tool to
Study the Temporal and Spatial Variability of Ecosystem-Scale Carbon Dioxide, Water Vapor, and Energy Flux Densities,
Bulletin of the American Meteorological Society, 82, 2415-2434, [https://doi.org/10.1175/1520-0477\(2001\)082<2415:FANTTS>2.3.CO;2](https://doi.org/10.1175/1520-0477(2001)082<2415:FANTTS>2.3.CO;2), 2001.

635 Baldocchi, D., Valentini, R., Running, S., Oechel, W., and Dahlman, R.: Strategies for measuring and modelling carbon
dioxide and water vapour fluxes over terrestrial ecosystems, Global Change Biology, 2, 159-168,
<https://doi.org/10.1111/j.1365-2486.1996.tb00069.x>, 1996.

640 Boden, T. A., Krassovski, M., and Yang, B.: The AmeriFlux data activity and data system: an evolving collection of data
management techniques, tools, products and services, Geosci. Instrum. Method. Data Syst., 2, 165-176, [10.5194/gi-2-165-2013](https://doi.org/10.5194/gi-2-165-2013), 2013.

645 Bojinski, S., Verstraete, M., Peterson, T. C., Richter, C., Simmons, A. J., and Zemp, M.: The Concept of Essential Climate
Variables in Support of Climate Research, Applications, and Policy, Bulletin of the American Meteorological Society, 95,
1431-1443, <https://doi.org/10.1175/BAMS-D-13-00047.1>, 2014.

Boukabara, S.-A., Krasnopolsky, V., Penny, S. G., Stewart, J. Q., McGovern, A., Hall, D., Ten Hove, J. E., Hickey, J.,
Huang, H.-L. A., Williams, J. K., Ide, K., Tissot, P., Haupt, S. E., Casey, K. S., Oza, N., Geer, A. J., Maddy, E. S., and
Hoffman, R. N.: Outlook for Exploiting Artificial Intelligence in the Earth and Environmental Sciences, Bulletin of the
American Meteorological Society, 102, E1016-E1032, <https://doi.org/10.1175/BAMS-D-20-0031.1>, 2021.

650 Buck, J. J. H., Bainbridge, S. J., Burger, E. F., Kraberg, A. C., Casari, M., Casey, K. S., Darroch, L., del Rio, J., Metfies, K.,
Delory, E., Fischer, P. F., Gardner, T., Heffernan, R., Jirka, S., Kokkinaki, A., Loebel, M., Buttigieg, P. L., Pearlman, J. S.,
and Schewe, I.: Ocean Data Product Integration Through Innovation-The Next Level of Data Interoperability, Frontiers in
Marine Science, 6, [10.3389/fmars.2019.00032](https://doi.org/10.3389/fmars.2019.00032), 2019.

655 Buisán, S. T., Smith, C. D., Ross, A., Kochendorfer, J., Collado, J. L., Alastrué, J., Wolff, M., Roulet, Y.-A., Earle, M. E.,
Laine, T., Rasmussen, R., and Nitu, R.: The potential for uncertainty in Numerical Weather Prediction model verification
when using solid precipitation observations, Atmospheric Science Letters, 21, e976, <https://doi.org/10.1002/asl.976>, 2020.



660 Casati, B., Robinson, T., Lemay, F., Køltzow, M., Haiden, T., Mekis, E., Lespinas, F., Fortin, V., Gascon, G., Milbrandt, J.,
and Smith, G.: Performance of the Canadian Arctic Prediction System during the YOPP Special Observing Periods,
Atmosphere-Ocean, 1-27, <https://doi.org/10.1080/07055900.2023.2191831>, 2023.

CF Standard Name Table: <http://cfconventions.org/Data/cf-standard-names/current/build/cf-standard-name-table.html>, last
665 access: 18 July 2023.

Clothiaux, E. E., Miller, M. A., Perez, R. C., Turner, D. D., Moran, K. P., Martner, B. E., Ackerman, T. P., Mace, G. G.,
Marchand, R. T., Widener, K. B., Rodriguez, D. J., Uttal, T., Mather, J. H., Flynn, C. J., Gaustad, K. L., and Ermold, B.: The
ARM Millimeter Wave Cloud Radars (MMCRs) and the Active Remote Sensing of Clouds (ARSCL) Value Added Product
670 (VAP), ARM user facility, Pacific Northwest National Laboratory, Richland, WA, United States, 56 pp.,
<https://doi.org/10.2172/1808567>, 2001.

Cox, C. J., Morris, S. M., Uttal, T., Burgener, R., Hall, E., Kutchenreiter, M., McComiskey, A., Long, C. N., Thomas, B. D.,
and Wendell, J.: The De-Icing Comparison Experiment (D-ICE): a study of broadband radiometric measurements under
675 icing conditions in the Arctic, *Atmos. Meas. Tech.*, 14, 1205-1224, <https://doi.org/10.5194/amt-14-1205-2021>, 2021.

Cox, C. J., Gallagher, M., Shupe, M. D., Persson, P. O. G., Solomon, A., Fairall, C. W., Ayers, T., Blomquist, B., Brooks, I.
M., Costa, D., Grachev, A., Gottas, D., Hutchings, J. K., Kutchenreiter, M., J. Leach, J., Morris, S. M., Morris, V., Osborn,
J., Pezoa, S., Preusser, A., Riihimaki, L., and Uttal, T.: Continuous observations of the surface energy budget and
680 meteorology over the Arctic sea ice during MOSAiC, *Scientific Data*, <https://doi.org/10.1038/s41597-023-02415-5>, 2023.

Data Citation Synthesis Group: Joint Declaration of Data Citation Principles, FORCE11, San Diego CA,
<https://doi.org/10.25490/a97f-egykh>, 2014.

685 DataCite Metadata Working Group: DataCite Metadata Schema Documentation for the Publication and Citation of Research
Data and Other Research Outputs. Version 4.4, DataCite e.V., 82 pp., <https://doi.org/10.14454/3w3z-sa82>, 2021.

Day, J. J., Svensson, G., Casati, B., Uttal, T., Khalsa, S.-J., Bazile, E., Akish, E., Azouz, N., Ferrighi, L., Frank, H.,
Gallagher, M., Godøy, Ø., Hartten, L. M., Huang, L. X., Holt, J., Di Stefano, M., Suomi, I., Mariani, Z., Morris, S.,
690 O'Connor, E., Pirazzini, R., Remes, T., Fadeev, R., Solomon, A., Tjernström, J., and Tolstykh, M.: The YOPP Site Model
Intercomparison Project (YOPPsiteMIP) Phase 1: Project Overview and Arctic Winter Forecast Evaluation, Geoscientific
Model Development, https://editor.copernicus.org/EGUsphere/ms_records/egusphere-2023-1550, submitted.



- Eaton, B., Gregory, J., Drach, B., Taylor, K., Hankin, S., Blower, J., Caron, J., Signell, R., Bentley, P., Rappa, G., Höck, H.,
695 Pamment, A., Juckes, M., Raspaud, M., Horne, R., Whiteaker, T., Blodgett, D., Zender, C., Lee, D., Hassell, D., Snow, A.
D., Kölling, T., Allured, D., Jelenak, A., Soerensen, A. M., Gaultier, L., and Herlédan, S.: NetCDF Climate and Forecast
(CF) Metadata Conventions Version 1.10, <http://cfconventions.org/Data/cf-conventions/cf-conventions-1.10/cf-conventions.html>, 2022.
- 700 Essential Climate Variables: <https://public.wmo.int/en/programmes/global-climate-observing-system/essential-climate-variables>, last access: 13 September 2023.
- Fratini, G. and Mauder, M.: Towards a consistent eddy-covariance processing: an intercomparison of EddyPro and TK3,
Atmos. Meas. Tech., 7, 2273-2281, 10.5194/amt-7-2273-2014, 2014.
- 705 Gettelman, A., Geer, A. J., Forbes, R. M., Carmichael, G. R., Feingold, G., Posselt, D. J., Stephens, G. L., van den Heever, S. C., Varble, A. C., and Zuidema, P.: The future of Earth system prediction: Advances in model-data fusion, Science Advances, 8, eabn3488, doi:10.1126/sciadv.abn3488, 2022.
- 710 Global Telecommunication System (GTS): <https://community.wmo.int/en/activity-areas/global-telecommunication-system-gts>, last access: 16 July 2023.
- Goessling, H. F., Jung, T., Klebe, S., Baeseman, J., Bauer, P., Chen, P., Chevallier, M., Dole, R., Gordon, N., Ruti, P., Bradley, A., Bromwich, D. H., Casati, B., Chechin, D., Day, J. J., Massonnet, F., Mills, B., Renfrew, I. A., Smith, G., and
715 Tatusko, R.: Paving the Way for the Year of Polar Prediction, Bulletin of the American Meteorological Society, 97, ES85-ES88, <https://doi.org/10.1175/BAMS-D-15-00270.1>, 2016.
- Grachev, A. A., Persson, P. O. G., Uttal, T., Akish, E. A., Cox, C. J., Morris, S. M., Fairall, C. W., Stone, R. S., Lesins, G., Makshatas, A. P., and Repina, I. A.: Seasonal and latitudinal variations of surface fluxes at two Arctic terrestrial sites, Climate
720 Dynamics, 51, 1793-1818, 10.1007/s00382-017-3983-4, 2018.
- Hanisch, R., Chalk, S., Coulon, R., Cox, S., Emmerson, S., Sandoval, F. J. F., Forbes, A., Frey, J., Hall, B., Hartshorn, R., Heus, P., Hodson, S., Hosaka, K., Hutzschenreuter, D., Kang, C.-S., Picard, S., and White, R.: Stop squandering data: make units of measurement machine-readable, Nature, 605, 222-224, <https://doi.org/10.1038/d41586-022-01233-w>, 2022.
- 725 Hartten, L. M. and Khalsa, S. J. S.: The H-K Variable SchemaTable developed for the YOPPsiteMIP, Zenodo.com, <https://doi.org/10.5281/zenodo.6255666>, 2022.



Hassell, D., Gregory, J., Blower, J., Lawrence, B. N., and Taylor, K. E.: A data model of the Climate and Forecast metadata
730 conventions (CF-1.6) with a software implementation (cf-python v2.1), *Geosci. Model Dev.*, 10, 4619-4646, 10.5194/gmd-
10-4619-2017, 2017.

Hogan, R. J. and O'Connor, E. J.: Facilitating cloud radar and lidar algorithms: the Cloudnet Instrument Synergy/Target
735 Categorization product, 14 pp., <http://www.met.rdg.ac.uk/~swrhgnrj/publications/categorization.pdf>, 2004.

Holloway, C. E., Petch, J. C., Beare, R. J., Bechtold, P., Craig, G. C., Derbyshire, S. H., Donner, L. J., Field, P. R., Gray, S.
L., Marsham, J. H., Parker, D. J., Plant, R. S., Roberts, N. M., Schultz, D. M., Stirling, A. J., and Woolnough, S. J.:
Understanding and representing atmospheric convection across scales: recommendations from the meeting held at
Dartington Hall, Devon, UK, 28–30 January 2013, *Atmospheric Science Letters*, 15, 348-353,
740 <https://doi.org/10.1002/asl2.508>, 2014.

Holtstlag, A. A. M., Svensson, G., Baas, P., Basu, S., Beare, B., Beljaars, A. C. M., Bosveld, F. C., Cuxart, J., Lindvall, J.,
Steenefeld, G. J., Tjernström, M., and Van De Wiel, B. J. H.: Stable Atmospheric Boundary Layers and Diurnal Cycles:
Challenges for Weather and Climate Models, *Bulletin of the American Meteorological Society*, 94, 1691-1706,
745 <https://doi.org/10.1175/BAMS-D-11-00187.1>, 2013.

Illingworth, A. J., Hogan, R. J., O'Connor, E. J., Bouniol, D., Brooks, M. E., Delanoé, J., Donovan, D. P., Eastment, J. D.,
Gaussiat, N., Goddard, J. W. F., Haeffelin, M., Baltink, H. K., Krasnov, O. A., Pelon, J., Piriou, J.-M., Protat, A.,
Russchenberg, H. W. J., Seifert, A., Tompkins, A. M., van Zadelhoff, G.-J., Vinit, F., Willén, U., Wilson, D. R., and
750 Wrench, C. L.: Cloudnet: Continuous Evaluation of Cloud Profiles in Seven Operational Models Using Ground-Based
Observations, *Bulletin of the American Meteorological Society*, 88, 883-898, <https://doi.org/10.1175/BAMS-88-6-883>,
2007.

Jones, M. B., Budden, A. E., Mecum, B., Clark, J., Brun, J., Lowndes, J., and McLean, E.: Data Science Training for Arctic
755 Researchers, Arctic Data Center, <https://doi.org/10.18739/A24746R2N>, 2020.

Jung, T., Gordon, N. D., Bauer, P., Bromwich, D. H., Chevallier, M., Day, J. J., Dawson, J., Doblas-Reyes, F. J., Fairall, C.,
Goessling, H. F., Holland, M., Inoue, J., Iversen, T., Klebe, S., Lemke, P., Losch, M., Makshtas, A., Mills, B., Nurmi, P.,
Perovich, D., Reid, P., Renfrew, I. A., Smith, G., Svensson, G., Tolstykh, M., and Yang, Q.: Advancing Polar Prediction
760 Capabilities on Daily to Seasonal Time Scales, *Bulletin of the American Meteorological Society*, 97, 1631-1647,
<https://doi.org/10.1175/BAMS-D-14-00246.1>, 2016.



Kaimal, J. C. and Finnigan, J. J.: Atmospheric Boundary Layer Flows: Their Structure and Measurement, Oxford University Press, New York, ISBN 9780195062397, 1994.

765

Kochendorfer, J., Nitu, R., Wolff, M., Mekis, E., Rasmussen, R., Baker, B., Earle, M. E., Reverdin, A., Wong, K., Smith, C. D., Yang, D., Roulet, Y. A., Meyers, T., Buisan, S., Isaksen, K., Brækkan, R., Landolt, S., and Jachcik, A.: Testing and development of transfer functions for weighing precipitation gauges in WMO-SPICE, *Hydrol. Earth Syst. Sci.*, 22, 1437-1452, <https://doi.org/10.5194/hess-22-1437-2018>, 2018.

770

Kochendorfer, J., Earle, M., Rasmussen, R., Smith, C., Yang, D., Morin, S., Mekis, E., Buisan, S., Roulet, Y.-A., Landolt, S., Wolff, M., Hoover, J., Thériault, J. M., Lee, G., Baker, B., Nitu, R., Lanza, L., Colli, M., and Meyers, T.: How Well Are We Measuring Snow Post-SPICE?, *Bulletin of the American Meteorological Society*, 103, E370-E388, <https://doi.org/10.1175/BAMS-D-20-0228.1>, 2022.

775

Køltzow, M., Casati, B., Haiden, T., and Valkonen, T.: Verification of Solid Precipitation Forecasts from Numerical Weather Prediction Models in Norway, *Weather and Forecasting*, 35, 2279-2292, <https://doi.org/10.1175/WAF-D-20-0060.1>, 2020.

780

Lavergne, T., Kern, S., Aaboe, S., Derby, L., Dybkjaer, G., Garric, G., Heil, P., Hendricks, S., Holfort, J., Howell, S., Key, J., Lieser, J. L., Maksym, T., Maslowski, W., Meier, W., Muñoz-Sabater, J., Nicolas, J., Özsoy, B., Rabe, B., Rack, W., Raphael, M., de Rosnay, P., Smolyanitsky, V., Tietsche, S., Ukita, J., Vichi, M., Wagner, P., Willmes, S., and Zhao, X.: A New Structure for the Sea Ice Essential Climate Variables of the Global Climate Observing System, *Bulletin of the American Meteorological Society*, 103, E1502-E1521, <https://doi.org/10.1175/BAMS-D-21-0227.1>, 2022.

785

Long, C. N. and Shi, Y.: The QCRad Value Added Product: Surface Radiation Measurement Quality Control Testing, Including Climatology Configurable Limits, PNNL, Richland, Washington, United States, 70 pp., <https://doi.org/10.2172/1019540>, 2006.

790

Long, C. N. and Shi, Y.: An Automated Quality Assessment and Control Algorithm for Surface Radiation Measurements, 2, 23-37, <http://dx.doi.org/10.2174/1874282300802010023>, 2008.

Mahrt, L. T. and Sun, J.: The Subgrid Velocity Scale in the Bulk Aerodynamic Relationship for Spatially Averaged Scalar Fluxes, *Monthly Weather Review*, 123, 3032-3041, [https://doi.org/10.1175/1520-0493\(1995\)123<3032:TSVSIT>2.0.CO;2](https://doi.org/10.1175/1520-0493(1995)123<3032:TSVSIT>2.0.CO;2),
795 1995.



Matrosov, S. Y., Shupe, M. D., and Uttal, T.: High temporal resolution estimates of Arctic snowfall rates emphasizing gauge and radar-based retrievals from the MOSAiC expedition, *Elementa: Science of the Anthropocene*, 10, <https://doi.org/10.1525/elementa.2021.00101>, 2022.

800

Matsui, N., Long, C. N., Augustine, J., Halliwell, D., Uttal, T., Longenecker, D., Niebergall, O., Wendell, J., and Albee, R.: Evaluation of Arctic broadband surface radiation measurements, *Atmos. Meas. Tech.*, 5, 429-438, [10.5194/amt-5-429-2012](https://doi.org/10.5194/amt-5-429-2012), 2012.

805 Mauder, M., Foken, T., Clement, R., Elbers, J. A., Eugster, W., Grünwald, T., Heusinkveld, B., and Kolle, O.: Quality control of CarboEurope flux data – Part 2: Inter-comparison of eddy-covariance software, *Biogeosciences*, 5, 451-462, [10.5194/bg-5-451-2008](https://doi.org/10.5194/bg-5-451-2008), 2008.

810 Monin, A. S. and Obukhov, A.: Basic laws of turbulent mixing in the surface layer of the atmosphere, *Contrib. Geophys. Inst. Acad. Sci. USSR*, 151, e187, 1954.

Morris, S. and Akish, E.: A-M Variable & Attribute Template Table developed for the YOPPsiteMIP, *Zenodo.com*, <https://doi.org/10.5281/zenodo.6780400>, 2022.

815 Morris, S. and Uttal, T.: Datagrams: Diagrammatic Metadata for Humans, *Bulletin of the American Meteorological Society*, 103, E1343-E1350, <https://doi.org/10.1175/BAMS-D-21-0219.1>, 2022.

820 Morris, S. M., Mariani, Z., Uttal, T., Akish, E., Crawford, R., Huang, L., Day, J., Tjernström, J., Godøy, Ø., Ferrighi, L., Hartten, L. M., Holt, J., Cox, C. J., O'Connor, E., Pirazzini, R., Maturilli, M., Prakash, G., Mather, J., Strong, K., Fogal, P., Kustov, V., Svensson, G., Gallagher, M., and Vasel, B.: Special Observing Period (SOP) Data for the Year of Polar Prediction site Model Intercomparison Project (YOPPsiteMIP), *Earth System Science Data*, https://editor.copernicus.org/ESSD/ms_records/essd-2023-279, submitted.

825 Nature Editorial : Time to recognize authorship of open data, *Nature*, 608, 8, <https://doi.org/10.1038/d41586-022-00921-x>, 2022.

Neang, A. B., Sutherland, W., Beach, M. W., and Lee, C. P.: Data Integration as Coordination: The Articulation of Data Work in an Ocean Science Collaboration, *Proc. ACM Hum.-Comput. Interact.*, 4, Article 256, <https://doi.org/10.1145/3432955>, 2021.



830

Nicolaus, M., Perovich, D. K., Spreen, G., Granskog, M. A., von Albedyll, L., Angelopoulos, M., Anhaus, P., Arndt, S., Belter, H. J., Bessonov, V., Birnbaum, G., Brauchle, J., Calmer, R., Cardellach, E., Cheng, B., Clemens-Sewall, D., Dadic, R., Damm, E., de Boer, G., Demir, O., Dethloff, K., Divine, D. V., Fong, A. A., Fons, S., Frey, M. M., Fuchs, N., Gabarró, C., Gerland, S., Goessling, H. F., Gradinger, R., Haapala, J., Haas, C., Hamilton, J., Hannula, H.-R., Hendricks, S., Herber, A., Heuzé, C., Hoppmann, M., Høyland, K. V., Huntemann, M., Hutchings, J. K., Hwang, B., Itkin, P., Jacobi, H.-W., Jaggi, M., Jutila, A., Kaleschke, L., Katlein, C., Kolabutin, N., Krampe, D., Kristensen, S. S., Krumpfen, T., Kurtz, N., Lampert, A., Lange, B. A., Lei, R., Light, B., Linhardt, F., Liston, G. E., Loose, B., Macfarlane, A. R., Mahmud, M., Matero, I. O., Maus, S., Morgenstern, A., Naderpour, R., Nandan, V., Niubom, A., Oggier, M., Oppelt, N., Pätzold, F., Perron, C., Petrovsky, T., Pirazzini, R., Polashenski, C., Rabe, B., Raphael, I. A., Regnery, J., Rex, M., Ricker, R., Riemann-Campe, K., Rinke, A., Rohde, J., Salganik, E., Scharien, R. K., Schiller, M., Schneebeli, M., Semmling, M., Shimanchuk, E., Shupe, M. D., Smith, M. M., Smolyanitsky, V., Sokolov, V., Stanton, T., Stroeve, J., Thielke, L., Timofeeva, A., Tonboe, R. T., Tavri, A., Tsamados, M., Wagner, D. N., Watkins, D., Webster, M., and Wendisch, M.: Overview of the MOSAiC expedition: Snow and sea ice, *Elementa: Science of the Anthropocene*, 10, <https://doi.org/10.1525/elementa.2021.000046>, 2022.

845 Nitu, R., Roulet, Y.-A., Wolff, M., Earle, M., Reverdin, A., Smith, C., Kochendorfer, J., Morin, S., Rasmussen, R., Wong, K., Alastrué, J., Arnold, L., Baker, B., Buisán, S., Collado, J. L., Colli, M., Collins, B., Gaydos, A., Hannula, H.-R., Hoover, J., Joe, P., Kontu, A., Laine, T., Lanza, L., Lanzinger, E., Lee, G., Lejeune, Y., Leppänen, L., Mekis, E., Panel, J.-M., Poikonen, A., Ryu, S., Sabatini, F., Theriault, J., Yang, D., Genthon, C., van den Heuvel, F., Hirasawa, N., Konishi, H., Motoyoshi, H., Nakai, S., Nishimura, K., Senese, A., and Yamashita, K.: WMO Solid Precipitation Intercomparison Experiment (SPICE) (2012 - 2015), World Meteorological Organization (WMO), Geneva, Switzerland, IOM No. 1, 1443 pp., <https://library.wmo.int/opac/>, 2018.

855 Ohmura, A., Dutton, E. G., Forgan, B., Fröhlich, C., Gilgen, H., Hegner, H., Heimo, A., König-Langlo, G., McArthur, B., Müller, G., Philipona, R., Pinker, R., Whitlock, C. H., Dehne, K., and Wild, M.: Baseline Surface Radiation Network (BSRN/WCRP): New Precision Radiometry for Climate Research, *Bulletin of the American Meteorological Society*, 79, 2115-2136, [https://doi.org/10.1175/1520-0477\(1998\)079<2115:BSRNBW>2.0.CO;2](https://doi.org/10.1175/1520-0477(1998)079<2115:BSRNBW>2.0.CO;2), 1998.

Papoutsoglou, E. A., Athanasiadis, I. N., Visser, R. G. F., and Finkers, R.: The benefits and struggles of FAIR data: the case of reusing plant phenotyping data, *Scientific Data*, 10, 457, <https://doi.org/10.1038/s41597-023-02364-z>, 2023.

860

Pierce, H. H., Dev, A., Statham, E., and Bierer, B. E.: Credit data generators for data reuse, *Nature*, 570, 30-32, <https://doi.org/10.1038/d41586-019-01715-4>, 2019.



865 PPP Steering Group & Co-authors: WWRP Polar Prediction Project Implementation Plan for the Year of Polar Prediction (YOPP), 80 pp., <https://www.polarprediction.net/about/implementation-and-science-plans/>, 2019.

Prakash, G., Shrestha, B., Younkin, K., Jundt, R., Martin, M., and Elliott, J.: Data Always Getting Bigger—A Scalable DOI Architecture for Big and Expanding Scientific Data, *Data*, 1, 11, <https://www.mdpi.com/2306-5729/1/2/11>, 2016.

870 Sardeshmukh, P. D., Compo, G. P., and Penland, C.: Need for Caution in Interpreting Extreme Weather Statistics, *Journal of Climate*, 28, 9166-9187, <https://doi.org/10.1175/JCLI-D-15-0020.1>, 2015.

875 Shupe, M. D., Rex, M., Blomquist, B., Persson, P. O. G., Schmale, J., Uttal, T., Althausen, D., Angot, H., Archer, S., Bariteau, L., Beck, I., Bilberry, J., Bucci, S., Buck, C., Boyer, M., Brasseur, Z., Brooks, I. M., Calmer, R., Cassano, J., Castro, V., Chu, D., Costa, D., Cox, C. J., Creamean, J., Crewell, S., Dahlke, S., Damm, E., de Boer, G., Deckelmann, H., Dethloff, K., Dütsch, M., Ebell, K., Ehrlich, A., Ellis, J., Engelmann, R., Fong, A. A., Frey, M. M., Gallagher, M. R., Ganzeveld, L., Gradinger, R., Graeser, J., Greenamyer, V., Griesche, H., Griffiths, S., Hamilton, J., Heinemann, G., Helmig, D., Herber, A., Heuzé, C., Hofer, J., Houchens, T., Howard, D., Inoue, J., Jacobi, H.-W., Jaiser, R., Jokinen, T., Jourdan, O., Jozef, G., King, W., Kirchgassner, A., Klingebiel, M., Krassovski, M., Krumpfen, T., Lampert, A., Landing, W., Laurila, T., 880 Lawrence, D., Lonardi, M., Loose, B., Lüpkes, C., Maahn, M., Macke, A., Maslowski, W., Marsay, C., Maturilli, M., Mech, M., Morris, S., Moser, M., Nicolaus, M., Ortega, P., Osborn, J., Pätzold, F., Perovich, D. K., Petäjä, T., Pilz, C., Pirazzini, R., Posman, K., Powers, H., Pratt, K. A., Preußner, A., Quéléver, L., Radenz, M., Rabe, B., Rinke, A., Sachs, T., Schulz, A., Siebert, H., Silva, T., Solomon, A., Sommerfeld, A., Spreen, G., Stephens, M., Stohl, A., Svensson, G., Uin, J., Viegas, J., Voigt, C., von der Gathen, P., Wehner, B., Welker, J. M., Wendisch, M., Werner, M., Xie, Z. Q., and Yue, F.: Overview of 885 the MOSAiC expedition: Atmosphere, *Elementa: Science of the Anthropocene*, 10, 10.1525/elementa.2021.00060, 2022.

Sprintall, J., Coles, V. J., Reed, K. A., Butler, A. H., Foltz, G. R., Penny, S. G., and Seo, H.: Using Process Studies to Improve Climate Modeling: Strategies for Success, *Bulletin of the American Meteorological Society*, 102, 523-526, <https://doi.org/10.1175/BAMS-D-19-0263.A>, 2021.

890

Stephens, G. L., Polcher, J., Zeng, X., van Oevelen, P., Poveda, G., Bosilovich, M., Ahn, M.-H., Balsamo, G., Duan, Q., Hegerl, G. C., Jakob, C., Lamptey, B., Leung, R., Piles, M., Su, Z., Dirmeyer, P., Findell, K. L., Verhoef, A., Ek, M., L'Ecuyer, T., Roca, R., Nazemi, A., Dominguez, F., Klocke, D., and Bony, S.: The First 30 Years of GEWEX, *Bulletin of the American Meteorological Society*, 104, E126-E157, <https://doi.org/10.1175/BAMS-D-22-0061.1>, 2023.

895



Stokes, G. M. and Schwartz, S. E.: The Atmospheric Radiation Measurement (ARM) Program: Programmatic Background and Design of the Cloud and Radiation Test Bed, *Bulletin of the American Meteorological Society*, 75, 1201-1222, [https://doi.org/10.1175/1520-0477\(1994\)075<1201:TARMPP>2.0.CO;2](https://doi.org/10.1175/1520-0477(1994)075<1201:TARMPP>2.0.CO;2), 1994.

900 Svensson, G., Casati, B., Day, J., Uttal, T., Godøy, Ø., and Hartten, L.: YOPPsiteMIP – The YOPP site Model Inter-comparison Project, Alfred-Wegener-Institut, Bremerhaven, 15 pp., https://www.polarprediction.net/fileadmin/user_upload/www.polarprediction.net/Home/Organization/Task_Teams/Atmospheric_Processes/YOPP_Supersite_common_model_output_rev8.pdf, 2020.

905 Taylor, K. E., Durack, P. J., Elkington, M., Guilyardi, E., Hassell, D., Lautenschlager, M., and Stockhause, M.: CMIP6 Participation Guidance for Modelers, <https://pcmdi.llnl.gov/CMIP6/Guide/modelers.html>, 2022.

Unidata: NetCDF User's Guide, version 1.1, <https://doi.org/10.26024/nw73-vm64>, 2023.

910 Unidata: UDUNITS-2, version 2.2.28 (2.2.28), UCAR/Unidata Program Center [software], <https://doi.org/10.5065/D6KD1WN0>, 2020.

Uttal, T., Starkweather, S., Drummond, J. R., Vihma, T., Makshtas, A. P., Darby, L. S., Burkhart, J. F., Cox, C. J., Schmeisser, L. N., Haiden, T., Maturilli, M., Shupe, M. D., De Boer, G., Saha, A., Grachev, A. A., Crepinsek, S. M.,
915 Bruhwiler, L., Goodison, B., McArthur, B., Walden, V. P., Dlugokencky, E. J., Persson, P. O. G., Lesins, G., Laurila, T., Ogren, J. A., Stone, R. S., Long, C. N., Sharma, S., Massling, A., Turner, D. D., Stanitski, D. M., Asmi, E., Aurela, M., Skov, H., Eleftheriadis, K., Virkkula, A., Platt, A., Førland, E. J., Iijima, Y., Nielsen, I. E., Bergin, M. H., Candlish, L., Zimov, N. S., Zimov, S. A., O'Neill, N. T., Fogal, F., Kivi, R., Konopleva-Akish, E. A., Verlinde, J., Kustov, V. Y., Vasel, B., Ivakhov, V. M., Viisanen, Y., and Intrieri, J. M.: International Arctic Systems for Observing the Atmosphere: An
920 International Polar Year Legacy Consortium, *Bulletin of the American Meteorological Society*, 97, 1033-1056, <https://doi.org/10.1175/BAMS-D-14-00145.1>, 2016.

Vannan, S., Downs, R. R., Meier, W., Wilson, B. E., and Gerasimov, I. V.: Data sets are foundational to research. Why don't we cite them?, *Eos*, 101, <https://doi.org/10.1029/2020EO151665>, 2020.

925

Vorosmarty, C., Rawlins, M., Hinzman, L., Francis, J., Serreze, M., Liljedahl, A., McDonald, K., Piasecki, M., and Rich, R.: Opportunities and Challenges in Arctic System Synthesis: A Consensus Report from the Arctic Research Community, New York, 84 pp., <https://www.arcus.org/publications/28459>, 2018.



- 930 Wei, Y., Shrestha, R., Pal, S., Gerken, T., Feng, S., McNelis, J., Singh, D., Thornton, M. M., Boyer, A. G., Shook, M. A., Chen, G., Baier, B. C., Barkley, Z. R., Barrick, J. D., Bennett, J. R., Browell, E. V., Campbell, J. F., Campbell, L. J., Choi, Y., Collins, J., Dobler, J., Eckl, M., Fiehn, A., Fried, A., Digangi, J. P., Barton-Grimley, R., Halliday, H., Klausner, T., Kooi, S., Kostinek, J., Lauvaux, T., Lin, B., McGill, M. J., Meadows, B., Miles, N. L., Nehrir, A. R., Nowak, J. B., Obland, M., O'Dell, C., Fao, R. M. P., Richardson, S. J., Richter, D., Roiger, A., Sweeney, C., Walega, J., Weibring, P., Williams, C. A.,
- 935 Yang, M. M., Zhou, Y., and Davis, K. J.: Atmospheric Carbon and Transport – America (ACT-America) Data Sets: Description, Management, and Delivery, *Earth and Space Science*, 8, e2020EA001634, <https://doi.org/10.1029/2020EA001634>, 2021.

- Wilkinson, M. D., Dumontier, M., Aalbersberg, I. J., Appleton, G., Axton, M., Baak, A., Blomberg, N., Boiten, J.-W., da
- 940 Silva Santos, L. B., Bourne, P. E., Bouwman, J., Brookes, A. J., Clark, T., Crosas, M., Dillo, I., Dumon, O., Edmunds, S., Evelo, C. T., Finkers, R., Gonzalez-Beltran, A., Gray, A. J. G., Groth, P., Goble, C., Grethe, J. S., Heringa, J., 't Hoen, P. A. C., Hooft, R., Kuhn, T., Kok, R., Kok, J., Lusher, S. J., Martone, M. E., Mons, A., Packer, A. L., Persson, B., Rocca-Serra, P., Roos, M., van Schaik, R., Sansone, S.-A., Schultes, E., Sengstag, T., Slater, T., Strawn, G., Swertz, M. A., Thompson, M., van der Lei, J., van Mulligen, E., Velterop, J., Waagmeester, A., Wittenburg, P., Wolstencroft, K., Zhao, J., and Mons,
- 945 B.: The FAIR Guiding Principles for scientific data management and stewardship, *Scientific Data*, 3, 160018, [10.1038/sdata.2016.18](https://doi.org/10.1038/sdata.2016.18), 2016.

- Wilson, J., Jung, T., Bazile, E., Bromwich, D., Casati, B., Day, J., De Coning, E., Eayrs, C., Grumbine, R., Ioue, J., Khalsa, S. J. S., Kristiansen, J., Lamers, M., Liggett, D., Olsen, S. M., Perovich, D., Renfrew, I. A., Smolyanitsky, V., Svensson, G.,
- 950 Sun, Q., Uttal, T., and Yang, Q.: The YOPP Final Summit: Assessing Past and Forecasting Future Polar Prediction Research, *Bulletin of the American Meteorological Society*, 104, E660-E665, <https://doi.org/10.1175/BAMS-D-22-0282.1>, 2023.

- Wolff, M. A., Isaksen, K., Petersen-Øverleir, A., Ødemark, K., Reitan, T., and Brækkan, R.: Derivation of a new continuous adjustment function for correcting wind-induced loss of solid precipitation: results of a Norwegian field study, *Hydrol. Earth*
- 955 *Syst. Sci.*, 19, 951-967, <https://doi.org/10.5194/hess-19-951-2015>, 2015.

- World Meteorological Organization (WMO): Manual on the Global Telecommunication System: Annex III to the WMO Technical Regulations, 2015, WMO (Series), no. 386, Secretariat of the World Meteorological Organization, Geneva, Switzerland, 197 pp., https://library.wmo.int/doc_num.php?explnum_id=10469, ISBN 978-92-63-10386-4, 2020.

960

- Xie, S., McCoy, R. B., Klein, S. A., Cederwall, T., Wiscombe, W. J., Jensen, M. P., Johnson, K. L., Clothiaux, E. E., Gaustad, K. L., Long, C. N., Mather, J. H., McFarlane, S. A., Shi, Y., Golaz, J.-C., Lin, Y., Hall, S. D., McCord, R. A., Palanisamy, G., and Turner, D. D.: CLOUDS AND MORE: ARM Climate Modeling Best Estimate Data: A New Data

<https://doi.org/10.5194/egusphere-2023-2413>
Preprint. Discussion started: 5 December 2023
© Author(s) 2023. CC BY 4.0 License.



Product for Climate Studies, Bulletin of the American Meteorological Society, 91, 13-20,
965 <https://doi.org/10.1175/2009BAMS2891.1>, 2010.

Zuo, G., Dou, Y., and Lei, R.: Discrimination Algorithm and Procedure of Snow Depth and Sea Ice Thickness Determination
Using Measurements of the Vertical Ice Temperature Profile by the Ice-Tethered Buoys, Sensors, 18, 4162,
<https://www.mdpi.com/1424-8220/18/12/4162>, 2018.

970