

A machine learning approach to the geomorphometric detection of ribbed moraines in Norway

Thomas J. Barnes¹, Thomas V. Schuler¹, Simon Filhol¹, Karianne S. Lilleøren¹

¹Department of Geosciences, University of Oslo, 0316, Oslo, Norway

5 *Correspondence to:* Thomas J. Barnes (thomas.barnes@geo.uio.no)

Abstract.

Machine learning is a powerful yet underutilised tool in geomorphology, commonly used for image-based pattern recognition. Analysing new high-resolution (1 – 10 m) elevation datasets, we investigate its usefulness for detecting discrete geomorphological features. This study develops a machine learning-based method for identifying ribbed moraines in digital elevation data and progresses to test its performance versus time consuming, manual methods. Ribbed moraines share geomorphometric characteristics with other glacial landforms, hence represent a valuable test of our new methodology in terms of differentiating between similar features, and wider for detection of landforms with similar characteristics. Furthermore, mapping ribbed moraines may provide valuable indications of their origin, a topic of debate within glacial geomorphology. To automatically detect ribbed moraines, we extract simple morphometrics from high-resolution digital elevation model data and mask regions where ribbed moraines are unlikely to form. We then test several machine learning algorithms before examining the best performer (K-means clustering) on three study areas in Norway of 15 km². Our results demonstrate balanced accuracy of 65 – 75 % when validating versus ground-truth. The performance depends on the availability of high-resolution elevation data in Norway, needed to resolve the spatial scale of the target (10-100m). We find the method effective at detecting both fields of ribbed moraines as well as individual ribbed moraines. We propose pathways for future implementation of this method on a large-scale and for increasing the detail of information gained about detected landforms. In conclusion, we demonstrate K-means clustering as a promising method for detecting ribbed moraines, with great potential to reduce the time needed to produce landform maps.

10
15
20
25
30
35

1 Introduction

1.1 Geomorphological mapping

Mapping of landforms has traditionally been a manual process, either through direct field observations or manual analysis of remotely sensed data (Smith & Clark, 2005; Verstappen, 2011; Evans, 2012; Sommerkorn, 2020). More recently, semi-automated methods have been developed, where computational analysis of remote sensing data is interpreted by the operator (Guitet et al 2013), giving rise to subjectivity and human error (Saha et al., 2011; Eisank et al., 2014; Sommerkorn, 2020). Often, the restriction of data availability, quality and resolution have been the primary limitations leading to the maintenance of traditional approaches, as the resolution of digital elevation models (DEM) has typically been limited to 30–300 m (Saha et al., 2011; Iwahashi et al., 2018) thus inhibiting the detection of metre-scale features. Additionally, until recently, sub-10 m datasets have been afflicted with patchy coverage (UKEA, 2023), limiting large-scale high-resolution digital mapping of smaller (sub-data resolution) landforms.

1.2 Machine learning as a solution

In recent years, automated landform mapping has taken a machine learning-based pattern recognition approach, starting with a DEM and extracting morphometrics from input data, such as slope, aspect, convexity and surface texture (Clubb et al., 2019; Eisank et al., 2014; Saha et al., 2011; Iwahashi et al., 2018). Thus far, machine-learning approaches focused on large-scale geomorphological feature detection, such as sedimentary basins or terrain classification due to resolution restrictions (Kong et al., 2020; Iwahashi et al., 2018). Yet, the increasing availability of high-resolution (<10 m) datasets (e.g. Kartverket, 2021; UKEA, 2023) enable individual feature mapping for large areas. Norway has a national high-resolution (0.6 - 1 m) DEM and orthophotography data coverage (Kartverket, 2021). In conjunction with this, the development of new and more robust machine learning methods, coupled with the effectiveness of older methods with new, high-resolution data (clustering and segmentation, Gentleman & Carey, 2008) suggest new approaches for small-scale landform detection are now possible.


As machine-learning is both fast, low on labour intensiveness once set-up and minimises human error (Gentleman & Carey, 2008), it is clear that it may be possible to address prior limitations to landform mapping. However, as high-resolution data availability is a recent development, only few studies, for example Corr et al. (2022), used a supervised random forest (RF) algorithm to detect supraglacial lakes in Antarctica. Aydda et al. (2019) detected dune forms using three different unsupervised machine learning algorithms, including K-means clustering (KM). Here, we propose combining new, high-resolution data and machine-learning, to overcome previous limitations in geomorphological mapping.

1.3 Study aims

In this study, we develop a machine-learning algorithm to detect specific small-scale geomorphological landforms in high-resolution DEM data, more specifically ribbed moraine. They usually have horizontal extents at a 10-100 m scale and are common in Norway (Dunlop & Clark, 2006; Dunlop et al., 2008; Hättestrand & Kleman, 1999; Finlayson & Bradwell, 2008). Furthermore, they typically form in shallow depressions in inland regions close to the former Fennoscandian ice divide (Sollid & Sørbel, 1994; Sarala, 2006; Fredin et al., 2013; Sommerkorn, 2020, Patton et al., 2016; 2017; Butcher, 2022). With this information, we can define potential study regions, and validation dataset. Ribbed moraines are subglacial ridges transverse to the glacial flow direction (Fig. 1) (Dunlop & Clark, 2006), and were documented across Canada and Fennoscandia in the mid-1900s (Hoppe, 1952; Frödin, 1954; Cowan, 1968; Lundqvist, 1969). They also are known to be present near morphologically similar landforms such as drumlins and hummocks, suggesting related formational processes. This spatial relationship raises the prospect of a “bedform continuum” present beneath ice sheets (Aario, 1977; Lundqvist, 1989; 1997; Dunlop & Clark, 2006; Ely et al., 2016; Möller & Dowling, 2018). Further, this spatial relationship allows us to determine

whether our approach can differentiate between different landforms with similar morphologies and spatial extents – this is possible through iteration with ribbed-moraine specific parameters across an area containing “ribbed moraine” and “non-ribbed moraine” landforms. In addition, we note scientific interest in specifically mapping ribbed moraines, the origin of which is still a matter of debate (Möller, 2006; Lindén et al., 2008; Boulton et al., 1987; Fisher & Shaw, 1992; Dunlop et al., 2008; 80 Sollid & Sørbel 1994).

Here, we develop and test two simple machine-learning algorithms: KM and RF (Gentleman & Carey, 2008) applied on new high-resolution datasets to design a computationally efficient, accurate and transferable method for automatically mapping ribbed moraine (Fig. 1). We define our performance metrics in terms of: efficiency – the processing speed of the method; effectiveness – the accuracy of the method; and transferability – how well the method performs in different terrain types on a regional/ country-wide scale. We further discuss the potential extensions to detect other landforms such as eskers, drumlins, megaripples etc.



90 **Figure 1: Aerial imagery of ribbed moraine at the Vinstre study area site (Fig. 3d) with (a) showing a general overview, and (b) showing a close-in view of ribbed moraine landforms. Ribbed moraine at this site appear to be partially anastomosed and form in a linear track (Dunlop & Clark, 2006). Aerial imagery from Norwegian national mapping data (Kartverket, 2021a, c).**

2 Methods

2.1 Machine learning

95 There are several machine learning approaches used previously in geomorphological research that may be useful for automatic feature detection. Many others exist, including U-net, a deep learning image segmentation method (Ronneberger et al., 2015), however, a main consideration of this study is to improve the time-efficiency of landform detection. Despite rapid segmentation once trained, U-net takes large amounts of time to train compared to random forest and K-means clustering (KM). Thus, we identify random forest and KM as two simple and lightweight methods that are used for similar image 100 segmentation problems. The first and simplest approach is unsupervised machine learning (Gentleman & Carey, 2008). Unsupervised machine learning produces a segmented output from input data and does not require training data (a dataset designed to describe what the algorithm should look for). The algorithm identifies clusters based on similarity of data properties (Gentleman & Carey, 2008). One such method is KM, often used in image segmentation (Burney & Tariq, 2014). Secondly, supervised machine learning methods such as Random Forest (RF) require a training dataset, consisting of labelled data to

105 provide a true reference for training the algorithm (Gentleman & Carey, 2008). RF has been used to great success for instance to map supraglacial lakes from satellite imagery (Discherl et al., 2020; Corr et al., 2022). Random Forest has additionally been proven as a valuable method for glacial and geomorphological work, for example mapping glacier change at high latitudes (Ali et al., 2023) and more directly in geomorphological mapping of glacial and hillslope process produced landforms in Switzerland (Giaccone et al., 2022).

110 As ribbed moraines typically have spatial scales of 10s – 100s of metres, (Hättestrand & Kleman, 1999; Dunlop & Clark, 2006), we select both RF and KM as suitable candidate methods for detecting them in high-resolution Digital Elevation Model (DEM) data (Fig. 2). We will compare the output from the RF approach, the KM approach and a series of more complex algorithms. We then compare the results to manually derived ground truth data (Fig. 2 b-d) to evaluate the performance of each method.

115 **2.1.1 Random forest**

The RF algorithm uses a series of decision trees (i.e. a forest) for classification, aiming to optimize agreement with training data (Gentleman & Carey, 2008). Within each decision tree, sequences of boolean questions subdivide the data (Breiman, 2001); the exact sequence and starting point of these questions is randomly altered for each tree of the forest. In this study, we test the RF method with 500 iterations using a majority vote method for final classification. Our RF method is trained
120 on a small area of known ribbed moraines to the north of the Vinstre study area. Training data here has not been used in any other component of the study and is derived from ribbed moraines mapped by Sommerkorn (2020).

2.1.2 K-means clustering


As an unsupervised algorithm, KM infers the defining parameters of each cluster based on the input data, and a series of 'K centroid' points randomly scattered throughout the dataset. The algorithm clusters most alike points around centroids
125 through repeated iterations of the model (Fig. 3; Gentleman & Carey, 2008). KM is commonly applied in fields such as photography and medicine (Burney & Tariq, 2014; Ng et al., 2020) but so far, has been rarely used in geomorphology (Lv et al., 2013). The ideal number of K-centroids is determined after the first iteration of the model (Likas et al., 2003), by plotting the sum of squared distances within the dataset against cluster count on a chart and selecting the 'elbow' point in the plotted line (Marutho et al., 2018). After the defining of the optimal number of K-centroids, the model is ready for data output (Fig.
130 3).

2.1.3 Outline of composite methods

We use different combinations of the above-described algorithms, for three additional 'composite' methods. These are designed to determine whether the limitations of individual algorithms can be overcome by combining RF and KM, at the expense of time-efficiency. Combined approaches are common within machine learning studies, as they can often improve
135 performance while avoiding micromanagement of the hyperparameters (Liu et al., 2021; Bhattacharjee et al., 2022).

The first composite method takes the output of a KM iteration to train the RF algorithm, rather than providing manually delineated training data. This method was selected as one of three combination methods with the aim of minimising subjectivity from outputs, allowing RF to determine features from an automated output, rather than a human-defined input. We term this method "KM-trained RF".

140



145 **Figure 2:** (a) Map showing the location of the study areas throughout mainland Norway. Karasjok (b) is shown in blue, Femundsmarka (c) is shown in green and Vinstre (d) is shown in magenta. Each location shows a submap (b, c, d) displaying outlines of ribbed moraines (yellow) used as ground truth in this study (Sommerkorn, 2020). Additionally, we used an area bordering the north of the site displayed in (d) for training the supervised methods. Basemap made of Eurostat’s GISCO administrative borders for Norway (GISCO, 2020). All maps projected in UTM 33N EPSG:5556, grid projection in (a) uses decimal degrees; b, c, d use UTM 33N m.

150 The second method is defined as the “OR” method, where both KM and RF are independently performed and the resulting classifications are merged by union, i.e. the final classification is positive if either KM or RF or both yield a positive. This method is designed to increase functionality if one or both methods are particularly strict on feature detection, or if each method functions more effectively in different areas.

155 The third method is defined as the “AND” method, where both KM and RF are independently performed and the resulting classifications are merged by intersection, i.e. the final classification is positive only if both KM or RF agree. We designed this method to increase functionality in situations where one or both methods have high rates of false positives.




Figure 3: Flowcharts describing (a) the basic principles behind the RF algorithm. (b) The basic principles behind the operation of a KM methodology. The model is first run to determine the number of K points required to represent the input data, and then run a second time to produce an output.

160 2.2 Method application

2.2.1 Study areas

We selected three study areas representing different landscapes present in Norway for evaluation (Fig. 2). The landscapes included are high mountain (Vinstre), heath/moorland (Karasjok) and semi-mountainous marshland (Femundsmarka). Each study area covers 15 x 15 km, with data at a 1 m resolution (Kartverket, 2021). The first study area is south of lake Vinstre in central southern Norway (Fig. 2d). This region is representative for inland Norway, with a high mountain plateau interspersed with deep glacially formed valleys (Strøm, 1948). Furthermore, there exists high-quality ribbed moraine mapping in and around the Vinstre study site (Fig. 2, 4; Sollid & Sørbel, 1994; Sommerkorn, 2020), which we use for training and validation of our methods. The second study area covers a subsection of the Femundsmarka National Park, near the Norway-Sweden border (Fig. 2c). This site is 30 km to the southwest of Lake Rogen, where ribbed moraines were initially identified and described (Lundqvist, 1969). A third study region lies in Finnmark, north of Karasjok, situated in a heathland and thus is largely different to the former two, both in form with large rolling hills (similar in morphology to ribbed moraines), and in geology, with much less exposed bedrock.

We trained each supervised method on a small 15 x 5 km subregion bordering the north of the Vinstre study area to determine the relevant parameters and morphometrics (Fig. 4). We initially tested each method on the Vinstre study region, as the terrain in the Vinstre region is highly complex and typical of inland Norway, making it useful for ensuring the algorithm functioned well in similarly complex regions.

2.2.2 Map data

We use Norway's 1 m DEM and resample it to 10 m using bilinear interpolation for consistency instead of using an available 10 m product which was produced from various and uncertain sources (Fig. 2) (Kartverket, 2021b). We resample data for three purposes (a) because ribbed moraine occur at the scale of tens of metres, sub-10 m data is unnecessarily detailed (b) because without resampling the 1 m data captures too much detail in the ground surface, leading to unnecessarily detailed morphometrics, and (c) to limit processing time, as we aim to produce a time-efficient methodology. The surface geology data is obtained as vector data from the Norwegian Geological Survey, NGU (NGU, 2022), and lake vector data from the Norwegian

Water Resource and Energy Directorate, NVE (NVE, 2023)- We used nearest neighbour interpolation to convert these vector
185 datasets into 10 m raster data, matching our resampled DEM. We obtained orthophotographs from Norway in pictures, (NiB)
(Kartverket, 2021a, c) and use this for ground-truth production and qualitative output validation. In addition, we use existing
geological maps and mapped ribbed moraine ridges from Sommerkorn (2020).

2.2.2 Derived data

We use our raw data for two purposes: to produce training/testing datasets, and to generate morphometric and mask
190 attributes for our machine learning algorithms. We manually delineated training/testing ground truth data using a combination
of previous moraine maps, the 1 m DEM and orthophotographs. We find that ribbed moraines in marsh areas are easily
identifiable in orthophoto imagery due to their distinct contrast in colour to their surroundings, however in forests they are not
detectable due to tree cover (Dunlop & Clark, 2006).

We derived morphometric and masking data from the DEM using several modules: the RichDEM python module
195 (Horn, 1981; Zevenbergen & Thorne, 1987; Barnes, 2016), SAGA (Conrad et al., 2015) and the opencv python module
(Bradski, 2000). We also produced several filtered DEMs from the input 1 m DEM including: a 400 m low pass filter, a 30 m
high pass filter, and a 30–400 m bandpass filter designed to isolate the size scale of ribbed moraines as defined by Dunlop and
Clark, (2006). For our topographic wetness index (Boehner et al., 2002) data, we calculated indices using local values derived
from the input DEM rather than global.

200 2.3 Morphometrics

We first select morphometrics based on the basic observable characteristics of ribbed moraines. In addition to the
filtered elevation data (30 – 400 m; Hättestrand & Kleman, 1999; Dunlop & Clark, 2006) we generated: slope, general slope,
curvature, planform curvature, profile curvature, spatial distribution (wavelength) and aspect. In addition, we derived the
topographic wetness index (Boehner et al., 2002, Conrad et al., 2015) and topographic position index (Guisan et al., 1999) as
205 metrics with the potential to add value to the segmentation algorithm. Each morphometric was identified from prior literature
(e.g. Eisank et al., 2014), and qualitatively tested for its ability to isolate ribbed moraine. With the combination of best-
performing morphometrics we suggest the possibility of differentiating ribbed moraines from other landforms.

Secondly, we selected masking values, to remove as much noise from the input data as possible and avoid obvious
misclassifications. This also aids our accuracy metrics by improving the balance of our data, as both RF and KM work most
210 successfully on balanced data, rather than unbalanced data (Pedregosa et al., 2011). Here, the best-balanced data is defined as
data where there is roughly a 50/50 split in the count of pixels per binary output category. Hence, masking values are used to
remove as many pixels as possible where ribbed moraines do not or cannot form, balancing the dataset somewhat. From
consulting literature, we defined areas where ribbed moraines cannot form: mountain peaks (Hättestrand & Kleman, 1999),
steep slopes (Sommerkorn, 2020) and where surface bedrock is present (Sommerkorn, 2020). We also defined lakes as areas
215 in which ribbed moraines are not detectable using DEM data. For each of these limitations, we produced a mask layer. Large-
scale (km<) peaks and ridges are masked from the dataset by setting a threshold on a laplacian of gaussian filter. The laplacian
of gaussian is a second-order derivative edge detection filter, which identifies regions that show sudden steep changes in
intensity (Kong et al., 2013). We mask steep slopes using a generalised slope layer generated from a low-pass filtered DEM
(400 m <), slopes inclined greater than the thresholding value (e.g. Table 2) were masked out of the dataset. The low-pass
220 filtered DEM is used together with a slope angle threshold to mask out the general valley slope angle. Additionally, we mask
mountains of a set wavelength from the testing to omit plateau-edge convex surfaces from analysis using a simple sinusoidal
laplacian filter (Kong et al., 2013), referred to as a “mountain wavelength mask”. In this case, mountains with a wavelength
of greater than the threshold value were filtered out of the dataset. To mask out areas of exposed bedrock, we used national

225 surface geological map of Norway (NGU, 2022) and mask all areas defined as exposed bedrock. Finally, we used NVE's lake
database to mask out lakes (NVE, 2023).

230 Method outputs were visually compared to the locations of ribbed moraines as defined in Sommerkorn (2020). We
made qualitative visual comparisons here as the changes made by different morphometrics were large, with numerical analysis
coming as part of later predictor tuning. We made our visual comparisons against high-resolution orthophotography
(Kartverket, 2021a, c), and Norway's 1 m resolution national DEM (Kartverket, 2021b). We then adjusted the algorithm to
produce the most visually accurate output with the fewest inputs. Hence, in developing the method, we removed aspect as it
made no notable contribution to the output. We also conflated both planform and profile curvature into the total curvature
metric, as we observed no difference in the output from any singular curvature parameter. In testing the value of the literature-
defined morphometrics, we found topographic wetness index to be highly useful, but topographic position index to add no
value to the output in any combination.

235 **2.4 Workflow in practice**

We implemented the series of segmentation algorithms on the combination of raw and derived data. We began by
pre-processing the datasets to fit with each chosen study area, defined by the borders of the associated 1 m resolution DEM
tile (Fig. 2). Once pre-processed, we ran each test algorithm on the datasets, for Vinstre, Femundsmarka and Karasjok. We
also carried out a second iteration of RF using KM-produced training data, and combined outputs from KM and RF to produce
240 each of our composite method outputs. Once we had completed each method, we prepared output datasets for statistical
analysis via a standard confusion matrix (Fig. 5).

The first metric to test each method was efficiency – defined as the required CPU time. Then, we measured the
effectiveness of each method in detecting ribbed moraines. For each of the five methods, we derived the confusion matrix,
stating the numbers of true positive, true negative, false positive and false negative pixels, comparing model outputs to the
245 manually delineated ground truth data (Fig. 5; Hong & Oh, 2021).




Figure 4: Illustration of morphometrics used in this study, with red line indicating the transect taken for subchart figures. (a) to (f) show clustering values, whilst (g) to (i) show masking values. (a) Shows elevation from the 1 m resampled to 10 m DEM data (Kartverket, 2021), (b) shows elevation with km-scale features filtered out, (c) shows the output of the general slope (low frequency slope) filter, (d) shows the local slope values used for clustering, (e) shows curvature values used for clustering, (f) shows topographic wetness index (TWI), (g) shows masking values used for excluding regions where ribbed moraine cannot form, (h) shows curvature derived from a laplacian of gaussian filtered DEM for masking mountain peaks (i) shows a similar plot at 1 km resolution for excluding large wavelength features.

250

255

2.5 Comparative Analysis

Ground Truth \ Output	Ribbed Moraine	Not Ribbed Moraine
Ribbed Moraine	True Positive	False Positive
Not Ribbed Moraine	False Negative	True Negative

Figure 5: Binary confusion matrix diagram showing actual vs. predicted values on a 2 x 2 grid.

In addition to producing a visual representation of each method's effectiveness (Fig. 4), we calculated accuracy metrics: balanced accuracy (BA (Equation 4); Brodersen et al., 2010), and F-Score (Sokolova et al., 2006), each of which represents different sides of the confusion matrix, and each ranging between 0 and 1. We selected BA and F-score over a standard accuracy metric (Equation 1) as standard accuracy is skewed positively in unbalanced datasets. As our dataset comprises regions where ribbed moraine do not make up 50% of the mapped area, our data can be considered unbalanced, hence we make use of BA. To calculate BA, we use Equation 4.

$$Accuracy = \frac{(True\ Positive + True\ Negative)}{(True\ Positive + False\ Negative + False\ Positive + True\ Negative)} \quad (1)$$

$$Recall = \frac{True\ Positive}{(True\ Positive + False\ Negative)} \quad (2)$$

$$Specificity = \frac{True\ Negative}{(True\ Negative + False\ Positive)} \quad (3)$$

$$BA = \frac{(Recall + Specificity)}{2} \quad (4)$$

BA accounts for the imbalance between classes, and it works effectively in identifying the influence that positives have on the accuracy of a dataset, hence a higher BA score represents a dataset with good positive returns versus Ground Truth. F-score is computed as the harmonic mean of precision (Equation 5) and recall (Equation 6).

$$Precision = \frac{True\ Positive}{(True\ Positive + False\ Positive)} \quad (5)$$

$$F\text{-score} = 2 \cdot \left(\frac{(Precision \cdot Recall)}{(Precision + Recall)} \right) \quad (6)$$

F-score ignores true negatives, and like BA works well on imbalanced datasets. F-score typically will be higher when there are lower false positive and false negative, as it pays less attention to true positives and true negatives than BA. Therefore, in working with both metrics, we gain a detailed understanding of the accuracy of our output.

2.6 Final algorithm adjustments

Once we identified which algorithm provided the best combination of effectiveness and efficiency, we improved the performance by adjusting the models' parameters. We took two approaches to make these adjustments: systematic and intuitive. For systematic improvements, we adjust parameter values in small increments over a physically plausible range: general slope mask threshold values are adjusted by 0.01 from 0.65 – 0.75, laplacian curvature mask threshold values are

adjusted by 0.005 from 0.04 – 0.08, general slope kernel size is adjusted by 10 m from 300 m – 400 m, and mountain
285 wavelength masking is adjusted by 100 m from 1 km to 2 km (Table 2). These tweaks thus provide an objective analysis of
outputs based on pure accuracy scores.

For our intuitive approach, we changed inputs with the aim to maximize visual agreement between the classification
and the evaluation data. This approach served to health-check our dataset, ensuring that our method specifically detected ribbed
moraines rather than “detection by accident”. We used both approaches as a means of performing a robust sensitivity analysis,
290 ensuring that specific landforms were detected. In certain situations, many pixels of ribbed moraines may be identified, but
many conjoining pixels could also be detected as “ribbed moraine”, thus only detecting ribbed moraine fields, rather than
discrete landforms. Hence, while we aim to use the systematic analysis to determine the parameters with the highest statistical
accuracy, we used the intuitive analysis to determine the parameters which lead to the highest accuracy scores while still
detecting discrete landforms rather than fields. The intuitive analysis involved changing the same parameters as in the
295 systematic tweaks, whilst also changing cluster counts by ± 1 , and minor adjustments at the scale of 10 m to the filtering kernel
size used on the DEM.

3.0 Results

3.1 Comparative method results

Our segmentation methods were quick to run (10 – 60 s) and produced reasonable outputs on a mid-range laptop
300 (Intel i5-1135G7, 2.40GHz, 2420 MHz, 4 core. Intel Iris Xe Graphics. 16GB 3200MHz RAM). However, the outputs varied
greatly between methods and study areas. As such, we separated our results between each segmentation algorithm and study
area, and then outlined the mean results between each study area for each method.

3.1.1 Vinstre

In our test and evaluation of the five chosen methods at Vinstre, we find reasonable outputs for each method, showing
305 the capabilities of detecting ribbed moraines in the terrain of this region (Fig. 6). However, each algorithm varies through its
false positives and false negative detection rates. The standard accuracy (Equation 1) values (Table 1) show accuracy to be
over 88% for all methods, with a mean of 0.926. However, due to the natural imbalance of our data, this metric is largely
dominated by the abundance of negatives. We expected some exaggerated false positive detection, yet some methods are better
than others at omitting this. In addition, some of the methods detect more false positives than others, which is more difficult
310 to exclude from the dataset. For example, we can see that KM and OR outputs show consistent false positives on the rounded
peaks in the Vinstre study area's south; while the KM-trained RF algorithm shows a widespread pattern of random false
positive detection (Table 1). The RF and AND methods yield fewer positive pixels overall, which leads to lower accuracy
scores as only the banks or ridges of ribbed moraines are detected, rather than full features.

Statistically, we find the KM and OR methods to have the highest and most consistent accuracy metrics, suggesting
315 that these methods have the most use in ribbed moraine detection. We see this with KM returning BA of 0.75 σ (standard
deviations) above the mean, and F-scores of 0.89 σ above the mean (Table 1). The OR output echoes this, with BA and F
values of 1.16 and 1.00 σ above the mean respectively (Table 1, Fig. 6). On the other hand, while the AND approach is the
second most effective method in terms of overall accuracy, we see poor results in balanced metrics, giving values of 1.20 σ
below the mean value. Furthermore, the KM-trained RF method shows the least promising visual output, the least consistent
320 and near-least promising accuracy output, where accuracy = 1.34 σ below the mean, BA = 0.57 σ below the mean, and F-
score = 0.41 σ above the mean (Table 1). Therefore, as an overall ranking, we rate the OR method as the best performing in
the Vinstre region, while the KM-trained RF method returns the least successful results.

3.1.2 Femundsmarka

On comparing results from Femundsmarka, we find a similar pattern to Vinstre, with high false positive and negatives common between KM-trained RF. Additionally, RF and AND classifiers, showed distinct features in qualitative detection. False positive detection of rounded hills in the Femundsmarka region is reminiscent of the Vinstre region in the KM and OR classifiers. We also rank each method the same as at the Vinstre site, whilst also showing that the variation in performance from the mean is similar between both sites. We find major differences are only present within the KM-trained RF method and the RF method in terms of variability. For these, KM-trained RF returns values of 1.27, 0.55 and 0.25 σ below the mean for Accuracy, F-score and BA, respectively. RF on the other hand, returns 1.04, 0.09 and 0.79 σ above the mean. In this case, we see that the KM-trained RF method consistently trended towards below-mean performance, whereas RF consistently performs above-average.

Table 1: Performance scores for each study area (Fig. 6); metrics are rounded to 3 decimals. As BA and F-score are combinations of recall and specificity, (Equations 2 and 3) we only include the BA and F-score outputs in this table.

Vinstre				
METHOD	Accuracy	F-Score	BA	
KM	0.915	0.229	0.679	
RF	0.963	0.163	0.550	
KM+RF	0.875	0.147	0.653	
OR	0.913	0.244	0.698	
AND	0.965	0.112	0.531	
<i>Average</i>	<i>0.926</i>	<i>0.179</i>	<i>0.622</i>	
<i>Std_Dev</i>	<i>0.038</i>	<i>0.056</i>	<i>0.076</i>	

Femundsmarka				
METHOD	Accuracy	F-Score	BA	
KM	0.925	0.214	0.639	
RF	0.963	0.182	0.556	
KM+RF	0.903	0.148	0.614	
OR	0.925	0.238	0.661	
AND	0.964	0.104	0.529	
<i>Average</i>	<i>0.936</i>	<i>0.177</i>	<i>0.600</i>	
<i>Std_Dev</i>	<i>0.026</i>	<i>0.053</i>	<i>0.056</i>	

Karasjok				
METHOD	Accuracy	F-Score	BA	
KM	0.881	0.059	0.666	
RF	0.981	0.135	0.566	
KM+RF	0.824	0.029	0.617	
OR	0.881	0.059	0.666	
AND	0.984	0.091	0.536	
<i>Average</i>	<i>0.910</i>	<i>0.075</i>	<i>0.611</i>	
<i>Std_Dev</i>	<i>0.070</i>	<i>0.040</i>	<i>0.059</i>	

Averages				
METHOD	Accuracy	F-Score	BA	Time (s)
KM	0.907	0.168	0.662	18
RF	0.969	0.160	0.558	1085
KM+RF	0.867	0.108	0.628	1103
OR	0.906	0.181	0.675	1433
AND	0.971	0.102	0.532	1493
<i>Average</i>	<i>0.924</i>	<i>0.144</i>	<i>0.611</i>	<i>1026</i>
<i>Std_Dev</i>	<i>0.045</i>	<i>0.036</i>	<i>0.063</i>	<i>594</i>

3.1.3 Karasjok

Performance patterns in the Karasjok region depart from those in the previous areas. There is greater variability between algorithms and a general increase in misclassifications. The RF and AND methods perform well in Femundsmarka and Vinstre, while the KM, KM-trained RF and OR methods produce outputs with speckled pixels. Many false positives are present, particularly in the southern part of the Karasjok region. It is likely that these are a result of different landscape morphologies between Vinstre, Femundsmarka and Karasjok, where the latter is a moorland environment, versus the prior two, which are more typical of inland Norway (Hjort et al., 2014). Statistically, we also see poor performance KM rates below the mean standard accuracy and F-score metrics by 0.40σ , whilst maintaining a high BA score of 0.93σ above average (Table 1). In short, this means KM is effective at detecting true positives but classifies too many false positives. RF is opposed to this, with accuracy and F-score returning values of 1.01 and 1.50σ above average respectively, versus 0.76σ below average for BA. The OR method returns identical values to KM, as RF under-classifies features in the Karasjok region, thus allowing KM to dominate the OR output in this area.

Despite the differences in the Karasjok region, we still see similarities between the Karasjok region and the prior two, with both the KM-trained RF and AND approaches having inconsistent variability from the mean, suggesting their lack of value as methods. Due to the locality of the false detections found in the KM and OR methods, it is possible that they are detecting features common to this region, which likely have a similar form to ribbed moraines found elsewhere in Norway.

3.1.4 Overall performance

Studying the overall performances of each method across the different study sites, we find similar results in the first two regions, where KM ranks consistently second in BA, and OR ranks highest. KM performance scores are both above average by 0.38σ and 0.81σ , while all other methods show some inconsistency between accuracy metrics. For example, the OR method ranks 1.02 – 1.03σ above the mean BA but ranks 0.40σ below the mean performance. Hence, while KM does not rank as highly as OR in BA, its output is more consistently accurate. As a result, we can discount the combined, RF and AND methods, as they all consistently rate third or worse in accuracy metrics, showing their lack of segmentation effectiveness.

In addition to accuracy, we determined the CPU time required for each method (Table 1) as a measure of efficiency. On a mid-range laptop (Intel i5-1135G7, 2.40GHz, 2420 MHz, 4 core. Intel Iris Xe Graphics. 16GB 3200MHz RAM), KM takes only 18 seconds on average per iteration, 1.67σ faster than the mean rate, while every other approach takes two orders of magnitude more, at over 1000 seconds. When considering our two most statistically effective methods, we find KM is notably faster than the OR methodology. Due to its much lower computational cost, we selected KM as the method of choice, despite its slightly lower performance.

3.2 K-means refinement

Upon producing a baseline set of predictors for the KM methodology, we conducted many predictor adjustments as a sensitivity analysis. These values aided in determining two final “best-fit” methods based on: statistics through a systematic approach; and statistics combined with qualitative observations made on the output maps.

370




Figure 6: Comparison of input to output in the three study regions. (a, c, e) show ground truth data, while (b, d, f) show accuracy maps denoting confusion matrix (Fig. 5) values for the clustered output of the KM method. Outputs are superimposed on a hillshade of the input DEMs (Kartverket, 2021b).

3.2.1 Systematic analysis

The systematic analysis yields optimal predictor values, in terms of average BA scores for each iteration per region. For example, the optimal value of the general slope threshold is 0.71, where BA values are 0.76, 0.61 and 0.72 for the Vinstre, Femundsmarka and Karasjok sites respectively. The optimum BA scores 1.7 σ above average for the range of tested values (Table 2).

In addition to our best outputs, we note interesting values and themes throughout the systematic analysis. In general, we found each region to have similar patterns of maximal performance. The only commonality between our three sites under the general slope kernel value tweaks was with a parameter value of 300 m (where the kernel size for slope smoothing is 300 m resolution), where all scores were above 0.64 BA. The pattern of scores calculated using 300 m general slope kernel, (0.74 Vinstre, 0.65 Femundsmarka, 0.73 Karasjok) is common throughout this parameter's output. While Femundsmarka generally follows the pattern of values in the other two datasets, changes are more muted, with values ranging between 0.61 and 0.67 for general slope kernel, general slope threshold and laplacian curvature threshold, versus changes of double this magnitude in the Vinstre and Karasjok regions. Furthermore, while there is general agreement throughout the data, Vinstre appears to be at odds in optimal values to Karasjok for accuracy scores in laplacian curvature threshold tests, with optimal values having no relation.

Table 2: Table showing predictor values from systematic analysis with BA as the key accuracy score. BA Range shows the range from minimum to maximum values of BA output. Each predictor was tested on an individual basis.

	Location	Value	Value Range	BA Global Average	BA Range	BA Standard Deviation
General Slope Threshold	Vinstre	0.71	0.65 – 0.75	0.71	0.606	1.7 σ above mean
	Femundsmarka	0.71			–	
	Karasjok	0.71			0.757	
Laplacian Curvature Threshold	Vinstre	0.07	0.04 – 0.08	0.67	0.608	0.9 σ above mean
	Femundsmarka	0.07			–	
	Karasjok	0.07			0.745	
General Slope Kernel	Vinstre	300 m	0.3 km – 0.4 km	0.71	0.601	1.8 σ above mean
	Femundsmarka	300 m			–	
	Karasjok	300 m			0.774	
Mountain Mask	Vinstre	1000 m	1 km – 2 km	0.76	0.538	2.4 σ above mean
	Femundsmarka	1000 m			–	
	Karasjok	1000 m			0.791	

The tweaks made to the mountain wavelength mask present much more varied results than the other tests; but a consistent optimal or near-optimal value. This lack of consistency is likely due to the different wavelengths of mountains in each study area. But additionally, it appears that a consistent wavelength for mountain features is roughly 1 km between trough and peak. Our results clearly show this, with maximum BA values of 0.79 (Vinstre), 0.76 (Femundsmarka) and 0.73 (Karasjok). This high performance also mean that the mountain mask parameter is the most influential parameter on statistical detection, improving the mean BA across the board by 0.10 (3.6 σ above the mean improvement). On the other hand, the least influential parameter is laplacian curvature threshold, with an improvement in accuracy of only 0.03, only 0.9 σ above the mean BA score. This result, however, is consistent with the lack of agreement between datasets when tweaking the laplacian curvature threshold value.

3.2.2 Intuitive approach

Results from our intuitive analysis show several outputs with high BA. Each of these approaches yields BA scores > 0.70 at Vinstre, and a mean of > 0.69 across all regions. Yet, visual evaluation reveals that settings in which BA score is > 0.70 can lead to moraine overclassification (Fig. 6). Thus, despite high BA, this validates our health-check, demonstrating that

higher statistical performance does not necessarily produce the best performing output for discrete feature detection. This potentially comes from the complexity of the terrain in which we iterate the method, with areas of higher relief generally performing worse than lower relief when using geomorphometric methods (Hjort et al., 2014). In short: we find $BA < 0.70$ leads to discrete detection of ribbed moraines, whilst $BA > 0.70$ delineates ribbed moraine fields, filling the gaps between features. Hence, we outline two potential approaches: (1) That high accuracy should be the focus, aiming to detect fields of ribbed moraines as in previous studies, and (2) that discrete ribbed moraines should be the focus, aiming to detect individual ribbed moraines on a large scale.

We also note several patterns between our analysis approaches. Firstly, we find less agreement between all three regions in the statistical output of our trial-and-error intuitive approach versus the systematic approach. Despite this, we also find there is consistent agreement in our most effective discrete landform detection method, with scores varying by $< 1 \sigma$. Interestingly, we find that again, Femundsmarka shows the lowest variability in BA score, with a standard deviation of 0.29σ , versus a standard deviation of 0.50σ for Vinstre, and 0.56σ for Femundsmarka.

4 Discussion

4.1 Pre-predictor tuning

The high level of agreement with ground truth in the Vinstre region show the KM algorithm's potential. Through F-score and BA, we find the KM method to be most successful in the Vinstre study area, likely due to the method being originally initiated on the complex, mountainous terrain of inland Norway. This is promising because it shows the method performs well at detecting ribbed moraine in regions of high relief, one of the most common areas for ribbed moraine to form (Sollid & Sørbel, 1994; Sommerkorn, 2020). Hence, we assert this method is transferrable throughout central Norway and other similarly mountainous regions. Yet, we find the main shortcoming of the pre-tuning method in the Vinstre study region to be over-detection of riverbanks, palaeochannels and other, morphologically similar landforms. This is potentially in-part due to the interpolation of 1 m to 10 m data averaging morphometric signals.

We additionally find good performance in the Femundsmarka study area, with an F-score of 0.21. This is most likely due to the partial similarity to the Vinstre study area, with the presence of some high mountain terrain. However, complexity in the local relief increases, including three terrain types: moorland, lake and high mountain, versus the consistent high mountain terrain of the Vinstre study area. While lakes are masked out, the mixed semi-mountainous marshland of Femundsmarka likely contributes to the lower accuracy scores due to transferability issues between different reliefs (Hjort et al., 2014), as Femundsmarka's results show the algorithm attempting to classify lakeshore and rounded moorland hills as ribbed moraine (Fig. 6d). This reflects the riverbank detection in Vinstre, and thus is most likely due to the similarity of morphology, particularly in relation to slope and curvature (Dunlop & Clark, 2006; Lindén et al., 2008; Möller, 2006). Additionally, Femundsmarka shows greater rates of false negatives than Vinstre, with many ribbed moraines in the south of the study area (Fig. 6d) being unclassified. The cause of the false negatives is unclear, but they could be attributed to the lower overall difference in elevation between ribbed moraines and their surroundings (30-40 m North, 20-25 m South) in southern Femundsmarka. Despite some of the limitations, we consider our initial results from Femundsmarka promising.

With an F-score of 0.06, Karasjok has a greater rate of false positives than other regions (Fig. 6). This may be a result of the landscape being predominantly comprised of rounded hills of similar appearance to ribbed moraine, similarly to areas of Femundsmarka. BA is 0.67, in comparison to the 0.64 and 0.68 for Femundsmarka and Vinstre, respectively. Hence, the primary issue with the Karasjok region is over-detection (increased false positives). We propose the issues of false positive detection we observe in Karasjok to be a result of inherent transferability issues that arise in geomorphometry. Hjort et al. (2014) describes the difficulty in extrapolating geomorphometric methods from high to low relief areas, as region specific conditions can have a strong impact on the relative importance of different variables. In the case of Karasjok, the change in

relief likely leads to a greater influence of the curvature metric. Alternatively, this could be a product of the interpolation
450 between 1 m and 10 m leading to an inaccurate curvature value for channel features.

The overdetection we observe in our study regions could be limited by predictor tuning (Moradi Fard et al., 2020) or through classifying detected features post-detection. For this study, we choose to take a consistent approach and undergo predictor tuning to improve our methodology, due to the initially promising output from our approach.

4.2 Finalised outputs

455 4.2.1 Computation considerations

For all iterations, we found that the relatively light computing load opens the possibility to apply the method to larger scales such as mapping ribbed moraines all over Norway. In addition, the parallelized implementation of K-means named “minibatchkmeans” (used in this study) (Pedregosa et al., 2011) further increases the processing speed on multicore servers.

4.2.2 Performance

460 In computing BA, we find reasonable (0.59) to good (0.78) scores depending on iteration and location. Mean BA scores vary by 0.11 (0.60 to 0.71), which we expect given the varied terrain types. This outlines issues in transferring between terrain types and maintaining performance. For example, during our systematic analysis, we set the general slope threshold to 0.75, and find BA values of 0.63 (Vinstre), 0.62 (Femundsmarka) and 0.73 (Karasjok), demonstrating better performance in the Karasjok moorland than in high mountain terrain. Due to discrepancies between what can be considered valuable in a
465 qualitative versus quantitative results, BAs below 0.75 do not suggest poor method performance, as values between 0.65 to 0.70 show the greatest performance in this study for detecting discrete ribbed moraines. It is, therefore, important to consider that the algorithm is not designed to detect ribbed moraine but “ribbed moraine-like” features. This means there will always be a degree of over/under classification, particularly in complex post-glacial and real-world landscapes due to the variability in ribbed moraine shape (Dunlop & Clark, 2006). We will also see over/under classification due to geomorphological features
470 of similar shape within our study areas, as ribbed moraines do not develop in isolation (e.g. drumlins, channels, mega-scale glaciolineations; Ely et al., 2016). Hence, over/underdetection can be perceived positively, as output features can be classified post-detection using parameters such as orientation versus glacial streamflow, with ribbed moraines as perpendicular to, and drumlinoid forms streamlined with flow directions (Dunlop & Clark, 2006; Ely et al., 2016).

Secondly, we find that BAs below 0.70 may be of greater value than those above 0.70 if we are aiming to identify
475 discrete features. Our results show that BAs of more than 0.70 present more true positives, resulting in good statistical results, but many more false positives are also present, as spaces between ribbed moraines are detected as ribbed moraine. This results in areas between ribbed moraine being filled up, leading to positive detection of moraine features, but discrete ribbed moraines are poorly captured. Hence, we consider a range of results as successful: BA 0.65 to 0.75 depending on whether the aim is to classify fields or discrete landforms. This is due to BA alone not being the ideal parameter for measuring the value of our
480 output – but instead we require complementary assessment, through qualitative analysis as a means of determining output value, on top of the ideal range of BA values. We thus believe both sets of BA scores are reasonably accurate, particularly as our raw accuracy never drops below 0.82, representing an 82% per-pixel success rate.

4.2.3 Transferability

The transferability of this algorithm depends on one major component: the wide-scale landscape type and relief,
485 showing similar issues with transferability to previous works (Hjort et al., 2014). We find the method works well when applied to similar landscape types, as seen in Femundsmarka and Vinstre. When transferred to different landscape types and reliefs, (i.e. between high mountain, moorland, and mixed marsh-mountain terrain) we observe poorer detection of ribbed moraines. As a result, this algorithm is transferable for most landscapes where ribbed moraines occur in inland Norway (high mountain),

however, for regions exhibiting differences to the high mountain regions, such as moorland and coastal regions, we can tune
490 the model accordingly at the scale of each DEM tile. This is relevant when considering expansion to other regions of the world,
as much of the Canadian shield is landscape close in format to Karasjok – which would indicate a need for predictor tuning
(Dunlop & Clark, 2006; Dunlop et al., 2008).

4.2.4 Detection of new features

While analysing our outputs, we found evidence of ribbed moraines detected by the algorithm that were not present
495 in ground truth data. These were commonly found in the Femundsmarka region, in areas adjacent to manually marked ribbed
moraine (Fig. 7). The detected landforms occur in small clusters, with a similar pattern to ribbed moraine as described in
Dunlop & Clark (2006). We suggest that this may be due to a series of factors, including the presence of forest and farmland,
obscuring ribbed moraine in stereo–imagery or poor DEM data used for manual production of ground truth data. Hence, we
suggest that while the method has some limitations, there are also clear advantages as the method uses LiDAR data, which
500 does not exhibit the same issues in differentiating between landforms under forest cover as spectral imagery.





Figure 7: Comparison of ground truth to detected pixels on (a) hillshade and (b) Orthophoto imagery (Kartverket, 2021a, c). Here, GT refers to “Ground Truth”.



510 **Figure 8: Detected features versus LGM ice flow direction in the Karasjok region. (a) shows values $\pm 25^\circ$ from orientation perpendicular to mean flow direction (blue arrows) during the FIS (Fennoscandian Ice Sheet) (Patton et al., 2016; 2017) and (b) shows values not oriented perpendicular to average flow angle during FIS (Patton et al., 2016; 2017). Both figures superimposed on 10 m DEM and hillshade (Kartverket, 2021b).**

To explore possibilities for further refinement, we calculated the orientation of each detected polygon using a simple bounding box method and determined the orientation of its longest axis. We compare these orientations with modelled flow direction of the FIS by comparing feature orientation to the mean flow direction of the ice sheet between 11kya and 25kya (Patton et al., 2016; 2017). Fig. 8 shows polygons perpendicular or streamlined ($\pm 25^\circ$) to glacial flow, helpful in distinguishing landforms oriented along (drumlinoid) from those oriented transverse (ribbed moraines) to flow direction (. Yet, we note limitations with hummock forms (Ely et al., 2016), as they have no dominant orientation and could end up in any group, demonstrating a need for a more robust classification method, potentially based on feature aspect ratio. We also note this method has challenges detecting the orientation of anastomosing features, a common type of ribbed moraine (Dunlop & Clark, 2006). Another potential limitation of this method is the low resolution of modelled glacial flow, which is in the order of 520 kilometres, wider than many glacial valleys.

4.3 Avenues for method improvement

We identified two areas of improvements. First, we suggest improving automation, which would allow scaling to larger areas. For example, through inclusion of automated predictor tuning for landscape types in the model parameters (parameter profiles). Automation would allow for application on a country scale with minimal input through the implementation of these parameter profiles, minimising false positive and false negative identification in varied landscapes. We envision four landscape types for Norway: high mountain (our original approach), low mountain, moorland (Karasjok) and coastal. We believe these four categories would cover the main landscape types common across regions where ribbed moraines are present. These categories would likely prove sufficient in the regions outside Norway where ribbed moraines are known to exist (e.g. Canada, Ireland, United Kingdom, Sweden, Finland), however regional differences in scale and geology may require landscape type parameterisation. 530

Second, we identify the importance of implementing a post-detection classification of landforms. Our results show that overclassification is common, particularly as there are many landforms with similar geomorphological properties to ribbed moraines (Fig. 6). We suggest future works include investigation into landform classification, as it would allow for identifying specific landforms including drumlinoid forms, ribbed moraine and others, improving the value of the methodology. This would potentially aid the continuum hypothesis of Ely et al. (2016), by showing landform transitions from ribbed moraine. Hence, we consider this as the next logical step in method development. In addition, we suggest that with post-detection 535

classification, this method could be used on a wider basis than only ribbed moraine, due to its ability to easily detect specific geomorphologies based on relatively basic morphometric traits.

5 Conclusions

540 To develop a method for automated mapping of landforms throughout Norway, we tested two machine learning algorithms and three composite approaches. We determined ribbed moraine as a suitable example landform, which would also be scientifically interesting to identify. Through testing, we settled on using an unsupervised K-means clustering algorithm for moraine detection, thus requiring no training data. Ground truth data for our testing was produced through manual analysis of high-resolution elevation model (Kartverket, 2021) derivatives, and high-resolution aerial/ satellite imagery (Kartverket, 545 2021).

Our results demonstrate unsupervised machine learning as sufficient for the automated detection of geomorphological features through a simple KM approach, rather than the need for complex supervised machine learning methods. We also demonstrate that minimal data are needed for this approach, with only high-resolution DEM derivatives, superficial geology and a lake mask required for our methodology to function. We evidence this in initial testing, where a supervised RF method 550 averaged poorer performance than an unsupervised KM approach. In addition, we find that we can differentiate between identification of discrete and fields of landforms. Hence, we find this method to be scalable, in that different output resolutions are possible.

This study indicates the value of automated machine learning for landform detection as a means of minimising time spent delineating features manually and in-field. We detect ribbed moraines throughout our study areas in relation to ground truth data and detect a small number of previously unmapped ribbed moraines in one of these. Thus, we show the value of an 555 objective and systematic method using new, high-resolution data in detecting features. Furthermore, we detect many additional features in all study areas, which have similar morphologies to ribbed moraine, but powerful refinement can be achieved by considering orientation with respect to the former ice flow direction.

In summary, this study demonstrates that unsupervised machine learning is a viable and efficient method for the 560 automated detection of ribbed moraines and similar features based on modern high-resolution DEM (Kartverket, 2021b). With our promising results, we identify a future path for such methodologies in geomorphology, as a means of updating geomorphological maps and producing new geomorphological maps where we have high-resolution data to input (e.g. dune mapping on Mars). Thus, we intend to develop this work into mapping ribbed moraine and adjacent features throughout Norway, so as to aid in developing our understanding of the geological history of Fennoscandia.

565 Code availability statement

Finalised code is available in the repository Aeteia/Ribbed-Moraine at <https://doi.org/10.5281/zenodo.7991094>, (Barnes & Filhol, 2023).

Data availability statement

570 The supporting data produced for this paper are openly available in the repository Aeteia/Ribbed-Moraine at <https://doi.org/10.5281/zenodo.7991094>, (Barnes & Filhol, 2023). The input data for this paper are freely available and can be found in the Norwegian national geospatial data archive at the following links: NiB (Kartverket, 2021a, c) <https://kartkatalog.geonorge.no/metadata/norge-i-bilder/e7cd5f9b-20e1-4f59-b379-64828cd38062>, national 1 m DEM (Kartverket, 2021b) <https://kartkatalog.geonorge.no/metadata/hoeyde-dtm1/0442c622-6639-4024-86be-846de8c15bb2>, lake database (NVE, 2023) <https://kartkatalog.geonorge.no/metadata/innsjoedatabase/823b8639-9a49-41bf-8571-3608435eb149>,

575 surface geology map (NGU, 2022) <https://kartkatalog.geonorge.no/metadata/loesmasser/3de4ddf6-d6b8-4398-8222-f5c47791a757>, Administrative borders (GISCO, 2020) <https://gisco-services.ec.europa.eu/distribution/v2/countries/>.

Author contributions

T. Barnes was the primary contributor to this paper in terms of each section of the CRediT guidelines. S. Filhol provided significant support in methodological development and refinement, including programming support and conceptualisation of method development – further S. Filhol provided review and editing support. T. V. Schuler provided conceptualisation and review / editing support, with methodological development guidance and project supervision. K. S. Lilleøren provided review / editing support, project supervision and methodology conceptualisation.

Competing interests

The authors declare no competing interests in the production of this manuscript.

585 Acknowledgements

We acknowledge L. S. Schmidt for her assistance in discussing possible ways of implementing our ideas in a programming sense, and computational support. We also acknowledge E. M. Lund for spellchecking the document.

Financial support

This study has been funded by Universitetet i Oslo. S. Filhol was funded from ClimaLand, an EEA/EU collaboration grant 590 between Norway and Romania 2014-2021, project code RO-NO-2019-0415,1290 contract no. 30/2020.

595

600

605

6 References

- Aario, R.: Classification and terminology of morainic landforms in Finland, *Boreas*, 6, 87–100, <https://doi.org/https://doi.org/10.1111/j.1502-3885.1977.tb00338.x>, 1977.
- 610 Anundsen, K.: Evidence of ice movement over southwest Norway indicating an ice dome over the coastal district of west Norway, *Quat. Sci. Rev.*, 9, 99–116, [https://doi.org/10.1016/0277-3791\(90\)90007-W](https://doi.org/10.1016/0277-3791(90)90007-W), 1990.
- Aydda, A., Althuwaynee, O. F., and Pokharel, B.: An easy method for barchan dunes automatic extraction from multispectral satellite data, *IOP Conf. Ser. Earth Environ. Sci.*, 419, <https://doi.org/10.1088/1755-1315/419/1/012015>, 2020.
- 615 Barnes, R.: RichDEM: terrain Analysis Software, <https://github.com/r-barnes/richtdem>, 2016.
- Barnes, T., Filhol, S., Ribbed-Moraine, <https://doi.org/10.5281/zenodo.7991094>, 2023
- Bhattacharjee, A., Murugan, R., and Goel, T.: A hybrid approach for lung cancer diagnosis using optimized random forest classification and K-means visualization algorithm, *Health Technol. (Berl.)*, 12, 787–800, <https://doi.org/10.1007/s12553-022-00679-2>, 2022.
- 620 Boulton, G. S., Horsfield, B., Clarke, C.: Large scale glacial geology and geomorphology of Canada and their implications for the history of the Laurentide ice sheet. INIQUA XII Int. Congr. Progr. Abstr., Ottawa, Ont. 135, 1987
- Bradski, G.: The OpenCV Library, Dr. Dobb's J. Softw. Tools, 2000.
- Breiman, L.: Random Forests, *Mach. Learn.*, 45, 5–32, <https://doi.org/10.1023/A:1010933404324>, 2001.
- Brodersen, K. H., Ong, C. S., Stephan, K. E., and Buhmann, J. M.: The balanced accuracy and its posterior distribution, *Proc. – Int. Conf. Pattern Recognit.*, 3121–3124, <https://doi.org/10.1109/ICPR.2010.764>, 2010.
- 625 Burney, S. M. A. and Tariq, H.: K-means Cluster Analysis for Image Segmentation, *Int. J. Comput. Appl.*, 96, 1–8, <https://doi.org/10.5120/16779-6360>, 2014.
- Butcher, F. E. G., Hughes, A. L. C., Ely, J. C., Christopher, D., Lewington, E. L. M., Boyes, B. M., Scoffield, A. C., Howcutt, S., and Dowling, T. P. F.: Reconstructing the flow evolution of the Fennoscandian Ice Sheet using new high-resolution digital elevation models., 45, 787263, 2021.
- 630 Böhner, J., Köthe, R., Conrad, O., Gross, J., Ringeler, A., and Selige, T.: Soil regionalisation by means of terrain analysis and process parameterisation, *Eur. Soil Bur.*, 213–222, 2001.
- Clubb, F. J., Bookhagen, B., and Rheinwalt, A.: Clustering River Profiles to Classify Geomorphic Domains, *J. Geophys. Res. Earth Surf.*, 124, 1417–1439, <https://doi.org/10.1029/2019JF005025>, 2019.
- 635 Conrad, O., Bechtel, B., Bock, M., Dietrich, H., Fischer, E., Gerlitz, L., Wehberg, J., Wichmann, V., and Böhner, J.: System for Automated Geoscientific Analyses (SAGA) v. 2.1.4, *Geosci. Model Dev.*, 8, 1991–2007, <https://doi.org/10.5194/gmd-8-1991-2015>, 2015.
- Corr, D., Leeson, A., McMillan, M., Zhang, C., Barnes, T.: An inventory of supraglacial lakes and channels across the West Antarctic Ice Sheet, *Earth System Science Data*, 14, 209–228, <https://doi.org/10.5194/essd-14-209-2022>, 2022.
- 640 Cowan, W.: Ribbed moraine: till fabric analysis and origin, *Can. J. Earth Sci.*, 5, 1145–1159, 1968.
- Dirscherl, M., Dietz, A. J., Kneisel, C., and Kuenzer, C.: Automated mapping of antarctic supraglacial lakes using a machine learning approach, *Remote Sens.*, 12, 1–27, <https://doi.org/10.3390/rs12071203>, 2020.
- Dunlop, P. and Clark, C. D.: The morphological characteristics of ribbed moraine, *Quat. Sci. Rev.*, 25, 1668–1691, <https://doi.org/10.1016/j.quascirev.2006.01.002>, 2006.
- 645 Dunlop, P., Clark, C. D., and Hindmarsh, R. C. A.: Bed ribbing instability explanation: Testing a numerical model of Ribbed moraine formation arising from couple flow of ice and subglacial sediment, *J. Geophys. Res. Earth Surf.*, 113, 1–15, <https://doi.org/10.1029/2007JF000954>, 2008.
- Eisank, C., Smith, M., and Hillier, J.: Assessment of multiresolution segmentation for delimiting drumlins in digital elevation models, *Geomorphology*, 214, 452–464, <https://doi.org/10.1016/j.geomorph.2014.02.028>, 2014.

- 650 Ely, J. C., Clark, C. D., Spagnolo, M., Stokes, C. R., Greenwood, S. L., Hughes, A. L. C., Dunlop, P., and Hess, D.: Do subglacial bedforms comprise a size and shape continuum?, *Geomorphology*, 257, 108–119, <https://doi.org/10.1016/j.geomorph.2016.01.001>, 2016.
- Evans, I. S.: Geomorphometry and landform mapping: What is a landform?, *Geomorphology*, 137, 94–106, <https://doi.org/10.1016/j.geomorph.2010.09.029>, 2012.
- 655 Finlayson, A. G. and Bradwell, T.: Morphological characteristics, formation and glaciological significance of Rogen moraine in northern Scotland, *Geomorphology*, 101, 607–617, <https://doi.org/10.1016/j.geomorph.2008.02.013>, 2008.
- Fisher, T. G., Shaw, J.: A depositional model for Rogen moraine with examples from the Avalon Peninsula, Newfoundland, *Canadian Journal of Earth Sciences*, 29,4, 669–686, <https://doi.org/10.1139/e92-058>, 1992.
- Fredin, O., Bergström, B., Eilertsen, R., Hansen, L., Longva, O., Nesje, A., and Sveian, H.: Glacial landforms and Quaternary landscape development in Norway, *Quat. Geol. Norw. Geol. Surv. Norw. Spec. Publ.*, 13, 5–25, 2013.
- 660 Frödin, G.: De sista skedena av Centraljämtlands glaciala historia, *Geogr. Uppsala Univ.*, 24, 1–251, 1954.
- Gentleman, R. and Carey, V.: Unsupervised Machine Learning, in: *Bioconductor Case Studies*, Cambridge, 137–157, <https://doi.org/10.1007/978-0-387-77240-0>, 2008.
- GISCO, Countries, 2020 – Administrative Units – Dataset, available at: <https://gisco-services.ec.europa.eu/distribution/v2/countries/>, 2020.
- 665 Guisan, Antoine, Weiss, Stuart B., and Weiss, A. D.: GLM versus CCA Spatial Modeling of Plant Species Distribution Author (s): Reviewed work (s): GLM versus CCA spatial modeling of plant species distribution, *Plant Ecol.*, 143, 107–122, 1999.
- Guitet, S., Cornu, J. F., Brunaux, O., Betbeder, J., Carozza, J. M., and Richard–Hansen, C.: Landform and landscape mapping, French Guiana (South America), *J. Maps*, 9, 325–335, <https://doi.org/10.1080/17445647.2013.785371>, 2013.
- 670 Hättestrand, C. and Kleman, J.: Ribbed moraine formation, *Quat. Sci. Rev.*, 18, 43–61, 1999.
- Hoppe, G.: Glacial Morphology and inland ice recession in northern Sweden, *Geogr. Ann.*, 41, 193–212, 1952.
- Hjort, J., Ujanen, J., Parviainen, M., Tolgensbakk, M., Etzelmüller, B.: Transferability of geomorphological distribution models: Evaluation using solifluction features in subarctic and Arctic regions., *Journal of Geomorphology* 204, 165–176, <https://dx.doi.org/10.1016/j.geomorph.2013.08.002>, 2014
- 675 Hong, C. S. and Oh, T. G.: TPR–TNR plot for confusion matrix, *Commun. Stat. Appl. Methods*, 28, 161–169, <https://doi.org/10.29220/CSAM.2021.28.2.161>, 2021.
- Horn, B. K. P.: Hill shading and the reflectance map. *Proceedings of the IEEE* 69, 14–47. <https://doi.org/10.1109/PROC.1981.11918>, 1981.
- 680 Iwahashi, J., Kamiya, I., Matsuoka, M., and Yamazaki, D.: Global terrain classification using 280 m DEMs: segmentation, clustering, and reclassification, *Progress in Earth and Planetary Science*, 1–31 pp., <https://doi.org/10.1186/s40645-017-0157-2>, 2018.
- Kartverket,Norge i Bilder, Norwegian mapping authority, Geovekst og kommunene (Nasjonal data), available at: http://opencache.statkart.no/gatekeeper/gk/gk.open_nib_utm33_wmts_v2?SERVICE=WMTS&REQUEST=GetCapabilities. 2021a.
- 685 Kartverket, Høydemodell DTM 1 m, Accessed: 2022.01.05, available at:<https://hoydedata.no/LaserInnsyn2/>. 2021b
- Kartverket, Norge I bilder WMS–Ortofoto. Trond Ola Ulvolden. <http://data.europa.eu/88u/dataset/dcee8bf4-fdf3-4433-a91b-209c7d9b0b0f> (Original work published 2016). 2021c.
- Kong, H., Akakin, H. C., & Sarma, S. E. A generalized laplacian of gaussian filter for blob detection and its applications. *IEEE Transactions on Cybernetics*, 43(6), 1719–1733. <https://doi.org/10.1109/TSMCB.2012.2228639>. 2013.
- 690 Kong, Y., Zeng, J., Wang, J., and Fang, Y.: A basin recognition method by landform classification and geometrical feature discrimination, *AIP Adv.*, 11, <https://doi.org/10.1063/5.0031695>, 2021.

- Likas, A., Vlassis, N., and J. Verbeek, J.: The global k-means clustering algorithm, *Pattern Recognit.*, 36, 451–461, [https://doi.org/10.1016/S0031-3203\(02\)00060-2](https://doi.org/10.1016/S0031-3203(02)00060-2), 2003.
- 695 Lindén, M., Möller, P., and Adrielsson, L.: Ribbed moraine formed by subglacial folding, thrust stacking and lee-side cavity infill, *Boreas*, 37, 102–131, <https://doi.org/10.1111/j.1502-3885.2007.00002.x>, 2008.
- Liu, C., Gu, Z., and Wang, J.: A Hybrid Intrusion Detection System Based on Scalable K-Means+ Random Forest and Deep Learning, *IEEE Access*, 9, 75729–75740, <https://doi.org/10.1109/ACCESS.2021.3082147>, 2021.
- Lundqvist, J.: Problems of the So-called Rogen Moraine, *Swedish Geol. Surv., Series C*, 32, 1969.
- 700 Lundqvist, J.: Rogen (ribbed) moraine-identification and possible origin, *Sediment. Geol.*, 62, 281–292, [https://doi.org/10.1016/0037-0738\(89\)90119-X](https://doi.org/10.1016/0037-0738(89)90119-X), 1989.
- Lundqvist, J.: Rogen moraine - An example of two-step formation of glacial landscapes, *Sediment. Geol.*, 111, 27–40, [https://doi.org/10.1016/S0037-0738\(97\)00004-3](https://doi.org/10.1016/S0037-0738(97)00004-3), 1997.
- Lv, Z., Liu, T., Shi, C., Benediktsson, J. A., and Du, H.: Novel Land Cover Change Detection Method Based on k-Means Clustering and Adaptive Majority Voting Using Bitemporal Remote Sensing Images, *IEEE Access*, 7, 34425–34437, <https://doi.org/10.1109/ACCESS.2019.2892648>, 2019.
- 705 Marutho, D., Hendra Handaka, S., Wijaya, E., and Muljono: The Determination of Cluster Number at k-mean Using Elbow Method and Purity Evaluation on Headline News, *Proc. - 2018 Int. Semin. Appl. Technol. Inf. Commun. Creat. Technol. Hum. Life, iSemantic 2018*, 533–538, <https://doi.org/10.1109/ISEMANTIC.2018.8549751>, 2018.
- 710 Moradi Fard, M., Thonet, T., and Gaussier, E.: Deep k-means: Jointly clustering with k-means and learning representations, *Pattern Recognit. Lett.*, 138, 185–192, <https://doi.org/10.1016/j.patrec.2020.07.028>, 2020.
- Möller, P.: Melt-out till and ribbed moraine formation, a case study from south Sweden, *Sediment. Geol.*, 232, 161–180, <https://doi.org/10.1016/j.sedgeo.2009.11.003>, 2010.
- Möller, P. and Dowling, T. P. F.: Equifinality in glacial geomorphology: instability theory examined via ribbed moraine and drumlins in Sweden, *Gff*, 140, 106–135, <https://doi.org/10.1080/11035897.2018.1441903>, 2018.
- 715 NVE, Innsjødatabase, Accessed 2023.01.01, available at: <https://www.nve.no/kart/kartdata/vassdragsdata/innsjodatabase/>. 2023.
- Ng, H. P., Ong, S. H., Foong, K. W. C., Goh, P. S., & Nowinski, W. L., Medical image segmentation using K-means clustering and improved watershed algorithm. *Proceedings of the IEEE Southwest Symposium on Image Analysis and Interpretation, 2006*, 61–65. <https://doi.org/10.1109/ssiai.2006.1633722>. 2006.
- 720 NGU, løsmasse, Accessed, 2022.08.01, available at: https://geo.ngu.no/kart/losmasse_mobil/. 2022.
- Patton, H., Hubbard, A., Andreassen, K., Auriac, A., Whitehouse, P. L., Stroeven, A. P., Shackleton, C., Winsborrow, M., Heyman, J., Hall, A. M.: Deglaciation of the Eurasian ice sheet complex, *Quat. Sci. Rev.*, 169, 148–172, <https://doi.org/10.1016/j.quascirev.2017.05.019>, 2017
- 725 Patton, H., Hubbard, A., Andreassen, K., Winsborrow, M., and Stroeven, A. P.: The build-up, configuration, and dynamical sensitivity of the Eurasian ice-sheet complex to Late Weichselian climatic and oceanic forcing, *Quat. Sci. Rev.*, 153, 97–121, <https://doi.org/10.1016/j.quascirev.2016.10.009>, 2016.
- Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M., Prettenhofer, P., Weiss, R., Dubourg, V., Vanderplas, J., Passos, A., Cournapeau, D., Brucher, M., Perrot, M., and Duchesnay, É.: Scikit-learn: Machine Learning in Python, *J. Mach. Learn. Res.*, 12, 2826–2830, 2011.
- 730 Ronneberger, O., Fischer, P., Brox, T.: U-Net: Convolutional Networks for Biomedical Image Segmentation. In: Navab, N., Hornegger, J., Wells, W., Frangi, A. (eds) *Medical Image Computing and Computer-Assisted Intervention – MICCAI 2015*. MICCAI 2015. Lecture Notes in Computer Science(), vol 9351. Springer, Cham. https://doi.org/10.1007/978-3-319-24574-4_28, 2015.

- 735 Saha, K., Wells, N. A., and Munro-Stasiuk, M.: An object-oriented approach to automated landform mapping: A case study of drumlins, *Comput. Geosci.*, 37, 1324–1336, <https://doi.org/10.1016/j.cageo.2011.04.001>, 2011.
- Sarala, P.: Ribbed moraine stratigraphy and formation in southern Finnish Lapland, *J. Quat. Sci.*, 21, 387–398, <https://doi.org/10.1002/jqs.995>, 2006.
- Singh, P.: Unsupervised Machine Learning, 161–181 pp., https://doi.org/10.1007/978-1-4842-4961-1_7, 2019.
- 740 Smith, M. J. and Clark, C. D.: Methods for the visualization of digital elevation models for landform mapping, *Earth Surf. Process. Landforms*, 30, 885–900, <https://doi.org/10.1002/esp.1210>, 2005.
- Sokolova, M., Japkowicz, N., and S, S.: Beyond Accuracy, F-Score and ROC: A Family of Discriminant Measures for Performance Evaluation, *Adv. Artif. Intell.*, 4304 LNAI, 1015–1021, 2006.
- Sollid, J. L. and Sorbel, L.: Distribution of Glacial Landforms in Southern Norway in Relation to the Thermal Regime of the Last Continental Ice Sheet, *Geogr. Ann. Ser. A, Phys. Geogr.*, 76, 25, <https://doi.org/10.2307/521317>, 1994.
- 745 Sommerkorn, J.: Distribution of ribbed moraines and their connection to subglacial water in the Oppland region of Norway, Universitetet i Oslo, 2020.
- Strøm, K. M.: The Geomorphology of Norway, *Geogr. Journal*, 112, 19–23, 1948.
- United Kingdom Environment Agency (UKEA). LiDAR Composite DTM 2022 – 1m, accessed 20.03.2023, available at: <https://www.data.gov.uk/dataset/01b3ee39-da3f-47b6-83da-dc98e73a461f/lidar-composite-dtm-2022-1m>. 2022.
- 750 Verstappen, H. T., Chapter Two – Old and New Trends in Geomorphological and Landform Mapping *Developments in Earth Surface Processes*, 15, pp. 13–38, <https://doi.org/10.1016/B978-0-444-53446-0.00002-1>. 2011.
- Zevenbergen, W. L., & Thorne, R. C. Quantitative analysis of land surface topography. *Earth Surface Processes and Landforms* 1, 12, 47–56. 1987.

755