

# Evaluation of downward and upward solar irradiances simulated by the Integrated Forecasting System of ECMWF using airborne observations above Arctic low-level clouds

Hanno Müller<sup>1</sup>, André Ehrlich<sup>1</sup>, Evelyn Jäkel<sup>1</sup>, Johannes Röttenbacher<sup>1</sup>, Benjamin Kirbus<sup>1</sup>, Michael Schäfer<sup>1</sup>, Robin J. Hogan<sup>2,3</sup>, and Manfred Wendisch<sup>1</sup>

<sup>1</sup>Leipzig Institute for Meteorology (LIM), Leipzig University, Leipzig, Germany

<sup>2</sup>European Centre for Medium-Range Weather Forecasts, Reading, United Kingdom

<sup>3</sup>Department of Meteorology, University of Reading, Reading, United Kingdom

**Correspondence:** Hanno Müller (hanno.mueller@uni-leipzig.de)

**Abstract.** The simulations of upward and downward irradiances by the Integrated Forecasting System (IFS) of the European Centre for Medium-Range Weather Forecasts are compared to broadband solar irradiance measurements from the Arctic CLOUD Observations Using airborne measurements during polar Day (ACLOUD) campaign. For this purpose, offline radiative transfer simulations with the ecRad radiation scheme using the operational IFS output were performed. The simulations of the downward solar irradiance agree within the measurement uncertainty. However, the IFS underestimates the reflected solar irradiances above sea ice significantly by  $-35 \text{ W m}^{-2}$ . Above open ocean, the agreement is closer with an overestimation of  $28 \text{ W m}^{-2}$ . A sensitivity study using measured surface and cloud properties is performed with ecRad to quantify the contributions of the surface albedo, cloud fraction, ice and liquid water path and cloud droplet number concentration to the observed bias. It shows that the IFS sea ice albedo climatology underestimates the observed sea ice albedo, causing more than 50 % of the bias. Considering the higher variability of in situ observations in the parameterization of the cloud droplet number concentration leads to a smaller bias of  $-27 \text{ W m}^{-2}$  above sea ice and a larger bias of  $48 \text{ W m}^{-2}$  above open ocean by increasing the range from  $36\text{-}69 \text{ cm}^{-3}$  to  $36\text{-}200 \text{ cm}^{-3}$ . Above sea ice, realistic surface albedos, cloud droplet number concentrations and liquid water paths contribute most to a bias improvement. Above open ocean, realistic cloud fractions and liquid water paths are most important to reduce the model-observation differences.

## 1 Introduction

The Arctic climate has changed more rapidly than the rest of the globe during the recent decades. One clear sign is the reduction of the sea ice extent of the Arctic ocean, particularly in September each year (Serreze and Meier, 2019). Another indicator is the increase of the near-surface air temperature that is in the Arctic more than twice as large as for the whole globe (Rantanen et al., 2022; Wendisch et al., 2023). The ongoing changes of the Arctic climate system emphasize a need for adaptations of forecast models to Arctic-specific peculiarities (Jung et al., 2016). Improved prediction systems for the Arctic would not only be a direct benefit for the future Arctic with its allowed shipping routes (Smith and Stephenson, 2013) but also an indirect benefit

for the forecast in Northern Hemisphere midlatitudes at longer lead times. This is caused by the linkage between the Arctic and midlatitudes that was investigated by e.g. Jung et al. (2014), Cohen et al. (2014), Overland et al. (2015) and Lawrence et al. (2019).

25 In the Arctic, Numerical Weather Prediction (NWP) models, such as the Integrated Forecasting System (IFS) by the European Centre for Medium-Range Weather Forecasts (ECMWF), often appear more uncertain compared to other regions on the globe (Bauer et al., 2016). The reasons for the lower predictive skills in the Arctic are various and often linked to the particularities of the Arctic climate system. One obvious issue in the Arctic results from sparse observational coverage, which limits data assimilation (Bauer et al., 2016; Jung and Matsueda, 2016; Lawrence et al., 2019; Ortega et al., 2022). Furthermore, modeling  
30 of the sea ice cover is a major obstacle for correctly representing the Arctic surface energy budget but is still uncertain due to the complexity of sea ice dynamics (Day et al., 2022). The representation of low-level Arctic clouds and especially mixed-phase clouds has been identified as another major source of uncertainty (Forbes and Ahlgrimm, 2014). As shown by Morrison and Pinto (2006), especially cloud microphysical schemes cause uncertainties in the cloud phase and precipitation.

Low-level clouds occur frequently in the Arctic (e.g. Eastman and Warren, 2010; Mioche et al., 2015) and show a pronounced  
35 longevity above sea ice and open ocean (Shupe et al., 2006; Verlinde et al., 2007). Their radiative properties are controlled by a complex system of coupled microphysical and dynamical processes that may differ depending on the surface conditions (Morrison et al., 2012; Wendisch et al., 2019). Especially for optical thin clouds, with a liquid water path (LWP) less than  $30 \text{ g m}^{-2}$ , the cloud radiative effect changes significantly for only small changes of cloud properties (Shupe and Intrieri, 2004). Thus, these clouds potentially introduce major model uncertainties. To constrain the effect of Arctic low-level clouds on the  
40 atmospheric radiation budget, it is necessary to identify shortcomings of microphysical parameterizations in NWP models to properly predict snow rate and cloud properties (Solomon et al., 2009). A substantial underprediction of cloud LWP together with an overprediction of cloud ice water path (IWP) was revealed by Solomon et al. (2009) and indicated an unrealistic growth of ice particles in the Weather Research Forecast model. Solomon et al. (2023) showed that contemporary models have difficulties to represent the radiative impact of Arctic clouds and still struggle to keep liquid water at low temperatures.  
45 Additionally, the change of the surface type (sea ice or open ocean) when clouds move on or off the sea ice initiates air mass transformations and changes of the cloud dynamics. This transition can result in cloud formation or cloud dissipation and is still poorly represented in NWP models (Pithan et al., 2018; Wendisch et al., 2021).

Two different concepts to evaluate clouds and their radiative effects in NWP have been applied in the past. The first approach applies model inter-comparisons (e.g. Klocke and Rodwell, 2014) to identify model uncertainties. This technique is not able  
50 to quantify potential model biases compared to reality. The second approach makes use of observed cloud properties. This approach was applied in the past decades to evaluate the representation of clouds in global NWP models using ground-based long-term cloud observations within the framework of Atmospheric Radiation Measurement (ARM) sites (e.g. Yang et al., 2006; Morcrette et al., 2012) or within the Cloudnet (Illingworth et al., 2007) framework (e.g. Hogan et al., 2009; Sinclair et al., 2022). Due to the nature of observations at a fixed location, only a few of these studies target specific Arctic sites  
55 (e.g. Klein et al., 2009; Morrison et al., 2009; Zhao and Wang, 2010; Forbes and Ahlgrimm, 2014). For an evaluation of the cloud representation in the central Arctic over the Arctic ocean, satellite observations can be utilized, which provide a

spatially broader view and come in polar regions together with frequent overpasses by polar-orbiting satellites, but cause difficulties in the data assimilation of microwave soundings above sea ice (Lawrence et al., 2019). Compared to the long-term observations of ARM, Cloudnet and satellites, shipborne observations provide short-term observations covering slowly varying different locations. Airborne observations bridge the gap between the ground-based or shipborne observations and the satellite observations and can provide in situ observations of cloud particle properties.

The efforts to improve model representations of Arctic clouds conducted in the past decades covered diverse aspects and quantities. The parameterization and representation of the sea ice albedo in various models was evaluated by e.g. Liu et al. (2007) and Karlsson and Svensson (2013), who identified the model sea ice albedo to determine both the sign and the amount of its cloud radiative effect. Low-level cloud fractions were assessed in reanalyses by Walsh et al. (2009) and found to be underestimated in summer, which leads to a bias in the solar radiation flux, while Sotiropoulou et al. (2016) evaluated the improvement of the representation of the vertical structure of mixed-phase clouds in IFS by the change from a diagnostic to a prognostic parameterization of mixed-phase clouds. Integrated microphysical quantities like LWP and IWP were investigated by Gu et al. (2021) who evaluated these quantities in Arctic reanalyses and found a mean underestimation of both LWP and IWP over the Arctic region compared to satellite observations. The representation of cloud droplet number concentrations in different models was evaluated by Geoffroy et al. (2010), Brenguier et al. (2011) and McCusker et al. (2023), who showed a slight improvement of the overestimation of the liquid cloud mass mixing ratio in low-levels clouds in the Met Office Unified Model (UM) by using representative cloud droplet number concentrations. Stevens et al. (2018) concluded from their model-intercomparison of cloud condensation nuclei-limited tenuous Arctic clouds that an appropriate treatment of the cloud droplet size distribution within models is important to account for aerosol-cloud interactions. Regarding the IFS, Beesley et al. (2000) evaluated the ECMWF model with observations collected during the Surface Heat Budget of the Arctic Ocean (SHEBA) campaign (Uttal et al., 2002) and identified a much larger observed fraction of liquid water clouds. Tjernström et al. (2021) evaluated the IFS with observations from the Arctic Ocean 2018 (AO2018) expedition (Vüllers et al., 2021) and revealed too high (near-)surface air temperatures in the IFS. McCusker et al. (2023) evaluated clouds during AO2018 within the IFS that overestimated cloud occurrence below 3 km. Forbes and Ahlgrimm (2014) revealed an underestimation of IFS cloud top albedo compared to observations from the Clouds and the Earth's Radiant Energy System (CERES) project. The bias is linked to an underestimation of liquid water content (LWC) near cloud tops, which results from the parameterization of the cloud phase based on diagnostic air temperature.

However, these evaluations are often based on remote sensing products, which themselves include major uncertainties mostly resulting from several assumptions in the retrieval algorithm, e.g. viewing geometries, instrument sensitivity or the ice crystal shape (Wendisch, 2005). Therefore, Formenti and Wendisch (2008) recommended to compare NWP models in the observational space of radiation, e.g. solar and thermal infrared radiation, radar and lidar reflectivities. Huang et al. (2017) used this approach to evaluate different global reanalyses like ECMWF Reanalysis - Interim and Climate Forecast System Reanalysis by the National Centers for Environmental Prediction, Matsui et al. (2014) and Berry et al. (2019) to evaluate Earth system models. Observations of airborne solar spectral irradiance have been used by Wolf et al. (2020) in combination with along-track radiative transfer simulations of the operational ecRad radiation scheme of ECMWF and a benchmark radiative transfer model.

Their analysis indicated that IFS underestimates the IWC in a frontal cloud system close to Iceland and that differences in the absorbing spectral band indicate deficiencies in the ecRad ice crystal optical properties. For the Arctic CLOUD Observations Using airborne measurements during polar Day campaign (ACLOUD; Wendisch et al., 2019), Kretzschmar et al. (2020) applied similar measurements and found a pronounced underestimation of the negative cloud radiative effect in the ICON model. This bias was traced back to the cloud condensation nuclei activation in the microphysical scheme. For a specific cloud case observed during ACLOUD, Ruiz-Donoso et al. (2020) investigated the thermodynamic phase of mixed-phase clouds as modeled by ICON large-eddy simulations and found that measured spectral radiances reveal an underestimation of the modeled ice crystal number concentration. Jäkel et al. (2019b) used ACLOUD observations to analyze the performance of the sea ice albedo scheme used in a regional coupled climate model and found an underestimation of the variability of the sea ice albedo caused by a biased surface albedo parameterization dependence on surface temperature. So far, the comprehensive ACLOUD data set has not been used for any IFS evaluation. While efforts have been made to include sea ice dynamics in IFS to tackle the high sea ice variability close to the sea ice edge (Keeley and Mogensen), the sea ice albedo in IFS is still based on climatological values with shortcomings identified by Pohl et al. (2020) using satellite observations.

In this paper, airborne radiation data from the ACLOUD campaign are used to evaluate the representation of Arctic low-level clouds and sea ice albedo in the IFS. The comparison is based on the observational space of solar irradiance and additionally implements active cloud remote sensing and in situ cloud microphysical observations. Section 2 describes the comparison strategy of observations and radiative transfer simulations, which are analyzed in Sect. 3. Based on the additional measurements, a sensitivity study is presented in Sect. 4. It aims at reducing the differences between modeled and observed irradiances by improving the surface albedo, cloud fraction and macro- and microphysical properties (LWP, IWP and cloud droplet number concentration) with actually measured data. The contributions of each individual parameter to the overall model uncertainty are summarized in the conclusions in Sect. 5.

## 2 Data and methods

### 2.1 Airborne observations

Airborne observations of the ACLOUD campaign (Ehrlich et al., 2019), which took place in May/June 2017 around Svalbard, Norway, provide a comprehensive data set of in situ and remote sensing observations for model evaluation and have been used in this analysis. Two upward and downward looking CMP22 pyranometers (spectral range of 300 nm to 3000 nm) aboard the Polar 5 aircraft (Wesche et al., 2016) measured the broadband solar irradiance (Stapf et al., 2019), which is referred to as solar irradiance in this study. The uncertainty of the CMP22 irradiance is typically about 2 % according to characterizations by Vuilleumier et al. (2014) for ground-based operations. However, airborne operation of the CMP22 in Arctic conditions may increase these uncertainties depending on solar zenith angle and environmental conditions, as described by Ehrlich et al. (2023) on a high altitude aircraft and Su et al. (2008) in a laboratory study and on active stabilization performance (Wendisch et al., 2001). The data are corrected for the aircraft specific operation as summarized in Ehrlich et al. (2019) including corrections for

aircraft attitude and instrument inertia. For the conditions during ACLOUD a maximum uncertainty of 3 % in regular straight  
125 flight sections is assumed.

Spectral solar irradiances are measured on Polar 5 with the Spectral Modular Airborne Radiation measurement system (SMART; Jäkel et al., 2019a; Wendisch et al., 2001) covering the spectral range between 345 nm and 2150 nm with a spectral resolution of 3-15 nm (Ehrlich et al., 2019). Additional remote sensing observations aboard Polar 5 include the Airborne Mobile Aerosol Lidar (AMALi) system (Stachlewska et al., 2010) and the Microwave Radar/radiometer for Arctic Clouds (MiRAC;  
130 Kliesch and Mech, 2019; Mech et al., 2019). AMALi is sensitive to both liquid and ice layers, while MiRAC is particularly sensitive to ice layers in the clouds. The cloud top altitude is derived from AMALi using a robust backscatter gradient approach that is retrieval-independent and is only based on the chosen instrument threshold (Kulla et al., 2021). For MiRAC a threshold of  $-30$  dBZ equivalent reflectivity factor is applied for identifying cloud particles above the height of 150 m. Altitudes below were not considered to exclude ground clutter. The uncertainty of the radar detector is given by Mech et al. (2019) as 0.5 dBZ,  
135 which only slightly affects the uncertainty in detecting cloud layers.

The Polar 6 aircraft was equipped with numerous in situ cloud probes. This study makes use of cloud particle number concentrations measured by the Small Ice Detector Mark 3 (SID-3; Schnaiter and Järvinen; Hirst et al., 2001; Vochezer et al., 2016). To characterize the surface conditions, sea ice concentration satellite data are retrieved from the Advanced Microwave Scanning Radiometer 2 (AMSR2) measurements (Melsheimer and Spreen, 2019; Spreen et al., 2008).

## 140 **2.2 Radiative transfer simulations**

### **2.2.1 Integrated Forecasting System**

The results of the simulations presented in this paper have been achieved with the 'Atmospheric Model high resolution' configuration (HRES) of the IFS of the ECMWF. Model cycle 43r1 was operational during the time of ACLOUD. A detailed description of the IFS can be found at <https://www.ecmwf.int/en/publications/ifs-documentation>. To evaluate the short term  
145 forecast and not the model initialisation, the 00 UTC runs with hourly forecast steps are used, issued about 12 hours before each flight. Prognostic variables from the IFS are available at 137 model levels. About 32 levels lay below a typical flight altitude of 3000 m with the highest vertical resolution of about 30 m close to the ground. The spacing between grid points of the longitude-latitude-grid is  $0.07^\circ$  both along the longitude and the latitude axis, resulting in a horizontal resolution of 1.4 km to 7.8 km in the campaign region. The surface type is classified as open ocean/sea ice when both the sea ice concentrations by  
150 AMSR2 and IFS are below 0.01 %/above 60 %.

### **2.2.2 ecRad radiation scheme**

The prognostic variables air pressure, air and skin temperature, specific humidity and cloud fraction from the IFS serve as direct input to the ecRad radiation scheme (Hogan and Bozzo, 2018). The ecRad version 1.4.0 is applied in an offline mode, which allows to run sensitivity studies. In addition, the required quantities liquid/ice cloud mass mixing ratios are calculated  
155 as sums of specific cloud liquid/ice water content and specific rain/snow water content. Similarly, the effective radii are not

prognostic variables in the IFS and need to be calculated consistently to the IFS. The definition of ice cloud effective radius follows the parameterization by Sun and Rikus (1999) and Sun (2001). The definition of liquid cloud effective radius in the IFS is based on the parameterization by Martin et al. (1994) with an adjustment by Wood (2000).

160 The IFS distinguishes over open ocean between a spectrally constant surface albedo value of 0.06 for diffuse radiation and a solar zenith angle dependent surface albedo given by Taylor et al. (1996) for direct radiation. Here, the open ocean albedo is approximated with the diffuse albedo only. The surface albedo used in ecRad is composed of this open ocean albedo and the sea ice albedo, which is based on the one-dimensional sea ice model by Ebert and Curry (1993) providing a monthly mean climatology of the spectral surface albedo in six solar bands (boundaries at 0.185, 0.250, 0.440, 0.690, 1.190, 2.38 and 4.0  $\mu\text{m}$ ). This climatology is interpolated to the day of the specific flight. The surface type composition is obtained from the prognostic sea ice cover in the IFS.

170 Mass mixing ratios of  $\text{CH}_4$ ,  $\text{CO}$ ,  $\text{NO}_2$  and eleven different hydrophilic and hydrophobic aerosol species from the Copernicus Atmosphere Monitoring Service (CAMS) global reanalysis (EAC4) were extracted from the CAMS Atmosphere Data Store (Inness et al., 2019). Similarly, volume mixing ratios of  $\text{CO}_2$  (Chevallier et al., 2010, 2019) and  $\text{N}_2\text{O}$  (Thompson et al., 2011) from the CAMS global inversion-optimised greenhouse gas fluxes and concentrations product were included. Ozone concentrations were obtained from operational ozone soundings above Ny-Ålesund, Svalbard. The TOA solar irradiance of  $1360.8 \text{ W m}^{-2}$  (Kopp and Lean, 2011) is adjusted to the Earth-Sun-distance from noon of every flight day.

175 In the operational configuration, ecRad uses the McICA radiative transfer solver (Pincus et al., 2003). However, this solver does not provide spectrally resolved irradiances across the vertical column, which is needed for a direct comparison in flight altitude. Therefore, the operational solver is replaced by the Tripleclouds solver (Shonk and Hogan, 2008). A comparison of surface irradiances (not shown here) showed that both solvers do not significantly differ. The exponential-random cloud overlap assumption is applied in the Tripleclouds solver. Cloud overlap is parameterized by the overlap decorrelation length, which is calculated after Shonk et al. (2010), Eq. 13. The aerosol scattering properties are based on the IFS version cycle 43r1, in combination with the operational aerosol type classification from cycle 43r3. For the ice crystal optical properties, the operational parameterization from Fu (1996) and Fu et al. (1998) is chosen. The used gas absorption model is based on the Rapid Radiative Transfer Model for GCMs (RRTM-G; Mlawer et al., 1997) and defines the spectral resolution of ecRad in terms of 14 shortwave bands. Running ecRad in the described configuration provides spectral upward and downward irradiances at the interfaces of the 137 full levels in the 14 shortwave bands, which are then integrated to broadband irradiances.

### 2.3 Considering the scale mismatch

185 For the comparison between measurements and simulations, the aircraft is assumed to artificially fly through the model grid space. For this purpose, the different spatial scales of airborne observations and simulations have to be considered. The mean horizontal grid spacing of the simulations is in the range of 4.6 km. The time Polar 5 needs to fly between two grid points accounts for about 60s with an average speed of  $80 \text{ m s}^{-1}$ . Therefore, the airborne data are averaged over 60s and ecRad is run every 60s at the mean aircraft location during the corresponding averaging track interval. The ecRad input is extracted from IFS according to the closest grid box to the mean position of Polar 5 and to the nearest 1 hour IFS time step to the 60s interval.

190 This results in a maximum time offset of 30 minutes between simulation and observation. Temporal interpolation of the IFS output was deliberately omitted to avoid smeared states in ecRad input variables. Similarly, without interpolation, the ecRad output at the closest model level to the flight altitude is selected for the comparison.

The statistical comparison between observations and simulations is done using frequency distributions of solar irradiance. This additionally accounts for spatial and temporal mismatches, which would be present in a point-by-point comparison. The frequency distributions are compared using two quantities. On the one hand, the deviation  $\Delta F$  of their mean values is calculated  
 195 via

$$\Delta F = \bar{P}_{\text{ecRad}} - \bar{P}_{\text{obs}}, \quad (1)$$

where  $\bar{P}_{\text{ecRad}}$  and  $\bar{P}_{\text{obs}}$  are the means of the number frequency distributions of solar irradiances  $P_{\text{ecRad}}$  and  $P_{\text{obs}}$  from ecRad and the observations. On the other hand, the Hellinger distance  $\mathcal{H}$  (Hellinger, 1909) is used as a metric to include the shape of  
 200 the frequency distributions in the comparison and is calculated following


$$\mathcal{H}(P_{\text{obs}}, P_{\text{ecRad}}) = \frac{1}{\sqrt{2}} \sqrt{\sum_{i=1}^n (\sqrt{p_{\text{obs},i}} - \sqrt{p_{\text{ecRad},i}})^2}, \quad (2)$$

with  $P_{\text{obs}} = (p_{\text{obs},1}, \dots, p_{\text{obs},n})$  and  $P_{\text{ecRad}} = (p_{\text{ecRad},1}, \dots, p_{\text{ecRad},n})$ . The index identifies the center of each bin.  $\mathcal{H}$  ranges from 0 to 1, where a value of 0 corresponds to identical distributions and a value of 1 characterizes fully independent distributions. In the following, both described quantities are accompanied by arrows ( $\uparrow, \downarrow$ ), indicating the upward or downward  
 205 direction.

### 3 Comparison of simulated and measured solar irradiances

A comparison is carried out between simulated and measured solar irradiances in order to quantify the representation and its uncertainty of Arctic low-level clouds in the IFS. To achieve this, the analysis is limited to scenes when no higher clouds located between the flight level of Polar 5 and top of atmosphere (TOA) are present. This condition needs to hold for both observations  
 210 and simulations and guarantees, that the reflected upward solar irradiance is only affected by possible clouds below the aircraft and not contaminated by attenuation of the incoming irradiance. Scenes are identified as cloud-free above Polar 5 when the standard deviation of the CMP22 downward solar irradiance within a 60s interval does not exceed the mean value by 0.7 %. Cloud-free conditions in the IFS are given, when the sum of the fraction of cloud cover in all model levels above the aircraft flight level is below 0.02. These thresholds reliably exclude mid-level and cirrus clouds from the analysis. The analysis is  
 215 further limited to periods when all remote sensing instruments provided data so that the retrieval of cloud fractions and of LWP above open ocean is available. The filtering results in 501 scenes (60s intervals) above sea ice and 210 scenes (60s intervals) above open ocean contributed by nine out of 19 research flights, which are shown as flight tracks in Fig. 1. All scenes lay west of Svalbard with the majority above sea ice with a relatively high sea ice concentration.

Figure 2 shows the frequency distributions of upward and downward solar irradiances measured by the CMP22 pyranometer  
 220 and simulated by ecRad separated for sea ice and open ocean. The downward irradiances cover the range from  $440 \text{ W m}^{-2}$  to




**Figure 1.** Sections of flight tracks of nine research flights that are included in the analysis after filtering. The mean sea ice concentration during the ACLOUD campaign derived from AMSR2 is shown in the background layer.

670  $W m^{-2}$ . The lower irradiances above open ocean result from the larger solar zenith angles during these flight sections in the morning after takeoff and in the afternoon before landing. There is good agreement between the simulated and observed distribution of downward irradiances with  $\Delta F^\downarrow = -14 W m^{-2}$  above sea ice (Fig. 2a) and  $\Delta F^\downarrow = -8 W m^{-2}$  above open ocean (Fig. 2b), which is within the 3% maximum uncertainty of the CMP22 measurements. The corresponding  $\mathcal{H}^\downarrow$  are calculated with 64 bins of  $10 W m^{-2}$  width from  $35 W m^{-2}$  to  $665 W m^{-2}$  and are 0.42 above sea ice and 0.37 above open ocean.

Observations of upward irradiance above sea ice surfaces range between  $300 W m^{-2}$  and mainly  $530 W m^{-2}$  (Fig. 2c). The simulations show a similar amount of low irradiances but end abruptly at  $450 W m^{-2}$ . This upper limit in the IFS seems to be limited to clouds over sea ice. While the distribution of upward irradiances above sea ice is relatively narrow due to the high albedo of the sea ice reducing the cloud radiative effect, the distribution of the upward irradiances above the open ocean with its dark surface and low surface albedo is broader (Fig. 2d). It covers a range of irradiances from  $150 W m^{-2}$  to  $470 W m^{-2}$  in the simulations and from  $30 W m^{-2}$  to mainly  $510 W m^{-2}$  in the observations. The low values of the measurements result from a combination of higher solar zenith angles, which are frequently present above open ocean, and scenes without any clouds below the Polar 5 where the dark open ocean absorbs the major part of the incoming solar radiation. High values correspond to cloudy scenes reflecting a large amount of the incoming solar irradiance (Fig. 2d). Over ocean, higher upward irradiances are





**Figure 2.** Distribution of (a,b) downward and (c,d) upward solar irradiances for above Polar 5 cloud-free scenes measured by the CMP22 and simulated by ecRad above (a,c) sea ice and (b,d) open ocean. The values in the corner indicate the difference of the mean irradiances between ecRad and CMP22, the corresponding  $\mathcal{H}^{\uparrow,\downarrow}$  and the number of included scenes.

simulated, despite the lower surface albedo. The means above sea ice show a bias of  $\Delta F^\uparrow = -35 W m^{-2}$  with a  $\mathcal{H}^\uparrow$  of 0.48 and above ocean a bias of  $\Delta F^\uparrow = 28 W m^{-2}$  with a  $\mathcal{H}^\uparrow$  of 0.36.

While the magnitudes of  $\Delta F^\downarrow$  are not significant,  $\Delta F^\uparrow$  exceed the measurement uncertainty and suggest that either surface or cloud properties are not represented correctly in the IFS.

**Table 1.** Overview of performed sets of ecRad simulations indicating which parameter was adjusted by which source.

ecRad run	adjusted parameter	source
1 (reference)	-	IFS cy43r1
2	$\alpha$	SMART
3	$f_{\text{cloud}}$	AMALi/MiRAC
4	IWP	non-observation based
5	LWP	MiRAC
6	LWP	non-observation based
7	$N_d$	SID-3

#### 240 4 Sensitivity study

There are numerous possible contributors to the observed bias of the reflected solar irradiance. In principle, the radiative transfer and thus, the reflected solar irradiance is mostly affected by the surface albedo, the cloud fraction and the optical depth of the cloud, neglecting the minor impact of atmospheric gases and aerosols. Following Kokhanovsky (2004), the optical depth  $\tau$  is related to the cloud properties LWP or LWC, particle effective radius  $r_{\text{eff}}$  and density of water  $\rho_w$  via


$$245 \quad \tau = \frac{3}{2} \cdot \frac{1}{\rho_w} \cdot \frac{\text{LWP}}{r_{\text{eff}}} = \frac{3}{2} \cdot \frac{1}{\rho_w} \cdot \frac{\int_z \text{LWC} dz}{r_{\text{eff}}}, \quad (3)$$

with a vertical integration over the altitude  $z$ . However, the optical depth is neither a direct user variable in IFS nor in ecRad. According to the IFS documentation, the mean liquid effective radius  $r_{\text{eff}}$  is parameterized following a variation from Martin et al. (1994) by

$$r_{\text{eff}} = \left( \frac{3 E_d (\text{LWC} + \text{RWC})}{4 \pi \rho_w k N_d} \right)^{\frac{1}{3}}, \quad (4)$$

250 where  $E_d$  is an enhancement factor considering an increased dispersion of the droplet size spectrum (Wood, 2000), LWC and RWC are the liquid and rain water content,  $k$  is a factor depending on the relative dispersion of the cloud droplet spectrum set to 0.77 above ocean and  $N_d$  is the cloud droplet number concentration.  $N_d$  is parameterized via the aerosol number and mass concentrations as a function of prognostic 10-m wind speed accounting for the injection of sea spray aerosols from the ocean (Martin et al., 1994; Boucher and Lohmann, 1995; Lowenthal et al., 2004; Erickson et al., 1986; Genthon, 1992).

255 These dependencies of the cloud radiative properties and the reflected irradiance finally suggest a sensitivity study testing the contribution of the individual parameters to the observed bias  $\Delta F^\uparrow$ . For the sensitivity runs, the IFS input to ecRad is adjusted for surface albedo, cloud fraction, LWP, IWP and  $N_d$  individually based on observations where possible. The sources of the observed parameters are listed in Table 1 and described below. The reference case is identical to the simulations shown in Sect. 3, where the operational IFS output is fed to ecRad to simulate the solar irradiances.



**Figure 3.** (a) Time series of modes of measured sea ice albedo in the 400-690 nm, 690-1190 nm and 1190-2155 nm band (circles) and IFS sea ice albedo climatology (solid lines). The dashed lines show the parameterization for the measurements. (b) Distribution of upward solar irradiances for above Polar 5 cloud-free scenes above sea ice measured by the CMP22 (blue) and simulated by ecRad with the adjusted sea ice albedo (orange) together with the reference simulations (grey).

In the area covered by the ACLOUD campaign the surface albedo conditions were rather constant above open ocean but more variable above sea ice, which was affected by the melt season. Therefore, this sensitivity run is limited to the observations over sea ice. A realistic constraint of the sea ice albedo is deduced from SMART albedometer measurements from low-level flight sections. Wavelength ranges from 400-690 nm, 690-1190 nm and 1190-2380 nm are chosen for a wavelength band approach.

265 Figure 3a shows sea ice albedos from measurements of all below-cloud flight sections at flight altitudes below 300 m over sea ice ( $\alpha_{obs}$ , mode values) together with the IFS sea ice albedo climatology in the different bands. The influence of the season is obvious, as the measured sea ice albedo values decrease, mainly because of snow metamorphism to larger grain sizes due to the increase in skin temperature, accumulating liquid water in the snow layer and formation of the surface scattering layer (Rosenburg et al., 2023). The IFS sea ice albedo climatology assumes a slower melting season in bands 2, 3 and 5. It

270 underestimates the surface albedo in bands 2 and 3 at the beginning and overestimates it at the end of the campaign, while there is an underestimation in bands 4 and 5 during the whole campaign. These findings support the shortcomings identified by Pohl et al. (2020) with climatologically fixed transitions between the dry snow, melting snow and bare sea ice albedo from Ebert and Curry (1993).

The impact of the faster sea ice albedo reduction and the underestimation of the sea ice albedo on the irradiances is investigated by adjusting the sea ice albedo climatology in a set of ecRad simulations. Linear regressions of the measured sea ice

275 albedo in the three SMART wavelength ranges are used to estimate  $\alpha_{obs}$  at each flight day. The following adjustments are

**Table 2.** Differences  $\Delta F^\uparrow$  of the mean simulated and observed upward solar irradiance distributions of ecRad simulations and CMP22 observations and their corresponding  $\mathcal{H}^\uparrow$  for all sets of simulations.

ecRad run		sea ice		open ocean	
		$\Delta F^\uparrow$ ( $\text{W m}^{-2}$ )	$\mathcal{H}^\uparrow$	$\Delta F^\uparrow$ ( $\text{W m}^{-2}$ )	$\mathcal{H}^\uparrow$
1	(reference)	-35	0.48	28	0.36
2	$\alpha$	-16	0.35	28	0.36
3	$f_{\text{cloud}}$	-35	0.48	18	0.35
4	IWP -50 %	-35	0.48	27	0.35
4	IWP +50 %	-35	0.48	30	0.38
5	LWP <sub>obs</sub>	-	-	-28	0.42
6	LWP -50 %	-45	0.54	-5	0.39
6	LWP +50 %	-28	0.47	47	0.39
7	$N_d$	-26	0.41	48	0.38

made to the spectral albedo bands in ecRad:

$$\alpha_{\text{IFS,Band } 2/3} = \alpha_{\text{IFS},250-440\text{ nm}/440-690\text{ nm}} = \alpha_{\text{obs},400-690\text{ nm}} = 1.757 - 0.005 \cdot \text{DOY} \quad (5)$$

$$\alpha_{\text{IFS,Band } 4} = \alpha_{\text{IFS},690-1190\text{ nm}} = \alpha_{\text{obs},690-1190\text{ nm}} = 1.896 - 0.007 \cdot \text{DOY} \quad (6)$$


$$280 \quad \alpha_{\text{IFS,Band } 5} = \alpha_{\text{IFS},1190-2380\text{ nm}} = \alpha_{\text{obs},1190-2155\text{ nm}} = 1.367 - 0.007 \cdot \text{DOY}, \quad (7)$$

where DOY is the day of the year. Bands 2 and 3 are considered together due to the sparse SMART coverage of band 2, bands 1 and 6 are kept unchanged as they lie out of the SMART wavelength range.

The results of the modified ecRad run are shown in Fig. 3b and compared to the reference run and the observations. Due to the higher sea ice albedo especially for band 4, the sensitivity run results on average in higher upward solar irradiances. In comparison to the reference distribution, the adjusted distribution emerges with  $320 \text{ W m}^{-2}$  at a  $10 \text{ W m}^{-2}$  higher upward irradiance, but ends  $20 \text{ W m}^{-2}$  higher at  $470 \text{ W m}^{-2}$ . The distribution itself is shifted to higher values throughout all upward irradiances with the highest peak located between  $420-430 \text{ W m}^{-2}$ . The bias  $\Delta F^\uparrow$  decreases accordingly from  $-35 \text{ W m}^{-2}$  to  $-16 \text{ W m}^{-2}$ . The corresponding  $\mathcal{H}^\uparrow$  decreases from 0.48 to 0.35. These and all following values from the subsequent ecRad runs are summarized in Table 2. Thus, the replacement of the original sea ice albedo reduces the gap between the simulated and the observed irradiances by more than 50 %. This indicates, that the representation of the sea ice albedo in the IFS causes one major part of the disagreement. Another major part may be caused by the representation of clouds.

## 4.2 Cloud fraction

The cloud fraction of the IFS is compared to airborne remote sensing observations. A lidar-based cloud mask from the AMALi cloud top altitude product (Kulla et al., 2021) is used in combination with a radar-based cloud identification from MiRAC. Merging both types of cloud identification leads to a remote sensing based cloud fraction  $f_{\text{cloud,RS}}$  that accounts for the



**Figure 4.** Two-dimensional frequency distributions of IFS cloud fractions and observational cloud fractions based on AMALi and MiRAC above (a) sea ice and (b) open ocean with the upward irradiance differences between ecRad simulations and CMP22 observations as circle colors.


different sensitivities of radar and lidar. This cloud fraction is calculated from 60s flight sections. As these flight sections cover only a cross section of a grid box, there is an imbalance between observed variability of cloud fraction and the grid box mean. Especially low cloud fractions are less likely in IFS. Separated for sections above sea ice and open ocean, Fig. 4 compares the combinations of observed and forecasted cloud fractions combined with the corresponding mean upward irradiance differences between ecRad simulations and observations. An ideal representation of the observed clouds in the IFS would entail all data circles to lie on the dashed diagonal line with white color indicating no bias in the observed and simulated solar irradiances. However, especially above open ocean, the remote sensing cloud fraction covers the whole range from cloud-less to overcast conditions, while the IFS shows only little variability with cloud fractions ranging between 60 % and 100 %. The data below the dashed diagonal correspond to an overestimation of the cloud fraction by the IFS, which causes the positive irradiance differences to dominate. In theory, the data above the 1:1 line in Fig. 4b correspond to an underestimation of the cloud fraction by the IFS, coming along with negative irradiance differences indicated by orange colors, and vice versa. Data that do not follow this theory are likely caused by the possible imbalance of cloud fractions induced by comparing the entire grid box with the aircraft transect. Data points, where the cloud fractions agree are mostly observed for overcast conditions. In this case, the bias of the ecRad simulations is negative above sea ice and positive above open ocean.

A set of ecRad simulations is performed where the prognostic cloud fraction  $f_{\text{cloud, IFS, level}}$  is replaced by the observations taking into account the vertical distribution of the clouds. As a basic approach, the IFS cloud profiles are kept constant. To account for maximum overlap, this approach ensures that the cloud fraction at the level where it has the profile's maximum cloud fraction is replaced by the remote sensing cloud fraction. All other levels are scaled accordingly, adopting the original

shape of the cloud fraction profile. This is realized by replacing  $f_{\text{cloud,IFS,level}}$  with  $f'_{\text{cloud,level}}$  calculated via

$$315 \quad f'_{\text{cloud,level}} = f_{\text{cloud,RS}} \cdot \frac{f_{\text{cloud,IFS,level}}}{f_{\text{cloud,IFS,max}}}, \quad (8)$$

where  $f_{\text{cloud,IFS,max}}$  is the maximum cloud fraction of all 137 model levels.



**Figure 5.** Distribution of upward solar irradiances above (a) sea ice measured by the CMP22 (blue) and simulated by ecRad with the adjusted cloud fractions (orange) together with the reference simulations (grey) and above (b) open ocean measured by the CMP22 (green) and simulated by ecRad with the adjusted cloud fractions (black) together with the reference simulations (grey).

Figure 5 compares the irradiance distributions from the ecRad simulations with the replaced cloud fraction to the reference run and the observations. Above sea ice, the simulated irradiance distribution does not change significantly. Due to the high surface albedo small changes of  $f_{\text{cloud}}$  do not significantly reduce the reflected radiation. Thus,  $\Delta F^{\uparrow}$  remains at  $-35 \text{ W m}^{-2}$  with  $\mathcal{H}^{\uparrow}$  remaining at 0.48. Above open ocean, occurrences of upward solar irradiance between  $200 \text{ W m}^{-2}$  and  $460 \text{ W m}^{-2}$  are mainly lower compared to the reference distribution. This reduction enables a new mode occurring between  $40 \text{ W m}^{-2}$  and  $50 \text{ W m}^{-2}$ , so the replacement by the observed cloud fraction results in a higher amount of data with low reflected irradiances. This results from the overestimation of prognostic  $f_{\text{cloud}}$  that may be linked to broken cloud conditions that cannot be resolved by the IFS.  $\Delta F^{\uparrow}$  is reduced by 36 % from  $28 \text{ W m}^{-2}$  to  $18 \text{ W m}^{-2}$  with a corresponding  $\mathcal{H}^{\uparrow}$  decrease from 0.36 to 0.35 (see  
325 Table 2).

### 4.3 Macro- and microphysical cloud properties

#### 4.3.1 Ice water path

The low-level clouds observed during ACLOUD are mostly of mixed-phase character although dominated by liquid droplets (Ruiz-Donoso et al., 2020; Klingebiel et al., 2023). To test the relevance of the representation of ice crystals in the IFS to the  
330 cloud-reflected solar irradiance, no direct observations are available from ACLOUD. Therefore, the prognostic IWP in terms

of the specific cloud ice water content is both increased and reduced on a theoretical basis by 50 % in two sets of simulations. This increase (reduction) of IWP is propagated to an increase (reduction) of the ice effective radius according to Sun and Rikus (1999) and Sun (2001). Over sea ice, the simulated upward irradiance did not change. Above open ocean, the mean simulated irradiance is only increased by  $1 \text{ W m}^{-2}$  when the IWP is increased by 50 % and is only reduced by  $2 \text{ W m}^{-2}$  when the IWP  
 335 is reduced by 50 %. Thus, more cloud ice increases the bias of the irradiance simulations and less cloud ice reduces the bias. These small effects confirm the relatively low IWP during ACLOUD reported by Klingebiel et al. (2023) and indicate that the cloud droplets dominate the cloud radiative properties. Here, ice crystals may not directly cause the bias between simulated and observed irradiances. Similarly, ice crystal shape and size will not significantly impact the irradiance reflected by these liquid dominated clouds (Ehrlich et al., 2008). Also the choice of the ice optics parameterization within ecRad can impact the  
 340 reflected irradiance (as shown by e.g. Wolf et al. (2020)). However, a model-observations-comparison in this regard would require an agreement between the IWC in the IFS and the observations. This agreement cannot be verified with the Polar 5 instrumentation.


### 4.3.2 Liquid water path

To adjust the prognostic LWP in the IFS with observations, LWP measurements derived from passive microwave remote  
 345 sensing observations on Polar 5 are applied. However, the LWP product by the passive 89 GHz channel from MiRAC (Kliesch and Mech, 2019) is only available above open ocean. Above sea ice with its high emissivity the retrieval sensitivity is not sufficient, which is why this sensitivity study is firstly limited to open ocean. To confront the observed LWP above open ocean with the IFS output, the prognostic liquid cloud mass mixing ratio is converted to LWC and vertically integrated between the surface and Polar 5 flight altitude to the IFS LWP.

350 The combinations of the observed and the prognostic LWP are shown for the 210 scenes in Fig. 6. Observations are again 60s averages of LWP including cloud-free data and IFS values are the grid box mean all-sky LWP. They reveal the tendency of the IFS to predict too high LWP, while the observations rarely show a LWP above  $150 \text{ g m}^{-2}$ . This point-by-point mismatch indicates that cloud heterogeneity is high for the observed clouds. This is typical for low-level clouds over open ocean (Schäfer et al., 2018), especially when linked to cold air outbreaks. Similar to the discussion in Sect. 4.2, this mismatch partly results  
 355 from the imbalance of Polar 5 sampling along a straight flight leg and IFS providing grid box means. The exact position of the horizontal cloud structures cannot be forecasted precisely and also may change within the time offset between observations and IFS output. However, as shown in Fig. 6, the differences of irradiance correlate with the mismatch in LWP. Even within one single research flight both overestimations and underestimations of the observed LWP by a factor of two occur.

A set of ecRad simulations is performed with adjusted LWP. The specific cloud liquid water content  $q_{\text{liq}}$  in ecRad is replaced  
 360 with

$$q'_{\text{liq}} = q_{\text{liq}} \cdot \frac{\text{LWP}_{\text{obs}}}{\text{LWP}_{\text{IFS}}}, \quad (9)$$




**Figure 6.** Combinations of prognostic IFS LWP and observed LWP based on MiRAC with classes of absolute frequency as circle size and upward solar irradiance differences between ecRad simulations and CMP22 observations as circle colors.

where  $LWP_{\text{obs}}$  is the ACLOUD LWP and  $LWP_{\text{IFS}}$  the prognostic integrated LWP, so that the LWC profile shape is kept. The liquid effective radius is recalculated respectively, considering the changed LWC in Eq. 4. After adjusting,  $r_{\text{eff}}$  is limited to 4–30  $\mu\text{m}$  to match the IFS constraints again.  $N_d$  remains fixed.

365 The distributions of the upward solar irradiance for the observations, the reference simulations and the adjusted simulations with replaced LWP and  $r_{\text{eff}}$  are shown in Fig. 7. Compared to the reference distribution, the adjusted distribution emerges already at 80  $\text{W m}^{-2}$  instead of 150  $\text{W m}^{-2}$  and ends at upward irradiances 20  $\text{W m}^{-2}$  lower than before. The main mode ranges between 230  $\text{W m}^{-2}$  and 290  $\text{W m}^{-2}$  instead of between 320  $\text{W m}^{-2}$  and 340  $\text{W m}^{-2}$ . Adjusting the LWP based on observations leads to a change in the correct direction by reducing the upward solar irradiances. Compared to the observations, 370 this impact is too strong resulting in a conversion of the overestimation to an underestimation. The adjustment overcompensates the reference bias between the simulated and the observed distribution of upward irradiances with  $\Delta F^\uparrow = -28 \text{ W m}^{-2}$  and the corresponding  $\mathcal{H}^\uparrow$  increase is included in Table 2.

To quantify the impact of LWP uncertainties not only for clouds above open ocean but also above sea ice, the prognostic LWP is both increased and reduced artificially by 50% in two sets of simulations, going along with the according increase and decrease of  $r_{\text{eff}}$ .  $N_d$  remains fixed.  $\Delta F^\uparrow$  changes above sea ice from  $-35 \text{ W m}^{-2}$  to  $-45 \text{ W m}^{-2}$  by the reduction of the LWP and to  $-28 \text{ W m}^{-2}$  by the increase of the LWP. This qualitatively matches the findings from Solomon et al. (2023) that IFS produces too small LWP in the central Arctic. The bias  $\Delta F^\uparrow$  above open ocean changes from 28  $\text{W m}^{-2}$  in the reference case to  $-5 \text{ W m}^{-2}$  by the reduction of the LWP and to 47  $\text{W m}^{-2}$  by doubling the LWP. This indicated that the adjustments of the IFS need to be different above sea ice compared to open ocean to match the observations during the ACLOUD campaign. 380 Above sea ice, the increase of the LWP and the implicit change of  $r_{\text{eff}}$  improves  $\Delta F^\uparrow$  (Fig. 8a) by a slightly higher emergence






**Figure 7.** Distribution of upward solar irradiances above open ocean measured by the CMP22 (green) and simulated by ecRad (black) with the adjusted LWP and liquid effective radius based on the MiRAC LWP scaling of the prognostic LWP together with the reference simulations (grey).


of the upward irradiance distribution at  $320 \text{ W m}^{-2}$ , by slightly higher irradiances over a wide range of the distribution and by keeping the same end of the distribution as in the reference case. Above open ocean, the decrease of the LWP and the implied  $r_{\text{eff}}$  improves  $\Delta F^\uparrow$  (Fig. 8b) by shifting the entire distribution to  $20\text{--}30 \text{ W m}^{-2}$  lower upward irradiances. Possible reasons for the different LWP change directions may lie in differences of cloud physics between sea ice and open ocean. Above  
 385 open ocean, especially during cold air outbreaks, turbulent surface fluxes of sensible and latent heat are magnitudes larger than above sea ice (e.g. Hartmann et al., 1997) and cloud droplet number concentrations are known to differ (Young et al., 2016). The changes shown by the sensitivity study match qualitatively the findings of Moser et al. (2023) with higher  $N_d$  above sea ice than above open ocean, assuming a linear relation between LWC and  $N_d$  as found by Leaitch et al. (2016) and Dionne et al. (2020). Although the bias  $\Delta F^\uparrow$  above open ocean is largely reduced,  $\mathcal{H}^\uparrow$  is increased to 0.39 due to significant changes in the  
 390 shape of the irradiance distribution, especially at the upper end where the sharp cut-off of the highest irradiances is reduced from  $470 \text{ W m}^{-2}$  to  $440 \text{ W m}^{-2}$  because of the lower LWP.

### 4.3.3 Cloud droplet number concentration

The cloud droplet number concentration affects the cloud radiative properties by occurring in Eq. 4. The parameterized  $N_d$  in the IFS is compared to in situ observations available from the SID-3 cloud probe aboard Polar 6 (Schnaiter and Järvinen).  
 395 However, the flight track of Polar 5 does not always match the path of Polar 6. Nevertheless on a statistical basis, Polar 5



**Figure 8.** Distribution of upward solar irradiances measured by the CMP22 (blue) and simulated by ecRad (a) above sea ice with a 50% increased LWP and (b) above open ocean with a 50% decreased LWP with subsequent adjustments to the liquid effective radius (orange) together with the reference simulations (grey).



**Figure 9.** Distributions of cloud droplet number concentrations measured by the SID-3 and parameterized within IFS above (a) sea ice and (b) open ocean averaged over 60s intervals.

and Polar 6 sampled the same cloud and air mass regimes. Figure 9 shows the result of the  $N_d$  parameterization described at the beginning of Sect. 4 together with the in situ observations averaged over 60s intervals for the filtered scenes. The number concentrations in the IFS are within a narrow range between  $36 \text{ cm}^{-3}$  and  $69 \text{ cm}^{-3}$  with slightly higher concentrations above sea ice due to slightly higher prognostic wind speeds leading to a stronger injection of sea spray aerosols to the atmosphere, which can act as cloud condensation nuclei. The in situ observations show a much broader range up to  $230 \text{ cm}^{-3}$ . The observed low values of  $N_d$  mostly result from cloud edges or cloud-free flight sections and are not comparable to the mean grid box values of the IFS. However, the high  $N_d$  of above  $200 \text{ cm}^{-3}$  measured by SID-3 are not captured by the IFS. The findings from Moser et al. (2023) for two different aircraft campaigns in the Arctic with higher  $N_d$  above sea ice compared to open ocean are different from the ALOUD observations, which may be attributed to a different season and different dominating air masses.


To investigate the impact of more realistic cloud droplet number concentrations on the reflected solar irradiance, a new set of ecRad simulations is performed with adjusted  $N_d$ . The lower boundary  $N_{d,\text{obs},\text{min}}$  is fixed to  $N_{d,\text{IFS},\text{min}} = 36 \text{ cm}^{-3}$  to account for the IFS grid box size, which cannot resolve cloud edges with only a few cloud droplets. The upper boundary  $N_{d,\text{obs},\text{max}}$  is set to  $200 \text{ cm}^{-3}$ , excluding only the highest values of the distribution's tail. The initial  $N_{d,\text{IFS}}$  appearing as cloud droplet number concentration in Eq. 4 is replaced by

$$N'_{d,\text{IFS}} = (N_{d,\text{IFS}} - N_{d,\text{IFS},\text{min}}) \cdot \frac{N_{d,\text{obs},\text{max}} - N_{d,\text{obs},\text{min}}}{N_{d,\text{IFS},\text{max}} - N_{d,\text{IFS},\text{min}}} + N_{d,\text{obs},\text{min}}, \quad (10)$$

where  $N'_{d,\text{IFS}}$  is the adjusted cloud droplet number concentration.  $N_{d,\text{IFS},\text{min}}$  ( $N_{d,\text{IFS},\text{max}}$ ) is the minimum (maximum) cloud droplet number concentration from the IFS parameterization and  $N_{d,\text{obs},\text{min}}$  ( $N_{d,\text{obs},\text{max}}$ ) the minimum (maximum) cloud droplet number concentrations derived from in situ observations. The liquid effective radius is recalculated respectively, considering the changed  $N_d$  in Eq. 4, while LWC remains fixed. Figure 10 shows the result of these adjustments. In general, the increase of  $N_d$  increases the reflected solar irradiance. Above sea ice, the maximum values of upward solar irradiance reach  $460 \text{ W m}^{-2}$  instead of  $450 \text{ W m}^{-2}$  while the minimum remains unchanged. In between, the distribution is shifted to slightly higher irradiances. Above open ocean, the entire distribution of adjusted upward solar irradiances is shifted to higher irradiances, with the minimum and maximum ranging  $10 \text{ W m}^{-2}$  higher.  $\Delta F^\uparrow$  decreases by scaling  $N_d$  above sea ice to  $-27 \text{ W m}^{-2}$ , but increases above open ocean to  $48 \text{ W m}^{-2}$ .  $\mathcal{H}^\uparrow$  changed to 0.41 above sea ice and to 0.38 above open ocean accordingly. Above sea ice, the  $N_d$  parameterization may be optimized by a higher variability. Above open ocean, this larger variability of  $N_d$  increases the overestimation by ecRad. A minor issue are observed differences of  $N_d$  between sea ice and open ocean surface, which are not taken into account by the parameterization in IFS, but can occur by different sea salt aerosol production mechanisms above sea ice (blowing snow) and open ocean (wave breaking), as described by e.g. Confer et al. (2023).

## 5 Conclusions

Airborne observations of broadband solar irradiance measured above Arctic low-level clouds during the ALOUD airborne campaign in May/June 2017 were used to evaluate the corresponding solar irradiances simulated by the IFS of the ECMWF. For this purpose, the ecRad radiative transfer scheme embedded in IFS was run in an offline mode using the output of the



**Figure 10.** Distribution of upward solar irradiances above (a) sea ice measured by the CMP22 (blue) and simulated by ecRad with the adjusted number concentration and liquid effective radius (orange) together with the reference simulations (grey) and above (b) open ocean measured by the CMP22 (green) and simulated by ecRad with the adjusted number concentrations and liquid effective radius (black) together with the reference simulations (grey).

corresponding IFS 00 UTC runs as input. While there is agreement within the observational uncertainty between the measured and simulated downward solar irradiance, larger differences exceeding the pyranometer’s uncertainty are found for the upward  
 430 solar irradiance. In a sensitivity study constrained by surface and cloud properties observed during ACLOUD, this bias was attributed to issues of the IFS in representing sea ice albedo and low-level, liquid-dominated mixed-phase clouds. The impacts of different surface and cloud properties were quantified. The limitations of the sea ice model by Ebert and Curry (1993) to represent the change of sea ice albedo during the melting season cause more than 50 % of the observed bias. A comparison to airborne observations reveals an underestimation of the sea ice albedo by the IFS, especially in the wavelength band from  
 435 690 nm to 1190 nm. Implementing the measured sea ice albedo values into ecRad decreases the bias between the simulations and the observations to  $-16 \text{ W m}^{-2}$ .

A misrepresentation of cloud fraction is assessed by active cloud remote sensing. The observed cloud fraction does not change  $\Delta F^\uparrow$  above sea ice, but reduces the bias from  $28 \text{ W m}^{-2}$  to  $18 \text{ W m}^{-2}$  above open ocean where the observations show lower cloud fractions and the difference between the dark ocean and clouds is particularly large. The impact of cloud ice was  
 440 quantified by artificially changing the IWP of the IFS output. The sensitivity of the upward solar irradiance to variations of the IWP of the underlying clouds is nearly negligible with the largest impact of  $-2 \text{ W m}^{-2}$  above open ocean by reducing the IWP by 50 %. The cloud optical properties strongly depend on the LWP of the clouds. Confronting the prognostic LWP with airborne observations (above open ocean only), reduces the positive  $\Delta F^\uparrow$  strongly and overcompensates it with a bias of  $-28 \text{ W m}^{-2}$ . To estimate the effect of a misrepresentation of LWP also above sea ice, a non-observation based sensitivity  
 445 study was performed. Increasing the LWP by 50 %,  $\Delta F^\uparrow$  improves above sea ice to  $-28 \text{ W m}^{-2}$  and by decreasing the LWP by 50 %,  $\Delta F^\uparrow$  improves to  $-5 \text{ W m}^{-2}$ .

Airborne in situ observations have shown that the range of  $N_d$  in the IFS is significantly smaller than measured. This affects the cloud radiative properties simulated by the IFS. Adjusting  $N_d$ , which occurs in the parameterization of the liquid effective radius from Martin et al. (1994) within a range of  $36\text{--}69\text{ cm}^{-3}$ , to a broader range of number concentrations found in the observations ( $36\text{--}200\text{ cm}^{-3}$ ), results in a bias reduction above sea ice to  $-27\text{ W m}^{-2}$  and in a bias enlargement to  $48\text{ W m}^{-2}$  above open ocean.

The sensitivity study identifies the misrepresentation of the surface albedo as the largest contributor to the bias above sea ice. The sea ice albedo values in the IFS are applied as representative constant albedo values of dry snow, melting snow and bare sea ice for fixed times of the year. Replacing these with a sea ice albedo parameterization that considers mixtures of different sea ice types and their specific albedos depending on parameters like the surface temperature may improve the ability of the IFS in correctly simulating the upward solar irradiances in the Arctic (Jäkel et al., 2019b, 2023). The uncertainties of cloud radiative effects in the IFS significantly depend on the surface type below the clouds. With large contributions to the bias improvement given by realistic cloud droplet number concentrations and liquid water paths above sea ice and by realistic cloud fractions and liquid water paths above open ocean, a large amount of the bias could be attributed to the representation of cloud micro- and macrophysical properties during the ACLOUD campaign.

*Data availability.* All ACLOUD-related data sets used in this study are publicly available on the PANGAEA Data Publisher (<https://www.pangaea.de/>) and referenced in Sect. 2.1. The IFS output used in this study was downloaded directly from the ECMWF servers using the Meteorological Archival and Retrieval System (registration required). Aerosol,  $\text{CH}_4$ ,  $\text{CO}$  and  $\text{NO}_2$  data were downloaded from the Copernicus Atmosphere Monitoring Service (CAMS) Atmosphere Data Store (ADS) on <https://ads.atmosphere.copernicus.eu/cdsapp#!/dataset/cams-global-reanalysis-eac4>.  $\text{CO}_2$  and  $\text{N}_2\text{O}$  data were downloaded from the Copernicus Atmosphere Monitoring Service (CAMS) Atmosphere Data Store (ADS) on <https://ads.atmosphere.copernicus.eu/cdsapp#!/dataset/cams-global-greenhouse-gas-inversion>. This study contains modified Copernicus Atmosphere Monitoring Service information (2023); neither the European Commission nor ECMWF is responsible for any use that may be made of the Copernicus information or data it contains. Ozone soundings used in this study were performed by the Alfred-Wegener-Institute, Helmholtz Centre for Polar- and Marine Research; the ozone data used in this publication were obtained from Peter von der Gathen as part of the Network for the Detection of Atmospheric Composition Change (NDACC) and are available through the NDACC website [www.ndacc.org](http://www.ndacc.org).

*Author contributions.* HM performed the analysis, the radiative transfer simulations and drafted the manuscript. AE and MW contributed to conception and design of the study. RH and JR contributed to the set up of ecRad. All authors contributed to the discussion of the results. All authors contributed to reviewing and editing of the manuscript.

*Competing interests.* The authors declare that they have no conflict of interest.

*Acknowledgements.* We gratefully acknowledge the funding by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – project number 268020496 – TRR 172, within the Transregional Collaborative Research Center “Arctic Amplification: Climate Relevant Atmospheric and Surface Processes, and Feedback Mechanisms (AC)<sup>3</sup>”.

## References

- 480 Bauer, P., Magnusson, L., Thépaut, J.-N., and Hamill, T. M.: Aspects of ECMWF model performance in polar areas, *Quarterly Journal of the Royal Meteorological Society*, 142, 583–596, <https://doi.org/10.1002/qj.2449>, 2016.
- Beesley, J. A., Bretherton, C. S., Jakob, C., Andreas, E. L., Intrieri, J. M., and Uttal, T. A.: A comparison of cloud and boundary layer variables in the ECMWF forecast model with observations at Surface Heat Budget of the Arctic Ocean (SHEBA) ice camp, *Journal of Geophysical Research*, 105, 12 337–12 349, <https://doi.org/10.1029/2000JD900079>, 2000.
- 485 Berry, E., Mace, G. G., and Gettelman, A.: Using A-Train Observations to Evaluate Cloud Occurrence and Radiative Effects in the Community Atmosphere Model during the Southeast Asia Summer Monsoon, *Journal of Climate*, 32, 4145–4165, <https://doi.org/10.1175/JCLI-D-18-0693.1>, 2019.
- Boucher, O. and Lohmann, U.: The sulfate-CCN-cloud albedo effect: A sensitivity study with two general circulation models, *Tellus B: Chemical and Physical Meteorology*, 47, 281–300, <https://doi.org/10.3402/tellusb.v47i3.16048>, 1995.
- 490 Brenguier, J.-L., Burnet, F., and Geoffroy, O.: Cloud optical thickness and liquid water path – does the  $k$  coefficient vary with droplet concentration?, *Atmospheric Chemistry and Physics*, 11, 9771–9786, <https://doi.org/10.5194/acp-11-9771-2011>, 2011.
- Chevallier, F., Ciais, P., Conway, T. J., Aalto, T., Anderson, B. E., Bousquet, P., Brunke, E. G., Ciattaglia, L., Esaki, Y., Fröhlich, M., Gomez, A., Gomez-Pelaez, A. J., Haszpra, L., Krummel, P. B., Langenfelds, R. L., Leuenberger, M., Machida, T., Maignan, F., Matsueda, H., Morguá, J. A., Mukai, H., Nakazawa, T., Peylin, P., Ramonet, M., Rivier, L., Sawa, Y., Schmidt, M., Steele, L. P., Vay, S. A., Vermeulen, A. T., Wofsy, S., and Worthly, D.: CO<sub>2</sub> surface fluxes at grid point scale estimated from a global 21 year reanalysis of atmospheric measurements, *Journal of Geophysical Research*, 115, <https://doi.org/10.1029/2010JD013887>, 2010.
- 495 Chevallier, F., Remaud, M., O’Dell, C. W., Baker, D., Peylin, P., and Cozic, A.: Objective evaluation of surface- and satellite-driven carbon dioxide atmospheric inversions, *Atmospheric Chemistry and Physics*, 19, 14 233–14 251, <https://doi.org/10.5194/acp-19-14233-2019>, 2019.
- 500 Cohen, J., Screen, J. A., Furtado, J. C., Barlow, M., Whittleston, D., Coumou, D., Francis, J., Dethloff, K., Entekhabi, D., Overland, J., and Jones, J.: Recent Arctic amplification and extreme mid-latitude weather, *Nature Geoscience*, 7, 627–637, <https://doi.org/10.1038/ngeo2234>, 2014.
- Confer, K. L., Jaeglé, L., Liston, G. E., Sharma, S., Nandan, V., Yackel, J., Ewert, M., and Horowitz, H. M.: Impact of Changing Arctic Sea Ice Extent, Sea Ice Age, and Snow Depth on Sea Salt Aerosol From Blowing Snow and the Open Ocean for 1980–2017, *J. Geophys. Res.-Atmos.*, 128, e2022JD037 667, <https://doi.org/10.1029/2022JD037667>, 2023.
- 505 Day, J. J., Keeley, S., Arduini, G., Magnusson, L., Mogensen, K., Rodwell, M., Sandu, I., and Tietsche, S.: Benefits and challenges of dynamic sea ice for weather forecasts, *Weather and Climate Dynamics*, 3, 713–731, <https://doi.org/10.5194/wcd-3-713-2022>, 2022.
- Dionne, J., von Salzen, K., Cole, J., Mahmood, R., Leitch, W. R., Lesins, G., Folkins, I., and Chang, R. Y.-W.: Modelling the relationship between liquid water content and cloud droplet number concentration observed in low clouds in the summer Arctic and its radiative effects, *Atmospheric Chemistry and Physics*, 20, 29–43, <https://doi.org/10.5194/acp-20-29-2020>, 2020.
- 510 Eastman, R. and Warren, S. G.: Interannual Variations of Arctic Cloud Types in Relation to Sea Ice, *Journal of Climate*, 23, 4216–4232, <https://doi.org/10.1175/2010JCLI3492.1>, 2010.
- Ebert, E. E. and Curry, J. A.: An intermediate one-dimensional thermodynamic sea ice model for investigating ice-atmosphere interactions, *Journal of Geophysical Research*, 98, 10 085, <https://doi.org/10.1029/93JC00656>, 1993.

- 515 Ehrlich, A., Wendisch, M., Bierwirth, E., Herber, A., and Schwarzenböck, A.: Ice crystal shape effects on solar radiative properties of Arctic mixed-phase clouds—Dependence on microphysical properties, *Atmospheric Research*, 88, 266–276, <https://doi.org/10.1016/j.atmosres.2007.11.018>, 2008.
- Ehrlich, A., Wendisch, M., Lüpkes, C., Buschmann, M., Bozem, H., Chechin, D., Clemen, H.-C., Dupuy, R., Eppers, O., Hartmann, J., Herber, A., Jäkel, E., Järvinen, E., Jourdan, O., Kästner, U., Kliesch, L.-L., Köllner, F., Mech, M., Mertes, S., Neuber, R., Ruiz-  
520 Donoso, E., Schnaiter, M., Schneider, J., Stapf, J., and Zanatta, M.: A comprehensive in situ and remote sensing data set from the Arctic CLOUD Observations Using airborne measurements during polar Day (ACLOUD) campaign, *Earth System Science Data*, 11, 1853–1881, <https://doi.org/10.5194/essd-11-1853-2019>, 2019.
- Ehrlich, A., Zöger, M., Giez, A., Nenakhov, V., Mallaun, C., Maser, R., Rösenthaller, T., Luebke, A. E., Wolf, K., Stevens, B., and Wendisch, M.: A new airborne broadband radiometer system and an efficient method to correct dynamic thermal offsets, *Atmospheric Measurement  
525 Techniques*, 16, 1563–1581, <https://doi.org/10.5194/amt-16-1563-2023>, 2023.
- Erickson, D. J., Merrill, J. T., and Duce, R. A.: Seasonal estimates of global atmospheric sea-salt distributions, *Journal of Geophysical Research*, 91, 1067, <https://doi.org/10.1029/JD091iD01p01067>, 1986.
- Forbes, R. M. and Ahlgrimm, M.: On the Representation of High-Latitude Boundary Layer Mixed-Phase Cloud in the ECMWF Global Model, *Monthly Weather Review*, 142, 3425–3445, <https://doi.org/10.1175/MWR-D-13-00325.1>, 2014.
- 530 Formenti, P. and Wendisch, M.: Combining Upcoming Satellite Missions and Aircraft Activities: Future Challenges for the EUFAR Fleet, *Bulletin of the American Meteorological Society*, 89, 385–388, <https://doi.org/10.1175/BAMS-89-3-385>, 2008.
- Fu, Q.: An Accurate Parameterization of the Solar Radiative Properties of Cirrus Clouds for Climate Models, *Journal of Climate*, 9, 2058–2082, [https://doi.org/10.1175/1520-0442\(1996\)009<2058:AAPOTS>2.0.CO;2](https://doi.org/10.1175/1520-0442(1996)009<2058:AAPOTS>2.0.CO;2), 1996.
- Fu, Q., Yang, P., and Sun, W. B.: An Accurate Parameterization of the Infrared Radiative Properties of Cirrus Clouds for Climate Models, *Journal of Climate*, 11, 2223–2237, [https://doi.org/10.1175/1520-0442\(1998\)011<2223:AAPOTI>2.0.CO;2](https://doi.org/10.1175/1520-0442(1998)011<2223:AAPOTI>2.0.CO;2), 1998.  
535
- Genthon, C.: Simulations of desert dust and sea-salt aerosols in Antarctica with a general circulation model of the atmosphere, *Tellus B*, 44, 371–389, <https://doi.org/10.1034/j.1600-0889.1992.00014.x>, 1992.
- Geoffroy, O., Brenguier, J.-L., and Burnet, F.: Parametric representation of the cloud droplet spectra for LES warm bulk microphysical schemes, *Atmospheric Chemistry and Physics*, 10, 4835–4848, <https://doi.org/10.5194/acp-10-4835-2010>, 2010.
- 540 Gu, M., Wang, Z., Wei, J., and Yu, X.: An assessment of Arctic cloud water paths in atmospheric reanalyses, *Acta Oceanologica Sinica*, 40, 46–57, <https://doi.org/10.1007/s13131-021-1706-5>, 2021.
- Hartmann, J., Kottmeier, C., and Raasch, S.: Roll Vortices and Boundary-Layer Development during a Cold Air Outbreak, *Bound.-Lay. Meteorol.*, 84, 45–65, <https://doi.org/10.1023/A:1000392931768>, 1997.
- Hellinger, E.: Neue Begründung der Theorie quadratischer Formen von unendlichvielen Veränderlichen, *Journal für die reine und angewandte  
545 Mathematik*, 1909, 210–271, <https://doi.org/10.1515/crll.1909.136.210>, 1909.
- Hirst, E., Kaye, P. H., Greenaway, R. S., Field, P., and Johnson, D. W.: Discrimination of micrometre-sized ice and super-cooled droplets in mixed-phase cloud, *Atmospheric Environment*, 35, 33–47, [https://doi.org/10.1016/S1352-2310\(00\)00377-0](https://doi.org/10.1016/S1352-2310(00)00377-0), 2001.
- Hogan, R. J. and Bozzo, A.: A Flexible and Efficient Radiation Scheme for the ECMWF Model, *Journal of Advances in Modeling Earth Systems*, 10, 1990–2008, <https://doi.org/10.1029/2018MS001364>, 2018.
- 550 Hogan, R. J., O’Connor, E. J., and Illingworth, A. J.: Verification of cloud-fraction forecasts, *Quarterly Journal of the Royal Meteorological Society*, 135, 1494–1511, <https://doi.org/10.1002/qj.481>, 2009.



- Huang, Y., Dong, X., Xi, B., Dolinar, E. K., Stanfield, R. E., and Qiu, S.: Quantifying the Uncertainties of Reanalyzed Arctic Cloud and Radiation Properties Using Satellite Surface Observations, *Journal of Climate*, 30, 8007–8029, <https://doi.org/10.1175/JCLI-D-16-0722.1>, 2017.
- 555 Illingworth, A. J., Hogan, R. J., O'Connor, E. J., Bouniol, D., Brooks, M. E., Delanoé, J., Donovan, D. P., Eastment, J. D., Gaussiat, N., Goddard, J. W. F., Haeffelin, M., Baltink, H. K., Krasnov, O. A., Pelon, J., Piriou, J.-M., Protat, A., Russchenberg, H. W. J., Seifert, A., Tompkins, A. M., van Zadelhoff, G.-J., Vinit, F., Willén, U., Wilson, D. R., and Wrench, C. L.: Cloudnet, *Bulletin of the American Meteorological Society*, 88, 883–898, <https://doi.org/10.1175/BAMS-88-6-883>, 2007.
- Inness, A., Ades, M., Agustí-Panareda, A., Barré, J., Benedictow, A., Blechschmidt, A.-M., Dominguez, J. J., Engelen, R., Eskes, H., Flemming, J., Huijnen, V., Jones, L., Kipling, Z., Massart, S., Parrington, M., Peuch, V.-H., Razinger, M., Remy, S., Schulz, M., and Suttie, M.: The CAMS reanalysis of atmospheric composition, *Atmospheric Chemistry and Physics*, 19, 3515–3556, <https://doi.org/10.5194/acp-19-3515-2019>, 2019.
- 560 Jäkel, E., Ehrlich, A., Schäfer, M., and Wendisch, M.: Aircraft measurements of spectral solar up- and downward irradiances in the Arctic during the ACLOUD campaign 2017, <https://doi.org/10.1594/PANGAEA.899177>, 2019a.
- 565 Jäkel, E., Stapf, J., Wendisch, M., Nicolaus, M., Dorn, W., and Rinke, A.: Validation of the sea ice surface albedo scheme of the regional climate model HIRHAM–NAOSIM using aircraft measurements during the ACLOUD/PASCAL campaigns, *The Cryosphere*, 13, 1695–1708, <https://doi.org/10.5194/tc-13-1695-2019>, 2019b.
- Jäkel, E., Becker, S., Sperzel, T. R., Niehaus, H., Spreen, G., Tao, R., Nicolaus, M., Dorn, W., Rinke, A., Brauchle, J., and Wendisch, M.: Observations and modeling of areal surface albedo and surface types in the Arctic, *EGUsphere*, pp. 1–30, <https://doi.org/10.5194/egusphere-2023-1337>, 2023.
- 570 Jung, T. and Matsueda, M.: Verification of global numerical weather forecasting systems in polar regions using TIGGE data, *Quarterly Journal of the Royal Meteorological Society*, 142, 574–582, <https://doi.org/10.1002/qj.2437>, 2016.
- Jung, T., Kasper, M. A., Semmler, T., and Serrar, S.: Arctic influence on subseasonal midlatitude prediction, *Geophysical Research Letters*, 41, 3676–3680, <https://doi.org/10.1002/2014GL059961>, 2014.
- 575 Jung, T., Gordon, N. D., Bauer, P., Bromwich, D. H., Chevallier, M., Day, J. J., Dawson, J., Doblas-Reyes, F., Fairall, C., Goessling, H. F., Holland, M., Inoue, J., Iversen, T., Klebe, S., Lemke, P., Losch, M., Makshtas, A., Mills, B., Nurmi, P., Perovich, D., Reid, P., Renfrew, I. A., Smith, G., Svensson, G., Tolstykh, M., and Yang, Q.: Advancing Polar Prediction Capabilities on Daily to Seasonal Time Scales, *Bulletin of the American Meteorological Society*, 97, 1631–1647, <https://doi.org/10.1175/BAMS-D-14-00246.1>, 2016.
- Karlsson, J. and Svensson, G.: Consequences of poor representation of Arctic sea-ice albedo and cloud-radiation interactions in the CMIP5 model ensemble, *Geophysical Research Letters*, 40, 4374–4379, <https://doi.org/10.1002/grl.50768>, 2013.
- 580 Keeley, S. and Mogensen, K.: Dynamic sea ice in the IFS, <https://doi.org/10.21957/4SKA25FURB>.
- Klein, S. A., McCoy, R. B., Morrison, H., Ackerman, A. S., Avramov, A., de Boer, G., Chen, M., Cole, J. N. S., Del Genio, A. D., Falk, M., Foster, M. J., Fridlind, A., Golaz, J.-C., Hashino, T., Harrington, J. Y., Hoose, C., Khairoutdinov, M. F., Larson, V. E., Liu, X., Luo, Y., McFarquhar, G. M., Menon, S., Neggers, R. A. J., Park, S., Poellot, M. R., Schmidt, J. M., Sednev, I., Shipway, B. J., Shupe, M. D.,
- 585 Spangenberg, D. A., Sud, Y. C., Turner, D. D., Veron, D. E., von Salzen, K., Walker, G. K., Wang, Z., Wolf, A. B., Xie, S., Xu, K.-M., Yang, F., and Zhang, G.: Intercomparison of model simulations of mixed-phase clouds observed during the ARM Mixed-Phase Arctic Cloud Experiment. I: single-layer cloud, *Quarterly Journal of the Royal Meteorological Society*, 135, 979–1002, <https://doi.org/10.1002/qj.416>, 2009.

- 590 Kliesch, L.-L. and Mech, M.: Airborne radar reflectivity and brightness temperature measurements with POLAR 5 during ACLOUD in May and June 2017, <https://doi.org/10.1594/PANGAEA.899565>, 2019.
- Klingebiel, M., Ehrlich, A., Ruiz-Donoso, E., Risse, N., Schirmacher, I., Jäkel, E., Schäfer, M., Wolf, K., Mech, M., Moser, M., Voigt, C., and Wendisch, M.: Variability and properties of liquid-dominated clouds over the ice-free and sea-ice-covered Arctic Ocean, *Atmospheric Chemistry and Physics*, 23, 15 289–15 304, <https://doi.org/10.5194/acp-23-15289-2023>, 2023.
- 595 Klocke, D. and Rodwell, M. J.: A comparison of two numerical weather prediction methods for diagnosing fast-physics errors in climate models, *Quarterly Journal of the Royal Meteorological Society*, 140, 517–524, <https://doi.org/10.1002/qj.2172>, 2014.
- Kokhanovsky, A.: Optical properties of terrestrial clouds, *Earth-Science Reviews*, 64, 189–241, [https://doi.org/10.1016/S0012-8252\(03\)00042-4](https://doi.org/10.1016/S0012-8252(03)00042-4), 2004.
- Kopp, G. and Lean, J. L.: A new, lower value of total solar irradiance: Evidence and climate significance, *Geophysical Research Letters*, 38, n/a–n/a, <https://doi.org/10.1029/2010GL045777>, 2011.
- 600 Kretzschmar, J., Stapf, J., Klocke, D., Wendisch, M., and Quaas, J.: Employing airborne radiation and cloud microphysics observations to improve cloud representation in ICON at kilometer-scale resolution in the Arctic, *Atmospheric Chemistry and Physics*, 20, 13 145–13 165, <https://doi.org/10.5194/acp-20-13145-2020>, 2020.
- Kulla, B. S., Mech, M., Risse, N., and Ritter, C.: Cloud top altitude retrieved from Lidar measurements during ACLOUD at 1 second resolution, <https://doi.org/10.1594/PANGAEA.932454>, 2021.
- 605 Lawrence, H., Bormann, N., Sandu, I., Day, J., Farnan, J., and Bauer, P.: Use and impact of Arctic observations in the ECMWF Numerical Weather Prediction system, *Quarterly Journal of the Royal Meteorological Society*, 145, 3432–3454, <https://doi.org/10.1002/qj.3628>, 2019.
- Leaitch, W. R., Korolev, A., Aliabadi, A. A., Burkart, J., Willis, M. D., Abbatt, J. P. D., Bozem, H., Hoor, P., Köllner, F., Schneider, J., Herber, A., Konrad, C., and Brauner, R.: Effects of 20–100 nm particles on liquid clouds in the clean summertime Arctic, *Atmospheric Chemistry and Physics*, 16, 11 107–11 124, <https://doi.org/10.5194/acp-16-11107-2016>, 2016.
- 610 Liu, J., Zhang, Z., Inoue, J., and Horton, R. M.: Evaluation of snow/ice albedo parameterizations and their impacts on sea ice simulations, *International Journal of Climatology*, 27, 81–91, <https://doi.org/10.1002/joc.1373>, 2007.
- Lowenthal, D. H., Borys, R. D., Choulaton, T. W., Bower, K. N., Flynn, M. J., and Gallagher, M. W.: Parameterization of the cloud droplet–sulfate relationship, *Atmospheric Environment*, 38, 287–292, <https://doi.org/10.1016/j.atmosenv.2003.09.046>, 2004.
- 615 Martin, G. M., Johnson, D. W., and Spice, A.: The Measurement and Parameterization of Effective Radius of Droplets in Warm Stratocumulus Clouds, *Journal of the Atmospheric Sciences*, 51, 1823–1842, [https://doi.org/10.1175/1520-0469\(1994\)051<1823:TMAPOE>2.0.CO;2](https://doi.org/10.1175/1520-0469(1994)051<1823:TMAPOE>2.0.CO;2), 1994.
- Matsui, T., Santanello, J., Shi, J. J., Tao, W.-K., Wu, D., Peters-Lidard, C., Kemp, E., Chin, M., Starr, D., Sekiguchi, M., and Aires, F.: Introducing multisensor satellite radiance-based evaluation for regional Earth System modeling, *Journal of Geophysical Research: Atmospheres*, 119, 8450–8475, <https://doi.org/10.1002/2013JD021424>, 2014.
- 620 McCusker, G. Y., Vüllers, J., Achtert, P., Field, P., Day, J. J., Forbes, R., Price, R., O’Connor, E., Tjernström, M., Prytherch, J., Neely III, R., and Brooks, I. M.: Evaluating Arctic clouds modelled with the Unified Model and Integrated Forecasting System, *Atmospheric Chemistry and Physics*, 23, 4819–4847, <https://doi.org/10.5194/acp-23-4819-2023>, 2023.
- Mech, M., Kliesch, L.-L., Anhäuser, A., Rose, T., Kollias, P., and Crewell, S.: Microwave Radar/radiometer for Arctic Clouds (MiRAC): first insights from the ACLOUD campaign, *Atmospheric Measurement Techniques*, 12, 5019–5037, <https://doi.org/10.5194/amt-12-5019-2019>, 2019.
- 625

- Melsheimer, C. and Spreen, G.: AMSR2 ASI sea ice concentration data, Arctic, version 5.4 (NetCDF) (July 2012 - December 2019), <https://doi.org/10.1594/PANGAEA.898399>, 2019.
- 630 Mioche, G., Jourdan, O., Ceccaldi, M., and Delanoë, J.: Variability of mixed-phase clouds in the Arctic with a focus on the Svalbard region: a study based on spaceborne active remote sensing, *Atmospheric Chemistry and Physics*, 15, 2445–2461, <https://doi.org/10.5194/acp-15-2445-2015>, 2015.
- Mlawer, E. J., Taubman, S. J., Brown, P. D., Iacono, M. J., and Clough, S. A.: Radiative transfer for inhomogeneous atmospheres: RRTM, a validated correlated-k model for the longwave, *Journal of Geophysical Research*, 102, 16 663–16 682, <https://doi.org/10.1029/97JD00237>, 1997.
- 635 Morcrette, C. J., O’Connor, E. J., and Petch, J. C.: Evaluation of two cloud parametrization schemes using ARM and Cloud-Net observations, *Quarterly Journal of the Royal Meteorological Society*, 138, 964–979, <https://doi.org/10.1002/qj.969>, 2012.
- Morrison, H. and Pinto, J. O.: Intercomparison of Bulk Cloud Microphysics Schemes in Mesoscale Simulations of Springtime Arctic Mixed-Phase Stratiform Clouds, *Monthly Weather Review*, 134, 1880–1900, <https://doi.org/10.1175/MWR3154.1>, 2006.
- Morrison, H., McCoy, R. B., Klein, S. A., Xie, S., Luo, Y., Avramov, A., Chen, M., Cole, J. N. S., Falk, M., Foster, M. J., Del Genio, A. D.,  
 640 Harrington, J. Y., Hoose, C., Khairoutdinov, M. F., Larson, V. E., Liu, X., McFarquhar, G. M., Poellot, M. R., von Salzen, K., Shipway, B. J., Shupe, M. D., Sud, Y. C., Turner, D. D., Veron, D. E., Walker, G. K., Wang, Z., Wolf, A. B., Xu, K.-M., Yang, F., and Zhang, G.: Intercomparison of model simulations of mixed-phase clouds observed during the ARM Mixed-Phase Arctic Cloud Experiment. II: Multilayer cloud, *Quarterly Journal of the Royal Meteorological Society*, 135, 1003–1019, <https://doi.org/10.1002/qj.415>, 2009.
- Morrison, H., de Boer, G., Feingold, G., Harrington, J., Shupe, M. D., and Sulia, K.: Resilience of persistent Arctic mixed-phase clouds,  
 645 *Nature Geoscience*, 5, 11–17, <https://doi.org/10.1038/ngeo1332>, 2012.
- Moser, M., Voigt, C., Jurkat-Witschas, T., Hahn, V., Mioche, G., Jourdan, O., Dupuy, R., Gourbeyre, C., Schwarzenboeck, A., Lucke, J., Boose, Y., Mech, M., Borrmann, S., Ehrlich, A., Herber, A., Lüpkes, C., and Wendisch, M.: Microphysical and thermodynamic phase analyses of Arctic low-level clouds measured above the sea ice and the open ocean in spring and summer, *Atmospheric Chemistry and Physics*, 23, 7257–7280, <https://doi.org/10.5194/acp-23-7257-2023>, 2023.
- 650 Ortega, P., Blockley, E. W., Køltzow, M., Massonnet, F., Sandu, I., Svensson, G., Acosta Navarro, J. C., Arduini, G., Batté, L., Bazile, E., Chevallier, M., Cruz-García, R., Day, J. J., Fichet, T., Flocco, D., Gupta, M., Hartung, K., Hawkins, E., Hinrichs, C., Magnusson, L., Moreno-Chamarro, E., Pérez-Montero, S., Ponsoni, L., Semmler, T., Smith, D., Sterlin, J., Tjernström, M., Välisuo, I., and Jung, T.: Improving Arctic Weather and Seasonal Climate Prediction: Recommendations for Future Forecast Systems Evolution from the European Project APPLICATE, *B. Am. Meteorol. Soc.*, 103, E2203–E2213, <https://doi.org/10.1175/BAMS-D-22-0083.1>, 2022.
- 655 Overland, J., Francis, J. A., Hall, R., Hanna, E., Kim, S.-J., and Vihma, T.: The Melting Arctic and Midlatitude Weather Patterns: Are They Connected?\*, *Journal of Climate*, 28, 7917–7932, <https://doi.org/10.1175/JCLI-D-14-00822.1>, 2015.
- Pincus, R., Barker, H. W., and Morcrette, J.-J.: A fast, flexible, approximate technique for computing radiative transfer in inhomogeneous cloud fields, *Journal of Geophysical Research*, 108, n/a–n/a, <https://doi.org/10.1029/2002JD003322>, 2003.
- Pithan, F., Svensson, G., Caballero, R., Chechin, D., Cronin, T. W., Ekman, A. M. L., Neggers, R., Shupe, M. D., Solomon, A., Tjernström,  
 660 M., and Wendisch, M.: Role of air-mass transformations in exchange between the Arctic and mid-latitudes, *Nature Geoscience*, 11, 805–812, <https://doi.org/10.1038/s41561-018-0234-1>, 2018.
- Pohl, C., Istomina, L., Tietsche, S., Jäkel, E., Stapf, J., Spreen, G., and Heygster, G.: Broadband albedo of Arctic sea ice from MERIS optical data, *The Cryosphere*, 14, 165–182, <https://doi.org/10.5194/tc-14-165-2020>, 2020.

- Rantanen, M., Karpechko, A. Y., Lipponen, A., Nordling, K., Hyvärinen, O., Ruosteenoja, K., Vihma, T., and Laaksonen, A.:  
665 The Arctic has warmed nearly four times faster than the globe since 1979, *Communications Earth & Environment*, 3, 1–10,  
<https://doi.org/10.1038/s43247-022-00498-3>, 2022.
- Rosenburg, S., Lange, C., Jäkel, E., Schäfer, M., Ehrlich, A., and Wendisch, M.: Retrieval of snow layer and melt pond proper-  
ties on Arctic sea ice from airborne imaging spectrometer observations, *Atmospheric Measurement Techniques*, 16, 3915–3930,  
<https://doi.org/10.5194/amt-16-3915-2023>, 2023.
- 670 Ruiz-Donoso, E., Ehrlich, A., Schäfer, M., Jäkel, E., Schemann, V., Crewell, S., Mech, M., Kulla, B. S., Kliesch, L.-L., Neuber, R., and  
Wendisch, M.: Small-scale structure of thermodynamic phase in Arctic mixed-phase clouds observed by airborne remote sensing during  
a cold air outbreak and a warm air advection event, *Atmospheric Chemistry and Physics*, 20, 5487–5511, <https://doi.org/10.5194/acp-20-5487-2020>, 2020.
- Schäfer, M., Loewe, K., Ehrlich, A., Hoose, C., and Wendisch, M.: Simulated and observed horizontal inhomogeneities of optical thickness  
675 of Arctic stratus, *Atmospheric Chemistry and Physics*, 18, 13 115–13 133, <https://doi.org/10.5194/acp-18-13115-2018>, 2018.
- Schnaiter, M. and Järvinen, E.: SID-3 1Hz size distribution of cloud particles during the ACLOUD campaign in 2017,  
<https://doi.org/10.1594/PANGAEA.900261>.
- Serreze, M. C. and Meier, W. N.: The Arctic’s sea ice cover: trends, variability, predictability, and comparisons to the Antarctic, *Annals of  
the New York Academy of Sciences*, 1436, 36–53, <https://doi.org/10.1111/nyas.13856>, 2019.
- 680 Shonk, J. K. P. and Hogan, R. J.: Tripleclouds: An Efficient Method for Representing Horizontal Cloud Inhomogeneity in 1D Radiation  
Schemes by Using Three Regions at Each Height, *Journal of Climate*, 21, 2352–2370, <https://doi.org/10.1175/2007JCLI1940.1>, 2008.
- Shonk, J. K. P., Hogan, R. J., Edwards, J. M., and Mace, G. G.: Effect of improving representation of horizontal and vertical cloud structure  
on the Earth’s global radiation budget. Part I: Review and parametrization, *Quarterly Journal of the Royal Meteorological Society*, 136,  
1191–1204, <https://doi.org/10.1002/qj.647>, 2010.
- 685 Shupe, M. D. and Intrieri, J. M.: Cloud Radiative Forcing of the Arctic Surface: The Influence of Cloud Properties, Surface Albedo, and  
Solar Zenith Angle, *Journal of Climate*, 17, 616–628, [https://doi.org/10.1175/1520-0442\(2004\)017<0616:CRFOTA>2.0.CO;2](https://doi.org/10.1175/1520-0442(2004)017<0616:CRFOTA>2.0.CO;2), 2004.
- Shupe, M. D., Matrosov, S. Y., and Uttal, T.: Arctic Mixed-Phase Cloud Properties Derived from Surface-Based Sensors at SHEBA, *Journal  
of the Atmospheric Sciences*, 63, 697–711, <https://doi.org/10.1175/JAS3659.1>, 2006.
- Sinclair, V. A., Ritvanen, J., Urbancic, G., Statnaia, I., Batrak, Y., Moiseev, D., and Kurppa, M.: Boundary-layer height and surface stability  
690 at Hyytiälä, Finland, in ERA5 and observations, *Atmospheric Measurement Techniques*, 15, 3075–3103, <https://doi.org/10.5194/amt-15-3075-2022>, 2022.
- Smith, L. C. and Stephenson, S. R.: New Trans-Arctic shipping routes navigable by midcentury, *Proceedings of the National Academy of  
Sciences of the United States of America*, 110, E1191–5, <https://doi.org/10.1073/pnas.1214212110>, 2013.
- Solomon, A., Morrison, H., Persson, O., Shupe, M. D., and Bao, J.-W.: Investigation of Microphysical Parameterizations of Snow  
695 and Ice in Arctic Clouds during M-PACE through Model–Observation Comparisons, *Monthly Weather Review*, 137, 3110–3128,  
<https://doi.org/10.1175/2009MWR2688.1>, 2009.
- Solomon, A., Shupe, M. D., Svensson, G., Barton, N. P., Batrak, Y., Bazile, E., Day, J. J., Doyle, J. D., Frank, H. P., Keeley, S., Remes, T.,  
and Tolstykh, M.: The winter central Arctic surface energy budget: A model evaluation using observations from the MOSAiC campaign,  
*Elem Sci Anth*, 11, <https://doi.org/10.1525/elementa.2022.00104>, 2023.
- 700 Sotiropoulou, G., Sedlar, J., Forbes, R., and Tjernström, M.: Summer Arctic clouds in the ECMWF forecast model: an evaluation of cloud  
parametrization schemes, *Quarterly Journal of the Royal Meteorological Society*, 142, 387–400, <https://doi.org/10.1002/qj.2658>, 2016.

- Spreen, G., Kaleschke, L., and Heygster, G.: Sea ice remote sensing using AMSR-E 89-GHz channels, *Journal of Geophysical Research*, 113, <https://doi.org/10.1029/2005JC003384>, 2008.
- 705 Stachlewska, I. S., Neuber, R., Lampert, A., Ritter, C., and Wehrle, G.: AMALi – the Airborne Mobile Aerosol Lidar for Arctic research, *Atmospheric Chemistry and Physics*, 10, 2947–2963, <https://doi.org/10.5194/acp-10-2947-2010>, 2010.
- Stapf, J., Ehrlich, A., Jäkel, E., and Wendisch, M.: Aircraft measurements of broadband irradiance during the ACLOUD campaign in 2017, <https://doi.org/10.1594/PANGAEA.900442>, 2019.
- 710 Stevens, R. G., Loewe, K., Dearden, C., Dimitrellos, A., Possner, A., Eirund, G. K., Raatikainen, T., Hill, A. A., Shipway, B. J., Wilkinson, J., Romakkaniemi, S., Tonttila, J., Laaksonen, A., Korhonen, H., Connolly, P., Lohmann, U., Hoose, C., Ekman, A. M. L., Carslaw, K. S., and Field, P. R.: A model intercomparison of CCN-limited tenuous clouds in the high Arctic, *Atmospheric Chemistry and Physics*, 18, 11 041–11 071, <https://doi.org/10.5194/acp-18-11041-2018>, 2018.
- Su, W., Dutton, E., Charlock, T. P., and Wiscombe, W.: Performance of Commercial Radiometers in Very Low Temperature and Pressure Environments Typical of Polar Regions and of the Stratosphere: A Laboratory Study, *Journal of Atmospheric and Oceanic Technology*, 25, 558–569, <https://doi.org/10.1175/2007JTECHA1005.1>, 2008.
- 715 Sun, Z.: Reply to comments by Greg M. McFarquhar on ‘Parametrization of effective sizes of cirrus-cloud particles and its verification against observations’. (October B, 1999, 125, 3037–3055), *Quarterly Journal of the Royal Meteorological Society*, 127, 267–271, <https://doi.org/10.1002/qj.49712757116>, 2001.
- Sun, Z. and Rikus, L.: Parametrization of effective sizes of cirrus-cloud particles and its verification against observations, *Quarterly Journal of the Royal Meteorological Society*, 125, 3037–3055, <https://doi.org/10.1002/qj.49712556012>, 1999.
- 720 Taylor, J. P., Edwards, J. M., Glew, M. D., Hignett, P., and Slingo, A.: Studies with a flexible new radiation code. II: Comparisons with aircraft short-wave observations, *Quarterly Journal of the Royal Meteorological Society*, 122, 839–861, <https://doi.org/10.1002/qj.49712253204>, 1996.
- Thompson, R. L., Bousquet, P., Chevallier, F., Rayner, P. J., and Ciais, P.: Impact of the atmospheric sink and vertical mixing on nitrous oxide fluxes estimated using inversion methods, *Journal of Geophysical Research*, 116, <https://doi.org/10.1029/2011JD015815>, 2011.
- 725 Tjernström, M., Svensson, G., Magnusson, L., Brooks, I. M., Prytherch, J., Vuilliers, J., and Young, G.: Central Arctic weather forecasting: Confronting the ECMWF IFS with observations from the Arctic Ocean 2018 expedition, *Quarterly Journal of the Royal Meteorological Society*, 147, 1278–1299, <https://doi.org/10.1002/qj.3971>, 2021.
- 730 Uttal, T., Curry, J. A., McPhee, M. G., Perovich, D. K., Moritz, R. E., Maslanik, J. A., Guest, P. S., Stern, H. L., Moore, J. A., Turenne, R., Heiberg, A., Serreze, M. C., Wylie, D. P., Persson, O. G., Paulson, C. A., Halle, C., Morison, J. H., Wheeler, P. A., Makshtas, A., Welch, H., Shupe, M. D., Intrieri, J. M., Stamnes, K., Lindsey, R. W., Pinkel, R., Pegau, W. S., Stanton, T. P., and Grenfeld, T. C.: Surface Heat Budget of the Arctic Ocean, *Bulletin of the American Meteorological Society*, 83, 255–275, [https://doi.org/10.1175/1520-0477\(2002\)083<0255:SHBOTA>2.3.CO;2](https://doi.org/10.1175/1520-0477(2002)083<0255:SHBOTA>2.3.CO;2), 2002.
- 735 Verlinde, J., Harrington, J. Y., McFarquhar, G. M., Yannuzzi, V. T., Avramov, A., Greenberg, S., Johnson, N., Zhang, G., Poellot, M. R., Mather, J. H., Turner, D. D., Eloranta, E. W., Zak, B. D., Prenni, A. J., Daniel, J. S., Kok, G. L., Tobin, D. C., Holz, R., Sassen, K., Spangenberg, D., Minnis, P., Tooman, T. P., Ivey, M. D., Richardson, S. J., Bahrmann, C. P., Shupe, M., DeMott, P. J., Heymsfield, A. J., and Schofield, R.: The Mixed-Phase Arctic Cloud Experiment, *Bulletin of the American Meteorological Society*, 88, 205–222, <https://doi.org/10.1175/BAMS-88-2-205>, 2007.

- Vochezer, P., Järvinen, E., Wagner, R., Kupiszewski, P., Leisner, T., and Schnaiter, M.: In situ characterization of mixed phase clouds using the Small Ice Detector and the Particle Phase Discriminator, *Atmospheric Measurement Techniques*, 9, 159–177, <https://doi.org/10.5194/amt-9-159-2016>, 2016.
- 740 Vuilleumier, L., Hauser, M., Félix, C., Vignola, F., Blanc, P., Kazantzidis, A., and Calpini, B.: Accuracy of ground surface broadband short-wave radiation monitoring, *Journal of Geophysical Research: Atmospheres*, 119, 13,838–13,860, <https://doi.org/10.1002/2014JD022335>, 2014.
- Vüllers, J., Achtert, P., Brooks, I. M., Tjernström, M., Prytherch, J., Burzik, A., and Neely III, R.: Meteorological and cloud conditions during the Arctic Ocean 2018 expedition, *Atmospheric Chemistry and Physics*, 21, 289–314, <https://doi.org/10.5194/acp-21-289-2021>, 2021.
- 745 Walsh, J. E., Chapman, W. L., and Portis, D. H.: Arctic Cloud Fraction and Radiative Fluxes in Atmospheric Reanalyses, *Journal of Climate*, 22, 2316–2334, <https://doi.org/10.1175/2008JCLI2213.1>, 2009.
- Wendisch, M.: Impact of cirrus crystal shape on solar spectral irradiance: A case study for subtropical cirrus, *Journal of Geophysical Research*, 110, <https://doi.org/10.1029/2004JD005294>, 2005.
- 750 Wendisch, M., Müller, D., Schell, D., and Heintzenberg, J.: An Airborne Spectral Albedometer with Active Horizontal Stabilization, *Journal of Atmospheric and Oceanic Technology*, 18, 1856–1866, [https://doi.org/10.1175/1520-0426\(2001\)018<1856:AASAWA>2.0.CO;2](https://doi.org/10.1175/1520-0426(2001)018<1856:AASAWA>2.0.CO;2), 2001.
- Wendisch, M., Macke, A., Ehrlich, A., Lüpkes, C., Mech, M., Chechin, D., Dethloff, K., Velasco, C. B., Bozem, H., Brückner, M., Clemen, H.-C., Crewell, S., Donth, T., Dupuy, R., Ebell, K., Egerer, U., Engelmann, R., Engler, C., Eppers, O., Gehrman, M., Gong, X., Gottschalk, M., Gourbeyre, C., Griesche, H., Hartmann, J., Hartmann, M., Heinold, B., Herber, A., Herrmann, H., Heygster, G., Hoor, P., Jafariserajehlou, S., Jäkel, E., Järvinen, E., Jourdan, O., Kästner, U., Kecorius, S., Knudsen, E. M., Köllner, F., Kretzschmar, J., Lelli, L., Leroy, D., Maturilli, M., Mei, L., Mertes, S., Mioche, G., Neuber, R., Nicolaus, M., Nomokonova, T., Notholt, J., Palm, M., van Pinxteren, M., Quaas, J., Richter, P., Ruiz-Donoso, E., Schäfer, M., Schmieder, K., Schnaiter, M., Schneider, J., Schwarzenböck, A., Seifert, P., Shupe, M. D., Siebert, H., Spreen, G., Stapf, J., Stratmann, F., Vogl, T., Welti, A., Wex, H., Wiedensohler, A., Zanatta, M., and Zeppenfeld, S.: The Arctic Cloud Puzzle: Using ALOUD/PASCAL Multiplatform Observations to Unravel the Role of Clouds and Aerosol Particles in Arctic Amplification, *Bulletin of the American Meteorological Society*, 100, 841–871, <https://doi.org/10.1175/BAMS-D-18-0072.1>, 2019.
- 760 Wendisch, M., Tegen, I., Neggers, R. A. J., and Spreen, G.: Glimpsing the Ins and Outs of the Arctic Atmospheric Cauldron, *Eos*, 102, <https://eos.org/features/glimpsing-the-ins-and-outs-of-the-arctic-atmospheric-cauldron>, 2021.
- Wendisch, M., Brückner, M., Crewell, S., Ehrlich, A., Notholt, J., Lüpkes, C., Macke, A., Burrows, J. P., Rinke, A., Quaas, J., Maturilli, M., Schemann, V., Shupe, M. D., Akansu, E. F., Barrientos-Velasco, C., Bärfuss, K., Blechschmidt, A.-M., Block, K., Bougoudis, I., Bozem, H., Böckmann, C., Bracher, A., Bresson, H., Bretschneider, L., Buschmann, M., Chechin, D. G., Chylik, J., Dahlke, S., Deneke, H., Dethloff, K., Donth, T., Dorn, W., Dupuy, R., Ebell, K., Egerer, U., Engelmann, R., Eppers, O., Gerdes, R., Gierens, R., Gorodetskaya, I. V., Gottschalk, M., Griesche, H., Gryanik, V. M., Handorf, D., Harm-Altstädter, B., Hartmann, J., Hartmann, M., Heinold, B., Herber, A., Herrmann, H., Heygster, G., Höschel, I., Hofmann, Z., Hölemann, J., Hünnerbein, A., Jafariserajehlou, S., Jäkel, E., Jacobi, C., Janout, M., Jansen, F., Jourdan, O., Jurányi, Z., Kalesse-Los, H., Kanzow, T., Käthner, R., Kliesch, L. L., Klingebiel, M., Knudsen, E. M., Kovács, T., Körtker, W., Krampe, D., Kretzschmar, J., Kreyling, D., Kulla, B., Kunkel, D., Lampert, A., Lauer, M., Lelli, L., von Lerber, A., Linke, O., Löhnert, U., Lonardi, M., Losa, S. N., Losch, M., Maahn, M., Mech, M., Mei, L., Mertes, S., Metzner, E., Mewes, D., Michaelis, J., Mioche, G., Moser, M., Nakoudi, K., Neggers, R., Neuber, R., Nomokonova, T., Oelker, J., Papakonstantinou-Presvelou, I., Pätzold, F., Pefanis, V., Pohl, C., van Pinxteren, M., Radovan, A., Rhein, M., Rex, M., Richter, A., Risse, N., Ritter, C., Rostosky, P., Rozanov, V. V., Donoso, E. R., Saavedra Garfias, P., Salzmann, M., Schacht, J., Schäfer, M., Schneider, J., Schnierstein, N., Seifert, P., Seo, S., Siebert,
- 775

- H., Soppa, M. A., Spreen, G., Stachlewska, I. S., Stapf, J., Stratmann, F., Tegen, I., Viceto, C., Voigt, C., Vountas, M., Walbröl, A., Walter, M., Wehner, B., Wex, H., Willmes, S., Zanatta, M., and Zeppenfeld, S.: Atmospheric and Surface Processes, and Feedback Mechanisms Determining Arctic Amplification: A Review of First Results and Prospects of the (AC)3 Project, *Bulletin of the American Meteorological Society*, 104, E208–E242, <https://doi.org/10.1175/BAMS-D-21-0218.1>, 2023.
- 780 Wesche, C., Steinhage, D., and Nixdorf, U.: Polar aircraft Polar5 and Polar6 operated by the Alfred Wegener Institute, *Journal of large-scale research facilities JLSRF*, 2, <https://doi.org/10.17815/jlsrf-2-153>, 2016.
- Wolf, K., Ehrlich, A., Mech, M., Hogan, R. J., and Wendisch, M.: Evaluation of ECMWF Radiation Scheme Using Aircraft Observations of Spectral Irradiance above Clouds, *Journal of the Atmospheric Sciences*, 77, 2665–2685, <https://doi.org/10.1175/JAS-D-19-0333.1>, 2020.
- Wood, R.: Parametrization of the effect of drizzle upon the droplet effective radius in stratocumulus clouds, *Quarterly Journal of the Royal Meteorological Society*, 126, 3309–3324, <https://doi.org/10.1002/qj.49712657015>, 2000.
- 785 Yang, F., Pan, H.-L., Krueger, S. K., Moorthi, S., and Lord, S. J.: Evaluation of the NCEP Global Forecast System at the ARM SGP Site, *Monthly Weather Review*, 134, 3668–3690, <https://doi.org/10.1175/MWR3264.1>, 2006.
- Young, G., Jones, H. M., Choulaton, T. W., Crosier, J., Bower, K. N., Gallagher, M. W., Davies, R. S., Renfrew, I. A., Elvidge, A. D., Darbyshire, E., Marengo, F., Brown, P. R. A., Ricketts, H. M. A., Connolly, P. J., Lloyd, G., Williams, P. I., Allan, J. D., Taylor, J. W.,
- 790 Liu, D., and Flynn, M. J.: Observed microphysical changes in Arctic mixed-phase clouds when transitioning from sea ice to open ocean, *Atmospheric Chemistry and Physics*, 16, 13 945–13 967, <https://doi.org/10.5194/acp-16-13945-2016>, 2016.
- Zhao, M. and Wang, Z.: Comparison of Arctic clouds between European Center for Medium-Range Weather Forecasts simulations and Atmospheric Radiation Measurement Climate Research Facility long-term observations at the North Slope of Alaska Barrow site, *Journal of Geophysical Research*, 115, <https://doi.org/10.1029/2010JD014285>, 2010.