

Fig. S1 The total N fertilizer input amount under different treatments. Different letters indicate significant differences at the level of $P < 0.05$. N+PK, straw return plus regular inorganic N-P-K fertilizers; 0.75N+PK, straw return plus regular inorganic P-K with 25% N fertilizer reduction; 0.5N+PK, straw return plus regular inorganic P-K with 50% N fertilizer reduction; PK, straw return plus regular inorganic P-K without N fertilizer.




Fig. S2 The soil C:N ratio (a), N:P ratio (b) and C:P ratio (c) under different treatments. Different letters indicate significant differences at the level of $p < 0.05$. N+PK, straw return plus regular inorganic N-P-K fertilizers; 0.75N+PK, straw return plus regular inorganic P-K with 25% N fertilizer reduction; 0.5N+PK, straw return plus regular inorganic P-K with 50% N fertilizer reduction; PK, straw return plus regular inorganic P-K without N fertilizer.




Fig. S3 Non-metric multidimensional scaling ordination showing the fungi (a) and bacteria (b) under different N input levels; significant differences in sample clustering are measured by ANOSIM. N+PK, straw return plus regular inorganic N-P-K fertilizers; 0.75N+PK, straw return plus regular inorganic P-K with 25% N fertilizer reduction; 0.5N+PK, straw return plus regular inorganic P-K with 50% N fertilizer reduction; PK, straw return plus regular inorganic P-K without N fertilizer.

Table S1 The yields and soil chemical properties under different treatments of bulk soil during the experimental process

| Year | Treatment | Yield (t ha ⁻¹) | pH | SOC (g kg ⁻¹) | Total N (g kg ⁻¹) | Total P (g kg ⁻¹) |
|------|-----------|-----------------------------|-------------|---------------------------|-------------------------------|-------------------------------|
| 2019 | N+PK | 11.17±0.73 a | 7.29±0.14 a | 17.60±2.10 a | 0.93±0.02 a | 0.65±0.08 a |
| | 0.75N+PK | 10.91±0.29 a | 7.31±0.20 a | 15.87±3.12 ab | 0.94±0.08 a | 0.61±0.12 a |
| | 0.5N+PK | 9.81±0.32 b | 7.39±0.18 a | 14.01±1.42 b | 0.89±0.10 a | 0.64±0.03 a |
| | PK | 9.93±0.39 b | 7.51±0.20 a | 13.58±0.15 b | 0.86±0.05 a | 0.55±0.27 a |
| 2020 | N+PK | 11.39±0.33 a | 7.27±0.10 a | 17.11±1.95 a | 0.93±0.07 a | 0.69±0.07 a |
| | 0.75N+PK | 12.00±1.19 a | 7.28±0.14 a | 17.01±1.77 a | 0.88±0.21 a | 0.63±0.10 a |
| | 0.5N+PK | 9.88±0.84 b | 7.27±0.11 a | 12.45±0.16 b | 0.86±0.01 ab | 0.67±0.07 a |
| | PK | 9.84±0.44 b | 7.48±0.16 a | 11.76±0.82 b | 0.81±0.03 b | 0.59±0.05 a |
| 2021 | N+PK | 11.41±0.05 ab | 7.25±0.21 a | 21.08±1.82 a | 1.37±0.11 a | 0.67±0.05 a |
| | 0.75N+PK | 11.65±0.06 a | 7.28±0.14 a | 20.95±1.27 a | 1.27±0.02 a | 0.66±0.19 a |
| | 0.5N+PK | 10.08±0.08 bc | 7.25±0.02 a | 14.01±2.01 b | 1.15±0.10 a | 0.49±0.10 b |
| | PK | 8.89±0.13 c | 7.39±0.10 a | 13.33±1.18 b | 1.10±0.28 a | 0.47±0.08 b |

The results show means ± standard deviations (n = 3). Different lowercase letters after values indicate significant differences between each treatment in the same year, P < 0.05. SOC, soil organic carbon; N+PK, straw return plus regular inorganic N-P-K fertilizers; 0.75N+PK, straw return plus regular inorganic P-K with 25% N fertilizer reduction; 0.5N+PK, straw return plus regular inorganic P-K with 50% N fertilizer reduction; PK, straw return plus regular inorganic P-K without N fertilizer.

Table S2 The basic chemical properties of initial and treated straw under different N input levels after straw return

| Treatment | Straw C (g kg^{-1}) | Total N (g kg^{-1}) | Total P (g kg^{-1}) | Total K (g kg^{-1}) |
|-----------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|
| Initial | 485.77 \pm 25.21 a | 6.72 \pm 0.36 c | 2.01 \pm 0.12 a | 21.00 \pm 0.13 a |
| N+PK | 428.86 \pm 17.82 b | 17.20 \pm 0.51 ab | 1.36 \pm 0.11 b | 1.48 \pm 0.16 bc |
| 0.75N+PK | 429.00 \pm 30.21 b | 16.72 \pm 0.45 b | 1.45 \pm 0.10 b | 1.18 \pm 0.14 c |
| 0.5N+PK | 427.72 \pm 29.96 b | 18.15 \pm 1.03 a | 1.81 \pm 0.17 a | 1.86 \pm 0.10 b |
| PK | 446.36 \pm 2.42 b | 18.33 \pm 0.53 a | 1.88 \pm 0.35 a | 1.77 \pm 0.46 b |

The results show means \pm standard deviations ($n = 3$). Different lowercase letters after values indicate significant differences between each treatment, $P < 0.05$. SOC, soil organic carbon; N, nitrogen; P, phosphorus; K, potassium; N+PK, straw return plus regular inorganic N-P-K fertilizers; 0.75N+PK, straw return plus regular inorganic P-K with 25% N fertilizer reduction; 0.5N+PK, straw return plus regular inorganic P-K with 50% N fertilizer reduction; PK, straw return plus regular inorganic P-K without N fertilizer.

Table S3 The abundances of fungal and bacterial abundances across different N fertilizer level treatments after straw return

| Treatment | Fungi abundance ($\times 10^7$ copies g $^{-1}$ soil) | Bacteria abundance ($\times 10^7$ copies g $^{-1}$ soil) | Fungi: Bacteria ratio |
|-----------|---|--|--------------------------|
| N+PK | 0.63 \pm 0.16 bc | 3.15 \pm 0.30 a | 0.20 \pm 0.04 b |
| 0.75N+PK | 0.85 \pm 0.09 a | 2.88 \pm 0.24 ab | 0.30 \pm 0.05 a |
| 0.5N+PK | 0.57 \pm 0.04 c | 2.87 \pm 0.42 ab | 0.20 \pm 0.03 b |
| PK | 0.39 \pm 0.05 d | 2.17 \pm 0.43 b | 0.18 \pm 0.02 c |

The results show means \pm standard deviations (n = 3). Different lowercase letters after values indicate significant differences between each treatment, $P < 0.05$. N+PK, straw return plus regular inorganic N-P-K fertilizers; 0.75N+PK, straw return plus regular inorganic P-K with 25% N fertilizer reduction; 0.5N+PK, straw return plus regular inorganic P-K with 50% N fertilizer reduction; PK, straw return plus regular inorganic P-K without N fertilizer.

Table S4 The fungal and bacterial alpha diversity under different N input levels after straw return

| | Treatment | Chao1 | Richness |
|----------|-----------|------------------|-----------------|
| Fungi | N+PK | 1118.60±71.84 a | 883.00±38.57 a |
| | 0.75N+PK | 1117.82±67.17 a | 796.33±28.45 ab |
| | 0.5N+PK | 1063.37±84.82 a | 781.00±33.87 ab |
| | PK | 1054.50±22.29 a | 772.33±27.54 b |
| Bacteria | N+PK | 5917.52±149.48 a | 4475.00±87.11 a |
| | 0.75N+PK | 5920.19±197.47 a | 4396.67±27.43 a |
| | 0.5N+PK | 5881.07±152.30 a | 4398.33±32.35 a |
| | PK | 5672.76±82.25 a | 4241.00±64.55 b |

The results show means ± standard deviations (n = 3). Different lowercase letters after values indicate significant differences between each treatment, $P < 0.05$. N+PK, straw return plus regular inorganic N-P-K fertilizers; 0.75N+PK, straw return plus regular inorganic P-K with 25% N fertilizer reduction; 0.5N+PK, straw return plus regular inorganic P-K with 50% N fertilizer reduction; PK, straw return plus regular inorganic P-K without N fertilizer.

Table S5 The relative abundances of keystone taxa across different N fertilizer level treatments after straw return

| Treatment | FOTU22 (Module 3) | BOTU21 (Module 1) | BOTU6346 (Module 3) | BOTU240 (Module 2) |
|-----------|----------------------|----------------------|------------------------|-----------------------|
| N+PK | 388.33 | 453.67 | 63.67 | 44.33 |
| 0.75N+PK | 536.67 | 638 | 147 | 79 |
| 0.50N+PK | 367 | 303.33 | 150 | 82.66 |
| PK | 109.33 | 421 | 18.33 | 63 |

The results show means (n = 3). Different lowercase letters after values indicate significant differences between each treatment, $P < 0.05$. N+PK, straw return plus regular inorganic N-P-K fertilizers; 0.75N+PK, straw return plus regular inorganic P-K with 25% N fertilizer reduction; 0.5N+PK, straw return plus regular inorganic P-K with 50% N fertilizer reduction; PK, straw return plus regular inorganic P-K without N fertilizer. FOTU, the OTU in fungi; BOTU, the OTU in Bacteria.