

1 The anomalous thundery month of June 1925 in SW IberiaSpain:

2 description and synoptic analysis

3 Francisco Javier Acero¹, Manuel Antón¹, Alejandro Jesús Pérez Aparicio^{1,2}, Nieves Bravo-Paredes¹,
4 Víctor Manuel Sánchez Carrasco¹, María Cruz Gallego¹, José Agustín García¹, Marcelino Núñez³, Irene
5 Tovar³, Javier Vaquero-Martínez⁴, José Manuel Vaquero¹

6 ¹Departamento de Física, Universidad de Extremadura, Badajoz, Spain

7 ²Earth Remote Sensing Laboratory (EaRSLab) and Institute of Earth Sciences – ICT (Polo de Évora), Instituto de Investigação
8 e Formação Avançada (IIFA), Universidade de Évora, Évora, Portugal

9 ³Agencia Estatal de Meteorología (AEMET), Badajoz, Spain

10 ⁴Departamento de Didáctica de las Ciencias Experimentales y de las Matemáticas, Universidad de Extremadura, Cáceres,
11 Spain

12 *Correspondence to:* J.M. Vaquero (jvaquero@unex.es)

13 **Abstract.** In a routine search for meteorological events with a great impact on society in the Extremadura region (SW interior
14 of Iberia) using newspapers, the month of June 1925 was detected as exceptional due to the large number of thunderstorms
15 associated with significant loss of human lives and material resources.~~In a routine search for meteorological events with a~~
16 ~~great impact on society in the Extremadura region (SW interior of Iberia) using newspapers, the month of June 1925 was~~
17 ~~detected as exceptional due to the large number of electrical storms that occurred and the significant impacts that caused, with~~
18 ~~serious losses in human lives and material resources.~~ This extraordinary month underwent a detailed examination from various,
19 complementary perspectives. Firstly, we reconstructed the history of the events, considering the most impacted locations and
20 the resulting damages. Periodical publications, especially the widely circulated “Extremadura” newspaper in 1925, were
21 pivotal in this regard. Secondly, we scrutinized monthly meteorological variables (precipitation, temperature, and cloudiness)
22 using the lengthiest available data series in Iberia. This aimed to underscore the exceptional characteristics of June 1925.
23 Lastly, we analyzed the synoptic situation of the thunderstorm events by employing 20CR reanalysis data. This approach
24 allowed us to comprehend, from a synoptic perspective, the exceptional nature of this month. Thereby, a combination of a
25 negative North Atlantic Oscillation (NAO) situation, elevated Convective Available Potential Energy (CAPE) values, and
26 abundant total water vapor availability in the region was revealed.~~This anomalous month was analyzed in detail from different,~~
27 ~~complementary perspectives: (i) the reconstruction of the history of the events, taking into account the most affected places~~
28 ~~and the most damaging impacts, from periodical publications (especially the “Extremadura” newspaper, which was the~~
29 ~~newspaper with the largest circulation in the region in 1925); (ii) the study of monthly meteorological variables (precipitation,~~
30 ~~temperature and cloudiness) of the longest series available in Iberia to highlight the exceptional nature of June 1925; and (iii)~~
31 ~~the analysis of the synoptic situation of the thunderstorms events using 20CR reanalysis data to understand from a synoptic~~

32 ~~point of view the exceptionality of this month, with a combination of a negative North Atlantic Oscillation (NAO) situation,~~
33 ~~high Convective Available Potential Energy (CAPE) values, and available water in the area.~~

34 **1 Introduction**

35 Thunderstorms are essential phenomena to understand the climate system (Markson, 2007; Rycroft et al., 2008). In addition
36 to their scientific interest, thunderstorms have important consequences in our society since they produce a huge variety of
37 dangers and problems such as heavy rain, lightning, large hail, tornadoes, etc. (Holle, 2016; Antonescu et al., 2017; Prein and
38 Holland, 2018). The scattered nature of all these phenomena has made their study and prediction difficult until a few decades
39 ago when large databases were available for the scientific community (see, for example, Taszarek et al., 2021).

40 The most affected area by thunderstorms in the Iberian Peninsula is located in the northeast, especially in the mountainous
41 regions of the Pyrenees (north Catalonia and Aragon) and the Iberian system (south Aragón). A climatology of stormy days
42 and electrical discharges was recently published by Núñez Mora et al. (2019). In the scientific literature, several exceptional
43 thunderstorm events in these areas of northeast Iberia can be found. For example, several authors have studied thunderstorms
44 that have produced exceptional episodes of hail, such as the events that occurred in July 2001 (Tudurí et al., 2003), in
45 September 2004 (Ceperuelo et al., 2006) or in June 2006 (Montanyà et al., 2009). In addition, other exceptional cases have
46 been studied, such as the severe thunderstorm on October 4th, 2007, that affected the island of Mallorca (Ramis et al., 2009) or
47 the convective system that affected Catalonia on March 21st, 2012, which produced a tornado (Bech et al., 2015). In all these
48 cases, convective activity was very intense, although both the patterns in the general circulation of the atmosphere and the
49 different local aspects can be very different. Climatological studies on ~~thunderstorms~~ in the rest of Iberia are scarcer. For
50 example, Ezcurra et al. (2008) studied the rain characteristics of ~~thunderstorms~~~~electrical storms~~ in northern Iberia during the
51 five-year period 1992-1996. The establishment of lightning detection networks allowed scientists to carry out interesting
52 studies for periods of around 10 years (Rivas Soriano et al., 2005; Santos et al., 2013). In addition, other studies have analyzed
53 the impact of ~~electrical-thunderstorms~~ ~~storms~~ on social and economic aspects, such as wildfires (García Ortega et al., 2011).

54 In this context, we discovered a notable set of news about thunderstorms in the Spanish historical press during the month of
55 June 1925. These journalistic reports strongly caught our attention since the geographical area where they occurred, the interior
56 of southwest Iberia, is one of the regions of Iberia with fewer days of thunderstorms per year and the consequences described
57 by journalists were exceptional. Therefore, the objectives of this article are (i) to make a detailed description of detrimental
58 effects on lives, goods and infrastructures of that extremely stormy month of June 1925 in southwest Spain from news collected
59 in newspapers, (ii) to carry out an evaluation of the observed meteorological data (precipitation, temperature, and cloudiness),
60 even though these events occurred almost a century ago, and (iii) to analyze the synoptic situation that caused these exceptional
61 thunderstorms.

64 The historical press of the region of Extremadura (southwest of Iberia) has been consulted to obtain information about the
 65 meteorological events. In particular, we analyzed the newspaper “Extremadura”, which led us to discover the unusual period
 66 of thunderstorms that occurred in 1925 affecting this region. The newspaper “Extremadura” was the most important newspaper
 67 in the region at that time, together with the newspaper “Hoy” which appeared later in 1933. Subsequently, the newspaper
 68 virtual library of the Spanish Government (www.prensahistorica.mcu.es) has also been consulted for the period between May
 69 15th to July 15th 1925. The main Extremadura newspapers consulted in this library have been: “La Montaña” and “Correo de
 70 la Mañana”. In addition, one national newspaper “La Correspondencia de España” has been analyzed. Eleven reports of
 71 thunderstorm events in Extremadura were found in the newspaper “Extremadura”, nine in the newspaper “La Montaña”, nine
 72 in the newspaper “Correo de la Mañana” and two in the newspaper “Correspondencia de España”. Some characteristic
 73 examples of the news reports found can be seen in Figure 1 and some basic information about them are listed in Table 1. From
 74 all of them, a database has been created describing each event, its location, the date of the event and the publication of the
 75 news, as well as information on the impact of the event such as economics impacts, human losses, and injured people.

66



67

78 **Figure 1: News clippings from the newspapers “Extremadura”, “Correo de la Mañana” and “La Montaña” (courtesy of the Central**
 79 **Library of the University of Extremadura).**

80

81

82

83

Table 1. Date, newspaper name, title, and a summary of the news that are reproduced in Figure 1 (from left to right).

<u>Date and newspaper name</u>	<u>Title</u>	<u>Summary</u>
<u>15/06/1925</u> <u>La Montaña</u>	<u>La tormenta de esta tarde ha sido de primera clase y de gran aparato “escénico”</u> <u>[This afternoon's thunderstorm was first class and had great "scenic" effects]</u>	<u>There was heavy rain and deafening thunder in the Cáceres area. It was similar to the thunderstorm that occurred on June 7.</u>
<u>15/06/1925</u> <u>La Montaña</u>	<u>Furiosa tormenta. Un joven muere ahogado, sin que aparezca su cadaver</u> <u>[Raging thunderstorm. A young man drowns, but his body is still unavailable]</u>	<u>Raging thunderstorm in Zarza de Granadilla. A shepherd drowns while crossing the “Aldevara” stream. The body is not found, despite the efforts of law enforcement and family members.</u>
<u>11/06/1925</u> <u>La Montaña</u>	<u>La tormenta del miércoles</u> <u>[Wednesday's thunderstorm]</u>	<u>A violent thunderstorm. The worst damage was in Malpartida de Cáceres, with three people injured by lightning.</u>
<u>09/06/1925</u> <u>Correo de la mañana</u>	<u>Horrorosa tormenta</u> <u>[Horrible thunderstorm]</u>	<u>Formidable thunderstorm in Segura de León: streets and houses are flooded, roads and highways are impassable, and there is a great impact on agricultural activities.</u>
<u>11/06/1925</u> <u>Correo de la mañana</u>	<u>De Zafra. Dos ahogados</u> <u>[From Zafra. Two drowned]</u>	<u>A huge thunderstorm caused the Peñaranda stream to rise. Two people drowned at Don Adrián's flour mill, where they were caught by a strong flood.</u>

2.2 Meteorological data and reanalysis

The Spanish Meteorological Agency (Agencia Estatal de Meteorología, AEMET) provided the records for the time series construction of the three meteorological variables analyzed in this work: precipitation (P), temperature (T) and cloudiness (N).

The relationship between the thunderstorm events and rainfall has been studied from 64 accumulated monthly precipitation series homogenized by AEMET (Luna et al., 2012). These time series cover 158 years from 1851 to 2008. Moreover, daily rainfall time series for seven locations placed over Extremadura region were used to analyze the short-term variability of precipitation in this region during June 1925.

With the goal to check the relationship between the thunderstorm events and temperature during June 1925, daily temperature records have been analyzed in this work using 20 long and reliable Spanish series covering the period 1850–2003 (Brunet et al., 2006).

96 The cloudiness observed in June 1925 over Spain was analyzed in this work by means of the so-called parameter of cloudiness
97 in 39 stations. Thus, the parameter of cloudiness (PC) used in our work to characterize the cloudiness is defined (in percentage)
98 as:


$$99 \text{PC} = 50 + 50 \cdot ((O - C)/N) \tag{1}$$

100 where O and C are the number of overcast and cloudless days, respectively, and N is the number of days in a given period
101 (month, season, year). We have used the data provided by Sánchez Lorenzo et al. (2012) who inferred monthly series of the
102 variable given by equation 1 from the number of cloudless and overcast days recorded every month in 39 Spanish stations
103 since 1866. For that, those authors recovered monthly series of cloudless and overcast days since 1865 from different volumes
104 of the publications entitled “Resumen de las observaciones meteorológicas efectuadas en la Península”, edited by AEMET,
105 from 1865 to 1950. Sánchez Lorenzo et al. (2012) inferred monthly series (in percentage) of this parameter since 1866 in these
106 39 Spanish stations from the number of cloudless and overcast days recorded every month. For that, these authors recovered
107 monthly series of cloudless and overcast days since 1865 from different volumes of the publications entitled “Resumen de las
108 observaciones meteorológicas efectuadas en la Península”, edited by AEMET, from 1865 to 1950.

109 Figure 2 shows the distribution of P, T and N stations in the Iberia Peninsula (circumferences and circles). In addition, this
110 plot also displays the location of seven P stations with daily data (inverted triangles) placed over the Extremadura region.

111 Additionally, the utilization of the latest version (version 3) of the NOAA/CIRES/DOE 20th Century Reanalysis (V3) data
112 (provided by the NOAA PSL, Boulder, Colorado, USA, from their website at <https://psl.noaa.gov>) was implemented (Compo
113 et al., 2011; Slivinski et al., 2019). This has been made possible by the latest data assimilation systems and several sets of
114 historical meteorological observations. This particular dataset is well-suited for the intended analysis as it offers a continuous
115 three- dimensional depiction of numerous meteorological variables dating back to 1836, encompassing a significantly longer
116 period compared to the standard NCEP/NCAR (since 1948) or ECMWF (since 1958) Reanalysis datasets. In particular, 20CR
117 uses an ensemble filter data assimilation method. Therefore, a direct estimation of the most likely state of the global atmosphere
118 (for each three-hour period). Moreover, there also is an estimation of the uncertainties in that reanalysis. Additionally, the
119 utilization of the latest version (version 3) of the NOAA/CIRES/DOE 20th Century Reanalysis data (provided by the NOAA
120 PSL, Boulder, Colorado, USA, from their website at <https://psl.noaa.gov>) was implemented (Compo et al., 2011; Slivinski et
121 al., 2019). This particular dataset is well suited for the intended analysis as it offers a continuous three dimensional depiction
122 of numerous meteorological variables dating back to 1871, encompassing a significantly longer period compared to the
123 standard NCEP/NCAR (since 1948) or ECMWF (since 1958) Reanalysis datasets. Evaluating the performance of the 20CR
124 reanalysis in the historical part is not a simple task since it is impossible to make comparisons with other reanalyses and can
125 only be done by comparison with independent observations (Slivinski et al., 2021). Some comparison exercises carried out
126 have been satisfactory. In particular, in our study area, the 20CR results were satisfactory for the extreme precipitation event
127 of autumn 1876 in the Guadiana River basin (Trigo et al., 2014).

128



129

130 **Figure 2: Map of Iberia with the borders of the region of Extremadura and its two provinces. The observatories are marked with**
 131 **blue circumferences (monthly cloudiness data, N), red circumferences (monthly precipitation data, P) or black dots (daily**
 132 **temperature data, T). Moreover, observatories with daily precipitation data in the region of Extremadura are shown with yellow**
 133 **inverted triangles.**

134

135 3 Historical description of the stormy month of June 1925

136 This episode of thunderstorms that occurred in June 1925 had a great impact throughout Extremadura. Figure 3 shows the
 137 position and name of the multiple towns and villages located at the north, center and mainly south of Extremadura where
 138 different kinds of damages caused by the thunderstorms were reported. Extremadura exhibits a diverse orography, significantly
 139 influencing its hydrological patterns. The region has mountainous terrain, such as the Sierra de Gata and Sierra de San Pedro
 140 (in the north and west, respectively), with mountains above 1000 m height, which act as natural barriers to moist air masses
 141 from the Atlantic. Conversely, the plains in the south, like La Serena or La Campiña provide fertile ground for agriculture and
 142 livestock. Moreover, there are several important rivers in Extremadura. The main rivers are the Guadiana and the Tajo, which
 143 flow from east to west. Other smaller rivers are the Alagón, Tiétar, Zújar, Salor, Ardila and Guadiato. These rivers play a
 144 crucial role in the regions climate as they serve as conduits for moisture and influence local weather patterns. The region's
 145 orography influences the air mass movement, specially in the norther mountainous areas, where orographic lift leads to higher

146 precipitation levels. Of course, the rivers contribute to the region's humidity levels, enhancing cloud formation and
147 precipitation.

148 The regional Extremadura newspapers included wide information on the thunderstorms of June 1925 and their impact on the
149 region. An overview of the thunderstorms and their impacts according to the newspaper reports is presented below.

150 The largest city where reports of thunderstorms have been found is Cáceres. This is the most important city in the province of
151 Cáceres, one of the two provinces of the region of Extremadura. According to reports in the newspapers “La Montaña” and
152 “Extremadura” there was a heavy thunderstorm in Cáceres on June 7th, another one on June 10th, a third one around June 14th–
153 15th and a fourth one on June 19th. In three of them (June 7th, 10th, and 14th–15th) there was flooding of streets and houses.
154 Furthermore, the thunderstorm on June 7th lasted for two hours, ~~during which~~ during which there were several lightning strikes,
155 one of which many lightning struck, one of which ec caused a a widespread power blackout generalized power blackout in the
156 city. On the other hand, on June 10th the thunderstorm lasted only ten minutes, but it was of great intensity with torrential rain
157 and huge hailstones that severely damaged the countryside. The center of these two thunderstorms was the area of the city of
158 Cáceres, with no rainfall in the surrounding area.

159 In other places, deaths were reported during some thunderstorms, such as ~~it the storm that~~ occurred in the Zafra, Villalba,
160 Bienvenida and La Lapa zone on June 10th, where a total of four people died, two of them drowned due to the enormous
161 flooding of the Peñaranda riverbank and the other two were struck by lightning in the hut where they were sheltering from the
162 thunderstorm, according to the newspapers “Extremadura” and “Correo de la Mañana”. Another death occurred in Zarza de
163 Granadilla when a man was swept away by the current while trying to ford a stream on June 10th, as reported in the newspaper
164 “La Montaña”. The death of a child who drowned when she was swept away by a stream in the thunderstorm in Berlanga is
165 also to be regretted, according to the news item of June 22nd in the newspaper “Correspondencia de España”, where it is also
166 stated that lightning killed three people in Llerena. The newspaper “Extremadura” reports that, in the village of Montemolín,
167 there were fifteen consecutive days of thunderstorms, killing a man when he was struck by lightning. The same newspaper
168 also reports that another person died from the same cause in the thunderstorm that occurred in Montánchez on June 8th.
169 However, the event with the highest number of deaths was the thunderstorm on June 18th in Higuera de Vargas according to
170 the newspaper “Correo de la Mañana”, in which five people died when they were struck by lightning while sheltering in a hut.


171 As well as the fatalities, there were several injured people and deceased animals. Moreover, besides all these dead people there
172 were many injured people and dead animals. For example, in that same hut in Higuera de Vargas, apart from the death of those
173 five people, four people were injured and eight pigs that were in the vicinity died. Moreover, according to the reports from the
174 newspaper “La Montaña”, there were also two people injured in the thunderstorm of June 10th in Cáceres ~~because they received~~
175 ~~an electric shock when they stumbled into a telephone cable that had come off.~~ Two people suffered burns when they were
176 struck by lightning in Malpartida de Cáceres and three donkeys were killed by the lightning according to the same newspaper.
177 In addition, many animals drowned in different locations. ~~In Cáceres, twelve hens and six sheeps disappeared by the water. In~~
178 ~~Zafra, the overflowing of the river Bodión swept away animals on June 10th, which also happened in Montemolín when the~~

179 ~~streams overflowed, according to the reports of the newspaper “Extremadura”. Furthermore, many animals also perished due~~
180 ~~to lightning strikes. That was the case of fifty one hens and one donkey in Segura de León.~~

181 Another of the most frequent impacts of the thunderstorms were the floods ~~and overflowings~~ that occurred in many places.
182 According to the news reported in the newspaper “Correo de la Mañana”, in Segura de León a strong thunderstorm around
183 June 7th–8th caused the flooding of a multitude of houses and streets. In addition, the strong flow of water caused the
184 watercourses to break in several places, sweeping away animals, devastating the fields, and leaving the trunks of holm oaks
185 bare due to the impact of the stones carried by the current. The same newspaper reports that further north, in Ribera del Fresno,
186 there were also major floods due to a thunderstorm on June 16th. The most insignificant stream was transformed into a mighty
187 river and the streets carried so much water that it was impossible to cross them. In some houses the water reached a height of
188 one meter, collapsing walls and sweeping away everything in its path. A few days later, in the same area, the newspaper
189 “Extremadura” reported a major thunderstorm on June 25th in the village of Hinojosa del Valle, during which the whole village
190 was ~~floodedflooded~~, and several houses were destroyed. In addition, it is reported that a stream overflowed its banks in Jerez
191 de los Caballeros due to another thunderstorm on June 21st. It must not be forgotten the overflowing of the Bodi3n river, the
192 Pe3nara riverbank and the Guadiana river in the thunderstorm on June 10th in the Zafra area mentioned above.

193 It is worth mentioning the damage caused to infrastructures by the intense thunderstorms. There were collapsed bridges, such
194 as the one over the river V3iar during the thunderstorm on June 6th in the area of Montemol3n according to the newspaper
195 “Correo de la Mañana”. Another bridge fell over the Tagus River due to the thunderstorm on June 7th in the area of Santiago
196 del Carbajo according to the newspaper “La Monta3a”. In addition, it is reported that traffic between Santiago del Carbajo and
197 a nearby village called Herrera de Alc3ntara was interrupted. The collapse of houses and walls was also very common in many
198 towns during these thunderstorms, as occurred in Segura de Le3n, C3ceres, Malpartida de C3ceres, Hinojosa del Valle, and
199 Ribera del Fresno.

200 Crop and field damages were extensive in many of the locations where thunderstorms developed, leading to a major economic
201 impact due to the region’s dependence on agriculture at that time. For example, a thunderstorm in Alconera on June 7th
202 destroyed crops and trees, leaving only the subsoil in many places, according to the newspaper “Correo de la Mañana”.
203 Something similar happened according to reports from the newspaper “Extremadura” on June 10th in Aldeanueva del Camino
204 and on June 18th in Eljas, where the water and hail caused considerable damage to the orchards.




206
207
208
209

Figure 3: Geographical distribution of the Extremadura locations affected by the storms occurred in **June 1925** according to the documentary sources consulted in this work. Color shows the number of deaths directly related **to the thunderstorm events to the 1925 thunderstorm events** extracted from the documentary sources (black dots means no deaths reported).

210 **4 Assessing the observed instrumental data**

211 As this episode of thunderstorms in June 1925 led to hard impacts throughout Extremadura, it is necessary to analyze the
212 behavior of rainfall in this month. For this purpose, daily rainfall data in seven locations over Extremadura were used. Figure
213 4 shows daily rainfall in June 1925 for these observatories. The local character of precipitation during thunderstorms is
214 revealed. Most observatories recorded precipitation between June 2nd and 6th, Cornalvo (in the center of the study area) being
215 the one with the highest values. During the rest of the month, thunderstorms and precipitation are more isolated, appearing in
216 some observatories while there was no rain in others. Thunderstorms with rainfall higher than 20 mm ~~day⁻¹/day~~
217 on June ~~7th~~, 2nd-8th, 13th, 16th and 18th.

218



219

220 **Figure 4: Daily rainfall recorded in seven observatories placed over Extremadura in the month of June 1925.**

221

222 In order to analyze if the accumulated rainfall in the month of June of 1925 was remarkable, Figure 5 shows the ranking of
223 that month compared to the remaining 157 June months for the time series of each observatory in peninsular Spain. The eight
224 observatories marked in red represent the places where June 1925 was the first or the second wettest June and are placed in
225 the southwest. In this same area, for most of the observatories, rainfall recorded in June 1925 is among the ten rainiest months
226 of June for the whole time period. On the contrary, there are four observatories in the northwest showing that June 1925 was
227 one of the driest months of June.


228 For the three meteorological variables analyzed in this work (precipitation, temperature, and cloudiness), the standardized
229 anomalies between June 1925 and the average of June of the corresponding variable have been estimated as follows:

230
$$Y = \frac{X_{June1925} - \bar{X}_{June}}{std(X_{June})}, \quad (24)$$

231 being $X_{June1925}$ the value for the variable in June 1925, \bar{X}_{June} and $std(X_{June})$ the mean and the standard deviation of the variable
 232 for the month of June for the whole time series, respectively. In this section, variables such as rainfall, temperature, and
 233 cloudiness are analyzed.

234 Figure 6 (left panel) shows the rainfall anomalies for sixty-four times series located over peninsular Spain. Note that, in order
 235 to allow a better interpretation of the spatial behavior of the results, the anomalies were spatially interpolated by a kriging
 236 procedure. The highest anomalies are located over the southwest of Spain, with the study area showing anomalies over 3, i.e.,
 237 in June 1925 it rained between 3 and 4 times more than normal in a month of June. For these observatories, June 1925 shows
 238 the highest accumulated rainfall of the 158 years. The rainfall anomalies decrease towards the north and northeast of Spain.


239



240

241 **Figure 5: Spatial distribution of the rankings representing the accumulated rainfall in the month of June 1925 among the other June**
 242 **months in the 158 years (1851 to 2008) that make up the complete time series for each observatory. Red numbers represent the**
 243 **observatories where June 1925 is the first or the second wettest June.**

244



245
246 **Figure 6: Rainfall (left), temperature (center) and cloudiness (right) anomalies for June 1925.**


247
248 When studying the relationship between temperature and thunderstorm events, it can be expected that the temperature will be
249 lower than usual in a month as rainy as the one that occurred in the study area. Figure 6 (central panel) shows the monthly
250 temperature anomalies for our time series. Anomalies showing a colder-than-average June 1925 lie in the southwest although
251 they are weak. Similarly as for the rainfall, the temperature anomalies decrease towards the northeast of Spain. Moreover,
252 Figure 6 (right panel) shows the spatial variability of the cloudiness monthly anomalies for June 1925 with respect to the
253 average for the 1866-2010 period in Spain. A clear dependence on latitude can be seen, with negative cloudiness anomalies
254 for all northern locations and positive anomalies for the central and southern sites. It can be seen a clear dependence on latitude,
255 with negative cloudiness anomalies for all northerner locations and positive anomalies for the central and southerner sites. In
256 addition, it is appreciated that the central and southwestern regions of Spain present the highest cloudiness anomalies. Several
257 locations exhibit extremely high cloudiness values in June 1925 compared to all months of June between 1866 and 2010. For
258 example, June 1925 was an absolute cloudiness record in Madrid, Cuenca, and Granada. It marked the second maximum value
259 in Badajoz, Toledo, and Málaga.

260 **5 Synoptic analysis leading to the June 1925 events**

261 In addition to the analysis of temperature, precipitation and cloudiness series, the synoptic situation of each day of June 1925
262 is analyzed in order to understand the reason for the stormy events during the month. For this purpose, the 20CR reanalysis
263 data were used to carry out the analysis. The wind vector (streamlines) and the geopotential height at 250 mb-hPa for each day
264 of June 1925 are plotted in Figure 7. Jet streams are a core of strong westerly winds located in the upper levels of the
265 atmospheretroposphere. Therefore, the jet stream is easily identified in Figure 7. In summer, the polar jet stream is weaker
266 than in winter, and this favors a wavier flow. The polar jet stream in the first days of June reached 50 m/s and the flow began
267 to ripple (Figure 7). The wave broke on the third day of June bringing on a cut-off low located over the southwest of the Iberian

268 Peninsula. During the next few days, the polar jet stream continued wavy, and an anticyclone began to form poleward of the
269 cut-off low. This situation can be assimilated to a blocking system (Barriopedro et al., 2010; Lupo, 2021).


270 The cut-off low pressure system was one of the prominent patterns during June 1925 and the corresponding convection
271 increased precipitation that was very intense locally. This could also explain the increase in cloudiness and lower temperatures
272 than usual for the month of June in this region. Note that the persistent trough and cut-off low pattern shown at 250 hPa and
273 also at 500 hPa is compatible with a strong low level southern flow (700 hPa or 850 hPa) over the area of study, especially
274 about the province of Badajoz, where there is usually a flow from the south and southwest at low levels. However, orographic
275 reinforcement of precipitation is not typical in the south of the province of Badajoz, since the mountains, even if they were
276 aligned perpendicular to the flow, are not high enough. This effect is well known upwind of the southern flow, in the Sierra de
277 los Caballeros (the peak of Tentudía 1104 m and the western summit of Los Bonaes 1053 m), but the locations affected by
278 the storms in 1925 (Figure 3) are all in the lee of the aforementioned flow. In fact, the entire province of Badajoz, except for
279 the southern mountains, can be considered geographically as a large valley of the Guadiana River, open to the west-southwest.
280 That is why this orographic forcing of precipitation does not occur here. Perhaps the specific orography in locations such as
281 Jerez de los Caballeros, Higuera de Vargas, La Lapa, etc., could have had some influence not on the precipitation but on its
282 channeling and could have generated some local effects such as flooding or overflows.



283

284

Figure 7: Wind vector (streamlines) and geopotential height at 250 ~~mb~~hPa for each day of June 1925.



286

287 **Figure 8: Synoptic situation of June 2nd (left), June 10th (center), and June 18th (right) showing an example of pattern types #5, #18,**
 288 **and #21, respectively, according to the classification by Santos et al. (2019). Geopotential height at 500 ~~mb~~-hPa is represented in top**
 289 **panels and SLP in bottom panels.**

290

291 Synoptic pattern classifications are a useful analytical tool for understanding the weather of a region. We will use the synoptic
 292 pattern classification established by Font-Tullot (1983, 2000) to analyze the synoptic situation of each day of June 1925.
 293 Specifically, we will use the newfangled pattern classification carried out by Santos et al. (2019), which updates and improves
 294 the well-known Font-Tullot classification for the Iberian region. This synoptic classification consists of 23 different patterns.
 295 Santos et al. (2015) used the ERA40 reanalyses to review the objective classification of Ribalaygua-Batalla and Borén-Iglesias
 296 (1995). Moreover, the subjective classification of Font-Tullot (1983) was recovered in detail, proposing 23 synoptic patterns,
 297 illustrated with situations of 23 specific dates, from the 1970s-1980s.~~Specifically, we will use the newfangled pattern~~
 298 ~~classification carried out by Santos et al. (2019), which updates and improves the well-known Font-classification for the Iberian~~
 299 ~~region. This synoptic classification consists of 23 different patterns.~~

300 The geopotential height at 500 ~~mb~~-hPa and the Sea Level Pressure (SLP) are analyzed for each day in order to identify which
 301 pattern corresponds to each day. Table 4-2 shows the seven patterns identified for June 1925. Five different patterns are
 302 identified between 1st and 22nd and all are associated with thunderstorms (except the pattern #16, not associated with
 303 thunderstorms, and #21, uncertain) by Santos et al. (2019). The most common patterns are #5 (Azores anticyclone and
 304 peninsular thermal depression), #18 (Ibero-African barometric trough), and #21 (barometric dam). Figure 8 shows an example

of these three patterns showing the SLP (bottom panels) and the geopotential height at 500 ~~mb~~-hPa (top panels). Patterns #5, #18, and #21 are represented in Figure 8 left (June 2nd), center (June 10th), and right (June 18th), respectively. Pattern #5 is associated with storms between May and September, being more frequent in July and August. In addition, pattern #18 is common in June and is associated with fair ealm-weather, although it could be cut-off lows in southern Spain. Finally, pattern #21 is associated with ealm-fair weather with occasional storms, especially in northern Iberia. Between days 23 and 30 June 1925, the most common pattern was #10. This pattern is associated with cold and dry weather in southern Spain. As it can be seen in Section 3 and Figure 4, most of the stormy and rainy days occurred between days 1 and 22. In fact, as discussed in Section 4 in relation to Figure 4, thunderstorms with rainfall higher than 20 mm day⁻¹ were recorded on June 2nd-8th, 13th, 16th and 18th. All these days, except for June 8th, are associated with patterns that could be compatible with thunderstorm or rain (see last column in Table 1). As evident from Section 3 and Figure 4, most stormy and rainy days occurred from day 1 to 22. Consequently, the synoptic analysis conducted in this section aligns with the observations documented in the newspapers. Therefore, the synoptic analysis carried out corresponds to what was recorded in the newspapers.

Table 12: Patterns identified in June 1925 according to the classification by Santos et al. (2019).


Pattern	Brief description	Days	Storm or rain
#5	Azores anticyclone and peninsular thermal depression	1-3, 6, 7, 28, 29	Yes
#8	Atlantic anticyclone and peninsular thermal depression	4, 5	Yes
#10	Gulf of Genoa depression	24-27	No
#16	British-Scandinavian anticyclone	8, 9	No
#18	Ibero-African barometric trough	10-13	Yes
#20	Summer peninsular cold depression	23	Yes
#21	Barometric dam	14-22	Uncertain

Lastly, we have generated synoptic charts of the main meteorological fields, as well as different composites of the monthly mean values and anomalies regarding the climatological period covered by the 20CR reanalysis. A summary of our results is

322 presented in Figure 9, which is made up of six panels. The top two panels show SLP while the middle two panels depict
323 Convective Available Potential Energy (CAPE) and the bottom two panels display total precipitable water. The panels on the
324 right present the composite means of the variables indicated for June 1925 while the panels on the left exhibit the composite
325 anomaly.

326 The top panels of Figure 9 show a typical negative North Atlantic Oscillation (NAO) situation with low pressures west of the
327 British Isles and negative SLP anomalies in southwestern Iberia. The middle panels of Figure 9 reveal that western Iberia had
328 high CAPE values in the context of the Atlantic and Mediterranean region, with positive mean anomalies in western Iberia
329 during June 1925 (the values shown correspond to the composite mean of the entire month). Finally, the bottom panels present
330 high values of precipitable water in the entire atmosphere in southwestern Iberia with the highest values of the anomaly over
331 the region of Extremadura. Note that these monthly anomalies are calculated from the composite mean value (climatology
332 time period selected for the calculus is 1981-2010). Therefore, the exceptional month of June 1925 in Extremadura was
333 characterized by a combination of negative NAO situation, high CAPE values, and ~~total water vapor available~~available water
334 in this area. In any case, note that Figure 9 shows the largest CAPE in Spain for June 1925 was not located exactly in the south-
335 western Spain but in north-western Spain and northern Portugal. It seems the 20CR reanalysis for such early times gives us
336 significant patterns although perhaps the exact location of the details is a little displaced.

337



338

339 **Figure 9: Composite mean (right panels) and composite anomaly (left panels) of SLP, CAPE and precipitable water entire**
 340 **atmosphere for June 1925 in the study area (top, middle, and bottom panels, respectively) from 20CR Reanalysis.**

341

342 6 Conclusions

343 Thunderstorms are crucial for understanding the climate system and have significant societal implications due to their various
344 hazards. The northeastern region of the Iberian Peninsula, particularly the mountainous areas of the Pyrenees and the Iberian
345 system, is highly affected by thunderstorms. Studies have examined exceptional thunderstorm events in this region, including
346 episodes of hail and severe thunderstorms. Climatological studies on storms in Iberia are limited but have explored rain
347 characteristics and the impact on social and economic aspects such as wildfires. A notable set of news reports from June 1925
348 in the interior Southwest of Iberia drew our attention due to the region's infrequent storms and exceptional consequences
349 described by journalists. In this study, we have provided a detailed description of the detrimental effects during that stormy
350 month. Moreover, we have evaluated instrumental data from almost a century ago and have analyzed the synoptic situation
351 that caused these exceptional thunderstorms.

352 The thunderstorms that occurred in June 1925 had a significant impact throughout Extremadura, Spain. Numerous towns and
353 villages in the north, center, and south of Extremadura reported various damages caused by the thunderstorms. The city of
354 Cáceres experienced multiple storms in June, with flooding of streets and houses on the 7th, 10th, and 14th–15th. The
355 thunderstorms in Cáceres were characterized by heavy rain, lightning, and large hailstones that caused power outages and
356 severe damage to the countryside. Other areas such as Zafra, Villalba, Bienvenida, La Lapa, Zarza de Granadilla, and Berlanga
357 also reported deaths and injuries from lightning strikes, flooding, and stream currents. Animals were affected as well, with
358 several cases of dead animals due to lightning strikes or drowning. Flooding and overflowing of rivers and streams were
359 widespread, leading to damaged houses, streets, and fields. Bridges, houses and walls collapsed, and crops and orchards
360 suffered extensive damage. The economic impact on agriculture was significant due to the destruction of crops and trees. These
361 storms had a profound impact on the region, causing loss of lives, injuries, infrastructure damage, and economic losses.

362 During the thunderstorms in June 1925 in Extremadura, the behavior of rainfall in the region was analyzed. Daily rainfall data
363 from seven locations in Extremadura were examined, revealing the local nature of precipitation during thunderstorms. The
364 highest values of precipitation were recorded between June 2nd and 6th, with Cornalvo station experiencing the most significant
365 rainfall. The rest of the month there were more isolated thunderstorms and varying precipitation patterns across the
366 observatories. Several days, including June 7th, 8th, 13th, 16th, and 18th, had thunderstorms with rainfall exceeding 20 mm/day.
367 To determine if the accumulated rainfall in June 1925 was exceptional compared to other June months, a ranking analysis was
368 conducted. Eight observatories in the southwestern region of peninsular Spain marked in red in Figure 5 had either the wettest
369 or second-wettest June on record in 1925. Most observatories in this area ranked among the top 10 rainiest Junes throughout
370 the entire dataset. In contrast, four observatories in the northwest indicated that June 1925 was one of the driest Junes. We also
371 examined standardized anomalies for precipitation, temperature, and cloudiness in June 1925 compared to the long-term
372 averages (1850-2003). The rainfall anomalies were highest in the southwest, indicating that June 1925 had 3 to 4 times more
373 rainfall than the average for a June month. The anomalies decreased towards the north and northeast of Spain. Temperature
374 anomalies were lower than average in the rainy study area, with colder temperatures observed in the southwest. Cloudiness

375 anomalies showed a clear dependence on latitude, with negative anomalies in northern locations and positive anomalies in
376 central and southern regions. Central and southwestern Spain had the highest cloudiness anomalies, with several locations
377 experiencing extremely high cloudiness compared to all other months of June from 1866 to 2010. Overall, June 1925 in
378 Extremadura had significant rainfall, lower temperatures than usual, and increased cloudiness, particularly in the southwestern
379 region.

380 We have analyzed the synoptic situation in June 1925 to understand the occurrence of stormy events during that month. The
381 20CR reanalysis data were used to examine the wind vector and geopotential height at 250 ~~mb~~-hPa for each day of June 1925.
382 The presence of a polar jet stream and its waviness was observed, indicating a wavy flow pattern. The daily synoptic situations
383 during this month show patterns associated with thunderstorms and rainfall in most of the days. Synoptic charts and composites
384 of monthly meteorological fields for June 1925 were also generated. Our analysis suggests a negative NAO situation, with low
385 pressures west of the British Isles and negative sea SLP anomalies in southwestern Iberia. Moreover, we have found high
386 CAPE values in western Iberia, with positive mean anomalies during June 1925, and high values of precipitable water in
387 southwestern Iberia, particularly in Extremadura. Overall, the exceptional month of June 1925 in southwest Iberia was
388 characterized by a combination of a negative NAO situation, high CAPE values, and abundant available water in the region.
389 The analysis carried out in this article sheds light on the most extreme convective processes that can occur over southwest
390 Iberia. The interest in these processes is enormous due to their catastrophic consequences.

391 **Data availability**

392 All raw data used in this study are public.

393 **Author contributions**

394 JMV planned the research; NB-P, IT, and JMV extracted the information from the newspapers; FJA, MA, NB-P, MCG, JAG,
395 MN, and JMV made the formal analysis of the data; FJA, MA, MCG, JAG, MN, IT, and JMV wrote the manuscript draft;
396 FJA, MA, AJPA, NB-P, VMSC, MCG, JAG, MN, IT, JV-M, and JMV reviewed and edited the manuscript.

397 **Competing interest**

398 The authors declare that they have no conflict of interest.

399 **Acknowledgments**

400 This research was supported by the Economy and Infrastructure Counseling of the Junta of Extremadura through project
401 IB20080. A.J.P. Aparicio thanks Universidad de Extremadura and Ministerio de Universidades of the Spanish Government
402 for the award of a postdoctoral fellowship Margarita Salas para la formación de jóvenes doctores (MS-11).

403

404 **References**

405 Antonescu, B., Schultz, D. M., Holzer, A., and Groenemeijer, P.: Tornadoes in Europe: an underestimated threat, *Bull. Am.*
406 *Meteorol. Soc.*, 98, 713–728, 2017.

407 Barriopedro, D., García-Herrera, R., and Trigo, R.M.: Application of blocking diagnosis methods to General Circulation
408 Models. Part I: A novel detection scheme, *Climate Dynamics*, 35(7), 1373–1391, doi:10.1007/s00382-010-0767-5, 2010.

409 Bech, J., Arús, J., Castán, S., Pineda, N., Rigo, T., Montanyà, J., and van der Velde, O.: A study of the 21 March 2012 tornadic
410 quasi linear convective system in Catalonia, *Atmospheric Research*, 158–159, 192–209, doi: 10.1016/j.atmosres.2014.08.009,
411 2015

412 Brunet, M., Saladie, O., Jones, P., Sigró, J., Aguilar, E., Moberg, A., Lister, D., Walther, A., López, D., and Almarza, C.: The
413 development of a new dataset of Spanish daily adjusted temperature series (SDATS) (1850–2003), *Int. J. Climatol.*, 26, 1777–
414 1802, doi: 10.1002/joc.1338, 2006.

415 Ceperuelo, M., Llasat, M.C., López, L., García-Ortega, E., and Sánchez, J.L.: Study of 11 September 2004 hailstorm event
416 using radar identification of 2-D systems and 3-D cells, *Advances in Geosciences*, 7, 215–222, doi: 10.5194/adgeo-7-215-
417 2006, 2006.

418 Compo, G.P., Whitaker, J.S., Sardeshmukh, P.D., Matsui, N., Allan, R.J., Yin, X., et al.: The twentieth century reanalysis
419 project, *Q. J. R. Meteorol. Soc.*, 137, 1–28. doi:10.1002/qj.776, 2011.

420 Ezcurra, A., Saenz, J., Ibarra-Berastegi, G., and Areitio, J.: Rainfall yield characteristics of electrical storm observed in the
421 Spanish Basque Country area during the period 1992–1996, *Atmospheric Research*, 89, 233–242, doi:
422 10.1016/j.atmosres.2008.02.011, 2008.

423 Font-Tullot, I.: *Climatología de España y Portugal*, Instituto Nacional de Meteorología, Madrid, 1983.

424 Font-Tullot, I.: *Climatología de España y Portugal*. Universidad de Salamanca, Salamanca, 2000.

425 García-Ortega, E., Trobajo, M.T., López, L., and Sánchez, J. L.: Synoptic patterns associated with wildfires caused by lightning
426 in Castile and Leon, Spain, *Nat. Hazards Earth Syst. Sci.*, 11, 851–863, doi:10.5194/nhess-11-851-2011, 2011.

427 Holle, R. L.: A summary of recent national-scale lightning fatality studies, *Wea. Clim. Soc.*, 8, 35–42. ~~2006~~2016.

428 Luna, M.Y., Guijarro, J.A., and López, J.A.: A monthly precipitation database for Spain (1851–2008): Reconstruction,
429 homogeneity and trends, *Advances in Science and Research*, 8, 14, doi:10.5194/asr-8-1-2012, 2012.

430 Lupo, A.R.: Atmospheric blocking events: a review, *Annals of the New York Academy of Sciences*, 1504(1), 5–24.
431 <https://doi.org/10.1111/nyas.14557>, 2021.

432 Markson, R.: The global circuit intensity: Its measurement and variation over the last 50 years, *Bull. Amer. Meteor. Soc.*, 88,
433 1–19, 2007.

434 Montanyà, J., Soula, S., Pineda, N., van der Velde, O., Clapers, P., Solà, G., Bech, J., and Romero, D.: Study of the total
435 lightning activity in a hailstorm, *Atmospheric Research*, 91, 430–437, 2009.

436 Núñez Mora, J.Á., Riesco Martín, J., Mora García, M.A.: *Climatología de descargas eléctricas y de días de tormenta en España*.
437 Agencia Estatal de Meteorología, Madrid, 2019.

438 Prein, A.F. and Holland, G. J.: Global estimates of damaging hail hazard, *Weather Clim. Extremes*, 22, 10–23, 2018.

439 Ramis, C., Romero, R., and Homar, V.: The severe thunderstorm of 4 October 2007 in Mallorca: an observational study, *Nat.*
440 *Hazards Earth Syst. Sci.*, 9, 1237–1245, 2009.

441 [Ribalaygua-Batalla, J., Borén-Iglesias, R.: Clasificación de patrones espaciales de precipitación diaria sobre la España](#)
442 [peninsular y Baleárica. Informe N° 3 del Servicio de Análisis e Investigación del Clima, Instituto Nacional de Meteorología,](#)
443 [Madrid, 1995.](#)

444 Rivas Soriano, L., de Pablo, F., and Tomas, C.: Ten-year study of cloud-to-ground lightning activity in the Iberian Peninsula,
445 *Journal of Atmospheric and Solar-Terrestrial Physics* 67, 1632–1639, 2005.

446 Rycroft, M.J., Harrison, R.G., Nicoll, K.A., and Mareev, E.A.: An Overview of Earth’s Global Electric Circuit and
447 Atmospheric Conductivity. In: Leblanc, F., Aplin, K.L., Yair, Y., Harrison, R.G., Lebreton, J.P., and Blanc, M. (eds): *Planetary*
448 *Atmospheric Electricity. Space Sciences Series of ISSI, vol 30.* Springer, New York, NY. doi: 10.1007/978-0-387-87664-1_6
449 (2008)

450 Sanchez-Lorenzo, A., Calbó, J., and Wild, M.: Increasing cloud cover in the 20th century: review and new findings in Spain.
451 *Clim. Past*, 8, 1199–1212, doi:10.5194/cp-8-1199-2012, 2012.

452 Santos, C., Subías, A., and Roa, A.: Recuperación de la clasificación sinóptica de Font: reconstrucción con el reanálisis ERA40,
453 AEMET, Madrid, 2019.

454 Santos, J.A., Reis, M. A., De Pablo, F., Rivas-Soriano, L., Leite, S.M.: Forcing factors of cloud-to-ground lightning over
455 Iberia: regional-scale assessments, *Nat. Hazards Earth Syst. Sci.*, 13, 1745–1758. doi:10.5194/nhess-13-1745-2013, 2013.

456 Slivinski, L.C., Compo, G.P., Whitaker, J.S., Sardeshmukh, P.D., Giese, B.S., McColl, C., Allan, R. et al.: Towards a more
457 reliable historical reanalysis: Improvements for version 3 of the Twentieth Century Reanalysis system, *Quarterly Journal of*
458 *the Royal Meteorological Society*, 145, 2876-2908, doi: 10.1002/qj.3598, 2019.

459 [Slivinski, L.C. et al.: An Evaluation of the Performance of the Twentieth Century Reanalysis Version 3. *Journal of Climate*](#)
460 [34, 1417. <https://doi.org/10.1175/JCLI-D-20-0505.1>, 2021.](#)

461 Taszarek, M., Allen, J.T., Marchio, M. et al.: Global climatology and trends in convective environments from ERA5 and
462 rawinsonde data, *npj Clim. Atmos. Sci.*, 4, 35, doi: 10.1038/s41612-021-00190-x, 2021.

463 [Trigo, R.M. et al.: The record precipitation and flood event in Iberia in December 1876: description and synoptic analysis.](#)
464 [*Front. Earth Sci.* 2, 3. doi: 10.3389/feart.2014.00003, 2014.](#)

465 Tudurí, E., Romero, R., López, L., García, E., Sánchez, J.L., and Ramis, C.: The 14 July 2001 hailstorm in northeastern Spain:
466 diagnosis of the meteorological situation, *Atmospheric Research*, 67–68, 541– 558. 2003.