



# 1 **The anomalous thundery month of June 1925 in SW Iberia:** 2 **description and synoptic analysis**

3 Francisco Javier Acero<sup>1</sup>, Manuel Antón<sup>1</sup>, Alejandro Jesús Pérez Aparicio<sup>1,2</sup>, Nieves Bravo-Paredes<sup>1</sup>,  
4 Víctor Manuel Sánchez Carrasco<sup>1</sup>, María Cruz Gallego<sup>1</sup>, José Agustín García<sup>1</sup>, Marcelino Núñez<sup>3</sup>, Irene  
5 Tovar<sup>3</sup>, Javier Vaquero-Martínez<sup>4</sup>, José Manuel Vaquero<sup>1</sup>

6 <sup>1</sup>Departamento de Física, Universidad de Extremadura, Badajoz, Spain

7 <sup>2</sup>Earth Remote Sensing Laboratory (EaRSLab) and Institute of Earth Sciences – ICT (Polo de Évora), Instituto de Investigação  
8 e Formação Avançada (IIFA), Universidade de Évora, Évora, Portugal

9 <sup>3</sup>Agencia Estatal de Meteorología (AEMET), Badajoz, Spain

10 <sup>4</sup>Departamento de Didáctica de las Ciencias Experimentales y de las Matemáticas, Universidad de Extremadura, Cáceres,  
11 Spain

12 *Correspondence to:* J.M. Vaquero (jvaquero@unex.es)

13 **Abstract.** In a routine search for meteorological events with a great impact on society in the Extremadura region (SW interior  
14 of Iberia) using newspapers, the month of June 1925 was detected as exceptional due to the large number of electrical storms  
15 that occurred and the significant impacts that caused, with serious losses in human lives and material resources. This anomalous  
16 month was analyzed in detail from different, complementary perspectives: (i) the reconstruction of the history of the events,  
17 taking into account the most affected places and the most damaging impacts, from periodical publications (especially the  
18 “Extremadura” newspaper, which was the newspaper with the largest circulation in the region in 1925); (ii) the study of  
19 monthly meteorological variables (precipitation, temperature and cloudiness) of the longest series available in Iberia to  
20 highlight the exceptional nature of June 1925; and (iii) the analysis of the synoptic situation of the thunderstorms events using  
21 20CR reanalysis data to understand from a synoptic point of view the exceptionality of this month, with a combination of a  
22 negative North Atlantic Oscillation (NAO) situation, high Convective Available Potential Energy (CAPE) values, and  
23 available water in the area.

## 24 **1 Introduction**

25 Thunderstorms are essential phenomena to understand the climate system (Markson, 2007; Rycroft et al., 2008). In addition  
26 to their scientific interest, thunderstorms have important consequences in our society since they produce a huge variety of  
27 dangers and problems such as heavy rain, lightning, large hail, tornadoes, etc. (Holle, 2016; Antonescu et al., 2017; Prein and  
28 Holland, 2018). The scattered nature of all these phenomena has made their study and prediction difficult until a few decades  
29 ago when large databases were available for the scientific community (see, for example, Taszarek et al., 2021).



30 The most affected area by thunderstorms in the Iberian Peninsula is located in the northeast, especially in the mountainous  
31 regions of the Pyrenees (north Catalonia and Aragon) and the Iberian system (south Aragón). A climatology of stormy days  
32 and electrical discharges was recently published by Núñez Mora et al. (2019). In the scientific literature, several exceptional  
33 thunderstorm events in these areas of northeast Iberia can be found. For example, several authors have studied thunderstorms  
34 that have produced exceptional episodes of hail, such as the events that occurred in July 2001 (Tudurí et al., 2003), in  
35 September 2004 (Ceperuelo et al., 2006) or in June 2006 (Montanyà et al., 2009). In addition, other exceptional cases have  
36 been studied, such as the severe thunderstorm on October 4<sup>th</sup>, 2007, that affected the island of Mallorca (Ramis et al., 2009) or  
37 the convective system that affected Catalonia on March 21<sup>st</sup>, 2012, which produced a tornado (Bech et al., 2015).  
38 Climatological studies on storms in the rest of Iberia are scarcer. For example, Ezcurra et al. (2008) studied the rain  
39 characteristics of electrical storms in northern Iberia during the five-year period 1992-1996. The establishment of lightning  
40 detection networks allowed scientists to carry out interesting studies for periods of around 10 years (Rivas Soriano et al., 2005;  
41 Santos et al., 2013). In addition, other studies have analyzed the impact of electrical storms on social and economic aspects,  
42 such as wildfires (García Ortega et al., 2011).

43 In this context, we discovered a notable set of news about thunderstorms in the Spanish historical press during the month of  
44 June 1925. These journalistic reports strongly caught our attention since the geographical area where they occurred, the interior  
45 of southwest Iberia, is one of the regions of Iberia with fewer days of thunderstorms per year and the consequences described  
46 by journalists were exceptional. Therefore, the objectives of this article are (i) to make a detailed description of detrimental  
47 effects on lives, goods and infrastructures of that extremely stormy month of June 1925 in southwest Spain from news collected  
48 in newspapers, (ii) to carry out an evaluation of the observed meteorological data (precipitation, temperature, and cloudiness),  
49 even though these events occurred almost a century ago, and (iii) to analyze the synoptic situation that caused these exceptional  
50 thunderstorms.

## 51 **2 Datasets and methodology**

### 52 **2.1 Historical sources**

53 The historical press of the region of Extremadura (southwest of Iberia) has been consulted to obtain information about the  
54 meteorological events. In particular, we analyzed the newspaper “Extremadura”, which led us to discover the unusual period  
55 of thunderstorms that occurred in 1925 affecting this region. The newspaper “Extremadura” was the most important newspaper  
56 in the region at that time, together with the newspaper “Hoy” which appeared later in 1933. Subsequently, the newspaper  
57 virtual library of the Spanish Government ([www.prensahistorica.mcu.es](http://www.prensahistorica.mcu.es)) has also been consulted for the period between May  
58 15<sup>th</sup> to July 15<sup>th</sup> 1925. The main Extremadura newspapers consulted in this library have been: “La Montaña” and “Correo de  
59 la Mañana”. In addition, one national newspaper “La Correspondencia de España” has been analyzed. Eleven reports of  
60 thunderstorm events in Extremadura were found in the newspaper “Extremadura”, nine in the newspaper “La Montaña”, nine  
61 in the newspaper “Correo de la Mañana” and two in the newspaper “Correspondencia de España”. Some characteristic

62 examples of the news reports found can be seen in Figure 1. From all of them, a database has been created describing each  
63 event, its location, the date of the event and the publication of the news, as well as information on the impact of the event such  
64 as economics impacts, human losses, and injured people.



66  
67 **Figure 1: News clippings from the newspapers “Extremadura”, “Correo de la Mañana” and “La Montaña” (courtesy of the Central**  
68 **Library of the University of Extremadura).**


## 69 2.2 Meteorological data and reanalysis

71 The Spanish Meteorological Agency (Agencia Estatal de Meteorología, AEMET) provided the records for the time series  
72 construction of the three meteorological variables analyzed in this work: precipitation (P), temperature (T) and cloudiness (N).  
73 The relationship between the thunderstorm events and rainfall has been studied from 64 accumulated monthly precipitation  
74 series homogenized by AEMET (Luna et al., 2012). These time series cover 158 years from 1851 to 2008. Moreover, daily  
75 rainfall time series for seven locations placed over Extremadura region were used to analyze the short-term variability of  
76 precipitation in this region during June 1925.

77 With the goal to check the relationship between the storm events and temperature during June 1925, daily temperature records  
78 have been analyzed in this work using 20 long and reliable Spanish series covering the period 1850–2003 (Brunet et al., 2006).  
79 The cloudiness observed in June 1925 over Spain was analyzed in this work by means of the so-called parameter of cloudiness  
80 in 39 stations. Sánchez Lorenzo et al. (2012) inferred monthly series (in percentage) of this parameter since 1866 in these 39  
81 Spanish stations from the number of cloudless and overcast days recorded every month. For that, these authors recovered  
82 monthly series of cloudless and overcast days since 1865 from different volumes of the publications entitled “Resumen de las  
83 observaciones meteorológicas efectuadas en la Península”, edited by AEMET, from 1865 to 1950.



84 Figure 2 shows the distribution of P, T and N stations in the Iberia Peninsula (circumferences and circles). In addition, this  
85 plot also displays the location of seven P stations with daily data (inverted triangles) placed over the Extremadura region.  
86 Additionally, the utilization of the latest version (version 3) of the NOAA/CIRES/DOE 20th Century Reanalysis (V3) data  
87 (provided by the NOAA PSL, Boulder, Colorado, USA, from their website at <https://psl.noaa.gov>) was implemented (Compo  
88 et al., 2011; Slivinski et al., 2019). This particular dataset is well-suited for the intended analysis as it offers a continuous three-  
89 dimensional depiction of numerous meteorological variables dating back to 1871, encompassing a significantly longer period  
90 compared to the standard NCEP/NCAR (since 1948) or ECMWF (since 1958) Reanalysis datasets.  
91



92  
93 **Figure 2: Map of Iberia with the borders of the region of Extremadura and its two provinces. The observatories are marked with**  
94 **blue circumferences (monthly cloudiness data, N), red circumferences (monthly precipitation data, P) or black dots (daily**  
95 **temperature data, T). Moreover, observatories with daily precipitation data in the region of Extremadura are shown with yellow**  
96 **inverted triangles.**

97



98 **3 Historical description of the stormy month of June 1925**

99 This episode of thunderstorms that occurred in June 1925 had a great impact throughout Extremadura. Figure 3 shows the  
100 position and name of the multiple towns and villages located at the north, center and mainly south of Extremadura where  
101 different kinds of damages caused by the storms were reported.

102 The regional Extremadura newspapers included wide information on the thunderstorms of June 1925 and their impact on the  
103 region. An overview of the storms and their impacts according to the newspaper reports is presented below.

104 The largest city where reports of thunderstorms have been found is Cáceres. This is the most important city in the province of  
105 Cáceres, one of the two provinces of the region of Extremadura. According to reports in the newspapers “La Montaña” and  
106 “Extremadura” there was a heavy thunderstorm in Cáceres on June 7<sup>th</sup>, another one on June 10<sup>th</sup>, a third one around June 14<sup>th</sup>–  
107 15<sup>th</sup> and a fourth one on June 19<sup>th</sup>. In three of them (June 7<sup>th</sup>, 10<sup>th</sup>, and 14<sup>th</sup>–15<sup>th</sup>) there was flooding of streets and houses.  
108 Furthermore, the thunderstorm on June 7<sup>th</sup> lasted for two hours, during which many lightning struck, one of which caused a  
109 generalized power blackout in the city. On the other hand, on June 10<sup>th</sup> the thunderstorm lasted only ten minutes, but it was of  
110 great intensity with torrential rain and huge hailstones that severely damaged the countryside. The center of these two  
111 thunderstorms was the area of the city of Cáceres, with no rainfall in the surrounding area.

112 In other places, deaths were reported during some thunderstorms, such as the storm that occurred in the Zafra, Villalba,  
113 Bienvenida and La Lapa zone on June 10<sup>th</sup>, where a total of four people died, two of them drowned due to the enormous  
114 flooding of the Peñaranda riverbank and the other two were struck by lightning in the hut where they were sheltering from the  
115 thunderstorm, according to the newspapers “Extremadura” and “Correo de la Mañana”. Another death occurred in Zarza de  
116 Granadilla when a man was swept away by the current while trying to ford a stream on June 10<sup>th</sup>, as reported in the newspaper  
117 “La Montaña”. The death of a child who drowned when she was swept away by a stream in the thunderstorm in Berlanga is  
118 also to be regretted, according to the news item of June 22<sup>nd</sup> in the newspaper “Correspondencia de España”, where it is also  
119 stated that lightning killed three people in Llerena. The newspaper “Extremadura” reports that, in the village of Montemolín,  
120 there were fifteen consecutive days of thunderstorms, killing a man when he was struck by lightning. The same newspaper  
121 also reports that another person died from the same cause in the thunderstorm that occurred in Montánchez on June 8<sup>th</sup>.  
122 However, the event with the highest number of deaths was the storm on June 18<sup>th</sup> in Higuera de Vargas according to the  
123 newspaper “Correo de la Mañana”, in which five people died when they were struck by lightning while sheltering in a hut.

124 Moreover, besides all these dead people there were many injured people and dead animals. For example, in that same hut in  
125 Higuera de Vargas, apart from the death of those five people, four people were injured and eight pigs that were in the vicinity  
126 died. Moreover, according to the reports from the newspaper “La Montaña”, there were also two people injured in the storm  
127 of June 10<sup>th</sup> in Cáceres because they received an electric shock when they stumbled into a telephone cable that had come off.  
128 Two people suffered burns when they were struck by lightning in Malpartida de Cáceres and three donkeys were killed by the  
129 lightning according to the same newspaper. In addition, many animals drowned in different locations. In Cáceres, twelve hens  
130 and six sheeps disappeared by the water. In Zafra, the overflowing of the river Bodi6n swept away animals on June 10<sup>th</sup>, which






131 also happened in Montemolín when the streams overflowed, according to the reports of the newspaper “Extremadura”.  
132 Furthermore, many animals also perished due to lightning strikes. That was the case of fifty one hens and one donkey in Segura  
133 de León.

134 Another of the most frequent impacts of the storms were the floods and overflowings that occurred in many places. According  
135 to the news reported in the newspaper “Correo de la Mañana”, in Segura de León a strong thunderstorm around June 7<sup>th</sup>–8<sup>th</sup>  
136 caused the flooding of a multitude of houses and streets. In addition, the strong flow of water caused the watercourses to break  
137 in several places, sweeping away animals, devastating the fields, and leaving the trunks of holm oaks bare due to the impact  
138 of the stones carried by the current. The same newspaper reports that further north, in Ribera del Fresno, there were also major  
139 floods due to a storm on June 16<sup>th</sup>. The most insignificant stream was transformed into a mighty river and the streets carried  
140 so much water that it was impossible to cross them. In some houses the water reached a height of one meter, collapsing walls  
141 and sweeping away everything in its path. A few days later, in the same area, the newspaper “Extremadura” reported a major  
142 storm on June 25<sup>th</sup> in the village of Hinojosa del Valle, during which the whole village was flooded and several houses were  
143 destroyed. In addition, it is reported that a stream overflowed its banks in Jerez de los Caballeros due to another storm on June  
144 21<sup>st</sup>. It must not be forgotten the overflowing of the Bodión river, the Peñara riverbank and the Guadiana river in the  
145 thunderstorm on June 10<sup>th</sup> in the Zafra area mentioned above.

146 It is worth mentioning the damage caused to infrastructures by the intense thunderstorms. There were collapsed bridges, such  
147 as the one over the river VÍar during the thunderstorm on June 6<sup>th</sup> in the area of Montemolín according to the newspaper  
148 “Correo de la Mañana”. Another bridge fell over the Tagus River due to the thunderstorm on June 7<sup>th</sup> in the area of Santiago  
149 del Carbajo according to the newspaper “La Montaña”. In addition, it is reported that traffic between Santiago del Carbajo and  
150 a nearby village called Herrera de Alcántara was interrupted. The collapse of houses and walls was also very common in many  
151 towns during these thunderstorms, as occurred in Segura de León, Cáceres, Malpartida de Cáceres, Hinojosa del Valle, and  
152 Ribera del Fresno.

153 Crop and field damages were extensive in many of the locations where thunderstorms developed, leading to a major economic  
154 impact due to the region’s dependence on agriculture at that time. For example, a thunderstorm in Alconera on June 7<sup>th</sup>  
155 destroyed crops and trees, leaving only the subsoil in many places, according to the newspaper “Correo de la Mañana”.  
156 Something similar happened according to reports from the newspaper “Extremadura” on June 10<sup>th</sup> in Aldeanueva del Camino  
157 and on June 18<sup>th</sup> in Eljas, where the water and hail caused considerable damage to the orchards.

158




159  
160  
161  
162

**Figure 3: Geographical distribution of the Extremadura locations affected by the storms occurred in 1925 according to the documentary sources consulted in this work. Color shows the number of deaths directly related to the 1925 thunderstorm events extracted from the documentary sources (black dots means no deaths reported).**

#### 163 4 Assessing the observed instrumental data

164 As this episode of thunderstorms in June 1925 led to hard impacts throughout Extremadura, it is necessary to analyze the  
165 behavior of rainfall in this month. For this purpose, daily rainfall data in seven locations over Extremadura were used. Figure  
166 4 shows daily rainfall in June 1925 for these observatories. The local character of precipitation during thunderstorms is  
167 revealed. Most observatories recorded precipitation between June 2<sup>nd</sup> and 6<sup>th</sup>, Cornalvo (in the center of the study area) being  
168 the one with the highest values. During the rest of the month, thunderstorms and precipitation are more isolated, appearing in  
169 some observatories while there was no rain in others. Thunderstorms with rainfall higher than 20 mm/day were recorded on  
170 June 7<sup>th</sup>, 8<sup>th</sup>, 13<sup>th</sup>, 16<sup>th</sup> and 18<sup>th</sup>.

171



172  
 173 **Figure 4: Daily rainfall recorded in seven observatories placed over Extremadura in the month of June 1925.**

174  
 175 In order to analyze if the accumulated rainfall in the month of June of 1925 was remarkable, Figure 5 shows the ranking of  
 176 that month compared to the remaining 157 June months for the time series of each observatory in peninsular Spain. The eight  
 177 observatories marked in red represent the places where June 1925 was the first or the second wettest June and are placed in  
 178 the southwest. In this same area, for most of the observatories, rainfall recorded in June 1925 is among the ten rainiest months  
 179 of June for the whole time period. On the contrary, there are four observatories in the northwest showing that June 1925 was  
 180 one of the driest months of June.

181 For the three meteorological variables analyzed in this work (precipitation, temperature, and cloudiness), the standardized  
 182 anomalies between June 1925 and the average of June of the corresponding variable have been estimated as follows:

183 
$$Y = \frac{X_{June1925} - \underline{X}_{June}}{std(X_{June})}, \quad (1)$$


184 being  $X_{June1925}$  the value for the variable in June 1925,  $\underline{X}_{June}$  and  $std(X_{June})$  the mean and the standard deviation of the variable  
 185 for the month of June for the whole time series, respectively. In this section, variables such as rainfall, temperature, and  
 186 cloudiness are analyzed.

187 Figure 6 (left panel) shows the rainfall anomalies for sixty-four times series located over peninsular Spain. Note that, in order  
 188 to allow a better interpretation of the spatial behavior of the results, the anomalies were spatially interpolated by a kriging  
 189 procedure. The highest anomalies are located over the southwest of Spain, with the study area showing anomalies over 3, i.e.,






190 in June 1925 it rained between 3 and 4 times more than normal in a month of June. For these observatories, June 1925 shows  
191 the highest accumulated rainfall of the 158 years. The rainfall anomalies decrease towards the north and northeast of Spain.  
192



193  
194 **Figure 5: Spatial distribution of the rankings representing the accumulated rainfall in the month of June 1925 among the other June**  
195 **months in the 158 years that make up the complete time series for each observatory. Red numbers represent the observatories where**  
196 **June 1925 is the first or the second wettest June.**

197



198  
199 **Figure 6: Rainfall (left), temperature (center) and cloudiness (right) anomalies for June 1925.**

200  
201 When studying the relationship between temperature and thunderstorm events, it can be expected that the temperature will be  
202 lower than usual in a month as rainy as the one that occurred in the study area. Figure 6 (central panel) shows the temperature  
203 anomalies for our time series. Anomalies showing a colder-than-average June 1925 lie in the southwest although they are




204 weak. Similarly as for the rainfall, the temperature anomalies decrease towards the northeast of Spain. Moreover, Figure 6  
205 (right panel) shows the spatial variability of the cloudiness anomalies for June 1925 with respect to the average for the 1866-  
206 2010 period in Spain. It can be seen a clear dependence on latitude, with negative cloudiness anomalies for all northerner  
207 locations and positive anomalies for the central and southerner sites. In addition, it is appreciated that the central and  
208 southwestern regions of Spain present the highest cloudiness anomalies. Several locations exhibit extremely high cloudiness  
209 values in June 1925 compared to all months of June between 1866 and 2010. For example, June 1925 was an absolute  
210 cloudiness record in Madrid, Cuenca, and Granada. It marked the second maximum value in Badajoz, Toledo, and Málaga.

## 211 **5 Synoptic analysis leading to the June 1925 events**

212 In addition to the analysis of temperature, precipitation and cloudiness series, the synoptic situation of each day of June 1925  
213 is analyzed in order to understand the reason for the stormy events during the month. For this purpose, the 20CR reanalysis  
214 data were used to carry out the analysis. The wind vector (streamlines) and the geopotential height at 250 mb for each day of  
215 June 1925 are plotted in Figure 7. Jet streams are a core of strong westerly winds located in the upper levels of the atmosphere.  
216 Therefore, the jet stream is easily identified in Figure 7. In summer, the polar jet stream is weaker than in winter, and this  
217 favors a wavier flow. The polar jet stream in the first days of June reached 50 m/s and the flow began to ripple (Figure 7). The  
218 wave broke on the third day of June bringing on a cut-off low located over the southwest of the Iberian Peninsula. During the  
219 next few days, the polar jet stream continued wavy, and an anticyclone began to form poleward of the cut-off low. This situation  
220 can be assimilated to a blocking system (Barriopedro et al., 2010; Lupo, 2021).

221




222

223

**Figure 7: Wind vector (streamlines) and geopotential height at 250 mb for each day of June 1925.**

224





225

226 **Figure 8: Synoptic situation of June 2<sup>nd</sup> (left), June 10<sup>th</sup> (center), and June 18<sup>th</sup> (right) showing an example of pattern types #5, #18,**  
 227 **and #21, respectively, according to the classification by Santos et al. (2019). Geopotential height at 500 mb is represented in top**  
 228 **panels and SLP in bottom panels.**

229

230 Synoptic pattern classifications are a useful analytical tool for understanding the weather of a region. We will use the synoptic  
 231 pattern classification established by Font (1983, 2000) to analyze the synoptic situation of each day of June 1925. Specifically,  
 232 we will use the newfangled pattern classification carried out by Santos et al. (2019), which updates and improves the well-  
 233 known Font classification for the Iberian region. This synoptic classification consists of 23 different patterns.

234 The geopotential height at 500 mb and the Sea Level Pressure (SLP) are analyzed for each day in order to identify which  
 235 pattern corresponds to each day. Table 1 shows the seven patterns identified for June 1925. Five different patterns are identified  
 236 between 1<sup>st</sup> and 22<sup>nd</sup> and all are associated with thunderstorms (except the pattern #16, not associated with thunderstorms, and  
 237 #21, uncertain) by Santos et al. (2019). The most common patterns are #5 (Azores anticyclone and peninsular thermal



238 depression), #18 (Ibero-African barometric trough), and #21 (barometric dam). Figure 8 shows an example of these three  
 239 patterns showing the SLP (bottom panels) and the geopotential height at 500 mb (top panels). Patterns #5, #18, and #21 are  
 240 represented in Figure 8 left (June 2<sup>nd</sup>), center (June 10<sup>th</sup>), and right (June 18<sup>th</sup>), respectively. Pattern #5 is associated with storms  
 241 between May and September, being more frequent in July and August. In addition, pattern #18 is common in June and is  
 242 associated with calm weather, although it could be cut-off lows in southern Spain. Finally, pattern #21 is associated with calm  
 243 weather with occasional storms, especially in northern Iberia. Between days 23 and 30 June 1925, the most common pattern  
 244 was #10. This pattern is associated with cold and dry weather in southern Spain. As it can be seen in Section 3 and Figure 4,  
 245 most of the stormy and rainy days occurred between days 1 and 22. Therefore, the synoptic analysis carried out corresponds  
 246 to what was recorded in the newspapers.

247

248 **Table 1:** Patterns identified in June 1925 according to the classification by Santos et al. (2019).

Pattern	Brief description	Days	Storm or rain
#5	Azores anticyclone and peninsular thermal depression	1-3, 6, 7, 28, 29	Yes
#8	Atlantic anticyclone and peninsular thermal depression	4, 5	Yes
#10	Gulf of Genoa depression	24-27	No
#16	British-Scandinavian anticyclone	8, 9	No
#18	Ibero-African barometric trough	10-13	Yes
#20	Summer peninsular cold depression	23	Yes
#21	Barometric dam	14-22	Uncertain

249

250 Lastly, we have generated synoptic charts of the main meteorological fields, as well as different composites of the monthly  
 251 mean values and anomalies regarding the climatological period covered by the 20CR reanalysis. A summary of our results is  
 252 presented in Figure 9, which is made up of six panels. The top two panels show SLP while the middle two panels depict  
 253 Convective Available Potential Energy (CAPE) and the bottom two panels display total precipitable water. The panels on the




254 right present the composite means of the variables indicated for June 1925 while the panels on the left exhibit the composite  
255 anomaly.

256 The top panels of Figure 9 show a typical negative North Atlantic Oscillation (NAO) situation with low pressures west of the  
257 British Isles and negative SLP anomalies in southwestern Iberia. The middle panels of Figure 9 reveal that western Iberia had  
258 high CAPE values in the context of the Atlantic and Mediterranean region, with positive mean anomalies in western Iberia  
259 during June 1925. Finally, the bottom panels present high values of precipitable water in the entire atmosphere in southwestern  
260 Iberia with the highest values of the anomaly over the region of Extremadura. Therefore, the exceptional month of June 1925  
261 in Extremadura was characterized by a combination of negative NAO situation, high CAPE values, and available water in this  
262 area.

263





264  
265  
266  
267

**Figure 9: Composite mean (right panels) and composite anomaly (left panels) of SLP, CAPE and precipitable water entire atmosphere for June 1925 in the study area (top, middle, and bottom panels, respectively) from 20CR Reanalysis.**



## 268 **6 Conclusions**

269 Thunderstorms are crucial for understanding the climate system and have significant societal implications due to their various  
270 hazards. The northeastern region of the Iberian Peninsula, particularly the mountainous areas of the Pyrenees and the Iberian  
271 system, is highly affected by thunderstorms. Studies have examined exceptional storm events in this region, including episodes  
272 of hail and severe thunderstorms. Climatological studies on storms in Iberia are limited but have explored rain characteristics  
273 and the impact on social and economic aspects such as wildfires. A notable set of news reports from June 1925 in the interior  
274 Southwest of Iberia drew our attention due to the region's infrequent storms and exceptional consequences described by  
275 journalists. In this study, we have provided a detailed description of the detrimental effects during that stormy month.  
276 Moreover, we have evaluated instrumental data from almost a century ago and have analyzed the synoptic situation that caused  
277 these exceptional thunderstorms.

278 The thunderstorms that occurred in June 1925 had a significant impact throughout Extremadura, Spain. Numerous towns and  
279 villages in the north, center, and south of Extremadura reported various damages caused by the thunderstorms. The city of  
280 Cáceres experienced multiple storms in June, with flooding of streets and houses on the 7<sup>th</sup>, 10<sup>th</sup>, and 14<sup>th</sup>–15<sup>th</sup>. The  
281 thunderstorms in Cáceres were characterized by heavy rain, lightning, and large hailstones that caused power outages and  
282 severe damage to the countryside. Other areas such as Zafra, Villalba, Bienvenida, La Lapa, Zarza de Granadilla, and Berlanga  
283 also reported deaths and injuries from lightning strikes, flooding, and stream currents. Animals were affected as well, with  
284 several cases of dead animals due to lightning strikes or drowning. Flooding and overflowing of rivers and streams were  
285 widespread, leading to damaged houses, streets, and fields. Bridges, houses and walls collapsed, and crops and orchards  
286 suffered extensive damage. The economic impact on agriculture was significant due to the destruction of crops and trees. These  
287 storms had a profound impact on the region, causing loss of lives, injuries, infrastructure damage, and economic losses.

288 During the thunderstorms in June 1925 in Extremadura, the behavior of rainfall in the region was analyzed. Daily rainfall data  
289 from seven locations in Extremadura were examined, revealing the local nature of precipitation during thunderstorms. The  
290 highest values of precipitation were recorded between June 2<sup>nd</sup> and 6<sup>th</sup>, with Cornalvo station experiencing the most significant  
291 rainfall. The rest of the month there were more isolated thunderstorms and varying precipitation patterns across the  
292 observatories. Several days, including June 7<sup>th</sup>, 8<sup>th</sup>, 13<sup>th</sup>, 16<sup>th</sup>, and 18<sup>th</sup>, had thunderstorms with rainfall exceeding 20 mm/day.  
293 To determine if the accumulated rainfall in June 1925 was exceptional compared to other June months, a ranking analysis was  
294 conducted. Eight observatories in the southwestern region of peninsular Spain marked in red in Figure 5 had either the wettest  
295 or second-wettest June on record in 1925. Most observatories in this area ranked among the top 10 rainiest Junes throughout  
296 the entire dataset. In contrast, four observatories in the northwest indicated that June 1925 was one of the driest Junes. We also  
297 examined standardized anomalies for precipitation, temperature, and cloudiness in June 1925 compared to the long-term  
298 averages (1850-2003). The rainfall anomalies were highest in the southwest, indicating that June 1925 had 3 to 4 times more  
299 rainfall than the average for a June month. The anomalies decreased towards the north and northeast of Spain. Temperature  
300 anomalies were lower than average in the rainy study area, with colder temperatures observed in the southwest. Cloudiness



301 anomalies showed a clear dependence on latitude, with negative anomalies in northern locations and positive anomalies in  
302 central and southern regions. Central and southwestern Spain had the highest cloudiness anomalies, with several locations  
303 experiencing extremely high cloudiness compared to all other months of June from 1866 to 2010. Overall, June 1925 in  
304 Extremadura had significant rainfall, lower temperatures than usual, and increased cloudiness, particularly in the southwestern  
305 region.

306 We have analyzed the synoptic situation in June 1925 to understand the occurrence of stormy events during that month. The  
307 20CR reanalysis data were used to examine the wind vector and geopotential height at 250 mb for each day of June 1925. The  
308 presence of a polar jet stream and its waviness was observed, indicating a wavy flow pattern. The daily synoptic situations  
309 during this month show patterns associated with thunderstorms and rainfall in most of the days. Synoptic charts and composites  
310 of monthly meteorological fields for June 1925 were also generated. Our analysis suggests a negative NAO situation, with low  
311 pressures west of the British Isles and negative sea SLP anomalies in southwestern Iberia. Moreover, we have found high  
312 CAPE values in western Iberia, with positive mean anomalies during June 1925, and high values of precipitable water in  
313 southwestern Iberia, particularly in Extremadura. Overall, the exceptional month of June 1925 in southwest Iberia was  
314 characterized by a combination of a negative NAO situation, high CAPE values, and abundant available water in the region.  
315 The analysis carried out in this article sheds light on the most extreme convective processes that can occur over southwest  
316 Iberia. The interest in these processes is enormous due to their catastrophic consequences.

#### 317 **Data availability**

318 All raw data used in this study are public.

#### 319 **Author contributions**

320 JMV planned the research; NB-P, IT, and JMV extracted the information from the newspapers; FJA, MA, NB-P, MCG, JAG,  
321 MN, and JMV made the formal analysis of the data; FJA, MA, MCG, JAG, MN, IT, and JMV wrote the manuscript draft;  
322 FJA, MA, AJPA, NB-P, VMSC, MCG, JAG, MN, IT, JV-M, and JMV reviewed and edited the manuscript.

#### 323 **Competing interest**

324 The authors declare that they have no conflict of interest.

#### 325 **Acknowledgments**

326 This research was supported by the Economy and Infrastructure Counseling of the Junta of Extremadura through project  
327 IB20080. A.J.P. Aparicio thanks Universidad de Extremadura and Ministerio de Universidades of the Spanish Government  
328 for the award of a postdoctoral fellowship Margarita Salas para la formación de jóvenes doctores (MS-11).

329

#### 330 **References**

331 Antonescu, B., Schultz, D. M., Holzer, A., and Groenemeijer, P.: Tornadoes in Europe: an underestimated threat, *Bull. Am.*  
332 *Meteorol. Soc.*, 98, 713–728, 2017.



- 333 Barriopedro, D., García-Herrera, R., and Trigo, R.M.: Application of blocking diagnosis methods to General Circulation  
334 Models. Part I: A novel detection scheme, *Climate Dynamics*, 35(7), 1373–1391, doi:10.1007/s00382-010-0767-5, 2010.
- 335 Bech, J., Arús, J., Castán, S., Pineda, N., Rigo, T., Montanyà, J., and van der Velde, O.: A study of the 21 March 2012 tornadic  
336 quasi linear convective system in Catalonia, *Atmospheric Research*, 158–159, 192-209, doi: 10.1016/j.atmosres.2014.08.009,  
337 2015
- 338 Brunet, M., Saladie, O., Jones, P., Sigró, J., Aguilar, E., Moberg, A., Lister, D., Walther, A., López, D., and Almarza, C.: The  
339 development of a new dataset of Spanish daily adjusted temperature series (SDATS) (1850-2003), *Int. J. Climatol.*, 26, 1777-  
340 1802, doi: 10.1002/joc.1338, 2006.
- 341 Ceperuelo, M., Llasat, M.C., López, L., García-Ortega, E., and Sánchez, J.L.: Study of 11 September 2004 hailstorm event  
342 using radar identification of 2-D systems and 3-D cells, *Advances in Geosciences*, 7, 215-222, doi: 10.5194/adgeo-7-215-  
343 2006, 2006.
- 344 Compo, G.P., Whitaker, J.S., Sardeshmukh, P.D., Matsui, N., Allan, R.J., Yin, X., et al.: The twentieth century reanalysis  
345 project, *Q. J. R. Meteorol. Soc.*, 137, 1–28. doi:10.1002/qj.776, 2011.
- 346 Ezcurra, A., Saenz, J., Ibarra-Berastegi, G., and Areitio, J.: Rainfall yield characteristics of electrical storm observed in the  
347 Spanish Basque Country area during the period 1992–1996, *Atmospheric Research*, 89, 233-242, doi:  
348 10.1016/j.atmosres.2008.02.011, 2008.
- 349 Font-Tullot, I.: *Climatología de España y Portugal*, Instituto Nacional de Meteorología, Madrid, 1983.
- 350 Font-Tullot, I.: *Climatología de España y Portugal*. Universidad de Salamanca, Salamanca, 2000.
- 351 García-Ortega, E., Trobajo, M.T., López, L., and Sánchez, J. L.: Synoptic patterns associated with wildfires caused by lightning  
352 in Castile and Leon, Spain, *Nat. Hazards Earth Syst. Sci.*, 11, 851–863, doi:10.5194/nhess-11-851-2011, 2011.
- 353 Holle, R. L.: A summary of recent national-scale lightning fatality studies, *Wea. Clim. Soc.*, 8, 35–42. 2006.
- 354 Luna, M.Y., Guijarro, J.A., and López, J.A.: A monthly precipitation database for Spain (1851-2008): Reconstruction,  
355 homogeneity and trends, *Advances in Science and Research*, 8, 14, doi:10.5194/asr-8-1-2012, 2012.
- 356 Lupo, A.R.: Atmospheric blocking events: a review, *Annals of the New York Academy of Sciences*, 1504(1), 5–24.  
357 <https://doi.org/10.1111/nyas.14557>, 2021.
- 358 Markson, R.: The global circuit intensity: Its measurement and variation over the last 50 years, *Bull. Amer. Meteor. Soc.*, 88,  
359 1–19, 2007.
- 360 Montanyà, J., Soula, S., Pineda, N., van der Velde, O., Clapers, P., Solà, G., Bech, J., and Romero, D.: Study of the total  
361 lightning activity in a hailstorm, *Atmospheric Research*, 91, 430–437, 2009.
- 362 Núñez Mora, J.Á., Riesco Martín, J., Mora García, M.A.: *Climatología de descargas eléctricas y de días de tormenta en España*.  
363 Agencia Estatal de Meteorología, Madrid, 2019.
- 364 Prein, A.F. and Holland, G. J.: Global estimates of damaging hail hazard, *Weather Clim. Extremes*, 22, 10–23, 2018.
- 365 Ramis, C., Romero, R., and Homar, V.: The severe thunderstorm of 4 October 2007 in Mallorca: an observational study, *Nat.*  
366 *Hazards Earth Syst. Sci.*, 9, 1237–1245, 2009.



- 367 Rivas Soriano, L., de Pablo, F., and Tomas, C.: Ten-year study of cloud-to-ground lightning activity in the Iberian Peninsula,  
368 *Journal of Atmospheric and Solar-Terrestrial Physics* 67, 1632–1639, 2005.
- 369 Rycroft, M.J., Harrison, R.G., Nicoll, K.A., and Mareev, E.A.: An Overview of Earth’s Global Electric Circuit and  
370 Atmospheric Conductivity. In: Leblanc, F., Aplin, K.L., Yair, Y., Harrison, R.G., Lebreton, J.P., and Blanc, M. (eds): *Planetary*  
371 *Atmospheric Electricity*. Space Sciences Series of ISSI, vol 30. Springer, New York, NY. doi: 10.1007/978-0-387-87664-1\_6  
372 (2008)
- 373 Sanchez-Lorenzo, A., Calbó, J., and Wild, M.: Increasing cloud cover in the 20th century: review and new findings in Spain.  
374 *Clim. Past*, 8, 1199–1212, doi:10.5194/cp-8-1199-2012, 2012.
- 375 Santos, C., Subías, A., and Roa, A.: *Recuperación de la clasificación sinóptica de Font: reconstrucción con el reanálisis ERA40*,  
376 AEMET, Madrid, 2019.
- 377 Santos, J.A., Reis, M. A., De Pablo, F., Rivas-Soriano, L., Leite, S.M.: Forcing factors of cloud-to-ground lightning over  
378 Iberia: regional-scale assessments, *Nat. Hazards Earth Syst. Sci.*, 13, 1745–1758. doi:10.5194/nhess-13-1745-2013, 2013.
- 379 Slivinski, L.C., Compo, G.P., Whitaker, J.S., Sardeshmukh, P.D., Giese, B.S., McColl, C., Allan, R. et al.: Towards a more  
380 reliable historical reanalysis: Improvements for version 3 of the Twentieth Century Reanalysis system, *Quarterly Journal of*  
381 *the Royal Meteorological Society*, 145, 2876-2908, doi: 10.1002/qj.3598, 2019.
- 382 Taszarek, M., Allen, J.T., Marchio, M. et al.: Global climatology and trends in convective environments from ERA5 and  
383 rawinsonde data, *npj Clim. Atmos. Sci.*, 4, 35, doi: 10.1038/s41612-021-00190-x, 2021.
- 384 Tudurí, E., Romero, R., López, L., García, E., Sánchez, J.L., and Ramis, C.: The 14 July 2001 hailstorm in northeastern Spain:  
385 diagnosis of the meteorological situation, *Atmospheric Research*, 67–68, 541– 558. 2003.