

# The ocean's biological and preformed carbon pumps in future steady-state climate scenarios

Benoît Pasquier<sup>1</sup>, Mark Holzer<sup>1</sup>, and Matthew A. Chamberlain<sup>2</sup>

<sup>1</sup>School of Mathematics and Statistics, University of New South Wales, Sydney, NSW, Australia

<sup>2</sup>Commonwealth Scientific and Industrial Research Organisation, Hobart, TAS, Australia

**Correspondence:** Benoît Pasquier (b.pasquier@unsw.edu.au) and Mark Holzer (mholzer@unsw.edu.au)

**Abstract.** The future of the marine carbon cycle is vitally important for climate and for the fertility of the oceans. However, predictions of future biogeochemistry are challenging because a myriad of processes needs parameterization and the future evolution of the physical ocean state is uncertain. Here, we embed a data-constrained model of the carbon cycle in steady circulations that correspond to perpetual 2090s conditions as simulated for the RCP4.5 and RCP8.5 scenarios. Focusing on steady-state changes from preindustrial conditions allows us to capture the response of the system on all timescales, not just on the sub-centennial timescales of typical transient simulations. We find that biological production experiences only modest declines (by 8–12 %) because the reduced nutrient supply due to a more sluggish future circulation and strongly shoaled mixed layers is counteracted by warming-stimulated growth. Organic-matter export declines by 15–25 % due to reductions in both biological production and export ratios, the latter being driven by warming-accelerated shallow respiration and reduced subduction of dissolved organic matter. The future biological pump cycles a 30–70 % larger regenerated inventory accumulated over longer sequestration times, while preformed DIC is shunted away from biological utilization to outgassing. The regenerated and preformed DIC inventories both increase by amounts of similar magnitude. We develop a conceptually new partitioning of preformed DIC to quantify the ocean's preformed carbon pump and its future changes. Near-surface paths of preformed DIC become more important in the future as weakened ventilation isolates the deep ocean. Thus, while regenerated DIC cycling becomes slower, preformed DIC cycling speeds up in the future.

## 1 Introduction

The ocean carbon cycle is a key control on future climate (e.g., Revelle and Suess, 1957; Broecker, 1982; Arias et al., 2021) and the sustained biological productivity of the ocean is essential for future food security (e.g., Golden, 2016; FAO, 2018). Anthropogenic forcing of the climate system will have profound effects on the future carbon cycle and has already begun to impact ocean biogeochemistry (e.g., Riebesell et al., 2009). In recent decades, dissolved oxygen has declined (e.g., Helm et al., 2011; Whitney et al., 2007), the ocean is acidifying (e.g., Doney et al., 2009), and the upper ocean has warmed and stratified (e.g., Li et al., 2020). The question of how the ocean carbon cycle will evolve in the future is thus of intense interest and has stimulated numerous studies (for a small selection see, e.g., Sarmiento and Le Quéré, 1996; Sarmiento et al., 1998; Matear and Hirst, 1999; Plattner et al., 2001; Riebesell et al., 2009; Bopp et al., 2013; Bernardello et al., 2014; Ito et al., 2015; Hauck et al., 2015; Moore et al., 2018; Henson et al., 2022; Wilson et al., 2022; Liu et al., 2023).

Carbon sequestration in the ocean occurs primarily through two mechanisms (Volk and Hoffert, 1985): (i) organic-matter production in the surface ocean is followed by the export of biogenic particles and dissolved organic carbon (DOC) to depth (the biological pump) and (ii) CO<sub>2</sub> is highly soluble in seawater (Weiss, 1974) and newly dissolved inorganic carbon (DIC) is physically transported into the ocean interior with the circulation (the solubility pump). In response to climate change, both the biological and solubility pumps are expected to change. This will have major consequences for the radiative forcing of the atmosphere (without the biological pump, preindustrial atmospheric *p*CO<sub>2</sub> concentrations would have been about 200 ppm higher; see, e.g., Volk and Hoffert, 1985; Holzer et al., 2021b), and for future food security as biological production is expected to diminish (e.g., Kwiatkowski et al., 2020).

For the 21st century and beyond there is broad agreement among models of the fifth and sixth phases of the Coupled Model Intercomparison Project (CMIP5 and CMIP6) that the ocean will sequester more carbon (Bopp et al., 2013; Ito et al., 2015; Hauck et al., 2015; Henson et al., 2022; Wilson et al., 2022; Liu et al., 2023). There is also broad agreement that despite future reductions in productivity and export (e.g., Bopp et al., 2013), the biological pump will cycle a larger pool of regenerated carbon in the future as a slowing overturning circulation increases the residence time of carbon in the deep ocean (e.g., Ito et al., 2015; Wilson et al., 2022; Liu et al., 2023).

However, the future evolution of the ocean carbon cycle remains uncertain (Riebesell et al., 2009; Henson et al., 2022) for many reasons: Future radiative forcing depends on uncertain socioeconomic factors (Meinshausen et al., 2020), numerous physical and complex biogeochemical processes need to be parameterized and calibrated (e.g., Gent et al., 1995; Kriest, 2017; Kriest et al., 2020), and ocean models must be properly spun up so that future changes are not biased by initial drift (e.g., Irving et al., 2021). Moreover, typical centennial simulations are unable to probe the response of the carbon cycle to future conditions on all its natural timescales, which exceed a millennium (Primeau, 2005; Holzer and Primeau, 2010; Holzer et al., 2021a). Recognizing the importance of the slow deep circulation for carbon cycling, some studies have explored the response of the ocean's carbon pumps to the year 2300 (e.g., Moore et al., 2018; Liu et al., 2023), but even that only probes a fraction of the full spectrum of timescales that will shape the long-term evolution of ocean biogeochemistry.

The central question of our work here is: What would the equilibrium steady-state ocean carbon cycle be if the ocean circulation and thermodynamic state at the end of the 21st century were frozen in time? Steady states are advantageous because they probe all the timescales of the system and avoid the complications of transience. To reduce uncertainty due to poorly constrained biogeochemical parameters, we employ a data-constrained model of the carbon cycle with optimized parameters (Pasquier et al., 2023). To avoid the computational costs of model spin-up, we directly solve for equilibrium using steady-state transport matrices built from the ocean circulation as simulated for the 2090s under the RCP4.5 and RCP8.5 future scenarios by the Australian Community Climate and Earth System Simulator (ACCESS1.3; Bi et al., 2013). (Representative Concentration Pathways RCP4.5 and RCP8.5 represent intermediate and worst-case scenarios for future global warming (Meinshausen et al., 2011).)

To comprehensively track all carbon through the ocean, we quantify future change in terms of the biological carbon pump and in terms of what we call the preformed carbon pump. A quantification of the preformed pump is made possible by a novel partition of preformed DIC according to its sources and sinks. The preformed pump defined here allows us to track all preformed DIC back in time to its sources when atmospheric CO<sub>2</sub> entered the ocean, when organic matter was remineralized in the euphotic zone, or when aphotically regenerated DIC resurfaced. In addition, we track preformed DIC forward in time to its sinks, when it will outgas to the atmosphere or be utilized biologically. In this way we quantify the timescales and flow rates (the “plumbing”) of not just the biological pump but also of the preformed pump, as well as the interaction between these pumps and the atmosphere and how these change in the future under idealized steady-state scenarios.

## 2 Methods

### 2.1 Ocean Circulation Models

To build our circulation models, we use physical ocean states from ACCESS1.3 climate-model simulations (Bi et al., 2013) for the preindustrial ocean and for two future climate scenarios. For the preindustrial ocean, we build on the work of Chamberlain et al. (2019) and Pasquier et al. (2023) and use the 1990s average of the circulation, thermodynamic, and forcing fields from the “historical” ACCESS1.3 runs submitted to CMIP5 (Taylor et al., 2012). We refer to this 1990s state as “preindustrial” because the carbon cycle was optimized against DIC observations corrected for anthropogenic DIC. The circulation itself and the other tracers used as constraints are appropriate for the 1990s, but because the preindustrial-to-1990s changes are dwarfed by the centennial changes analysed here, we do not make a further distinction with a true preindustrial state.

For the future ocean states, we use the 2090s average for the ACCESS1.3 CMIP5 runs for the RCP4.5 and RCP8.5 scenarios. We use RCP8.5 (which represents the worst-case scenario for future global warming; Meinshausen et al., 2011) because it provides the strongest perturbation, and the more likely intermediate RCP4.5 to assess the sensitivity of our results to climate-change scenario. We prescribe atmospheric CO<sub>2</sub> ratios at 278, 536, and 886 ppm for the preindustrial-, RCP4.5-, and RCP8.5-based states analyzed below.

The ocean’s advective–diffusive flux-divergence operator is discretized on the numerical grid and organized into “transport matrices” for each state following the approach of Chamberlain et al. (2019). The horizontal advective fluxes across grid-

cell faces are taken from the averaged ACCESS1.3 fields and the vertical fluxes are calculated from mass conservation by integrating up from the seafloor. To speed up numerical solutions, we coarse-grain the grid by lumping horizontally neighboring grid cells  $2 \times 2$  as done by Pasquier et al. (2023). Using the original resolution without re-optimizing parameters degrades the biogeochemical model's match with observations with little improvement in finer-scale features. The coarse-grained grid has a nominal horizontal resolution of  $2^\circ \times 2^\circ$  (finer in latitude near the equator) and 50 depth levels with layer thicknesses that increase from 10 m for the surface layer to 335 m for the deepest layer. We prescribe background horizontal and vertical diffusivities of  $500 \text{ m}^2 \text{ s}^{-1}$  and  $10^{-5} \text{ m}^2 \text{ s}^{-1}$ , respectively, and a mixed-layer vertical diffusivity of  $0.1 \text{ m}^2 \text{ s}^{-1}$ . The transport matrix for the preindustrial state is identical to the ACCESS-M matrix used by Pasquier et al. (2023).

We use the transport matrices to embed our biogeochemical model (PCO2, Pasquier et al., 2023) in the corresponding circulations. The transport matrices and atmospheric  $p\text{CO}_2$  are held fixed in time, and we solve for steady state using an efficient iterative Newton solver. We thus calculate the equilibrium biogeochemical state under perpetual preindustrial and 2090s RCP4.5 and RCP8.5 conditions. While this avoids spin-up issues, we emphasize that our solutions for the future cannot be interpreted as predictions for the 2090s. Instead, our steady-state solutions allow us to probe all the system's natural timescales by determining what the asymptotically long-term adjustment of the carbon cycle would look like if the circulations were frozen in time.

## 2.2 Biogeochemistry Model

We use PCO2, a simple model of the ocean's carbon, nutrient, and oxygen cycles (Pasquier et al., 2023). Here, we only detail the features and parameterizations directly relevant to the present work. Crucially, PCO2 mechanistically couples the carbon, phosphorus, and oxygen cycles capturing important nonlinear interactions and feedbacks in the response of the biogeochemistry to climate change. These include the effect of nutrient supply and temperature on biological production and the effect of temperature and oxygen on bacterial respiration.

Pasquier et al. (2023) objectively optimized the parameters of the PCO2 model for the preindustrial state against the observed concentrations of phosphate ( $\text{PO}_4$ ), DIC, oxygen, and total alkalinity, with DIC observations corrected for anthropogenic carbon. The optimization tends to correct biases from the embedding circulation as well as biases from the biogeochemistry model (see Section 4 for model caveats). We use the same biogeochemical parameters for both preindustrial and future states to capture the response of the carbon cycle to changes in physical ocean state without any changes in plankton physiology.

In a nutshell, PCO2 makes the following simplifying approximations: Phosphate is the only limiting nutrient, which avoids the complexities of the nitrogen cycle and micronutrients such as iron, and the marine ecosystem is approximated by a single phytoplankton species with an implicitly parameterized mortality to avoid the considerable complexities of higher trophic levels. These simplifications are justified a fortiori by the good fit to the observations.

We focus on DIC, which in steady state obeys

$$\mathcal{T}[\text{DIC}] = -U + R + J_{\text{atm}}, \quad (1)$$

where  $\mathcal{T}[\text{DIC}] = \nabla \cdot (\mathbf{u} - \mathbf{K} \nabla)[\text{DIC}]$  is the flux divergence of DIC due to advection (with velocity  $\mathbf{u}$ ) and eddy diffusion (diffusivity tensor  $\mathbf{K}$ ) and the local sources and sinks are on the right. The biological utilization rate per unit volume  $U$  (further detailed below) converts DIC to DOC, fast- and slow-sinking particulate organic carbon ( $\text{POC}_f$  and  $\text{POC}_s$ ), or particulate inorganic carbon (PIC). Organic matter is then remineralized back to DIC through respiration at rates  $R_{\text{DOC}}$ ,  $R_{\text{POC}_f}$ ,  $R_{\text{POC}_s}$ , while PIC is dissolved to DIC at rate  $D_{\text{PIC}}$ . In  $\text{PCO}_2$ , biogenic particles are only transported by gravitational settling, which dominates advective–diffusive transport. All the particles that reach the bottom-most grid box are either remineralized or dissolved there. In Eq. (1),  $R$  denotes the total DIC regeneration rate, i.e.,  $R = R_{\text{DOC}} + R_{\text{POC}_f} + R_{\text{POC}_s} + D_{\text{PIC}}$ . In addition, DIC also has local sources or sinks  $J_{\text{atm}}$  through  $\text{CO}_2$  with the atmosphere (described further below).

The biological DIC uptake rate  $U$  is nonlinearly colimited by temperature, light, and nutrient availability, the latter being additionally modulated by variable C:P uptake stoichiometry parameterized here in terms of phosphate concentration. Specifically we parameterize  $U$  as

$$U = \underbrace{\frac{p_{\text{max}}}{\tau} e^{\kappa T}}_{\beta} \underbrace{\left( \frac{\text{PAR}}{\text{PAR} + k_I} \right)^2}_{\lambda} \underbrace{\left( \frac{[\text{PO}_4]}{[\text{PO}_4] + k_P} \right)^2 \frac{1}{m [\text{PO}_4] + b}}_{\alpha}, \quad (2)$$

where  $\tau = 30$  d is a growth timescale,  $p_{\text{max}} = 23.4 \mu\text{M}$  is a scale for phytoplankton concentration,  $T$  is the water temperature,  $\kappa = 0.063 \text{ K}^{-1}$  sets the  $e$ -folding temperature for growth and mortality, and  $k_I = 10 \text{ W m}^{-2}$  and  $k_P = 3.14 \mu\text{M}$  are the half-saturation constraints for photosynthetically active radiation (PAR) and  $\text{PO}_4$ . The last fraction in Eq. (2) is the C:P uptake ratio parameterized in terms of  $[\text{PO}_4]$  with slope  $m = 6.9 \text{ mmolP molC}^{-1} \mu\text{M}^{-1}$  and intercept  $b = 6.0 \text{ mmolP molC}^{-1}$  (Galbraith and Martiny, 2015). (See Table 1 of Pasquier et al. (2023) for the values of all model parameters.) In Eq. (2) we grouped terms into factors  $\beta$  (temperature-related),  $\lambda$  (light-related), and  $\alpha$  (nutrient-related). With these definitions, the future-minus-preindustrial change  $\Delta U$  can be decomposed into contributions from the changes in each of these factors by writing  $U + \Delta U = (\beta + \Delta\beta)(\lambda + \Delta\lambda)(\alpha + \Delta\alpha)$ . Throughout,  $\Delta X$  denotes the future-minus-preindustrial change in  $X$  with  $X$  being the preindustrial value.

At the surface, carbon enters and exits the ocean through  $\text{CO}_2$  air–sea exchange following the parameterization of Wanninkhof (2014). Specifically, in Eq. (1),  $J_{\text{atm}} = w_0 K_0 (p\text{CO}_2^{\text{atm}} - p\text{CO}_2^{\text{ocn}})/z_0$  is the source/sink of  $[\text{DIC}]$  due to air–sea exchange in the surface layer, where  $w_0$  is the gas-transfer velocity,  $K_0$  is the  $\text{CO}_2$  solubility,  $p\text{CO}_2^{\text{atm}}$  is the atmospheric partial pressure of  $\text{CO}_2$  at the sea surface,  $p\text{CO}_2^{\text{ocn}}$  is the seawater equivalent partial pressure, and  $z_0$  is the thickness of the top model layer ( $K_0$  and  $p\text{CO}_2^{\text{ocn}}$  are computed using the MATLAB CO2SYS package; Lewis and Wallace, 1998; van Heuven et al., 2011). Note that in our model, carbon can only enter or exit the ocean through air–sea exchange so that in steady state  $\int J_{\text{atm}}(\mathbf{r}) d^3\mathbf{r} = 0$ .

### 2.3 Tracking preformed DIC from its sources to its sinks

We partition the DIC concentration into its usual preformed and regenerated components, but explicitly identify the sources and sinks of preformed DIC. Because the re-emergence of regenerated DIC into the euphotic zone is one of the sources of preformed DIC, we first consider regenerated DIC.

To track regenerated DIC, we label DIC during regeneration in the aphotic zone (mask  $\Omega_{\text{aph}} = 1$  in the aphotic zone and 0 otherwise) and immediately unlabel it on entry into the euphotic zone (mask  $\Omega_{\text{eup}} = 1 - \Omega_{\text{aph}}$ ). This unlabelling is conveniently accomplished by fast relaxation to zero with timescale  $\tau_0 = 1$  s. The regenerated DIC concentration  $C_{\text{reg}}$  thus obeys

$$\mathcal{T} C_{\text{reg}} = \Omega_{\text{aph}} R - \Omega_{\text{eup}} C_{\text{reg}} / \tau_0. \quad (3)$$

150  $C_{\text{reg}}$  is then straightforwardly partitioned according to regeneration mechanism (DOC,  $\text{POC}_f$ , or  $\text{POC}_s$  respiration, or PIC dissolution) by replacing  $R$  in Eq. (3) with the corresponding respiration or dissolution rate. The flow rates, residence times, and pathways of the mechanism-partitioned  $C_{\text{reg}}$  are then used to quantify the “plumbing” of the biological pump as in the work of Pasquier et al. (2023).

To track preformed DIC, we define a new labelling tracer that is allowed to roam over the entire ocean, including the euphotic  
155 zone, and for which we diagnose explicit euphotic sources and sinks. This contrasts sharply with the traditional approach where concentrations in the surface ocean (taken as the euphotic zone in many models or above the maximum mixed-layer depth in data-based analyses) are defined as preformed and then propagated into the ocean interior (e.g., Ito and Follows, 2005). Our preformed labelling tracer has the exact same concentrations as obtained by the traditional approach (i.e., by propagating euphotic concentrations), but identifying its sources and sinks makes it possible to quantify the transport of preformed DIC  
160 within the surface ocean, which is not possible with the traditional approach.

Preformed DIC has three sources and two sinks. The sources are gross  $\text{CO}_2$  ingassing at rate  $J_{\text{atm}}^\downarrow = w_0 K_0 p\text{CO}_2^{\text{atm}} / z_0$ , regeneration within the euphotic zone at rate  $\Omega_{\text{eup}} R$ , and emergence of aphotically regenerated DIC into the euphotic zone. When aphotically regenerated DIC enters the euphotic zone, its regenerated label is replaced by the preformed label so that the corresponding rate of labelling preformed DIC (“newly” preformed) is equal to the rate of unlabelling regenerated DIC given  
165 by  $\Omega_{\text{eup}} C_{\text{reg}} / \tau_0$  in Eq. (3). The sinks are biological utilization, removing the preformed label at rate  $U$ , and gross outgassing, removing the preformed label at rate  $J_{\text{atm}}^\uparrow = w_0 K_0 p\text{CO}_2^{\text{ocn}} / z_0$ . The preformed DIC concentration  $C_{\text{pre}}$  thus obeys

$$\mathcal{T} C_{\text{pre}} = J_{\text{atm}}^\downarrow + \Omega_{\text{eup}} R + \Omega_{\text{eup}} C_{\text{reg}} / \tau_0 - J_{\text{atm}}^\uparrow - U. \quad (4)$$

Casting the equation for preformed DIC in this way has major advantages over the traditional boundary-value approach. For the first time, we will be able (i) to partition preformed DIC according to its source and sink mechanisms, (ii) to track the  
170 transport of DIC through the surface ocean, and (iii) to quantify bulk flow rates of preformed DIC without the complications of diffusive one-way fluxes being singular (further discussion in Section 4).

To partition  $C_{\text{pre}}$  according to specified source origin ( $s$ ) and sink destination (loss,  $l$ ), we employ linear labelling tracers (e.g., Holzer et al., 2014; Pasquier and Holzer, 2018; Holzer and DeVries, 2022, details in Appendix A). This provides the preformed concentration  $C_{\text{pre}}^{s \rightarrow l}$ , which we use to quantify the “plumbing” of the ocean’s preformed carbon pump and how it  
175 changes in the future.

### 3 Results

#### 3.1 The future biological pump

The key components of the biological carbon pump are organic-matter production, export of organic matter, and sequestration of regenerated DIC in the aphotic interior. We now examine how each of these components change from their preindustrial values and what the main mechanisms are that drive these changes.

##### 3.1.1 Changes in organic-matter production


Organic-matter production in the euphotic zone, modelled here by Eq. (2) as DIC uptake  $U$ , is a key metric of ocean health and its future is of great importance for food security (e.g., Costello et al., 2020). The circulation of our future states is known to be more sluggish (Holzer et al., 2020) and the question is how this will affect future production.

Production appears to be remarkably resilient: The globally integrated production decreases by 8 % and 12 % for the RCP4.5 and RCP8.5 scenarios, respectively. Given the large physical and thermodynamic changes in our future scenarios, this resilience points to strong compensations between competing mechanisms. The nutrient supply does decline as expected from the more sluggish circulation: The global euphotic phosphate inventories decline by 12 % and 19 % for the two scenarios (Fig. B1a–e). These declines are enhanced by increased Southern Ocean trapping (e.g., Primeau et al., 2013) due to the slower future circulation and decreased ventilation. The decreased nutrient supply reduces biological carbon uptake despite increases in C:P uptake ratios, as parameterized following Galbraith and Martiny (2015). This is counteracted by warming, which exponentially enhances nutrient and carbon uptake rates. The euphotic zone warms globally by 1.5 °C and 2.7 °C (Fig. C1) for RCP4.5 and RCP8.5 although the North Atlantic contains a patch of prominent cooling (e.g., Caesar et al., 2018).

Production declines because of lower nutrient supply and despite warming-enhanced growth. To quantify the drivers of production change  $\Delta U$ , we decompose it algebraically into contributions from changes in nutrient limitation ( $\Delta\alpha$ ), temperature growth factor ( $\Delta\beta$ ), and light limitation ( $\Delta\lambda$ ) (see Eq. (2); Fig. 1). The contributions from nutrient limitation only are  $\beta\lambda\Delta\alpha = -18 \text{ PgC yr}^{-1}$  and  $-28 \text{ PgC yr}^{-1}$  for RCP4.5 and RCP8.5, while the corresponding contributions from the temperature factor only are  $\Delta\beta\lambda\alpha = +11 \text{ PgC yr}^{-1}$  and  $+22 \text{ PgC yr}^{-1}$ . The spatial correlations between  $\Delta\alpha$  and  $\Delta\beta$  (Fig. 1 purple bars) are negative as  $\Delta\alpha$  and  $\Delta\beta$  are of opposite sign and thus reinforce the production decrease. The other terms in the decomposition are less than about  $1 \text{ PgC yr}^{-1}$  in magnitude. Our finding that Southern Ocean nutrient trapping and reduced nutrient supply to the euphotic zone are the primary drivers of production decline is consistent with projections for the next few centuries (e.g., Moore et al., 2018).

##### 3.1.2 Changes in export ratio

Export production,  $J_{\text{ex}}$ , is the rate with which organic carbon taken up at a given location is exported and respired in the aphotic interior (e.g., Primeau et al., 2013; Kwon et al., 2022). The export ratio  $f = J_{\text{ex}}/U$ , is thus a measure of the efficiency of export and is useful for understanding future changes in organic-matter export.  $f$  generally decreases in the future (Fig. D2)



**Figure 1.** (a) Carbon uptake  $U$  for preindustrial conditions (leftmost grey bar) and the RCP4.5-based future state (rightmost grey bar). The colored bars show the future-minus-preindustrial changes  $\Delta U$  due to changes in the factors  $\alpha$  (nutrients),  $\lambda$  (light), and  $\beta$  (temperature) (see Eq. (2)). These changes are plotted as a waterfall chart with each contribution (each bar) starting where the previous one ends. The values of the contributions are indicated in units of PgC yr<sup>-1</sup>. (b) As (a) for RCP8.5.

with production-weighted global means reduced from a preindustrial value of 0.32 to 0.30 and 0.28 for RCP4.5 and RCP8.5, respectively. While export ratios are expected to decline overall with warming, it is understood that locally many factors other than temperature can play an important role (e.g., Laws et al., 2000, 2011; Cael and Follows, 2016). Because of the nonlocal and nonlinear coupling of the mechanisms that set export ratios in our model, a formal decomposition into factors of changes  $\Delta$  is not useful.

Warming stimulates euphotic POC respiration (decreasing export ratios) but causes particles to sink faster through less viscous water (increasing export ratios). For POC, the more rapid sinking does not fully compensate the increased respiration rates (at least for our optimized parameters), so that the net effect is an overall reduction in POC export ratio (consistent with, e.g., Cael and Follows, 2016). For PIC (only dissolution, no respiration), the effect is faster sinking increasing the PIC export ratio, but this has little impact because the preindustrial PIC export ratio is already close to 1. Global euphotic deoxygenation (Fig. B1f–j) is driven by reduced oxygen solubility and reduced photosynthesis (except in the Weddell and Ross Seas due to future sea-ice loss), which slows euphotic POC respiration thereby increasing POC export ratios. DOC export



ratios are expected to decrease with the circulation slowdown because DOC is exported by water transport. Zonal integrals of  
220 the contributions from each export pathway (DOC, POC<sub>s</sub>, POC<sub>f</sub>, and PIC) are shown in Fig. D3.


### 3.1.3 Changes in export production

We now consider changes in carbon export production  $J_{\text{ex}}$  itself. Figure 2 shows  $J_{\text{ex}}$  for the preindustrial state and for the two  
future scenarios along with the corresponding future-minus-preindustrial change,  $\Delta J_{\text{ex}}$ . Globally integrated,  $J_{\text{ex}}$  decreases  
by 14 % and 24 % in the RCP4.5 and RCP8.5 scenarios, respectively. This overall decrease in export production is driven by  
225 the combined action of changes in temperature, PAR, and euphotic nutrient and oxygen concentrations, themselves driven by  
changes in ocean circulation and air–sea exchange.

Large reductions in  $J_{\text{ex}}$  (Fig. 2g–j) occur approximately in regions of large preindustrial export production with the Southern  
Ocean being most affected. In the Ross Sea, a large decrease in  $[\text{PO}_4]$  results in a significant decrease in uptake and export. This  
is because the unrealistic deep mixed layer of the preindustrial ACCESS circulation (Bi et al., 2013), which brings nutrient-  
230 rich deep waters to the surface, disappears in the future states (Fig. C2). By contrast, in the Weddell Sea, export production  
increases despite the disappearance of the unrealistically deep mixed layer because retreating sea ice allows for additional  
photosynthesis. There are also patches of export increases of similar magnitude in the Sea of Okhotsk, in the mid-latitude  
southern Atlantic and Indian Ocean, and in the North Atlantic. The patterns of  $\Delta J_{\text{ex}}$  broadly correlate with those of  $\Delta U$  but  
some notable differences (Figs. 2 and D1) suggest that changes in organic-matter production alone do not suffice to explain  
235 changes in export production.

To quantify the role of changes in organic-matter production ( $\Delta U$ ) and changes in export ratio ( $\Delta f$ ) in shaping future export  
production, we decompose  $\Delta J_{\text{ex}}$  into contributions from  $\Delta U$  and  $\Delta f$  and their spatial correlation. We find that in the global  
mean changes in export ratio alone  $U \Delta f$  contribute about 50 % more to  $\Delta J_{\text{ex}}$  than changes in production alone  $f \Delta U$  (Fig 3).  
The cross-term contribution  $\Delta f \Delta U$  captures the spatial correlation between  $\Delta U$  and  $\Delta f$  and is positive but about an order  
240 of magnitude smaller (Fig 3c,f). The declines in future export production are thus driven by both changes in production and  
export ratios, but the latter make larger contributions globally, particularly for the RCP8.5 scenario.


The spatial patterns of the contributions  $U \Delta f$  and  $f \Delta U$  to the export-production changes are very different in character,  
with  $U \Delta f$  being negative almost everywhere (Fig. 3a,d) and  $f \Delta U$  having both signs (Fig. 3b,e). In the Northern Hemisphere,  
many features of  $U \Delta f$  mirror (with opposite sign) the patterns of the surface temperature and oxygen changes (Figs. C1 and  
245 B1i,j) suggesting important contributions from both. This is not the case in the Southern Ocean, where changes in export ratios  
are likely driven by changes in DOC export driven in turn by the large circulation changes. The pattern of  $f \Delta U$  shows that  
 $\Delta U$  drives both decreases (negative) and increases (positive over about 40 % of the ocean) in export production. The changes  
in organic-matter production itself  $\Delta U$  are due to the competing effects of changes in nutrient concentrations, temperature, and  
light as documented for the global integral above (Fig. 1).  $f \Delta U$  is largest in magnitude at high latitudes where it is dominated  
250 by the pattern of  $\Delta U$  with zonal bands of alternating sign indicating meridional shifts in organic-matter production. Overall, the  
contributions  $U \Delta f$  tend to dominate near the equator while the contributions  $f \Delta U$  from changes in organic-matter production  
are more important at high latitudes, where  $f$  is larger (Fig. D2).



**Figure 2.** (a) Map of export production,  $J_{\text{ex}}(\mathbf{r})$ , vertically integrated, in the preindustrial state. (b) Zonal integral of (a). (c–d) As (a–b) for the future RCP4.5-based state. (e–f) As (c–d) for RCP8.5. (g–j) As (c–f) for  $\Delta J_{\text{ex}}$ .

### 3.2 Global changes in carbon sequestration

The sequestration strengths of the biological and preformed carbon pumps are conveniently quantified by the sizes of the regenerated and preformed inventories. Table 1 lists the total, regenerated, and preformed DIC inventories ( $\mu_{\text{tot}}$ ,  $\mu_{\text{reg}}$ , and  $\mu_{\text{pre}}$ ) for the preindustrial and future states and the corresponding preindustrial-to-future changes. The ocean's total carbon



**Figure 3.** (a,b,d,e) Maps of vertically integrated contributions from changes in export ratio  $U \Delta f$  (a,d) and carbon uptake  $f \Delta U$  (b,e) to the changes in export production  $\Delta J_{ex}$ , in the RCP4.5 (a,b) and RCP8.5 (d,e) scenarios. (c,f) Corresponding zonal integrals of  $U \Delta f$  (blue),  $f \Delta U$  (orange), and  $\Delta J_{ex}$  (black). Also shown is the zonal integral of  $\Delta f \Delta U$  (gray).

**Table 1.** Total, regenerated, and preformed DIC inventories for the preindustrial, RCP4.5, and RCP8.5 states in units of PgC. Changes relative to the preindustrial state are also shown.

State	$\mu_{tot}$	$\Delta\mu_{tot}$	$\mu_{reg}$	$\Delta\mu_{reg}$	$\mu_{pre}$	$\Delta\mu_{pre}$
Preindustrial	35500		1900		33600	
Future RCP4.5	37200	+1700	2500	+600	34800	+1200
Future RCP8.5	38400	+2900	3300	+1400	35100	+1500

inventory increases by 5% and 8% for the RCP4.5 and RCP8.5 scenarios, respectively. Contributions to the changes from regenerated and preformed DIC are on the same order despite being driven by distinct mechanisms, which we will now explore in detail.

260 To trace the pathways of regenerated DIC, we plot in Fig. 4 the zonal mean  $C_{reg}$  and its future-minus-preindustrial change,  $\Delta C_{reg}$ . The zonal-mean  $\Delta C_{reg}$  (Fig. 4j–o) shows large increases throughout the entire deep ocean of up to about 100  $\mu\text{M}$  and 200  $\mu\text{M}$  for the RCP4.5 and RCP8.5 scenarios, respectively. The increases are most prominent in the deep Southern Ocean, where a slower circulation increasingly traps regenerated DIC. For the RCP4.5-based scenario, the Southern Ocean increase

is limited in the Atlantic sector because the unrealistically deep mixed layer, which short-circuits Southern Ocean nutrient trapping (Pasquier et al., 2023), has not completely shoaled. The slight decreases at low latitudes and thermocline depths are likely due to decreased tropical organic-matter production.


The regenerated DIC inventory  $\mu_{\text{reg}}$  is governed by the product of globally integrated export production  $\Phi_{\text{ex}} = \int J_{\text{ex}}(\mathbf{r}) d^3\mathbf{r}$  and bulk sequestration time  $\Gamma_{\text{reg}} = \mu_{\text{reg}}/\Phi_{\text{ex}}$  for which the respired exported organic matter is allowed to accumulate in the aphotic ocean (e.g., DeVries et al., 2012; Holzer et al., 2021b). We find that the fraction of DIC that is regenerated increases from its preindustrial values of 5.4 % to 6.7 % (a 25 % increase) for RCP4.5 and to 8.5 % (a 60 % increase) for RCP8.5 (Fig. 5). We will now show that these increases are driven primarily by changes in the circulation pathways that return regenerated DIC to the euphotic zone.

The pie charts of Fig. 5 show the fractional contributions to the global regenerated DIC inventory from the export of DOC, POC<sub>s</sub>, POC<sub>f</sub>, and PIC, for the preindustrial state and each future scenario. The contributions from each of these export mechanisms are remarkably impervious to change with future-minus-preindustrial differences within 1–3 % for both RCP4.5 and RCP8.5. This is surprising because DOC, POC<sub>s</sub>, POC<sub>f</sub>, and PIC do not remineralize at the same location and particularly not at the same depth.


To demonstrate that bulk sequestration time is the key control on the biological pump we partitioned the regenerated DIC inventory into separate pools according to regeneration mechanism and calculated the flow-rate through each pool (equal to the volume-integrated regeneration rate). Each pool may be considered to be a “pipe” in the “plumbing” of the biological pump, with the bulk sequestration time for each pipe simply being the ratio of inventory in the pipe to the corresponding flow rate. These pipes are depicted as horizontal bars in Figure 5 whose length, width, and area quantify the bulk sequestration time (yr), flow rate (PgC yr<sup>-1</sup>), and inventory (PgC), respectively (as done previously for the optimized preindustrial state by Pasquier et al., 2023).

Figure 5 shows that for each export mechanism (DOC, POC<sub>s</sub>, POC<sub>f</sub>, and PIC), export production declines and sequestration time increases. The increases in sequestration time are consistent with an overall slowdown of the circulation and with longer re-exposure times for regenerated DIC to return to the euphotic zone (Fig. C3). Across all mechanisms, export production rates decline by 8–16 % and 11–27 % for RCP4.5 and RCP8.5. These export changes affect POC<sub>s</sub> the most because the pattern of its export-ratio changes correlate well with preindustrial production (see Figs. D3b,f and D1) and PIC the least because PIC dissolution in our model is independent of temperature, oxygen, or circulation. Across all mechanisms, re-exposure times increase by 40–50 % and 100–160 % for RCP4.5 and RCP8.5. Re-exposure times of DIC regenerated from slow-sinking POC<sub>s</sub> are the most affected, likely because POC<sub>s</sub> is respired at an average depth of about 200 m, where the relative changes in re-exposure times are the largest (Fig. C3).

The effect of increased bulk sequestration times wins over the effect of decreased export production rates. This is shown in Table 2, where the change in each pipe’s regenerated DIC inventory  $\Delta\mu_{\text{reg}}$  is formally decomposed into contributions from  $\Delta\Gamma_{\text{reg}}$  and  $\Delta\Phi_{\text{ex}}$ , including cross terms. Overall, regenerated inventories increase by 25–40 % and 60–90 % across all regeneration mechanisms for the RCP4.5 and RCP8.5 scenarios. In both scenarios, the circulation changes quantified here by  $\Delta\Gamma_{\text{reg}}$  are the dominant control: The increase in regenerated inventory due to longer re-exposure times alone ( $\Phi_{\text{ex}} \Delta\Gamma_{\text{reg}}$ ) is  $\sim 4$ –6 times



**Figure 4.** (a–c) Zonal mean  $C_{reg}$  for the preindustrial state in the Atlantic (a), Pacific (b), and Indian Ocean (c). For all zonal means shown in this work, the Atlantic basin excludes the Gulf of Mexico and the Caribbean, and the Pacific basin excludes the Sea of Japan so that the averages are more cleanly interpretable. (d–f) As (a–c) for the future RCP4.5-based state. (g–i) As (d–f) for RCP8.5. (j–o) As (d–i) for  $\Delta C_{reg}$ .



**Figure 5.** (a) Regenerated DIC contributions from each mechanism  $k = \text{DOC}, \text{POC}_s, \text{POC}_f,$  and  $\text{PIC}$  (blue, orange, red, and grey) for the preindustrial state represented as both a pie chart and a bar chart. The pie chart quantifies the fractional contribution  $\mu_{\text{reg}}^k / \mu_{\text{reg}}$ . Pump sequestration strength, as quantified by  $\mu_{\text{reg}} / \mu_{\text{DIC}}$ , is indicated above the pie chart. Each bar can be interpreted as a pipe flowing from left to right and whose area represents the regenerated DIC inventory,  $\mu_{\text{reg}}^k$ , which equals globally integrated export production,  $\Phi_{\text{ex}}^k$  (pipe width,  $y$ -axis) multiplied by bulk residence time,  $\Gamma_{\text{reg}}^k$  (pipe length,  $x$ -axis). (b) As (a) for the future RCP4.5 state. (c) As (b) for RCP8.5.

larger than the decrease due to smaller export production alone ( $\Gamma_{\text{reg}} \Delta \Phi_{\text{ex}}$ ). In addition, the cross term  $\Delta \Gamma_{\text{reg}} \Delta \Phi_{\text{ex}}$ , which captures the spatial correlations between  $\Delta \Phi_{\text{ex}}$  and  $\Delta \Gamma_{\text{reg}}$ , makes contributions on the same order as  $\Gamma_{\text{reg}} \Delta \Phi_{\text{ex}}$ , underlining the highly non-linear response of the biological pump in both scenarios.

**Table 2.** Contributions to the change in the regenerated carbon inventory  $\Delta\mu_{\text{reg}}^k$  from changes in bulk residence time  $\Gamma_{\text{reg}}^k$  and export production  $\Phi_{\text{ex}}^k$  for each pump mechanism  $k$  (values in PgC rounded to two significant digits).


Scenario	Mechanism $k$	$\Delta\mu_{\text{reg}}^k$	=	$\Phi_{\text{ex}}^k \Delta\Gamma_{\text{reg}}^k$	+	$\Gamma_{\text{reg}}^k \Delta\Phi_{\text{ex}}^k$	+	$\Delta\Gamma_{\text{reg}}^k \Delta\Phi_{\text{ex}}^k$
RCP4.5	total	+580		+950		-240		-120
	DOC	+73		+130		-38		-16
	POC <sub>s</sub>	+140		+230		-59		-38
	POC <sub>f</sub>	+210		+380		-110		-52
	PIC	+160		+210		-35		-18
RCP8.5	total	+1400		+2200		-390		-460
	DOC	+190		+320		-63		-66
	POC <sub>s</sub>	+320		+570		-98		-160
	POC <sub>f</sub>	+490		+860		-180		-190
	PIC	+340		+440		-48		-51

### 3.3 The preformed carbon pump

To assess the future state of the biological pump, it is useful to place it in the context of overall carbon sequestration by also considering the preformed DIC pool (e.g., Ito et al., 2015). Our approach differs conceptually from previous quantifications of the abiotic carbon cycle in terms of the solubility pump (e.g., Volk and Hoffert, 1985), where one typically focuses on the sub-surface pathways of preformed DIC, and efficiency is defined relative to complete surface saturation. By contrast, our new preformed DIC tracer allows us to quantify not only interior but also euphotic DIC pathways, how they change in the future, and how they affect the ocean’s entire DIC inventory. The “preformed carbon pump” considered here is thus the natural counterpart of the biological pump.

We partition the ocean’s preformed DIC into pools that connect specified sources of preformed DIC (ingassing, euphotic regeneration, or emergence of aphotically regenerated DIC) to specified sinks (outgassing or biological uptake). Each pool may again be considered to be a “pipe” of the ocean’s preformed DIC pump with the flow rate through each pool determining the corresponding bulk residence time. To the best of our knowledge, this is the first time that preformed DIC has been partitioned in this way (see Methods and Appendix A for mathematical details).

Figure 6 shows the preformed DIC inventory  $\mu_{\text{pre}}$ , partitioned into “pipes” that connect each source  $s$  to each loss  $l$ , for the preindustrial and future states. As in Fig. 5, the inventory  $\mu_{\text{pre}}^{s \rightarrow l}$  (area) of each pipe is the product of globally integrated flow rate  $\Phi_{\text{pre}}^{s \rightarrow l}$  ( $y$ -axis width) and bulk residence time  $\Gamma_{\text{pre}}^{s \rightarrow l}$  ( $x$ -axis length). When interpreting Figure 6 it is useful to keep in mind the following constraints on the flow rates: (i) The emergence rate of aphotically regenerated DIC into the preformed pool is equal to the export production  $J_{\text{ex}}$  as source and sink balance in steady state. (ii) The biological utilization rate of preformed DIC is equal to the total nutrient uptake  $U$  as euphotic DIC is preformed by definition. (iii) The euphotic regeneration rate is



**Figure 6.** Preformed DIC pipes for (a) the preindustrial state, (b) the future RCP4.5-based state, and (c) the future RCP8.5-based state, partitioned according to source (left) and sink (right). The area, width ( $y$ -axis), and length ( $x$ -axis) of each  $s \rightarrow l$  pipe represent the inventory  $\mu_{\text{pre}}^{s \rightarrow l}$  (white text), flow rate  $\Phi_{\text{pre}}^{s \rightarrow l}$  (black text at the end of each pipe), and bulk residence time  $\Gamma_{\text{pre}}^{s \rightarrow l}$ , respectively. For comparison, the dotted rectangle in the bottom left corner of each panel indicates the total regenerated DIC pipe (all mechanisms summed/averaged from Fig. 5), also represented as export  $\Phi_{\text{ex}}$  (width)  $\times$  bulk residence time  $\Gamma_{\text{reg}}$  (length).



equal to the rate of un-exported production  $U - J_{\text{ex}} = (1 - f)U$  in steady state. Given the novelty of this diagnostic, we first take a look at the preindustrial state before examining future changes.

### 3.3.1 The preindustrial preformed pump

The (vertical) widths of the preformed pipes (Fig. 6) that terminate in biological utilization represent the rate with which preformed DIC fuels photosynthetic organic-matter production. In the preindustrial ocean, the fraction of newly preformed DIC destined for biological uptake is remarkably large at 60–65 % across all source processes. Here, “newly” preformed DIC denotes preformed DIC at the time of labelling it as “preformed”, i.e., at its source. For instance, of the 63 PgC yr<sup>-1</sup> ingassed from the atmosphere, 38 PgC yr<sup>-1</sup> support biological production, while the remainder outgasses without ever interacting with biology. In terms of inventories, 60 % of all preformed DIC is in transit to biological uptake while the rest is destined for outgassing. Conversely, of the total biological DIC uptake that occurs, 36 % is supplied by ingassing, 44 % by euphotic regeneration, and 20 % by upwelling of aphotically regenerated DIC.

To better understand the transport pathways of preformed DIC, we quantify the amount that enters the aphotic ocean. To that end, we partitioned the DIC in each preformed pipe according to whether or not it enters the aphotic interior during its source-to-sink transit (Eqs. (A9) and (A10)). This partition (not shown in Fig. 6) revealed that regardless of source–sink pair more than 90 % of preformed DIC that has been newly injected into the euphotic zone (“newly preformed”) will eventually explore the aphotic interior with a bulk first-contact time with the aphotic interior of less than half a year (roughly in agreement with the findings of Bopp et al., 2015). The preformed DIC pipes thus consist almost entirely of pathways that probe the aphotic interior.

Figure 6a also shows the bulk preindustrial residence times in each preformed DIC pipe, which range from 150 yr to 280 yr, depending on pipe. In comparison, the bulk residence time for regenerated DIC is only about 60 yr, which might seem surprising given that DIC regeneration occurs at depth. However, preformed DIC can roam over the euphotic zone for times on the order of a century without being biologically utilized or outgassed, while regenerated DIC immediately loses its regenerated label on contact with the euphotic zone. Furthermore, not all deep-to-surface paths are slow: Analysis of nutrient cycling in a data-assimilated circulation (Pasquier and Holzer, 2016) showed that nutrients regenerated at intermediate depths (and hence DIC) are likely to return to the surface through short diffusive vertical pathways while deep regenerated DIC is predominantly transported into Circumpolar Deep Water (CDW), which provides a relatively fast upwelling conduit back to the surface. For the ACCESS-M PCO<sub>2</sub> model, the CDW conduit is additionally short-circuited regionally in the Southern Ocean by the parent model’s unrealistic deep mixing (Fig. C2 and also Holzer et al., 2020).

### 3.3.2 The future preformed pump


In our perpetual future scenarios, the global preformed DIC inventory  $\mu_{\text{pre}}$  increases from its preindustrial value by  $\sim 1200$  PgC and 1500 PgC for the RCP4.5 and RCP8.5 scenarios, respectively (Table 3). While this is only about 5 % of the total preformed inventory, the absolute changes are of the same order of magnitude as (and larger than) the changes in the regenerated DIC pool ( $\sim 600$  PgC and 1400 PgC for the two scenarios).

**Table 3.** Contributions to the response of the preformed carbon inventory  $\Delta\mu_{\text{pre}}^{s \rightarrow l}$  from changes in bulk residence time  $\Delta\Gamma_{\text{pre}}^{s \rightarrow l}$  and changes in flow rate  $\Delta\Phi_{\text{pre}}^{s \rightarrow l}$  (values in PgC and rounded to two significant digits).

Scenario	Connection (source $s \rightarrow$ loss $l$ )	$\Delta\mu_{\text{pre}}^{s \rightarrow l}$	$\Phi_{\text{pre}}^{s \rightarrow l}$	$\Delta\Gamma_{\text{pre}}^{s \rightarrow l}$	$\Gamma_{\text{pre}}^{s \rightarrow l}$	$\Delta\Phi_{\text{pre}}^{s \rightarrow l}$	$\Delta\Gamma_{\text{pre}}^{s \rightarrow l}$	$\Delta\Phi_{\text{pre}}^{s \rightarrow l}$
RCP4.5	total (all $\rightarrow$ all)	+1200	-7800	+11000				-2300
	ingassing $\rightarrow$ outgassing	+6600	-1200	+9900				-2100
	ingassing $\rightarrow$ bio. uptake	+170	-1900	+2800				-680
	euphotic regen. $\rightarrow$ outgassing	+540	-910	+1800				-370
	euphotic regen. $\rightarrow$ bio. uptake	-3200	-1700	-2000				+520
	$C_{\text{reg}}$ emergence $\rightarrow$ outgassing	+2.4	-740	+940				-190
	$C_{\text{reg}}$ emergence $\rightarrow$ bio. uptake	-2900	-1300	-2100				+510
RCP8.5	total (all $\rightarrow$ all)	+1500	-15000	+28000				-12000
	ingassing $\rightarrow$ outgassing	+12000	-2300	+26000				-11000
	ingassing $\rightarrow$ bio. uptake	-1000	-3600	+4700				-2100
	euphotic regen. $\rightarrow$ outgassing	+320	-1700	+3300				-1200
	euphotic regen. $\rightarrow$ bio. uptake	-4800	-3000	-3200				+1400
	$C_{\text{reg}}$ emergence $\rightarrow$ outgassing	-940	-1600	+1300				-590
	$C_{\text{reg}}$ emergence $\rightarrow$ bio. uptake	-4300	-2600	-3300				+1600

The plumbing of the preformed DIC pump is reshaped entirely in our future scenarios. Preformed DIC is rerouted from biological utilization to outgassing. The strong increase in gross outgassing rate (by roughly a factor of 2 and 3 for RCP4.5 and RCP8.5) reflects the corresponding factor of 2 and 3 increases in atmospheric  $p\text{CO}_2$ . The roles of reduced  $\text{CO}_2$  solubility in warmer surface waters and changes in gas-exchange coefficients due to wind changes are dwarfed by the prescribed  $p\text{CO}_2$  increases. The increase in outgassing rate reduces the fraction of newly preformed DIC destined to biological utilization from a preindustrial 60–65 % to 40–50 % and 30–35 % in the two future scenarios (ranges are given across source mechanisms). Similarly, the fraction of biological production supported by newly ingassed DIC increases from 35 % in the preindustrial era to 50 % and 65 % in the future. Across all source–sink pairs, the ingassing-to-outgassing flow rates increase the most (roughly by factors 3 and 6 for RCP4.5 and RCP8.5) because of the dramatic change in atmospheric  $p\text{CO}_2$ . Note that, had we prescribed the total (i.e., oceanic plus atmospheric) carbon inventory instead of only the atmospheric  $p\text{CO}_2$ , the increase in atmospheric  $p\text{CO}_2$  and in the ingassing/outgassing fluxes in Fig. 6 would have been significantly smaller.

The re-routing of preformed DIC points to the biological pump becoming *less* efficient in the future: Regenerated DIC is much less likely to pass another time through the biological pump before outgassing to the atmosphere. Specifically, only about 40 % (RCP4.5, 30 % for RCP8.5) of preformed DIC sourced from regenerated organic matter is destined to biological utilization in the future compared to 60 % preindustrially. In the sense that the number of life-time passages through the biological pump is a measure of carbon pump efficiency (Holzer et al., 2021b), the future biological pump becomes *less*



**Figure 7.** Zonal means of the change of the normalized path density of preformed DIC from each source mechanism  $s$  (rows) to each loss mechanism  $l$  (columns),  $\Delta [C_{\text{pre}}^{s \rightarrow l} / \bar{C}_{\text{pre}}^{s \rightarrow l}]$ , for the RCP4.5-based state (a–f) and the RCP8.5-based state (g–l). Note that the normalization by the global mean  $\bar{C}_{\text{pre}}^{s \rightarrow l}$  is necessary to uncover changes in pattern, given the large overall changes in the magnitude of  $\bar{C}_{\text{pre}}^{s \rightarrow l}$ .

370 efficient. Using the vertical attenuation of the POC flux as a measure of pump efficiency (e.g., Buesseler et al., 2020; Volk and Hoffert, 1985), we similarly find that the efficiency of POC transfer to 1000 m below the euphotic zone decreases by order 5 % in the future. Hence, the pump becomes *stronger* in the sense that it sequesters a larger regenerated DIC pool in the future, but it does so *less* efficiently in the sense of reduced export rate and lifetime biological pump passages.

A key feature of the future preformed pump is that the bulk residence times in its “pipes” are shorter than in the preindustrial state. Preformed DIC pipes thus respond oppositely to regenerated DIC pipes: While the flow of regenerated DIC slows down with the more sluggish future circulation, the flow of preformed DIC actually speeds up in the future: Bulk preformed residence times across all the pipes decrease by 20–25 % and 40–50 % for the RCP4.5 and RCP8.5 scenarios. This at first counterintuitive behavior can be understood in terms of the future deep ocean becoming more isolated due to weaker future ventilation. Faster

375

surface paths are thus expected to become more important contributors to the overall transport than the slower deep paths.

380 To confirm and quantify this, we calculated the path density (e.g., Holzer and Primeau, 2013; Pasquier and Holzer, 2016) for source- and sink-partitioned preformed DIC. For a given source–sink pair this path density is simply the concentration of DIC in transit from specified source to specified sink. Figure 7 shows the future-minus-preindustrial changes in the global zonal-mean normalized preformed DIC path densities. (The path densities were normalized by the corresponding global preformed DIC inventory to reveal changes in pattern.) For all source–sink pairs, the future path density becomes surface-intensified with

385 strongly increased upper-ocean transport away from high latitudes. This is consistent with changes in the fraction of newly preformed DIC that will remain entirely within the euphotic zone before outgassing or biological utilization (Eqs. (A9)–(A10); not shown), which increases from about 5 % in the preindustrial state to 8 % and 12 % in the RCP4.5- and RCP8.5-based states.

#### 4 Discussion

We have analyzed the ocean carbon cycle for idealized steady-state future conditions corresponding to a perpetual 2090s

390 ocean state as predicted for the RCP4.5 and RCP8.5 scenarios. Our approach tracks all carbon in the ocean by considering the biologically pumped regenerated carbon pool, the preformed carbon pool, and the exchanges between these pools and the atmosphere. By focusing on the steady-state equilibrium response of the carbon cycle, we gain insight into how the system adjusts on all timescales, including the long millennial timescales of deep ventilation (e.g., Primeau and Holzer, 2006). This is important to capture, e.g., strengthened Southern Ocean nutrient trapping, which is driven by circulation slowdown and

395 manifests in transient simulations only well after the 21st century (Moore et al., 2018). In contrast, 21st-century transient simulations can only probe the response on centennial and sub-centennial timescales.

To track preformed DIC not only in the interior but also in the surface ocean, we introduced a conceptually new partition of preformed DIC according to the sources that inject DIC into the euphotic zone and the sinks that remove it. This represents a significant advance over the traditional view of preformed DIC as being merely the propagated surface DIC concentration (e.g.,

400 Ito and Follows, 2005), with which it is impossible to track preformed DIC transport in the surface ocean. Our new approach allows us to paint a quantitative picture of the “plumbing” of the ocean’s preformed carbon pump and how it connects the atmosphere to the biological pump.

Our new partition of preformed tracers will be useful beyond the scope of the present work. For example, a key quantity of interest in the ocean’s nutrient cycles is the fraction of production in one region supported by the supply of nutrients that

405 are newly preformed in another region. This cannot be accurately quantified using the traditional concept of preformed tracers because of the singular diffusive return flux of locally labelled preformed tracers (Hall and Holzer, 2003; Primeau and Holzer, 2006). Previous approaches have sidestepped this difficulty by either perturbing the system in the origin region and attributing the nonlinear response elsewhere to contributions from the origin region (e.g., Sarmiento et al., 2004), using interior volumes to label nutrient origin (e.g., Palter et al., 2010), or labelling nutrient origin according to last biological utilization or regeneration

410 (e.g., Holzer and Primeau, 2013; Pasquier and Holzer, 2016). By contrast, our new partition of preformed tracers has non-

diffusive sinks and is hence not subject to diffusive singularities making it the ideal tool for tracking preformed connectivity in the ocean.

A number of caveats must be kept in mind:

- 415 i. Our future states are steady and thus not predictions of future transient evolution. However, our idealized steady states do reveal the key mechanisms driving the system's responses to future change.
- 420 ii. Our steady circulations do not capture seasonal covariances between physical, thermodynamic, and biological variations (e.g., Riebesell et al., 2009). While this could be addressed with a cyclo-stationary model (e.g., Bardin et al., 2014), doing so would greatly increase complexity and computational cost. We expect seasonality to mostly affect the upper few hundred meters of the water column as was the case for Huang et al. (2021) who built a seasonally-varying ocean circulation inverse model (CYCLOCIM). We therefore do not expect the absence of explicit seasonality to affect the qualitative character of our results, as most of the circulation-driven changes analyzed occur at depth.
- 425 iii. Most quantitative aspects of our results are likely model specific and to some extent imprinted by circulation biases. While the ACCESS1.3 parent model is state-of-the-art, it produces Antarctic Bottom Water (AABW) through unrealistically deep convection in the Southern Ocean (Bi et al., 2013). However, the future reduction in AABW seen here is robust across CMIP5 models and also expected from recent observations (de Lavergne et al., 2014). We therefore do not expect our results to be sensitive to this unrealistic model feature. Although a model with more realistic AABW formation might produce quantitatively different results (e.g., production could increase due to a smaller decline in mixed layer depths), the qualitative links between changes and their driving mechanisms should be robust and model-independent (e.g., mixed-layer shoaling driving intensified nutrient trapping; Liu et al., 2023).
- 430 iv. The parent ACCESS1.3 model does not include meltwater from terrestrial ice sheets. Including ice-sheet meltwater would further reduce Southern Ocean ventilation (Purich et al., 2018; Purich and England, 2023; Li et al., 2023). This would likely strengthen the already dominating role of the circulation in driving changes in the biological and preformed carbon pumps.
- 435 v. Our model may not capture potentially important effects from biogeochemical mechanisms that are not explicitly parameterized (e.g., Henson et al., 2022). These include changes in community composition from adaptation and evolution (e.g., Boyd, 2015; Passow and Carlson, 2012; Doney et al., 2009; Lomas et al., 2022), changes in nitrogen and/or iron limitation (e.g., Thornton et al., 2009; Jickells et al., 2005), or changes in early diagenesis and sediment fluxes (e.g., Griffiths et al., 2017; Sweetman et al., 2017). However, at least some of our model's mechanistic shortcomings are partially compensated by having optimized biogeochemical parameters (Pasquier et al., 2023).
- 440 vi. Using the 1990s and 2090s decadal means from a single climate-model run does not capture decadal variability. However, we expect decadal variability to be small compared to the centennial changes considered here. While important for precise predictions, decadal variability is not important in our idealized steady-state framework for which any reasonable future state suffices.

A number of factors made the ACCESS-embedded PCO<sub>2</sub> model a natural choice for this study, which builds on previous  
445 publications. While we could have chosen a different circulation model, all models have biases, and none of the other models’  
transport matrices have been assessed in as much detail as those for ACCESS1.3 (Chamberlain et al., 2019; Holzer et al.,  
2020). Furthermore, Pasquier et al. (2023) optimized PCO<sub>2</sub> embedded in the ACCESS1.3 transport matrix (“ACCESS-M  
PCO<sub>2</sub>”) and quantified the effects of circulation biases on the biological pump. Importantly, our optimized preindustrial state  
fits observations of DIC, total alkalinity, dissolved O<sub>2</sub>, and PO<sub>4</sub> better than most CMIP5 and CMIP6 models (see, e.g., Bao and  
450 Li, 2016; Fu et al., 2022; Planchat et al., 2023).

How do the steady-state responses analyzed here compare with the trends seen in transient simulations? On one hand, we  
would expect a closer correspondence with the long-term behavior of multi-century simulations, where the system’s slow  
processes have had a chance to begin to assert themselves. On the other hand, even the long-term evolution of the ocean  
will likely be characterized by transience (Schmittner et al., 2008) and an overturning slowdown may eventually recover once  
455 dynamical equilibrium becomes re-established (e.g., Bi et al., 2001; Jansen and Nadeau, 2019). Simulation time is thus key  
when comparing to steady-state responses and one must be careful with the interpretation of transient simulations. For example,  
an analysis of a transient 21st-century circulation slowdown may interpret a decreased preformed DIC inventory as a weakening  
of the solubility pump (e.g., Raven and Falkowski, 1999; Liu et al., 2023). However, at steady-state, a more sluggish circulation  
is expected to enhance the solubility pump and increase the preformed DIC inventory by allowing larger spatial gradients to  
460 form (Murnane et al., 1999; Toggweiler et al., 2003; DeVries, 2022).

Our estimates of future DIC increases by 1700–2900 PgC roughly agree with estimates from multi-century simulations. For  
example, using 350-yr simulations with 2× and 4× preindustrial *p*CO<sub>2</sub>, Sarmiento and Le Quéré (1996) found increased DIC  
inventories ranging 1000–2000 PgC. Similarly, in simulations with *p*CO<sub>2</sub> prescribed at 550 and 1000 μatm, Plattner et al. (2001)  
found increases ranging 1000–1500 PgC by the year 2500. More recently, Liu et al. (2023) found an increase of about 1000 PgC  
465 by 2300 for the RCP8.5 scenario. For CMIP6 models under the SSP2-4.5 and SSP5-8.5 scenarios (which nominally match  
RCP4.5 and RCP8.5; Riahi et al., 2017; Arias et al., 2021), the increase in ocean carbon sequestration is about 400–500 PgC  
by the year 2100 (Liu et al., 2023), of which only about 100 PgC is regenerated DIC as there has been insufficient time for it to  
accumulate at depth.

The qualitative agreement of our results with the long-term behavior of transient simulations underlines the common driving  
470 mechanisms that are at work. Across almost all CMIP5 and CMIP6 models (e.g., Bopp et al., 2013; Hauck et al., 2015;  
Kwiatkowski et al., 2020; Arora et al., 2020; Wilson et al., 2022; Liu et al., 2023), a slower future circulation reduces nutrient  
supply, which reduces production and export despite warming-accelerated growth, and increases the sequestration time of  
regenerated DIC. At the same time, higher future atmospheric *p*CO<sub>2</sub> drives more carbon into the ocean. While the PCO<sub>2</sub>  
model explicitly represents only three (dependence of viscosity on *T* and dependence of respiration on *T* and O<sub>2</sub>) of twelve  
475 export-controlling mechanisms identified by Henson et al. (2022), our results broadly agree with the corresponding expected  
future effects in terms of both magnitude and direction: The dominant effect is that warming reduces export through enhanced  
shallow respiration, followed by faster particle sinking in less viscous seawater and decreased deep respiration due to reduced  
oxygen. Our results also roughly agree with the changes in the controls on carbon export and biological utilization identified by

Boyd (2015). However, their one-dimensional water-column analysis cannot capture the effects of circulation changes, which  
480 are a key control on the future carbon cycle identified here, while our analysis cannot capture changes in plankton community  
composition identified as a key driver by Boyd. PCO<sub>2</sub> may implicitly capture the effects of such changes through its optimized  
nonlinear production parameterization.

Our findings challenge the hypothesis of Liu et al. (2023) that a future circulation slowdown will reduce the capacity of  
the ocean to take up anthropogenic CO<sub>2</sub> on multi-century timescales. Our analysis shows that the circulation slowdown is the  
485 dominant driver of the large regenerated DIC increases in the long-term steady state through increased residence times in the  
deep ocean. A slower circulation may also increase the preformed DIC inventory by increasing DIC gradients thereby increas-  
ing the efficiency of the solubility pump (e.g., Murnane et al., 1999; Toggweiler et al., 2003), but this effect is overwhelmed  
by the large future increases in atmospheric *p*CO<sub>2</sub>. The bulk of the DIC inventory is located in the deep ocean and returns to  
the surface with a distribution of transit times that has a mean exceeding a millennium close to the seafloor rather than a few  
490 centuries (e.g., Fig. C3 or Primeau, 2005). The correlation for the year 2100 between carbon uptake and decreased overturning  
across CMIP6 models found by Liu et al. can therefore only capture the relatively short-timescale response of preformed DIC,  
and the correlation may thus very well change sign by 2300 or a few centuries later, as more time becomes available to fill  
abyssal waters with additional DIC, assuming that the slow circulation persists.

Our decomposition of the change in regenerated DIC inventory into contributions from changes in export and circulation is  
495 consistent with the results of Liu et al. (2023), but our analysis identifies important nonlinearities that appear to be absent in  
their simulation to the year 2300. Broadly consistent with our findings, they found that the circulation slowdown contributes  
an increase in  $\mu_{\text{reg}}$  below 2000 m that is roughly 5× larger than the reduction due to weakened export. However, in stark  
contrast with our finding of a large correlation between changes in regeneration and re-exposure times (roughly 20–30 % of the  
total change, see Table 1), Liu et al. argue that the interaction between export and circulation is negligible. That the nonlinear  
500 interaction terms would fortuitously collapse to zero because of model specifics or experimental design is unlikely. Based on  
our analyses, it is more likely that this difference stems predominantly from the fact that our future steady states capture the  
response associated with much longer timescales, allowing for the system’s nonlinear interactions to develop fully.

## 5 Conclusions

We investigated the steady-state response of the ocean’s carbon pumps to idealized future scenarios. This was done by embed-  
505 ding a relatively simple biogeochemical model into the average ocean state as predicted by the ACCESS climate model for the  
2090s under the RCP4.5 and RCP8.5 scenarios. We then solved for the steady state of the biogeochemistry keeping the ocean  
state and prescribed atmospheric *p*CO<sub>2</sub> fixed.

Most features of the carbon cycle’s response are already manifest in the RCP4.5-based scenario. For many quantities, the  
response for RCP8.5 is approximately twice that for RCP4.5 roughly tracking the change in atmospheric *p*CO<sub>2</sub>. One way in  
510 which the system’s nonlinearities manifest is in the spatial correlations between key driving mechanisms, which are 3–8 times

higher for RCP8.5 than for RCP4.5. These nonlinearities can either strengthen (e.g., biological uptake) or weaken (e.g., export production) the overall response.

Our analysis focused on the mechanisms that drive the response to climate change, brought to light here by applying novel diagnostics of carbon sequestration and transport made possible by the steady-state framework. We partitioned the ocean's DIC into regenerated and preformed pools, further partitioned according to sources and sinks. Each partitioned DIC pool, together with its source-to-sink flow rate defining the residence time in each pool, may be considered to be an advective–diffusive “pipe”. These pipes constitute the “plumbing” of the biological and preformed carbon pumps. Whereas preformed DIC is usually considered as determined by a concentration boundary condition, here we introduced a new partitioning of preformed DIC that allows us to track not only its interior transport, but also its surface pathways. To the best of our knowledge, this made it possible for the first time to track DIC from its injection into the preformed pool (by ingassing, euphotic regeneration, or upwelling of regenerated DIC) to its exit from the preformed pool through outgassing or biological uptake.

Our main conclusions are:

1. Biological productivity is remarkably resilient in the face of large environmental change. For our perpetual 2090s scenarios, production declines by only  $\sim 10\%$  from preindustrial values even for RCP8.5. These declines are driven by reduced nutrient supply but partially compensated by surface warming, which exponentially enhances organic-matter production and the recycling of nutrients within the euphotic zone. The overall nutrient supply declines because of intensified Southern Ocean nutrient trapping and because decreased ventilation and strong mixed-layer shoaling slows the resurfacing of nutrients from depth.
2. Export production declines by  $14\%$  and  $24\%$  in the RCP4.5- and RCP8.5-based steady-state scenarios, respectively. These declines are driven not only by reduced organic-matter production but also by reduced export ratios. The reduction in export ratios is driven primarily by enhanced shallow POC respiration due to surface warming and by decreased DOC export due to reduced ventilation. Decreased respiration from deoxygenation and faster POC sinking in less viscous warmer water tends to increase export ratios but is of secondary importance.
3. The future biological pump cycles a larger regenerated DIC inventory but does so less efficiently. The regenerated DIC inventory increases by  $30\%$  ( $\sim 600$  PgC) and  $70\%$  ( $\sim 1400$  PgC) for the RCP4.5- and RCP8.5-based scenarios, respectively. These steady-state increases capture millennial response timescales and are driven primarily by reduced deep ventilation that increases the sequestration time in the aphotic ocean. This allows more regenerated DIC to accumulate at depth and to participate in Southern Ocean trapping. Reduced export production only compensates for about half of the increase in regenerated inventories, underlining the key importance of circulation changes. Biological pump efficiency is decreased in the sense that resurfacing regenerated DIC contributes less to organic-matter production.
4. The preformed carbon pump is completely replumbed in our future scenarios. Preformed DIC is largely rerouted from supporting biological production to outgassing primarily because dissolved  $\text{CO}_2$  at the surface, and hence gross ingassing and outgassing, increase 2–3 times in approximate proportion to the prescribed atmospheric  $p\text{CO}_2$  increases. Reduced



biological production and decreased CO<sub>2</sub> solubility in warmer waters play only a secondary role. Because the inventory of preformed DIC in the euphotic zone increases only by a few percent due to carbonate buffering, increased ingassing and outgassing rates correspond to preformed DIC having shorter residence times in the euphotic zone. As a consequence, the fraction of newly preformed DIC that supports biological production shrinks from about 60 % in the preindustrial to roughly 40 % in the future. However, the fraction of biological production supported by newly ingassed DIC increases from a preindustrial 36 % to more than 50 % in the future, again because of the large increase in ingassing.

5. Preformed DIC is cycled more rapidly in our future scenarios despite the overall slower circulation. Across all source and sink mechanisms, the residence times of preformed DIC in the entire ocean decrease by 20–50 % in the future, while the global preformed inventory increases by only a few percent (~ 5 % or 1500 PgC for the RCP8.5 case), implying faster source-to-sink flow rates. This is driven by fast shallow preformed transport pathways becoming more important relative to slow deep pathways as reduced ventilation isolates the deep ocean. The fraction of newly preformed DIC that never enters the aphotic ocean roughly doubles from its preindustrial value of 5 % in both future scenarios.

Our analysis reveals a complex multifaceted response of the carbon cycle to changes in ocean state, even for our relatively simple biogeochemistry model and idealized scenarios. On one hand, a more sluggish future circulation slows down biological cycling but the regenerated carbon inventory increases due to longer sequestration times. On the other hand, preformed DIC shoals with faster source-to-sink flow due to the increased isolation of the deep ocean. While the response of the biological and preformed pumps are driven by different mechanisms with widely different response timescales, the regenerated and preformed DIC inventories both increase by similar absolute amounts when the full spectrum of response timescales is captured in steady state.

*Code and data availability.* The MATLAB and Julia code for this work will be made available upon acceptance of this article. The figures were created in Julia (Bezanson et al., 2017) with the Makie.jl plotting package (Danisch and Krumbiegel, 2021). The transport matrices were built from the historical, RCP4.5, and RCP8.5 ACCESS1.3 CMIP5 model runs available at <https://esgf.nci.org.au/projects/esgf-nci/>. This output also includes temperature, salinity, photosynthetically available radiation (PAR), sea-ice, and wind fields.

## Appendix A: Preformed DIC source-to-loss partition

To partition preformed DIC according to source and loss mechanism, we consider the following linear labelling tracer equation whose solution is identical to that of Eq. (4):

$$(T + l_{\text{atm}} + l_{\text{bio}}) C_{\text{pre}} = J_{\text{atm}}^{\downarrow} + \Omega_{\text{eup}} R + \Omega_{\text{eup}} C_{\text{reg}} / \tau_0, \quad (\text{A1})$$

where  $l_{\text{atm}} = J_{\text{atm}}^{\uparrow} / [\text{DIC}]$  and  $l_{\text{bio}} = U / [\text{DIC}]$  are equivalent linear loss rates for preformed DIC corresponding to the inverse timescales of local outgassing and biological uptake, respectively. Equation (A1) exploits the fact that labels are removed in proportion to their fractional abundance even when the removal process is nonlinear (e.g., Holzer and DeVries, 2022). The

Green function  $G_{\text{pre}}$  for Eq. (A1) obeys

$$575 \quad (\mathcal{T}_{\mathbf{r}} + l_{\text{atm}}(\mathbf{r}) + l_{\text{bio}}(\mathbf{r})) G_{\text{pre}}(\mathbf{r}|\mathbf{r}') = \delta(\mathbf{r} - \mathbf{r}'), \quad (\text{A2})$$

where  $\delta(\mathbf{r} - \mathbf{r}')$  is the three-dimensional Dirac distribution and the subscript in  $\mathcal{T}_{\mathbf{r}}$  indicates that the differential operator  $\mathcal{T}$  is applied at point  $\mathbf{r}$ . The Green function  $G_{\text{pre}}(\mathbf{r}|\mathbf{r}')$  allows us to cleanly partition  $C_{\text{pre}}$  according to every source and sink. Specifically, the preformed DIC contributed by any source  $s$  is given by

$$C_{\text{pre}}^s = \int G_{\text{pre}}(\mathbf{r}|\mathbf{r}') s(\mathbf{r}') d^3\mathbf{r}', \quad (\text{A3})$$

580 which in matrix form is equivalent to

$$C_{\text{pre}}^s = (\mathbf{T} + \mathbf{L}_{\text{atm}} + \mathbf{L}_{\text{bio}})^{-1} \mathbf{s} \quad (\text{A4})$$

where  $\mathbf{T}$  is the advective–diffusive transport matrix,  $\mathbf{L}_{\text{atm}}$  and  $\mathbf{L}_{\text{bio}}$  are diagonal matrices representing the local loss rates  $l_{\text{atm}}$  and  $l_{\text{bio}}$ , and  $\mathbf{s}$  is a column vector of the source  $s$ .

To quantify the fraction of  $C_{\text{pre}}$  destined for a given loss mechanism (outgassing or uptake), we take the adjoint Green  
585 function  $\tilde{G}_{\text{pre}}$  defined as the solution to

$$(\tilde{\mathcal{T}}_{\mathbf{r}} + l_{\text{atm}}(\mathbf{r}) + l_{\text{bio}}(\mathbf{r})) \tilde{G}_{\text{pre}}(\mathbf{r}|\mathbf{r}') = \delta(\mathbf{r} - \mathbf{r}'), \quad (\text{A5})$$

where  $\tilde{\mathcal{T}}$  is the adjoint of  $\mathcal{T}$  with respect to the volume-weighted inner product. The fraction of preformed DIC destined for loss process  $l$ , where  $l = l_{\text{bio}}$  or  $l_{\text{atm}}$ , is given by

$$f_{\text{pre}}^l = \int \tilde{G}_{\text{pre}}(\mathbf{r}|\mathbf{r}') l(\mathbf{r}') d^3\mathbf{r}'. \quad (\text{A6})$$

590 In matrix form, Eq. (A8) becomes

$$\mathbf{f}_{\text{pre}}^l = (\tilde{\mathbf{T}} + \mathbf{L}_{\text{atm}} + \mathbf{L}_{\text{bio}})^{-1} \mathbf{l}, \quad (\text{A7})$$

where  $\tilde{\mathbf{T}} = \mathbf{V}^{-1} \mathbf{T}^T \mathbf{V}$  is the adjoint of  $\mathbf{T}$  with respect to the volume-weighted inner product ( $\mathbf{V}$  is a diagonal matrix of grid-box volumes) and  $\mathbf{l}$  is the vector representing the grid-point values of  $l$ .

The preformed DIC concentration due to source mechanism  $s$  and destined for loss mechanism  $l$  is thus given by

$$595 \quad C_{\text{pre}}^{s \rightarrow l} = f_{\text{pre}}^l C_{\text{pre}}^s. \quad (\text{A8})$$

The inventory of the  $s \rightarrow l$  preformed DIC pipe, i.e., the amount of preformed DIC in transit from source  $s$  to loss  $l$ , is given by  $\mu_{\text{pre}}^{s \rightarrow l} = \int C_{\text{pre}}^{s \rightarrow l}(\mathbf{r}) d^3\mathbf{r}$ . Similarly, the global flow rate of the  $s \rightarrow l$  preformed DIC pipe is given by  $\Phi_{\text{pre}}^{s \rightarrow l} = \int f_{\text{pre}}^l(\mathbf{r}) s(\mathbf{r}) d^3\mathbf{r}$ .

The bulk transit time is then simply given by  $\Gamma_{\text{pre}}^{s \rightarrow l} = \mu_{\text{pre}}^{s \rightarrow l} / \Phi_{\text{pre}}^{s \rightarrow l}$ .

To better diagnose the physical pathways of preformed DIC, we further partition the DIC in the preformed pipes into DIC  
600 that remains in the euphotic zone throughout the preformed lifetime and DIC that explores the aphotic interior. To this end, we introduce a labelling tracer for euphotic-only DIC,  $C_{\text{pre}}^{\text{eup}}$ , which is governed by

$$(\mathcal{T} + l_{\text{atm}} + l_{\text{bio}} + l_{\text{aph}}) C_{\text{pre}}^{\text{eup}} = J_{\text{atm}}^{\downarrow} + \Omega_{\text{eup}} R + \Omega_{\text{eup}} C_{\text{reg}} / \tau_0, \quad (\text{A9})$$


which is similar to Eq. (A1) but where  $l_{\text{aph}} = \Omega_{\text{aph}}/\tau_0$  additionally unlabels any preformed DIC that leaves the euphotic zone. Taking the difference with the full preformed DIC from Eq. (A1) gives the concentration  $C_{\text{pre}}^{\text{aph}}$  of preformed DIC that will explore the aphotic interior and obeys

$$(\mathcal{T} + l_{\text{atm}} + l_{\text{bio}}) C_{\text{pre}}^{\text{aph}} = l_{\text{aph}} C_{\text{pre}}^{\text{eup}}. \quad (\text{A10})$$

In practice, Eqs. (A9) and (A10) are again solved in matrix form.

## Appendix B: Changes in tracer fields

610 Nutrient supply (here  $\text{PO}_4$ ) is the dominant driver of the response of production (Fig. 1), while oxygen concentrations control the respiration rates in the euphotic zone and thus export ratios. Figure B1 shows the euphotic-zone mean  $[\text{PO}_4]$  and  $[\text{O}_2]$  in the preindustrial and future states, along with the corresponding change. Phosphate concentrations decrease almost everywhere, with a particularly strong decrease near the Weddell Sea and the Ross Sea, where the mixed layer that was unrealistically deep preindustrially has shoaled considerably (Fig. C2). For euphotic-zone  $[\text{O}_2]$ , we also see a global decline except in the Weddell and Ross seas, where oxygen was strongly undersaturated preindustrially because of the unrealistic MLDs.




**Figure B1.** (a) Mean euphotic  $[\text{PO}_4]$  for the preindustrial state. (b) As (a) for the RCP4.5-based state. (c) As (b) for RCP8.5. (d–e) As (b–c) for  $\Delta[\text{PO}_4]$ . (f–j) As (a–e) for  $\text{O}_2$ .

## 615 Appendix C: Thermodynamic Fields and Circulation Features


Through its control on phytoplankton and bacterial growth rates, temperature change is a key driver of production and export changes. Figure C1 shows the changes in euphotic-zone temperature from the preindustrial values for the RCP4.5 and RCP8.5 scenarios. (The temperature fields used here are decadal means for the 1990s and 2090s of the parent ACCESS1.3 model, consistent with our methodology for extracting the circulations.) Temperature strongly increases almost everywhere except  
620 in the highest latitudes of the Southern Ocean and in the subpolar North Atlantic “cold blob” where temperatures decrease (thought to be driven partly by the Atlantic overturning circulation slowdown; see, e.g., Cheng et al., 2022).

The mixed layer plays a key role in supplying nutrients to the euphotic zone. Figure C2 shows the mixed layer depths (MLDs) in the preindustrial and future states. (As for temperature, the MLD fields were taken from 1990s and 2090s averages of ACCESS1.3 simulations.) The unrealistically deep MLDs of the 1990s (used here as the preindustrial state) dramatically  
625 shoal in the 2090s, partially shutting off the model’s deep-water formation (Bi et al., 2013) although the deep Atlantic MLD near the Weddell Sea has not entirely subsided in the RCP4.5 scenario (Fig. C2b).

The dominant control on the sequestration strength of the biological pump is the mean time  $\Gamma^\uparrow$  to return water to the euphotic layer (termed re-exposure time; Primeau, 2005; DeVries and Holzer, 2019; Holzer et al., 2020), because it sets the mean residence (or sequestration) time of regenerated DIC. Figure C3 shows the Atlantic, Pacific, and Indian Ocean zonal  
630 mean  $\Gamma^\uparrow$  for the preindustrial and future states. (Re-exposure times are calculated as in the work by Holzer et al. (2020) who considered only the RCP8.5 scenario.)  $\Gamma^\uparrow$  increases greatly (by up to 600 and 1200 years for the RCP4.5 and RCP8.5 scenarios, respectively). At first glance the RCP8.5 response may appear to be proportional to the RCP4.5 response, but on closer inspection different patterns are visible, particularly in the Atlantic sector. These differences are consistent with the corresponding differences in MLD (Fig. C2).




**Figure C1.** (a) Change in vertical mean euphotic temperature,  $\Delta T$ , for the RCP4.5 scenario. (b) As (a) for RCP8.5.



**Figure C2.** (a) Mixed layer depth (MLD) for the preindustrial state. (b) As (a) for the future RCP4.5 state. (c) As (b) for RCP8.5. (d–e) As (b–c) for the change in mixed layer depth,  $\Delta$ MLD.






**Figure C3.** (a–c) Mean water re-exposure time,  $\Gamma^\uparrow$ , for the preindustrial circulation, zonally averaged over the Atlantic (a), Pacific (b), and Indian Ocean (c). (d–f) As (a–c) for the RCP4.5 state. (g–i) As (d–f) for RCP8.5. (j–o) As (d–i) for  $\Delta\Gamma^\uparrow$ .


## 635 **Appendix D: Biological production and export production**

Figure D1 shows the production,  $U$ , for the preindustrial and future states, along with the corresponding changes. Similarly, Fig. D2 shows the corresponding export ratios,  $f$ . As shown by Fig. 3, important contributions from export ratios occur in places where  $\Delta f$  and  $U$  are both large, and important contributions from production occur where  $f$  and  $\Delta U$  are both large.


The changes in export production vary with mechanism, i.e.,  $\text{POC}_f$ ,  $\text{POC}_s$ , DOC, or PIC. The zonally integrated changes  
640 for each mechanism are shown in Fig. D3. For  $\text{POC}_s$  export, changes are strongly dominated by export ratios, particularly in the tropics likely because very little  $\text{POC}_s$  makes it through the euphotic zone ( $f \approx 0$ ) and thus  $\Delta J_{\text{ex}} \approx U \Delta f$ . In contrast, PIC export is entirely controlled by changes in production ( $\Delta J_{\text{ex}} \approx f \Delta U$ ; Fig. D3) because in our model its export ratio is unaffected by any environmental change ( $f \approx 0.97$  and  $\Delta f \approx 0.001$ ).



**Figure D1.** (a) Map of carbon uptake,  $U(\mathbf{r})$ , vertically integrated, in the preindustrial state. (b) Zonal integral of (a). (c–d) As (a–b) for the future RCP4.5-based state. (e–f) As (c–d) for RCP8.5. (g–j) As (c–f) for  $\Delta U$ .



**Figure D2.** (a) Map of export ratio,  $f$ , in the preindustrial state. (b) Zonal mean of (a). (c–d) As (a–b) for the future RCP4.5-based state. (e–f) As (c–d) for RCP8.5. (g–j) As (c–f) for  $\Delta f$ .



**Figure D3.** (a–d) Change in DOC (a),  $\text{POC}_s$  (b),  $\text{POC}_f$  (c), and PIC (d) zonally integrated export production,  $\Delta J_{\text{ex}}$  (black), for the RCP4.5-based state, decomposed into contributions from  $U \Delta f$  (blue),  $f \Delta U$  (orange), and  $\Delta f \Delta U$  (gray). The globally integrated contribution to  $\Delta J_{\text{ex}}$  is indicated in each panel. (e–h) As (a–d) for the RCP8.5-based state.

*Author contributions.* BP and MH designed the study, analyzed the data, and wrote the manuscript with input from all co-authors. BP wrote  
645 the code, performed the experiments, and produced the figures with input from all co-authors. The funding was acquired by MH.

*Competing interests.* The authors declare that they have no conflict of interest.

*Acknowledgements.* We thank Yi Liu for helpful discussions. This work was supported by Australian Research Council grant DP210101650 (to MH) and undertaken with the assistance of resources and services from the National Computational Infrastructure (NCI), which is supported by the Australian Government.

650 **References**

- Arias, P., Bellouin, N., Coppola, E., Jones, R., Krinner, G., Marotzke, J., Naik, V., Palmer, M., Plattner, G.-K., Rogelj, J., Rojas, M., Sillmann, J., Storelvmo, T., Thorne, P., Trewin, B., Achuta Rao, K., Adhikary, B., Allan, R., Armour, K., Bala, G., Barimalala, R., Berger, S., Canadell, J., Cassou, C., Cherchi, A., Collins, W., Collins, W., Connors, S., Corti, S., Cruz, F., Dentener, F., Dereczynski, C., Di Luca, A., Diongue Niang, A., Doblas-Reyes, F., Dosio, A., Douville, H., Engelbrecht, F., Eyring, V., Fischer, E., Forster, P., Fox-Kemper, B.,  
655 Fuglestedt, J., Fyfe, J., Gillett, N., Goldfarb, L., Gorodetskaya, I., Gutierrez, J., Hamdi, R., Hawkins, E., Hewitt, H., Hope, P., Islam, A., Jones, C., Kaufman, D., Kopp, R., Kosaka, Y., Kossin, J., Krakovska, S., Lee, J.-Y., Li, J., Mauritsen, T., Maycock, T., Meinshausen, M., Min, S.-K., Monteiro, P., Ngo-Duc, T., Otto, F., Pinto, I., Pirani, A., Raghavan, K., Ranasinghe, R., Ruane, A., Ruiz, L., Sallée, J.-B., Samset, B., Sathyendranath, S., Seneviratne, S., Sörensson, A., Szopa, S., Takayabu, I., Tréguier, A.-M., van den Hurk, B., Vautard, R., von Schuckmann, K., Zaehle, S., Zhang, X., and Zickfeld, K.: Technical Summary, in: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J., Maycock, T., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., pp. 33–144, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/9781009157896.002>, 2021.
- Arora, V. K., Katavouta, A., Williams, R. G., Jones, C. D., Brovkin, V., Friedlingstein, P., Schwinger, J., Bopp, L., Boucher, O., Cadule, P.,  
665 Chamberlain, M. A., Christian, J. R., Delire, C., Fisher, R. A., Hajima, T., Ilyina, T., Joetzjer, E., Kawamiya, M., Koven, C. D., Krasting, J. P., Law, R. M., Lawrence, D. M., Lenton, A., Lindsay, K., Pongratz, J., Raddatz, T., Séférian, R., Tachiiri, K., Tjiputra, J. F., Wiltshire, A., Wu, T., and Ziehn, T.: Carbon–Concentration and Carbon–Climate Feedbacks in CMIP6 Models and Their Comparison to CMIP5 Models, *Biogeosciences*, 17, 4173–4222, <https://doi.org/10.5194/bg-17-4173-2020>, 2020.
- Bao, Y. and Li, Y.: Simulations of Dissolved Oxygen Concentration in CMIP5 Earth System Models, *Acta Oceanologica Sinica*, 35, 28–37,  
670 <https://doi.org/10.1007/s13131-016-0959-x>, 2016.
- Bardin, A., Primeau, F. W., and Lindsay, K.: An offline implicit solver for simulating prebomb radiocarbon, *Ocean Modelling*, 73, 45–58, <https://doi.org/10.1016/j.ocemod.2013.09.008>, 2014.
- Bernardello, R., Marinov, I., Palter, J. B., Galbraith, E. D., and Sarmiento, J. L.: Impact of Weddell Sea deep convection on natural and anthropogenic carbon in a climate model, *Geophysical Research Letters*, 41, 7262–7269, <https://doi.org/10.1002/2014GL061313>, 2014.
- 675 Bezanson, J., Edelman, A., Karpinski, S., and Shah, V. B.: Julia: A Fresh Approach to Numerical Computing, *SIAM Review*, 59, 65–98, <https://doi.org/10.1137/141000671>, 2017.
- Bi, D., Budd, W. F., Hirst, A. C., and Wu, X.: Collapse and Reorganisation of the Southern Ocean Overturning under Global Warming in a Coupled Model, *Geophysical Research Letters*, 28, 3927–3930, <https://doi.org/10.1029/2001GL013705>, 2001.
- Bi, D., Marsland, S. J., Uotila, P., O’Farrell, S., Fiedler, R. A. S., Sullivan, A., Griffies, S. M., Zhou, X., and Hirst, A. C.: ACCESS-OM: The  
680 Ocean and Sea Ice Core of the ACCESS Coupled Model, *Australian Meteorological and Oceanographic Journal*, 63, 213–232, 2013.
- Bopp, L., Resplandy, L., Orr, J. C., Doney, S. C., Dunne, J. P., Gehlen, M., Halloran, P., Heinze, C., Ilyina, T., Séférian, R., Tjiputra, J., and Vichi, M.: Multiple stressors of ocean ecosystems in the 21st century: projections with CMIP5 models, *Biogeosciences*, 10, 6225–6245, <https://doi.org/10.5194/bg-10-6225-2013>, 2013.
- Bopp, L., Lévy, M., Resplandy, L., and Sallée, J. B.: Pathways of anthropogenic carbon subduction in the global ocean, *Geophysical Research  
685 Letters*, 42, 6416–6423, <https://doi.org/10.1002/2015GL065073>, 2015.

- Boyd, P. W.: Toward quantifying the response of the oceans' biological pump to climate change, *Frontiers in Marine Science*, 2, <https://doi.org/10.3389/fmars.2015.00077>, 2015.
- Broecker, W. S.: Glacial to Interglacial Changes in Ocean Chemistry, *Progress in Oceanography*, 11, 151–197, [https://doi.org/10.1016/0079-6611\(82\)90007-6](https://doi.org/10.1016/0079-6611(82)90007-6), 1982.
- 690 Buesseler, K. O., Boyd, P. W., Black, E. E., and Siegel, D. A.: Metrics that matter for assessing the ocean biological carbon pump, *Proceedings of the National Academy of Sciences*, 117, 9679–9687, <https://doi.org/10.1073/pnas.1918114117>, 2020.
- Caël, B. B. and Follows, M. J.: On the temperature dependence of oceanic export efficiency, *Geophysical Research Letters*, 43, 5170–5175, <https://doi.org/10.1002/2016GL068877>, 2016.
- Caesar, L., Rahmstorf, S., Robinson, A., Feulner, G., and Saba, V.: Observed Fingerprint of a Weakening Atlantic Ocean Overturning  
695 Circulation, *Nature*, 556, 191–196, <https://doi.org/10.1038/s41586-018-0006-5>, 2018.
- Chamberlain, M. A., Matear, R. J., Holzer, M., Bi, D., and Marsland, S. J.: Transport matrices from standard ocean-model output and quantifying circulation response to climate change, *Ocean Modelling*, 135, 1–13, <https://doi.org/10.1016/j.ocemod.2019.01.005>, 2019.
- Cheng, L., von Schuckmann, K., Abraham, J. P., Trenberth, K. E., Mann, M. E., Zanna, L., England, M. H., Zika, J. D., Fasullo, J. T., Yu, Y., Pan, Y., Zhu, J., Newsom, E. R., Bronselaer, B., and Lin, X.: Past and Future Ocean Warming, *Nature Reviews Earth & Environment*, 3,  
700 776–794, <https://doi.org/10.1038/s43017-022-00345-1>, 2022.
- Costello, C., Cao, L., Gelcich, S., Cisneros-Mata, M. Á., Free, C. M., Froehlich, H. E., Golden, C. D., Ishimura, G., Maier, J., Macadam-Somer, I., Mangin, T., Melnychuk, M. C., Miyahara, M., de Moor, C. L., Naylor, R., Nøstbakken, L., Ojea, E., O'Reilly, E., Parma, A. M., Plantinga, A. J., Thilsted, S. H., and Lubchenco, J.: The Future of Food from the Sea, *Nature*, 588, 95–100, <https://doi.org/10.1038/s41586-020-2616-y>, 2020.
- 705 Danisch, S. and Krumbiegel, J.: Makie.JI: Flexible High-Performance Data Visualization for Julia, *Journal of Open Source Software*, 6, 3349, <https://doi.org/10.21105/joss.03349>, 2021.
- de Lavergne, C., Palter, J. B., Galbraith, E. D., Bernardello, R., and Marinov, I.: Cessation of Deep Convection in the Open Southern Ocean under Anthropogenic Climate Change, *Nature Climate Change*, 4, 278–282, <https://doi.org/10.1038/nclimate2132>, 2014.
- DeVries, T.: The Ocean Carbon Cycle, *Annual Review of Environment and Resources*, 47, 317–341, <https://doi.org/10.1146/annurev-environ-120920-111307>, 2022.
- 710 DeVries, T. and Holzer, M.: Radiocarbon and Helium Isotope Constraints on Deep Ocean Ventilation and Mantle-<sup>3</sup>He Sources, *Journal of Geophysical Research: Oceans*, 124, 3036–3057, <https://doi.org/10.1029/2018JC014716>, 2019.
- DeVries, T., Primeau, W. F., and Deutsch, C.: The sequestration efficiency of the biological pump, *Geophysical Research Letters*, 39, <https://doi.org/10.1029/2012GL051963>, 2012.
- 715 Doney, S. C., Fabry, V. J., Feely, R. A., and Kleypas, J. A.: Ocean Acidification: The Other CO<sub>2</sub> Problem, *Annual Review of Marine Science*, 1, 169–192, <https://doi.org/10.1146/annurev.marine.010908.163834>, PMID: 21141034, 2009.
- FAO: The State of World Fisheries and Aquaculture 2018 - Meeting the sustainable development goals, Food and Agriculture Organization of the United Nations FAO, ISBN 978-92-5-130562-1, 2018.
- Fu, W., Moore, J. K., Primeau, F., Collier, N., Ogunro, O. O., Hoffman, F. M., and Randerson, J. T.: Evaluation of Ocean Biogeochemistry and Carbon Cycling in CMIP Earth System Models With the International Ocean Model Benchmarking (IOMB) Software System, *Journal of Geophysical Research: Oceans*, 127, e2022JC018965, <https://doi.org/10.1029/2022JC018965>, 2022.
- 720 Galbraith, E. D. and Martiny, A. C.: A simple nutrient-dependence mechanism for predicting the stoichiometry of marine ecosystems, *Proceedings of the National Academy of Sciences*, 112, 8199–8204, <https://doi.org/10.1073/pnas.1423917112>, 2015.



- Gent, P. R., Willebrand, J., McDougall, T. J., and McWilliams, J. C.: Parameterizing eddy-induced tracer transports in ocean circulation models, *Journal of physical oceanography*, 25, 463–474, [https://doi.org/10.1175/1520-0485\(1995\)025<0463:PEITTI>2.0.CO;2](https://doi.org/10.1175/1520-0485(1995)025<0463:PEITTI>2.0.CO;2), 1995.
- 725 Golden, C.: Fall in fish catch threatens human health, *Nature*, 534, 317–320, 2016.
- Griffiths, J. R., Kadin, M., Nascimento, F. J. A., Tamelander, T., Törnroos, A., Bonaglia, S., Bonsdorff, E., Brüchert, V., Gårdmark, A., Järnström, M., Kotta, J., Lindegren, M., Nordström, M. C., Norkko, A., Olsson, J., Weigel, B., Žydelis, R., Blenckner, T., Niiranen, S., and Winder, M.: The Importance of Benthic–Pelagic Coupling for Marine Ecosystem Functioning in a Changing World, *Global Change*
- 730 *Biology*, 23, 2179–2196, <https://doi.org/10.1111/gcb.13642>, 2017.
- Hall, T. M. and Holzer, M.: Advective-diffusive mass flux and implications for stratosphere-troposphere exchange, *Geophysical Research Letters*, 30, <https://doi.org/10.1029/2002GL016419>, 2003.
- Hauck, J., Völker, C., Wolf-Gladrow, D. A., Laufkötter, C., Vogt, M., Aumont, O., Bopp, L., Buitenhuis, E. T., Doney, S. C., Dunne, J., Gruber, N., Hashioka, T., John, J., Le Quéré, C., Lima, I. D., Nakano, H., Séférian, R., and Totterdell, I.: On the Southern
- 735 *Ocean CO<sub>2</sub> uptake and the role of the biological carbon pump in the 21st century*, *Global Biogeochemical Cycles*, 29, 1451–1470, <https://doi.org/10.1002/2015GB005140>, 2015.
- Helm, K. P., Bindoff, N. L., and Church, J. A.: Observed decreases in oxygen content of the global ocean, *Geophysical Research Letters*, 38, <https://doi.org/10.1029/2011GL049513>, 2011.
- Henson, S. A., Laufkötter, C., Leung, S., Giering, S. L. C., Palevsky, H. I., and Cavan, E. L.: Uncertain Response of Ocean Biological Carbon
- 740 *Export in a Changing World*, *Nature Geoscience*, 15, 248–254, <https://doi.org/10.1038/s41561-022-00927-0>, 2022.
- Holzer, M. and DeVries, T.: Source-labeled anthropogenic carbon reveals a large shift of preindustrial carbon from the ocean to the atmosphere, *Global Biogeochemical Cycles*, 36, e2022GB007405, <https://doi.org/10.1029/2022GB007405>, 2022.
- Holzer, M. and Primeau, F. W.: Improved Constraints on Transit Time Distributions from Argon 39: A Maximum Entropy Approach, *Journal of Geophysical Research: Oceans*, 115, <https://doi.org/10.1029/2010JC006410>, 2010.
- 745 Holzer, M. and Primeau, F. W.: Global teleconnections in the oceanic phosphorus cycle: patterns, paths, and timescales, *Journal of Geophysical Research: Oceans*, 118, 1775–1796, <https://doi.org/10.1002/jgrc.20072>, 2013.
- Holzer, M., Primeau, F. W., DeVries, T., and Matear, R.: The Southern Ocean silicon trap: Data-constrained estimates of regenerated silicic acid, trapping efficiencies, and global transport paths, *Journal of Geophysical Research: Oceans*, 119, 313–331, <https://doi.org/10.1002/2013JC009356>, 2014.
- 750 Holzer, M., Chamberlain, M. A., and Matear, R. J.: Climate-Driven Changes in the Ocean’s Ventilation Pathways and Time Scales Diagnosed From Transport Matrices, *Journal of Geophysical Research: Oceans*, 125, e2020JC016414, <https://doi.org/10.1029/2020JC016414>, e2020JC016414 10.1029/2020JC016414, 2020.
- Holzer, M., DeVries, T., and de Lavergne, C.: Diffusion controls the ventilation of a Pacific Shadow Zone above abyssal overturning, *Nature Communications*, 12, 4348, <https://doi.org/10.1038/s41467-021-24648-x>, 2021a.
- 755 Holzer, M., Kwon, E. Y., and Pasquier, B.: A New Metric of the Biological Carbon Pump: Number of Pump Passages and Its Control on Atmospheric pCO<sub>2</sub>, *Global Biogeochemical Cycles*, 35, e2020GB006863, <https://doi.org/10.1029/2020GB006863>, e2020GB006863 2020GB006863, 2021b.
- Huang, Q., Primeau, F. W., and DeVries, T.: CYCLOCIM: A 4-D variational assimilation system for the climatological mean seasonal cycle of the ocean circulation, *Ocean Modelling*, 159, 101762, <https://api.semanticscholar.org/CorpusID:234316061>, 2021.
- 760 Irving, D., Hobbs, W., Church, J., and Zika, J.: A Mass and Energy Conservation Analysis of Drift in the CMIP6 Ensemble, *Journal of Climate*, 34, 3157 – 3170, <https://doi.org/10.1175/JCLI-D-20-0281.1>, 2021.

- Ito, T. and Follows, M. J.: Preformed phosphate, soft tissue pump and atmospheric CO<sub>2</sub>, *Journal of Marine Research*, 63, 813–839, 2005.
- Ito, T., Bracco, A., Deutsch, C., Frenzel, H., Long, M., and Takano, Y.: Sustained growth of the Southern Ocean carbon storage in a warming climate, *Geophysical Research Letters*, 42, 4516–4522, <https://doi.org/10.1002/2015GL064320>, 2015.
- 765 Jansen, M. F. and Nadeau, L.-P.: A Toy Model for the Response of the Residual Overturning Circulation to Surface Warming, *Journal of Physical Oceanography*, 49, 1249–1268, <https://doi.org/10.1175/JPO-D-18-0187.1>, 2019.
- Jickells, T. D., An, Z. S., Andersen, K. K., Baker, A. R., Bergametti, G., Brooks, N., Cao, J. J., Boyd, P. W., Duce, R. A., Hunter, K. A., Kawahata, H., Kubilay, N., laRoche, J., Liss, P. S., Mahowald, N., Prospero, J. M., Ridgwell, A. J., Tegen, I., and Torres, R.: Global Iron Connections Between Desert Dust, Ocean Biogeochemistry, and Climate, *Science*, 308, 67–71, <https://doi.org/10.1126/science.1105959>,  
770 2005.
- Kriest, I.: Calibration of a simple and a complex model of global marine biogeochemistry, *Biogeosciences*, 14, 4965–4984, <https://doi.org/10.5194/bg-14-4965-2017>, 2017.
- Kriest, I., Kähler, P., Koeve, W., Kvale, K., Sauerland, V., and Oschlies, A.: One size fits all? Calibrating an ocean biogeochemistry model for different circulations, *Biogeosciences*, 17, 3057–3082, <https://doi.org/10.5194/bg-17-3057-2020>, 2020.
- 775 Kwiatkowski, L., Torres, O., Bopp, L., Aumont, O., Chamberlain, M., Christian, J. R., Dunne, J. P., Gehlen, M., Ilyina, T., John, J. G., Lenton, A., Li, H., Lovenduski, N. S., Orr, J. C., Palmieri, J., Santana-Falcón, Y., Schwinger, J., Séférian, R., Stock, C. A., Tagliabue, A., Takano, Y., Tjiputra, J., Toyama, K., Tsujino, H., Watanabe, M., Yamamoto, A., Yool, A., and Ziehn, T.: Twenty-First Century Ocean Warming, Acidification, Deoxygenation, and Upper-Ocean Nutrient and Primary Production Decline from CMIP6 Model Projections, *Biogeosciences*, 17, 3439–3470, <https://doi.org/10.5194/bg-17-3439-2020>, 2020.
- 780 Kwon, E. Y., Holzer, M., Timmermann, A., and Primeau, F. W.: Estimating three-dimensional carbon-to-phosphorus stoichiometry of exported marine organic matter, *Global Biogeochemical Cycles*, 36, e2021GB007154, <https://doi.org/10.1029/2021GB007154>, 2022.
- Laws, E. A., Falkowski, P. G., Smith Jr., W. O., Ducklow, H., and McCarthy, J. J.: Temperature effects on export production in the open ocean, *Global Biogeochemical Cycles*, 14, 1231–1246, <https://doi.org/10.1029/1999GB001229>, 2000.
- Laws, E. A., D'Sa, E., and Naik, P.: Simple equations to estimate ratios of new or export production to total production from  
785 satellite-derived estimates of sea surface temperature and primary production, *Limnology and Oceanography: Methods*, 9, 593–601, <https://doi.org/10.4319/lom.2011.9.593>, 2011.
- Lewis, E. R. and Wallace, D. W. R.: Program Developed for CO<sub>2</sub> System Calculations, ORNL/CDIAC-105. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, TN., <https://doi.org/10.15485/1464255>, 1998.
- Li, G., Cheng, L., Zhu, J., Trenberth, K. E., Mann, M. E., and Abraham, J. P.: Increasing Ocean Stratification over the Past Half-Century,  
790 *Nature Climate Change*, 10, 1116–1123, <https://doi.org/10.1038/s41558-020-00918-2>, 2020.
- Li, Q., England, M. H., Hogg, A. M., Rintoul, S. R., and Morrison, A. K.: Abyssal Ocean Overturning Slowdown and Warming Driven by Antarctic Meltwater, *Nature*, 615, 841–847, <https://doi.org/10.1038/s41586-023-05762-w>, 2023.
- Liu, Y., Moore, J. K., Primeau, F., and Wang, W. L.: Reduced CO<sub>2</sub> Uptake and Growing Nutrient Sequestration from Slowing Overturning Circulation, *Nature Climate Change*, 13, 83–90, <https://doi.org/10.1038/s41558-022-01555-7>, 2023.
- 795 Lomas, M. W., Bates, N. R., Johnson, R. J., Steinberg, D. K., and Tanioka, T.: Adaptive Carbon Export Response to Warming in the Sargasso Sea, *Nature Communications*, 13, 1211, <https://doi.org/10.1038/s41467-022-28842-3>, 2022.
- Matear, R. J. and Hirst, A. C.: Climate change feedback on the future oceanic CO<sub>2</sub> uptake, *Tellus B*, 51, 722–733, <https://doi.org/10.1034/j.1600-0889.1999.t01-1-00012.x>, 1999.

- Meinshausen, M., Smith, S. J., Calvin, K., Daniel, J. S., Kainuma, M. L. T., Lamarque, J.-F., Matsumoto, K., Montzka, S. A., Raper, S. C. B.,  
800 Riahi, K., Thomson, A., Velders, G. J. M., and van Vuuren, D. P.: The RCP Greenhouse Gas Concentrations and Their Extensions from  
1765 to 2300, *Climatic Change*, 109, 213, <https://doi.org/10.1007/s10584-011-0156-z>, 2011.
- Meinshausen, M., Nicholls, Z. R. J., Lewis, J., Gidden, M. J., Vogel, E., Freund, M., Beyerle, U., Gessner, C., Nauels, A., Bauer, N., Canadell,  
J. G., Daniel, J. S., John, A., Krummel, P. B., Luderer, G., Meinshausen, N., Montzka, S. A., Rayner, P. J., Reimann, S., Smith, S. J., van den  
Berg, M., Velders, G. J. M., Vollmer, M. K., and Wang, R. H. J.: The shared socio-economic pathway (SSP) greenhouse gas concentrations  
805 and their extensions to 2500, *Geoscientific Model Development*, 13, 3571–3605, <https://doi.org/10.5194/gmd-13-3571-2020>, 2020.
- Moore, J. K., Fu, W., Primeau, F. W., Britten, G. L., Lindsay, K., Long, M., Doney, S. C., Mahowald, N., Hoffman, F.,  
and Randerson, J. T.: Sustained climate warming drives declining marine biological productivity, *Science*, 359, 1139–1143,  
<https://doi.org/10.1126/science.aao6379>, 2018.
- Murnane, R. J., Sarmiento, J. L., and Le Quéré, C.: Spatial distribution of air-sea CO<sub>2</sub> fluxes and the interhemispheric transport of carbon by  
810 the oceans, *Global Biogeochemical Cycles*, 13, 287–305, <https://doi.org/10.1029/1998GB900009>, 1999.
- Palter, J. B., Sarmiento, J. L., Gnanadesikan, A., Simeon, J., and Slater, R. D.: Fueling export production: nutrient return pathways from the  
deep ocean and their dependence on the Meridional Overturning Circulation, *Biogeosciences*, 7, 3549–3568, <https://doi.org/10.5194/bg-7-3549-2010>, 2010.
- Pasquier, B. and Holzer, M.: The plumbing of the global biological pump: Efficiency control through leaks, pathways, and time scales,  
815 *Journal of Geophysical Research: Oceans*, 121, 6367–6388, <https://doi.org/10.1002/2016JC011821>, 2016.
- Pasquier, B. and Holzer, M.: The number of past and future regenerations of iron in the ocean and its intrinsic fertilization efficiency,  
*Biogeosciences*, 15, 7177–7203, <https://doi.org/10.5194/bg-15-7177-2018>, 2018.
- Pasquier, B., Holzer, M., Chamberlain, M. A., Matear, R. J., Bindoff, N. L., and Primeau, F. W.: Optimal parameters for the ocean’s nu-  
trient, carbon, and oxygen cycles compensate for circulation biases but replumb the biological pump, *Biogeosciences*, 20, 2985–3009,  
820 <https://doi.org/10.5194/bg-20-2985-2023>, 2023.
- Passow, U. and Carlson, C. A.: The Biological Pump in a High CO<sub>2</sub> World, *Marine Ecology Progress Series*, 470, 249–271,  
<https://doi.org/10.3354/meps09985>, 2012.
- Planchat, A., Kwiatkowski, L., Bopp, L., Torres, O., Christian, J. R., Butenschön, M., Lovato, T., Séférian, R., Chamberlain, M. A., Aumont,  
O., Watanabe, M., Yamamoto, A., Yool, A., Ilyina, T., Tsjujino, H., Krumhardt, K. M., Schwinger, J., Tjiputra, J., Dunne, J. P., and Stock,  
825 C.: The Representation of Alkalinity and the Carbonate Pump from CMIP5 to CMIP6 Earth System Models and Implications for the  
Carbon Cycle, *Biogeosciences*, 20, 1195–1257, <https://doi.org/10.5194/bg-20-1195-2023>, 2023.
- Plattner, G.-K., Joos, F., Stocker, T. F., and Marchal, O.: Feedback mechanisms and sensitivities of ocean carbon uptake under global  
warming, *Tellus B: Chemical and Physical Meteorology*, 53, 564–592, <https://doi.org/10.3402/tellusb.v53i5.16637>, 2001.
- Primeau, F. W.: Characterizing Transport between the Surface Mixed Layer and the Ocean Interior with a Forward and Adjoint Global Ocean  
830 Transport Model, *Journal of Physical Oceanography*, 35, 545–564, <https://doi.org/10.1175/JPO2699.1>, 2005.
- Primeau, F. W. and Holzer, M.: The Ocean’s Memory of the Atmosphere: Residence-Time and Ventilation-Rate Distributions of Water  
Masses, *Journal of Physical Oceanography*, 36, 1439–1456, <https://doi.org/10.1175/JPO2919.1>, 2006.
- Primeau, F. W., Holzer, M., and DeVries, T.: Southern Ocean nutrient trapping and the efficiency of the biological pump, *Journal of Geo-  
physical Research: Oceans*, 118, 2547–2564, <https://doi.org/10.1002/jgrc.20181>, 2013.
- 835 Purich, A. and England, M. H.: Projected Impacts of Antarctic Meltwater Anomalies over the Twenty-First Century, *Journal of Climate*, 36,  
2703–2719, <https://doi.org/10.1175/JCLI-D-22-0457.1>, 2023.

- Purich, A., England, M. H., Cai, W., Sullivan, A., and Durack, P. J.: Impacts of Broad-Scale Surface Freshening of the Southern Ocean in a Coupled Climate Model, *Journal of Climate*, 31, 2613–2632, <https://doi.org/10.1175/JCLI-D-17-0092.1>, 2018.
- 840 Raven, J. A. and Falkowski, P. G.: Oceanic sinks for atmospheric CO<sub>2</sub>, *Plant, Cell & Environment*, 22, 741–755, <https://doi.org/10.1046/j.1365-3040.1999.00419.x>, 1999.
- Revelle, R. and Suess, H. E.: Carbon Dioxide Exchange Between Atmosphere and Ocean and the Question of an Increase of Atmospheric CO<sub>2</sub> during the Past Decades, *Tellus*, 9, 18–27, <https://doi.org/10.3402/tellusa.v9i1.9075>, 1957.
- Riahi, K., van Vuuren, D. P., Kriegler, E., Edmonds, J., O'Neill, B. C., Fujimori, S., Bauer, N., Calvin, K., Dellink, R., Fricko, O., Lutz, W., Popp, A., Cuaresma, J. C., KC, S., Leimbach, M., Jiang, L., Kram, T., Rao, S., Emmerling, J., Ebi, K., Hasegawa, T., Havlik, P., 845 Humpenöder, F., Da Silva, L. A., Smith, S., Stehfest, E., Bosetti, V., Eom, J., Gernaat, D., Masui, T., Rogelj, J., Strefler, J., Drouet, L., Krey, V., Luderer, G., Harmsen, M., Takahashi, K., Baumstark, L., Doelman, J. C., Kainuma, M., Klimont, Z., Marangoni, G., Lotze-Campen, H., Obersteiner, M., Tabeau, A., and Tavoni, M.: The Shared Socioeconomic Pathways and their energy, land use, and greenhouse gas emissions implications: An overview, *Global Environmental Change*, 42, 153–168, <https://doi.org/10.1016/j.gloenvcha.2016.05.009>, 2017.
- 850 Riebesell, U., Körtzinger, A., and Oschlies, A.: Sensitivities of marine carbon fluxes to ocean change, *Proceedings of the National Academy of Sciences*, 106, 20602–20609, <https://doi.org/10.1073/pnas.0813291106>, 2009.
- Sarmiento, J. L. and Le Quéré, C.: Oceanic Carbon Dioxide Uptake in a Model of Century-Scale Global Warming, *Science*, 274, 1346–1350, <https://doi.org/10.1126/science.274.5291.1346>, 1996.
- Sarmiento, J. L., Hughes, T. M. C., Stouffer, R. J., and Manabe, S.: Simulated Response of the Ocean Carbon Cycle to Anthropogenic 855 Climate Warming, *Nature*, 393, 245–249, <https://doi.org/10.1038/30455>, 1998.
- Sarmiento, J. L., Gruber, N., Brzezinski, M. A., and Dunne, J. P.: High-Latitude Controls of Thermocline Nutrients and Low Latitude Biological Productivity, *Nature*, 427, 56–60, <https://doi.org/10.1038/nature02127>, 2004.
- Schmittner, A., Oschlies, A., Matthews, H. D., and Galbraith, E. D.: Future changes in climate, ocean circulation, ecosystems, and biogeochemical cycling simulated for a business-as-usual CO<sub>2</sub> emission scenario until year 4000 AD, *Global Biogeochemical Cycles*, 22, 860 <https://doi.org/10.1029/2007GB002953>, 2008.
- Sweetman, A. K., Thurber, A. R., Smith, C. R., Levin, L. A., Mora, C., Wei, C.-L., Gooday, A. J., Jones, D. O. B., Rex, M., Yasuhara, M., Ingels, J., Ruhl, H. A., Frieder, C. A., Danovaro, R., Würzberg, L., Baco, A., Grupe, B. M., Pasulka, A., Meyer, K. S., Dunlop, K. M., Henry, L.-A., and Roberts, J. M.: Major Impacts of Climate Change on Deep-Sea Benthic Ecosystems, *Elementa: Science of the Anthropocene*, 5, 4, <https://doi.org/10.1525/elementa.203>, 2017.
- 865 Taylor, K., Stouffer, R., and Meehl, G.: An Overview of CMIP5 and the experiment design, *Bull. Amer. Meteor. Soc.*, 93, 485–498, <https://doi.org/10.1175/BAMS-D-11-00094.1>, 2012.
- Thornton, P. E., Doney, S. C., Lindsay, K., Moore, J. K., Mahowald, N., Randerson, J. T., Fung, I., Lamarque, J.-F., Feddesma, J. J., and Lee, Y.-H.: Carbon-nitrogen interactions regulate climate-carbon cycle feedbacks: results from an atmosphere-ocean general circulation model, *Biogeosciences*, 6, 2099–2120, <https://doi.org/10.5194/bg-6-2099-2009>, 2009.
- 870 Toggweiler, J. R., Gnanadesikan, A., Carson, S., Murnane, R., and Sarmiento, J. L.: Representation of the carbon cycle in box models and GCMs: 1. Solubility pump, *Global Biogeochemical Cycles*, 17, <https://doi.org/10.1029/2001GB001401>, 2003.
- van Heuven, S., Pierrot, D., Rae, J., Lewis, E., and Wallace, D.: CO<sub>2</sub>SYS v1.1, MATLAB program developed for CO<sub>2</sub> system calculations, ORNL/CDIAC-105b. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, U.S. DoE, Oak Ridge, TN., 2011.

- Volk, T. and Hoffert, M. I.: Ocean Carbon Pumps: Analysis of Relative Strengths and Efficiencies in Ocean-Driven Atmospheric CO<sub>2</sub> Changes, in: *The Carbon Cycle and Atmospheric CO<sub>2</sub>: Natural Variations Archean to Present*, edited by Sundquist, E. T. and Broecker, W. S., pp. 99–110, American Geophysical Union (AGU), ISBN 9781118664322, <https://doi.org/10.1029/GM032p0099>, 1985.
- 875
- Wanninkhof, R.: Relationship between wind speed and gas exchange over the ocean revisited, *Limnology and Oceanography: Methods*, 12, 351–362, <https://doi.org/10.4319/lom.2014.12.351>, 2014.
- Weiss, R. F.: Carbon Dioxide in Water and Seawater: The Solubility of a Non-Ideal Gas, *Marine Chemistry*, 2, 203–215, [https://doi.org/10.1016/0304-4203\(74\)90015-2](https://doi.org/10.1016/0304-4203(74)90015-2), 1974.
- 880
- Whitney, F. A., Freeland, H. J., and Robert, M.: Persistently Declining Oxygen Levels in the Interior Waters of the Eastern Subarctic Pacific, *Progress in Oceanography*, 75, 179–199, <https://doi.org/10.1016/j.pocean.2007.08.007>, 2007.
- Wilson, J. D., Andrews, O., Katavouta, A., de Melo Virissimo, F., Death, R. M., Adloff, M., Baker, C. A., Blackledge, B., Goldsworth, F. W., Kennedy-Asser, A. T., Liu, Q., Sieradzan, K. R., Vosper, E., and Ying, R.: The biological carbon pump in CMIP6 models: 21st century trends and uncertainties, *Proceedings of the National Academy of Sciences*, 119, e2204369 119, <https://doi.org/10.1073/pnas.2204369119>, 2022.
- 885