

**~~Inclination~~ Land inclination controls CO<sub>2</sub> and N<sub>2</sub>O fluxes, but not CH<sub>4</sub> uptake, from a temperate upland forest soil**

Lauren M. Gillespie<sup>1</sup>, Nathalie Y. Triches<sup>2</sup>, Diego Abalos<sup>3</sup>, Peter Finke<sup>4</sup>, Sophie Zechmeister-Boltenstern<sup>1</sup>, Stephan Glatzel<sup>5</sup>, Eugenio Díaz-Pinés<sup>1</sup>

<sup>1</sup> Institute of Soil Research, University of Natural Resources and Life Sciences, Vienna (BOKU), Peter-Jordan-Straße 82, 1190, Vienna, Austria.

<sup>2</sup> Department of Environment, Faculty of Bioscience Engineering, Ghent University, Coupure links 653, B-9000, Ghent, Belgium. Present address: Max Planck Research Institute for Biogeochemistry, Hans-Knöll-Straße 10, 07745, Jena, Germany.

<sup>3</sup> Department of Agroecology, iCLIMATE, Aarhus University, Blichers Allé 20, Tjele, 8830, Denmark.

<sup>4</sup> Department of Environment, Research group of soilscape genesis, Ghent University, Coupure links 653, B-9000, Ghent, Belgium.

<sup>5</sup> University of Vienna, Department of Geography and Regional Research, Universitätsstraße 7, 1010 Vienna, Austria.

**Abstract**

Inclination and spatial variability in soil and litter properties influence soil greenhouse gas (GHG) fluxes, and thus on-going climate change, but their relationship in forest ecosystems is poorly understood. To elucidate this, we explored the effect of inclination, distance to a stream, soil moisture, soil temperature, and other soil and litter properties on soil-atmosphere fluxes of carbon dioxide (CO<sub>2</sub>), methane (CH<sub>4</sub>), and nitrous oxide (N<sub>2</sub>O) with automated static chambers in a temperate upland forest in Eastern Austria. We hypothesised that soil CO<sub>2</sub> emissions and CH<sub>4</sub> uptake are higher in sloped locations with lower soil moisture content, whereas soil N<sub>2</sub>O emissions are higher in flat, wetter locations. During the measurement period, soil CO<sub>2</sub> emissions were significantly higher on flat locations ( $p < 0.05$ ), and increased with increasing soil temperature ( $p < 0.001$ ) and decreasing soil moisture ( $p < 0.001$ ). The soil acted as a CH<sub>4</sub>

**Commenté [LG1]:** Please give a more specific title. The word “Inclination” can have multiple meaning. Please make it a bit clearer in the title what inclination is being referring to. Land inclination?? Slope of the land??

**Response:** We will use “Land inclination...” in the title to remove ambiguity.

27 sink, and CH<sub>4</sub> uptake was not significantly related to inclination. However, CH<sub>4</sub> uptake was  
28 significantly higher at locations furthest away from the stream compared to at the stream ( $p <$   
29 0.001), and positively related to litter weight and soil C content ( $p < 0.01$ ). N<sub>2</sub>O fluxes were  
30 significantly higher on flat locations and further away from the stream ( $p < 0.05$ ), and increased  
31 with increasing soil moisture ( $p < 0.001$ ), soil temperature ( $p < 0.001$ ) and litter depth ( $p <$   
32 0.05). Overall, this study underlines the importance of inclination and the resulting soil and  
33 litter properties in predicting GHG fluxes from forest soils and therefore their potential source-  
34 sink balance.

35  
36 **Keywords:** slope inclination, soil greenhouse gas fluxes, carbon dioxide, methane, nitrous  
37 oxide, soil moisture, forest litter

38  
39 **Introduction**

40 Forests play a crucial role in the global climate by emitting and consuming the greenhouse gases  
41 (GHGs) carbon dioxide (CO<sub>2</sub>), methane (CH<sub>4</sub>), and nitrous oxide (N<sub>2</sub>O) (IPCC, 2022). They  
42 store a large amount of carbon (C) in the vegetation and soil organic matter and can be effective  
43 CO<sub>2</sub> sinks (Pan et al., 2011). Soil microorganisms also take up atmospheric C through the  
44 oxidation of CH<sub>4</sub> during methanotrophy (Le Mer and Roger, 2001; Hiltbrunner et al., 2012).  
45 However, forest soils also emit substantial quantities of CO<sub>2</sub> (Webster et al., 2008), which, in  
46 aerobic conditions, is mainly released by root respiration and microbial respiration during  
47 decomposition (Cronan, 2018; Zechmeister-Boltenstern et al., 2018). ~~Nitrous oxide (N<sub>2</sub>O)~~ is  
48 produced by soil microorganisms, mainly during nitrification and denitrification (Butterbach-  
49 Bahl et al., 2013). In aerobic conditions, bacteria convert ammonium to nitrite and further to  
50 nitrate during nitrification. In anoxic conditions, nitrate is then used as an alternative electron  
51 acceptor instead of O<sub>2</sub> and reduced to N<sub>2</sub> during denitrification (Butterbach-Bahl et al., 2014).  
52 Under most conditions, these processes occur simultaneously and usually result in a net

**Commenté [LG2]:** Abstract  
In the keywords, why is topography included? Topography  
has not been discussed in this paper

**Response:** We thank Reviewer 2 for catching this, it will be  
replaced with 'slope inclination'.

**Commenté [LG3]:** L44: don't need to repeat N<sub>2</sub>O as it has  
already been stated above in line #38

**Response:** We thank Reviewer 2 for pointing this out. It will  
be removed.

atmospheric emission of N<sub>2</sub>O (Ambus, 1998). Conversely, net N<sub>2</sub>O uptake has been reported from forest soils, especially since monitoring instrumentation has become sensitive enough to measure very low fluxes (Savage et al., 2014; Subke et al., 2021). Net N<sub>2</sub>O uptake (from the atmosphere into the soil) is a complex process closely tied to N<sub>2</sub>O consumption (within the soil) that is driven principally by denitrifying bacteria (Liu et al., 2022).

Temporal and spatial variations in soil CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O fluxes are driven mostly by changes in soil temperature and soil moisture (Raich and Potter, 1995; Davidson et al., 1998; Le Mer and Roger, 2001; Butterbach-Bahl et al., 2014). Rising temperatures accelerate microbial activities and, consequently, the production and emission of N<sub>2</sub>O and CO<sub>2</sub> (Butterbach-Bahl et al., 2013). Elevated soil respiration could lead to a depletion of O<sub>2</sub>, which also results in increased N<sub>2</sub>O from denitrification (Butterbach-Bahl et al., 2013). Contrarily, CH<sub>4</sub> uptake appears to be less sensitive to temperature changes than CO<sub>2</sub> and N<sub>2</sub>O fluxes (Hanson and Hanson, 1996). Soil moisture has a major influence on all GHG fluxes by regulating O<sub>2</sub> and substrate availability to soil microorganisms and influencing the diffusion of gases within the soil matrix (Butterbach-Bahl et al., 2014; Schimel, 2018). Indeed, soil microbial activity decreases as soils become water saturated ~~saturated due to O<sub>2</sub> limitation~~ (Davidson et al., 2012). Soil moisture further affects fluxes since diffusion coefficients of GHG in air are approximately 10<sup>4</sup> times larger than in water (Marrero and Mason, 1972).

Inclination and distance to a water source influence some of the most important drivers of soil GHG fluxes. For example, soil moisture content changes on small scales at different inclinations through accumulation, runoff, and leaching of precipitation water (Burt and Butcher, 1985; Lookingbill and Urban, 2004; Lin et al., 2006). Inclination also modifies other important drivers of soil GHG fluxes, such as the hydrological transport of nutrients (Hairston and Grigal, 1994), litter accumulation (Butler et al., 1986), soil aeration, soil texture, soil pH, and substrate availability (soil C and N), usually resulting in a high GHG spatial variability (e.g., Fierer and Jackson, 2006; Thomas and Packham, 2007). Flat locations by a water source

**Commenté [LG4]:** 1.1. 50: Could you please add some information about the processes behind N<sub>2</sub>O uptake by soils?

**Response:** We propose to add:

“Net N<sub>2</sub>O uptake (from the atmosphere into the soil) is a complex process closely tied to N<sub>2</sub>O consumption (within the soil) that is driven principally by denitrifying bacteria (Liu et al. 2022).”

Although we do not have space to fully explain soil N<sub>2</sub>O uptake and N<sub>2</sub>O consumption in this study, since it was not the primary focus, the readers will now be able to find additional explanations in the newly added reference.

Liu, H., Li, Y., Pan, B. *et al.* Pathways of soil N<sub>2</sub>O uptake, consumption, and its driving factors: a review. *Environ Sci Pollut Res* **29**, 30850–30864 (2022). <https://doi.org/10.1007/s11356-022-18619-y>

**Commenté [LG5]:** L62-63: Please revise this statement.

**Response:** We will replace “saturated due to O<sub>2</sub> limitation” by “water saturated”.

are also at higher risk to be influenced by flooding and subsequent changes to the soil properties and soil microbial community (Ou et al., 2019; Unger et al., 2009). Forest litter in particular can have a major impact on the exchange of GHGs by adding nutrients to the soil, acting as a physical barrier (i.e., holding gases in the soil rather than releasing them into the atmosphere) or influencing the water and heat exchange between soil and atmosphere (Leitner et al., 2016; Walkiewicz et al., 2021).

Studies on the effect of inclination on GHG fluxes from temperate upland forest soils are particularly rare. Some studies reported higher soil CO<sub>2</sub> emissions on sloped compared to flat locations, associated with warmer air and soil temperatures and lower soil moisture contents, favouring faster diffusion rates though not so low as to impede microbial activity (Yu et al., 2008; Warner et al., 2018). Conversely, no effect of topography on soil CO<sub>2</sub> emissions has also been reported in a laboratory study from a montane tropical forest (Arias-Navarro et al., 2017). With regard to CH<sub>4</sub>, relatively little is known on how inclination and its influence on chemical and physical soil properties may affect CH<sub>4</sub> fluxes (Warner et al., 2018). Soil CH<sub>4</sub> uptake is highly variable in space and time and appears to be highest on dry slopes (Hiltbrunner et al., 2012; Yu et al., 2021), even though it is assumed that temperate upland forest soils take up CH<sub>4</sub> irrespective of the inclination (Lamprea Pineda et al., 2021). Effects of inclination on N<sub>2</sub>O fluxes are also contradictory. Some studies show increased N<sub>2</sub>O emissions with higher soil water content at flat locations (Davidson et al., 2000; Lamprea Pineda et al., 2021), whereas others show a higher emission in aerated soils on slopes (Yu et al., 2008, 2021). Assessing the impact of inclination on soil GHGs therefore remains a challenging task.

In this study, we aim to improve the understanding of the effects of inclination and distance to a stream on the emission and uptake of GHGs in a temperate upland forest soil in Eastern Austria. We monitored soil CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O fluxes with automated chambers over six months for two different inclinations and at four distances from a stream in a deciduous forest. We tested three hypotheses: 1) Soil CO<sub>2</sub> emissions are higher in sloped than flat locations

**Commenté [LG6]:** This study shows the effect of slope (of the land) and distance (to the stream) on GHGs in a temperate forest soil. GHG emissions are modified by the local land conditions, slopes, and topography, and it is very important to take into account these factors when looking at landscape-scale emissions. The authors used recent technological analyzers known for their high precision and sensitivity to demonstrate how CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O fluxes vary within a short space. However, there are a number of issues that should be addressed. The main aspects that need to be revisited are the interpretation of the results, and how Inclination and distance are regarded as factors for the changes in the emissions. Distance by itself is not a factor causing the differences in the emissions across the plots, rather the changes in the soil properties, which are of course not, modified by the distance itself.

**Response:** In this study, we argue that inclination and distance to a water source can be used as proxies for the variations in soil moisture, soil temperature, and other soil properties (Line 66-77). Distance to the stream and inclination influence soil water retention, which also influences soil temperature, as well as litter retention (e.g. litter descending from sloped locations and accumulating at the flat locations) and nutrient leaching.

**Commenté [LG7]:** L91-92: The impact of topographic variation hasn't been studied so much with regard to GHG emissions. Is it due to the difficult nature of the task or the general assumption that the slope has no impact on GHG emissions?

**Response:** Indeed, it has not been studied much in the past, and we believe that to a large extent this may be due to the challenging logistics required for it, in terms of equipment that captures short-term changes in gaseous emissions, appropriate chambers for the task, and a suitable site. Our state-of-the-art instrumentation and a comprehensively monitored site allowed us to overcome these challenges and provide, as the reviewer indicates, one of the few available datasets documenting the impact of topographic variation on GHG emissions in a forest soil.

because of the inclination *per se* and the lower soil moisture content at sloped locations; 2) Soil CH<sub>4</sub> uptake is higher in sloped than flat locations because of the inclination *per se* and the lower soil moisture content at sloped locations; and 3) Soil N<sub>2</sub>O emissions are lower in sloped than flat locations because of the inclination *per se* and the higher soil moisture content at flat locations.

## Methods

### Study site and experimental design

This study was conducted within the framework of the “Long-Term Ecosystem and socio-ecological Research Infrastructure - Carbon, Water and Nitrogen” (LTER-CWN) project (further information is available at <https://www.lter-austria.at/en/cwn-sites/>). The BOKU University Forest “Rosalia Lehrforst” (47°42′25.35″ N / 16°16′36.62″ E) is one of the associated sites and served as the site for our study (see Fürst et al., 2021) for more information). At the site, European beech (*Fagus sylvatica* L.) and Norway spruce (*Picea abies* (L.) H. Karst.) are the dominant tree species, but alluvial forest species (*Alnus* spp. Mill, *Fraxinus excelsior* L.) are also present next to the study location. The elevation is around 400 m a.s.l. and the dominant soil type is pseudo-gleyic Cambisol (Schad, 2016).

We used the GasFluxTrailer (explained below) to measure soil GHG fluxes from 17 June to 24 November 2020. We positioned 16 chambers linearly in groups of four at four different distances from a small forest stream: 0.5 m, 5 m, 10 m, and 15 m (Fig. S1). Adjacent trees to the chambers were *F. sylvatica* and *P. abies*. These distances served as first treatment effect and are hereafter referred to as chamber group (CG): CG0.5, CG5, CG10, and CG15. These distances were chosen because they were expected to cover a decreasing soil moisture gradient from CG0.5 towards CG15. To measure this gradient, one Em50 (METER Group, Inc. Pullman, WA; USA) was connected to four ECH2O 5 TM volumetric water content and temperature sensors (METER Group). One sensor was installed per CG approximately one meter away from

#### Commenté [LG8]: 2.1. 98-102: Hypotheses:

a) In the hypotheses you speak about “inclination *per se*”. Could you please explain, what is exactly meant with “*per se*”? For me inclination is mainly a proxy for other directly influencing site or soil parameters like soil moisture of temperature.

**Response:** We agree with Reviewer 1 that the three “*per se*”’s in the hypotheses do not add any meaningful information and lead to confusion. We will remove them.

b) Hypothesis No. 3 addresses potential impact on “N<sub>2</sub>O emissions”. In the following results and discussion sections the term “N<sub>2</sub>O fluxes” is primarily used instead of “N<sub>2</sub>O emission”.

**Response:** We thank Reviewer 1 for bringing this to our attention. We will make sure to use ‘emissions’ and ‘uptake’ when appropriate and only use ‘fluxes’ when referring to both emissions and uptake.

c) In the results section data on N<sub>2</sub>O uptake are provided but there is no related hypothesis.

**Response:** N<sub>2</sub>O uptake had only recently been measured reliably thanks to advancement in equipment and is generally minimal in comparison with emissions. We believe it important to report on this N<sub>2</sub>O uptake since its existence is interesting, but it does not merit a dedicated hypothesis since its occurrence is infrequent and unlikely to play a dominant role in soil greenhouse gas fluxes.

**Commenté [LG9]:** The authors may look at the historical conditions at the site. The measurements are set up close to a watercourse, but the possible floods and their consequences have not been adequately discussed in this manuscript. Frequent inundation can lead to varying soil properties, and drying-rewetting enhances decomposition.

**Response:** We can add further detail on how the flat locations by the stream are at higher risk to be influenced by flooding and how that may have influenced soil properties. We can also add that there were no inundations or significant drying/rewetting events during the measurements.

**Commenté [LG10]:** L115: GasFluxTrailer is a platform. This statement sounds GasFluxTrailer was used to measure GHGs, but the trailer is the platform to position your gas analyzer.

**Response:** We refer to the GasFluxTrailer as the combination of the gas chambers, the multiplexer, and the gas analysers, as a unit. It is thanks to this unit that the GHG fluxes are measured.

**Commenté [LG11]:** 3.116-117: Some additional information about the investigated forest sites would be helpful here. Are all locations covered by the same tree species or what about the exposition of the slopes?

**Response:** We agree and will add the exposition of the slope and the adjacent tree species.

the outer chamber (Fig S.1) one Em50 data logger was installed per CG approximately one meter away from the outer chamber measuring soil temperature and soil moisture (see Fig. S1). The four Em50 data loggers were each connected to an ECH<sub>2</sub>O 5™ Volumetric Water Content and Temperature sensor. As a second treatment effect, the distances were also chosen so that

the CGs were set up at two different inclinations. CG0.5 and CG5 were located at flat (average 1°; the slope at these distances did not exceed 2°), CG10 and CG15 at sloped locations (average 35°; west-facing).

For meteorological information, we used the precipitation (OTT Pluvio L weighing rain gauge) and air temperature (air temperature and humidity sensor TR1) data recorded at 30 min intervals by the weather station “Mehlbeerleiten”, located approximately 100 m north-west of the site (Diaz-Pines and Gasch, 2021; Fürst et al., 2021).

#### *Gas flux measurements: GasFluxTrailer*

An automated and mobile measuring system was used, termed the GasFluxTrailer. It consists of a mobile trailer estimating soil-atmosphere GHG exchange rates of CH<sub>4</sub>, CO<sub>2</sub>, and N<sub>2</sub>O. The GasFluxTrailer connects with the chambers, and it controls the sampling of each individual chamber (i.e., the opening and closing and gas sampling) and recording of the gas concentrations. The 16 automated, static, non-steady-state, non-flow-through chambers (Pumpanen et al., 2004) with an area of 0.5 m × 0.5 m and height of 0.15 m are made of stainless-steel and placed on stainless-steel frames of the same area. They are equipped with fans to ensure homogenous air mixing. The gas analysers are a G2301 (PICARRO Inc., Santa Clara, USA), measuring concentrations of CO<sub>2</sub> and CH<sub>4</sub>, and a G5131i (PICARRO Inc.), estimating N<sub>2</sub>O concentrations. The software used to run automatic sequences is the IDASw Recorder 4.5.0., developed by the Institute of Meteorology and Climate Research Atmospheric Environmental Research (IMK-IFU) in Germany.

**Commenté [LG12]:** L123: The manufacturing company name, and country is missing

**Response:** Thank you for pointing this out. This part should read: “to measure this gradient, one Em50 (METER Group, Inc. Pullman, WA; USA) was connected to four ECH<sub>2</sub>O 5™ volumetric water content and temperature sensors (METER Group). One sensor was installed per CG approximately one meter away from the outer chamber (see Fig S.1)”

**Commenté [LG13]:** L133/134: Again here. The trailer is being mentioned as a system estimating the gas exchanges. This may confuse the readers. The gas samples are analyzed by the two picarro analyzers.

**Response:** Although the GHG quantification was conducted by the PICARRO analysers, we refer to the GasFluxTrailer as the unit needed for the overall flux estimation, since the trailer does not just contain the analysers but also all equipment required for the automated measurements, including for example the software used to control the entire sampling process, i.e. opening and closing of the chambers, transport of the gas from the chambers, etc.

**Commenté [LG14]:** L142-144: Chambers closing and opening simultaneously or successively

**Response:** The closing and opening is done successively. This information will be added to the text.

*Field and laboratory measurements*

We inserted the chamber frames 5 cm deep into the soil approximately one month before the measuring campaign to avoid additional soil CO<sub>2</sub> release from cut roots, affecting our measurements (Davidson et al., 2002). For each measurement estimate, a chamber was closed for 10 min, which, thanks to the highly sensitive instruments used here, was sufficient time to measure gas concentrations changes, including low N<sub>2</sub>O fluxes (Harris et al., 2021). The closing and opening was done successively; thus ~~One full cycle of all 16 chambers took 160 minutes.~~

We calculated fluxes with a linear regression approach according to Butterbach-Bahl et al. (2011). This was justified with short chamber closure times and a relatively large chamber size (Hutchinson and Mosier, 1981). Positive flux values indicate gas emission from the soil, and negative values indicate net uptake. To ensure the system was running and working correctly, we controlled the GHG flux measurements on-site every week and three-four times a week remotely. There were no inundations or significant drying/rewetting events during the observation period.

Close to each of the 16 chambers, a litter and soil sample was collected in December 2021. The litter depth was measured first, before disturbing the litter and topsoil by placing a 0.2 m × 0.2 m frame on the ground at this location. The litter was then collected within this frame, dried at 65°C for 7 days, and weighed. After litter collection and removal of organic layer, two soil cores (stainless steel core, 7 cm diameter, 7 cm depth) were taken from the topsoil mineral layer for analyses of pH, C and N content, and soil texture. C and N contents (%) were determined by dry combustion on 1.6 mg of soil using the Austrian standard ÖNORM L 1080 (ÖNORM, 2013). Particle size analysis was conducted using the pipette method on 10 g of soil according to the Austrian standard ÖNORM L 1061 (ÖNORM, 2002), after the organic material had been burned off in an oven at 550 °C, to determine soil texture (%). In short, sieved soil (<2 mm) is agitated in a volume of water, and a pipette is used to sample a defined volume at a defined depth at specific times after which the samples are dried to determine clay and silt contents.

**Commenté [LG15]:** L200: This statement should be moved from here to the above section (Field measurements, L149-152).

**Response:** This will be done.

183 The remaining soil is then sieved (63 µm) to determine sand content. Soil pH was measured on  
 184 5 g of soil with 0.01 M CaCl<sub>2</sub> using the Austrian standard ÖNORM L 1083 (ÖNORM, 2006).  
 185 Because the soil was relatively rocky, we calculated the soil bulk density (BD, g cm<sup>-3</sup>) including  
 186 the coarse (stone) fraction as:

$$BD \text{ with stones} = \frac{\text{dry soil weight}}{\text{core volume}}$$

187 where dry soil weight is the weight of the soil in the core after oven drying in g and core volume  
 188 is the volume of the core in cm<sup>3</sup>.  
 190 We calculated the total porosity (Φ) using the bulk density and an estimated soil particle density,  
 191 obtained by a weighted average of the specific weights of mineral material (2.65 g cm<sup>-3</sup>) and  
 192 organic matter (1.45 g cm<sup>-3</sup>). We took into account the organic matter content because it was  
 193 relatively high, i.e., between 8 and 27 %.

#### 195 *Data processing and statistics*

196 We quality-controlled the CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O flux data using the determination coefficient (R-  
 197 squared, R<sup>2</sup>) values between GHG concentrations and the time after chamber closure. For CO<sub>2</sub>  
 198 and CH<sub>4</sub>, we filtered the data with R<sup>2</sup> > 0.8 and a visual plausibility check based on expert  
 199 knowledge. For N<sub>2</sub>O, R<sup>2</sup> > 0.8 was applied only if fluxes were > 5 µg N<sub>2</sub>O-N m<sup>-2</sup> h<sup>-1</sup>. For low  
 200 flux rates (< 5 µg N<sub>2</sub>O-N m<sup>-2</sup> h<sup>-1</sup>), we did not remove values with R<sup>2</sup> < 0.8 if corresponding CO<sub>2</sub>  
 201 fluxes were valid. We kept these measurements in the dataset, because the low R<sup>2</sup> values were  
 202 due to fluxes below the detection limit of the system; however, the measurement itself remained  
 203 valid as indicated by plausible CO<sub>2</sub> fluxes, and as elaborated in Parkin et al. (2012). Through  
 204 this quality control, we found that two chambers did not produce any reliable measurements  
 205 from 24 September onwards. August data for all chambers was excluded due to malfunctioning  
 206 of the equipment that was not initially detected. Furthermore, all the data from one chamber  
 207 (chamber 13) were also not used for the analysis because of a failure in the chamber gas

**Commenté [LG16]:** 4.172/1. 237/Tab. 1: It is mentioned that the investigated soils have a larger stone content, which is also confirmed by the data in Tab. 1. However, bulk density – explicitly including the coarse soil fraction – with values clearly below 1.0 g cm<sup>-3</sup> is very low and values of 0.15 or 0.12 g cm<sup>-3</sup> seem to be unrealistic. The reason for this low bulk density should be explained and data checked (see comment on Tab. 1 below).

**Response:** Thanks for spotting this; there was indeed an issue with the values in Table 1 (see our response to Reviewer 1's comment 6; the corrected Table 1 can be found in the Supplement document) and the 0.15 and 0.12 g cm<sup>-3</sup> were incorrect. This has been corrected in the table at the end of our comments to the reviewers. We consider bulk densities below 1.0 g cm<sup>-3</sup> (0.6-0.8 in our case) are realistic and common for forest soils (Beguin et al. 2017; Llek et al. 2017), even considering the stone content (7-13 % vol).

Beguin, J., Fuglstad, G. A., Mansuy, N., & Paré, D. (2017). Predicting soil properties in the Canadian boreal forest with limited data: Comparison of spatial and non-spatial statistical approaches. *Geoderma*, 306, 195-205.  
 Ilek, A., Kucza, J., & Szostek, M. (2017). The effect of the bulk density and the decomposition index of organic matter on the water storage capacity of the surface layers of forest soils. *Geoderma*, 285, 27-34.



208 sampling. After data quality screening, there were 125 measurement days included in analysis  
209 for CO<sub>2</sub> and CH<sub>4</sub>, and 85 days for N<sub>2</sub>O.

210 All statistical analyses were performed with R (version 4.0.4; R Core Team, 2022). All data  
211 was visually and statistically checked for normality (Levene's test) and homoscedasticity before  
212 testing for statistical differences. Since the original data was not normally distributed, CO<sub>2</sub> and  
213 N<sub>2</sub>O fluxes were log-transformed. ~~One full cycle of all 16 chambers took 160 minutes.~~ To  
214 homogenise the data from the gas flux analysers and the soil temperature and soil moisture  
215 sensors, we rounded all gas flux data to 3-hour intervals (00:00, 03:00, 06:00, 09:00, 12:00,  
216 15:00, 18:00, 21:00), corresponding to the approximate gas flux measurement cycle duration.  
217 Soil temperature and soil moisture data was available every 30 min and was thus also  
218 aggregated for the same 3-hour intervals. For the statistical analyses, we ran linear mixed-effect  
219 models (LMM) using the "lmer" function from the lme4 package (version 1.1-27; Bates et al.,  
220 2015), the "lmerTest" package (version 3.1-3; Kuznetsova et al., 2017), and the "optimx"  
221 function from the optimx package (version 2021-6.12; Nash and Varadhan, 2011). Models were  
222 selected according to the guidelines of Zuur et al. (2009). For the null models, soil temperature,  
223 soil moisture, and inclination or distance from the stream (i.e., 0.5 m to 15 m away from the  
224 stream, CG0.5 – CG15) were included as fixed effects, with an interaction between soil  
225 temperature and soil moisture. Sampling date and chamber number were included as random  
226 effects. Sampling date was included as a random variable since we were not exploring temporal  
227 changes and since there were multiple observations per day. Inclination and distance were not  
228 included in the same model because they were highly correlated. We therefore separated our  
229 treatments in "inclination" and "distance", resulting in two LMM models per GHG. We then  
230 created a model, using the original model structure, including each soil or litter characteristic  
231 individually as an additional explanatory variable. The model Akaike Information Criteria  
232 (AIC)s were then compared using ANOVA. Finally, we selected the model with the lowest AIC  
233 value if it was significantly different from the null model. This was done for each gas-

**Commenté [LG17]:** 5.218: Please write the term abbreviated as "AIC" in full when the abbreviation is used for the first time.

**Response:** Ok, this will be added.

inclination or distance combination. To obtain the conditional and marginal  $R^2$  of the models, the “r2\_nakagawa” function from the performance package was used (version 0.7.3; Nakagawa et al., 2017).

## Results

Over the measurement period (June–November 2020, 161 days), the mean air temperature was 12.30°C and cumulative precipitation was 561 mm. The average volumetric water content, here referred to as ‘soil moisture’, was  $0.22 \pm 0.07 \text{ m}^3 \text{ m}^{-3}$ , with wetter soils in flat ( $0.28 \pm 0.04 \text{ m}^3 \text{ m}^{-3}$ ) compared to sloped locations ( $0.17 \pm 0.02 \text{ m}^3 \text{ m}^{-3}$ ; Fig. S2). The mean soil temperature was  $12.85 \pm 2.62^\circ\text{C}$ , with no significant difference between flat and sloped locations. The wettest and warmest location was at CG5 ( $0.31 \pm 0.03 \text{ m}^3 \text{ m}^{-3}$  and  $13.62 \pm 2.54^\circ\text{C}$ ; Fig. S2). Changes in soil moisture and soil temperature were strongly related to variation of precipitation and air temperature (Fig. S3). Furthermore, the interaction between soil moisture and soil temperature was significant in all models ( $p < 0.001$ ), showing a decrease in soil moisture with increasing soil temperature. Litter depth and weight were much lower at CG0.5 than at all other CGs (Table 1). Soil N and C contents and organic matter content were lowest at CG0.5 and highest at CG10, but C:N ratios were similar at all CGs (Table 1). Bulk density was low ( $0.6\text{--}0.8 \text{ g cm}^{-3}$ ) at all distances. Soil pH was considerably higher at CG0.5 compared to all other CGs (Table 1). The soil in flat locations was sandier, whereas the sloped locations were more clayey (Table 1).

**Table 1:** Average value and standard error of litter and soil parameters at each distance from the stream. “CG” indicates chamber group, with the numbers 0.5, 5, 10, and 15 defining the distance to the stream (m). Different letters indicate differences between distances (Dunn multiple comparison test after Kruskal–Wallis test,  $p < 0.05$ ) for each variable.

Variable	Unit	Chamber group			
		CG0.5	CG5	CG10	CG15

**Commenté [LG18]:** 6.234–239/Tab. 1: Please check this section and the data provided in Tab. 1: According to Tab. 1 litter depth and weight has lowest values at GC5 not at GC0.5. The same is true for soil C and N content. The data shown in Tab. 1 obviously need some corrections: E.g., pH values of 0.65 or 0.34 in soils (as shown for GC5 and GC15, respectively) are extremely unlikely and also C:N ratios of 1.35 or even 0.81 are not really realistic. Most probably, some of the average values and values of standard error have been exchanged in single columns and for some of the parameters.

**Response:** We are very thankful that Reviewer 1 brought this to our attention. There was indeed an issue with the updated table. This has now been corrected which can be found in the Supplement document. The values are now consistent with the observations made in the text.

**Commenté [LG19]:** L238: According to the results in Table 1, the soils at CG0.5 and CG10 are sandier compared to the two locations, CG5 and CG15. And the clay contents of all distances are very low.

**Response:** We are very thankful that Reviewer 2 brought this to our attention. There was indeed an issue with the updated table. This has now been corrected which can be found in the Supplement document. The values are now consistent with the observations made in the text.

**Commenté [LG20]:** 7.172/1. 237/Tab. 1: It is mentioned that the investigated soils have a larger stone content, which is also confirmed by the data in Tab. 1. However, bulk density – explicitly including the coarse soil fraction – with values clearly below  $1.0 \text{ g cm}^{-3}$  is very low and values of  $0.15$  or  $0.12 \text{ g cm}^{-3}$  seem to be unrealistic. The reason for this low bulk density should be explained and data checked (see comment on Tab. 1 below).

**Response:** Thanks for spotting this; there was indeed an issue with the values in Table 1 (see our response to Reviewer 1’s comment 6; the corrected Table 1 can be found in the Supplement document) and the  $0.15$  and  $0.12 \text{ g cm}^{-3}$  were incorrect. This has been corrected in the table at the end of ...

**Commenté [LG21]:** High standard deviations are highly visible in the soil properties of the site, particularly at CG5 and CG15, as presented in Table 1, which indicates high uncertainty and less confidence in the results or the number of samples.

**Response:** There was an issue with the values in Table 1, which has now been corrected and can be found in the Supplement document. In this corrected table the standard deviations are much lower.

**Commenté [LG22]:** L240: Table 1: Litter depth, litter weight, soil C, porosity, organic matter, soil pH, sand content, silt content and clay content at CG5 and CG15 have very high standard deviations indicating high spatial variability and thus uncertainty. First, why such big variability have occurred within such small area? Second, why didn’t you attempt to increase the number of sampling points to reduce the variability? Moreover, in none of the sections of this manuscript have I seen explanations for why these variabilities have occurred.

**Response:** There was an issue when Table 1 values were updated. This has now been corrected and can be found in ...


Litter depth	cm	$4.4 \pm 0.7^a$	$7.0 \pm 1.2^{ab}$	$8.5 \pm 1.0^b$	$8.0 \pm 1.4^b$
Litter weight	g m <sup>-2</sup>	$147.7 \pm 23.1^a$	$311.8 \pm 47.0^{ab}$	$358.5 \pm 100.0^{ab}$	$622.2 \pm 362.1^b$
Soil N content	%	$0.25 \pm 0.06^a$	$0.39 \pm 0.09^{ab}$	$0.6 \pm 0.26^b$	$0.42 \pm 0.18^{ab}$
Soil C content	%	$4.12 \pm 0.78^a$	$6.35 \pm 1.65^{ab}$	$10.15 \pm 4.8^b$	$7.85 \pm 4.29^{ab}$
Soil C:N ratio	-	$16.56 \pm 1.35^a$	$16.24 \pm 0.81^a$	$17.07 \pm 1.81^a$	$18.23 \pm 1.99^a$
Bulk density*	g cm <sup>-3</sup>	$0.81 \pm 0.15^a$	$0.73 \pm 0.12^a$	$0.6 \pm 0.11^a$	$0.81 \pm 0.08^a$
Volumetric stone content	%	$7.59 \pm 8.4^a$	$7.84 \pm 2.57^a$	$10.79 \pm 2.78^a$	$13.16 \pm 2.24^a$
Porosity†	-	$0.75 \pm 0.01^a$	$0.79 \pm 0.03^{ab}$	$0.87 \pm 0.04^b$	$0.80 \pm 0.02^{ab}$
Organic material (OM)	%	$9.25 \pm 1.4^a$	$13.87 \pm 3.73^{ab}$	$20.86 \pm 8.01^b$	$16.70 \pm 7.02^{ab}$
Soil pH	-	$5.57 \pm 0.65^a$	$4.00 \pm 0.34^{ab}$	$4.01 \pm 0.34^{ab}$	$3.78 \pm 0.31^b$
Sand content	%	$598.970 \pm 7.5^a$	$52.0 \pm 9.5^a$	$40.6 \pm 3.7^a$	$41.6 \pm 4.4^a$
Silt content	%	$38.5 \pm 7.7^a$	$45.1 \pm 8.5^a$	$53.1 \pm 4.5^a$	$52.0 \pm 5.0^a$
Clay content	%	$2.5 \pm 0.3^a$	$2.9 \pm 1.4^a$	$6.3 \pm 1.4^b$	$6.5 \pm 0.7^{ab}$

\*with coarse material

†without coarse material

#### Soil CO<sub>2</sub> emissions

The average soil CO<sub>2</sub> emissions during the observation period were  $116.2 \pm 61.5$  mg CO<sub>2</sub>-C m<sup>-2</sup> h<sup>-1</sup>, with flat and sloped locations emitting  $113.6 \pm 66.7$  and  $118.6 \pm 56.3$  mg CO<sub>2</sub>-C m<sup>-2</sup> h<sup>-1</sup>, respectively (Table 2, Fig. 1a). The soil CO<sub>2</sub> emission pattern was bell-curved with increasing distance from the stream, with the lowest emissions at CG0.5, the highest emissions at CG5 and CG10, and relatively low emissions at CG15 as compared to CG10 (Table 2, Fig. 1a). Our analysis showed a significant inclination effect on soil CO<sub>2</sub> emissions ( $p < 0.05$ ); furthermore, we found a significant difference between emissions at CG0.5 and CG5 ( $p < 0.001$ ), as well as between CG0.5 and CG10 ( $p < 0.05$ , Table 2).



**Figure 1:** a.  $\text{CO}_2$  emissions ( $\text{mg CO}_2\text{-C m}^{-2} \text{ h}^{-1}$ ), b.  $\text{CH}_4$  uptake ( $\mu\text{g CH}_4\text{-C m}^{-2} \text{ h}^{-1}$ ), c.  $\text{N}_2\text{O}$  flux ( $\mu\text{g N}_2\text{O-N m}^{-2} \text{ h}^{-1}$ ), and d.  $\text{N}_2\text{O}$  uptake ( $\mu\text{g N}_2\text{O-N m}^{-2} \text{ h}^{-1}$ ) at four distances from a stream: 0.5 m, 5 m, 10 m, and 15 m (i.e., Chamber Groups: CG0.5, CG5, CG10, and CG15). Blue indicates flat locations, and green indicates sloped locations. Statistical significances are from the ‘distance model’ (linear mixed model, LMM) for the differences between the four distances and the ‘inclination model’ for the differences between the flat and slope positions associated with each gas (Table 1, 2, 3); no LMM was run for  $\text{N}_2\text{O}$  uptake. Non-significance is indicated by ‘NS’ and  $p$ -values are coded as  $p < 0.1$  ‘.’,  $p < 0.05$  ‘\*’,  $p < 0.01$  ‘\*\*’, and  $p < 0.001$  ‘\*\*\*’.

**Commenté [LG23]:** L254: It seems average fluxes are reported here, but cumulative fluxes are generally a better approach to compare fluxes of different treatments. Why is average flux preferred over the cumulative flux?

**Response:** We chose to report average fluxes, oppose to cumulative fluxes, due to the data gaps, notably in August, that were not at the same moments between the three GHGs. In addition, we wish to avoid readers who skim the article to assume that cumulative values cover the entire year and comparing it with their or other studies.


**Table 2:** LMM results exploring the relationship between inclination (flat compared to slope) or distance (m), soil moisture ( $\text{m}^3 \text{m}^{-3}$ ), soil temperature ( $^{\circ}\text{C}$ ), soil moisture:soil temperature interaction, soil pH, and volumetric stone content on soil  $\text{CO}_2$  emissions ( $\text{mg CO}_2\text{-C m}^{-2} \text{h}^{-1}$ ). Soil pH and volumetric stone content are included because the LMM models including these variables had AIC values statistically smaller than the null model.  $R^2\text{m}$  indicates marginal  $R^2$ , and  $R^2\text{c}$  indicates conditional  $R^2$  values.  $P$ -values are coded as:  $p < 0.05$  ‘\*’,  $p < 0.01$  ‘\*\*’, and  $p < 0.001$  ‘\*\*\*’.

<b>CO<sub>2</sub> emissions</b>	<b>R<sup>2</sup>c=</b>	0.91	<b>R<sup>2</sup>m=</b>	0.28	<b>AIC=</b>	-9475.99
<b>Inclination – pH</b>	Estimate	Std. Error	df	t value	Pr(> t )	
Soil moisture	-1.48	0.18	11330.00	-8.22	< 2E-16	***
Soil temperature	0.06	4.55E-03	9060.00	14.08	< 2E-16	***
Inclination (slope)	-0.41	0.17	12.20	-2.42	0.03	*
Moisture:temperature	-0.05	0.01	11410.00	-4.35	1.40E-05	***
Soil pH	-0.41	0.12	12.00	-3.33	6.02E-03	**
<b>CO<sub>2</sub> emissions</b>	<b>R<sup>2</sup>c=</b>	0.91	<b>R<sup>2</sup>m=</b>	0.42	<b>AIC=</b>	-9474.05
<b>Distance – stone content</b>	Estimate	Std. Error	df	t value	Pr(> t )	
Soil moisture	-1.49	0.18	11300.00	-8.26	< 2E-16	***
Soil temperature	0.06	4.55E-03	9060.00	14.07	< 2E-16	***
Distance 5 m	0.86	0.16	10.10	5.52	2.49E-04	***
Distance 10 m	0.43	0.16	10.10	2.76	0.02	*
Distance 15 m	0.14	0.16	10.10	0.86	0.41	
Moisture:temperature	-0.05	0.01	11400.00	-4.35	1.39E-05	***
Volumetric stone content	0.02	0.01	10.00	1.76	0.11	

Both model results showed a significant negative correlation between soil  $\text{CO}_2$  emissions and soil moisture ( $p < 0.001$ , Table 2). This pattern was more distinct looking at the CGs at the different distances (Fig. 2a). A significant positive correlation between  $\text{CO}_2$  emissions and soil temperature was found ( $p < 0.001$ , Table 2, Fig. 2b). The interaction between soil moisture and temperature, namely soil moisture decreasing with increasing soil temperature, was shown to correlate negatively with  $\text{CO}_2$  emissions ( $p < 0.001$ , Table 2). According to “inclination” model results,  $\text{CO}_2$  emissions also decreased with increasing soil pH when comparing flat to sloped locations ( $p < 0.01$ , Table 2).

**Commenté [LG24]:** L273-278: There is no need to mention the significance of the main factors (soil moisture and temperature) when the interaction between the two is significant.

**Response:** We humbly disagree, it is very possible for the main factors to not be significant while the interaction between the two is. We therefore chose to leave this information in the text.



**Figure 2:** Relationship between soil CO<sub>2</sub> emissions (mg CO<sub>2</sub>-C m<sup>-2</sup> h<sup>-1</sup>) and **a.** soil moisture (m<sup>3</sup> m<sup>-3</sup>), and **b.** soil temperature (°C) by distance from the stream (0.5 m, 5 m, 10 m, 15 m). Flat locations are indicated in blue (0.5 m and 5 m) and sloped locations in green (10 m and 15 m). The fitted lines show the linear regression on geometrically distributed data using the “geom\_smooth” function (method = “lm”) from ggplot2. The R<sup>2</sup> for these regressions are shown in Table 2.

**Commenté [LG25]:** L282-287: R<sub>2</sub>s in Table 2 represent marginal and conditional R<sub>2</sub> as described by the authors. However, for each regression represented in Figure 2, no R<sub>2</sub> values are shown. The R<sub>2</sub> and P-values should be shown in each figure.

**Response:** We also consider that it would be ideal to include this information in the figures, and we wished to do so. However, the difficulty is making it evident to the reader what the values are referring to. We were unable to find a clear way to indicate what values were from the distance model and which were from inclination model and then the p-values associated to each explanatory variable and interactions. For CO<sub>2</sub> emissions, for example, we would need to include 14 values (4 R<sup>2</sup> values and 10 p-values) plus labelling on the figure, which would make it illegible. Moreover, distributing them amongst the different panels would make it very difficult to understand what they were referring to, and the reader would be dependent on the value labels. Although not ideal, we believe it is much easier for the reader to indicate in the figure legend to find the r<sub>2</sub> and p-values in the tables.

302

303

304 *Soil CH<sub>4</sub> uptake*

305 The soil showed an average CH<sub>4</sub> uptake of  $88.5 \pm 58.0 \mu\text{g CH}_4\text{-C m}^{-2} \text{ h}^{-1}$ , with uptake 180 %

306 higher in sloped as compared to flat locations ( $126.9 \pm 51.3$  and  $45.0 \pm 25.3 \mu\text{g CH}_4\text{-C m}^{-2} \text{ h}^{-1}$ ,

307 respectively; Fig. 1b). Average CH<sub>4</sub> uptake increased by approximately  $40 \mu\text{g CH}_4\text{-C m}^{-2} \text{ h}^{-1}$

308 per 5 m distance further away from the stream (Fig. 1b). However, the “inclination” model

309 showed only marginally significant differences between the CH<sub>4</sub> uptake at flat and sloped

310 locations ( $p < 0.1$ , Table 3). Litter weight was positively correlated with the CH<sub>4</sub> uptake at flat

311 and sloped locations ( $p < 0.001$ ). The “distance” model showed a significant difference between

312 the locations at the stream (CG0.5) and furthest away (CG15;  $p < 0.001$ , Table 3) and a positive

313 correlation between soil C content and CH<sub>4</sub> uptake at all CGs ( $p < 0.01$ , Table 3).

314

315 **Table 3:** LMM results exploring the relationship between inclination (flat compared to slope)

316 or distance (m), soil moisture ( $\text{m}^3 \text{ m}^{-3}$ ), soil temperature ( $^{\circ}\text{C}$ ), soil moisture:soil temperature

317 interaction, litter weight (g), and soil C content effects on soil CH<sub>4</sub> uptake ( $\mu\text{g CH}_4\text{-C m}^{-2} \text{ h}^{-1}$ ).

318 Litter weight and soil C content are included because the LMM models including these

319 variables had AIC values statistically smaller than the null model.  $R^2\text{m}$  indicates marginal  $R^2$ ,

320 and  $R^2\text{c}$  indicates conditional  $R^2$  values.  $P$ -values are coded as:  $p < 0.1$  ‘.’,  $p < 0.05$  ‘\*’,  $p < 0.01$

321 ‘\*\*’, and  $p < 0.001$  ‘\*\*\*’.

<b>CH<sub>4</sub> uptake</b>	$R^2\text{c}=$	0.97	$R^2\text{m}=$	0.67	AIC=	88007.79
<b>Inclination – Litter weight</b>	Estimate	Std. Error	df	t value	Pr(> t )	
Soil moisture	173.06	12.81	11318.95	13.51	< 2E-16	***
Soil temperature	-2.52	0.33	10140.69	-7.71	1.43E-14	***
Inclination (slope)	30.51	15.49	12.11	1.97	0.07	.
Moisture:temperature	-14.73	0.80	11406.27	-18.34	< 2E-16	***
Litter weight	0.80	0.16	12.00	4.92	3.54E-4	***
<b>CH<sub>4</sub> uptake</b>	$R^2\text{c}=$	0.97	$R^2\text{m}=$	0.70	AIC=	87987.56


<b>Distance – Soil C content</b>	Estimate	Std. Error	df	t value	Pr(> t )	
Soil moisture	172.71	12.81	11313.21	13.48	< 2E-16	***
Soil temperature	-2.52	0.33	10139.66	-7.71	1.41E-14	***
Distance 5 m	31.93	18.74	10.02	1.70	0.12	
Distance 10 m	24.10	22.20	10.02	1.09	0.30	
Distance 15 m	93.49	19.82	10.02	4.72	8.15E-04	***
Moisture:temperature	-14.73	0.80	11406.02	-18.34	< 2E-16	***
Soil C content	7.82	2.04	10.00	3.83	3.3E-03	**

Both “inclination” and “distance” model results show a significant, positive correlation between soil moisture and CH<sub>4</sub> uptake ( $p < 0.001$ ), and a significant, negative correlation between soil temperature and CH<sub>4</sub> uptake ( $p < 0.001$ , Table 3). These patterns could, however, not be confirmed visually (Fig. 3). Like for CO<sub>2</sub> emissions, the soil moisture:soil temperature interaction, namely soil moisture decreasing with increasing soil temperature, was significant ( $p < 0.001$ , Table 3). According to the “inclination” model results, litter weight was positively correlated with the CH<sub>4</sub> uptake at flat and sloped locations ( $p < 0.001$ ). The “distance” model showed that higher soil C content resulted in a higher CH<sub>4</sub> uptake at all CGs ( $p < 0.01$ , Table 3).

**Commenté [LG26]:** L311-312: This is because the interaction is significant. If the interaction is significant, it is difficult to separate the variance due to the main effects.

**Response:** Thanks for pointing this out, this caveat is underlined in the discussion (L453-456).





**Figure 3:** Relationship between CH<sub>4</sub> uptake (μg CH<sub>4</sub>-C m<sup>-2</sup> h<sup>-1</sup>) and **a.** soil moisture (m<sup>3</sup> m<sup>-3</sup>), and **b.** soil temperature (°C) by distance from the stream (0.5 m, 5 m, 10 m, 15 m). Flat locations are indicated in blue (0.5 m and 5 m) and sloped locations in green (10 m and 15 m). The fitted lines show the linear regression on geometrically distributed data using the “geom\_smooth” function (method = “lm”) from ggplot2. The R<sup>2</sup> for these regressions are shown in Table 3.

**Commenté [LG27]:** L319-323: Figure 3: Please see the above two comments.

**Response:** Please see the response to the comment about Figure 2.


340 The soil had an average N<sub>2</sub>O ~~flux-emission~~ of  $5.9 \pm 6.3 \mu\text{g N}_2\text{O-N m}^{-2} \text{ h}^{-1}$ , with flat locations  
341 having 120% higher fluxes than sloped ( $8.4 \pm 7.2$  and  $3.8 \pm 4.5 \mu\text{g N}_2\text{O-N m}^{-2} \text{ h}^{-1}$ , respectively;  
342 Fig. 1c). The “inclination” model results showed significantly decreasing N<sub>2</sub>O ~~fluxes-emissions~~  
343 on sloped locations compared to flat locations ( $p < 0.05$ , Table 3). This was supported by the  
344 “distance” model results, with significantly decreasing ~~fluxes-emissions~~ from CG0.5 towards  
345 CG15 (Fig. 1c, Table 4).

346 **Table 4:** LMM results exploring the relationship between inclination (flat compared to sloped)  
347 or distance (m), soil moisture ( $\text{m}^3 \text{ m}^{-3}$ ), soil temperature ( $^{\circ}\text{C}$ ), soil moisture:soil temperature  
348 interaction, and litter depth (cm) on soil N<sub>2</sub>O ~~fluxes-emissions~~ ( $\mu\text{g N}_2\text{O-N m}^{-2} \text{ h}^{-1}$ ). Litter depth  
349 is included because the LMM model including this variable had an AIC value statistically  
350 smaller than the null model.  $R^2\text{m}$  indicates marginal  $R^2$ , and  $R^2\text{c}$  indicates conditional  $R^2$  values.  
351  $P$ -values are coded as:  $p < 0.05$  ‘\*’,  $p < 0.01$  ‘\*\*’, and  $p < 0.001$  ‘\*\*\*’.

<b>N<sub>2</sub>O <del>fluxemissions</del></b>	$R^2\text{c} =$	0.79		$R^2\text{m} =$	0.21	AIC =	4993.94
<b>Inclination</b>	Estimate	Std. Error	df	t value	Pr(> t )		
Soil moisture	7.75	0.62	7660.60	12.46	< 2E-16	***	
Soil temperature	0.16	0.01	3119.98	11.42	< 2E-16	***	
Inclination (slope)	-0.62	0.23	13.61	-2.71	0.02	*	
Moisture:temperature	-0.58	0.04	7445.77	-14.07	< 2E-16	***	
<b>N<sub>2</sub>O <del>fluxemissions</del></b>	$R^2\text{c} =$	0.80		$R^2\text{m} =$	0.39	AIC =	4995.59
<b>Distance – Litter depth</b>	Estimate	Std. Error	df	t value	Pr(> t )		
Soil moisture	7.74	0.62	7650.00	12.45	< 2E-16	***	
Soil temperature	0.16	0.01	3120.00	11.40	< 2E-16	***	
Distance 5 m	-0.82	0.35	10.10	-2.35	0.04	*	
Distance 10 m	-1.51	0.45	10.00	-3.36	7.24E-03	**	
Distance 15 m	-1.81	0.42	10.10	-4.36	1.42E-03	**	
Moisture:temperature	-0.58	0.04	7440.00	-14.04	< 2E-16	***	
Litter depth	0.25	0.09	9.99	2.70	0.02	*	

352 We found significant positive correlations between N<sub>2</sub>O ~~fluxes-emissions~~ and both soil moisture  
353 and soil temperature in both the “inclination” and “distance” model ( $p < 0.001$ , Table 3). The  
354 correlation between N<sub>2</sub>O ~~fluxes-emissions~~ and soil moisture appeared bell-curved at CG5 and  
355

356 CG10 (Fig. 4a). The correlation between N<sub>2</sub>O ~~fluxes-emissions~~ and soil temperature appeared  
357 bell-curved at CG10 (Fig. 4b). As for CO<sub>2</sub> and CH<sub>4</sub> fluxes, the soil moisture:soil temperature  
358 interaction resulted in significantly decreasing N<sub>2</sub>O ~~fluxes-emissions~~ across all CGs and both  
359 the flat and sloped locations. Similar to the “inclination” model results for CH<sub>4</sub> uptake, the N<sub>2</sub>O  
360 “distance” model showed that a higher litter depth resulted in increasing N<sub>2</sub>O ~~fluxes-emissions~~  
361 at all CGs ( $p < 0.05$ ).



**Figure 4:** Relationship between  $\text{N}_2\text{O}$  fluxes ( $\mu\text{g N}_2\text{O-N m}^{-2} \text{ h}^{-1}$ ) and **a.** soil moisture ( $\text{m}^3 \text{ m}^{-3}$ ), and **b.** soil temperature ( $^{\circ}\text{C}$ ) by distance from the stream (0.5 m, 5 m, 10 m, 15 m). Flat locations are indicated in blue (0.5 m and 5 m) and sloped locations in green (10 m and 15 m). The fitted lines show the linear regression on geometrically distributed data using the “geom\_smooth” function (method = “lm”) from ggplot2. The  $R^2$  for these regressions are shown in Table 4.

Over the 85-day measurement period, we detected episodes of N<sub>2</sub>O uptake at eleven chambers. The measured uptake rates averaged  $0.51 \pm 0.48 \mu\text{g N}_2\text{O-N m}^{-2} \text{h}^{-1}$ . N<sub>2</sub>O uptake occurred predominantly in sloped locations (number of observations: 65 sloped, 16 flat), notably at CG15 (50 observations; Fig. 1d), and predominantly later in the measurement period (September to November).

## Discussion

### *Soil CO<sub>2</sub> emissions*

The soil CO<sub>2</sub> emissions estimated in this study are similar to those from studies in nearby forests, with  $115.7 \text{ mg CO}_2\text{-C m}^{-2} \text{h}^{-1}$  and  $113.0 \text{ mg CO}_2\text{-C m}^{-2} \text{h}^{-1}$  emitted in Rosalia (Leitner et al., 2016) and at Schottenwald, near Vienna, respectively (Hahn et al., 2000). The values we measured are only slightly lower than the average soil CO<sub>2</sub> emission from 18 different forest ecosystems amongst Europe (Janssens et al., 2001). However, other studies in comparable beech and spruce stands in France (Epron et al., 1999) and Germany have found values up to 50% lower (Luo et al., 2012). Apart from differences in measurement methods and seasons, it is very likely that most of the differences can be explained by variations in soil moisture (e.g., Hanson et al., 1993) and temperature (e.g., Lloyd and Taylor, 1994), as discussed in the following section.

### *Effect of inclination and distance to a stream on soil CO<sub>2</sub> emissions*

~~Our Model~~ results showed a significant negative effect of inclination, with lower soil CO<sub>2</sub> emissions on sloped locations. ~~This isese results contradict contrary to~~ our first hypothesis, ~~as well as and to the findings of~~ studies from temperate and boreal forests in North America (Creed et al., 2013; Warner et al., 2018), where soil CO<sub>2</sub> emissions were highest in sloped locations compared to ridge and flat locations, ~~while a subtropical forest in Puerto Rico showed only a weak relation between CO<sub>2</sub> fluxes and topographic variation~~ (Quebbeman et al., 2022).

**Commenté [LG28]:** Lastly, the authors compared their findings with earlier studies in the discussion section, which is good despite incomplete year measurement in this study, but the authors should also look at other studies with similar objectives where GHG emissions are investigated with respect to the slope of the land or with reference to streams or rivers.

**Response:** Although we touched on this in the introduction, it is true we could better develop this in the discussion section. We will compare our findings to similar studies investigating these aspects such as:

- Arias-Navarro et al., 2017, Geophys. Res. Biogeosciences
- Davidson et al., 2000, Bioscience
- Hiltbrunner et al., 2012, Glob. Chang. Biol.
- Lamprea Pineda et al., 2021, Biogeosciences
- Quebbeman et al. 2022, Ecosystems
- Warner et al., 2018, Biogeochemistry
- Yu et al., 2008, Glob. Chang. Biol.
- Yu et al., 2021, Sci. Total Environ.

**Commenté [LG29]:** L374-375: According to Figure 1a, the lowest CO<sub>2</sub> emission is at CG0.5 followed by CG15, which is on the sloped location. Therefore, this statement is not true. The CO<sub>2</sub> emission at the flat area is not significantly different from the CG15 and also the major differences between two distances occur within the flat area (CG0.5 and CG5). Thus, the values presented in Figure 1 won't enable us to conclude slope as a factor influencing the CO<sub>2</sub> emission while the most significant difference is observed within the flat locations. Distance can also not be a factor affecting the CO<sub>2</sub> emission.

**Response:** The confusion likely originates from the fact of having two positions on flat locations (CG0.5 and CG5) and two on sloped ones (CG10 and CG15). Overall, the fluxes from flat locations were significantly lower than sloped locations as indicated by the inclination model. We agree with Reviewer 2 that we can soften this statement about the influence of inclination and immediately call attention to the fact that the highest values were at the middle distances as written on line 378.

We propose:  
 “Model results showed a significant negative effect of inclination, with lower soil CO<sub>2</sub> emissions on sloped locations, which contrary to our first hypothesis and to the findings of studies from temperate and boreal forests in North America (Creed et al., 2013; Warner et al., 2018) where soil CO<sub>2</sub> emissions were highest in sloped locations compared to ridge and flat locations. However, our results suggest that higher CO<sub>2</sub> emissions at flat locations were mainly driven by CG5, where we observed the highest CO<sub>2</sub> emissions. Being at the foot of the slope...”

However, our results suggest that higher CO<sub>2</sub> emissions at flat locations were mainly driven by CG5, where we observed the highest CO<sub>2</sub> emissions. Moreover, we found significantly higher emissions at CG5 and CG10 compared to CG0.5, suggesting that not only inclination, but also the distance to the stream had a major impact on CO<sub>2</sub> emissions. The higher CO<sub>2</sub> emissions found at flat locations were most likely driven by CG5, where we recorded the highest CO<sub>2</sub> emissions. Being at the foot hill of the slope, CG5 likely received a large water and nutrient input from the steep slope as compared to the other distances and had optimal conditions for soil microbial activity, thus resulting in peak soil CO<sub>2</sub> emissions from microbial respiration. A soil texture favourable to microbial activity (enough clay to retain moisture and enough sand to allow sufficient volatile substrate and O<sub>2</sub> access) could lead to such a peak, but the clay content was not significantly different between CG0.5, CG5, and CG15 nor was the sand significantly different at any distance. The effect of soil moisture on CO<sub>2</sub> emissions was different across the CGs: at CG10, where we recorded the second-highest emissions, soil moisture was as low as at CG15. It is possible that the high porosity at CG10 enabled an easier diffusion of CO<sub>2</sub> from the soil matrix to the atmosphere. However, even though we found highest emissions at the wettest CG, our overall results showed higher CO<sub>2</sub> emissions with decreasing soil moisture, probably due to the negative correlation between soil moisture and soil temperature. Indeed, the strong interaction between soil moisture and temperature, seen in the model results for all three gases, restricts our ability to draw firm conclusions for these variables individually. Consistent over all CGs, we found that CO<sub>2</sub> emissions increased with increasing soil temperature, in agreement with findings from, e.g., temperate Norway spruce and beech forests in Europe (Epron et al., 1999; Hahn et al., 2000; Buchmann, 2000; Luo et al., 2012), where most temporal variations in the soil CO<sub>2</sub> flux could be explained by soil temperature. The spatial variability of soil moisture and soil temperature itself may be an effect of a different slope, its exposition and the direction from where the rain comes. This influences the amount of rain reaching the soil surface and the evapotranspiration of the forest, which results in a differing water balance. Compared to sites

**Commenté [LG30]:** L380-381: CG5 receiving water from the steep slope cannot favour microbial activity by itself. Is the water carrying nutrients and organic matter? Then, this might lead to changes in the microbial activity. The authors haven't said anything about the water coming from the stream. The plots are located very close to the stream and there is a high possibility that there is an interaction between the stream water and the nearby plots.

**Response:** We disagree that water cannot stimulate microbial activity by itself; several drying and rewetting studies support the influence of water content on microbial activity. We agree though that water accumulation could have also assisted in higher nutrient content, and we will add this to the text:

"Being at the foot hill of the slope, CG5 likely received a larger water and nutrient input from the steep slope as compared to the other distances and had optimal conditions for soil microbial activity"

in North America (Creed et al., 2013; Warner et al., 2018) and Germany (Buchmann, 2000), and considering the exposition of the slope (Finke 2022, personal communication), our site is likely drier.

We suggest that the effect of inclination and distance to the stream were closely interacting with indirect effects on soil properties and resulted in different soil CO<sub>2</sub> emissions than we expected, notably at CG5. For example, CO<sub>2</sub> emissions were significantly lower at CG0.5 than all other CGs, and soil pH was the highest at this distance, probably due to the close proximity to the forest stream with a higher pH value or root-mediated changes in the pH (Hinsinger et al., 2003; Fürst et al., 2021). Higher soil pH (> 5) can increase soil CO<sub>2</sub> fluxes by stimulating autotrophic respiration from living roots and heterotrophic respiration from soil microorganisms (Reth et al., 2005; Aciego Pietri and Brookes, 2008). However, our model results suggest ~~decreasing~~ increasing CO<sub>2</sub> emissions with low soil pH values. We suggest that this is due to the chemistry in the soil, namely the dominating carbonate species (Finke 2022, personal communication). At a low soil pH, carbonic acid (H<sub>2</sub>CO<sub>3</sub>) dominates over carbonate (CO<sub>3</sub><sup>2-</sup>), and carbonic acid might release CO<sub>2</sub>. At high pH, carbonate dominates, which can hinder CO<sub>2</sub> emissions. We encourage researchers to analyse their sites covering a wider range of microbial communities, roots, and soil nutrients, which might give further insight on whether soil pH directly or indirectly influences soil CO<sub>2</sub> emissions on a topological and moisture gradient. Overall, inclination likely had an indirect effect on the CO<sub>2</sub> emissions at our study site through its influence on soil moisture and soil properties at the base of the slope (GC5) where the highest emissions were measured, and distance to the stream, as well as the resulting indirect effect on soil properties, are the main drivers of soil CO<sub>2</sub> emissions at our study site.

#### CH<sub>4</sub> uptake

The soil CH<sub>4</sub> uptake at our site was considerably higher than values reported from other studies in the same forest (Leitner et al., 2016), in forests near Vienna (Hahn et al., 2000), and in

**Commenté [LG31]:** L412-414: This statement contradicts to the model results mention in L410, where decreasing CO<sub>2</sub> was associated with low pH value.

**Response:** We thank Reviewer 2 for pointing this out. The statement on line L410 is incorrect, there is a negative relationship between soil CO<sub>2</sub> emissions and pH, which means higher CO<sub>2</sub> emissions are correlated with lower pH values. This will be corrected. The results are therefore consistent with the statement on L412-414.

**Commenté [LG32]:** L417-419: The results showed the main drivers of the CO<sub>2</sub> emissions are neither the slope nor the distance from the stream. All measured results showed high spatial variability with no particular pattern to slopes or distances of the plots.

**Response:** We understand Reviewer 2's concern that this statement may be too strong for the results shown. We propose: "We conclude that inclination likely had an indirect effect on the CO<sub>2</sub> emissions at our study site through its influence on soil moisture and soil properties at the base of the slope (GC5) where the highest emissions were measured."

Germany (Born et al., 1990; Brumme and Borken, 1999). These differing values support the findings in forest ecosystems across Northern Europe, where temperate forest soils showed CH<sub>4</sub> uptake rates with a widely varying range between 1-165 µg CH<sub>4</sub>-C m<sup>-2</sup> h<sup>-1</sup> (Smith et al., 2000). The uptake on our sloped locations (126.9 ± 51.3 µg CH<sub>4</sub>-C m<sup>-2</sup> h<sup>-1</sup>) falls on the upper end of this range. Different measurement methods, involving the use of manual chambers and gas chromatography in nearby plots (see Leitner et al., 2016) compared to automated chambers and laser-based gas analysers in our study, could explain the dissimilar values obtained in the same forest ecosystem. In addition, the measurement period of this study did not cover the entire year, which may give rise to the differences between this study and previous studies conducted at the same site. As for soil CO<sub>2</sub> emissions, spatial variability resulting from the exposition of the slope, and the differences in soil moisture and soil temperature, might be other reason for our high values. Because the soils at our site are relatively dry, this might have favoured the uptake of soil CH<sub>4</sub>.

#### *Effect of inclination and distance to a stream on soil CH<sub>4</sub> uptake*

Opposite to our second hypothesis, soil CH<sub>4</sub> uptake was not significantly correlated with inclination. This is opposite to the findings of other studies that did find an inclination effect. However, the studies are not in agreement as to where uptake is higher: in a subtropical forest in Puerto Rico, higher CH<sub>4</sub> uptake on ridges was found as compared to in valleys (Quebbeman et al., 2022); in a temperate forest in Maryland, USA, CH<sub>4</sub> uptake was higher in transition zones than uplands, and valley bottoms were occasionally large net sources (Warner et al., 2018); and in a tropical forest in China, hillslopes were found to be hotspots for CH<sub>4</sub> uptake, while the slope foot and groundwater discharge zone contributed less (Yu et al., 2021). ~~However~~ Nonetheless, soil CH<sub>4</sub> uptake was significantly higher at CG15 compared to CG0.5, suggesting that the distance to the stream did have an effect on CH<sub>4</sub> uptake; the two other distances were potentially not far enough from the stream for them to have a significant effect

**Commenté [LG33]:** L428-431: These differences may also arise from the differences in annual climate conditions such as temperature and precipitation. Please keep in mind that this study hasn't completed the full year measurements, which may give rise to the differences between this and previous studies conducted at the same sites. This needs to be explored.

**Response:** We agree with Reviewer 2 that the differences in climate conditions and different time periods measured could be better underlined. We will underline this more in the text.



on the soil moisture, soil temperature, and soil parameters that would lead to an effect on the CH<sub>4</sub> uptake.

With significant positive correlations between both litter weight and soil C content with CH<sub>4</sub> uptake, we suggest that soil C content and litter regulated CH<sub>4</sub> uptake over distance.

In agreement with our findings, Warner et al. (2018) found a higher CH<sub>4</sub> uptake on locations with high C content in a temperate forest landscape in Maryland, USA. Litter can hinder water from precipitation to easily enter into the soil (Walkiewicz et al., 2021). Since there was more litter on sloped than on flat locations, the litter could have stored the rainfall water, thus keeping the mineral soils underneath drier at sloped locations, as has been reported in other studies

(Borken and Beese, 2006; Wang et al., 2013). We therefore suggest that inclination modulated the soil CH<sub>4</sub> uptake through its influence on weight and depth of the litter layer, and that inclination *per se* was not the main driver of CH<sub>4</sub> uptake at our site. Instead, the weight and depth of the litter layer and the soil C content had the largest effect on the CH<sub>4</sub> uptake.

In our study, both models showed higher CH<sub>4</sub> uptake rates with increasing soil moisture and decreasing soil temperature. This does not only contradict findings from other forests (e.g.,

Adamsen and King, 1993; Castro et al., 1995) but cannot be distinguished visually (Fig. 3). It is possible that our models produced ambiguous results ~~of~~for soil moisture and temperature,

because they were unavoidably associated in our studied *in situ* system; both variables are influenced by inclination and distance to a stream concurrently and this thus limits our ability to draw firm conclusions about either variable separately. ~~and~~Running a LMM with one variable or the other did not help resolve this ambiguity. A long-term study in Högwald Forest,

a Germany forest, also found that soil moisture and soil temperature only weakly correlated with CH<sub>4</sub> uptake and were not able to find a suitable empirical model for CH<sub>4</sub> (Luo et al., 2012).

The lack of clear relationships between soil moisture and soil temperature with CH<sub>4</sub> uptake confirms that litter and soil C content were the best predictors of CH<sub>4</sub> uptake at our site.

Soil N<sub>2</sub>O fluxes

**Commenté [LG34]:** L438: Is it really distance that has an effect on CH<sub>4</sub> uptake? Based on table 3, distances of 5 m and 10 m are not significant, even though 15 m shows significance. Soil moisture and temperature seem to be the major factor controlling the CH<sub>4</sub> uptake rate.

**Response:** It is possible that the two other distances were not far enough from the stream for them to have a significant effect, and Figure S2 shows soil moisture and soil texture to be influenced by either inclination and/or distance. We agree though that we can better re-iterate the influences of distance and inclination on soil moisture and soil temperature.

**Commenté [LG35]:** L448-450: In L296, it is mentioned that CH<sub>4</sub> is marginally affected by inclination by referring to Table 3. However, inclination is mentioned here a non-driver of CH<sub>4</sub> uptake. Please be consistent when the results are interpreted.

**Response:** In the results, we report on data and model results, while in the discussion we interpret these results. We do not find that the indicated sentences contradict what is stated in the results but interprets and concludes based on the ensemble of results.

**Commenté [LG36]:** L451-452: High CH<sub>4</sub> uptake was associated with decreasing soil moisture rather than increasing?

**Response:** The model results shown in Table 3 indicate a positive correlation between CH<sub>4</sub> uptake and soil moisture. This result was certainly unexpected for us as well. As we discuss in L451-460, we believe that factors other than soil moisture may have had a strong influence on CH<sub>4</sub> uptake during our study.

**Commenté [LG37]:** L453-456: The model generates what has been given to it. If the data is valid and a correct procedure is followed, the model will produce the right output. Being able to correctly interpret the model result is also critical. Interpreting the main effects separately while the interaction is significant may lead to a wrong conclusion. L458-460: Please see the comment above.

**Response:** Thank you for pointing out this. We believe that the point the referee is underlining is precisely what we are trying to explain in the indicated lines: it is soil moisture and temperature combined that they need to be looked up, because they are unavoidable associated. We will rewrite this sentence in the revised manuscript to clarify the message we are trying to convey.

The soil N<sub>2</sub>O ~~fluxes-emissions~~ from our site were very similar to the rates reported 200 m further upslope from this study (Leitner et al., 2016) and in deciduous forests near Vienna (Pilegaard et al., 2006), with values between 5.4 and 6.4 µg N<sub>2</sub>O-N m<sup>-2</sup> h<sup>-1</sup>, respectively. They are also comparable to the average N<sub>2</sub>O ~~fluxes-emissions~~ from soils in seven European coniferous forests (Pilegaard et al., 2006), but lower than N<sub>2</sub>O ~~flux-emission~~ estimates in forests subjected to high N deposition rates in Europe (Hahn et al., 2000; Luo et al., 2012; Gundersen et al., 2012), suggesting that N deposition was not a significant driver for the N<sub>2</sub>O ~~fluxes-emissions~~ at our study site. In addition to data on low N<sub>2</sub>O emissions, we provide a new dataset from a temperate upland forest soil with reliable N<sub>2</sub>O uptake measurements, highlighting the possibility of upland forest soils acting as N<sub>2</sub>O sink (Wrage et al., 2004; Savage et al., 2014). With the GasFluxTrailer being a robust, state-of-the-art instrument and a total of 7670 N<sub>2</sub>O flux observations, 81 observations indicating uptake, we are confident that the N<sub>2</sub>O uptake we measured is not instrumental noise (see Cowan et al., 2014).

#### *Effect of inclination and distance to a stream on soil N<sub>2</sub>O ~~fluxes-emissions~~*

In agreement with our third hypothesis, N<sub>2</sub>O ~~fluxes-emissions~~ were significantly lower in sloped locations with lower soil moisture content, which was also found by other forest soil studies in France (Vilain et al., 2012), Kenya (Arias-Navarro et al., 2017), Australia (Butterbach-Bahl et al., 2004), and Ecuador (Lamprea Pineda et al., 2021); although, this is opposite to the findings in forests in China (Yu et al., 2021) and in Puerto Rico (Quebbeman et al., 2022). Furthermore, N<sub>2</sub>O ~~fluxes-emissions~~ in flat positions increased with increasing soil temperature. Our findings therefore could support the hypothesis that inclination *per se* influences N<sub>2</sub>O ~~fluxes-emissions~~ from temperate upland forest soils. However, this soil temperature effect should be interpreted with caution considering the concurrent, significant soil moisture:soil temperature interaction, which could influence the significance of individual effects. N<sub>2</sub>O ~~fluxes-emissions~~ further decreased significantly with increasing distance to the stream. The decrease of N<sub>2</sub>O ~~fluxes~~

emissions from CG0.5 to CG15 might also be a consequence of the higher litter depth at these distances. The quantity and quality of the litter input has been shown to influence N<sub>2</sub>O ~~fluxes~~ emissions from forests (Ambus et al., 2006; Pilegaard et al., 2006; Walkiewicz et al., 2021), especially when coniferous needle litter is compared with deciduous leaf litter. Moreover, tree species have been found to exert a strong control on N cycling in forests (Lovett et al., 2004). We suggest that the thick, mostly deciduous leaf litter layer provided a physical barrier that hindered rainfall water to easily reach the soil matrix and thus affected N<sub>2</sub>O ~~fluxes~~ emissions indirectly by reducing soil moisture, which is in line with what we suggested for the CH<sub>4</sub> uptake. Our conclusions, however, are not consistent with a study conducted at another site in Rosalia, where removal of litter led to lowered N<sub>2</sub>O ~~fluxes~~ emissions (Leitner et al., 2016). This site was, however, a pure mature beech stand. Because it is unclear how much of the total soil N<sub>2</sub>O emissions resulted from the litter layer, we suggest that further studies repeat litter removal versus control experiments to quantify the magnitude of N<sub>2</sub>O emissions resulting from litter. We propose that for our site, a large fraction of the N remained stored in the litter layer and was not released as N<sub>2</sub>O (Eickenscheidt and Brumme, 2013).

## Conclusion

With the state-of-the-art technology used in this study, our dataset allows a detailed look at the influence of inclination, distance to a stream, soil moisture, soil temperature, and other soil and litter properties on soil CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O fluxes in a temperate upland forest in Eastern Austria. Our study provides evidence of the complex interactions between inclination and distance to a stream, and the resulting small-scale changes of soil and litter properties within an upland forest ecosystem. We suggest that soil CO<sub>2</sub> emissions were likely indirectly affected by inclination through its influence on soil moisture and soil properties and distance to the stream, as well as the changes in soil properties caused by inclination and distance. Contrary to our expectations, soil CO<sub>2</sub> emissions were lower in sloped locations with where lower soil moisture

content was lower. Our study site was a large CH<sub>4</sub> sink over the whole measurement period, with higher soil CH<sub>4</sub> uptake rates on the locations furthest away from the stream. Because inclination was not significantly related to the uptake of CH<sub>4</sub>, we suggest that it was not a direct driver of CH<sub>4</sub> uptake at our site. Instead of soil moisture, which is commonly cited as the main driver of CH<sub>4</sub> fluxes, we found that soil C content and litter depth and weight were likely the main drivers of CH<sub>4</sub> uptake. Our study showed a clear, significant influence of inclination and distance to the stream on soil N<sub>2</sub>O fluxes-emissions from a temperate upland forest ecosystem, which was to some extent regulated by litter depth. We showed that the impact of inclination and distance to a stream on GHG fluxes is driven by multiple direct and indirect effects, highlighting the need to consider small-scale differences as controlling factor for future GHG flux studies to improve future GHG balance predictions in forest ecosystems.

#### **Statements and Declarations**

#### **Funding**

This work has been financially supported by the Austrian Research Promotion Agency (FFG), project “LTER-CWN: Long-Term Ecosystem Research Infrastructure for Carbon, Water and Nitrogen” (858024), by the Austrian Climate Research Program, 12th Call (KR19AC0K17557, “EXAFOR”) and by the European Commission (H2020 Project eLTER PLUS, Number 871128)

#### **Conflicts of interest/Competing interests**

The authors have no conflicts of interest to declare.

#### **Authors' contributions**

E.D.-P designed the experiment and collected and pre-processed the data.

L.G. and N.T. analysed the data, assisted by D.A, P.F, and E.D-P.

577 L.G. conducted the final statistics and finalised the figures.

578 N.T. and L.G wrote the manuscript with extensive comments from E.D.-P.

579 D.A, P.F, S.G. and S.Z.-B. edited the manuscript.

580 E.D.-P., S.Z.-B., and S.G., were responsible for infrastructure development in the LTER-

581 CWN project.

582 S.Z.-B. and S.G. were responsible for funding acquisition.

583

#### 584 **Availability of data and material**

585 The gas flux data and soil and litter parameter data are stored in the online repository B2Share

586 and can be shared upon request. The meteorological data is open access on the online

587 repository B2Share: <http://doi.org/10.34730/f883fa7ae62648debd6e172448cfbc9b>.

588

#### 589 **Code availability**

590 The code can be provided upon request.

591

592

#### 593 **References**

594 Aciego Pietri, J. C. and Brookes, P. C.: Nitrogen mineralisation along a pH gradient of a silty  
595 loam UK soil, *Soil Biol Biochem*, 40, 797–802, <https://doi.org/10.1016/j.soilbio.2007.10.014>,  
596 2008.

597 Adamsen, A. P. S. and King, G. M.: Methane Consumption in Temperate and Subarctic  
598 Forest Soils: Rates, Vertical Zonation, and Responses to Water and Nitrogen, *Appl Environ*  
599 *Microbiol*, 59, 485–490, <https://doi.org/10.1128/aem.59.2.485-490.1993>, 1993.

600 Ambus, P.: Nitrous oxide production by denitrification and nitrification in temperate forest,  
601 grassland and agricultural soils, *Eur J Soil Sci*, 49, 495–502, <https://doi.org/10.1046/j.1365-2389.1998.4930495.x>, 1998.

603 Ambus, P., Zechmeister-Boltenstern, S., and Butterbach-Bahl, K.: Sources of nitrous oxide  
604 emitted from European forest soils, *Biogeosciences*, 3, 135–145, <https://doi.org/10.5194/bg-3-135-2006>, 2006.

606 Arias-Navarro, C., Díaz-Pinés, E., Klatt, S., Brandt, P., Rufino, M. C., Butterbach-Bahl, K.,  
607 and Verchot, L. V: Spatial variability of soil N<sub>2</sub>O and CO<sub>2</sub> fluxes in different topographic  
608 positions in a tropical montane forest in Kenya, *J Geophys Res Biogeosci*, 122, 514–527,  
609 <https://doi.org/10.1002/2016JG003667>, 2017.

610 Bates, D., Mächler, M., Bolker, B., and Walker, S.: Fitting Linear Mixed-Effects Models  
 611 using lme4, J Stat Softw, 67, 1–48, 2015.  
 612 Borken, W. and Beese, F.: Methane and nitrous oxide fluxes of soils in pure and mixed stands  
 613 of European beech and Norway spruce, Eur J Soil Sci, 57, 617–625,  
 614 <https://doi.org/10.1111/j.1365-2389.2005.00752.x>, 2006.  
 615 Born, M., Dorr, H., and Levin, I.: Methane consumption in aerated soils of the temperate  
 616 zone, Tellus B, 42, 2–8, <https://doi.org/10.1034/j.1600-0889.1990.00002.x>, 1990.  
 617 Brumme, R. and Borken, W.: Site variation in methane oxidation as affected by atmospheric  
 618 deposition and type of temperate forest ecosystem, Global Biogeochem Cycles, 13, 493–501,  
 619 <https://doi.org/10.1029/1998GB900017>, 1999.  
 620 Buchmann, N.: Biotic and abiotic factors controlling soil respiration rates in *Picea abies*  
 621 stands, Soil Biol Biochem, 32, 1625–1635, [https://doi.org/10.1016/S0038-0717\(00\)00077-8](https://doi.org/10.1016/S0038-0717(00)00077-8),  
 622 2000.  
 623 Burt, T. P. and Butcher, D. P.: Topographic controls of soil moisture distributions, Journal of  
 624 Soil Science, 36, 469–486, <https://doi.org/10.1111/j.1365-2389.1985.tb00351.x>, 1985.  
 625 Butler, J., Goetz, H., and Richardson, J. L.: Vegetation and Soil-Landscape Relationships in  
 626 the North Dakota Badlands, American Midland Naturalist, 116, 378,  
 627 <https://doi.org/10.2307/2425746>, 1986.  
 628 Butterbach-Bahl, K., Kock, M., Willibald, G., Hewett, B., Buhagiar, S., Papen, H., and Kiese,  
 629 R.: Temporal variations of fluxes of NO, NO<sub>2</sub>, N<sub>2</sub>O, CO<sub>2</sub>, and CH<sub>4</sub> in a tropical rain forest  
 630 ecosystem, Global Biogeochem Cycles, 18, n/a-n/a, <https://doi.org/10.1029/2004GB002243>,  
 631 2004.  
 632 Butterbach-bahl, K., Kiese, R., and Liu, C.: Chapter eighteen - Measurements of Biosphere–  
 633 Atmosphere Exchange of CH<sub>4</sub> in Terrestrial Ecosystems, in: Methods in Enzymology, vol.  
 634 495, Elsevier Inc., 271–287, [https://doi.org/10.1016/B978-0-12-386905-](https://doi.org/10.1016/B978-0-12-386905-0.00018-8)  
 635 [0.00018-8](https://doi.org/10.1016/B978-0-12-386905-0.00018-8), 2011.  
 636 Butterbach-Bahl, K., Baggs, E. M., Dannenmann, M., Kiese, R., and Zechmeister-  
 637 Boltensern, S.: Nitrous oxide emissions from soils: how well do we understand the processes  
 638 and their controls?, Philosophical Transactions of the Royal Society B: Biological Sciences,  
 639 368, 20130122, <https://doi.org/10.1098/rstb.2013.0122>, 2013.  
 640 Butterbach-Bahl, K., Diaz-Pines, E., and Dannenmann, M.: Soil Trace Gas Emissions and  
 641 Climate Change, in: Global Environmental Change, Springer Netherlands, Dordrecht, 325–  
 642 334, [https://doi.org/10.1007/978-94-007-5784-4\\_4](https://doi.org/10.1007/978-94-007-5784-4_4), 2014.  
 643 Castro, M. S., Steudler, P. A., Melillo, J. M., Aber, J. D., and Bowden, R. D.: Factors  
 644 controlling atmospheric methane consumption by temperate forest soils, Global Biogeochem  
 645 Cycles, 9, 1–10, <https://doi.org/10.1029/94GB02651>, 1995.  
 646 Cowan, N. J., Famulari, D., Levy, P. E., Anderson, M., Reay, D. S., and Skiba, U. M.:  
 647 Investigating uptake of N<sub>2</sub>O in agricultural soils using a high-precision dynamic chamber  
 648 method, Atmos Meas Tech, 7, 4455–4462, <https://doi.org/10.5194/amt-7-4455-2014>, 2014.  
 649 Creed, I. F., Webster, K. L., Braun, G. L., Bourbonnière, R. A., and Beall, F. D.:  
 650 Topographically regulated traps of dissolved organic carbon create hotspots of soil carbon  
 651 dioxide efflux in forests, Biogeochemistry, 112, 149–164, [https://doi.org/10.1007/s10533-](https://doi.org/10.1007/s10533-012-9713-4)  
 652 [012-9713-4](https://doi.org/10.1007/s10533-012-9713-4), 2013.  
 653 Cronan, C. S.: Ecosystem Biogeochemistry, Springer International Publishing, Cham,  
 654 <https://doi.org/10.1007/978-3-319-66444-6>, 2018.  
 655 Davidson, E. A., Keller, M., Erickson, H. E., Verchot, L. V., and Veldkamp, E.: Testing a  
 656 Conceptual Model of Soil Emissions of Nitrous and Nitric Oxides, Bioscience, 50, 667,  
 657 [https://doi.org/10.1641/0006-3568\(2000\)050\[0667:tacmos\]2.0.co;2](https://doi.org/10.1641/0006-3568(2000)050[0667:tacmos]2.0.co;2), 2000.  
 658 Davidson, E. A., Savage, K., Verchot, L. V., and Navarro, R.: Minimizing artifacts and biases  
 659 in chamber-based measurements of soil respiration, Agric For Meteorol, 113, 21–37,  
 660 [https://doi.org/10.1016/S0168-1923\(02\)00100-4](https://doi.org/10.1016/S0168-1923(02)00100-4), 2002.

Davidson, E. A., Samanta, S., Caramori, S. S., and Savage, K.: The Dual Arrhenius and Michaelis-Menten kinetics model for decomposition of soil organic matter at hourly to seasonal time scales, *Glob Chang Biol*, 18, 371–384, <https://doi.org/10.1111/j.1365-2486.2011.02546.x>, 2012.

Davidson, EriC. A., Belk, E., and Boone, R. D.: Soil water content and temperature as independent or confounded factors controlling soil respiration in a temperate mixed hardwood forest, *Glob Chang Biol*, 4, 217–227, <https://doi.org/10.1046/j.1365-2486.1998.00128.x>, 1998.

Diaz-Pines, E. and Gasch, J.: Rosalia Lehrforst Austria - Meteorological Data 2001-2020, <https://doi.org/10.23728/b2share.681966be29a34f3ebc6015ac255ab143>, 2021.

Eickenscheidt, N. and Brumme, R.: Contribution of <sup>15</sup>N-labelled leaf litter to N turnover, nitrous oxide emissions and N sequestration in a beech forest during eleven years, *Plant Soil*, 362, 67–77, <https://doi.org/10.1007/s11104-012-1245-0>, 2013.

Epron, D., Farque, L., Lucot, É., and Badot, P.-M.: Soil CO<sub>2</sub> efflux in a beech forest: dependence on soil temperature and soil water content, *Ann For Sci*, 56, 221–226, <https://doi.org/10.1051/forest:19990304>, 1999.

Fierer, N. and Jackson, R. B.: The diversity and biogeography of soil bacterial communities, *Proc Natl Acad Sci U S A*, 103, 626–631, <https://doi.org/10.1073/pnas.0507535103>, 2006.

Fürst, J., Nachtnebel, H. P., Gasch, J., Nolz, R., Stockinger, M. P., Stumpp, C., and Schulz, K.: Rosalia: an experimental research site to study hydrological processes in a forest catchment, *Earth Syst Sci Data*, 13, 4019–4034, <https://doi.org/10.5194/essd-13-4019-2021>, 2021.

Gundersen, P., Christiansen, J. R., Alberti, G., Brüggemann, N., Castaldi, S., Gasche, R., Kitzler, B., Klemetsson, L., Lobo-do-Vale, R., Moldan, F., Rütting, T., Schleppi, P., Weslien, P., and Zechmeister-Boltenstern, S.: The response of methane and nitrous oxide fluxes to forest change in Europe, *Biogeosciences*, 9, 3999–4012, <https://doi.org/10.5194/bg-9-3999-2012>, 2012.

Hahn, M., Gartner, K., and Zechmeister-Boltenstern, S.: Greenhouse gas emissions (N<sub>2</sub>O, CO<sub>2</sub> and CH<sub>4</sub>) from three forest soils near Vienna (Austria) with different water and nitrogen regimes, *Die Bodenkultur*, 2, 115–125, 2000.

Hairston, A. B. and Grigal, D. F.: Topographic variation in soil water and nitrogen for two forested landforms in Minnesota, USA, *Geoderma*, 64, 125–138, [https://doi.org/10.1016/0016-7061\(94\)90093-0](https://doi.org/10.1016/0016-7061(94)90093-0), 1994.

Hanson, P. J., Wullschlegel, S. D., Bohlman, S. A., and Todd, D. E.: Seasonal and topographic patterns of forest floor CO<sub>2</sub> efflux from an upland oak forest, *Tree Physiol*, 13, 1–15, <https://doi.org/10.1093/treephys/13.1.1>, 1993.

Hanson, R. S. and Hanson, T. E.: Methanotrophic bacteria, *Microbiol Rev*, 60, 439–471, <https://doi.org/10.1128/mr.60.2.439-471.1996>, 1996.

Harris, E., Diaz-Pines, E., Stoll, E., Schlöter, M., Schulz, S., Duffner, C., Li, K., Moore, K. L., Ingrisch, J., Reinthaler, D., Zechmeister-Boltenstern, S., Glatzel, S., Brüggemann, N., and Bahn, M.: Denitrifying pathways dominate nitrous oxide emissions from managed grassland during drought and rewetting, *Sci Adv*, 7, eabb7118, <https://doi.org/10.1126/sciadv.abb7118>, 2021.

Hiltbrunner, D., Zimmermann, S., Karbin, S., Hagedorn, F., and Niklaus, P. A.: Increasing soil methane sink along a 120-year afforestation chronosequence is driven by soil moisture, *Glob Chang Biol*, 18, 3664–3671, <https://doi.org/10.1111/j.1365-2486.2012.02798.x>, 2012.

Hinsinger, P., Plassard, C., Tang, C., and Jaillard, B.: Origins of root-mediated pH changes in the rhizosphere and their responses to environmental constraints: A review, *Plant Soil*, 248, 43–59, <https://doi.org/10.1023/A:1022371130939>, 2003.

710 Hutchinson, G. L. and Mosier, A. R.: Improved Soil Cover Method for Field Measurement of  
 711 Nitrous Oxide Fluxes, *Soil Science Society of America Journal*, 45, 311–316,  
 712 <https://doi.org/10.2136/sssaj1981.03615995004500020017x>, 1981.  
 713 IPCC: IPCC, 2022: Climate Change 2022: Impacts, Adaptation, and Vulnerability.  
 714 Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental  
 715 Panel on Climate Change, Cambridge University Press. Cambridge University Press,  
 716 Cambridge, UK and New York, <https://doi.org/10.1017/9781009325844>, 2022.  
 717 Janssens, I. A., Lankreijer, H., Matteucci, G., Kowalski, A. S., Buchmann, N., Epron, D.,  
 718 Pilegaard, K., Kutsch, W., Longdoz, B., Grünwald, T., Montagnani, L., Dore, S., Rebmann,  
 719 C., Moors, E. J., Grelle, A., Rannik, Ü., Morgenstern, K., Oltchev, S., Clement, R.,  
 720 Guðmundsson, J., Minerbi, S., Berbigier, P., Ibrom, A., Moncrieff, J., Aubinet, M.,  
 721 Bernhofer, C., Jensen, N. O., Vesala, T., Granier, A., Schulze, E.-D., Lindroth, A., Dolman,  
 722 A. J., Jarvis, P. G., Ceulemans, R., and Valentini, R.: Productivity overshadows temperature  
 723 in determining soil and ecosystem respiration across European forests, *Glob Chang Biol*, 7,  
 724 269–278, <https://doi.org/10.1046/j.1365-2486.2001.00412.x>, 2001.  
 725 Kuznetsova, A., Brockhoff, P. B., and Christensen, R. H. B.: lmerTest Package: Tests in  
 726 Linear Mixed Effects Models, *J Stat Softw*, 82, <https://doi.org/10.18637/jss.v082.i13>, 2017.  
 727 Lamprea Pineda, P. A., Bauters, M., Verbeeck, H., Baez, S., Barthel, M., Bodé, S., and  
 728 Boeckx, P.: Ideas and perspectives: Patterns of soil CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O fluxes along an  
 729 altitudinal gradient - A pilot study from an Ecuadorian neotropical montane forest,  
 730 *Biogeosciences*, 18, 413–421, <https://doi.org/10.5194/bg-18-413-2021>, 2021.  
 731 Leitner, S., Sae-Tun, O., Kranzinger, L., Zechmeister-Boltenstern, S., and Zimmermann, M.:  
 732 Contribution of litter layer to soil greenhouse gas emissions in a temperate beech forest, *Plant*  
 733 *Soil*, 403, 455–469, <https://doi.org/10.1007/s11104-015-2771-3>, 2016.  
 734 Lin, H. S., Kogelmann, W., Walker, C., and Bruns, M. A.: Soil moisture patterns in a forested  
 735 catchment: A hydrogeological perspective, *Geoderma*, 131, 345–368,  
 736 <https://doi.org/10.1016/j.geoderma.2005.03.013>, 2006.  
 737 Liu, H., Li, Y., Pan, B., Zheng, X., Yu, J., Ding, H., and Zhang, Y.: Pathways of soil N<sub>2</sub>O  
 738 uptake, consumption, and its driving factors: a review, *Environmental Science and Pollution*  
 739 *Research*, 29, 30850–30864, <https://doi.org/10.1007/s11356-022-18619-y>, 2022.  
 740 Lloyd, J. and Taylor, J. A.: On the Temperature Dependence of Soil Respiration, *Funct Ecol*,  
 741 8, 315, <https://doi.org/10.2307/2389824>, 1994.  
 742 Lookingbill, T. and Urban, D.: An empirical approach towards improved spatial estimates of  
 743 soil moisture for vegetation analysis, *Landsc Ecol*, 19, 417–433,  
 744 <https://doi.org/10.1023/B:LAND.0000030451.29571.8b>, 2004.  
 745 Lovett, G. M., Weathers, K. C., Arthur, M. A., and Schultz, J. C.: Nitrogen cycling in a  
 746 northern hardwood forest: Do species matter?, *Biogeochemistry*, 67, 289–308,  
 747 <https://doi.org/10.1023/B:BIOG.0000015786.65466.f5>, 2004.  
 748 Luo, G. J., Brüggemann, N., Wolf, B., Gasche, R., Grote, R., and Butterbach-Bahl, K.:  
 749 Decadal variability of soil CO<sub>2</sub>, NO, N<sub>2</sub>O, and CH<sub>4</sub> fluxes at the Höglwald Forest, Germany,  
 750 *Biogeosciences*, 9, 1741–1763, <https://doi.org/10.5194/bg-9-1741-2012>, 2012.  
 751 Marrero, T. R. and Mason, E. A.: Gaseous Diffusion Coefficients, *J Phys Chem Ref Data*, 1,  
 752 3–118, <https://doi.org/10.1063/1.3253094>, 1972.  
 753 Le Mer, J. and Roger, P.: Production, oxidation, emission and consumption of methane by  
 754 soils: A review, *Eur J Soil Biol*, 37, 25–50, [https://doi.org/10.1016/S1164-5563\(01\)01067-6](https://doi.org/10.1016/S1164-5563(01)01067-6),  
 755 2001.  
 756 Nakagawa, S., Johnson, P. C. D., and Schielzeth, H.: The coefficient of determination R<sup>2</sup> and  
 757 intra-class correlation coefficient from generalized linear mixed-effects models revisited and  
 758 expanded, *J R Soc Interface*, 14, 20170213, <https://doi.org/10.1098/rsif.2017.0213>, 2017.  
 759 Nash, J. C. and Varadhan, R.: Unifying Optimization Algorithms to Aid Software System  
 760 Users: optimx for R, *J Stat Softw*, 43, <https://doi.org/10.18637/jss.v043.i09>, 2011.



761 ÖNORM: ÖNORM L 1061. Physical determination of grain size distribution of soils less than  
 762 2 mm particle size, Austrian Standards Institute, Vienna, 2002.  
 763 ÖNORM: ÖNORM L 1083. Chemical analyses of soils - Determination of acidity (pH value),  
 764 Austrian Standards Institute, Vienna, 2006.  
 765 ÖNORM: ÖNORM L 1080. Chemical Analyses of Soils—Determination of Organic Carbon  
 766 by Dry Combustion with and without Consideration of Carbonates, Austrian Standards  
 767 Institute, Vienna, 2013.  
 768 Ou, Y., Rousseau, A. N., Wang, L., Yan, B., Gumiere, T., and Zhu, H.: Identification of the  
 769 alteration of riparian wetland on soil properties, enzyme activities and microbial communities  
 770 following extreme flooding, *Geoderma*, 337, 825–833,  
 771 <https://doi.org/10.1016/j.geoderma.2018.10.032>, 2019.  
 772 Pan, Y., Birdsey, R. A., Fang, J., Houghton, R., Kauppi, P. E., Kurz, W. A., Phillips, O. L.,  
 773 Shvidenko, A., Lewis, S. L., Canadell, J. G., Ciais, P., Jackson, R. B., Pacala, S. W.,  
 774 McGuire, A. D., Piao, S., Rautiainen, A., Sitch, S., and Hayes, D.: A Large and Persistent  
 775 Carbon Sink in the World's Forests, *Science* (1979), 333, 988–993,  
 776 <https://doi.org/10.1126/science.1201609>, 2011.  
 777 Parkin, T. B., Venterea, R. T., and Hargreaves, S. K.: Calculating the Detection Limits of  
 778 Chamber-based Soil Greenhouse Gas Flux Measurements, *J Environ Qual*, 41, 705–715,  
 779 <https://doi.org/10.2134/jeq2011.0394>, 2012.  
 780 Pilegaard, K., Skiba, U., Ambus, P., Beier, C., Brüggemann, N., Butterbach-Bahl, K., Dick,  
 781 J., Dorsey, J., Duyzer, J., Gallagher, M., Gasche, R., Horvath, L., Kitzler, B., Leip, A.,  
 782 Pihlatie, M. K., Rosenkranz, P., Seufert, G., Vesala, T., Westrate, H., and Zechmeister-  
 783 Boltenstern, S.: Factors controlling regional differences in forest soil emission of nitrogen  
 784 oxides (NO and N<sub>2</sub>O), *Biogeosciences*, 3, 651–661, <https://doi.org/10.5194/bg-3-651-2006>,  
 785 2006.  
 786 Pumpanen, J., Kolari, P., Ilvesniemi, H., Minkinen, K., Vesala, T., Niinistö, S., Lohila, A.,  
 787 Larmola, T., Morero, M., Pihlatie, M., Janssens, I., Yuste, J. C., Grünzweig, J. M., Reth, S.,  
 788 Subke, J.-A., Savage, K., Kutsch, W., Østreng, G., Ziegler, W., Anthoni, P., Lindroth, A., and  
 789 Hari, P.: Comparison of different chamber techniques for measuring soil CO<sub>2</sub> efflux, *Agric  
 790 For Meteorol*, 123, 159–176, <https://doi.org/10.1016/j.agrformet.2003.12.001>, 2004.  
 791 Quebbeman, A. W., Menge, D. N. L., Zimmerman, J., and Uriarte, M.: Spatial Variation in  
 792 Soil Greenhouse Gas Fluxes in a Subtropical Forest, *Ecosystems*, 25,  
 793 <https://doi.org/10.1007/s10021-021-0067>, 2022.  
 794 Raich, J. W. and Potter, C. S.: Global patterns of carbon dioxide emissions from soils, *Global  
 795 Biogeochem Cycles*, 9, 23–36, <https://doi.org/10.1029/94GB02723>, 1995.  
 796 R Core Team: R: A language and environment for statistical computing. R Foundation for  
 797 Statistical Computing, R Foundation for Statistical Computing, Vienna, Austria, 2022.  
 798 Reth, S., Reichstein, M., and Falge, E.: The effect of soil water content, soil temperature, soil  
 799 pH-value and the root mass on soil CO<sub>2</sub> efflux – A modified model, *Plant Soil*, 268, 21–33,  
 800 <https://doi.org/10.1007/s11104-005-0175-5>, 2005.  
 801 Savage, K., Phillips, R., and Davidson, E.: High temporal frequency measurements of  
 802 greenhouse gas emissions from soils, *Biogeosciences*, 11, 2709–2720,  
 803 <https://doi.org/10.5194/bg-11-2709-2014>, 2014.  
 804 Schad, P.: The International Soil Classification System WRB, Third Edition, 2014, in:  
 805 Springer Water, Springer International Publishing, 563–571, [https://doi.org/10.1007/978-3-319-24409-9\\_25](https://doi.org/10.1007/978-3-319-24409-9_25), 2016.  
 806 Schimel, J. P.: Life in Dry Soils: Effects of drought on soil microbial communities and  
 807 processes, *Annu Rev Ecol Evol Syst*, 49, 409–432, <https://doi.org/10.1146/annurev-ecolsys-110617-062614>, 2018.  
 808 Smith, K. A., Dobbie, K. E., Ball, B. C., Bakken, L. R., Sitaula, B. K., Hansen, S., Brumme,  
 809 R., Borken, W., Christensen, S., Priemé, A., Fowler, D., Macdonald, J. A., Skiba, U.,  
 811

Klemedtsson, L., Kasimir-Klemedtsson, A., Degórska, A., and Orlanski, P.: Oxidation of atmospheric methane in Northern European soils, comparison with other ecosystems, and uncertainties in the global terrestrial sink, *Glob Chang Biol*, 6, 791–803, <https://doi.org/10.1046/j.1365-2486.2000.00356.x>, 2000.

Subke, J.-A., Kutzbach, L., and Risk, D.: Soil Chamber Measurements, in: *Springer Handbook of Atmospheric Measurements*, Springer International Publishing, 1603–1624, [https://doi.org/10.1007/978-3-030-52171-4\\_60](https://doi.org/10.1007/978-3-030-52171-4_60), 2021.

Thomas, P. A. and Packham, J. R.: Ecology of woodlands and forests: Description, dynamics and diversity, 1–528 pp., <https://doi.org/10.1017/CBO9780511805578>, 2007.

Unger, I. M., Motavalli, P. P., and Muzika, R.-M.: Changes in soil chemical properties with flooding: A field laboratory approach, *Agric Ecosyst Environ*, 131, 105–110, <https://doi.org/10.1016/j.agee.2008.09.013>, 2009.

Vilain, G., Garnier, J., Passy, P., Silvestre, M., and Billen, G.: Budget of N<sub>2</sub>O emissions at the watershed scale: role of land cover and topography (the Orgeval basin, France), *Biogeosciences*, 9, 1085–1097, <https://doi.org/10.5194/bg-9-1085-2012>, 2012.

Walkiewicz, A., Rafalska, A., Bulak, P., Bieganski, A., and Osborne, B.: How Can Litter Modify the Fluxes of CO<sub>2</sub> and CH<sub>4</sub> from Forest Soils? A Mini-Review, *Forests*, 12, 1276, <https://doi.org/10.3390/f12091276>, 2021.

Wang, Y., Wang, H., Ma, Z., Dai, X., Wen, X., Liu, Y., and Wang, Z.-L.: The litter layer acts as a moisture-induced bidirectional buffer for atmospheric methane uptake by soil of a subtropical pine plantation, *Soil Biol Biochem*, 66, 45–50, <https://doi.org/10.1016/j.soilbio.2013.06.018>, 2013.

Warner, D. L., Vargas, R., Seyfferth, A., and Inamdar, S.: Transitional slopes act as hotspots of both soil CO<sub>2</sub> emission and CH<sub>4</sub> uptake in a temperate forest landscape, *Biogeochemistry*, 138, 121–135, <https://doi.org/10.1007/s10533-018-0435-0>, 2018.

Webster, K. L., Creed, I. F., Bourbonnière, R. A., and Beall, F. D.: Controls on the heterogeneity of soil respiration in a tolerant hardwood forest, *J Geophys Res*, 113, G03018, <https://doi.org/10.1029/2008JG000706>, 2008.

Wrage, N., Lauf, J., del Prado, A., Pinto, M., Pietrzak, S., Yamulki, S., Oenema, O., and Gebauer, G.: Distinguishing sources of N<sub>2</sub>O in European grasslands by stable isotope analysis, *Rapid Communications in Mass Spectrometry*, 18, 1201–1207, <https://doi.org/10.1002/rcm.1461>, 2004.

Yu, K., Faulkner, S. P., and Baldwin, M. J.: Effect of hydrological conditions on nitrous oxide, methane, and carbon dioxide dynamics in a bottomland hardwood forest and its implication for soil carbon sequestration, *Glob Chang Biol*, 14, 798–812, <https://doi.org/10.1111/j.1365-2486.2008.01545.x>, 2008.

Yu, L., Zhu, J., Ji, H., Bai, X., Lin, Y., Zhang, Y., Sha, L., Liu, Y., Song, Q., Dörsch, P., Mulder, J., and Zhou, W.: Topography-related controls on N<sub>2</sub>O emission and CH<sub>4</sub> uptake in a tropical rainforest catchment, *Science of The Total Environment*, 775, 145616, <https://doi.org/10.1016/j.scitotenv.2021.145616>, 2021.

Zechmeister-Boltenstern, S., Díaz-Pinés, E., Spann, C., Hofmann, K., Schneckner, J., and Reinsch, S.: Soil—The Hidden Part of Climate, in: *Soil and Climate*, CRC Press, Boca Raton, FL : CRC Press, Taylor & Francis Group, 2018. | Series: Advances in soil science, 11–60, <https://doi.org/10.1201/b21225-2>, 2018.

Zuur, A. F., Ieno, E. N., Walker, N., Saveliev, A. A., and Smith, G. M.: Mixed effects models and extensions in ecology with R, Springer New York, New York, NY, <https://doi.org/10.1007/978-0-387-87458-6>, 2009.