

1 Tipping points in ocean and atmosphere circulations

2 Sina Loriani^{1,2}, Yevgeny Aksenov³, David Armstrong McKay^{4,5,6}, Govindasamy Bala⁷, Andreas Born⁸,
3 Cristiano Mazur Chiessi⁹, Henk A. Dijkstra¹⁰, Jonathan F. Donges^{1,2,5}, Sybren Drijfhout^{11,12,13}, Matthew
4 H. England^{14,15}, Alexey V. Fedorov^{16,17}, Laura Jackson¹⁸, Kai Kornhuber^{19,20}, Gabriele Messori^{21,22,23},
5 Francesco S.R. Pausata²⁴, Stefanie Rynders³, Jean-Baptiste Sallée²⁵, Bablu Sinha³, Steven C.
6 Sherwood²⁶, Didier Swingedouw²⁷, Thejna Tharammal²⁸

- 7 1. Earth Resilience Science Unit and Earth System Analysis, Potsdam Institute for Climate Impact Research, Member
8 of the Leibniz Association, Telegrafenberg 31A, 14473 Potsdam
- 9 2. Integrative Earth System Science, Max Planck Institute of Geoanthropology, Jena, Germany
- 10 3. Marine Systems Modelling, National Oceanography Centre, European Way, Southampton, SO14 3ZH, United
11 Kingdom
- 12 4. Global Systems Institute, University of Exeter, Exeter, UK
- 13 5. Stockholm Resilience Centre, Stockholm University, Stockholm, Sweden
- 14 6. Geography, School of Global Studies, University of Sussex, Brighton, UK
- 15 7. Center for Atmospheric and Oceanic Sciences, Indian Institute of Science, Bengaluru-560012, Karnataka, India
- 16 8. Department of Earth Science, University of Bergen and Bjerknes Centre for Climate Research
- 17 9. School of Arts, Sciences and Humanities, University of São Paulo, Av. Arlindo Bettio 1000, 03828-000 São Paulo
18 SP, Brazil
- 19 10. Institute for Marine and Atmospheric research Utrecht, Department of Physics, Utrecht University, Netherlands
- 20 11. Royal Netherlands Meteorological Institute (KNMI), De Bilt, Netherlands
- 21 12. Ocean and Earth Science, University of Southampton, Southampton, UK
- 22 13. Faculty of Science, Utrecht University, Utrecht, Netherlands
- 23 14. Centre for Marine Science and Innovation (CMSI)
- 24 15. ARC Australian Centre for Excellence in Antarctic Science, University of New South Wales, Sydney, NSW,
25 Australia
- 26 16. Dept of Earth and Planetary Sciences, Yale University, New Haven, CT, USA
- 27 17. LOCEAN-IPSL, Sorbonne University, Paris, France
- 28 18. Hadley Centre, Met Office, Fitzroy Road, Exeter, UK
- 29 19. International Institute for Applied Systems Analysis, Laxenburg, Austria
- 30 20. Lamont-Doherty Earth Observatory, Columbia University, New York, USA
- 31 21. Dept. of Earth Sciences, Uppsala University, Uppsala, Sweden
- 32 22. Swedish Centre for Impacts of Climate Extremes (climes), Uppsala University, Uppsala, Sweden
- 33 23. Dept. of Meteorology, Stockholm University, Stockholm, Sweden
- 34 24. Department of Earth and Atmospheric Sciences, University of Quebec in Montreal, Montreal, Quebec, Canada
- 35 25. Sorbonne Université, Laboratoire d'Océanographie et du Climat, CNRS/IRD/MNHN, Paris, France
- 36 26. Climate Change Research Centre, UNSW Sydney, Kensington NSW 2052 Australia
- 37 27. Environnements et Paléoenvironnements Océaniques et Continentaux (EPOC) Univ. Bordeaux, CNRS, Bordeaux
38 INP, EPOC, UMR 5805, F-33600 Pessac, France
- 39 28. Interdisciplinary Centre for Water Research, Indian Institute of Science, Bangalore 560012, India

40 *Correspondence:* Sina Loriani (sina.loriani@pik-potsdam.de)

Abstract. Continued anthropogenic pressures on the Earth system hold the potential to disrupt established circulation patterns in the ocean and atmosphere. In this narrative review, we investigate tipping points in these systems by assessing scientific evidence for feedbacks that may drive self-sustained change beyond critical forcing thresholds, drawing on insights from expert elicitation. The literature provides multiple strands of evidence for oceanic tipping points in the Atlantic Meridional Overturning Circulation (AMOC), the North Atlantic Subpolar Gyre (SPG), and the Antarctic Overturning Circulation, which may collapse under warmer and ‘fresher’ (i.e. less salty) conditions. A slowdown or collapse of these oceanic circulations would have far-reaching consequences for the rest of the climate system and could lead to strong impacts on human societies and the biosphere. Among the atmospheric circulation systems considered, a few lines of evidence suggest the West African monsoon as a tipping system. Its abrupt changes in the past have led to vastly different vegetation states of the Sahara (e.g. “green Sahara” states). Despite multiple potential sources of destabilisation, evidence about tipping of the monsoon systems over South America and Asia is limited. Although theoretically possible, there is currently little indication for tipping points in tropical clouds or mid-latitude atmospheric circulations. Similarly, tipping towards a more extreme or persistent state of the El Niño-Southern Oscillation (ENSO) is currently not fully supported by models and observations. While the tipping thresholds for many of these systems are uncertain, tipping could have severe socio-environmental consequences. Stabilising Earth’s climate (along with minimising other environmental pressures, like aerosol pollution and ecosystem degradation) is critical for reducing the likelihood of reaching tipping points in the ocean-atmosphere system.

1 Introduction

1.1 Tipping Points in circulation systems

The Earth’s oceans and atmosphere shape both the climate and weather of the planet. Human-driven climate change is causing and will continue to cause far-reaching and long-term changes in the circulation of the oceans and atmosphere. The effect of rising greenhouse gas concentrations is to trap additional heat in the Earth system, driving atmospheric and ocean warming (with the latter accounting for more than 90 per cent of the heat trapped so far, IPCC AR6 WG1 Ch9). There is increasing evidence for changes in key circulation patterns, such as a slowing of the Atlantic Meridional Overturning Circulation (AMOC) (Dima and Lohmann 2010; Caesar et al., 2018; Rahmstorf et al. 2015; Zhu et al., 2023), and trends in mid-latitude weather patterns (Faranda et al., 2023; Horton et al., 2015; Coumou et al., 2018). A growing body of research suggests potential tipping points in some of these systems. Tipping points are defined as critical forcing thresholds (e.g. levels of global warming or deforestation), beyond which the system switches to a qualitatively different state due to small perturbations (Lenton et al., 2008; Wang et al., 2023; Matthews et al., 2021). Here we focus on the large sub-class of systems where this change is driven by positive feedbacks in a self-sustained fashion (Armstrong McKay et al. 2022; Lenton et al., 2023). Systems with these features are called *tipping systems* (or *tipping elements* in the

73 context of large-scale Earth system components), and undergo a fundamental state transition at a rate largely determined by
74 the system itself (Armstrong McKay et al., 2022; NRC; 2002). This often occurs abruptly and/or irreversibly, with severe
75 consequences for the Earth and human systems (IPCC AR6 WG1 Ch4). Prominent examples involve potential widespread
76 dieback of the Amazon forest system, collapse of the ice sheets on Greenland and Antarctica, coral bleaching and permafrost
77 melting synchronised over large spatial extents. Similarly, indications for tipping can be found across the circulations in the
78 ocean and atmosphere: Paleoclimate proxy data suggest deep water formation in the North Atlantic has abruptly shifted to a
79 weaker or completely ‘off’ state during previous glacial cycles, with major climatic consequences – a pattern supported by
80 some models (Böhm et al. 2015; IPCC AR6 Ch8, 2021: Ch9 2021). Some model simulations have also shown abrupt
81 changes in the Antarctic deep water formation (Lago and England, 2019; Li et al., 2023). It has also been suggested that the
82 Indian Summer Monsoon could shift to an alternative state as a result of aerosol emissions, counter to the general trend of
83 monsoon strengthening with warming (Levermann et al. 2009; IPCC AR6 WG1 Ch10), as well as potential shifts in
84 circulations on the Southern Hemisphere to El Niño-like mean conditions (Fedorov et al, 2006). Additionally, numerical
85 simulations point to the possibility of tipping behaviour in atmospheric blocking, in the form of a self-sustaining,
86 feedback-driven shift (Drijfhout et al., 2013). Furthermore, there have been discussions on potential large-scale
87 reorganisation of tropical circulation and cloud systems (Schneider et al., 2019, Caballero and Carlson, 2018).

88 1.2 Assessment approach

89 In this review article, we assess the available literature on potential tipping points across the outlined wide range of
90 circulation systems. This work contributes to recent efforts aimed to categorise the breadth of Earth system components with
91 respect to their potential for tipping across the different domains, and has served as the corresponding chapter for the
92 recently published Global Tipping Points Report (Lenton et al., 2023), complementary to chapters on potential tipping points
93 in the cryosphere and biosphere. Our assessment draws on several sources of evidence: Direct observations via in-situ
94 measurements or remote sensing allow for near-time monitoring of the systems, and can provide indications that the system
95 is weakening or even showing early warning signs of approaching a tipping point. However, these observational time series
96 are often (too) short to observe nonlinear behaviour in potential tipping systems. Observational and paleoclimate proxies
97 (i.e. indirect reconstructions) greatly extend data availability to longer time scales and offer historical analogues of tipping
98 dynamics. They are complemented by numerical models of varying complexity, which are used to study system-wide
99 feedbacks in a consistent setup. Concerning oceanic and atmospheric circulations, the different generations of
100 Atmosphere-Ocean General Circulation Models (AOGCMs) up to state-of-the-art coupled Earth System models have been
101 providing great insights into the dynamics of these systems, not least via standardised assessments in the context of the
102 Coupled Model Intercomparison Project (CMIP, Eyring et al., 2016). Eventually, however, these models are limited in
103 resolution and can only resolve potential tipping dynamics if the model captures the respective underlying feedback.
104 Therefore, conceptual and low-order models are indispensable to guide the understanding of potential tipping mechanisms

and, potentially, to identify modelling gaps. In summary, all the outlined lines of evidence complement one another with their respective strengths and drawbacks, and increase confidence in a system's assessment where they converge (Boers et al., 2022).

For our assessment, we therefore draw on a rating system where confidence increases with both robustness (high robustness = multiple, consistent, independent lines of high-quality evidence) and agreement of evidence, as developed and employed in the IPCC context (Mastrandrea et al., 2010). To this end, a system is **classified as tipping system** in this work with

- **high confidence (+++):** Multiple, independent lines of evidence consistently indicate the presence of feedback loops that can drive self-perpetuating change beyond a threshold, leading to a state shift in that system. E.g., there are strong paleo analogues, consistent tipping behaviour in models across the hierarchy, and, if applicable (when the tipping threshold is expected at present-day or soon-to-be-reached forcing levels), proxy and direct observations are compatible with the expected tipping dynamics.
- **medium confidence (++):** Multiple, independent lines of evidence indicate the presence of such feedback loops, however there are uncertainties in timing, magnitude or feedback strength. E.g., there are tipping dynamics in some models, and paleo records hint at dynamics compatible with tipping. Support from observations is limited or contested.
- **low confidence (+):** Singular lines of evidence indicate the presence of such feedback loops, and tipping dynamics only emerge in specific models or under constrained assumptions. E.g., tipping is in principle conceivable via conceptual models, but there are no clear or only weak paleo analogues, and limited demonstration of tipping dynamics in numerical models.

Similarly, we classify a system as **not being a tipping system** if there is evidence indicating the lack of feedback loops that can drive self-perpetuating change beyond a threshold (with low/medium/high confidence). It is **unclear** whether the system is a tipping system if there is conflicting or limited evidence about the existence of such feedback loops. Note that in this approach, irreversibility and abruptness are useful indicators for tipping dynamics, but the defining criterion is a feedback loop beyond a forcing threshold as outlined above, extending the IPCC definition of a tipping point ("*A critical threshold beyond which a system reorganizes, often abruptly and/or irreversibly*"; Matthews et al., 2021) by a mechanistic understanding (see also Armstrong McKay et al., 2022) reflecting the notion of a catastrophic bifurcation with a vanishing stable attractor (Dijkstra, 2013). Furthermore, it is important to emphasise that our assessment reflects the confidence in determining whether a system *can* tip (under present-day or broadly plausible future conditions) — not the confidence that it *will* tip. The latter depends on the trajectory of external forcing and the position of the critical thresholds, both of which are subject to considerable uncertainties.

Although tipping point research is expanding, systematic model intercomparison studies dedicated to tipping dynamics remain scarce—limited so far to isolated efforts like the North Atlantic Hosing Modelling Intercomparison Project, (NAHosMIP; Jackson et al., 2023) or the Ice Sheet Model Intercomparison Project (ISMIP; Nowicki et al., 2016) targeting specific aspects of one system. As a result, current assessments like ours can offer rather qualitative confidence statements

as outlined above, while highlighting the need for future work to establish a more comprehensive quantitative foundation. In the future, emerging initiatives like the Tipping Points Modelling Intercomparison Project (TIPMIP; Winkelmann et al., 2025) could enable confidence ratings based on the fraction of models that exhibit abrupt shifts in setups dedicated to scrutinise tipping dynamics. For now, this review contextualises the current state of science through the lens of self-reinforcing feedbacks beyond critical forcing thresholds, with a focus on consistency and qualitative agreement across lines of evidence.




Figure 1: Potential tipping systems in ocean and atmosphere circulations considered in this review. +/++/+++ denotes that a system is classified as a tipping system with low/medium/high confidence; “?” marks conflicting or limited evidence.


148 2 Ocean Circulation

149 2.1 Atlantic Meridional Overturning Circulation (AMOC)

150 The Atlantic Meridional Overturning Circulation (AMOC) refers to a three-dimensional circulation present in the Atlantic
151 (Figure 2a) whereby warm upper ocean waters ('upper branch') move northward from the tip of Southern Africa to the
152 northern North Atlantic, where they cool, sink and return southwards as cold deep waters ('lower branch'). The AMOC
153 shapes the climate of the whole Earth, influencing, for example, the 1-2°C temperature difference between the Northern and
154 Southern Hemispheres, and the location and strength of rainfall across all tropical regions (Buckley and Marshall, 2016,
155 Feulner et al., 2013, Marshall et al., 2014).

156

157



158 **Figure 2: Overview of the major oceanic circulation systems in the North Atlantic. a) The surface currents (orange pathways) are**
159 **connected to deep ocean currents (blue) through sites where dense (cold, salty) water sinks, driving the overturning circulation**
160 **(pink shaded areas). b) One critical feedback is the salt-advection feedback, in which a weakening of the circulation strength**
161 **reduces the northward advection of salt, which in turn weakens the circulation.**

162 Fresh, warm water is less dense than cold, salty water. In the future, surface waters in the northern North Atlantic will
163 become less dense, because of anthropogenic warming and freshening from ice melt and increased regional precipitation.
164 This will weaken deep water formation in that region, which will disrupt the AMOC, causing it to weaken significantly or
165 even collapse completely. AMOC strength has only been observed directly since 2004 (Srokosz and Bryden, 2013), with
166 more uncertain reconstructions based on observations such as surface temperature, which extend back in time before 2004
167 ('observational proxies'), or from paleoclimate archives such as ocean sediment cores which extend back to prehistoric
168 times ('paleoclimate proxies') (Caesar et al. 2018, 2021, Moffa-Sánchez et al., 2019). The lack of a sufficiently long

169 observational record is a major hurdle for robust understanding of the AMOC, particularly since the North Atlantic is a
170 region of high variability on interannual to decadal timescales (Boer 2000). Current observations show that it is freshening
171 at subpolar latitudes (50-65°N), most strongly in the upper 100m, and warming, most strongly between 100-500 m water
172 depth (IPCC AR6, 2021). Both trends act to reduce AMOC strength. Greenland Ice Sheet melt is accelerating and releasing
173 extra fresh water into the North Atlantic (The IMBIE Team, 2020). In addition, Arctic sea ice is reducing in surface extent
174 and thickness (Serreze and Meier, 2019) and overall Arctic river discharge is increasing (Holmes et al., 2021), adding fresh
175 water to the Arctic which can then leak into the North Atlantic.

176 Limited direct observations of AMOC strength make current trends uncertain, but there are some indirect signs of ongoing
177 weakening. Observational and paleoclimate proxies suggest the AMOC may have weakened by around 15 per cent over the
178 past 50 years (Caesar et al. 2018) and may be at its weakest in 1,000 years (Caesar et al., 2021). However, the proxy data
179 used in these studies have large uncertainties, and agreement amongst reconstructions is contentious (Moffa-Sanchez et al.,
180 2019, Killbourne et al., 2022, Caesar et al., 2022, Terhaar et al., 2025). It remains therefore difficult to confidently discern
181 potential recent trends from natural variability (Bonnet et al. 2021, Latif et al., 2022) since natural variability is still able to
182 explain the trend over the last century. What is clear is that if the observed cooling trend in the subpolar gyre is only due to
183 external forcing, then CMIP6 climate models do not capture correctly such a negative trend in their ensemble mean (cf.
184 Figure 2 of Qasmi, 2022).

185 The IPCC's most recent assessment is that the AMOC has weakened relative to 1850-1900, but with low confidence due to
186 disagreement among reconstructions (Moffa-Sanchez et al., 2019, Killbourne et al., 2022) and models forced response
187 (IPCC AR6 Ch9, 2021). For the future, the IPCC projects that it is very likely that the AMOC will decline in the 21st
188 century (however with low confidence on timings and magnitude) (IPCC AR6 Ch4, 2021). There is medium confidence
189 (about 5 on a scale of 1 to 10) that a collapse would not happen before 2100, though a collapse is judged to be as likely as
190 not by 2300. Hence the possibility of an AMOC collapse within the next century is very much left open by the latest IPCC
191 report.

192 2.1.1 Evidence for tipping dynamics

193 The AMOC has been proposed as a 'global core' climate tipping system with medium confidence by Armstrong McKay et
194 al. (2022). Paleorecords indicate it may have abruptly switched between stronger and weaker modes during recent glacial
195 cycles (Figure 3). Most of the time (including the warm Holocene of the past 12,000 years) the AMOC is in a strong, warm
196 mode, but during peak glacials it sometimes shifted to a weak, cold mode instead (Böhm et al., 2014). It may also
197 occasionally have collapsed entirely to an 'off' mode during some 'Heinrich' events, in which iceberg outbursts from the
198 North American Laurentide Ice Sheet temporarily blocked Atlantic overturning for several centuries (Alley et al., 1999;
199 Rahmstorf et al. 2002; Weijer et al. 2019). While more evidence exists for switches between a warm and cold on-mode in
200 the paleo records, some of these data support switches to an off-mode, possibly occurring through a two-step process from
201 warm-on to cold-on to off (Weijer et al., 2019).

202 In two previous censuses of climate model projections, a shut-down of the AMOC was observed in a small minority of
203 simulations (Drijfhout et al., 2015; Sgubin et al., 2017). This shut-down took many decades and was preceded by decreases

in subpolar surface air and ocean temperatures and increased sea ice cover. Ultimately, deep mixing ceased to occur, cutting the connection between the surface and the deep ocean. Although only a few climate models showed a shutdown of the AMOC, there are concerns that the AMOC may be too stable in CMIP-type climate models (Mecking et al., 2017, Liu et al., 2017), and hence may underestimate the likelihood of AMOC collapse (IPCC AR6 WG1 Ch9).




Figure 3: Different AMOC modes and paleo-evidence. The diagrams on the left show two AMOC modes as indicated by sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ in paleo records. NADW: North Atlantic Deep Water; AABW: Antarctic Bottom Water, adapted from Böhm et al., 2015. B AMOC slowdown events during the last 70,000 years as recorded by sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ data (McManus et al., 2004; Böhm et al., 2015) from the Western North Atlantic (Bermuda Rise, ca. 34°N). Sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ from the Bermuda Rise is a proxy for AMOC strength that assesses the southward flowing North Atlantic Deep Water between ca. 3,500 and 4,500m water depth. The top of the panel marks the timing of past major events of freshwater discharge to the high latitudes of the North Atlantic that decreased AMOC strength (Sarnthein et al., 2003; Carlson et al., 2013; Sanchez Goñi and Harrison, 2009). The red shading highlights past AMOC slowdown events. There is also evidence of AMOC shifts during the last interglacial period, 116,000-128,000 years ago (Galaasen et al. 2014).

Some recent studies have suggested that early warning signals' indicating destabilisation can be detected in reconstructed 'fingerprints' of AMOC strength over the 20th Century (Boers, 2021), and if a tipping point is assumed then the collapse threshold could be reached during the 21st Century (Ditlevsen & Ditlevsen, 2023). These studies used observational proxies for temperature and salinity from the Northeastern subpolar North Atlantic, which are used as indirect AMOC fingerprints rather than direct measurements of AMOC strength. This gives long enough data to analyse for early warning signals, but using indirect proxies adds uncertainty. For instance, the models used by Boers (2021) and Ditlevsen & Ditlevsen (2023) to project collapse are highly simplified, using an indirect 1D metric for the AMOC with a double-well as its stability

landscape assumed, and do not take into account the low-frequency variability of the AMOC. After fitting the timeseries of the metric to various statistical properties of a 1D variable approaching a tipping point (switching from one well to the other in its stability landscape), there are substantial uncertainties with this methodology (see also Michel et al., 2023 highlighting potential false warnings) in particular to determine tipping times (Ben Yami et al. 2024). Despite these caveats, the results amount to a serious warning that the AMOC might be en route to tipping, supported by further potential early warning signals found from analysis of Northern Hemisphere paleoproxies (Michel et al., 2022), and as derived from a physics-based signal (van Westen et al., 2024). However, the claim that we might expect tipping within a few decades is not substantiated enough.

AMOC stability is strongly linked to the ‘salt-advection feedback’ (Stommel, 1961, and see Figure 2b). The AMOC imports salt into the Atlantic and transports it from the South Atlantic to the northern North Atlantic. If the AMOC weakens then less salt is transported to the northern North Atlantic, the surface waters freshen, which inhibits sinking, and the AMOC weakens further. The AMOC collapses seen in models (Drijfhout et al., 2015, Sgubin et al., 2017) were driven by this salt-advection feedback. However, the strength of this feedback, and the timescale over which it operates are governed by processes whose effects are quite uncertain. Although Figure 2a shows typical pathways of surface and deep water through the Atlantic, these pathways are an average picture over many decades. Individual water parcels may get caught up in basin-scale surface or deep recirculations, smaller-scale eddies and meandering currents. There is no definitive evidence though from models or observations that these small-scale processes systematically impact the salt advection feedback.

Additionally, changes in the AMOC have other impacts on salinity – for instance through affecting evaporation and precipitation patterns (Jackson 2013, Weijer et al, 2019). These other feedbacks can temporarily mask, and may even overcome, the salt-advection feedback, potentially changing the stability of the AMOC (Jackson, 2013, Gent, 2018). It is difficult to characterise these processes and feedbacks from observations alone due to insufficient data coverage both in time and space, so we are dependent on numerical models. However, many studies have used reduced complexity models, which may not capture all the potential feedbacks, and even the current generation of climate models have quite low spatial resolution and do not characterise narrow currents, eddies and processes such as horizontal and vertical mixing very well. Indeed, it has been shown recently (Swingedouw et al. 2022) that the inclusion of fine-scale processes of the order of a few kilometres can increase the spread of Greenland meltwater from coastal waters towards the centre of the Labrador Sea, which might more rapidly affect deep convection processes through reduction in surface salinity.

Armstrong McKay et al (2022) estimated with low confidence a global warming threshold for AMOC collapse of $\sim 4^{\circ}\text{C}$ ($1.4\text{--}8^{\circ}\text{C}$). In our view, the range is a better indication of the uncertainty in the different model responses rather than a relationship to global warming, as the likelihood is probably less dependent on temperature, but strongly depends on salinity changes and the strength of opposing feedbacks on the freshwater budget. Studies with climate models have found that adding freshwater can cause the AMOC to collapse and not recover in some models. Since many climate models might be biased towards stability, however, these studies use an unphysically large amount of freshwater to explore the sensitivity (Jackson et al., 2023). Although adding freshwater causes a collapse, they show the threshold is dependent on the strength of

the AMOC and deep convection, rather than on the amount of freshwater added (Jackson and Wood 2018, Jackson et al, 2023). AMOC collapse may also be more sensitive to the rate of freshwater forcing than the total magnitude (Lohmann & Ditlevsen, 2021).




Figure 4: AMOC hysteresis in a CMIP model, adapted from Heinze et al. (2023). CMIP6 overshoot experiments (using UKESM; Jones et al. 2020) showing hysteresis - different states of the AMOC (vertical axis) for the same atmospheric CO2 concentration (horizontal axis). Possible causes are delayed or non-linear response to forcing or possibly bistability of AMOC. The AMOC strength is measured in ‘Sverdrups’ (Sv; i.e. a flow of 1 million cubic metres per second); colours from yellow to blue show model years from 2015 to 2100 respectively.

Hysteresis and bistability both refer to systems which can adopt one of two or more statistically stationary states for the same external forcing, such as CO2 concentration (see Figure 4 and Boucher et al., 2012). Commonly, this is explored by approaching the same external conditions with different trajectories in model simulations, e.g. increasing and reversing the forcing to study reversibility. Bistability involving a full collapse of the AMOC by artificially flooding the North Atlantic with freshwater has been demonstrated (or strongly implied) in theoretical models (Stommel, 1961) and climate models of reduced complexity (Rahmstorf et al., 2005; Hawkins et al., 2011). These types of numerical experiments study bistability through forcings that change slowly enough for the system to equilibrate, typically requiring long simulations and thus coarse model resolution for reasonable computational performance. Alternatively shorter experiments, where forcing is applied temporarily, have been conducted with more complex models. These have demonstrated weak AMOC states which persist for at least 100 years in about half of a test group of CMIP6-type models (Jackson et al., 2023) and in a

276 high-resolution ocean-atmosphere coupled climate model (Mecking et al., 2016). In simpler (coarser resolution models) full
 277 hysteresis experiments have been carried out and those unequivocally support the existence of bi-stability (Weijer et al
 278 2019). Only recently has AMOC tipping (Van Westen et al., 2025) and a bistable regime (Van Westen et al., 2023) been
 279 found in a CMIP-type model in response to gradually changing increasing freshwater release in the North Atlantic. Hence,
 280 AMOC tipping has been found over a whole hierarchy of ocean-climate models, still only excluding very high-resolution
 281 (strongly eddying) climate models, for which appropriate experiments have not yet been conducted.

282 Besides a bi-stable regime bounded by two bifurcation points, a mono-stable off- and on-mode exist on each side of the two
 283 bifurcation points. In present-day climate models through tuning the AMOC always is characterised by the warm on-mode
 284 and either resides in the monostable or bistable regime. Whether or not the AMOC is stable under pre-industrial and
 285 present-day conditions must be model-dependent, depending on details of the background state, probably to a large extent
 286 determined by the freshwater balance in the Atlantic and how northward salinity or freshwater transport by the ocean
 287 circulation balanced by the surface forcing. It is not yet understood why transitions to an off-state or much weakened
 288 AMOC state forced by the same freshwater hosing occurs in some models and not in others (Jackson et al., 2023). However,
 289 as previously mentioned, there is evidence that the present generation of climate models is too stable due to model biases in
 290 the distribution of ocean salinity (Liu et al., 2017, Mecking et al 2017). Not only is it difficult to prove system bistability, the
 291 complexity of the system and interaction with multiple drivers make it hard to assess collapse thresholds. It may be that
 292 realistic freshwater input is not sufficient to cause the transition, or that changing CO₂ alters the underlying system stability,
 293 thus increasing the critical freshwater threshold (Wood et al., 2019). Nevertheless, overshoot scenarios, where the CO₂ trend
 294 is assumed to reverse at some point in the future, provide some useful information about reversibility of the AMOC on
 295 human timescales (e.g. decades to centuries). Figure 4 shows how the AMOC changes in the UKESM climate model under
 296 an overshoot emission scenario exceeding and returning to 500ppm. Even if CO₂ concentrations return to 500ppm by 2100
 297 the AMOC is still only 77 per cent of the strength it was in 2050 also at 500ppm, exhibiting irreversible weakening on these
 298 timescales.

299 The timescale of AMOC tipping was estimated by Armstrong McKay et al (2022) to be 15-300 years, however this range is
 300 very dependent on the strength of the freshwater forcing applied in experiments, which in many cases is unrealistically large
 301 as compared to projected melting of the Greenland Ice Sheet and increase in precipitation and river runoff. Moreover, the
 302 assessment is also potentially impacted by the models being unrealistically stable. With a realistic forcing scenario, the
 303 timescale will depend on the feedbacks. A basin-wide salt advection feedback may have a century timescale, while the same
 304 salt feedback acting on a regional scale (between North Atlantic subtropical and subpolar gyre) is only a few decades. The
 305 latter feedback then may induce fast tipping to a (probably) metastable state that on the longer term is stabilised or
 306 destabilised by the basin-wide Atlantic salt feedback. Even faster timescales associated with tipping to such a hypothetical
 307 metastable state are possible when deep mixing is capped off by sudden increases in sea ice cover, or freshwater halocline.
 308 This fast tipping is then induced by convective feedbacks after which the advective salt feedbacks may further stabilise or
 309 destabilise the convection collapse (Rahmstorf et al., 2001; Kuhlbrodt et al., 2001).

AMOC collapse would lead to cooling over most of the Northern Hemisphere, particularly strong (up to 10°C relative to preindustrial) over Western and Northern Europe. In addition, a southward shift of the intertropical convergence zone would occur, impacting monsoon systems globally and causing large changes in storminess and rainfall patterns (Jackson et al, 2015). A collapse of the AMOC would influence sea level rise along the boundaries of the North Atlantic, modify Arctic sea ice and permafrost distribution (Schwinger et al., 2022; Bulgin et al, 2023), reduce oceanic carbon uptake (Rhein et al., 2016) and potentially lead to ocean deoxygenation (Kwiatkowski et al., 2020) and severe disruption of marine ecosystems (including changes in the North Atlantic Subpolar Gyre, see below), impacting North Atlantic fish stocks (e.g. Heinze et al., 2021).

2.1.2 Assessment and knowledge gaps

Although the AMOC does not always behave like a tipping system in many ocean/climate models, paleoceanographic evidence strongly points to its capability for tipping or at least to shift to another state that can be quasi-stable for many centuries (Figure 3). Tipping is also suggested in a recent study of several CMIP6 models (Jackson et al, 2023) and in another study which found that removing model salinity biases strongly increased the likelihood of tipping (Liu et al., 2017). This does not necessarily mean that tipping is likely in a future climate, since some of these scenarios specified unrealistic inputs of freshwater or GHG emissions. Nonetheless, although the likelihood for collapse is considered small compared to the likelihood of AMOC decline, the potential impacts of AMOC tipping make it an important risk to consider in framing mitigation targets, for instance.

The latest AR6 assessment states that we have only medium confidence that an AMOC collapse will not happen before 2100 (IPCC AR6 WG1 Ch9). This uncertainty is due to models having strong ocean salinity biases, absence of meltwater release from the Greenland Ice Sheet in climate change scenarios, and the possible impact of eddies and other unresolved ocean processes on freshwater pathways. However, a recent study with the PAGES2K database of climate reconstructions of the past 2,000 years suggests, using statistical methods based on dynamical systems theory, that the AMOC variability might be slowing down (Michel et al., 2022), as do the studies of Boers (2021) and Ditlevsen and Ditlevsen (2023). If an AMOC tipping does exist, as suggested in a number of models (e.g. Boulton et al. 2014), this might mean that the AMOC system might be approaching this tipping point. AR6 also concluded that reported recent weakening in both historical model simulations and observation-based reconstructions of the AMOC have low confidence. Direct AMOC observations have not been made for long enough to separate a long-term weakening from short-term variability. Another recent study suggests that we will need to wait until at least 2028 to obtain a robust statistical signal of AMOC weakening (Lobelle et al. 2020). Thus, the coming years will be crucial for detection of an AMOC weakening potentially leading to longer-term instability. There are substantial uncertainties around how the AMOC evolves over long timescales, because of a lack of direct observations. More paleo-reconstructions of AMOC strength, ocean surface temperature, and other AMOC-related properties with high temporal resolution, using appropriate proxies and careful chronological control performed for key past periods (e.g. last millennium, millennial-scale climate change events, previous interglacials), hold great potential to improve

our understanding about the AMOC as a tipping point. Indeed, this might help to test early warning metrics coming from dynamical system theory in some past real-case conditions. These reconstructions might be based on several sources of proxy data to gain robustness (e.g. sortable silt in the deep ocean, $^{231}\text{Pa}/^{230}\text{Th}$ in the deep ocean, radiocarbon in the deep versus surface ocean, surface and intermediate ocean fingerprints). Other open issues are to: (i) reconcile disagreements between paleo-reconstructions and model simulations, and (ii) develop improved fingerprints for creating historical reconstructions and monitoring the AMOC.

Current climate models suffer from imperfect representation of some important processes (such as eddies and mixing) and from biases which can impact the AMOC response to forcings. Hence we need to assess how important these issues are for representing AMOC stability, in particular, to understand how different feedbacks vary across models and are affected by modelling deficiencies. Given these issues, a robust assessment of the likelihood of an AMOC collapse is difficult, but based on the evidence presented, we assess that the AMOC features tipping dynamics with medium confidence. One potential way forward, given these uncertainties, is in developing observable precursors to a collapse that could be monitored.

2.2 North Atlantic Subpolar Gyre (SPG)

The North Atlantic Subpolar Gyre (SPG) is an oceanic cyclonic (counter-clockwise in the northern hemisphere) flow to the south of Greenland (Figure 5), driven by wind and buoyancy forcing. The latter is linked to a site of deep ocean convection in the Labrador-Irminger Seas, i.e. sinking of the subsurface ocean waters to great depths, contributing to the AMOC (Figures 2, 5, 6). Some studies have suggested that the SPG could have a tipping point where the convection collapses, having profound impacts on the SPG system.

There are indications for change in the SPG, as observations show that Labrador Sea Water (LSW) formed during oceanic deep convection events after 2014 was less dense than the LSW formed between 1987 and 1994 (Yashayaev and Loder, 2016), potentially influencing the AMOC. Moreover, the observed ‘warming hole’ over the North Atlantic can be explained by AMOC slowdown (Drijfhout et al 2012, Caesar et al 2018, also see AMOC above) and has also been linked to SPG weakening in CMIP6 models (Sgubin et al 2017, Swingedouw et al 2021). In these models, a collapse of the oceanic convection causes a localised North Atlantic regional surface air temperature drop of $\sim 2\text{--}3^\circ\text{C}$. This cooling moderates warming over north-west Europe and eastern Canada in global warming scenarios, although it is smaller and less widespread than that associated with AMOC collapse.




Figure 5: Map showing the maximum ocean mixing depth in the North Atlantic (light to dark blue), showing deep water convection sites driving the AMOC and SPG east and south of Greenland respectively (with the Labrador-Irminger Seas convection area bordered by yellow). The pale arrows show surface water currents, with the anti-clockwise subpolar gyre occurring within the red bordered area. From Sgubin et al. (2017).

A northward-shift of the atmospheric jet stream, which is predicted to take place with SPG weakening, means more weather extremes in Europe (which may be linked to the unusual cooling and heat waves in recent years) (Osman et al., 2021) and southward shift of the intertropical convergence zone (ITCZ, see Figure 1) (Sgubin et al. 2017; Swingedouw et al. 2021). Models suggest potential impact of the SPG collapse on the European weather, precipitation regime and climate (Swingedouw et al., 2021). This change in the physical system may trigger changes in ecosystems with detrimental consequences for the North Atlantic spring bloom and the overall Atlantic marine primary productivity. Neither of these reverse to the preindustrial state even when the emissions do by the 2100s as the models show (Yool et al., 2015; Heinze et al., 2021). This would impose a strong impact on fisheries and biodiversity (Swingedouw et al., 2021), with expected wide societal implications (e.g., Holm et al., 2022). Last but not least, a transition between two SPG stable states has been suggested to explain the onset of the so-called ‘Little Ice Age’ in which colder conditions prevailed in Europe during the 16th-19th centuries (Lehner et al. 2013; Schleussner et al., 2015; Michel et al. 2022).

Ventilation of Labrador Sea Water (LSW) is accompanied by an uptake of oxygen. Starting in 2014, the convection in the Labrador Sea became more intense and reached depths of 1,500m and below. Consequently, oxygen in LSW is in general increased, but this increase did not penetrate the densest part of this water mass (Rhein et al., 2017). The oxygen concentrations in the deepest part of the LSW (around 2,000m) have decreased in the formation region and along the main export pathways (southward and eastward crossing the Mid-Atlantic Ridge) for more than 20 years. Most of the oxygen

390 from the export of newly formed LSW has been consumed north of the equator (Koelling et al., 2022), and the long-term
 391 oxygen decline along the southward LSW pathway might have impacts on ecosystems in the tropics and subtropics over
 392 longer timescales (e.g., Heinze et al., 2021).


393 The potential shutting-down of winter convection in the Labrador Sea (see Figure 6a,b and Swingedouw et al., 2021) will
 394 also stop the production of Labrador Slope Water (LSLW). This water is next to the Labrador Sea continental slope and is
 395 lighter and less deep than LSW. It contributes to AMOC and the Gulf Stream and can influence variability of the Atlantic
 396 climate system overall (New et al., 2021). The LSLW is rich in nutrients and oxygen too, thereby affecting the ecosystems
 397 on the North American continental shelf and shelf slope (e.g. Claret et al., 2018) and might affect tropical and subtropical
 398 marine ecosystems on a timescale of several decades. Furthermore, the SPG takes up large amounts of atmospheric carbon
 399 and exports it to the deep ocean (Henson et al., 2022).

400 Shallowing of the SPG (Sgubin et al., 2017; Swingedouw et al., 2021) would directly increase regional CO₂ uptake but
 401 negatively impact marine biology, for instance threatening the habitat of cold-water corals in the area due to higher acidity
 402 with more CO₂ dissolved in the water (Fröb et al., 2019; Fontela et al., 2020; Garcia-Ibanez et al., 2021). Weakening or
 403 collapse of the SPG would reduce the amount of carbon-depleted intermediate water being upwelled and newly
 404 carbon-enriched water being convected, reducing export of anthropogenic CO₂ to the deep ocean (Halloran et al. 2015;
 405 Ridge and McKinley 2020), which in turn might lead to an increase of atmospheric CO₂ concentration on the long term
 406 (Schmittner et al. 2007). Declining SPG strength may also be reducing the currently high phytoplankton productivity in this
 407 area (Osman et al., 2019; Henson et al. 2022), reducing the amount of biologically fixed carbon to deeper water too.


408 Changes in the overall Atlantic ocean circulation (AMOC and SPG) can impact the spread of Atlantic water into the Arctic
 409 and affect marine ecosystems there. Summer sea ice decline reduces light limitation, rendering Arctic ecosystems more
 410 similar to the present North Atlantic (Yool et al., 2015). Increased seasonal phytoplankton blooms will deplete nutrients in
 411 the ocean, but increased inputs from rivers and coastal erosion can alleviate this, with Arctic primary production (i.e. the
 412 turnover photosynthesising plankton biomass) projected to increase by about 30-50 per cent in this century. Invasive species
 413 can also extend further into the Arctic habitat due to warming and current changes, e.g. in the Barents Sea and from the
 414 Pacific (Kelly et al., 2020; Neukermans et al., 2018; Oziel et al., 2020; Terhaar et al., 2021).

415 In the North Atlantic, the AMOC can be defined as north-going warm ‘limb’ and saline upper waters and south-going,
 416 colder, denser deep water ‘limb’ (Frajka-Williams et al. 2019). In contrast, in the Subpolar North Atlantic and the SPG, the
 417 AMOC features a third ‘limb’ of a cold, fresh western boundary current with the origin in the Arctic Ocean and Nordic Seas
 418 (Bacon et al., 2022). This is likely linked with the deep convection and winter oceanic mixing in the Labrador, Irminger and
 419 Iceland seas, injecting waters into the deep, southward-flowing limb of the AMOC (Bower et al. 2019). Changes in SPG
 420 circulation are associated with the shallowing of the oceanic mixed layer and convection (Figure 6a,b) in the SPG and link
 421 the predicted future weakening of the North Atlantic subtropical gyre and a strengthening of the Nordic Seas gyre, pointing
 422 to the influences of the upstream changes in the Arctic on the North Atlantic (Swingedouw et al., 2021).


a MLD in winter 2020-30




b MLD change by winter 2040-50



c SST change by summer 2040-50



d Precipitation change by winter 2040-50



423

424 Figure 6: Projected SPG changes. a) Winter ocean mixed layer depth (MLD) as indicator of ocean convection in winter 2020-30
425 (January-March). b) Changes in projected MLD by winter 2040-50 with respect to the 2020-30 baseline. c) Change in summer sea
426 surface temperature (SST) and d) winter total atmospheric precipitation, respectively, projected by winter 2040-50.
427 NEMO-MEDUSA 1/4 degree high resolution model results using ssp370 CMIP6 scenario 2015-2099. High-resolution simulations
428 are courtesy of A. Coward, A. Yool, K. Popova and S. Kelly, NOC. Please see Supplementary Information for more analysis, and
429 Swingedouw et al. (2021) for the IPCC CMIP6 model results.

430 2.2.1 Evidence for tipping dynamics

431 Potential convection instability in the Labrador and Irminger Seas and the wider SPG is believed to be linked to lightening
432 of the upper ocean waters due to reduced salinity (e.g., due to increased precipitation, Figure 6d), thus increasing
433 ‘stratification’ – i.e. reduced mixing between layers of the water column. Warming (Figure 6c) also plays a role and could
434 contribute to convection collapse (Armstrong McKay et al., 2022). Freshening and warming make surface waters more
435 buoyant and thus harder to sink, which, beyond a threshold, can abruptly propel a self-sustained convection collapse
436 (Drijfhout et al. 2015, Sgubin et al., 2017). This process can result in two alternative stable SPG states (Levermann and
437 Born, 2007), with or without deep convection (Armstrong McKay et al., 2022). Similar to the AMOC, SPG stability is also
438 strongly linked to the salt-advection feedback. When the SPG is ‘on’, it brings dense salty waters from the North Atlantic
439 drift into the Irminger and Labrador Seas, allowing deep sinking and convection to occur (Born & Stocker, 2014; Born et al.,
440 2016). When convection decreases due to stratification, the SPG weakens, fresher North Atlantic waters flows eastwards,
441 and the convection is further weakened, which eventually leads to convection collapse in some models. SPG collapse leads
442 to cooling across the SPG region, and so will impact marine biology and bordering regions.

443 A freshwater anomaly is currently building up in the Beaufort gyre – a pile-up of fresh water at the surface of the Beaufort
444 Sea in the Arctic – due to increased freshwater input from rivers, sea ice and snow melting as well as the prevailing
445 clockwise (anticyclonic in the northern hemisphere) winds over the sea (Haine, et al.. 2015; Reagan et al., 2019; Kelly et al.,
446 2020). There is a considerable risk that this freshwater excess might flush into the SPG, disrupting the AMOC (Zhang and
447 Thomas, 2021). The most recent changes in Beaufort gyre size and circulation (Lin et al. 2023) suggest flushing might occur
448 very soon or has already started. The SPG system has recently experienced its largest freshening for the last 120 years in its
449 eastern side due to changes in the atmospheric circulation (Holliday et al. 2020). In contrast, so far there is only limited
450 evidence of Arctic freshwater fluxes impacting freshwater accumulation in the Labrador Sea (Florindo-Lopez et al., 2021).
451 An increased freshwater input into SPG water mass formation regions from melting of Greenland’s glaciers can also inhibit
452 deep water formation and reduce the SPG and AMOC (Dukhovskoy et al., 2021).

453 Although SPG changes are apparently linked to the AMOC, the SPG collapse can occur much faster than AMOC collapse,
454 on the timescale of only a few decades (Armstrong McKay et al., 2022), and independently (Sgubin et al., 2017). Armstrong
455 McKay et al. (2022) estimated global warming threshold of $\sim 1.8^{\circ}\text{C}$ (1.1 to 3.8°C) for the SPG collapse (high confidence)
456 based on climate models from CMIP5 and CMIP6. Abrupt future SPG collapse is diverse in the CMIP6 models, occurring
457 as early as the 2040s (~ 1 to 2°C) but in only a subset of models. However, as these models better represent some key
458 processes, the chance of SPG collapse is estimated at 36-44 per cent (Sgubin et al., 2017; Swingedouw et al., 2021).

459 2.2.2 Assessment and knowledge gaps

460 Similar to Armstrong McKay et al. (2022), the SPG is classified as a tipping system with medium confidence. A global
461 warming threshold for tipping that could be passed within the next few decades, and an estimated tipping timescale of years

462 to a few decades, raise reasons for concern. Furthermore, cessation of deep water production from other sources in the
463 Labrador and Nordic Seas and the Arctic could also present other potential tipping points in the future North Atlantic
464 (Sgubin et al., 2017).

465 **2.3 Southern Ocean circulation**

466 Two main tipping points in the Southern Ocean have been discussed in the past, which both could have large and global
467 climate consequences. The first is the slowdown and collapse of the Antarctic Overturning Circulation; the second is the
468 abrupt change in ocean circulation on the Antarctic continental shelf, leading to suddenly rising ocean temperature in contact
469 with the Antarctic ice shelves fringing the ice sheet.




Figure 7: Circulations and potential tipping systems in the Southern Ocean. Adapted from Li, England et al (2023) and IPCC SROCC Fig CB7.1

Along with the AMOC, the Antarctic overturning circulation constitutes the second branch of the global ocean overturning circulation linking the surface to the abyssal ocean (Figure 7), forming Antarctic Bottom Water (AABW) through sinking of the shelf waters around the Antarctic continent. A key mechanism is brine rejection from sea-ice formation: very salty water that is left behind when ocean water freezes, which causes the ambient liquid water to become heavier and sink. This is maintained by offshore winds blowing away from the Antarctic continent, pushing sea ice away from the coast and forming areas of open water (so-called polynyas) supporting brine rejection. The formation of AABW sustains the operation of the lower branch of the Antarctic overturning circulation (Figure 7 and Abernathey et al., 2016).

In contrast to our understanding of the AMOC, any changes related to the future of the Antarctic Overturning Circulation has remained at low or medium confidence due to a persistent lack of process understanding (Fox-Kemper et al., 2021; Heuzé et al., 2021; Purich and England 2023). However, evidence of its ongoing decline has escalated in recent years, both from observations (Gunn et al., 2023; Zhou et al., 2023; including record low sea ice extent in 2022-2023) and numerical models (Lago & England, 2019; Liu et al. 2022; Li et al, 2023), linked to the changes in melt water, wind trends, sea ice transport and water mass formation (Holland et al., 2012).

Change or collapse in the Antarctic Overturning Circulation has the potential for widespread climate and ecosystem implications within this century. The Southern Ocean surface temperature is set by a delicate balance between ocean overturning strength, upper ocean stratification (the degree of mixing between ocean layers), air sea ice fluxes and sea ice cover. The Antarctic Overturning Circulation affects cloud feedbacks and has been shown to be a key regulator of Earth's

492 global energy balance, so much so that it is the main control on the timing at which the 2°C global warming threshold will
493 be reached for a given emission scenario (Bronselae et al., 2018, Dong et al., 2022; Shin et al., 2023).
494 Reduced Antarctic overturning can also shift global precipitation patterns, resulting in drying of the Southern Hemisphere
495 and wetting of the Northern Hemisphere (Bronselae et al., 2018). Reduced Antarctic overturning also reduces the efficiency
496 of the global ocean carbon sink, leaving more nutrient-rich water at the seafloor (Liu et al., 2022), and also affects global
497 ocean heat storage (Li et al., 2023). Amplifying feedbacks to further shelf water warming and ice melt are also possible
498 (Bronselae et al., 2018; Purich and England 2023; Li et al., 2023).

499 2.3.1 Evidence for tipping dynamics

500 Different generation climate models consistently project a slowing or collapse of the Antarctic overturning under a warming
501 climate (Heuzé et al., 2015, 2021; Lago and England, 2019; Meredith et al., 2019; Fox-Kemper et al., 2021; Liu et al.,
502 2022). However, our confidence in these models to assess change in Antarctic overturning is limited due to known
503 limitations in the representation of dense water formation (Purich and England 2023). Limitations come also from the lack
504 of representation of increased Antarctic ice sheet meltwater in most models (Fox-Kemper et al., 2021). Armstrong McKay et
505 al. (2022) identified the Antarctic Overturning Circulation as a potential but uncertain tipping system in the climate system,
506 but gaps in process understanding meant a threshold remained uncertain. They estimated the Antarctic Overturning
507 Circulation to be prone to collapse at a global warming level of 1.75-3°C based on Lago & England (2019).

508

509 Specifically designed model experiments aiming to bridge some of these limitations, in combination with evidence from
510 observed changes (Gunn et al., 2023; Purkey and Johnson, 2013), suggest that we are currently heading toward a decline and
511 possible collapse of the Antarctic Overturning Circulation (Li et al., 2023, Zhou et al., 2023). The rapidity of this decline
512 might even be underestimated, according to recent observations (Gunn et al., 2023). The sensitivity of the overturning
513 circulation to increases in upper ocean stratification is also consistent with paleo evidence. Observations from marine
514 sediments suggest that AABW formation was vulnerable to freshwater fluxes during past interglacials (Hayes et al., 2014;
515 Huang et al., 2020; Turney et al., 2020) and that AABW formation was strongly reduced (Skinner et al., 2010; Gottschalk et
516 al., 2016; Jaccard et al., 2016) or possibly totally curtailed (Huang et al., 2020) during the Last Glacial Maximum and earlier
517 transient cold intervals.

518 Local water mass characteristics and associated circulation regimes on the Antarctic continental shelf are setting the rate of
519 ice shelf melt rates in ice ‘cavities’, the regions of ocean water covered by floating ice shelves. Relatively warm water
520 reaching the continental shelf in west Antarctica causes high basal melt rates with severe consequences for the ice shelf, ice
521 sheet dynamics, and sea level rise (Naughten et al., 2023). In contrast, the largest ice shelf cavities in the Weddell and Ross
522 Seas are not exposed to this relatively warm water, and consequently have melt rates orders of magnitude smaller than in
523 West Antarctica. Despite this, the Weddell and Ross Sea ice shelf cavities have been shown to exhibit tipping behaviour
524 (Hellmer et al., 2012; 2017; Siahaan et al. 2022). Models show that they are prone to sudden warming of their cavity under

525 future climate change, dramatically increasing basal melting with important consequences for global sea level rise (Hellmer
526 et al., 2012; 2017; Siahhaan et al. 2022). Once tipped into a warm state, such cavities could be irreversibly maintained in such
527 a state, even when forcing is reduced (Hellmer et al., 2017). However, it remains unclear what threshold would need to be
528 crossed to tip those cavities from a cold to warm state, and it may only occur under extreme climate change scenarios.

529 2.3.2 Assessment and knowledge gaps


530 In summary, the combination of process-based understanding and observational, modelling and paleoclimate evidence
531 suggests that the Antarctic Overturning Circulation will continue to decline in the 21st Century. There is increasing evidence
532 for positive amplifying feedback loops that can lead to a collapse of the overturning, with widespread global climate and
533 ecosystem consequences. Closely linked to this is a potential tipping in continental shelf water temperature, driven by
534 amplifying meltwater feedbacks once a regional temperature threshold is crossed. We therefore classify the Southern Ocean
535 Circulation as a tipping system with medium confidence. However, its potential tipping thresholds remain uncertain.

536 3. Atmospheric Circulations: Monsoons

537 Monsoon circulations are large-scale seasonal changes in the direction and strength of prevailing winds over South Asia,
538 East Asia, Africa, Australia and the Americas. Historically, monsoons were seen as large-scale sea breeze circulations driven
539 by land-sea heating differences due to seasonal changes in incoming solar radiation (Figure 8). Currently, a perspective of a
540 global monsoon has emerged (Trenberth et al., 2000; Wang & Ding, 2008), whereby the monsoon systems are seen as
541 interconnected and driven by localised seasonal and more extreme migrations of the ITCZ (Gadgil, 2018; Geen et al., 2020,
542 and references within). Monsoon regions in the world experience heavy precipitation in the summer months, and the global
543 monsoon system is an integral part of the global hydrological cycle, contributing ~31 per cent of total precipitation over the
544 globe (Wang and Ding, 2008).

545

546



547

548 **Figure 8: Monsoon systems.** Shown is the total precipitation difference between Northern hemisphere summer (June-August, JJA)
 549 and winter months (December-February, DJF), highlighting the dominant precipitation patterns over South America (SAM), West
 550 Africa (WAM), India (ISM) and East Asia (EASM). Based on data from ERA5 (Hersbach et al., 2023), with monthly averages over
 551 1980-2010.

552 There is a recent intensification trend in global monsoon precipitation, mainly due to enhanced northern hemisphere summer
 553 monsoon (Wang et al., 2012). It will likely continue in the future (high confidence, IPCC AR6, by $\sim 1\text{-}3\%$ per $^{\circ}\text{C}$ warming)
 554 because of increased water vapour related to warming driven by increased CO_2 in the atmosphere (Hsu et al., 2013; Lee and
 555 Wang, 2014; Chen et al., 2020; Moon and Ha, 2020; Wang et al., 2020), although a few studies conversely show that climate
 556 warming may lead to a weakened global monsoon circulation (Hsu et al., 2012, 2013). Climate simulations also project
 557 expansion of global monsoon domain areas with increasing CO_2 (Wang et al., 2020; Paik et al., 2023) and increased
 558 frequency of monsoon precipitation extremes in the 21st Century (Chevuturi et al., 2018; Ali et al., 2020, Modi and Mishra,
 559 2019; Moon and Ha, 2020; Katzenberger et al., 2021).

560 Monsoon precipitation is vital for agrarian populations and livelihoods in vast areas of South Asia, Africa, and South
 561 America, and changes to it could expose almost two thirds of the global population to disastrous effects (Wang et al., 2021).
 562 Hence it is crucial to understand the dynamics and potential nonlinear changes or tipping behaviour of monsoon systems
 563 under a changing climate. Here the ‘tipping’ of monsoon systems refers to a significant, feedback-driven shift in the
 564 precipitation state of the monsoon, with implications for the regional and global climate and ecosystems. In this discussion
 565 we assess if monsoon systems show any evidence of nonlinear (tipping or abrupt) responses to climate forcings based on
 566 available literature.

567 3.1 Indian summer monsoon (ISM)

568 During the summer season over South Asia (June-September), winds from the south west carry large amounts of water
569 vapour from the Indian Ocean to the Indian subcontinent and cause heavy precipitation in the region, providing ~80 per cent
570 of the total annual precipitation (Figure 8). ISM precipitation shows considerable intra-seasonal, interannual, and decadal
571 variability, many times with precipitation extremes (leading to droughts, floods) during the season, and years and decades
572 with above and below (in drought years) normal precipitation. Indian monsoon variability is strongly influenced by
573 ocean-atmosphere interactions such as El Niño Southern Oscillation (ENSO), Indian Ocean Dipole events (irregular changes
574 in the temperature gradients in the Indian Ocean, Cherchi et al., 2021; Chowdary et al., 2021; Hrudya et al., 2021), and
575 cooler temperatures in the North Atlantic (Borah et al., 2020).

576 ISM precipitation declined in the second half of the 20th Century, attributed mainly to human-driven aerosol loading
577 (Bollasina et al., 2011) and strong Indian Ocean warming (Roxy et al., 2015). Recent studies (Jin and Wang, 2017) suggest
578 the ISM has revived since 2002, linked to enhanced warming over the Indian subcontinent due to reduced low clouds,
579 resulting in an increased land-ocean thermal gradient. Future projections suggest increases in the ISM precipitation in future
580 warming scenarios (by 5.3 per cent per celsius of global warming according to CMIP6 models, Katzenberger et al., 2021)
581 and a longer monsoon duration (Moon and Ha, 2020).

582 3.1.1 Evidence for tipping dynamics

583 Many periods of abrupt ISM transitions have been identified in past monsoon records in association with high-latitude
584 climate events (Schulz et al., 1998; Morrill et al., 2003) such as during Heinrich events (glacial outbursts that temporarily
585 shut down the AMOC) (McManus et al., 2004; Stager et al., 2011), the Younger Dryas (a temporary return to more intense
586 glacial conditions between 12,900–11,700 years ago, Carlson et al., 2013; Cai et al., 2008), and several periods during the
587 more recent Holocene (Gupta et al., 2003; Berkelhammer et al., 2012; Yan and Liu, 2019). However, the mechanisms of
588 such abrupt transitions are not clearly understood. Efforts have been made to identify any Indian monsoon tipping
589 mechanisms using simplified models (Zickfeld et al., 2005; Levermann et al., 2009).

590 An internal feedback mechanism, a “positive moisture advection feedback” (Zickfeld et al., 2005; Levermann et al., 2009;
591 Schewe et al., 2012), has been suggested as responsible for abrupt transitions simulated using these analytical models. In this
592 feedback, the atmospheric temperature gradient between the land and cooler ocean in summer leads to the onshore transport
593 of moist air (advection), which then rises, forms clouds, and condenses into rain. The phase transition from vapour to liquid
594 warms the surrounding air (through the release of latent heat, or ‘diabatic heating’), increasing the land-ocean temperature
595 gradient, and sustaining this monsoon circulation. Any forcing that weakens this pressure gradient can therefore lead to
596 monsoon destabilisation (Zickfeld et al., 2005). If monsoon winds weaken, advection and condensation reduce, and the
597 threshold for a monsoon tipping is reached when the diabatic heating fails to balance the heat advection away from the
598 region (Levermann et al., 2009).

Contrarily, follow-up studies (Boos and Storelvmo, 2016) challenge occurrence of any tipping in these simplified models, and rule out any abrupt monsoon responses to human-driven forcings in the future, and instead attribute past monsoon shifts to rapid forcings or vegetation feedbacks. Simplified models omit key aspects and feedbacks in the monsoon system (specifically, static stability of the troposphere in the models that simulated the monsoon tipping, Boos and Storelvmo, 2016; Kumar and Seshadri, 2023). Hence, more studies using models that represent the complexities of the monsoon and paleoclimate data are required for a clearer picture on any non-linear changes in the monsoon system.

Apart from climate change, aerosols pose another significant human-driven pressure on the Earth system. Aerosols influence the Earth's radiative budget, climate, and hydrological cycle by reflecting or absorbing solar radiation, changing the optical properties of clouds, and also by acting as cloud condensation nuclei. An increase in anthropogenic aerosols has been attributed as the major reason for the decline of Northern Hemispheric summer monsoon strength from the 1950s to 1980s (Cao et al., 2022), due to its dimming effect.

A large increase in regional aerosol loading over South and East Asia (>0.25 Aerosol Optical Depth, AOD, Steffen et al., 2015) could potentially switch the Asian regional monsoon systems to a drier state. Further, hemispheric asymmetries in the aerosol loading (>0.15 AOD Rockström et al., 2023), due to volcanic eruptions, human sources, or intentional geoengineering, could lead to hemispheric temperature asymmetries and changes in the location of the ITCZ, significantly disrupting regional monsoons over West Africa and South Asia (Haywood et al., 2013; Rockström et al., 2023; Richardson et al., 2023). However, there is no direct evidence of aerosols causing a tipping of the monsoon systems, and uncertainties in threshold estimates are large due to complex aerosol microphysics and aerosol-cloud interactions. Hence, systematic observational and modelling approaches would be needed to reduce the uncertainties, as well as additional assessments of interhemispheric asymmetries in the aerosol distribution.

3.1.2 Assessment and knowledge gaps

The Indian summer monsoon system was earlier classified as one of the Earth's tipping systems (Lenton et al., 2008), based on the threshold behaviour of the ISM in the past and the moisture-advection feedback (Levermann et al., 2009), which was refuted by later studies (Boos and Storelvmo, 2016; Seshadri, 2017). Most recently, Armstrong McKay et al. (2022) categorise ISM as an “uncertain potential [climate] tipping element” as global warming is not likely to cause tipping behaviour directly in ISM precipitation.

Based on this current literature, the chances for ISM exhibiting a tipping behaviour towards a new low-precipitation state under climate change are uncertain, warranting extensive studies on the subject. However, potential tipping behaviour in the AMOC (relation to global monsoon described in West African monsoon below) or increase in the interhemispheric asymmetry of aerosol loading in the atmosphere beyond potential threshold levels could lead to large disruptions to monsoon systems. This could cause calamitous effects on millions of people in the monsoon regions, even in the absence of tipping.

631 3.2 West African monsoon (WAM)

632 The West African monsoon (WAM) controls hydroclimatic conditions, vegetation, and mineral-dust emissions of northern
633 tropical and subtropical Africa, up to the semi-arid Sahel region at the southern edge of the Sahara Desert (Figure 8). The
634 strength of the monsoon shows large variations over a range of timescales from interannual to decadal and longer. Albedo
635 (reflectivity of the Earth's surface) changes caused by human-driven land-cover changes and desertification (Charney et al.,
636 1975; Charney, 1975; Otterman, 1974) can affect rainfall: a less vegetated surface with higher albedo increases radiative
637 loss, thereby reducing temperature and suppressing the rising and condensation of moist air into rainfall (i.e. convective
638 precipitation). Variations of sea surface temperatures (SSTs) in different oceanic basins can also drive interannual and
639 decadal variability in WAM precipitation (Rodríguez-Fonseca et al., 2015). Other major factors that affect WAM variability
640 are land surface variability such as variations in soil moisture (Giannini et al., 2013; Zeng et al., 1999), vegetation (Charney
641 et al., 1975; Kucharski et al., 2013; Otterman, 1974; Wang et al., 2004; Xue, 1997), high-latitude cooling (Collins et al.,
642 2017), and dust emissions (Konare et al., 2008; Solomon et al., 2008; Zhao et al., 2011). Charney et al. (1975) proposed the
643 vegetation-climate positive feedback mechanism to explain the decadal precipitation variability in the WAM. The feedback
644 mechanism is explained as desertification leads to increased albedo, surface cooling, and increased stability in the
645 atmosphere, hence, reduced convection and precipitation, thereby amplifying the initial response. It has since been
646 suggested that additional factors such as evapotranspiration, dust emissions, and water vapour recycling related to the
647 vegetation further contribute to the vegetation-climate feedback (Rachmayani et al., 2015; Pausata et al., 2016; Yu et al.,
648 2017; Chandan and Peltier, 2020).

649 3.2.1 Evidence for tipping dynamics

650 Paleoclimate records underscore dramatic variations of the WAM in the more distant past, such as the periodic expansion of
651 vegetation into the Sahara Desert during the so-called 'African humid periods' (AHPs) and are linked to the emergence of
652 ancient cultures along the Nile. The vegetation-climate feedback likely played a significant role in amplifying and sustaining
653 the wetter conditions and facilitating a green Sahara during AHPs (Pausata et al., 2016). Another example is the drought
654 200–300 years ago, which caused the water level of Lake Bosumtwi in Ghana to fall by almost four times as much as it did
655 during the drought of the 1970s and 1980s. Large past variations of the WAM, such as those during the AHPs, raise the
656 question of whether present day anthropogenic global warming could have potentially significant impacts on the WAM.
657 Although the nature and magnitude of radiative forcing were different during the AHPs than they are now (i.e. an external
658 change in insolation due to orbital forcing versus an internal change from increased greenhouse gases), the fact that the
659 AHPs occurred under a globally warmer climate than the pre-industrial period invites questions.
660 Some paleoclimate archives show WAM precipitation changes that took place over several centuries (deMenocal et al.,
661 2000; McGee et al., 2013), i.e. an order of magnitude faster than the orbital forcing. However, others show a much more
662 gradual change (e.g. Kröpelin et al. 2008) with a time-varying withdrawal of the WAM from North to South following the

insolation changes (Shanahan et al., 2015). Because of geographic variability of the African landscape and African monsoon circulation, abrupt changes can occur in several, but not all, regions at different times during the transition from the humid to arid climate (Dallmeyer et al., 2021).

By inducing latitudinal movements of the ITCZ, change in the Atlantic Meridional Overturning Circulation (AMOC) is considered to play a role in shifts of global monsoon systems (Stager et al., 2011). Paleoclimate evidence suggests that glacial meltwater-induced weakening of the AMOC during Heinrich events in the last glacial period led to abrupt Asian and African monsoon weakening (Mohtadi et al., 2014; 2016). Similarly, the Younger Dryas led to a cool and dry state over Northern Hemisphere tropical monsoon regions, as well as the 8.2-kyr event during the Holocene (Alley and Ágústsdóttir, 2005). North Atlantic fresh water-hosing simulations using climate models (Lewis et al., 2010; Pausata et al., 2011; Kageyama et al., 2013) confirm these shifts in ITCZ can occur as a result of substantial glacial meltwater release. These influences of AMOC on the monsoon systems have also been studied in the context of the South American monsoon (see below). Hence, a collapse of AMOC has the potential to cause disruptions to the regional monsoon systems and other tropical precipitation systems over Asia, Africa, and South America (Gupta et al., 2003; IPCCAR6; Ben-Yami et al., 2024). Additionally, location and intensity of the tropospheric jet streams have been identified as processes modulating the strength and position of the African rainbelt (e.g., Farnsworth et al., 2011, Nicholson and Dezfuli, 2013).

3.2.2 Assessment and knowledge gaps

Abrupt changes in one region can be induced by abrupt changes in other regions, a process sometimes referred to as “induced tipping” (Pausata et al., 2020). The AHP transition of the Sahara was slow with respect to timescales of individual humans and local ecosystems, but regionally rapid with respect to changes in the driver. Based on the record of large past variations of WAM precipitation patterns (including collapse), and the existence of positive amplifying feedbacks, we classify WAM as a tipping system with low confidence. This is in line with previous assessments (Armstrong McKay et al., 2022), in which a lower tipping threshold of 2°C global warming was estimated but attributed low confidence due to limited model resolution of vegetation shifts, and model disagreements in future trends. The timescale of abrupt shifts is estimated to range from decades as observed in CMIP5 models (Drijfhout et al., 2015) to centuries based on paleorecords (Hopcroft and Valdes, 2021; Shanahan et al., 2015). Potential additional destabilisation through AMOC weakening and atmospheric aerosol loading, and the far-reaching implications of WAM tipping call for intensified research efforts on this system.

3.3 South American Monsoon (SAM)

The South American Monsoon (SAM) system is characterised by strong seasonality in precipitation, even though it does not show a reversal of low-level winds like the Asian monsoon (Zhou and Lau, 1998; Vera et al., 2006; Liebmann and Mechoso, 2011; Carvalho et al., 2012). Studies on the SAM system are relatively few compared to the Asian and African monsoon systems, as it was not classified as a monsoon system until a couple of decades ago (Zhou and Lau, 1998).

694 A mature SAM system (from December to February) shows features such as enhanced northeastern trade winds, increased
695 land-ocean thermal gradient, and the development of an active convective zone (the South Atlantic Convergence Zone)
696 (Figure 8; Zhou and Lau, 1998). The SAM system affects vast areas of tropical South America all the way to southern
697 Brazil, and provides more than 50 % of the annual precipitation to these regions (Vera et al., 2006) including most of the
698 Amazon rainforest. SAM precipitation varies from interannual to orbital timescales (Chiessi et al., 2009; Liebmann and
699 Mechoso, 2011; Carvalho and Cavalcanti, 2016; Hou et al., 2020).

700 The influence of anthropogenic climate change on the SAM precipitation is ambiguous (Douville et al., 2021), and many
701 CMIP5/CMIP6 models are noted for their poor representation of SAM precipitation (Jones and Carvalho, 2013; Douville et
702 al., 2021). IPCC AR6 finds high confidence in delayed onset of the SAM precipitation since the 1970s associated with
703 climate change, which could worsen with increased CO₂ levels (Douville et al., 2021). However, the projected future change
704 in total SAM precipitation is uncertain, as the models show low agreement on the projections (Douville et al., 2021).

705 3.3.1 Evidence for tipping dynamics

706 Orbital timescale changes (i.e. over 10s of 1000s of years) in SAM precipitation seem to be largely controlled by changes in
707 insolation and respond linearly to it (Cruz et al., 2005; Hou et al., 2020). Millennial-scale changes (i.e. over 1000s years) in
708 the SAM are thought to be associated with variations in the strength of the AMOC as described for the West African
709 monsoon (see above). In particular, paleo evidence indicates that an increase in tropical South American precipitation to the
710 south of the equator followed a weakening of the AMOC related to Heinrich stadials (Mulitza et al., 2017; Zhang et al.,
711 2017; Campos et al., 2019). Similarly, meltwater flux from the Laurentide Ice Sheet during the Younger Dryas probably led
712 to wet anomalies over tropical South America to the south of the equator (McManus et al., 2004; Broecker et al., 2010;
713 Venancio et al., 2020; Brovkin et al., 2021). Earth system model projections of AMOC collapse impacts on tropical South
714 American rainfall are model dependent, but generally find a reduction in rainfall over northernmost South America and an
715 increase over tropical South America to the south of the equator (Bellomo et al., 2023; Nian et al., 2023; Orihuela-Pinto et
716 al., 2022a; Liu et al., 2020).

717 Deforestation of the Amazon rainforest may force a tipping dynamic in the SAM system. In a non-linear model for moisture
718 transport, deforestation over 30-50 % of the Amazon rainforest led to a tipping point in the SAM system, causing
719 precipitation reductions of up to 40 % in non-forested parts of the western Amazon (Boers et al., 2017). This reduction in
720 SAM precipitation is caused by the breakdown of a positive amplifying feedback mechanism present in Amazonian
721 precipitation that involves latent heat of condensation over the Amazon rainforest due to transpiration (i.e. water lost from
722 plants) and water vapour transport from the Atlantic. Reduced transpiration due to deforestation can no longer sufficiently
723 provide water vapour to sustain the latent heat required, thereby reducing the inflow of oceanic water vapour, and leading to
724 a SAM tipping in this model (Boers et al., 2017). In addition to deforestation, transpiration over northern Amazon may be
725 reduced due to an increase in seasonal tropical vegetation related to an AMOC slowdown (Akabane et al., 2024), further
726 curtailing moisture availability for the SAM.

727 3.3.2 Assessment and knowledge gaps

728 A combination of climate change and deforestation could lead to substantial changes in the SAM system, affecting many
729 millions of people. Additionally, a decrease in AMOC strength could potentially trigger major changes in tropical South
730 American precipitation (see Wunderling et al., 2023). However, the current scarcity of research in the subject limits our
731 ability to fully understand and assess the tipping potential of the SAM system, and we classify the possibility of SAM
732 tipping to be uncertain.

733 3.4 East Asian Summer Monsoon (EASM)

734 Beyond the major tropical monsoon systems, there are other subtropical monsoon systems which ostensibly share similar
735 characteristics, such as the East Asian Summer Monsoon (EASM), which is a part of the South Asian Monsoon. Unlike the
736 tropical monsoon systems which are driven by the ITCZ movement and characterised by heavy summer rainfall, the
737 precipitation and wind patterns of the EASM are linked to mid-latitude frontal systems and the jet stream, and its interaction
738 with the Tibetan Plateau (Molnar et al., 2010; Son et al., 2019). A recent study on CMIP6 projections of EASM by
739 Katzenberger and Levermann (2024) finds that the future precipitation and interannual variability in the EASM region is
740 projected to consistently increase under various climate change scenarios. They suggest these changes are primarily driven
741 by thermodynamic responses to global warming, rather than major shifts in atmospheric circulation patterns. Further, the
742 same study finds an increased frequency of extreme wet seasons based on the CMIP6 simulations.

743 3.4.1 Evidence for tipping dynamics

744 Speleothem and loess records from China, and analyses based on these records (Wang et al., 2008; Thomas et al., 2015;
745 Rousseau et al., 2023) reveal that the EASM has undergone periods of abrupt changes on millennial timescales. These
746 changes are more sudden than the change in Northern Hemisphere summer insolation that is thought to be the primary
747 driver, and this suggests a potential tipping behaviour in the EASM. Schewe et al. (2012) proposed a mechanism involving
748 moisture advection feedback to explain these abrupt changes in the past, and suggests that a minimum threshold of specific
749 humidity over the ocean is essential to sustain monsoon precipitation over the land. The changes in insolation may trigger
750 changes in sea-surface temperature and evaporation, hence specific humidity over the ocean. According to the conceptual
751 model proposed by Schewe et al. (2012), gradual, insolation-driven changes in oceanic humidity lead to abrupt transitions-
752 the system suddenly switches between states that can or cannot sustain a strong monsoon circulation whenever the oceanic
753 specific humidity crosses the threshold. Although paleoclimate records show evidence of past abrupt shifts in the EASM, the
754 mechanism detailed in Schewe et al. (2012) was contested by a later study by Boos and Storelvmo (2016). They highlight
755 that the underlying theory in Schewe et al. (2012) omitted the static stability of the troposphere. Modeling experiments by
756 Boos and Storelvmo (2016), primarily examining decadal timescales, find a more linear monsoon response to forcings. They

757 suggest that the mechanisms behind orbital-scale abrupt changes might involve very large forcings or slow-acting
758 components like ice sheets and large-scale vegetation changes.

759

760 Further, like the Indian, African, and South American monsoons, the EASM could be affected by a potential collapse of the
761 AMOC (Sun et al., 2012; Ben-Yami et al., 2024). Freshwater hosing experiments using climate models that mimic AMOC
762 slowdown show increased temperature gradients between the equator and the poles, and suggest northern westerlies acting
763 as a pathway in transmitting climate change signals from the North Atlantic to the Asian monsoon regions (Sun et al., 2012).

764 3.4.2 Assessment and knowledge gaps

765 While paleoclimate records provide compelling evidence for EASM's tipping behaviour in the past, a deeper understanding
766 of the underlying mechanisms and internal feedback is crucial for assessing its vulnerability to future tipping points. We
767 therefore classify this system as uncertain.

768 3.5 Summary

769 The major monsoon systems discussed here share some similar, underlying dynamics, as they are complex phenomena
770 emerging from the interplay of global and regional factors. As such, questions around potential tipping largely relate to the
771 susceptibility of changes in the global monsoon to anthropogenic forcing, e.g. either directly via interhemispheric
772 differences in aerosol loading or indirectly via climate-change driven AMOC collapse. In principle, regional feedbacks may
773 maintain or significantly shift a monsoon regime but there is insufficient evidence across the discussed systems, with
774 exception of the West African monsoon. This system does not only feature strong positive feedbacks involving vegetation
775 and dust, but has also reportedly collapsed abruptly several times in the past. Overall, we therefore classify tipping of the
776 West African monsoon as possible with low confidence, and South American, East Asian and Indian monsoons as uncertain.

777 4. Global atmospheric circulations

778 4.1 Tropical clouds, circulation, and climate sensitivity

779 Clouds play an important role in the climate system, as they contribute to the regulation of Earth's energy budget. In general,
780 the altitude and composition of clouds determines the amount of shortwave radiation to reach the planet's surface, as well as
781 the longwave radiation (heat) escaping to space. For example, low, thick clouds have a net cooling effect as they deflect
782 large fractions of incoming sunlight, while also having a weak greenhouse warming effect due to their low altitude (i.e.
783 having a comparable temperature to the ground). A changing climate, which causes changes in temperature, humidity, and
784 circulation patterns, affects the formation and dynamics of clouds. This, in turn, can influence the climate and how much
785 warming results from increased atmospheric CO₂ concentrations (i.e. 'climate sensitivity').

786 4.1.1 Evidence for tipping dynamics

787 Literature on cloud-induced tipping points is very limited. Yet cloud-forming processes exhibit strong hysteresis on weather
788 timescales. Indeed, a cloud droplet forms when water starts to stick to a particle after a certain level of humidity (in which a
789 so-called hygroscopic aerosol particle crosses a humidity tipping point into an unstable condensational growth phase); and
790 precipitation, once initiated, is a self-reinforcing cascade where larger particles fall faster and hence grow faster by
791 collisions. Coupling of these micro-scale processes to atmospheric dynamics can lead to spontaneous and irreversible
792 transitions at the intermediate mesoscale – in particular, the transition of shallow cloud layers from closed to open-cell
793 geometries (honeycomb-like cloud patterns formed by convecting air) (Feingold et al. 2015), driven by a self-sustaining
794 process involving precipitation-induced downdraft, subsequent updraft and new cloud formation (Yamaguchi and Feingold,
795 2015). Another example is self-aggregation of deep convection, apparently driven by radiative and moisture feedback
796 loops (Müller et al. 2022). Both of these significantly decrease cloud cover and albedo, potentially enabling climate
797 interactions. Could further coupling out to planetary scales produce climate-relevant tipping behaviour? Complicating this
798 question is the fact that cloud-related processes are not well represented in current climate models, limiting their ability to
799 guide us.

800 The most discussed possibility has been the extreme case of a global climate runaway. If the atmosphere became sufficiently
801 opaque to infrared (i.e. if it became harder for longwave heat energy to escape due to overcast high cloud, very high
802 humidity, or CFC-like greenhouse gases filling in spectral absorption windows), the planet could effectively lose its ability
803 to cool to space, producing a Venus-like runaway. Although general circulation models (GCMs) and paleoclimate evidence
804 suggest climate sensitivity rises as climate warms (Sherwood et al., 2020), calculations show virtually no chance of runaway
805 warming on Earth at current insolation levels (LeConte et al. 2013).

806 A more plausible scenario is unexpectedly strong global positive amplifying radiative feedback from clouds and high
807 climate sensitivity. Although presumably reversible, this would be serious. With respect to high clouds, suggested missing
808 feedbacks (due to novel microphysical or aggregation mechanisms) have generally been negative (e.g. Mauritsen and
809 Stevens 2015). Low clouds are a greater concern: one recent study using a multiscale atmospheric model found a strong and
810 growing positive amplifying feedback from rapid disappearance of these clouds (Schneider et al. 2019), highlighting the
811 possibility of nonlinear cloud behaviour and surprises (Bloch-Johnson et al., 2015, Caballero and Huber 2013). Although
812 various observations generally weigh against high-end climate sensitivities above 4°C per CO₂ doubling, they cannot rule
813 them out (Sherwood et al., 2020).

814 A final possibility is surprising reorganisations of tropospheric circulation. Innovative atmospheric models (Carlson and
815 Caballero 2016, Seeley and Wordsworth 2021) and geologic evidence (Tziperman and Farrell, 2009, Caballero and Huber
816 2010) have suggested possible “super-MJO” (the ‘Madden-Julian Oscillation’ being the dominant mode of ‘intraseasonal’
817 variability in the tropical Indo-Pacific, and is characterised by the eastward spread of enhanced or suppressed tropical
818 rainfall lasting less than a season) and/or reorganisation of the tropical atmospheric circulation in a warmer climate due to

cloud-circulation coupling. These scenarios are supported by little evidence, but if they did occur they could massively alter hydrology in many regions. Poor representation of tropical low clouds has also likely inhibited coupled model simulations of decadal variability or regional trends (Bellomo et al. 2014, Myers et al. 2018), raising the possibility that even if clouds cannot drive tipping points they might amplify other tipping points in ways that are missing from current models.


4.1.2 Assessment and knowledge gaps

In summary, concern about cloud-driven tipping points is relatively low. Cloud feedbacks will however likely affect the strength of climate responses, including for many tipping points. For example, they could potentially amplify variability, and current models may not be capturing this well. High climate sensitivity from strongly positive cloud feedbacks also cannot be ruled out.

4.2 El Niño-Southern Oscillation (ENSO)

The El Niño-Southern Oscillation (ENSO) is the dominant interannual mode of variability in Earth's climate. It originates in the tropical Pacific, where it affects sea surface temperatures (SST), trade winds, rainfall, and many other climate variables. El Niño events typically happen every 3 to 5 years (hence the term 'interannual'). The tropical Pacific average climate is characterised by a strong east-west gradient along the equator of about 5-6°C, with warmer SSTs in the west and colder SSTs in the east maintained by easterly Pacific trade winds. During El Niño, the warm phase of this oscillation, this gradient weakens, while during La Niña, its cold phase, it intensifies (examples are shown in Figure 9a). Both phases of this oscillation have far-reaching impacts on global climate and weather patterns, ecosystems, and human health (e.g. McPhaden et al. 2020).

The impacts of ENSO become especially pronounced during the strongest events, often referred to as extreme El Niños (Figure 9b). At their peak, these events can eliminate the east-west ocean temperature gradient along the equator, leading to a temporary collapse of the trade winds. Additionally, an extreme El Niño causes an increase in global mean surface temperature of up to 0.25 °C (e.g. Hu and Fedorov 2017), contributing to the prevalence of heat waves around the globe. While only a few El Niño events reach large magnitudes, the global impacts of these events result in billions of dollars in damage (e.g. Callahan and Mankin 2023).



844

845 **Figure 9: ENSO warm and cold phases and observational record. a** Examples of strong La Niña (top) and El Niño (bottom) events
 846 seen in the tropical Pacific surface temperature (SST) distribution, with characteristic strong and weak SST gradient along the
 847 equator, respectively. **b** ENSO record since the 1980s. Observed SST variations in the eastern equatorial Pacific since 1980. Note
 848 the three extreme events of the past four decades (1982, 1997 and 2015) and the weakening of ENSO variability between years
 849 2000 and 2015. Temperature is averaged for the NINO3 region (5°C-5°N, 150°W-90°W) in the eastern equatorial Pacific. Based on
 850 NOAA Extended Reconstructed SST V5 data (Huang et al., 2017).

851 4.2.1 Evidence for tipping dynamics

852 Extensive research conducted since the 1980s has significantly advanced our understanding of the physics behind El Niño,
 853 leading to improved predictive capabilities of climate models (L’Heureux et al., 2017). ENSO is now recognised as a
 854 large-scale, irregular, internal oscillatory mode of variability within the tropical climate system influenced by atmospheric
 855 noise (Timmermann et al., 2018). The spatial pattern of ENSO is determined by ocean-atmosphere feedbacks, while its
 856 timescale is determined by ocean dynamics. In particular, it is a sequence of self-reinforcing feedbacks between SSTs,
 857 changes in zonal surface winds, equatorial upwelling, zonal equatorial currents, and ocean thermocline depth that promotes
 858 the growth of El Niño anomalies and contributes to the positive Bjerknes feedback (McPhaden et al. 2020) .

859

860 Coral-based proxy data indicate that the amplitude and frequency of ENSO events has gradually increased during the
 861 Holocene (Grothe et al., 2019; Lawman et al., 2022), possibly due to an increase in strong El Niño events. All extreme El
 862 Niños in the observational record (1982, 1997, and 2015) occurred during the accelerated growth of global mean
 863 temperatures. This raises the question whether this trend is indicative of upcoming changes in the tropical Pacific to
 864 conditions with more frequent extreme El Niño events.

In the context of tipping points, the question arises: is there a critical threshold with an abrupt and/or irreversible transition to such a new state? While there are no indications so far for a tipping point in ENSO behavior, several recent studies (e.g. Cai et al. 2018, 2022; Heede and Fedorov 2023a) have indeed suggested that El Niño magnitude and impacts may intensify under global warming (Figure 10), even though there is still no model consensus on the systematic future change in ENSO as IPCC AR6 and the results in Figure 10 suggest.




Figure 10: Overview of projected changes in extreme El Niño events in CMIP6 climate models. The bar chart shows the time-mean frequency of extreme El Niño events (the number of events per decade) for several idealised and more realistic global warming experiments (abrupt-4xCO₂, 1pctCO₂, SSP5-8.5 and SSP1-2.6) next to the pre-industrial Control simulation (piControl). From Heede and Fedorov 2023a (cropped).

It is projected that the eastern equatorial Pacific will warm faster than the western part of the basin, leading to an eastern equatorial Pacific (EEP) warming pattern or El Niño-like mean conditions, associated with weaker Pacific trade winds. Most climate model future projections exhibit this pattern (e.g. DiNezio et al. 2009; Xie et al. 2010; Heede and Fedorov 2021), and increased ENSO variability is prevalent in models that simulate stronger nonlinear (Bjerknes) feedbacks (Cai et al. 2022). A recent comprehensive study of CMIP6 models and scenarios concluded that although a common mechanism to explain a change in ENSO activity across models is missing, its increase under warming scenarios is robust (Heede and Fedorov, 2023a).

Furthermore, during the warm Pliocene epoch approximately 3-5 million years ago when global surface temperature were ~3°C above pre-industrial, the east-west SST gradient was indeed reduced (Wara et al. 2005; Fedorov et al. 2006, 2013, 2015; Tierney et al. 2019). This state is often referred to as ‘permanent El Niño-like’ conditions, which does not indicate ENSO changes but rather a consistent mean decrease in the east-west SST gradient. While debates on this topic are ongoing, estimates for this gradient reduction range from 1.5 to 4°C, depending on the time interval, proxy data, and the definition of this gradient.

4.2.2 Assessment and knowledge gaps

Therefore, there is a general expectation of a future reduction in the Pacific's east-west SST gradient by the end of the 21st century. Together with other contributing factors, such as the strengthening of the Madden-Julian Oscillation (MJO), the

890 dominant intraseasonal mode in the tropical Indo-Pacific, Arnold and Randall 2015), this reduction is expected to amplify
891 ENSO (Heede and Fedorov 2023a). Additionally, a warmer atmosphere can hold more water vapor, which could result in
892 stronger precipitation and heating anomalies in the atmosphere, leading to greater remote impacts of El Niño events.
893 Consequently, the collective evidence implies an increase of El Niño magnitude and impacts under global warming. There
894 is, however, insufficient indication for a critical transition associated with an abrupt or irreversible regime shift towards a
895 new, more extreme, or persistent ENSO state, such that ENSO is not considered a tipping system with medium confidence
896 (see also Armstrong McKay et al., 2022). However, it is well connected to other Earth system components (e.g. affecting
897 tropical monsoon rainfall), thereby possibly playing a role in tipping cascades linking different tipping elements via global
898 teleconnections (Wunderling et al., 2023).

899 Notably, the projections of a future EEP warming pattern, weaker mean trade winds, and stronger El Niño events contradict
900 the recent decadal trends in the tropical climate over the past 30 years or so. In fact, since the early 1990s, the Pacific trade
901 winds have strengthened, and the eastern equatorial Pacific has become colder (e.g. Ma and Zhou. 2016; Seager et al. 2022;
902 Wills et al. 2022; Heede and Fedorov 2023b). Whether these trends reflect an ocean thermostat-like response to global
903 warming, internal variability of the system, or both, remains an open question. Similarly, the magnitude of ENSO events has
904 been generally weaker since the 2000s compared to the 1980s and 1990s (Fig. 1B; also Capotondi et al. 2015 or Fedorov et
905 al. 2020).

906 Therefore, debates on the future of the tropical Pacific and ENSO revolve around the question of when the transition to a
907 mean EEP pattern and weaker trade winds may occur, likely leading to a stronger El Niño and more frequent extreme
908 events. Simulations with global climate models including strongly eddying ocean components (Wieners et al. 2019; Chang
909 et al., 2020) and the recently developing 2023-2024 El Niño are expected to help reduce persistent model tropical biases in
910 SST, precipitation, and ocean thermocline, and to resolve some of the unresolved issues.

911 4.3 Mid-latitude atmospheric dynamics

912 The mid-latitude atmospheric circulation is characterised by a band of strong westerly winds, with the largest velocities at an
913 altitude of 7-12 km. In the Northern Hemisphere, these form the so-called polar ‘jet-stream’. The jet serves as a separation
914 of cold air masses at high-latitudes in the North from temperate air masses further south. Large meanders in the jet are
915 referred to as Planetary, or Rossby waves. In most cases, these waves move over large distances and decline over timescales
916 of a few days. When persisting for a prolonged time over the same location (referred to as ‘quasi-stationary’ waves) they can
917 lead to high-impact climate extremes, including temperature extremes or heavy precipitation. An example is the record
918 breaking heatwave of 2021 in the North American Pacific Northwest (Philip et al., 2021).




Figure 11: Potential changes in mid-latitude atmospheric circulations under global warming, exemplary for the Northern Hemisphere. Reduction of AMOC, atmospheric blocking events, arctic warming and other drivers can modify the jet stream. Potential consequences are a northward shift, slowdown and enhanced meandering, related to increases in extreme weather phenomena.

Atmospheric features such as blocks (quasi-stationary high-pressure regions that divert, or “block” the large-scale atmospheric flow on timescales of several days to weeks) are intimately linked to these persistent meanders in the jet. A widely discussed effect of climate change is a poleward shift of the mid-latitude jet, although this may be season and location dependent (Oudar et al. 2020), and smaller than previously thought (Curtis et al., 2020) (Figure 11).

4.3.1 Evidence for tipping dynamics

In climate models, the magnitude of the jet’s meridional shift strongly depends on the reduction of the overturning circulation in the Atlantic (AMOC). Models with a strong AMOC reduction in the future tend to project a stronger poleward shift of the jet than models with a weaker AMOC reduction, which sometimes even display an equatorward shift (Bellomo et al., 2021). This is because a strong weakening of the AMOC might amplify the latitudinal temperature gradient that drives the strength and position of the zonal jet. As a consequence, the extent of AMOC change is a large contributor to atmospheric circulation uncertainty in regional climate change projections (Bellomo et al., 2021).

Furthermore, it has been suggested that the mid-latitude flow might weaken, leading to more persistent and slower moving weather patterns (Coumou et al. 2015, Kornhuber & Tamarin-Brodsky 2021). A possible driver is Arctic amplification, namely the fact that the Arctic is warming more rapidly than the rest of the planet, driven by a variety of mechanisms (Previdi et al., 2021) including temperature- and sea ice-related feedbacks (Dai et al., 2019). Arctic amplification reduces the equator–pole temperature contrast, and could result in a weakening and enhanced meandering of the jet stream (e.g. Francis

939 and Vavrus, 2015). This, in turn, could potentially lead to an even larger reduction in the latitudinal temperature contrast,
940 further weakening the jet and so on. While Arctic amplification is most evident during winter, such increase in waviness
941 may also be occurring during the summer season (Coumou et al., 2018). However, evidence that the occurrence of
942 large-amplitude atmospheric waves is increasing is debated (e.g. Screen and Simmonds, 2013; Blackport and Screen, 2020;
943 Riboldi et al., 2020), and mechanisms which would reduce blocking in the future have also been proposed (e.g. Kennedy et
944 al., 2016).

945 As part of this debate, it has been proposed that several weather extremes in recent decades were associated with a
946 quasi-stationary, quasi-resonant wave pattern. This results from the interaction of climatological waves that are perpetually
947 forced by orography (mountain geography) and land-sea contrasts with transient meanders of the jet stream (Petoukhov et
948 al., 2013), given a set of favourable conditions (White et al. 2022). Petoukhov et al. (2013) also hypothesised that Arctic
949 amplification and the associated weakened, wavier jet may provide increasingly favourable conditions for the occurrence of
950 quasi-resonance. This can result in circulation features which accelerate regional extreme weather occurrence trends, for
951 example heatwave trends in Europe (Rousi et al. 2022), although the direction of causality is debated (Wirth and Polster,
952 2021). If recent extreme events are indeed associated with a resonance mechanism that only is triggered when the jet crosses
953 a certain threshold in waviness, a tipping point might be involved. However, it is uncertain whether this would be associated
954 with hysteresis and irreversibility or would just be a reversible, but abrupt, shift of the atmosphere towards enhanced
955 large-amplitude mid-latitude waves.

956 More generally, there is no robust evidence that continued climate change and Arctic amplification will lead to a tipping
957 towards a wavy-jet state, systematically higher amplitude and/or more frequent planetary waves, or blocks. Equally, there is
958 no robust evidence that these hypothetical changes would be self-sustaining. Indeed, while a number of large changes in
959 atmospheric dynamical features may occur under climate change, these are typically discussed as gradual changes, without
960 explicit hysteresis or tipping behaviour. Similarly, there is no robust evidence pointing to tipping-like behaviour in the jet
961 stream's latitudinal location, although gradual, long-term shifts may occur.

962 It should nonetheless be noted that atmospheric circulation responses to climate change are characterised by large model
963 uncertainty and are possibly biased by the relatively low resolution of global climate models compared to e.g.
964 weather-prediction models (Shepherd, 2019). Indeed, there is indication that for models to exhibit tipping in atmospheric
965 blocking, they need to involve a sensitive sea-ice model and sufficient resolution to capture the sea-ice-atmosphere-AMOC
966 feedback driving the transition (Drijfhout et al., 2013).

967 4.3.2 Assessment and knowledge gaps

968 Although theoretically possible, there is thus no robust evidence for tipping point behaviour in mid-latitude atmospheric
969 circulations in the near future. At the same time, a number of relevant physical processes are currently debated or
970 ill-constrained, not least by large model uncertainty. We thus evaluate the mid-latitude atmosphere as not displaying tipping
971 points, with low confidence.

972 The mid-latitude large-scale circulation itself may nonetheless be affected by tipping behaviour of other components of the
 973 Earth system to which it is coupled. An example is the Greenland ice sheet, whose topography is important for determining
 974 the atmospheric circulation in the North Atlantic sector (White et al., 2019). A collapse of the Greenland ice sheet may
 975 trigger tipping-like changes in the atmospheric circulation in the region. The mid-latitude large-scale circulation may also
 976 contribute to tipping behaviour elsewhere in the Earth System. For example, joint non-tipping changes in mid-latitude
 977 atmospheric dynamics, the associated surface climate, and other components of the Earth system, may lead to tipping point
 978 behaviour in vegetation (Lloret and Batllori, 2021). Similarly, drier soils, a strengthened land-atmosphere coupling, and a
 979 contribution from large-scale circulation anomalies recently triggered a shift to hotter and drier conditions in inner East Asia
 980 (Zhang et al., 2020). Both the vegetation and dryness changes could in turn feed back onto the atmospheric circulation.
 981 Due to such feedbacks and interactions between the atmospheric circulation and other components of the Earth system, and
 982 due to its role in weather and climate extremes, an improved understanding of the physical processes underlying changes in
 983 mid-latitude atmospheric dynamics under recent and future climate change appears pivotal in a tipping point context. Large
 984 model uncertainty in projecting abrupt regional atmospheric circulation changes conditioned by changes in the ocean,
 985 cryosphere, or land surface would lend itself eminently for a storyline approach (Zappa and Shepherd, 2017). Tipping of the
 986 atmospheric circulation, and associated weather extremes, would then be conditioned by threshold behaviour in other,
 987 connected systems.
 988 Finally, we argue for the need to investigate whether recent, record-breaking weather extremes can be explained by the
 989 gradual changes in likelihood distribution due to the recent and ongoing climate change, or whether they are signs of
 990 abruptly changing likelihood distributions. Such a shift in the distribution of extremes could be diagnosed using extreme
 991 value theory. Although a shift cannot be associated with a global tipping point, it would suggest that the extreme value
 992 distribution of (a) certain type(s) of extreme weather did witness regional tipping, whether or not reversible, in the sense of a
 993 large nonlinear change in response to a small and gradual change in forcing, potentially driven by self-sustaining feedbacks.

994 **5. Discussion and Outlook**

995 In general, the circulation systems discussed in this review are driven by gradients in, e.g., temperature, salinity or pressure,
 996 and interconnect various parts of the Earth system. They are therefore very susceptible to the multitude of changes that
 997 humans are causing to all planetary spheres, disrupting established equilibria and dynamics. Alongside the significant input
 998 of additional energy to the Earth system via greenhouse gas emissions, large-scale land-use change, injection of aerosols to
 999 the atmosphere and other drivers directly and indirectly impact the atmospheric and oceanic circulation systems, all of which
 1000 show signs of change (IPCC AR6, 2021).
 1001 In this review, we have investigated the potential for tipping in these systems, anchored to a tipping definition of strong
 1002 positive, reinforcing feedbacks that lead to a large change beyond a forcing threshold (following Armstrong McKay et al. ,
 1003 2022). Synthesis Table 1 summarises our findings, listing key drivers of change and destabilisation for each circulation

1004 system, as well as ensuing impacts following a state shift. Importantly, we highlight a number of positive feedbacks that
1005 could give rise to self-reinforcing change beyond a forcing threshold. Finally, we collate evidence for signature tipping
1006 characteristics like hysteresis and abruptness, and classify each potential tipping system along the criteria defined in the
1007 introduction.

1008

1009 We classify the oceanic overturning circulations - the subpolar gyre (SPG), the Atlantic Meridional Overturning Circulation
1010 (AMOC) and Antarctic Overturning Circulation - as tipping systems with medium confidence. There are signs of a
1011 weakening overturning in the AMOC and of the SPG circulation, and a projected decline of all three under a warming
1012 climate. One common main driver of this change is the freshwater influx, primarily from melting of the ice sheets,
1013 disrupting the buoyancy conditions at the overturning sites. Other contributing factors include warming oceans, and changes
1014 in wind and precipitation patterns. While process understanding, modelling and paleo evidence suggest a possible bistability
1015 in these systems, the assessment of exact tipping thresholds is difficult. Current climate models suffer from imperfect
1016 representation of some important processes (such as eddies and mixing) and from biases which can impact the response to
1017 forcings. In particular, the representation of deep convection in the Southern Ocean is deficient, with convection occurring
1018 in the open ocean rather than shelf seas (Heuzé et al., 2021; Purich and England, 2021). It has further been shown that the
1019 model representation of SPG tipping can be dependent on biases in stratification in the North Atlantic (Sgubin et al., 2017).
1020 For the AMOC, various studies have suggested that it tends to be too stable in many climate models because of biases in
1021 simulating salinity dynamics and the intertropical convergence zone (ITCZ) (Liu et al., 2017; Mecking et al 2017,
1022 Swingedouw et al. 2022). As observational time scales are short compared to the natural variability timescales of the oceanic
1023 overturning circulations, paleo reconstructions have proven indispensable for understanding potential long-term changes.
1024 Indeed, the early warning methodology, increasingly applied to the North Atlantic (Boers, 2021; Ditlevsen and Ditlevsen,
1025 2023; Michel et al., 2022), is highly dependent on these historical proxies. Albeit being subject to ongoing debate (Michel et
1026 al., 2023), these warnings add to the reasons for concern regarding the possible tipping of ocean circulation systems.

1027 Moving to the monsoon systems, our review highlights that there are multiple drivers beyond (direct or indirect)
1028 anthropogenic climate change that influence these systems. Land-use change on large scales and aerosol emissions have
1029 direct impacts on the monsoon dynamics, potentially through positive feedback loops. For example, the vegetation-albedo
1030 coupling is identified as a critical feedback having led to past abrupt shifts in the West African monsoon (WAM), linked to
1031 drastic changes in the Sahara vegetation states. However, limitations in modelling vegetation shifts and model disagreements
1032 on future trends call for improved assessments. We thus consider the WAM as a tipping system, yet with low confidence.
1033 Evidence for tipping is limited for the other monsoon systems under consideration over Asia and South America. Hence, we
1034 attribute an uncertain status to them. Studies on the Indian summer monsoon (ISM) tipping so far have employed simplified
1035 analytical models that fail to represent the full spectrum of feedbacks and processes within this complex monsoon system
1036 (Boos and Storelvmo, 2016; Kumar and Seshadri, 2023). Similar arguments apply to the East Asian summer monsoon
1037 (EASM). Current climate models further inadequately capture South American monsoon (SAM) precipitation, and the

1038 impact of climate change on SAM precipitation remains uncertain (Jones and Carvalho, 2013; Douville et al., 2021). Further
 1039 research utilising paleoclimate evidence, climate model reconstructions of past monsoons, and comprehensive modelling of
 1040 the system's responses to various forcing thresholds are required to clarify any non-linear changes within both the Indian
 1041 and South American monsoon systems. Critically, the dynamics of the global monsoon are substantially influenced by the
 1042 AMOC, the position of the ITCZ and interhemispheric aerosol loadings, such that a global, coupled approach is promising
 1043 for more insights into potential tipping of the monsoon systems.

1044 Finally, we have considered global atmospheric circulation systems. Although theoretically possible, there is no robust
 1045 evidence for tipping point behaviour in mid-latitude atmospheric circulations. Nonetheless, there are some physical
 1046 mechanisms, such as enhanced jet stream meandering and planetary wave resonance, which may lead to tipping-like
 1047 behaviour. However, their physical interpretation and relevance for tipping are debated. In more general terms, the
 1048 atmospheric circulation responses to climate change are characterised by large model uncertainty (Shepherd, 2019), and
 1049 mid-latitude circulations may be affected by tipping behaviour of other components of the Earth system (Orihuela-Pinto et
 1050 al., 2022). We thus categorise the mid-latitude atmosphere as not displaying tipping points, albeit with low confidence.

1051 Similarly, there is insufficient indication for a state shift in El Niño Southern Oscillation (ENSO) towards a state with more
 1052 persistent or extreme El Niño events. An increase of El Niño magnitude and impacts is expected under global warming,
 1053 however, not abruptly or irreversibly. For these reasons and a lack of a comprehensive mechanism across models to explain
 1054 a change in future ENSO activity (Heede and Fedorov, 2023a), we classify ENSO as no tipping system with medium
 1055 confidence. Residual uncertainties remain due to divergence of model predictions, and known model biases in SST,
 1056 precipitation and ocean thermocline (Wieners et al. 2019; Chang et al., 2020). With similar confidence we can rule out
 1057 cloud-driven tipping points. Although large scale reorganisations are in principle conceivable, conjectures of strong positive
 1058 feedbacks leading to extreme climate sensitivity, e.g. runaway warming or rapid disappearance of low clouds, are deemed
 1059 unlikely based on current models and understanding (LeConte et al. 2013). In summary, we rate the concern about tipping of
 1060 these global circulation systems as low with medium to low confidence. This, however, does not rule out non-linear
 1061 amplification of trends or interactions with other tipping systems in the Earth system, like the high-latitude cryosphere.

1062 The review presented here was conducted by an expert group, drawing on multiple lines of evidence, including
 1063 observational data, coupled models of low and high complexity and paleo records. It presents a broad overview of the
 1064 present state of knowledge on the subject of circulation tipping points, while acknowledging the challenges and limitations
 1065 discussed below

1066 One general difficulty in assessing tipping points is the inherent need to map out and quantify relevant amplifying and
 1067 dampening feedback loops. Naturally, this requires understanding and modelling nonlinearities of the underlying systems,
 1068 which becomes increasingly challenging conceptually when the relevant feedbacks involve many diverse systems (as they
 1069 typically do for coupled ocean and atmosphere circulations). This conceptual issue translates into a numerical one, where
 1070 model complexity needs to be traded-off against computational runtimes. Partly for these reasons, there is currently only a

1071 limited evidence base on systematic assessments of destabilising drivers on potential tipping systems, such as NAHosMIP
 1072 exploring freshwater impacts on the AMOC (Jackson et al., 2023).

1073 In recent years, however, there has been a surge of numerical experiments with models of ever-increasing complexity and
 1074 resolution (e.g., Liu et al., 2017, Mecking et al 2017, Chang et al., 2020, Li et al., 2023). These experiments shed light on so
 1075 far under-represented important feedback processes and small-scale dynamics, and identify potential model biases towards
 1076 simulating too stable conditions. Model resolution will affect the type of instabilities which can be represented. For example,
 1077 on ocean circulations, once the spatial resolution of an ocean model is smaller than the internal Rossby deformation radius,
 1078 baroclinic instabilities can be represented leading to ocean eddies. The presence of these eddies will also lead to
 1079 modification of the mean flow and associated transports, and hence affect its stability properties. However, this has only
 1080 been systematically analysed for very idealised cases, such as wind-driven ocean circulation (Berloff and McWilliams,
 1081 1999). Moreover, there was no relationship found between the presence of weak AMOC states (that are stable for at least a
 1082 hundred years) and ocean resolution in a set of CMIP6 GCMs (Jackson et al 2023). Therefore, although higher resolution
 1083 captures important dynamics, its effect on model stability remains an active area of investigation. Model complexity, if
 1084 meant as to include additional processes, can strongly affect the stability of an ocean flow. For example, by including a
 1085 sea-ice component into an ocean model, the multi-stability regime of the AMOC will be affected (Van Westen and Dijkstra,
 1086 2023; Van Westen et al., 2024). Along similar lines, recent work demonstrates that high-resolution modelling of ocean
 1087 eddies impacts mid-latitude atmospheric patterns and storm tracks (Zhou and Cheng, 2022; Czaja et al., 2019), albeit with
 1088 no explicit investigation on potential tipping.

1089 Building on this progress, with respect to tipping in ocean and atmosphere circulations, future research should intensify its
 1090 efforts to address these key research questions:

1091

- 1092 (1) Are circulation systems susceptible to tipping? Where tipping is in principle conceivable, is its absence in
 1093 numerical simulations a deficit of the model or can physical stabilising feedbacks be identified? This involves
 1094 bridging the gap between physical process understanding and conceptual models (which clearly allow for tipping)
 1095 on one hand, and high-complexity and -resolution numerical models that might be biased towards stability on the
 1096 other hand. Paleo reconstructions provide an essential resource for this, along with continued observations at high
 1097 spatial and temporal resolutions.
- 1098 (2) At which levels of forcing are tipping thresholds expected based on systematic uncertainty assessment? Such an
 1099 analysis should take into account different forms of uncertainties such as on model initial conditions,
 1100 parameterisation and model structure as well as in data products from Earth observations and paleoclimate
 1101 reconstructions. New methodologies need to be developed to propagate these uncertainties transparently towards
 1102 providing reliable and interpretable estimates of tipping threshold distributions.
- 1103 (3) What are the impacts of tipping point transgressions? Tipping of any of the discussed systems would entail
 1104 significant consequences for Earth and human systems. As for the AMOC, a state shift can cause disruptions in

1105 many other systems across the globe, including shifts in precipitation patterns in the Northern hemisphere and in
 1106 the tropics with significant expected impacts on agriculture, human settlements and other infrastructure adapted to
 1107 pre-tipping climate conditions. An understanding of the tipping dynamics and the anticipated biophysical states
 1108 after tipping can serve as foundation for standardised impact modelling, or for storyline approaches on extreme
 1109 climate change.
 1110
 1111 An initiative designed to explicitly investigate these research questions in a systematic way is the now established Tipping
 1112 Points Modelling Intercomparison Project ([TIPMIP](#); Winkelmann et al., 2025). To that end, it is designed to study the
 1113 occurrence of tipping across a range of biophysical systems, investigating the critical thresholds and aspects like
 1114 irreversibility, rate-induced tipping and spatio-temporal scales. TIPMIP builds on domain-specific endeavours (e.g.
 1115 NAHosMIP, ISMIP, ...), thereby complementing general modelling initiatives addressing parts of these questions in ongoing
 1116 programs (e.g. CMIP). In addition, updated expert elicitations on e.g. tipping interactions (Kriegler et al., 2008) can serve as
 1117 a valuable complement to numerical modelling efforts. Overall, the multi-faceted questions on the tipping potential of
 1118 circulation systems may be answered by studying the hierarchy of models. This way, the effects of small-scale processes
 1119 (model resolution) and additional large-scale feedback processes (model complexity) on the tipping behaviour can be
 1120 systematically assessed (Dijkstra, 2013). Relatedly, theoretical advances shed light on new properties to be studied in tipping
 1121 systems such as a shifted focus from stable long-term attractors to the change in relative basin widths when studying
 1122 rate-induced tipping (Ashwin et al., 2012; Feudel, 2023). The former is anyways often-times regarded as oversimplification
 1123 that does not fully hold up to time-varying, continuous drivers (Bathiany et al., 2018) or in systems with spatial degrees of
 1124 freedom (Rietkerk et al., 2004). Steps forward can build on insights from hydrodynamic theory such as critical transitions in
 1125 turbulence on multiple scales (van Kan and Alexakis, 2020; van Kan, 2024) or transitions between multiple- stable states in
 1126 aquaplanets (Brunetti et al., 2019; Ragon et al., 2022). Of particular importance in the context of circulation systems is the
 1127 role of interactions between different parts of the Earth system, and the implications for reciprocal (in-)stability and aspects
 1128 like early warning signals (Klose et al., 2021). Overall, it is needed to advance not only resolution and complexity of
 1129 numerical models but also to further pursue fundamental, conceptual research in nonlinear Earth system dynamics for a
 1130 better understanding of the underlying mechanisms and relevant uncertainties in climate tipping points. Finally, an improved
 1131 understanding of how to monitor tipping systems is needed, to eventually establish early warning systems for mitigation and
 1132 adaptation purposes (Swingedouw et al., 2020; Wood et al., 2024).

1133 Table 1. Summary of evidence for tipping dynamics, key drivers, and biophysical impacts in each system considered in this
 1134 review. Key: +++ Yes (high confidence), ++ Yes (medium confidence), + Yes (low confidence), - - - No (high confidence), -
 1135 - No (medium confidence), - No (low confidence) Primary drivers are bolded, DC: Direct Climate driver (via direct
 1136 impact of emissions on radiative forcing); CA: Climate-Associated driver (including second-order & related effects of
 1137 climate change); NC: Non-Climate driver, PF: positive (amplifying) feedback, NF: negative (damping) feedback. Drivers
 1138 can enhance (↑) the tipping process or counter it (↓)

Key drivers	Key biophysical impacts	Key feedbacks	Abrupt / large rate change?	Critical threshold(s)?	Irreversible? (timescale)	Tipping system
OCEAN CIRCULATIONS						
Atlantic Meridional Overturning Circulation (AMOC) Shutdown/collapse						
<ul style="list-style-type: none"> DC: ocean warming (↑) DC: precipitation increase (↑) CA: Greenland ice sheet meltwater increase (↑, primary in the future) CA: Arctic river discharge increase (↑) CA: sea ice extent & thickness decrease (↑) 	<ul style="list-style-type: none"> Cooling over Northern Hemisphere (up to 10°C over W/N Europe) Warming of the surface South Atlantic Change in precipitation and weather patterns over Europe Change in location and strength of rainfall in all tropical regions Reduced efficiency of global carbon sink, and ocean acidification Reduced support for primary production in Atlantic oceans Deoxygenation in the North Atlantic Change in sea level in the North Atlantic Modification of sea ice and arctic permafrost distribution Increase in winter storminess in western Europe Southward migration of the Atlantic subtropical gyres Reduced land productivity in western Europe Increased wetland in some tropical areas and associated methane emission Change in rainforest response in drying regions 	<ul style="list-style-type: none"> Salt-advection n (↑) Sea-ice melting (↑) Heat transport (↓) Temperature (↑) Surface heat flux (↑) Collapse of convection in the Labrador and Irminger Seas (↑) 	Feedback-dependent: Century (basin-wide salt advection feedback), Few decades (North Atlantic salt-advection feedback), < few decades (sudden increase in sea-ice cover in all convective regions)	Salinity change/freshwater/AMOC strength Thresholds likely path-dependent (depending on rate and spatial pattern)	++ (centuries)	++
North Atlantic Subpolar Gyre (SPG) Collapse						

Key drivers	Key biophysical impacts	Key feedbacks	Abrupt / large rate change?	Critical threshold(s)?	Irreversible? (timescale)	Tipping system
<ul style="list-style-type: none"> DC: ocean warming (↑) DC: precipitation increase (↑) CA: Greenland ice sheet meltwater increase (↑, primary in the future) CA: Arctic river discharge increase (↑) CA: sea ice extent & thickness decrease (↑) 	<ul style="list-style-type: none"> Increase in summer heat waves frequency Collapse of the North Atlantic spring bloom and the Atlantic marine primary productivity Increase in regional ocean acidification Regional long-term oxygen decline Impact on marine ecosystems in the tropics and subtropics 	<ul style="list-style-type: none"> Salt-advection (↑) Sea-ice melting (↑) Heat transport (↓) Temperature (↑) Surface heat flux (↑) Collapse of convection in the Labrador and Irminger Seas (↑) 	Years to few decades	Salinity change/freshwater Global warming 1.1-3.8°C	++ (decades)	++
Southern Ocean circulation Antarctic Overturning Collapse / Rapid continental shelf warming						
<ul style="list-style-type: none"> DC: ocean warming (↑) CA: Antarctic ice sheet meltwater increase (↑) CA: wind trends (↑) CA: Sea ice formation (↑) DC: precipitation increase (↑) 	<ul style="list-style-type: none"> Modification of Earth's global energy balance, timing of reaching 2°C global warming Reduced efficiency of global carbon sink Change in global heat storage Reduced support for primary production in world's oceans Drying of Southern Hemisphere Wetting of Northern Hemisphere Modification of regional albedo, shelf water temperatures Potential feedbacks to further ice shelf melt 	<ul style="list-style-type: none"> Density-stratification (↑) Meltwater-warming (↑) 	++ (AABW formation & abyssal overturning shutdown within decades)	Salinity change/freshwater	++ (cavity warming reversion would need 20th-century atm conditions + reduced meltwater input)	++
ATMOSPHERIC CIRCULATIONS: MONSOONS						
Indian summer monsoon (ISM) Collapse / Shift to low-precipitation state						

Key drivers	Key biophysical impacts	Key feedbacks	Abrupt / large rate change?	Critical threshold(s)?	Irreversible? (timescale)	Tipping system
<ul style="list-style-type: none"> NA: increased summer insolation (↓) DC: increased water vapour in atmosphere (↓) CA: Indian Ocean Dipole events (?) CA: ENSO change (?) CA: North Atlantic cold SST (↑) NC: aerosol loading (↑) CA: Indian Ocean warming (↑) CA: low cloud reduction (↓) 	<ul style="list-style-type: none"> Massive change in precipitation Change in tropical and subtropical climates Biodiversity loss and ecosystem degradation 	<ul style="list-style-type: none"> Moisture-advection (↓) 	Decades to centuries	Regional AOD level over Indian subcontinent (>0.25) Interhemispheric AOD difference (>0.15) AMOC slowdown (unknown threshold)	Uncertain; likely decades to centuries	unknown
West African monsoon (WAM) Collapse or abrupt strengthening						
<ul style="list-style-type: none"> DC: increased water vapour in atmosphere NA: increased summer insolation NC: land-cover change CA: desertification CA: AMOC slowdown CA: regional SST variations CA: High latitude cooling CA/NC: regional soil moisture variation CA/NC: regional vegetation variation NA: dust emissions 	<ul style="list-style-type: none"> Massive change in precipitation Change in tropical and subtropical climates Biodiversity loss and ecosystem degradation 	<ul style="list-style-type: none"> Vegetation-albedo (↑) 	Decades to centuries	Insolation changes in the Northern Hemisphere summers and surface albedo changes (unknown threshold) Interhemispheric asymmetry in AOD (>0.15) AMOC slowdown (unknown threshold)	Decades to centuries	+
South American monsoon (SAM) Collapse or abrupt strengthening						
<ul style="list-style-type: none"> DC: increased water vapour in atmosphere NA: increased summer insolation CA: AMOC slowdown NC: Amazon deforestation 	<ul style="list-style-type: none"> Massive change in precipitation Change in tropical and subtropical climates Biodiversity loss and ecosystem degradation 	<ul style="list-style-type: none"> Vegetation-moisture (?) 	Decades	Interhemispheric asymmetry in AOD (>0.15) Extent of Amazon deforestation (30-50%) AMOC slowdown (unknown threshold)	Uncertain; likely decades to centuries	unknown

GLOBAL ATMOSPHERIC CIRCULATIONS
Tropical clouds, circulation and climate sensitivity Shift to different large-scale configuration

Key drivers	Key biophysical impacts	Key feedbacks	Abrupt / large rate change?	Critical threshold(s)?	Irreversible? (timescale)	Tipping system
<ul style="list-style-type: none"> DC: atmospheric warming (†) DC: ocean warming (†) 	<ul style="list-style-type: none"> Massive alteration of hydrology in many regions Impact on ambient atmospheric-oceanic phenomena such as ENSO Strong intensification of global climate change 	<ul style="list-style-type: none"> Cloud-moisture-radiation (†) 	Unknown	Unknown	Unknown	--
El Niño Southern Oscillation (ENSO) Shift to more extreme or persistent state						
<ul style="list-style-type: none"> DC: east vs west Pacific warming (†) DC: increased water vapour in atmosphere (†) DC: weaker trade winds (†) CA: MJO strengthening (†) 	<ul style="list-style-type: none"> Temporary trade wind collapse during El Niño phase Increase in global mean surface temperatures during El Niño phase Modification of global atmospheric circulation Modification of worldwide patterns of weather variability 	<ul style="list-style-type: none"> Bjerknes (†) (SST-trade winds-ocean thermocline) 	No evidence (gradual)	No evidence (gradual)	No evidence	--
Mid-latitude atmospheric dynamics Shift to wavy-jet state / more frequent or extreme planetary waves or blocks / latitude shift						
<ul style="list-style-type: none"> CA: AMOC slowdown CA: Midlatitude flow weakening (†) DC: Arctic amplification (†) 	<ul style="list-style-type: none"> More persistent and slower moving weather patterns Increase in extreme events on Northern hemisphere 	<ul style="list-style-type: none"> Debated: Waviness quasi-resonance (†) 	No evidence	Potentially waviness threshold, beyond which quasi-resonance kicks in	No evidence	-

1139

1140

1141

1142 *Code and data availability.* There is no data or code that has been produced for this review article.

1143

1144 *Author contributions.* SL and D.I.A.M have designed the overall structure and scope. All authors have contributed to the
1145 writing. SL and YA produced the figures with input from the authors. All authors have reviewed and edited the final version
1146 of the manuscript.

1147

1148 *Competing Interests.* Some authors are members of the editorial board of the journal Earth System Dynamics. Otherwise, the
1149 authors have no other competing interests to declare.

1150

1151 *Acknowledgements.* This review article has been carried out within the framework of the Global Tipping Points Report 2023.
1152 We would like to thank Norman Steinert for developing the original concept of the tipping systems overview figure (Fig 1)
1153 and for providing the basemap. S.L. and D.I.A.M. acknowledge financial support from the Earth Commission. The work of
1154 the Earth Commission, a program of Future Earth, is made possible through the support of the Global Commons Alliance, a
1155 sponsored project of Rockefeller Philanthropy Advisors. Furthermore, D.I.A.M. acknowledges financial support via the
1156 Global Tipping Points Report project, funded by the Bezos Earth Fund. Y.A., S.R. and B.S. acknowledge funding from the
1157 project COMFORT (grant agreement no. 820989) under the European Union’s Horizon 2020 research and innovation
1158 programme, and from the EC Horizon Europe project OptimESM “Optimal High Resolution Earth System Models for
1159 Exploring Future Climate Changes”, grant 101081193 and UKRI grant 10039429, CANARI (NE/W004984/1),
1160 UKRI/NERC HighLight Topic Projects “Interacting ice Sheet and Ocean Tipping - Indicators, Processes, Impacts and
1161 Challenges (ISOTIPIC)”, grant NE/Y503320/1, and “Consequences of Arctic Warming for European Climate and Extreme
1162 Weather” (ArctiCONNECT, NE/V004875/1). Y.A. and S.R. also acknowledge funding support from the UK NERC projects
1163 LTS-M BIOPOLE (NE/W004933/1). The work of Y.A. is also funded by the UKRI (grant 10038003) as part of the EPOC
1164 project (Explaining and Predicting the Ocean Conveyor; grant number: 101059547). Views and opinions expressed are
1165 however those of the author(s) only and do not necessarily reflect those of the European Union. Neither the European Union
1166 nor the granting authority can be held responsible for them. Y.A. and S.R. acknowledge the use of the ARCHER UK
1167 National Supercomputing and JASMIN. For the use of the data from the NEMO-MEDUSA 1/4 degree high resolution
1168 model simulations in this paper we acknowledge authorship of Drs Andrew Coward, Andrew Yool, Katya Popova and
1169 Stephen Kelly from the National Oceanography Centre UK (NOC). D.S. acknowledges funding from the project TipESM,
1170 EU grant 10111337673 and Advanced Research + Invention Agency (ARIA)’s UK projects SORTED and AdvanTIP
1171 (SCOP-PR01-P003 - Advancing Tipping Point Early Warning AdvanTip). For the EU projects the work reflects only the
1172 authors’ view; the European Commission and their executive agency are not responsible for any use that may be made of the
1173 information the work contains. C.M.C. acknowledges the financial support from FAPESP (grants 2018/15123-4 and
1174 2019/24349-9) and CNPq (grant 305285/2025-4). HD acknowledges financial support from the European Research Council
1175 through the ERC-AdG project TAOC (project 101055096). J.F.D. acknowledges financial support from the European
1176 Research Council Advanced Grant project ERA (Earth Resilience in the Anthropocene, ERC-2016-ADG-743080) and by
1177 the project CHANGES funded by the German Federal Ministry for Education and Research (BMBF) within the framework
1178 ‘PIK_Change’ under grant 01LS2001A. M.H.E. acknowledges support from the Australian Research Council (ARC), Grant
1179 Nos. SR200100008 and DP190100494. A.V.F. acknowledges support from the ARCHANGE project of the “Make our
1180 planet great again” program (ANR-18-MPGA-0001), France, and from National Aeronautics and Space Administration
1181 (80NSSC21K0558), National Oceanic and Atmospheric Administration (NA20OAR4310377), National Science Foundation
1182 (AGS-205309), and Department of Energy (DE-SC0023134, DE-SC0024186), USA. LJ was supported by the Met Office
1183 Hadley Centre Climate Programme funded by DSIT and by TiPES. This project is TiPES contribution no. 262, and this
1184 project has received funding from the European Union’s Horizon 2020 research and innovation programme under grant

1185 agreement no. 820970. GM acknowledges funding from the Swedish Research Council Vetenskapsrådet (proj. No.
 1186 2022-06599). T.T acknowledges funding from the Department of Science and Technology, India through the ‘INSPIRE
 1187 Faculty Fellowship’. This is ClimTip contribution 40, the ClimTip project has received funding from the European Union's
 1188 Horizon Europe research and innovation programme under grant agreement No. 101137601: Funded by the European
 1189 Union. Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the
 1190 European Union or the European Climate, Infrastructure and Environment Executive Agency (CINEA). Neither the
 1191 European Union nor the granting authority can be held responsible for them. JBS was supported by OCEAN ICE, which is
 1192 funded by the European Union, Horizon Europe Funding Programme for research and innovation under grant agreement Nr.
 1193 101060452.

1194 References

- 1195 Abernathey, R. P., Cerovecki, I., Holland, P. R., Newsom, E., Mazloff, M., and Talley, L. D.: Water-mass transformation
 1196 by sea ice in the upper branch of the Southern Ocean overturning, *Nature Geosci.*, 9, 596–601,
 1197 <https://doi.org/10.1038/ngeo2749>, 2016.
- 1198 Akabane, T. K., Chiessi, C. M., Hirota, M., Bouimetarhan, I., Prange, M., Mulitza, S., Bertassoli Jr, D. J., Häggi, C.,
 1199 Staal, A., Lohmann, G., Boers, N., Daniau, A. L., Oliveira, R. S., Campos, M. C., Shi, X., and De Oliveira, P. E.: Weaker
 1200 Atlantic overturning circulation increases the vulnerability of northern Amazon forests, *Nat. Geosci.*, 17, 1284–1290,
 1201 <https://doi.org/10.1038/s41561-024-01578-z>, 2024.
- 1202 Ali, H., Modi, P., and Mishra, V.: Increased flood risk in Indian sub-continent under the warming climate, *Weather and*
 1203 *Climate Extremes*, 25, 100212, <https://doi.org/10.1016/j.wace.2019.100212>, 2019.
- 1204 Alley, R. B. and Ágústssdóttir, A. M.: The 8k event: cause and consequences of a major Holocene abrupt climate change,
 1205 *Quaternary Science Reviews*, 24, 1123–1149, <https://doi.org/10.1016/j.quascirev.2004.12.004>, 2005.
- 1206 Alley, R. B., Clark, P. U., Keigwin, L. D., and Webb, R. S.: Making Sense of Millennial-Scale Climate Change, in:
 1207 *Mechanisms of Global Climate Change at Millennial Time Scales*, American Geophysical Union (AGU), 385–394,
 1208 <https://doi.org/10.1029/GM112p0385>, 1999.
- 1209 Armstrong McKay, D. I., Staal, A., Abrams, J. F., Winkelmann, R., Sakschewski, B., Loriani, S., Fetzer, I., Cornell, S.
 1210 E., Rockström, J., and Lenton, T. M.: Exceeding 1.5°C global warming could trigger multiple climate tipping points,
 1211 *Science*, 377, eabn7950, <https://doi.org/10.1126/science.abn7950>, 2022.
- 1212 Arnold, N. P. and Randall, D. A.: Global-scale convective aggregation: Implications for the Madden-Julian Oscillation,
 1213 *Journal of Advances in Modeling Earth Systems*, 7, 1499–1518, <https://doi.org/10.1002/2015MS000498>, 2015.

- 1214 Ashwin, P., Wieczorek, S., Vitolo, R., and Cox, P.: Tipping points in open systems: bifurcation, noise-induced and
1215 rate-dependent examples in the climate system, *Phil. Trans. R. Soc. A.*, 370, 1166–1184,
1216 <https://doi.org/10.1098/rsta.2011.0306>, 2012.
- 1217 Bacon, S., Gould, W. J., and Jia, Y.: Open-ocean convection in the Irminger Sea, *Geophysical Research Letters*, 30,
1218 <https://doi.org/10.1029/2002GL016271>, 2003.
- 1219 Bathiany, S., Scheffer, M., van Nes, E. H., Williamson, M. S., and Lenton, T. M.: Abrupt Climate Change in an
1220 Oscillating World, *Sci Rep*, 8, 5040, <https://doi.org/10.1038/s41598-018-23377-4>, 2018.
- 1221 Bellomo, K., Clement, A., Mauritsen, T., Rädcl, G., and Stevens, B.: Simulating the Role of Subtropical Stratocumulus
1222 Clouds in Driving Pacific Climate Variability, *Journal of Climate*, 27, 5119–5131,
1223 <https://doi.org/10.1175/JCLI-D-13-00548.1>, 2014.
- 1224 Bellomo, K., Angeloni, M., Corti, S., and von Hardenberg, J.: Future climate change shaped by inter-model differences
1225 in Atlantic meridional overturning circulation response, *Nat Commun*, 12, 3659,
1226 <https://doi.org/10.1038/s41467-021-24015-w>, 2021.
- 1227 Bellomo, K., Meccia, V. L., D’Agostino, R., Fabiano, F., Larson, S. M., von Hardenberg, J., and Corti, S.: Impacts of a
1228 weakened AMOC on precipitation over the Euro-Atlantic region in the EC-Earth3 climate model, *Clim Dyn*, 61,
1229 3397–3416, <https://doi.org/10.1007/s00382-023-06754-2>, 2023.
- 1230 Ben-Yami, M., Good, P., Jackson, L. C., Crucifix, M., Hu, A., Saenko, O., Swingedouw, D., and Boers, N.: Impacts of
1231 AMOC Collapse on Monsoon Rainfall: A Multi-Model Comparison, *Earth’s Future*, 12, e2023EF003959,
1232 <https://doi.org/10.1029/2023EF003959>, 2024a.
- 1233 Ben-Yami, M., Morr, A., Bathiany, S., and Boers, N.: Uncertainties too large to predict tipping times of major Earth
1234 system components from historical data, *Science Advances*, 10, ead14841, <https://doi.org/10.1126/sciadv.adl4841>, 2024b.
- 1235 Berkelhammer, M., Sinha, A., Stott, L., Cheng, H., Pausata, F. s. r., and Yoshimura, K.: An Abrupt Shift in the Indian
1236 Monsoon 4000 Years Ago, in: *Climates, Landscapes, and Civilizations*, American Geophysical Union (AGU), 75–88,
1237 <https://doi.org/10.1029/2012GM001207>, 2012.
- 1238 Berloff, P. S. and McWilliams, J. C.: Large-Scale, Low-Frequency Variability in Wind-Driven Ocean Gyres, *J. Phys.*
1239 *Oceanogr.*, 29, 1925–1949, [https://doi.org/10.1175/1520-0485\(1999\)029<1925:LSLFVI>2.0.CO;2](https://doi.org/10.1175/1520-0485(1999)029<1925:LSLFVI>2.0.CO;2), 1999.
- 1240 Blackport, R. and Screen, J. A.: Insignificant effect of Arctic amplification on the amplitude of midlatitude atmospheric
1241 waves, *Science Advances*, 6, eaay2880, <https://doi.org/10.1126/sciadv.aay2880>, 2020.
- 1242 Bloch-Johnson, J., Pierrehumbert, R. T., and Abbot, D. S.: Feedback temperature dependence determines the risk of high
1243 warming, *Geophysical Research Letters*, 42, 4973–4980, <https://doi.org/10.1002/2015GL064240>, 2015.

- 1244 Boer, G. J.: A study of atmosphere-ocean predictability on long time scales, *Climate Dynamics*, 16, 469–477,
1245 <https://doi.org/10.1007/s003820050340>, 2000.
- 1246 Boers, N.: Observation-based early-warning signals for a collapse of the Atlantic Meridional Overturning Circulation,
1247 *Nat. Clim. Chang.*, 11, 680–688, <https://doi.org/10.1038/s41558-021-01097-4>, 2021.
- 1248 Boers, N., Marwan, N., Barbosa, H. M. J., and Kurths, J.: A deforestation-induced tipping point for the South American
1249 monsoon system, *Sci Rep*, 7, 41489, <https://doi.org/10.1038/srep41489>, 2017.
- 1250 Boers, N., Ghil, M., and Stocker, T. F.: Theoretical and paleoclimatic evidence for abrupt transitions in the Earth system,
1251 *Environ. Res. Lett.*, 17, 093006, <https://doi.org/10.1088/1748-9326/ac8944>, 2022.
- 1252 Böhm, E., Lippold, J., Gutjahr, M., Frank, M., Blaser, P., Antz, B., Fohlmeister, J., Frank, N., Andersen, M. B., and
1253 Deininger, M.: Strong and deep Atlantic meridional overturning circulation during the last glacial cycle, *Nature*, 517,
1254 73–76, <https://doi.org/10.1038/nature14059>, 2015.
- 1255 Bollasina, M. A., Ming, Y., and Ramaswamy, V.: Anthropogenic Aerosols and the Weakening of the South Asian
1256 Summer Monsoon, *Science*, 334, 502–505, <https://doi.org/10.1126/science.1204994>, 2011.
- 1257 Bonnet, R., Boucher, O., Deshayes, J., Gastineau, G., Hourdin, F., Mignot, J., Servonnat, J., and Swingedouw, D.:
1258 Presentation and Evaluation of the IPSL-CM6A-LR Ensemble of Extended Historical Simulations, *Journal of Advances*
1259 *in Modeling Earth Systems*, 13, e2021MS002565, <https://doi.org/10.1029/2021MS002565>, 2021.
- 1260 Boos, W. R. and Storelvmø, T.: Near-linear response of mean monsoon strength to a broad range of radiative forcings,
1261 *Proceedings of the National Academy of Sciences*, 113, 1510–1515, <https://doi.org/10.1073/pnas.1517143113>, 2016.
- 1262 Borah, P. J., Venugopal, V., Sukhatme, J., Muddebihal, P., and Goswami, B. N.: Indian monsoon derailed by a North
1263 Atlantic wavetrain, *Science*, 370, 1335–1338, <https://doi.org/10.1126/science.aay6043>, 2020.
- 1264 Born, A. and Stocker, T. F.: Two Stable Equilibria of the Atlantic Subpolar Gyre, *Journal of Physical Oceanography*, 44,
1265 246–264, <https://doi.org/10.1175/JPO-D-13-073.1>, 2014.
- 1266 Born, A., Stocker, T. F., and Sandø, A. B.: Transport of salt and freshwater in the Atlantic Subpolar Gyre, *Ocean*
1267 *Dynamics*, 66, 1051–1064, <https://doi.org/10.1007/s10236-016-0970-y>, 2016.
- 1268 Boucher, O., Halloran, P. R., Burke, E. J., Doutriaux-Boucher, M., Jones, C. D., Lowe, J., Ringer, M. A., Robertson, E.,
1269 and Wu, P.: Reversibility in an Earth System model in response to CO₂ concentration changes, *Environ. Res. Lett.*, 7,
1270 024013, <https://doi.org/10.1088/1748-9326/7/2/024013>, 2012.
- 1271 Boulton, C. A., Allison, L. C., and Lenton, T. M.: Early warning signals of Atlantic Meridional Overturning Circulation
1272 collapse in a fully coupled climate model, *Nat Commun*, 5, 5752, <https://doi.org/10.1038/ncomms6752>, 2014.

- 1273 Bower, A., Lozier, S., Biastoch, A., Drouin, K., Foukal, N., Furey, H., Lankhorst, M., R  hs, S., and Zou, S.: Lagrangian
1274 Views of the Pathways of the Atlantic Meridional Overturning Circulation, *Journal of Geophysical Research: Oceans*,
1275 124, 5313–5335, <https://doi.org/10.1029/2019JC015014>, 2019.
- 1276 Broecker, W. S., Denton, G. H., Edwards, R. L., Cheng, H., Alley, R. B., and Putnam, A. E.: Putting the Younger Dryas
1277 cold event into context, *Quaternary Science Reviews*, 29, 1078–1081, <https://doi.org/10.1016/j.quascirev.2010.02.019>,
1278 2010.
- 1279 Bronselaer, B., Winton, M., Griffies, S. M., Hurlin, W. J., Rodgers, K. B., Sergienko, O. V., Stouffer, R. J., and Russell,
1280 J. L.: Change in future climate due to Antarctic meltwater, *Nature*, 564, 53–58,
1281 <https://doi.org/10.1038/s41586-018-0712-z>, 2018.
- 1282 Brovkin, V., Brook, E., Williams, J. W., Bathiany, S., Lenton, T. M., Barton, M., DeConto, R. M., Donges, J. F.,
1283 Ganopolski, A., McManus, J., Praetorius, S., de Vernal, A., Abe-Ouchi, A., Cheng, H., Claussen, M., Crucifix, M.,
1284 Gallop  n, G., Iglesias, V., Kaufman, D. S., Kleinen, T., Lambert, F., van der Leeuw, S., Liddy, H., Loutre, M.-F., McGee,
1285 D., Rehfeld, K., Rhodes, R., Seddon, A. W. R., Trauth, M. H., Vanderveken, L., and Yu, Z.: Past abrupt changes, tipping
1286 points and cascading impacts in the Earth system, *Nat. Geosci.*, 14, 550–558,
1287 <https://doi.org/10.1038/s41561-021-00790-5>, 2021.
- 1288 Brunetti, M., Kasparian, J., and V  rard, C.: Co-existing climate attractors in a coupled aquaplanet, *Clim Dyn*, 53,
1289 6293–6308, <https://doi.org/10.1007/s00382-019-04926-7>, 2019.
- 1290 Buckley, M. W. and Marshall, J.: Observations, inferences, and mechanisms of the Atlantic Meridional Overturning
1291 Circulation: A review, *Reviews of Geophysics*, 54, 5–63, <https://doi.org/10.1002/2015RG000493>, 2016.
- 1292 Bulgin, C. E., Mecking, J. V., Harvey, B. J., Jevrejeva, S., McCarroll, N. F., Merchant, C. J., and Sinha, B.: Dynamic
1293 sea-level changes and potential implications for storm surges in the UK: a storylines perspective, *Environ. Res. Lett.*, 18,
1294 044033, <https://doi.org/10.1088/1748-9326/acc6df>, 2023.
- 1295 Caballero, R. and Carlson, H.: Surface Superrotation, *Journal of the Atmospheric Sciences*, 75, 3671–3689,
1296 <https://doi.org/10.1175/JAS-D-18-0076.1>, 2018.
- 1297 Caballero, R. and Huber, M.: Spontaneous transition to superrotation in warm climates simulated by CAM3,
1298 *Geophysical Research Letters*, 37, <https://doi.org/10.1029/2010GL043468>, 2010.
- 1299 Caballero, R. and Huber, M.: State-dependent climate sensitivity in past warm climates and its implications for future
1300 climate projections, *Proceedings of the National Academy of Sciences*, 110, 14162–14167,
1301 <https://doi.org/10.1073/pnas.1303365110>, 2013.

- 1302 Caesar, L., Rahmstorf, S., Robinson, A., Feulner, G., and Saba, V.: Observed fingerprint of a weakening Atlantic Ocean
1303 overturning circulation, *Nature*, 556, 191–196, <https://doi.org/10.1038/s41586-018-0006-5>, 2018.
- 1304 Caesar, L., McCarthy, G. D., Thornalley, D. J. R., Cahill, N., and Rahmstorf, S.: Current Atlantic Meridional
1305 Overturning Circulation weakest in last millennium, *Nat. Geosci.*, 14, 118–120,
1306 <https://doi.org/10.1038/s41561-021-00699-z>, 2021.
- 1307 Caesar, L., McCarthy, G. D., Thornalley, D. J. R., Cahill, N., and Rahmstorf, S.: Reply to: Atlantic circulation change
1308 still uncertain, *Nat. Geosci.*, 15, 168–170, <https://doi.org/10.1038/s41561-022-00897-3>, 2022.
- 1309 Cai, B., Edwards, R. L., Cheng, H., Tan, M., Wang, X., and Liu, T.: A dry episode during the Younger Dryas and
1310 centennial-scale weak monsoon events during the early Holocene: A high-resolution stalagmite record from southeast of
1311 the Loess Plateau, China, *Geophysical Research Letters*, 35, <https://doi.org/10.1029/2007GL030986>, 2008.
- 1312 Cai, W., Wang, G., Dewitte, B., Wu, L., Santoso, A., Takahashi, K., Yang, Y., Carréric, A., and McPhaden, M. J.:
1313 Increased variability of eastern Pacific El Niño under greenhouse warming, *Nature*, 564, 201–206,
1314 <https://doi.org/10.1038/s41586-018-0776-9>, 2018.
- 1315 Cai, W., Santoso, A., Collins, M., Dewitte, B., Karamperidou, C., Kug, J.-S., Lengaigne, M., McPhaden, M. J., Stuecker,
1316 M. F., Taschetto, A. S., Timmermann, A., Wu, L., Yeh, S.-W., Wang, G., Ng, B., Jia, F., Yang, Y., Ying, J., Zheng, X.-T.,
1317 Bayr, T., Brown, J. R., Capotondi, A., Cobb, K. M., Gan, B., Geng, T., Ham, Y.-G., Jin, F.-F., Jo, H.-S., Li, X., Lin, X.,
1318 McGregor, S., Park, J.-H., Stein, K., Yang, K., Zhang, L., and Zhong, W.: Changing El Niño–Southern Oscillation in a
1319 warming climate, *Nat Rev Earth Environ*, 2, 628–644, <https://doi.org/10.1038/s43017-021-00199-z>, 2021.
- 1320 Cai, W., Ng, B., Wang, G., Santoso, A., Wu, L., and Yang, K.: Increased ENSO sea surface temperature variability under
1321 four IPCC emission scenarios, *Nat. Clim. Chang.*, 12, 228–231, <https://doi.org/10.1038/s41558-022-01282-z>, 2022.
- 1322 Callahan, C. W. and Mankin, J. S.: Persistent effect of El Niño on global economic growth, *Science*, 380, 1064–1069,
1323 <https://doi.org/10.1126/science.adf2983>, 2023.
- 1324 Campos, M. C., Chiessi, C. M., Prange, M., Mulitza, S., Kuhnert, H., Paul, A., Venancio, I. M., Albuquerque, A. L. S.,
1325 Cruz, F. W., and Bahr, A.: A new mechanism for millennial scale positive precipitation anomalies over tropical South
1326 America, *Quaternary Science Reviews*, 225, 105990, <https://doi.org/10.1016/j.quascirev.2019.105990>, 2019.
- 1327 Cao, J., Wang, H., Wang, B., Zhao, H., Wang, C., and Zhu, X.: Higher Sensitivity of Northern Hemisphere Monsoon to
1328 Anthropogenic Aerosol Than Greenhouse Gases, *Geophysical Research Letters*, 49, e2022GL100270,
1329 <https://doi.org/10.1029/2022GL100270>, 2022.
- 1330 Capotondi, A., Wittenberg, A. T., Newman, M., Lorenzo, E. D., Yu, J.-Y., Braconnot, P., Cole, J., Dewitte, B., Giese, B.,
1331 Guilyardi, E., Jin, F.-F., Karnauskas, K., Kirtman, B., Lee, T., Schneider, N., Xue, Y., and Yeh, S.-W.: Understanding

- 1332 ENSO Diversity, *Bulletin of the American Meteorological Society*, 96, 921–938,
1333 <https://doi.org/10.1175/BAMS-D-13-00117.1>, 2015.
- 1334 Carlson, A. E.: PALEOCLIMATE | The Younger Dryas Climate Event, in: *Encyclopedia of Quaternary Science* (Second
1335 Edition), edited by: Elias, S. A. and Mock, C. J., Elsevier, Amsterdam, 126–134,
1336 <https://doi.org/10.1016/B978-0-444-53643-3.00029-7>, 2013.
- 1337 Carlson, H. and Caballero, R.: Enhanced MJO and transition to superrotation in warm climates, *Journal of Advances in*
1338 *Modeling Earth Systems*, 8, 304–318, <https://doi.org/10.1002/2015MS000615>, 2016.
- 1339 Carvalho, L. M. V., Jones, C., Posadas, A. N. D., Quiroz, R., Bookhagen, B., and Liebmann, B.: Precipitation
1340 Characteristics of the South American Monsoon System Derived from Multiple Datasets, *Journal of Climate*, 25,
1341 4600–4620, <https://doi.org/10.1175/JCLI-D-11-00335.1>, 2012.
- 1342 de Carvalho, L. M. V. and Cavalcanti, I. F. A.: The South American Monsoon System (SAMS), in: *The Monsoons and*
1343 *Climate Change: Observations and Modeling*, edited by: de Carvalho, L. M. V. and Jones, C., Springer International
1344 Publishing, Cham, 121–148, https://doi.org/10.1007/978-3-319-21650-8_6, 2016.
- 1345 Chandan, D. and Peltier, W. R.: African Humid Period Precipitation Sustained by Robust Vegetation, Soil, and Lake
1346 Feedbacks, *Geophysical Research Letters*, 47, e2020GL088728, <https://doi.org/10.1029/2020GL088728>, 2020.
- 1347 Chang, P., Zhang, S., Danabasoglu, G., Yeager, S. G., Fu, H., Wang, H., Castruccio, F. S., Chen, Y., Edwards, J., Fu, D.,
1348 Jia, Y., Laurindo, L. C., Liu, X., Rosenbloom, N., Small, R. J., Xu, G., Zeng, Y., Zhang, Q., Bacmeister, J., Bailey, D. A.,
1349 Duan, X., DuVivier, A. K., Li, D., Li, Y., Neale, R., Stössel, A., Wang, L., Zhuang, Y., Baker, A., Bates, S., Dennis, J.,
1350 Diao, X., Gan, B., Gopal, A., Jia, D., Jing, Z., Ma, X., Saravanan, R., Strand, W. G., Tao, J., Yang, H., Wang, X., Wei, Z.,
1351 and Wu, L.: An Unprecedented Set of High-Resolution Earth System Simulations for Understanding Multiscale
1352 Interactions in Climate Variability and Change, *Journal of Advances in Modeling Earth Systems*, 12, e2020MS002298,
1353 <https://doi.org/10.1029/2020MS002298>, 2020.
- 1354 Charney, J., Stone, P. H., and Quirk, W. J.: Drought in the Sahara: A Biogeophysical Feedback Mechanism, *Science*,
1355 187, 434–435, <https://doi.org/10.1126/science.187.4175.434>, 1975.
- 1356 Charney, J. G.: Dynamics of deserts and drought in the Sahel, *Quarterly Journal of the Royal Meteorological Society*,
1357 101, 193–202, <https://doi.org/10.1002/qj.49710142802>, 1975.
- 1358 Chen, Z., Zhou, T., Zhang, L., Chen, X., Zhang, W., and Jiang, J.: Global Land Monsoon Precipitation Changes in
1359 CMIP6 Projections, *Geophysical Research Letters*, 47, e2019GL086902, <https://doi.org/10.1029/2019GL086902>, 2020.

- 1360 Cherchi, A., Terray, P., Ratna, S. B., Sankar, S., Sooraj, K. P., and Behera, S.: Chapter 8 - Indian Ocean Dipole influence
 1361 on Indian summer monsoon and ENSO: A review, in: Indian Summer Monsoon Variability, edited by: Chowdary, J.,
 1362 Parekh, A., and Gnanaseelan, C., Elsevier, 157–182, <https://doi.org/10.1016/B978-0-12-822402-1.00011-9>, 2021.
- 1363 Chevuturi, A., Klingaman, N. P., Turner, A. G., and Hannah, S.: Projected Changes in the Asian-Australian Monsoon
 1364 Region in 1.5°C and 2.0°C Global-Warming Scenarios, *Earth's Future*, 6, 339–358,
 1365 <https://doi.org/10.1002/2017EF000734>, 2018.
- 1366 Chiessi, C. M., Mulitza, S., Pätzold, J., Wefer, G., and Marengo, J. A.: Possible impact of the Atlantic Multidecadal
 1367 Oscillation on the South American summer monsoon, *Geophysical Research Letters*, 36,
 1368 <https://doi.org/10.1029/2009GL039914>, 2009.
- 1369 Claret, M., Galbraith, E. D., Palter, J. B., Bianchi, D., Fennel, K., Gilbert, D., and Dunne, J. P.: Rapid coastal
 1370 deoxygenation due to ocean circulation shift in the northwest Atlantic, *Nature Clim Change*, 8, 868–872,
 1371 <https://doi.org/10.1038/s41558-018-0263-1>, 2018.
- 1372 Collins, J. A., Prange, M., Caley, T., Gimeno, L., Beckmann, B., Mulitza, S., Skonieczny, C., Roche, D., and Schefuß,
 1373 E.: Rapid termination of the African Humid Period triggered by northern high-latitude cooling, *Nat Commun*, 8, 1372,
 1374 <https://doi.org/10.1038/s41467-017-01454-y>, 2017.
- 1375 Coumou, D., Lehmann, J., and Beckmann, J.: The weakening summer circulation in the Northern Hemisphere
 1376 mid-latitudes, *Science*, 348, 324–327, <https://doi.org/10.1126/science.1261768>, 2015.
- 1377 Coumou, D., Di Capua, G., Vavrus, S., Wang, L., and Wang, S.: The influence of Arctic amplification on mid-latitude
 1378 summer circulation, *Nat Commun*, 9, 2959, <https://doi.org/10.1038/s41467-018-05256-8>, 2018.
- 1379 Cruz, F. W., Burns, S. J., Karmann, I., Sharp, W. D., Vuille, M., Cardoso, A. O., Ferrari, J. A., Silva Dias, P. L., and
 1380 Viana, O.: Insolation-driven changes in atmospheric circulation over the past 116,000 years in subtropical Brazil, *Nature*,
 1381 434, 63–66, <https://doi.org/10.1038/nature03365>, 2005.
- 1382 Curtis, P. E., Ceppi, P., and Zappa, G.: Role of the mean state for the Southern Hemispheric jet stream response to CO₂
 1383 forcing in CMIP6 models, *Environ. Res. Lett.*, 15, 064011, <https://doi.org/10.1088/1748-9326/ab8331>, 2020.
- 1384 Czaja, A., Frankignoul, C., Minobe, S., and Vannière, B.: Simulating the Midlatitude Atmospheric Circulation: What
 1385 Might We Gain From High-Resolution Modeling of Air-Sea Interactions?, *Curr Clim Change Rep*, 5, 390–406,
 1386 <https://doi.org/10.1007/s40641-019-00148-5>, 2019.
- 1387 Dai, A., Luo, D., Song, M., and Liu, J.: Arctic amplification is caused by sea-ice loss under increasing CO₂, *Nat*
 1388 *Commun*, 10, 121, <https://doi.org/10.1038/s41467-018-07954-9>, 2019.

1389 Dallmeyer, A., Claussen, M., Lorenz, S. J., Sigl, M., Toohey, M., and Herzschuh, U.: Holocene vegetation transitions
1390 and their climatic drivers in MPI-ESM1.2, *Climate of the Past*, 17, 2481–2513, <https://doi.org/10.5194/cp-17-2481-2021>,
1391 2021.

1392 deMenocal, P., Ortiz, J., Guilderson, T., Adkins, J., Sarnthein, M., Baker, L., and Yarusinsky, M.: Abrupt onset and
1393 termination of the African Humid Period:: rapid climate responses to gradual insolation forcing, *Quaternary Science*
1394 *Reviews*, 19, 347–361, [https://doi.org/10.1016/S0277-3791\(99\)00081-5](https://doi.org/10.1016/S0277-3791(99)00081-5), 2000.

1395 Dijkstra, H. A.: *Nonlinear climate dynamics*, Cambridge university press, Cambridge, 2013.

1396 Dima, M. and Lohmann, G.: Evidence for Two Distinct Modes of Large-Scale Ocean Circulation Changes over the Last
1397 Century, *Journal of Climate*, 23, 5–16, <https://doi.org/10.1175/2009JCLI2867.1>, 2010.

1398 DiNezio, P. N., Clement, A. C., Vecchi, G. A., Soden, B. J., Kirtman, B. P., and Lee, S.-K.: Climate Response of the
1399 Equatorial Pacific to Global Warming, *Journal of Climate*, 22, 4873–4892, <https://doi.org/10.1175/2009JCLI2982.1>,
1400 2009.

1401 Ditlevsen, P. and Ditlevsen, S.: Warning of a forthcoming collapse of the Atlantic meridional overturning circulation, *Nat*
1402 *Commun*, 14, 4254, <https://doi.org/10.1038/s41467-023-39810-w>, 2023.

1403 Dong, Y., Pauling, A. G., Sadai, S., and Armour, K. C.: Antarctic Ice-Sheet Meltwater Reduces Transient Warming and
1404 Climate Sensitivity Through the Sea-Surface Temperature Pattern Effect, *Geophysical Research Letters*, 49,
1405 e2022GL101249, <https://doi.org/10.1029/2022GL101249>, 2022.

1406 Douville, H., Raghavan, K., Renwick, J., Allan, R. P., Arias, P. A., Barlow, M., Cerezo-Mota, R., Cherchi, A., Gan, T. Y.,
1407 Gergis, J., Jiang, D., Khan, A., Pokam Mba, W., Rosenfeld, D., Tierney, J., and Zolina, O.: Chapter 6: Water Cycle
1408 Changes, in: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth*
1409 *Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, 2021.

1410 Drijfhout, S., Oldenborgh, G. J. van, and Cimadoribus, A.: Is a Decline of AMOC Causing the Warming Hole above the
1411 North Atlantic in Observed and Modeled Warming Patterns?, *Journal of Climate*, 25, 8373–8379,
1412 <https://doi.org/10.1175/JCLI-D-12-00490.1>, 2012.

1413 Drijfhout, S., Gleeson, E., Dijkstra, H. A., and Livina, V.: Spontaneous abrupt climate change due to an atmospheric
1414 blocking–sea-ice–ocean feedback in an unforced climate model simulation, *Proceedings of the National Academy of*
1415 *Sciences*, 110, 19713–19718, <https://doi.org/10.1073/pnas.1304912110>, 2013.

1416 Drijfhout, S., Bathiany, S., Beaulieu, C., Brovkin, V., Claussen, M., Huntingford, C., Scheffer, M., Sgubin, G., and
1417 Swingedouw, D.: Catalogue of abrupt shifts in Intergovernmental Panel on Climate Change climate models, *Proceedings*
1418 *of the National Academy of Sciences*, 112, E5777–E5786, <https://doi.org/10.1073/pnas.1511451112>, 2015.

1419 Druckenmiller, M. L., Moon, T. A., Thoman, R. L., Ballinger, T. J., Berner, L. T., Bernhard, G. H., Bhatt, U. S., Bjerke,
 1420 J. W., Box, J. E., Brown, R., Cappelen, J., Christiansen, H. H., Decharme, B., Derksen, C., Divine, D., Drozdov, D. S.,
 1421 Chereque, A. E., Epstein, H. E., Farquharson, L. M., Farrell, S. L., Fausto, R. S., Fettweis, X., Fioletov, V. E., Forbes, B.
 1422 C., Frost, G. V., Gargulinski, E., Gerland, S., Goetz, S. J., Grabinski, Z., Grooß, J.-U., Haas, C., Hanna, E.,
 1423 Hanssen-Bauer, I., Hendricks, S., Holmes, R. M., Ialongo, I., Isaksen, K., Jain, P., Johnsen, B., Kaleschke, L., Kholodov,
 1424 A. L., Kim, S.-J., Korsgaard, N. J., Labe, Z., Lakkala, K., Lara, M. J., Loomis, B., Luoju, K., Macander, M. J.,
 1425 Malkova, G. V., Mankoff, K. D., Manney, G. L., McClelland, J. W., Meier, W. N., Mote, T., Mudryk, L., Müller, R.,
 1426 Nyland, K. E., Overland, J. E., Park, T., Pavlova, O., Perovich, D., Petty, A., Phoenix, G. K., Raynolds, M. K., Reijmer,
 1427 C. H., Richter-Menge, J., Ricker, R., Romanovsky, V. E., Scott, L., Shapiro, H., Shiklomanov, A. I., Shiklomanov, N. I.,
 1428 Smeets, C. J. P. P., Smith, S. L., Soja, A., Spencer, R. G. M., Starkweather, S., Streletskiy, D. A., Suslova, A., Svendby,
 1429 T., Tank, S. E., Tedesco, M., Tian-Kunze, X., Timmermans, M.-L., Tømmervik, H., Tretiakov, M., Tschudi, M.,
 1430 Vakhutinsky, S., As, D. van, Wal, R. S. W. van de, Veraverbeke, S., Walker, D. A., Walsh, J. E., Wang, M., Webster, M.,
 1431 Winton, Ø., Wood, K., York, A., et al.: The Arctic, *Bulletin of the American Meteorological Society*, 102, S263–S316,
 1432 <https://doi.org/10.1175/BAMS-D-21-0086.1>, 2021.

1433 Dukhovskoy, D. S., Yashayaev, I., Chassignet, E. P., Myers, P. G., Platov, G., and Proshutinsky, A.: Time Scales of the
 1434 Greenland Freshwater Anomaly in the Subpolar North Atlantic, *Journal of Climate*, 34, 8971–8987,
 1435 <https://doi.org/10.1175/JCLI-D-20-0610.1>, 2021.

1436 Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled
 1437 Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geoscientific Model
 1438 Development*, 9, 1937–1958, <https://doi.org/10.5194/gmd-9-1937-2016>, 2016.

1439 Faranda, D., Messori, G., Jezequel, A., Vrac, M., and Yiou, P.: Atmospheric circulation compounds anthropogenic
 1440 warming and impacts of climate extremes in Europe, *Proceedings of the National Academy of Sciences*, 120,
 1441 e2214525120, <https://doi.org/10.1073/pnas.2214525120>, 2023.

1442 Fedorov, A. V., Dekens, P. S., McCarthy, M., Ravelo, A. C., deMenocal, P. B., Barreiro, M., Pacanowski, R. C., and
 1443 Philander, S. G.: The Pliocene Paradox (Mechanisms for a Permanent El Niño), *Science*, 312, 1485–1489,
 1444 <https://doi.org/10.1126/science.1122666>, 2006.

1445 Fedorov, A. V., Brierley, C. M., Lawrence, K. T., Liu, Z., Dekens, P. S., and Ravelo, A. C.: Patterns and mechanisms of
 1446 early Pliocene warmth, *Nature*, 496, 43–49, <https://doi.org/10.1038/nature12003>, 2013.

1447 Fedorov, A. V., Burls, N. J., Lawrence, K. T., and Peterson, L. C.: Tightly linked zonal and meridional sea surface
 1448 temperature gradients over the past five million years, *Nature Geosci*, 8, 975–980, <https://doi.org/10.1038/ngeo2577>,
 1449 2015.

- 1450 Fedorov, A. V., Hu, S., Wittenberg, A. T., Levine, A. F. Z., and Deser, C.: ENSO Low-Frequency Modulation and Mean
1451 State Interactions, in: El Niño Southern Oscillation in a Changing Climate, American Geophysical Union (AGU),
1452 173–198, <https://doi.org/10.1002/9781119548164.ch8>, 2020.
- 1453 Feingold, G., Koren, I., Yamaguchi, T., and Kazil, J.: On the reversibility of transitions between closed and open cellular
1454 convection, *Atmospheric Chemistry and Physics*, 15, 7351–7367, <https://doi.org/10.5194/acp-15-7351-2015>, 2015.
- 1455 Feudel, U.: Rate-induced tipping in ecosystems and climate: the role of unstable states, basin boundaries and transient
1456 dynamics, *Nonlin. Processes Geophys.*, 30, 481–502, <https://doi.org/10.5194/npg-30-481-2023>, 2023.
- 1457 Feulner, G., Rahmstorf, S., Levermann, A., and Volkwardt, S.: On the Origin of the Surface Air Temperature Difference
1458 between the Hemispheres in Earth’s Present-Day Climate, *Journal of Climate*, 26, 7136–7150,
1459 <https://doi.org/10.1175/JCLI-D-12-00636.1>, 2013.
- 1460 Florindo-López, C., Bacon, S., Aksenov, Y., Chafik, L., Colbourne, E., and Holliday, N. P.: Arctic Ocean and Hudson
1461 Bay Freshwater Exports: New Estimates from Seven Decades of Hydrographic Surveys on the Labrador Shelf, *Journal*
1462 *of Climate*, 33, 8849–8868, <https://doi.org/10.1175/JCLI-D-19-0083.1>, 2020.
- 1463 Fontela, M., Pérez, F. F., Mercier, H., and Lherminier, P.: North Atlantic Western Boundary Currents Are Intense
1464 Dissolved Organic Carbon Streams, *Frontiers in Marine Science*, 7, 2020.
- 1465 Fox-Kemper, B., Hewitt, H. T., Xiao, C., Aðalgeirsdóttir, G., Drijfhout, S. S., Edwards, T. L., Golledge, N. R., Hemer,
1466 M., Kopp, R. E., Krinner, G., Mix, A., Notz, D., Nowicki, S., Nurhati, I. S., Ruiz, L., Sallée, J.-B., Slangen, A. B. A.,
1467 and Yu, Y.: Chapter 9: Ocean, Cryosphere and Sea Level Change, in: *Climate Change 2021: The Physical Science Basis.*
1468 *Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*,
1469 edited by: Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb,
1470 L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, O.,
1471 Yu, R., and Zhou, B., Cambridge University Press, 2021.
- 1472 Frajka-Williams, E., Ansorge, I. J., Baehr, J., Bryden, H. L., Chidichimo, M. P., Cunningham, S. A., Danabasoglu, G.,
1473 Dong, S., Donohue, K. A., Elipot, S., Heimbach, P., Holliday, N. P., Hummels, R., Jackson, L. C., Karstensen, J.,
1474 Lankhorst, M., Le Bras, I. A., Lozier, M. S., McDonagh, E. L., Meinen, C. S., Mercier, H., Moat, B. I., Perez, R. C.,
1475 Piecuch, C. G., Rhein, M., Srokosz, M. A., Trenberth, K. E., Bacon, S., Forget, G., Goni, G., Kieke, D., Koelling, J.,
1476 Lamont, T., McCarthy, G. D., Mertens, C., Send, U., Smeed, D. A., Speich, S., van den Berg, M., Volkov, D., and
1477 Wilson, C.: Atlantic Meridional Overturning Circulation: Observed Transport and Variability, *Frontiers in Marine*
1478 *Science*, 6, 2019.
- 1479 Francis, J. A. and Vavrus, S. J.: Evidence for a wavier jet stream in response to rapid Arctic warming, *Environ. Res.*
1480 *Lett.*, 10, 014005, <https://doi.org/10.1088/1748-9326/10/1/014005>, 2015.

- 1481 Fröb, F., Olsen, A., Becker, M., Chafik, L., Johannessen, T., Reverdin, G., and Omar, A.: Wintertime fCO₂ Variability in
1482 the Subpolar North Atlantic Since 2004, *Geophysical Research Letters*, 46, 1580–1590,
1483 <https://doi.org/10.1029/2018GL080554>, 2019.
- 1484 Gadgil, S.: The monsoon system: Land–sea breeze or the ITCZ?, *J Earth Syst Sci*, 127, 1,
1485 <https://doi.org/10.1007/s12040-017-0916-x>, 2018.
- 1486 Galaasen, E. V., Ninnemann, U. S., Irvah, N., Kleiven, H. (Kikki) F., Rosenthal, Y., Kissel, C., and Hodell, D. A.: Rapid
1487 Reductions in North Atlantic Deep Water During the Peak of the Last Interglacial Period, *Science*, 343, 1129–1132,
1488 <https://doi.org/10.1126/science.1248667>, 2014.
- 1489 García-Ibáñez, M. I., Bates, N. R., Bakker, D. C. E., Fontela, M., and Velo, A.: Cold-water corals in the Subpolar North
1490 Atlantic Ocean exposed to aragonite undersaturation if the 2 °C global warming target is not met, *Global and Planetary*
1491 *Change*, 201, 103480, <https://doi.org/10.1016/j.gloplacha.2021.103480>, 2021.
- 1492 Geen, R., Bordoni, S., Battisti, D. S., and Hui, K.: Monsoons, ITCZs, and the Concept of the Global Monsoon, *Reviews*
1493 *of Geophysics*, 58, e2020RG000700, <https://doi.org/10.1029/2020RG000700>, 2020.
- 1494 Gent, P. R.: A commentary on the Atlantic meridional overturning circulation stability in climate models, *Ocean*
1495 *Modelling*, 122, 57–66, <https://doi.org/10.1016/j.ocemod.2017.12.006>, 2018.
- 1496 Giannini, A., Salack, S., Lodoun, T., Ali, A., Gaye, A. T., and Ndiaye, O.: A unifying view of climate change in the
1497 Sahel linking intra-seasonal, interannual and longer time scales, *Environ. Res. Lett.*, 8, 024010,
1498 <https://doi.org/10.1088/1748-9326/8/2/024010>, 2013.
- 1499 Gottschalk, J., Skinner, L. C., Lippold, J., Vogel, H., Frank, N., Jaccard, S. L., and Waelbroeck, C.: Biological and
1500 physical controls in the Southern Ocean on past millennial-scale atmospheric CO₂ changes, *Nat Commun*, 7, 11539,
1501 <https://doi.org/10.1038/ncomms11539>, 2016.
- 1502 Grothe, P. R., Cobb, K. M., Liguori, G., Di Lorenzo, E., Capotondi, A., Lu, Y., Cheng, H., Edwards, R. L., Southon, J.
1503 R., Santos, G. M., Deocampo, D. M., Lynch-Stieglitz, J., Chen, T., Sayani, H. R., Thompson, D. M., Conroy, J. L.,
1504 Moore, A. L., Townsend, K., Hagos, M., O'Connor, G., and Toth, L. T.: Enhanced El Niño–Southern Oscillation
1505 Variability in Recent Decades, *Geophysical Research Letters*, 47, e2019GL083906,
1506 <https://doi.org/10.1029/2019GL083906>, 2020.
- 1507 Gunn, K. L., Rintoul, S. R., England, M. H., and Bowen, M. M.: Recent reduced abyssal overturning and ventilation in
1508 the Australian Antarctic Basin, *Nat. Clim. Chang.*, 13, 537–544, <https://doi.org/10.1038/s41558-023-01667-8>, 2023.
- 1509 Gupta, A. K., Anderson, D. M., and Overpeck, J. T.: Abrupt changes in the Asian southwest monsoon during the
1510 Holocene and their links to the North Atlantic Ocean, *Nature*, 421, 354–357, <https://doi.org/10.1038/nature01340>, 2003.

1511 Haine, T. W. N., Curry, B., Gerdes, R., Hansen, E., Karcher, M., Lee, C., Rudels, B., Spreen, G., de Steur, L., Stewart, K.
 1512 D., and Woodgate, R.: Arctic freshwater export: Status, mechanisms, and prospects, *Global and Planetary Change*, 125,
 1513 13–35, <https://doi.org/10.1016/j.gloplacha.2014.11.013>, 2015.

1514 Halloran, P. R., Booth, B. B. B., Jones, C. D., Lambert, F. H., McNeall, D. J., Totterdell, I. J., and Völker, C.: The
 1515 mechanisms of North Atlantic CO₂ uptake in a large Earth System Model ensemble, *Biogeosciences*, 12, 4497–4508,
 1516 <https://doi.org/10.5194/bg-12-4497-2015>, 2015.

1517 Hawkins, E., Smith, R. S., Allison, L. C., Gregory, J. M., Woollings, T. J., Pohlmann, H., and de Cuevas, B.: Bistability
 1518 of the Atlantic overturning circulation in a global climate model and links to ocean freshwater transport, *Geophysical*
 1519 *Research Letters*, 38, <https://doi.org/10.1029/2011GL047208>, 2011.

1520 Hayes, C. T., Martínez-García, A., Hasenfratz, A. P., Jaccard, S. L., Hodell, D. A., Sigman, D. M., Haug, G. H., and
 1521 Anderson, R. F.: A stagnation event in the deep South Atlantic during the last interglacial period, *Science*, 346,
 1522 1514–1517, <https://doi.org/10.1126/science.1256620>, 2014.

1523 Haywood, J. M., Jones, A., Bellouin, N., and Stephenson, D.: Asymmetric forcing from stratospheric aerosols impacts
 1524 Sahelian rainfall, *Nature Clim Change*, 3, 660–665, <https://doi.org/10.1038/nclimate1857>, 2013.

1525 Heede, U. K. and Fedorov, A. V.: Eastern equatorial Pacific warming delayed by aerosols and thermostat response to
 1526 CO₂ increase, *Nat. Clim. Chang.*, 11, 696–703, <https://doi.org/10.1038/s41558-021-01101-x>, 2021.

1527 Heede, U. K. and Fedorov, A. V.: Colder Eastern Equatorial Pacific and Stronger Walker Circulation in the Early 21st
 1528 Century: Separating the Forced Response to Global Warming From Natural Variability, *Geophysical Research Letters*,
 1529 50, e2022GL101020, <https://doi.org/10.1029/2022GL101020>, 2023a.

1530 Heede, U. K. and Fedorov, A. V.: Towards understanding the robust strengthening of ENSO and more frequent extreme
 1531 El Niño events in CMIP6 global warming simulations, *Clim Dyn*, 61, 3047–3060,
 1532 <https://doi.org/10.1007/s00382-023-06856-x>, 2023b.

1533 Heinze, C., Blenckner, T., Martins, H., Rusiecka, D., Döscher, R., Gehlen, M., Gruber, N., Holland, E., Hov, Ø., Joos, F.,
 1534 Matthews, J. B. R., Rødven, R., and Wilson, S.: The quiet crossing of ocean tipping points, *Proceedings of the National*
 1535 *Academy of Sciences*, 118, e2008478118, <https://doi.org/10.1073/pnas.2008478118>, 2021.

1536 Hellmer, H. H., Kauker, F., Timmermann, R., Determann, J., and Rae, J.: Twenty-first-century warming of a large
 1537 Antarctic ice-shelf cavity by a redirected coastal current, *Nature*, 485, 225–228, <https://doi.org/10.1038/nature11064>,
 1538 2012.

1539 Hellmer, H. H., Kauker, F., Timmermann, R., and Hattermann, T.: The Fate of the Southern Weddell Sea Continental
 1540 Shelf in a Warming Climate, *Journal of Climate*, 30, 4337–4350, <https://doi.org/10.1175/JCLI-D-16-0420.1>, 2017.

1541 Henson, S. A., Laufkötter, C., Leung, S., Giering, S. L. C., Palevsky, H. I., and Cavan, E. L.: Uncertain response of
 1542 ocean biological carbon export in a changing world, *Nat. Geosci.*, 15, 248–254,
 1543 <https://doi.org/10.1038/s41561-022-00927-0>, 2022.

1544 Heuzé, C.: North Atlantic deep water formation and AMOC in CMIP5 models, *Ocean Science*, 13, 609–622,
 1545 <https://doi.org/10.5194/os-13-609-2017>, 2017.

1546 Heuzé, C.: Antarctic Bottom Water and North Atlantic Deep Water in CMIP6 models, *Ocean Science*, 17, 59–90,
 1547 <https://doi.org/10.5194/os-17-59-2021>, 2021a.

1548 Heuzé, C.: Antarctic Bottom Water and North Atlantic Deep Water in CMIP6 models, *Ocean Science*, 17, 59–90,
 1549 <https://doi.org/10.5194/os-17-59-2021>, 2021b.

1550 Heuzé, C., Heywood, K. J., Stevens, D. P., and Ridley, J. K.: Changes in Global Ocean Bottom Properties and Volume
 1551 Transports in CMIP5 Models under Climate Change Scenarios, *Journal of Climate*, 28, 2917–2944,
 1552 <https://doi.org/10.1175/JCLI-D-14-00381.1>, 2015.

1553 Holland, P. R. and Kwok, R.: Wind-driven trends in Antarctic sea-ice drift, *Nature Geosci*, 5, 872–875,
 1554 <https://doi.org/10.1038/ngeo1627>, 2012.

1555 Holliday, N. P., Bersch, M., Berx, B., Chafik, L., Cunningham, S., Florindo-López, C., Hátún, H., Johns, W., Josey, S.
 1556 A., Larsen, K. M. H., Mulet, S., Oltmanns, M., Reverdin, G., Rossby, T., Thierry, V., Valdimarsson, H., and Yashayaev,
 1557 I.: Ocean circulation causes the largest freshening event for 120 years in eastern subpolar North Atlantic, *Nat Commun*,
 1558 11, 585, <https://doi.org/10.1038/s41467-020-14474-y>, 2020.

1559 Hopcroft, P. O. and Valdes, P. J.: Paleoclimate-conditioning reveals a North Africa land-atmosphere tipping point, *Proc*
 1560 *Natl Acad Sci U S A*, 118, e2108783118, <https://doi.org/10.1073/pnas.2108783118>, 2021.

1561 Horton, D. E., Johnson, N. C., Singh, D., Swain, D. L., Rajaratnam, B., and Diffenbaugh, N. S.: Contribution of changes
 1562 in atmospheric circulation patterns to extreme temperature trends, *Nature*, 522, 465–469,
 1563 <https://doi.org/10.1038/nature14550>, 2015.

1564 Hou, A., Bahr, A., Raddatz, J., Voigt, S., Greule, M., Albuquerque, A. L., Chiessi, C. M., and Friedrich, O.: Insolation
 1565 and Greenhouse Gas Forcing of the South American Monsoon System Across Three Glacial-Interglacial Cycles,
 1566 *Geophysical Research Letters*, 47, e2020GL087948, <https://doi.org/10.1029/2020GL087948>, 2020.

1567 Hrudya, P. H., Varikoden, H., and Vishnu, R.: A review on the Indian summer monsoon rainfall, variability and its
 1568 association with ENSO and IOD, *Meteorol Atmos Phys*, 133, 1–14, <https://doi.org/10.1007/s00703-020-00734-5>, 2021.

1569 Hsu, P., Li, T., Luo, J.-J., Murakami, H., Kitoh, A., and Zhao, M.: Increase of global monsoon area and precipitation
 1570 under global warming: A robust signal?, *Geophysical Research Letters*, 39, <https://doi.org/10.1029/2012GL051037>,
 1571 2012.

1572 Hsu, P., Li, T., Murakami, H., and Kitoh, A.: Future change of the global monsoon revealed from 19 CMIP5 models,
 1573 *Journal of Geophysical Research: Atmospheres*, 118, 1247–1260, <https://doi.org/10.1002/jgrd.50145>, 2013.

1574 Hu, S. and Fedorov, A. V.: The extreme El Niño of 2015–2016 and the end of global warming hiatus, *Geophysical*
 1575 *Research Letters*, 44, 3816–3824, <https://doi.org/10.1002/2017GL072908>, 2017.

1576 Huang, H., Gutzjahr, M., Eisenhauer, A., and Kuhn, G.: No detectable Weddell Sea Antarctic Bottom Water export during
 1577 the Last and Penultimate Glacial Maximum, *Nat Commun*, 11, 424, <https://doi.org/10.1038/s41467-020-14302-3>, 2020.

1578 Jaccard, S. L., Galbraith, E. D., Martínez-García, A., and Anderson, R. F.: Covariation of deep Southern Ocean
 1579 oxygenation and atmospheric CO₂ through the last ice age, *Nature*, 530, 207–210, <https://doi.org/10.1038/nature16514>,
 1580 2016.

1581 Jackson, L. C.: Shutdown and recovery of the AMOC in a coupled global climate model: The role of the advective
 1582 feedback, *Geophysical Research Letters*, 40, 1182–1188, <https://doi.org/10.1002/grl.50289>, 2013.

1583 Jackson, L. C. and Wood, R. A.: Hysteresis and Resilience of the AMOC in an Eddy-Permitting GCM, *Geophysical*
 1584 *Research Letters*, 45, 8547–8556, <https://doi.org/10.1029/2018GL078104>, 2018.

1585 Jackson, L. C., Kahana, R., Graham, T., Ringer, M. A., Woollings, T., Mecking, J. V., and Wood, R. A.: Global and
 1586 European climate impacts of a slowdown of the AMOC in a high resolution GCM, *Clim Dyn*, 45, 3299–3316,
 1587 <https://doi.org/10.1007/s00382-015-2540-2>, 2015.

1588 Jackson, L. C., Alastrué de Asenjo, E., Bellomo, K., Danabasoglu, G., Haak, H., Hu, A., Jungclaus, J., Lee, W., Meccia,
 1589 V. L., Saenko, O., Shao, A., and Swingedouw, D.: Understanding AMOC stability: the North Atlantic Hosing Model
 1590 Intercomparison Project, *Geoscientific Model Development*, 16, 1975–1995, <https://doi.org/10.5194/gmd-16-1975-2023>,
 1591 2023.

1592 Jin, Q. and Wang, C.: A revival of Indian summer monsoon rainfall since 2002, *Nature Clim Change*, 7, 587–594,
 1593 <https://doi.org/10.1038/nclimate3348>, 2017.

1594 Jones, C. and Carvalho, L. M. V.: Climate Change in the South American Monsoon System: Present Climate and CMIP5
 1595 Projections, *Journal of Climate*, 26, 6660–6678, <https://doi.org/10.1175/JCLI-D-12-00412.1>, 2013.

1596 Jones, C., Liddicoat, S., and Wiltshire, A.: MOHC UKESM1.0-LL model output prepared for CMIP6 CDRMIP
 1597 esm-ssp534-over, <https://doi.org/10.22033/ESGF/CMIP6.12203>, 2020.

1598 Kageyama, M., Merkel, U., Otto-Bliesner, B., Prange, M., Abe-Ouchi, A., Lohmann, G., Ohgaito, R., Roche, D. M.,
 1599 Singarayer, J., Swingedouw, D., and X Zhang: Climatic impacts of fresh water hosing under Last Glacial Maximum
 1600 conditions: a multi-model study, *Climate of the Past*, 9, 935–953, <https://doi.org/10.5194/cp-9-935-2013>, 2013.

1601 Kan, A. van: Phase Transitions in Anisotropic Turbulence, <https://doi.org/10.48550/arXiv.2408.02844>, 25 January 2025.

1602 Kan, A. van and Alexakis, A.: Critical transition in fast-rotating turbulence within highly elongated domains, *Journal of*
 1603 *Fluid Mechanics*, 899, A33, <https://doi.org/10.1017/jfm.2020.443>, 2020.

1604 Katzenberger, A. and Levermann, A.: Consistent increase in East Asian Summer Monsoon rainfall and its variability
 1605 under climate change over China in CMIP6, *Earth System Dynamics*, 15, 1137–1151,
 1606 <https://doi.org/10.5194/esd-15-1137-2024>, 2024.

1607 Katzenberger, A., Schewe, J., Pongratz, J., and Levermann, A.: Robust increase of Indian monsoon rainfall and its
 1608 variability under future warming in CMIP6 models, *Earth System Dynamics*, 12, 367–386,
 1609 <https://doi.org/10.5194/esd-12-367-2021>, 2021.

1610 Kelly, S. J., Popova, E., Aksenov, Y., Marsh, R., and Yool, A.: They Came From the Pacific: How Changing Arctic
 1611 Currents Could Contribute to an Ecological Regime Shift in the Atlantic Ocean, *Earth’s Future*, 8, e2019EF001394,
 1612 <https://doi.org/10.1029/2019EF001394>, 2020.

1613 Kennedy, D., Parker, T., Woollings, T., Harvey, B., and Shaffrey, L.: The response of high-impact blocking weather
 1614 systems to climate change, *Geophysical Research Letters*, 43, 7250–7258, <https://doi.org/10.1002/2016GL069725>, 2016.

1615 Kilbourne, K. H., Wanamaker, A. D., Moffa-Sanchez, P., Reynolds, D. J., Amrhein, D. E., Butler, P. G., Gebbie, G.,
 1616 Goes, M., Jansen, M. F., Little, C. M., Mette, M., Moreno-Chamarro, E., Ortega, P., Otto-Bliesner, B. L., Rossby, T.,
 1617 Scourse, J., and Whitney, N. M.: Atlantic circulation change still uncertain, *Nat. Geosci.*, 15, 165–167,
 1618 <https://doi.org/10.1038/s41561-022-00896-4>, 2022.

1619 Klose, A. K., Wunderling, N., Winkelmann, R., and Donges, J. F.: What do we mean, ‘tipping cascade’?, *Environ. Res.*
 1620 *Lett.*, 16, 125011, <https://doi.org/10.1088/1748-9326/ac3955>, 2021.

1621 Koelling, J., Atamanchuk, D., Karstensen, J., Handmann, P., and Wallace, D. W. R.: Oxygen export to the deep ocean
 1622 following Labrador Sea Water formation, *Biogeosciences*, 19, 437–454, <https://doi.org/10.5194/bg-19-437-2022>, 2022.

1623 Konare, A., Zakey, A. S., Solmon, F., Giorgi, F., Rauscher, S., Ibrah, S., and Bi, X.: A regional climate modeling study of
 1624 the effect of desert dust on the West African monsoon, *Journal of Geophysical Research: Atmospheres*, 113,
 1625 <https://doi.org/10.1029/2007JD009322>, 2008.

1626 Kornhuber, K. and Tamarin-Brodsky, T.: Future Changes in Northern Hemisphere Summer Weather Persistence Linked
 1627 to Projected Arctic Warming, *Geophysical Research Letters*, 48, e2020GL091603,
 1628 <https://doi.org/10.1029/2020GL091603>, 2021.

1629 Kriegler, E., Hall, J. W., Held, H., Dawson, R., and Schellnhuber, H. J.: Imprecise probability assessment of tipping
 1630 points in the climate system, *Proceedings of the National Academy of Sciences*, 106, 5041–5046,
 1631 <https://doi.org/10.1073/pnas.0809117106>, 2009.

1632 Kröpelin, S., Verschuren, D., Lézine, A.-M., Eggermont, H., Cocquyt, C., Francus, P., Cazet, J.-P., Fagot, M., Rumes, B.,
 1633 Russell, J. M., Darius, F., Conley, D. J., Schuster, M., von Suchodoletz, H., and Engstrom, D. R.: Climate-Driven
 1634 Ecosystem Succession in the Sahara: The Past 6000 Years, *Science*, 320, 765–768,
 1635 <https://doi.org/10.1126/science.1154913>, 2008.

1636 Kucharski, F., Zeng, N., and Kalnay, E.: A further assessment of vegetation feedback on decadal Sahel rainfall
 1637 variability, *Clim Dyn*, 40, 1453–1466, <https://doi.org/10.1007/s00382-012-1397-x>, 2013.

1638 Kuhlbrodt, T., Titz, S., Feudel, U., and Rahmstorf, S.: A simple model of seasonal open ocean convection, *Ocean
 1639 Dynamics*, 52, 36–49, <https://doi.org/10.1007/s10236-001-8175-3>, 2001.

1640 Kumar, S. K. and Seshadri, A. K.: Origins and suppression of bifurcation phenomena in lower-order monsoon models,
 1641 *Earth System Dynamics Discussions*, 1–19, <https://doi.org/10.5194/esd-2022-30>, 2022.

1642 Kwiatkowski, L., Torres, O., Bopp, L., Aumont, O., Chamberlain, M., Christian, J. R., Dunne, J. P., Gehlen, M., Ilyina,
 1643 T., John, J. G., Lenton, A., Li, H., Lovenduski, N. S., Orr, J. C., Palmieri, J., Santana-Falcón, Y., Schwinger, J., Séférian,
 1644 R., Stock, C. A., Tagliabue, A., Takano, Y., Tjiputra, J., Toyama, K., Tsujino, H., Watanabe, M., Yamamoto, A., Yool, A.,
 1645 and Ziehn, T.: Twenty-first century ocean warming, acidification, deoxygenation, and upper-ocean nutrient and primary
 1646 production decline from CMIP6 model projections, *Biogeosciences*, 17, 3439–3470,
 1647 <https://doi.org/10.5194/bg-17-3439-2020>, 2020.

1648 Lago, V. and England, M. H.: Projected Slowdown of Antarctic Bottom Water Formation in Response to Amplified
 1649 Meltwater Contributions, *Journal of Climate*, 32, 6319–6335, <https://doi.org/10.1175/JCLI-D-18-0622.1>, 2019.

1650 Latif, M., Sun, J., Visbeck, M., and Hadi Bordbar, M.: Natural variability has dominated Atlantic Meridional
 1651 Overturning Circulation since 1900, *Nat. Clim. Chang.*, 12, 455–460, <https://doi.org/10.1038/s41558-022-01342-4>, 2022.

1652 Lawman, A. E., Di Nezio, P. N., Partin, J. W., Dee, S. G., Thirumalai, K., and Quinn, T. M.: Unraveling forced responses
 1653 of extreme El Niño variability over the Holocene, *Science Advances*, 8, eabm4313,
 1654 <https://doi.org/10.1126/sciadv.abm4313>, 2022.

- 1655 Leconte, J., Forget, F., Charnay, B., Wordsworth, R., and Pottier, A.: Increased insolation threshold for runaway
1656 greenhouse processes on Earth-like planets, *Nature*, 504, 268–271, <https://doi.org/10.1038/nature12827>, 2013.
- 1657 Lee, J.-Y. and Wang, B.: Future change of global monsoon in the CMIP5, *Clim Dyn*, 42, 101–119,
1658 <https://doi.org/10.1007/s00382-012-1564-0>, 2014.
- 1659 Lehner, F., Born, A., Raible, C. C., and Stocker, T. F.: Amplified Inception of European Little Ice Age by Sea
1660 Ice–Ocean–Atmosphere Feedbacks, *Journal of Climate*, 26, 7586–7602, <https://doi.org/10.1175/JCLI-D-12-00690.1>,
1661 2013.
- 1662 Lenton, T. M., Held, H., Kriegler, E., Hall, J. W., Lucht, W., Rahmstorf, S., and Schellnhuber, H. J.: Tipping elements in
1663 the Earth’s climate system, *Proceedings of the National Academy of Sciences*, 105, 1786–1793,
1664 <https://doi.org/10.1073/pnas.0705414105>, 2008.
- 1665 Levermann, A. and Born, A.: Bistability of the Atlantic subpolar gyre in a coarse-resolution climate model, *Geophysical*
1666 *Research Letters*, 34, <https://doi.org/10.1029/2007GL031732>, 2007.
- 1667 Levermann, A., Schewe, J., Petoukhov, V., and Held, H.: Basic mechanism for abrupt monsoon transitions, *Proceedings*
1668 *of the National Academy of Sciences*, 106, 20572–20577, <https://doi.org/10.1073/pnas.0901414106>, 2009a.
- 1669 Levermann, A., Schewe, J., Petoukhov, V., and Held, H.: Basic mechanism for abrupt monsoon transitions, *Proceedings*
1670 *of the National Academy of Sciences*, 106, 20572–20577, <https://doi.org/10.1073/pnas.0901414106>, 2009b.
- 1671 Lewis, S. C., LeGrande, A. N., Kelley, M., and Schmidt, G. A.: Water vapour source impacts on oxygen isotope
1672 variability in tropical precipitation during Heinrich events, *Clim. Past*, 6, 325–343,
1673 <https://doi.org/10.5194/cp-6-325-2010>, 2010.
- 1674 L’Heureux, M. L., Tippett, M. K., Kumar, A., Butler, A. H., Ciaso, L. M., Ding, Q., Harnos, K. J., and Johnson, N. C.:
1675 Strong Relations Between ENSO and the Arctic Oscillation in the North American Multimodel Ensemble, *Geophysical*
1676 *Research Letters*, 44, 11,654–11,662, <https://doi.org/10.1002/2017GL074854>, 2017.
- 1677 Li, Q., England, M. H., Hogg, A. M., Rintoul, S. R., and Morrison, A. K.: Abyssal ocean overturning slowdown and
1678 warming driven by Antarctic meltwater, *Nature*, 615, 841–847, <https://doi.org/10.1038/s41586-023-05762-w>, 2023.
- 1679 Liebmann, B. and Mechoso, C. R.: The south american monsoon system, in: *The Global Monsoon System*, vol. Volume
1680 5, WORLD SCIENTIFIC, 137–157, https://doi.org/10.1142/9789814343411_0009, 2011.
- 1681 Lin, P., Pickart, R. S., Heorton, H., Tsamados, M., Itoh, M., and Kikuchi, T.: Recent state transition of the Arctic Ocean’s
1682 Beaufort Gyre, *Nat. Geosci.*, 16, 485–491, <https://doi.org/10.1038/s41561-023-01184-5>, 2023.

- 1683 Liu, W., Xie, S.-P., Liu, Z., and Zhu, J.: Overlooked possibility of a collapsed Atlantic Meridional Overturning
1684 Circulation in warming climate, *Science Advances*, 3, e1601666, <https://doi.org/10.1126/sciadv.1601666>, 2017.
- 1685 Liu, W., Fedorov, A. V., Xie, S.-P., and Hu, S.: Climate impacts of a weakened Atlantic Meridional Overturning
1686 Circulation in a warming climate, *Science Advances*, 6, eaaz4876, <https://doi.org/10.1126/sciadv.aaz4876>, 2020.
- 1687 Lloret, F. and Batllori, E.: Climate-Induced Global Forest Shifts due to Heatwave-Drought, in: *Ecosystem Collapse and
1688 Climate Change*, edited by: Canadell, J. G. and Jackson, R. B., Springer International Publishing, Cham, 155–186,
1689 https://doi.org/10.1007/978-3-030-71330-0_7, 2021.
- 1690 Lobelle, D., Beaulieu, C., Livina, V., Sévellec, F., and Frajka-Williams, E.: Detectability of an AMOC Decline in Current
1691 and Projected Climate Changes, *Geophysical Research Letters*, 47, e2020GL089974,
1692 <https://doi.org/10.1029/2020GL089974>, 2020.
- 1693 Lohmann, J. and Ditlevsen, P. D.: Risk of tipping the overturning circulation due to increasing rates of ice melt,
1694 *Proceedings of the National Academy of Sciences*, 118, e2017989118, <https://doi.org/10.1073/pnas.2017989118>, 2021.
- 1695 Ma, S. and Zhou, T.: Robust Strengthening and Westward Shift of the Tropical Pacific Walker Circulation during
1696 1979–2012: A Comparison of 7 Sets of Reanalysis Data and 26 CMIP5 Models, *Journal of Climate*, 29, 3097–3118,
1697 <https://doi.org/10.1175/JCLI-D-15-0398.1>, 2016.
- 1698 Marshall, J., Donohoe, A., Ferreira, D., and McGee, D.: The ocean’s role in setting the mean position of the
1699 Inter-Tropical Convergence Zone, *Clim Dyn*, 42, 1967–1979, <https://doi.org/10.1007/s00382-013-1767-z>, 2014.
- 1700 Mastrandrea, M. D., Mach, K. J., Plattner, G.-K., Edenhofer, O., Stocker, T. F., Field, C. B., Ebi, K. L., and Matschoss, P.
1701 R.: The IPCC AR5 guidance note on consistent treatment of uncertainties: a common approach across the working
1702 groups, *Climatic Change*, 108, 675, <https://doi.org/10.1007/s10584-011-0178-6>, 2011.
- 1703 J.B.R. Matthews, Vincent Möller, Renée van Diemen, Jan S. Fuglestad, Valérie Masson-Delmotte, Carlos Méndez,
1704 Sergey Semenov, and Andy Reisinger: IPCC, 2021: Annex VII: Glossary, 2021.
- 1705 Mauritsen, T. and Stevens, B.: Missing iris effect as a possible cause of muted hydrological change and high climate
1706 sensitivity in models, *Nature Geosci*, 8, 346–351, <https://doi.org/10.1038/ngeo2414>, 2015.
- 1707 McGee, D., deMenocal, P. B., Winckler, G., Stuut, J. B. W., and Bradtmiller, L. I.: The magnitude, timing and abruptness
1708 of changes in North African dust deposition over the last 20,000yr, *Earth and Planetary Science Letters*, 371–372,
1709 163–176, <https://doi.org/10.1016/j.epsl.2013.03.054>, 2013.
- 1710 McManus, J. F., Francois, R., Gherardi, J.-M., Keigwin, L. D., and Brown-Leger, S.: Collapse and rapid resumption of
1711 Atlantic meridional circulation linked to deglacial climate changes, *Nature*, 428, 834–837,
1712 <https://doi.org/10.1038/nature02494>, 2004a.

1713 McManus, J. F., Francois, R., Gherardi, J.-M., Keigwin, L. D., and Brown-Leger, S.: Collapse and rapid resumption of
 1714 Atlantic meridional circulation linked to deglacial climate changes, *Nature*, 428, 834–837,
 1715 <https://doi.org/10.1038/nature02494>, 2004b.

1716 McPhaden, M. J. (Ed.): *El Niño southern oscillation in a changing climate*, First edition., Wiley-American Geophysical
 1717 Union, Hoboken, NJ, 2020.

1718 Mecking, J. V., Drijfhout, S. S., Jackson, L. C., and Graham, T.: Stable AMOC off state in an eddy-permitting coupled
 1719 climate model, *Clim Dyn*, 47, 2455–2470, <https://doi.org/10.1007/s00382-016-2975-0>, 2016.

1720 Mecking, J. V., Drijfhout, S. S., Jackson, L. C., and Andrews, M. B.: The effect of model bias on Atlantic freshwater
 1721 transport and implications for AMOC bi-stability, *Tellus A: Dynamic Meteorology and Oceanography*, 69, 1299910,
 1722 <https://doi.org/10.1080/16000870.2017.1299910>, 2017.

1723 Michel, S., Dijkstra, H., Guardamagna, F., Jaques-Dumas, V., van Westen, R., and von der Heydt, A.: Deep learning
 1724 reconstruction of Atlantic Meridional Overturning Circulation strength validates ongoing twenty-first century decline.,
 1725 <https://www.researchsquare.com/article/rs-3377545/v1>, 2023.

1726 Michel, S. L. L., Swingedouw, D., Ortega, P., Gastineau, G., Mignot, J., McCarthy, G., and Khodri, M.: Early warning
 1727 signal for a tipping point suggested by a millennial Atlantic Multidecadal Variability reconstruction, *Nat Commun*, 13,
 1728 5176, <https://doi.org/10.1038/s41467-022-32704-3>, 2022.

1729 Moffa-Sánchez, P., Moreno-Chamarro, E., Reynolds, D. J., Ortega, P., Cunningham, L., Swingedouw, D., Amrhein, D.
 1730 E., Halfar, J., Jonkers, L., Jungclauss, J. H., Perner, K., Wanamaker, A., and Yeager, S.: Variability in the Northern North
 1731 Atlantic and Arctic Oceans Across the Last Two Millennia: A Review, *Paleoceanography and Paleoclimatology*, 34,
 1732 1399–1436, <https://doi.org/10.1029/2018PA003508>, 2019.

1733 Mohtadi, M., Prange, M., Oppo, D. W., De Pol-Holz, R., Merkel, U., Zhang, X., Steinke, S., and Lückge, A.: North
 1734 Atlantic forcing of tropical Indian Ocean climate, *Nature*, 509, 76–80, <https://doi.org/10.1038/nature13196>, 2014.

1735 Mohtadi, M., Prange, M., and Steinke, S.: Palaeoclimatic insights into forcing and response of monsoon rainfall, *Nature*,
 1736 533, 191–199, <https://doi.org/10.1038/nature17450>, 2016.

1737 Molnar, P., Boos, W. R., and Battisti, D. S.: Orographic Controls on Climate and Paleoclimate of Asia: Thermal and
 1738 Mechanical Roles for the Tibetan Plateau, *Annual Review of Earth and Planetary Sciences*, 38, 77–102,
 1739 <https://doi.org/10.1146/annurev-earth-040809-152456>, 2010.

1740 Moon, S. and Ha, K.-J.: Future changes in monsoon duration and precipitation using CMIP6, *npj Clim Atmos Sci*, 3,
 1741 1–7, <https://doi.org/10.1038/s41612-020-00151-w>, 2020.

- 1742 Morrill, C., Overpeck, J. T., and Cole, J. E.: A synthesis of abrupt changes in the Asian summer monsoon since the last
1743 deglaciation, *The Holocene*, 13, 465–476, <https://doi.org/10.1191/0959683603hl639ft>, 2003.
- 1744 Mulitza, S., Chiessi, C. M., Schefuß, E., Lippold, J., Wichmann, D., Antz, B., Mackensen, A., Paul, A., Prange, M.,
1745 Rehfeld, K., Werner, M., Bickert, T., Frank, N., Kuhnert, H., Lynch-Stieglitz, J., Portillo-Ramos, R. C., Sawakuchi, A.
1746 O., Schulz, M., Schwenk, T., Tiedemann, R., Vahlenkamp, M., and Zhang, Y.: Synchronous and proportional deglacial
1747 changes in Atlantic meridional overturning and northeast Brazilian precipitation, *Paleoceanography*, 32, 622–633,
1748 <https://doi.org/10.1002/2017PA003084>, 2017.
- 1749 Muller, C., Yang, D., Craig, G., Cronin, T., Fildier, B., Haerter, J. O., Hohenegger, C., Mapes, B., Randall, D., Shamekh,
1750 S., and Sherwood, S. C.: Spontaneous Aggregation of Convective Storms, *Annual Review of Fluid Mechanics*, 54,
1751 133–157, <https://doi.org/10.1146/annurev-fluid-022421-011319>, 2022.
- 1752 Myers, T. A., Mechoso, C. R., Cesana, G. V., DeFlorio, M. J., and Waliser, D. E.: Cloud Feedback Key to Marine
1753 Heatwave off Baja California, *Geophysical Research Letters*, 45, 4345–4352, <https://doi.org/10.1029/2018GL078242>,
1754 2018.
- 1755 Naughten, K. A., Holland, P. R., and De Rydt, J.: Unavoidable future increase in West Antarctic ice-shelf melting over
1756 the twenty-first century, *Nat. Clim. Chang.*, 13, 1222–1228, <https://doi.org/10.1038/s41558-023-01818-x>, 2023.
- 1757 Neukermans, G., Oziel, L., and Babin, M.: Increased intrusion of warming Atlantic water leads to rapid expansion of
1758 temperate phytoplankton in the Arctic, *Global Change Biology*, 24, 2545–2553, <https://doi.org/10.1111/gcb.14075>, 2018.
- 1759 New, A. L., Smede, D. A., Czaja, A., Blaker, A. T., Mecking, J. V., Mathews, J. P., and Sanchez-Franks, A.: Labrador
1760 Slope Water connects the subarctic with the Gulf Stream, *Environ. Res. Lett.*, 16, 084019,
1761 <https://doi.org/10.1088/1748-9326/ac1293>, 2021.
- 1762 Nian, D., Bathiany, S., Ben-Yami, M., Blaschke, L., Hirota, M., Rodrigues, R., and Boers, N.: The combined impact of
1763 global warming and AMOC collapse on the Amazon Rainforest, <https://www.researchsquare.com/article/rs-2673317/v1>,
1764 31 May 2023.
- 1765 Nicholson, S. E. and Dezfuli, A. K.: The Relationship of Rainfall Variability in Western Equatorial Africa to the Tropical
1766 Oceans and Atmospheric Circulation. Part I: The Boreal Spring, *Journal of Climate*, 26, 45–65,
1767 <https://doi.org/10.1175/JCLI-D-11-00653.1>, 2013.
- 1768 Nowicki, S. M. J., Payne, A., Larour, E., Seroussi, H., Goelzer, H., Lipscomb, W., Gregory, J., Abe-Ouchi, A., and
1769 Shepherd, A.: Ice Sheet Model Intercomparison Project (ISMIP6) contribution to CMIP6, *Geosci. Model Dev.*, 9,
1770 4521–4545, <https://doi.org/10.5194/gmd-9-4521-2016>, 2016.

- 1771 Orihuela-Pinto, B., England, M. H., and Taschetto, A. S.: Interbasin and interhemispheric impacts of a collapsed Atlantic
1772 Overturning Circulation, *Nat. Clim. Chang.*, 12, 558–565, <https://doi.org/10.1038/s41558-022-01380-y>, 2022.
- 1773 Osman, M. B., Das, S. B., Trusel, L. D., Evans, M. J., Fischer, H., Grieman, M. M., Kipfstuhl, S., McConnell, J. R., and
1774 Saltzman, E. S.: Industrial-era decline in subarctic Atlantic productivity, *Nature*, 569, 551–555,
1775 <https://doi.org/10.1038/s41586-019-1181-8>, 2019.
- 1776 Osman, M. B., Coats, S., Das, S. B., McConnell, J. R., and Chellman, N.: North Atlantic jet stream projections in the
1777 context of the past 1,250 years, *Proceedings of the National Academy of Sciences*, 118, e2104105118,
1778 <https://doi.org/10.1073/pnas.2104105118>, 2021.
- 1779 Otterman, J.: Baring High-Albedo Soils by Overgrazing: A Hypothesized Desertification Mechanism, *Science*, 186,
1780 531–533, <https://doi.org/10.1126/science.186.4163.531>, 1974.
- 1781 Oudar, T., Cattiaux, J., and Douville, H.: Drivers of the Northern Extratropical Eddy-Driven Jet Change in CMIP5 and
1782 CMIP6 Models, *Geophysical Research Letters*, 47, e2019GL086695, <https://doi.org/10.1029/2019GL086695>, 2020.
- 1783 Oziel, L., Baudena, A., Ardyna, M., Massicotte, P., Randelhoff, A., Sallée, J.-B., Ingvaldsen, R. B., Devred, E., and
1784 Babin, M.: Faster Atlantic currents drive poleward expansion of temperate phytoplankton in the Arctic Ocean, *Nat*
1785 *Commun*, 11, 1705, <https://doi.org/10.1038/s41467-020-15485-5>, 2020.
- 1786 Paik, S., An, S.-I., Min, S.-K., King, A. D., and Shin, J.: Hysteretic Behavior of Global to Regional Monsoon Area
1787 Under CO₂ Ramp-Up and Ramp-Down, *Earth's Future*, 11, e2022EF003434, <https://doi.org/10.1029/2022EF003434>,
1788 2023.
- 1789 Pausata, F. S. R., Li, C., Wettstein, J. J., Kageyama, M., and Nisancioglu, K. H.: The key role of topography in altering
1790 North Atlantic atmospheric circulation during the last glacial period, *Climate of the Past*, 7, 1089–1101,
1791 <https://doi.org/10.5194/cp-7-1089-2011>, 2011.
- 1792 Pausata, F. S. R., Messori, G., and Zhang, Q.: Impacts of dust reduction on the northward expansion of the African
1793 monsoon during the Green Sahara period, *Earth and Planetary Science Letters*, 434, 298–307,
1794 <https://doi.org/10.1016/j.epsl.2015.11.049>, 2016.
- 1795 Petoukhov, V., Rahmstorf, S., Petri, S., and Schellnhuber, H. J.: Quasiresonant amplification of planetary waves and
1796 recent Northern Hemisphere weather extremes, *Proceedings of the National Academy of Sciences*, 110, 5336–5341,
1797 <https://doi.org/10.1073/pnas.1222000110>, 2013.
- 1798 Philip, S. Y., Kew, S. F., van Oldenborgh, G. J., Anslow, F. S., Seneviratne, S. I., Vautard, R., Coumou, D., Ebi, K. L.,
1799 Arrighi, J., Singh, R., van Aalst, M., Pereira Marghidan, C., Wehner, M., Yang, W., Li, S., Schumacher, D. L., Hauser,
1800 M., Bonnet, R., Luu, L. N., Lehner, F., Gillett, N., Tradowsky, J. S., Vecchi, G. A., Rodell, C., Stull, R. B., Howard, R.,

- 1801 and Otto, F. E. L.: Rapid attribution analysis of the extraordinary heat wave on the Pacific coast of the US and Canada in
1802 June 2021, *Earth System Dynamics*, 13, 1689–1713, <https://doi.org/10.5194/esd-13-1689-2022>, 2022.
- 1803 Previdi, M., Smith, K. L., and Polvani, L. M.: Arctic amplification of climate change: a review of underlying
1804 mechanisms, *Environ. Res. Lett.*, 16, 093003, <https://doi.org/10.1088/1748-9326/ac1c29>, 2021.
- 1805 Purich, A. and England, M. H.: Historical and Future Projected Warming of Antarctic Shelf Bottom Water in CMIP6
1806 Models, *Geophysical Research Letters*, 48, e2021GL092752, <https://doi.org/10.1029/2021GL092752>, 2021.
- 1807 Purich, A. and England, M. H.: Projected Impacts of Antarctic Meltwater Anomalies over the Twenty-First Century,
1808 *Journal of Climate*, 36, 2703–2719, <https://doi.org/10.1175/JCLI-D-22-0457.1>, 2023.
- 1809 Purkey, S. G. and Johnson, G. C.: Antarctic Bottom Water Warming and Freshening: Contributions to Sea Level Rise,
1810 Ocean Freshwater Budgets, and Global Heat Gain, *Journal of Climate*, 26, 6105–6122,
1811 <https://doi.org/10.1175/JCLI-D-12-00834.1>, 2013.
- 1812 Qasmi, S. and Ribes, A.: Reducing uncertainty in local temperature projections, *Science Advances*, 8, eabo6872,
1813 <https://doi.org/10.1126/sciadv.abo6872>, 2022.
- 1814 Rachmayani, R., Prange, M., and Schulz, M.: North African vegetation–precipitation feedback in early and
1815 mid-Holocene climate simulations with CCSM3-DGVM, *Climate of the Past*, 11, 175–185,
1816 <https://doi.org/10.5194/cp-11-175-2015>, 2015.
- 1817 Ragon, C., Lembo, V., Lucarini, V., V  rard, C., Kasparian, J., and Brunetti, M.: Robustness of Competing Climatic
1818 States, *Journal of Climate*, 35, 2769–2784, <https://doi.org/10.1175/JCLI-D-21-0148.1>, 2022.
- 1819 Rahmstorf, S.: A simple model of seasonal open ocean convection, *Ocean Dynamics*, 52, 26–35,
1820 <https://doi.org/10.1007/s10236-001-8174-4>, 2001.
- 1821 Rahmstorf, S.: Ocean circulation and climate during the past 120,000 years, *Nature*, 419, 207–214,
1822 <https://doi.org/10.1038/nature01090>, 2002.
- 1823 Rahmstorf, S., Crucifix, M., Ganopolski, A., Goosse, H., Kamenkovich, I., Knutti, R., Lohmann, G., Marsh, R., Mysak,
1824 L. A., Wang, Z., and Weaver, A. J.: Thermohaline circulation hysteresis: A model intercomparison, *Geophysical*
1825 *Research Letters*, 32, <https://doi.org/10.1029/2005GL023655>, 2005.
- 1826 Rahmstorf, S., Box, J. E., Feulner, G., Mann, M. E., Robinson, A., Rutherford, S., and Schaffernicht, E. J.: Exceptional
1827 twentieth-century slowdown in Atlantic Ocean overturning circulation, *Nature Clim Change*, 5, 475–480,
1828 <https://doi.org/10.1038/nclimate2554>, 2015.

- 1829 Rhein, M., Steinfeldt, R., Kieke, D., Stendardo, I., and Yashayaev, I.: Ventilation variability of Labrador Sea Water and
1830 its impact on oxygen and anthropogenic carbon: a review, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 375, 20160321, <https://doi.org/10.1098/rsta.2016.0321>, 2017.
- 1832 Riboldi, J., Lott, F., D’Andrea, F., and Rivi re, G.: On the Linkage Between Rossby Wave Phase Speed, Atmospheric
1833 Blocking, and Arctic Amplification, *Geophysical Research Letters*, 47, e2020GL087796,
1834 <https://doi.org/10.1029/2020GL087796>, 2020.
- 1835 Richardson, K., Steffen, W., Lucht, W., Bendtsen, J., Cornell, S. E., Donges, J. F., Dr ke, M., Fetzer, I., Bala, G., von
1836 Bloh, W., Feulner, G., Fiedler, S., Gerten, D., Gleeson, T., Hofmann, M., Huiskamp, W., Kumm, M., Mohan, C.,
1837 Nogu s-Bravo, D., Petri, S., Porkka, M., Rahmstorf, S., Schaphoff, S., Thonicke, K., Tobian, A., Virkki, V.,
1838 Wang-Erlandsson, L., Weber, L., and Rockstr m, J.: Earth beyond six of nine planetary boundaries, *Science Advances*,
1839 9, eadh2458, <https://doi.org/10.1126/sciadv.adh2458>, 2023.
- 1840 Ridge, S. M. and McKinley, G. A.: Ocean carbon uptake under aggressive emission mitigation, *Biogeosciences*, 18,
1841 2711–2725, <https://doi.org/10.5194/bg-18-2711-2021>, 2021.
- 1842 Rietkerk, M., Dekker, S. C., de Ruiter, P. C., and van de Koppel, J.: Self-Organized Patchiness and Catastrophic Shifts in
1843 Ecosystems, *Science*, 305, 1926–1929, <https://doi.org/10.1126/science.1101867>, 2004.
- 1844 Rockstr m, J., Gupta, J., Qin, D., Lade, S. J., Abrams, J. F., Andersen, L. S., Armstrong McKay, D. I., Bai, X., Bala, G.,
1845 Bunn, S. E., Ciobanu, D., DeClerck, F., Ebi, K., Gifford, L., Gordon, C., Hasan, S., Kanie, N., Lenton, T. M., Loriani, S.,
1846 Liverman, D. M., Mohamed, A., Nakicenovic, N., Obura, D., Ospina, D., Prodani, K., Rammelt, C., Sakschewski, B.,
1847 Scholtens, J., Stewart-Koster, B., Tharammal, T., van Vuuren, D., Verburg, P. H., Winkelmann, R., Zimm, C., Bennett, E.
1848 M., Bringezu, S., Broadgate, W., Green, P. A., Huang, L., Jacobson, L., Ndehedehe, C., Pedde, S., Rocha, J., Scheffer,
1849 M., Schulte-Uebbing, L., de Vries, W., Xiao, C., Xu, C., Xu, X., Zafra-Calvo, N., and Zhang, X.: Safe and just Earth
1850 system boundaries, *Nature*, 1–10, <https://doi.org/10.1038/s41586-023-06083-8>, 2023.
- 1851 Rodr guez-Fonseca, B., Mohino, E., Mechoso, C. R., Caminade, C., Biasutti, M., Gaetani, M., Garcia-Serrano, J., Vizzy,
1852 E. K., Cook, K., Xue, Y., Polo, I., Losada, T., Druyan, L., Fontaine, B., Bader, J., Doblas-Reyes, F. J., Goddard, L.,
1853 Janicot, S., Arribas, A., Lau, W., Colman, A., Vellinga, M., Rowell, D. P., Kucharski, F., and Voldoire, A.: Variability
1854 and Predictability of West African Droughts: A Review on the Role of Sea Surface Temperature Anomalies, *Journal of*
1855 *Climate*, 28, 4034–4060, <https://doi.org/10.1175/JCLI-D-14-00130.1>, 2015.
- 1856 Rousi, E., Kornhuber, K., Beobide-Arsuaga, G., Luo, F., and Coumou, D.: Accelerated western European heatwave
1857 trends linked to more-persistent double jets over Eurasia, *Nat Commun*, 13, 3851,
1858 <https://doi.org/10.1038/s41467-022-31432-y>, 2022.

1859 Rousseau, D.-D., Bagniewski, W., and Sun, Y.: Detection of abrupt changes in East Asian monsoon from Chinese loess
1860 and speleothem records, *Global and Planetary Change*, 227, 104154, <https://doi.org/10.1016/j.gloplacha.2023.104154>,
1861 2023.

1862 Roxy, M. K., Ritika, K., Terray, P., Murtugudde, R., Ashok, K., and Goswami, B. N.: Drying of Indian subcontinent by
1863 rapid Indian Ocean warming and a weakening land-sea thermal gradient, *Nat Commun*, 6, 7423,
1864 <https://doi.org/10.1038/ncomms8423>, 2015.

1865 Sanchez Goñi, M. F. and Harrison, S. P.: Millennial-scale climate variability and vegetation changes during the Last
1866 Glacial: Concepts and terminology, *Quaternary Science Reviews*, 29, 2823–2827,
1867 <https://doi.org/10.1016/j.quascirev.2009.11.014>, 2010.

1868 Sarinthein, M., Stattegger, K., Dreger, D., Erlenkeuser, H., Grootes, P., Haupt, B. J., Jung, S., Kiefer, T., Kuhnt, W.,
1869 Pflaumann, U., Schäfer-Neth, C., Schulz, H., Schulz, M., Seidov, D., Simstich, J., Kreveld, S. van, Vogelsang, E.,
1870 Völker, A., and Weinelt, M.: Fundamental Modes and Abrupt Changes in North Atlantic Circulation and Climate over
1871 the last 60 ky — Concepts, Reconstruction and Numerical Modeling, in: *The Northern North Atlantic: A Changing*
1872 *Environment*, edited by: Schäfer, P., Ritzrau, W., Schlüter, M., and Thiede, J., Springer, Berlin, Heidelberg, 365–410,
1873 https://doi.org/10.1007/978-3-642-56876-3_21, 2001.

1874 Schewe, J., Levermann, A., and Cheng, H.: A critical humidity threshold for monsoon transitions, *Climate of the Past*, 8,
1875 535–544, <https://doi.org/10.5194/cp-8-535-2012>, 2012.

1876 Schleussner, C.-F., Divine, D. V., Donges, J. F., Miettinen, A., and Donner, R. V.: Indications for a North Atlantic ocean
1877 circulation regime shift at the onset of the Little Ice Age, *Clim Dyn*, 45, 3623–3633,
1878 <https://doi.org/10.1007/s00382-015-2561-x>, 2015.

1879 Schmittner, A., Brook, E. J., and Ahn, J.: Impact of the ocean’s Overturning circulation on atmospheric CO₂, in: *Ocean*
1880 *Circulation: Mechanisms and Impacts—Past and Future Changes of Meridional Overturning*, American Geophysical
1881 Union (AGU), 315–334, <https://doi.org/10.1029/173GM20>, 2007.

1882 Schneider, T., Kaul, C. M., and Pressel, K. G.: Possible climate transitions from breakup of stratocumulus decks under
1883 greenhouse warming, *Nat. Geosci.*, 12, 163–167, <https://doi.org/10.1038/s41561-019-0310-1>, 2019.

1884 Schulz, von Rad, Erlenkeuser, and von Rad: Correlation between Arabian Sea and Greenland climate oscillations of the
1885 past 110,000 years, *Nature*, 393, 54–57, <https://doi.org/10.1038/31750>, 1998.

1886 Schwinger, J., Asaadi, A., Goris, N., and Lee, H.: Possibility for strong northern hemisphere high-latitude cooling under
1887 negative emissions, *Nat Commun*, 13, 1095, <https://doi.org/10.1038/s41467-022-28573-5>, 2022.

- 1888 Screen, J. A. and Simmonds, I.: Exploring links between Arctic amplification and mid-latitude weather, *Geophysical*
1889 *Research Letters*, 40, 959–964, <https://doi.org/10.1002/grl.50174>, 2013.
- 1890 Seager, R., Henderson, N., and Cane, M.: Persistent Discrepancies between Observed and Modeled Trends in the
1891 Tropical Pacific Ocean, *Journal of Climate*, 35, 4571–4584, <https://doi.org/10.1175/JCLI-D-21-0648.1>, 2022.
- 1892 Seeley, J. T. and Wordsworth, R. D.: Episodic deluges in simulated hothouse climates, *Nature*, 599, 74–79,
1893 <https://doi.org/10.1038/s41586-021-03919-z>, 2021.
- 1894 Serreze, M. C. and Meier, W. N.: The Arctic’s sea ice cover: trends, variability, predictability, and comparisons to the
1895 Antarctic, *Annals of the New York Academy of Sciences*, 1436, 36–53, <https://doi.org/10.1111/nyas.13856>, 2019.
- 1896 Seshadri, A. K.: Energetics and monsoon bifurcations, *Clim Dyn*, 48, 561–576,
1897 <https://doi.org/10.1007/s00382-016-3094-7>, 2017.
- 1898 Sgubin, G., Swingedouw, D., Drijfhout, S., Mary, Y., and Bennabi, A.: Abrupt cooling over the North Atlantic in modern
1899 climate models, *Nat Commun*, 8, 14375, <https://doi.org/10.1038/ncomms14375>, 2017.
- 1900 Shanahan, T. M., McKay, N. P., Hughen, K. A., Overpeck, J. T., Otto-Bliesner, B., Heil, C. W., King, J., Scholz, C. A.,
1901 and Peck, J.: The time-transgressive termination of the African Humid Period, *Nature Geosci*, 8, 140–144,
1902 <https://doi.org/10.1038/ngeo2329>, 2015.
- 1903 Shepherd, A., Ivins, E., Rignot, E., Smith, B., van den Broeke, M., Velicogna, I., Whitehouse, P., Briggs, K., Joughin, I.,
1904 Krinner, G., Nowicki, S., Payne, T., Scambos, T., Schlegel, N., A. G., Agosta, C., Ahlstrøm, A., Babonis, G., Barletta, V.
1905 R., Bjørk, A. A., Blazquez, A., Bonin, J., Colgan, W., Csatho, B., Cullather, R., Engdahl, M. E., Felikson, D., Fettweis,
1906 X., Forsberg, R., Hogg, A. E., Gallee, H., Gardner, A., Gilbert, L., Gourmelen, N., Groh, A., Gunter, B., Hanna, E.,
1907 Harig, C., Helm, V., Horvath, A., Horwath, M., Khan, S., Kjeldsen, K. K., Konrad, H., Langen, P. L., Lecavalier, B.,
1908 Loomis, B., Luthcke, S., McMillan, M., Melini, D., Mernild, S., Mohajerani, Y., Moore, P., Mottram, R., Mouginit, J.,
1909 Moyano, G., Muir, A., Nagler, T., Nield, G., Nilsson, J., Noël, B., Otsuka, I., Pattie, M. E., Peltier, W. R., Pie, N.,
1910 Rietbroek, R., Rott, H., Sandberg Sørensen, L., Sasgen, I., Save, H., Scheuchl, B., Schrama, E., Schröder, L., Seo, K.-W.,
1911 Simonsen, S. B., Slater, T., Spada, G., Sutterley, T., Talpe, M., Tarasov, L., van de Berg, W. J., van der Wal, W., van
1912 Wesse, M., Vishwakarma, B. D., Wiese, D., Wilton, D., Wagner, T., Wouters, B., Wuite, J., and The IMBIE Team:
1913 Mass balance of the Greenland Ice Sheet from 1992 to 2018, *Nature*, 579, 233–239,
1914 <https://doi.org/10.1038/s41586-019-1855-2>, 2020.
- 1915 Shepherd, T. G.: Storyline approach to the construction of regional climate change information, *Proceedings of the Royal*
1916 *Society A: Mathematical, Physical and Engineering Sciences*, 475, 20190013, <https://doi.org/10.1098/rspa.2019.0013>,
1917 2019.

- 1918 Sherwood, S. C., Webb, M. J., Annan, J. D., Armour, K. C., Forster, P. M., Hargreaves, J. C., Hegerl, G., Klein, S. A.,
1919 Marvel, K. D., Rohling, E. J., Watanabe, M., Andrews, T., Braconnot, P., Bretherton, C. S., Foster, G. L., Hausfather, Z.,
1920 von der Heydt, A. S., Knutti, R., Mauritsen, T., Norris, J. R., Proistosescu, C., Rugenstein, M., Schmidt, G. A., Tokarska,
1921 K. B., and Zelinka, M. D.: An Assessment of Earth's Climate Sensitivity Using Multiple Lines of Evidence, *Reviews of*
1922 *Geophysics*, 58, e2019RG000678, <https://doi.org/10.1029/2019RG000678>, 2020.
- 1923 Shin, S.-J., Yeh, S.-W., An, S.-I., Keenlyside, N., Xie, S.-P., and Park, J.-H.: Southern Ocean Control of 2°C Global
1924 Warming in Climate Models, *Earth's Future*, 11, e2022EF003212, <https://doi.org/10.1029/2022EF003212>, 2023.
- 1925 Siahayan, A., Smith, R. S., Holland, P. R., Jenkins, A., Gregory, J. M., Lee, V., Mathiot, P., Payne, A. J., Ridley, J. K., and
1926 Jones, C. G.: The Antarctic contribution to 21st-century sea-level rise predicted by the UK Earth System Model with an
1927 interactive ice sheet, *The Cryosphere*, 16, 4053–4086, <https://doi.org/10.5194/tc-16-4053-2022>, 2022.
- 1928 Skinner, L. C., Fallon, S., Waelbroeck, C., Michel, E., and Barker, S.: Ventilation of the Deep Southern Ocean and
1929 Deglacial CO₂ Rise, *Science*, 328, 1147–1151, <https://doi.org/10.1126/science.1183627>, 2010.
- 1930 Solmon, F., Mallet, M., Elguindi, N., Giorgi, F., Zakey, A., and Konaré, A.: Dust aerosol impact on regional precipitation
1931 over western Africa, mechanisms and sensitivity to absorption properties, *Geophysical Research Letters*, 35,
1932 <https://doi.org/10.1029/2008GL035900>, 2008.
- 1933 Son, J.-H., Seo, K.-H., and Wang, B.: Dynamical Control of the Tibetan Plateau on the East Asian Summer Monsoon,
1934 *Geophysical Research Letters*, 46, 7672–7679, <https://doi.org/10.1029/2019GL083104>, 2019.
- 1935 Srokosz, M., Baringer, M., Bryden, H., Cunningham, S., Delworth, T., Lozier, S., Marotzke, J., and Sutton, R.: Past,
1936 Present, and Future Changes in the Atlantic Meridional Overturning Circulation, *Bulletin of the American*
1937 *Meteorological Society*, 93, 1663–1676, <https://doi.org/10.1175/BAMS-D-11-00151.1>, 2012.
- 1938 Stager, J. C., Ryves, D. B., Chase, B. M., and Pausata, F. S. R.: Catastrophic Drought in the Afro-Asian Monsoon Region
1939 During Heinrich Event 1, *Science*, 331, 1299–1302, <https://doi.org/10.1126/science.1198322>, 2011.
- 1940 Steffen, W., Richardson, K., Rockström, J., Cornell, S. E., Fetzer, I., Bennett, E. M., Biggs, R., Carpenter, S. R., de Vries,
1941 W., de Wit, C. A., Folke, C., Gerten, D., Heinke, J., Mace, G. M., Persson, L. M., Ramanathan, V., Reyers, B., and
1942 Sörlin, S.: Planetary boundaries: Guiding human development on a changing planet, *Science*, 347, 1259855,
1943 <https://doi.org/10.1126/science.1259855>, 2015.
- 1944 Stommel, H.: Thermohaline Convection with Two Stable Regimes of Flow, *Tellus*, 13, 224–230,
1945 <https://doi.org/10.1111/j.2153-3490.1961.tb00079.x>, 1961.
- 1946 Sun, Y., Clemens, S. C., Morrill, C., Lin, X., Wang, X., and An, Z.: Influence of Atlantic meridional overturning
1947 circulation on the East Asian winter monsoon, *Nature Geosci*, 5, 46–49, <https://doi.org/10.1038/ngeo1326>, 2012.

- 1948 Swingedouw, D., Ifejika Speranza, C., Bartsch, A., Durand, G., Jamet, C., Beaugrand, G., and Conversi, A.: Early
1949 Warning from Space for a Few Key Tipping Points in Physical, Biological, and Social-Ecological Systems, *Surv*
1950 *Geophys*, 41, 1237–1284, <https://doi.org/10.1007/s10712-020-09604-6>, 2020.
- 1951 Swingedouw, D., Bily, A., Esquerdo, C., Borchert, L. F., Sgubin, G., Mignot, J., and Menary, M.: On the risk of abrupt
1952 changes in the North Atlantic subpolar gyre in CMIP6 models, *Annals of the New York Academy of Sciences*, 1504,
1953 187–201, <https://doi.org/10.1111/nyas.14659>, 2021.
- 1954 Swingedouw, D., Houssais, M.-N., Herbaut, C., Blaizot, A.-C., Devilliers, M., and Deshayes, J.: AMOC Recent and
1955 Future Trends: A Crucial Role for Oceanic Resolution and Greenland Melting?, *Frontiers in Climate*, 4, 2022.
- 1956 Terhaar, J., Torres, O., Bourgeois, T., and Kwiatkowski, L.: Arctic Ocean acidification over the 21st century co-driven by
1957 anthropogenic carbon increases and freshening in the CMIP6 model ensemble, *Biogeosciences*, 18, 2221–2240,
1958 <https://doi.org/10.5194/bg-18-2221-2021>, 2021.
- 1959 Thomas, Z. A., Kwasniok, F., Boulton, C. A., Cox, P. M., Jones, R. T., Lenton, T. M., and Turney, C. S. M.: Early
1960 warnings and missed alarms for abrupt monsoon transitions, *Clim. Past*, 11, 1621–1633,
1961 <https://doi.org/10.5194/cp-11-1621-2015>, 2015.
- 1962 Tierney, J. E., Haywood, A. M., Feng, R., Bhattacharya, T., and Otto-Bliesner, B. L.: Pliocene Warmth Consistent With
1963 Greenhouse Gas Forcing, *Geophysical Research Letters*, 46, 9136–9144, <https://doi.org/10.1029/2019GL083802>, 2019.
- 1964 Timmermann, A., An, S.-I., Kug, J.-S., Jin, F.-F., Cai, W., Capotondi, A., Cobb, K. M., Lengaigne, M., McPhaden, M. J.,
1965 Stuecker, M. F., Stein, K., Wittenberg, A. T., Yun, K.-S., Bayr, T., Chen, H.-C., Chikamoto, Y., Dewitte, B., Dommenget,
1966 D., Grothe, P., Guilyardi, E., Ham, Y.-G., Hayashi, M., Ineson, S., Kang, D., Kim, S., Kim, W., Lee, J.-Y., Li, T., Luo,
1967 J.-J., McGregor, S., Planton, Y., Power, S., Rashid, H., Ren, H.-L., Santoso, A., Takahashi, K., Todd, A., Wang, G.,
1968 Wang, G., Xie, R., Yang, W.-H., Yeh, S.-W., Yoon, J., Zeller, E., and Zhang, X.: El Niño–Southern Oscillation
1969 complexity, *Nature*, 559, 535–545, <https://doi.org/10.1038/s41586-018-0252-6>, 2018.
- 1970 Trenberth, K. E., Stepaniak, D. P., and Caron, J. M.: The Global Monsoon as Seen through the Divergent Atmospheric
1971 Circulation, *Journal of Climate*, 13, 3969–3993,
1972 [https://doi.org/10.1175/1520-0442\(2000\)013<3969:TGMASST>2.0.CO;2](https://doi.org/10.1175/1520-0442(2000)013<3969:TGMASST>2.0.CO;2), 2000.
- 1973 Turney, C. S. M., Fogwill, C. J., Golledge, N. R., McKay, N. P., van Sebille, E., Jones, R. T., Etheridge, D., Rubino, M.,
1974 Thornton, D. P., Davies, S. M., Ramsey, C. B., Thomas, Z. A., Bird, M. I., Munksgaard, N. C., Kohno, M., Woodward,
1975 J., Winter, K., Weyrich, L. S., Rootes, C. M., Millman, H., Albert, P. G., Rivera, A., van Ommen, T., Curran, M., Moy,
1976 A., Rahmstorf, S., Kawamura, K., Hillenbrand, C.-D., Weber, M. E., Manning, C. J., Young, J., and Cooper, A.: Early
1977 Last Interglacial ocean warming drove substantial ice mass loss from Antarctica, *Proceedings of the National Academy*
1978 *of Sciences*, 117, 3996–4006, <https://doi.org/10.1073/pnas.1902469117>, 2020.

- 1979 Tziperman, E. and Farrell, B.: Pliocene equatorial temperature: Lessons from atmospheric superrotation,
1980 *Paleoceanography*, 24, <https://doi.org/10.1029/2008PA001652>, 2009.
- 1981 Van Westen, R. M. and Dijkstra, H. A.: Asymmetry of AMOC Hysteresis in a State-Of-The-Art Global Climate Model,
1982 *Geophysical Research Letters*, 50, e2023GL106088, <https://doi.org/10.1029/2023GL106088>, 2023.
- 1983 Venancio, I. M., Shimizu, M. H., Santos, T. P., Lessa, D. O., Portilho-Ramos, R. C., Chiessi, C. M., Crivellari, S.,
1984 Mulitza, S., Kuhnert, H., Tiedemann, R., Vahlenkamp, M., Bickert, T., Sampaio, G., Albuquerque, A. L. S., Veiga, S.,
1985 Nobre, P., and Nobre, C.: Changes in surface hydrography at the western tropical Atlantic during the Younger Dryas,
1986 *Global and Planetary Change*, 184, 103047, <https://doi.org/10.1016/j.gloplacha.2019.103047>, 2020.
- 1987 Vera, C., Higgins, W., Amador, J., Ambrizzi, T., Garreaud, R., Gochis, D., Gutzler, D., Lettenmaier, D., Marengo, J.,
1988 Mechoso, C. R., Nogues-Paegle, J., Dias, P. L. S., and Zhang, C.: Toward a Unified View of the American Monsoon
1989 Systems, *Journal of Climate*, 19, 4977–5000, <https://doi.org/10.1175/JCLI3896.1>, 2006.
- 1990 Wang, B. and Ding, Q.: Global monsoon: Dominant mode of annual variation in the tropics, *Dynamics of Atmospheres
1991 and Oceans*, 44, 165–183, <https://doi.org/10.1016/j.dynatmoce.2007.05.002>, 2008.
- 1992 Wang, B., Liu, J., Kim, H.-J., Webster, P. J., and Yim, S.-Y.: Recent change of the global monsoon precipitation
1993 (1979–2008), *Clim Dyn*, 39, 1123–1135, <https://doi.org/10.1007/s00382-011-1266-z>, 2012.
- 1994 Wang, B., Jin, C., and Liu, J.: Understanding Future Change of Global Monsoons Projected by CMIP6 Models, *Journal
1995 of Climate*, 33, 6471–6489, <https://doi.org/10.1175/JCLI-D-19-0993.1>, 2020.
- 1996 Wang, B., Biasutti, M., Byrne, M. P., Castro, C., Chang, C.-P., Cook, K., Fu, R., Grimm, A. M., Ha, K.-J., Hendon, H.,
1997 Kitoh, A., Krishnan, R., Lee, J.-Y., Li, J., Liu, J., Moise, A., Pascale, S., Roxy, M. K., Seth, A., Sui, C.-H., Turner, A.,
1998 Yang, S., Yun, K.-S., Zhang, L., and Zhou, T.: Monsoons Climate Change Assessment, *Bulletin of the American
1999 Meteorological Society*, 102, E1–E19, <https://doi.org/10.1175/BAMS-D-19-0335.1>, 2021.
- 2000 Wang, G., Eltahir, E. A. B., Foley, J. A., Pollard, D., and Levis, S.: Decadal variability of rainfall in the Sahel: results
2001 from the coupled GENESIS-IBIS atmosphere-biosphere model, *Climate Dynamics*, 22, 625–637,
2002 <https://doi.org/10.1007/s00382-004-0411-3>, 2004.
- 2003 Wang, S., Foster, A., Lenz, E. A., Kessler, J. D., Stroeve, J. C., Anderson, L. O., Turetsky, M., Betts, R., Zou, S., Liu, W.,
2004 Boos, W. R., and Hausfather, Z.: Mechanisms and Impacts of Earth System Tipping Elements, *Reviews of Geophysics*,
2005 61, e2021RG000757, <https://doi.org/10.1029/2021RG000757>, 2023.
- 2006 Wang, Y., Cheng, H., Edwards, R. L., Kong, X., Shao, X., Chen, S., Wu, J., Jiang, X., Wang, X., and An, Z.: Millennial-
2007 and orbital-scale changes in the East Asian monsoon over the past 224,000 years, *Nature*, 451, 1090–1093,
2008 <https://doi.org/10.1038/nature06692>, 2008.

- 2009 Wara, M. W., Ravelo, A. C., and Delaney, M. L.: Permanent El Niño-like conditions during the Pliocene warm period,
2010 Science, 309, 758–761, <https://doi.org/10.1126/science.1112596>, 2005.
- 2011 Weijer, W., Cheng, W., Drijfhout, S. S., Fedorov, A. V., Hu, A., Jackson, L. C., Liu, W., McDonagh, E. L., Mecking, J.
2012 V., and Zhang, J.: Stability of the Atlantic Meridional Overturning Circulation: A Review and Synthesis, Journal of
2013 Geophysical Research: Oceans, 124, 5336–5375, <https://doi.org/10.1029/2019JC015083>, 2019.
- 2014 van Westen, R. M., Kliphuis, M., and Dijkstra, H. A.: Collapse of the Atlantic Meridional Overturning Circulation in a
2015 Strongly Eddying Ocean-Only Model, Geophysical Research Letters, 52, e2024GL114532,
2016 <https://doi.org/10.1029/2024GL114532>, 2025.
- 2017 White, R. H., Hilgenbrink, C., and Sheshadri, A.: The Importance of Greenland in Setting the Northern Preferred
2018 Position of the North Atlantic Eddy-Driven Jet, Geophysical Research Letters, 46, 14126–14134,
2019 <https://doi.org/10.1029/2019GL084780>, 2019.
- 2020 White, R. H., Kornhuber, K., Martius, O., and Wirth, V.: From Atmospheric Waves to Heatwaves: A Waveguide
2021 Perspective for Understanding and Predicting Concurrent, Persistent, and Extreme Extratropical Weather, Bulletin of the
2022 American Meteorological Society, 103, E923–E935, <https://doi.org/10.1175/BAMS-D-21-0170.1>, 2022.
- 2023 Wieners, C. E., Dijkstra, H. A., and de Ruijter, W. P. M.: The interaction between the Western Indian Ocean and ENSO
2024 in CESM, Clim Dyn, 52, 5153–5172, <https://doi.org/10.1007/s00382-018-4438-2>, 2019.
- 2025 Winkelmann, R., Dennis, D. P., Donges, J. F., Loriani, S., Klose, A. K., Abrams, J. F., Alvarez-Solas, J., Albrecht, T.,
2026 Armstrong McKay, D., Bathiany, S., Blasco Navarro, J., Brovkin, V., Burke, E., Danabasoglu, G., Donner, R. V., Drüke,
2027 M., Georgievski, G., Goelzer, H., Harper, A. B., Hegerl, G., Hirota, M., Hu, A., Jackson, L. C., Jones, C., Kim, H.,
2028 Koenigk, T., Lawrence, P., Lenton, T. M., Liddy, H., Licón-Saláiz, J., Menthon, M., Montoya, M., Nitzbon, J., Nowicki,
2029 S., Otto-Bliesner, B., Pausata, F., Rahmstorf, S., Ramin, K., Robinson, A., Rockström, J., Romanou, A., Sakschewski,
2030 B., Schädel, C., Sherwood, S., Smith, R. S., Steinert, N. J., Swingedouw, D., Willeit, M., Weijer, W., Wood, R., Wyser,
2031 K., and Yang, S.: The Tipping Points Modelling Intercomparison Project (TIPMIP): Assessing tipping point risks in the
2032 Earth system, EGUsphere, 1–52, <https://doi.org/10.5194/egusphere-2025-1899>, 2025.
- 2033 Wills, R. C. J., Dong, Y., Proistosescu, C., Armour, K. C., and Battisti, D. S.: Systematic Climate Model Biases in the
2034 Large-Scale Patterns of Recent Sea-Surface Temperature and Sea-Level Pressure Change, Geophysical Research Letters,
2035 49, e2022GL100011, <https://doi.org/10.1029/2022GL100011>, 2022.
- 2036 Wirth, V. and Polster, C.: The Problem of Diagnosing Jet Waveguidability in the Presence of Large-Amplitude Eddies,
2037 Journal of the Atmospheric Sciences, 78, 3137–3151, <https://doi.org/10.1175/JAS-D-20-0292.1>, 2021.

- 2038 Wood, R. A., Rodríguez, J. M., Smith, R. S., Jackson, L. C., and Hawkins, E.: Observable, low-order dynamical controls
2039 on thresholds of the Atlantic meridional overturning circulation, *Clim Dyn*, 53, 6815–6834,
2040 <https://doi.org/10.1007/s00382-019-04956-1>, 2019.
- 2041 Wood, R. A., Baker, J. A., Beaugrand, G., Boutin, J., Conversi, A., Donner, R. V., Frenger, I., Goberville, E., Hayashida,
2042 H., Koeve, W., Kvale, K., Landolfi, A., Maslowski, W., Oschlies, A., Romanou, A., Somes, C. J., Stocker, T. F., and
2043 Swingedouw, D.: Opportunities for Earth Observation to Inform Risk Management for Ocean Tipping Points, *Surv*
2044 *Geophys*, <https://doi.org/10.1007/s10712-024-09859-3>, 2024.
- 2045 Wunderling, N., Von Der Heydt, A., Aksenov, Y., Barker, S., Bastiaansen, R., Brovkin, V., Brunetti, M., Couplet, V.,
2046 Kleinen, T., Lear, C. H., Lohmann, J., Roman-Cuesta, R. M., Sinet, S., Swingedouw, D., Winkelmann, R., Anand, P.,
2047 Barichivich, J., Bathiany, S., Baudena, M., Bruun, J. T., Chiessi, C. M., Coxall, H. K., Docquier, D., Donges, J. F.,
2048 Falkena, S. K. J., Klose, A. K., Obura, D., Rocha, J., Rynders, S., Steinert, N. J., and Willeit, M.: Climate tipping point
2049 interactions and cascades: A review, *Climate change/Other interactions/Earth system and climate modeling*,
2050 <https://doi.org/10.5194/egusphere-2023-1576>, 2023.
- 2051 Xie, S.-P., Deser, C., Vecchi, G. A., Ma, J., Teng, H., and Wittenberg, A. T.: Global Warming Pattern Formation: Sea
2052 Surface Temperature and Rainfall, *Journal of Climate*, 23, 966–986, <https://doi.org/10.1175/2009JCLI3329.1>, 2010.
- 2053 Xue, Y.: Biosphere feedback on regional climate in tropical North Africa, *Quarterly Journal of the Royal Meteorological*
2054 *Society*, 123, 1483–1515, <https://doi.org/10.1002/qj.49712354203>, 1997.
- 2055 Yamaguchi, T. and Feingold, G.: On the relationship between open cellular convective cloud patterns and the spatial
2056 distribution of precipitation, *Atmos. Chem. Phys.*, 15, 1237–1251, <https://doi.org/10.5194/acp-15-1237-2015>, 2015.
- 2057 Yan, M. and Liu, J.: Physical processes of cooling and mega-drought during the 4.2 ka BP event: results
2058 from TraCE-21ka simulations, *Climate of the Past*, 15, 265–277, <https://doi.org/10.5194/cp-15-265-2019>, 2019.
- 2059 Yashayaev, I. and Loder, J. W.: Recurrent replenishment of Labrador Sea Water and associated decadal-scale variability,
2060 *Journal of Geophysical Research: Oceans*, 121, 8095–8114, <https://doi.org/10.1002/2016JC012046>, 2016.
- 2061 Yool, A., Popova, E. E., and Coward, A. C.: Future change in ocean productivity: Is the Arctic the new Atlantic?, *Journal*
2062 *of Geophysical Research: Oceans*, 120, 7771–7790, <https://doi.org/10.1002/2015JC011167>, 2015.
- 2063 Yu, Y., Notaro, M., Wang, F., Mao, J., Shi, X., and Wei, Y.: Observed positive vegetation-rainfall feedbacks in the Sahel
2064 dominated by a moisture recycling mechanism, *Nat Commun*, 8, 1873, <https://doi.org/10.1038/s41467-017-02021-1>,
2065 2017.
- 2066 Zappa, G. and Shepherd, T. G.: Storylines of Atmospheric Circulation Change for European Regional Climate Impact
2067 Assessment, *Journal of Climate*, 30, 6561–6577, <https://doi.org/10.1175/JCLI-D-16-0807.1>, 2017.

- 2068 Zeng, N., Neelin, J. D., Lau, K.-M., and Tucker, C. J.: Enhancement of Interdecadal Climate Variability in the Sahel by
2069 Vegetation Interaction, *Science*, 286, 1537–1540, <https://doi.org/10.1126/science.286.5444.1537>, 1999.
- 2070 Zhang, P., Jeong, J.-H., Yoon, J.-H., Kim, H., Wang, S.-Y. S., Linderholm, H. W., Fang, K., Wu, X., and Chen, D.:
2071 Abrupt shift to hotter and drier climate over inner East Asia beyond the tipping point, *Science*, 370, 1095–1099,
2072 <https://doi.org/10.1126/science.abb3368>, 2020.
- 2073 Zhang, R. and Thomas, M.: Horizontal circulation across density surfaces contributes substantially to the long-term
2074 mean northern Atlantic Meridional Overturning Circulation, *Commun Earth Environ*, 2, 1–12,
2075 <https://doi.org/10.1038/s43247-021-00182-y>, 2021.
- 2076 Zhang, Y., Chiessi, C. M., Mulitza, S., Sawakuchi, A. O., Häggi, C., Zabel, M., Portilho-Ramos, R. C., Schefuß, E.,
2077 Crivellari, S., and Wefer, G.: Different precipitation patterns across tropical South America during Heinrich and
2078 Dansgaard-Oeschger stadials, *Quaternary Science Reviews*, 177, 1–9, <https://doi.org/10.1016/j.quascirev.2017.10.012>,
2079 2017.
- 2080 Zhao, C., Liu, X., Ruby Leung, L., and Hagos, S.: Radiative impact of mineral dust on monsoon precipitation variability
2081 over West Africa, *Atmos. Chem. Phys.*, 11, 1879–1893, <https://doi.org/10.5194/acp-11-1879-2011>, 2011.
- 2082 Zhou, G. and Cheng, X.: Impacts of Oceanic Fronts and Eddies in the Kuroshio-Oyashio Extension Region on the
2083 Atmospheric General Circulation and Storm Track, *Adv. Atmos. Sci.*, 39, 22–54,
2084 <https://doi.org/10.1007/s00376-021-0408-4>, 2022.
- 2085 Zhou, J. and Lau, K.-M.: Does a Monsoon Climate Exist over South America?, *Journal of Climate*, 11, 1020–1040,
2086 [https://doi.org/10.1175/1520-0442\(1998\)011<1020:DAMCEO>2.0.CO;2](https://doi.org/10.1175/1520-0442(1998)011<1020:DAMCEO>2.0.CO;2), 1998.
- 2087 Zhou, S., Meijers, A. J. S., Meredith, M. P., Abrahamsen, E. P., Holland, P. R., Silvano, A., Sallée, J.-B., and Østerhus,
2088 S.: Slowdown of Antarctic Bottom Water export driven by climatic wind and sea-ice changes, *Nat. Clim. Chang.*, 13,
2089 701–709, <https://doi.org/10.1038/s41558-023-01695-4>, 2023.
- 2090 Zhu, C., Liu, Z., Zhang, S., and Wu, L.: Likely accelerated weakening of Atlantic overturning circulation emerges in
2091 optimal salinity fingerprint, *Nat Commun*, 14, 1245, <https://doi.org/10.1038/s41467-023-36288-4>, 2023.
- 2092 Zickfeld, K., Knopf, B., Petoukhov, V., and Schellnhuber, H. J.: Is the Indian summer monsoon stable against global
2093 change?, *Geophysical Research Letters*, 32, <https://doi.org/10.1029/2005GL022771>, 2005.