

# Short communication: Resolving the discrepancy between U–Pb age estimates for the ‘Likhall’ ~~zircon~~-bed, a key level in the Ordovician timescale

André N. Paul<sup>1\*</sup>, Anders Lindskog<sup>2</sup>, Urs Schaltegger<sup>1</sup>

5 1) Department of Earth Sciences, Université de Genève, Switzerland

2) Department of Geology, Lund University, Sweden

\*corresponding author, e-mail: Andre.Paul@unige.ch

## Abstract

The ‘Likhall’ ~~zircon~~-bed is a rare case of a single-age zircon population from a carbonate rock, which in this case is contextualised with remarkable biotic and environmental changes as well as meteorite bombardment of Earth after an asteroid breakup in space. Published chemical-abrasion, high-precision isotope-dilution, thermal ionization mass spectrometry (CA-ID-TIMS) U–Pb age estimates disagree at the typical precision of <0.1% for a <sup>206</sup>Pb/<sup>238</sup>U date, which has led to discrepancies in the interpretation of the timing of events and their possible cause–effect relationships. We evaluate here the relative strengths and weaknesses, and discrepancies in the so far published datasets, propose strategies to overcome them and present a new U-Pb dataset with improved precision and accuracy. Ultimately, we find that domains of residual Pb-loss are a significant source of age-offset between previously published data, amplified by differences in data evaluation strategies. Our new dataset benefits from an improved chemical abrasion protocol resulting in a more complete mitigation of decay-damage induced grain portions, and points to a weighted mean age estimate of 466.37±0.14/0.18/0.53 Ma for the ‘Likhall’ zircon population. This age is intermediate between previous estimates, but outside of analytical uncertainty, and provides a firm tie point for the Ordovician timescale.

## 1 Introduction

The Ordovician ‘Likhall’ ~~bed hosts zircons, are hosted within a fossiliferous and is an rich~~ ‘orthoceratite limestone’ ~~at outcropping at~~ Kinnekulle, Sweden, locally referred to as ‘Täljsten’, which is an important marker interval in the regional stratigraphy (see below). These strata record remarkable changes in paleoenvironmental conditions and biodiversity (Lindskog and Eriksson, 2017; Servais and Harper, 2018; Lindskog et al., 2017). Besides being unusually rich in prismatic zircon of apparently single age, ~~thus providing means for timescale calibration,~~ the ‘Likhall’ bed coincides with an interval with uniquely abundant L-chondritic ‘fossil’ meteorites and purportedly related chromite grains (Lindskog et al., 2017; Liao et al., 2020; and references therein). This has been linked to the breakup of an asteroid in space and ~~the~~ temporal overlap between excess meteoritic matter and prominent climatic changes and biodiversity peaks has spurred the controversial hypothesis that ~~Ordovician~~

30 ~~biodiversification was instigated by~~ meteorite bombardment fundamentally affected the Ordovician climate and instigated  
biodiversification (Schmitz et al. 2008, 2019; cf. Lindskog et al. 2017; Rasmussen et al. 2021). Considering absolute/numerical  
time perspectives. ~~However,~~ two individual U–Pb geochronological studies have arrived at significantly different age estimate  
of zircon from the ‘Likhall’ bed ~~this rock~~ – 467.50±0.28 (Lindskog et al. 2017) and 465.18±0.17 Ma (Liao et al. 2020) – each  
favouring reaching contrasting conclusions for the absolute timing of the L-chondrite breakup event, ~~(non-)correlation of the~~  
35 ~~meteorite bombardment with biodiversification,~~ and with differing implications for the Ordovician event stratigraphy and  
timescales (particularly the Darriwilian Stage).

While the occurrence of an apparently single U–Pb zircon age population in a carbonate rock is already mysterious/uncommon,  
the percent difference between these two previously published  $^{206}\text{Pb}$ - $^{238}\text{U}$  ages amounts to 0.4%, which exceeds the expected  
reproducibility level of, e.g. for example, natural zircon reference materials (0.1%; Schaltegger et al., 2021). Thus, we have to  
40 evaluate the accuracy of the published U–Pb ages on the ‘Likhall’ zircons before we can determine a more robust age estimate.  
A significant effort has recently been undertaken by the U-Pb community to reduce inter-lab bias in isotope dilution – thermal  
ionisation mass spectrometry (ID-TIMS), via the use of precisely calibrated EARTHTIME (ET) isotopic tracers (Condon et  
al., 2015; McLean et al., 2015), community-wide shared data reduction (Bowring et al., 2012; McLean et al., 2011) and sample  
preparation procedures (Widmann et al., 2019). However, the two previously published ‘Likhall’ zircon datasets were  
45 produced employing different isotopic tracers, instrumentation and chemical abrasion procedures: A mixed ET  $^{202}\text{Pb}$ - $^{205}\text{Pb}$ -  
 $^{233}\text{U}$ - $^{235}\text{U}$  tracer was used in the Lindskog et al. (2017) study, while it was an in-house  $^{202}\text{Pb}$ - $^{205}\text{Pb}$ - $^{233}\text{U}$ - $^{236}\text{U}$  tracer for the Liao  
et al. (2020) study. Further differences concern the measurement procedure of the U isotope composition (on TIMS as  $\text{UO}_2$   
for the first, on a multi collector – inductively coupled plasma – mass spectrometer (MC-ICP-MS) as a metal for the latter).  
Most importantly, chemical abrasion procedures also differ, being annealing at 900°C for 3 days and leaching at 180°C for  
50 12hrs (Lindskog et al., 2017) and annealing at 900°C for 48h and leaching at 190°C for 15hrs (Liao et al., 2020).

The variation of the chemical abrasion procedure parameters has profound effects on the U–Pb zircon isotopic systematics.  
The duration as well as the temperature of the ~~chemical abrasion/partial dissolution~~ has/ may have significant impact on the  
potential to effectively remove structurally damaged domains, which typically produce anomalously young dates (Mattinson,  
2005; Huyskens et al., 2016; Keller et al., 2019; Widmann et al., 2019). Both of the aforementioned studies of the ‘Likhall’  
55 bed utilised temperature lower than the most recent recommendation for the chemical abrasion procedure (McKanna et al.,  
2023a, 2023b; Widmann et al., 2019), which raises concerns about remnant Pb-loss domains present in the zircons analysed.  
Different chemical abrasion procedures s may be variably effective in removing visible inclusions s (e.g. apatite and/or melt  
inclusions) in the analysed zircon crystals.

The impact of tracer uncertainty is significantly lower than the 0.4% discrepancy between the absolute ages obtained from  
‘Likhall’ bed. Similarly, the variation of the instrumental setup for U isotope analysis does not seem to introduce significant  
off-sets, although we cannot evaluate this with certainty, due to limited amount of comparable data. The  $^{206}\text{Pb}/^{238}\text{U}$  age of the  
natural reference zircon material Temora reported by Liao et al. (2020) from the mixed TIMS–MC-ICP-MS analysis  
60 (417.19±0.15 Ma) is in agreement with the most recent estimates of 417.310 ± 0.074 Ma (von Quadt et al., 2016) and

417.353±0.052 Ma (Schaltegger et al., 2021). Reported values of the ET100 solution from the lab utilised in the Liao et al. (2020) is 100.169±0.047 Ma, indistinguishable from the reference value of 100.173±0.003 Ma (Schaltegger et al., 2021) and our value produced (100.1678 ± 0.0046 Ma) during the period of data collection of this study.

Thus, we can reasonably assume that the difference in the chemical abrasion procedure is the main source for the ~~poor~~ reproducibility/significant difference between the two previously published studies of ‘Likhall’ zircons. Furthermore, there is a difference in data interpretation strategy of the two published datasets. Lindsog et al. (2017) put the emphasis on the largest, statistically valid, weighted mean age plateau, whereas Liao et al. (2020) chose the youngest cluster of zircon U–Pb analyses. In the following, we will first explore different interpretation strategies of the existing data, such as *i*) looking for the largest statistically valid plateau, *ii*) the youngest cluster, *iii*) the duration of time recorded by the youngest and oldest zircon U–Pb date ( $\Delta t$ ) and *iv*) the youngest concordant single zircon U–Pb analysis. Ideally, applying the same interpretation methodology to both datasets should reduce the discrepancy of the absolute U–Pb ages. Furthermore, we add another set of U–Pb dates from the same material but apply the chemical abrasion procedure of Widmann et al. (2019), which should more effectively mitigate Pb-loss. Ultimately, we will suggest a revised age for the ‘Likhall’ bed.

## 2 Methods

Single grains of ‘Likhall’ zircon crystals free of visible inclusions and cracks were hand-picked ~~under-using~~ a binocular microscope at a magnification of  $\times 20$  to  $\times 40$  from the same petri dish as the material analyzed by Lindsog et al. (2017). The size of individual ~~fragments-zircons~~ was variable, with length ranging from  $\sim 50\mu\text{m}$  to  $\sim 300\mu\text{m}$ . The chemical abrasion procedures were 48hrs of annealing at  $900^\circ\text{C}$ , 12hrs partial dissolution at  $210^\circ\text{C}$ , as defined as optimal in Widmann et al. (2019). Afterwards, we observe that some zircons had either fragmented or embayment’s at the core predominantly, or some channels had more efficiently progressed inwards, without any systematic preference identifiable. Individual zircon grains were washed in 3ml Savillex beakers in an ultrasonic bath,  $4\times$  in  $7\text{N HNO}_3^-$ , transferred into individual  $200\mu\text{L}$  Savillex microcapsules, along with 2–3 drops of  $\text{HF}_{\text{conc}}$  and 3.9–5.5 mg of a mixed  $^{202}\text{Pb}$ - $^{205}\text{Pb}$ - $^{233}\text{U}$ - $^{235}\text{U}$  tracer solution (ET2535, Condon et al., 2015; McLean et al., 2015), and dissolved at  $210^\circ\text{C}$  in a Parr bomb for 48hrs. After dissolution, samples were dried down on a hotplate at  $120^\circ\text{C}$ , re-dissolved in  $3\text{N HCl}$ , and then U and Pb were separated using a single column anion exchange chemistry. Uranium and Pb were loaded on outgassed, zone-refined, single Re filaments with a silica-gel/phosphoric acid emitter solution (Gerstenberger and Haase, 1997).

Uranium and Pb isotopic compositions were measured on an IsotopX Phoenix TIMS at the University of Geneva. Lead was measured in dynamic mode using a Daly detector, U was measured as an oxide in static mode using Faraday cups coupled to  $10^{12}\ \Omega$  resistance amplifiers. Measured isotopic ratios were corrected for interferences of  $^{238}\text{U}^{18}\text{O}^{16}\text{O}$  on  $^{235}\text{U}^{16}\text{O}_2$  using a  $^{18}\text{O}/^{16}\text{O}$  composition of 0.00205, based on repeat measurements of the U500 standard. Mass fractionation of U was corrected using a double isotope tracer with a  $^{235}\text{U}/^{233}\text{U}$  of  $0.99506\pm 0.005$ . Mass fractionation of Pb is calculated and corrected using a  $^{202}\text{Pb}/^{205}\text{Pb}$  ratio of  $0.99923913\pm 0.00026555$  ( $1\sigma$ ) (Condon et al., 2015). Zircon Pb analyses were corrected for laboratory

blanks, with a  $^{206}\text{Pb}/^{204}\text{Pb}$  of  $17.10 \pm 0.21$ , a  $^{207}\text{Pb}/^{204}\text{Pb}$  of  $15.07 \pm 0.11$  and a  $^{208}\text{Pb}/^{204}\text{Pb}$  of  $36.17 \pm 0.25$  (all  $1\sigma$ ), based on repeat measurements of total procedural blanks for the zircon U-Pb column chemistry. All data were processed using the Tripoli v. 4.10 and Redux v. 3.7.1 U-Pb software (Bowring et al., 2012; McLean et al., 2011). Weighted mean U-Pb age uncertainties are reported at the  $2\sigma$  level in the format  $A \pm X/Y/Z$ , where A is the weighted mean age, X is analytical uncertainty, Y is analytical and tracer uncertainty combined, and Z is analytical, tracer, and decay constant uncertainties combined (Schoene et al., 2006). Thorium abundances for each grain, and subsequently  $\text{Th}/\text{U}_{\text{zircon}}$ , were calculated using the abundance of  $^{208}\text{Pb}$  within the crystal and the  $^{206}\text{Pb}/^{238}\text{U}$  age to calculate radiogenic in-growth (McLean et al., 2011). All U-Pb dates are corrected for initial  $^{230}\text{Th}$  disequilibrium, assuming a Th/U ratio of the magma of  $3.5 \pm 1$ .

Repeat analyses of the ET100 solution ( $^{206}\text{Pb}/^{238}\text{U}$  date:  $100.173 \pm 0.003$  Ma; Schaltegger et al., 2021) yielded a value of  $100.1678 \pm 0.0046$  Ma (MSWD = 3.2, n = 32/40), during the period of data collection. One batch comprised of eight consecutive ET100 samples was rejected, due to an anomalously young average age (batch internally consistent).

### 4.3 Geological Setting

The Baltoscandian region was largely covered by an epeiric sea throughout much of the Ordovician, leaving behind a sedimentary record mainly in the form of mudstones, shales, and limestones (e.g., Nielsen et al., 2023, and references therein). In the Lower–Middle Ordovician, the continent Baltica was at a relatively high latitudinal position and any relief of the regional Precambrian basement was near-completely smoothed out. These conditions limited terrigenous input into the basin, and sedimentation rates were typically low (a few mm/ka). The regional intra-cratonic Ordovician succession is therefore relatively thin. Volcanic dust may have contributed a relatively large proportion of the non-carbonate sedimentary materials (Lindstrom, 1974). The Middle Ordovician (Dapingian–Darriwilian) was mainly characterized by widespread deposition of cool-water carbonate sediments, in Sweden commonly referred to as the ‘orthoceratite limestone’ (e.g., Lindskog and Eriksson, 2017; Lindskog et al., 2023).

The Ordovician succession of Baltoscandia contains numerous bentonite beds of varying thickness and lateral distribution, the most prominent of which occur in the Upper Ordovician (e.g., Ballo et al., 2019; Bergström, 1989; and references therein). Few of the bentonite beds have been isotopically dated, and even fewer so using modern techniques such as CA-ID-TIMS. Some carbonate beds in the ‘orthoceratite limestone’ contain abundant zircon, and the crystal characteristics and U–Pb age data of the grains indicate a volcanic origin (Lindskog et al. 2017; Liao et al. 2020). Thus, the zircon-rich beds arguably represent ‘crypto-tephra’ (McLaughlin et al., 2023).

The table-mountain Kinnekulle in the province of Västergötland, south-central Sweden, hosts a relatively expanded ‘orthoceratite limestone’ succession (e.g., Lindskog and Eriksson, 2017). These rocks have been the target of several studies of, e.g., paleontology, sedimentology, and geochemistry (e.g., Ahlberg et al., 2023, and references therein). Our sample materials derive from the Thorsberg quarry on eastern Kinnekulle (WGS84 coordinates 58.579167, 13.429444), from a

130 distinct, gray-colored interval traditionally referred to as the ‘Täljsten’ (‘carving stone’) by local quarrymen. In more detail, the sample derives from a specific bed referred to as ‘Likhall’ (‘corpse slab’), which has been shown to contain very abundant zircon grains (Lindskog et al. 2017; Liao et al. 2020). The biostratigraphic context of this c. 10 cm thick bed is very well known, and its base coincides with that of the geographically widely distributed *Yangtzeplacognathus crassus* conodont Zone in the middle Darriwilian (for more details, see Lindskog et al. 2017).

### 3.4 Results

135 A total of 22 zircons from the zircon-rich ‘Likhall’ bed were analysed for their U–Pb isotopic composition. The U and Pb isotopic data are presented in Table 1, interpreted U–Pb dates are illustrated in Concordia space in Fig. 1, revealing a large spread in  $^{206}\text{Pb}/^{238}\text{U}$  dates as well as variable discordance. The new data presented here comprise 15 analytically concordant and 7 discordant  $^{206}\text{Pb}/^{238}\text{U}$  zircon analyses, which range in  $^{206}\text{Pb}/^{238}\text{U}$  dates from  $468.8\pm 1.2$  to  $462.43\pm 0.27$  Ma. The ratio of radiogenic to common lead ( $\text{Pb}^*/\text{Pb}_c$ ) ranges from 11.1 to 79.7, with variable amounts of  $\text{Pb}_c$  (0.20 to 1.09 pg). The  $^{207}\text{Pb}/^{206}\text{Pb}$  dates range from  $461\pm 10$  to  $490\pm 15$  Ma.

140 The compilation of previously published zircon U–Pb analyses of the ‘Likhall’ bed comprises 17 analyses from Lindskog et al. (2021) and 21 analyses from Liao et al. (2020), which are now complemented by our 22 new analyses in this study. The  $^{206}\text{Pb}/^{238}\text{U}$  dates of all these datasets are plotted in Fig. 2, along with the weighted mean  $^{206}\text{Pb}/^{238}\text{U}$  age for each of the datasets following the selection criteria discussed below. All data are available in Table 1 and Supplement (Table S1).




Fig. 1) Concordia diagram generated in ETRedux v. 3.7.1, using colour coding for bands of  $Pb^*/Pb_c$  as discussed in the text for the error ellipses as a function of the  $Pb^*/Pb_c$  value. Red-Orange colours indicates higher  $Pb^*/Pb_c > 50$ , whereas blue colours indicates low  $Pb^*/Pb_c$  values between 15-50, grey error ellipses indicate  $Pb^*/Pb_c < 15$ . Ellipses marked by an asterisk are interpreted to be affected by Pb-loss. Black outline indicates analyses considered for the weighted mean age, where the selection criterion is  $Pb^*/Pb_c > 50$  and analytical concordance (part or full ellipse overlaps with the uncertainty band of the Concordia curve). Grey band illustrates the uncertainty band of the Concordia curve.

150



165 ~~sedimentation rates were typically low (a few mm/ka). The regional intra-cratonic Ordovician succession is therefore relatively thin. Volcanic dust may have contributed a relatively large proportion of the non-carbonate sedimentary materials (Lindstrom, 1974). The Middle Ordovician (Dapingian-Darriwilian) was mainly characterized by widespread deposition of cool-water carbonate sediments, in Sweden commonly referred to as the ‘orthoceratite limestone’ (e.g., Lindskog and Eriksson, 2017). The Ordovician succession of Baltoscandia contains numerous bentonite beds of varying thickness and lateral distribution, the~~  
170 ~~most prominent of which occur in the Upper Ordovician (e.g., Ballo et al., 2019; Bergström, 1989; and references therein). Few of the bentonite beds have been isotopically dated, and even fewer so using modern techniques such as CA-ID-TIMS. Some carbonate beds in the ‘orthoceratite limestone’ contain abundant zircon, and the crystal characteristics and U–Pb age data of the grains indicate a volcanic origin (Lindskog et al. 2017; Liao et al. 2020). Thus, the zircon-rich beds arguably represent ‘crypto-tephra’ (McLaughlin et al., 2023).~~

175 ~~The table mountain Kinnekulle in the province of Västergötland, south central Sweden, hosts a relatively expanded ‘orthoceratite limestone’ succession (e.g., Lindskog and Eriksson, 2017). These rocks have been the target of several studies of, e.g., paleontology, sedimentology, and geochemistry (e.g., Ahlberg et al., 2023, and references therein). Our sample materials derive from the Thorsberg quarry on eastern Kinnekulle (WGS84 coordinates 58.579167, 13.429444), from a distinct, gray-colored interval traditionally referred to as the ‘Täljsten’ (‘carving stone’) by local quarrymen. In more detail,~~  
180 ~~the sample derives from a specific bed referred to as ‘Likhall’ (‘corpse slab’), which has shown to contain very abundant zircon grains (Lindskog et al. 2017; Liao et al. 2020). The biostratigraphic context of this c. 10 cm thick bed is very well known, and its base coincides with that of the geographically widely distributed *Yangtzeplacognathus crassus* conodont Zone in the middle Darriwilian (for more details, see Lindskog et al. 2017).~~

## 5 Discussion

### 185 5.1 Radiogenic Pb/common Pb ratio ( $Pb^*/Pb_c$ ) as a selection criterion:

The color-coding of the data ellipses in Fig. 1 indicates that the majority of the analyses plot onto or near ~~the~~ concordia within its uncertainty band. Analyses with low  $Pb^*/Pb_c$  ( $<50$ ) have significantly higher scatter in both  $^{206}Pb/^{238}U$  and  $^{207}Pb/^{235}U$  ratios. Subsequently, we only use high  $Pb^*/Pb_c$  ( $>50$ ) analyses for our age interpretation and will also apply this strategy to the previously published datasets. The accuracy of  $^{206}Pb/^{238}U$  dates is highest ~~at~~  $Pb^*/Pb_c >15$ –~~20~~ (Schaltegger et al., 2021), which  
190 is the case for most of our analyses that underwent chemical abrasion for 12hrs at 210°C, and for some of the analyses of Liao et al. (2020) abraded for 15hrs at 190°C (Fig. 3). Contrastingly, a major part of the Lindskog et al. (2017) dates abraded for 12hrs at 180°C plot below this threshold at ~~very~~-variable  $^{206}Pb/^{238}U$  dates. Considering the significant differences in the range of  $Pb^*/Pb_c$  reported in the three datasets, we have also evaluated  $Pb^*/Pb_c$  bands ranging  $<15$ ,  $15$ - $50$  in addition to  $>50$  discussed above.


195 When considering only analyses with  $Pb^*/Pb_c <15$ , the Lindskog et al (2017), Liao et al. (2020) and this study datasets would result in weighted mean ages of  $468.40 \pm 0.50$  Ma (MSWD = 2.7, n = 10),  $465.92 \pm 0.21$  Ma (MSWD = 3.6, n = 14) and



467.67±0.84 Ma (MSWD = 0.18, n = 2). Improvement of the weighted mean ages between the studies can be observed when considering Pb\*/Pbc band 15-50, where the studies result in 467.12±0.59 Ma (MSWD = 0.46, n = 6), 466.68±0.18 Ma (MSWD = 5.4, n = 6) and 466.79±0.15 Ma (MSWD = 1.4, n = 11) (discordant data not considered, Table S1). In our new dataset, the discrepancy between the proposed U-Pb age when comparing Pb\*/Pbc bands 15-50 and >50 is 0.09%, which is the expected reproducibility of natural reference materials (Schaltegger et al. 2021). Potential sources of this discrepancy may be i) residual Pb-loss, ii) inaccurate Pb-blank correction iii) natural variability and duration of magmatic zircon growth in a magma chamber. Where possible, high Pb\*/Pbc data should be preferred for interpretation, to reduce effects from potential i) residual Pb-loss and ii) inaccurate Pb-blank correction.

205 Adopting the Pb\*/Pbc (>50 in this study) as a criterion for data selection thus is an important step towards greater accuracy. The data of Lindskog et al. (2017) shows significant correlation between Pb\* and Pb<sub>c</sub> (Fig. 4), at overall high Pb<sub>c</sub> values (up to ~6 pg), which makes us believe that some indiscernible mineral inclusions were not removed during chemical abrasion. Chemical abrasion at higher temperature ~~ideally~~ should access and dissolve these (McKanna et al., 2023a), if the hypothesis of un-resolved inclusions is correct, and would reduce the total Pb<sub>c</sub>, ~~if it primarily originated from inclusions observed~~.

210 Subsequently, we would then use only ~~low blank or~~ high Pb\*/Pbc (>50) analyses to make an age interpretation. The high Pb<sub>c</sub> concentrations of the Lindskog et al. (2017) analyses (up to ~~5.996~~ pg), suggest that the conditions during the chemical abrasion procedure were not sufficient to remove the inclusions that are clearly visible in the grain separate (Lindskog et al., 2017), leading to generally low Pb\*/Pbc. The higher dependency on the Pb<sub>c</sub> correction can potentially result in over- or under-correction, which results in younger or older dates calculated. This is further explored in section 5.3 further on.




215


**Fig. 3)** Comparison of calculated  $^{238}\text{U}/^{206}\text{Pb}$  zircon dates with their respective  $\text{Pb}^*/\text{Pb}_c$  values. Low  $\text{Pb}^*/\text{Pb}_c$  values typically indicate stronger effects of blank correction on the calculated  $^{238}\text{U}/^{206}\text{Pb}$  dates (Schaltegger et al. 2021). Colours indicate the different studies and differences in their chemical abrasion procedure. Samples marked with an asterisk are discordant in concordia space (see Fig. 1).

220

Lindskog et al. 2017



Liao et al. 2020



this study




Fig. 4) Comparison of  $Pb_c$  vs  $Pb^*$  of a) the Lindskog et al. (2017) study, b) the Liao et al. (2020) study and c) this study. Neither study exhibits simple correlation without outlier rejection. A positive correlation between  $Pb_c$  and  $Pb^*$  may indicate presence of inclusions (mineral and/or melt), which were not removed during the chemical abrasion process.

225

## 5.2 Residual Pb-loss in chemically abraded natural zircon:

~~In natural zircons  $\alpha$  decay damage induced partial Pb-loss is one reason for normally discordant U–Pb data (Mezger and Krogstad, 1997). This~~ The effect of Pb-loss may be mitigated by removing decay damaged (metamict) portions of the zircon grains using the chemical abrasion procedure (Mattinson, 2005; Mezger and Krogstad, 1997). Subsequently, if all Pb-loss domains were removed, the isotopic analysis should yield an ~~analytically~~ concordant result for both the  $^{206}Pb$ – $^{238}U$  and  $^{207}Pb$ – $^{235}U$  decay series, respectively. ~~Such information can then be related to the age of eruption or solidification, while the range of U–Pb may be a measure of the duration of crystallisation of zircon in a magma.~~ However, the chemical abrasion may not have removed 100% of metamict portions and some domains with partial Pb-loss may still be present in the grain (so-called residual Pb-loss). This may be the case even if the ~~optimal-most recent~~ calibration of the chemical abrasion procedure (12hrs at 210°C, Widmann et al., 2019) is utilized. ~~It should also be noted that chemical abrasion calibration may be affected by geochemical composition, parent isotope zonation, accessibility of damaged domains for the chemical agent and other effects.~~ Previous work has shown that natural zircon reference materials (Temora and GJ-1) treated at 180°C and 210°C for 12hrs retains excess scatter in their U–Pb systematics, restricting repeatability confidence to ca. 0.1% of the absolute age, while synthetic solutions offer a repeatability at a precision of up to 0.01% in the same study (Schaltegger et al., 2021). ~~It is however not clear if excess scatter in Temora and GJ-1 derives from inherited older components, magmatic processes or residual Pb-loss for these materials (Schaltegger et al. 2021).~~ When comparing the effects of different chemical abrasion temperatures, Huyskens et al. (2016) found that temperatures of 190°C and lower may yield incomplete removal of Pb-loss domains, in agreement with later experiments (e.g. Widmann et al., 2019; Schaltegger et al., 2021). Such an effect of incomplete removal of metamict domains biased by Pb-loss may possibly be detected through analytical discordance between the two decay schemes (Condon et al. 2024), provided that the analytical precision, especially of the  $^{207}Pb/^{235}U$  decay series, was sufficient, ~~assuming that no inheritance is present (possible in the new dataset, not resolvable in Lindskog et al. 2017 dataset).~~

Between the previously published age estimates for the ‘Likhall’ bed by Lindskog et al. (2017) ( $467.50 \pm 0.28$  Ma) and Liao et al. (2020) ( $465.18 \pm 0.17$  Ma), there is a discrepancy of ca. 0.4% between the proposed U–Pb ages (Fig. 2). The two studies differ in their chemical abrasion protocols: Lindskog et al. (2017) utilised a 180°C and 12hrs procedure, whereas Liao et al. (2020) utilised a 190°C and 15hrs procedure. Both of these protocols diverge from the Widmann et al. (2019) parameters and we must consider the potential that these analyses included relict domains of Pb-loss, that may bias the U–Pb dates towards younger dates for both studies. Curiously, for the first case, the lower T abrasion resulted in (on average) older interpreted zircon U–Pb data (Lindskog et al., 2017), an effect we will explore in the next section discussing blank corrections.

Both the Lindskog et al. (2017) and Liao et al. (2020) datasets show very little evidence for normal discordance, as analyses which are ~~technically~~-discordant are typically older than the interpreted U–Pb age, suggesting inheritance rather than Pb-loss. ~~Normal-Negative~~ discordance can be masked if the analytical precision is insufficient, in particular when the measured  $^{207}\text{Pb}$  intensity is low. In our new data we observe two discordant analyses that are by ~2–4 Ma (Fig. 1) younger than the previous age estimates (Lindskog et al., 2017; Liao et al., 2020). We therefore infer that the data of Liao et al. (2020) and Lindskog et al. (2017) were affected by Pb-loss domains and/or relict inclusions that were not penetrated during chemical abrasion (expressed through elevated  $\text{Pb}_c$  and/or younger dates), which may also not be excluded for our new dataset with absolute certainty. Relict inclusions have an unknown amount of  $\text{Pb}_c$  and its composition cannot be assessed by ID-TIMS data. This could result in erroneous blank composition corrections, which may result in older U–Pb dates. Therefore, we base our discussion and interpretation on those data that are the least affected by Pb blank correction.

### 5.3 Lead blank isotopic composition correction effects on the spread of zircon U–Pb dates:

One of the fundamental assumptions in zircon U–Pb geochronology is that zircon does not incorporate Pb during crystallisation and therefore does not contain initial/~~common-Pb~~ (which can be monitored through analysis of  $^{204}\text{Pb}$ ) (Watson et al., 1997). Therefore, all zircon U–Pb analyses can be corrected for the presence of common Pb through measurement of  $^{204}\text{Pb}$  during data acquisition, assuming that all  $\text{Pb}_c$  is derived from the procedural blank, by adopting the mean isotopic composition from repeat analysis of procedural blank measurements. The uncertainty of this mean blank isotopic composition is propagated into the U–Pb date calculation (Schmitz and Schoene, 2007). An accurate and precise correction of blank Pb thus results in a more precise and accurate U–Pb dates. Conversely, inaccurate blank corrections may result in apparently too old or young U–Pb dates, if the  $\text{Pb}^*/\text{Pb}_c$  is low (Schaltegger et al., 2021).

In the Lindskog et al. (2017) dataset, a significant proportion of the analytical uncertainty is controlled by the Pb blank composition correction, as  $\text{Pb}^*/\text{Pb}_c$  ratios range from 2 to 22, while the Liao et al. (2020) dataset ranges from 2 to 30 (plus one high value at 67). In our new dataset, we observe  $\text{Pb}^*/\text{Pb}_c$  from 11 to 78. When we compare the  $^{206}\text{Pb}/^{238}\text{U}$  zircon U–Pb dates with their  $\text{Pb}^*/\text{Pb}_c$  (Fig. 3), we observe that the Lindskog et al. (2017) data exhibit a slightly negative correlation, whereas the Liao et al. (2020) are ~~scatteringscattered~~. However, we observe in the Liao et al. (2020) data that the youngest analyses are consistently associated with lower  $\text{Pb}^*/\text{Pb}_c$ , with the exception of the analysis with the highest  $\text{Pb}^*/\text{Pb}_c$  which is also relatively young (Fig. 3). In our newly acquired dataset, correlation between  $^{206}\text{Pb}/^{238}\text{U}$  zircon U–Pb date with its  $\text{Pb}^*/\text{Pb}_c$  is absent, suggesting that blank correction does not introduce significant bias (Fig. 3).

The strong correlation between  $\text{Pb}^*$  and  $\text{Pb}_c$  ( $R^2$  of 0.52 after rejecting two outliers;  $\text{Pb}^*$  max = 55.33 pg and  $\text{Pb}_c$  max = 5.24 pg) in the Lindskog et al. (2017) dataset is concerning (Fig. 4), as it implies that the analysed zircons contained inclusions that were not removed during the chemical abrasion procedure. Evidence for potential inclusions (before chemical abrasion) is provided by imaging of zircon crystals analysed by Lindskog et al. (2017) and Liao et al. (2020), matching our observations during mineral selection. If we assume that larger zircons contain more  $\text{Pb}^*$  and a larger volume of  $\text{Pb}_c$ -bearing inclusions, this would explain why the Lindskog et al. (2017) dataset exhibits a negative correlation between  $\text{Pb}^*/\text{Pb}_c$  and  $^{206}\text{Pb}/^{238}\text{U}$  dates, as

they are more strongly affected by an inaccurate blank correction. The slope of the correlation between  $Pb^*/Pb_c$  and  $^{206}Pb/^{238}U$  dates is controlled by the difference between the “true”  $Pb_c$  composition and the assigned  $Pb_c$  composition for blank correction. In the Liao et al. (2020) dataset, correlation between  $Pb^*$  and  $Pb_c$  is absent, but measured  $Pb^*/Pb_c$  are comparable to those of  
290 Lindskog et al. (2017; Fig. 4). Our new dataset does not show any correlation between  $Pb^*$  and  $Pb_c$  ( $Pb^*$  max = 59.80 pg and  $Pb_c$  max = 1.09 pg; Fig. 4) nor between  $Pb^*/Pb_c$  and  $^{206}Pb/^{238}U$  date (Fig. 3).

Some of the low  $Pb^*$  zircons in our new data also have elevated blanks, causing scatter in the low  $Pb^*$  analyses systematics, whereas all the high  $Pb^*$  analyses associate with slightly higher blanks. We therefore consider it likely that the  $Pb_c$  is primarily controlled by un-resolved inclusion (transparent, mineral and/or melt). These observations are in line with the observations by  
295 McKanna et al. (2023a) and the Lindskog et al. (2017) data (Fig 4a), that chemical abrasion at high temperatures is necessary to effectively remove inclusions that are deeply seated within the zircon crystal. Factors such as location of inclusions within or outside of high U, damaged domains may impact the efficiency of the chemical abrasion procedure.

#### 5.4 The impact of the interpretation strategy on U–Pb zircon ages

300 Correlation between U–Pb dates and  $Pb^*/Pb_c$  has implications for interpreting the absolute age of the ‘Likhall’ bed. Several different interpretation strategies exist, such as i) weighted mean of a subset of data (e.g. Lindskog et al. 2017), ii) youngest cluster of overlapping analyses at  $2\sigma$  (e.g. Liao et al., 2020), iii) considering the entire range of concordant zircon U–Pb analyses as autocrystic growth within the magma chamber (Samperton et al., 2015), iv) considering the youngest concordant analysis as best proxy for the timing of eruption and v) applying a stochastic (Bayesian) sampling approach (Keller et al., 2018,  
305 Traylor et al. 2021). We discuss in the following the impact of interpretation scenarios i–iv on the suggested age for the ‘Likhall’ bed (v is beyond the scope of this study, but we report model results in Table S2, which we find to be too heavily dependent on the input data itself.) Analytical and mineralogical effects such as using variable or inaccurate blank isotopic composition and the presence of  $Pb_c$ -rich inclusions, together with the presence of residual Pb-loss and minor inheritance of old radiogenic Pb (potentially from inherited cores), add to any of the further discussed discrepancies.

##### 310 5.4.1 i) Subset interpretation

Lindskog et al. (2017) preferred a data interpretation based on the statistically most robust weighted mean age, representing the largest number of statistically valid analyses, which results in  $467.50 \pm 0.28$  Ma (MSWD = 1.4, n = 9, published value rejecting two younger and six older, concordant analyses). The underlying assumption is that some un-resolved Pb-loss may affect the youngest analyses and that inheritance may explain the oldest range of zircons analysed (Samperton et al., 2015).  
315 Applying the same strategy to the Liao et al. (2020) dataset, we obtain  $466.34 \pm 0.20$  Ma (MSWD = 0.99, n = 12), rejecting five younger and three older analyses. For our new dataset, the result would be  $466.875 \pm 0.074$  Ma (MSWD = 0.87, n = 9), rejecting

3 discordant analyses, 6 younger analyses and 5 older analyses. The maximum difference between the absolute ages amounts to 0.25%, which is better than the difference of 0.4% of the previously published values. The lowest discrepancy is achieved between Liao et al. (2020) and our dataset, with a difference of 0.11%.

#### 320 **5.4.2 ii) Youngest cluster interpretation**

Liao et al. (2020) preferred an interpretation based on the youngest, statistically valid age cluster, which resulted in 465.18±0.17 Ma (MSWD = 1.3, n = 5, published value). The underlying assumption is that all Pb-loss is effectively removed and that inheritance or protracted growth in the magma chamber is responsible for the older analyses. For the Linds kog et al. (2017) data, this approach results in 466.96±0.30 Ma (MSWD = 1.5, n = 6). For our new dataset, the corresponding result is 325 466.43±0.11 Ma (MSWD = 5, n = 5). The resulting maximum spread is 0.38%, close to the published discrepancy of 0.4%. The lowest discrepancy is achieved between Linds kog et al. (2017) and our dataset, with a difference of 0.11%.

#### **5.4.3 iii) range of concordant zircon U–Pb dates**

The duration of zircon crystallisation in a magma chamber is of interest for studies involving magma chamber dynamics and crustal evolution and we here compare the  $\Delta t$  defined as spread between youngest and oldest (concordant) zircon U–Pb analysis. The underlying assumption is that all concordant analyses reflect growth from the same magma chamber (Samperton et al., 2015). The  $\Delta t$  for the Linds kog dataset is 5.35 Ma, Liao et al. (2020) is 3.8 Ma (rejecting two outliers marked by Liao et al. (2020)) and our new dataset is 1.82 Ma (rejecting 3 young discordant data points) – results which differ significantly from each other. One possible explanation could be the unresolved inclusions and resultant low  $Pb^*/Pb_c$ , which can result in calculated too young or too old dates, augmenting natural spreads in zircon U–Pb dates. The maximum difference between the 335 three datasets is 194%, suggesting that interpretation is problematic with respect to the duration of magma chamber activity.

#### **5.4.4 iv) youngest concordant zircon U–Pb analysis**

The youngest concordant zircon U–Pb analysis interpretation can be useful in volcanic samples. Here, the base assumption is that zircon continuously crystallises in the magma, and the youngest zircon represents the best proxy to the timing of eruption. In this regard, for Liao et al. (2020) the youngest concordant zircon U–Pb analysis is 465.07±0.17 Ma, for Linds kog et al., 340 (2017) it is 466.03±1.06 Ma and for our new dataset it is 466.05±0.29 Ma. The maximum difference between the three datasets is 0.21%, better than the discrepancy of the published U–Pb dates. There is a near indistinguishable difference between the Linds kog et al. (2017) and our youngest zircon U–Pb analyses (0.004%).

### **5.5 Which strategy to choose?**

It becomes clear from Fig. 2 that the three datasets were obtained under different analytical regimes in different labs. Among 345 the differences is an improvement in analytical precision (increase of  $Pb^*/Pb_c$  through blank reduction). This will help to

narrow down the duration of zircon growth in the magma system prior to eruption and leads to more reliable identification of autocrystic vs. antecrystic zircon. Identifying Pb-loss remains challenging in the older datasets, which utilized partial dissolution procedures at lower temperature and/or of shorter duration. ~~Any assessment is particularly difficult when supporting information such as blank isotopic composition is absent.~~ Furthermore, low Pb\*/Pb<sub>c</sub> obscure discordance between the two decay series, due to elevated <sup>207</sup>Pb/<sup>235</sup>U uncertainty. ~~The two datasets produced with the EARTHTIME isotopic tracer ET2535 show better comparability despite divergent chemical abrasion procedures, pointing to a systematic effect of different tracer calibration as well. However, t~~The largest difference in ~~ageage, however,~~ is caused by the varying approaches to data interpretation. If the same strategy is chosen, the discrepancy between the proposed U–Pb age in the two previously published studies and the present one significantly decreases.

350 We can conclude that Pb\*/Pb<sub>c</sub> is of fundamental importance for the ‘Likhall’ zircon datasets. If we, for example, consider only analyses that are characterised by Pb\*/Pb<sub>c</sub> >50 – i.e., a value where zircon dates are barely affected by blank correction – we would have to reject the entire Lindskog et al. (2017) dataset and all but one analysis from the Liao et al. (2020) study, and it leaves only 8 out of 22 analyses from our new dataset, of which only 3 form a cluster of youngest, overlapping grains. Adopting this reduced dataset for interpretation yields the following results: i) the largest number of overlapping U–Pb dates  
360 forms a weighted mean of 466.76±0.12 Ma, ii) the youngest cluster is 466.37±0.14 Ma, iii) the Δt reduces to 780 kyrs, which is more consistent with predictions of thermal models for magma chambers (e.g., Caricchi et al., 2016; Weber et al., 2020), and iv) the youngest concordant zircon analysis remains unchanged at 466.05±0.29 Ma. The maximum difference between interpretations i), ii) and iv) is 0.15%, close to desired reproducibility value of 0.1% for natural reference materials (Schaltegger et al., 2021). More rigorous filtering based on Pb\*/Pb<sub>c</sub> thus yields a more coherent dataset between the three studies, however,  
365 at low n-value.

Therefore, we propose that the weighted mean age of the youngest cluster of three high-Pb\*/Pb<sub>c</sub> analyses at 466.37±0.14/0.18/0.53 Ma (analytical/ +tracer/ +decay constant uncertainty; MSWD=1.9, n=3) has the highest probability for being an accurate age estimate for the ‘Likhall’ bed.

## 5.6 Implications for the Ordovician timescale and the absolute timing of events

370 Our new age for the ‘Likhall’ bed provides a carefully considered and well constrained tie point in the overall Ordovician timescale and event-stratigraphic framework. In the Geological Time Scale 2020 (GTS2020), the level corresponding to ‘Likhall’, i.e., the basal *Yangtzeplacognathus crassus* conodont Zone, is placed at c. 469 Ma (Goldman et al., 2020, figure 20.3). This is well outside our new age estimate of 466.37±0.14/0.18/0.53 Ma (~~as well as and also those of those of~~ Lindskog et al., 2017 and Liao et al., 2020), ~~which may and thus~~ warrants ~~some~~ adjustment of the Ordovician timescale – especially ~~so~~  
375 ~~as given that~~ the range of the *Y. crassus* Zone in the GTS2020 scheme does not even overlap with our age estimate. The revised age for ‘Likhall’ further suggests that the timing of the L-chondrite breakup event in space, as interpreted based on cosmic-ray exposure age data from ‘fossil’ meteorites ~~and chromite abundance~~, should be placed at c. 467.4 Ma (cf. Korochantseva et al. 2007; Heck et al. 2008; Lindskog et al. 2017; Liao et al. 2020; and references therein).



We refrain from extending our data interpretations until reliable U-Pb dates from both older and younger strata are available. Nonetheless, it is apparent from the relative timing in the rock record that the purported meteorite bombardment significantly postdates the onset of climate change and biodiversity spurts in the Middle Ordovician (e.g., Lindskog et al. 2017, 2023; Rasmussen et al. 2019). These stratigraphic offsets are unlikely to be bridged by numerical dating, but continued efforts to refine the Ordovician timescale will provide essential temporal context to the long-term development of the geobiosphere.

## 6 Conclusions

- 1) Our new high-precision  $^{206}\text{Pb}/^{238}\text{U}$  data set of the ‘Likhall’ zircon population represents an analytical improvement over previously published results: (a) our new data show higher  $\text{Pb}^*/\text{Pb}_c$  and therefore are less affected by the choice of isotopic composition used for blank correction. (ab) higher analytical precision allows for identification of normal discordance in young analyses, (bc) the presence of residual Pb-loss despite application of the currently considered optimal parameters for chemical abrasion (210°C for 12hrs; Widmann et al., 2019) suggests that previous data acquired at lower chemical abrasion temperatures were (likely more) affected by Pb-loss domains; ~~(c) our new data show higher  $\text{Pb}^*/\text{Pb}_c$  and therefore are less affected by the choice of isotopic composition used for blank correction.~~ The elevated  $\text{Pb}_c$  in the former studies is suggested to be due to incomplete removal of  $\text{Pb}_c$ -rich inclusions during chemical abrasion.
- 2) The choice of the data interpretation strategy is the main reason for the discrepancies between datasets from the three different laboratories (Lindskog et al., 2017; Liao et al., 2020; and the present study), particularly in cases where elevated  $\text{Pb}_c$  causes decreased analytical precision. Further reduction of the reproducibility between these studies is caused by the use of different tracer solutions and their respective calibrations.
- 3) The difference in interpreted age drastically ~~decreased~~decreases when only data subsets with high  $\text{Pb}^*/\text{Pb}_c$  are considered. For the data sets considered here, this implies an empirical  $\text{Pb}^*/\text{Pb}_c$  threshold of 50. For analyses with  $\text{Pb}^*/\text{Pb}_c > 50$ , the weighted mean (youngest cluster) U–Pb zircon age for the ‘Likhall’ bed is  $466.37 \pm 0.14/0.18/0.53$  (analytical/+tracer/+decay constant uncertainties), suggesting that previously published U–Pb ages are inaccurate.
- 4) The efficient removal of Pb-loss domains and micro inclusions is of paramount importance for achieving an accurate U–Pb date. Considering the presence of discordant, young analyses in our Ordovician-age dataset despite chemical abrasion conditions that are considered to be optimal, there is incentive to further develop the chemical abrasion procedure.
- 5) Considering biostratigraphic aspects, compared to GTS2020, the revised age of the ‘Likhall’ bed necessitates significant internal adjustment of the Ordovician timescale.

## Author Contributions

Conceptualization: André N. Paul, Anders Lindskog, Urs Schaltegger

Data Curation & Formal Analysis: André N. Paul

Funding Acquisition: Anders Lindskog and Urs Schaltegger

410 Investigation: André N. Paul, Anders Lindskog, Urs Schaltegger

Visualization: André N. Paul and Anders Lindskog

Writing – Original Draft Preparation: André N. Paul

Writing – Review & Editing: André N. Paul, Anders Lindskog, Urs Schaltegger

### **Declaration of Competing Interests**

415 The authors declare that they have no conflict of interest.

### **Acknowledgments**

A.L. acknowledges funding from the Birgit and Hellmuth Hertz' Foundation and the Royal Physiographic Society of Lund. This study benefitted from SNFNS grant CRSII5\_180253 awarded to U.S.

### **References**

- 420 Ahlberg, P., Calner, M., Lehnert, O., Wickström, L., and Lindskog, A.: Regional geology of Västergötland Province, Sweden, in: Field guide for the ISOS 14 post-conference excursion. Geologiska Föreningen Specialpublikation, 9–12, 2023.
- Ballo, E. G., Augland, L. E., Hammer, Ø., and Svensen, H. H.: A new age model for the Ordovician (Sandbian) K-bentonites in Oslo, Norway, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 520, 203–213, <https://doi.org/10.1016/J.PALAEO.2019.01.016>, 2019.
- 425 Bergström, S. M.: Use of graphic correlation for assessing event-stratigraphic significance and trans-Atlantic relationships of Ordovician K-bentonites, *Proc. Acad. Sci. Est. SSR*, 38, 55–59, 1989.
- Bowring, J. F., McLean, N. M., and Bowring, S. A.: Engineering cyber infrastructure for U-Pb geochronology: Tripoli and U-Pb-Redux, *Geochemistry, Geophys. Geosystems*, 12, <https://doi.org/10.1029/2010GC003479>, 2012.
- Caricchi, L., Simpson, G., and Schaltegger, U.: Estimates of volume and magma input in crustal magmatic systems from zircon  
430 geochronology: The effect of modeling assumptions and system variables, *Front. Earth Sci.*, 4, 48, <https://doi.org/10.3389/FEART.2016.00048/BIBTEX>, 2016.
- Condon, D. J., Schoene, B., McLean, N. M., Bowring, S. A., and Parrish, R. R.: Metrology and traceability of U-Pb isotope dilution geochronology (EARTHTIME Tracer Calibration Part I), *Geochim. Cosmochim. Acta*, 164, 464–480, <https://doi.org/10.1016/j.gca.2015.05.026>, 2015.
- 435 Gerstenberger, H. and Haase, G.: A highly effective emitter substance for mass spectrometric Pb isotope ratio determinations, *Chem. Geol.*, 136, 309–312, [https://doi.org/10.1016/S0009-2541\(96\)00033-2](https://doi.org/10.1016/S0009-2541(96)00033-2), 1997.

- Goldman, D., Sadler, P. M., Leslie, S. A., Agterberg, F. P., and Gradstein, F. M.: The Ordovician Period, in: *Geologic Time Scale 2020*, edited by: Gradstein, F. M., Ogg, J. G., Schmitz, M. D., and Ogg, G. M., Elsevier, 631–694, <https://doi.org/10.1016/C2020-1-02369-3>, 2020.
- 440 [Heck, P.R., Schmitz, B., Baur, H. & Wieler, R. 2008: Noble gases in fossil micrometeorites and meteorites from 470 Myr old sediments from southern Sweden, and new evidence for the L-chondrite parent body breakup event. \*Meteoritics and Planetary Science\* 43, 517–528.](#)
- Huyskens, M. H., Zink, S., and Amelin, Y.: Evaluation of temperature-time conditions for the chemical abrasion treatment of single zircons for U-Pb geochronology, *Chem. Geol.*, 438, 25–35, <https://doi.org/10.1016/J.CHEMGEO.2016.05.013>, 2016.
- 445 Keller, B. C., Boehnke, P., Schoene, B., and Harrison, T. M.: Stepwise chemical abrasion-isotope dilution-thermal ionization mass spectrometry with trace element analysis of microfractured Hadean zircon, *Geochronology*, 1, 85–97, <https://doi.org/10.5194/GCHRON-1-85-2019>, 2019.
- Keller, C. B., Schoene, B., and Samperton, K. M.: A stochastic sampling approach to zircon eruption age interpretation, *Geochemical Perspect. Lett.*, 8, 31–35, <https://doi.org/10.7185/GEOCHEMLET.1826>, 2018.
- 450 [Liao, S., Huyskens, M. H., Yin, Q.-Z., Schmitz, B. \(2020\). Absolute dating of the L-chondrite parent body breakup with high-precision U–Pb zircon geochronology from Ordovician limestone. \*Earth and Planetary Science Letters\*, 547, 116442. <https://doi.org/10.1016/j.epsl.2020.116442>](#)
- Lindskog, A. and Eriksson, M. E.: Megascopic processes reflected in the microscopic realm: sedimentary and biotic dynamics of the Middle Ordovician “orthoceratite limestone” at Kinnekulle, Sweden, *GFF*, 139, 163–183, <https://doi.org/10.1080/11035897.2017.1291538>, 2017.
- Lindskog, A., Costa, M. M., Rasmussen, C. M., Connelly, J. N., and Eriksson, M. E.: Refined Ordovician timescale reveals no link between asteroid breakup and biodiversification, *Nat. Commun.* 2017 81, 8, 1–8, <https://doi.org/10.1038/ncomms14066>, 2017.
- 460 [Lindskog, A., Young, S.A., Bowman, C.N., Kozik, N.P., Newby, S.M., Eriksson, M.E., Pettersson, J., Molin, E. & Owens, J.D. 2023: Oxygenation of the Baltoscandian shelf linked to Ordovician biodiversification. \*Nature Geoscience\* 16, 1047–1053](#)
- Lindstrom, M.: Volcanic contribution to Ordovician pelagic sediments, *J. Sediment. Res.*, 44, 287–291, <https://doi.org/10.1306/74D72A13-2B21-11D7-8648000102C1865D>, 1974.
- Mattinson, J. M.: Zircon U–Pb chemical abrasion (“CA-TIMS”) method: Combined annealing and multi-step partial dissolution analysis for improved precision and accuracy of zircon ages, *Chem. Geol.*, 220, 47–66, <https://doi.org/10.1016/J.CHEMGEO.2005.03.011>, 2005.
- McKanna, A. J., Koran, I., Schoene, B., and Ketcham, R. A.: Chemical abrasion: the mechanics of zircon dissolution, *Geochronology*, 5, 127–151, <https://doi.org/10.5194/gchron-5-127-2023>, 2023a.
- McKanna, A. J., Schoene, B., and Szymanowski, D.: Geochronological and Geochemical Effects of Zircon Chemical  
470 Abrasion: Insights from Single-Crystal Stepwise Dissolution Experiments , *Geochronology*, 2023b.

- McLaughlin, P. I., Normore, L., Sell, B. K., and Ramezani, J.: Ordovician tephra distribution, tephrochronology and geochronology, *Geol. Soc. London, Spec. Publ.*, 532, 79–90, <https://doi.org/10.1144/SP532-2022-267>, 2023.
- McLean, N. M., Bowring, J. F., and Bowring, S. A.: An algorithm for U-Pb isotope dilution data reduction and uncertainty propagation, *Geochemistry, Geophys. Geosystems*, 12, 0–18, <https://doi.org/10.1029/2010GC003478>, 2011.
- 475 McLean, N. M., Condon, D. J., Schoene, B., and Bowring, S. A.: Evaluating uncertainties in the calibration of isotopic reference materials and multi-element isotopic tracers (EARTHTIME Tracer Calibration Part II) - ScienceDirect, *Geochim. Cosmochim. Acta*, 164, 481–501, 2015.
- Mezger, K. and Krogstad, E. J.: Interpretation of discordant U-Pb zircon ages: An evaluation, *J. Metamorph. Geol.*, 15, 127–140, <https://doi.org/10.1111/J.1525-1314.1997.00008.X>, 1997.
- 480 Nielsen, A. T., Ove, J., Ebbestad, R., Hammer, Ø., Alexander, D., Harper, T., Lindskog, A., Mac, C., Rasmussen, Ø., and Stouge, S.: The Ordovician of Scandinavia: a revised regional stage classification, *Geol. Soc. London, Spec. Publ.*, 532, 267–315, <https://doi.org/10.1144/SP532-2022-157>, 2023.
- von Quadt, A., Jörn-Frederik Wotzlaw, Yannick Buret, E. Large, S. J., Irena Peytcheva, and Anne Trinquier: High-precision zircon U/Pb geochronology by ID-TIMS using new  $10^{-13}$  ohm resistors, *J. Anal. At. Spectrom.*, 31, 658–665, <https://doi.org/10.1039/C5JA00457H>, 2016.
- 485 [Rasmussen, J.A., Thibault, N. & Rasmussen, C.M.Ø. 2021: Middle Ordovician astrochronology decouples asteroid breakup from glacially-induced biotic radiations. \*Nature Communications\* 12, 6430.](#)
- Samperton, K. M., Schoene, B., Cottle, J. M., Brenhin Keller, C., Crowley, J. L., and Schmitz, M. D.: Magma emplacement, differentiation and cooling in the middle crust: Integrated zircon geochronological–geochemical constraints from the Bergell
- 490 Intrusion, Central Alps, *Chem. Geol.*, 417, 322–340, <https://doi.org/10.1016/J.CHEMGEO.2015.10.024>, 2015.
- Schaltegger, U., Ovtcharova, M., Gaynor, S. P., Schoene, B., Wotzlaw, J.-F., Davies, J. F. H. L., Farina, F., Greber, N. D., Szymanowski, D., and Chelle-Michou, C.: Long-term repeatability and interlaboratory reproducibility of high-precision ID-TIMS U–Pb geochronology, *J. Anal. At. Spectrom.*, 36, 1466–1477, <https://doi.org/10.1039/D1JA00116G>, 2021.
- Schmitz, M. D. and Schoene, B.: Derivation of isotope ratios, errors, and error correlations for U-Pb geochronology using  $^{205}\text{Pb}$ - $^{235}\text{U}$ -( $^{233}\text{U}$ )-spiked isotope dilution thermal ionization mass spectrometric data, *Geochemistry, Geophys. Geosystems*, 8, <https://doi.org/10.1029/2006GC001492>, 2007.
- 495 [Schmitz, B., Harper, D.A.T., Peucker-Ehrenbrink, B., Stouge, S., Alwmark, C., Cronholm, A., Bergström, S.M., Tassinari, M. & Xiaofeng W. 2008: Asteroid breakup linked to the Great Ordovician Biodiversification Event. \*Nature Geoscience\* 1, 49-53.](#)
- 500 [Schmitz, B., Farley, K.A., Goderis, S., Heck, P.R., Bergström, S.M., Boschi, S., Claeys, P., Debaille, V., Dronov, A., van Ginneken, M., Harper, D.A.T., Iqbal, F., Friberg, J., Liao, S., Martin, E., Meier, M.M.M., Peucker-Ehrenbrink, B., Soens, B., Wieler, R. & Terfelt, F. 2019: An extraterrestrial trigger for the mid-Ordovician ice age: Dust from the breakup of the L-chondrite parent body. \*Science Advances\* 5, eaax4184.](#)
- Schoene, B., Crowley, J. L., Condon, D. J., Schmitz, M. D., and Bowring, S. A.: Reassessing the uranium decay constants for geochronology using ID-TIMS U-Pb data, *Geochim. Cosmochim. Acta*, 70, 426–445,

505 <https://doi.org/10.1016/j.gca.2005.09.007>, 2006.

Servais, T. and Harper, D. A. T.: The Great Ordovician Biodiversification Event (GOBE): definition, concept and duration, *Lethaia*, 51, 151–164, <https://doi.org/10.1111/LET.12259>, 2018.

Watson, E. B., Cherniak, D. J., Hanchar, J. M., Harrison, T. M., and Wark, D. A.: The incorporation of Pb into zircon, *Chem. Geol.*, 141, 19–31, 1997.

510 Weber, G., Caricchi, L., Arce, J. L., and Schmitt, A. K.: Determining the current size and state of subvolcanic magma reservoirs, *Nat. Commun.* 2020 111, 11, 1–14, <https://doi.org/10.1038/s41467-020-19084-2>, 2020.

Widmann, P., Davies, J. H. F. L., and Schaltegger, U.: Calibrating chemical abrasion: Its effects on zircon crystal structure, chemical composition and UPb age, *Chem. Geol.*, 511, 1–10, <https://doi.org/10.1016/J.CHEMGEO.2019.02.026>, 2019.

515