# Analysing the sensitivity of a blowing snow model (SnowPappus) to precipitation forcing, blowing snow, and spatial resolution.

Ange Haddjeri<sup>1</sup>, Matthieu Baron<sup>1</sup>, Matthieu Lafaysse<sup>1</sup>, Louis Le Toumelin<sup>1</sup>, César Deschamps-Berger<sup>2,4</sup>, Vincent Vionnet<sup>3</sup>, Simon Gascoin<sup>4</sup>, Matthieu Vernay<sup>1</sup>, and Marie Dumont<sup>1</sup>

<sup>1</sup>Univ. Grenoble Alpes, Université de Toulouse, Météo-France, CNRS, CNRM, Centre d'Études de la Neige, Grenoble, France
 <sup>2</sup>Instituto Pirenaico de Ecología, Consejo Superior de Investigaciones Científicas (IPE-CSIC), Zaragoza, Spain
 <sup>3</sup>Meteorological Research Division, Environment and Climate Change Canada, Dorval, QC, Canada
 <sup>4</sup>Centre d'Études Spatiales de la Biosphère, CESBIO, Univ. Toulouse, CNES/CNRS/INRAE/IRD/UPS, Toulouse, France
 Correspondence: ange.haddjeri@umr-cnrm.fr

Abstract. Accurate snow cover modeling is a high stake for mountain regions. Alpine snow evolution and spatial variability result from a multitude of complex processes including interactions between wind and snow. The SnowPappus blowing snow model was designed to add blowing snow modeling capabilities to the SURFEX/Crocus simulation system for applications across large spatial and temporal extents. This paper presents the very-first spatialized evaluation of this simulation system

- 5 over a 902 km<sup>2</sup> domain in the French Alps. Here we compare snow cover simulations to the spatial distribution of snow height obtained from Pleiades Pléiades satellites stereo-imagery and to Snow Melt-Out Dates snow melt-out dates from Sentinel-2 and Landsat 8 time series over three snow seasons. We analyzed the sensitivity of the simulations to three different precipitation datasets and two horizontal resolutions. The evaluations are presented as a function of elevation and landform types. The results show that the SnowPappus model forced with high-resolution wind fields enhances the snow cover spatial variability at high
- 10 elevations allowing a better agreement with observations between observed and simulated spatial distributions above 2500 m and near peaks and ridges. Model improvements are not obvious at low to medium altitudes where precipitation errors are the prevailing uncertainty. Our study illustrates the necessity to consider error contributions from blowing snow, precipitation forcings, and unresolved subgrid variability for robust evaluations of spatialized snow simulations. Despite the significant effect of the unresolved spatial scales of snow transport, 250 m horizontal resolution snow simulations using SnowPappus are found 15 to be a promising avenue for large-scale modeling of alpine snowpacks.
  - 1 Introduction

Snow cover in mountainous terrains is characterized by an important variability at multiple temporal and spatial scales (Pomeroy and Gray, 1995; Clark et al., 2011; Anderson et al., 2014; Mott et al., 2018). This variability results from a large diversity of slopes, aspects, and elevations which induce a high variability of precipitation, wind, temperature, and radiation.

20 Accurate snowpack modeling is key to describing this high variability in hydrological applications, climate projections, and hazard forecasting in mountainous terrains (IPCC, 2022; Morin et al., 2020).

The spatial variability of precipitation amount and phase at different scales is known to be one of the main sources of snowpack variability at the mountain range scale (1-100 km) (Clark et al., 2011). These patterns are specific to the regional topography and atmospheric flow, with regions of increased or decreased precipitation (Colle et al., 2013). At the slope scale

- 25 (a few hundred meters), preferential deposition and snowfall enhancement are the main processes responsible for snowfall variability (before snowflake settlement) (Mott et al., 2018). The preferential deposition is the result of the interaction of the near-surface flow field with particle trajectories, creating areas of local accumulation (Lehning et al., 2008). Snowfall enhancement is a process in which surface flows are responsible for locally increased air moisture, leading to the formation or maintenance of low-level clouds, ultimately enhancing solid precipitations through a seeder-feeder mechanism (Bergeron,
- 30 1965; Choularton and Perry, 1986; Minder et al., 2011). Post-depositional processes finally, determine the snow variability at the slope scale and below. Blowing snow transport, snow sublimation, snow redistribution by avalanches, snow compaction, and melt are the main processes at work (Winstral et al., 2002; Bernhardt and Schulz, 2010; Mott et al., 2018). Blowing snow transport produces a mass transfer of snow from windward areas to leeward deposition zones. Additional mass loss occurs in those events due to the sublimation of suspended snow Liston and Sturm (1998); Yang et al. (2010). The heterogeneity of snow
- 35 mass loss due to snow melt (Brauchli et al., 2017) and surface sublimation (Pomeroy et al., 1998; Strasser et al., 2008) also contribute to snowpack variability in alpine terrain.

The multi-scale variability of the alpine snowpack makes it challenging to represent in numerical models. Various approaches have been developed for that purpose. Numerical Weather Prediction (NWP) models at the kilometer scale or higher resolution models can explicitly represent the orographic precipitation patterns, and in some cases local low-level cloud for-

- 40 mation, triggering snowfall enhancement (Lehning et al., 2006; Vionnet et al., 2017; Wang and Huang, 2017; Monteiro et al., 2022). However, the quantification of the precipitation is still impacted by important uncertainties with errors in precipitation amounts and phases, localization, and timing increased with coarse grid size (Clark et al., 2011; Ménard et al., 2019; Lundquist et al., 2019). Downscaling tools can be used to better represent the local meteorology from NWP models (Sen Gupta and Tarboton, 2016; Marsh et al., 2023; Bernhardt et al., 2010; Mital et al., 2022). Eventually, the representation of post-depositional
- 45 processes, notably-

Due to the complex intertwining of snow variability processes, in particular blowing snow transport, benefits from the community found benefits in the development of dedicated high-resolution models coupled to (e.g. Vionnet et al., 2014; Sharma et al., 2021) or forced by atmospheric models (e.g., Lehning et al., 2006; Liston et al., 2007; Marsh et al., 2020; Sharma et al., 2021; Baron et al., 2024

- 50 (e.g., Lehning et al., 2006; Liston et al., 2007; Marsh et al., 2020; Baron et al., 2024; Quéno et al., 2023). The spatial evaluation of this last-type of system, dedicated to modeling part of the observed snow spatial variability is a challenge in itself. The snow modeling community has long been evaluating this kind of model locally. Evaluations can be carried out using direct measurements of the variable of interest, for example for blowing snow modeling, Vionnet et al. (2014) evaluated directly the simulated blowing snow fluxes of the Meso-NH/Crocus system against locally measured fluxes from a Snow Particle Counters
- 55 (Sato et al., 1993). Amory et al. (2021) also compares simulated snow transport occurrence and mass fluxes to on-field observations in Antarctica. Similarly, Baron et al. (2024) compares the simulated snow transport occurrence and transport mass

fluxes from the SnowPappus model to field observations. One of the main drawbacks of this method is the low number of direct snow transport observations found in literature, combined with the very high spatial variability of snow transport fluxes. As a result, spatialized snow simulations are more classically evaluated using measurements of snow height and Snow Water

- 60 Equivalent (SWE). Prasad et al. (2001) and Liston et al. (2007) compare snow simulations done using the SnowTran-3D system (Liston and Sturm, 1998; Liston et al., 2007) with SWE measurements from manual snow surveys. Marsh et al. (2020) compare simulated SWE from the Canadian Hydrological Model (CHM) to point observations. Mott et al. (2008) evaluate the Alpine3D system (Lehning et al., 2006) comparing simulation with an interpolated map of snow height measurements. The use of interpolation methods in Mott et al. (2008) is justified by the high measurement density of the study area. However, in the
- 65 majority of other study areas, in situ measurements are way too sparse to characterize the complex spatial variability of alpine snow cover (Pepin et al., 2015; Bales et al., 2006; Vernay et al., 2022; Pomeroy et al., 2009). This situation explains the recent use of snow remote sensing (unmanned aerial vehicle, satellite) as a means of spatial evaluation for large-scale snow cover models. Those methods are still limited by the low availability of remotely sensed variables, mainly snow cover fraction and derived variables or snow height. For example, Vionnet et al. (2021) compare CHM snow simulations (Marsh et al., 2020) with
- 70 airborne Lidar snow height maps as well as Sentinel 2 snow cover maps. Very recently, (Quéno et al., 2023) used Lidar aquired snow height maps to evaluate the ability of the FSM2oshd framework (Quéno et al., 2023) of representing snow accumulation and erosion areas.

In the above-cited studies using the snow height, SWE, or snow presence related variables, the evaluation experiments are generally carried out using a single set of meteorological inputs. However, it is known that meteorological inputs explain an

- 75 important part of the variability (Clark et al., 2011; Colle et al., 2013; Mott et al., 2018) and uncertainty Raleigh et al. (2015); Günther et al. (2019) in simulated snow depth height and SWE. While the evaluations of model inputs and simulated processes are usually conducted separately, they are often interdependent. Thus, it is difficult to determine if potential errors come from the model or the input variable. For instance, a good simulation of snow height requires both accurate precipitation input and a robust snow evolution model. In order to To robustly assess the value of a distributed snow modelin the presence of
- 80 uncertainty in the meteorological forcing , this uncertainty has to be accounted for to raise robust conclusions, the uncertainty in meteorological forcing must be considered.

To improve the snow spatial variability of the French snow modeling system, Baron et al. (2024) have developed a novel explicit blowing snow transport model, SnowPappus, coupled with the Crocus physical snow simulation model (Vionnet et al., 2012). This model explicitly represents the vertically integrated saltation and suspension mass fluxes, as a function of wind

speed and surface snowpack properties simulated by Crocus. This system is implemented on a regular grid to simulate snowpack evolution at a mountain range scale (approx.  $100000 \text{ km}^2$ ), for multiple snow seasons. Baron et al. (2024) focused on the evaluation of the model at point scale, demonstrating its ability to accurately simulate blowing-snow fluxes and occurrence at observation stations. However, spatial evaluations of simulations on large distributed domains are currently missing.

The goal of this paper is to present a spatial evaluation of the SnowPappus blowing snow simulation framework (Baron et al., 2024), considering the uncertainty of precipitation estimates through different data sources that exhibit contrasted spatial patterns. The evaluation is based on Pleiades snow depth-Pléiades snow height maps (Deschamps-Berger et al., 2020) and Sentinel-2/Landsat 8 snow melt-out dates (Gascoin et al., 2019) and covers three consecutive snow seasons. We discuss the relative influence of precipitation sources, and blowing snow implementation on the simulated snow cover variability and how their interactions can affect the evaluation of a distributed snow model. By using two contrasted horizontal resolution

95

simulations, we also analyze how the unresolved spatial scales of blowing snow can affect simulation results at a 250 m spatial scale. Finally, we emphasize the main challenges to be solved for more advanced evaluations of spatialized snow simulations.

#### 2 Data and Methods

#### 2.1 Study area

The simulation study site is located east of Grenoble in the French Alps (Fig. 1). It covers 902 km<sup>2</sup> including the GrandesRousses and Arves massifs. This area exhibits a complex topography, with elevations ranging from 700 up to 3900 meters a.s.l., a wide range of snow and temperature conditions, and different types of landscape features such as valleys, forests, alpine pastures, lakes, and glaciers. In this area, the local knowledge of precipitation flow patterns tells us that most winter storms come from North-Western flows usually giving increased precipitation on western slopes. The study area includes the Col du Lac Blanc observatory (Guyomarc'h et al., 2019) where Baron et al. (2024) evaluated the blowing snow fluxes and occurrence simulated by SnowPappus. It counts 41 distinct ice patches or glaciers of various sizes as defined in the RGI

Consortium (2017), forested areas, and three hydroelectric dams which underline the hydrological importance of the area. As our simulation system is only able to represent snow in open areas, the forests, glaciers, lakes, and rivers in the study zone are masked in our simulations and observation datasets. For the forest mask, the BD FORET® V2 dataset has been used (IGN©, 2021a) with a masking threshold of 25% of forested sub-pixel area. Waterways are masked following data from BD

110 TOPO® (IGN©, 2021b). In the valley, the urban areas around the city of Bourg d'Oisans, Allemont, and Saint Michel de Maurienne are also discarded from the analysis.

#### 2.2 Grid generation and spatial resolution

The simulations are based on two grid resolutions, 30 m and 250 m. Both simulation grids are built using as reference the French 5 m RGE ALTI® digital elevation model (DEM) from IGN© (2021c). The 5 m high-resolution DEM is resampled us-

115 ing the average method to 30 m and 250 m horizontal resolutions using GDAL/OGR contributors (2023). At 250 m resolution, the 902 km<sup>2</sup> full simulation area is composed of 14 443 grid points. At 30 m horizontal resolution, the test zone is composed of 1 005 699 simulation points.

A geomorphons classification (or landform) is performed on the 250 m resolution DEM, for a more detailed analysis of simulation domain features. Geomorphons introduced by Jasiewicz and Stepinski (2013) are defined as the fundamental structural elements of a landscape. Here, we used the Whitebox Geospatial (2023) classification tool, an open-source software from Lindsay (2014). The algorithm is based on a line-of-sight analysis with similarities to the more classical topographic posi-



**Figure 1.** Map of the simulation domain (red) and evaluation areas (<u>Pleiades Pléiades</u> green and blue, <u>Sentinel 2 snow melt out date in</u> red). Points of interest are shown as well as forest, glacier, and lake masks. The classification of 250 m pixels is visible with a 300 m step elevation in shades of blue. The summer aerial photography base map is from IGN© (2022).

tion index (TPI). This method identifies a set of topographic patterns corresponding to specific terrain attributes and landform types. The advantage of this type of classification is that it allows topographic information to be conveyed mimicking the
result of a classification process carried out by a human analyst. Another advantage of the geomorphons classification is that it adapts to the surrounding terrain and can lead to the identification of landform elements regardless of their scale. Jasiewicz and Stepinski (2013) identify the 10 most common landform elements used for geomorphons classification (peak (summit), ridge, shoulder, spur (convex), slope, hollow (concave), footslope, valley, pit (depression) and flat, illustration can be found in Fig. 3 of Jasiewicz and Stepinski (2013)). This allows us to generate a simple, intuitive, and scale-independent landform map of our simulation domain (Jasiewicz and Stepinski, 2013). Figure 2 illustrates the geomorphons classification result (called landform

classification in the following) of each 250 m grid cell of our simulation domain in terms of the 10 most common landform

elements. Figure 3 illustrates the elevation distribution of the landform elements of our simulation domain. No landform feature corresponding to Jasiewicz and Stepinski (2013) 'Shoulder', 'Footslope', and 'Flat' are found in our simulation domains. Finally, Appendix Fig. A1 demonstrates the good representativeness of the landform features frequency of our domain compared to the entire French Alps and Pyrenees.

For our analysis, each 250 m simulation grid cell is classified in terms of landform types (see Fig. 2) and into 300 m elevation bands (see Fig. 1).



**Figure 2.** Map showing the result of the landform classification performed on our 250 m simulation domain. Each grid cell is classified in terms of the 10 most common landforms described by Jasiewicz and Stepinski (2013). (Peak (summit), ridge, shoulder, spur (convex), slope, hollow (concave), footslope, valley, pit (depression), flat) In our domain, "flat", "footslope", and "shoulder" landforms are not present. The summer aerial photography base map is from IGN© (2022).



Figure 3. Distribution of simulation pixels in terms of 300m step elevation bands and the landform classification. The number of pixels corresponding to each class is shown.

# 2.3 Crocus snow model

140

This paper uses the well-established 1D snow model Crocus (Brun et al., 1989; Vionnet et al., 2012), used operationally by the French weather forecast agency (Météo-France), in support of avalanche hazard forecasting (Morin et al., 2020). This snow model relies on a dynamical layering of the snowpack and an explicit representation of snow metamorphism and microstructure up to 50 snow layers (Vionnet et al., 2012). The Crocus snow model is coupled to the ISBA ground and land cover model and is embedded in the SURFEX modeling framework (Decharme et al., 2011; Masson et al., 2013). The snowpack is simulated on a squared grid of 250 m or 30 m spacing. Simulations are run over 3 years, starting from 02 August 2017 06:00 to 02 June 2020 06:00, using a 15-minute time step. The initial ground conditions are taken from a 10-year (2007-2017) simulation driven by the SAFRAN reanalysis (Vernay et al., 2022).

#### 2.4 SnowPappus blowing snow model

The SnowPappus blowing snow model has been specially developed to describe snow variability due to blowing snow when the Crocus model is applied in the range of 25 m to 250 m horizontal resolutions (Baron et al., 2024). SnowPappus model simulates

150 blowing snow occurrence, horizontal transport fluxes, and blowing snow sublimation rate on each grid cell as a function of 2D atmospheric forcing and snow surface properties. Then, a blowing snow mass balance is used to quantify eroded or accumulated snow amounts and modify the simulated snow profiles accordingly. Model methods and parameterizations used to represent the different blowing snow processes are described in detail and discussed against existing literature in Baron et al. (2024).

SnowPappus can run over multiple snow seasons and very large domains , in the (about 10<sup>5</sup> km<sup>2</sup> (at 250 m resolution or 10<sup>6</sup>
 simulation points), in a reasonable computing time (less than half a day on a single computing node, depending on simulation outputs).

#### 2.5 Meteorological forcings

#### 2.5.1 SAFRAN simulations

As most spatialized applications of the Crocus snow model over the French Alps, our simulations are forced by the SAFRAN

- 160 meteorological reanalysis (Vernay et al., 2022). The SAFRAN analysis system uses as a background, vertical profiles of air temperature, humidity, and the precipitation fields of the ARPEGE NWP system and assimilates near-surface meteorological observations to generate, with an hourly time step, the typical sets of inputs required by a snow model such as Crocus (Durand et al., 1993). The SAFRAN reanalysis provides these meteorological inputs over 23 climatologically homogeneous areas (named massifs) covering the whole French Alps (Vernay et al., 2022). Here, only the SAFRAN reanalysis corresponding
- to the Grandes-Rousses massif was used to build the simulation meteorological forcing. For a distributed use, the SAFRAN analysis is linearly interpolated as a function of elevation, for each simulation point (as in Vionnet et al. (2016)) to match the gridded geometry of our simulation domain. Solar radiations are projected according to the slope inclination and aspect of each pixel and masked in the case of shadows from the surrounding topography similarly to Revuelto et al. (2018) and Deschamps-Berger et al. (2022). In SAFRAN, all meteorological variables are assumed to be constant inside a massif for a
- 170 given elevation. Whereas large-scale biases are low thanks to the assimilation of numerous gauges observations (Vernay et al., 2022), a large part of the spatial variability of precipitation remains unresolved under this assumption (Vionnet et al., 2016; Quéno et al., 2016; Vionnet et al., 2019; Deschamps-Berger et al., 2022). Therefore, other precipitation estimates are also considered in this study, respectively from AROME NWP system and ANTILOPE radar-based analysis. An illustration of the mean daily SAFRAN precipitation spatial variability can be seen in Appendix Fig. B1 (a). This map informs us of the spatial
- 175 and inter-model variability and of the absolute precipitation values of each precipitation dataset.

#### 2.5.2 AROME precipitation simulations

AROME is the high-resolution non-hydrostatic NWP system operated by Meteo-France for short-range forecasts. It uses a 1.3 km grid spacing (Seity et al., 2011). Due to high temperature and radiative biases of AROME in mountain areas (Quéno et al., 2020; Gouttevin et al., 2023), and to focus our study on the uncertainty of precipitation spatial patterns rather than on

180 all meteorological forcing uncertainties, we only consider in our simulations only the precipitation amount from AROME the total precipitation amount (liquid and solid) from AROME in our simulations. Thus, the precipitation phase is set to solid if the SAFRAN air temperature is lower than 274.15 K and liquid if the air temperature is above 274.15 K. Although more advanced and continuous phase functions are available in the literature (Froidurot et al., 2014; Vionnet et al., 2022), this choice was made for consistency between all simulations, as this threshold is also used by the SAFRAN system. SAFRAN analysis data

are used for all the other forcing variables. An illustration of the mean daily AROME precipitations can be seen in Appendix

185

Fig. B1 (d).

#### 2.5.3 ANTILOPE precipitation simulations

ANTILOPE is an hourly precipitation estimation analysis product (rain rate), combining radar and rain/snow gauges with 0.01° (1 km) resolution (Champeaux et al., 2009). ANTILOPE precipitation fields are the kriging result of available gauge
measurements, using radar-estimated precipitation fields as external drift. In a mountainous context, the skill of the product can be affected by a partial or total mask of the radar beams and other common radar measurement errors (Faure, 2017; Yu et al., 2018), although the use of observations improves the precipitation estimates compared to raw radar precipitation. Since ANTILOPE doesn't discriminate solid from liquid precipitations, we used information from SAFRAN to classify rainfall and snowfall as well as all the other forcing variables. The precipitation phase is set to solid if the SAFRAN air temperature is lower than 274.15 K and liquid if the air temperature is above 274.15 K. SAFRAN analysis data are used for all the other forcing variables. An illustration of the mean daily ANTILOPE precipitations can be seen in Appendix Fig. B1 (c).

#### 2.6 DEVINE wind downscaling model

Wind fields are important drivers of turbulent and mass exchanges at the surface of the snowpack. Wind fields also profoundly determine the onset and evolution of drifting and blowing snow episodes. The interaction between mesoscale winds and local
topography induces strong modifications both in terms of speed and direction that deeply influence the spatial variability of wind fields at the local mountain scale (Whiteman, 2000). Neither interpolated NWP systems operating with a km resolution (Seity et al., 2011) nor massif-scale SAFRAN reanalyses (Vernay et al., 2022) would permit us to take into account the local influence of terrain on wind fields. As the local scale wind patterns are crucial to determine high-resolution snow patterns (Musselman et al., 2015), we downscaled AROME wind fields (1.3 km grid spacing) down to a 30 m grid using the DEVINE downscaling model (Le Toumelin et al., 2022). This method leverages a convolutional neural network to emulate the behavior

of the complex atmospheric model ARPS, previously run on a large number of synthetic topographies and in turn increase the spatial resolution of wind fields in complex terrain.

The DEVINE downscaling model has been previously evaluated over a large number of in-situ stations (Le Toumelin et al., 2022, 2023), including stations located in our study domain. It has been shown that DEVINE wind outputs can represent wind

210 acceleration along ridges and summits, deceleration windward, and some deflection around topographic obstacles but can not represent more complex wind patterns such as thermal winds or recirculation areas.

#### 2.7 Model experiments

To disentangle the impacts of the simulation of blowing snow from the precipitation forcing, 250 meter resolution simulations were run with and without the SnowPappus transport module and with the 3 different precipitation datasets, resulting in 6 different 250 m simulations. All simulations utilize the default options, the GM98 parametrization of blowing snow occurrence, and the flux limiter is activated (Guyomarc'h and Mérindol, 1998; Baron et al., 2024). In addition, blowing snow sublimation is activated using SBSM-like-Simplified Blowing Snow Model-like equations (Essery et al., 1999; Baron et al., 2024). To investigate the role of unresolved transport variability of snow transport, 2 additional simulations were done at 30 m resolution with and without the SnowPappus transport module (see HR standing for High Resolution). Due to the higher numerical cost of these simulations and the increased complexity of downscaling and validating weather forcing at this resolution, 30 m resolution simulations were only run with the SAFRAN precipitation fields.

As the 30 m resolution simulations are used in this paper to evaluate the contribution of unresolved spatial variability in 250 m simulations, we do not evaluate 30 m simulations at their native resolution but after a resampling on the 250 m resolution grid using ESMF first-order conservative method (Earth System Modeling Framework et al., 2023).

225

Regardless of the simulation configuration, 10-year SAFRAN spinup simulations (without SnowPappus) were performed to produce the ground's initial condition.

Experiment name	Precipitation forcing	Blowing snow transport mode	Computing resolution
Safran	SAFRAN	No transport	250m
Safran with transport	SAFRAN	Transport	250m
Safran HR	SAFRAN	No transport	30m
Safran HR with transport	SAFRAN	Transport	30m
Arome	AROME	No transport	250m
Arome with transport	AROME	Transport	250m
Antilope	Antilope	No transport	250m
Antilope with transport	Antilope	Transport	250m

A summary of all the simulation experiments is found in Table 1:

**Table 1.** Overview of the different simulation experiments, their model configurations and naming. Important note, the computing resolution is different from the evaluation resolution. All simulations are evaluated at 250m resolution (see Sect. 2.8).

#### 2.8 Reference dataset and evaluation methods

#### 2.8.1 Sentinel-2 snow Snow melt-out dates

- We use the annual snow melt-out dates (SMOD) distributed by Theia (level 3 product of the Snow collection Gascoin et al. (2019)). This product has a 20 m spatial resolution and indicates the last date of the longest continuous snow period. It is obtained by the linear interpolation in the time dimension of all the single-date (level 2) snow cover area products between 1 September and 31 August (Gascoin et al., 2019). These products are generated from Sentinel-2 and Landsat 8 images. The SMOD is given in days starting from September 1st. The Sentinel-2 SMOD obtained for our simulation domain is resampled
- 235 from the 20 m horizontal resolution-We resampled the SMOD to the 250 m resolution grid of our model using the Earth System Modeling Framework et al. (2023) (ESMF) first-order conservative method.

We use the same definition to compute the SMOD from the model output. To compute the longest continuous snow period, the snow height is first averaged by day and then a snow height threshold of 20 cm is applied, as this value was found to be optimal in the sensitivity analysis of Deschamps-Berger et al. (2022) at the same horizontal resolution. For this threshold, the

240 accuracy is found to be 91.5% (See Fig. 10 of Gascoin et al., 2019). The forest, glaciers, and lakes on the domain are masked out both for the Sentinel-2 observations and simulations using the mask defined in Fig. 1 and Sect. 2.1.

# 2.8.2 The Pleiades Pléiades stereo-imagery

In order to evaluate more directly the spatial variability of the simulated snow height, we used snow height maps derived from Pleiades\_Pléiades\_stereo-images (Deschamps-Berger et al., 2020; Marti et al., 2016). In both cases, snow height is defined

- 245 following a vertical axis. Pleiades The Pléiades raw images have a 0.5 m horizontal resolution and are acquired on request. Snow height maps are obtained by combining two surface elevations of the same area, one with snow and one snow-free. Each surface elevation map is derived from pairs or triplets of Pleiades images. A reference snow-free Pléiades images. The reference snow-off surface elevation is acquired during the summercalculated from Pléiades stereo-images acquired on 28 September 2016 for the Galibier area and 24 October 2018 for the Grandes Rousses area. A winter image acquisition gives
- 250 a second elevation map with snow on the ground. The difference between the two elevation fields provides a snow height map. The generated snow height map has a horizontal resolution of 2m. Two observation zones have been selected based on available images. The first one covers 66 km<sup>2</sup> around the Col du Lac Blanc experimental site in the Grand-Rousse mountains. The second observation zone covers 168 km<sup>2</sup> around the Lautaret experimental site in the Galibier mountains. Two dates with snow on the ground have been analyzed for each observation zone. For the Grandes Rousses area, two images are available
- 255 for a different year but at the same period of the season (Pleiades Pléiades 1 on 13 May 2019 and Pleiades Pléiades 2 on 4 May 2020). For the Galibier area, the two images give a view at different times of the same season (Pleiades Pléiades 3 on 23 January 2018 and Pleiades Pléiades 4 on 16 March 2018).

Because of <u>Due to</u> the method used to reconstruct the snow height map, usual <u>Pleiades Pléiades</u> observations contain a few no-data pixels where the quality of the observation is lower due to cliffs, steep slopes, clouds, or shaded areas. The snow height map was filtered to exclude values out of the [-0.5 m; 20 m] range. According to previous <u>Pleiades</u> Pléiades snow

maps evaluations (Deschamps-Berger et al., 2020), averaging 2 m Pleiades Pléiades observation resolution to 250 m reduces the vertical snow height standard error between to approximately 0.3 and 0.4 m. Negative snow heights are kept as removing them would slightly positively bias the averages. For comparison with snowpack simulations, the 2 m resolution Pleiades snow depth Pléiades snow height map is resampled to the 250 m horizontal resolution grid using a conservative method, similarly to

265 (Deschamps-Berger et al., 2020). The entire 250 m pixel is set to No Data if more than 70% of the 2 m pixels is No Data-, this threshold ensures the minimum representativeness of the pixel's snow height.

The forest, glaciers, and lakes on the domain are masked both for the <u>Pleiades Pléiades</u> observations and simulations using the mask defined in Sect. 2.1. An additional mask of the No Data pixels in the <u>Pleiades Pléiades</u> data is applied to the simulations for accurate comparison.

# 270 2.8.3 Observation data summary

All observations are summarized in Table 2. For the comparison with simulations, the observed SMOD and snow height are grouped by altitude in various elevation bands. The reference altitude used for the observation is the 250 m DEM described in Sect. 2.2. This elevation band classification is one of the most often used in the literature because the elevation is one of the main topographic drivers of snow cover variability (Grünewald et al., 2014; Vionnet et al., 2022; Monteiro et al., 2022; Tong et al., 2009; Deschamps-Berger et al., 2022; Vionnet et al., 2017). However, for a more advanced analysis of the snow cover

spatial structure, we also present our results grouped according to the landforms presented in Sect. 2.2.

275

280

Satellite observation	Observed area	Variable	Observation date	Native resolution	Aggregated resolution
Pléiades 1	Grandes Rousses region (GR)	Snow height (m)	13 May 2019	2 m	250 m
Pléiades 2	Grandes Rousses region (GR)	Snow height (m)	4 May 2020	2 m	250 m
Pléiades 3	Galibier region (Glb)	Snow height (m)	23 January 2018	2 m	250 m
Pléiades 4	Galibier region (Glb)	Snow height (m)	16 March 2018	2 m	250 m
Sentinel 2 A	Entire simulation domain	SMOD (days)	1 September 2017 to 1 september 2018	20 m	250 m
Sentinel 2 B	Entire simulation domain	SMOD (days)	1 September 2018 to 1 September 2019	20 m	250 m

**Table 2.** Summary of satellite observations used in this study. The observed area, variables, resolutions, and dates are detailed for each observation. The aggregated resolution reminds the resampling operation done on the observation data before comparison with simulations.

#### 2.9 Synoptic scores

To summarise the simulation and observation comparison, a combination of three scores has been chosen. The mean bias, standard deviation ratio, and Spatial Probability Score value (SPS) are used to quantify the similarity between the simulated and observed snow height distribution.

The mean bias (MB) defined in Eq. 1 describes the mean bias between the simulation and the observation over a given area.

$$MB(S) = \langle S \rangle - \langle \mathbb{O} \rangle \tag{1}$$

where  $\langle S \rangle$  represents the mean value of the simulation distribution and  $\mathbb{O}$  the observation distribution.

The standard deviation ratio (or  $\sigma$  ratio in the following) (Eq. 2) is the ratio between the simulated standard deviation and 285 the observed one.

$$\boldsymbol{\sigma} \operatorname{ratio}(\mathbf{S}) = \frac{\sigma(\mathbf{S})}{\sigma(\mathbf{O})}$$
(2)

with  $\sigma(S)$  the simulated standard deviation and  $\sigma(\mathbb{O})$  the observed standard deviation.

The Spatial Probability Score (SPS) is a distance metric between two Cumulative Distribution Functions (CDF) defined in Eq. 3 as the quadratic discrepancy measure between the simulated CDF, noted F and  $P_{obs}$  the empirical observed CDF.

290 
$$\underbrace{SPS}_{\mathbb{S}}(F) = \int_{\mathbb{S}} \left( F(x) - P_{obs}(x) \right)^2 dS.$$
(3)

with  $x \in S$ , and S the volume of possible values of the indicator of interest. This score is derived from the Continuous Ranked Probability Score (CRPS) primarily used to evaluate probabilistic forecasts in ensemble forecasting (Candille and Talagrand, 2005). Here, it is used to compare two spatial distributions, similarly to the description in Goessling and Jung (2018), although using a non-binary probability field for the perfect observation. In short, this metric quantifies the similarity of two probability densities. It is close to the Wasserstein distance (Rüschendorf, 1985) used in (Vionnet et al., 2021) although here of order 2. A

295

low SPS value means a small distance between the observed and the simulated distributions. Two identical distributions give an SPS score of 0. The SPS unit is identical to the unit of the target variable (m for snow height, days for SMOD).

$$\underline{SPS}(F) = \int_{\mathbb{S}} \left(F(x) - P_{obs}(x)\right)^2 \cdot \underline{dS}$$

with  $x \in S$ , and S the volume of possible values of the indicator of interest.

# 300 3 Results

#### 3.1 Spatial variability of the snow height

This section compares observed and simulated snow height from the 8 experiments described in Table 1 to the 4 Pleiades Pleiades observations (Table 2).

#### 3.1.1 Grandes Rousses area

305 Map of simulated and observed snow height on the 13 May 2019 10:00. (a) Safran 250 m simulation with snow transport and without snow transport in the left submap. (b) Arome 250 m simulation with snow transport and without snow transport and without snow transport in the left submap. (c) Antilope 250 m simulation with snow transport and without snow transport in the left submap. (d) The Pleiades 1 observed snow height resampled to 250 m. The absence of snow in simulations is indicated in grey. For d) the summer aerial photography base map is from IGN© (2022).

- 310 First, An example of simulated and observed snow height variability can be found in Fig. 4illustrates snow height simulations and observations. In this figure, snow height is simulated on 13 May 2019. It 2019 for the Safran, Arome, and Antilope precipitation forcings, and observed for the Grandes Rousses area on the same date. For each precipitation forcing, the subfigure shows from one side the impact of the SnowPappus blowing snow model on the simulated snow height, where we can see areas of increased and decreased snow height, most visible along the ridges and peaks, and from another side, the major impact of
- 315 precipitation forcing on the resulting snow height, more specifically for the AROME experiment Fig. 4 b)in each left sub-maps the same simulation without snow transport. It can be observed that the primary areas of model snow accumulation are located in regions of high observed snow height. However, the opposite is not true, many areas with high observed snow accumulation are not present in the simulations. Additional maps for the 13 May 2019 comparing resolutions can be found in Appendix Fig. C1.

# 320 3.1.1 Grandes Rousses area

In this section, we compare the two Pléiades snow heights images of the Grandes Rousses to simulations. For a quantitative evaluation of these snow height maps, we summarized simulations and observations in terms of spatial distributions. For each Pleiades Pléiades subdomain, simulated snow heights are discretized using 300 m elevation bands. Each snow height distribution describes the internal snow height spatial variability within each elevation band. The associated spatial distributions obtained in the Grandes Rousses region (green box in Fig. 1) are compared in Fig. 5 (a-b) for 13 May 2019 (same data as Fig.

- And 4 May 2020. The skill of these distributions is summarized through synthetic scores presented in Fig. 6 (a-b-c-d-e-f).
   Figure 5 (a) shows a major impact of the precipitation input on the simulated snow height distribution, regardless of the use of snow transport in simulations. Precipitation inputs impact the simulated snow height distribution's shape, variance, and mean value. Simulation with AROME precipitations leads to the highest snow height on average, for all elevation bands and
- the two evaluation dates. This result is especially notable for Fig. 5 (a) at the elevation band of 2500 to 2800 m on the 13 May 2019 evaluation. The AROME precipitation gives a mean snow height of about 3 m representing a positive bias compared to the Pleiades Pléiades observations mean (1.87 m), consistently with Fig. 6 (a). The standard deviation of the AROME simulations overestimates PleiadesPléiades' standard deviation at low elevations on both dates Fig. 6 (b-e). It is more realistic between 2500 and 2800 m, and dependent on snow transport at higher elevations.

335

325

Simulations with the ANTILOPE precipitation, on the other hand, have a negative snow height bias compared to the observation. For the 2500 to 2800 m elevation band, the ANTILOPE simulations give a mean snow height of only 1.30 m. This negative bias tends to increase with altitude as illustrated in Fig. 6 (a). The standard deviation of ANTILOPE simulations is closer to observations than AROME simulations. Similar results are found for 4 of May 2020 (Fig. 5 (b), Fig. 6 (d-e-f)). For

340 the SAFRAN and SAFRAN HR simulations, results are consistent between the simulation resolution (250m and 30m) but with varying skill between both evaluation years. The mean simulated snow height for the 2500 to 2800m elevation band (1.90 m) is consistent with observations on 13 May 2019. At all elevations, the SAFRAN and SAFRAN HR simulations show a lower bias compared to observations than simulations forced by AROME and ANTILOPE. However, the SAFRAN mean value of

# a) Safran



c) Antilope



b) Arome



Figure 4. Spatial distribution of Maps illustrating raw simulation outputs and Pléiades observed snow height on the 13 May 2019 10:00. (ma) for 300Safran 250 m elevation bands simulation with snow transport and without snow transport in Pleiades observations-the left sub-map. (grayb) and all Arome 250 m simulation configurations over with snow transport and without snow transport in the Grandes Rousses mountains on 13 May 2019 (Pleiades 1)left sub-map. (ac) Antilope 250 m simulation with snow transport and 4 May 2020 (Pleiades 2)without snow transport in the left sub-map. (bd) - The different colors represent the different precipitation forcing and horizontal resolutions while hatching is used Pléiades 1 observed snow height resampled to distinguish between the activation and deactivation 250 m. The absence of the SnowPappus blowing snow schemein simulations is indicated in grey. For d) the summer aerial photography base map is from IGN© (2022).

4 May 2020 (around 2 m) underestimates the observed snow height. A similar result is found at both resolutions and with or
 345 without snow transport. At most elevations, the standard deviation is similar to ANTILOPE forcing and close to the Pleiades
 Pléiades observations.

The representation of snow transport is found to only impact snow height bias for the upper elevation band (with a reduction of the mean snow height). The fact that this impact is beneficial or detrimental depends on the choice of the meteorological

- 350 forcing. Below this elevation band, the impact of snow transport on the bias is found shallow to nonexistent. Figure 5 (a) also shows that for the 3 upper elevation bands (above 2500m), SnowPappus (non-hatched violin plots) increases the simulated standard deviation on average by 63.7% between snow height distribution with and without snow transport (hatched violin plots). Figure 5 (b) (above 2500m) shows a similar standard deviation increase of 67.7% for the next year when adding snow transport in the simulations. The impact of snow transport on the simulated snow height is noticeable for all precipitation inputs.
- 355 Figure 6 (b-e) shows that the increase of standard deviation (above 2500m) introduced by the transport module is sometimes overestimated comparatively to the observation ( $\sigma$  ratio >1), depending on the date and precipitation forcing. Below 2500 m, the snow height distributions with and without snow transport, for a given precipitation input, are broadly similar.

Finally, SPS scores (Fig. 6 (c-f)) suggest that model performances are mostly modified by SnowPappus at the highest pixels of our domain and that simulation improvements or degradation depend on combinations of model setups (transport/no transport and precipitation forcings). Thus, no optimal precipitation forcing can be identified systematically.

#### 3.1.2 Galibier area

The second snow height evaluation (Fig. 7 and Fig. 8), is done in the Galibier Mountains area (blue box in Fig. 1). It uses two snow height maps derived from Pleiades Pléiades stereo-images acquired in January and March 2018, in which snow

- 365 cover extends to lower elevations than in the Grandes Rousses snow height maps. At both dates, a positive bias appears for the three lowest elevation bands, regardless of the forcing and decreasing with elevation (Fig. 8 (a-d)). For instance, the Pleiades Pléiades observation provides a mean snow height of 1.25 m at the 1900-2200 m elevation band on 23 January 2018, while the simulation's mean values are all above 1.74 m of snow. At these elevations, the distribution standard deviation (0.71 m in the Pleiades observations) standard deviation is underestimated by all simulations (from 0.71 m for the reference Pléiades observation and the pleiades observation of the pleiades observation observation observation of the pleiades observation observation observation of the pleiades observation observati
- 370 <u>observation</u>, 0.16 m for SAFRAN to and, 0.32 m for AROME), without any significant impact of the snow transport module (see std ratio Fig. 8 (b-e)).

For the two upper elevation bands, the main features are consistent with the evaluations on the Grandes Rousses area: at both dates, the AROME simulations exhibit the highest mean snow height corresponding to a positive bias compared to observations, while the ANTILOPE simulations exhibit the lowest snow height corresponding to a negative bias. Contrary to the previous

- 375 section, SAFRAN simulations also overestimate snow height in the Galibier area. For all precipitation forcing, the standard deviation of simulations without transport (between 0.11 m and 0.25 m in January at the upper level) is much smaller than in the observed spatial distribution (0.87 m). When the snow transport is activated, a more realistic spatial variability is obtained, although it is still underestimated below 2800 m elevation. This conclusion remains valid on 16 March (Fig. 7 (b)). As for the Grandes Rousses evaluation, the simulations with the AROME precipitation have the highest spatial variability in snow height
- at all elevations and for the two dates. Conversely, the simulations with SAFRAN have the smallest standard deviation across elevation.

For the SPS score (Fig. 8 (c-f)), the ANTILOPE forcing leads to the simulated snow height the closest to the observations produces very accurate scores ( $\leq 0.1$  m) between 2200 and 2800 m, and for both dates above 2200 m, contrary, in contrast to the results obtained over the Grandes Rousses area. For the lowest elevation band, AROME simulation with snow transport

385

is the closest to observations with an SPS score of 0.4 and 0.3 m in January and March. The simulations forced by SAFRAN precipitation provide the spatial distributions the furthest to <u>Pleiades Pléiades</u> observations. This is an opposite conclusion to the results obtained on the Grandes Rousses region. Therefore, the evaluations on this second region confirm the difficulty of identifying optimal precipitation forcing for all regions, elevations, and dates.

#### 3.1.3 Summary: Relative impacts of snow transport and precipitation forcing

- 390 When comparing the simulation performance between both simulation regions and all dates, common and different patterns arise. The impact of the simulation of snow transport on the bias is restricted to the upper altitude, close to mountain summits. For all simulations, higher average snow height are obtained without transport than with transport. However, the added value of snow transport can not be established on this criteria as the bias highly depends on the precipitation forcing.
- In all simulations, lower snow heights are obtained with ANTILOPE precipitation and higher snow heights with AROME precipitation, with increasing differences with elevation. However, upon examining the various bias scores (as shown in Fig. 6, 8, 10 (a-d), it is clear that no precipitation forcing leads to the simulation of the most unbiased snow heights for all areas and evaluation dates produces a snow height simulation that is systematically less biased than the others over the different evaluation dates and areas. Across all elevation ranges, the mean driver of simulation bias appears to be the precipitation dataset choice.
- 400 The addition of snow transport increases and improves the simulated standard deviation , of snow height of 0.25 and reduces the SPS value closer to the observation. The impact of transport on scores is mainly above 2500 m and with a slope increasing with elevation. For these upper elevation bands, the snow transport prevails in the choice of the precipitation dataset on this criteria. For the lower elevation bands, the main driver of simulation standard deviation appears to be the choice of the precipitation dataset. The simulations using AROME precipitation have the biggest standard deviation value of all simulations.

# 405 3.1.4 Impact of subgrid variability

The effect of unresolved subgrid variability can be analyzed by comparing SAFRAN and SAFRAN HR simulations. Simulation maps illustrating the expected subgrid variability can be found in Appendix Fig. C1 (a-b-c). At the 30 m grid resolution, the spatial variability of the topography (SAFRAN HR without transport vs. SAFRAN without transport) reduces the snow height and increases the 250 m scale spatial variance at high elevations for the Grandes Rousses evaluations. The impact is smaller for the Galibiar assessment

410 for the Galibier assessment.

Resolving snow transport at a 30 m grid resolution (later averaged to 250 m) reduces the impact of snow transport on simulations compared to 250 m simulations (Fig. 6, 8, orange dashed and continuous lines are less distant than almost always either less distant or equivalent to the green dashed and continuous lines). As snow transport tends to decrease snow height and increase spatial variance, the impact of resolving the 30 m scale snow transport has an opposite effect on the resolution of

the 30 m scale topography. Finally, SAFRAN HR and SAFRAN simulations are relatively close in terms of snow height bias,  $\sigma$  ratio, and SPS. Therefore, despite the significant impact of the simulation resolution at the process level, it is not obvious that the 250 m simulations would provide less accurate snow height spatial distributions than simulations computed at higher resolution and resampled on a 250 m grid.



Figure 5. Spatial distribution of snow height (m) for 300 m elevation bands in Pléiades observations (gray) and all simulation configurations over the Grandes Rousses mountains on 13 May 2019 (Pléiades 1)(a) and 4 May 2020 (Pléiades 2)(b). The different colors represent the different precipitation forcing and horizontal resolutions while hatching is used to distinguish between the activation and deactivation of the SnowPappus blowing snow scheme. Each violin contains a small inner black box representing the first and third quartile, and a white dot representing the median distribution value. The number of samples for each subgroup is denoted by N.



**Figure 6.** Scores comparing simulations with the Grandes Rousses <u>Pleiades Pléiades</u> observations. (a-b-c) compares simulations with the 13 May 2019 <u>Pleiades Pléiades 1</u> image ; (d-e-f) is based on the 4 May 2020 <u>Pleiades Pléiades 2</u> image. (a-d) Bias (m) ; (b-e) Standard deviation ratio (no unit) ; (c-f) SPS (m). All scores are presented as a function of 300 m elevation intervals.



**Figure 7.** Spatial distribution of Snow height (m) for 300 m elevation bands in Pleiades Pléiades observations (gray) and all simulation configurations over the Galibier on 23 January 2018 (Pleiades Pléiades 3)(a) and 16 March 2018 (Pleiades Pléiades 4)(b). Similar legend as in Fig. 5. Each violin contains a small inner black box representing the first and third quartile, and a white dot representing the median distribution value. The number of samples for each subgroup is denoted by N.



**Figure 8.** Scores comparing simulations with the Galibier <u>Pleiades Pléiades</u> observations. (a-b-c) compares simulations with the 23 January 2018 <u>Pleiades Pléiades</u> 3 image ; (d-e-f) is based on the 16 March 2018 <u>Pleiades Pléiades</u> 4 image. (a-d) Bias (m) ; (b-e) Standard deviation ratio (no unit) ; (c-f) SPS (m). All scores are presented as a function of 300 m elevation intervals.

#### 3.2 Spatial variability of Snow Melt-Out Date (SMOD)

420 The evaluations of the SMOD allow us to extend the evaluations over our entire simulation domain. It also provides complementary information on the entire snow season than the use of snow height snapshots. The associated spatial SMOD distributions for the 2017-2018 and 2018-2019 years can be found Fig. 9 (a-b). The synthetic summary of simulation performances is given by Fig. 10 (a-b-c-d-e-f).

#### 3.2.1 Differences between years

- 425 There is a notable difference in distribution variance when comparing the evaluation results for the two years are some notable differences between the 2017-2018 and 2018-2019 snow season SMOD (Fig. 9). In For the two lower elevation bands, there are on average 15 days (Fig. 9 (b), a) and 20 days (Fig. 9 b) SMOD difference between the two years for the observation. A similar SMOD difference is found on average for the simulations. However, for the 2018-2019 modeled SMOD clearly shows a wider variability at every altitude band and for each simulation configuration compared to 2017-2018 snow season (Fig. 9
- 430 (a)). Below 2200m, the SMOD spatial variability is also higher in b), we see the observations and simulations distribution have a much bigger variance than the previous season. Continuing the analysis of the upper elevation bands, we see that we observe an earlier SMOD for the 2018-2019 than in season and every the elevation bands. This observation is even more pronounced for the 3 upper elevation bands. We also note that the distribution variability is greater for this year. For the 2017-2018 in Pleiades observations but with less pronounced differences than in simulations, snow season at elevations between 2500 and
- 435 3400 m, there was an average SMOD difference of 18 days between observations and simulations. In the subsequent season, this difference increased to an average of 39 days. In contrast, during the 2018-2019 snow season, the simulations that included snow transport were, on average, closer to the observed variance than for the other seasons.

# 3.2.2 Impact of precipitation forcing

Above 2500 m, the simulation with transport and the ANTILOPE precipitation has the largest distribution variance and lower
mean SMOD value. As a result, this is the only simulation that underestimates the mean SMOD in 2018-2019 (Fig. 10 (d)) and overestimates its spatial variability (Fig. 10 (e)).

For the other precipitation forcing, in 2017-2018 the SMOD is overestimated at low elevations indicating that the snow meltout occurred too late in the season compared to observation (Fig. 9 b). Conversely, SMOD is underestimated at high elevations with a simulated melting up to 20 days before the observed one. In 2018-2019, SMOD was overestimated at all elevations. On

445 average, the spatial variance of SMOD is underestimated except for simulations with AROME the ANTILOPE forcing at the two lowest elevation bands upper elevation where the  $\sigma$  ratio is found elose to much above 1. It is interesting to note that in the lowest elevation band Fig. 9 (b), an AROME pixel never has a snow height greater than the SMOD threshold during the year (see Sect. 2.8.1) and is therefore given a 0-day SMOD value, contrary to what happens for all the other simulations and the observation.

- The SPS values of SMOD (Fig. 10 (c-f)) finally provided the opposite conclusion between both years with the simulations forced by AROME being the closest to the observed spatial distribution at all elevations in 2017-2018 but the furthest in 2018-2019 and conversely the simulations forced by ANTILOPE the furthest to observations in 2017-2018 and the closest in 2018-2019. This result confirms, using SMOD, the conclusion of Sect. 3.1 that no precipitation forcing can be systematically identified as optimal for all evaluation periods and criteria
- 455 Analysis of wind speed and direction over the three simulation years indicates that the values remained consistent. Examination of the yearly solid precipitation of the SAFRAN forcing reveals that the 2018-2019 season had 21% less snowfall compared to 2017-2018. This result is consistent with a smaller amount of solid precipitation leading to a reduced amount of snow accumulation and the observed earlier SMOD for the 2018-2019 snow season.

#### 3.2.3 Impact of snow transport

- 460 Figure 9 and Fig. 10 (b) and (e) show an increase in SMOD standard deviation when activating snow transport in simulations, consistent with the increase of snow height variance shown in Sect. 3.1. The standard deviation of simulated SMOD with transport is found to increase with altitude. This increasing spatial variance of SMOD with altitude is also observed with Sentinel 2 imagery, whereas simulation without transport keeps a standard deviation rather similar. Nevertheless, the impact of precipitation forcing on SMOD spatial variability is at least as high as the impact of snow transport. Therefore, the added value
- 465 of snow transport is still difficult to establish on that criteria, although simulations based on SAFRAN and AROME forcing would suggest that the SMOD spatial variability is improved by the snow transport module.

The addition of snow transport also decreases the SMOD for all 250 m resolution simulations, with an increasing impact on altitude. Therefore, concerning the SMOD bias, the impact of snow transport can be either beneficial in 2018-2019 or detrimental in 2017-2018 depending on the sign of the general bias of the simulation without transport.

470 Except with ANTILOPE in 2017-2018, simulating snow transport always results in improved SPS of SMOD. The detrimental impact of snow transport at high elevations in 2017-2018 for ANTILOPE forcing is probably linked with the much smaller estimates of precipitation by ANTILOPE (Appendix Fig. B1), for which the reasons are discussed in Sect. 4.2.

#### 3.2.4 Impact of subgrid variability

The addition of snow transport in the 30 m resolution SAFRAN HR simulations increases the SMOD for the two evaluation 475 years while on the contrary snow transport decreases the SMOD in the 250 m resolution SAFRAN simulation. In terms of spatial variance, the increase of spatial variance by snow transport is higher at 250 m resolution than at 30 m resolution. The direct impact of resolving a finer topography in simulations without transport (differences between orange and green dashed lines) is less significant than on snow height evaluations.

24



Figure 9. Snowpack Melt-Out Date (SMOD) depending on elevation and simulation configuration. The simulations are compared with Sentinel-2 SMOD observations of the full simulation domain, for the 2017-2018 (Sentinel 2 A) and 2018-2019 years (Sentinel 2 B). Each violin contains a small inner black box representing the first and third quartile, and a white dot representing the median distribution value. The number of samples for each subgroup is denoted by N.



Figure 10. Scores comparing simulated SMOD with Sentinel 2 imagery. (a-b-c) 2017-2018 year (Sentinel 2 A); (d-e-f) 2018-2019 year (Sentinel 2 B). (a-d) Bias (days); (b-e) Standard deviation ratio (no unit); (c-f) SPS (days). All scores are presented as a function of 300 m elevation intervals.

# 3.3 Impact of landform types

480 In this section, the spatial simulated distributions of snow heights and SMOD are compared with satellite observations while grouped by landform types instead of elevation bands, as described in Sect. 2.2 and Fig. 2.



**Figure 11.** Scores comparing simulations with the Galibier <u>Pleiades Pléiades</u> observations. (a-b-c) compares simulations with the 23 January 2018 <u>Pleiades Pléiades</u> 3 image ; (d-e-f) is based on the 16 March 2018 <u>Pleiades Pléiades</u> 4 image. (a-d) Bias (m) ; (b-e) Standard deviation ratio (no unit) ; (c-f) SPS. All scores are presented as a function of landform pixel classifications.

Figure 11 compares the January and March 2018 Galibier Pleiades Pleiades images to snow height simulations grouped by landform types. Appendix Fig. D1 presents the same comparison using 13 May 2019 and 4 May 2020 Grandes Rousses Pleiades Pleiades images. The comparison of simulation with and without transport shows that most of the snow height bias

485

difference (Fig. 11 (a) and (d) is found on pixels classified as 'Peak (summit)' or 'Ridges'. However, this impact on bias can be either beneficial or detrimental depending on the bias of the simulation without transport, highly dependent on the precipitation forcing, similar to the results described in the analyses based on elevation bands.

Simulations with snow transport show lower mean snow height in pixels classified as 'Peak (summit)' and 'Ridges' compared to simulations without transport. Conversely, simulations without snow transport exhibit lower snow height on 'Spur (Convex)'. 'Slopes', and 'Hollow (Concave)'. This behavior can be found in each 250 m simulation, regardless of the simulation forcing

490

(Fig. 11, Fig. D1) except for the SAFRAN HR simulations.

It can be noticed that the ordering between the different precipitation inputs does not depend on the landforms classes while it was quite dependent on elevation bands.

For all precipitation datasets, the standard deviation of simulations without transport is much smaller than in the observed 495 spatial distribution. The landforms where the activation of transport has the most impact on snow height variance are (by order of importance): 'Peak (summit)', 'Ridges', 'Spur (Convex)', 'Slopes' and 'Hollow (Concave)'. Compared to the observed variance, this behavior is more realistic when snow transport is activated, Fig. 11 (b) and (e) for all precipitation forcings. While the simulated distribution variance better matches the order of magnitude of the observed one for 'Peak (summit)', 'Ridges', and 'Spur (Convex)', the simulated distribution variance of 'Slopes' and 'Hollow (Concave)' is seen to be still half of

500 the observed one. As for the evaluation using elevation bands, the simulations with the AROME precipitation have the highest spatial variability for the two dates and each landform type. Conversely, the simulations with SAFRAN and no snow transport have the smallest standard deviation across landforms.

The improvement of the SPS score Fig. 11 (c) and (f) due to the representation of snow transport is major on 'Peaks (summits)' and 'Ridges' and noticeable on 'Spur (convex)' and 'Slope' classes. This improvement is obtained for all precipitation

505 forcings (even in the ANTILOPE case where the bias has deteriorated), therefore mainly driven by the improvement in snow height variance. The SAFRAN HR simulations with and without transport have the lowest (better) SPS values for the two evaluation dates and most of the landform types while it is not the least biased. A closer look at the simulation's Directly examining the SAFRAN HR cumulative distribution function (not shown here) shows reveals that the shape of the cumulative distribution is indeed more similar to the observed one (albeit biased) observation, despite being biased. This conclusion differs from the results obtained using a grid cell classification by elevation bands. 510

Figure 12 presents the two-year evaluation using Sentinel 2 SMOD and landforms. Figure 12 (a-d) show the impact of SnowPappus blowing snow simulations on simulated SMOD. As for snow height in Fig. 11, the impact on SMOD is mainly limited most significant in areas classified as 'Peak (summit)' or 'Ridge'. SMOD difference when using SnowPappus is found to go up to 20 days for 'Peak (summit)'. When comparing the two evaluation years, similar behavior is found, except the 2018-

515 2019 year appears uniformly more biased. Areas classified as 'Valley' and 'Pit' appear more biased for every precipitation forcing.



Figure 12. Scores comparing simulated SMOD with Sentinel 2 imagery. (a-b-c) 2017-2018 year (Sentinel 2 A) ; (d-e-f) 2018-2019 year (Sentinel 2 B). (a-d) Bias (days) ; (b-e) Standard deviation ratio (no unit) ; (c-f) SPS (days). All scores are presented as a function of landform pixel classification.

In Fig. 12 (b-e), the impact of snow transport is again mainly present on 'Peak (summit)' and 'Ridges' landforms, but with a high variability between both years and a lower impact when AROME precipitation forcing is used (with already a higher spatial variability even without transport). In 2017-2018, the impact of snow transport is always beneficial on the SMOD  $\sigma$ ratio, as obtained for snow heights, but for the second season, it depends on the precipitation forcing. The simulated standard deviation of AROME simulations appears close to the observed one for all landforms but pixels classified as 'Pit'. This better behavior of AROME was not illustrated in the evaluation done using the elevation bands. The other precipitation forcing appears underdispersed with a standard deviation radio around ratio of 0.5-0.75.

525

Looking at the SPS scores Fig. 12 (c-f), the impact of snow transport is less clear than for snow heights, similar to the results obtained by elevation bands. SPS values are smaller using landform classification than elevation bands. All simulations appear to have difficulties in representing SMOD distribution in pixels classified as 'Valley' and 'Pit' conversely to results of Fig. 11. The SPS scores are also more similar between the two evaluation years, contrary to the SPS score using elevation bands grouping. Figure 10 (c) shows a distance increase between all simulations and observations distribution between 2500-3100 m. This effect is not found in Fig. 12 (c). On the contrary to Fig. 11, the SAFRAN HR does not have the lowest simulations have the overall highest SPS values, and AROME simulations'  $\sigma$  ratios are found to be more realistic.

530

The impact of the group axis on results can be quantified by comparing identical data grouped differently. Fig. 13 presents the mean SPS values of the 8 simulations when grouped by elevation bands and landform types. We can see that SPS values are lower when grouped by landform types. We can see that grouping by elevation gives more homogeneous SPS scores for the different observation dates than grouping by landform where variation between dates is greater.

![](_page_29_Figure_5.jpeg)

Figure 13. Plots of average SPS values for the different simulations when grouped with elevation bands or landform types. This graph illustrates the impact of the grouping axis (elevation or landform types) on snow height and snow melt-out date.

# 535 3.4 Impact of the SnowPappus model

Correlation between observations and the different simulations can be seen in Fig. 14. This figure illustrates scatter plots of snow height for the four different Pléiades observations to the 8 different simulations (SAFRAN, SAFRAN HR, AROME, ANTILOPE, with and without activation of SnowPappus). For each simulation, we see more dispersed scatter plots for simulations with snow transport (more points at high and low snow heights). It can be noted that the mean difference in

- 540 slope caused by the addition of snow transport is small (<0.02). Overall, AROME simulations have the steeper slopes. To quantify how correlated the simulation results are to the observations, we use the Pearson correlation coefficient. In Table 3 and Table 4, we see Pearson correlation coefficients between Pléiades observations and simulations for each of the 8 simulations. Correlation is computed two times, for the entire area (all elevations) and restricted to elevation higher than 2700 m.
- 545 Looking at the Pearson correlation for the entire Pléiades domain, the addition of snow transport in simulations decreases the correlation coefficient in 15 of 16 experiments. Restricting the correlation to pixels to elevation above 2700 m (pixels with high snow transport probability) the addition of snow transport in simulations increases the Pearson correlation for each experiment. This result is found statistically significant (with significance set at p-value <0.05) in 15 out of 16 experiments.</p>

We note that for the entire domain, the variability of the correlation score for a single observation is greater according to the source of precipitation than for the addition of the snow transport process in the simulations.

- To better disentangle the impact of precipitation-forcing variability and the addition of snow transport, this result can be supplemented with Fig. 15, representing the mean SPS value of the 6 different 250 m simulations forced with SAFRAN, ANTILOPE, AROME, for elevations above 2700 m. In this figure, the addition of snow transport to the simulations clearly shows improvements in the mean SPS score on snow height and SMOD above 2700 m. SnowPappus lowers SPS values of
- 555 respectively 0.06 m and 0.85 days above 2700 m.

![](_page_31_Figure_0.jpeg)

**Figure 14.** Observed vs simulated snow height scatter plots for the four different Pléiades observations and the different precipitation forcings. For each precipitation forcing, the simulation with snow transport is in blue and without transport in red. Linear regression line synthesizes distribution changes when adding snow transport. The respective Pearson correlation coefficient can be found in Tables 3 and 4.

Forcing	Snow transport	Pléiades 1 2019-05-13	Pléiades 1 2019-05-13 Z>2700 m	Pléiades 2 2020-05-20	Pléiades 2 2020-05-20 Z>2700 m
Safran	Yes	0.744	0.173	0.766	0.184
Safran	No	0.751	0.027	0.795	<u>0.062</u>
Safran HR	Yes	0.739	0.190	0.803	0.221
Safran HR	No	0.741	0.097	0.820	0.173
Arome	Yes	0.826	0.169	0.820	0.211
Arome	No	0.835	0.064	0.850	0.127
Antilope	Yes	0.592	0.174	0.672	0.198
Antilope	No	0.591	0.045	0.693	0.105

Table 3. Pearson correlation coefficients between observed and simulated snow height (Grandes Rousses). For each observation/simulation pair, a second correlation is computed only with pixel elevations (Z) above 2700 m. Underlined values represent p-value >0.05. Bold values correspond to the best correlation value between simulation with and without the SnowPappus model.

Forcing	Snow transport	Pléiades 3 2018-01-23	Pléiades 3 2018-01-23 Z>2700m	Pléiades 4 2018-03-16	Pléiades 4 2018-03-16 Z>2700
Safran	Yes	0.492	0.192	0.586	0.346
Safran	No	0.599	<u>0.068</u>	0.675	0.142
Safran HR	Yes	0.535	0.110	0.626	0.353
Safran HR	No	0.598	0.086	0.664	0.157
Arome	Yes	0.549	0.189	0.716	0.414
Arome	No	0.600	0.025	0.744	0.355
Antilope	Yes	0.437	0.181	0.566	0.422
Antilope	No	0.509	0.020	0.606	0.266

**Table 4.** Pearson correlation coefficients between observed and simulated snow height (Galibier). For each observation/simulation pair, a second correlation is computed only with pixel elevations (Z) above 2700m. <u>Underlined values</u> represent p-value >0.05. **Bold** values correspond to the best correlation value between simulation with and without the SnowPappus model.

![](_page_33_Figure_0.jpeg)

Figure 15. Plots of average SPS values for the three different 250 m forcings (SAFRAN, ANTILOPE, AROME) for pixels with elevations above 2700 m. This graph illustrates the impact of the SnowPappus transport model on snow height and snow melt-out date.

# 4 Discussion

#### 4.1 Sources The Added value of snow spatial variability the SnowPappus model

In this study, we compare snow height maps simulated with and without transport using various sources for the precipitation

- 560 forcing. The combined impact of precipitation forcing variability and wind-blown redistribution <u>using the SnowPappus model</u> is visible, on snow height, in Fig. 5 and Fig. 7 and SMOD in Fig. 9. We have found that the main driver of the simulated snow height distribution mean value is the precipitation forcing. The precipitation forcing spatial variability across low to medium altitudesis also found to be the main driver of simulated snow height distribution standard deviation. On the contrary, for the upper altitudes and especially around For the higher altitudes, particularly around the summit pixels, the standard deviation of
- 565 the simulated snow height distribution and the and snow melt-out date distribution is mainly primarily determined by blowing snow transport.

In our simulations, The blowing snow transport impact is mostly restricted to 'Peak (summit)' and 'Ridges' landforms (Fig. 11, Fig. D1, Fig. 12) and more generally to the upper elevation bands (above 28002700-2800 m). In addition, the snow height variance is seen to increase closer to the observation using the SnowPappus' simulated snow transport, regardless of the pre-

- 570 cipitation bias. Looking at the Pléiades bias values in Fig. 6 and Fig. 8 (a) and (d), blowing snow reduces the absolute mean bias value of snow height in all simulations. This is probably a combined effect of increased snow density and mass loss from increased sublimation. This result is consistent with Vionnet et al. (2021). In Sect. 3.4, results show the SnowPappus model consistently improves the correlation between Pléiades observations and simulations above 2700 m. This improvement was found to be significant in 15 out of 16 experiments. In our simulation domain, the elevation of 2700 m and above correspond
- 575 to areas where our snow transport model has a significant impact on the simulated snow height. For the Pléiades domain as a whole, the correlation coefficient is modestly reduced with the addition of snow transport. These results are a generalization of the correlation analysis done in Baron et al. (2024) and are consistent with the previous conclusions. However, the added value of SnowPappus can not be established only on this criteria as we find the simulated bias and standard deviation dependent on the precipitation forcing. To better disentangle the effects of precipitation forcing variability and the addition of snow transport, the
- 580 previous conclusions can be supplemented with Fig. 15, which shows the mean SPS value of the 6 different 250 m simulations forced with SAFRAN, ANTILOPE, AROME for elevations above 2700 m. Based on these results, it can be concluded that the addition of the SnowPappus model to our 250 m simulation improves the simulated snow distribution closer to the observed distribution for both snow height and SMOD.

In the literature, Bernhardt et al. (2012) studied the influence of lateral snow redistribution using SnowTran-3D (Liston et al.,

- 585 2007) and found an enhancement in the simulated snow spatial pattern using lateral snow redistribution. Using the Canadian Hydrological Model (Marsh et al., 2020), Vionnet et al. (2021) find snowpack simulation without lateral snow redistribution unable to capture the spatial variability of snow cover in alpine terrain. The On the opposite, the addition of lateral snow redistribution is found to provide the best a better estimate of snow height across elevation and increase the simulated snow height distribution variance. Very recently, Quéno et al. (2023) evaluated FSM2trans snow redistribution model (Quéno et al., 2023)
- 590 and find the addition of snow redistribution necessary at hectometric or finer resolution to better represent snowpack hetero-

genity. Snow heights simulated using lateral snow redistribution are found to have significantly better variability. Additionally, (Quéno et al., 2023) conducted a model sensibility analysis to horizontal resolution and <u>found\_established</u> that although the most realistic simulation pattern <u>was\_is</u> found at 25 m resolution, the addition of lateral snow redistribution at coarser resolution is <u>found</u> also having a positive impact on snow distribution and variance. Hence, we estimate that the results obtained in this

study on the impact of snow transport are the SnowPappus snow transport model compatible with the results found in the

595

literature.

In this study, we tested the sensitivity of the spatial snow height variability found in our simulation to the

# 4.2 Sensitivity to precipitation forcings

In our simulations, the mean and median snow height and SMOD values are primarily driven by the spatial variability of precipitation , snow transport, and spatial resolution, but numerous other processes shape snow spatial variability in the field (as reviewed in Mott et al. (2018)). Those processes occur at different scales, going from the regional scale to the slope and lower scale. It is worth noticing that neither preferential deposition nor the seeder-feeder mechanism is represented in our simulations . Looking at the Pleiades bias values in forcing, as shown in Fig. 5, Fig. 7, and Fig. 9. Precipitation forcing is also found to be the main driver of simulated snow height distribution standard deviation across low to medium elevations. As the

- 605 Crocus snow model is known to produce realistic snow heights and snow water equivalent when forced by well-controlled meteorological forcing (Lafaysse et al., 2017; Menard et al., 2021), the large biases of snow height and SMOD observed in the snow simulations presented most probably reflect intrinsic biases of the different precipitation dataset. In Fig. 6 and Fig. 8 (a) and (d), blowing snow reduces the absolute mean bias value of snow height in all simulations. This is probably a combined effect of increased snow density and mass loss from increased sublimation. This result is consistent with Vionnet et al. (2021)
- In for a simulation with a precipitation dataset leading to an already negatively biased snow height (like ANTILOPE), the addition of snow transport further degrades the overall snow height simulation bias. Similar behavior is found on the SMOD in Fig. 10 (a) and (d) with an exception for the 30m simulation where it increases the absolute SMOD bias. Thus the beneficial or detrimental impact of blowing snow transport on mean snow height or SMOD bias is mainly dependent on the precipitation bias. This result is consistent with Raleigh et al. (2015); Schlögl et al. (2016); Günther et al. (2019), also finding the prevalence of meteorological forcing errors in snow simulations.

#### 4.3 On the precipitation forcings

As the Crocus snow model is known to produce realistic snow heights and snow water equivalent when forced by well-controlled meteorological forcing (Lafaysse et al., 2017; Menard et al., 2021), the large biases of snow height and SMOD obtained in the simulations presented in this paper probably largely reflect intrinsic biases of the different precipitation dataset. More specifi-

620 cally, when looking at the simulation forcing with the ANTILOPE precipitation source, we found that across all elevation bands, the ANTILOPE forcings have the lowest snow height and SMOD values. Furthermore, a A negative bias trend is seen with elevation (Fig. B1). The ANTILOPE precipitation product is primarily based on radar reflectivity measurements, technique known to be more challenging in mountainous terrains (Yu et al., 2018; Faure et al., 2019; Foresti et al., 2018; Germann et al., 2022). Meteorological radars are based on the backscatter measurement of several conic-shaped microwave electromagnetic

- 625 impulses produced by the radar itself at several elevation angles (conic-shaped radar beams). When a mountain intercepts a radar beam, the reflected signal (called ground clutter) is unusable and rejected by the ANTILOPE algorithm. If the mountain intercepts less than 70% of the beam section, the signal reflected further away is simply corrected to account for the loss of information but is still used for precipitation rate estimation. Precipitation estimation over pixels affected by ground clutters thus comes from beams that can be several thousands of meters higher than the mountain, often above the base of precipitation
- 630 clouds. This leads to an underestimation of the precipitation rate over mountain ridges (Appendix Fig. B1), due to ground clutter. One possible compromise would consist of using the observed spatial variability from ANTILOPE while correcting the elevation-dependent precipitation bias.

In contrast to ANTILOPE, our simulations with the AROME precipitation source lead, on average, to the highest snow height across the elevation bands. It is particularly visible in Fig. 4 (b), where the snow height patterns squared patterns can

- 635 be seen and come from the 1.3km AROME pixels forming distinctive precipitation patterns (see Appendix Fig. B1 (d). This behavior of excessive solid precipitation at high elevations is known and coherent with other snow cover studies using AROME as the precipitation source (Quéno et al., 2016; Vionnet et al., 2019; Monteiro et al., 2022). Multifactorial causes were discussed by Monteiro et al. (2022); Gouttevin et al. (2023). The variable biases between the western (Grandes Rousses) and eastern (Galibier) evaluation areas suggest that orographic enhancement of precipitation might be especially overestimated by
- 640 AROME on the areas the most exposed to the prevailing westerly flows.

In contrastFinally, the two SAFRAN simulations that are the only ones to assimilate precipitation gauge observations show a good agreement in terms of SMOD and snow height biases but are spatially under-dispersed by design. The SAFRAN meteorological reanalysis (Vernay et al., 2022) uses one vertical profile per climatologically homogeneous area, interpolated to all simulation points with altitude (see Sect. 2.5.1). Within one of these areas, the spatial variability is thus limited to a vertical gradient as described in Vionnet et al. (2016). This also explains why opposite snow height biases are obtained with this forcing

between western (Grandes Rousses) and eastern (Galibier) areas.

#### 4.3 Limitations of the studySensitivity to horizontal resolution

This study highlights the difficulties in disentangling precipitation and blowing snow errors in snow simulation. Our results show that precipitation forcing is the largest contributor to snow height error across elevation. The utmost care must be taken to estimate the precipitation field as accurately as possible. Our numerical simulations naturally have other types of errors and

650

limitations.

645

#### 4.3.1 Sensitivity to horizontal resolution

Numerical simulations rely on numerical discretization and parameterization. The choice of the spatial resolution a spatial resolution inevitably introduces errors. This choice depends on a trade-off between scientific arguments, computing resources, and external factors. As described in Baron et al. (2024), the 250 m resolution choice of the SnowPappus scheme was motivated

and external factors. As described in Baron et al. (2024), the 250 m resolution choice of the SnowPappus scheme was motivated by the desire for applicability in large-extent systems (50000 to 100000 km<sup>2</sup>, multi-decadal simulations). Numerical resolution choices impact the numerical resolution choice impacts simulated slope inclinations and aspects and can lead to inaccuracy in the radiative balance (Baba et al., 2019). The snow radiative balance is particularly heterogeneous in mountainous areas due to complex topography leading and can lead to significant changes in snow properties and melting rates. Comparing

660

simulations computed at 250 and 30 m resolution provides an analysis of the unresolved subgrid variability of our simulations. Direct. A direct comparison between Appendix Fig. C1 (a-b-c) illustrates reveals the expected unresolved subgrid variability. Snow height patterns on the 250 m simulations (Appendix Fig. C1(a)) exhibit steeper variations than the 250 m re-gridded maps obtained from 30 m simulations (SAFRAN HR, Appendix Fig. C1 (b)) where the snow patterns are found smoother. To understand the origin of this subgrid variability. Appendix Fig. C1 (c) illustrates the raw 30 m simulation where snow patterns 665 are more sophisticated. As expected (Baba et al., 2019), resolving the 30 m topography in simulation without snow transport increases the spatial variance at high elevations and reduces the 250 m averaged snow height (Sect. 3.1.4). On the other hand, increasing Increasing the resolution in simulation with snow transport increases the SMOD (Sect. 3.2.4) but reduces the expected snow height variability gain compared to 250 m simulations. The complex behavior of resolution can be summarized by comparing the mean SPS value across all elevations of the SAFRAN and SAFRAN HR simulations, as in Appendix Fig. E1.

670 This figure shows a very small mean SPS change with resolution. The mean difference is found to be lower than 0.01 m for the Pléiades observations and lower than a day (0.98) for SMOD. Until now, the resolution-dependence of snow transport models was only documented at higher resolutions (Grünewald et al., 2013; Marsh et al., 2020; Mott et al., 2010; Schneiderbauer and Prokop, 2011) but results recently submitted suggest that the resolution-sensitivity of SnowPappus in this range of spatial scales is similar to what is obtained from another snow transport model at similar resolutions (Quéno et al., 2023).

675 Ouéno et al. (2016); Bellaire et al. (2014); Schirmer and Jamieson (2015) Quéno et al. (2016); Bellaire et al. (2014) and Schirmer and Jamieson (2015) also emphasize the relationship between the available resolution of the atmospheric forcing and the resulting snow simulation leading to the recommendation of forcing spatial resolution of 1 km or less, although evaluations of these data are very challenging at this scale. A much lower resolution is needed to capture the relationship between topography wind field and snow accumulation (Dadic et al., 2010) or for a complete 680 representation of wind-driven processes (Mott et al., 2008, 2014; Bernhardt et al., 2010; Vionnet et al., 2017).

Additional errors in snow-wind interaction come from process modeling. In our simulation, only the blowing snow is represented, the preferential deposition and snowfall enhancement are not represented. Blowing snow transport parameterization also suffers from uncertainties. The parameterizations used in our model were developed for flat areas and the question remains on using them in complex turbulent mountain topography. As discussed by Aksamit and Pomeroy (2018), turbulence gust wind,

685 and eddies are some of the main small-scale contributors to blowing snow. In any case, the DEVINE downscaled wind does not yet take into account wind turbulence and recirculation zone. Furthermore, further studies are needed to quantify the relative impact of recirculation zones over lee deceleration at the 250 m spatial resolution used in this work.

In our simulations, not all cover types and processes are represented. Forests and glaciers are masked out. When comparing simulations to observations, bias on the border of masked areas likely occurs due to unrealistic snow interactions simulated at 690 the borders of such masked areas. Soil properties are also an important parameter influencing snowpack simulations but they are often overlooked. Our simulations use the 1km resolution ECOCLIMAP landcover database (Champeaux et al., 2005) and

38

the 1km Harmonized World Soil Database (HWSD) v1.10 (Nachtergaele et al., 2009) as the reference for soil properties. The spatial variability of soil properties in mountains is significant and the 1km resolution of the HWSD database might not capture enough of the spatial variability, leading to spatial errors.

#### 695 4.3.1 Uncertainties and limitations of observations

In this work, simulations are evaluated against spatialized snow observations, giving a larger perspective than point snow observations. Indeed, although point scale snow observations are commonly used to evaluate spatialized simulations (Horton and Haegeli, 2, their spatial distribution and representativeness is quite low in mountains (Grünewald and Lehning, 2015; Pomeroy et al., 2009; Pepin et a , while satellite snow observations like Pleiades snow height and Sentinel 2 SMOD have a much better spatial coverage and

- 700 representativeness. Nonetheless, some errors and limitations also come from the observation side. The Pleiades observations are on-demand snapshots of snow height over a predefined extent and at a specific date. By design, this limits the usage of Pleiades images and the extent to which our conclusions stay valid. Pleiades observations of snow height can be affected by shadows, steep slopes, and the reference image used in the processing method. These limitations restrict the domain of observations and make it difficult to retrieve accurate snow height on glaciers. The Pleiades horizontal resolution of 2 m causes
- 705 challenges in comparison with coarser resolution simulation. Unsimulated sub-grid processes are captured by the observation, which complexifies the analysis. For the Sentinel 2 observations, the major limitation comes from the snow detection algorithm. The presence or absence of snow is not directly comparable to our simulation output. In Sect. 2.8.1, the presence or absence of snow is expressed as a function based on a given simulated snow height threshold. This raises the question of the sensitivity to this threshold (Gascoin et al., 2019; Hofmeister et al., 2022).

#### 710 4.3.1 Impact of the grid cells classification in spatial evaluations

#### 4.4 Impact of the grid cells classification in spatial evaluations

Classically in the snow modeling community, the quantitative spatial snow evaluations are performed by grouping simulation points (Grünewald et al., 2014; Vionnet et al., 2022; Monteiro et al., 2022; Tong et al., 2009; Deschamps-Berger et al., 2022; Vionnet et al., 2017) because (1) it allows to summarize summarizing large simulations dataset with more concise diagnostics
and (2) these evaluations are less demanding for models on large areas than pixel-to-pixel evaluations. In Sect. 3.3, results are presented using landform classification instead of elevation bands. The results are partly correlated because summits and ridges mainly cover the highest elevations of the domain (Fig. 3). Thus, we consistently obtain that snow transport mainly affects snow height and SMOD on the upper elevation bands and on for geomorphons 'Peaks (summit)' and 'Ridge'. However, some conclusions significantly differ between both space classifications, especially for the *σ* ratio score . Using landform

720 grouping, the intra-class variance includes a relatively large spatial variability due to elevation while using elevation grouping, the intra-class variance includes a relatively large spatial variability due to topographic features. This might suggest that these simulations better capture the altitudinal variability of precipitation than their topographic-dependent variability. This result clearly illustrates the sensitivity of some scores to the choice of the grid cell classification. For the SPS score, a and the SPS. A notable difference is found for the in the SPS snow height comparison. SPS scores are

- 725 found more homogeneous and smaller to show lower variability for the different precipitation forcings and overall smaller SPS values using landforms grouping than elevation bands (Fig. 11, Appendix Fig. D1 (c-f)). This result can be supplemented with Fig. 13 showing the mean SPS value being consistently better using landform groups. This suggests simulated and observed distributions are closer when grouped using landforms than elevation bands. Thus for Grandes Rousses and Galibier areas, we can again conclude that the observed snow height variability Indeed, for the landform grouping, each different landform
- 730 distribution includes a relatively large spatial variability (intra-class variance) mostly due to elevation is better captured than the variability due to landform.

, slope, aspect, and precipitation variability (Clark et al., 2011; Freudiger et al., 2017). On the other hand, using elevation grouping the intra-class variance is mostly caused by other processes than elevation. This might suggest that our simulations better capture the altitudinal variability of snow than the other processes and the topographic-dependent variability. This result clearly illustrates the sensitivity to the choice of grid cell classification groups of the scores used.

- The landform classification also provides complementary insights into the behavior of the SnowPappus model. In Fig. 11 and <u>Appendix</u> Fig. D1 (a-d), SnowPappus reduces the mean snow height on pixels classified as 'Peak (summit)' and 'Ridges' and increases the snow height on 'Spur (convex)', 'Slope' and 'Hollow (concave)' comparatively to simulations without snow transport. This behavior is the result of the wind spatial patterns obtained from Le Toumelin et al. (2023). It can be understood
- as snow on 'Peak (summit)' and 'Ridges' (ablation areas) being transported to pixels classified as 'Spur (convex)', 'Slope' or 'Hollow (concave)' (deposition areas) due to respective wind acceleration and deceleration on these pixels. This behaviour is similar between all 250 m resolution simulations while different patterns are obtained from the 30m SAFRAN HR simulation. Indeed, the finer spatial scales resolved at 30-m resolution produce more complex spatial patterns of wind fields and snow transport. When averaged at 250-m resolution, these simulations produce a more subtle influence of snow transport than
- simulations at the coarser 250-m resolution, with a less direct influence of the 250-m topography on the simulated snow patterns.

# 4.5 Limitations of simulations

735

In this study, we tested the sensitivity of simulated spatial snow height and SMOD variability to the variability of precipitation, snow transport, and spatial resolution, but numerous other processes shape snow spatial variability in the field, as reviewed

- 750 in Mott et al. (2018). Those processes occur at different scales, going from the regional scale to the slope and lower scale. It is worth noticing that neither preferential deposition nor the seeder-feeder mechanism is represented in our simulations. Additional errors in snow-wind interaction come from process modeling. In our simulation, only the blowing snow redistribution process is represented, the preferential deposition and snowfall enhancement are not. Blowing snow transport parameterization also suffers from uncertainties. The parameterizations used in our model were developed for flat areas and the question remains
- 755 on using them in complex turbulent mountain topography. As discussed by Aksamit and Pomeroy (2018), turbulence gust wind, and eddies are some of the main small-scale contributors to blowing snow. In any case, the DEVINE downscaled wind does not yet take into account wind turbulence and recirculation zone. Further studies are needed to quantify the relative impact of

recirculation zones over lee deceleration at the 250 m spatial resolution used in this work.

Moreover, in our simulations not all cover types and processes are represented. Forests and glaciers are masked out for

- 760 comparisons in addition to unusable observed pixels (due to topographic shadows or other observation limitations). The necessity of using these masks inevitably has an impact on the results by reducing the number of pixels in use for the evaluation. For masked forest pixels, we know that snow transport is strongly inhibited in forested areas leading to the creation of snow accumulation at the forest edges (Bernhardt et al., 2012). It is not possible to account for this phenomenon solely through post-processing masks. In our simulation domain, the main forested areas are located at low elevations where snow transport is
- 765 minimal or non-existent. On the other hand, glaciers are generally located in areas prone to blowing snow. Removing glaciers from the analysis significantly decreases the total number of pixels subject to snow transport at higher altitudes, but it does not introduce bias to pixels at the edges of the mask. Although the simulated and observed snow heights are not directly usable, it is important to note that snow transport is still simulated on masked glacier pixels and thus can contribute to neighbor pixels' snow balance.

# 770 4.6 Uncertainties and limitations of observations

In this work, simulations are evaluated against spatialized snow observations, giving a larger perspective than point snow observations. Indeed, although point scale snow observations are commonly used to evaluate spatialized simulations (Horton and Haegeli, 2, their spatial distribution and representativeness is quite low in mountains (Grünewald and Lehning, 2015; Pomeroy et al., 2009; Pepin et a , while satellite snow observations like Pléiades snow height and Sentinel 2 SMOD have a much better spatial coverage and

- 775 representativeness. Nonetheless, some errors and limitations also come from the observation side. The Pléiades observations are on-demand snapshots of snow height over a predefined extent and at a specific date. By design, this limits the usage of Pléiades images and the extent to which our conclusions stay valid. Pléiades observations of snow height can be affected by shadows, steep slopes, and the reference image used in the processing method. These limitations restrict the domain of observations and make it difficult to retrieve accurate snow height on glaciers. The Pléiades horizontal resolution of 2 m causes
- 780 challenges in comparison with coarser resolution simulation. Unsimulated sub-grid processes are captured by the observation, which complexifies the analysis (Fig. 4). For the Sentinel 2 observations, the major limitation comes from the snow detection algorithm. The presence or absence of snow is not directly comparable to our simulation output. In Sect. 2.8.1, the presence or absence of snow is expressed as a function based on a given simulated snow height threshold. This raises the question of the sensitivity to this threshold (Gascoin et al., 2019; Hofmeister et al., 2022).

# 785 5 Conclusions

We conducted a spatialized evaluation of the blowing snow model SnowPappus (Baron et al., 2024) joint with a sensitivity analysis to alpine precipitation variability on a 902 km<sup>2</sup> area representative of the French Alps and the Pyrenees in terms of landforms. We evaluated the simulated snow height and Snow Melt Out Date (SMOD) using <u>Pleiades Pléiades</u> and Sentinel 2 satellite products. The <u>Pleiades Pléiades</u> comparisons were conducted in two different areas, for <u>different times in the same</u>

- 790 snow season and in between two 4 different dates over 3 different snow seasons. Eight different snow simulations have been run over three snow seasons with three different precipitation forcings and two simulation resolutions (Table 1). We performed simulation analysis with the aim of disentangling simulation error contributions between from the SnowPappus blowing snow model, precipitation forcings, and the precipitation forcings variability, and from the unresolved subgrid variability. The main conclusions of this study are :
- Simulations without snow transport are found unable to capture the spatial variability of snow cover in alpine terrain. The addition of snow transport to simulations provides a better estimate SnowPappus' snow transport in the simulations results in a more physically realistic assessment of snow height across elevations and increases the and SMOD, regardless of precipitation forcing, and leads to simulated snow height and SMOD closer to the observations. Furthermore, the use of the SnowPappus model increases the variance of simulated snow height variance regardless of the regardless of precipitation forcing bias.
  - Precipitation errors overlook the bias and the standard deviation of are identified as the main source of bias and standard deviation in snow height and SMOD at low to medium altitudes. This result underlines that the greatest care must be taken to obtain the most accurate precipitation fields for snow simulation. It also suggests that assessments of spatial snow simulations that do not account for precipitation uncertainty are unlikely to provide informative insights regarding the accuracy of particular simulated snow processes.
  - Blowing snow transport impact on snow height, SMOD bias, and standard deviation prevail for high altitudes or pixels classified by the landform classifier as 'Peak (summit)' and 'Ridges'.
  - Simulated snow heights are found relatively close between the two computation resolutions (The changes in simulation snow height and SMOD mean SPS when increasing computation resolutions from 250 m, to 30 m) in terms of snow height bias,  $\sigma$  ratio, and SPS values. Changes are minimal. For Pléiades observations, the change is less than 0.01 m, and for SMOD, it is less than 1 day. These changes in spatial variance and bias when increasing resolution from 250 m to 30 m are are also lower than the simulated spatial variability due to precipitation inputs and the impact of blowing snow in the proper affected areas.
  - The addition of the SnowPappus blowing snow model to a 250 m Crocus snowpack simulation always consistently increases and improves the simulated snow spatial variability at high elevations in accordance with observations. However, improvement of snow height and SMOD biases are only obtained using precipitation forcing that does not suffer from strong negative biases.

The findings of this study show promising results of using the SnowPappus blowing snow model for large-scale modeling of alpine snowpack. Future work will focus on adding to the simulation workflow assimilation methods to improve precipitation estimates. Our results suggest that snow transport impact on snow height at 250 m resolution is significant at high elevations and therefore needs to be accounted for. However, unresolved processes at this spatial scale, parameterization uncertainty,

810

805

815

and evaluation challenges are responsible for large uncertainties that will need to be appropriately quantified in future works, especially with the goal of designing a snow ensemble assimilation frameworks (e.g. Cluzet et al., 2021; Deschamps-Berger et al., 2022).

- 830 //opensource.umr-cnrm.fr/projects/snowtools\_git/wiki/Install\_SURFEX. The version used in this work is tagged as SnowPappus-v1.0. A user manual, describing the SURFEX namelist options related to SnowPappus is available at https://doi.org/10.5281/zenodo.7681340. More general information about SURFEX use can be found at https://github.com/UMR-CNRM/snowtools.

The code and model weights used for DEVINE are available freely at https://github.com/louisletoumelin/neural\_network\_and\_devine/. The WhiteboxTools Open Core software used to compute the landforms classification is freely available at https://www.whiteboxgeo.com/.

- The Earth System Modeling Framework (ESMF) software is freely available at https://earthsystemmodeling.org/.
  The GDAL software is also freely available https://doi.org/10.5281/zenodo.8340595.
  The DEM used in this study originates from IGN® website. Tiles can be downloaded freely at https://geoservices.ign.fr/rgealti.
  Forest mask originates from IGN® and can be downloaded freely at https://geoservices.ign.fr/bdforet.
  Waterways, lakes, and cities' spatial extent originate from IGN® and can be downloaded freely at https://geoservices.ign.fr/bdtopo.
  840 The glacier masks can be downloaded freely from the Randolf glacier inventory at http://glims.colorado.edu/glacierdata/.
- The SAFRAN reanalysis dataset is freely available on the AERIS data center on the following link https://doi.org/10.25326/37#v2020.2. The AROME and ANTILOPE precipitation dataset can be requested online https://donneespubliques.meteofrance.fr/. The Sentinel 2 Level 3 dataset Theia level 3B snow products can be accessed at https://www.theia-land.fr/en/product/snow/. The raw Pleiades Pléiades images are not publicly available. The 250 m averaged Pleiades Pléiades images can be found at https://doi.

(The data accessibility and URL links status is confirmed on 24 Oct 2023 March 22th 2024)

*Author contributions.* A.H. ran the simulations, collected and processed the different data, wrote the paper, and made the spatial evaluations and the scientific choices. A.H. and M.B. developed jointly the SnowPappus code with equal contribution. M.B. helped write the Crocus and SnowPappus sections. M.L. supervised the work, helped a lot with the scientific choices, and provided technical support on Crocus and

850 SURFEX. M.L. and M.D. were extensively involved in the proofreading process and the scientific choices. L.LT provided the DEVINE downscaled wind forcing and helped in writing the wind DEVINE section. M.V. provided the ANTILOPE precipitation forcing and helped the writing of the ANTILOPE section. C.D.B. and S.G. provided the Pléiades images and the stereo-imagery treatment. C.D.B., S.G., and V.V. helped with the evaluation methodology, participated in theoretical discussions, and proofread the article.

<sup>845</sup> org/10.5281/zenodo.10037253.

*Competing interests.* One of the co-authors is a member of the editorial board of The Cryosphere. Other authors have no competing interests to declare.

Acknowledgements. This investigation was done using Meteo France's CNRM's, and CEN's computational and human resources. CNRM/-CEN is part of LabEx OSUG@2020. The main source of funding is the Region Auvergne Rhône-Alpes (France) SENSASS project and CNES TOSCA. We gratefully acknowledge Sabine Radanovics, Diego Monteiro, and Matthieu Fructus for their expertise.

![](_page_44_Figure_1.jpeg)

**Figure A1.** Landforms classification frequency (at 250 m) for our simulation domain and the entire French Alps and Pyrenees (as an area defined in Vernay et al. (2022)). We found a good agreement between our simulation pixel classification and the frequency of the French Alps and Pyrenees. We note that using 250 m resolution, no pixel corresponding to shoulder, footslope, and flat are classified as such.

# 860 Appendix B: Additional map of mean daily precipitation variability

![](_page_45_Figure_1.jpeg)

**Figure B1.** Map of the mean daily precipitation over 2017-2020 in (mm/day) for the 4 different forcings. This illustrates the spatial variability of each precipitation dataset (a. SAFRAN, b. SAFRAN HR, c. ANTILOPE, d. AROME). In each map, the mean solid fraction of precipitation is displayed as a contour plot (i.e. 1=only solid precipitation). The summer aerial photography base map is from IGN© (2022).

# Appendix C: Additional map of simulated snow height for Pleiades Pléiades 1, 13 May 2019

a) Safran

![](_page_46_Figure_2.jpeg)

c) Safran 30m

![](_page_46_Figure_4.jpeg)

b) Safran HR

![](_page_46_Figure_6.jpeg)

	] Simulation area ] Forest, glaciers, and lake mask
Sn	ow height (m)
	>0
	0.91
	1.82
	2.73
	3.64
	4.55
	5.46
	6.3
	7
F	-

**Figure C1.** Map of simulated snow height on the 13 May 2019 10:00. (a) Safran 250m simulation with snow transport and without snow transport in the left submap. (b) Safran HR simulation (30 m computed resampled to 250m) with snow transport and without snow transport in the left submap. (c) Raw Safran 30m simulation with snow transport and without snow transport in the left submap. The absence of snow is indicated in grey.

# Appendix D: Additional landform analysis

![](_page_47_Figure_1.jpeg)

**Figure D1.** Scores comparing simulations with the Grandes Rousses <u>Pleiades Pléiades</u> observations. (a-b-c) compares simulations with the 13 May 2019 <u>Pleiades Pléiades</u> 1 image; (d-e-f) is based on the 4 May 2020 <u>Pleiades Pléiades</u> 2 image. (a-d) Bias (m); (b-e)  $\sigma$  ratio (no unit); (c-f) SPS. The important standard deviation ratio values found for the 'Pit' landform type in (b) and (e) are explained because almost all pixels of this type have no observed snow whereas the simulations give snow on every pixel. This is particularly visible in Subfigure (d) and (e) where Antilope, Safran, and Safran HR have almost no simulated snow and a fair standard deviation ratio but Antilope. All scores are presented as a function of landform pixel classification.

![](_page_48_Figure_1.jpeg)

Figure E1. Plots of average SPS values for the SAFRAN and SAFRAN HR simulations when grouped with 300 m elevation bands. This graph illustrates the impact of the computing horizontal resolution on snow height and snow melt-out date.

#### References

- 865 Aksamit, N. O. and Pomeroy, J. W.: Scale Interactions in Turbulence for Mountain Blowing Snow, Journal of Hydrometeorology, 19, 305 320, https://doi.org/https://doi.org/10.1175/JHM-D-17-0179.1, 2018.
  - Amory, C., Kittel, C., Le Toumelin, L., Agosta, C., Delhasse, A., Favier, V., and Fettweis, X.: Performance of MAR (v3.11) in simulating the drifting-snow climate and surface mass balance of Adélie Land, East Antarctica, Geoscientific Model Development, 14, 3487–3510, https://doi.org/10.5194/gmd-14-3487-2021, 2021.
- 870 Anderson, B. T., McNamara, J. P., Marshall, H.-P., and Flores, A. N.: Insights into the physical processes controlling correlations between snow distribution and terrain properties, Water Resources Research, 50, 4545–4563, https://doi.org/https://doi.org/10.1002/2013WR013714, 2014.
  - Baba, M. W., Gascoin, S., Kinnard, C., Marchane, A., and Hanich, L.: Effect of Digital Elevation Model Resolution on the Simulation of the Snow Cover Evolution in the High Atlas, Water Resour. Res., 55, 5360–5378, https://doi.org/10.1029/2018WR023789, 2019.
- 875 Bales, R. C., Molotch, N. P., Painter, T. H., Dettinger, M. D., Rice, R., and Dozier, J.: Mountain hydrology of the western United States, Water Resources Research, 42, https://doi.org/10.1029/2005WR004387, 2006.
  - Baron, M., Haddjeri, A., Lafaysse, M., Le Toumelin, L., Vionnet, V., and Fructus, M.: SnowPappus v1.0, a blowing-snow model for largescale applications of Crocus snow scheme, Geoscientific Model Development Discussions, 17, 1297–1326, https://doi.org/10.5194/gmd-17-1297-2024, 2024.
- 880 Bellaire, S., Katurji, M., Schulmann, T., and Hobman, A.: Towards a high-resolution operational forecasting tool for the southern alps New Zealand, in: Proceedings, International Snow Science Workshop, Banff, 2014, https://arc.lib.montana.edu/snow-science/objects/ISSW14\_ paper\_P1.01.pdf, 2014.
  - Bergeron, T.: On the low-level redistribution of atmospheric water caused by orography, in: Proc. of the Int. Conf. on Cloud Physics, pp. 96–100, https://doi.org/10.1088/0031-9112/16/9/004, 1965.
- 885 Bernhardt, M. and Schulz, K.: SnowSlide: A simple routine for calculating gravitational snow transport, Geophysical Research Letters, 37, https://doi.org/10.1029/2010GL043086, 2010.
  - Bernhardt, M., Liston, G. E., Strasser, U., Zängl, G., and Schulz, K.: High resolution modelling of snow transport in complex terrain using downscaled MM5 wind fields, The Cryosphere, 4, 99–113, https://doi.org/10.5194/tc-4-99-2010, 2010.
- Bernhardt, M., Schulz, K., Liston, G. E., and Zängl, G.: The influence of lateral snow redistribution processes on snow melt and sublimation
   in alpine regions, Journal of Hydrology, 424–425, 196–206, https://doi.org/10.1016/i.ihydrol.2012.01.001, 2012.
  - Brauchli, T., Trujillo, E., Huwald, H., and Lehning, M.: Influence of slope-scale snowmelt on catchment response simulated with the Alpine3D model, Water Resources Research, 53, 10723–10739, https://doi.org/10.1002/2017WR021278, 2017.
    - Brun, E., Martin, E., Simon, V., Gendre, C., and Coleou, C.: An Energy and Mass Model of Snow Cover Suitable for Operational Avalanche Forecasting, Journal of Glaciology, 35, 333–342, https://doi.org/10.3189/S0022143000009254, 1989.
- 895 Candille, G. and Talagrand, O.: Evaluation of probabilistic prediction systems for a scalar variable, Quart. J. Roy. Meteorol. Soc., 131, 2131–2150, https://doi.org/10.1256/qj.04.71, 2005.
  - Champeaux, J. L., Masson, V., and Chauvin, F.: ECOCLIMAP: a global database of land surface parameters at 1 km resolution, Meteorological Applications: A journal of forecasting, practical applications, training techniques and modelling, 12, 29–32, 2005.

Champeaux, J.-L., Dupuy, P., Laurantin, O., Soulan, I., Tabary, P., and Soubeyroux, J.-M.: Les mesures de précipitations et l'estimation des
 lames d'eau à Météo-France : état de l'art et perspectives, La Houille Blanche, 95, 28–34, https://doi.org/10.1051/lhb/2009052, 2009.

50

- Choularton, T. W. and Perry, S. J.: A model of the orographic enhancement of snowfall by the seeder-feeder mechanism, Ouarterly Journal of the Royal Meteorological Society, 112, 335-345, https://doi.org/10.1002/qj.49711247204, 1986.
- Clark, M. P., Hendrikx, J., Slater, A. G., Kavetski, D., Anderson, B., Cullen, N. J., Kerr, T., Hreinsson, E. Ö., and Woods, R. A.: Representing spatial variability of snow water equivalent in hydrologic and land-surface models: A review, Water Resour. Res., 47, W07 539, https://doi.org/10.1029/2011WR010745, 2011.

905

- Cluzet, B., Lafaysse, M., Cosme, E., Albergel, C., Meunier, L.-F., and Dumont, M.: CrocO\_v1.0: a particle filter to assimilate snowpack observations in a spatialised framework, Geosci. Model Dev., 14, 1595–1614, https://doi.org/10.5194/gmd-14-1595-2021, 2021.
- Colle, B. A., Smith, R. B., and Wesley, D. A.: Theory, observations, and predictions of orographic precipitation, in: Mountain Weather Research and Forecasting: Recent Progress and Current Challenges, chap. 6, pp. 291-344. Springer, https://doi.org/10.1007/978-94-007-
- 910 4098-3 6, 2013.
  - Dadic, R., Mott, R., Lehning, M., and Burlando, P.: Wind influence on snow depth distribution and accumulation over glaciers, Journal of Geophysical Research: Earth Surface, 115, https://doi.org/10.1029/2009JF001261, 2010.
  - Decharme, B., Boone, A., Delire, C., and Noilhan, J.: Local evaluation of the Interaction between Soil Biosphere Atmosphere soil multilayer diffusion scheme using four pedotransfer functions, J. Geophys. Res., 116, D20 126, https://doi.org/10.1029/2011JD016002, 2011.
- 915 Deschamps-Berger, C., Gascoin, S., Berthier, E., Deems, J., Gutmann, E., Dehecq, A., Shean, D., and Dumont, M.: Snow depth mapping from stereo satellite imagery in mountainous terrain: evaluation using airborne laser-scanning data, The Cryosphere, 14, 2925–2940, https://doi.org/10.5194/tc-14-2925-2020, 2020.
- Deschamps-Berger, C., Cluzet, B., Dumont, M., Lafaysse, M., Berthier, E., Fanise, P., and Gascoin, S.: Improving the Spatial Distribution of Snow Cover Simulations by Assimilation of Satellite Stereoscopic Imagery, Water Resour. Res., 58, 920 https://doi.org/10.1029/2021WR030271, 2022.
  - Durand, Y., Brun, E., Mérindol, L., Guyomarc'h, G., Lesaffre, B., and Martin, E.: A meteorological estimation of relevant parameters for snow models, Ann. Glaciol., 18, 65–71, https://doi.org/10.3189/S0260305500011277, 1993.
    - Earth System Modeling Framework, University Corporation for Atmospheric Research, Massachusetts Institute of Technology, Geophysical Fluid Dynamics Laboratory, University of Michigan, National Centers for Environmental Prediction, Los Alamos National Laboratory,
- 925 Argonne National Laboratory, and NASA Goddard Space Flight Center: Earth System Modeling Framework (ESMF) Grid Remapping (Version 8.4.2), https://earthsystemmodeling.org/, 2023.
  - Essery, R., Li, L., and Pomeroy, J.: A distributed model of blowing snow over complex terrain, Hydrological Processes, 13, 2423–2438, https://doi.org/10.1002/(SICI)1099-1085(199910)13:14/15<2423::AID-HYP853>3.0.CO;2-U, 1999.

Faure, D.: Quality Analysis of the 2016 Quantitative Precipitation Estimates in the French Alps, in: 37th Conference on radar meteorology,

- 930 AMS, https://ams.confex.com/ams/38RADAR/webprogram/Paper320389.html, 2017.
  - Faure, D., Delrieu, G., and Gaussiat, N.: Impact of the Altitudinal Gradients of Precipitation on the Radar QPE Bias in the French Alps, Atmosphere, 10, https://doi.org/10.3390/atmos10060306, 2019.
- Foresti, L., Sideris, I. V., Panziera, L., Nerini, D., and Germann, U.: A 10-year radar-based analysis of orographic precipitation growth and decay patterns over the Swiss Alpine region, Quarterly Journal of the Royal Meteorological Society, 144, 2277–2301, 935 https://doi.org/https://doi.org/10.1002/qj.3364, 2018.
  - Freudiger, D., Kohn, I., Seibert, J., Stahl, K., and Weiler, M.: Snow redistribution for the hydrological modeling of alpine catchments, WIREs Water, 4, e1232, https://doi.org/10.1002/wat2.1232, 2017.

Froidurot, S., Zin, I., Hingray, B., and Gautheron, A.: Sensitivity of precipitation phase over the Swiss Alps to different meteorological variables, Journal of Hydrometeorology, 15, 685–696, https://doi.org/10.1175/JHM-D-13-073.1, 2014.

- 940 Gascoin, S., Grizonnet, M., Bouchet, M., Salgues, G., and Hagolle, O.: Theia Snow collection: high-resolution operational snow cover maps from Sentinel-2 and Landsat-8 data, Earth Syst. Sci. Data, 11, 493–514, https://doi.org/10.5194/essd-11-493-2019, 2019.
  - GDAL/OGR contributors: GDAL/OGR Geospatial Data Abstraction software Library (Version 3.6.2), Open Source Geospatial Foundation, https://doi.org/10.5281/zenodo.5884351, 2023.
- Germann, U., Boscacci, M., Clementi, L., Gabella, M., Hering, A., Sartori, M., Sideris, I. V., and Calpini, B.: Weather Radar in Complex
   Orography. Remote Sensing, 14, https://doi.org/10.3390/rs14030503, 2022.
  - Goessling, H. and Jung, T.: A probabilistic verification score for contours: Methodology and application to Arctic ice-edge forecasts, Quarterly Journal of the Royal Meteorological Society, 144, 735–743, https://doi.org/10.1002/qj.3242, 2018.
- Gouttevin, I., Vionnet, V., Seity, Y., Boone, A., Lafaysse, M., Deliot, Y., and Merzisen, H.: To the Origin of a Wintertime Screen-Level Temperature Bias at High Altitude in a Kilometric NWP Model, J. Hydrometeorol., 24, 53 71, https://doi.org/10.1175/JHM-D-21-0200.1, 2023.
  - Grünewald, T., Stötter, J., Pomeroy, J. W., Dadic, R., Moreno Baños, I., Marturià, J., Sproß, M., Hopkinson, C., Burlando, P., and Lehning, M.: Statistical modelling of the snow depth distribution in open alpine terrain, Hydrology and Earth System Sciences, 17, 3005–3021, https://doi.org/10.5194/hess-17-3005-2013, 2013.

- Grünewald, T. and Lehning, M.: Are flat-field snow depth measurements representative? A comparison of selected index sites with areal snow depth measurements at the small catchment scale, Hydrological Processes, 29, 1717–1728, https://doi.org/10.1002/hyp.10295, 2015.
- Guyomarc'h, G., Bellot, H., Vionnet, V., Naaim-Bouvet, F., Déliot, Y., Fontaine, F., Puglièse, P., Nishimura, K., Durand, Y., and Naaim, M.: A meteorological and blowing snow data set (2000–2016) from a high-elevation alpine site (Col du Lac Blanc, France, 2720 m a.s.l.),
- 960 Earth Syst. Sci. Data, 11, 57–69, https://doi.org/10.5194/essd-11-57-2019, 2019.
  - Guyomarc'h, G. and Mérindol, L.: Validation of an application for forecasting blowing snow, Annals of Glaciology, 26, 138–143, https://doi.org/10.3189/1998AoG26-1-138-143, 1998.
    - Günther, D., Marke, T., Essery, R., and Strasser, U.: Uncertainties in Snowpack Simulations—Assessing the Impact of Model Structure, Parameter Choice, and Forcing Data Error on Point-Scale Energy Balance Snow Model Performance, Water Resour. Res., 55, 2779–2800,
- 965 https://doi.org/10.1029/2018WR023403, 2019.

955

Hofmeister, F., Arias-Rodriguez, L. F., Premier, V., Marin, C., Notarnicola, C., Disse, M., and Chiogna, G.: Intercomparison of Sentinel-2 and modelled snow cover maps in a high-elevation Alpine catchment, Journal of Hydrology X, 15, 100123, https://doi.org/10.1016/j.hydroa.2022.100123, 2022.

Horton, S. and Haegeli, P.: Using snow depth observation to provide insight into the quality of regional-scale snowpack simulations for
avalanche forecasting, EGUsphere, 2022, 1–28, https://doi.org/10.5194/egusphere-2022-237, 2022.

IGN©: BD Foret®V2, https://geoservices.ign.fr/bdforet, 2021a.

IGN©: BD Topo®, https://geoservices.ign.fr/bdtopo, 2021b.

IGN©: RGE ALTI®5M, https://geoservices.ign.fr/rgealti, 2021c.

IGN©: BD Ortho®, https://geoservices.ign.fr/bdortho, 2022.

975 IPCC: The Ocean and Cryosphere in a Changing Climate, Cambridge University Press, https://doi.org/10.1017/9781009157964, 2022.

Grünewald, T., Bühler, Y., and Lehning, M.: Elevation dependency of mountain snow depth, The Cryosphere, 8, 2381–2394, https://doi.org/10.5194/tc-8-2381-2014, 2014.

- Jasiewicz, J. and Stepinski, T. F.: Geomorphons—a pattern recognition approach to classification and mapping of landforms, Geomorphology, 182, 147–156, https://doi.org/10.1016/j.geomorph.2012.11.005, 2013.
- Lafaysse, M., Cluzet, B., Dumont, M., Lejeune, Y., Vionnet, V., and Morin, S.: A multiphysical ensemble system of numerical snow modelling, The Cryosphere, 11, 1173–1198, https://doi.org/10.5194/tc-11-1173-2017, 2017.
- 980 Le Toumelin, L., Gouttevin, I., Helbig, N., Galiez, C., Roux, M., and Karbou, F.: Emulating the adaptation of wind fields to complex terrain with deep-learning, Artif. Intell. Earth Syst., pp. 1 – 39, https://doi.org/10.1175/AIES-D-22-0034.1, 2022.
  - Le Toumelin, L., Gouttevin, I., Galiez, C., and Helbig, N.: A two-folds deep learning strategy to correct and downscale winds over mountains, Nonlinear Processes in Geophysics, 2023, 1–32, https://doi.org/10.5194/npg-31-75-2024, 2023.
- Lehning, M., Völksch, I., Gustafsson, D., Nguyen, T. A., Stähli, M., and Zappa, M.: ALPINE3D: a detailed model of moun tain surface processes and its application to snow hydrology, Hydrological Processes: An International Journal, 20, 2111–2128,
  - Lehning, M., Löwe, H., Ryser, M., and Raderschall, N.: Inhomogeneous precipitation distribution and snow transport in steep terrain, Water Resources Research, 44, https://doi.org/10.1029/2007WR006545, 2008.

Lindsay, J.: The whitebox geospatial analysis tools project and open-access GIS, in: Proceedings of the GIS Research UK 22nd Annual

- 990 Conference, The University of Glasgow, pp. 16–18, https://www.gla.ac.uk/media/Media\_401757\_smxx.pdf, 2014.
  - Liston, G. and Sturm, M.: Α snow-transport model for complex terrain. J. Glaciol.. 44. 498-516, https://doi.org/10.3189/S0022143000002021, 1998.
    - Liston, G. E., Haehnel, R. B., Sturm, M., Hiemstra, C. A., Berezovskaya, S., and Tabler, R. D.: Simulating complex snow distributions in windy environments using SnowTran-3D, J. Glaciol., 53, 241–256, https://doi.org/10.3189/172756507782202865, 2007.
- 995 Lundquist, J., Hughes, M., Gutmann, E., and Kapnick, S.: Our Skill in Modeling Mountain Rain and Snow is Bypassing the Skill of Our Observational Networks, Bulletin of the American Meteorological Society, 100, 2473–2490, https://doi.org/10.1175/BAMS-D-19-0001.1, 2019.
- Marsh, C. B., Pomeroy, J. W., and Wheater, H. S.: The Canadian Hydrological Model (CHM) v1.0: a multi-scale, multi-extent, variable-complexity hydrological model design and overview, Geosci. Model Dev., 13, 225–247, https://doi.org/10.5194/gmd-13-225-2020, 2020.
  - Marsh, C. B., Vionnet, V., and Pomeroy, J. W.: Windmapper: An Efficient Wind Downscaling Method for Hydrological Models, Water Resources Research, 59, e2022WR032683, https://doi.org/10.1029/2022WR032683, 2023.
  - Marti, R., Gascoin, S., Berthier, E., de Pinel, M., Houet, T., and Laffly, D.: Mapping snow depth in open alpine terrain from stereo satellite imagery, The Cryosphere, 10, 1361–1380, https://doi.org/10.5194/tc-10-1361-2016, 2016.
- 1005 Masson, V., Le Moigne, P., Martin, E., Faroux, S., Alias, A., Alkama, R., Belamari, S., Barbu, A., Boone, A., Bouyssel, F., Brousseau, P., Brun, E., Calvet, J.-C., Carrer, D., Decharme, B., Delire, C., Donier, S., Essaouini, K., Gibelin, A.-L., Giordani, H., Habets, F., Jidane, M., Kerdraon, G., Kourzeneva, E., Lafaysse, M., Lafont, S., Lebeaupin Brossier, C., Lemonsu, A., Mahfouf, J.-F., Marguinaud, P., Mokhtari, M., Morin, S., Pigeon, G., Salgado, R., Seity, Y., Taillefer, F., Tanguy, G., Tulet, P., Vincendon, B., Vionnet, V., and Voldoire, A.: The SURFEXv7.2 land and ocean surface platform for coupled or offline simulation of Earth surface variables and fluxes, Geosci. Model Dev.,
- 1010 6, 929–960, https://doi.org/10.5194/gmd-6-929-2013, 2013.

https://doi.org/10.1002/hyp.6204, 2006.

Ménard, C. B., Essery, R., Barr, A., Bartlett, P., Derry, J., Dumont, M., Fierz, C., Kim, H., Kontu, A., Lejeune, Y., Marks, D., Niwano, M., Raleigh, M., Wang, L., and Wever, N.: Meteorological and evaluation datasets for snow modelling at 10 reference sites: description of in situ and bias-corrected reanalysis data, Earth Syst. Sci. Data, 11, 865–880, https://doi.org/10.5194/essd-11-865-2019, 2019. Menard, C. B., Essery, R., Krinner, G., Arduini, G., Bartlett, P., Boone, A., Brutel-Vuilmet, C., Burke, E., Cuntz, M., Dai, Y., Decharme,

- B., Dutra, E., Fang, X., Fierz, C., Gusev, Y., Hagemann, S., Haverd, V., Kim, H., Lafaysse, M., Marke, T., Nasonova, O., Nitta, T., Niwano, M., Pomeroy, J., Schädler, G., Semenov, V. A., Smirnova, T., Strasser, U., Swenson, S., Turkov, D., Wever, N., and Yuan, H.: Scientific and Human Errors in a Snow Model Intercomparison, Bulletin of the American Meteorological Society, 102, E61 E79, https://doi.org/10.1175/BAMS-D-19-0329.1, 2021.
- Minder, J. R., Durran, D. R., and Roe, G. H.: Mesoscale controls on the mountainside snow line, Journal of the atmospheric sciences, 68, 2107–2127, https://doi.org/10.1175/JAS-D-10-05006.1, 2011.
  - Mital, U., Dwivedi, D., Brown, J. B., and Steefel, C. I.: Downscaled hyper-resolution (400m) gridded datasets of daily precipitation and temperature (2008–2019) for the East–Taylor subbasin (western United States), Earth System Science Data, 14, 4949–4966, https://doi.org/10.5194/essd-14-4949-2022, 2022.
- Monteiro, D., Caillaud, C., Samacoïts, R., Lafaysse, M., and Morin, S.: Potential and limitations of convectionpermitting CNRM-AROME climate modelling in the French Alps, International Journal of Climatology, 42, 7162–7185,
  - https://doi.org/https://doi.org/10.1002/joc.7637, 2022.
    Morin, S., Horton, S., Techel, F., Bavay, M., Coléou, C., Fierz, C., Gobiet, A., Hagenmuller, P., Lafaysse, M., Ližar, M., Mitterer, C., Monti, F., Müller, K., Olefs, M., Snook, J. S., van Herwijnen, A., and Vionnet, V.: Application of physical snowpack models in support of operational avalanche hazard forecasting: A status report on current implementations and prospects for the future, Cold Reg. Sci. Tech.,
- 1030 170, 102 910, https://doi.org/10.1016/j.coldregions.2019.102910, 2020.
  - Mott, R., Faure, F., Lehning, M., Löwe, H., Hynek, B., Michlmayer, G., Prokop, A., and Schöner, W.: Simulation of seasonal snow-cover distribution for glacierized sites on Sonnblick, Austria, with the Alpine3D model, Annals of Glaciology, 49, 155–160, https://doi.org/10.3189/172756408787814924, 2008.
- Mott, R., Schirmer, M., Bavay, M., Grünewald, T., and Lehning, M.: Understanding snow-transport processes shaping the mountain snowcover, The Cryosphere, 4, 545–559, https://doi.org/10.5194/tc-4-545-2010, publisher: Copernicus GmbH, 2010.
  - Mott, R., Scipión, D., Schneebeli, M., Dawes, N., Berne, A., and Lehning, M.: Orographic effects on snow deposition patterns in mountainous terrain, Journal of Geophysical Research: Atmospheres, 119, 1419–1439, https://doi.org/10.1002/2013JD019880, 2014.
    - Mott, R., Vionnet, V., and Grünewald, T.: The Seasonal Snow Cover Dynamics: Review on Wind-Driven Coupling Processes, Front. Earth Sci., 6, 197, https://doi.org/10.3389/feart.2018.00197, 2018.
- 1040 Mott, R., Winstral, A., Cluzet, B., Helbig, N., Magnusson, J., Mazzotti, G., Quéno, L., Schirmer, M., Webster, C., and Jonas, T.: Operational snow-hydrological modeling for Switzerland, Frontiers in Earth Science, 11, 1228 158, https://doi.org/10.3389/feart.2023.1228158, 2023. Musselman, K. N., Pomeroy, J. W., Essery, R. L., and Leroux, N.: Impact of windflow calculations on simulations of alpine snow accumula
  - tion, redistribution and ablation, Hydrological Processes, 29, 3983–3999, https://doi.org/10.1002/hyp.10595, 2015.
- Nachtergaele, F., Velthuizen, H., Verelst, L., and Wiberg, D.: Harmonized world soil database (hwsd), Food and Agriculture Organization of the United Nations, Rome, 2009.
  - Pepin, N., Bradley, R. S., Diaz, H. F., Baraer, M., Caceres, E. B., Forsythe, N., Fowler, H., Greenwood, G., Hashmi, M. Z., Liu, X. D., Miller, J. R., Ning, L., Ohmura, A., Palazzi, E., Rangwala, I., Schöner, W., Severskiy, I., Shahgedanova, M., Wang, M. B., Williamson, S. N., Yang, D. Q., and Mountain Research Initiative EDW Working Group: Elevation-dependent warming in mountain regions of the world, Nature Climate Change, 5, 424–430, https://doi.org/10.1038/nclimate2563, number: 5 Publisher: Nature Publishing Group, 2015.
- 1050 Pomeroy, MacDonald, M., DeBeer, C., and Brown, T.: Modeling Alpine Snow Hydrology in the Canadian Rocky Mountains, https: //westernsnowconference.org/node/747, 2009.

Pomeroy, J. and Gray, D.: Snow accumulation, relocation and management, vol. 144, https://hdl.handle.net/10388/15161, 1995.

Pomeroy, J., Gray, D., Shook, K., Toth, B., Essery, R., Pietroniro, A., and Hedstrom, N.: An evaluation of snow accumulation and ablation processes for land surface modelling, Hydrological Processes, 12, 2339–2367, https://doi.org/10.1002/(SICI)1099-1085(199812)12:

1055 15<2339::AID-HYP800>3.0.CO;2-L, 1998.

1070

1085

- Prasad, R., Tarboton, D. G., Liston, G. E., Luce, C. H., and Seyfried, M. S.: Testing a blowing snow model against distributed snow measurements at Upper Sheep Creek, Idaho, United States of America, Water Resources Research, 37, 1341–1356, https://doi.org/10.1029/2000WR900317, 2001.
- Quéno, L., Karbou, F., Vionnet, V., and Dombrowski-Etchevers, I.: Satellite-derived products of solar and longwave irradiances used for
- 1060 snowpack modelling in mountainous terrain, Hydrology and Earth System Sciences, 24, 2083–2104, https://doi.org/10.5194/hess-24-2083-2020, 2020.
  - Quéno, L., Vionnet, V., Dombrowski-Etchevers, I., Lafaysse, M., Dumont, M., and Karbou, F.: Snowpack modelling in the Pyrenees driven by kilometric-resolution meteorological forecasts, The Cryosphere, 10, 1571–1589, https://doi.org/10.5194/tc-10-1571-2016, 2016.
- Quéno, L., Mott, R., Morin, P., Cluzet, B., Mazzotti, G., and Jonas, T.: Snow redistribution in an intermediate-complexity snow hydrology
   modelling framework, EGUsphere, p. 1–32, https://doi.org/10.5194/egusphere-2023-2071, 2023.
  - Raleigh, M. S., Lundquist, J. D., and Clark, M. P.: Exploring the impact of forcing error characteristics on physically based snow simulations within a global sensitivity analysis framework, Hydrol. Earth Syst. Sci., 19, 3153–3179, https://doi.org/10.5194/hess-19-3153-2015, 2015.

Revuelto, J., Lecourt, G., Lafaysse, M., Zin, I., Charrois, L., Vionnet, V., Dumont, M., Rabatel, A., Six, D., Condom, T., et al.: Multi-Criteria Evaluation of Snowpack Simulations in Complex Alpine Terrain Using Satellite and In Situ Observations, Remote Sens., 10, 1171, https://doi.org/10.3390/rs10081171, 2018.

- RGI Consortium: Randolph Glacier inventory-a dataset of global glacier outlines: Version 6.0, https://doi.org/https://doi.org/10.7265/4m1f-gd79, 2017.
  - Rüschendorf, L.: The Wasserstein distance and approximation theorems, Probability Theory and Related Fields, 70, 117–129, https://doi.org/10.1007/BF00532240, 1985.
- 1075 Sato, T., Kimura, T., Ishimaru, T., and Maruyama, T.: Field test of a new snow-particle counter (SPC) system, Annals of Glaciology, 18, 149–154, https://doi.org/10.3189/S0260305500011411, 1993.
  - Schirmer, M. and Jamieson, B.: Verification of analysed and forecasted winter precipitation in complex terrain, The Cryosphere, 9, 587–601, https://doi.org/10.5194/tc-9-587-2015, 2015.
  - Schlögl, S., Marty, C., Bavay, M., and Lehning, M.: Sensitivity of Alpine3D modeled snow cover to modifications in
- 1080 DEM resolution, station coverage and meteorological input quantities, Environmental Modelling & Software, 83, 387–396, https://doi.org/10.1016/j.envsoft.2016.02.017, 2016.
  - Schneiderbauer, S. and Prokop, A.: The atmospheric snow-transport model: SnowDrift3D, Journal of Glaciology, 57, 526–542, https://doi.org/10.3189/002214311796905677, 2011.
  - Seity, Y., Brousseau, P., Malardel, S., Hello, G., Bénard, P., Bouttier, F., Lac, C., and Masson, V.: The AROME-France convective scale operational model, Monthly Weather Review, 129, 976–991, https://doi.org/10.1175/2010MWR3425.1, 2011.
  - Sen Gupta, A. and Tarboton, D. G.: A tool for downscaling weather data from large-grid reanalysis products to finer spatial scales for distributed hydrological applications, Environmental Modelling & Software, 84, 50–69, https://doi.org/10.1016/j.envsoft.2016.06.014, 2016.

Sharma, V., Gerber, F., and Lehning, M.: Introducing CRYOWRF v1.0: Multiscale atmospheric flow simulations with advanced snow cover

- 1090 modelling, Geoscientific Model Development Discussions, 2021, 1–46, https://doi.org/10.5194/gmd-2021-231, 2021.
  - Strasser, U., Bernhardt, M., Weber, M., Liston, G., and Mauser, W.: Is snow sublimation important in the alpine water balance?, The Cryosphere, 2, 53–66, https://doi.org/10.5194/tc-2-53-2008, 2008.
    - Tong, J., Déry, S., and Jackson, P.: Topographic control of snow distribution in an alpine watershed of western Canada inferred from spatiallyfiltered MODIS snow products, Hydrology and Earth System Sciences, 13, 319–326, https://doi.org/10.5194/hess-13-319-2009, 2009.
- 1095 Vernay, M., Lafaysse, M., Monteiro, D., Hagenmuller, P., Nheili, R., Samacoïts, R., Verfaillie, D., and Morin, S.: The S2M meteorological and snow cover reanalysis over the French mountainous areas: description and evaluation (1958–2021), Earth System Science Data, 14, 1707–1733, https://doi.org/10.5194/essd-14-1707-2022, 2022.
  - Vionnet, V., Brun, E., Morin, S., Boone, A., Martin, E., Faroux, S., Le-Moigne, P., and Willemet, J.-M.: The detailed snowpack scheme Crocus and its implementation in SURFEX v7.2, Geosci. Model. Dev., 5, 773–791, https://doi.org/10.5194/gmd-5-773-2012, 2012.
- 1100 Vionnet, V., Martin, E., Masson, V., Guyomarc'h, G., Naaim-Bouvet, F., Prokop, A., Durand, Y., and Lac, C.: Simulation of wind-induced snow transport and sublimation in alpine terrain using a fully coupled snowpack/atmosphere model, The Cryosphere, 8, 395–415, https://doi.org/10.5194/tc-8-395-2014, 2014.
  - Vionnet, V., Martin, E., Masson, V., Lac, C., Naaim Bouvet, F., and Guyomarc'h, G.: High-Resolution Large Eddy Simulation of Snow Accumulation in Alpine Terrain, J. Geophys. Res. Atmos., 122, 11,005–11,021, https://doi.org/10.1002/2017JD026947, 2017.
- 1105 Vionnet, V., Six, D., Auger, L., Dumont, M., Lafaysse, M., Quéno, L., Réveillet, M., Dombrowski Etchevers, I., Thibert, E., and Vincent, C.: Sub-kilometer precipitation datasets for snowpack and glacier modeling in alpine terrain, Front. Earth Sci., 7, 182, https://doi.org/10.3389/feart.2019.00182, 2019.
  - Vionnet, V., Marsh, C. B., Menounos, B., Gascoin, S., Wayand, N. E., Shea, J., Mukherjee, K., and Pomeroy, J. W.: Multi-scale snowdriftpermitting modelling of mountain snowpack, The Cryosphere, 15, 743–769, https://doi.org/10.5194/tc-15-743-2021, 2021.
- 1110 Vionnet, V., Verville, M., Fortin, V., Brugman, M., Abrahamowicz, M., Lemay, F., Thériault, J. M., Lafaysse, M., and Milbrandt, J. A.: Snow Level From Post-Processing of Atmospheric Model Improves Snowfall Estimate and Snowpack Prediction in Mountains, Water Resour. Res., 58, e2021WR031778, https://doi.org/10.1029/2021WR031778, 2022.
  - Vionnet, V., Dombrowski-Etchevers, I., Lafaysse, M., Quéno, L., Seity, Y., and Bazile, E.: Numerical weather forecasts at kilometer scale in the French Alps : evaluation and applications for snowpack modelling, J. Hydrometeor., https://doi.org/10.1175/JHM-D-15-0241.1, 2016.
- 1115 Wang, Z. and Huang, N.: Numerical simulation of the falling snow deposition over complex terrain, Journal of Geophysical Research: Atmospheres, 122, 980–1000, https://doi.org/10.1002/2016JD025316, 2017.
  - Whitebox Geospatial: WhiteboxTools open core, Whitebox Geospatial Inc, https://www.whiteboxgeo.com/manual/wbt\_book/available\_tools/geomorphometric\_analysis.html#Geomorphons, 2023.
- Whiteman, C. D.: Mountain meteorology: fundamentals and applications, Oxford University Press, https://doi.org/10.1093/oso/9780195132717.001.0001, 2000.
  - Winstral, A., Elder, K., and Davis, R. E.: Spatial snow modeling of wind-redistributed snow using terrain-based parameters, Journal of hydrometeorology, 3, 524–538, https://doi.org/10.1175/1525-7541(2002)003<0524:SSMOWR>2.0.CO;2, 2002.
  - Yang, J., Yau, M., Fang, X., and Pomeroy, J.: A triple-moment blowing snow-atmospheric model and its application in computing the seasonal wintertime snow mass budget, Hydrol. Earth Sys. Sci., 14, 1063–1079, https://doi.org/10.5194/hess-14-1063-2010, 2010.
- 1125 Yu, N., Gaussiat, N., and Tabary, P.: Polarimetric X-band weather radars for quantitative precipitation estimation in mountainous regions, Quarterly Journal of the Royal Meteorological Society, 144, 2603–2619, https://doi.org/10.1002/qj.3366, 2018.