



Suppressed atmospheric chemical aging of cooking organic aerosol particles in wintertime conditions

Wenli Liu¹, Longkun He², Yingjun Liu², Keren Liao², Qi Chen², Mikinori Kuwata¹

¹Department of Atmospheric and Oceanic Sciences, Laboratory for Climate and Ocean-Atmosphere Studies, Peking University, Beijing, 100871, China

²State Key Joint Laboratory of Environmental Simulation and Pollution Control, BIC-ESAT and IJRC, College of Environmental Sciences and Engineering, Peking University, Beijing, 100871, China

Correspondence to: Mikinori Kuwata (kuwata@pku.edu.cn)

Abstract.

Cooking organic aerosol (COA) is one of the major constituents of particulate matter in urban areas. COA is oxidized by atmospheric oxidants such as ozone, changing its physical, chemical and toxicological properties. However, atmospheric chemical lifetimes of COA and its tracers such as oleic acid are typically longer than that have been estimated by laboratory studies. We tackled the issue by considering temperature. Namely, we hypothesize that increased viscosity of COA at ambient temperature accounts for its prolonged atmospheric chemical lifetimes in wintertime. Laboratory generated COA particles from cooking oil were exposed to ozone in an aerosol flow tube reactor for the temperature range of -20 °C ~ 35 °C. The pseudo-second order chemical reaction rate constants (k_2) decreased by orders of magnitude for lower temperatures. The temperature dependence of k_2 was fit well by considering diffusion-limited chemical reaction mechanism, suggesting that reduced viscosity was responsible for the decrease in chemical reactivity. In combination with the observed global surface temperature, the atmospheric chemical lifetimes of COA were estimated to be much longer in wintertime (>1 hour) than that in summertime (a few minutes) for temperate and boreal regions. Our present study demonstrates that the oxidation lifetimes of COA particles will need to be parameterized as a function of temperature in the future for estimating environmental impacts and fates of this category of particulate matter.

1 Introduction

Organic aerosol (OA) is a major component of atmospheric particles (Zhang et al., 2011; Laskin et al., 2019). OA plays important roles in the climate and air quality (Menon et al., 2008; Liu et al., 2013; Robinson et al., 2018; Liu et al., 2019). One of the key sources of OA in urban areas is cooking (Rogge et al., 1991; Hildemann et al., 1991; Crippa et al., 2013; Lee et al., 2015; Sun et al., 2016; Daellenbach et al., 2017; Guo et al., 2020; Huang et al., 2021). Atmospheric abundance of COA is typically estimated by factor analysis of aerosol mass spectra or by employing chemical tracers (Zhang et al., 2011; Huang et al., 2021). Hildemann et al. identified that cooking organic aerosol (COA) accounted for 21% of mass loading of OA in Los Angeles, USA (Hildemann et al., 1991). COA was also identified in European cities (Crippa et al., 2013; Daellenbach et al., 2017). In the case of the Asian region, high abundances of COA have been observed especially at cities in China, including

Beijing, Shanghai, Guangzhou, and Hong Kong (Lee et al., 2015; Sun et al., 2016; Guo et al., 2020; Huang et al., 2021). The mass fractions of COA are highly variable (8-33%), which might be explained by difference in local source distributions and uncertainties in source apportionment (Zheng et al., 2023).

35 Processes involved in COA emissions include meat cooking operations and stir/deep frying by vegetable oils (Zhao et al., 2007; Weitkamp et al., 2008). In both cases, fatty acids such as oleic, linoleic, palmitic, and stearic acids were identified as the major marker compounds for COA (Song et al., 2023; Zhao et al., 2007). As COA contains unsaturated fatty acids such as oleic and linoleic acids, oxidation of COA occurs following chemical reactions with ozone, OH, and NO₃ (Weitkamp et al., 2008; Li et al., 2020; Wang and Yu, 2021).

40 The chemical aging processes of COA and its surrogates change their hygroscopicity and toxicity (Liu et al., 2021; Wang et al., 2021). For instance, ozonolysis of oleic and linoleic acids particles makes them to be highly active as cloud condensation nuclei (CCN) (Broekhuizen et al., 2004; Shilling et al., 2007). Oxidative potential of COA enhances following oxidation (Wang et al., 2021). In addition, COA aging processes need to be well understood for appropriately appointing sources of OA (Reyes-Villegas et al., 2018). It should be noted that the COA factor from factor analysis of the aerosol mass spectrometric
45 measurements may contain both fresh and aged COA, while the profiles of molecular level chemical tracers may change due to atmospheric oxidation (Huang et al., 2021).

The discrepancy in the atmospheric chemical lifetimes of COA/corresponding marker compounds and estimations of the corresponding values by laboratory experiments have been discussed for a few decades (Robinson et al., 2006; Rudich et al., 2007; Weitkamp et al., 2008; Wang and Yu, 2021). Atmospheric chemical lifetimes of oleic and linoleic acids estimated by
50 laboratory experiments are typically in the order of minutes (Morris et al., 2002; Knopf et al., 2005; Hung and Tang, 2010; Liu et al., 2023). On the other hand, a field observation in Los Angeles, USA suggested that the atmospheric chemical lifetime of oleic acid was much longer than the time scale (Rogge et al., 1991; Zahardis and Petrucci, 2007). Recent observational studies in the Yangtze River Delta region in China suggested that the nighttime chemical lifetimes of oleic acid from cooking in the region was in the order of hours (Wang and Yu, 2021).

55 The discrepancy among the field and laboratory chemical lifetimes of COA markers has typically been associated with chemical composition of COA (Rudich et al., 2007). Namely, mixing of saturated fatty acids from cooking such as stearic and palmitic acids makes COA to be highly viscous or solid at room temperature, reducing chemical reactivity (Knopf et al., 2005). Chemical reactions of COA with ozone also lead to phase transition of liquid to semi-solid or solid phases, potentially due to the formation of high-molecular-weight chemical species (Xu et al., 2022).

60 Recent laboratory studies started suggesting that temperature may also be an important factor for regulating chemical reactivity of COA (Li et al., 2020; Kaur Kohli et al., 2023; Liu et al., 2023). Reactive uptake coefficient (γ) of ozone by canola oil film on a coated flow tube reactor decreased by an order of magnitude at the melting point (Li et al., 2020). Ozonolysis of oleic acid particles that contain ammonium sulfate seeds became unmeasurably slow by an aerosol flow tube reactor for the temperature range below its melting point (Liu et al., 2023). Considering that ambient temperatures during wintertime in many
65 cities in the world are significantly lower than a typical laboratory condition (20 ~ 25 °C), temperature could also play an




important role in regulating the atmospheric chemical lifetimes of COA. However, previous temperature dependent oxidation experiments of COA were conducted with organic films or droplets on substrates, meaning that phase transition process was influenced by heterogeneous nucleation (Hung and Tang, 2010; Li et al., 2020; Liu et al., 2023).

In this study, we have conducted laboratory experiments for testing the role of temperature on chemical reactivity of COA.

70 The laboratory study was conducted using an aerosol flow tube for the temperature range of $-20\text{ }^{\circ}\text{C}$ to $35\text{ }^{\circ}\text{C}$. The COA particles for the experiments were produced by heating canola oil, Chinese-style hot pot soup base, and lard. The experimental results were used for evaluating changes in chemical lifetimes of COA as a function of temperature.

2 Materials and Methods

Figure 1 shows the experimental setup. The experimental setup consisted of three components, including particle generator, 75 aerosol flow tube, and online instruments for quantifying the chemical composition and size distribution of aerosol particles. Purified air produced by the zero-air generator (Model 747–30, AADCO Instruments, Inc.) was employed for the setup. Table S1 summarizes the list of experimental runs. Further details about the experimental setup can be found in our previous publication (Liu et al., 2023).



80 **Figure 1. Schematic diagram of the experimental setup. Particles were measured using the SV-TAG and OPC for Exp. #8, 15 and 22. The ACSM and SEMS were employed for other experiments.**

2.1 Generation of COA particles

Three types of cooking materials, including canola oil (product ID: 10057184957713, Jinlongyu Group Co., Ltd.), Chinese-style hot pot soup base (Sichuan spicy hot pot, product ID: 858421, Haidilao) and lard (produced by a cafeteria in the campus 85 of Peking University (PKU)), were employed for the experiment. Lard was extracted by heating pork fat using a Chinese frying pan for around half hour following addition of a trace amount of vegetable oil and water. The samples were stored in a refrigerator until usage.

COA particles were generated by heating cooking oils to $180 \pm 2\text{ }^{\circ}\text{C}$ using a U-tube wrapped with six layers of aluminium foils and a heating tape inside. A thermocouple was installed for measuring the temperature of the outer wall of the U-tube.

90 Approximately 3 ml of samples were put in the bottom of the U-tube. The hot pot soup base contained some solid components, including spices. Only the supernatant liquid was employed for the experiment. A small amount of lard was heated during the



sample preparation process, as it forms solid phase at room temperature. The air flow rate passed through the U-tube was controlled by a mass flow controller (MFC, Alicat Scientific, Inc.) at 2 lpm. Nucleation and condensation of the heated samples occurred in an air-cooled condenser. An impactor (Model 8008, Brechtel Manufacturing, Inc.) was used to removed large
95 particles. Volatile organic compounds (VOCs) were removed by a diffusion dryer (AGS-Dryer, Brechtel Manufacturing, Inc.) containing activated charcoal for avoiding formation of secondary organic aerosol particles during the ozonolysis experiments. Particles were stored in a 100 l stainless-steel container for three hours for growing them by coagulation prior to conducting oxidation experiments. The mode diameters of particles following coagulation were 300 – 400 nm (Fig. S1). The size ranges were comparable to the ambient COA particles in Beijing (Ma et al., 2023).

100 2.2 Temperature-controlled aerosol flow tube

An aerosol flow tube, which was installed in a temperature-controlled box, was employed for the ozonolysis reaction. The aerosol flow tube consisted of a borosilicate glass tube (5 cm in inner diameter, and 1 m in length) and a movable injector (1/4 inch stainless steel tubing supported by 3/8 inch stainless steel tubing). The positions of the movable injector were changed for investigating oxidation kinetics for the reaction time of 0 to 60 seconds at room temperature. Temperature of the box for
105 the aerosol flow tube was controlled for the range of -20 °C to 35 °C by a chest freezer and a temperature-controlled liquid circulator. The temporal variation of temperature was ± 0.2 °C at -20 °C.

Ozone was produced using an ultraviolet lamp (UV-M, Beijing Tonglin Technology Co., Ltd). Ozone concentration was continuously monitored by an ozone analyzer (Model 49i, Thermo Fisher Scientific, Inc.). Ozone concentration was adjusted to be 450 ppb and 7 ppm for normal and high ozone exposure experiments, respectively. A diffusion dryer containing the
110 ozone destruction catalyst (F800, Beijing Tonglin Technology Co., Ltd) was connected immediately after the flow tube for terminating chemical reactions.

2.3 Particle characterization

Chemical composition of particles was monitored using the time-of-flight aerosol chemical speciation monitor (ToF-ACSM, Aerodyne Research, Inc.) equipped with the PM_{2.5} aerodynamic lens and capture vaporizer (Zheng et al., 2020). The mass-to-
115 charge (m/z) range of the mass spectra was 12 to 210. The multilinear engine (ME-2) solver was employed for factor analysis of the aerosol mass spectra (Budisulistiorini et al., 2021; Liu et al., 2023). A scanning electrical mobility system (SEMS, Brechtel Manufacturing, Inc.), which consisted of the combination of the differential mobility analyzer (DMA) and mixing-based condensation particle counter (MCPC), measured size distribution of particles. Size distributions of particles were recorded every 3 minutes in 60 bins for the diameter range of 10 and 600 nm.

120 In addition, the semi-volatile thermal-desorption aerosol gas chromatography (SV-TAG, Aerodyne Research Inc. & Aerosol Dynamic Inc.) was also used for experiments at room temperature for molecular level chemical analysis (Li et al., 2022; Liu et al., 2023). Standards for major fatty acids in COA, including linoleic, oleic, palmitic, and stearic acids (Table S2) were employed for calibrating the instrument. The SV-TAG was operated in ‘bypass’ mode for sampling both gas- and particle-



125 phase species. The oven temperature of the concentrating trap and flow rate for the gas chromatography were changed for
126 identifying compounds (Fig. S2) and for separating fatty acids of interest for kinetics analysis (Fig. S3). Identification of
127 detected chemical species were conducted using MassHunter (Agilent). An optical aerosol counter (OPC, Model 11-D,
GRIMM Aerosol Technik Ainring) was employed together with the SV-TAG for mass concentration measurements.

3 Results and discussion

3.1 Chemical composition of COA particles

130 Figure 2 shows the gas chromatogram for COA particles from the hot pot soup base. Chromatograms for all types of COA are
131 shown in Fig. S2. The SV-TAG identified more than 500 chemical species in COA particles. The number of identified organic
132 molecular formula was primarily dominated by oxygenated compounds such as carboxylic acids. Nitrogen-containing
133 compounds and hydrocarbons were also detected (i.e., amides and alkanes). In addition, there were some chemical species
134 containing sulfur, phosphorus, or halogen atoms. Examples include benzenesulfonamide, phosphinic acid, and bromofluorene.
135 Peaks of plasticizers (phthalate) were likely originated from Tygon tubings and residuals in zero air (Liu et al., 2023).




Figure 2: Gas chromatogram of hot pot COA particles prior to ozone exposure.

Oleic, linoleic, palmitic, and stearic acids were the four dominant fatty acids, accounting for ~20% of the total particle mass. Fatty acids were likely released from thermal degradation of triglycerides and phospholipids (Abdullahi et al., 2013; Zhao et al., 2015). Figure 3 shows the mass fractions of these four fatty acids in investigated COA. Canola oil particles were rich in unsaturated fatty acids (i.e., linoleic and oleic acids), while COA generated from lard contained the highest fractions of saturated fatty acids (i.e., palmitic and stearic acids). High abundance of linoleic acid in COA particles from hot pot might be originated from chicken oil in the sample (Song et al., 2023). Some other fatty acids with even carbon number, such as dodecanoic, myristic, hexadecenoic, linoelaidic, and arachidic acids were also detected, though abundances of signals for these
145 compounds were minor.





Figure 3: Mass fractions of four major fatty acids in canola oil, hot pot and lard COA particles.

Some other minor organic species were also detected, such as sterols, tocopherols, and polycyclic aromatic hydrocarbons (PAHs). Sterols widely exist in vegetable and animal tissues (Sikorski and Kolakowska, 2010). The SV-TAG detected sterols such as cholesterol, campesterol, sitosterol, stigmasterol and fucosterol. The main dietary source of tocopherol is vegetable oil (Jiang et al., 2001). Distinct signals from β , γ , and δ tocopherols were found in three types of COA particles. Usage of vegetable oil during the extraction process might have contributed to the existence of this compound in lard COA. PAHs such as benzo(a)pyrene, benz(a)anthracene, pyrene and fluoranthene were detected, although their abundances were limited. Capsaicin was non-negligible for only hot pot COA particles, likely originated from species.

Figure 4 shows the ToF-ACSM mass spectra of three types of COA particles. The major peaks for the mass spectra included $m/z = 41, 55, 67$ and 91 , as have been reported in previous COA studies (Mohr et al., 2009; He et al., 2010; Kaltsonoudis et al., 2017). Ratios of f_{55}/f_{57} ($f_{55}/f_{57} = 2.4 \sim 3.2$) were consistent with previous COA measurements (Zhang et al., 2011; Kaltsonoudis et al., 2017). High-molecular weight ions ($m/z > 100$) that have been observed for oleic acid ($m/z = 178, 191$ and 202) were also detected (Hu et al., 2018; Liu et al., 2023). These dominant peaks were most likely hydrocarbon ions, as have been reported by previous COA studies (He et al., 2010; Kaltsonoudis et al., 2017; Hu et al., 2018).

Generally, mass spectra for the three types of COA were similar ($r^2 > 0.95$). Similarities in aerosol mass spectra for COA from various sources have also previously been reported (He et al., 2010). The mass spectrum of hot pot COA particles exhibited relatively large differences from the other two types of COA. Fractions of signals at $m/z = 105, 107, 117$, and 137 were 2~5 times higher in hot pot COA particles.



165

Figure 4: Mass spectra of (a) canola oil Exp. #1; (b) hot pot Exp. #9; (c) lard Exp. #20. COA particles (black bars). Mass spectra of particles following 2.7×10^{-5} atm s of ozone exposure are shown in red dots. The signal fractions of m/z of 101~210 are enhanced by 30, 20 and 40 times as noted in the figure.

170 3.2 Chemical characteristics of oxidized COA particles

Ozonolysis of COA particles changed their chemical compositions (Fig. S3). Especially, abundances of unsaturated fatty acids decreased following oxidation, as C=C double bonds in unsaturated compounds are highly reactive with ozone (Zahardis and Petrucci, 2007). Abundance of azelaic acid, which formed following ozonolysis of oleic acid, increased following oxidation. These changes were ubiquitously observed for all types of COA. There were no significant changes in mass concentration of saturated fatty acids by oxidation.

175

The ToF-ACSM data also demonstrated reductions in high molecular weight ions (i.e., $m/z = 165, 191,$ and 202) following oxidation, likely due to fragmentations of unsaturated acids by ozonolysis. As a result, the contributions of smaller molecular ions such as $m/z = 43$ and 44 increased (Fig. 4). These data were qualitatively in line with the result of the SV-TAG measurements that the oxidation processes induced fragmentation of organic compounds in COA.



180 3.3 Estimation of the pseudo-second order reaction rate constants

The ME-2 analysis was conducted for quantitatively estimating the mass fraction of COA that had been oxidized using the aerosol mass spectra. The details about the ME-2 analysis are provided in Text S1. Briefly, two-factor solutions were obtained for each set of experiment. A factor that corresponded to COA prior to ozone exposure was denoted as ‘fresh COA’, while another one was named as ‘oxidized COA’. The value for the degree of freedom (*a-value*) of the mass spectra of particles prior to ozone exposure in the aerosol flow tube was set to be 0.2. The mass spectra range of $m/z = 41$ to 210 were employed for the analysis. The mass spectra for the high ozone exposure at 25 °C were analyzed together with each experimental dataset to constrain the profiles for reaction products. The resulting factor profiles of two factors for each type of COA are summarized in Fig. S4.

Figure S5 shows the change in the fractional contributions of the ‘fresh COA’ factor (f_{fresh}) following ozone exposure for the lard experiments. The values of f_{fresh} exponentially decreased after COA particles were exposed to ozone. The process was fit by the following equation by assuming the pseudo-second order reaction with ozone:

$$f_{fresh} = f_{fresh_0} \exp(-k_2 P_{O_3} t) \quad (1)$$

The term of f_{fresh_0} indicates the mass fraction of ‘fresh COA’ prior to ozone exposure. The retrieved value of f_{fresh_0} was 1.00 ± 0.02 for all the cases, agreeing well with the expectation that ‘fresh COA’ was the sole component prior to ozone exposure. P_{O_3} corresponds to the partial pressure of ozone. The residence time of COA particles in the reactor is t . $P_{O_3} t$ corresponds to ozone exposure. The value of k_2 corresponds to the pseudo-second order reaction rate constant for the chemical reaction of Fresh COA + O₃ → Oxidized COA. It should be noted that f_{fresh} was occasionally larger than f_{fresh_0} when the chemical reaction was extremely slow due to uncertainties associated with the experiment and data analysis. In these cases, k_2 was forced to be zero in the following analysis.

In the case of the SV-TAG data, the oxidation processes of unsaturated fatty acids (i.e., oleic and linoleic acids) were parameterized by employing a saturated fatty acid (stearic acid) as a tracer (Wang and Yu, 2021).

$$\frac{[f_{unsaturated_fatty_acid}]}{[f_{stearic_acid}]} = \frac{[f_{unsaturated_fatty_acid_0}]}{[f_{stearic_acid_0}]} \exp(-k_2 P_{O_3} t) \quad (2)$$

In the above equation, $f_{unsaturated_fatty_acid_0}$, $f_{unsaturated_fatty_acid}$, $f_{stearic_acid_0}$, and $f_{stearic_acid}$ correspond to abundances of unsaturated fatty acids prior to ozone exposure, unsaturated fatty acids following ozone exposure, stearic acid prior to ozone exposure, stearic acid following ozone exposure.

The obtained values of k_2 for 25 °C are compared in Table 1, along with the corresponding values for oleic acid particles retrieved from our previous study (Liu et al., 2023). The mean diameter of the oleic acid particles was ~400 nm, which was slightly larger than COA particles. Generally, the values of k_2 were in the range of 0.05 ~ 0.09 ppb⁻¹ h⁻¹. The values of k_2 of COA particles obtained by the ME-2 analysis of the ToF-ACSM data and molecular level measurements of oleic acid by the SV-TAG agreed within the difference of 10~40%, suggesting that the time scale for chemical aging of COA and oleic acid agreed with in the range of uncertainty.



Table 1. Comparison of obtained values of k_2 ($\text{ppb}^{-1} \text{h}^{-1}$) for oleic acid in particles from the SV-TAG measurements and COA particles from the ACSM measurements at 25 °C.

| Type | k_2 for oleic acid | k_2 for COA particles |
|-------------------------|----------------------|-------------------------|
| Oleic Acid ^a | 0.067 | 0.078 ± 0.008 |
| Canola Oil | 0.053 | 0.086 ± 0.005 |
| Hot Pot | 0.055 | 0.061 ± 0.003 |
| Lard | 0.049 | 0.056 ± 0.003 |

^a Retrieved from our previous study (Liu et al., 2023)

215 The similarities in the values of k_2 by the ME-2 analysis and SV-TAG were reasonable, considering that unsaturated fatty acids were the major chemical species that were responsible for the oxidation process. Oxidation of linoleic acid is known to be 1.6 times faster than that by oleic acid (Moise and Rudich, 2002; Hearn and Smith, 2004). However, it only occupied a small portion, except for hot pot COA particles. A previous study for reactive uptake coefficient of ozone by canola oil film ($\gamma = 0.6 \times 10^{-3}$) (Li et al., 2020) also reported that the chemical reaction process was comparable with that by oleic acid ($\gamma = 0.8 \times 10^{-3}$)
220 (Thornberry and Abbatt, 2004). Differences in experimental procedures could also account for the discrepancy between the two studies.

3.4 Temperature dependence in k_2

Figure 5 summarizes the obtained values of k_2 plotted against temperature. In general, k_2 decreased for lower temperatures. On the other hand, the values of k_2 for elevated temperature ranges ($T > 30$ °C) were higher than that for room temperature ($T \sim$
225 25 °C). For the temperature range above 25 °C, the retrieved values of k_2 for all types of COA and oleic acid particles were only different by less than 3 times. For -10 °C $< T < 10$ °C, the values of k_2 were different by more than an order of magnitude, depending on COA types. All types of COA were not highly reactive for $T < -15$ °C.




Figure 5: Dependence of k_2 on temperature. Data for oleic acid in laboratory were retrieved from our previous study (Liu et al., 2023). The values of k_2 for Exp. #7, 10 and 21 were unmeasurably slow for our experimental setup. Thus, the corresponding data are shown in open symbols at the bottom.

The change of k_2 with temperature was gradual, rather than abrupt (Fig. 5 and Fig. S6). The gradual reduction in chemical reactions for reduced temperature range has previously been reported for films that experienced glass transitions using coated flow tube experimental setup (Li et al., 2020). The observed relationships between k_2 and temperature exhibited transitions at around -15 °C (canola oil), -5 °C (hot pot), and 10 °C (lard). Below these temperatures, the values of k_2 were not highly sensitive to temperature/unmeasurably small. On the other hand, above these temperature ranges, changes in k_2 with temperature were more pronounced. In the case of chemical systems that experience glass transition, similar changes in γ were also observed using the coated flow tube (Li et al., 2020; Moise and Rudich, 2002). Namely, γ was almost constant for the temperature range below the glass transition temperature, as surface reactions dominated for the region. However, γ increased almost linearly with temperature for the warmer region. A similar transition was also likely occurring to the COA particles. COA particles that contained higher fractions of saturated fatty acids have smaller k_2 for a certain temperature, as well as higher transition temperature. Previous laboratory studies demonstrated that mixing of saturated fatty acids such as lauric, myristic, stearic, and palmitic acids reduced chemical reactivity of oleic acid particles at room temperature (Hearn and Smith, 2005; Katrib et al., 2005; Knopf et al., 2005). We hypothesize that enhanced abundance of saturated fatty acids increased transition temperatures of COA particles, though further studies that directly quantify viscosity or glass transition would still be needed for confirming the idea.

4 Atmospheric implications

Our laboratory experiments in the present study demonstrated that pseudo-second order reaction rate constants k_2 between COA and ozone were highly sensitive to temperature, likely due to changes in viscosity. For estimating the potential impact



250 of temperature on atmospheric chemical lifetimes of COA, the experimentally obtained k_2 - T relationships were phenomenologically fit by equations. The relationships do not follow the Arrhenius equation well (Fig. S7), suggesting that the major factor that influences the k_2 - T relationships is not related to the activation energy of the chemical reaction. In the case of diffusion-controlled chemical reaction processes, the overall reaction rate constant is known to be related to viscosity (η) (Krenos, 2001). The temperature dependence in η can be evaluated by the Vogel–Fulcher–Tammann (VFT) equation;

255
$$\ln \eta = A + \frac{B}{T - T_0} \quad (3)$$

In the above equation, A , B , and T_0 are empirically obtained parameters. By considering the VFT equation, k_2 values were fit by the following equation under the assumption that the oxidation processes of COA were controlled by viscosity change.

$$\ln k_2 = \alpha_1 + \frac{\alpha_2}{T + \alpha_3} \quad (4)$$

In the above equation, α_1 , α_2 , and α_3 are parameters that are empirically optimized. The fitting results for the three types of laboratory generated COA are summarized in Fig. S8. The corresponding optimized parameters are tabulated in Table S3.

260 Figure 6 shows the distributions of estimated atmospheric chemical lifetimes of COA with respect to ozonolysis in June and December. The figure was created by assuming that chemical reactivity of ambient COA was comparable to that from canola oil. Monthly mean surface air temperature data were obtained from the website of the Physical Sciences Laboratory of NOAA (<https://psl.noaa.gov/mddb2/makePlot.html?variableID=1603>) for June and December of 2021 (Fig. S9). Ozone concentration was assumed to be 30 ppb. Qualitatively similar estimations were also obtained for other types of COA (Fig. S10) and ozone concentrations (Fig. S11).






Figure 6: Estimated chemical lifetimes of COA during (a) June and (b) December. The fit function for the experimental data of canola oil COA was employed for the calculation by assuming the ozone concentration of 30 ppb.

270 Temperature was between 10 ~ 30 °C for most areas in June, except for Greenland and the Tibetan plateau. As k_2 only changed by less than twice for the temperature range, the estimated atmospheric chemical lifetimes were shorter than one hour and did not exhibit an extensive variation.

In December, temperatures of the tropical and southern hemisphere regions were high. Temperatures in high-latitude regions in the northern hemisphere were lower than 0 °C, resulting in atmospheric chemical lifetimes of longer than 10 hours. In the
275 case of the mid-latitude regions, COA may survive in the atmosphere for the time scale of 1 ~ 10 hours, consistent with recent observational studies in some cities in China (Wang and Yu, 2021; Li et al., 2023).

The above discussion only considered chemical reactions with ozone under dry conditions, while COA particles were also highly reactive with other atmospheric oxidants such as OH and NO₃ radicals (Li et al., 2020). In addition, the discussion ignored changes in chemical reactivity of COA particles that might occur following initial reactions (Hosny et al., 2016;
280 Berkemeier et al., 2021). Potential influence of hygroscopic growth of partially aged COA on its chemical reactivity was also ignored. Mixing state of COA particles that could influence chemical reactivity at low temperature was also not considered (Liu et al., 2023; Ma et al., 2023).

As atmospheric chemical lifetimes of oleic and linoleic acids are short, online measurement techniques are required for estimating their atmospheric chemical lifetimes from ambient data. So far, such studies have only been conducted in a limited
285 number of cities. Further observational evidence will be needed for evaluating the roles of environmental conditions such as temperature on atmospheric chemical lifetimes of COA. Especially, wintertime observations in cold regions would be important. We suggest that future laboratory and field studies of COA will warrant underpinning its atmospheric fate and environmental impacts.

290 5 Conclusions

We conducted a series of laboratory experiments to investigate the influence of temperature on chemical reactivity of COA. Three types of COA particles were generated by heating canola oil, hot pot soup base, and lard. The produced COA particles were exposed to ozone in an aerosol flow tube reactor under the temperature range of -20~35°C. Changes in chemical composition of COA particles were characterized by the ToF-ACSM and SV-TAG. The experimental results were used for
295 calculating the chemical reaction rate constants k_2 . The observed temperature dependences in k_2 were fit well by the VFT-equation, suggesting that reduced viscosity was responsible for the decrease in chemical reactivity. The parameterized temperature-dependent k_2 were used for estimating the chemical lifetimes of COA particles on the global scale, in combination with observed global surface temperature data. During summertime, atmospheric lifetimes of COA are short (< 1 hour) for most regions, while the corresponding value is longer than 10 hours in high-latitude regions in the northern hemisphere during
300 wintertime. This study demonstrated suppressed chemical reactivity of COA particles under low temperature needs to be considered in simulating COA fate.



Data availability

The data in this study are available from the corresponding author upon request.

Author contributions

305 Conceptualization - MK and WL; Experiment investigation – WL, LH and KL; Data analysis – WL, LH and MK; Resources and Funding acquisition – MK, YL and QC; Writing – original draft preparation – WL and MK; Writing – review & editing – LH, YL, QC.

Competing interests

One of the co-authors is a member of the editorial board of Atmospheric Chemistry and Physics.

310 **Acknowledgments**

We acknowledge Mr. Zhao at Jiayuan Canteen in Peking University for providing the lard sample for the study.

Financial support

The study was funded by the National Natural Science Foundation of China (42175121, 42150610485, and 92044303).



References

- 315 Abdullahi, K. L., Delgado-Saborit, J. M., and Harrison, R. M.: Emissions and indoor concentrations of particulate matter and its specific chemical components from cooking: A review, *Atmos. Environ.*, 71, 260-294, <http://doi.org/10.1016/j.atmosenv.2013.01.061>, 2013.
- Berkemeier, T., Mishra, A., Mattei, C., Huisman, A. J., Krieger, U. K., and Pöschl, U.: Ozonolysis of oleic acid aerosol revisited: multiphase chemical kinetics and reaction mechanisms, *ACS Earth Space Chem.*, 5, 3313-3323, <http://doi.org/10.1021/acsearthspacechem.1c00232>, 2021.
- 320 Broekhuizen, K. E., Thornberry, T., Kumar, P. P., and Abbatt, J. P. D.: Formation of cloud condensation nuclei by oxidative processing: Unsaturated fatty acids, *J. Geophys. Res.: Atmos.*, 109, D24206, <https://doi.org/10.1029/2004JD005298>, 2004.
- Budisulistiorini, S. H., Chen, J., Itoh, M., and Kuwata, M.: Can online aerosol mass spectrometry analysis classify secondary organic aerosol (SOA) and oxidized primary organic aerosol (OPOA)? A case study of laboratory and field studies of Indonesian biomass burning, *ACS Earth Space Chem.*, 5, 3511-3522, <https://doi.org/10.1021/acsearthspacechem.1c00319>, 2021.
- 325 Crippa, M., DeCarlo, P. F., Slowik, J. G., Mohr, C., Heringa, M. F., Chirico, R., Poulain, L., Freutel, F., Sciare, J., Cozic, J., Di Marco, C. F., Elsasser, M., Nicolas, J. B., Marchand, N., Abidi, E., Wiedensohler, A., Drewnick, F., Schneider, J., Borrmann, S., Nemitz, E., Zimmermann, R., Jaffrezo, J. L., Prévôt, A. S. H., and Baltensperger, U.: Wintertime aerosol chemical composition and source apportionment of the organic fraction in the metropolitan area of Paris, *Atmos. Chem. Phys.*, 13, 961-981, <http://doi.org/10.5194/acp-13-961-2013>, 2013.
- 330 Daellenbach, K. R., Stefenelli, G., Bozzetti, C., Vlachou, A., Fermo, P., Gonzalez, R., Piazzalunga, A., Colombi, C., Canonaco, F., Hueglin, C., Kasper-Giebl, A., Jaffrezo, J. L., Bianchi, F., Slowik, J. G., Baltensperger, U., El-Haddad, I., and Prevot, A. S. H.: Long-term chemical analysis and organic aerosol source apportionment at nine sites in central Europe: source identification and uncertainty assessment, *Atmos. Chem. Phys.*, 17, 13265-13282, <http://doi.org/10.5194/acp-17-13265-2017>, 2017.
- 335 Guo, J. C., Zhou, S. Z., Cai, M. F., Zhao, J., Song, W., Zhao, W. X., Hu, W. W., Sun, Y. L., He, Y., Yang, C. Q., Xu, X. Z., Zhang, Z. S., Cheng, P., Fan, Q., Hang, J., Fan, S. J., Wang, X. M., and Wang, X. M.: Characterization of submicron particles by time-of-flight aerosol chemical speciation monitor (ToF-ACSM) during wintertime: aerosol composition, sources, and chemical processes in Guangzhou, China, *Atmos. Chem. Phys.*, 20, 7595-7615, <http://doi.org/10.5194/acp-20-7595-2020>, 2020.
- 340 He, L. Y., Lin, Y., Huang, X. F., Guo, S., Xue, L., Su, Q., Hu, M., Luan, S. J., and Zhang, Y. H.: Characterization of high-resolution aerosol mass spectra of primary organic aerosol emissions from Chinese cooking and biomass burning, *Atmos. Chem. Phys.*, 10, 11535-11543, <http://doi.org/10.5194/acp-10-11535-2010>, 2010.
- 345 Hearn, J. D. and Smith, G. D.: Kinetics and product studies for ozonolysis reactions of organic particles using aerosol CIMS, *J. Phys. Chem. A*, 108, 10019-10029, <https://doi.org/10.1021/jp0404145>, 2004.
- Hearn, J. D. and Smith, G. D.: Measuring rates of reaction in supercooled organic particles with implications for atmospheric aerosol, *Phys. Chem. Chem. Phys.*, 7, 2549-2551, <https://doi.org/10.1039/b506424d>, 2005.
- 350 Hildemann, L. M., Markowski, G. R., and Cass, G. R.: Chemical-composition of emissions from urban sources of fine organic aerosol, *Environ. Sci. Technol.*, 25, 744-759, <http://doi.org/10.1021/es00016a021>, 1991.
- Hosny, N. A., Fitzgerald, C., Vyšniauskas, A., Athanasiadis, A., Berkemeier, T., Uygur, N., Pöschl, U., Shiraiwa, M., Kalberer, M., Pope, F. D., and Kuimova, M. K.: Direct imaging of changes in aerosol particle viscosity upon hydration and chemical aging, *Chem. Sci.*, 7, 1357-1367, <http://dx.doi.org/10.1039/C5SC02959G>, 2016.
- 355 Hu, W., Day, D. A., Campuzano-Jost, P., Nault, B. A., Park, T., Lee, T., Croteau, P., Canagaratna, M. R., Jayne, J. T., Worsnop, D. R., and Jimenez, J. L.: Evaluation of the new capture vaporizer for aerosol mass spectrometers: Characterization of organic aerosol mass spectra, *Aerosol Sci. Technol.*, 52, 725-739, <https://doi.org/10.1080/02786826.2018.1454584>, 2018.
- Huang, D. D., Zhu, S., An, J., Wang, Q., Qiao, L., Zhou, M., He, X., Ma, Y., Sun, Y., Huang, C., Yu, J. Z., and Zhang, Q.: Comparative assessment of cooking emission contributions to urban organic aerosol using online molecular tracers and Aerosol Mass Spectrometry measurements, *Environ. Sci. Technol.*, 55, 14526-14535, <https://doi.org/10.1021/acs.est.1c03280>, 2021.
- 360 Hung, H.-M. and Tang, C.-W.: Effects of temperature and physical state on heterogeneous oxidation of oleic acid droplets with ozone, *J. Phys. Chem. A*, 114, 13104-13112, <https://doi.org/10.1021/jp105042w>, 2010.



- Jiang, Q., Christen, S., Shigenaga, M. K., and Ames, B. N.: γ -Tocopherol, the major form of vitamin E in the US diet, deserves more attention, *Am. J. Clin. Nutr.*, 74, 714-722, <https://doi.org/10.1093/ajcn/74.6.714>, 2001.
- 365 Kaltsonoudis, C., Kostenidou, E., Louvaris, E., Psichoudaki, M., Tsiligiannis, E., Florou, K., Liangou, A., and Pandis, S. N.: Characterization of fresh and aged organic aerosol emissions from meat charbroiling, *Atmos. Chem. Phys.*, 17, 7143-7155, <https://doi.org/10.5194/acp-17-7143-2017>, 2017.
- Katrib, Y., Biskos, G., Buseck, P. R., Davidovits, P., Jayne, J. T., Mochida, M., Wise, M. E., Worsnop, D. R., and Martin, S. T.: Ozonolysis of mixed oleic-acid/stearic-acid particles: reaction kinetics and chemical morphology, *J. Phys. Chem. A*, 109, 10910-10919, <https://doi.org/10.1021/jp054714d>, 2005.
- 370 Kaur Kohli, R., Reynolds, R. S., Wilson, K., and Davies, J. F.: Exploring the influence of particle phase in the ozonolysis of oleic and elaidic acid, *Aerosol Sci. Technol.*, 1-21, <https://doi.org/10.1080/02786826.2023.2226183>, 2023.
- Knopf, D. A., Anthony, L. M., and Bertram, A. K.: Reactive uptake of O₃ by multicomponent and multiphase mixtures containing oleic acid, *J. Phys. Chem. A*, 109, 5579-5589, <https://doi.org/10.1021/jp0512513>, 2005.
- 375 Krenos, J. R.: Chemical kinetics and reaction dynamics, Dover Publications, <https://doi.org/10.1021/ed078p1466>, 2001.
- Laskin, A., Moffet, R. C., and Gilles, M. K.: Chemical imaging of atmospheric particles, *Acc. Chem. Res.*, 52, 3419-3431, <http://doi.org/10.1021/acs.accounts.9b00396>, 2019.
- Lee, B. P., Li, Y. J., Yu, J. Z., Louie, P. K. K., and Chan, C. K.: Characteristics of submicron particulate matter at the urban roadside in downtown Hong Kong—Overview of 4 months of continuous high-resolution aerosol mass spectrometer measurements, *J. Geophys. Res.: Atmos.*, 120, 7040-7058, <https://doi.org/10.1002/2015JD023311>, 2015.
- 380 Li, J., Forrester, S. M., and Knopf, D. A.: Heterogeneous oxidation of amorphous organic aerosol surrogates by O₃, NO₃, and OH at typical tropospheric temperatures, *Atmos. Chem. Phys.*, 20, 6055-6080, <https://doi.org/10.5194/acp-20-6055-2020>, 2020.
- 385 Li, R., Zhang, K., Li, Q., Yang, L., Wang, S., Liu, Z., Zhang, X., Chen, H., Yi, Y., Feng, J., Wang, Q., Huang, L., Wang, W., Wang, Y., Yu, J. Z., and Li, L.: Characteristics and degradation of organic aerosols from cooking sources based on hourly observations of organic molecular markers in urban environments, *Atmos. Chem. Phys.*, 23, 3065-3081, <https://acp.copernicus.org/articles/23/3065/2023/>, 2023.
- Li, Y., He, L., Xie, D., Zhao, A., Wang, L., Kreisberg, N. M., Jayne, J., and Liu, Y.: Strong temperature influence and indiscernible ventilation effect on dynamics of some semivolatile organic compounds in the indoor air of an office, *Environ. Int.*, 165, 107305, <https://www.ncbi.nlm.nih.gov/pubmed/35635961>, 2022.
- 390 Liu, J., Zhang, F., Xu, W., Chen, L., Ren, J., Jiang, S., Sun, Y., and Li, Z.: A large impact of cooking organic aerosol (COA) on particle hygroscopicity and CCN activity in urban atmosphere, *J. Geophys. Res.: Atmos.*, 126, e2020JD033628, <https://doi.org/10.1029/2020jd033628>, 2021.
- Liu, S., Xing, J., Zhao, B., Wang, J., Wang, S., Zhang, X., and Ding, A.: Understanding of aerosol–climate interactions in China: aerosol impacts on solar radiation, temperature, cloud, and precipitation and its changes under future climate and emission scenarios, *Curr. Pollut. Rep.*, 5, 36-51, <https://doi.org/10.1007/s40726-019-00107-6>, 2019.
- 395 Liu, W., Liao, K., Chen, Q., He, L., Liu, Y., and Kuwata, M.: Existence of crystalline ammonium sulfate nuclei affects chemical reactivity of oleic acid particles through heterogeneous nucleation, *J. Geophys. Res.: Atmos.*, 128, e2023JD038675, <https://doi.org/10.1029/2023JD038675>, 2023.
- 400 Liu, X., Gu, J., Li, Y., Cheng, Y., Qu, Y., Han, T., Wang, J., Tian, H., Chen, J., and Zhang, Y.: Increase of aerosol scattering by hygroscopic growth: Observation, modeling, and implications on visibility, *Atmos. Res.*, 132-133, 91-101, <https://doi.org/10.1016/j.atmosres.2013.04.007>, 2013.
- Ma, T., Furutani, H., Duan, F., Ma, Y., Toyoda, M., Kimoto, T., Huang, T., and He, K.: Real-time Single-Particle characteristics and aging of cooking aerosols in urban Beijing, *Environ. Sci. Technol. Lett.*, 10, 404-409, <https://doi.org/10.1021/acs.estlett.3c00053>, 2023.
- 405 Menon, S., Unger, N., Koch, D., Francis, J., Garrett, T., Sednev, I., Shindell, D., and Streets, D.: Aerosol climate effects and air quality impacts from 1980 to 2030, *Environ. Res. Lett.*, 3, 024004, <http://doi.org/10.1088/1748-9326/3/2/024004>, 2008.
- Mohr, C., Huffman, J. A., Cubison, M. J., Aiken, A. C., Docherty, K. S., Kimmel, J. R., Ulbrich, I. M., Hannigan, M., and Jimenez, J. L.: Characterization of primary organic aerosol emissions from meat cooking, trash burning, and motor vehicles with high-resolution aerosol mass spectrometry and comparison with ambient and chamber observations, *Environ. Sci. Technol.*, 43, 2443-2449, <https://doi.org/10.1021/es8011518>, 2009.



- Moise, T. and Rudich, Y.: Reactive uptake of ozone by aerosol-associated unsaturated fatty acids: kinetics, mechanism, and products, *J. Phys. Chem. A*, 106, 6469-6476, <https://doi.org/10.1021/jp025597e>, 2002.
- 415 Morris, J. W., Davidovits, P., Jayne, J. T., Jimenez, J. L., Shi, Q., Kolb, C. E., Worsnop, D. R., Barney, W. S., and Cass, G.: Kinetics of submicron oleic acid aerosols with ozone: A novel aerosol mass spectrometric technique, *Geophys. Res. Lett.*, 29, 71-71-71-74, <https://doi.org/10.1029/2002gl014692>, 2002.
- Reyes-Villegas, E., Bannan, T., Le Breton, M., Mehra, A., Priestley, M., Percival, C., Coe, H., and Allan, J. D.: Online chemical characterization of food-cooking organic aerosols: implications for source apportionment, *Environ. Sci. Technol.*, 52, 5308-5318, <https://doi.org/10.1021/acs.est.7b06278>, 2018.
- 420 Robinson, A. L., Donahue, N. M., and Rogge, W. F.: Photochemical oxidation and changes in molecular composition of organic aerosol in the regional context, *J. Geophys. Res.: Atmos.*, 111, D03302, <https://doi.org/10.1029/2005JD006265>, 2006.
- Robinson, E. S., Gu, P., Ye, Q., Li, H. Z., Shah, R. U., Apte, J. S., Robinson, A. L., and Presto, A. A.: Restaurant impacts on outdoor air quality: elevated organic aerosol mass from restaurant cooking with neighborhood-scale plume extents, *Environ. Sci. Technol.*, 52, 9285-9294, <https://doi.org/10.1021/acs.est.8b02654>, 2018.
- 425 Rogge, W. F., Hildemann, L. M., Mazurek, M. A., Cass, G. R., and Simonelt, B. R. T.: Sources of fine organic aerosol .1. charbroilers and meat cooking operations, *Environ. Sci. Technol.*, 25, 1112-1125, <https://doi.org/10.1021/es00018a015>, 1991.
- Rudich, Y., Donahue, N. M., and Mentel, T. F.: Aging of organic aerosol: Bridging the gap between laboratory and field studies, *Annu. Rev. Phys. Chem.*, 58, 321-352, <https://doi.org/10.1146/annurev.physchem.58.032806.104432>, 2007.
- 430 Shilling, J. E., King, S. M., Mochida, M., Worsnop, D. R., and Martin, S. T.: Mass spectral evidence that small changes in composition caused by oxidative aging processes alter aerosol CCN properties, *J. Phys. Chem. A*, 111, 3358-3368, <https://doi.org/10.1021/jp068822r>, 2007.
- Sikorski, Z. Z. E. and Kolakowska, A.: *Chemical, Biological, and Functional Aspects of Food Lipids*, CRC Press 2010.
- Song, K., Guo, S., Gong, Y., Lv, D., Wan, Z., Zhang, Y., Fu, Z., Hu, K., and Lu, S.: Non-target scanning of organics from cooking emissions using comprehensive two-dimensional gas chromatography-mass spectrometer (GC×GC-MS), *Appl. Geochemistry*, 151, <https://doi.org/10.1016/j.apgeochem.2023.105601>, 2023.
- 435 Sun, Y. L., Du, W., Fu, P. Q., Wang, Q. Q., Li, J., Ge, X. L., Zhang, Q., Zhu, C. M., Ren, L. J., Xu, W. Q., Zhao, J., Han, T. T., Worsnop, D. R., and Wang, Z. F.: Primary and secondary aerosols in Beijing in winter: sources, variations and processes, *Atmos. Chem. Phys.*, 16, 8309-8329, <https://doi.org/10.5194/acp-16-8309-2016>, 2016.
- Thornberry, T. and Abbatt, J. P. D.: Heterogeneous reaction of ozone with liquid unsaturated fatty acids: detailed kinetics and gas-phase product studies, *Phys. Chem. Chem. Phys.*, 6, 84-93, <https://doi.org/10.1039/b310149e>, 2004.
- 440 Wang, Q. and Yu, J. Z.: Ambient measurements of heterogeneous ozone oxidation rates of oleic, elaidic, and linoleic acid using a relative rate constant approach in an urban environment, *Geophys. Res. Lett.*, 48, e2021GL095130, <https://doi.org/10.1029/2021gl095130>, 2021.
- Wang, S., Takhar, M., Zhao, Y., Al Rashdi, L. N. S., and Chan, A. W. H.: Dynamic oxidative potential of organic aerosol from heated cooking oil, *ACS Earth Space Chem.*, 5, 1150-1162, <https://doi.org/10.1021/acsearthspacechem.1c00038>, 2021.
- 445 Weitkamp, E. A., Hartz, K. E., Sage, A. M., Donahue, N. M., and Robinson, A. L.: Laboratory measurements of the heterogeneous oxidation of condensed-phase organic molecular markers for meat cooking emissions, *Environ. Sci. Technol.*, 42, 5177-5182, <https://doi.org/10.1021/es800181b>, 2008.
- Xu, S., Mahrt, F., Gregson, F. K. A., and Bertram, A. K.: Possible effects of ozone chemistry on the phase behavior of skin oil and cooking oil films and particles indoors, *ACS Earth Space Chem.*, 6, 1836-1845, <http://doi.org/10.1021/acsearthspacechem.2c00092>, 2022.
- 450 Zahardis, J. and Petrucci, G. A.: The oleic acid-ozone heterogeneous reaction system: products, kinetics, secondary chemistry, and atmospheric implications of a model system - a review, *Atmos. Chem. Phys.*, 7, 1237-1274, <http://doi.org/10.5194/acp-7-1237-2007>, 2007.
- 455 Zhang, Q., Jimenez, J. L., Canagaratna, M. R., Ulbrich, I. M., Ng, N. L., Worsnop, D. R., and Sun, Y.: Understanding atmospheric organic aerosols via factor analysis of aerosol mass spectrometry: a review, *Anal. Bioanal. Chem.*, 401, 3045-3067, <https://doi.org/10.1007/s00216-011-5355-y>, 2011.
- Zhao, X. Y., Hu, Q. H., Wang, X. M., Ding, X., He, Q. F., Zhang, Z., Shen, R. Q., Lü, S. J., Liu, T. Y., Fu, X. X., and Chen, L. G.: Composition profiles of organic aerosols from Chinese residential cooking: case study in urban Guangzhou, south China, *J Atmos Chem*, 72, 1-18, <Go to ISI>://WOS:000352975400001, 2015.
- 460



- Zhao, Y. L., Hu, M., Slanina, S., and Zhang, Y. H.: Chemical compositions of fine particulate organic matter emitted from Chinese cooking, *Environ. Sci. Technol.*, 41, 99-105, <https://doi.org/10.1021/es0614518>, 2007.
- 465 Zheng, Y., Cheng, X., Liao, K., Li, Y., Li, Y. J., Huang, R.-J., Hu, W., Liu, Y., Zhu, T., Chen, S., Zeng, L., Worsnop, D. R., and Chen, Q.: Characterization of anthropogenic organic aerosols by TOF-ACSM with the new capture vaporizer, *Atmos. Meas. Tech.*, 13, 2457-2472, <https://doi.org/10.5194/amt-13-2457-2020>, 2020.
- Zheng, Y., Miao, R., Zhang, Q., Li, Y., Cheng, X., Liao, K., Koenig, T. K., Ge, Y., Tang, L., Shang, D., Hu, M., Chen, S., and Chen, Q.: Secondary Formation of Submicron and Supermicron Organic and Inorganic Aerosols in a Highly Polluted Urban Area, *J. Geophys. Res.: Atmos.*, 128, e2022JD037865, <https://doi.org/10.1029/2022JD037865>, 2023.